















SWEET'S HORTUS BRITANNICUS;

OR,

A CATALOGUE OF PLANTS,

INDIGENOUS, OR CULTIVATED IN THE GARDENS

OF

GREAT BRITAIN;

ARRANGED ACCORDING TO THEIR

NATURAL ORDERS,

WITH

REFERENCES TO THE LINNEAN CLASS AND ORDER TO WHICH EACH GENUS BELONGS;

THE WHOLE BROUGHT UP TO THE PRESENT TIME,

AND CONTAINS

Above 34,000 Plants;

By far the greatest Number ever before published in any Garden Catalogue of this or any other Country; with their Generic and Specific Names, Habitats, when introduced to this Country, Times of Flowering, Colours of the Flowers, Accentuations; and numerous other Arrangements, for the first Time introduced, with all the most useful Synonyms.

SECOND EDITION.

241891

BY ROBERT SWEET, F.L.S.

Author of Hortus Suburbanus Londinensis; Botanical Cultivator; Geraniaceæ; Cistineæ; British Flower Garden; Flora Australasica; Florist's Guide; the British Warblers, &c.

LONDON:

JAMES RIDGWAY, 169, PICCADILLY.

1830.

feel every instant around and about me. I am not at all
surprised to see you so gratified by your visit. I hope to have
them again - but as you can only have one day.

and after many months of indecision he came to the conclusion that he must go back to his old business and start a new one. He was a very poor businessman and he had to work hard to earn his living. He had to give up his old business and start a new one.

TILLING, PRINTER, GROSVENOR ROW, CHELSEA.

THE rapid sale of the last edition, at once shows the preference given to the superiority of the Natural Arrangement, over that of the Linnean, for a Garden Catalogue, as it brings together all the plants that are nearest related to each other, and which have a general similarity by which any person, who already has a tolerable knowledge of plants, may give a good guess at first sight to what family, or Order, any plant or specimen they possess may belong to, or is related ; this gives a pleasing idea, that is not to be met with by the Linnean Botanist, as he only looks at the Stamens and Styles to see what Class and Order it belongs to, without ever thinking to what else it is allied ; but we still consider the addition of the Linnean Classes and Orders, of great use, as they are so readily attained by the young Botanist, though he should always keep the natural arrangement in view.

The pains we have taken with the present edition may be readily supposed, by the length of time it has taken us to get through it ; and had it not been for the kindness and liberality of A. B. Lambert, Esq. who always admitted us to his Library with pleasure, we should certainly not have been able to have made it so complete ; he is also so kind as to give us information of new plants or importations that he hears or knows of, for which we now return him our sincere thanks.

Mr. David Don, Librarian of the Linnean Society, we feel much obliged for his readiness in procuring us any books or specimens we wish to examine, and also inform us of any new publications that we may not have heard of.

The Linnean collection of specimens being now in the Linnean Society's collection, has enabled us to clear up many doubts and mistakes, by examining Linnaeus's original specimens, several of which we intend setting to rights by publishing figures of them in our British Flower Garden, many of the very common plants having been confused with others, or were considered in a doubtful state.

To Robert Barclay, Esq. of Bury Hill, we also feel greatly obliged for his kindness in sending us specimens of the numerous new plants that are continually flowering in his delightful garden, where rarities are without end ; and we must not forget his gardener, Mr. D. Cameron, to whose merit may be attached the highly cultivated state of the plants, and to whom we are indebted for the names of many that have not yet flowered, and that we should not have otherwise known were in the country.

We have also to return our thanks to A. H. Haworth, Esq. for his liberality in presenting us with all his papers as soon as published, of the new and rare succulent plants and bulbs, published in Taylor's Philosophical Magazine.

To the Honourable and Reverend William Herbert we are indebted for the arrangement of the genus *Crinum*, and for the names of the hybrid species of *Amaryllideæ* raised by him, and those that he possesses, which have been raised in other collections, and the references by whom, or at what place they were raised, which is added in the column of Native, and the two parents from which they originated is given, where known: the whole of which are marked with W. H.; he has also been so kind as to furnish us with a list of many other new plants, that have flowered with him, and that had not before been known in our collections.

We also return our thanks to Mr. G. Charlwood, for his readiness in lending us any Botanical publications that we did not ourselves possess, his collection consisting of many rare and valuable botanical works.

To Mr. William Anderson, the worthy Curator of the Apothecaries' Company's Garden, at Chelsea, we are also much indebted, for his kindness in allowing us to copy his catalogues of newly introduced plants, and for his observations on their habit and culture; and to numerous other Gardeners and Nurserymen we return our best thanks, for their kindness and attention to forward our researches.

We flatter ourselves that our Subscribers will be well satisfied with the present edition, as it contains all the new arrangements and improvements up to the present time, that no other Catalogue, however new, can boast of; we have also been careful what we have inserted, not to put in thousands of plants that were never in the country, nor likely ever to be, a plan sometimes resorted to, to swell the size of the book, and to show a great number of names, but we believe there would be a difficulty to find the plants, should they be sought for. As for its being entirely free from errors, we will not assert, as we have no doubt but some may be found, but none, we believe, of much consequence. We now conclude, with our sincerest thanks to our numerous Subscribers, and hope we shall continue to merit their favour and approbation, as we shall always do our utmost endeavours to give them satisfaction.

REFERENCE TO THE BOOKS,

IN WHICH THE FIGURES REFERRED TO, ARE PUBLISHED.

- Abb. ins.* The Natural History of the rarer Lepidopterous Insects of Georgia, collected from the observations of J. Abbott, by J. E. Smith. 1797. 4to.
- Act. Stock.* Kongl. Vetenskaps Academiens Handlingar. Stockholm, 1739. 8vo.
- A.D.C.* Decandolle (Alphonso.) Monographie des Campanulees. Paris. 1830.
- All. auct.* Allioni (Carolus.) Auctuarium ad floram Pedemontanam. 4to. Taurini. 1789.
- All. m. taur.* Car. Allionii Miscellanea Philosophico-mathematica Societatis privatæ Taurinensis.
- All. ped.* Car. Allionii Flora Pedemontana.
- Alp. Ægypt.* Alpinus (Prosper.) De Plantis Ægypti liber. 1 vol. 4to.
- Alp. ex.* Alpinus (Prosper.) De Plantis Exoticis, libri duo. 4to. 1629.
- Amm. ruth.* Amman (Johan.) Stirpium rario- rum in imperio Rutheno sponte prove- nientium Icones et Descriptiones. 4to. Petropoli. 1739.
- Amoen. ac.* Caroli Linnæi Amoenitates aca- demicæ. 8vo.
- And. bot. rep.* Andrews (Henry.) The Botanist's repository for new and rare plants. 4to. 1797. et seq.
- Ana. geran.* Andrews (Henry.) Geraniums, or a monograph of the genus Geranium. 1805. seq. 4to.
- And. heaths.* Andrews (Henry.) Coloured engravings of Heaths, with botanical de- scriptions. fol. and 8vo. 1802, 1809, et seq.
- Ann. bot.* Annals of Botany, by C. Konig and J. Sims. 1805. 8vo.
- Ann. mus.* Annales du Museum d'Histoire naturelle, par les Professeurs de cet établissemens. 1802. seq. 4to.
- Ap. to B. Field's trav.* Appendix to Baron Field's Travels. 4to.
- Ard. mem.* Memorie di osservazioni, e di sperienze sopra la coltura, e gli usi di varie piante, da P. Arduino. Padova. 1766. 4to.
- Ard. sag. pad.* Arduino (Pietro.) Saggi di Padova.
- Ard. spec.* Arduino (Pietro.) Animadversio- num botanicarum Specimen. 4to. Patavii. 1759.
- Asso. arrag.* De Asso (Ignatius.) Synopsis Stirpium indigenarum Arragoniæ. 4to. Marsiliæ. 1779.
- Aub. Thou. obs.* Aubert Du Petit Thouars Plantes des îles de l'Afrique australie. 4to. 1804.
- Aub. gui.* Histoire des plantes de la Guiane Francoise, par M. Fusée Aublet. 1775. 4to.
- Balb. act. taur.* } Miscellanea Botanica. 4to.
Balb. misc. taur. } ex Actis Academæ Tau-
rinensis exserta. 1804.
- Balb. h. taur.* Horti Academicæ Taurinensis stirpium minus cognitarum aut forte no-
varum icones et descriptiones. Taurini. 1811.
- Banks ic. Kæmp.* Banks (Josephus.) icones selectæ Plantarum quas in Japonia col-
legit et delineavit E. Kæmpfer. fol. 1791.
- Banks. rel. Hous.* Reliquia Houstounianæ. 4to. 1781.
- Barr. ic.* } Plantæ per Galliam, Hispaniam,
Barr. rar. } et Italiam, observatae, Iconibus
æneis exhibitæ a Jac. Barreliero. 1714. fol.
- Bart. med. bot.* Barton (Benj. Smith.) Collec-
tions for an essay towards a materia me-
dica of the United States. 1 fasc. in 8vo.
Philadelphia. 1798.
- Bartram* (William.) Travels through North and South Ca-
Bartr. trav. } rolinia, Georgia, and Florida. 8vo. Philadelphia. 1791.
- Bauer. n. hol.* Bauer (Ferdinand.) Illustra-
tiones floræ novæ Hollandiæ. fol. 1813.
- Bauh. hist.* Bauhin (Johannes.) Historia Plantarum universalis. fol. 1651.
- Bauh. prodr.* Caspar Bauhini προδρόμος Thea-
tri Botanici. Basileæ. 1671. 4to.
- Beau. fl. ou.* Palisot de Beauvois flore des royaumes d'Oware et de Benin. fol. 1805.
- Bell. act. taur.* Bellardi (Ludovico.) Memoires de l'Academie Royale des Sciences de Turin. 4to. 1782.
- Berg. act. holm.* Bergius (Pehr. Jonas.) Kongl. Vetenskaps Academien Handlingar. 8vo. 1739.
- Berg. cap.* Petr. Jon. Bergii descriptiones Plantarum ex Capite bonæ spei. 1769. 8vo.
- Berg. phyt.* Bergeret Phytonomatotechnie universelle. fol.
- Berr. in As. res.* Berry in Asiatic Researches, or Transactions of the Society instituted in Bengal. 4to. 1788.
- Besl. eyst.* Besler (Basilus.) Hortus Eystet- tensis. fol. 1612.
- Bieb. cent.* Bieberstein (Marshall von.) Centuria plantarum rariorū Rossiae meridionalis. Charkoviae.
- Big. med. bot.* Bigelow (Jac.) American Me-
dical Botany. 4to. 1817.
- Biv. Bern. st. rar.* Bivona Bernardi. Stir-
pium rariorum minusque cognitarum in Sicilia provenientium Descriptiones. 4to. 1813.
- Biv. cent. pl.* Bivona Bernardi (Antonin.) Sicularum Plantarum Centuria prima. 8vo. 1806.

- Blackw.** Blackwell (Elizabeth.) A curious Herbal, containing 500 Cuts of the useful Plants. fol. 1737.
- Blum. fl. jav.** Blume (C. L.) Flora Java, &c. fol. 1829.
- Boc. Mus.** Boccone (Paolo.) Museo di Piante rare della Sicilia, Malta, Corsica, Italia, Piemonte e Germania. 4to. 1697.
- Boc. sic. rar.** Boccone (Paolo.) Icones et Descriptiones rariorū Plantarū Siciliæ, Melitæ, Galiae et Italiæ. 4to. 1674.
- Boer. lugd.** Herm. Boerhaave Index alter Plantarum quæ in horto Academiæ Lugduno Batavæ aluntur. 4to. 1720.
- Boiss. fl. eur.** De Boissieu (C. V.) Flore d'Europe. 8vo. 1805.
- Bolt. fil.** Bolton (James.) Filices Britannicæ. 4to. 1785.
- Bonp. mon.** Bonpland (Aimé.) Monographie des Melastomes et autres Plantes de cet ordre. fol. 1809.
- Bonp. nav.** Bonpland (Aimé.) Description des Plantes rares cultivées à Malmaison et à Navarre. fol. 1813.
- Bory. {
Bory. it. } de St. Vincent (J. B. M.) A Voyage en Espagne a travers les royaumes de Galice, Leon, etc. 8vo.
- Bot. mag.** Curtis's Botanical Magazine, continued by J. Sims, J. Bellenden Ker, and W. J. Hooker. 8vo.
- Bot. reg.** Edwards (Sydenham.) The Botanical Register descr. by J. Bellenden Ker. 8vo. 1815. continued by J. Lindley.
- Bot. rep.** Andrews (Henry.) The Botanist's Repository for new and rare Plants. 4to. 1797.
- Braam ic. chin.** Braam's Icones Chinensis.
- Bradl. Suc.** Bradley (Richard.) Historia Plantarum succulentarum. 4to. 1716.
- Braun. Salz.** De Braune (Franz. Ant.) Salzburgische flora. 8vo. 1791.
- Breyn. cent.** Breynius (Jacobus.) Exoticarum Plantarum centuria. fol. 1678.
- Breyn. ic.** Breynii (J.) Icones rariorū et exoticarum Plantarum. 4to. 1739.
- Brong. rham.** Brongniart (Adolph.) Dissertationes des Rhamneos. 4to. Paris.
- Brot. phyt.** Brotero (Felix Avellar.) Phytographia Lusitanicæ selectior. fol. 1801.
- Brown. gen. rem.** Brown (Robert.) General Remarks, Geographical and Systematical, on the Botany of Terra Australis. 4to. 1814.
- Br. Jam.** Browne (Patrick.) The Civil and Natural History of Jamaica. fol. 1756.
- Bruses. tr.** Bruce (James.) Travels to discover the source of the Nile. 4to. 1790.
- Buchoz. ic. col.** Buchoz (Pierre Joseph.) Collection des fleures qui se cultivent tant dans les Jardins de la chine que dans ceux d'Europe. fol.
- Bull. herb.** Bulliard. Herbier de la France. fol. 1780.
- Burm. afr.** Burmann (Johannes.) Rariorum Africanarum Plantarum. 4to.
- Burm. dec.** {
} 1738.
- Burm. Ind.** Nic. Laur. Burmanni Flora India. 4to. 1768.
- Burm. zeyl.** Burmann (Johannes.) Thesaurus Zeylanicus. 4to. 1737.
- Buxb. cent.** Buxbaum (Joh. Christ.) Plantarum minus cognitarum Centuria quinque. 4to. 1728.
- Camer. hort.** Camerarius (Alexander.) Hortus Medicus et Philosophicus. 4to. 1588.
- Cates. car.** Catesby (Mark.) The Natural History of Carolina & Florida. fol. 1741.
- Cav. diss.** Cavanilles (Ant. Jos.) Monadelphiæ classis Dissertationes decem. 4to. 1785.
- Cav. ic.** Cavanilles (Ant. Jos.) Icones et Descriptiones Plantarum quæ aut sponte in Hispania crescent aut in hortis hospitantur. fol. 1791.
- Chabré. sci.** Chabrey (Dominicus.) Omnium Stirpium Sciagraphia. fol. 1677.
- Choisy. p. hyp.** {
} Choisy (J. D.) Prodromus d'une monographie de la famille des Hypericinées. 4to. 1821.
- Chois. selag.** Choisy (J. D.) Memoire sur la famille des Selaginees. 4to. Geneva. 1823.
- Clus. hist.** Clusius seu l'Ecluse (Charles de.) Rariorum Plantarum Historia. fol. 1601.
- Col. ephr.** Columna (Fabius.) Minus cognitarum stirpium Ephrasis. 4to. 1616.
- Col. phyt.** Columna (Fabius.) Phytobasanos. 4to. 1592.
- Col. h. rip.** Colla Hortus Ripulensis. 4to.
- Com. hort.** Commelyn (Casp.) Horti Medici Amstelodamensis rariorum Plantarum descriptiones et icones. fol. 1703.
- Comm. petrop.** Commelyn (Casp.) Commentarii Academiacæ Scientiarum imperialis Petropolitanae. 4to. 1728.
- Comm. prael.** Commelyn (Casp.) Praeludia Botanica ad publicas Plantarum demonstrationes. 4to. 1703.
- Comm. rar.** Commelyn (Casp.) Horti Medici Amstelodamensis Plantæ rariores et exoticæ. 4to. 1706.
- Cook. et.** Cook (J.) Voyages, 2 vols. 4to. 1777.
- Corn. can.** Cornuti (Jacob.) Canadensium Plantarum aliarumque nondum editarum historia. 4to. 1635.
- Crantz. Aust.** Crantz (Hen. Joh. Nepom.) Stirpium austriacarum. 8vo. 1762.
- Cup. pan.** Cupani (Franciscus.) Catalogus plantarum Sicularum noviter detectarum. Panormi. 1652.
- Curt. lond.** Curtis (William.) A catalogue of the plants growing wild in the environs of London. fol. 1774. et seq.
- Cyril. ic. fasc.** {
} Cyrilii (Dom.) Plantarum rariorū regni Neapolitani
- Cyr. neap.** {
} fasciculus. 1778. fol.
- Dalech. lug.** Dalechamps (Jacques.) Historia generalis plantarum. fol. 1586.
- Dard. trait. ren.** D'ardenne (Jean Paul Rome.) Traite des Renoncules. 8vo. 1746.
- DC. an. mus.** De Candolle (Augustin Pyramus.) Annales du Museum d'histoire naturelle. 4to. 1802.

- DC. astr.** De Candolle (Augustin Pyramus.) *Astragalologia.* 4to. et fol. 1802.
- DC. dis. cact.** De Candolle (Aug. Pyr.) Revue de la famille des Cactees. 4to. Paris. 1829.
- DC. h. mons.** De Candolle (Augustin Pyramus.) Catalogus plantarum horti botanici Monspeliensis addito observationum circa species novas aut non satis cognitas fasciculo. 8vo. 1813.
- DC. ic. gall. rar.** De Candolle (Augustin Pyramus.) Icones plantarum Galliae rariores. 4to. 1808.
- DC. leg. mem.** De Candolle (Augustin Pyramus.) Memoires sur la famille des Leguminosae. 4to. 1826.
- DC. mem. mus.** De Candolle (Augustin Pyramus.) Memoires du Museum d'histoire naturelle. 4to. 1815.
- DC. mem. umb.** De Candolle (Aug. Pyr.) Memoire sur la famille des Ombellifères. 4to. Paris. 1829.
- DC. plant. grass.** De Candolle (Augustin Pyramus.) Plantarum historia succulentarum. Fasc. 28. 1799.
- Debr. flor. nov.** De Bry (Joannes Theodor.) *Florilegium novum.* fol. 1612.
- Delile.** (Aline Rafeneau.) { Moires botaniques extraits de la Description de l'Egypte. fol. 1813.
- De la Roch. dis.** De La Roche (Daniel.) Specimens botanicum inaugurae sistens descriptiones plantarum aliquot novarum. 4to. 1766.
- Deless. ic. sel.** Delessert (Benj.) Icônes selectæ plantarum in system. univers. descriptarum. 4to. 1820.
- Descrip. de Egyp.** Description de l'Egypte ou Recueil des observations et des recherches faites pendant l'expédition de l'armée Francaise. 4to. et fol. 1810.
- Desf. act. par.** Desfontaines (Rene Louiche.) Mémoires de l'Académie Royale des Sciences. 4to. 1666.
- Desf. atl.** Desfontaines (Rene Louiche.) Flora atlantica. 4to. 1798.
- Desv. j. bot.** Desvaux. Journal de Botanique. 8vo. 1808.
- D. et H.** Dreves et Hayne. Choix de Plantes de l'Europe. fasc. 4. 1802.
- Dill. elt.** Dillenius (Joh. Jac.) Hortus elthamensis. fol. 1732.
- Dill. musc.** Dillenius (Joh. Jac.) Historia muscorum. 4to. 1741.
- Dodart mem.** Dodart (Denys.) Mémoires pour servir à l'Histoire des Plantes. fol. 1676.
- Dod. pemp.** Dodart (Denys.) Stirpium historiæ pemptades. fol. 1583.
- Du Fresn. Val.** Dufresne (Pierre.) Histoire naturelle et medicale de la famille des Valerianées. 4to. 1811.
- Duh. arb.** Duhamel du Monceau (Henry Louis.) Traité des arbres et arbustes qui cultivent en France en pleine terre. 4to. 1755.
- Duham. ed. nov.** Duhamel du Monceau (Henry Louis.) Traites des arbres et arbustes qui se cultivent en pleine terre en France. fol. 1801.
- Dun. mon.** Dunal (Michel Felix.) Monographie de la Famille des Anonacees. 4to. 1817.
- Dunal. solan.** Dunal (Michel Felix.) Histoire naturelle medicale et économique des Solanum et des genres qui ont été confondus avec eux. 4to. 1813.
- Ehret pict.** Ehret (Georg. Dion.) Planta et Papilionæ rariores. fol. 1748.
- Eng. bot.** English Botany. By J. E. Smith; the Figures by J. Sowerby. 8vo.
- Exot. bot.** Exotic Botany. By J. E. Smith; the Figures by J. Sowerby. 8vo.
- Ferr. hesp.** Ferrari (Joh. Bapt.) Hesperides, sive de malorum aureorum cultura et usu. fol. 1646.
- Feuillee.** (Louis.) Journal des observations Physiques, Mathématiques et Botaniques faites dans l'Amérique méridionale. 4to. 1814.
- Fl. dan.** Icônes plantarum sponte nascentium in regnis Daniæ et Norvegiae, etc. fol. 1761.
- Fl. lap.** Linnaeus s. von Linne (Carolus.) Flora Lapponica. 8vo. 1737.
- Flor. græc.** Flora Græca J. E. Smith. fol. 1552.
- Flor. mex. ic. ined.** Sesse et Mocinno. Flora Mexicana Icônes ineditæ.
- Fl. per.** Ruiz (Hippol.) et Pavon (Jos.) Flora Peruviana et Chilensis. 8vo. 1798.
- Flugg. an. mus.** Flugge (Johan.) Annales du Muséum d'Histoire naturelle. 4to.
- Fraser mon.** Fraser (John.) A short history of the Agrostis cornucopiae. fol. 1784.
- Forsk. egypt.** (Forskaol (Petrus.) Flora Ægyptiaca. 4to. 1775.
- Forsk. dis.** (Forskaol (Petrus.) Flora Ægyptiaca arabica. 4to. 1775.
- Forst. gen.** Forster (George.) Characteres generum Plantarum quas in itinere ad insulas maris australis, etc. 4to. 1776.
- Fourc. j. bot.** Fourcroy (Ant. Franc.) La Médecine éclairée par les Sciences physiques, ou Journal des Découvertes, etc. 8vo. 1791.
- Froel. gent.** Froelich (Jos. Aloys.) De Gentiana libellus. 8vo. 1796.
- Fuchs. hist.** Fuchs (Leonhard.) De historia Stirpium commentarii insignes. fol. 1542.
- Gaert. carp.** Gaertner (Car. Frid.) Supplementum Carpologiæ. 4to. 1805.
- Gaert. fr.** (Gaertner (Josephus.) De fructibus et seminibus plantarum. 4to. 1788.
- Gaert. sem.** (Gaertner (Josephus.) De fructibus et seminibus plantarum. 4to. 1788.
- Garidel.** (Pierre Joseph.) Histoire des Plantes qui naissent en Province et principalement aux Environs d'Aix. fol. 1715.
- Gay dis. las.** Gay dissertation Lasiopetalæ.
- Gerard. gal.** (Gerard (John.) Flora Galloprovincialis. 8vo. 1761.
- Ger. emac.** Gerard (John.) The Herbal, enlarged by Th. Johnson. fol. 1633.
- Ger. herb.** Gerard (John.) The Herbal, or general history of Plants. fol. 1797.
- Gilib. h. pl.** Gilibert (Joh. Em.) Histoire des plantes d'Europe, ou éléments de botanique pratique. 3 vol. in 8vo. Lyon. 1806.
- Giorn. pis.** Giornale Pisano. 8vo. 1801.
- Giseck. ic.** Giseke (Paul Dieter.) et Schulze (J. Dom.) Icônes plantarum. 1777.

- Gmel. fl. bad.* Gmelin (Carol. Christ.) Flora Badensis-Alsatica. 8vo. 1808.
- Gmel. sib.* Gmelin (Joh. Georg.) Flora sibirica. 4to. 1747.
- Gou. fl. mons.* Gouan (Anton.) Flora Mons-peliacæ. 8vo. 1765.
- Gouan. ill.* Gouan (Anton.) Illustrationes botanicæ. fol. 1773.
- Hacq. pl. alp. car.* Hacquet (Balth.) Plantæ alpinæ Carniolicae. 4to. 1782.
- Hall. all.* Haller (Albert.) Allii genus naturale constitutum. 4to. 1745.
- Hall. helv.* Haller (Albert.) Historia Stirpium indigenarum Helvetiae. fol. 1768.
- Hayn. term. bot.* Hayne (Fried. Gottl.) et Willdenow (Carl. Ludv.) Terminii botanici iconibus illustrati. 4to. 1799.
- Hayne in ust. an.* Hayne (Fried. Gottl.) in Usteri Annalen der Botanik. 8vo. 1791.
- H. et B. n. gen.* Humboldt, Bonpland et Kunth Nova Plantarum genera et species, etc. 4to. 1815.
- H. et B. pl. aeq.* Von Humboldt (Alexand. Fr. Henr.) et Bonpland (Amie) Plantes equinoxiales. fol. 1808.
- Herbert. append.* Herbert (William.) A Botanical arrangement of bulbous roots. 8vo. 1821.
- Herm. lugd.* Hermann (Paulus.) Flora Lugduno-Batavæ flores. 8vo. 1690.
- Herm. parad.* Hermann (Paulus.) Paradisi Batavi prodromus. 12mo. 1689.
- Hills. veg. syst.* Hill (John.) Outlines of a system of vegetable generation. 8vo. 1758.
- Hoff. c. gott.* Hoffmann (Georg. Franz.) Programma Hortus Goettingensis. fol. 1793.
- Hoff. deut. fl.* Hoffmann (Georg. Franz.) Deutschlands flora, ou Flore de l'Allemagne. 12mo. 1791.
- Hoff. phyt.* Hoffmann (Georg. Franz.) Phytographische Blätter. 8vo.
- Hoffm. salic.* Hoffmann (Georg. Franz.) Historia salicium iconibus illustro. fol. 1785.
- Hoff. umb.* Hoffmann (Georg. Franz.) Plantarum umbelliferarum genera. 8vo. 1816.
- Hoff. et Lk. fl. lus.* Hoffmannsegg et Link.
- Hoff. f. port.* Flore portugaise. fol. 1806.
- Hook. ex. flor.* Hooker (W. J.) Exotic flora. 8vo.
- H. et G. fil.* Hooker (W. J.) and Greville (W. K.) Icones Filicum. fol. London. 1827. et seq.
- Hop. in. sturm. fl. sus.* Hoppe in Sturm Deutschland flora. 4to. 1798.
- Horn. fl. dan.* Hornemann (J. W.) Icones Plantarum sponte nascentium in regnis Daniæ et Norvegiae, etc. fol. 1761.
- Hort. Kew.* Aiton (Willam.) Hortus Kewensis, editio prima. 8vo. 1789.
- Hort. trans.* Transactions of the Horticultural Society of London. 4to.
- Host. gram.* Host (Nicol. Thomas.) Icones et descriptions Graminum austriacorum. fol. 1801.
- Hout. pfanz.* Houttuyn (Mart.) Des ritters von Linne Pflanzen system nach der anleitung des Houttuynschen werks. 8vo. 1777.
- Hout. syst.* Houttuyn (Mart.) Des ritters von Linne Pflanzen system nach der anleitung des Houttuynschen werks. 14 vol. 8vo. 1777—1788.
- Icon. Kämpf.* Kämpfer (Engelbert.) Icones selectæ plantarum. fol. 1791.
- Icon. Hort. Kew.* Delineations of exotic Plants cultivated in the Royal Garden at Kew. Published by W. T. Aiton.
- Isn. act. par.* Isnard. Histoire de l'Academie Royale des Sciences, avec les Mémoires de Mathematique et de Physique.
- Jacq. amer.* Jacquin (Nic. Jos.) selectarum Stirpium Americanarum Historia. fol. 1763.
- Jacq. am. pict.* Jacquin (N. J.) Stirpium Americanarum Historia.
- Jacq. aust.* Von Jacquin (Nicol. Jos.) Flora Austriacæ Icones. fol. 1773.
- Jacq. coll.* Von Jacquin (Nicol. Jos.) Collectanea ad Botanicam, etc. Spectantia. 4to. 1786.
- Jacq. f. ecl.* Von Jacquin (Jos. Franc.) Eclatæ Botanicae. fol. 1811.
- Jacq. frag.* Von Jacquin (Nicol. Jos.) Fragmenta Botanica. fol. 1809.
- Jacq. h. vind.* Von Jacquin (Nicol. Jos.) Hortus Botanicus vindobonensis. fol. 1770.
- Jacq. ic.* Von Jacquin (Nicol. Jos.) Icones Plantarum rariorum. fol. 1781.
- Jacq. misc.* Von Jacquin (Nicol. Jos.) Miscellanea Austriaca ad Botanicam, etc. Spectantia. 4to. 1778.
- Jacq. obs.* Von Jacquin (Nicol. Jos.) Observations botanicae. fol. 1764.
- Jacq. ox.* Von Jacquin (Nicol. Jos.) Oxalis monographia iconibus illustrata. 4to. 1792.
- Jacq. schoenb.* Von Jacquin (Nicol. Jos.) Plantarum rariorum horti cæsari Schœnbrunnensis. fol. 1797.
- Jacq. stap.* Von Jacquin (Nicol. Jos.) Stapelia cultæ. fol. 1806.
- J. mem. ac. sci. par.* Jussieu, Mémoires de la Société d'Histoire Naturelle de Paris. 4to. 1799.
- Journ. sc.* Ker (J. B.) Journal of Science and the Arts. 8vo.
- Jung. ic. rar.* Junghans (Phil. Casp.) Icones Plantarum ad vitam impressæ. fol. 1787.
- Juss. an.* Jussieu, Annales du Museum d'Histoire Naturelle. 4to. 1802.
- Juss. rut.* Jussieu (Adrien.) Mémoires sur les Rutaceæ. Paris. 1825. 4to.
- Kämp. am.* Kämpfer (Engelbert.) Amænitates exoticae. 4to. 1712.
- Kalm. it.* Kalm (Pehr.) Resatil norra America. 8vo. 1753.
- Ker's review.* Ker (J. B.) Review of the Genera Amaryllis, Crinum, and Pancratium, in the Journal of Science and the Arts. 8vo.
- Ker's strelitz.* Ker (H. B.) Coloured Figures of the Genus Strelitzia. fol.
- Knight's prot.* Knight (Joseph.) Horticultural Essays on the natural order of Proteæ. 4to. 1809.
- Knor. del.* Knorr (Georg. Wolfgang.) Deliciae naturæ selectæ oder auserlesene. fol. 1766.
- Knorr. thes.* Knorr (Georg. Wolfgang.) The-

- taurus rei herbariae hortensisque universalis.* fol. 1770.
- Kon. an. bot.* Annals of Botany, by C. Konig and J. Sims. 2 vol. 8vo. I. 1805. II. 1806.
- Krock. siles.* Krockier (Ant. Joh.) Flora Silesiaca renovata. 2 vol. 8vo.
- Kth. mim.* Kunth (Car. Sig.) Mimoses et autres plantes legumineuses du nouveau continent. fol. 1819.
- Lab. nov. hol.* Labillardiere (Jac. Jul.) Nova Hollandiae plantarum specimen. fol. 1804.
- Lab. ic. pl. Syr.* Labillardiere (Jac. Jul.) Icones plantarum Syriæ rariorium. 4to. 1791.
- Lab. voy.* Labillardiere (Jac. Jul.) Relation du Voyage a la recherche de La Peyrouse. 4to. 1798.
- Lamb. cinch.* Lambert (Aylm. Bourk.) Description of the Cinchona. 4to. 1797.
- Lamb. pin.* Lambert (Aylm. Bourk.) A Description of the genus Pinus. fol. 1803.
- Lam. ill.* Monet De La Mark (Jean Bapt.) Illustration des Genres. 1791.
- Lap. pyr. ic. ined.* Picot De La Peyrouse (Philippe.) Figures de la Flore des Pyrenees. fol. 1795.
- Laroch. eryng.* De La Roche (Francois.) Eryngiorum nec non generis novi alepideæ historia. fol. 1808.
- Ledeb. fl. alt.* Ledebour. Flora Altaica. fol. 1829.
- Leers. fl. herb.* Leers (Joh. Dan.) Flora Herbornensis. 8vo. 1789.
- Lehm. nicot.* Lehman (Joh. Georg. Chr.) Generis Nicotianarum historia. 4to. 1818.
- L. Herit. ger.* L'Heritier (Charles Louis.) Geraniologia sive Erodii, etc. historia. fol. 1787.
- L. Her. sert. ang.* L'Heritier (Charles Louis.) Sertum anglicum seu Plantæ rariores, etc. fol. 1788.
- L. Her. stirp.* L'Heritier (Charles Louis.) Stirpes novæ aut minus cognitæ. fol. 1784.
- Lind. als.* Von Lindern (Fr. Balth.) Hortus Alsaticus Plantas in Alsatia, nobili designans. 12mo. 1747.
- Lindl. coll.* Lindley (John.) Collectanea Botanica. fol.
- Lindl. digit.* Lindley (John.) Digitalium Monographia. fol.
- Lindl. ros.* Lindley (John.) Rosarum Monographia, or a Botanical History of Roses. 8vo. 1820.
- Lin. amœn. ac.* Linnaeus s. Von Linne (Carolus.) Amœnitates academicae seu diss. antehac seorsim editæ. 8vo. 1749.
- Lin. fl. lap.* Linnaeus (Carolus.) Flora Lapponica. 1 vol. 8vo. 1737.
- Linn. hort. clif.* Linnaeus (Carolus.) Hortus Cliffortianus. 1 vol. fol. 1737.
- Lin. f. dec.* Linnaeus Filius (Carolus.) Plantarum rariores horti Upsaliensis decas 1. 1762. fol. 1763.
- Linn. fil. fasc.* Linnaeus Filius (Carolus.) Plantarum rariores horti Upsaliensis. fasc. 1. fol. 1767.
- Linn. sup.* Linnaeus Filius (Carolus.) Supplementum Plantarum. 1 vol. 8vo. 1781.
- Linn. trans.* Transactions of the Linnean Society of London. vol. 1. to 16. 4to.
- Lk. et Ott. abb.* Link et Otto. Abbildungen und Beschreibungen seltener Pflanzen im Berl. Garten. Berlin. 1821 et seq.
- Lk. et Ott. dis. cac.* Link et Otto. Cacti et Echinocacti. icones.
- Link. et Ot. h. ber.* Link et Otto. Hortus Berlinensis. fol.
- Lobel. adv.* De Lobel seu Lobelius (Mathias.) Adversariorum altera pars. 1 vol. fol. 1605.
- Lob. ic.* Lobelius (Mathias.) Stirpium Icones. 1 vol. 4to. 1591.
- Lodd. bot. cab.* Botanical Cabinet, by Messrs. Loddiges'. 4to.
- Loef. it. rar.* Loefling (Petr.) Iter Hispaniæ cum, eller Resa til spanska, etc. 1 vol. 8vo. 1758.
- Loes. prus.* Loeselius (Joh.) Flora Prussica. 1 vol. 4to. 1703.
- Loisel. gal.* Loiseleur-Deslonchamps (J. L. A.) Flora Gallica. 2 vol. 12mo. I. 1806, II. 1807.
- Magn. b. mons.* Magnol (Petrus.) Botanicon Monspeliense. 1 vol. 12mo. 1686.
- Marcg. bras.* Marcgravius (Georg.) Historia rerum naturalium Brasiliæ. 1 vol. fol. 1648.
- Mart. cent.* Martyn (John.) Historia Plantarum rariores rariorū centuriæ 1 de-
- Mart. dec.* cas 1-5. fol. 1728.
- Mart. fl. rust.* Martyn (Thomas.) Flora rustica. 4 vol. 8vo. 1792-4.
- Mart. n. pl. bras.* Martius (C. F. P.) Nova Genera et Species Plantarum quas in itinere, par Brasiliam. ann. 1817-1820. 4to.
- Mass. Stap.* Masson (Francis.) Stapeliae novæ fasc. fol. 1796.
- Math. com.* Mathiolus (P. Andr.) Commentarii in libros Dioscoridis de Medica Materia.
- Meerb. ic.* Meerburg (Nicol.) Plantarum selectarum Icones pictæ. fol. 1798.
- Menz. pug.* Mentzel (Christ.) Pugilus Plantarum rariores. fol. 1682.
- Merian. surin.* Merian (Maria Sibylla.) De metamorphosibus insectorum Surinamensium. 1 vol. fol. 1726.
- Mich. amer.* Michaux (André.) Flora Boreali Americana. 2 vol. 8vo. 1803.
- Mich. arb.* Michaux fils (André Franc.) Histoire des arbres forestiers de l'Amérique septentrionale. 3 vol. 4to. 1810-1813.
- Mich. querc.* Michaux (André.) Histoire des chênes de l'Amérique septentrionale. 1 vol. fol. 1801.
- Mich. gen.* Michelii (Petr. Anto.) Nova Plantarum genera. 1 vol. fol. 1729.
- Mill. ic.* Figures of Plants described in the Gardener's Dictionary, by Philip Miller. 2 vol. fol. 1760.
- J. Mill. ic.* Seven coloured Plates of Plants, by John Miller.
- J. F. Mill. ic.* Sixty coloured Plates of Plants and Animals, by Joh. Fred. Miller. fol.
- Miss Laur. ros.* A Collection of Roses from Nature, by Miss Lawrence (now Mrs. Kearse). 90 plates.
- Moret. fl. vicent.* Moretti (Gius.) Notizia

BOOKS QUOTED.

- supra diverse piante da aggiungarsi alla flora Vicentina. 4to. 1813.
- Moris. hist.* } Morison (Robert J.) Plantarum
Moris. s. } historia universalis Oxoniensis.
Moris. ox. } 2 vol. fol. 1680.
- Moris. umb.* Morison (Robert.) Plantarum umbelliferarum distributio nova. 1 vol. fol. 1672.
- Munt. phyt. cur.* Munting (Abrah.) Phytographia curiosa. 1 vol. fol. 1702, 1713, 1727.
- Murr. com.* } Murray (Joh. Andr.) Commentarii Societatis regiae
Mur. c. got. } Scientiarum Goettingensis. 4to.
- Nectoux voy.* Nectoux (H.) Voyage dans la haute Egypte, avec des observations sur les diverses espèces de sené. fol. 1808.
- N. et Mart. ac. b.* Nees and Martius, in Mémoires de l'Académie royale des Sciences de Berlin. 4to. ab ann. 1770 ad 1816.
- Nees. hor. ber.* Nees von Esenbeck (G. G.) Horae physicae Berolinensis collectæ ex symbolis virorum doctorum Link, Rudolphi, etc. 1 vol. fol. 1820.
- Niss. act. par.* Nissolle in Histoire de l'Académie Royale des Sciences, avec les Mémoires de Mathématique et de Physique. 4to.
- Nois. jard. fr.* Noisette (Louis.) Le Jardin fruitier. 2 fasc. in 4to. Paris. 1813.
- Nov. act. pet.* Nova Acta Academæ Scientiarum Imperialis Petropolitanae.
- Nov. com. pet.* Novi Commentarii Academæ Scientiarum Imperialis Petropolitanae.
- Oed. fl. dan.* Von Oeder (Georg. Christ.) Icones plantarum sponte nascentium in regnis Daniæ et Norvegiæ, etc. fol.
- Ort. dec.* De Ortega (Cas Gomez.) Novarum aut rariorum Plantarum Horti Reg. Botan. Matrit. Descriptionum Decades. 4to.
- Osb. it.* Osbeck (Pehr.) Dagboek oef' er en ostendyck resa. 1 vol. 8vo.
- Palisot d'owar.* } Palisot de Beauvois. Flore
Pal. fl. d'ow. } des royaumes d'Oware et
of Benin. fol. 1805 et seq.
- Pall. act. pet.* Pallas (Peter Simon.) In Acta Academæ scientiarum imperialis Petropolitanae. 4to.
- Pall. it. ed. gal.* Voyage dans l'Empire Russe. 8 vol. 8vo.
- Pall. it. ed. germ.* Pallas (P. S.) Reise durch verschiedene provinzen des Russischen Reichs. 3 vol. 4to. 1771—1776.
- Pall. ross.* Pallas (P. S.) Flora Rossica. fol. 2 vol. 1784, 1788.
- Par. lond.* } Salisbury (Rich. Anth.) Parasal.
Sal. par. lond. } dissus Londinensis. 2 vol. 4to.
- Park. par.* Parkinson (John.) Paradisi in sole paradisus terrestris, or a garden of flowers. 1 vol. fol. 1629.
- Park. theat.* Parkinson (John.) Theatrum botanicum. 1 vol. fol. 1640.
- Patters. iter.* A Narrative of four journeys into the country of the Hottentots and Cafafria. By Will, Patterson. 4to. 1789.
- Pet. eng. herb.* } Petiver (James.) Herbarii
Pet. h. brit. } Britannici Catalogus. fol.
- Pet. gaz.* Petiver (James.) Gazophylacium naturæ et artis. fol. 1702, 1704.
- Pet. Th. veg. afr.* Aubert du Petit Thouars. Plantes des îles de l'Afrique australe. 3 fasc. in 4to. Paris. 1804—6.
- Pio viol.* Pio (Joh. Bapt.) De Violâ specimen. 4to. 1813.
- Pis. bras.* Piso (Guil.) Historia naturalis Brasiliæ. 1 vol. fol. 1648.
- Piso. lib.* Pisonis (Guil.) De India utriusque re naturali et medica libri 14. fol. 1658.
- Plenck. ic.* Plenck (Jos. Jac.) Icônes plantarum medicinalium. fasc. fol. 1803 et seq.
- Pl. rar. hung.* } Waldstein (Franc.) et Kitai-
bel (Paul.) Descriptiones et W. et K. hung. } Icônes plantarum rariorum Hungariae. 3 vol. fol.
- Pluk. alm.* Plukenett (Leonard.) Almagestum botanicum sive phyt. onomasticon. 1 vol. 4to. 1796.
- Pluk. am.* Plukenett (L.) Amalteum botanicum. 1 vol. 4to. 1705.
- Pluk. mant.* Plukenett (L.) Almagesti botanici Mantissa. 1 vol. 4to. 1700.
- Pluk. phyt.* Plukenett (L.) Phytographia sive Stirpium illustriorum, etc. 4 vol. 4to. 1691.
- Plum. amer.* Plumier (Charles.) Description des Plantes de l'Amérique. 1 vol. fol. 1693 et 1713.
- Plum. ed. B.* } Plante Americanæ à C. Plu-
Pl. ed. Burm. } mier detectæ et à J. Bur-
mann editæ. 5 fasc. fol. 1755.
- Plum. fil.* Plumier (C.) Traité des fougères de l'Amérique. 1 vol. fol. 1705.
- Plum. gen.* Plumier (C.) Nova plantarum Americanarum genera. 1 vol. 4to. 1703.
- Plum. ic.* Plumier (C.) Plantarum Americanarum fasc. 10, continentis plantes quas olim C. Plumieri detexit et depinxit, Edidit Jo. Burmannus.
- Pohl. pl. bras.* Pohl Plantæ Brasiliæ. 4to. 1830.
- Pollich Palat.* Pollich (Joh. Adam.) Historia plantarum in Palatinatu sponte nascentium. 3 vol. 8vo. 1776.
- Poit. et T. fl.* Poiteau et Turpin. Flore Parisienne. 6 livr. fol. 1808.
- Pursh amer.* Flora Americae Septentrionalis; or a systematic arrangement and description of the plants of North America. By Frederick Pursh. 2 vol. 8vo.
- Quer. hisp.* Querry Martiner (Joseph.) Flora Espanola. 4 vol. 4to. 1762—1764.
- Raii syn.* } Ray (Joh.) Synopsis Stirpium Bri-
Ray. syn. } tannicarum. 1 vol. 8vo.
- Rauw. it.* Rauwolf (Leonth.) Etlicher Schoner Ausslandischer Kreuter, deren in Seiner Rayss, in die Morgenlander Gethon, Gedacht wirt, lebendige Contrafactur.
- Red. lil.* } Redoute (P. J.) Les Lilacées.
Redoute. lil. } 8 vol. fol. 1802—1816.
- Redout. ros.* Redoute (P. J.) Les Roses. 4to. et fol. 1817 et seq.
- Reich. ac.* Reichenbach (Ludov.) Monographia generis Aconiti et Delphinii. Lipsiæ. 1820. fol.
- Reich. ic.* Reichenbach (Ludov.) Icônes et descriptions plantarum. 4to. Lipsiæ. 1822 et seq.
- Relig. Houst.* Reliquiæ Houstounianæ, in Bibliotheca Jos. Banks asservatis. 4to. 1781.

- Renealm. spec.* Pauli Renealni Specimen Historiae Plantarum. 1 vol. 4to. 1611.
- Retz. obs.* Retzius (And. Joh.) Observations Botanicæ. 6 fasc. fol. 1774—1791.
- Rheed. mal.* Van Rheede (Henricus.) Hortus Indicus Malabaricus. 12 vol. fol.
- Rhe.hort. mal.* tus Indicus Malabaricus. 12 vol. fol.
- Rivin. monop.* Rivinus (Aug. Quirinus.) Ordo Plantarum flore irregulari monopetalo. 1 vol. fol. 1690.
- Riv. tetr.* Rivinus (A. Q.) Ordo Plantarum flore irregulari tetrapetalo. 1 vol. fol. 1691.
- Riv. pent.* Rivinus (A. Q.) Ordo Plantarum flore irregulari pentapetalo. 1 vol. fol. 1699.
- Rob. ic.* Three hundred and nineteen plates of plants; engraved by Nic. Robert. A. Bosse, and Lud. de Chatillon. fol.
- Rod. Scub. cer.* Scubler (Gust.) Diss. sistens descriptiones cerealium cultorum. Tubing. 1818. 4to.
- Ræm. arch.* Römer (Joh. Jac.) Archiv für die Botanik. 3 vol. 4to. 1796, 1799, 1805.
- Ræm. script.* Römer (J. J.) Scriptores de Plantis Hispanicis, Lusitanicis, et Brasiliensis. 1 vol. 8vo. 1796.
- Ræm. et Ust. mag.* Römer (Joh. Jac.) et Usteri (Paul.) Magazin für die Botanik. 4to. 1789—1790. Zurich.
- Rosc. scitam.* Roscoe's Figures of the order of Scitamineæ. fol.
- Rosc. fl. ill. sea.* Roscoe (Mrs. M.) Floral Illustrations of the Seasons. Liverpool. Royal 4to. 1829 et seq.
- Roth. abh.* Roth (Alb. Wilh.) Botanische abhandlungen und beobachtungen. 4to. 1787.
- Rottb. gram.* Descriptiones et Icones Plantarum rariorum, etc.
- Rottb. desc. rar.* 1 vol. fol. 1773.
- Roxb. corom.* Roxburgh (William.) Plants of the coast of Coromandel, published under the direction of Sir Joseph Banks. fol.
- Rudg. gui.* Rudge (Edward.) Plantarum Guianeæ rariorum Icones et desc. fol. 1805.
- Ruiz. et P. prod.* Ruiz (Hippol.) et Pavon (Jos.) Floræ Peruvianaæ et Chilensis prodromus. fol.
- Rumph. amb.* Rumphius (Georg. Everh.) Herbarium Amboinense. 6 vol. fol. 1750-1755.
- Russ. alep.* Russell (Alex.) Natural History of Aleppo. 1 vol. 4to. 1756. ed. 2. 1794.
- Sabb. hort.* Sabbati (Liberatus.) Hortus Romanus. fol. vol. 3. 8vo. 1772, 1774, 1805.
- Sal. stirp. rar.* Salisbury (R. A.) Icones stirpium rariorum. 1 fasc. fol. 1791.
- San. viag.* Santi (Georg.) Viaggi al monte Amiata. 1 vol. 8vo. 1795. II. 1798. III. 1806.
- Savi fl. pis.* Savi (Gaetano.) Flora Pisana. 2 vol. 8vo. 1798.
- Scheuch. gram.* Scheuchzer (Johan.) Agrostographia, sive Graminum, Juncorum, Cyperorum, iisque affinium historia.
- Scheuch. it.* Scheuchzer (Joh. Jac.) Itineraria Alpina. 4to. ed. 1. 1702—1708. Ed. 2; 1 vol. 4to. 1723.
- Schk. caric.* Schkuhr (Christ.) Histoire des Carex ou Laiches. 1 vol. 8vo. 1802.
- Schkahr filic.* Schkuhr (Christ.) Abbildungen der Farrnkräuter. 4to.
- Schk. hand.* Schkuhr (C.) Botanisches handbuch. 3 vol. 8vo.
- Schm. arb.* Schmidt (Franz.) Österreichs allegemene Baumzucht 2 Band. fol. 1792—1794.
- Schmid. arb.* Schmidel (Cas. Christ.) Icones Plantarum. 1 vol. fol. 1762.
- Schousb. maroc.* Schousboe (P. K. A.) Jagttæ gelser over voextriget i Marocco. 1 vol. 4to. 1800.
- Schrader. d. hal.* Schrader (Henr. Adolph.) De Halophytis Pallasiæ. 4to. 1810.
- Schrader. gen. nov.* Schrader (H.A.) Nova Plantarum genera. 1 fasc. fol. 1797.
- Schrader. germ.* Schrader (H.A.) Flora Germanica. 8vo. 1806.
- Schr. s. han.* Schrader (H.A.) Sertum Hannoveranum. 4 fasc. fol. 1795, 1796.
- Schrader. verb.* Schrader (H.A.) Monographia generis verbasci. 4to. 1813.
- Schrader. ver.* Schrader (H.A.) Commentatio de Veronicais spicatis. 8vo. 1803.
- Schrank h. mon.* Schrank (Fr. von Paula.) Plantæ rariores horti Monacensis. 1817, 1818, 1819.
- Schreb. a. ups.* Von Schreber (Joh. Christ. Dan.) Acta litteraria et scientiarum Upsaliæ aut ab academia Upsaliensi publicata. 2 vol. 1720, 1816.
- Sch. gram.* Schreber (J. C. D.) Beschreibung der Græser. fol. 1769—1779.
- Scop. carn.* Scopoli (Joh. Ant.) Flora Carniolica. 8vo. 1760. ed. 2. 2 vol. 8vo. 1772.
- Scop. del.* Scopoli (J.A.) Deliciae floræ Insulæ bracæ. 3 vol. fol. 1786, 1788.
- Seba. mus.* Seba (Alb.) Locupletissimi Rerum naturalium Thesauri Descriptio. 4 vol. fol. 1734—1765.
- Second. d. chen.* De Secondat (M.) Mémoires sur l'Histoire naturelle du Chêne. 1785. fol.
- Segu. ver.* Seguier (Joh. Franc.) Plantæ Véronenses. 3 vol. 8vo. 1745.
- Ser. m. helv.* Seringe (N. C.) Musée Helvétique d'histoire naturelle. 4to. 1818, 1820.
- Shaw afric.* Shaw (Thomas.) Travels in Barbary and the Levant. 1 vol. 1738.—Supplement. 1746.
- Sinc. gr. wob.* Sinclair (George.) Hortus gramineus Woburnensis. 1 vol. 8vo. 1825.
- Sloane.* Sloane (Hans.) A Voyage to the Islands Madeira, Barbadoes, Nieves, St. Christopher's, and Jamaica; with the Natural History of the last of those Islands. 2 vol. fol. 1707.
- Sm. exot. bot.* Smith (J. Edward.) Exotic Botany. 8vo. 1804—1808.
- Sm. ic. ined.* Smith (J. E.) Plantarum Icones hactenus ineditæ. fasc. 1-3. 1789, 1791. fol.
- Smith. ic. pict.* Smith (J. E.) Icones pictæ Plantarum rariorum. 3 fasc. fol. 1790—1793.
- Sm. nov. hol.* Smith (J. E.) A specimen of the Botany of New Holland. 1 vol. 4to. 1793.

BOOKS QUOTED.

- Smith spic.* Smith (J. E.) *Spicilegium Botanicum.* 2 fasc. fol. 1791, 1792.
- Smith. tracts.* Smith (J. E.) *Tracts relating to Natural History.* 1 vol. 8vo. 1798.
- Sole's mints.* Sole (William.) *Menthæ Britannicæ; being a new botanical arrangement of all the British Mints hitherto discovered.* 1 vol. fol. 1798.
- Sonner. it.* Sonnerat (P.) *Voyage aux Indes orientales et à la Chine.* 2 vol. 4to.
- Sonner. n. guin.* Voyage à la nouvelle Guinée. 1 vol. 4to. 1776.
- Sparm. a. st.* Sparrman (Andr.) *Kongl. Vetenskaps Academiens Handlingar.* Stockholm. 1739. seq. 8vo.
- Spr. anl.* Sprengel (Kurt.) *Anleitung zur Kentmiss der Gewachse.* 2 theil. 8vo. 1817, 1818.
- Spreng. crypt.* Sprengel (K.) *Introduction to the study of Cryptogamous Plants, translated from the German.* 8vo. 1813.
- Spr. fl. hall.* Sprengel (K.) *Floræ halensis Tentamen novum.* 1 vol. 8vo. 1806.
- Spreng. umb.* Sprengel (K.) *Species umbelliferarum minus cognitarum illustratae.* 1 vol. 4to. 1818.
- Sternb. saxif.* Sternberg (Graf. Casp.) *Revisio Saxifragarum Iconibus illustrata.* 1 vol. fol. 1810.
- Stær. lib. de stram.* Stærck (Ant.) *Libellus de Stramonio, Hyosiamo. Aconito.* 8vo. 1763.
- Sturm. deut.* Sturm (Jacob.) *Deutschland flora.* 4to.
- Swartz ic.* Swartz (Olof.) *Icones Plantarum Indiae occidentalis.* 1 fasc. fol. 1794.
- Swart. obs.* Swartz (O.) *Observationes botanicae.* 1 vol. 8vo. 1791.
- Sw. syn. fil.* Swartz (O.) *Synopsis filicum earum genera et species compl.* 1 vol. 8vo. 1806.
- Swt. br. fl. gar.* Sweet (Robert.) *The British Flower Garden, published in monthly numbers, containing full and accurate coloured figures and descriptions, with the best mode of cultivation, of the most choice and curious plants, that may be cultivated in the open air of Great Britain.* 3 vol. royal 8vo.
- Swt. br. fl. gar. s. 2.* Sweet (R.) *British Flower Garden, series 2, number 1 to 17, et seq.*
- Swt. cist.* Sweet (R.) *Cistinæ. The natural Order of Cistus, or Rock-Rose, complete in one volume, containing 112 beautifully coloured figures of the different species and varieties of this handsome family of plants.*
- Swt. flor. aust.* Sweet (R.) *Flora Australasica, or a selection of the most beautiful and interesting plants, natives of New Holland, and the South Sea Islands, containing 56 beautifully coloured figures from the living plants.* 1 vol. royal 8vo. London, 1827-1828.
- Swt. flor. guid.* Sweet (R.) *The Florist's Guide, or Cultivator's Directory, containing elegantly coloured figures of the choicest flowers cultivated by Florists, in-*
- cluding Tulips, Hyacinths, Ranunculus', Carnations, Pinks, Auriculas, Polyanthus's, Georgina's, and Roses. 1827 et seq. royal 8vo. London.
- Sweet ger.* Sweet (R.) *Geraniaceæ, or the natural Order of Geraniums, published in monthly parts, with accurate coloured figures and descriptions of the different kinds, and their mode of culture.* royal 8vo. Five volumes complete.
- Swert. flor.* Swertia (Eman.) *Florilegium tractans de variis floribus.* 1 vol. fol. 1612.
- Tab. ic.* Tabernæmontanus (Jac. Theod.) *Eicones Plantarum cur N. Bassæo.* 4to. 1590.
- Tabern. kraut.* Tabernæmontanus (Jac.Theod.) *Kräuterbuch.* fol. 1588.
- Ten. flor. med.* Tenore (Michel.) *Saggio sulle qualità medicinali delle plante Napolitanæ.* 8vo. Napoli. 1808.
- Tenor. fl. nap.* Tenore (Michel.) *Flora Napo-litana.* fol. 1811.
- Ten. mem. croc.* Tenore (M.) *Mémoires sui Crochi.* Naples.
- Thunb. eric.* Thunberg (Car. Petr.) *de Erica.* 4to.
- Thunb. ir.* Thunberg (Car. Petr.) *Dissertatio de Iris.* 4to. 1782.
- Thunb. jap.* Thunberg (Car. Petr.) *Flora Japonica.* 8vo. 1784.
- Thunb. voy.* Thunberg (Car. Petr.) *Voyage au Japon, etc.* 8vo. 1796.
- Till. pis.* Tilli (Mich. Ang.) *Catalogus horti Pisani.* fol. 1723.
- Tourn. voy.* Pitton De Tournefort (Joseph.) *Relation d'un voyage du Levant.* 4to. 1717.
- Trat. arch.* Trattinick (Leop.) *Archiv der Gewächskunde.* 4to. 1811.
- Trattin. fl. aust.* Trattinick (Leop.) *Flora Austriae sicca.* fol. 1792.
- Tratt. tab.* Trattinick (Leop.) *Observationes botanicae Tabularium rei herbariae illus-trates.* 4to. 1811.
- Tratt. thes. bot.* Trattinick (Leop.) *Thesaurus Botanicus.* fol. 1819.
- Trev. delph.* Treviranus (Lud. Christ.) *De Delphinio et Aquilegia observationes.* 4to. 1817.
- Trew. ehret.* Trew (Christ. Jac.) *Plantæ selectæ ab Ehret pictæ.* fol. 1750.
- Trew. pl. rar.* Trew (Christ. Jac.) *Plantæ rariiores.* ed. J. C. Keller. fol. 1763.
- Trin. ic. gram.* Trinius (Car. Bern.) *Icones Graminæ.*
- Triumf. obs.* Triumfetti (Joh. Bapt.) *Observationes de ortu et vegetatione Plantarum.* 4to. 1635.
- Ust. N. ann.* Usteri (Paul.) *Neue annalen der Botanik.* 8vo. 1794.
- Vahl. ecl.* Vahl (Martinus.) *Eclogæ Ameri-canæ.* fol. 1796.
- Vahl. ic. rar.* Vahl (Martinus.) *Icones Plantarum in eclogis descriptarum.* fol. 1798.
- Vahl. symb.* Vahl (Martinus.) *Symbolæ botanicae.* fol. 1790.
- Vail. par.* Vaillant (Sebastien.) *Botanicon Parisiense.* fol. 1727.

- Vent. cels.* Ventenat (Etienne Pierre.) Description des plantes nouvelles ou peu connues du jardin de J. M. Cels. fol. 1800.
- Vent. choix.* Ventenat (Etienne Pierre.) Choix des plantes. fol. 1803.
- Vent. malm.* Ventenat (Etienne Pierre.) Jardin de la Malmaison. fol. 1803.
- Vill. c. stras.* Villars (D.) Catalogue methodique du jardin de Strasbourg. 8vo. 1807.
- Vill. dauph.* { Villars (D.) Histoire des Plantes du Dauphine. 8vo. 1786.
- Viv. frag.* { Viviani (Dom.) Floræ Italicae fragmenta. 4to. 1808.
- Vivian. fr. fl. it.* { licæ fragmenta. 4to. 1808.
- Volck. norib.* Volkamer (Jo. Georg.) Flora Noribergensis. 4to. 1700.
- Wahl. helv.* Wahlenberg (Georg.) De vegetatione et clymate Helvetiae septentrionalis specimen. 8vo. 1813.
- Wahl. lap.* Wahlenberg (Georg.) Flora Lapponica. 8vo. 1812.
- Wal. as. res.* Wallich in Asiatic Researches, or Transactions of the Society instituted in Bengal. 4to. 1788.
- Wall. pl. as.* } Wallich (Nathanael.) Plantæ asiaticæ rariores, ou description et figures d'un choix de plantes inédites de l'Inde orientale. fol. 1829 et seq.
- Wall. pl. ind.* } Walther (Aug. Frid.) Designatio plantarum horti ejus. 8vo. 1735.
- Walth. hort.* Von Wangenheim (Fried Ad. Jul.) Anpflanzung North Amerikanischer Holzarten.
- Wats. dend. brit.* Watson (W.) Dendrologia Britannica. 2 vol. 8vo.
- Weig. obs.* Weigel (Christ. Ehret.) Observations Botanicæ. 4to. 1772.
- Wei. et N. rub.* Weihe (Aug.) et Nees (C. G.) Rubi germanici.
- Weinm. phyt.* Weinmann (Joh. Guil.) Phytanthoza Iconographica. fol. 1737.
- Wendl. ac.* Wendland (H. Lud.) Commentatis de Acaciis aphyllis. 4to. 1820.
- Wendl. col.* Wendland (H. Lud.) Collectio Plantarum tum exoticarum quam indigenarum. 4to. 1805.
- Wendl. er.* Wendland (Joh. Christ.) Ericarum Icones et descriptions. 4to. 1798.
- Wendl. h. herr.* Wendland (Joh. Christ.) Hortus Herrenhusanus. fol. 1798.
- Wend. s. han.* Wendland. (J. C.) Sertum Hanoveranum. 4 fasc. fol.
- Willd. arb.* Willdenow (Car. Lud.) Berlinische Baumzucht. 8vo. 1797.
- W. ber. mag.* Willdenow (Car. Lud.) Berlinischer magazin oder gesammlete schriften, etc. 4to. 1765.
- Willd. hort. ber.* Willdenow (Car. Lud.) Hortus Berolinensis. fol. 1806.
- Woodv. med. bot.* Woodville (William.) Medical Botany. 3 vol. 4to. 1790.
- Zanon. hist.* Zanoni (Jacobus.) Rariorum stirpium historia. Ed. C. Monti. 1 vol. fol. 1742.
- Zorn. ic.* Zorn (Joh.) Icones Plantarum medicinalium. 8vo. 1779.

ABBREVIATIONS.

- A. B. Annals of Botany.
A. DC. Alphonso De Candolle. Monographie des Campanulees.
Afz. Afzelius, a Swedish Professor.
A. G. Andrews's Geraniums.
A. H. Andrews's Heaths.
A. R. Andrews's Botanist's Repository.
All. Allioni, an Italian Botanist.
Aub. Aublet, a French Traveller in Guiana.
B. Mr. Robert Brown, a celebrated English Botanist.
B. A. Barton's Flora Boreali Americana. 2 vol. 4to.
Bal. Balbis, a French Botanist.
Bau. Baumgarten.
B. C. Loddiges's Botanical Cabinet.
B. D. Bartling. *Diosmeæ descriptæ et illustræ.* 1824. 8vo.
Be. Bertoloni, an Italian Botanist.
Bes. Besser, a Russian Botanist.
BF. Bancroft.
B. F. G. Sweet's British Flower Garden.
Bhi. Bernhardi, a German Botanist.
B. M. Curtis's Botanical Magazine.
B. O. Beauvois. *Flora d'Oware et de Benin.*
Boj. Bojer, Professor of Botany in the Isle of France.
B. P. Brown's Prodromus Floræ novæ Hollandiæ.
B. R. Botanical Register.
Bro. Brogniart's Rhamneæ.
B. T. Burchell's Travels in Africa.
C. (Cav.) Cavanille's, a Spanish Botanist.
C. C. Colvill's Catalogue.
C. I. Cavanille's Icones.
C. L. T. Colebrooke, in Linnean Transactions.
C. S. Choisy. *Mémoire sur la Famille des Selaginées.*
Cyr. Cyrilli, an Italian Botanist.
D. David Douglas, on the Plants of the West Coast of America.
DC. Decandolle (Augustin Pyramus.)
D. D. David Don, Librarian of the Linnean Society, Soho Square.
Desf. Desfontaines, a French Botanist.
Desv. (Dv.) Desvaux, a French Botanist.
DF. Dufresne's Valerianees.
D. L. (Lht.) David Lockhart, Botanical Gardener at Trinidad.
Dod. Dodonaeus, a Dutch Botanist.
D. P. David Don's Prodromus Floræ Nepalensis.
E. Ehrenberg, a Traveller in Egypt, &c.
E. B. Smith's English Botany.
E. C. Elliott's Flora of Carolina.
E. F. Smith's English Flora.
Ex. B. Smith's Exotic Botany.
F. Fischer, Curator of the Botanic Garden at St. Petersburgh.
F. D. Flora Danica.
F. E. Forskaol Flora Ægyptiaco-Arabica.
F. G. Flora Graeca.
F. I. Flora Indica.
G. Gärtner, a celebrated German Carpologist.
G. B. Gray's Flora Britannica.
G. D. George Don, an English Botanist.
Gm. Gmelin, a Russian Botanist.
Gr. Graham, Regius Professor of Botany at Edinburgh.
G. R. Georgi, Geographische, etc. Bechreibung des Russischen Reichs.
H. Hornemann, a Danish Botanist.
H. A. Hooker's Flora Boreali Americana.
H. B. Hortus Bengalensis.
H. E. Hortus Epsomensis.
H. E. F. Hooker's Exotic Flora.
H. F. Hooker and Greville's Ferns.
H. G. Hoare's Geraniaceæ.
H. K. Hortus Kewensis.
H. M. Hooker's Miscellany.
H. M. N. Haworth's Miscellanea Naturalia.
H. P. Idem, in Taylor's Philosophical Magazine.
H. R. Idem, Saxifragæ and Revisions of Succulent Plants.
H. S. Idem, Synopsis Plantarum Succulentarum.
H. S. S. Idem, Supplementum Plantarum Succulentarum.
H. S. L. Hortus Suburbanus Londinensis.
H. T. Transactions of the Horticultural Society.
H. U. Hoffman's Umbelliferæ.
J. Jussieu.
Jac. Jacquin.
J. E. (J. F.) Jacquin (Filius) Eclogæ Botanicae.
J. I. Jacquin's Icones.
J. S. Jacquin's Stapeliæ.
K. F. Kaulfuss. Filices.
K. S. Kunth's Synopsis Plantarum.
L. Linnæus.
L. A. Lehman's Asperifoliae.
L. A. Ledebour's Flora Altaica.
Lam. Lamarck.
L. C. Loddiges's Catalogue.
L. col. Lindley's Collectanea Botanica.
Led. Ledebour, a Russian Botanist.
L. en. Link's Enumeratio Plantarum.
LG. et Lag. Lagasca, a Spanish Botanist.
L'H. L'Heritier.
L. O. Link et Otto, Abbildungen et Beschreibungen seltener Pflanzen in Berlin Garten.
L. P. Lehman's Potentillæ.
L. R. Lindley, Rosarum Monographia.
L. T. Transactions of the Linnean Society.
M. Michaux. Flora Boreali Americana.
M. B. Marschall de Bieberstein, Flora Tauro-Caucasica.
M. C. Mackay's Catalogue.
Med. Medicus, a German Botanist of the last Century.
Mh. Mönch, a German Botanist.

- Mil. Miller, an English Gardener and Botanist.
 M. N. Martius. Flora of Brazil.
 M. P. Martius. Genera et Species Palmarum.
 N. Nuttall's Genera of North American Plants.
 N. A. Nee's Asteræ.
 N. L. F. News of Literature and Fashion.
 N. P. Nestler's Potentillæ.
 Ot. Otto, Curator of the Botanic Garden at Berlin.
 P. B. Pohl's Flora Brasiliana.
 Ph. Pursh. Flora Americana Septentrionalis.
 P. L. Salisbury in Paradisus Londinensis.
 Poir. Poiret, a French Botanist.
 Pr. Presl, a Bohemian Botanist.
 P. S. Persoon's Synopsis Plantarum.
 R. (Rox.) Roxburgh.
 R. A. Reichenbach. Monographie Generis Aconiti et Delphinii.
 Raf. Rafinesque Schmalz.
 R. B. Roth's neue Beitrage zur Botanik.
 R. C. Roxburgh's Plants of the Coast of Coromandel.
 R. O. Richard's Orchideæ.
 R. P. Ruiz et Pavon. Flora Peruviana.
 R. S. Reemer et Schultes Systema Vegetabilium.
 R. U. (Rch.) Reichenbach Ubersicht der gat-tung Aconitum.
 S. Schulze's Systema Vegetabilium.
 S. C. Sweet's Cistineæ.
 Sch. (Schl.) Schleicher, a Swiss Collector.
 Sck. Schrank, a Bavarian Botanist.
 Scop. Scopoli, an Italian Botanist.
 Scr. Schrader, a German Botanist.
 Set. Schott, a Traveller in Brazil.
 S. F. A. Sweet's Flora Australasica.
 S. F. G. Sweet's Florist's Guide.
 S. G. Sweet's Geraniaceæ.
 Sie. Sieber, a Botanical Collector.
 S. M. Sole's Mints.
 S. O. Seidl et Opiz.
 S. P. Salisbury's Prodromus.
 S. S. Sprengel's Systema Vegetabilium.
 S. T. Savi. Observations in varias Trifoliorum Species.
 Sw. Swartz, a Swedish Botanist.
 T. C. Tenore. Catalogus Plantarum Horti Regii Neapolitanæ.
 T. N. Idem, Flora Neapolitana.
 T. P. Idem, Flora Neapolitanæ Prodromi.
 T. S. Idem Synopsis Floræ Neapolitanæ.
 Th. Thunberg, a Swedish Botanical Traveller.
 Tv. Treviranus, a German Botanist.
 V. Ventenat, a French Botanist.
 V. M. Ventenat's Jardin de Malmaison.
 Viv. Viviani, an Italian Botanist.
 W. Willdenow. Species Plantarum.
 W. A. Idem, Historia Amaranthorum.
 W. en. Idem, Enumeratio Plantarum Horti Bot. Berolinensis.
 Wall. Wallich, Curator of the Botanic Garden at Calcutta.
 W. C. Wendland. Collectio Plantarum.
 W. D. Watson's Dendrologia Britannica.
 W. H. William Herbert. Observations on the Amaryllideæ, &c.
 W. K. Waldstein and Kitaibel. Descrip-tions et icones plantarum rariorum Hungariae.
1. 2. Flowering the first and second month of the year.
 3. 6. Flowering from the third to the sixth month, and so on according to the different figures.
 1. 12. Flowering all the year round, or at various seasons of the year, not having any settled period.
 — Denotes the same as the line above it.
 ... Uncertain, or not properly known.
 H. *l.* Hardy large tree.
 H. *h.* Hardy Shrub, or small tree.
 H. *th.* Hardy Evergreen Shrub.
 F. *h.* Frame Shrub, or requiring the protec-tion of a Frame or Mats in severe frosty weather, but to be exposed when the weather is mild.
 G. *h.* Greenhouse Shrub, requiring the pro-tection of a Greenhouse in Winter.
 S. *h.* Shrubs requiring to be kept in the Stove or Hothouse in Winter, but may be removed to the Greenhouse in Summer.
 D. S. *h.* Dry Stove Shrub, a plant requiring very little water.
 D. G. *h.* Dry Greenhouse Shrub, or a plant that requires but little water in Winter; many of which will succeed well in a good frame or pit, with the covering of a mat or two on the lights in severe frost, but to be continually exposed to the air in mild weather.
h. Suffruticose, a dwarf Shrub, or a soft wooded Shrub.
 4. Herbaceous, perennial, dying down in Winter, and shooting up afresh the following Spring.
h. Climbing Shrub, a Shrub that re-quires support, either with a trellis, stick, or line, or any other convenient support.
 4. Climbing, perennial, herbaceous plant, requiring the same sort of support as the last.
 w. *h.* Water Shrub, or one that prefers grow-ing in water, or in moist situations.
 w. 4. Perennial herbaceous plant, requiring to be grown in water, or in a moist situa-tion.
 3. Biennial, seldom surviving more than two years.
 ○. Annual, requiring to be sown yearly.
 b. Bulbous-rooted.
 H. Hardy, surviving through the Winter without protection.
 F. Frame, not quite hardy, but requiring the protection of a frame or mats in severe weather.
 G. Greenhouse, requiring the protection of a Greenhouse in Winter, but to be well supplied with air in mild weather.
 I. Intermediate between Stove and Green-house.
 S. Stove, those require the protection of a Stove or Hothouse in Winter; the tem-perature should never be allowed to get below 60 degrees of Fahrenheit's Ther-mometer.
 C. B. S. Cape of Good Hope.
 N. S. W. New South Wales.
 N. Holl. New Holland.
 V. Diem. Isl. Van Dieman's Island.

THE BRITISH FLOWER GARDEN, & ORNAMENTAL SHRUBBERY.

Series the Second.

Publishing, in Monthly Numbers, at Three Shillings each ; and each Number containing Four full and highly coloured Figures, with Dissections of the most essential parts, of the most beautiful, rare, curious, or interesting Herbaceous Plants, or Flowering Shrubs, that may be cultivated in the open Borders, Shrubberies, or by the side of Walls, or Fences, in the Gardens of this Country ; with a full account of the best and most successful mode of Management in Culture and Propagation ; the Soils and Situations most suitable for them ; the Protection necessary, if any be required in Winter ; their Scientific and English Names, with full Descriptions ; Natural and Linnean Classes and Orders ; Derivations of the Generic Names ; and all other information that may be considered of importance to the Botanist, Cultivator, or Amateur ; and the most beautiful and interesting subjects are chosen. Twenty-five Numbers form each Volume, and Three Volumes complete each Series.

Sets of the First Series of **SWEET'S BRITISH FLOWER GARDEN**, complete, in Three Volumes, may still be obtained, each Volume containing One Hundred elegantly coloured Figures ; with a Systematical, Alphabetical, and English Index to each, List of Books quoted, &c. Price 3*l. 16*s.** each. Single Numbers, at 3*s.* each ; any back Numbers may be obtained to complete Sets.

The First Series is published by W. SIMPKIN and R. MARSHALL, Stationers' hall-court, Ludgate Street ; and may be procured, by Order, from all Booksellers.

"The British Flower Garden ought to be in the hands of all Cultivators of Hardy Flowering Plants." *Loudon's Gardener's Magazine.*

"The Figures, drawn and coloured from living specimens, by E. D. Smith, under the Author's inspection, are correct portraits of the subjects intended, and the plates are full and well executed. At the bottom of the plates are given coloured dissections of the different parts of the flower, so that, by that means, the parts of fructification are well explained."—"We recommend the British Flower Garden to the Cultivators and admirers of hardy Flowering plants ; and we have no doubt but they will feel themselves much interested by its perusal." *News of Literature and Fashion.*

SWEET'S GERANIACEÆ.

Now complete, in Five Volumes, each containing One Hundred beautifully coloured Figures of the choicest and most distinct Species and Hybrid Productions of this greatly admired Tribe. From this Work may be selected a collection of the sorts most suited to the taste of any Lady or Gentleman, who wishes to make a collection of this beautiful Family.

SWEET'S CISTINEÆ.

An Account of the Family of CISTUS, or ROCK-ROSE ; illustrated with handsomely coloured Figures ; now complete, in Twenty-eight Numbers, at 3*s.* each ; containing One Hundred and Twelve Figures and Descriptions of this handsome and generally admired Tribe of Plants, with Descriptions of the other Species that could not be obtained ; with a full account of the best Method for their Cultivation and Propagation ; or any thing else belonging to them, considered of importance.

THE FLORIST'S GUIDE ; or CULTIVATOR'S DIRECTORY.

Publishing, in Monthly Numbers, at 3*s.* each ; and Two Volumes will complete the Work. Each Number contains Four highly finished coloured Figures, drawn from the living Plants of the Flowers most esteemed by Florists—as Tulips, Ranunculus', Auriculas, Hyacinths, Carnations, Pinks, Roses, &c. ; with the most approved Methods of Cultivation, Directions for raising new Varieties from Seeds, &c.

FLORA AUSTRALASICA.

Published in one handsome Royal Octavo Volume ; containing Fifty-six elegantly coloured Figures, of the most beautiful and curious Plants, Natives of New Holland, and the South Sea Islands, (generally called Botany Bay Plants.) They are well adapted for a Greenhouse, or Conservatory ; and many will survive the Winters, in the open ground, with a very little protection. Price 2*l. 2*s.* 6*d.** in Fourteen Numbers, Index, &c. ; or single Numbers, at 3*s.* each.

In the Press, and soon will be published, a New Edition of
SWEET'S HOOTHOUSE AND GREENHOUSE MANUAL ;
Or BOTANICAL CULTIVATOR.

Containing numerous Genera not before enumerated in any former edition ; with Directions for their Treatment in Cultivation and Propagation ; Treatment of Plants in Rooms, and all those grown in the Hothouses, Greenhouses, Frames, or in the open Ground of the Gardens, or Plantations in this Country ; with a full account of the Management of Bulbs, and the Parasitical and other Orchideous Plants.

SWEET'S BRITISH WARBLERS :

Or an Account of the Summer Birds of Passage belonging to the Family of Sylvia ; illustrated with beautiful coloured Figures of the natural size, chiefly taken from living Birds in the Author's collection ; with full Directions for their Treatment, as practised by the Author, showing how all the fine singing Birds belonging to this tribe, may be kept in confinement in as good health as any common birds whatever, and will continue in full song nearly all the year ; with an account of the best Method of catching them, or rearing them from the nest ; and numerous other observations concerning their migration, food, habit when wild, or any thing relating to their history, that may be considered of importance. Published in Three Parts, and containing Sixteen Figures. Part 1, containing Six Figures, 6*s. 6*d.** ; Part 2, Five Figures, 5*s.* ; and Part 3, Five Figures, 5*s.* The Figures are of the Nightingale, Blackcap, Larger and Lesser Whitethroats, Greater Pettychaps, or Garden Warbler ; Willow and Wood Wrens ; Dartford, Reed, and Sedge Warblers ; Winchat, Stonechat, Wheatear, Redstart, &c.

CLASSIS 1.

DICOTYLEDONEÆ seu EXOGENÆ.

* SUBCLASSIS 1. THALAMIFLORÆ.

ORDO I.

RANUNCULACEÆ. DC. syst. nat. 127.

Tribus 1. CLEMATIDEÆ.

Syst. Name.	Color.	Eng. Name.	Native.	Introd.	Flower.	Habit.	Ref. to Figures.
CLEMATIS. W.		VIRGIN'S BOWER.	Polyandria Polygynia. L.				
1 erécta. DC.	(wh.)	upright.	Austria.	1597.	6. 8. H. ȝ.	Jacq. aust. 3. t. 291.	
γ hispánica.	(wh.)	Spanish.					
2 flámmula. DC.	(wh.)	sweet-scented.	S. Europe.	1596.	7. 10. H. h.~	Lob. ic. 627. f. 1.	
α rotundifolia.	(wh.)	round-leaved.	—	—	—	H. h.~	Ten. fl. neap. v. 1. t. 48.
β vulgaris.	(wh.)	oblong-leaved.	—	—	—	H. h.~	
γ marítima.	(wh.)	narrow-leaved.	—	—	—	H. h.~	Zanic. ist. 69. t. 129.
3 Massoniâna. DC.	(wh.)	Masson's.	C. B. S.	G. h.~	
4 orientális. DC.	(ye.)	oriental.	Levant.	1731.	7. 10. H. h.~	Dill. elt. t. 119. f. 145.	
5 glâuca. w.	(ye.)	glaucous.	Siberia.	7. 8. H. h.~	Willd. arb. 65. t. 4. f. 1.	
6 paniculâta. DC.	(wh.)	panicked.	Japan.	1789.	6. 8. H. h.~	Houtt. pfanz. 7. f. 2.	
7 chinénica. DC.	(wh.)	Chinese.	China.	1820.	H. h.~	Retz. obs. 2. n. 53. t. 2.
8 terniflòra. DC.	(wh.)	three-flowered.	—	1826.	H. h.~	
9 Vitálba. DC.	(wh.)	Traveller's Joy.	England.	7. 9. H. h.~	Eng. bot. 612.	
10 virginíâna. DC.	(wh.)	Virginian.	N. America.	1767.	6. 8. H. h.~	Wats. dend. brit. t. 74.	
11 brasiliâna. DC.	(wh.)	Brazilian.	Brazil.	1820.	S. h.~	Deless. ic. sel. 1. t. 1.
12 dioica. DC.	(wh.)	Jamaica.	Jamaica.	1733.	5. 6. S. h.~	Slo. hist. 1. t. 128. f. 1.	
13 americâna. DC.	(wh.)	West-Indian.	W. Indies.	S. h.~	
14 coriacea. DC.	(wh.)	leathery-ld.	N. S. W.	1821.	G. h.~	
15 stenosépala. DC.	(wh.)	narrow-sepaled.	—	1826.	5. 8. G. h.~		
16 aristâta. DC.	(wh.)	awned-anther'd.	—	1812.	5. 8. G. h.~	Bot. reg. 238.	
17 hedysarifòlia. DC.	(wh.)	Hedysarum-ld.	Nepaul.	1819.	7. 2. G. h.~	Bot. reg. 599.	
18 biternâta. DC.	(wh.)	biternate.	Japan.	1825.	7. 9. H. h.~		
19 triternâta. DC.	(wh.)	triternate-ld.	—	1806.	H. h.~	
20 dominica. DC.	(wh.)	St. Domingo.	St. Domingo.	S. h.~	Jac. am. p. 2. t. 261. f. 244.
21 brachiâta. DC.	(wh.)	armed.	C. B. S.	1804.	10. 12. G. h.~	Bot. reg. 97.	
22 grandifòria. DC.	(wh.)	great-flowered.	Sierra Leon.	1822.	S. h.~	
23 dahurica. DC.	(pu.)	Dahurian.	Dahuria.	1822.	6. 8. H. h.~		
24 zanzibârica.	(wh.)	Zanzibar.	Africa.	1826.	S. h.~	
25 angustifòlia. DC.	(wh.)	narrow-leaved.	Siberia.	1787.	5. 9. H. ȝ.	Jacq. ic. 1. t. 104.	
26 diversifòlia. DC.	(pu.)	various-leaved.	—	9. 10. H. h.~	Deless. ic. sel. 1. t. 4.	
27 Viôrna. DC.	(pu.)	leathery-flow'd.	N. America.	1730.	6. 9. H. h.~	Jacq. ecl. 1. t. 32.	
28 cylindrica. DC.	(bl.)	long-flowered.	—	1802.	7. 9. H. h.~	Bot. mag. 1160.	
Viôrna. A.R. 71. nec aliorum.							
29 Símsii.	(pu.)	Sims's.	—	1812.	6. 9. H. h.~	Bot. mag. 1816.	
cordata. B.M. nec aliorum.							
30 reticulâta. DC.	(pu.)	netted.	—	—	H. h.~	Wats. dend. brit. t. 72.	
31 Japonica. DC.	(pu.)	Japan.	Japan.	1826.	H. h.~	
32 integrifòlia. DC.	(bl.)	entire-leaved.	Hungary.	1596.	6. 8. H. ȝ.	Bot. mag. 65.	
β ? angustifòlia.	(bl.)	narrowentire ld.	Siberia.	—	H. ȝ.	
γ ? elongâta. DC.	(bl.)	long-leaved.	—	1823.	—	H. ȝ.	Tratt. tab. 4. t. 178.

RANUNCULACEÆ.

33 ochroleuca. DC.	(st.) silky.	N. America.	1767.	6. 7. H. ȝ.	Bot. mag. 1175.
34 gentianoides. DC.	(wh.) gentian-like.	V. Diem. Isl.	1825.	... G. ȝ.	Deless. ic. sel. 1. t. 5.
35 flórida. DC.	(wh.) large-flowered.	Japan.	1776.	4. 9. H. ȝ.	Bot. mag. 834.
β flóre pleno	(wh.) double-flowered.				H. ȝ.
36 Viticella. DC.	(pu.) smooth-seeded.	Spain.	1569.	6. 9. H. ȝ.	Bot. mag. 565.
α purpurea.	(pu.) purple-flowered.				
β cærulea.	(bl.) blue-flowered.				
γ pulchella. P.S.	(bl.) double-flowered.			H. ȝ.	Chabr. sci. p. 117. f. 5.
37 críspā. B.M.	(pu.) curled-flower'd.	N. America.	1726.	7. 9. H. ȝ.	Bot. mag. 1892.
38 parviflóra. DC.	(wh.) small-flowered.	1822.	6. 8. H. ȝ.	DC. pl. rar. gen. t. 12.
campaniflóra. Lodd. bot. cab. 987. non DC.					
39 cirrhosa. DC.	(pu.) tendriled.	S. Europe.	1596.	3. 4. H. ȝ.	Flor. græc. t. 517.
40 pedicellata.	(st.) pediced.	Majorca.	— H. ȝ.	Bot. mag. 1070.
cirrhosa. B.M. nec aliorum. cirrhosa β . pedicellata. DC.					
41 semitriloba. DC.	(st.) three-lobed.	Spain.	9. 11. H. ȝ.	
42 calycina. H.K.	(st.) evergreen.	Minorca.	1783.	4. 6. H. ȝ.	Bot. mag. 959.
baleárica. DC.					
ATRAG'ENE. W.	ATRAG'ENE.	Polyandria Polygynia. L.			
1 austriaca. B.M.	(bl.) Austrian.	Austria.	1792.	5. 7. H. ȝ.	Bot. rep. 180.
alpina. s.s. Clématis alpina. DC.					
2 sibirica. B.M.	(wh.) Siberian.	Siberia.	1753.	3. 5. H. ȝ.	Bot. mag. 1951.
Clématis sibirica. DC.					
3 ohotensis. s.s.	(wh.) linear-petaled.	1822.	— H. ȝ.	
Clématis ohotensis. DC.					
4 americana. B.M. (pu.) American.		N. America.	1797.	5. 6. H. ȝ.	Bot. mag. 887.
Clématis verticillaris. DC.					
5 occidentalis. s.s.	shining-leaved.	1824.	— H. ȝ.	
NARAV'ELIA. DC.	NARAV'ELIA.	Polyandria Polygynia. L.			
zeylánica. DC. Ceylon.	Ceylon.	1796.	— S. ȝ.	
β Roxburghii. DC. (ye.) Roxburgh's.		Coromandel.	—	S. ȝ.	Roxb cor. 2. t. 188.

Tribus II. ANEMONEÆ.

THALI'CTRUM. DC. MEADOW-RUE. Polyandria Polygynia. L.

1 contortum. DC.	(wh.) crook-seeded.	Siberia.	1796.	6. 7. H. ȝ.	
2 aquilegiólium. DC.	.. Columbine-ld.	Europe.	1731.	5. 7. H. ȝ.	
α álbum.	(wh.) white-stamened.	—	H. ȝ.	Jacq. aust. t. 318.
β atropurpureum. (pu.) purple.	—	—	H. ȝ.	Bot. mag. 1818.
γ formosum. B.M. (pu.) Sabine's.	—	—	H. ȝ.	Bot. mag. 2025.
3 clavatum. DC.	(wh.) club-stamened.	N. America.	1812.	6. 8. H. ȝ.	
4 mexicanum. DC. (gr.) Mexican.		Mexico.	1826.	7. 9. H. ȝ.	Hern. mex. 236. ic.
5 corynéllum. DC.	(wh.) purple-stemm'd.	N. America.	— H. ȝ.	
6 polygamum. s.s. (ye.) polygamous.	—	5. 7. H. ȝ.		
7 revolutum. DC.	(ye.) revolute	1806.	6. 7. H. ȝ.	
pubescens. Ph. non. DC.					
8 dioicum. DC.	(ye.) diecious.	1759.	— H. ȝ.	Deles. ic. sel. 1. t. 8.
9 carolinianum. DC. (ye.) Carolina.	—	— H. ȝ.		
10 purpurascens. DC.	(pu.) dwarf purple.	1699.	— H. ȝ.	
11 petaloideum. DC.	(wh.) Dahurian.	Dahuria.	1799.	— H. ȝ.	Deles. ic. sel. 1. t. 9.
12 alpinum. DC.	(gr.) Alpine.	Britain.	5. 7. H. ȝ.	Eng. bot. 262.
13 foetidum. DC.	(wh.) foetid.	Europe.	1640.	— H. ȝ.	W. et K. hung. 2. t. 174.
14 pubescens. DC.	(ye.) pubescent.	—	H. ȝ.	
15 acutilobum. DC.	(ye.) acute-lobed.	Siberia.	— H. ȝ.	Deles. ic. sel. 1. t. 10.
16 divergens. s.s.	(gr.) divergent.	1820.	6. 7. H. ȝ.	
17 Schweiggeri. s.s. (ye.) Schweigger's.	1826.	— H. ȝ.	
18 calabricum. s.s.	(ye.) Calabrian.	Sicily.	1802.	6. 8. H. ȝ.	Moris. 1.s.9.t. 20. f. 16.
19 squarrósum. DC.	(ye.) squarrose	Siberia.	1806.	— H. ȝ.	
20 sibéricum. DC.	(ye.) Siberian.	1775.	— H. ȝ.	Gärt. fruct. 1. t. 74.
21 minus. DC.	(ye.) lesser.	Britain.	6. 7. H. ȝ.	Eng. bot. t. 11.
22 collinum. DC.	(ye.) hill.	Germany.	1823.	— H. ȝ.	
23 saxatile. DC.	(ye.) rock.	Europe.	5. 7. H. ȝ.	
24 ambiguum. s.s.	(ye.) ambiguous.	Switzerland.	1820.	— H. ȝ.	
25 elatum. DC.	(ye.) tall.	Hungary.	1794.	6. 8. H. ȝ.	Jacq. vind. 3. t. 95.
26 mājus. DC.	(ye.) greater.	England.	— H. ȝ.	Eng. bot. 611.
27 nūtans. DC.	(ye.) nodding.	Italy.	— H. ȝ.	
28 médium. DC.	(ye.) middle.	Hungary.	1789.	— H. ȝ.	Jacq. vind. 3. t. 96.

RANUNCULACEÆ.

3

29	concínnum. DC.	(wh.) neat.	6.	H. ¼.	
30	flexuòsum. Jac.	(ye.) flexuous.	6.	H. ¼.	
31	trig'num. DC.	(ye.) three-styled.	Dahuria.	1823.	—	H. ¼.
32	glaucéscens. DC.	(pu.) loose-flowered.	Russia.	—	6.	H. ¼.
33	galio'ides. DC.	(ye.) Galium-leaved.	Europe.	1816.	5. 7.	H. ¼.
34	angustifòlium. DC.	(ye.) narrow-leaved.	Germany.	1739.	6. 7.	H. ¼.
35	divaricátum. s.s.	(ye.) spreading.	—	—	H. ¼.
36	cynapifòlium. F.	(ye.) Fool's-parsleyd	Siberia.	1821.	—	H. ¼.
37	oligospérnum. F.	(ye.) few-seeded.	—	—	H. ¼.
38	diffusum. Sch.	(ye.) diffuse.	—	—	H. ¼.
39	lúcidum. DC.	(ye.) shining.	Spain.	1739.	5. 7.	H. ¼.
40	rosmarinifòlium. s.s.	(ye.) Rosemary-ld	Italy.	1826.	6. 7.	H. ¼.
41	nigricans. DC.	(ye.) black.	Austria.	1798.	5. 7.	H. ¼.
42	flavum. DC.	(ye.) common.	Britain.	—	—	H. ¼.
43	vaginátum. Desf.	(ye.) sheathed.	Europe.	—	—	H. ¼.
44	símplex. DC.	(ye.) simple-stalked.	Sweden.	1778.	5. 6.	H. ¼.
45	cinereum. DC.	(ye.) ash-coloured.	—	—	H. ¼.
46	glàcum. DC.	(ye.) glaucous	S. Europe.	1798.	6. 7.	H. ¼.
47	rugosum. DC.	(ye.) rugged.	N. America.	1774.	7.	H. ¼.
48	ranunculinum. DC	(ye.) Ranunculus-ld.	1806.	6. 7.	H. ¼.
49	tuberòsum. DC.	(wh.) tuberous-rooted.	Spain.	1713.	6.	H. ¼.
50	anemonoides. DC.	(wh.) Anemone-like.	N. America.	1768.	4. 5.	H. ¼.
	Anemone thalictroides. W.					
	β flòre plèno.	double-flowered.	1822.	—	H. ¼.
PULSATILLA. W.en. PASQUE-FLOWER. Polyandria Polygynia.						
1	vernális. s.s.	(pu.) spring.	N. Europe.	1752.	4.	H. ¼.
	Hallèri. s.s.	(bl.) Haller's.	Switzerland.	1816.	4. 5.	H. ¼.
3	cérnua. s.s.	(bl.) drooping.	Japan.	1806.	4. 6.	H. ¼.
4	pátnes. s.s.	(va.) spreading.	Siberia.	1752.	6. 7.	H. ¼.
	β ochroleuca.	(st.) straw-coloured.	—	5. 7.	H. ¼.
5	vulgáris. s.s.	(bl.) common.	England.	—	—	H. ¼.
	β rúbra. DC.	(re.) red-flowered.	Europe.	—	—	H. ¼.
6	intermédia.	(bl.) long-petal'd.	Switzerland.	1820.	—	H. ¼.
	Anemone intermédia. Schult. longipétala. Schleich.					
7	praténis. s.s.	(pu.) meadow.	Germany.	1731.	5.	H. ¼.
8	obsolèta.	(bl.) pale-flowered.	—	—	H. ¼.
9	albâna. s.s.	(wh.) white-flowered.	Caucasus.	1827.	—	H. ¼.
10	dahurica. DC.	(fl.) Dahirian.	Dahuria.	1823.	—	H. ¼.
11	alpina. s.s.	(wh.) Alpine.	Europe.	1658.	7.	H. ¼.
12	mícrantha.	(wh.) small-flowered.	1827.	—	H. ¼.
	Anemone alpina. Jacq. alpina β micránttha. DC.					
13	apiifòlia.	(ye.) Parsley-leaved.	—	—	H. ¼.
	Anemone alpina δ flavescentis. DC.					
14	sulphurea.	(st.) sulphur-coloured.	—	—	H. ¼.
	Anemone sulphurea. L. apiifolia. w.					
ANEM'ONE. W.en. ANEM'ONE. Polyandria Polygynia. L.						
1	capénsis. DC.	(pu.) Cape.	C.B.S.	1795.	3. 7.	G. ½.
	Atragène capénsis. W.					
2	coronária. DC.	(va.) Poppy.	Levant.	1596.	1. 12.	H. ¼.
	β flòre-plèno.	(va.) double flowered.	—	—	
3	pavonina. DC.	(re.) acute-petaled.	—	4. 5.	H. ¼.
	α coccinea.	(sc.) scarlet.	—	—	H. ¼.
	β lilacina.	(li.) lilac.	—	—	H. ¼.
	γ flòre-plèno.	(sc.) double scarlet.	—	—	H. ¼.
4	stellata. DC.	(pu.) star.	Italy.	1597.	—	H. ¼.
	horténsia. B.M. 123.					
	α versicolor.	(v.) various-coloured.	—	—	H. ¼.
	β purpurea.	(pu.) purple.	—	—	H. ¼.
5	palmata. DC.	(ye.) palmated.	Portugal.	1597.	5. 6.	H. ¼.
	β flòre-plèno.	(ye.) double-flowered.	—	—	H. ¼.
6	parvifòra. DC.	(wh.) small-flowered.	Labrador.	1823.	3. 5.	H. ¼.
	cuneifòlia. Ph.					
7	caroliniána. DC.	(pu.) slender.	Carolina.	—	5.	H. ¼.
	tenella. Ph.					
8	apennina. DC.	(bl.) blue mountain.	England.	3. 4.	H. ¼.
9	coerulea. DC.	(bl.) blue.	Siberia.	1827.	—	H. ¼.
10	uralénsis. DC.	(bl.) Uralian.	—	—	H. ¼.
11	baldénsis. DC.	(wh.) Strawberry-like.	Switzerland.	1792.	5.	H. ¼.
	fragifera. Jacq.			B 2		

RANUNCULACEÆ.

12 nemorðsa. DC.	(wh.) wood.	Britain.	3. 5. H. 4.	Eng. bot. 355.
β florē-plēno.	(wh.) double-flowered.	—	—	H. 4.	Debry.fl.nov.t.29 et 56.
γ quinquefolia. DC.	(w.) five-leaved.	N. America.	1812.	— H. 4.	
13 Fischeriāna. DC.	(wh.) Fischer's.	Siberia.	1827.	4. 5. H. 4.	
14 lancifolia. DC.	(wh.) lance-leaved.	N. America.	1823.	5. 7. H. 4.	Lob. ic. 281. f. 1.
15 trifolia. DC.	(wh.) three-leaved.	France.	1597.	4. 5. H. 4.	Eng. bot. 1484.
16 ranunculoides. DC.	(ye.) yellow wood.	England.	3. 4. H. 4.	Deless. ic. sel. 1. t. 15.
17 refléxa. DC.	(wh.) reflex-flowered.	Siberia.	1824.	4. 5. H. 4.	Bot. mag. 54.
18 sylvestris. DC.	(wh.) Snowdrop.	Europe.	1596.	4. 5. H. 4.	— 2167.
19 álba. DC.	(wh.) five-sepaled.	Dahuria.	1818.	4. 5. H. 4.	
ochoténensis. F.					
20 virginiana. DC.	(wh.) Virginian.	N. America.	1722.	5. 6. H. 4.	Herm. par. t. 18.
21 pensylvánica. DC.	(wh.) Pennsylvanian.	—	1766.	— H. 4.	
22 dichótoma. DC.	(wh.) forked.	—	1768.	— H. 4.	Lin. f. dec. 29. t. 15.
23 narcissiflora. DC.	(wh.) Narcissus-fl'd.	Europe.	1773.	5. H. 4.	Bot. mag. 1120.
24 umbellata. DC.	(wh.) umbel flower'd.	Levant.	1824.	— H. 4.	Deless. ic. sel. 1. t. 18.
25 sibirica. DC.	(wh.) Siberian.	Siberia.	1804.	6. H. 4.	
HEPATICÆ. DC.	HEPATICA. Polyandria Polygynia. L.				
1 acutiloba. DC.	(bl.) acute-lobed.	N. America.	1826.	3. 4. H. 4.	Bot. reg 387.
2 americana. DC.	(bl.) American.	—	1819.	3. 4. H. 4.	Eng. bot. t. 51.
3 triloba. DC.	(va.) three-lobed.	Europe.	1573.	2. 4. H. 4.	
α cærulea.	(bl.) blue.				
β cæruleo-plena.	(bl.) double-blue.				
γ rùbra.	(re.) red.	—	—	—	Bot. mag. 10.
δ rùbro-plena.	(re.) double-red.	—	—	—	
ε álba.	(wh.) red-anthered white.	—	—	—	
ζ nivea.	(wh.) snowy-white.	—	—	—	
HYDRA'STIS. DC.	YELLOW-ROOT. Polyandria Polygynia. L.				
canadensis. DC.	(wh.) Canadian.	Canada.	1759	5. 6. H. 4.	Pict. hort. par. 37. t. 17.
KNOWLTONIA. DC.	KNOWLTONIA. Polyandria Polygynia. L.				
1 rígida. DC.	(gr.) thick-leaved.	C. B. S.	1780.	3. 5. G. 4.	Vent. malm. p. 22. t. 22.
2 vesicatória. DC.	(gr.) blistering.	—	1691.	2. 4. G. 4.	Bot. mag. 775.
3 gráclis. DC.	(gr.) slender.	—	1820.	— G. 4.	Deles. ic. sel. 1. t. 19.
4 hirsuta. DC.	(gr.) hairy-leaved.	—	—	— G. 4.	Burm. afr. t. 51.
5 daucifolia. DC.	(gr.) Carrot-leaved.	—	1822.	— G. 4.	
AD'ONIS. DC.	AD'ONIS. Polyandria Polygynia. L.				
1 autumnalis. DC.	(sc.) Pheasant's-eye. Britain.	5. 10. H. ⊙.	Eng. bot. 308.	
2 flava. DC.	(ye.) yellow.	France.	6. 7. H. ⊙.	Tab. ic. 790. f. 1.
3 micrantha. DC.	(ye.) small-flowered.	S. France.	1826.	5. 6. H. ⊙.	
4 citrina. DC.	(st.) Citron-colour'd.	—	1822.	5. 6. H. ⊙.	Jac. aust. 4. t. 355.
5 flámmea. DC.	(fl.) flame-coloured.	Austria.	1800.	6. 7. H. ⊙.	Weinm. phyt. t. 27.
6 æstivális. DC.	(sc.) tall.	S. Europe.	1629.	— H. ⊙.	
miniata. Jac. aust. t. 354.					
7 vernalis. DC.	(ye.) vernal.	Europe.	1629.	3. 4. H. 4.	Bot. mag. 134.
8 apennina. w. en.	(ye.) Apennine.	—	— H. 4.	Mentz. pug. t. 3. f. 1.
vernális β Mentzelii. DC.					
9 pyrenáica. DC.	(ye.) Pyrenean.	Pyrenees.	1820.	6. 7. H. 4.	Deless. ic. sel. 1. t. 21.
10 sibirica. s.s.	(ye.) Siberian.	Siberia.	1827.	4. 5. H. 4.	
11 davúrica. s.s.	(ye.) Davurian.	—	—	— H. 4.	Reich. ic. t. 321.
12 distorta. T.P.	(ye.) distorted.	Naples.	—	— H. 4.	Ten. fl. med. p. 448. t. 1.

Tribus III. RANUNCULEÆ.

MYOS'URUS. DC.	MOUSE-TAIL. Pentandria Polygynia. L.				
mínimus. DC.	(gr.) small.	Britain.	4. 5. H. ⊙.	Eng. bot. 435.
CERATOCE'PHALUS.	CERATOCE'PHALUS. Pentandria Polygynia.				
1 falcatus. DC.	(ye.) sickle-seeded.	S. Europe.	1739.	5. H. ⊙.	Jac. aust. t. 48.
2 orthoceras. DC.	(ye.) straight-seeded.	Siberia.	1823.	— H. ⊙.	Deles. ic. sel. 1. t. 23.
RANU'NCULUS. DC.	CROW-FOOT. Polyandria Polygynia. L.				
1 hederaceus. DC.	(wh.) Ivy-leaved.	Britain.	5. 8. H.w. 4.	Eng. bot. 2003.
2 tripartitus. DC.	(wh.) three-parted.	—	— H.w. 4.	DC. ic. g. rar. 1. t. 49.
β obtusiflorus. DC.	(w.) blunt-flowered.	—	—	— H.w. 4.	Pet. eng. herb. t. 39.f.1.
3 aquátillis. DC.	(wh.) water.	—	4. 8. H.w. 4.	Eng. bot. 101.
α heterophyllus. DC.	(w.) various-leaved.	—	— H.w. 4.	Tab. ic. 54. f. 2.
β peltatus. DC.	(wh.) peltate-leaved.	—	— H.w. 4.	Lob. ic. 2. p. 35. f. 2.

RANUNCULACEÆ.

4	pantothrix. DC. (wh.)	divided-leaved. Britain.	4. 8. H.w. 4.	Lob. ic 791. f. 1.
α	capillaceus. DC. (wh.)	fine-leaved.	— H.w. 4.	Pet. eng. herb. t. 39.f.4.
β	cæspitosus. DC. (wh.)	tufted.	— H.w. 4.	_____ t. 39. f. 3.
γ	peucedanifolius. DC. (wh.)	Fennel-l'd.	— H.w. 4.	_____ t. 39. f. 2.
	flutiilis. w.				
5	rutæfolius. DC. (wh.)	Rue-leaved.	Austria.	1759. 5. 7. H. 4.	Jac. coll. 1. t. 6 et 7.
6	glaciælis. DC. (wh.)	two-flowered.	Lapland.	1775. 6. 8. H. 4.	Flor. dan. t. 19.
7	Séguier. DC. (wh.)	Seguier's.	Switzerland.	1820. — H. 4.	Vill. dauph. 4. t. 49.
8	alpétris. DC. (wh.)	Alpine.	Scotland.	Eng. bot. 2390.
9	crenatus. DC. (wh.)	crenate-leaved.	Hungary.	1822. — H. 4.	W. et K. hung. 1. t. 10.
10	aconitifolius. DC. (wh.)	Aconite-leaved.	Europe.	1596. 5. 6. H. 4.	J. Ger. hist. 951. f. 4.
	α humiliis. DC. (wh.)	dwarf.	— H. 4.	Lob. adv. 300. f. 2.
	β crassicaulis. DC. (wh.)	thick-stalked.	— H. 4.	
11	platanifolius. w. (wh.)	Plane-tree-l'd.	Germany.	1769. 6. 7. H. 4.	Flor. dan. t. 111.
	β flôre-pleno.	double-flowered.	— H. 4.	Bot. mag. 204.
	aconitifolius. B.M. 204.				
12	lacerus. DC. (wh.)	torn-leaved.	Piedmont.	Bell. act. taur. 5. t. 8.
13	pallidus. DC. (st.)	pale-yellow.	Hybrid.	Schrank. hort. mont. 57.
14	pyrenæus. DC. (wh.)	Pyrenean.	Pyrenees.	Deles. ic. sel. 1. t. 27.f.B.
	β bupleurifolius. DC. (wh.)	one-flowered.	Jac. misc. 1. t. 18. f. 1.
	γ plantagineus. DC. (wh.)	many-flowered.	All ped.n. 1445.t.76.f.1.
15	angustifolius. DC. (wh.)	narrow-leaved.	1822. — H. 4.	Deles. ic. sel. 1. t. 27.f.C.
16	amplexicaulis. DC. (w.)	Plantain-leav'd.	1633. 4. 5. H. 4.	Bot. mag. 266.
17	parnassifolius. DC. (w.)	Parnassia-leavd.	1769. 6. 7. H. 4.	_____ 386.
18	gramineus. DC. (ye.)	grass-leaved.	Wales.	Eng. bot. 2306.
	β flôre-pleno.	double-flowered.	Lob. ic. 671. f. 2.
19	bupleurioides. DC. (ye.)	Hare's-ear-l'd.	Portugal.	1826. — H. 4.	
20	Lingua. DC. (ye.)	great-spear-wort.	Britain.	Eng. bot. 100.
21	Flâmmula. DC. (ye.)	less.spear-wort.	_____ 387.
	β serrata. DC. (ye.)	saw-leaved.	Lob. ic. 670. f. 2.
	γ ovata. DC. (ye.)	oval-leaved.	
22	réptans. DC. (ye.)	leastspear-wort.	Flor. dan. t. 108.
23	filiformis. M.	filiform.	N. America.	1816. — H.w. 4.	
24	uliginosus. DC. (ye.)	marsh.	Teneriffe.	1826. 6. 7. H.w. \odot .	Deles. ic. sel. 1. t. 29.
25	bonariensis. DC. (ye.)	Buenos Ayres.	S. America.	1819. — H.w. \odot .	W. et K. pl. hung. 1.t.45.
26	polypillus. DC. (ye.)	many-leaved.	Hungary.	Vill. dauph. 4. t. 49.
27	ophioglossifolius. (ye.)	fistulous-stalk'd.	S. Europe.	1822. 6. H.w. \odot .	
28	nodiflorus. DC. (ye.)	knot-flowered.	France.	1714. 5. 7. H.w. \odot .	Bot. mag. 2171.
	β dentatus. DC. (ye.)	toothed.	— H.w. \odot .	Jac. vind. 3. t. 31.
29	salsuginosus. DC. (ye.)	Russian.	Russia.	1821. — H. 4.	
	<i>ruthenicus</i> . Jac. vind. 3. t. 31.				
30	Cymbalariae. Ph. (ye.)	Ivy-like.	N. America.	1823. — H.w. 4.	
31	bullatæ. DC. (ye.)	blistered.	S. Europe.	1640. 5. 8. H. 4.	Clus. hist. 1. f. 2.
	α parviflorus. DC. (ye.)	small-flowered.	— H. 4.	_____ 1. f. 1.
	β grandiflorus. DC. (ye.)	large-flowered.	— H. 4.	Corn. can. 95. ic.
	γ flôre-pleno.	double-flowered.	— H. 4.	Barr. ic. t. 581.
32	charophyllos. DC. (ye.)	Chervil-leaved.	
	α vulgaris. DC. (ye.)	common.	Flor. græc. t. 521.
	β gregarius DC. (ye.)	villous.	Desf. ic. pict. h. par. t. 74.
33	adscendens. DC. (ye.)	ascending.	Portugal.	1826. 5. 6. H. 4.	
34	millefoliatu. DC. (ye.)	numerous-l'd.	Sicily.	1824. 4. 7. H. 4.	Bot. mag. 2229.
35	fumariaefolius. DC. (ye.)	Fumitory-leav'd.	Jacq. aust. t. 222.
36	oxyspermus. DC. (ye.)	awned-seeded.	Tauria.	1822. 5. 6. H. 4.	Swt. br. fl. gar. 94.
37	pedatus. DC. (ye.)	pedate.	Hungary.	1806. 5. 6. H. 4.	Besl. eyst. 1. t. 13. f. 1.
38	ill'yricus. DC. (ye.)	Illyrian.	Europe.	1596. — H. 4.	DC. ic. gall. rar. t. 50.
39	monspeliacus. DC. (ye.)	Montpelier.	Montpelier.	1816. — H. 4.	Dard. trait. ren. 1. t. 6.
	α angustilobus. DC. (y.)	narrow-lobed.	Mill. ic. t. 216.
	β cuneatus. DC. (ye.)	wedge-lobed.	Swt. flor. guid. 61.
	γ rotundifolius. DC. (y.)	round-leaved.	_____ 54.
40	asiaticus. DC. (va.)	garden.	Levant.	1596. — H. 4.	_____ 12.
	<i>A vulgaris</i> . DC. (va.)	common.	_____ 28.
	<i>a pectus</i> . S.F.G. (r.w.)	Quillafla.	_____ 68.
	β incomparabilis. (r.w.)	Rose incomparable.	_____ 6.
	γ Victoriañus. (bh.)	Princess Alexandrina Victoria.	_____ 34.
	δ mixtus. (r.y.)	Mélange des beautés.	_____ 37.
	ϵ Juliusianus. (p.y.)	Julius.	_____ 18.
	ζ Galitzini. (p.y.)	Prince Galitzin.	
	η Douglasii. (p.y.)	Gadwin Douglas.	
	θ aureus. (ye.)	Beroth.	
	ι Burnsii. (p.w.)	Burns the Poet.	

RANUNCULACEÆ.

κ áter.	(d.p.) <i>Vereatre.</i>	5. 6.	H. ȝ.	Swt. flor. guid. 43.
λ <i>hæmánthus.</i>	(cr.) <i>Xanthus.</i>	H. ȝ.	78.
μ <i>Agrícoli.</i>	(s.p.) <i>Agricola.</i>	H. ȝ.	73.
ν <i>Clarenceánum.</i>	(s.p.) <i>Duke of Clarence.</i>	H. ȝ.	83.
ξ <i>Carlosiánum.</i>	(s.p.) <i>Carlos.</i>	H. ȝ.	90.
<i>B sanguineus.</i> DC. (va.)	<i>Turban.</i>	Levant.	1596.	Lob. ic. 672. f. 2.
<i>C tenuilobus.</i> DC. (va.)	slender-lobed.	H. ȝ.	Debr. flor. nov. t. 31.
α <i>aterrimus.</i>	(da.) <i>Viola le vrai noir.</i>	H. ȝ.	Swt. flor. guid. 60.
β <i>Nomiasi.</i>	(pu.) <i>Nomias.</i>	H. ȝ. 24.
γ <i>variegátus.</i>	(w.r.) <i>Oeillet parfait.</i>	H. ȝ. 40.
δ <i>cárus.</i>	(w.r.) <i>Cara.</i>	H. ȝ. 86.
41 <i>créticus.</i> DC.	(ye.) <i>Cretan.</i>	Candia.	1659.	4. 5.	H. ȝ.
42 <i>cortusifolius.</i> DC.	(ye.) <i>Cortusa-leaved.</i>	Canaries.	1826.	4. 6.	H. ȝ.
43 <i>Thòra.</i> DC.	(ye.) kidney-leaved.	AlpsEurope.	1710.	5. 6.	H. ȝ.
44 <i>scutátus.</i> DC.	(ye.) shield-leaved.	Hungary.	1822.	4. 5.	H. ȝ.
45 <i>brevifolius.</i> DC.	(ye.) short-leaved.	Naples.	1826.	H. ȝ.
46 <i>hybridus.</i> DC.	(ye.) hybrid.	Austria.	1822.	6. 7.	H. ȝ.
47 <i>auricomus.</i> DC.	(ye.) Goldilocks.	Britain.	3. 5.	H. ȝ.
48 <i>cassíbicus.</i> DC.	(ye.) Cassubian.	Siberia.	1794.	6. 7.	H. ȝ.
49 <i>abortivus.</i> DC.	(ye.) three-flowered.	N. America.	1713.	5. 8.	H. ȝ.
50 <i>sclerátus.</i> DC.	(ye.) Celery-leaved.	Britain.	H.w. ⊙.
51 <i>lappónicus.</i> DC.	(ye.) Lapland.	Lapland.	1827.	4. 6.	H. ȝ.
52 <i>hyperbóreus.</i> DC.	(ye.) northern.	Siberia.	1822.	6. 8.	H. ȝ.
53 <i>frigidus.</i> DC.	(ye.) frigid.	1826.	H. ȝ.
54 <i>nívialis.</i> DC.	(ye.) snowy.	Lapland.	1775.	H. ȝ.
55 <i>Sabini.</i> s.s.	(ye.) Sabine's.	N. America.	1827.	H. ȝ.
56 <i>árticus.</i> s.s.	(ye.) arctic.	H. ȝ.
57 <i>Púrshii.</i> s.s.	(ye.) Pursh's.	H. ȝ.
58 <i>affinís.</i> s.s.	(ye.) likened.	H. ȝ.
59 <i>montáñus.</i> DC.	(ye.) mountain.	Austria.	H. ȝ.
60 <i>Villárisi.</i> DC.	(ye.) Villars's.	France.	5. 7.	H. ȝ.
61 <i>Gouàni.</i> DC.	(ye.) Gouan's.	Pyrenees.	1818.	H. ȝ.
62 <i>ácris.</i> DC.	(ye.) upright.	Britain.	6. 7.	H. ȝ.
β <i>flòre pleno.</i>	(ye.) double-flowered.	H. ȝ.
γ <i>svyláticus.</i> DC.	(ye.) wood.	H. ȝ.
δ <i>multifidus.</i> DC.	(ye.) multifid.	H. ȝ.
63 <i>Brútius.</i> DC.	(ye.) Neapolitan.	Naples.	1828.	H. ȝ.
64 <i>Stevéni.</i> DC.	(ye.) Steven's.	Poland.	1825.	H. ȝ.
65 <i>caucásicus.</i> DC.	(ye.) Caucasian.	Caucasus.	6. 8.	H. ȝ.
66 <i>rúfulus.</i> DC.	(ye.) rufous-haired.	Portugal.	H. ȝ.
67 <i>polyánthemos.</i> DC.	(ye.) many-flowered.	N. Europe.	1596.	5. 6.	H. ȝ.
68 <i>nemorósus.</i> DC.	(ye.) wood.	Switzerland.	1818.	5. 7.	H. ȝ.
69 <i>Breygniánius.</i> s.s.	(ye.) Breygnius's.	H. ȝ.
$\ddot{\alpha}$ <i>reáreus.</i> Schl. <i>villoso</i> s.	St. Am. fl. ag. 227. bon. t. 5.
70 <i>lanuginósus.</i> DC.	(ye.) woolly-leaved.	S. Europe.	1683.	6. 7.	H. ȝ.
71 <i>obtusifolius.</i> DC.	(ye.) blunt-leaved.	Spain.	1825.	H. ȝ.
72 <i>tuberósus.</i> DC.	(ye.) tuberous.	Pyrenees.	1824.	5. 7.	H. ȝ.
73 <i>napellifolius.</i> DC.	(ye.) Napellus-leav'd.	Greece.	1825.	H. ȝ.
74 <i>platyspermus.</i> DC.	(ye.) flat-seeded.	Siberia.	1827.	H. ȝ.
75 <i>disséctus.</i> DC.	(ye.) cut-leaved.	1823.	H. ȝ.
76 <i>rèpens.</i> DC.	(ye.) creeping.	Britain.	5. 9.	H. ȝ.
β <i>flòre pleno.</i>	(ye.) double-flowered.	H. ȝ.
γ <i>glábratus.</i> DC.	(ye.) shining-leaved.	H.w. ȝ.
77 <i>lappáceus.</i> DC.	(ye.) New Holland.	N. S. W.	1822.	G. ȝ.
78 <i>plebeius.</i> DC.	(ye.) plebeian.	1823.	5. 7.	G. ȝ.
79 <i>hirtus.</i> DC.	(ye.) hairy.	N. Zealand.	H. ȝ.
80 <i>pensylvánicus.</i> DC.	(ye.) Pensylvanian.	N. America.	1785.	6. 7.	H. ȝ.
<i>canadénis.</i> Jac.	Jacq. ie. 1. t. 105.
81 <i>fasciculáris.</i> DC.	(ye.) fascicled.	1823.	4. 5.	H. ȝ.
82 <i>marylándicus.</i> DC.	(ye.) Maryland.	1811.	5. 7	H. ȝ.
83 <i>carolinianus.</i> DC.	(ye.) Carolina.	1826.	5. 7.	H. ȝ.
84 <i>tomentósus.</i> DC.	(ye.) tomentose.	1820.	H. ȝ.
85 <i>bulbósus.</i> DC.	(ye.) bulbous.	Britain.	4. 6	H. ⊙.
β <i>flòre pleno.</i>	(ye.) double-flowered.	Eng. bot. 515.
86 <i>bracteátus.</i> Sch.	(ye.) bracted.	Switzerland.	1817.	Lob. ic. 666. f. 2.
87 <i>Philonótis.</i> DC.	(ye.) pale hairy.	Britain.	6. 10.	H. ⊙.
α <i>hirsútus.</i> E.B.	(ye.) hairy upright.	Curt. lond. 2. t. 40.
β <i>intermedius.</i> DC.	(ye.) intermediate.	France.	Eng. bot. t. 1504.
γ <i>párvulus.</i> DC.	(ye.) little upright.	England.	Cöl. ecphr. t. 316.
88 <i>tuberculátus.</i> DC.	(ye.) tubercled.	1821.	5. 6.	H. ȝ.

RANUNCULACEÆ.

7

89	Hornemánni. DC. (ye.)	Hornemann's. N. America.	1820.	5. 7. H. ¼.		
	tuberösus. Horn. non Lapeyr.					
90	arvénensis. DC.	(ye.) corn.	Britain.	—	H. ⊖.	Eng. bot. 135.
91	muricátus. DC.	(ye.) prickly seeded.	S. Europe.	1683.	7. 8. H. ⊖.	Lam. ill. t. 498.
92	echinátus. v.	(ye.) American.	N. America.	—	6. 7. H. ⊖.	Vent. cels. t. 73.
93	parviflórus. DC.	(ye.) small-flowered.	England.	—	5. 6. H. ⊖.	Eng. bot. 120.
94	trilóbis. DC.	(ye.) three-lobed.	S. Europe.	1824.	5. 7. H. ⊖.	Desf. atl. 1. t. 113.
95	sessíflórus. DC.	(ye.) sessile-flowered.	N. S. W.	1822.	—	H. ⊖.
96	laciniátus. DC.	(ye.) jagged.	Transylvania.	1826.	6. 7. H. ¼.	
97	crassícaulis. H.	(ye.) thick-stemmed.	—	—	—	H. ¼.
FIC'ARIA. DC.		PILEWORT.	Polyandria Polygynia. L.			
ranunculóides. DC.	(ye.) vernal.	Britain.	—	3. 5. H. ¼.	Eng. bot. 584.	
vérna. p.s. Ranúnculus Ficária. w.						
β flòre-pléno.	(ye.) double-flowered.	—	—	—	H. ¼.	

Tribus IV. HELLEBOREÆ.

CA'LTHA. DC.		MARSH-MARYGOLD.	Polyandria Polygynia L.			
1 palústris. DC.	(ye.) common.	Britain.	—	3. 5. H. w. ¼.		Eng. bot. 506.
β flòre-pléno.	(ye.) double-flowered.	—	—	—	H. w. ¼.	Tab. ic. 751. f. 1.
2 minor. Mill.	(ye.) small.	—	—	—	H. w. ¼.	— 750. f. 2.
3 radicans. DC.	(ye.) creeping.	Scotland.	—	—	H. w. ¼.	Linn. trans. 8. t. 17.
4 intégríma. DC.	(ye.) entire-leaved.	N. America.	1827.	5. 7. H. w. ¼.		
5 parnassiífolia. DC.	(ye.) Parnassia-leavd.	—	1825.	—	H. w. ¼.	
	ficarioides. Ph.					
6 ártica. s.s.	(ye.) northern.	—	1827.	—	H. w. ¼.	
7 flabellifólia. DC.	(ye.) fan-leaved.	—	1818.	7. 8. H. w. ¼.	Pur. fl. am. sept. 2. t. 17.	
8 nátans. DC.	(wh.) floating.	Siberia.	1823.	4. 5. H. w. ¼.	Gmel. sib. 4. t. 82.	
TRO'LLIUS. DC.		GLOBE-FLOWER.	Polyandria Polygynia. L.			
1 europæus. DC.	(ye.) European.	Britain.	—	5. 6. H. ¼.	Eng. bot. t. 28.	
	a altissimus. DC.	(ye.) tall.	—	—	H. ¼.	
β hùmilis. DC.	(ye.) ducarf.	—	—	—	H. ¼.	
2 napellifólius. DC.	(ye.) intermediate.	—	—	—	H. ¼.	
3 caucásicus. DC.	(ye.) Caucasian.	Caucasus.	—	—	H. ¼.	
4 pátlus. DC.	(ye.) spreading.	Siberia.	—	5. 7. H. ¼.		
5 Ledebourii. s.s.	(ye.) Ledebour's.	—	1827.	—	H. ¼.	
6 asiáticus. DC.	(ye.) Asiatic.	—	1759.	—	H. ¼.	
7 americanus. DC.	(ye.) American.	N. America.	1805.	5. 7. H. ¼.		
	láxus. Ph.	Gaissénia vérna. Raf.				
ERA'NTHIS. DC.		WINTER-ACONITE.	Polyandria Polygynia. L.			
1 hyémalis. DC.	(ye.) common.	Europe.	1596.	1. 3. H. ¼.	Bot. mag. 3.	
2 sibirica. DC.	(ye.) Siberian.	Siberia.	1824.	2. 4. H. ¼.		
HELLE'BORUS. DC.		HELLEBORE.	Polyandria Polygynia. L.			
1 niger. DC.	(wh.) Christmas-Rose.	Europe.	1596.	1. 3. H. ¼.		
	a latifolius.	(wh.) broad-leaved.	—	—	H. ¼.	Bot. mag. 8.
β angustifolius.	(wh.) narrow-leaved.	—	—	—	H. ¼.	
2 purpurascens.	(pu.) purplish.	Hungary.	1817.	3. 4. H. ¼.	Pl. rar. hung. 2. t. 101.	
3 Bocconi. T. P.	(gr.) Boccone's.	Sicily.	1826.	—	Bocc. mus. 2. t. 11. f. 2.	
4 odórus. DC.	(gr.) sweet-scented.	Hungary.	—	—		
5 víridis. DC.	(gr.) green-flowered.	Britain.	—	4. 6. H. ¼.	Eng. bot. 200.	
6 atrorúbens. DC.	(pu.) dark-red.	Hungary.	1822.	1. 3. H. ¼.	Pl. rar. hung. 3. t. 271.	
7 dumetórum. DC.	(gr.) bush.	—	1817.	3. 4. H. ¼.	Swt. br. fl. gar. 109.	
8 feétidus. DC.	(gr.) Bear's-foot.	England.	—	2. 4. H. ¼.	Eng. bot. 613.	
9 lívidus. DC.	(br.) three-leaved.	Corsica.	1710.	1. 5. H. ¼.	Bot. mag. 72.	
	β integrilóbis. DC. (br.)	entire-lobed.	—	—	H. ¼.	
CO'PTIS. DC.		CO'PTIS.	Polyandria Polygynia. L.			
	trifolia. DC.	(wh.) three-leaved.	N. America.	1782.	6. 7. H. ¼.	Lodd. bot. cab. t. 173.
ISOP'YRUM. DC.		ISOP'YRUM.	Polyandria Polygynia. L.			
1 thalictróides. DC.	(wh.) Meadow-Ruel'd.	Italy.	1759.	3. 4. H. ¼.	Jac. aust. 2. t. 105.	
2 fumarióides. DC.	(wh.) Fumitory-l'd.	Siberia.	1741.	6. 7. H. ⊖.	Schkuhr hand. 2. t. 153.	
3 grandiflórum. F.	(wh.) large-flowered.	Altay.	1804.	—	H. ¼.	

GARIDE'LLA. DC.	GARIDE'LLA. Decandria Trigynia. L.	Nigellástrum. DC. (wh.)	Nigella-leaved. S. Europe. 1736.	6. 7. H. ⊙	Bot. mag. 1266.
NIGE'LLA. DC.	FENNEL-FLOWER. Polyandria Pentagynia. L.				
1 corniculàta. DC.	(st.) horned.	1822.	6. 9. H. ⊙.		Bot. mag. 1264.
2 orientális. DC.	(ye.) yellow. Syria. 1699.	—	— H. ⊙.	—	1265.
3 hispánica. DC.	(bl.) Spanish. Spain. 1629.	—	— H. ⊙.	Bull. herb. t. 126.	
4 arvén̄s. DC.	(wh.) field. Europe. 1683.	—	— H. ⊙.	Flor. græc. t. 511.	
5 satív̄a. DC.	(wh.) small. Egypt. 1548.	—	— H. ⊙.	Math. com. 581. f. IV.	
β crética. DC.	(wh.) Cretan. Crete. —	—	— H. ⊙.	Lob. ic. 741. f. 1.	
γ citrīna. DC.	(wh.) yellow-seeded. —	—	— H. ⊙.	Bot. mag. 22.	
6 damascéna. DC.	(wh.) common. S. Europe. 1570.	—	— H. ⊙.	Clus. hist. 2. p. 208. f. 1.	
β flôre-plêno.	(wh.) double-flowered. —	—	— H. ⊙.		
7 coartát̄a. DC.	(wh.) dwarf. —	1793.	— H. ⊙.		
AQUIL'EGIA. DC.	COLUMBINE. Polyandria Pentagynia. L.				
1 vulgāris. DC.	(va.) common. Britain.	5. 7.	H. 4.	Eng. bot. 297.	
β corniculàta. DC. (va.)	downward-spur'd	—	— H. 4.	Barri. ic. 614. 615. 616. &c.	
γ invérsa. DC.	upward-spurred.	—	— H. 4.	— 613.	
δstellat̄a. DC.	(va.) starry.	—	— H. 4.	— 619. 620. &c.	
• degén̄er. DC.	(gr.) green-flowered.	—	— H. 4.	— 608.	
2 viscósa. DC.	(bl.) clammy. S. Europe. 1752.	5. 6.	H. 4.	Gouan. ill. t. 19.	
3 sibírica. DC.	(bl.) Siberian. Siberia.	—	— H. 4.	Deles. ic. sel. 1. t. 47.	
4 glandulosâ. F.	(bl.) glandulous. Altay. 1818.	—	— H. 4.		
a díscolor. DC.	(b.w.) two-coloured. —	—	— H. 4.		
β cóncolor. DC.	(bl.) one-coloured. —	—	— H. 4.		
5 alpina. DC.	(bl.) Alpine. Switzerland. 1731.	—	— H. 4.		
6 pyrenaica. DC.	(bl.) Pyrenean. Pyrenees. 1821.	—	— H. 4.		
7 canadénsis. DC.	(re.) Canadian. N. America. 1640.	4. 5.	H. 4.	Swt. br. fl. gar. 3. t. 218.	
8 formòsa. F.	(re.) handsome. Kamtschatka. 1824.	—	— H. 4.	Trev. delph. 23. t. 2.	
9 viridiflòra. DC.	(gr.) green-flowered. Siberia. 1780.	5. 6.	H. 4.	Bot. mag. 246.	
10 dahúrica. DC.	(pu.) Dahurian. Dahuria. 1821.	—	— H. 4.		
11 Fischeriâna. DC.	(pu.) Fischer's. —	—	— H. 4.	Jac. ic. 1. t. 102.	
12 atropurpurea. DC.	(pu.) dark-purple. Siberia.	5. 6.	H. 4.	Deless. ic. sel. 1. t. 49.	
13 hy'bida. DC.	(p.y.) hybrid. —	—	— H. 4.		
DELPHI'NIUM. DC.	LARKSPUR. Polyandria Trigynia. L.			Botan. regist. 922.	
1 Ajácis. DC.	(va.) Rocket. Tauria. 1573.	6. 9.	H. ⊙.	Bot. mag. 1221.	
β flôre-plêno.	(va.) double-flowered. —	—	— H. ⊙.		
2 Consolida. DC.	(va.) field. England.	—	— H. ⊙.		
β flôre-plêno.	(va.) double-flowered. —	—	— H. ⊙.		
3 pubescens. DC.	(bl.) pubescent. S. France.	—	— H. ⊙.		
4 Aconiti. DC.	(pu.) Aconite-like. Levant. 1801.	—	— H. ⊙.		
5 ambiguum. DC.	(bl.) doubtful. Barbary. 1759.	7. 9.	H. ⊙.		
6 júnceum. DC.	(bl.) Rush-like. Italy. 1629.	—	— H. ⊙.		
peregrinum. Lin. non Lam.					
7 grâcile. DC.	(re.) slender. Spain. 1826.	—	— H. ⊙.		
8 cardiopétalum. DC.	(bl.) heart-petaled. Pyrenees. 1822.	—	— H. ⊙.		
peregrinum. Lam. non Lin.					
9 vîrgatum. DC.	(bl.) twiggy. Syria. 1824.	6. 9.	H. ⊙.		
10 grandiflorum. DC.	(bl.) great-flowered. Siberia. 1741.	6. 9.	H. 4.		
β chinéns. F.	(bl.) Chinese. China. 1816.	—	— H. 4.		
γ álbum.	(wh.) white-flowered. —	—	— H. 4.		
δ flôre-plêno.	(bl.) double-flowered. —	—	— H. 4.		
11 cheilánthum. DC.	(bl.) hairy-leaved. Dahuria. 1818.	—	— H. 4.		
12 vîrscens. N.	(wh.) greenish. N. America. 1827.	—	— H. 4.		
13 pumiceum. DC.	(pu.) dark-purple. Tartary. 1785.	7. 9.	H. 4.		
14 albiflòrum. DC.	(wh.) white-flowered. Armenia. 1823.	—	— H. 4.		
15 ochroleúcum. DC.	(st.) sulphur-coloured. Iberia. 1817.	—	— H. 4.		
16 hy'bridum. DC.	(bl.) hairy. Siberia. 1794.	6. 9.	H. 4.		
17 fissum. P. S.	(bl.) cloven. Hungary. 1816.	—	— H. 4.		
18 velutinum. DC.	(bl.) velvetty. Italy. 1824.	—	— H. 4.		
19 pentágynum. DC.	(bl.) five-styled. Portugal. 1820.	—	— H. 4.		
20 Menziesii. DC.	(bl.) Menzies'. N. America. 1826.	6. 8.	H. 4.		
21 élégans. DC.	(bl.) elegant. —	1741.	— H. 4.		
β multipléx.	(bl.) common-double. —	—	— H. 4.		
22 tricórne. DC.	(bl.) three-horned. —	1806.	4. 6. H. 4.		
23 amœnum. DC.	(bl.) pretty. Siberia.	—	— H. 4.	Lodd. bot. cab. 306.	
24 azûreum. DC.	(bl.) azure. Carolina. 1805.	5. 7.	H. 4.	Gmel. sib. 4. t. 77.	
25 exaltát̄um. DC.	(bl.) tall. N. America. 1758.	7. 8.	H. 4.	Deless. ic. sel. 1. t. 60.	
26 urceolát̄um. DC.	(bl.) hollow-leaved. —	1806.	6. 7. H. 4.	Mill. ic. t. 250. f. 2?	
				Jac. ic. 1. t. 101.	

27	revolutum. DC.	(bl.) revolute.	7. 9. H. 24.		
28	mesoleucum. DC.	(bl.) white-lipped.	1824.	H. 24.	Swt. br. fl. gar. ic.	
29	palmatifidum. DC.	(bl.) palmate.	Siberia.	6. 9. H. 24.	Gmel. sib. 4. t. 79?	
	α hispidum. DC.	(bl.) hispid.	H. 24.	— 4. t. 75?	
	β glabellum. DC.	(bl.) smooth.	H. 24.		
30	intermedium. DC.	(bl.) intermediate.	Europe.	1710	7. 8. H. 24.	Gmel. sib. 4. t. 80?
	α pilosissimum. DC.	(bl.) villous.	H. 24.	Pl. rar. hung. 3. t. 246.	
	β alpinum. DC.	(bl.) Alpine.	1819.	H. 24.		
	γ leptostachyum. DC.	(bl.) slender-spiked.	H. 24.		
	δ ranunculifolium. DC.	Ranunculus-la.	H. 24.		
	ϵ laxum. DC.	(bl.) loose-flowered.	H. 24.		
31	cuneatum. DC.	(bl.) wedge-leaved.	Siberia.	1815.	H. 24.	Mill. ic. t. 119?
32	villosum. DC.	(bl.) villous.	1819.	H. 24.	Bot. reg. 327.
33	dyctiocarpum. DC.	(bl.) netted-capsuled.	Siberia.	H. 24.	
34	laxiflorum. DC.	(bl.) loose-flowered.	6. 8. H. 24.		
35	montanum. DC.	(bl.) mountain.	Pyrenees.	H. 24.	
	β bracteatum. DC.	(bl.) large-bracted.	Switzerland.	H. 24.	
36	dasycarpum. DC.	(bl.) thick-capsuled.	Caucasus.	1824.	H. 24.	
37	speciosum. DC.	(bl.) large-flowered.	1817.	8. 10. H. 24.	Deless. ic. sel. 1. t. 62.
38	spurium. F.	(bl.) spurious.	Siberia.	1817.	6. 9. H. 24.	
39	ucranicum. Bes.	(bl.) Ucranian.	1818.	H. 24.	
40	pseudoperegrinum. F.	(bl.) Fischer's.	H. 24.	
41	crassicaule. Gm.	(bl.) thick-stemmed.	1822.	H. 24.	
42	pallidum. F.	(pa.) pale-flowered.	Siberia.	1818.	H. 24.	Trev. delph. t. 1. et 2.
43	flexuosum. DC.	(bl.) flexuose.	Caucasus.	1817.	7. 9. H. 24.	
44	discolor. F.	(v.) two-coloured.	Siberia.	1819.	6. 8. H. 24.	
45	triste. DC.	(da.) dark-flowered.	1822.	H. 24.	
46	Requiemii. DC.	(pa.) Requien's.	S. Europe.	1819.	6. 7. F. 3.	
47	pictum. DC.	(pa.) painted.	1816.	H. 24.	Deles. ic. sel. 1. t. 63.
	Staphysagria. DC.	(bl.) Stavesacre.	1596.	5. 6. F. 3.	Swt. br. fl. gar. 2. 123.
						Flor. græc. t. 508.
ACONITUM. DC. MONKSHOOD.						
1	A'nthora. L.	(st.) wholesome.	Pyrenees.	1596.	6. 8. H. 24.	Rch. ac. p. 61. t. 1. f.A.
	β grandiflorum. R.U.	(st.) large-flower'd.	H. 24.	— p. 63. t. 1. f. B.
	γ glabriiflorum. R.U.	(st.) smooth-flowd.	H. 24.	
2	eulophum. R.U.	(st.) eatable.	Switzerland.	1824.	H. 24.	Rch. ac. p. 69. t. 5.
	anthoroidium. DC.	non R.U.				
3	nemorosum. R.U.	(ye.) wood.	Caucasus.	1822.	H. 24.	— p. 71. t. 6.
4	Dec ndollii. R.U.	(ye.) Decandolle's.	Pyrenees.	H. 24.	— p. 67. t. 3.
5	Jacquinii. R.U.	(ye.) Jacquin's.	Austria.	H. 24.	— p. 65. t. 2.
	A'nthora. Jacq.	aust. t. 3s2.				
6	Pallasii. R.U.	(st.) Pallas's.	Siberia.	1824.	H. 24.	— p. 72. t. 6. f. A.B.
7	inclinatum.	(st.) inclined.	H. 24.	Ser. m. helv. I. t. 15. f. 2.
	anthroidium.	Rch. ac. 68. t. 4. non DC.				
8	Koelleanum. R.U.	(bl.) Koelle's.	Germany.	5. 7. H. 24.	Rch. ac. t. 11. f. 1.
9	tauricum. R.U.	(bl.) Taurian.	Tauria.	1752.	6. 8. H. 24.	— t. 12. f. 2—3.
10	firmum. R.U.	(bl.) firm.	Germany.	1822.	H. 24.	— t. 14. f. 2.
11	acutum. R.U.	(bl.) acute.	H. 24.	— t. 14. f. 1.
12	latum. R.U.	(bl.) great-flowered.	Sweden.	1823.	H. 24.	— t. 13. f. 2.
13	Clusii. R.U.	(bl.) Clusius's.	Austria.	6. 7. H. 24.	Mor. his. 3.s.12.t.3.f.17.
14	amœnum. R.U.	(bl.) delightful.	Sweden.	1824.	6. 8. H. 24.	Rch. ac. 93. t. 14. f. 1.
15	Hoppii. R.U.	(bl.) Hoppe's.	Carinthia.	H. 24.	
16	oligocarpum. R.U.	(bl.) two-podded.	Styria.	1823.	H. 24.	
17	calibotrys. R.U.	(bl.) long-spiked.	Moravia.	1824.	6. 7. H. 24.	Rch. ac. 98. t. 16. f. 1.
18	tenuifolium. R.U.	(bl.) slender-leaved.	Styria.	H. 24.	
19	angustifolium. R.U.	(bl.) narrow-leaved.	Siberia.	H. 24.	Rch. ac. 95. t. 15. f. 2.
20	eustachyum. R.U.	(bl.) loose-spiked.	M. Baldo.	H. 24.	— t. 15. f. 3.
21	Napellus. R.U.	(bl.) common.	Europe.	5. 7. H. 24.	Lob. ic. 679. f. 1.
22	compactum. R.U.	(bl.) compact.	Switzerland.	1823.	H. 24.	
	β rubellum. R.U.	(re.) red-flowered.	H. 24.	
23	Halleri. R.U.	(bl.) Haller's.	1824.	6. 7. H. 24.	Chabr. sci. p. 531. f. 2.
	β humile. R.U.	(bl.) dwarf.	H. 24.	
	γ ramosum. R.U.	(bl.) branching.	1822.	H. 24.	
	δ bicolor. R.U.	(w.b.) two-coloured.	H. 24.	
24	virgatum. R.U.	(bl.) slender-branch'd.	1824.	H. 24.	
25	venustum. R.U.	(bl.) beautiful.	1823.	6. 8. H. 24.	
26	laxiflorum. R.U.	(bl.) loose-flowered.	1824.	H. 24.	
27	Funkii. R.U.	(bl.) Funk's.	Germany	H. 24.	
28	Braunii. R.U.	(bl.) Braune's	6. 7. H. 24.	
29	trichocarpum. R.U.	(bl.) three-podded.	H. 24.	

30 pubescens. R.U. (<i>bl.</i>) pubescent.	Switzerland.	5. 7.	H. 24.
31 Mielichóferi. R.U. (<i>bl.</i>) Mielichofer's.	Germany.	1823.	6. 8.	H. 24.
32 pláticatum. R.U. (<i>bl.</i>) plaited.	Sweden.	1824.	—	H. 24.
33 Koehléri. R.U. (<i>bl.</i>) Koehler's	—	1823.	—	H. 24.
34 elátum. R.U. (<i>bl.</i>) tall.	Germany.	—	H. 24.
35 álbidum. R.U. (<i>wh.</i>) whitish.	1824.	—	H. 24.
36 neubergéne. R.U. (<i>bl.</i>) Styrian. <i>Napélas.</i> Jacq. aust. 4. t. 381.	Styria.	—	H. 24.
37 Meyéri. R.U. (<i>bl.</i>) Meyer's.	Germany.	1824.	—	H. 24.
38 Bernhardiànum. (<i>bl.</i>) Bernhardi's.	1823.	—	H. 24.
39 stríctum. R.U. (<i>bl.</i>) straight. <i>spicatum.</i> Donn. h. cant. ed. 10. p. 209.	Siberia.	—	H. 24.
40 Sprengeli. R.U. (<i>bl.</i>) Sprengel's.	1823.	6. 8.	H. 24.
41 Willdenowii. R.U. (<i>bl.</i>) Willdenow's.	Carniola.	—	—	H. 24.
42 éminens. R.U. (<i>bl.</i>) eminent.	—	1822.	—	H. 24.
43 Schleichéri. R.U. (<i>bl.</i>) Schleicher's. β <i>comosum.</i> R.U. (<i>bl.</i>) tufted.	Switzerland.	5. 6.	H. 24.
44 microstachyum. R.U. (<i>b.</i>) small-spiked.	Hungary.	1822.	6. 8.	H. 24.
45 formòsum. R.U. (<i>bl.</i>) handsome.	Germany.	1824.	—	H. 24.
46 rigidum. R.U. (<i>bl.</i>) rigid.	Sweden.	—	—	H. 24.
47 commutatum. R.U. (<i>bl.</i>) changed.	Switzerland.	—	—	H. 24.
48 lágum. R.A. (<i>bl.</i>) loose.	Germany.	1823.	6. 8.	H. 24.
49 autumnále. R.A. (<i>bl.</i>) autumnal.	—	—	—	H. 24.
50 ampliflórum. R.U. (<i>bl.</i>) large-flowered.	Austria.	1823.	—	H. 24.
51 máximum. DC. (<i>bl.</i>) large.	Kamtschatka.	—	7. 8.	H. 24.
52 biflorum. DC. (<i>va.</i>) two-flowered. α <i>luteo-caeruleum</i> (<i>y.b.</i>) blue and yellow. β <i>caeruleum.</i> (<i>bl.</i>) blue-flowered.	Altay.	1826.	6. 8.	H. 24.
53 ferox. DC. (<i>pu.</i>) poison-rooted. <i>virðsum.</i> D.P.	Nepaul.	1820.	—	H. 24.
54 productum. R.U. (<i>bl.</i>) long-lipped.	Siberia.	1824.	7. 8.	H. 24.
55 semigaleatum. R.U. (<i>bl.</i>) half-helmeted.	Kamtschatka.	—	—	H. 24.
56 uncinatum. DC. (<i>pu.</i>) American. β <i>Michauxianum.</i> (<i>pu.</i>) Michaux's	N. America.	1768.	—	H. 24.
57 villósum. R.U. (<i>bl.</i>) villous.	Siberia.	1825.	—	H. 24.
58 fláccidum. R.U. (<i>bl.</i>) flaccid. <i>ciliare</i> β <i>polytrichum.</i> DC.	—	—	H. 24.
59 volubile. R.U. (<i>bl.</i>) twining.	—	1799.	—	H. 24.
60 tortuosum. DC. (<i>bl.</i>) twisting.	—	1812.	—	H. 24.
61 eriostemum. DC. (<i>bl.</i>) woolly-stamen'd.	Siberia.	1824.	6. 8.	H. 24.
62 recognitum. R.U. (<i>bl.</i>) recognised.	—	—	—	H. 24.
63 ambiguum. R.U. (<i>bl.</i>) ambiguous.	Siberia.	1826.	—	H. 24.
64 tóxicum. R.U. (<i>bl.</i>) poisonous.	Germany.	—	—	H. 24.
65 reflexum. R.U. (<i>bl.</i>) reflexed.	Carniola.	1826.	6. 8.	H. 24.
66 cérunum. R.U. (<i>bl.</i>) drooping-flow'd.	Germany.	1824.	6. 7.	H. 24.
67 acuminatum. R.U. (<i>bl.</i>) acuminate.	M. Baldo.	1824.	—	H. 24.
68 parviflórum. R.U. (<i>bl.</i>) small-flowered.	Switzerland.	—	—	H. 24.
69 hebégynum. DC. (<i>bl.</i>) downy-podded.	—	—	—	H. 24.
70 paniculatum. R.U. (<i>bl.</i>) paniced.	—	5. 7.	H. 24.
71 laciniósom. Sch. (<i>bl.</i>) jagged.	—	1822.	—	H. 24.
72 mólle. R.U. (<i>bl.</i>) soft.	—	1824.	—	H. 24.
73 gibbosum. DC. (<i>bl.</i>) gibbous.	Caucasus.	1823.	—	H. 24.
74 hamátum. R.U. (<i>bl.</i>) hooked.	Sweden.	1804.	6. 8.	H. 24.
75 palmatifidum. R.U. (<i>bl.</i>) palmatifid.	M. Carpathian.	1819.	—	H. 24.
76 pyramidale. R.U. (<i>bl.</i>) pyramidal. α <i>densiflórum.</i> R.U. (<i>bl.</i>) close-flowered. β <i>elongatum.</i> R.U. (<i>bl.</i>) elongated.	Bohemia.	1596.	—	H. 24.
γ <i>bicolor.</i> R.U. (<i>w.b.</i>) two-coloured.	—	—	—	H. 24.
77 Sterkiánium. R.U. (<i>bl.</i>) Sterk's.	S. Europe.	1824.	—	H. 24.
β <i>laxiflórum.</i> R.U. (<i>bl.</i>) loose-flowered.	—	—	—	H. 24.
γ <i>bicolor.</i> R.U. (<i>w.b.</i>) variegated.	—	—	—	H. 24.
78 versicolor. R.U. (<i>va.</i>) various-colored.	Switzerland.	—	—	H. 24.
β <i>bicolor.</i> R.U. (<i>w.b.</i>) white and blue.	—	—	—	H. 24.
79 décorum. R.U. (<i>bl.</i>) neat.	Pyrenees.	1824.	—	H. 24.
80 exaltatum. R.U. (<i>bl.</i>) lofty.	—	—	—	H. 24.
81 macránthum. R.U. (<i>bl.</i>) large-blossom'd.	Germany.	1826.	—	H. 24.
82 Cámmarum. R.U. (<i>pu.</i>) purple.	Europe.	1752.	—	H. 24.
83 illinitum. R.U. (<i>bl.</i>) besmeared.	Switzerland.	1822.	—	H. 24.
84 Breiterianum. R.U. (<i>bl.</i>) Breiter's.	—	1821.	—	H. 24.

Clus. hist. 2. p. 96. f. 1.

Rch. ac. t. 17. f. 1.

Mor. his. 3.s.12.t.3.f.19.

Rch. ac. t. 18. f. 2.

Rch. ac. t. 18. f. 3.

Rch. ac. 97. t. 15. f. 4.

— t. 17. f. 2.

Ser. mus. 1.t.15.f.31.32.

— 1. t. 15.f.45.46.

Rch. ac. 74. t. 7. f. 1. 2.

Ser. mus. 1.t.15.f.43.44.

Rch. ac. 75. t. 7. f. 3.

— 77. t. 8.

Bot. mag. 1119.

Deles. ic. sel. 1. t. 65.

Ser. mus. 1.t.15.f.28.29.

Chabr. sci. 531. f. 6.

Chabr. sci. 531. f. 3.

Ser. m. 1. t. 15.f. 35.36.

— 1. t. 15. f. 14. 15.

Stor. lib. ac. 69. ic.

Blackw. ic. 561.

Lodd. bot. cab. 794.

Ser. m. 1. t. 15. f. 37. 38.

Jacq. aust. 5. t. 424.

85 <i>Ottoniænum</i> . R. U. (<i>bl.</i>)	Otto's.	1823.	6. 8. H. 2.	
86 <i>speciosum</i> . R. U. (<i>bl.</i>)	specious.	1804.	7. 8. H. 2.	Ser. m. 1. t. 15. f. 26. 27.
87 <i>bulbiferum</i> . R. U. (<i>bl.</i>)	bulb-bearing.	1824.	6. 8. H. 2.	
88 <i>leucanthum</i> . R. U. (<i>w.</i>)	white-flowered.	—	H. 2.	
89 <i>lasiocarpum</i> . R. U. (<i>bl.</i>)	woolly-podded.	Hungary.	1822.	—	H. 2.
90 <i>gracile</i> . R. U. (<i>bl.</i>)	slender.	Germany.	1824.	6. 7. H. 2.	
91 <i>rhynchanthum</i> . R. U.	beaked.	Hungary.	1804.	6. 8. H. 2.	
β <i>bicolor</i> . R. U. (<i>w.b.</i>)	two-coloured.	—	H. 2.	
92 <i>nasutum</i> . R. U. (<i>bl.</i>)	large-nosed.	Caucasus.	1823.	—	H. 2.
93 <i>rostratum</i> . R. U. (<i>pu.</i>)	beaked.	Switzerland.	—	H. 2.	Lodd. bot. cab. 203.
94 <i>variegatum</i> . DC. (<i>p.w.</i>)	variegated.	S. Europe.	1597.	—	Clus. hist. 2. p. 98. f. 1.
95 <i>mixtum</i> . R. U. (<i>p.w.</i>)	mixed.	—	H. 2.	Rivin. pentap. ic.
96 <i>álbum</i> . R. U. (<i>wh.</i>)	white.	Levant.	1752.	7. 8. H. 2.	Ser. m. 1. t. 15. f. 22. 23.
97 <i>japonicum</i> . DC. (<i>va.</i>)	Japan.	Japan.	1790.	6. 9. H. 2.	
α <i>cárneum</i> . (<i>fl.</i>)	flesh-coloured.	—	—	H. 2.	
β <i>cæruleum</i> . (<i>bl.</i>)	blue-flowered.	—	—	H. 2.	
98 <i>boreale</i> . R. U. (<i>st.</i>)	Siberian.	Siberia.	6. 8. H. 2.	
99 <i>barbatum</i> . DC. (<i>st.</i>)	bearded.	—	1807.	—	Swt. br. fl. gar. v. 2. 164.
100 <i>hispidum</i> . DC. (<i>st.</i>)	hispid.	—	1826.	—	Gmel. sib. 4. t. 81.
<i>Gmelini</i> . R. U.					
101 <i>squarrósum</i> . DC. (<i>st.</i>)	squarrose.	—	1823.	—	H. 2.
102 <i>pallidum</i> . R. U. (<i>st.</i>)	pale-flowered.	—	1823.	—	H. 2.
103 <i>zoóctonum</i> . R. U. (<i>st.</i>)	beast-bane.	Germany.	1824.	—	H. 2.
104 <i>melóctonum</i> . R. U. (<i>st.</i>)	Badger-bane.	Piedmont.	1823.	—	H. 2.
105 <i>pyrenaicum</i> . DC. (<i>st.</i>)	Pyrenean.	Pyrenees.	1739.	—	H. 2.
106 <i>galéontonum</i> . R. U. (<i>st.</i>)	Weasel-bane.	Hungary.	1823.	—	H. 2.
107 <i>moldávicum</i> . R. U. (<i>pu.</i>)	Moldavian.	Bohemia.	—	—	H. 2.
108 <i>ochroleúcum</i> . R. U. (<i>st.</i>)	pale-white.	Caucasus.	1794.	—	H. 2.
109 <i>ægòphorum</i> . R. U. (<i>st.</i>)	Goat-poison.	—	1822.	—	H. 2.
110 <i>perniciósum</i> . R. U. (<i>st.</i>)	pernicious.	France.	—	—	Bois. fl. eur. 2. t. 369.
111 <i>myóctonum</i> . R. U. (<i>st.</i>)	Mouse-bane.	Thuringia.	—	—	
112 <i>strictissimum</i> . R. U. (<i>st.</i>)	straight.	Barnth.	—	Cam. epit. 831. ic.
113 <i>altíssimum</i> . R. U. (<i>st.</i>)	tallest.	Germany.	—	Chabr. sci. p. 531. f. 1.
114 <i>réctum</i> . R. U. (<i>st.</i>)	upright.	Italy.	—	Tabern. ic. 583. f. 1.
115 <i>vulpária</i> . R. U. (<i>st.</i>)	Fox-bane.	Germany.	1824.	—	Barrel. ic. 599.
β <i>latifolium</i> . R. U. (<i>st.</i>)	broad-leaved.	Bavaria.	—	—	
116 <i>tragóctonum</i> . R. U. (<i>st.</i>)	Goat-bane.	Carinthia.	—	—	
117 <i>lupicida</i> . R. U. (<i>st.</i>)	Wolf-bane.	Switzerland.	1824.	—	
118 <i>arctóphonum</i> . R. U. (<i>st.</i>)	Bear-bane.	Styria.	1826.	—	
119 <i>lagóctonum</i> . R. U. (<i>st.</i>)	Hare-bane.	Bavaria.	—	—	
120 <i>Phthòra</i> . R. U. (<i>st.</i>)	destructive.	Hungary.	—	—	
121 <i>austrále</i> . R. U. (<i>pu.</i>)	Carpathian.	M. Carpathian.	1815.	—	Bot. mag. 2196.
<i>septentrionálē</i> β <i>carpáthicum</i> . B. M.					
122 <i>rubicündum</i> . F. (<i>pu.</i>)	rubicundi.	Russia.	1822.	—	Blackw. ic. 563.
123 <i>cynóctonum</i> . R. U. (<i>st.</i>)	Dog-bane.	Germany.	1824.	—	
124 <i>thelyphonum</i> . R. U. (<i>st.</i>)	Elwert's.	Switzerland.	—	—	Barrel. ic. 600.
125 <i>therióphonum</i> . R. U. (<i>st.</i>)	park-bane.	Italy.	6. 7. H. 2.	Clus. hist. 5. p. 94. ic.
126 <i>lycóctonum</i> . R. U. (<i>st.</i>)	great-yellow.	Germany.	1596.	7. 8. H. 2.	Trag. stirp. 247. ic.
127 <i>lupária</i> . R. U. (<i>st.</i>)	Wolf-wort.	—	—	H. 2.	Flor. dan. t. 123.
128 <i>septentrionálē</i> . R. U. (<i>bl.</i>)	northern.	N. Europe.	1800.	—	H. 2.

Tribus V. PÆONIACEÆ. DC. prodr. 1. 64.

CIMICI'FUGA. L.	BUG-WORT.	Polyandria Pentagynia. L.	
1 <i>fœ'tida</i> . w. (<i>wh.</i>)	stinking.	Siberia.	1777. 6. 7. H. 2.
2 <i>podocárpa</i> . (<i>wh.</i>)	American.	Carolina.	1822. 8. 9. H. 2.
<i>americana</i> . M. <i>Actæa podocarpa</i> . DC.			
3 <i>palmata</i> . Ph. (<i>wh.</i>)	palmated.	—	1812. 7. 8. H. 2.
4 <i>cordifolia</i> . Ph. (<i>wh.</i>)	heart-leaved.	—	6. 7. H. 2.
MACR'OTYS. Raf.	SNAKEROOT.	Polyandria Monogynia. L.	
<i>racemosa</i> . (<i>wh.</i>)	black-rooted.	N. America.	1732. 6. 7. H. 2.
<i>racemosa</i> . w. (<i>wh.</i>)			
<i>Actæa racemosa</i> . w.	<i>Cimicifuga serpentaria</i> . Ph.		
ACT'EA. Ph.	BANE-BERRIES.	Polyandria Monogynia. L.	
1 <i>spicata</i> . DC. (<i>wh.</i>)	common.	Britain. 4. 6. H. 2.
2 <i>rúbra</i> . W.en. (<i>wh.</i>)	red-berried.	N. America. — H. 2.
<i>brachypétala</i> . β <i>rúbra</i> . DC.			
			Eng. bot. 918.

RANUNCULACEÆ.

3 álba, Mill.	(wh.) white-berried.	N. America.	4. 6. H. ¼.	Corn. canad. t. 77.
brachypétala.	α álba.	DC.			
ZANTHOR'IZA. DC.	YELLOW-ROOT.	Pentandria Polygynia. L.			
apiifolia. DC.	(pu.) Parsley-leaved.	N. America.	1766.	2. 4. H. ½.	Bot. mag. 1736.
PÆONIA. DC.	PEONY.	Polyandria Digynia. L.			
1 Moután. H.K.	Chinese-tree.	China.	1789.	4. 6. H. ½.	
α papaverácea. L.T. (w.)	Poppy-flowered.	—	1806.	— H. ½.	Bot. rep. 463.
β Banksii. L.T. (bl.)	common.	—	1789.	— H. ½.	— 448.
γ rosea. L.T. (ro.)	rose-coloured.	—	1794.	— H. ½.	— 373.
2 édulis. P.L.	(va.) eatable-rooted.	—	5. 6. H. ¼.	
albiflora. L.T.					
α vestalis. L.T. (wh.)	virgin.	Siberia.	1784.	— H. ¼.	Andr. rep. 64.
β candida. L.T. (wh.)	pale-flowered.	—	— H. ¼.	
γ taratica. L.T. (bh.)	Tartarian.	Tartary.	— H. ¼.	
δ sibirica. L.T. (wh.)	Siberian.	Siberia.	— H. ¼.	
ε rubescens. L.T. (bh.)	pale-red.	—	— H. ¼.	
ζ uniflora. L.T. (wh.)	one-flowered.	—	— H. ¼.	
η Whitleji. L.T. (wh.)	double-white.	China.	1808.	6. 7. H. ¼.	Bot. mag. 1756.
θ Humei. L.T. (re.)	double-crimson.	—	1810.	— H. ¼.	Bot. reg. 630.
ι frágrans. L.T. (re.)	Rose-scented.	—	1805.	— H. ¼.	Bot. mag. 1768.
3 anomala. L.T. (re.)	jagged-leaved.	Siberia.	1788.	5. 6. H. ¼.	Bot. reg. 485.
laciñata. Pall. ross. 2. t. 85.					Bot. mag. 1754.
4 hybrida. DC. (re.)	intermediate.	Caucasus.	1822.	— H. ¼.	
5 tenuifolia. L.T. (cr.)	slender-leaved.	Siberia.	1765.	5. 6. H. ¼.	Bot. mag. 926.
6 officinalis. L.T. (va.)	official.	Europe.	1548.	5. 6. H. ¼.	Bot. mag. 1784.
α Sabini. L.T. (sc.)	Sabine's.	—	— H. ¼.	
β ròsica. L.T. (ro.)	rose-coloured.	—	— H. ¼.	
γ blanda. L.T. (bh.)	blush-flowered.	—	— H. ¼.	
δ rubra. L.T. (cr.)	double-red.	—	— H. ¼.	Lob. ic. 684.
ε carnescens. L.T. (fl.)	flesh-coloured.	—	— H. ¼.	
ζ albicans. L.T. (bh.)	double-blush.	—	— H. ¼.	
7 corallina. L.T. (cr.)	English.	England.	—	— H. ¼.	Park. par. f. 4.
8 daurica. L.T. (fl.)	Daurian.	Dauria.	1790.	— H. ¼.	Eng. bot. 1513.
triternata. DC. prodri.					Bot. mag. 1441.
9 lobata. DC. (sc.)	lobed-leaved.	Portugal.	1821.	— H. ¼.	
10 Russi. DC. (cr.)	crimson.	Sicily.	—	— H. ¼.	Swt. br. fl. gar. 70.
11 pùbens. B.M.	(re.) pubescent.	—	1821.	— H. ¼.	— v. 2. t. 122.
12 hùmilia. L.T.	(re.) dwarf.	Spain.	1633.	5. H. ¼.	Bot. mag. 2264.
13 décora. L.T. (re.)	comely.	Turkey.	—	— H. ¼.	— 1422.
α Pallasiæ. L.T. (re.)	Pallas's.	—	— H. ¼.	
β elatior. L.T. (re.)	tall.	—	— H. ¼.	
14 arietina. L.T. (re.)	rams-horned.	—	— H. ¼.	
15 crética. B.R. (bh.)	Cretan.	Crete.	—	— H. ¼.	Bot. reg. 819.
16 peregrina. L.T. (re.)	Turkish.	Levant.	1588.	5. 6. H. ¼.	Bot. mag. 1050.
α byzantina. L.T. (re.)	Levant.	—	— H. ¼.	
β compacta. L.T. (re.)	compact.	France?	—	— H. ¼.	
γ Grevillii. L.T. (re.)	Greville's.	—	— H. ¼.	
17 paradóxa. L.T. (re.)	paradoxical.	Levant.	—	— H. ¼.	
α simpliciflora. (re.)	single-flowered.	—	— H. ¼.	Swt. br. fl. gar. 19.
β fimbriata. L.T. (cr.)	double-flowered.	—	— H. ¼.	Bot. reg. 474.
18 mollis. L.T. (re.)	soft-leaved.	Siberia?	—	— H. ¼.	Swt. br. fl. gar. 113.
19 villosa. B.F.G. (wh.)	villous.	—	1822.	— H. ¼.	
sessiliflora. B.M. 2648.					

ORDO II.

DILLENIACEÆ. DC. syst. nat. 1. p. 395.

Tribus I. DELIMACEÆ. DC.

TETRA'CERA. DC.	TETRA'CERA.	Polygamia Dioecia.			
1 volubilis. DC.	(ye.)	climbing.	Barbadoes.	S. h. — Pluk. alm. 48. t. 146.f.1.
2 jamaicensis. DC.	(ye.)	Jamaica.	Jamaica.	1826.	S. h. —
3 obovata. DC.	(ye.)	obovate-leaved.	Senegal.	1822.	S. h. —
4 alnifolia. DC.	(ye.)	Alder-leaved.	Sierra Leone.	—	S. h. —
5 potatoria. Afz.	(ye.)	water-bearing.	—	—	S. h. —
DOLIOCA'RPU'S. DC.	DOLIOCA'RPU'S.	Polyandria Monogynia.			
1 Calínea. DC.	(wh.)	climbing.	Guiana.	1822.	S. h. — Aub. Guian. 1. t. 221.
Tetracera calinea. w.	Calínea scándens.	Aub.			

DEL'IMA. DC.	DEL'IMA.	Polyandria Monogynia.
sarmentosa. DC.	sarmentose.	Ceylon. 1822. S. h. Lam. ill. t. 475.
Tetrácerasarmentosa. w.		
CURATE'LLA. DC.	CURATE'LLA.	Polyandria Digynia.
1 americana. DC. (wh.) American.	Guiana. S. h. Aub. guian. 1. t. 232.
2 ? alata. v.	wing-petioled. S. h. Venten. choix. t. 49.

Tribus II. **DILLENEÆ. DC.**

PLEURA'NDRA. DC.	PLEURA'NDRA.	Polyandria Monogynia.
1 bracteata. DC	(ye.) elegant.	N. S. W. 1822. 4. 8. G. h. Deles. ic. sel. 1. t. 78.
2 niftida. DC.	(ye.) shining.	_____ _____ G. h.
3 cneódrum. DC.	(ye.) blunt-leaved.	_____ 1824. 5. 7. G. h.
4 scàbra. DC.	(ye.) rough-leaved.	_____ _____ G. h.
5 aciculàris. DC.	(ye.) needle-leaved.	_____ _____ 4. 7. G. h. Labill. nov. holl. 2.t.144.
6 strícta. DC.	(ye.) upright.	_____ 1822. _____ G. h.
7 calycina. DC.	(ye.) linear-leaved.	_____ _____ G. h.
CANDO'LLEA.	CANDO'LLEA.	Polyadelphia Polyandria.
cuneifòrmis. DC. (ye.) wedge-leaved.		N. Holland. 1823. 5. 8. G. h. Bot. mag. 2711.
HIBBERTIA. DC.	HIBBE'RRIA.	Polyandria Mono-Pentagynia.
1 grossulariaefolia. DC. (y.) Gooseberry-lid.		N. Holland. 1803. 3. 8. G. h. Bot. mag. 1218.
crenata. A.R. t. 472.		
2 volubilis. DC.	(ye.) twining.	N. S. W. 1796. 5. 10. G. h. Andr. rep. t. 126.
3 dentata. DC.	(ye.) tooth-leaved.	N. Holland. 1814. 1. 8. G. h. Bot. reg. 282.
4 saligna DC.	(ye.) Willow-leaved.	N. S. W. 1824. 5. 8. G. h.
5 virgata DC.	(ye.) twiggy.	_____ 1822. G. h.
6 fasciculata. DC.	(ye.) bushy.	_____ 1826. 5. 8. G. h.
7 linearis. DC.	(ye.) linear-leaved.	_____ _____ G. h.
8 monogyna. DC.	(ye.) one-styled.	_____ _____ 5. 8. G. h.
9 pedunculata. DC.	(ye.) long-pediced.	_____ _____ 4. 9. G. h. Bot. reg. 1001.
		corifolia. Bot. mag. 2672.

WO'R'MIA. DC.	WO'R'MIA.	Polyandria Pentagynia.
dentata. DC. (ye.) tooth-leaved.		Ceylon. 1821. S. h.
Dillénia dentata. w.		Thunb. linn. tr. 1. t. 20.

COLBERTIA. DC.	COLBE'R'TIA.	Polyandria Pentagynia.
1 coromandeliana. DC. (y.) Coromandel.		E. Indies. 1803. S. h.
Dillénia pentágynia. Roxb.		Roxb. cor. 1. t. 20.
2 scabrélla. D.P. rough.		Nepaul. 1826. S. h.
DILL'ENIA. DC.	DILL'ENIA.	Polyandria Monogynia. L.
1 speciosa. DC. (wh.) large-flowered.		E. Indies. 1800. S. h.
2 intègra. DC. entire-leaved.		Ceylon. 1826. S. h.
		Exot. bot. t. 2. 3.
		Thunb. linn. tr. 1. t. 18.

ORDO III.

WINTEREÆ. *Brown in DC. syst. 1. add. p. 548.*

ILL'ICIUM. DC.	ANISEED-TREE.	Polyandria Polygynia. L.
1 floridànum. DC. (cr.) red-flowered.	Florida.	1766. 4. 6. F. h. Bot. mag. 439.
2 anisatum. DC. (ye.) Chinese.	China. F. h. Lam. ill. t. 493. f. 2.
3 parviflòrum. DC. (ye.) small-flowered.	Florida.	1790. — F. h. Vent. cels. t. 22.
DR'IMYS. DC.	DR'IMYS.	Polyandria Tetra-Octogynia.
Winteri. DC. Winter-bark.	Magellan.	1825. G. h. Forst. gen. p. 84. t. 42.
Wintera aromàtica. w.	Winterana aromatica. Sol. med. obs. 5. p. 46. t. 1.	
TASMA'NNIA. DC.	TASMA'NNIA.	Dicæcia Polyandria.
dipétala. B. two-petaled.	N. Holland.	1824. 5. 8. G. h.

ORDO IV.

MAGNOLIACEÆ. *DC. syst. nat. v. 1. p. 439.*

MICH'ELIA. DC.	MICH'ELIA.	Polyandria Polygynia. L.
Champáca. DC. (ye.) sweet-scented.	E. Indies.	1779. S. h. Lam. ill. t. 493.

MAGN'OLIA. DC.	MAGN'OLIA. Polyandria Polygynia. L.	Carolina.	1734.	6. 10.	H. $\frac{1}{2}$.
1 grandiflora. DC. (<i>wh.</i>) Laurel-leaved.	Laurel-leaved.	Carolina.	1734.	6. 10.	H. $\frac{1}{2}$.
α rotundifolia. (<i>wh.</i>) round-leaved.	H. $\frac{1}{2}$.
β obovata. H.K. (<i>wh.</i>) obovate-leaved.	H. $\frac{1}{2}$.
γ elliptica. H.K. (<i>wh.</i>) elliptic-leaved.	H. $\frac{1}{2}$.
δ ferruginea. B.M. (<i>w.</i>) ferruginous.	H. $\frac{1}{2}$.
ϵ lanceolata. H.K. (<i>wh.</i>) Exmouth.	H. $\frac{1}{2}$.
2 glauca. DC. (<i>wh.</i>) glaucous-leaved.	N. America.	1688.	6. 9.	H. $\frac{1}{2}$.	Mill. ic. v. 2. t. 172. Tr. Eh. p. 8. t. 33-35.
3 longifolia. (<i>wh.</i>) narrow-leaved.	Bot. mag. 1952. Mich. arb. t. 2.
glauca β longifolia. Ph.					
4 Thompsoniana. (<i>wh.</i>) hybrid.	Hybrid.	1818.	—	H. $\frac{1}{2}$.	Mich. arb. t. 5.
5 umbrælla. DC. (<i>wh.</i>) Umbrella.	N. America.	1752.	5. 6.	H. $\frac{1}{2}$.	
α tripetala. w.					
6 acuminata. DC. (<i>pa.</i>) bluish-flowered.	1736.	5. 7.	H. $\frac{1}{2}$.	Bot. mag. 2427.
β Candolii. (<i>st.</i>) De Candolle's.	Sav.bi.ital.1819.n.47.ic.
7 auriculata. DC. (<i>wh.</i>) ear-leaved.	1786.	4. 5.	H. $\frac{1}{2}$.	Bot. mag. 1206.
8 pyramidata. DC. (<i>wh.</i>) small ear-leaved.	1811.	—	H. $\frac{1}{2}$.	Bot. reg. 407.
9 macrophylla. DC. (<i>wh.</i>) large-leaved.	1800.	6. 7.	H. $\frac{1}{2}$.	Bot. mag. 2189.
10 cordata. DC. (<i>st.</i>) heart-leaved.	1801.	—	H. $\frac{1}{2}$.	Bot. reg. 325.
11 conspicua. P.L. (<i>wh.</i>) Youlan.	China.	1789.	2. 5.	H. $\frac{1}{2}$.	Bot. mag. 1621.
<i>Yulan.</i> DC. Bonpl. nav. t. 20.					
12 Soulangeana. (<i>p.w.</i>) Soulange's.	Hybrid.	1826.	—	H. $\frac{1}{2}$.	Swt. br. fl. gar. 260.
13 Kòbus. DC. (<i>p.w.</i>) slender.	China.	1804.	3. 4.	H. $\frac{1}{2}$.	Par. lond. 87.
<i>tomentosa.</i> w. græciliſ. P.L.					
14 obovata. DC. (<i>p.w.</i>) obovate-leaved.	1790.	4. 6.	H. $\frac{1}{2}$.	Bot. mag. 390.
<i>purpurea.</i> B.M. <i>discolor.</i> V.					
15 fuscata. DC. (<i>br.</i>) brown-stalked.	1789.	—	G. $\frac{1}{2}$.	1008.
16 ammonœfolia. P.L. (<i>br.</i>) small-flowered.	1804.	—	G. $\frac{1}{2}$.	Par. lond. 5.
17 pumila. DC. (<i>wh.</i>) dwarf.	1786.	1. 12.	G. $\frac{1}{2}$.	Bot. mag. 977.
LIRIODE'NDRON. W. TULIP-TREE.	Polyandria Polygynia. L.				
<i>tulipifera.</i> w. (<i>ye.</i>) common.	N. America.	1663.	6. 7.	H. 5.	Bot. mag. 275.
α acutiloba. M. (<i>ye.</i>) acute-lobed.	—	—	H. 5.	
β obtusiloba. M. (<i>ye.</i>) blunt-lobed.	—	—	H. 5.	

ORDO V.

ANONACEÆ. DC. syst. nat. 1. p. 463.

AN'ONA. DC.	CUSTARD-APPLE. Polyandria Polygynia. L.	W. Indies.	1656.	6. 8.	S. $\frac{1}{2}$.
1 muricata. DC. (<i>st.</i>) Soursop.	Mexico.	1826.	S. $\frac{1}{2}$.	Jacq. obs. 1. t. 5.
2 purpurea. DC. (<i>pu.</i>) purple.	Cumana.	—	S. $\frac{1}{2}$.	Dunal. mon. p. 64. t. 2.
3 Humboldtii. DC. (<i>ye.</i>) Humboldt's.	W. Indies.	1824.	6. 8.	S. $\frac{1}{2}$.	— p. 64. t. 3.
4 laurifolia. DC. (<i>wh.</i>) laurel-leaved.	1731.	—	S. $\frac{1}{2}$.	Catesb. car. 2. t. 67.
5 palustris. DC. (<i>ye.</i>) shining-leaved.	Guiana.	1820.	—	S. $\frac{1}{2}$.	Sloan. hist. 2. t. 228. f. 1.
6 longifolia. DC. (<i>pu.</i>) long-leaved.	Cayenne.	1822.	—	S. $\frac{1}{2}$.	Aub. gui. 1. t. 248.
7 punetata. DC. (<i>st.</i>) spotted-fruited.	Guiana.	1803.	6. 8.	S. $\frac{1}{2}$.	— 1. t. 247.
8 paludosa. DC. (<i>gr.</i>) marsh.	S. America.	1731.	—	S. $\frac{1}{2}$.	— 1. t. 246.
9 squamosa. DC. (<i>st.</i>) sweetsop.	St. Thomas.	1823.	S. $\frac{1}{2}$.	Jacq. obs. 1. t. 6. f. 1.
10 cinerea. DC. (<i>st.</i>) cinereous.	S. America.	1739.	7. 8.	S. $\frac{1}{2}$.	Bot. mag. 2011.
11 Cherimolia. DC. (<i>wh.</i>) Cherimoli.	1690.	5. 7.	S. $\frac{1}{2}$.	Catesb. car. 2. t. 86.
<i>tripetala.</i> w. B.M.	Guiana.	S. $\frac{1}{2}$.	Rumph. amb. 1. t. 45.
12 reticulata. DC. (<i>st.</i>) netted.	Carolina.	1774.	7. 8.	S. $\frac{1}{2}$.	Catesb. car. 2. t. 64.
13 mucosa. DC. (<i>st.</i>) narrow-leaved.	Mauritius.	1825.	—	S. $\frac{1}{2}$.	Dunal. mon. p. 75. t. 6.
14 glabra. DC. (<i>st.</i>) smooth-fruited.	1824.	—	S. $\frac{1}{2}$.	— p. 76. t. 7.
15 grandiflora. DC. (<i>wh.</i>) great-flowered.	Ceylon.	—	—	S. $\frac{1}{2}$.	Deles. ic. sel. 1. t. 86.
16 amplexicaulis. DC. (<i>pu.</i>) stem-clasping.	Senegal.	1822.	—	S. $\frac{1}{2}$.	Gært. fr. 2. t. 125. f. 1.
17 asiatica. DC.	—	—	S. $\frac{1}{2}$.	
18 senegalensis. DC. (<i>st.</i>) Senegal.	—	—	S. $\frac{1}{2}$.	Dun. mon. t. 9.
MONOD'ORA. DC.	MONODORA. Polyandria Monogynia.	Jamaica.	—	—	Duh. arb. ed. 2. v. 2. t. 25.
myristica. DC. (<i>wh.</i>) Nutmeg.	—	—	S. $\frac{1}{2}$.	Dun. mon. t. 10.
ASIM'INA. DC.	ASIM'INA. Polyandria Polygynia.	—	—	Dun. mon. t. 11.
1 parviflora. DC. (<i>pu.</i>) small-flowered.	N. America.	1806.	4. 5.	H. $\frac{1}{2}$.	
2 triloba. DC. (<i>pu.</i>) trifid-fruited.	1736.	—	H. $\frac{1}{2}$.	
<i>Annona triloba.</i> w. Ph.					
3 pygmæa. DC. (<i>wh.</i>) dwarf.	1812.	—	H. $\frac{1}{2}$.	
<i>Orchidocarpum pygmaeum.</i> M.					
4 grandiflora. DC. (<i>wh.</i>) large-flowered.	Florida.	1820.	—	H. $\frac{1}{2}$.	
<i>Porcælia grandiflora.</i> P.S.					

UV'ARIA. DC.	UV'ARIA.	Polyandria Polygynia. L.					
1 zeylànica. DC. (pu.) Ceylon.	E. Indies.	1794.	S. h.	Lam. ill. t. 495. f. 2.		
2 Gærtneri. DC.	Gærtner's.	S. h.	t. 495. f. 3.	
3 lùtea. DC. (st.) yellow-fruited.	1822.	S. h.	Roxb. cor. 1. t. 36.		
4 tomentòsa. DC. (br.) woolly.	S. h.	I. t. 35.	
5 velutina. DC. (br.) velvetty.	E. Indies.	1825.	S. h.			
6 lucida. Boj.	shining.	E. Africa.	S. h.			
UN'ONA. DC.	UN'ONA.	Polyandria Polygynia. L.					
1 Nàrum. DC. (re.) Rheede's.	E. Indies.	S. h.	Rheed. mal. 2. t. 9.	
2 fuscâta. DC. (br.) brown-flower'd. Guiana.	1820.	S. h.				
3 acuminâta. DC. (br.) taper-pointed.	S. h.				
4 odorâta. DC. (st.) sweet-scented.	Java.	1804.	5. 7.	S. h.	Lam. ill. t. 495. f. 1.		
5 longifòlia. DC. (ye.) long-leaved.	E. Indies.	1820.	S. h.	Sonn. voy. 4. t. 131.		
6 aromâtica. DC. (pu.) aromatic.	Guiana.	S. h.	Aub. gui. 2. t. 243.		
7 æthiòpica. DC. (wh.) Ethiopian.	Sierra Leone.	1822.	S. h.	Math. com. 1. p. 434. ic.		
8 oxypétala. DC. (br.) sharp-petaled.	S. h.				
ARTAB'OTRYS. Br.	ARTAB'OTRYS.	Polyandria Polygynia. L.					
odoratissimus. B.R. (b.) sweet-scented.	China.	1758.	6. 7.	S. h.	Bot. reg. 423.	
<i>Annôna hexapétala. w.</i>	<i>Uvâria odoratissima. H.B.</i>						
XYL'OPIA. DC.	BITTER-WOOD.	Polyandria Polygynia. L.					
1 muricâta. DC. (br.) rough-fruited.	W. Indies.	1793.	S. h.	Brown. jam. 250. t. 5. f. 2.		
2 frutescens. DC. (br.) shrubby.	Guiana.	1823.	S. h.	Aub. gui. 1. t. 292.		
3 glâbra. DC. (br.) smooth-fruited.	W. Indies.	S. h.	Dun. mon. t. 19.		
GUATT'ERIA. DC.	GUATT'ERIA.	Polyandria Polygynia. L.					
1 cerasóides. DC. (br.) Cherry-fruited.	E. Indies.	1820.	S. h.	Roxb. cor. 1. t. 33.		
2 suberôsa. DC. (wh.) cork-barked.	S. h.	I. t. 34.		
3 rûfa. DC. (br.) rufous.	India.	4. 8.	S. h.	Bot. reg. 836.		
4 virgâta. DC. (wh.) Lance-wood.	Jamaica.	1793.	—	S. h.	Dun. mon. t. 31.		
<i>Uvâria lanceolata. w.</i>							
5 laurifòlia. DC. (wh.) Laurel-leaved.	1820.	S. h.	t. 32.	
<i>Uvâria laurifòlia. Sw.</i>							

ORDO VI.

MENISPERMACEÆ. DC. prodr. 1. p. 95.

Tribus II. MENISPERMEÆ. DC.

CO'CULUS. DC.	CO'CULUS.	Dioecia Hexandria.					
1 rotundifolius. DC. (gr.) round-leaved.	1820.	S. h.		
2 cordifolius. DC. (gr.) heart-leaved.	E. Indies.	1822.	S. h.	Rheed. mal. 7. t. 21.	
3 suberòsa. DC. (gr.) cork-barked.	1790.	S. h.	Gært. fr. 1. t. 70. f. 7.	
4 Plukènètii. DC. (gr.) Plukènet's.	S. h.	Pluk. man. 52.t 345.f.2.	
5 incânus. L.T. (gr.) hoary.	1820.	S. h.		
6 tomentòsus. L.T. (gr.) woolly-leaved.	S. h.		
7 críspus. L.T. (gr.) curled.	1822.	S. h.	Rumph. amb. 5 t. 44. f. 1.	
8 palmátus. DC. (gr.) palmated.	S. h.	Berr. in As. res. 10. t. 5.	
9 orbiculátus. DC. (gr.) orbicular-leaved.	1790.	S. h.	Rheed. mal. 11. t. 62.	
10 laurifolius. DC. (gr.) Laurel-leaved.	1820.	S. h.		
COSCI'NIUM. C.L.T.	COSCI'NIUM.	Dioecia Hexandria.					
fenestrâtum. L.T. (gr.) knotted.	Ceylon.	1820.	S. h.	Gært. fr. 1. t. 46. f. 5.	
TILIAC'ORA. C.L.T.	TILIAC'ORA.	Dioecia Hexandria.					
racemosa. L.T. (gr.) racemed.	E. Indies.	S. h.		
WENDLA'NDIA. W.	WENDLA'NDIA.	Heptandria Polygynia. W.					
populifòlia. w. (wh.) Poplar-leaved.	Carolina.	1759.	6. 7.	H. h.	Wend. h. herr. 3. t. 16.	
<i>Menispérnum Carolinum. L.</i>	<i>Cocculus Carolinus. DC.</i>						
CISSA'MPELOS. DC.	CISSA'MPELOS.	Dioecia Monadelphia. L.					
1 Paréira. DC. (gr.) velvet-leaf.	S. America.	1733.	7. 8.	S. h.	Lam. ill. t. 830.	
2 microcárpa. DC. (gr.) small-fruited.	W. Indies.	1823.	S. h.		
3 hirsuta. DC. (gr.) hairy.	Nepaul.	1820.	G. h.		
4 Mauritiâna. DC. (gr.) hispid-branch'd.	Mauritius.	S. h.		
5 Caapêba. DC. (gr.) nerved-leaved.	S. America.	1733.	7. 8.	S. h.	Plum. ic. 67. f. 2.	
6 ? capénis. DC. (gr.) Cape.	C. B. S.	1775.	G. h.		
MENISPER'MUM. DC.	MOON-SEED.	Dioecia Dodecadria. L.					
1 canadénse. DC. (st.) Canadian.	N. America.	1691.	6. 7.	H. h.	Bot. mag. 1910.	

2 dàuricum. DC.	(st.) Daurian.	Dahuria.	1823. H. $\frac{1}{2}$ —. Deles. ic. sel. 1. t. 100.
3 smilacinum. DC.	(st.) Smilax-leaved.	Carolina.	1776. G. $\frac{1}{2}$ —. Jacq. ic. 3. t. 629.
<i>Cissampelos smilacina</i> . W.				
4 Lyoni. Ph.	(st.) Lyon's.	N. America.	1812.	6. 7. H. $\frac{1}{2}$ —.
AB'UTA. DC.	AB'UTA.	<i>Menispermum</i> sp. w.		
rufescens. DC.	(pu.) brown-leaved.	Guiana.	1822. S. $\frac{1}{2}$ —. Aub. gui. 1. t. 250.

Tribus III. SCHIZANDREÆ. DC. prodri. 1. p. 104.

SCHIZA'NDRA. DC.	SCHIZA'NDRA.	Monœcia Pentandria.		
coccinea. DC.	(sc.) scarlet-flower'd.	Carolina.	1806.	6. 7. G. $\frac{1}{2}$ —. Bot mag. 1413.

ORDO VII.

BERBERIDEÆ. DC. syst. nat. v. 2. p. 1.

BE'RBERIS. W.

	BARBERRY.	Hexandria Monogynia.	L.	
1 vulgàris. DC.	(ye.) common.	England. 4. 5. H. $\frac{1}{2}$.	Eng. bot. 49.
α rùbra.	(ye.) red-fruited.	—	— H. $\frac{1}{2}$.	
β lùtea. DC.	(ye.) yellow-fruited.	—	— H. $\frac{1}{2}$.	
γ violacea. DC.	(ye.) violet-fruited.	—	— H. $\frac{1}{2}$.	
δ purpurea. DC.	(ye.) purple-fruited.	—	— H. $\frac{1}{2}$.	
ϵ nigra. DC.	(ye.) black-fruited.	Levant.	— H. $\frac{1}{2}$.	
ζ álba. DC.	(ye.) white-fruited.	—	— H. $\frac{1}{2}$.	
ϑ asperma. DC.	(ye.) stoneless.	—	— H. $\frac{1}{2}$.	
2 ibérica.	(ye.) Iberian.	Iberia.	1790. — H. $\frac{1}{2}$.	
<i>vulgàris</i> β <i>ibérica</i> . DC.				
3 emarginata. DC.	(ye.) emarginate.	Siberia.	1820. — H. $\frac{1}{2}$.	
4 canadensis. DC.	(ye.) Canadian.	Canada.	1759. 4. 6. H. $\frac{1}{2}$.	
5 sinensis. DC.	(ye.) China.	China.	1800. — H. $\frac{1}{2}$.	
6 aristata DC.	(ye.) awned.	Nepaul.	1820. 6. 8. H. $\frac{1}{2}$.	Wats. dend. brit. t. 26. Bot. mag. 2549.
<i>Chitria</i> . Bot. reg. 729.				
7 crética. DC.	(ye.) Cretan.	Crete.	1759. 5. 6. H. $\frac{1}{2}$.	Flor. græc. t. 342.
8 ilicifolia. DC.	(ye.) Holly-leaved.	Terra del Fuego.	1791. 7. 8. H. $\frac{1}{2}$.	
9 asiática. DC.	(ye.) Asiatic.	Nepaul.	1820. 6. 8. H. $\frac{1}{2}$.	Deles. ic. sel. 2. t. 1.
10 Wallichiana. DC.	(ye.) Wallich's.	—	— 6. 7. H. $\frac{1}{2}$.	
11 buxifolia. DC.	(ye.) Box-leaved.	Magellan.	1827. G. $\frac{1}{2}$.	Lam. ill. t. 253. f. 3.
12 empetrifolia. DC.	(ye.) Emptetrum-l.	—	— G. $\frac{1}{2}$.	
13 heterophylla. DC.	(ye.) various-leaved.	—	— 5. 6. H. $\frac{1}{2}$.	Hook. exot. fl. t. 14.
14 sibirica. DC.	(ye.) Siberian.	Siberia.	1790. 6. 7. H. $\frac{1}{2}$.	Bot. reg. 487.
15 inérmis. DC.	(ye.) spineless.	Magellan.	1827. G. $\frac{1}{2}$.	
16 pinnata. B.R.	(ye.) pinnate-leaved.	America.	1820. 3. 5. G. $\frac{1}{2}$.	Bot. reg. 702.
<i>Mahonia fascicularis</i> . DC.				
17 nepens. B.R.	(ye.) creeping-rooted.	N. America.	1822. 4. 6. H. $\frac{1}{2}$.	Bot. reg. 1176.
18 nervosa. P.	(ye.) nerved-leaved.	—	— H. $\frac{1}{2}$.	Pursh. fl. am. sept. 1.t.5.
NAND'INA. DC.	NAND'INA.	Hexandria Monogynia.		
domestica. DC. (wh.)	garden.	China.	1804. 6. 7. G. $\frac{1}{2}$.	
LEO'NTICE. DC.	LEO'NTICE.	Hexandria Monogynia.	L.	
1 Chrysogonium. DC.	(ye.) Oak-leaved.	Levant.	1740. 3. 6. H. $\frac{1}{2}$.	Barrel. ic. 1113.
2 Leontopetalum. DC.	(ye.) Lion's-leaf.	—	— 1597. 4. 5. H. $\frac{1}{2}$.	Lam. ill. t. 254. f. 1.
3 vesicaria. DC.	(ye.) bladder-capsul'd.	Siberia.	1822. — H. $\frac{1}{2}$.	Pal. a.p. 1779. 2.t.9.f.4.
4 altaica. DC.	(ye.) Altay.	—	— H. $\frac{1}{2}$.	— p. 257. t. 8.f.1.2.3.
5 thalicroides. DC.	(ye.) Columbine-l.	N. America.	1755. 5. H. $\frac{1}{2}$.	Mich. amer. 1. t. 21.
<i>Caulophyllum thalictroides</i> . M.				
EPIMEDIUM. DC.	BARRENWORT.	Tetrandria Monogynia.	L.	
alpinum. DC.	(pu.) Alpine.	England. 4. 5. H. $\frac{1}{2}$.	Eng. bot. 438.
DIPHYLL'EIA. DC.	DIPHYLL'EIA.	Hexandria Monogynia.	L.	
cymosa. DC.	(wh.) blue-berried.	N. America.	1812. 5. 6. H. $\frac{1}{2}$.	Bot. mag. 1666.

ORDO VIII.

PODOPHYLLEÆ. DC. syst. nat. 2. p. 31.

Tribus I. PODOPHYLLEÆ.

PODOPHYLLUM. DC. DUCK's-FOOT. Polyandria Monogynia. L.					
peltatum. DC. (wh.) May-apple.	N. America.	1664.	5.	H. 24.	Bot. mag. 1819.
JEFFERS'ONIA. DC. JEFFERS'ONIA. Octandria Monogynia.					
diphylla. DC. (wh.) two-leaved.	N. America.	1792.	5.	H. 24.	Bot. mag. 1513.

Podophyllum diphyllum. w.

Tribus II. HYDROPELTIDEÆ.

CABO'MBA. DC.	CABO'MBA. Hexandria Digynia. W.				
aquatica. DC. (ye.) water.	Carolina.	1823.	S.w. 24.	Aubl. gui. 1. t. 124.
<i>Néctris peltata.</i> Ph.					
HYDROPELTIS. DC. HYDROPELTIS.	Polyandria Polygynia.				
purpurea. DC. (pu.) purple.	N. America.	1798.	7. 8.	F.w. 24.	Bot. mag. 1147.
<i>Brasenia peltata.</i> Ph.					

ORDO IX.

NYMPHÆACEÆ. DC. syst. nat. 2. p. 34.

Tribus I. NELUMBONEÆ.

NELU'MBIUM. DC.	SACRED-BEAN. Polyandria Polygynia. L.				
1 speciosum. DC. (bh.) Indian.	India.	1787.	6. 8.	S.w. 24.	Bot. mag. 903.
<i>Cyamus Nelumbo.</i> Sm. Exot. bot. 1. t. 31. 32.					
2 Támara.	(ro.) Rheede's.	—	—	S.w. 24.	Rheed. mal. xi. t. 30.
3 cáspicum. F.	(ro.) Fischer's.	Volga.	1821.	— S.w. 24.	
4 luteum. DC. (ye.) yellow.	N. America.	1810.	—	F.w. 24.	Poit.a.m. 13. t. 29.f. 42.
5 jamaicense. DC. Jamaica.	W. Indies.	1823.	—	S.w. 24.	

Tribus II. NYMPHÆEEÆ.

EURY'ALE. DC.	EURY'ALE. Polyandria Monogynia. L.				
férox. DC. (re.) prickly.	India.	1809.	7. 9.	S.w. 24.	Bot. mag. 1447.
<i>Anneslea spinosa.</i> And. bot. rep. 618.					
NYMPHE'A. DC.	WATER-LILY. Polyandria Monogynia. L.				
1 scutifolia. DC. (bl.) Cape blue.	C. B. S.	1792.	6. 9.	S.w. 24.	Bot. mag. 552.
<i>cærulaea.</i> Bot. rep. 197.					
2 cœrulea. DC. (bl.) Egyptian blue.	Egypt.	1812.	—	S.w. 24.	Vent. malm. t. 6.
<i>stellata.</i> var. Bot. mag. 2058.					
3 cyanea. H.B.	(bl.) Indian-blue.	E. Indies.	1809.	— S.w. 24.	
<i>Cahlâra.</i> Donn. cant. ed. 7.					
4 stellata. DC.	(bl.) star-flowered.	—	1803.	— S.w. 24.	Andr. rep. 330.
5 edulis. DC.	(wh.) eatable.	—	— S.w. 24.	
<i>esculenta.</i> H.B.					
6 rûbra. DC.	(re.) red-flowered.	—	1803.	7. 8. S.w. 24.	Bot. mag. 1280.
7 rôsea.	(ro.) rose-coloured.	—	—	— S.w. 24.	— 1364.
<i>rûbra.</i> β <i>rôsea.</i> DC.					
8 pubescens. DC.	(wh.) Indian Lotus.	—	—	5. 8. S.w. 24.	Rheed. h. mal. xi. t. 26.
9 Lôtus. DC.	(wh.) Egyptian.	Egypt.	1802.	6. 9. S.w. 24.	Del. fl. egyp. ill. t. 60. f. 1.
10 thérmalis. DC.	(wh.) Hungarian.	Hungary.	—	— S.w. 24.	Bot. mag. 797.
<i>Lôtus.</i> And. bot. rep. 391.					

NYMPHÆACEÆ.

11	ámpla. DC.	(wh.) West-Indian.	W. Indies.	6. 9. S.w. 2.	Plum. mss. 123. t. 4.
12	versicolor. DC.	(bh.) changeable	E. Indies.	1807.	— S.w. 2.	Bot. mag. 1189.
13	reniformis. DC.	(wh.) kidney-leaved.	Carolina.	1823.	7. G.w. 2.	Deles. ic. sel. 2. t. 5.
14	álba. DC.	(wh.) white.	Britain.	6. 7. H.w. 2.	Eng. bot. 160.
15	odorata. DC.	(wh.) sweet-scented.	N. America.	1786.	7. H.w. 2.	Bot. mag. 819.
16	minor. DC.	(wh.) smaller odorata. β minor. B.M.	—	1812.	— H.w. 2.	— 1652.
17	nítida. DC.	(wh.) cup-flowered.	Siberia.	1809.	7. 8. H.w. 2.	Bot. mag. 1359.
18	pygmæa. DC.	(wh.) pygmy. <i>Castália pygmæa</i> . Sal. par. lond. 68.	China.	1805	5. 9. H.w. 2.	— 1525.
19	blanda. DC.	(wh.) handsome.	W. Indies.	1827. S.w. 2.	N'UPHAR. DC. N'UPHAR.
			Polyandria Monogynia. L.			
1	lútea. DC.	(ye.) common. <i>Nymphæa lútea</i> . w.	Britain.	6. 7. H.w. 2.	Eng. bot. 159.
2	púmila. DC.	(ye.) small-yellow. minima. E.B.	Scotland	— H.w. 2.	— 2292.
3	Kalmiàna. DC.	(ye.) Canadian.	N. America.	1807.	7. 8. H.w. 2.	Bot. mag. 1243.
4	ádvena. DC.	(ye.) stripe-flowered. <i>Nymphæa ádvena</i> , w.	—	1772.	— H.w. 2.	Bot. mag. 684.

ORDO X.

SARRACENIÆ.

SARRAC'ENIA. W.		SIDE-SADDLE-FLOWER.	Polyandria Monogynia. L.		
1	purpurea. w.	(pu.) purple.	N. America.	1640.	6. 7. H. 2.
2	rúbra. w.	(re.) red-flowered. psittácinæ. M?	—	1786.	— H. 2.
3	variolaris. w.	(st.) hook-leaved. adúnca. Ex. bot. t. 53. minor w.	—	1803.	— H. 2.
4	fláva. w.	(ye.) yellow.	—	1752.	— H. 2.

ORDO XI.

PAPAVERACEÆ. DC. syst. nat. 2. p. 67.

PAP'AVER. DC.		POPPY.	Polyandria Monogynia.		
1	nudicáule. DC.	(ye.) naked-stalked.	Siberia.	1730.	6. 8. H. 2.
	β gláratum. DC.	(ye.) smooth-stalked.	—	—	H. 2.
	γ radicátum. DC.	(ye.) small hairy.	Norway.	—	H. 2.
2	rúbro-aurantiacum.	F. red orange.	Dahuria.	1822.	— H. 2.
3	microcárpus. DC. (br.)	small-capsuled.	Kamtschatka.	—	H. 2.
4	pyréniacum. DC.	(ye.) Pyrenean. aurantiacum. Lois.	Pyrenees.	—	H. 2.
	α lúteum. DC.	(ye.) yellow-flowered.	—	—	H. 2.
	β puniceum. DC.	(re.) red-flowered.	—	—	H. 2.
5	alpinum. DC.	(wh.) Alpine.	Austria.	1759.	— H. 2.
6	hybridum. DC.	(re.) mongrel.	England.	6. 7. H. 2.
7	Argemône. DC.	(re.) rough. β marítimum. With one-flowered.	Britain.	H. 2.
	dùbium. DC.	(re.) smooth.	—	—	H. 2.
9	Rhœas. DC.	(re.) common-corn.	—	—	H. 2.
	α maculatum.	(re.) spot-petaled.	—	—	H. 2.
	β coccineum.	(sc.) scarlet-flowered.	—	—	H. 2.
	γ cárneum.	(fl.) flesh-coloured.	—	—	H. 2.
	δ álbum.	(wh.) white-flowered.	—	—	H. 2.
	ϵ álbo-marginátum.	white-edged.	—	—	H. 2.
	ζ coccíneo-marginátum.	scarlet-edged.	—	—	H. 2.

Bot. mag. 1633.

Bot. mag. t. 2344.

Seg. ver. 1. t. 4. f. 4.

Barr. ic. t. 764.

Swt. br. fl. gar. 247.

Eng. bot. 43.

— 643.

Eng. bot. 644.

— 645.

Curt. lond. t. 32.

Tab. ic. f. 1.

Barr. obs. ic. t. 763.

Wein. phyt. t. 788. f. d.

— t. 790. f. e.

— t. 790. f. b.

10	<i>lævigatūm.</i> DC.	(<i>re.</i>) slender.	Caucasus.	1823.	6. 7.	H. ⊙.	
11	<i>arenārium.</i> DC.	(<i>re.</i>) sand.	Siberia.	1825.	6. 8.	H. ⊙.	Vig. diss. 39.n.4.t.1.f.1.
12	<i>Roubiæ.</i> DC.	(<i>re.</i>) Roubieu's.	Montpelier.	1823.	—	H. ⊙.	Bot. reg. 134.
13	<i>floribundūm.</i> DC.	(<i>re.</i>) many-flowered.	Levant.	1815.	6. 7.	H. ♂.	Swt. br. fl. gar. 173.
14	<i>hórridūm.</i> DC.	(<i>co.</i>) prickly-stalk'd.	N. S. W.	1825.	6. 8.	H. ⊙.	Bot. mag. 57.
15	<i>orientálē.</i> DC.	(<i>re.</i>) oriental.	—	1714.	5. 6.	H. ♀.	Lind. coll. t. 23.
16	<i>bracteátūm.</i> DC.	(<i>re.</i>) bracted.	Caucasus.	1821.	5. 8.	H. ♀.	Swt. br. fl. gar. 172.
17	<i>setígerūm.</i> DC.	(<i>wh.</i>) bristle-pointed.	Italy.	1824.	—	H. ⊙.	Lam. ill. t. 451.
18	<i>somniférūm.</i> DC.	(<i>va.</i>) garden.	England.	7. 8.	H. ⊙.	Lob. ic. t. 274. f. 1.
	* <i>nigrum.</i> DC.	(<i>va.</i>) black-seeded.	—	H. ⊙.	Swert. flor. 2. t. 22. f. 1.
	β <i>flore-pleno.</i>	(<i>re.</i>) double-red.	—	H. ⊙.	Wein. phyt. t. 795. f. c.
	γ <i>fimbriatum.</i>	(<i>va.</i>) fringed-flowered.	—	H. ⊙.	Eng. bot. 2145.
	** <i>album.</i> DC.	(<i>wh.</i>) white-seeded.	England.	—	H. ⊙.	Swert. flor. 2. t. 22. f. 2.
	β <i>flore-pleno.</i>	(<i>wh.</i>) double-white.	—	—	H. ⊙.	
	γ <i>variegatum.</i>	(<i>v.</i>) variegated.	—	—	H. ⊙.	
19	<i>caucásicum.</i> DC.	(<i>re.</i>) Caucasean.	Caucasus.	1813.	6. 7.	H. ⊙.	Bot. mag. 1675.
20	<i>armeniācum.</i> DC.	(<i>re.</i>) Armenian.	Armenia.	1815.	—	H. ⊙.	
	<i>Argemone armeniacu.</i> w.						
	<i>ARGEMONE.</i> DC.	<i>ARGEMONE.</i>	Polyandria Monogynia.				
1	<i>mexicāna.</i> DC.	(<i>ye.</i>) Mexican.	Mexico.	1592.	7. 8.	H. ⊙.	Bot. mag. 243.
2	<i>ochroleúca.</i> B.F.G.(<i>st.</i>)	straw-coloured.	—	1827.	7. 10.	H. ⊙.	Swt. br. fl. gar. 242.
3	<i>albiflóra.</i> B.M.	(<i>wh.</i>) white-flowered.	Louisiana.	1820.	—	H. ♀.	Bot. mag. 2342.
4	<i>grandiflóra.</i> B.F.G.(<i>w.</i>)	large-flowered.	Mexico.	1827.	7. 10.	H. ♀.	Swt. br. fl. gar. 226.
	<i>MECONO'PSIS.</i> DC.	<i>MECONO'PSIS.</i>	Polyandria Monogynia.				
	<i>cámrifica.</i> DC.	(<i>st.</i>) Welsh.	Wales.	5. 8.	H. ♀.	Eng. bot. 66.
	<i>Papaver cámricum.</i> w.						
	<i>SANGUIN'ARIA.</i> DC.	<i>PUCCOON.</i>	Polyandria Monogynia.				
	<i>canadénsis.</i> DC.	(<i>wh.</i>) Bloodwort.	N. America.	1680.	3. 4.	H. ♀.	Bot. mag. 162.
	<i>MACLEA'YA.</i> B.	<i>MACLEA'YA.</i>	Polyandria Monogynia.				
	<i>cordátā.</i> B.	(<i>bh.</i>) heart-leaved.	China.	1795.	5. 8.	H. ♀.	Bot. mag. 1905.
	<i>Boccónia cordátā.</i> DC.						
	<i>BOCC'ONIA.</i> DC.	<i>Bocc'ONIA.</i>	Dodecandria Monogynia.				
1	<i>frutéscens.</i> DC.	(<i>gr.</i>) Tree Celandine.	W. Indies.	1739.	1. 4.	S. ♀.	Lodd. bot. cab. t. 83.
2	<i>integritifólia.</i> DC.	(<i>gr.</i>) entire-leaved.	Peru.	1822.	S. ♀.	H. et B. nov. pl. 1.t. 35.
	<i>ESCHSCHO'LTZIA.</i> D.	<i>ESCHSCHO'LTZIA.</i>	Polyandria Monogynia.				
	<i>californica.</i> DC.	(<i>ye.</i>) Menzies'.	California.	1826.	7. 10.	H. ♀.	Swt. br. fl. gar. 265.
	<i>HUNNEMA'NNIA.</i> B.F.G.	<i>HUNNEMA'NNIA.</i>	Polyandria Monogynia.				
	<i>fumariæfólia.</i> B.F.G.(<i>y.</i>)	Fumitory-leav'd Mexico.	1827.	7. 10.	H. ♀?	Swt. br. fl. gar. 276.	
	<i>RÖM'ERIA.</i> DC.	<i>RÖM'ERIA.</i>	Polyandria Monogynia.				
1	<i>hybrida.</i> DC.	(<i>re.</i>) violet-flowered.	England.	5. 7.	H. ⊙.	Eng. bot. 201.
	<i>Gláciūm violáceum.</i> H.K.						
2	<i>réfracta.</i> DC.	(<i>re.</i>) pendulous-fruit.	Tauria.	1823.	—	H. ⊙.	Deles. ic. sel. 2. t. 8.
	<i>GLAUCIUM.</i> DC.	<i>HORN-POPPY.</i>	Polyandria Monogynia.				
1	<i>flávum.</i> DC.	(<i>ye.</i>) yellow.	Britain.	6. 10.	H. ♂.	Eng. bot. t. 8.
	<i>luteum.</i> H.K.	<i>Chelidónium gláciūm.</i> L.					
2	<i>fulvum.</i> DC.	(<i>or.</i>) orange.	S. Europe.	1802.	8. 9.	H. ♀.	Swt. br. fl. gar. 35.
3	<i>tricolor.</i> s.s.	(<i>v.</i>) three-coloured.	Poland.	1828.	6. 9.	H. ⊙.	
4	<i>corniculátum.</i> DC.	(<i>re.</i>) various-colour'd	Europe.	6. 7.	H. ⊙.	Eng. bot. 1433.
	<i>a phænićeum.</i> DC.	(<i>re.</i>) red-flowered.	England.	—	H. ⊙.	
	<i>β flaviflórum.</i> DC.	(<i>ye.</i>) yellow-flowered.	Tauria.	1823.	—	H. ⊙.	
	<i>CHELID'ONIUM.</i> DC.	<i>CELANDINE.</i>	Polyandria Monogynia. L.				
1	<i>május.</i> DC.	(<i>ye.</i>) common.	Britain.	4. 10.	H. ♀.	Eng. bot. 1581.
2	<i>grandiflórum.</i> DC.	(<i>ye.</i>) large-flowered.	Dahuria.	1823.	—	H. ♀.	Mill. ic. 1. t. 92. f. 2.
3	<i>laciniátum.</i> DC.	(<i>ye.</i>) jagged.	S. Europe.	—	H. ♀.	
	<i>HYPE'COUN.</i> DC.	<i>HYPE'COUN.</i>	Tetrandria Digynia. L.				
1	<i>procumbens.</i> DC.	(<i>ye.</i>) procumbent.	S. Europe.	1596.	6. 7.	H. ⊙.	Swt. br. fl. gar. 3.t. 217.
2	<i>péndulum.</i> DC.	(<i>ye.</i>) pendulous.	S. France.	1640.	—	H. ⊙.	Mill. ic. t. 250. f. 1.
3	<i>erectum.</i> DC.	(<i>ye.</i>) erect.	Siberia.	1759.	5. 6.	H. ♂.	Amm. ruth. 58. t. 9.

ORDO XII.

FUMARIACEÆ. DC. syst. nat. 2. p. 105.

DICL'YTRA. DC.	DICL'YTRA. Diadelphia Hexandria. L.					
1 cucullària. DC. (wh.) naked stalked.	N. America. 1731.	6. 7.	H. ¼.	Bot. mag. 1127.		
<i>Fumaria cucullaria</i> . B.M.						
β divaricàta. DC. (wh.) spreading-spur'd.	N. America.	—	— H. ¼.			
2 bracteòsa. DC. (wh.) leafy-bracted.	—	1823.	— H. ¼.			
3 formòsa. DC. (re.) bluish.	—	1796.	— H. ¼.	Bot. mag. 1335.		
4 exímia. DC. (re.) choice.	—	1812.	— H. ¼.	Bot. reg. 51.		
<i>Fumaria eximia</i> . B.R.						
5 tenuifòlia. DC. (bh.) slender-leav'd.	Kamtschatka. 1824.	5. 7.	H. ¼.	Deles. ic. sel. 2. t. 9. f. B.		
6 canadén sis. DC. (pu.) Canadian.	Canada. 1823.	—	H. ¼.			
7 lachenalíæflòra. DC. (p.) Lachenalia-fld.	Siberia. 1824.	H. ¼.	Rudol. m. sc. pet. t. 1. t. 19.		
ADL'UMIA. DC.	ADL'UMIA. Diadelphia Hexandria. L.					
cirrhòsa. DC. (bh.) tendrilled.	N. America. 1778.	6. 9.	H. ½.	Vent. choix. t. 19.		
<i>Fumaria fungòsa</i> . w.						
CYSTICA'PNOS. DC.	CYSTICA'PNOS. Diadelphia Hexandria. L.					
africana. DC. (bh.) African.	C. B. S. 1696.	6. 8.	H. ⊙.	Wein. phyt. t. 521. f. c.		
<i>Fumaria vesicarià</i> . w.						
CORY'DALIS. DC.	CORY'DALIS. Diadelphia Hexandria. L.					
1 pauciflòra. DC. (pu.) few-flowered.	Siberia. 1823.	H. ¼.			
2 Marschalliana. DC. (st.) Marschall's.	Tauria. —	4. 5.	H. ¼.			
3 tuberosa. DC. (pu.) hollow-rooted.	Europe. 1596.	2. 4.	H. ¼.	Bot. mag. 232.		
β albiflòra. (wh.) white-flowered.	—	—	H. ¼.	2340.		
<i>Fumaria cava. albiflora</i> . B.M.						
4 fabacea. DC. (pu.) Bean-leaved.	Germany. 1815.	—	H. ¼.	Horn. fl. dan. t. 1394.		
5 caucásica. DC. (pu.) Caucasean.	Caucasus. 1823.	—	H. ¼.			
6 bulbosa. DC. (pu.) solid-rooted.	Britain.	—	H. ¼.	Eng. bot. 1471.		
<i>Fumaria sólida</i> . E.B. <i>Corydalis Halleri</i> . W. en.						
7 angustifòlia. DC. (pu.) narrow-leaved.	Iberia. 1823.	—	H. ¼.			
8 bracteàta. DC. (st.) large-bracted.	Siberia. 1822.	—	H. ¼.			
9 nòbilis. DC. (st.) great-flowered.	—	1783.	5.	H. ¼.	Bot. reg. 395.	
10 Geb'èri. F. (bh.) Altay.	Altay. 1827.	5. 6.	H. ¼?			
11 paoniæfòlia. DC. (pu.) Paeony-leaved.	Siberia. 1823.	—	H. ¼.	Gmel. sib. 4. t. 34.		
12 glànca. DC. (bh.) glaucous.	N. America. 1683.	7. 8.	H. ⊙.	Bot. mag. 179.		
<i>Fumaria sempervirens</i> . w.						
13 strícta. DC. (pu.) upright.	Siberia. 1825.	5. 7.	H. ¼.			
14 sibírica. DC. (ye.) Siberian.	—	—	H. ¼.	Amm. ruth. n. 173. t. 20.		
15 impatiens. DC. (ye.) impatient.	Dahuria. 1823.	—	H. ⊙.			
16 àurea. DC. (ye.) golden.	N. America. 1812.	5. 7.	H. ½.	Bot. reg. 66.		
<i>Fumaria aurea</i> . B.R.						
17 lùtea. P.S. (ye.) yellow.	England.	4. 10.	H. ¼.	Eng. bot. 588.		
18 capnóides. P.S. (wh.) white-flowered.	S. Europe. 1596.	5. 10.	H. ½.	Pluk. alm. t. 90. f. 2.		
19 acáulis. DC. (st.) stalkless.	Germany. 1825.	7. 9.	H. ⊙.	Jacq. ic. 3. t. 554.		
20 uralénsis. DC. (ye.) Fischer's.	Siberia. 1823.	H. ¼.			
21 breviflòra. DC. (st.) short-flowered.	Kamtschatka. —	5. 10.	H. ⊙.	Eng. bot. 103.		
22 claviculata. DC. (wh.) climbing.	Britain.	—	H. ⊙.			
SARCOCA'PNOS. DC.	SARCOCA'PNOS. Diadelphia Hexandria. L.					
enneaphyllea. DC. (bh.) nine-leaved.	Spain. 1714.	5. 7.	H. ¼.	Barr. ic. t. 42.		
<i>Fumaria enneaphyllea</i> . w.						
FUM'ARIA. DC.	FUMITORY. Diadelphia Hexandria. L.					
1 spicata. DC. (re.) narrow-leaved.	S. Europe. 1714.	4. 6.	H. ⊙.	Lob. ic. t. 757. f. 2.		
2 capreolata. DC. (wh.) climbing.	Europe.	5. 9.	H. ⊙.	DC. ic. rar. 1. t. 34.		
β Burchellii. DC. (wh.) <i>Burchell's.</i>	C. B. S. 1812.	—	H. ⊙.			
3média. DC. (pu.) intermediate.	Britain.	—	H. ⊙.	Eng. bot. 943.		
capreolata. E.B.						
4 prehésibilis. K. (pu.) short-leaved.	Hungary. 1821.	6. 9.	H. ⊙.			
5 alexandrina. E.	Egypt. 1826.	—	H. ⊙.			
6 officinális. DC. (re.) common.	—	—	H. ⊙.	Eng. bot. 589.		
β grandiflòra. DC. (re.) large-flowered.	C. B. S. 1818.	—	H. ⊙.			
7 micrantha. DC. (wh.) small-blossom'd.	Spain. 1824.	4. 8.	H. ⊙.			
8 parviflòra. DC. (wh.) small-flowered.	England.	8. 9.	H. ⊙.	Eng. bot. 590.		
9 Vaillantii. DC. (pu.) Vaillant's.	France.	5. 6.	H. ⊙.	Vail. bot. par. t. 10. f. 6.		
10 densiflòra. DC. (pu.) close-flowered.	Montpellier. 1823.	—	H. ⊙.			
β albida. DC. (wh.) white-flowered.	—	—	H. ⊙.			

ORDO XIII.

CRUCIFERÆ. DC. syst. nat. 2. p. 139.

SUBORDO I. PLEURORHIZEÆ. DC.

Tribus I. ARABIDEÆ seu Pleurorhizeæ Siliquosæ. DC.

MATHIOLA. DC.	STOCK. Tetradymania Siliquosa. L.	5. 8.	H. ♂.	
1 simplicicáulis.	(sc.) Brompton.	H. ♂.	Besl. eyst. 2. t. 3. f. 3.
α coccinea.	(sc.) scarlet.	H. ♂.	— 2. t. 3. f. 2.
β álba.	(wh.) white.	H. ♂.	Eng. bot. 1935.
2 incànæ. DC.	(va.) queen. England.	5. 11.	H. ♀.	Wein. phyt. t. 642. f. 9.
α coccinea.	(sc.) scarlet.	H. ♀.	Besl. eyst. 2. t. 3. f. 2.
β purpurea.	(pu.) purple.	H. ♀.	— 2. t. 3. f. 1.
γ álba.	(wh.) white.	H. ♀.	J. Bauh. hist. 2. f. 1.
δ multipléx.	(va.) double-flowered.	H. ♀.	Dalech. lug. 802. f. 1.
3 ánnua. DC.	(va.) ten-week. S. Europe. 1731. 5. 11.	H. ♂.		Wein. phyt. t. 642. f. 2.
α coccinea.	(sc.) scarlet.	H. ♂.	Moris. ox. s. 3. t. 8. f. 2.
β purpurea.	(pu.) purple.	H. ♂.	
δ rúbra.	(re.) red.	H. ♂.	
ε rósea.	(ro.) rose-coloured.	H. ♂.	
ζ cárnea.	(fl.) flesh-coloured.	H. ♂.	
η fuscata.	(br.) brown.	H. ♂.	
δ fúlva.	(br.) tawny.	H. ♂.	
ι lilacina.	(li.) lilac.	H. ♂.	
κ álba.	(wh.) white.	H. ♂.	
λ variegata.	(v.) variegated.	H. ♂.	
μ flôre plêno.	(va.) double-flowered.	H. ♂.	
glâbra. DC.	(va.) Wall-flower-ld.	H. ♀.	
α álba.	(wh.) white-flowered.	H. ♀.	
β purpurea.	(pu.) purple.	H. ♀.	
γ coccinea.	(sc.) scarlet.	H. ♀.	
δ flôre plêno.	(va.) double-flowered.	H. ♀.	
5 græca. DC.	(wh.) smooth-leaved. Greece.	H. ♂.	
6 fenestrâlis. DC.	(re.) window. Crete. 1759. 7. 8	F. ♀.		Jacq. vind. 2. t. 179.
7 sinuâta. DC.	(pu.) great sea. England.	H. ♂.	Eng. bot. 462.
8 acâulis. DC.	(pu.) stalkless. Egypt. 1823. —	H. ♂.		Pall. it. 1. app. 117.t.O.
9 torulôsa. DC.	(pu.) twisted-podded. C. B. S. 1816. —	G. ♀.		Bot. mag. 1711.
10 tatárica. DC.	(br.) Tartarian. Tartary. 1826. —	H. ♀.		
11 odoratissima. DC.	(br.) sweet-scented. Tauria. 1797. 6. 7.	G. ♀.		
β tanaciénsis. DC.	(br.) fragrant. Tanaïm. 1822. —	G. ♀.		
12 vâria. DC.	(br.) entire-leaved. Greece.	5. 7.	G. ♀.	Fl. græc. t. 636.
13 tristis. DC.	(br.) various-leaved. S. Europe. 1768. —	G. ♀.		Barr. ic. t. 803.
14 coronopifòlia. DC.	(pu.) Buckshorn-ld. 1825. —	H. ♀.		Flor. græc. t. 637.
15 lívida. DC.	(br.) livid. Egypt. 1828. —	H. ♂.		Deles. ic. sel. 2. t. 12.
16 longipétala. DC.	(br.) long-petaled. Bagdad. 1819. —	H. ♂.		Vent. cels. t. 93.
17 tricuspidâta. DC.	(pu.) three-forked. Barbary. 1739. 7. 10.	H. ♂.		Swt. br. fl. gar. 46.
18 parviflôra. DC.	(pu.) small-flowered. Morocco. 1799. —	H. ♂.		
CHEIRANTHUS. DC	WALL-FLOWER. Tetradymania Siliquosa. L.			
1 Chéiri. DC.	(ye.) garden. S. Europe. 1573. 4. 7.	H. ♀.		Schkuhr hand. 2. t. 184.
α vulgâris.	(ye.) common. —	—	H. ♀.	Besl. eyst. 2. t. 5. f. 3.
β flôre plêno.	(ye.) double-yellow. —	—	H. ♀.	— 2. t. 4. f. 2. 3.
γ grandiñorus.	(ye.) large-flowered. —	—	H. ♀.	— 2. t. 5. f. 2.
δ máximus.	(ye.) largest. —	—	H. ♀.	— 2. t. 5. f. 1.
ε serrátus.	(ye.) saw-leaved. —	—	H. ♀.	
ζ pátulus.	(ye.) double-spreading. —	—	H. ♀.	
ι ferrugíneus.	(p.y.) rusty-flowered. —	—	H. ♀.	
κ vârius.	(p.y.) various-coloured. —	—	H. ♀.	
λ sangúineus.	(p.y.) bloody. —	—	H. ♀.	
μ hæmânthus.	(p.y.) double-bloody. —	—	H. ♀.	
ν flâvescens.	(ye.) pale-yellow. —	—	H. ♀.	
ξ thyrsóideus.	(ye.) bunch-flowered. —	—	H. ♀.	

CRUCIFERÆ.

2 fruticulòsus. E.B. (<i>ye.</i>) common.	England.	4. 7. H. $\frac{1}{2}$.	Eng. bot. 1934.
3 alpinus. DC. (<i>ye.</i>) Alpine.	Norway. 1823.	— H. $\frac{1}{2}$.	Wahl. fl. lap. t. 12. f. 1.
4 ochroleucus. DC. (<i>st.</i>) pale yellow. <i>decumbens</i> . W. en.	Switzerland. 1820.	6. 8. H. $\frac{1}{2}$.	
5 tenuifolius. DC. (<i>ye.</i>) fine-leaved.	Madeira. 1777.	5. 6. F. $\frac{1}{2}$.	
6 mutabilis. DC. (<i>li.</i>) changeable.	—	3. 5. F. $\frac{1}{2}$.	Bot. mag. 195.
7 longifolius. V. (<i>li.</i>) long-leaved.	—	1815. 9. 12. F. $\frac{1}{2}$.	Vent. malm. t. 83.
8 scoparius. DC. (<i>va.</i>) rock.	Teneriffe. —	3. 8. F. $\frac{1}{2}$.	
<i>a</i> <i>purpurascens</i> . DC. (<i>p.</i>) <i>purplish.</i>	—	— F. $\frac{1}{2}$.	
β <i>æruginosus</i> . DC. (<i>br.</i>) <i>rusty.</i>	—	— F. $\frac{1}{2}$.	
γ <i>chamæleo</i> . DC. (<i>p.y.</i>) <i>Chameleon.</i>	—	— F. $\frac{1}{2}$.	
<i>Chéri</i> , var. <i>Chamæleon</i> . B.R.	—	— F. $\frac{1}{2}$.	Bot. reg. 219.
9 semperflorens. DC. (<i>w.</i>) ever-blowing.	Barbary. —	1. 12. F. $\frac{1}{2}$.	
10 frutescens. P.S. (<i>wh.</i>) frutescent.	—	3. 7. F. $\frac{1}{2}$.	
11 linifolius. DC. (<i>pu.</i>) flax-leaved.	Spain. —	3. 8. F. $\frac{1}{2}$.	
12 arborescens. Sie. (<i>pu.</i>) arborescent.	Levant. 1827.	4. 8. G. $\frac{1}{2}$.	
NASTURTIUM. DC. NASTURTIUM. officinale. DC. (<i>wh.</i>) Water Cress.	Tetradynamia Siliquosa. L.		
<i>Sisy'mbrium Nastúrtium</i> . W.	Britain.	5. 7. H.w. $\frac{1}{2}$.	Eng. bot. 855.
β <i>præcocius</i> . DC. (<i>wh.</i>) early.	—	— H.w. $\frac{1}{2}$.	
2 sylvestre. DC. (<i>ye.</i>) creeping.	— 6. 9. H.w. $\frac{1}{2}$.	Pet. h. brit. t. 47. f. 3.
3 anceps. DC. (<i>ye.</i>) flat-edged.	N. Europe. 1826.	— H.w. $\frac{1}{2}$.	Eng. bot. 2324.
4 palustre. DC. (<i>ye.</i>) marsh.	Britain.	— H.w. \odot .	Flor. dan. t. 984.
<i>Sisy'mbrium terrestre</i> . E.B.	Siberia. 1780.	5. 6. H. $\frac{1}{2}$.	Eng. bot. 1747.
5 sagittatum. DC. (<i>ye.</i>) arrow-leaved.	Carinthia. 1823.	— H. $\frac{1}{2}$.	
6 lippizense. DC. (<i>ye.</i>) Carinthian.	Pyrenees. 1775.	— H. $\frac{1}{2}$.	
7 pyrenaicum. DC. (<i>ye.</i>) Pyrenean.	Britain.	6. 8. H.w. $\frac{1}{2}$.	
8 amphibium. DC. (<i>ye.</i>) amphibious.	—	— H.w. $\frac{1}{2}$.	
<i>a</i> <i>indivisum</i> . DC. (<i>ye.</i>) <i>brd. wat.-radish.</i>	—	— H.w. $\frac{1}{2}$.	
β <i>variifolium</i> . DC. (<i>ye.</i>) <i>jag. wat.-radish.</i>	—	— H.w. $\frac{1}{2}$.	
9 benghalense. DC. (<i>wh.</i>) Bengal.	Bengal.	6. 10. H. \odot .	Jacq. ic. 1. t. 122.
10 clandestinum. DC. (<i>wh.</i>) small-flowered.	Brazil. 1821.	— H. \odot .	— 3. t. 505.
11 indicum. DC. (<i>gr.</i>) Indian.	E. Indies. —	— H. \odot .	All. fl. ped. t. 18. f. 1.
12 atrovirens. H. (<i>ye.</i>) dark green.	China. 1826.	— H. \odot .	Eng. bot. 1840.
LEPTOCA'RPEA. DC. LEPTOCA'RPEA. TETRADYNA. Siliquosa. L.	Tetradynamia Siliquosa. L.		Pet. h. brit. t. 49. f. 8.
<i>Loeselii</i> . DC. (<i>ye.</i>) Loeselius's.	Germany. 1683.	5. 8. H. \odot .	— t. 49. f. 10.
<i>Sisy'mbrium Loeselii</i> . W. <i>Turritis Loeselii</i> . H. K.			
NOTOCERAS. DC. NOTOCERAS. TETRADYNA. Siliquosa. L.			Jacq. aust. t. 324.
1 canariense. DC. (<i>ye.</i>) Canary.	Canaries. 1779.	8. 9. H. \odot .	Jacq. fil. eclog. t. 111.
<i>Ery'simum bicorne</i> . W.			
2 hispanicum. DC. (<i>ye.</i>) Spanish.	Spain. 1822.	3. 10. H. \odot .	Deles. ic. sel. 2. t. 17.
3 quadricorne. DC. (<i>ye.</i>) four-horned.	Siberia. 1824.	5. 10. H. \odot .	— 2. t. 16.
BARBAR'EA. DC. WINTER-CRESS. TETRADYNA. Siliquosa. L.			
1 vulgàris. DC. (<i>ye.</i>) common.	Britain.	5. 8. H. $\frac{1}{2}$.	Eng. bot. 443.
<i>Ery'simum Barbaræ</i> . W.			— 1129.
2 præcox. DC. (<i>ye.</i>) early.	England.	4. 10. H. $\frac{1}{2}$.	
3 taurica. DC. (<i>ye.</i>) Taurian.	Tauria. 1825.	7. 9. H. $\frac{1}{2}$.	
4 ibérica. DC. (<i>ye.</i>) Iberian.	Iberia. 1816.	5. 8. H. $\frac{1}{2}$.	
<i>Cheiranthus ibericus</i> . W. en.			
5 plantaginea. DC. <i>Plaintain-leav'd.</i> Levant.	1823.	— H. $\frac{1}{2}$.	Deles. ic. sel. 2. t. 19.
STEVENIA. DC. STEVENIA. TETRADYNA. Siliquosa. L.			
1 alyssoides. DC. (<i>wh.</i>) Alyssum-like.	Siberia.. 1823.	6. 8. H. \odot .	Deles. ic. sel. 2. t. 20.
2 cheiranthoides. DC. (<i>pu.</i>) Wall-flow.-like.	—	— H. $\frac{1}{2}$.	— 2. t. 21.
BRA'YA. DC. BRA'YA. TETRADYNA. Siliquosa. DC.			
1 alpina. DC. (<i>pu.</i>) Alpine.	Carinthia. 1821.	4. 5. H. $\frac{1}{2}$.	Deles. ic. sel. 2. t. 22.
2 glabella. DC. (<i>pu.</i>) smooth.	Arctic R. 1827.	— H. $\frac{1}{2}$.	
TURRITIS. DC. TOWER-MUSTARD. TETRADYNA. Siliquosa. L.			
1 glabra. DC. (<i>wh.</i>) long-podded.	England.	5. 6. H. \odot .	Eng. bot. 777.
β ramosa. DC. (<i>wh.</i>) branching.	— H. \odot .	Clus. hist. 2. p. 126. f. 2.
2 salsuginosa. DC. (<i>wh.</i>) shorter-podded.	Siberia. 1822.	— H. \odot .	Palit. 2. app. n. 114. t. V.
A'RABIS. DC. WALL-CRESS. TETRADYNA. Siliquosa. L.			
1 verna. DC. (<i>pu.</i>) vernal.	S. Europe. 1710.	4. 5. H. \odot .	Flor. græc. t. 641.
<i>Hesperis verna</i> . W.			
2 rosea. DC. (<i>ro.</i>) rose-coloured.	Calabria. 1827.	— H. $\frac{1}{2}$.	Deles. ic. sel. 2. t. 23.
3 alpina. DC. (<i>wh.</i>) Alpine.	Pyrenees. 1596.	3. 5. H. $\frac{1}{2}$.	Bot. mag. 226.
β Clusiànæ. DC. (<i>wh.</i>) <i>Clusiæ's.</i>	—	— H. $\frac{1}{2}$.	Clus. hist. 2. p. 125. f. 2.
γ nana. DC. (<i>wh.</i>) dwarf.	Hungary.	— H. $\frac{1}{2}$.	
4 álbida. DC. (<i>wh.</i>) early-flowered.	Tauria. 1798.	1. 5. H. $\frac{1}{2}$.	Jacq. f. ecl. 1. t. 71.
<i>caucásica</i> . B.M. 2046.			

5 undulatæ. L. en. (wh.) wave-leaved.	S. Europe.	1823.	5. 8.	H. ♂.	
6 thyrsoidæ. DC. (wh.) bunch-flowered.	Levant.	1826.	—	H. ♀.	Flor. græc. t. 642.
7 longifolia. DC. (wh.) long-leaved.	Persia.	1820.	—	H. ♀.	Deles. ic. sel. 2. t. 25.
8 mollis. DC. (wh.) soft.	Caucasus.	1825.	5. 7.	H. ♂.	
9 toxophylla. DC. (wh.) bow-leaved.	Volga.	1828.	—	H. ♂.	
10 auriculatæ. DC. (wh.) ear-leaved. aspera. All. ped. t. 2. f. 2.	S. Europe.	1805.	5. 6.	H. ♂.	Vill. dauph. 3. t. 37.
11 saxatilis. DC. (wh.) rock.	—	—	—	H. ♂.	
12 crispata. DC. (wh.) curled-leaved.	Carniola.	1816.	3. 4.	H. ♀.	
13 sagittata. DC. (wh.) arrow-leaved.	Europe.	5. 7.	H. ♂.	
14 hirsuta. DC. (wh.) hairy.	Britain.	—	H. ♂.	Eng. bot. 587.
15 stenopetalæ. W. en. (w.) narrow-petal'd.	1822.	6. 8.	H. ♂.	
16 curtisiliqua. DC. (wh.) short-podded.	Scania.	1825.	5. 7.	H. ♂.	
17 Allioniæ. DC. (wh.) Allion's.	Italy.	5. 7.	H. ♀.	
18 muralis. DC. (wh.) dwarf.	Switzerland.	1824.	5. 6.	H. ♀.	
19 stricta. DC. (wh.) Bristol.	England.	—	H. ♀.	Eng. bot. 614.
20 ciliata. DC. (wh.) ciliated.	Ireland.	6. 7.	H. ♂.	————— 1746.
Turritis alpina. E.B.					
21 incana. DC. (wh.) hoary.	Switzerland.	1816.	5. 6.	H. ♂.	
22 Thaliâna. DC. (wh.) common.	Britain.	4. 6.	H. ♂.	Eng. bot. 901.
23 serpyllifolia. DC. (wh.) Thyme-leaved.	Pyrenees.	1824.	6. 7.	H. ♂.	
24 procurrens. DC. (wh.) procurent.	Hungary.	—	5. 6.	H. ♀.	W. et. Kit. hun. 2.t.144.
25 præcox. DC. (wh.) early-flowering.	—	1822.	4. 6.	H. ♀.	
26 Schivereckiana. DC. (w.) Schivereck's.	Austria.	1827.	—	H. ♀.	
27 petraea. DC. (wh.) short-podded.	Europe.	5. 7.	H. ♀.	
α Crantziana. DC. (w.) Crantz's.	Germany.	—	H. ♀.	Crantz. aust. t. 3. f. 2.
β hispida. DC. (wh.) hispid.	Wales.	—	H. ♀.	Flor. dan. t. 386.
γ hastulata. DC. (wh.) hastulate.	Britain.	—	H. ♀.	Eng. bot. 409.
28 ambigua. DC. (wh.) ambiguous.	Siberia.	1824.	5. 7.	H. ♀.	
29 lyrata. DC. (wh.) lyre-leaved.	N. America.	1823.	5. 6.	H. ♀.	
30 arenosa. DC. (ro.) purple.	Europe.	1798.	6. 7.	H. ♀.	
31 Halleria. DC. (wh.) Haller's.	—	1820.	5.	H. ♂.	Schranck. m. 3. t. 256.
32 stolonifera. DC. (wh.) stoloniferous.	Carniola.	1824.	—	H. ♀.	W. et Kit. hun. 2.t.120.
33 ovirensis. DC. (ro.) reddish.	Carinthia.	—	5. 6.	H. ♀.	Scop. carn. ed. 2. t. 39.
34 cebennensis. DC. (re.) pale-violet.	S. France.	1824.	5. 7.	H. ♂.	Jacq. ic. 1. t. 125.
35 lasiloba. DC. (wh.) woolly-podded.	Mexico.	—	—	H. ♂.	Deles. ic. sel. 2. t. 26.
36 Turritæ. DC. (wh.) tower.	England.	4. 5.	H. ♂.	
37 pendula. DC. (wh.) pendulous.	Siberia.	1752.	5. 6.	H. ♂.	Eng. bot. 178.
38 Patriniana. DC. (wh.) Patrin's.	—	1826.	—	H. ♂.	Jacq. vind. 3. t. 34.
39 oxyota. DC. (wh.) sharp-eared.	—	—	—	H. ♂.	Deles. ic. sel. 2. t. 27.
40 luctuosa. DC. (wh.) shining-leaved.	Hungary.	1790.	6. 7.	H. ♀.	
41 canadensis. DC. (wh.) sickle-podded.	N. America.	1768.	5. 7.	H. ♀.	Deles ic. sel. 2. t. 28.
42 pumila. DC. (wh.) nodding. nuttans. B.M. 2219. cilariæ. W. en.	Switzerland.	1658.	4. 7.	H. ♀.	Jacq. fl. aust. t. 281.
43 bellidifolia. DC. (wh.) Daisy-leaved.	Pyrenees.	1773.	5. 6.	H. ♀.	————— 3. t. 280.
44 coerulea. DC. (bl.) blue.	—	1793.	6. 7.	H. ♀.	Sturm. deut. fl. ic.
45 collina. DC. (wh.) hill.	Naples.	1823.	—	H. ♀.	
46 dasycarpa. DC. (wh.) thick-podded.	Poland.	1826.	5. 7.	H. ♀.	
MACROP'ODIUM. DC. MACROP'ODIUM. Tetradymania Siliquosa. L.					
nivale. DC. (wh.) Siberian.	Siberia.	1796.	6. 9.	H. ♀.	Pall. it. 2. ap.n. 113.t.u.
Cardamine nivalis. Pall.					
CARDAMINE. DC. LADY'S SMOCK. Tetradymania Siliquosa. L.					
1 rhomboidæ. DC. (wh.) rhomboid.	N. America.	1825.	5. 7.	H. w. ♀.	Pluk. amal. t. 435. t. 6.
2 rotundifolia. DC. (wh.) round-leaved.	—	1823.	5. 6.	H. w. ♀.	
3 asarifolia. DC. (wh.) kidney-leaved.	Italy.	1710.	6. 7.	H. w. ♀.	Bot. mag. 1735.
4 bellidifolia. DC. (wh.) Daisy-leaved.	Scotland.	4. 6.	H. ♀.	Œd. fl. Dan. t. 20.
α petiolaris. DC. (wh.) long-petioled.	Lapland.	—	H. ♀.	Eng. bot. 2355.
β alpina. DC. (wh.) Alpine.	Scotland.	—	H. ♀.	Sturm. fl. germ. ic.
5 resedifolia. DC. (wh.) Rocket-leaved.	Germany.	1658.	7.	H. ♂.	Pluk. alm. t. 101. f. 5.
6 microphylla. DC. (wh.) small-leaved.	Siberia.	1828.	5. 7.	H. ♀.	
7 africana. DC. (wh.) African.	C. B. S.	1691.	5. 6.	G. ♀.	
8 trifolia. DC. (wh.) three-leaved.	Europe.	1629.	3. 6.	H. w. ♀.	Bot. mag. 452.
9 apnara. DC. (wh.) bitter.	Britain.	4. 5.	H. w. ♀.	Eng. bot. 1000.
10 prorepens. DC. (wh.) creeping.	Siberia.	1825.	4. 6.	H. w. ♀.	Eng. bot. 776.
11 pratensis. DC. (li.) Cuckoo-flower.	Britain.	—	H. w. ♀.	Clus. hist. 2. p. 129. f. 1.
β flôre pleno.	(li.) double-flowered.	—	—	H. w. ♀.	
12 dentata. DC. (wh.) toothed.	Poland.	1823.	—	H. w. ♀.	
13 hirsuta. DC. (wh.) hairy.	Britain.	1. 12.	H. ♂.	Eng. bot. 492.
14 sylvatica. DC. (wh.) wood.	Europe.	5. 6.	H. ♂.	Flor. dan. t. 735.

CRUCIFERÆ.

15 umbrosa. DC. (wh.) smooth.	Europe.	4. 6. H. ⊖.	
16 parviflora. DC. (wh.) small-flowered.	1822. 4. 5. H. ⊖.	Gmel. fl. sib. 3. t. 64.	
17 impatiens. DC. (wh.) impatient.	Britain.	Eng. bot. 80.	
18 latifolia. DC. (pu.) broad-leaved.	Spain. 1710. 6. 8. H. w. Ȑ.	Herm. par. p. 203. t. 69.	
19 macrophylla. DC. (li.) large-leaved.	Siberia. 1824. — H. Ȑ.		
20 chelidonia. DC. (pu.) Celandine-l'd.	Italy. 1739. 3. 5. H. ♂.	W. et Kit. hung. 2. t. 140.	
21 thalictroides. DC. (wh.) Thalictrum-l'd.	Switzerland. 1824. 6. 7. H. ♂.	All. ped. n. 951. t. 57. f. 1.	
22 glauca. DC. (wh.) glaucous.	Calabria. 1824. 5. 7. H. Ȑ.	Deles. ic. sel. 2. t. 31.	
23 occulta. DC. (gr.) hidden-flower'd.	China. — 5. 9. H. ⊖.		

PTERONEURUM. DC. PTERONEURUM. Tetradymania Siliquosa. L.

1 carnosum. DC. (wh.) fleshy.	Dalmatia. 1824. 6. 8. H. Ȑ.	W. et Kit. hung. 2. t. 129.
2 græcum. DC. (wh.) Grecian.	S. Europe. 1710. — H. ⊖.	Flor. græc. t. 631.

Cardamine græca. w.

DENTARIA. DC. TOOTH-WORT. Tetradymania Siliquosa. L.			
1 polyphylla. DC. (st.) many-leaved.	Hungary. 1817. 5. 6. H. Ȑ.	W. et Kit. hung. 2. t. 160.	
2 enneaphylla. DC. (wh.) nine-leaved.	Austria. 1656. — H. Ȑ.	Jacq. fl. aust. t. 316.	
3 glandulosa. DC. (pu.) glandular.	Hungary. 1815. — H. Ȑ.	W. et Kit. hung. 3. t. 272.	
4 lacinata. DC. (wh.) laciniate.	N. America. 1823. 4. 6. H. Ȑ.		
5 diphylla. DC. (wh.) two-leaved.	— 1810. 5. 6. H. Ȑ.	Bot. mag. 1465.	
6 trifolia. DC. (wh.) three-leaved.	Hungary. 1824. — H. Ȑ.	W. et K. hung. 2. t. 139.	
7 digitata. DC. (w.p.) fingered.	Switzerland. 1656. — H. Ȑ.	Bot. mag. 2202.	

pentaphylla. B.M. Cardamine pentaphylla. H. K. ed. 2.

8 pinnata. DC. (wh.) seven-leaved.	— 1683. — H. Ȑ.	Boiss. fl. eur. t. 449.
9 quinquefolia. DC. (wh.) five-leaved.	Caucasus. 1823. 4. 5. H. Ȑ.	Deles. ic. sel. 2. t. 33.
10 bulbifera. DC. (bh.) bulbiferous.	England. — H. Ȑ.	Eng. bot. 309.

Cardamine bulbifera. H. K. ed. 2.

11 tenuifolia. DC. (pu.) slender-leaved.	Siberia. 1825. 5. 6. H. Ȑ.	Gmel. sib. 3. n. 41. t. 65.
trifida. Lam. ill. t. 562. f. 2.		

PARRYA. B. PARRYA. Tetradymania Siliquosa.

1 arctica. B. (pu.) arctic.	Melvill Isl. 1827. 5. 6. H. Ȑ.	
2 arabisiflora. (pu.) Arabis-flower'd. Siberia.	1798. — H. Ȑ.	Linn. am. ac. 2. t. L. f. 20.

Arabis grandiflora. L. *Neurolema arabidisflorum*. DC.

PLATYPETALUM. B. PLATYPETALUM. Tetradymania Siliquosa.		
purpurascens. B. (pu.) purplish.	Melvill Isl. 1827. 5. 6. H. Ȑ.	

Tribus II. ALYSSINEÆ seu Pleurorhizeæ Latiseptæ. DC.

LUNARIA. DC. HONESTY. Tetradymania Siliculosa. L.		
1 rediviva. DC. (li.) perennial.	Germany. 1596. 4. 6. H. Ȑ.	Hill veg. syst. t. 38. f. 1.
2 biennis. DC. (w.p.) biennial.	— 1570. — H. ♂.	t. 38. f. 2.

ánnua. w.		
a violacea.	(pu.) violet-flowered.	— — — H. ♂.
β albiflora.	(wh.) white-flowered.	— — — H. ♂.

RICOTIA. DC. RICOTIA. Tetradymania Siliculosa. L.		
Lunaria. DC. Syrian.	Syria. 1757. 6. 7. H. ⊖.	Lam. ill. t. 561.

ægyptiacu. Bot. Reg. 49.		
FARSETIA. DC. FARSETIA. Tetradymania Siliculosa. L.		

1 aegyptiaca. DC. (br.) Stock-like.	Levant. 1788. 6. 7. G. Ȑ.	Desf. atl. 2. t. 160.
cheiranthoides. H. K.	Cheiranthus Farsetia. L.	

FIBIGIA. Med. FIBIGIA. Tetradymania Siliculosa.		
1 suffruticosa. (pu.) suffrutescent.	Persia. 1820. 4. 5. G. Ȑ.	Vent. cels. t. 19.

Lunaria suffruticosa.	Vent. Farsetia suffruticosa. DC.	
2 lunarioides. (ye.) oriental.	Greece. 1731. 6. 7. H. Ȑ.	Tourn. ic ed. ger. 1 t. 30.

3 clypeata. Med. (ye.) buckler-podded.	S. Europe. 1596. — H. ♂.	Schkuhr. han. 2. t. 181.
Alyssum clypeatum. L.	Farsetia clypeata. DC.	

BERTEROA. DC. BERTEROA. Tetradymania Siliculosa. L.		
1 incana. DC. (wh.) hoary.	Europe. 1640. 7. 9. H. ♂.	Horn. fl. dan. 1461.

Alyssum incanum. w.		
2 mutabilis. DC. (bh.) changeable.	Levant. 1802. 7. 8. H. Ȑ.	Vent. cels. t. 85.

3 obliqua. DC. (wh.) oblique-podded.	Naples. 1824. — H. Ȑ.	Flor. græc. t. 623.
AUBRIETIA. DC. AUBRIETIA. Tetradymania Siliculosa. L.		

1 deltoidea. DC. (pu.) spreading.	Levant. 1710. 3. 5. H. Ȑ.	Bot. mag. 126.
Farsetia deltoidea. H. K.	Alyssum deltoideum. L.	

2 purpurea. DC. (pu.) tufted.	— 1821. — H. Ȑ.	Swt. br. fl. gar. 207.
VESICARIA. VESICARIA. Tetradymania Siliculosa. L.		

1 utriculata. DC. (ye.) smooth.	Levant. 1739. 4. 6. H. Ȑ.	Bot. mag. 130.
Alyssum utriculatum. B.M.		

2 reticulata. DC. (ye.) veined-podded. 6. H. Ȑ.	Tour. voy. 2. p. 252. ic.
--	---------------	---------------------------

CRUCIFERÆ.

25

3 Ludoviciana. DC. (ye.) Missouri.	Missouri.	1825.	5. 6. H. ȝ.	
4 sinuata. DC. sinuate-leaved.	Spain.	1596.	4. 6. H. ♂.	Schkuhr. han. 2. t. 181.
5 crética. DC. Cretan.	Crete.	1739.	5. 8. H. ȝ.	Alp. ex. p. 117 et 118. ic.
Alyssum creticum. w.				
6 ártica. DC. (ye.) arctic.	Greenland.	1826.	5. 6. H. ȝ.	Horn. fl. dan. t. 1520.
7 arenosa. DC. (ye.) sand.	Arctic Reg.	—	6. 8. H. ȝ.	
SCHIVERECKIA. DC. SCHIVERECKIA.	Tetradynamia Siliculosa. L.			
podólica. DC. (wh.) canescent.	Podolia.	1821.	5. 7. H. ȝ.	Swt. br. fl. gar. t. 77.
KO'NIGA. B. Ko'NIGA.	Tetradynamia Siliculosa.			
1 marítima. DC. (wh.) sweet-scented.	S. Europe.	5. 10. H. ȝ.	Eng. bot. 1729.
Alyssum marítimum. E.B. halimifolium. B.M. 101.				
2 canariensis. (wh.) Canary.	Canaries.	1812.	— G. ℡.	
Alyssum marítimum. β canariense. DC.				
β variegata. (wh.) striped-leaved.	—	— G. ℡.	
ADYS'ETON. Scop. ADYS'ETON.	Tetradynamia Siliculosa.			
1 saxátilis. (ye.) rock.	Russia.	1710.	4. 5. H. ℡.	Bot. mag. 159.
Alyssum saxátilis. D.				
2 gemonense. (ye.) canescent.	Germany.	—	— H. ℡.	Jacq. ic. 3. t. 503.
3 orientále.	Crete.	—	— H. ℡.	Flor. græc. t. 625.
4 spathulatum.	Siberia.	1821.	4. 6. H. ℡.	Deles. ic. sel. 2. t. 37.
5 argénteum.	Switzerland.	—	5. 6. H. ℡.	All. ped. t. 54. f. 3.
6 Bertolónii.	Italy.	1825.	— H. ℡.	
7 murále.	Europe.	1801.	6. 8. H. ℡.	W. et Kit. hung. I. t. 6.
8 obtusifolium.	Siberia.	1812.	— H. ℡.	Deles. ic. sel. 2. t. 38.
9 serpylifolium.	S. Europe.	1823.	5. 7. H. ℡.	
10 rēpens.	Transylvania.	1827.	— H. ȝ.	
11 vernále.	(ye.) vernal.	1823.	4. 5. H. ȝ.	Schrank hort. mon. t. 96.
12 tortuosum.	Hungary.	1804.	6. 7. H. ȝ.	W. et Kit. hung. I. t. 91.
13 alpéstre.	Italy.	1777.	6. 10. H. ℡.	Flor. græc. t. 624.
14 nebrodēnse.	Sicily.	1827.	5. 6. H. ℡.	
15 Marschallianum.	Caucasus.	—	— H. ℡.	
16 montánum. sc.	Germany.	1713.	5. 8. H. ȝ.	Bot. mag. 419.
Alyssum montánum. B.M.				
17 cuneifolium.	(ye.) wedge-leaved.	Naples.	1825.	5. 8. H. ȝ.
18 diffusum.	(ye.) spreading.	—	—	H. ȝ.
19 Wulfeniánum.	(ye.) Wulfen's.	Carinthia.	1821.	— H. ȝ.
20 Fischeriánum.	(ye.) Fischer's.	Siberia.	1828.	— H. ȝ.
Alyssum Fischeriánum. DC.				
ALY'SSUM. B. MADWORT.	Tetradynamia Siliculosa. L.			
1 umbellatum. DC. (ye.) umbelled.	Tauria.	1828.	5. 8. H. ⊙.	
2 rostratum. DC. (ye.) beaked.	—	1824.	6. 9. H. ⊙.	Stev.m.ac.P.3.t.15.f.1.
3 micropétalum. DC. (ye.) small-petaled.	Iberia.	—	5. 8. H. ⊙.	Deles. ic. sel. 2. t. 39.
4 hirsutum. DC. (ye.) hairy.	—	—	— H. ⊙.	— 2. t. 40.
5 campestre. DC. (ye.) field.	Europe.	1768.	6. 8. H. ⊙.	Flor. græc. t. 626.
6 calycinum. DC. (ye.) persistent-cal'd.	—	1740.	— H. ⊙.	Jacq. aust. t. 338.
7 míminum. DC. (ye.) small.	—	1791.	— H. ⊙.	Tratt. thes. bot. t. 35.
ANODO'NTEA. DC. ANODO'NTEA.	Tetradynamia Siliculosa.			
1 edéntula.	(ye.) toothless.	Hungary.	1821.	5. 6. H. ♂.
Alyssum edéntulum. DC. Adyséton edéntulum. L. enum.				W. et K. hung. I. t. 92.
2 dasycarpa.	(ye.) thick-podded.	Siberia.	1827.	6. 10. H. ⊙.
3 rupestris.	(wh.) silvery rock.	Italy.	1824.	5. 10. H. ȝ.
4 halimifolia.	(wh.) Halimus-leaved.	—	—	H. ℡.
Alyssum halimifolium. DC.				
5 spinosa.	(wh.) thorny.	S. Europe.	1683.	6. 8. H. ℡.
6 macrocarpa.	(wh.) large-podded.	—	1825.	— H. ℡.
7 canescens.	(wh.) hoary.	Siberia.	1828.	4. 9. H. ℡.
MENTOCUS. DC. MENTOCUS.	Tetradynamia Siliculosa. L.			
linifolius. DC. (ye.) Flax-leaved.	Levant.	1822.	5. 6. H. ⊙.	Deles. ic. sel. 2. t. 42.
Alyssum linifolium. w.				
CLYPE'OLA. DC. TREACLE-MUSTARD.	Tetradynamia Siliculosa. L.			
1 Jonthláspi. DC. (ye.) common.	S. Europe.	1710.	5. 7. H. ⊙.	Desv.j. bot. 3. t. 25.f.7.
2 eriophora. DC. (wh.) woolly-podded.	Spain.	1820.	— H. ⊙.	— 3. t. 25.f.10.
PELTARIA. DC. PELTARIA.	Tetradynamia Siliculosa. L.			
alliacea. DC. (wh.) garlic-scented.	Austria.	1601.	5. 7. H. ȝ.	Jacq. aust. 2. t. 123.
PETROCAL'LIS. DC. PETROCAL'LIS.	Tetradynamia Siliculosa. L.			
pyrenaica. DC. (ro.) Pyrenean.	Pyrenees.	1759.	5. 6. H. ȝ.	Lodd. bot. cab. 635.
Draba pyrenaica. B.M. 713.				
DRABA. DC. WHITLOW-GRASS.	Tetradynamia Siliculosa. L.			
1 aizóides. DC. (ye.) sen-green.	Wales.	2. 4. H. ȝ.	Eng. bot. 1271.

CRUCIFERÆ.

2	brachystemon. DC. (<i>ye.</i>) short-stamened.	1731.	2. 4. H. ¼.	Bot. mag. 170.
	<i>aizoides</i> . B.M. <i>ciliaris</i> . H.K.				
3	aizoon. DC. (<i>ye.</i>) hairy-podded.	Hungary.	1821.	— H. ¼.	
4	cuspidata. DC. (<i>ye.</i>) sharp-pointed.	Tauria.	—	4. 5. H. ¼.	
5	bryoides. DC. (<i>ye.</i>) Bryum-like.	Caucasus.	—	3. 4. H. ¼.	
6	bruniaefolia. DC. (<i>ye.</i>) Brunia-leaved.	—	1826.	5. 6. H. ¼.	
7	ericæfolia. DC. (<i>ye.</i>) Heath-leaved.	—	1823.	— H. ¼.	
8	pilosa. DC. (<i>ye.</i>) pilose.	Siberia.	1825.	— H. ¼.	
9	alpina. DC. (<i>ye.</i>) Alpine.	Norway.	1823.	4. 5. H. ¼.	
10	glacialis. DC. (<i>ye.</i>) starry-haired.	Siberia.	1825.	5. 6. H. ¼.	
11	repens. DC. (<i>ye.</i>) creeping.	Caucasus.	1823.	— H. ¼.	
12	Gmelini. DC. (<i>ye.</i>) Gmelin's.	Siberia.	—	— H. ¼.	Gmel. sib. 3. t. 56. f. 2.
13	tridentata. DC. (<i>ye.</i>) three-toothed.	Caucasus.	1828.	— H. ¼.	
14	muriella. DC. (<i>wh.</i>) tufted.	Norway.	—	— H. ¼.	
15	incompta. DC. (<i>ye.</i>) straggling.	Caucasus.	—	6. H. ¼.	Wahl. fl. lap. t. 11. f. 2.
16	oblongata. DC. (<i>wh.</i>) oblong-leaved.	Baffin's-bay.	1820.	5. 6. H. ¼.	Deles. ic. sel. 2. t. 44.
17	corymbosa. DC. (<i>wh.</i>) corymbose.	—	—	— H. ¼.	
18	hirta. DC. (<i>wh.</i>) hairy.	Lapland.	5. 7. H. ¼.	
19	rupestris. DC. (<i>wh.</i>) rock.	Scotland.	— H. ¼.	Wahl. f. lap. t. 11. f. 3.
	<i>hirta</i> . E.B. non Linn.				Eng. bot. 1338.
20	nivalis. DC. (<i>wh.</i>) snowy.	Norway.	1824.	5. 6. H. ¼.	
	<i>stellata</i> . F.D. nec aliorum.				Flor. dan. t. 142.
21	lapponica. DC. (<i>wh.</i>) Lapland.	Lapland.	—	— H. ¼.	
22	helvetica. DC. (<i>wh.</i>) Swiss.	Switzerland.	1824.	— H. ¼.	
23	fladnizensis. DC. (<i>wh.</i>) fringe-leaved.	—	—	6. 8. H. ¼.	Jacq. misc. 1.t. 17. f. 1;
24	tomentosa. DC. (<i>wh.</i>) woolly-leaved.	—	—	5. 7. H. ¼.	Wahl. helv. p. 123.t. 3;
25	stellata. DC. (<i>wh.</i>) starry.	—	1820.	— H. ¼.	Jacq. obs. t. 4. f. 3.
	<i>hirta</i> . Jacq. aust. t. 432.				
26	contorta. DC. (<i>wh.</i>) twisted-podded.	Britain.	5. 6. H. ♂.	Eng. bot. t. 388.
	<i>incana</i> . E.B.				
27	confusa. DC. (<i>wh.</i>) confused.	N. Europe.	— H. ♂.	Flor. dan. t. 130.
	<i>incana</i> . F.D.				
28	arabisans. DC. (<i>wh.</i>) Arabis-like.	N. America.	1827.	— H. ♂.	
29	daurica. DC. (<i>wh.</i>) Daurian.	Dahuria.	1824.	— H. ♂.	
30	auraea. DC. (<i>ye.</i>) golden.	Greenland.	—	— H. ♂.	
31	lutea. DC. (<i>ye.</i>) yellow.	Caucasus.	1825.	5. 8. H. ♂.	
32	gracilis. GR. (<i>ye.</i>) slender.	N. America.	1827.	— H. ♂.	
33	nemoralis. DC. (<i>ye.</i>) wood.	Europe, &c.	1759.	— H. ♂.	
34	murialis. DC. (<i>wh.</i>) wall.	England.	4. 5. H. ♂.	
	<i>ERO'PHILA</i> . DC. <i>ERO'PHILA</i> .	Tetradynamia Siliculosa. L.			
1	americanæ. DC. (<i>wh.</i>) American.	N. America.	1823.	3. 4. H. ♂.	
2	vulgäris. DC. (<i>wh.</i>) common.	Britain.	— H. ♂.	
	<i>Draba verna</i> . Eng. bot. 586.				Flor. dan. t. 983.
3	præcox. DC. (<i>wh.</i>) round-podded.	Caucasus.	1824.	2. 4. H. ♂.	
	<i>KERNERA</i> . Med. <i>KE'RNERA</i> . Tetradynamia Siliculosa.				
1	saxatilis. (<i>wh.</i>) rock.	Europe.	1775.	5. 7. H. ¼.	
	<i>myagroides</i> . Med. <i>Cochlearia saxatilis</i> . DC. <i>Myagrum saxatile</i> . L.				
	β incisa. (<i>wh.</i>) cut-leaved.	—	—	— H. ¼.	
2	auriculata. (<i>wh.</i>) eared.	Pyrenees.	1824.	— H. ¼.	
	<i>Cochlearia auriculata</i> . DC.				
	<i>ARMOR'ACIA</i> . Plin. HORSE-RADISH. Tetradynamia Siliculosa.				
1	rusticana. Bau. (<i>wh.</i>) common.	England.	5. 6. H. ¼.	
	<i>Cochlearia Armoracia</i> . E.B.				Eng. bot. 2223.
2	macrocarpa. Bau. (<i>wh.</i>) large-fruited.	Hungary.	1806.	6. 8. H. ¼.	
	<i>COCHLE'ARIA</i> . DC. SCURVY-GRASS. Tetradynamia Siliculosa. L.				W. et K. hung. 2. t. 184.
1	glaucifolia. DC. (<i>wh.</i>) Woad-leaved.	S. Europe.	1648.	5. 7. H. ♂.	
2	anglica. DC. (<i>wh.</i>) English.	England.	5. H. ♂.	
3	officinalis. DC. (<i>wh.</i>) common.	Britain.	4. 5. H. ♂.	
	β minor. DC. lesser.	—	—	— H. ♂.	
	γ rotundifolia. DC. round-leaved.	—	—	— H. ♂.	
	<i>granlandica</i> . With. brit. 573. non Lin.				
4	pyrenæica. DC. (<i>wh.</i>) Pyrenean.	Pyrenees.	6. 8. H. ♂.	
5	grænländica. DC. (<i>wh.</i>) Greenland.	Scotland.	5. 6. H. ¼.	
6	dánica. DC. (<i>wh.</i>) Danish.	Britain.	—	— H. ♂.	
7	alpina. (<i>wh.</i>) Alpine.	Ireland.	5. 6. H. ¼.	
8	fenestrata. DC. (<i>wh.</i>) fenestrate.	Baffin's Bay.	1820.	4. 5. H. ¼.	
9	grandiflora. DC. (<i>wh.</i>) large-flowered.	Siberia.	1826.	5. 6. H. ¼.	
10	integrifolia. DC. (<i>wh.</i>) entire-leaved.	Altay.	1827.	— H. ¼.	
11	acaulis. DC. (<i>li.</i>) stalkless.	Portugal.	1823.	3. 4. H. ¼.	
	<i>Deles. ic. sel. 2. t. 48.</i>				
	<i>Eng. bot. 2403.</i>				
	<i>———— 696.</i>				
	<i>Jacq. ecl. t. 132.</i>				

Tribus III. THLASPIDEÆ seu Pleurorhizeæ Angustiseptæ. DC.

THLA'SPI. DC.	BASTARD-CRESS.	Tetradynamia Siliculosa. L.				
1 latifolium. DC. (wh.)	broad-leaved.	Caucasus.	1828.	4. 5. H. 4.	Deles. ic. sel. 2. t. 51.	
2 ceratocarpon. DC. (wh.)	horn-podded.	Siberia.	1779.	7. H. ⊙.	Scop. insub. 1. t. 4.	
3 arvense. DC. (wh.)	Penny-cress.	Britain.	6. 7. H. ⊙.	Eng. bot. 1659.	
4 collinum. DC. (wh.)	hill.	Iberia.	1824.	4. 9. H. ⊙.	Jacq. ic. 1. t. 121.	
5 alliaceum. DC. (wh.)	Garlic-scented.	S. Europe.	1714.	5. 7. H. ⊙.	Eng. bot. 2354.	
6 strictum.	(wh.) upright.	Persia.	1827.	4. 9. H. ⊙.	Jacq. aust. 3. t. 237.	
7 perforiatum. DC. (wh.)	perfoliate.	England.	4. 7. H. ⊙.	Eng. bot. t. 81.	
8 montanum. DC. (wh.)	mountain.	Europe.	5. 6. H. 4.		
9 alpestre. DC. (wh.)	Alpine.	England.	5. 7. H. 4.		
10 cochleariforme. DC. (wh.)	Scurvy-grass.	Dahuria.	1827.	6. 7. H. 4.		
CA'PSELLA. DC.	SHEPHERDS-PURSE.	Tetradynamia Siliculosa. L.				
1 Búrsa-pastoris. DC. (w.)	common.	Britain.	1. 12. H. ⊙.	Eng. bot. 1485.	
<i>Thlásphi Búrsa-pastoris. L.</i>						
β minor. DC. (wh.)	small.	—	— H. ⊙.	Tabern. ic. 199.	
γ integrifolia. DC. (w.)	entire-leaved.	—	— H. ⊙.	Moris. ox. 2. t. 20. f. 1.	
δ coronopifolia. DC. (w.)	<i>Buck's-horn-l'd.</i>	—	— H. ⊙.	Hill. veg. sys. 11. t. 42. f. 1.	
2 procumbens. B. (wh.)	procumbent.	Europe.	1823.	3. 5. H. ⊙.		
<i>Lepidium procumbens. L. Hutchinsia procumbens. DC.</i>						
HUTCHI'NSIA. DC.	HUTCHI'NSIA.	Tetradynamia Siliculosa. L.				
1 rotundifolia. DC. (bh.)	round-leaved.	Switzerland.	1759.	5. 7. H. 4.	Scop. carn. n. 805. t. 37.	
<i>Ibèris rotundifolia. L.</i>						
2 cepeafolia. DC. (pu.)	toothed-leaved.	Carinthia.	1824.	4. 5. H. 4.	Jacq. misc. 2. p. 28. t. 1.	
3 stylosa. DC. (ro.)	long-styled.	Naples.	1826.	3. 5. H. 4.	Bot. mag. 2772.	
4 hastulata. DC. (bh.)	halbert-leaved.	Siberia.	1827.	— H. 4.	Gmel. sib. 3. t. 5.	
5 calycina. DC. (wh.)	persistent-cal'd.	—	—	4. 5. H. 4.	Jacq. aust. 2. t. 137.	
6 alpina. DC. (wh.)	Alpine.	Pyrenees.	1775.	4. 6. H. 4.		
<i>Lepidium alpinum. L.</i>						
7 petræa. DC.	rock.	England.	3. 5. H. ♂.	Eng. bot. 111.	
<i>Lepidium petræum. L.</i>						
TEESD'ALIA. DC.	TEESD'ALIA.	Tetradynamia Siliculosa. L.				
1 Ibèris. DC.	Shepherd's-cress.	England.	5. 7. H. ⊙.	Eng. bot. 327	
<i>nudicáulis. H.K. Ibèris nudicáulis. L. Guepinia Ibèris. DC.</i>						
2 Lepidium. DC. (wh.)	equal-petal'd.	S. Europe.	5. 8. H. ⊙.	Magn. ic. n. car. 247.	
<i>Lepidium nudicáule. L. Teesdália reguláris. L.T. Guepinia Lepidium. Dv.</i>						
IB'ERIS. DC.	CANDY-TUFT.	Tetradynamia Siliculosa. L.				
1 semperflorens. DC. (w.)	broad-leaved.	Sicily.	1679.	1. 12. G. ℡.	Moris. ox. 2. t. 25. f. 5.	
2 gibraltarica. DC. (bh.)	Gibraltar.	Gibraltar.	1732.	5. 6. G. ℡.	Bot. mag. 124.	
3 saxatilis. DC. (wh.)	rock.	S. Europe.	1739.	4. 6. H. ℡.	Gou. fl. mons. 177. t. 1.	
4 corifolia. (wh.)	Coris-leaved.	—	—	— H. ℡.	Clus. hist. 2. p. 132. ic.	
<i>saxatilis β corifolia. Bot. mag. 1642.</i>						
5 pubescens. DC. (wh.)	pubescent.	—	1821.	— H. ℡.		
6 subvelutina. DC. (wh.)	velvety.	Spain.	1824.	— H. ℡.	Flor. græc. t. 620.	
7 sempervirens. DC. (wh.)	evergreen.	Crete.	1731.	— H. ℡.	All. ped. n. 920. t. 40. f. 3.	
8 Garrexiana. DC. (wh.)	Allioni's.	Pyrenees.	6. 8. H. ℡.		
9 conferta. DC. (wh.)	crowded-leaved.	Spain.	1824.	— H. ℡.	Eng. bot. 52.	
10 amara. DC. (wh.)	bitter.	England.	— H. ⊙.	Lob. ic. t. 217. f. 2.	
11 pinnata. DC. (wh.)	wing-leaved.	S. Europe.	1596.	— H. ⊙.	Swt. br. fl. gar. 50.	
12 odorata. DC. (wh.)	sweet-scented.	Crete.	1806.	— H. ⊙.		
13 Lagascana. DC. (wh.)	Lagasca's.	Spain.	1824.	— H. ⊙.	Bot. mag. 106.	
14 umbellata. DC. (pu.)	purple.	S. Europe.	1596.	— H. ⊙.	Garid. aix. t. 105.	
15 limifolia. DC. (pu.)	Flax-leaved.	—	1759.	— H. ♂.		
16 ciliata. DC. (wh.)	fringed.	—	—	6. 7. H. ♂.		
17 taurica. DC. (wh.)	Taurian.	Tauria.	1802.	5. 7. H. ♂.	All. auct. p. 15. t. 2. f. 1.	
18 violacea. DC. (pu.)	blunt-leaved.	—	1782.	6. 7. H. ⊙.	Berg. phyt. ic.	
19 nana. DC. (pu.)	dwarf.	S. Europe.	1822.	5. 7. H. ♂.	Swt. br. fl. gar. 88.	
20 spathulata. DC. (pu.)	spathulate.	Pyrenees.	1826.	6. 8. H. ⊙.		
21 Tenoreana. DC. (wh.)	Tenore's.	Naples.	—	5. 7. H. ℡.		
22 Prouti. DC. (wh.)	Sicilian.	Sicily.	1828.	— H. ℡.		
23 contrácta. DC. (wh.)	contracted.	Spain.	1824.	4. 6. H. ℡.		
BISCUTE'LLA. DC.	BUCKLER-MUSTARD.	Tetradynamia Siliculosa. L.				
1 auriculata. DC. (ye.)	ear-podded.	S. Europe.	1683.	6. 7. H. ⊙.	DC. diss. n. 1. t. 1. f. 2.	
2 erigerifolia. DC. (ye.)	Erigeron-leav'd.	Spain.	1824.	— H. ⊙.	Deles. ic. sel. 2. t. 55.	
3 hispida. DC. (ye.)	hispid.	S. France.	1822.	— H. ⊙.	Bot. mag. 2444.	
4 cichoriifolia. DC. (ye.)	Succory-leaved.	Pyrenees.	—	— H. 4.	DC. diss. n. 4. t. 2.	

CRUCIFERÆ.

5 lyrata. DC.	(ye.) lyre-leaved.	Spain.	1799.	6. 7. H. ⊙.	Bocc. sic. 45. t. 25.
6 raphanifolia. DC.	(ye.) Radish-leaved.	Sicily.	1824.	— H. ⊙.	
7 marítima. DC.	(ye.) sea.	Naples.	—	— H. ⊙.	Tenor. fl. nap. t. 61.
8 ciliata. DC.	(ye.) fringe-podded.	Italy.	1790.	— H. ⊙.	DC. ic. gal. rar. 1. t. 39.
coronopifolia. w. non. Lin.					
9 depræsa. DC.	(ye.) dwarf.	Egypt.	1823.	— H. ⊙.	
10 microcárpa. DC.	(ye.) small-podded.	Gibraltar.	—	5. 6. H. ⊙.	DC. diss. n. 12. t. 9. f. 2.
11 eriocárpa. DC.	(ye.) woolly-podded.	Spain.	1826.	5. 7. H. ⊙.	Col. ecphr. l.t. 284. f. 1.
12 Columnæ. DC.	(ye.) Columna's.	Apulia.	1824.	6. 7. H. ⊙.	Fl. græc. t. 629.
13 ápula. DC.	(ye.) spear-leaved.	Italy.	1710.	— H. ⊙.	Gært. fruct. 2. t. 141.
14 leiocárpa. DC.	(ye.) smooth-capsul'd.	Levant.	1824.	— H. ⊙.	
15 obovata. DC.	(ye.) obovate-leaved.	— H. ⊙.	
16 montâna. DC.	(ye.) mountain.	Spain.	1824.	3. 4. H. ⊙.	Cavan. ic. 2. t. 177.
17 lævigåta. DC.	(ye.) smooth-podded.	Europe.	1777.	— H. 4.	DC. ic. rar. 11. t. 38.
β alpestris. W.en.	(ye.) Hungarian.	Hungary.	1816.	— H. 4.	W. et Kit. hung. 3. t. 228.
18 lúcida. DC.	(ye.) glossy.	Italy.	1827.	4. 6. H. 4.	DC. diss. n. 20. t. 7.
19 coronopifolia. DC.	(ye.) Buck's-horn-l'd.	S. Europe.	1790.	— H. 4.	DC. diss. n. 22. t. 18.
20 ambigua. DC.	(ye.) ambiguous.	— H. 4.	— n. 23. t. 11. f. 1.
21 saxatilis. DC.	(ye.) rock.	1824.	— H. 4.	— n. 24. t. 10.
22 sempervires. DC.	(ye.) downy-leaved.	Spain.	1784.	— F. ½.	Barr. ic. t. 841.
MEGACA'RPÆA. DC.	MEGACA'RPÆA.	Tetradynamia Siliculosa. L.			
laciniatæ. DC.	(ye.) jagged-leaved.	Volga.	1821.	6. 7. H. 4.	DC. diss. bisc. n. 5. t. 3.
Biscutella megalocarpa. F.					

Tribus IV. EUCLIDIEÆ seu Pleurorhizeæ Nucamentaceæ. DC.

EUCLIDIUM. DC.	EUCLIDIUM.	Tetradynamia Siliculosa. L.			
1 syriacum. DC.	(wh.) Syrian.	Levant.	1778.	7. 8. H. ⊙.	Jacq. aust. 1. t. 6.
Bünias syriaca. w.	Anastatica syriaca.	Jacq.			
2 tatáricum. DC.	(wh.) Tartarian.	Tartary.	1823.	4. 5. H. ⊙.	Pall. it. 3. ap. t. u. f. 2.
Bünias tatárica. w.					
OCHTHODIUM. DC.	OCHTHODIUM.	Tetradynamia Siliculosa. L.			
ægyptiacum. DC.	(ye.) Egyptian.	Egypt.	1787.	8. H. ⊙.	Jacq vind. t. 145.
Rapistrum ægyptiacum. H.K.	Bünias ægyptiaca.	L.			

Tribus V. ANASTATICEÆ seu Pleurorhizeæ Septulatæ. DC.

ANASTATICA. DC.	ROSE OF JERICHO.	Tetradynamia Siliculosa. L.			
Hierochuntina. DC. (w.) common.		Levant.	1597.	5. 8. H. ⊙.	Jacq. vind. 1. t. 58.

Tribus VI. CAKILINEÆ seu Pleurorhizeæ Lomentaceæ. DC.

CAK'ILE. DC.	CAK'ILE.	Tetradynamia Siliculosa. L.			
1 marítima. DC.	(pu.) Sea Rocket.	Britain.	6. 9. H. ⊙.	Eng. bot. 231.
β sinuatifolia. DC.	(p.) sinuate-leaved.	— H. ⊙.	Horn. fl. dan. t. 1583.
2 americana. DC.	(li.) American.	America.	— H. ⊙.	Tussac fl. ant. 1. f. 17.
CHORISPORA. DC.	CHORISPORA.	Tetradynamia Siliquosa. L.			
1 tenella. DC.	(pu.) purple.	Tauria.	1780.	6. 7. H. ⊙.	Pall. it. 3. ap. t. L. f. 3.
2 strícta. DC.	(pu.) upright.	Siberia.	1825.	5. 7. H. ⊙.	
3 sibírica. DC.	(ye.) Siberian.	Siberia.	1822.	— H. ⊙.	Murr. com. gött. t. 11.
4 ibérica. DC.	(ye.) strong-scented.	Iberia.	—	— H. ⊙.	Deles. ic. sel. 2. t. 582.

Subordo II. NOTORHIZEÆ. DC. syst. 2. p. 438.

Tribus VII. SISYMBREÆ seu Notorhizeæ Siliquosæ. DC.

MALC'OMIA. DC.	MALC'OMIA.	Tetradynamia Siliquosa. L.			
1 africana. DC.	(pu.) African.	Africa.	1747.	6. 7. H. ⊙.	Buxb. cent. 4. t. 44.
2 taraxacifolia. DC.	(pu.) Dandelion-l'd.	1795.	— H. ⊙.	
3 láxa. DC.	(pu.) loose-branched.	Siberia.	1822.	— H. ⊙.	

4 Chia. DC.	(pu.) dwarf.-branching.	Chio.	1732.	5. 6. H. ⊖.	Swt. br. fl. gar. 40.
5 marítima. DC.	(p.w.) dwarf annual.	S. Europe.	1713.	— H. ⊖.	Bot. mag. 166.
Cheiránthus marítimus. B.M.					
6 arenaria. DC.	(pu.) sand.	Barbary.	1804.	— H. ⊖.	Desf. fl. atl. 2. t. 162.
Héspéris arenaria. Desf.					
7 parviflora. DC.	(pu.) small-flowered.	S. Europe.	1822.	6. 8. H. ⊖.	DC. ic. gall. rar. t. 35.
8 lyrata. DC.	(pu.) rigid.	Levant.	1824.	5. 8. H. ⊖.	Flor. græc. t. 635.
9 littorea. DC.	(li.) sea.	S. Europe.	1683.	6. 11. H. ♂.	Swt. br. fl. gar. 54.
0 pátula. DC.	(pu.) spreading.	Spain.	1825.	5. 7. H. ♀.	
1 lácera. DC.	(pu.) cut-leaved.	S. Europe.	—	— H. ⊖.	Desf. ann. mus. 11.t. 34.
Héspéris pinnatifida. Desf.					
HE'SPÉRIS. DC.	ROCKET. Tetradymania Siliquosa. L.				
1 tristis. DC.	(br.) night-smelling.	Europe.	1622.	4. 6. H. ♂.	
β major.	(st.) larger.	—	1827.	4. 7. H. ♂.	Bot. mag. 730.
2 frágans. F.	(pu.) evening-scented.	Siberia?	1821.	— H. ♂.	Swt. br. fl. gar. 61.
3 laciniata. DC.	(pu.) laciniated.	Italy.	1820.	5. 6. H. ♂.	All. ped. n. 985.t.82.f.1.
4 villosa. DC.	(pu.) villous.	Italy.	1827.	— H. ♂.	
5 grandiflora. B.M.	(pu.) large-flowered.	1818.	5. 8. H. ♂.	Bot. mag. 2683.
6 elata. H.	(pu.) tall.	1825.	— H. ♂.	W. et Kit. hung. 2.t.200.
7 runcinata. DC.	(bh.) runcinate.	Hungary.	1804.	5. 7. H. ♂.	
8 bituminosa. W.en.	(bh.) clammy.	1816.	— H. ♂.	Lob. ic. t. 323. f. 2.
9 matronalis. L.	(va.) common.	Europe.	1597.	5. 8. H. ♀.	Tabern. kraut. 692. f. 1.
α albibiflora. DC.	(wh.) single-white.	—	—	— H. ♀.	
β álbo-plena.	(wh.) double-white.	—	—	— H. ♀.	
γ purpurea. DC.	(pu.) single-purple.	—	—	— H. ♀.	
δ purpureo-plena. (pu.)	double-purple.	—	—	— H. ♀.	
ε variegata. DC.	(v.) dble. -variegated.	—	—	— H. ♀.	
ζ foliiflora. DC.	(gr.) green-flowered.	—	—	— H. ♀.	
10 inodóra. L.	(pu.) scentless.	Britain.	— H. ♀.	
11 sibírica. L.	(pu.) Siberian.	Siberia.	— H. ♀.	
12 heterophylla. DC.	(pu.) various-leaved.	Naples.	1825.	5. 7. H. ♀.	
13 áprica. DC.	(pu.) sun.	Siberia.	1823.	5. 6. H. ♀.	
14 ramosissima. DC.	(pu.) branched.	Barbary.	1827.	— H. ⊖.	Deles. ic. sel. 2. t. 62.
ANDREO'SKIA. DC.	ANDREO'SKIA. Tetradymania Siliquosa. L.				Desf. atl. 2. t. 161.
1 integrifolia. DC.	(wh.) entire-leaved.	Siberia.	1824.	6. 7. H. ⊖.	
Sisy'mbrium integrifolium. w. Héspéris glándulosa. P.S.					
2 pectinata DC.	(wh.) pectinated.	Dahuria.	1827.	5. 6. H. ♂.	
SISY'MBRIUM. DC.	SISY'MBRIUM. Tetradymania Siliquosa. L.				
1 officinale. DC.	(ye.) Hedge-mustard.	Britain.	5. 7. H. ⊖.	Eng. bot. 725.
Ery'simum officinale.	E.B.				
2 strictissimum. DC.	(ye.) spear-leaved.	Europe.	1658.	6. 8. H. ♀.	Jacq. aust. 2. t. 194.
Cheir'nia strictissima. L. en.					
3 júnceum. DC.	(ye.) rushy.	Hungary.	1804.	7. H. ♀.	W. et Kit. hung. 3.t.234.
Ery'simum júnceum. w. Cheir'nia júncea. L. en.					
4 hispánicum. DC.	(ye.) Spanish.	Spain.	1820.	5. 6. H. ♂.	Jacq. ic. rar. 1. t. 124.
5 pùmulum. DC.	(ye.) dwarf.	Caucasus.	—	4. 5. H. ⊖.	
6 obtusángulum. DC.	(ye.) blunt-angled.	Switzerland.	5. 8. H. ⊖.	Moris. ox. 2.s.3.t.5.f.10.
7 acutángulum. DC.	(ye.) acute-angled.	Pyrenees.	1791.	6. 7. H. ♂.	Jacq. vind. 3. t. 97.
sinapoides. H.K. Sinápis pyrenæica. Jacq.					
8 taraxacifolium. DC.	(y.) Dandelion-ld.	France.	1823.	5. 6. H. ♂.	DC. ic. rar. p.11. t. 37.
9 contortum. W.en.	(ye.) twisted.	Spain.	1824.	— H. ♂.	
10 affine. W.en.	(ye.) related.	1823.	5. 8. H. ♂.	
11 austriacum. DC.	(ye.) Austrian.	Austria.	1799.	6. 8. H. ⊖.	Jacq. aust. 3. t. 262.
12 eckartsbergense. w.	(y.) bending-podd'd.	—	—	— H. ♂.	
Tillieri. w.	(ye.) Tillier's.	Switzerland.	1824.	6. 8. H. ⊖.	
I'rio. DC.	(ye.) London Rocket.	England.	5. 8. H. ⊖.	Eng. bot. 1631.
15 glábrum. W.en.	(ye.) smooth.	1823.	— H. ⊖.	
16 gallicum. W.en.	(ye.) French.	France.	—	— H. ⊖.	
17 nítidum. DC.	(ye.) glossy.	6. 8. H. ⊖.	
18 subhastátum. DC.	(ye.) halbert-leaved.	Archipelago.	1822.	4. 5. H. ⊖.	
19 Colùmnæ. W.	(ye.) Columnna's.	Italy.	1796.	6. 7. H. ⊖.	Jacq. aust. t. 323.
20 altissimum. L.	(ye.) tall.	Siberia.	1759.	8. H. ⊖.	Walth. hort. t. 122.
21 orientàle. L.	(ye.) oriental.	Levant.	1739.	7. 8. H. ⊖.	
22 pannonicum. DC.	(st.) Hungarian.	Hungary.	1787.	— H. ⊖.	Jacq. ic. rar. 1. t. 123.
23 lyratum. DC.	(ye.) lyrate.	C. B. S.	5. 8. G. ♀.	Deles. ic. sel. 2. t. 64.
24 ásperum. DC.	(ye.) rough-podded.	S. Europe.	1778.	5. 6. H. ⊖.	J. Bauh. hist. 2. f. 3.
25 fugáx. DC.	(ye.) fugacious.	Spain.	1820.	7. 8. H. ♀.	
26 Sophia. DC.	(ye.) Flix-weed.	Britain.	7. 8. H. ⊖.	Eng. bot. 963.

CRUCIFERÆ.

27 péricum. DC.	(ye.) Persian.	Persia.	1820.	7. 8. H. ⊙.
28 Richardsóni.	(ye.) Richardson's.	N. America.	1828.	6. 7. H. ⊙.
<i>canescens</i> . Richardson,	<i>non Nutt.</i>			
29 brachycarpum. DC.	short-podded.	_____	_____	H. ⊙.
30 tripinnatum. DC.	(ye.) tripinnate.	C. B. S.	1817.	_____ H. ⊙.
31 milfolium. DC.	(ye.) Milfoil-leaved.	Canaries.	1779.	5. 9. H. ⊙.
32 tanacetifolium. DC.	(ye.) Tansy-leaved.	Italy.	1731	6. 7. H. ¼.
33 supinum. DC.	(wh.) trailing.	S. Europe.	1778.	_____ H. ⊙.
<i>Ery'simum supinum</i> .	L. en.			
34 hirsutum. DC.	(ye.) hairy-podded.	_____	1822.	_____ H. ⊙.
<i>Ery'simum hirsutum</i> .	L. en.			
35 polyceratum. DC.	(ye.) sinuate-leaved.	_____	1633.	6. 8. H. ⊙.
<i>Ery'simum polyceratum</i> .	L. en.			Jacq. vind. t. 79.
36 rígidum. DC.	(wh.) rigid.	Tauria.	1824.	6. 8. H. ⊙.
37 bursifolium. DC.	(wh.) various-leaved.	_____	1732.	_____ H. ⊙.
38 pinnatifidum. DC.	(wh.) pinnatifid.	_____	1822.	5. 8. H. ¼.
39 erysimoides. DC.	(wh.) Erysimum-like.	Spain.	1827.	10. 12. H. ¼.
40 contortuplicatum. (bh.)	twisted-podded.	Cumana.	_____	6. 8. H. ⊙.
<i>Heiráspis contortuplicatus</i> . Bieb.	<i>Cheiránthus contortuplicatus</i> . w.			
ALLIÁRIA. DC.	ALLIÁRIA.	Tetradynamia Siliquosa. L.		
1 officinalis. DC.	(wh.) garlic-scented.	Britain.	5. 6. H. ¼.
<i>Ery'simum officinale</i> .	E.B. <i>Hésporis Alliaria</i> . Lam.			Eng. bot. 796.
2 brachycarpa. DC.	(wh.) short-podded.	Iberia.	1827.	5. 7. H. ¼.
ERY'SIMUM. DC.	ERY'SIMUM.	Tetradynamia Siliquosa. L.		
1 siliculosum. DC.	(ye.) siliculose.	Tauria.	1824.	5. 6. H. ♂.
2 sículum. DC.	(ye.) Sicilian.	Sicily.	1824.	6. 7. H. ♂.
3 sessiliflorum. DC.	(ye.) sessile-flower'd.	Siberia.	1794.	_____ H. ¼.
<i>Cheiránthus quadrangulus</i> . w.	<i>Cheirínia sessiliflora</i> . L. en.			<i>Syrénia Lamárcii</i> . Andrz.
4 angustifolium. DC.	narrow-leaved.	Hungary.	1800.	7. 8. H. ♂.
<i>Cheirínia angustifolia</i> .	L. en.			W. et Kit. hung. t. 98.
5 cuspidatum. DC.	(ye.) sharp-pointed.	Caucasus.	1800.	5. 6. H. ♂.
<i>Syrénia Biebersteinii</i> .	Andrz.	<i>Cheiránthus cuspidatus</i> . w.		Buxb. cent. 2. t. 33. f. 1.
6 leptostylum. DC.	(ye.) slender-styled.	Caucasus.	1825.	_____ H. ♂.
7 exaltatum. DC.	(ye.) tall.	Poland.	_____	_____ H. ♂.
8 suffruticosum. DC.	(st.) suffruticose.	1822.	4. 5. H. ♂.
9 stríctum. DC.	(ye.) upright.	Austria.	1795.	6. 8. H. ♂.
<i>odoratum</i> . w. <i>hieracifolium</i> . Jacq.	<i>Cheirínia strícta</i> . L. en.			Jacq. aust. 1. t. 73.
10 virgátum. DC.	(ye.) twiggy.	Portugal.	1807.	6. 7. H. ♂.
11 longisiliquosum. DC.	(ye.) long-podded.	Switzerland.	1823.	_____ H. ♂.
12 hieracifolium. DC.	(ye.) Hawkweed-ld.	Europe.	_____	5. 6. H. ♂.
13 Marschallianum. (ye.)	Marschall's.	Poland.	1827.	_____ H. ♂.
14 pátilum. H.	(ye.) spreading.	1824.	_____ H. ♂.
15 púmilum. H.	(ye.) dwarf.	Switzerland.	_____	_____ H. ♂.
16 áureum. DC.	(ye.) golden.	Caucasus.	1824.	_____ H. ♂.
17 ibéricum. DC.	(ye.) Iberian.	Iberia.	1803.	5. 8. H. ♂.
<i>Cheiránthus armeniacus</i> . B.M.	<i>Cheiránthus ibéricus</i> . Bieb.			Deles. ic. sel. 2. t. 66.
18 cheiranthoides. DC.	(ye.) treacle.	Britain.	Bot. mag. 835.
19 repandum. DC.	(ye.) small-flowered.	S. Europe.	1772.	5. 6. H. ⊙.
20 helveticum. DC.	(ye.) Swiss.	Switzerland.	1793.	_____ H. ♂.
<i>Cheiránthus helveticus</i> . Jacq.	<i>Cheirínia helvetica</i> . L. en.			Eng. bot. 942.
21 canescens. DC.	(ye.) canescent.	S. Europe.	1731.	5. 7. H. ♂.
<i>diffusum</i> . w. <i>Cheiránthus alpinus</i> . Jacq.	aust. 1. t. 75.			Jacq. aust. 1. t. 22.
22 Andrzejowskianum. (ye.)	Andrzejowski's.	Tauria.	1828.	_____ H. ♂.
23 collinum. DC.	(ye.) hill.	Caucasus.	_____	_____ H. ♂.
24 leptophyllum. DC.	(ye.) slender-leaved.	Iberia.	1827.	_____ H. ♂.
25 versicolor. DC.	(va.) various-color'd.	Persia.	1823.	5. 7. H. ♂.
<i>Cheiránthus versicolor</i> . M.B.				Jacq. vind. t. 9.
26 lanceolatum. DC.	(ye.) spear-leaved.	_____	1597.	_____ H. ♂.
<i>Cheiránthus erysimoides</i> . Jacq.				Jacq. aust. t. 74.
27 fírmum.	(ye.) hoary-podded.	Switzerland.	1824.	_____ H. ♂.
<i>Cheirínia firma</i> . L. en.	<i>Cheiránthus firmus</i> . W. en.			Barr. ic. t. 884.
28 rhæticum. DC.	(ye.) Rhætian.	Rhætia.	_____	_____ H. ♂.
29 dùbium. DC.	(ye.) doubtful.	_____	_____ H. ♂.
30 longifolium. DC.	(ye.) long-leaved.	Algiers.	1822.	_____ H. ¼.
31 gráile. DC.	(ye.) slender.	Caucasus.	1824.	6. 8. H. ♂.
32 áperum. DC.	(ye.) rough.	Missouri.	_____	_____ H. ♂.
33 strigósum. DC.	(ye.) strigose.	Siberia.	1828.	_____ H. ♂.
34 Redóskii. DC.	(ye.) Redóski's.	_____	1825.	_____ H. ♂.

Jacq. ic. rar. 1. t. 127.
Mor. his. 2. s. 3. t. 6. f. 19.
Isn. act. Par. 1724. t. 18.

35 altissimum. DC. (ye.) tallest. <i>Cheirinia altissima.</i> L.en.	Limbourg. 1822. 6. 7 H.-♂.
36 sabulosum. F. (ye.) gravel.	Siberia. 1823. — H. ♂.
37 bicolor. DC. (v.) two-coloured.	Switzerland. 1827. — H. ♂.
CONRI'NGIA. L. en. CONRI'NGIA.	Tetradynamia Siliquosa. L.
1 alpina. L. en. (wh.) Alpine. <i>Brássica alpina.</i> L. <i>Ery'simum alpinum.</i> DC.	Europe. 1793. 5. 6. H. ♀. Vill. dauph. 3. t. 36.
2 perfoliata. L. en. (wh.) perfoliate. <i>Brássica orientalis.</i> E.B. <i>Ery'simum perfoliatum.</i> DC.	England. — H. ⊙. Eng. bot. 1804.
3 austriaca. (st.) Austrian. <i>Brássica austriaca.</i> Jacq. <i>Ery'simum austriacum.</i> DC.	Austria. — H. ⊙. Jacq. aust. t. 283.
STANLEYA. DC. STANLEYA.	Tetradynamia Siliquosa.
pinnatifida. DC. pinnatifid.	Missouri. 1812. 5. 6. H. ♀.
<i>Cleome pinnata.</i> Ph.	

Tribus VIII. CAMELINEÆ seu Notorhizeæ Latiseptæ. DC.

CAMEL'INA. DC. CAMEL'INA.	Tetradynamia Siliculosa. L.
1 sativa. DC. (ye.) Gold of pleasure. Britain. 5. 7. H. ⊙. Eng. bot. 1254.
<i>Alyssum sativum.</i> E.B. <i>Myagrum sativum.</i> L.	
α pilosa. DC. (ye.) hairy. — H. ⊙. Moris. ox. 2.s. 3.t. 21. f. 2.
β glabrata. DC. (ye.) smooth. — H. ⊙. Lind. als. 94. t. 1.
2 dentata. DC. (ye.) tooth-leaved.	Europe. 1806. 6. 7. H. ⊙. J. Bauh. hist. 2.p. 893. ic.
3 pinnatifida. H. (ye.) pinnatifid.	Spain. — — H. ⊙.
4 microcarpa. DC. (ye.) small-podded.	Poland. 1827. — — H. ⊙. Deles. ic. sel. 2. t. 69.
5 barbareafolia. DC. (ye.) Barbarea-ld.	Siberia. 1824. 5. 7. H. ♀. — — 2. t. 70.
6 austriaca. DC. (ye.) Austrian.	Austria. 1795. 6. 7. H. ⊙. Jacq. aust. 2. t. 111.
NE'SLIA. DC. NE'SLIA.	Tetradynamia Siliculosa.
paniculata. DC. (ye.) panicled.	Europe. 1683. 6. 8. H. ⊙. Flor. dan. t. 204.
<i>Rapistrum paniculatum.</i> Gært. <i>Myagrum paniculatum.</i> L.	

Tribus IX. LEPIDINEÆ seu Notorhizeæ Angustiseptæ. DC.

SENEBIE'RA. DC. SENEBIE'RA.	Tetradynamia Siliculosa. L.
1 pinnatifida. DC. (wh.) pinnatifid.	England. 7. 8. H. ⊙. Eng. bot. 248.
<i>Corónopus dídyma.</i> E.B. <i>Lepidium dídymum.</i> L.	
β incisa. W.en. (wh.) cut-leaved.	America. 1817. — — H. ⊙.
2 Corónopus. DC. (wh.) common.	Britain. 6. 10. H. ⊙. Eng. bot. 1660.
<i>Corónopus Ruéllii.</i> E.B. <i>Cochlearia Corónopus.</i> L.	
LEPI'DIUM. DC. PEPPERWORT.	Tetradynamia Siliculosa. L.
1 Drába DC. (wh.) Whitlow-grass.	Europe. 1596. 5. 6. H. ♀. Jacq. fl. aust. 4. t. 315.
<i>Cochlearia Drába.</i> Jacq.	
2 chalepense. DC. (wh.) Aleppo.	Aleppo. 1798. 5. 7. H. ⊙. Moris. ox. 2. s. 3. t. 25.
3 coronopifolium. DC. (w.) Buckshorn-ld.	Volga. 1823. — — H. ♀.
4 sativum. DC. (wh.) common Cress.	Persia. 1548. 5. 10. H. ⊙.
β críspum. DC. (wh.) curled Cress.	— — — — H. ⊙.
γ latifolium. DC. (wh.) broad-leaved.	— — — — H. ⊙.
5 campéstre. DC. (wh.) hoary field.	Britain. 6. 7. H. ⊙.
<i>Thlaspi campéstre.</i> E.B.	
6 hirtum. DC. (wh.) hairy.	— — — — H. ♂. — — 1803.
<i>Thlaspi hirtum.</i> E.B.	
7 spinosum. DC. (wh.) prickly.	Levant. 1787. 4. 9. H. ⊙.
8 virginicum. DC. (wh.) Virginian.	America. 1713. 6. 7. H. ⊙.
9 subulatum. DC. (wh.) awl-leaved.	Spain. 1739. 7. 8. F. ♂.
10 ruderale. DC. (wh.) narrow-leaved.	Britain. 6. 7. H. ⊙.
11 incisum. DC. (wh.) cut-leaved.	Tauria. 1824. 5. 6. H. ⊙.
12 vesicarium. DC. (wh.) bladdered.	Iberia. 1821. 4. 8. H. ⊙.
13 angulósum. DC. (wh.) angular-stalk'd.	Hungary. 1825. — — H. ⊙.
14 perfoliatum. DC. (wh.) various-leaved.	Austria. 1640. 4. 6. H. ⊙.
15 Cardámines. DC. (wh.) Spanish Cress.	Spain. 1789. 6. 7. F. ♂.
16 divaricatum. DC. (wh.) close-spiked.	C. B. S. 1774. 5. 8. G. ♂.
17 bonariense. DC. (wh.) Buenos Ayres.	S. America. 1732. 5. 6. H. ⊙.
18 pubescens. DC. (wh.) pubescent.	Para. 1820. 5. 8. H. ⊙.
19 cuneifolium. DC. (wh.) wedge-leaved.	N. S. W. 1820. 6. 8. H. ⊙.

CRUCIFERÆ.

20	piscidium. DC.	(wh.)	South Sea.	Society Isles.	1779.	9.	H. ⊙.	Mont. n.a.n.c. 6. t. 5.a.
21	desertorum. Led.	(wh.)	desert.	Siberia.	1827.	—	H. ⊙.	
22	decumbens. DC.	(wh.)	decumbent.	1820.	5. 8.	H. ⊙.	
23	oleraceum. DC.	(wh.)	eatable.	New Zealand.	—	H. ⊙.	
24	lyratum. DC.	(wh.)	lyre-leaved.	Levant.	1759.	6. 7.	H. ⊙.	Tourn. it. 2. t. 389.
25	latifolium. DC.	(wh.)	broad-leaved.	Britain.	—	H. ¼.	Eng. bot. 182.
26	affine. DC.	(wh.)	saw-leaved.	Siberia.	1823.	7. 8.	H. ¼.	W. et Kit. hung. 1. t. 4.
27	crassifolium. DC.	(wh.)	thick-leaved.	Hungary.	—	5. 6.	H. ¼.	Cav. ic. 2. t. 161. f. 2.
28	suffruticosum. DC.	(w.)	shrubby.	Spain.	1790.	7. 8.	F. ♂.	
	graminifolium. Cav.							
29	lineare. DC.	(wh.)	linear-leaved.	Spain.	—	F. ♂.	
30	Iberis. DC.	(wh.)	grass-leaved.	S. Europe.	1683.	6. 9.	H. ¼.	Moris. his. 2.s.3.t.21.f.1.
	graminifolia. H.K.							
31	capéense. DC.	(wh.)	Cape.	C. B. S.	1818.	6. 7.	G. ¼.	
BIVON'EA. DC.	BIVON'EA.			Tetradynamia Siliculosa. L.				
lutea. DC.	(ye.)	yellow.	Sicily.	1823.	4. 5.	H. ⊙.	Cup. pan. sic. 2. t. 256.	
Thlaspi luteum. Biv.								
ÆTHION'EMA. DC.	ÆTHION'EMA.			Tetradynamia Siliculosa. L.				
1	saxatilis. DC.	(w.p.)	rock.	S. Europe.	1759.	6. 7.	H. ⊙.	Jacq. aust 3. t. 236.
2	gracile. DC.	(wh.)	slender.	Carniola.	1822.	—	H. ♂.	Boc. mus. p. 79. t. 70.
3	Buxbaumii. DC.	(pu.)	Buxbaum's.	Iberia.	1820.	—	H. ⊙.	Buxb. cent. 1. t. 5. f. 1.
4	monospermum DC.	(p.)	one-seeded.	Spain.	1778.	7. 8.	H. ♂.	

Tribus X. ISATIDEÆ seu Notorhizeæ Nucamentaceæ. DC.

TAUSCH'ERIA. DC.	TAUSCH'ERIA.			Tetradynamia Siliculosa. L.				
1	lasiocarpa. DC.	(wh.)	villous-podded.	Siberia.	1824.	6. 7.	H. ⊙.	
2	gymnocarpa. DC.	(wh.)	smooth-podded.	—	—	5. 6.	H. ⊙.	
IS'ATIS. DC.	WOAD.			Tetradynamia Siliculosa. L.				
1	armena. DC.	(ye.)	Armenian.	Armenia.	1824.	6. 7.	H. ⊙.	Desv. j. 3. t. 25. f. 6.
2	latiflqua. DC.	(ye.)	broad-podded.	Iberia.	1827.	5. 7.	H. ♂.	
3	lusitánica. DC.	(ye.)	Portugal.	Portugal.	1739.	—	H. ⊙.	All. ped. t. 86. f. 2.
4	alpina. DC.	(ye.)	Alpine.	Italy.	1800.	—	H. ¼.	Trat. arch. 2. t. 68.
5	præcox. DC.	(ye.)	early-flowering.	Hungary.	1820.	4. 6.	H. ♂.	Deles. ic. sel. 2. t. 78.
6	littoralis. DC.	(ye.)	sea-side.	Tauria.	—	6. 7.	H. ⊙.	— 2. t. 79.
7	hebecarpa. DC.	(ye.)	pubescent-podded.	Tauria.	1827.	—	H. ⊙.	
8	dasyarpa. Led.	(ye.)	thick-podded.	Siberia.	—	5. 7.	H. ♂.	
9	tinctória. DC.	(ye.)	common.	England.	5. 7.	H. ♂.	Eng. bot. 97.
β	sativa. DC.	(ye.)	smooth-leaved.	—	H. ♂.	Blackw. herb. t. 246.
γ	hirsuta. DC.	(ye.)	hairy-leaved.	—	H. ♂.	
δ	microcarpa. DC.	(ye.)	small-podded.	—	H. ♂.	
10	campestris. DC.	(ye.)	field.	Podolia.	1823.	—	H. ⊙.	
11	bannática. DC.	(ye.)	Bannatic.	Hungary.	—	—	H. ⊙.	
12	táurica. DC.	(ye.)	Taurian.	Tauria.	1827.	—	H. ♂.	
13	oblongata. DC.	(ye.)	oblong-podded.	Siberia.	—	—	H. ♂.	
14	orientalis. DC.	(ye.)	oriental.	Levant.	—	H. ⊙.	
15	canescens. DC.	(ye.)	canescent.	S. Europe.	1825.	—	H. ♂.	Buxb. cent. 1. p. 4.t.5.
16	alépica. DC.	(ye.)	Aleppo.	Levant.	1739.	6. 7.	H. ⊙.	Scop. d. ins. 2. t. 16.
17	variegata. Led.	(ye.)	variegated.	Siberia.	1827.	—	H. ♂.	
MYAGRUM. DC.	MYAGRUM.			Tetradynamia Siliculosa. L.				
perfoliatum. DC.	(ye.)	perfoliate.	Europe.	1648.	6. 7.	H. ⊙.	Schkuhr. han. 2. t. 178.	
SOBOLEWSKIA. DC.	SOBOLEWSKIA.			Tetradynamia Siliculosa. L.				
lithóphila. DC.	(wh.)	large-fruited.	Iberia.	1823.	5. 8.	H. ♂.	Deles. ic. sel. 2. t. 80.	

Tribus XI. ANCHONIEÆ seu Notorhizeæ Lomentaceæ. DC.

GOLDBA'CHIA. DC.	GOLDBA'CHIA.			Tetradynamia Siliquosa.				
1	laevigata. DC.	(li.)	smooth.	Astracan.	1823.	5. 6.	H. ⊙.	Deles. ic. sel. 2. t. 81.
2	torulosa. DC.	(li.)	twisted-podded.	Levant.	—	H. ⊙.	
3	tetragona. Led.	(li.)	four-sided.	—	1825.	—	H. ⊙.	
STERI'GMA. DC.	STERI'GMA.			Tetradynamia Siliquosa.				
1	tomentosum. DC.	(ye.)	tomentose.	Caspia.	1823.	4. 5.	H. ♂.	Pall. it. 2. ap. t. K. f. 2.
	Cheiránthus tomentosus. w.							
2	torulosum. DC.	(ye.)	twisted-podded.	Iberia.	—	—	H. ♂.	Buxb. cent. 5. t. 52.f. 2.

Subordo III. ORTHOPLOCEÆ. DC. syst. 2. p. 581.

Tribus XII. BRASSICEÆ seu Orthoploceæ Siliquosæ. DC.

BRA'SSICA. DC.	CABBAGE. Tetrodynamia Siliquosa. L.	England.	4. 6. H. ♂.	
1 oleracea. DC. (va.) common.	England.	H. ♂.	Eng. bot. t. 637.
A. SYLVE'STRIS. DC. (st.) WILD.	H. ♂.	
B. ACE'PHALA. DC. (st.) CAVALIER.	H. ♂.	
α ramðsa. DC. (st.) thousand-headed.	H. ♂.	
β vulgáris. DC. (st.) upright.	H. ♂.	Moris. ox. s. 3. t. 1. f. 6.
* viridis. DC. (st.) green.	H. ♂.	Lob. ic. 243. f. 1.
** purpuráscens. DC. (st.) purple.	H. ♂.	Chabr. sciag. 270. f. 6.
γ quercifòlia. DC. (st.) oak-leaved.	H. ♂.	_____ 270. f. 1.
δ sabellíca. DC. (st.) BORECOLE.	H. ♂.	_____ 271. f. 1.
* pinnata. DC. (st.) winged-leaved.	H. ♂.	Lob. ic. 246. f. 2.
** purpuráscens. DC. (st.) purple.	H. ♂.	
*** versicolor. DC. (st.) variegated.	H. ♂.	
ε palmifòlia. DC. (st.) palm-leaved.	H. ♂.	
ζ costatâ. DC. (wh.) thick-ribbed.	H. ♂.	
* nepenthiformis. DC. (wh.) tubular-l.	H. ♂.	Hort. trans. 5. p. 13. t. 1.
C. BULLATA. DC. (st.) SAVOY.	H. ♂.	Weinm. phyt. t. 261.
α vulgáris. DC. (st.) common.	H. ♂.	Lob. ic. t. 244. f. 1.
* præcox. DC. (st.) early.	H. ♂.	
** humili. DC. (st.) dwarf.	H. ♂.	
*** turionénis. DC. (st.) Tourraine.	H. ♂.	
**** aurâta. DC. (st.) yellow.	H. ♂.	Lob. ic. t. 244. f. 2.
β oblonga. DC. (st.) winter.	H. ♂.	
γ majo. DC. (st.) large.	H. ♂.	
δ gemmifera. DC. (st.) Brussel's-sprouts.	H. ♂.	Lob. ic. 243. f. 2.
D. CAPIT'ATA. DC. (st.) HEADED.	H. ♂.	
α deprissa. DC. (st.) drum-head.	H. ♂.	
β sphæ'rica. DC. (st.) common.	H. ♂.	
* alba. DC. (st.) white.	H. ♂.	
** rûbra. DC. (st.) red.	H. ♂.	Dod. pempt. 621. f. 2.
γ obovata. DC. (st.) Sugarloaf.	H. ♂.	
δ ellíptica. DC. (st.) early york.	H. ♂.	
ε subacuta.	(st.) Battersea.	H. ♂.	
ζ cónica. DC. (st.) heart.	H. ♂.	
E. CA'ULO-R'APA. DC. TURNEP-CABBAGE.	H. ♂.	Dalech. lugd. 522. f. 3.
α communis. DC. (st.) flat-leaved.	4. 6. H. ♂.		Lob. ic. 246. f. 1.
* alba. DC. (st.) white-stemmed.	H. ♂.	
** purpuráscens. DC. (st.) violet-stemd.	H. ♂.	Weinm. phyt. t. 264.
β? críspa. DC. (st.) curl-leaved.	H. ♂.	
F. BOTR YTIS. DC. (st.) BROCCOLI.	H. ♂.	Lob. ic. 245. f. 1.
α cauliflora. DC. (st.) Cauliflower.	H. ♂.	Weinm. phyt. t. 256.
β asparagóides. DC. (st.) Broccoli.	H. ♂.	Dalech. lugd. 522. f. 1.
* communis. DC. (st.) white.	H. ♂.	Chabr. sciagr. 269. f. 3.
** violacea. DC. (w.s.) violet.	H. ♂.	
*** viridis. (w.s.) green.	H. ♂.	
**** ramðsa. (w.s.) branching.	H. ♂.	
***** autumnális. (st.) Cape.	H. ♂.	
2 campéstris. DC. (ye.) field.	England.	6. H. ♂.	Eng. bot. 2224.
A. OLE'FERA. DC. (ye.) COLE-SEED.	H. ♂.	Dod. pempt. 623. f. 1.
* autumnális. DC. (ye.) autumnal.	H. ♂.	
** præcox. DC. (ye.) early.	H. ♂.	
B. PA'BULARIA. DC. (ye.) SHEEP'S.	H. ♂.	
C. N'APO-BRA'SSICA. NAVEW.	H. ♂.	
α communis. DC. (ye.) common.	H. ♂.	
* alba. DC. (ye.) white-rooted.	H. ♂.	
** purpuráscens. DC. (ye.) purple-rooted.	H. ♂.	
β Rutabaga. DC. (st.) Swedish Turnep.	H. ♂.	
3 Ràpa. DC. (ye.) Turnep.	Britain.	4. 6. H. ♂.	Eng. bot. 2176.
A. DEPRE'SSA. DC. FLAT-ROOTED.	H. ♂.	Mart. fl. rust. t. 49. 50.
* alba. DC. (ye.) white-round.	H. ♂.	
** flavescens. DC. (ye.) yellow-round.	H. ♂.	
*** nigricans. DC. (ye.) dark-round.	H. ♂.	
**** punicea. DC. (ye.) red-round.	H. ♂.	
***** viridis. DC. (ye.) green-round.	H. ♂.	
***** præcox. (ye.) early Dutch.	H. ♂.	

B. OBLONGA. DC. (<i>ye.</i>)	OBLONG-ROOTED.	Britain.	4. 6. H. ♂.	Lob. ic. t. 197. f. 2.
C. OLEI' FERA. DC. (<i>y.</i>)	OIL-BEARING.	Dauphiny.	— H. ♂.	— t. 298. f. 1.
4 Nápus. DC.	<i>(ye.) Rape.</i>	Britain.	5. H. ♂.	Eng. bot. 2146.
A. OLEI' FERA. DC.	<i>(ye.) wild.</i>	—	— H. ♂.	Lob. ic. t. 200. f. 2.
B. ESCULE'NTA. DC.	<i>(ye.) esculent.</i>	—	— H. ♂.	— 200. f. 1.
α álba. DC.	<i>(ye.) white-rooted.</i>	—	— H. ♂.	Blackw. herb. t. 410.
β fláva. DC.	<i>(ye.) yellow-rooted.</i>	—	— H. ♂.	
γ nígricans. DC.	<i>(ye.) black-rooted.</i>	—	— H. ♂.	
5 pr'ecox. DC.	<i>(st.) early.</i>	S. Europe.	5. 6. H. ♂.	Cup. panph. 1. t. 132.
6 chinénsis. DC.	<i>(ye.) Chinese.</i>	China.	1770. 7. H. ♂.	Deles. ic. sel. 2. t. 86.
7 violacea. DC.	<i>(li.) violet-flowered.</i>	—	— H. ♂.	Ten. fl. nap. t. 62.
8 incána. DC.	<i>(st.) hoary.</i>	Naples.	1824. 4. 5. H. ♂.	Desf. fl. atl. 2. t. 165.
9 baleárica. DC.	<i>(st.) oak-leaved.</i>	Minorca.	— 5. 7. H. ♂.	Vill. dauph. 3. t. 39.
10 hy'brida. T.C.	<i>(st.) hybrid.</i>	Naples.	—	— 3. t. 36.
11 Gravínae. DC.	<i>(ye.) pubescent.</i>	Aprutia.	— 5. 6. H. ♀.	Eng. bot. 962.
12 pinnatifida. DC.	<i>(st.) pinnatifid.</i>	Spain.	— 9. 10. H. ♀.	Bull. herb. t. 331.
13 húmilia. DC.	<i>(ye.) dwarf.</i>	S. France.	— 4. 5. H. ♀.	Vill. dauph. 3. t. 36.
14 repándia. DC.	<i>(ye.) wavy.</i>	S. Europe.	1818. 6. 8. H. ♀.	Willd. hort. ber. t. 19.
15 Richérii. DC.	<i>(ye.) alpine.</i>	—	1825. 5. 8. H. ♀.	Gou. ill. p. 44. t. 20. f. A.
16 monénensis. DC.	<i>(st.) Isle of Man.</i>	Britain.	—	
17 Erucástrum. DC.	<i>(ye.) runcinate.</i>	S. Europe.	1790. —	
18 cheiránthos. DC.	<i>(ye.) Stock-like.</i>	—	—	
19 cheiranthiflora. DC.	<i>(y.) Wall-flowered.</i>	—	1820. —	
20 Tournefórtii. DC.	<i>(st.) Tournefort's.</i>	Spain.	—	
21 lævigáta. DC.	<i>(wh.) smooth.</i>	—	1825. —	
22 valentina. DC.	<i>(wh.) Spanish.</i>	—	—	
23 fruticulósa. DC.	<i>(st.) frutescent.</i>	S. Europe.	1820. 4. 5. H. ♂.	
24 elongáta. DC.	<i>(ye.) stalk-leaved.</i>	Hungary.	1801. 5. 10. H. ♂.	Barr. ic. t. 195. f. 1.
25 sabulária. DC.	<i>(ye.) gravel.</i>	Portugal.	1818. —	Cyr. pl. rar. 2. t. 1.
26 odoráta. Sck.	<i>(ye.) sweet-scented.</i>	—	1826. —	W. et K. hung. 1. t. 28.
SIN'APIS. DC.		MUSTARD. Tetradymania Siliquosa.	L.	Brot. phyt. p. 97. t. 43.
1 nigra. DC.	<i>(ye.) common.</i>	Britain.	—	
β torulósa. DC.	<i>(ye.) twisted-podded.</i>	—	4. 8. H. ♂.	Eng. bot. 969.
γ túrgida. DC.	<i>(ye.) turgid-podded.</i>	—	— H. ♂.	
δ villósa. DC.	<i>(ye.) villous-podded.</i>	—	— H. ♂.	
2 geniculáta. DC.	<i>(ye.) jointed.</i>	Barbary.	1824. —	
3 lævigáta. DC.	<i>(ye.) smooth.</i>	Spain.	1769. 6. 7. H. ♂.	
4 auriculáta. DC.	<i>(ye.) eared.</i>	—	— H. ♂.	
5 integrifolia. DC.	<i>(ye.) entire-leaved.</i>	E. Indies.	1804. 7. 8. H. ♂.	W. hort. ber. t. 14.
6 júncea. DC.	<i>(ye.) fine-leaved.</i>	China.	1710. 6. 7. H. ♂.	Jacq. h. vind. t. 171.
7 chinénsis. DC.	<i>(ye.) Chinese.</i>	—	— H. ♂.	Ard. spec. 1. t. 10.
8 brassicáta. DC.	<i>(ye.) Cabbage-leav'd.</i>	—	— H. ♂.	
9 cérnua. DC.	<i>(wh.) drooping.</i>	China.	1824. 7. 9. H. ♂.	
10 pubéscens. DC.	<i>(ye.) downy.</i>	Sicily.	1789. —	Ard. spec. 1. t. 9.
11 arvénsis. DC.	<i>(ye.) Charlock.</i>	Britain.	—	Eng. bot. 1748.
β incisa.	<i>(ye.) cut-leaved.</i>	Europe.	—	Lob. ic. 198. f. 2.
12 orientális. DC.	<i>(ye.) oriental.</i>	Levant.	1778. 6. 7. H. ♂.	Schk. hand. 1. t. 186.
13 taurica. DC.	<i>(ye.) Taurian.</i>	Tauria.	1823. 4. 8. H. ♂.	
14 subbipinnatifida. DC.	<i>(ye.) subbipinnatifid.</i>	Spain.	—	
15 Káber. DC.	<i>(ye.) Persian.</i>	Persia.	—	
16 Alliòni. DC.	<i>(ye.) Allioni's.</i>	Egypt.	1789. —	Jacq. h. vind. 2. t. 168.
17 túrgida. DC.	<i>(ye.) turgid-podded.</i>	—	— H. ♂.	
18 incána. DC.	<i>(ye.) hoary-jointed.</i>	S. Europe.	1771. 7. H. ♂.	Jacq. h. vind. 2. t. 169.
19 heterophylla. DC.	<i>(ye.) various-leaved.</i>	Spain.	1820. 5. 6. H. ♂.	
20 álba. DC.	<i>(ye.) white-seeded.</i>	Britain.	—	Eng. bot. 1677.
21 hispida. DC.	<i>(ye.) hispid.</i>	Spain.	1804. 6. 7. H. ♂.	
22 dissécta. DC.	<i>(ye.) cut-leaved.</i>	—	— 4. 6. H. ♂.	
23 hastáta. DC.	<i>(ye.) hastate-leaved.</i>	N. Holland.	1824. 7. 9. H. ♂.	
24 foliða. DC.	<i>(ye.) leafy.</i>	Levant.	1804. 5. 6. H. ♂.	
25 ápula. DC.	<i>(ye.) Apulian.</i>	Italy.	1824. 6. 8. H. ♂.	
26 ramósä. H.B.	<i>(ye.) branching.</i>	E. Indies.	1804. 7. 9. H. ♂.	
27 virgáta. H.	<i>(ye.) twiggy.</i>	—	1827. 6. 8. H. ♂.	
28 subulária.	<i>(ye.) awl-shaped.</i>	—	— H. ♂.	
29 frutéscens. DC.	<i>(st.) frutescent.</i>	Madeira.	1777. 12. 6. G. ♂.	
MORICA'NDIA. DC.	MORICA'NDIA.	Tetradymania Siliquosa. L.		
arvénsis. DC.	<i>(pu.) field.</i>	S. Europe.	1739. 6. H. ♀.	Swt. br. fl. gar. 278.
<i>Brássica arvénsis. L.</i>	<i>Túrritis arvénsis. H.K.</i>			
DIPLOTA'XIS. DC.	DIPLOTA'XIS.	Tetradymania Siliquosa. L.		
1 crassifolia. DC.	<i>(ye.) thick-leaved.</i>	Sicily.	1824. 5. 8. H. ♀.	

2 erucoides. DC.	(ye.) dwarf.	S. Europe.	1736.	6. 7. H. ⊙.	Jacq. vind. 2. t. 170.
	<i>Sinapis erucoides</i> . L.				
3 virgata. DC.	(ye.) slender.	Spain.	1820.	— H. ⊙.	
4 catholica. DC.	(ye.) Spanish.	—	—	4. 5. H. ⊙.	
	<i>Sisymbrium catholicum</i> . L.				
5 tenuifolia. DC.	(ye.) Wall rocket.	England.	7. 10. H. ⊙.	Eng. bot. 525.
	<i>Sisymbrium tenuifolium</i> . E.B. <i>Sinapis tenuifolia</i> . H.K.				
6 muralis. DC.	(ye.) sand.	England.	7. 9. H. ⊙.	Eng. bot. 1090.
	<i>Sisymbrium murale</i> . L. <i>Sinapis muralis</i> . H.K.				
7 viminea. DC.	(ye.) twiggy.	S. Europe.	5. 6. H. ⊙.	Mor. his. ox. 2. t. 5.f. 8.
8 saxatilis. DC.	(ye.) rock.	—	—	— H. ¼.	
9 ramosissima. DC.	(ye.) branching.	—	1822.	— H. ¼.	
ER'UCA. DC.	ER'UCA. Tetradymania Siliquosa. L.				
1 sativa. DC.	(wh.) stripe-flowered.	S. Europe.	1573.	7. H. ⊙.	Fl. græc. t. 646. et 647.
	<i>Brassica Erüca</i> . L. <i>Euzònum sativum</i> . L.en.				
2 hispida. DC.	(wh.) hispid.	Naples.	1824.	— H. ⊙.	
3 vesicaria. DC.	(wh.) bladdered.	Spain.	1820.	— H. ⊙.	Asso. syn. arrag. 88. t. 4.
	β <i>flaviflora</i> . DC. (ye.) yellow.	—	—	— H. ⊙.	

Tribus XIII. *VELLEÆ* seu Orthoploceæ Latiseptæ. DC.

VELLA. DC.	CRESS-ROCKET. Tetradymania Siliculosa. L.				
pseudocystis. DC. (y.)	shrubby.	Spain.	1759.	4. 5. H. ½.	Bot. reg. 293.
CARRICHT'ERA. DC.	CARRICHT'ERA. Tetradymania Siliculosa. L.				
Vellæ. DC.	(st.) annual.	England.	6. 7. H. ⊙.	Eng. bot. 1441.
	<i>Vellæ annua</i> . E.B.				
SUCCO'WIA. DC.	SUCCO'WIA. Tetradymania Siliculosa. L.				
baleàrica. DC.	(ye.) Minorca.	Minorca.	1781.	6. 7. H. ⊙.	Jacq. vind. 2. t. 144.
	<i>Bünias baleàrica</i> . L.				

Tribus XIV. *PSYCHINEÆ* seu Orthoploceæ Angustiseptæ. DC.

PSYCHINE. DC.	PSYCH'INE. Tetradymania Siliculosa. L.				
stylösa. DC.	(wh.) striped-flower'd. Barbary.	1822.	4. 5. H. ⊙.	Desf. fl. atl. 2. t. 148.	

Tribus XV. *ZILLEÆ* seu Orthoploceæ Nucamentaceæ. DC.

ZILLA. DC.	ZILLA. Tetradymania Siliculosa. L.				
myagroïdes. DC. (li.) spiny.	Egypt.	4. 5. H. ½.	Vent. malm. t. 16.	
	<i>Bünias spinosa</i> . Vent.				
CALEP'INA. DC.	CALEP'INA. Tetradymania Siliculosa. L.				
Corvini. DC.	(wh.) annual.	S. Europe,	1820.	4. 6. H. ⊙.	W. et Kit. hung. 2.t.107.
	<i>Bünias cochlearioides</i> . w.				

Tribus XVI. *RAPHANEÆ* seu Orthoploceæ Lomentaceæ. DC.

CRA'MBE. DC.	COLEWORT. Tetradymania Siliculosa. L.				
1 maritima. DC.	(wh.) Sea Kale.	Britain.	5. 6. H. ¼.	Eng. bot. 924.
2 pinnatifida. DC.	(wh.) smooth-winged.	Siberia.	1759.	6. 7. H. ¼.	Jacq. ic. 1. t. 128.
3 orientalis. DC.	(wh.) oriental.	Levant.	1752.	6. 7. H. ¼.	
4 tatárica. DC.	(wh.) Tartarian.	Hungary.	1789.	— H. ¼.	Jacq. ic. 1. t. 129.
5 áspera. DC.	(wh.) rough.	Tauria.	1823.	5. 6. H. ¼.	Deles. ic. sel. 2. t. 91.
6 júncea. DC.	(wh.) rushy.	Iberia.	1828.	— H. ¼.	
7 cordifolia. DC.	(wh.) heart-leaved.	Caucasus.	—	6. 7. H. ¼.	
8 hispánica. DC.	(wh.) Spanish.	Spain.	1683.	— H. ⊙.	Jacq. obs. 2. t. 41.
9 filiformis. DC.	(wh.) Patagonian.	Patagonia.	1796.	7. 8. H. ⊙.	Jacq. ic. 3. t. 504.
10 fruticosa. DC.	(wh.) Madeira.	Madeira.	1777.	5. 11. G. ½.	
11 strigosa. DC.	(wh.) strigose shrubb.	Canaries.	1779.	5. 6. G. ½.	L'. her. stirp. 1. t. 72.

RAPISTRUM. DC.	RAPISTRUM.	Tetradynamia Siliculosa. L.				
1 perénne. DC.	(ye.) perennial.	Europe.	1789.	7. H. ȝ.	Jacq. aust. 5. t. 414.	
<i>Myagrum perenne.</i> L.	<i>Cakile perennis.</i> H.K.					
2 rugosum. DC.	(ye.) wrinkled.		1739.	6. 7. H. ⊖.	All. ped. 1. t. 78.	
<i>Myagrum rugosum.</i> L.	<i>Cakile rugosa.</i> H.K.					
3 orientálē. DC.	(ye.) oriental.	Levant.	1795.	— H. ⊖.	Flor. græc. t. 612.	
ENARTHROCARPUS.	ENARTHROCARPUS.	Tetradynamia Siliquosa.				
1 lyrátus. DC.	(ye.) yellow-strip'd.	Egypt.	1826.	6. 8. H. ⊖.	Delil. fl. &g. 105. t. 36. f. 1.	
2 pterocárpus. DC. wing-podded.		1828. H. ⊖.	Deles. ic. sel. 2. t. 93.	
RAPHANUS. DC.	RADISH.	Tetradynamia Siliquosa. L.				
1 satívus. Mil.	(li.) common.	China.	1548.	5. 6. H. ⊖.	Fuchs. hist. 659. ic.	
α pr'æcox.	(li.) early Salmon.			— H. ⊖.		
β macrorhízus.	(li.) long Salmon.			— H. ⊖.		
γ purpùreus.	(li.) purple-rooted.			— H. ⊖.		
2 rotundus. Mil.	(li.) Turnep-rooted.			— H. ⊖.	Lob. ic. t. 201. f. 1.	
3 chinénsis. Mil.	(li.) oil-seeded.			— H. ⊖.		
4 niger. DC.	(li.) black Spanish.			— H. ⊖.	Lob. ic. 202. f. 1.	
α vulgàris. DC.	(li.) long-rooted.			— H. ⊖.	Wein. phyt. t. 860. f. c.	
β rotundus. DC.	(li.) round-rooted.			— H. ⊖.	— t. 860. f. b.	
γ griseus. DC.	(li.) grey.			— H. ⊖.		
5 orbicularis. Mil.	(li.) white Spanish.			— H. ⊖.		
6 caudátus. DC.	(pu.) long-podded.	Java.	1815.	5. 8. H. ⊖.	Lin. f. dec. 1. t. 10.	
7 Raphanístrum. DC.	(ea.) wild.	Britain.	4. 8. H. ⊖.	Eng. bot. 856.	
α álbum.	(wh.) white-flowered.		— H. ⊖.	Wein. phyt. t. 862. f. a.	
β purpuráscaens.	(pu.) purple-flowered.		— H. ⊖.	— t. 862. f. b.	
ζ flávum.	(ye.) yellow-flowered.		— H. ⊖.	J. Bauh. h. 2. p. 844. ic.	
8 Landra. DC.	(ye.) Italian.	Italy.	1827.	4. 6. H. ♂.	Deles. ic. sel. 2. t. 94.	
9 marítimus. DC.	(ye.) sea-side.	Britain.	5. 7. H. ♂.	Eng. bot. 1643.	

Subordo IV. SPIROLOBEÆ. DC. syst. 2. p. 670.

Tribus XVII. *BUNIADEÆ* seu Spirolobæ Nucamentaceæ. DC.

B'UNIAS. DC.	B'UNIAS.	Tetradynamia Siliculosa. L.			
1 Erucágo. DC.	(ye.) prickly-podded.	S. Europe.	1640.	6. 7. H. ⊖.	Jacq. aust. 4. t. 340.
2 áspéra. DC.	(ye.) rough.	Portugal.	1823.	6. 7. H. ⊖.	
3 orientális. DC.	(ye.) oriental.	Levant.	1795.	5. 7. H. ȝ.	Schk. han. 2. t. 189.

Tribus XVIII. *ERUCARIEÆ* seu Spirolobæ Lomentaceæ. DC.

ERUCARIA. DC.	ERUCARIA.	Tetradynamia Siliquosa. L.			
1 aléppica. DC.	(pu.) Aleppo.	Levant.	1680.	7. 8. H. ⊖.	Vent. h. cels. t. 64.
<i>Cordylocárpus lavigátus.</i> w.					
2 crassiñólia. DC.	(pu.) thick-leaved.	Egypt.	1828.	7. 9. H. ⊖.	Delil. pl. bot. t. 34. f. 1.

Subordo V. DIPLECOLOBEÆ. DC. syst. 2. p. 676.

Tribus XIX. *HELIOPHILEÆ* seu Diplecolobeæ Siliquosæ. L.

HELIOPHILA. DC.	HELIOPHILA.	Tetradynamia Siliquosa. L.			
1 filiformis. DC.	(pu.) awl-podded.	C. B. S.	1786.	7. 9. H. ⊖.	Lam. ill. t. 563. f. 3. ?.
2 dissécta. DC.	(bl.) cut-leaved.		1822.	6. 8. H. ⊖.	
3 tenuisiliqua. DC.	(bh.) slender-podded.			— H. ⊖.	Deles. ic. sel. 2. t. 96.
4 amplexicáulis. DC.	(bh.) stem-clasping.		1774.	6. 9. H. ⊖.	Jacq. frag. t. 64. f. 2.
5 rivális. DC.	(wh.) rivulet,		1819.	— H. ⊖.	
6 péndula. DC.	(wh.) pendulous		1792.	— H. ⊖.	Lam. ill. t. 562. f. 2.
<i>pinnata.</i> Vent. malm. t. 113. non. L.					

		C: B. S.	1819.	6. 9. H. ⊙.	
7	trifida. DC. (pu.) trifid-leaved. pinnatæ. L. non Vent.	_____	1824.	— H. ⊙.	Pluk. mant. t. 432. f. 2.
8	pusilla. DC. (wh.) small.	_____	—	H. ⊙.	
9	lepidioïdes. L.en (wh.) Cress-like.	_____	—	H. ⊙.	
10	diffusa. DC. (wh.) spreading.	_____	1820.	— H. ⊙.	
11	Peltaria. DC. (wh.) small-white.	_____	—	H. ⊙.	
12	pilosa. DC. (bl.) hairy. α integrifolia.DC.(bl.) entire-leaved. β incisa. DC. (bl.) cut-leaved. arabioides. B.M.	_____	1768.	— H. ⊙.	Jacq. ic. 3. t. 506. Bot. mag. 496.
13	stricta. B.M. (bl.) upright.	_____	1824.	— H. ⊙.	Bot. mag. 2526.
14	digitata. DC. (bl.) fingered.	_____	1819.	— H. ⊙.	Bot. reg. 838.
15	coronopifolia. DC. (bl.) Buck's-horn-ld.	_____	1778.	6. 7. H. ⊙.	Herm. lugd. t. 367.
16	pectinata. DC. (wh.) pectinated.	_____	1819.	— H. ⊙.	
17	fœniculacea. DC. (pu.) Fennel-leaved.	_____	1774.	6. 9. H. ⊙.	
18	crithmifolia. DC. (bh.) Samphire-ld.	_____	1816.	— H. ⊙.	
19	platysiliqua. DC. (bh.) flat-podded.	_____	1774.	7. 8. G. ḥ.	
20	linearifolia. DC. (bl.) linear-leaved.	_____	1819.	— G. ḥ.	
21	scoparia. DC. (bh.) Broom-like.	_____	1802.	— G. ḥ.	Deles. ic. sel. 2. t. 98.
22	incana. DC. (pu.) hoary.	_____	1774.	5. 8. G. ḥ.	
23	cleomoides. DC. (wh.) Cleome-like. Cheiranthus strictus. W.	_____	1802.	1. 12. G. ḥ.	Deles. ic. sel. 2. t. 99.

Tribus XX. SUBULARIEÆ seu Diplocolobeæ Latiseptæ. DC.

SUBUL'ARIA. DC. AWLWORT. Tetradynamia Siliculosa. L.
aquatica. DC. (wh.) water. Britain. 7. H.w. ⊙. Eng. bot. 732.

Subordo VI. SCHIZOPETALEÆ.

SCHIZOPETALUM. B.M. SCHIZOPETALUM. Tetradynamia Siliquosa. L.
Walkeri. B.M. (wh.) Walker's. Chili. 1822. 7. 9. H. ⊙. Brown bot. reg. 752.

ORDO XIV.

CAPPARIDEÆ. DC. prodr. 1. p. 237.

Tribus I. CLEOMEÆ. DC.

PERITOMA. DC	PERITOMA. Monadelphia Hexandria.				
serrulatum. DC. (pu.) saw-leaved.	Missouri.	1823.	6. 8. H. ⊙.		
Cleome serrulata. Ph.					
GYNANDRO'PSIS. DC. GYNANDRO'PSIS.	Monadelphia Hexandria.				
1 sessilifolia. DC. (wh.) sessile-leaved.	W. Indies.	1820.	6. 8. H. ⊙.		
2 triphylla. DC. (wh.) three-leaved.	—	—	— H. ⊙.		Herm. lugd. 565. ic
Cleome triphylla. L.?					
3 pentaphylla. DC.(wh.) five-leaved.	India.	1640.	— H. ⊙.		Jacq. vind. 1. t. 21.
Cleome pentaphylla. B.M. 1681.					
4 candelabrum. (wh.) chandelier.	W. Indies.	1824.	7. 9. H. ⊙.		Bot. mag. 2656.
Cleome candelabrum. B.M.					
5 mnicata. Scr. (wh.) bristly.	1828.	— S. ⊙.		
6 heterotricha. DC.(wh.) various-haired.	C. B. S.	1822.	— H. ⊙.		
CLEOME. DC.	CLEOME. Tetradynamia Siliquosa. L.				
1 gigantea. DC. (gr.) gigantic.	S. America.	1774.	6. 8. S. ḥ.		Jacq. obs. 4. t. 76.
2 heptaphylla. DC. (wh.) seven-leaved.	W. Indies.	1820.	— H. ⊙.		
3 spinosa. DC. (wh.) white-flowered.	—	1731.	5. 10. S. ḥ.		Marc. bras. t. 34.
4 pungens. DC. (bh.) pink-flowered.	S. America.	1812.	7. 8. S. ⊙.		W. hort. ber. 1. t. 18.
spinosa. B.M. 1640. nec aliorum.					
5 pubescens. B.M. (wh.) pubescent.	1815.	7. 8. H. ⊙.		Bot. mag. 1857.

CAPPARIDÆ.

6 ròsea. DC.	(ro.) rose-coloured.	Brazil.	1825.	5. 8.	S. ⊖.	Bot. reg. 960.
7 monophylla. DC.	(ye.) simple-leaved.	E. Indies.	1759.	—	H. ⊖.	Rheed. mal. 9. t. 34.
β zeylânica. DC.	(ye.) Ceylon.			—	H. ⊖.	Burm. zeyl. t. 100. f. 2.
8 procumbens. DC.	(wh.) procumbent.	W. Indies.	1798.	—	S. Ȑ.	Jacq. amer. t. 120.
9 violacea. DC.	(vi.) violet-coloured.	Portugal.	1776.	6. 7.	H. ⊖.	Schk. han. 2. t. 189. f. 1.
10 Dilleniâna. DC.	(ye.) Dillenius's.	Levant.	1732.	—	H. ⊖.	Dill. elt. t. 266. f. 345.
	ornithopodioides. W.					
11 ibérica. DC.	(ye.) Iberian.	Iberia.	1820.	—	H. ⊖.	Buxb. cent. 1. t. 9. f. 2.
12 virgâta. DC.	(ye.) twiggy.	Persia.	—	H. ⊖.	Lin. f. dec. t. 8.
13 arâbica. DC.	(ye.) Arabian.	Arabia.	1794.	—	H. ⊖.	
14 flava. DC.	(ye.) yellow.	N. Holland.	1825.	7. 10.	H. ⊖.	
15 polygamia. DC.	(wh.) polygamious.	W. Indies.	1822.	6. 7.	H. ⊖.	Slo. jam. t. 124. f. 1.
16 aculeata. DC.	(wh.) prickly.	S. America.	—	—	H. ⊖.	Mart. dec. t. 45.
17 Houstonia. DC.	(fl.) Houston's.	W. Indies.	1730.	—	S. ⊖.	
18 diffusa. DC.	(wh.) spreading.	Brazil.	1823.	—	H. ⊖.	
19 micrantha.	(wh.) small-flowered.	1824.	6. 10.	S. Ȑ.	
POLANISIA. DC.	POLANISIA.	Dodecadria Monogynia. L.				
1 Chelidônii. DC.	(ro.) Celandine-fld.	E. Indies.	1790.	7. 10.	S. ⊖.	
	Cleome Chelidônii. L.					
2 graveolens. DC.	(pu.) strong-scented.	N. America.	—	H. ⊖.	Corn. can. 131. ic.
3 viscosa. DC.	(ye.) viscid.	Ceylon.	1730.	—	S. ⊖.	Rheed. mal. 9. t. 23.
4 dodecandra. DC.	(pu.) three-leaved.	—	1795.	—	S. ⊖.	Burm. zeyl. t. 100. f. 1.
	Cleome dodecandra. L.					
5 uniglandulosa. DC.	(w.) one-glandled.	S. America.	1823.	—	H. ⊖.	Cav. ic. 4. t. 306.

Tribus II. CAPPAREÆ. DC.

CRATÆVA. DC.	CRATÆVA.	Dodecadria Monogynia. L.				
1 gynândra. DC.	(wh.) thin-leaved.	Jamaica.	1789.	S. Ȑ.	Pluk. alm. t. 147. f. 6.
2 Tâpia. DC.	(wh.) Garlick Pear.	S. America.	1752.	S. Ȑ.	Plum. gen. t. 21.
3 Roxburghii. B.	(wh.) Roxburgh's.	E. Indies.	1822.	S. Ȑ.	
	Câpparis trifoliata. H.B.					
RITCHIEA. B.	RITCHIEA.	Dodecadria Monogynia.				
	frâgrans.	(wh.) fragrant.	Sierra Leon.	1795.	6. 7.	S. Ȑ. — Bot. mag. 596.
	Cratæva fragans. B.M.	capparoides. A.R.	t. 176.			
NIEBUHRIA. DC.	NIEBUHRIA.	Polyandria Monogynia. L.				
1 cáfra. DC.	(wh.) Cafrarian.	C. B. S.	1820.	G. Ȑ.	
2 madagascariensis. DC.	(w.) Madagascar.	Madagascar.	1822.	S. Ȑ.	
3 oblongifolia. DC.	(wh.) oblong-leaved.	E. Indies.	1823.	S. Ȑ. —	
	Câpparis heterochita. H.B.					
BO'SCIA. DC.	BO'SCIA.	Dodecadria Monogynia. L.				
senegalensis. DC.	(wh.) Senegal.	Senegal.	1822.	S. Ȑ.	Lam. ill. t. 395. f. 2.
CA'PPARIS. DC.	CAPER-TREE.	Polyandria Monogynia. L.				
1 mariâna. DC.	(wh.) heart-leaved.	Timor.	1822.	S. Ȑ.	Jacq. Schoenb. t. 109.
2 rupéstris. DC.	(wh.) rock.	Crete.	7. 10.	F. Ȑ. —	Flor. græc. t. 487.
	spinosa. B.M.	291. nec aliorum.				
3 spinosa. DC.	(wh.) common.	S. Europe.	1596.	5. 8.	G. Ȑ. —	Flor. græc. t. 486.
4 ovata. W.	(wh.) oval-leaved.	Mauritania.	—	G. Ȑ. —	Boc. sic. t. 42. f. 3.
	Fontanætii. DC.					
5 herbacea. DC.	(wh.) herbaceous.	Tauria.	1819.	—	H. Ȑ.	
6 aegyptia. DC.	(wh.) Egyptian.	Egypt.	1822.	G. Ȑ.	Delil. fl. eg. t. 31. f. 3.
7 aphylla. DC.	(wh.) leafless.	E. Indies.	—	S. Ȑ.	
8 zeylânica. DC.	(wh.) Ceylon.	Ceylon.	—	S. Ȑ. —	
9 acuminata. DC.	(wh.) acuminate.	E. Indies.	1821.	S. Ȑ. —	Braam ic. chin. t. 29. ?
10 frondosa. DC.	(g.p.) large-leaved.	Carthagena.	1800.	S. Ȑ.	Jacq. amer. t. 104.
11 triflora. DC.	(wh.) three-flowered.	S. America.	S. Ȑ.	Jacq. amer. t. 98.
12 cynophallóphora. DC.	(wh.) Bay-leavd.	W. Indies.	1752.	S. Ȑ.	ed. pict. t. 146.
13 eustachiana. DC.	(pu.) purple-flower'd.	—	1822.	S. Ȑ.	— t. 99.
14 saligna. DC. Willow-leaved.	S. Cruz.	1807.	S. Ȑ.	Jacq. amer. t. 102.
15 linearis. DC.	(wh.) linear-leaved.	W. Indies.	1793.	S. Ȑ.	— t. 99.
16 verrucosa. DC.	(wh.) wart-podded.	Carthagena.	1823.	S. Ȑ.	Jacq. amer. t. 106.
17 pulcherrima. DC.	(st.) handsome.	—	S. Ȑ.	
	arboréscens. Mil.					
18 amygdalina. DC.	(wh.) Almond-leaved.	S. America.	1819.	G. Ȑ.	Plum. gen. 40. t. 16.

19 odoratissima. DC. (<i>pu.</i>) sweet-scented.	Caracas.	1814.	S. $\frac{1}{2}$.	Jacq. schœn. t. 110.
20 tenuisiliqua. DC. (<i>wh.</i>) slender-podded.	Carthagena.	1822.	S. $\frac{1}{2}$.	Jacq. amer. t. 105.
21 ferruginea. DC. (<i>wh.</i>) ferruginous.	Jamaica.	S. $\frac{1}{2}$.	— t. 100.
<i>octandra</i> . Jacq.				
22 Bréymia. DC. (<i>wh.</i>) Oleaster-leav'd.	W. Indies.	1752.	G. $\frac{1}{2}$.	Jacq. amer. t. 103.?
23 jamaicensis. DC. (<i>wh.</i>) Jamaica.	—	1793.	S. $\frac{1}{2}$.	— t. 101.
24 torulosa. DC. (<i>wh.</i>) twisting.	—	1819.	S. $\frac{1}{2}$.	
STEPHA'NIA. DC. STEPHANIA. Hexandria Monogynia.				
cleomoides. DC. (<i>o.y.</i>) Cleome-like.	Caracas.	1823.	S. $\frac{1}{2}$.	Jacq. schœn. t. 111.
Cápparis paradóxa. Jacq. <i>Steriphoma cleomoides</i> , s.s.				
MORIS'ONIA. DC. MORISONIA. Monadelphia Polyandria. L.				
americana. DC. (<i>wh.</i>) American.	Caribees.	1824.	S. $\frac{1}{2}$.	Jacq. amer. t. 97.

ORDO XV.

RESEDACEÆ. DC. theor. ed. 1. p. 214.

RE'SEDA. W.	RE'SEDA. Dodecandria Trigynia. L.			
1 Lutèola. w.	(<i>st.</i>) Dyers weed. Britain. 6. 7.	H. \odot .	Eng. bot. 320.
2 críspata. L. en.	(<i>st.</i>) curled-leaved. S. Europe.	1820. —	H. \odot .	
3 viréscens. s.s.	(<i>st.</i>) greenish. Spain.	—	H. \odot .	
4 canescens. s.s.	(<i>wh.</i>) hoary. Egypt.	1597. 5. 7.	H. $\frac{1}{2}$.	Clus. hist. 1. t. 295.
5 gláuca. s.s.	(<i>wh.</i>) glaucous. S. Europe.	1700. —	F. $\frac{1}{2}$.	Pluk. alm. t. 107. f. 2.
6 scoparia. s.s.	(<i>wh.</i>) Broom-like. Teneriffe.	1815. 8. 9.	G. $\frac{1}{2}$.	
7 dipétala. s.s.	(<i>wh.</i>) two-petaled. C. B. S.	1774. 8.	G. δ .	
8 limifolia. s.s.	(<i>wh.</i>) flax-leaved. S. Europe.	1820. 7. 8.	F. δ .	
9 subulata. Del.	(<i>wh.</i>) awl-leaved. Egypt.	1827. 6. 8.	H. δ .	
10 chinensis. s.s.	(<i>wh.</i>) Chinese. China.	1824. —	H. δ .	
11 sesamoides. s.s.	(<i>wh.</i>) spear-leaved. France.	1787. —	H. \odot .	All. ped. 2. t. 88. f. 3.
12 mediterránea. s.s. (<i>w.</i>)	Mediterranean. Palestine.	1791. 6. 9.	H. \odot .	Jacq. ic. 3. t. 475.
13 odorata. s.s.	(<i>st.</i>) Mignonette. Egypt.	1752. 6. 10.	H. \odot .	Bot. mag. 29.
β frutescens. B.R. (<i>st.</i>) tree mignonette.	—	G. $\frac{1}{2}$.	Bot. reg. 227.
14 Phyteuma. s.s. (<i>wh.</i>) trifid.	S. Europe.	—	6. 9. H. \odot .	Jacq. aust. 2. t. 132.
15 incisa. T.C.	(<i>wh.</i>) cut-leaved. Italy.	1825. —	H. \odot .	
16 ramosissima. s.s. (<i>wh.</i>) branching.	Spain.	1816. —	H. $\frac{1}{2}$.	
17 saxatilis. s.s. (<i>wh.</i>) rock.	—	—	H. $\frac{1}{2}$.	
<i>stricta</i> . P.S.	—	—	H. $\frac{1}{2}$.	
18 pruinosa. s.s. (<i>wh.</i>) papillose-ld.	Egypt.	1823. —	H. $\frac{1}{2}$.	
19 lútea. s.s. (<i>st.</i>) Base-rocket.	Britain. 7. 8.	H. δ .	Eng. bot. 321.
20 undata. s.s. (<i>wh.</i>) wave-leaved.	Spain.	1739. 6. 8.	H. $\frac{1}{2}$.	Barrel. rar. t. 587.
21 álba. s.s. (<i>wh.</i>) upright-white.	S. Europe.	1596. 5. 10.	H. δ .	Lobel. ic. 222.
22 fruticulosa. s.s. (<i>wh.</i>) frutescent.	Spain.	1794. 9.	G. $\frac{1}{2}$.	Jacq. ic. 3. t. 474.
23 bipinnata. s.s. (<i>wh.</i>) bipinnate.	—	1816. 6. 8.	G. $\frac{1}{2}$.	
24 myriophylla. T.C. (<i>st.</i>) numerous-ld.	Italy.	1825. —	H. $\frac{1}{2}$.	

ORDO XVI.

DATISCEÆ. Brown in app. to Denham and Clap. trav. p. 25.

DATI'SCA. L.	DATI'SCA. Diœcia Dodecandria. L.		
1 cannabina. L.	(<i>ye.</i>) Hemp-like.	Levant.	1640. 7. 9. H. $\frac{1}{2}$.
2 hírta. w.	(<i>ye.</i>) hairy.	Pensylvania.	1826. 9. 10. H. $\frac{1}{2}$.
			Schkuhr han. 3. t. 336.

ORDO XVII.

FLACOURTIANEÆ. DC. prodr. 1. p. 255.

Tribus I. *PATRISIEÆ*. DC.

RYAN'EA. DC. **RYAN'EA.** Polyandria Monogynia.
 speciosa. DC. (re) beautiful. Trinidad. 1822. S. h. Vahl. ecl. 1. p. 51. t. 9.
Patrisia pyrifera. P.S.

Tribus II. *FLACOURTIEÆ*. DC.

FLACOURTIA.	DC.	FLACOURTIA.	Dicoccia Polyandria.	L.
1 Ramóntchi.	DC. (<i>wh.</i>)	shining-leaved.	Madagascar.	1775. 6. 7. S. $\frac{1}{2}$.
2 sápida.	DC. (<i>wh.</i>)	esculent.	E. Indies.	1800. S. $\frac{1}{2}$.
3 inérmis.	DC. (<i>wh.</i>)	unarmed.	Moluccas.	1814. 10. 2. S. $\frac{1}{2}$.
4 sepiátria.	DC. (<i>wh.</i>)	obovate-leaved.	Coromandel.	1820. S. $\frac{1}{2}$.
5 cataphrácta.	DC. (<i>wh.</i>)	many-spined.	E. Indies.	1804. S. $\frac{1}{2}$.
6 rotundifólia.	H.B. (<i>wh.</i>)	round-leaved.	_____	1823. S. $\frac{1}{2}$.
7 flavescens.	DC. (<i>st.</i>)	yellow-flower'd.	Guinea.	1780. S. $\frac{1}{2}$.
8 rhamnóides.	DC. (<i>wh.</i>)	Rhamnus-like.	C. B. S.	1819. G. $\frac{1}{2}$.

Tribus III. KIGGELARIEÆ. DC.

KIGGEL'ARIA. DC.	KIGGEL'ARIA.	Dicæcia Decandria. L.				
1 africana. DC.	(wh.) African.	C. B. S.	1683.	5. 6.	G. h.	Lam. ill. t. 821.
2 integrifolia. DC.	(wh.) entire-leaved.		1819.	—	G. h.	Jacq. ic. rar. t. 628.
MELI'CYTUS. DC.	MELI'CYTUS.	Dicæcia Pentandria.				
ramiflorus. DC. branch-flow'rd.	N. Zealand.	1822.	G. h.	Lam. ill. t. 812. f. 1.

ORDO XVIII.

BIXINEÆ. *DC. prodr.* 1. p. 259.

BI'XA. DC.	ANOTTA.	Polyandria Monogynia. L.				
1 purpurea. (pu.)	purple.	E. Indies.	1820.	S. h.	
2 Orellana. DC.	(ro.) heart-leaved.	S. America.	1690.	5. 8.	S. h.	Bot. mag. 1456.
3 Urucurána. w.	(ro.) white-leaved.	Brazil.	1823.	S. h.	
LE'TIA. DC.	LE'TIA.	Polyandria Monogynia. L.				
Thámnia. DC.	shining-leaved.	Jamaica	S. h.
PRO'CRIA. DC.	PRO'CRIA.	Polyandria Monogynia. L.				Browne jam. t. 25. f. 2.
1 Crúcis. DC.	(gr.) oval-leaved.	Santa Cruz.	S. h.	Vahl symb. 3. t. 64.
2 serrata. DC.	(gr.) saw-leaved.	Montserrat.	1822.	S. h.	
3 theafórmis. DC.	(gr.) Tea-leaved.	Bourbon.	—	S. h.	Bory voy. 2. t. 24.
<i>Lúdia heterophylla.</i> Bory. non. Lam.						
LUDIA. DC.	LUDIA.	Polyandria Monogynia. L.				
1 heterophylla. DC. (gr.)	various-leaved.	Mauritius.	1822.	S. h.	Lam. ill. t. 466. f. 1. 2.
2 sessiliflora. DC.	(bl.) sessile-flower'd.	—	—	S. h.	Jacq. schœnb. 1. t. 112.
<i>tuberulata.</i> Jacq.						

ORDO XIX.

CISTINEÆ. DC. prodr. 1, p. 263.

CISTUS. DC. ROCK-ROSE. *Polyandria Monogynia*. L.
 1 *candidissimus*. DC. (*ro.*) Canary Island. Canaries. 1817. 6. 9. F. $\frac{1}{2}$. Swt. Cist. t. 3.
 2 *vaginatus*. DC. (*ro.*) oblong-leaved. Teneriffe. 1779. 4. 8. F. $\frac{1}{2}$. — t. 9.

CISTINEÆ.

3	álbidus. DC.	(ro.) white-leaved.	S. Europe.	1640.	5. 9.	H. Ȣ.	Swt. cist. t. 31.
4	incánus. DC.	(li.) hoary-leaved.	1596.	5. 8.	F. Ȣ.	t. 44.	
5	canéscens. s.c.	(pu.) narrow hoary.	F. Ȣ.	t. 45.	
6	villósus. DC.	(pu.) villous.	1640.	F. Ȣ.	t. 35.	
7	rotundifólius. s.c.	(pu.) round-leaved.	F. Ȣ.	t. 75.	
8	undulátus. DC.	(pu.) wave-leaved.	H. Ȣ.	t. 63.	
9	heterophyllus. DC.	(p.) various-leaved.	Algiers.	F. Ȣ.	t. 6.	
10	crispus. DC.	(pu.) curled-leaved.	S. Europe.	1656.	F. Ȣ.	t. 22.
11	créticus. DC.	(pu.) Cretan.	Levant.	1731.	F. Ȣ.	Flor. græc. t. 495.
12	purpureus. DC.	(pu.) purple.	6. 8.	F. Ȣ.	Swt. cist. t. 17.	
13	cymósus. DC.	(li.) cyme-flowered.	F. Ȣ.	t. 90.	
	incánus. Flor. græc. t. 494.	nec aliorum.					
14	parviflòrus. DC.	(li.) small-flowered.	1821.	F. Ȣ.	Swt. cist. t. 14.
15	Cupaniánum. s.s. (wh.)	heart-leaved.	Sicily.	6. 10.	H. Ȣ.	t. 70.
16	asperifólius. s.c. (wh.)	rough-leaved.	H. Ȣ.	t. 87.	
17	acutifólius. s.c. (wh.)	acute-leaved.	S. Europe.	5. 8.	H. Ȣ.	t. 78.
18	salvífolius. DC. (wh.)	Sage leaved.	Europe.	1548.	5. 9.	H. Ȣ.	t. 54.
19	obtusifólius. s.c. (wh.)	blunt-leaved.	Levant.	F. Ȣ.	t. 42.
20	oblongifólius. s.c. (wh.)	oblong-leaved.	6. 10.	H. Ȣ.	t. 67.
21	corbariánsis. DC. (wh.)	mountain.	S. France.	5. 6.	H. Ȣ.	t. 8.
22	platysépalus. s.c. (wh.)	broad-sepaled.	Levant.	6. 8.	F. Ȣ.	t. 47.
23	florentinus. DC. (wh.)	Florentine.	Italy.	6. 7.	H. Ȣ.	t. 59.
24	monspliénsis. DC. (w.)	Montpelier.	S. Europe.	1656.	F. Ȣ.	t. 27.
25	Lèdon. DC.	(wh.) many-fl. gum.	S. France.	1730.	6. 8.	H. Ȣ.	Duham. arb. 1. t. 66.
26	hirsútus. DC.	(wh.) hairy.	Spain.	1656.	H. Ȣ.	Swt. cist. t. 19.
27	psilosépalus. s.c. (wh.)	smooth-calyxed.	H. Ȣ.	t. 33.	
28	fáxus. DC.	(wh.) waved-leaved.	Spain.	1656.	H. Ȣ.	t. 12.
29	populifólius. s.c. (wh.)	Poplar-leaved.	5. 6.	H. Ȣ.	t. 23.
30	latifólius. s.c.	(wh.) broad-leaved.	Barbary.	F. Ȣ.	t. 15.
31	laurifólius. DC.	(wh.) Laurel-leaved.	S. Europe.	1731.	6. 8.	H. Ȣ.	t. 52.
32	cípricus. DC.	(wh.) common-gum.	Cyprus.	H. Ȣ.	t. 39.
	ladaníferus. Bot. mag. 112.	nec. aliorum.	C. undulatus. L. en?				
33	ladaníferus. DC.	(wh.) flat-leaved gum.	Spain.	1629.	F. Ȣ.	Swt. cist. t. 84.
α	albiflòrus. DC.	(wh.) white-flowered.	F. Ȣ.
β	maculátus. DC. (w.p.)	spot-flowered.	F. Ȣ.	t. 1.
34	Clùsii. DC.	(wh.) Clusi's.	Spain.	H. Ȣ.	t. 32.
	HELIA NTHEMUM. DC.	SUN-ROSE.	Polyandria Monogynia. L.				
1	Libanótis. DC.	(wh.) Rosemary-like.	Spain.	1752.	6.	F. Ȣ.	Barr. ic. 294.
2	umbellátum. DC.	(wh.) umbel-flowered.	S. Europe.	1731.	6. 8.	F. Ȣ.	Swt. cist. t. 5.
3	ocymóides. DC.	(ye.) Basil-like.	F. Ȣ.	t. 13.
	Cistus sampsuciðolius. Cav. ic. 1. t. 96.						
4	alyssóides. DC.	(ye.) Alyssum-like.	F. Ȣ.	Vent. choix. t. 20.
5	rugosum. DC.	(ye.) rugose.	Portugal.	F. Ȣ.	Swt. cist. t. 65.
6	scabrosum. DC.	(ye.) rough.	Italy.	1775.	F. Ȣ.	t. 81.
7	algarvénse. DC.	(ye.) Algarvian.	Portugal.	1800.	7. 8.	F. Ȣ.	t. 40.
	Cistus algarvénis. B.M. 627.						
8	formósus. DC.	(ye.) beautiful.	1780.	5. 7.	F. Ȣ.	Swt. cist. t. 50.
	Cistus formósus. B.M. 264.						
9	atriplicifólium. DC.	(ye.) Orach-leaved.	Spain.	1656.	6. 7.	F. Ȣ.	Barr. ic. t. 292.
10	microphyllum. s.c. (ye.)	small-leaved.	S. Europe.	6. 10.	F. Ȣ.	Swt. cist. t. 96.
11	lasiánthum. DC.	(ye.) downy.	Portugal.	1822.	7. 9.	F. Ȣ.	
12	involucrátum. DC.	(ye.) short-pedicled.	S. Europe.	1826.	6. 8.	F. Ȣ.	
13	cánidum. s.c.	(ye.) white-leaved.	Spain.	1822.	F. Ȣ.	Swt. cist. t. 25.
14	cheiranthóides. DC.	(ye.) Stock-like.	1800.	7. 8.	F. Ȣ.	t. 107.
15	halimifólium. s.c. (ye.)	Sea-Purslane-ld	1656.	F. Ȣ.	t. 4.
16	rosmarinifólium. DC.	(ye.) Rosemary-ld.	N. America.	1823.	H. Ȣ.	
17	glomerátum. DC.	(ye.) cluster-flower'd.	New Spain.	6. 9.	F. Ȣ.	Swt. cist. t.
18	ramulifórum. DC.	(ye.) branch-flowerg.	Carolina.	1823.	H. Ȣ.	
19	canadénse. DC.	(ye.) Canadian.	Canada.	1799.	H. Ȣ.	Swt. cist. t. 21.
20	caroliniánum. DC.	(ye.) Carolina.	Carolina.	1823.	F. Ȣ.	t. 99.
21	brasiliénse. DC.	(st.) Brazilian.	Brazil.	6. 8.	F. Ȣ.	t. 43.
22	polygalafólium. s.c. (ye.)	Milkwort-leavd.	F. Ȣ.	t. 11.
23	lignósus. s.c.	(ye.) woody-stem'm'd.	S. Europe.	1810.	F. Ȣ.	t. 46.
24	Tuberária. DC.	(ye.) Plantain-leavd.	1752.	F. Ȣ.	t. 18.
25	plantagínéum. DC.	(ye.) Plantain like.	1823.	H. ○.	
26	guttátum. DC.	(ye.) spotted-flowerd.	England.	6. 8.	H. ○.	Eng. bot. 544.
27	eriocáulon. DC.	(ye.) woolly-stalked.	Spain.	1823.	H. ○.	Swt. cist. t. 30.
28	punctátum. DC.	(ye.) punctated.	France.	1816.	H. ○.	t. 61.
29	lunulátum. DC.	(ye.) lunulate-mark'd.	Piedmont.	H. Ȣ.	All. auct. p. 30. t. 2. f. 3.

CISTINEÆ.

30	villòsum. DC.	(st.) villous.	Spain.	1823.	H. ⓧ.	
31	nilòticum. DC.	(st.) Nile.	Egypt.	H. ⓧ.	Swt. cist. t. 41.
32	ledifòlium. DC.	(st.) Ledum-leaved.	England.	H. ⓧ.	Cav. ic. t. 144.
33	intermèdium. DC.	(ye.) intermediate.	Spain.	1759.	H. ⓧ.	
	salicifòlium. Cav.					
34	denticulàtum. DC.	(ye.) tooth-leaved.	S. France.	1828.	H. ⓧ.	Swt. cist. t. 71.
35	salicifòlium. DC.	(ye.) Willow-leaved.	H. ⓧ.	
36	sangùineum. DC.	(ye.) bloody-stalked.	1823.	H. ⓧ.	
	retrofráctum. P.S.					
37	ægyptiacum. DC.	(ye.) Egyptian.	Egypt.	1764.	H. ⓧ.	Jacq. obs. 3. t. 68.
38	Lippii. DC.	(ye.) small-petaled.	1820.	F. ⓧ.	
39	ellipticum. DC.	(st.) elliptic-leaved.	Levant.	1827.	F. ⓧ.	Swt. cist. t. 108.
40	confértum. DC.	(ye.) close-flowered.	Teneriffe.	6. 7.	F. ⓧ.
41	canarière. DC.	(ye.) Canary.	Canaries.	1790.	F. ⓧ.	Jacq. ic. 1. t. 97.
42	ericoides. DC.	(ye.) Heath-leaved.	S. Europe.	6. 8.	Cav. ic. 2. t. 172.
43	Fumâna. DC.	(ye.) Heath-like.	1752.	F. ⓧ.	Swt. cist. t. 16.
44	procumbens. DC.	(ye.) procumbent.	F. ⓧ.	—— t. 68.
45	arábicum. DC.	(sc.) Arabian.	Levant.	1826.	F. ⓧ.	—— t. 97.
46	lævipes. DC.	(ye.) cluster-leaved.	1690.	G. ⓧ.	—— t. 24.
47	víride. DC.	(ye.) green-leaved.	Sicily.	1828.	F. ⓧ.	
48	juniperinum. DC.	(ye.) Juniper-leaved.	G. ⓧ.	Barr. ic. t. 443.
49	Barreliéri. DC.	(ye.) Barrelier's.	Italy.	1822.	F. ⓧ.	—— t. 416.
50	thymifòlium. DC.	(ye.) Thyme-leaved.	Spain.	1658.	F. ⓧ.	Swt. cist. t. 102.
51	glutinosum. DC.	(ye.) clammy.	1790.	F. ⓧ.	—— t. 83.
52	mólle. DC.	(ye.) soft.	F. ⓧ.	Cav. ic. 3. t. 262. f. 2.
53	origanifòlium. DC.	(ye.) Marjoram-ld.	1795.	F. ⓧ.	—— 3. t. 262. f. 1.
54	dichotomum. DC.	(ye.) forked.	F. ⓧ.	—— 3. t. 263. f. 1.
55	celándicum. DC.	(ye.) smooth-leaved.	Europe.	1816.	H. ⓧ.	Swt. cist. t. 85.
56	alpéstris. DC.	(ye.) alpine.	H. ⓧ.	—— t. 2.
57	penicillatum. DC.	(ye.) pencilled.	S. Europe.	H. ⓧ.	
58	obovatum. DC.	(ye.) obovate-leaved.	Spain.	1826.	F. ⓧ.	Barr. rar. t. 366.
59	ítalicum. DC.	(ye.) Italian.	Italy.	1779.	7. 9.	Swt. cist. t. 77.
60	vineále. DC.	(ye.) slender-trailing.	Europe.	1823.	H. ⓧ.	—— t. 56.
61	cánium. DC.	(ye.) hoary.	1772.	6. 7.	Barr. rar. t. 441.
62	marifòlium. DC.	(ye.) Marum-leaved	F. ⓧ.	Swt. cist. t. 71.
63	pulchellum. s.c.	(ye.) neat.	S. Europe.	H. ⓧ.	Cav. ic. 2. t. 142.
64	rotundifòlium. DC.	(ye.) round-leaved.	Spain.	1826.	F. ⓧ.	
	<i>Cistus nummularius. Cav. non L.</i>					
65	crassifòlium. DC.	(ye.) thick-leaved.	F. ⓧ.	Cav. ic. 2. t. 141.
66	cinéreum. DC.	(ye.) gray.	Spain.	F. ⓧ.	—— 2. t. 139.
67	squammátum. DC.	(ye.) scaly.	1815.	F. ⓧ.	Barr. ic. t. 288.
68	lavandulæfòlium. DC.	(ye.) Lavender-ld.	Lévant.	1739.	F. ⓧ.	Jacq. ic. 1. t. 96.
	β syriacum. DC.	(ye.) Syrian.	Syria.	F. ⓧ.	
69	stoéchadifòlium. DC.	(ye.) woolly-leaved.	Portugal.	1819.	F. ⓧ.	Swt. cist. t. 53.
70	croceum. DC.	(ye.) Saffron-coloured.	Spain.	F. ⓧ.	
71	nudicáule. DC.	(ye.) naked-stalked.	1826.	F. ⓧ.	Cav. ic. 3. t. 261.
72	gláucum. DC.	(ye.) glaucous-leaved.	1815.	6. 8.	Swt. cist. t. 20.
73	leptophyllum. DC.	(ye.) narrow-leaved.	7. 8.	—— t. 60.
74	serpyllifòlium. DC.	(ye.) Serpyllum-ld.	England.	5. 9.	—— t. 34.
75	vulgare. DC.	(ye.) common.	Britain.	H. ⓧ.	—— t. 64.
	β flòre pleno.	(ye.) double-flowered.	H. ⓧ.	—— t. 28.
76	surrejánum. DC.	(ye.) dotted-leaved.	England.	7. 10.	t. 34. f. 3.
	β tomentòsum.	(ye.) woolly-leaved.	H. ⓧ.	Viv. frag. 1. t. 8. f. 2.
77	ovatum. DC.	(ye.) ovate-leaved.	Geneva.	6. 8.	
78	lúcidum. H.	(ye.) glossy-leaved.	1827.	H. ⓧ.	Swt. cist. t. 105.
79	táuricum. s.c.	(p.y.) Taurian.	Tauria.	1824.	H. ⓧ.	—— t. 89.
80	Andersoni. s.c.	(ye.) Anderson's.	Hybrid.	1827.	5. 10.	—— t. 69.
81	grandifòrum. DC.	(ye.) large-flowered.	Pyrenees.	1800.	H. ⓧ.	Bot. mag. 1803.
82	sampsucifòlium.	(p.y.) rough dwarf.	Montpelier.	H. ⓧ.	
	<i>Cistus sampsucifòlius.</i> B.M. non Cav.					
83	nummularium. DC.	(ye.) Money-wort.	1752.	H. ⓧ.	Swt. cist. t. 80.
84	angustifòlium. DC.	(ye.) distinct-petaled.	1800.	H. ⓧ.	Jacq. vind. 3. t. 53.
85	hirtum. DC.	(ye.) bristly-calyxed.	Spain.	1759.	F. ⓧ.	Swt. cist. t. 109.
86	barbátum. P.S.	(ye.) bearded.	S. Europe.	1820.	H. ⓧ.	—— t. 73.
87	eriosépalon. s.c.	(p.y.) woolly-sepaled.	H. ⓧ.	—— t. 76.
88	rotatum. rotate.	1828.	F. ⓧ.	Swt. cist. t. 82.
89	racemósrum. DC.	(wh.) long-racemed.	S. Europe.	F. ⓧ.	—— t. 49.
90	pilosum. DC.	(wh.) hairy.	1731.	F. ⓧ.	—— t. 48.
91	lineáre. DC.	(wh.) linear-leaved.	1824.	6. 8.	F. ⓧ.

CISTINEÆ.

92	virgátum. DC. (p.r.) slender.	Barbary.	—	—	F. $\frac{1}{2}$.	Swt. cist. t. 79.
93	apenninum. DC. (wh.) Apennine.	Europe.	—	—	H. $\frac{1}{2}$.	— t. 62.
94	hispidum. DC. (wh.) hispid.	S. France.	—	—	H. $\frac{1}{2}$.	
95	pulveruléntum. DC. (w.) powdered.		—	—	H. $\frac{1}{2}$.	Swt. cist. t. 29.
96	canescens. s.c. (cr.) canescent.		—	—	H. $\frac{1}{2}$.	— t. 51.
97	rhodánthum. DC. (ro.) rose-flowered.	Spain.	—	—	H. $\frac{1}{2}$.	— t. 7.
98	diversifólium. s.c. (re.) various-leaved.		—	—	5. 9. F. $\frac{1}{2}$.	— t. 95.
	β multiplex. s.c. (re.) full-flowered.		—	—	F. $\frac{1}{2}$.	— t. 98.
99	cùpreum. s.c. (co.) copper-colour'd. Hybrid.		—	—	F. $\frac{1}{2}$.	t. 66.
100	versicolor. s.c. (p.r.) various-colour'd.		—	—	H. $\frac{1}{2}$.	— t. 26.
101	venústum. s.c. (sc.) charming.		—	—	H. $\frac{1}{2}$.	— t. 10.
	β flôre plêno.		—	—	H. $\frac{1}{2}$.	
102	variegátum. s.c. (r.w.) striped-flower'd. Hybrid.		—	—	F. $\frac{1}{2}$.	Swt. cist. t. 38.
103	lanceolátum. s.c. (w.) spear-leaved.		—	—	H. $\frac{1}{2}$.	— t. 100.
104	polifólium. DC. (wh.) white-mountain.	England.	—	5. 7.	H. $\frac{1}{2}$.	— t. 88.
105	confusum. s.c. (wh.) confused.	Europe.	—	—	H. $\frac{1}{2}$.	— t. 91.
106	macrâthon. s.c. (w.) great-flowered.		—	—	H. $\frac{1}{2}$.	— t. 103.
	β multiplex. s.c. (w.) full-flowered.		—	—	H. $\frac{1}{2}$.	— t. 104.
107	mutábile. DC. (p.r.) changeable.	Spain.	1795.	6. 7.	H. $\frac{1}{2}$.	— t. 106.
108	rôseum. DC. (ro.) rosy.	S. Europe.	—	—	H. $\frac{1}{2}$.	— t. 55.
	β multiplex. sc. (ro.) double-rose.		—	—	H. $\frac{1}{2}$.	— t. 86.
109	Milléri. s.c. (sa.) Miller's.		—	—	H. $\frac{1}{2}$.	— t. 101.
110	hyssopifólium. DC. (v.) Hyssop-leaved.	Italy.	—	—	F. $\frac{1}{2}$.	
	α crocátum. s.c. (sa.) saffron-colour'd.		—	—	F. $\frac{1}{2}$.	Swt. cist. t. 92.
	β cùpreum. s.c. (co.) copper-coloured.		—	—	F. $\frac{1}{2}$.	— t. 58.
	γ multiplex. s.c. (co.) full-flowered.		—	—	F. $\frac{1}{2}$.	— t. 72.
111	fe'tidum. DC. (wh.) Bryony-scented.		—	—	H. $\frac{1}{2}$.	Jacq. ic. 1. t. 98.
112	sulphureum. DC. (st.) sulphur-color'd.	Spain.	1815.	—	H. $\frac{1}{2}$.	Swt. cist. t. 37.
113	stramíneum. s.c. (st.) straw-coloured.		—	—	H. $\frac{1}{2}$.	— t. 93.
	β multiplex. s.c. (st.) full-flowered.		—	—	H. $\frac{1}{2}$.	— t. 94.
HUDS'ONIA. DC.		HUDS'ONIA. Dodecadenia Monogynia. L.				
1	ericóides. DC. (ye.) Heath-like.	N. America.	1805.	5. 7.	F. $\frac{1}{2}$.	Swt. cist. t. 36.
2	tomentosa. N. (ye.) tomentose.		1826.	6. 8.	F. $\frac{1}{2}$.	— t. 57.
LECH'EA. DC.		LECH'EA. Tri-Dodecadria Trigynia.				
1	villòsa. DC. (ye.) villous.	Canada.	1780.	7. 8.	H. $\frac{1}{2}$.	Lam. ill. t. 52. f. 2.
	májor. Ph. non Lin. minor L. non Ph.					
2	minor. DC. (ye.) lesser.		1802.	—	H. $\frac{1}{2}$.	Lam. ill. t. 52. f. 1.
3	thymifolia. DC. (ye.) Thyme-leaved.	N. America.	1823.	—	H. $\frac{1}{2}$.	

ORDO XX.

VIOLARIEÆ. DC. prodr. 1. p. 287.

Tribus I. VIOLEÆ. DC.

CALY'PTRION. DC.	CALY'PTRION.	Pentandria Monogynia.				
Aubléti. DC. (wh.) Aublet's.		S. America.	1820.	S. $\frac{1}{2}$.	Aub. gui. 2. t. 319.
<i>Viola Hybánthus</i> . Aub. <i>Jonidium Aubléti</i> . R.S.						
NOISETTIA. DC.	NOISETTIA.	Pentandria Monogynia.				
1 longifòlia. DC. (wh.) long-leaved.		Cayenne.	1822.	S. $\frac{1}{2}$.	Nov. g. am. t. 409. b.f. 2.
<i>Jonidium longifolium</i> . R.S.						
2 pyrifòlia. Mart. (wh.) Pear-leaved.	Brazil.	1822.	S. $\frac{1}{2}$.	Mart. n. pl. bras. ic.	
VIOLA. DC.	VIOLET.	Pentandria Monogynia.				
1 pedáta. DC. (bl.) cut-leaved.	N. America.	1759.	5. 8.	H. $\frac{1}{2}$.	Swt. br. fl. gar. t. 69.	
2 flabellifòlia. B.C. (b.p.) fan-leaved.		1822.	—	H. $\frac{1}{2}$.	Lod. bot. cab. 777.	
	<i>pedáta</i> β <i>atropurpurea</i> . DC. <i>pedáta</i> var. <i>bicolor</i> . Ph.					
3 digitáta. Ph. (bl.) finger-leaved.	Virginia.	1824.	—	H. $\frac{1}{2}$.		
4 palmáta. DC. (bl.) palmated.		1752.	5. 6.	H. $\frac{1}{2}$.	Bot. mag. 535.	
5 asarifolia. DC. (bl.) Asarum-leaved.		1823.	—	H. $\frac{1}{2}$.		
6 papilionácea. DC. (bl.) variegated.		1800.	—	H. $\frac{1}{2}$.		
7 obliqua. Ph. (bl.) oblique-flower'd.		1762.	—	H. $\frac{1}{2}$.		
8 euculláta. Ph. (bl.) hollow-leaved.		1772.	5. 7.	H. $\frac{1}{2}$.	Bot. mag. 1795.	
9 pinnáta. R.S. (pu.) winged-leaved.	Europe.	1752.	5. 6.	H. $\frac{1}{2}$.	All. m. taur. 3. t. 5. f. 2.	
10 multifida. R.S. (pu.) multifid.	Siberia.	—	—	H. $\frac{1}{2}$.	Gmel. sib. 4. t. 48. f. 4.	

VIOLARIEÆ.

11 variegata. DC.	(pu.)	variegated-l'd.	Dahuria.	1818.	5. 6.	H. Ȑ.
12 primulæfolia. DC.	(bl.)	Primrose-l'd.	N. America.	1783.	4. 6.	H. Ȑ.
13 lanceolata. DC.	(wh.)	spear-leaved.	—	1759.	6. 7.	F. Ȑ.
14 Patrini. DC.	(bl.)	Siberian.	Siberia.	...	—	H. Ȑ.
15 cæspitosa. D.P.	(pu.)	tufted.	Nepaul.	1825.	5. 8.	F. Ȑ.
<i>Patrini γ napaulensis. DC. primulæfolia. Buch. non L.</i>						
16 palmaris. D.P.	(ye.)	long-petioled.	—	—	—	F. Ȑ.
17 ovata. DC.	(bl.)	oval-leaved.	N. America.	1822.	—	H. Ȑ.
18 sagittata. DC.	(b.w.)	arrow-leaved.	—	1775.	—	H. Ȑ.
19 dentata. Ph.	(bl.)	large-toothed.	Penssylvania.	1828.	—	H. Ȑ.
20 betonicæfolia. DC.	(pu.)	Betony-leaved.	N. S. W.	1823.	—	F. Ȑ.
21 Fischeri.	(pu.)	Fischer's.	Siberia.	1823.	5. 7.	H. Ȑ.
<i>sudavis. F. non. M.B.</i>						
22 palustris. DC.	(pa.)	marsh.	Britain.	...	5. 6.	H. Ȑ.
23 blanda. DC.	(wh.)	white-flowered.	N. America.	1802.	5. 7.	H. Ȑ.
24 rotundifolia. DC.	(ye.)	round-leaved.	—	1800.	—	H. Ȑ.
25 sororia. W.	(pu.)	white-rooted.	—	1802.	4. 6.	H. Ȑ.
26 aspera. DC.	(wh.)	rough.	Nepaul.	1825.	5. 8.	F. Ȑ.
27 hirta. E.B.	(bl.)	hairy.	England.	...	—	H. Ȑ.
28 ambigua. R.S.	(pu.)	ambiguous.	Danube.	1822.	—	H. Ȑ.
29 campestris. M.B.	(pu.)	field.	Tauria.	1824.	—	H. Ȑ.
30 hirsuta. R.S.	(bl.)	hairy-stalked.	Russia.	1824.	1. 12.	H. Ȑ.
31 collina. R.S.	(li.)	hill.	Volhynia.	—	5. 8.	H. Ȑ.
32 cordata. S.S.	(bl.)	heart-leaved.	N. America.	—	5. 7.	H. Ȑ.
33 suavis. M.B.	(p.b.)	fragrant.	Siberia.	1823.	3. 5.	H. Ȑ.
34 odorata. DC.	(vi.)	sweet-scented.	Britain.	...	3. 10.	H. Ȑ.
<i>α violacea.</i> (vi.) violet-coloured.						
β rosea.	(ro.)	rose-coloured.	—	—	—	H. Ȑ.
γ alba.	(wh.)	white-flowered.	—	—	—	H. Ȑ.
δ caerulea.	(bl.)	blue-flowered.	—	—	—	H. Ȑ.
ε purpureo-plena. (pu.)	double-purple.	—	—	—	—	H. Ȑ.
ζ albo-plena. (wh.)	double-white.	—	—	—	—	H. Ȑ.
η caeruleo-plena. (bl.)	double-blue.	—	—	—	—	H. Ȑ.
θ pallido-plena. (p.b.)	Neapolitan.	Naples.	...	—	—	H. Ȑ.
35 uliginosa. DC.	(pu.)	fen.	Carniola.	1822.	4. 5.	H. Ȑ.
36 mirabilis. DC.	(w.l.)	broad-leaved.	Europe.	1732	6. 8.	H. Ȑ.
37 striata. DC.	(st.)	streaked.	N. America.	1772.	—	H. Ȑ.
38 ochroleuca. DC.	(st.)	sulphur-color'd.	—	—	—	H. Ȑ.
39 canina. DC.	(bl.)	dog's.	Britain.	...	4. 6.	H. Ȑ.
40 sylvestris. R.S.	(p.b.)	wood.	Hungary.	1823.	—	H. Ȑ.
41 Riviniana. s.s.	(bl.)	Rivini's.	Europe.	...	4. 6.	H. Ȑ.
42 ericotorum. L.en.	(bl.)	heath.	Germany.	1826.	—	H. Ȑ.
43 neglecta. R.S.	(p.b.)	neglected.	Bohemia.	1824.	—	H. Ȑ.
44 débilis. DC.	(p.b.)	weak.	N. America.	1820.	—	H. Ȑ.
45 arenaria. DC.	(bl.)	sand.	Europe.	—	—	H. Ȑ.
<i>Allionii. R.S. nummulariifolia. Schl.</i>						
46 glauca. M.B.	(pu.)	glauous.	Russia.	1820.	—	H. Ȑ.
47 pumila. DC.	(p.b.)	small.	Europe.	—	—	H. Ȑ.
48 flavicornis. E.F.	(bl.)	yellow-spurred.	Britain.	...	5. 7.	H. Ȑ.
49 montana. s.s.	(bl.)	mountain.	Europe.	1683.	5. 6.	H. Ȑ.
50 stricta. H.	(p.b.)	upright.	—	1822.	—	H. Ȑ.
<i>Hornemanniana. R.S.</i>						
51 persicifolia. s.s.	(bl.)	Peach-leaved.	S. Europe.	1818.	—	H. Ȑ.
52 laccea. R.S.	(pa.)	cream-coloured.	England.	...	—	H. Ȑ.
53 Ruppii. s.s.	(pa.)	Piedmont.	Piedmont.	1820.	—	H. Ȑ.
54 Broussonetiæna. R.S.	(pa.)	Broussonet's.	Teneriffe.	1825.	—	G. Ȑ.
55 arboræscens. DC.	(pa.)	shrubby.	S. Europe.	1779.	4. 5.	G. Ȑ.
56 epipsila. s.s.	(p.b.)	creeping.	Siberia.	1821.	4. 6.	H. Ȑ.
57 femina. F.	(bl.)	Siberian.	—	1827.	—	H. Ȑ.
58 Selkirkii. DC.	Selkirk's.	Montreal.	1823.	—	H. Ȑ.
59 pygmæa. DC.	(p.b.)	pigmy.	Andes.	1822.	7. 11.	H. Ȑ.
60 biflora. DC.	(ye.)	two-flowered.	Europe.	1752.	—	H. Ȑ.
61 Nuttallii. DC.	(ye.)	Nuttall's.	N. America.	1812.	5. 6.	H. Ȑ.
62 hastata. DC.	(ye.)	hastate.	—	1823.	5. 7.	H. Ȑ.
63 albiloba. L.en.	(wh.)	white-flowered.	—	—	—	H. Ȑ.
64 canadensis. DC.	(w.r.)	Canadian.	—	1783.	—	H. Ȑ.
65 pubescens. DC.	(ye.)	downy.	—	1772.	6. 7.	H. Ȑ.
66 eriocarpa. DC.	(ye.)	woolly-capsul'd.	—	1820.	—	H. Ȑ.
67 uniflora. DC.	(ye.)	Siberian.	Siberia.	1774.	—	H. Ȑ.

Swt. br. fl. gar. 2. t. 174.

Eng. bot. 444.
W. hort. ber. 1. t. 24.

W. hort. ber. 1. t. 72.

Eng. bot. 894.
W. et K. hung. t. 190.

Eng. bot. t. 619.

Jacq. fl. aust. t. 19.

Eng. bot. 620.

Pio viol. p. 20. t. 1. f. 2.

Vill. c. stras. t. 5.
Raii synops. t. 24. f. 1.
Bot. mag. 1595.Schkuhr hand. 3. t. 269.
Eng. bot. 445.
Lodd. bot. cab. 686.

Barr. ic. 568.

Flor. dan. 46.

Swt. br. fl. gar. 223.

102.

Gmel. sib. 4. t. 48. f. 5.

68 chrysántha. Ser. (<i>ye.</i>) golden-flowered.	1823.	5. 8. H. ȝ.	
69 nummularifólia. DC. (<i>b.</i>) money-wort. Piedmont. — — H. ȝ.			All. ped. n. 1640. t. 9. f. 4.
70 alpina. DC. (<i>pu.</i>) alpine. Austria. 1826. — — H. ȝ.			Jacq. obs. 1. t. II.
71 cornúta. DC. (<i>bl.</i>) horned. Pyrenees. 1776. 5. 6. H. ȝ.			Bot. mag. 791.
72 cenísia. DC. (<i>bl.</i>) Mount-Cenis. Switzerland. 1759. 6. 7. H. ȝ.			All. ped. 2. t. 22. f. 6.
73 valdériæ. R.S. (<i>pu.</i>) fringed-leaved. — — 1805. 5. 6. H. ȝ.			— — 2. t. 24. f. 3.
74 calcaráta. R.S. (<i>bl.</i>) spurred. Europe. 1752. 3. 6. H. ȝ.			Vill. cat. arg. t. 5.
75 Villarsiàna. R.S. (<i>bl.</i>) Villars's. S. France. — — H. ȝ.			Jacq. coll. 4. t. 11. f. 1.
76 Zóysii. R.S. (<i>ye.</i>) crenated. Europe. 7. 9. H. ȝ.			
77 amœ'na. DC. (<i>pu.</i>) purple. Scotland. 5. 8. H. ȝ.			
78 altáica. DC. (<i>ye.</i>) Tartarian. Altay. 1805. 3. 6. H. ȝ.			Bot. reg. 54.
grandiflòra. L. non DC. β purpurea. DC. (<i>pu.</i>) purple-flowered. Altay. 3. 6. H. ȝ.			Swt. br. fl. gar. ic.
79 prostrata. R.S. (<i>st.</i>) prostrate. Teneriffe. 1824. — — H. ♂.			
80 sudética. W.en. (<i>ye.</i>) large-flowered. Europe. 5. 8. H. ȝ.			
grandiflòra. DC. fl. fr. 5. p. 620.			
81 lutea. E.B. (<i>ye.</i>) large yellow. Britain. 5. 8. H. ȝ.			Eng. bot. 721.
82 saxátilis. R.S. (<i>st.</i>) rock. Bohemia. 1824. — — H. ȝ.			Swt. br. fl. gar. t.
83 rothomagénsis. DC. (<i>b.</i>) Rouen. France. 1783. — — H. ȝ.			Bot. mag. 1498.
84 tricolor. R.S. (<i>b.y.</i>) Heart's-ease. Britain. 4. 11. H. ♂.			(Ed. fl. dan. t. 623.)
85 bannática. R.S. (<i>st.</i>) intermediate. Hungary. 5. 8. H. ♂.			
86 arvénisia. R.S. (<i>st.</i>) corn. Britain. — — H. ♂.			
87 declináta. R.S. (<i>bl.</i>) declined. Hungary. 1820. — — H. ♂.			W. et K. hung. 3. t. 223.
88 Kitaibeliána. R.S. (<i>st.</i>) Kitaibel's. Switzerland. 1824. 4. 6. H. ♂.			
ERP'ETION B.F.G. SPURLESS VIOLET. Pentandria Monogynia.			
1 hederáceum. (<i>b.w.</i>) Ivy-leaved. V. Diem. Isl. 1825. 5. 9. F. ȝ.			Lab. nov. holl. 1. t. 91.
Viola hederacea. Lab.			
2 renífórmæ. B.F.G. (<i>b.w.</i>) kidney-leaved. N. S. W. 1823. — — F. ȝ.			Swt. br. fl. gar. 170.
Viola reniformis. B.			
SOLEA. DC. S'OLEA. Pentandria Monogynia.			
cónicolor. DC. (<i>gr.</i>) green-flowered. N. America. 1788. 6. 7. H. ȝ.			Linn. trans. 6. t. 28.
POMB'ALIA. DC. POMB'ALIA. Pentandria Monogynia.			
'Itubu. DC. (<i>w.h.</i>) white Ipecacuanha. Brazil. 1822. 7. 8. S. ȝ.			H. et. B. 5. t. 496. f. 2.
β calceolária. (<i>w.h.</i>) villous-stemmed. — — — — S. ȝ.			Bot. mag. 2453
Jonidiánum Ipecacuánha. β . B.M. 2453.			
JONIDIUM. DC. JONIDIUM. Pentandria Monogynia.			
1 stríctum. DC. (<i>w.h.</i>) upright. St. Domingo. 1820. S. ȝ.			
2 polygalæfólium. DC. (<i>st.</i>) Milkwort-lö. S. America. 1797. 4. 8. G. ȝ.			Vent. malm. 27.
3 verbenáceum. DC. (<i>p.b.</i>) Vervain-like. — — 1822. 6. 8. H. ♂.			H. et B. V. 379. t. 497.

Tribus II. ALSODINEÆ. DC.

ALSODE'IA. DC. ALSODE'IA. Pentandria Monogynia.			
1 arbórea. DC. (<i>w.h.</i>) tree. Madagascar. 1823. S. ȝ.			
2 latifólia. DC. (<i>w.h.</i>) broad-leaved. — — — — S. ȝ.			Pet.Th. veg. af. t. 18. f. 2.
CERA'NTHERA. DC. CERA'NTHERA. Pentandria Monogynia.			
dentáta. DC. (<i>w.h.</i>) toothed. Sierra Leone. 1822. S. ȝ.			Beauv. fl. d'ow. 2. t. 65.

Tribus III. SAUVAGEÆ. DC.

SAUVAG'ESIA. DC. SAUVAG'ESIA. Pentandria Monogynia.			
1 erécta. DC. (<i>w.h.</i>) upright. S. America. 1823. 7. 9. S. ♂.			Jacq. am. pict. t. 77.
2 nútans. DC. (<i>w.h.</i>) nodding. Madagascar. — — — — S. ♂.			

ORDO XXI.

DROSERACEÆ. DC. prodr. 1. p. 317.

DR'OSERA. DC. SUNDEW. Pentandria Pentagynia. L.			
1 acáulis. DC. (<i>w.h.</i>) stalkless. C. B. S. 1821. 7. 8. G.w. ȝ.			
2 pauciflòra. DC. (<i>w.h.</i>) few-flowered. — — — — G.w. ȝ.			

DROSERACEÆ.

3 cuneifolia. DC.	(w.r.) wedge-leaved.	C. B. S.	1824.	6. 8. G. w. ȝ.	
α alba.	(wh.) white-flowered.	—	—	— G. w. ȝ.	
β rubra.	(re.) red-flowered.	—	—	— G. w. ȝ.	
4 rotundifolia. DC.	(wh.) round-leaved.	Britain.	7. 8. H. w. ȝ.	
5 longifolia. L.	(wh.) intermediate.	England.	— H. w. ȝ.	
intermedia. DC.					Eng. bot. 868.
6 americana. W. en.	(wh.) American.	N. America.	1824.	— H. w. ȝ.	
7 anglica. DC.	(wh.) great.	—	— H. w. ȝ.	
longifolia. D. et H. 3. t. 75. A.					Eng. bot. 869.
8 linearis. DC.	(wh.) linear-leaved.	N. America.	1822.	5. 6. H. w. ȝ.	
9 filiformis. Ph.	(pu.) thready-leaved.	—	1811.	— H. w. ȝ.	
ALDROVA'NDA. DC.	ALDROVA'NDA.	Pentandria Pentagynia.	L.		
vesiculosa. DC.	(wh.) whorl-leaved.	S. Europe.	1826.	6. 8. H. w. ⓧ.	
B'YBLIS. DC.	B'YBLIS.	Pentandria Monogynia.			
liniflora. DC.	(bl.) blue-flowered.	N. S. W.	1803.	5. 6. G. w. ȝ.?	
DION'EA. DC.	DION'EA.	Decandria Monogynia.	L.		
Muscipula. DC.	(wh.) Venus Flytrap.	Carolina.	1768.	7. 8. G. ȝ.	
PARNA'SSIA. DC.	PARNA'SSIA.	Pentandria Tetragynia.	L.		
1 palustris. DC.	(wh.) marsh.	Britain.	7. 8. H. w. ȝ.	
2 parviflora. DC.	(wh.) small-flowered.	N. America.	5. 6. H. w. ȝ.	
3 caroliniana. DC.	(wh.) Carolina.	Carolina.	1802.	5. 8. H. w. ȝ.	
4 asarifolia. DC.	(wh.) Asarum-leaved.	N. America.	1812.	7. 8. H. w. ȝ.	
					Vent. malm. t. 39.

ORDO XXII.

POLYGALÆ. DC. prodr. 1. p. 231.

POLYGALA. DC.	MILKWORT.	Diadelphia Octandria. L.			
1 oppositifolia. DC.	(pu.) opposite-leav'd.	C. B. S.	1790.	1. 12. G. h.	Bot. reg. 636.
2 cordifolia. DC.	(pu.) heart-leaved.	—	1791.	— G. h.	
3 tetragona. DC.	(pu.) square-stalked.	—	1820.	— G. h.	
	attenuata. Lod. bot. cab. 1000. <i>oppositifolia</i> β major. B.R. 1146.				
4 nummularia. DC.	(pu.) Moneywort-ld.	C. B. S.	1812.	— G. h.	
5 latifolia. B.R.	(pu.) broad-leaved.	—	1820.	— G. h.	Bot. reg. 645.
	cordifolia. B.M. non DC.				
6 borboniaeæfolia. DC.	(p.) Borbonia-leav'd.	—	1790.	— G. h.	
	<i>oppositifolia</i> . B.M. non DC.				
7 grandiflora. B.C.	(pu.) large-flowered.	—	1825.	4. 9. G. h.	Lodd. bot. cab. 1227.
8 myrtifolia. DC.	(pu.) Myrtle-leaved.	—	1707.	5. 8. G. h.	Bot. reg. 669.
9 ligulâris. DC.	(pu.) ligulate-leaved.	—	1820.	— G. h.	— 637.
10 intermedia. DC.	(pu.) intermediate.	—	5. 10. G. h.	
11 bracteolata. DC.	(pu.) spear-leaved.	—	1713.	— G. h.	Bot. mag. 345.
12 Linkiana.	(pa.) slender-leaved.	—	1820.	— G. h.	
	<i>tenuifolia</i> . L. en. non w.				
13 umbellata.	(pu.) hispid-stalked.	—	—	— G. h.	Burm. afr. t. 73. f. 5.
14 Burmanni. DC.	(pu.) Burman's.	—	— G. h.	— t. 73. f. 4.
15 simplæx. DC.	(pu.) simple-stalked.	—	1820.	— G. h.	
16 affinis. DC.	(pu.) blunt-leaved.	—	1825.	6. 9. G. h.	
17 speciosa. DC.	(pu.) showy.	—	1814.	— G. h.	Bot. reg. 150.
18 teretifolia. DC.	(pu.) columnar-ld.	—	1791.	5. 8. G. h.	Andr. rep. 370.
19 Garcini. DC.	(pu.) slender-branch'd	—	5. 10. G. h.	Burm. afr. t. 73. f. 3.
20 major. DC.	(pu.) large Austrian.	Austria.	1739.	7. 8. H. ȝ.	Jacq. aust. 5. t. 413.
21 vulgaris. DC.	(va.) common.	Britain.	5. 7. H. ȝ.	Eng. bot. 76.
	α vera. DC.	(r.b.) true.	— H. ȝ.	Flor. dan. t. 516.
	β pubescens. DC.	(r.b.) pubescent.	— H. ȝ.	
	γ elata. DC.	(r.b.) tall.	— H. ȝ.	Schk. handb. t. 194.
	δ Verviana. DC.	(wh.) white-flowered.	— H. ȝ.	
	ε acutifolia. DC.	(bl.) blue-flowered.	— H. ȝ.	
	ζ angustifolia. DC.	(ro.) rose-coloured.	— H. ȝ.	
	η grandiflora. DC.	(re.) red-flowered.	— H. ȝ.	
22 amara. DC.	(bl.) bitter.	Europe.	1775.	— H. ȝ.	Bot. mag. 2437.
23 monspeliaca. DC.	(fl.) Montpelier.	Montpelier.	1823.	6. 9. H. ⓧ.	DC. ic. rar. 1. t. 9.
24 incarnata. DC.	(fl.) flesh-coloured.	N. America.	1812.	6. 7. H. ⓧ.	
25 cruciata. DC.	(ro.) four-leaved.	—	1739.	— H. ⓧ.	
26 brevifolia. DC.	(re.) short-leaved.	—	1824.	— H. ⓧ.	
27 fastigiata. DC.	(re.) close-branched.	—	—	— H. ⓧ.	

28 lùtea. DC.	(ye.) golden.	N. America.	1739.	6. 7. H. ⊖.	Pluk. am. t. 438. f. 6.
29 nàna. DC.	(gr.) dwarf.	_____	1815.	7. 8. H. ⊖.	
30 purpùrea. DC.	(ro.) purple. sanguínea. Mich.	_____	1739.	7. 9. H. ⊖.	
31 sanguínea. DC.	(cr.) bloody.	_____	_____	H. ⊖.	
32 ambigua. DC.	(pu.) ambiguous.	_____	1824.	6. 7. H. ⊖.	Pluk. am. t. 438. f. 4.
33 verticillàta. DC.	(w.r.) whorl-leaved.	_____	1739.	— H. ⊖.	Bot. reg. 761.
34 paniculàta. DC.	(pu.) paniced.	S. America.	1822.	7. 10. S. ⊖.	
35 graminifòlia. DC.	(st.) grass-leaved.	N. America.	—	6. 7. F. 4.	
36 ábla. N.	(wh.) white-flowered.	Louisiana.	1827.	6. 7. H. 4.	
37 rubélla. DC.	(re.) reddish.	N. America.	1828.	— H. 4.	
38 Sénega. DC.	(bh.) Rattlesnake-rt.	_____	1739.	— H. 4.	Bot. mag. 1051.
39 paucifòlia. DC.	(pu.) naked-stalked.	_____	1791.	5. 6. H. 4.	— 2852.
40 commutatâ. (pu.) frutescent.	purpùrea. H.K. nec aliorum.	_____	_____	H. 2.	
41 Chamæbúxus. DC. (w.)	Box-leaved.	Europe.	1658.	— H. 2.	Bot. mag. 316.
SALOM'ONIA. DC.	SALOM'ONIA.	Monadelphia	Tetrandria.		
cantoniénsis. DC. (wh.)	Canton.	China.	1824.	6. 8. H. ⊖.	
COMESPER'MA. DC.	COMESPER'MA.	Monadelphia	Octandria.		
1 ericina. DC.	(pu.) Heath-like.	N. S. W.	1822.	5. 8. G. 2.	
2 virgàta. DC.	(pu.) twiggy.	_____	1826.	— G. 2.	Lab. nov. holl. 2. t. 159.
3 coridifòlia. F.T.	(pu.) Coris-leaved.	_____	1822.	— G. 2.	
MURA'LTIÀ. DC.	MURA'LTIÀ.	Diadelphia	Octandria. L.		
1 Heistèria. DC.	(pu.) Furze-leaved.	C. B. S.	1787.	1. 12. G. 2.	Bot. mag. 340.
Poly'gala Heistèria. B.M.					
2 alopecuróides. DC. (p.)	Fox-tail.	_____	1800.	5. 8. G. 2.	Bot. mag. 1006.
Poly'gala alopecuróides. B.M.					
3 squarròsa. DC.	(li.) squarrose.	_____	1821.	— G. 2.	
4 stipulàcea. DC.	(pu.) stipuled.	_____	1801.	4. 9. G. 2.	Bot. mag. 1715.
5 juniperifòlia. DC. (re.)	Juniper-leaved.	_____	...	G. 2.	
6 diffusa. DC.	(li.) diffuse.	_____	1800.	1. 12. G. 2.	
7 vírgata. DC.	(pu.) twiggy.	_____	—	G. 2.	
8 linophylla. DC.	(pu.) Flax leaved.	_____	1812.	5. 8. G. 2.	
9 macrocèras. DC.	(pu.) long-horned.	_____	—	G. 2.	
10 míxta. DC.	(li.) Heath-leaved.	_____	1791.	1. 12. G. 2.	Bot. mag. 1714.
Poly'gala míxta. B.M.					
α rúbra.	(li.) pale red.	_____	—	G. 2.	
β ábla.	(wh.) white-flowered.	_____	—	G. 2.	
11 ciliàris. DC.	(li.) ciliated.	_____	1822.	5. 7. G. 2.	
12 hùmilis. DC.	(pu.) humble.	_____	1817.	5. 8. G. 2.	Lodd. bot. cab. 420.
Poly'gala hùmilis. B.C.					
13 pubéscent. DC.	(li.) pubescent.	_____	1823.	— G. 2.	
MU'NDIA. DC.	MU'NDIA.	Diadelphia	Octandria. L.		
spinòsa. DC.	(li.) spiny.	C. B. S.	1780.	1. 5. G. 2.	
α latifòlia. DC.	(li.) broad-leaved.	_____	—	G. 2.	
β angustifòlia. DC.	(li.) narrow-leaved.	_____	—	G. 2.	
SECURID'ACA. DC.	SECURID'ACA.	Diadelphia	Octandria. L.		
1 virgàta. DC.	(v.) slender.	Jamaica.	1822.	... S. 2.	Plum.ed.Bur.t.248.f.1
2 volubilis. DC.	(re.) climbing.	W. Indies.	1739.	... S. 2.	Jacq. amer.t.183.f.38.
3 erécta. DC.	(pu.) upright.	_____	1824	... S. 2.	— t.183.f.39.
4 paniculàta. DC.	(st.) paniced.	Cayenne.	1820.	... S. 2.	Lam. ill. t. 599. f. 2.

ORDO XXIII.

TREMANDREÆ. DC. prodr. 1. p. 343.

TETRATH'ECA. DC. TETRATH'ECA. Octandria Monogynia.

1 júncea. DC.	(w.p.) rushy.	N. S. W.	1803.	7. 8. G. 2.	Sm. nov. hol. 1. t. 2.
2 glandulòsa. DC.	(re.) glandular.	V. Diemen.	1822.	— G. 2.	Lab. nov. hol. 1. t. 123.
3 pilosa. DC.	(re.) hairy.	_____	—	G. 2.	— 1. t. 122.
glandulòsa. Ex. bot. 21.					
4 ericifòlia. DC.	(r.w.) Heath-leaved.	N. S. W.	—	G. 2.	Sm. exot. bot. t. 20.
5 rubiæóides. F.T.	(ro.) Rubia-like.	_____	1823.	6. 8. G. 2.	

ORDO XXIV.

PITTOSPOREÆ. DC. prodr. 1. p. 345.

BILLARDIE'RA. DC.	APPLE-BERRY.	Pentandria Monogynia. L.					
1 scàndens. DC.	(st.) downy-fruited.	N. S. W.	1790.	6. 8.	G. h. ↗.	Swt. fl. aust. t. 54.	
2 mutabilis. DC.	(p.s.) changeable.	_____	1795.	5. 9.	G. h. ↗.	Bot. mag. 1313.	
3 longiflòra. DC.	(st.) blue-fruited.	V. Diemen.	1810.	—	G. h. ↗.	_____ 1507.	
4 fusiformis. DC.	(bl.) long-fruited.	_____	1822.	—	G. h. ↗.	Lab. nov. holl. t. 90.	
5 angustifolia. DC.	(st.) narrow-leaved.	N. S. W.	_____	—	G. h. ↗.	_____	
PITTO'SPORUM. DC.	PITTO'SPORUM.	Pentandria Monogynia. L.					
1 coriaceum. DC.	(wh.) thick-leaved.	Madeira.	1787.	5. 6.	G. h.	Andr. rep. 151.	
2 viridiflorum. DC.	(gr.) green-flowered.	C. B. S.	1806.	—	G. h.	Bot. mag. 1684.	
3 Tobira. DC.	(wh.) glossy-leaved.	China.	1804.	3. 8.	G. h.	_____ 1396.	
4 undulatum. DC.	(wh.) wave-leaved.	N. S. W.	1789.	2. 6.	G. h.	Bot. reg. 16.	
5 revolutum. DC.	(st.) revolute-leaved.	_____	1795.	6. 7.	G. h.	_____ 186.	
6 tomentosum. DC.	(st.) woolly-leaved.	N. Holland.	1825.	6. 7.	G. h.	Swt. fl. aust. t. 33.	
7 fulvum. DC.	(st.) tawny.	N. S. W.	1824.	1. 5.	G. h.	_____ t. 25.	
8 tenuifolium. G.	(st.) Willow-leaved.	_____	1818.	—	G. h.	Gært. fr. l. t. 59.	
9 ferrugineum. DC.	(st.) rusty-leaved.	Guinea.	1787.	2. 5.	S. h.	Bot. mag. 2075.	
BURS'ARIA. DC.	BURS'ARIA.	Pentandria Monogynia. L.					
spinosa. DC.	(wh.) thorny.	N. S. W.	1793.	8. 12.	G. h.	Bot. mag. 1767.	
<i>Itea spinosa.</i> And. rep. t. 314.							
SEN'ACIA. DC.	SEN'ACIA.	Pentandria Monogynia. L.					
1 undulata. DC.	(wh.) wave-leaved.	Mauritius.	1785.	—	S. h.		
2 nepalensis. DC.	(wh.) flat-leaved.	Nepaul.	1826.	—	G. h.		

ORDO XXV.

FRANKENIACEÆ. DC. prodr. 1. p. 349.

FRANK'ENIA. DC.	SEA HEATH.	Hexandria Monogynia. L.					
1 pulverentá. DC.	(re.) powdery.	England.	7.	H. 4.	Eng. bot. 2222.	
2 nodiflòra. DC.	(re.) knotted-flow'r'd.	C. B. S.	1818.	6. 8.	G. 4.	Lam. ill. t. 262. f. 4.	
3 Nôthria. DC.	(re.) Cape.	_____	1816.	—	G. 4.	Berg. cap. 171. t. 1. f. 2.	
<i>Nôthria r��pens.</i> Berg.							
4 l��vis. DC.	(re.) smooth.	England.	7. 8.	H. 4.	Eng. bot. 205.	
5 intermédia. DC.	(re.) sea-side.	S. Europe.	1820.	6. 8.	H. 4.	Mich. gen. t. 22. f. 2.	
<i>hirsuta.</i> var. <i>cal��brica.</i> L.							
6 hispida. DC.	(re.) hairy.	Siberia.	1789.	—	H. 4.	Flor. gr��c. t. 313.	
<i>hirsuta.</i> var. <i>cr��tica.</i> L.							
7 ericifolia. DC.	(re.) Heath-leaved.	Canaries.	1815.	—	G. h.		
8 pauciflora. DC.	(ro.) few-flowered.	N. Holland.	1825.	8. 10.	G. h.		

ORDO XXVI.

CARYOPHYLLEÆ. DC. prodr. 1. p. 351.

Tribus I. SILENEÆ. DC.

GYPSO'PHILA. DC.	GYPSO'PHILA.	Decandria Digynia. L.					
1 Struthium. DC.	(wh.) fleshy-leaved.	Spain.	1729.	7. 8.	H. 4.	Barr. ic. 64. t. 119.	
2 fastigiata. DC.	(wh.) angular-leaved.	Europe.	1759.	6. 7.	H. 4.	Gmel. sib. 4. t. 61. f. 1.	
3 dichotoma. DC.	(wh.) forked.	Galicia.	1826.	6. 8.	H. 4.		
4 arenaria. DC.	(wh.) sand.	Hungary.	1801.	7. 8.	H. 4.	W. et. K. hung. t. 41.	
5 viscosa. DC.	(bh.) clammy.	Levant.	1773.	6. 8.	H. 4.	Mur. c. got. 1783. t. 3.	
6 altissima. DC.	(wh.) upright.	Siberia.	1759.	7. 8.	H. 4.	Gmel. sib. 4. t. 60.	
7 latifolia. F.	(wh.) broad-leaved.	_____	1824.	6. 9.	H. 4.		

8 <i>perfoliata</i> . DC. (wh.) perfoliate.	Spain.	1732.	6. 9.	H. ȝ.	Dill. elt. t. 276.	
β <i>tomentosa</i> . DC. (wh.) woolly.	Tauria.	—	—	H. ȝ.	Barr. ic. t. 1002.	
9 <i>scorzonerifolia</i> . DC. (w.) <i>Scorzonera</i> -ld.	Caucasus.	1823.	—	H. ȝ.		
10 <i>acutifolia</i> . DC. (wh.) sharp-leaved.	Siberia.	1821.	—	H. ȝ.		
11 <i>paniculata</i> . DC. (wh.) paniced.	Levant.	1759.	6. 7.	H. ȝ.	Jacq. aust. 5. t. app. 1	
12 <i>adscendens</i> . DC. (wh.) ascending.	Tauria.	1819.	—	H. ȝ.	Jacq. h. vind. 2. t. 138.	
13 <i>elegans</i> . DC. (bh.) elegant.	Iberia.	1820.	6. 8.	H. ⓧ.	Swt. br. fl. gar.	
14 <i>Stevéni</i> . DC. (wh.) Steven's.	Caucasus.	1817.	—	H. ȝ.		
15 <i>tenuifolia</i> . DC. (ro.) slender-leaved.	Pyrenees.	1824.	—	H. ȝ.		
16 <i>rēpens</i> . DC. (wh.) creeping.	1774.	7. 9.	H. ȝ.	Bot. mag. 1448.	
17 <i>dubia</i> . DC. (wh.) doubtful.	1815.	5. 9.	H. ȝ.		
18 <i>prostrata</i> . DC. (bh.) trailing.	Siberia.	1759.	7. 9.	H. ȝ.	Bot. mag. 1281.	
19 <i>murālis</i> . DC. (li.) wall.	Europe.	1739.	6. 10.	H. ⓧ.	Flor. græc. t. 381.	
20 <i>serotina</i> . W. en. (wh.) late-flowering.	Magdeburg.	1816.	—	H. ⓧ.		
21 <i>glomerata</i> . DC. (wh.) round-headed.	Tauria.	1804.	6. 7.	H. ȝ.		
22 <i>saxifraga</i> . DC. (bh.) small.	Germany.	1774.	7. 8.	H. ȝ.	Exot. bot. 2. t. 90.	
23 <i>rígida</i> . L. (li.) rigid.	France.	1769.	6. 8.	H. ȝ.	Flor. græc. t. 382.	
DIA'NTHUS. DC.	PINK. Decandria Digynia.	L.				
1 <i>prolifer</i> . DC. (ro.) proliferous.	England.	7. 8.	H. ⓧ.	Eng. bot. 956.	
2 <i>diminūtus</i> . W. (ro.) small-flowered.	S. Europe.	1771.	—	H. ⓧ.		
3 <i>Armèria</i> . DC. (re.) Deptford.	England.	7. 9.	H. ⓧ.	Eng. bot. 317.	
4 <i>pseudo-armèria</i> . (re.) canescens.	Tauria.	1820.	—	H. ȝ.	Bot. mag. 2288.	
5 <i>armerioïdes</i> . DC. (re.) Thrift-like.	New Jersey.	—	—	H. ⓧ.		
6 <i>barbatus</i> . DC. (va.) Sweet-William.	Germany.	1573.	6. 7.	H. ȝ.	Bot. mag. 207. 205	
β <i>flōre plēno</i> .	(va.) double.	—	—	H. ȝ.		
7 <i>latifolius</i> . DC. (cr.) broad-leaved.	5. 11.	H. ȝ.	Swt. br. fl. gar. t. 2.	
8 <i>aggregatū</i> . DC. (re.) clustered.	1820.	6. 8.	H. ȝ.	Thunb. jap. t. 23.	
9 <i>japonicus</i> . DC. (re.) Japanese.	China.	1804.	6. 10.	F. ȝ.		
10 <i>cephalotes</i> . DC. (re.) close-headed.	6. 9.	H. ȝ.		
11 <i>capitatus</i> . DC. (re.) headed.	Tauria.	1794.	6. 7.	H. ȝ.		
12 <i>polymorphus</i> . DC. (re.) variable.	1820.	6. 8.	H. ȝ.		
13 <i>diutinus</i> . L.en. (re.) pedicled.	Hungary.	—	—	H. ȝ.	Swt. br. fl. gar.	
14 <i>Balbisi</i> . DC. (re.) Balbis's.	Genoa.	1827.	7. 10.	H. ȝ.	Swt. br. fl. gar.	
<i>glaucophyllus</i> . H.						
15 <i>giganteus</i> . DC. (re.) gigantic.	Bulgaria.	—	6. 10.	H. ȝ.	Swt. br. fl. gar. 288.	
16 <i>ferrugineus</i> . DC. (st.) rusty.	Pyrenees.	1756.	7. 9.	H. ȝ.	Barr. ic. t. 497.	
17 <i>Carthusianorum</i> . DC. (re.) <i>Carthusian</i> .	Europe.	1573.	—	H. ȝ.	Flor. græc. t. 392.	
18 <i>afrorūbens</i> . All. (re.) dark-red.	Italy.	1802.	—	H. ȝ.	Jacq. ic. 3. t. 467.	
19 <i>nānus</i> .	(cr.) dwarf.	Switzerland.	1820.	—	H. ȝ.	
<i>Carthusianorum</i> γ <i>nanus</i> . DC.						
20 <i>arboreus</i> . DC. (re.) arborescent.	Crete.	—	—	F. ⚡.	Flor. græc. t. 406.	
21 <i>juniperinus</i> . DC. (re.) Juniper-leaved.	Greece.	1825.	—	F. ⚡.		
22 <i>fruticosus</i> . DC. (re.) frutescent.	Crete.	1815.	6. 9.	F. ⚡.	Flor. græc. t. 407.	
23 <i>suffruticosus</i> . DC. (re.) shrubby.	1804.	6. 7.	F. ⚡.		
24 <i>rupicola</i> . DC. (ro.) Sicilian.	Sicily.	1825.	6. 8.	H. ȝ.	Ten. fl. napol. t. 39.	
<i>Bisigniāni</i> . T.						
25 <i>carolinianus</i> . DC. (re.) Carolina.	N. America.	1811.	6. 9.	F. ȝ.		
26 <i>caryophylloïdes</i> . DC. (re.) Clove-like.	1827.	—	H. ȝ.		
27 <i>asper</i> . W. en. (re.) rough-stalked.	Switzerland.	1802.	7. 8.	H. ȝ.	Parad. lond. 62.	
28 <i>collinus</i> . W. en. (re.) hill.	Hungary.	1800.	7. 9.	H. ȝ.		
29 <i>umbellatus</i> . DC. (re.) umbelled.	1825.	—	H. ȝ.		
30 <i>campestris</i> . DC. (re.) field.	Tauria.	1815.	7. 8.	H. ȝ.	Bot. mag. 1876.	
31 <i>alpēstris</i> . DC. (re.) pasture.	1820.	—	H. ȝ.	Balb. act. taur. 7. t. 1.	
32 <i>nítidus</i> . DC. (re.) glossy.	Hungary.	1821.	6. 8.	H. ȝ.	W. et K. hung. 2. t. 191.	
33 <i>hírtus</i> . DC. (re.) hairy.	Dauphiny.	—	—	H. ȝ.	Vill. delph. 3. t. 46.	
34 <i>guttatus</i> . DC. (pu.) spotted.	Caucasus.	1816.	7. 9.	H. ȝ.		
35 <i>versicolor</i> . DC. (s.r.) various-colour'd	Russia.	1821.	—	H. ȝ.		
36 <i>pratensis</i> . DC. (st.) meadow.	Tauria.	—	6. 8.	H. ȝ.		
<i>ochroleucus</i> . L.en. non P. S. <i>tataricus</i> . F.						
37 <i>chinensis</i> . DC. (va.) China.	China.	1713.	7. 9.	H. ⚢.	Bot. mag. 28.	
β <i>flōre-plēno</i> .	(va.) double-flowered.	—	—	H. ⚢.		
38 <i>montanus</i> . DC. (pu.) mountain.	Caucasus.	1803.	6. 9.	H. ȝ.	Bot. mag. t. 1162.	
<i>discolor</i> . B.M.						
39 <i>ochroleucus</i> . DC. (st.) straw-coloured.	Levant.	1821.	—	H. ȝ.		
40 <i>Caryophyllus</i> . DC. (va.) Clove.	England.	6. 8.	H. ȝ.	Eng. bot. 214.	
β <i>multiplex</i> ,	(cr.) double Clove.	—	—	H. ȝ.		
γ <i>ruber-cinctus</i> .	(wh.) <i>Hogg's Beauty of Middlesex Picotee</i> .	—	—	H. ȝ.	Swt. flor. guid. 70.	
δ <i>inimitabilis</i> .	(w.c.) <i>Hird's Inimitable Picotee</i> .	—	—	H. ȝ.	— t. 80.	
ε <i>striatulus</i> .	(y.r.) <i>Rosalie de Rohan Picotee</i> .	—	—	H. ȝ.	— t. 25.	

CARYOPHYLLEÆ.

ζ purpùreο-cinctus. (y.p.)	Erasmus Picotee.	6. 8.	H. 2.	Swt. flor. guid. 59.
η Clíntòniae. (r.w.)	Hufton's Dutchess of Newcastle Carnation.	H. 2.	t. 14.	
θ triūmphans. (p.w.)	Pardo's Ace of Trumps Carnation.	H. 2.	t. 36.	
ι carùleo-purpùreus. (p.w.)	Cordon Bleu Carnation.	H. 2.	t. 64.	
κ Franklini. (r.p.w.)	Franklin's Queen of Hearts Carnation.	H. 2.	t. 20.	
λ pùellulus. (c.r.w.)	Pucelle de Gant Carnation.	H. 2.	t. 67.	
μ Leopoldi. (c.p.w.)	Ives's Prince Leopold Carnation.	H. 2.	t. 31.	
ν iricolor. (c.p.w.)	Cartwright's Rainbow Carnation.	H. 2.	t. 9.	
ξ Bacchus. (s.p.w.)	Davey's Bacchus Carnation.	H. 2.	t. 44.	
ο Tarrara. (s.p.w.)	Hall's Tarrara Carnation.	H. 2.	t. 21.	
π pulchella. (s.p.w.)	Franklin's Tartar Carnation.	H. 2.		
ρ fruticòsus. (va.)	tree-Carnation.	H. 2.		
ξ imbricatus. (d.r.)	Wheat-ear.	H. 2.		
41 sylvétris. (re.)	short-scaled.	Jurassa.	6. 7.	H. 2.
virgíneus. B.M. 1740, non L.					
42 longicáulis. DC.	(re.) long-stalked.	Naples.	1824.	—	H. 2.
43 monadéphus. DC. (wh.)	procumbent.	Levant.	—	H. 2.
procúmbens. P.S.					Vent. h. cels. t. 39.
44 suavéolens. DC.	(sp.) sweet-scented.	1823.	—	H. 2.
45 Poiretiànum. DC. (pu.)	Poiret's.	—	—	H. 2.
β flòre-plèno.	(pu.) double-flowering.	—	—	H. 2.
46 Liboschitziànum. DC.	Armenian.	Armenia.	1823.	6. 8.	H. 2.
petr'æus. M.B. non DC.					
47 sylváticus. DC.	(cr.) wood.	Ratisbon.	1815.	—	H. 2.
48 arbúscula. B.R.	(re.) China tree.	China.	1824.	7. 10.	H. 2.
49 pomeridiànum. DC. (st.)	afternoon.	Palestine.	1804.	—	H. 2.
50 leptopétalus. DC. (wh.)	narrow-petaled.	Caucasus.	1814.	—	H. 2.
51 pungens. DC.	(wh.) pungent.	Spain.	1781.	8. 10.	H. 2.
52 bicolor. DC.	(w.p.) two-coloured.	Tauria.	1818.	6. 8.	H. 2.
53 furcátus. DC.	(fl.) forked.	Piedmont.	1820.	—	H. 2.
54 virgíneus. DC.	(re.) virgin.	Montpelier.	1732.	—	H. 2.
55 deltóides. DC.	(re.) maiden.	Britain.	6. 10.	H. 2.
56 gláucus. L.	(wh.) glaucous.	—	—	—	H. 2.
57 crenátus. B.R.	(wh.) long-cupped.	C. B. S.	1817.	8.	G. 2.
58 rígidus. DC.	(pa.) rigid.	Tauria.	1802.	6. 10.	H. 2.
59 clavátus. DC.	(fl.) clubbed.	—	1824.	6. 8.	H. 2.
60 suávis. DC.	(fl.) sweet.	—	1820.	6. 7.	H. 2.
61 répens. DC.	(re.) creeping.	Siberia.	1825.	—	H. 2.
62 cæ'sius. DC.	(fl.) tufted.	Britain.	—	—	H. 2.
63 alpinus. DC.	(re.) Alpine.	Austria.	1759.	—	H. 2.
64 glacialis. DC.	(re.) acute-leaved.	France.	1820.	—	H. 2.
65 elegans. DC.	(re.) elegant.	Levant.	1825.	—	F. 2.
66 multipunctátus. DC.	(sp.) spotted.	—	—	6. 9.	F. 2.
67 Hornemánni. DC.	(fl.) Horneman's.	Italy.	1822.	6. 7.	H. 2.
68 Sternbergii. DC.	(re.) Sternberg's.	—	1825.	—	F. 2.
69 petr'æus. DC.	(wh.) rock.	Hungary.	1804.	7. 8.	H. 2.
β Simsii.	(wh.) Sims's.	—	—	—	H. 2.
70 ibéricus. DC.	(pu.) Iberian.	Iberia.	1820.	6. 8.	H. 2.
Willdenowii. L.en.					
71 ruthénicus. DC.	(re.) Russian.	Russia.	—	—	H. 2.
72 dentósus. F.	(re.) toothed.	Siberia.	1826.	—	H. 2.
73 Buchtménsis. F.	(re.) Buchtmorian.	Russia.	—	—	H. 2.
74 gálico. DC.	(wh.) French.	France.	—	—	H. 2.
75 albens. DC.	(wh.) white Cape.	C. B. S.	1787.	8.	G. 2.
76 serrulátus. DC.	(ro.) serrulate.	Tunis.	1826.	—	H. 2.
77 pulchéllus. P.S.	(re.) pretty.	—	—	—	H. 2.
78 dúbios. H.	(w.p.) doubtful.	—	—	5. 6.	H. 2.
præcox. W.en.					
79 plumárius. DC.	(wh.) common.	Europe.	1629.	6. 8.	H. 2.
β heròicus.	(w.p.) Styles's Hero Pink.	—	—	H. 2.
γ Cheeséanus.	(w.p.) Cheese's Miss Cheese.	—	—	H. 2.
δ Houseanus.	(w.p.) House's Woodland Beauty.	—	—	H. 2.
ε Uxbridgeanus.	(w.r.) Dry's Earl of Uxbridge	—	—	H. 2.
ζ Aclánðæ.	(w.p.) Knight's Lady Acland	—	—	H. 2.
η incl'yus.	(w.p.) Smith's Champion	—	—	H. 2.
θ Barrátti.	(w.p.) Barratt's Conqueror	—	—	H. 2.
ι regális.	(w.p.) Davey's Roi de pourpre	—	—	H. 2.
80 adulterínus.	(f.p.) hybrid.	Hybrid.	—	—	H. 2.
β Juliétti.	(f.p.) Davey's Juliet.	—	—	H. 2.

Swt. flor. guid. 17.

t. 13.

t. 47.

t. 55.

t. 40.

t. 27.

t. 32.

t. 33.

Swt. flor. guid. 8.

81 Penry'nae.	(pu.) Lady Penryns. Hybrid.	6. 8. H. ¼.	
β purpureus.	(pu.) Davey's Lady Penryns.—	— H. ¼.	Swt. flor. guid. 41.
82 hortensis. W.en.	(wh.) garden.	Europe.	1805. 6. 8. H. ¼.	
83 caucaseus. DC.	(re.) Caucasian.	Caucasus.	1803. 6. 9. H. ¼.	Bot. mag. 795.
84 saxatilis. DC.	(wh.) rock.	France.	— H. ¼.
85 squarrosum. DC.	(wh.) squarrose.	Tauria.	1823. 6. 8. H. ¼.	Bieb. cent. 1. t. 33.
86 Mussini. DC.	(wh.) Mussin's.	Caucasus.	1825. — H. ¼.	
87 frágans. B.M.	(wh.) fragrant.	1804. 6. 7. H. ¼.	Bot. mag. 2067.
88 punctatus. DC.	(sp.) spot-flowered.	1825. — H. ¼.	Lodd. bot. cab. 896.
89 serotinus. DC.	(wh.) late-flowering.	Hungary.	1804. 7. 8. H. ¼.	W. et K. hung. 2.t.172.
90 arenarius. DC.	(wh.) sand.	N. Europe. 8. H. ¼.	Bot. mag. 2038.
91 prostratus. DC.	(re.) prostrate.	C. B. S. 8. 9. G. ¼.	Jacq. scheenb. 3.t. 271.
92 fimbriatus. DC.	(re.) fringed.	Iberia.	1802. 6. 8. H. ½.	Bot. mag. 1069.
93 monspessulanus. DC.	(re.) Montpelier.	Pyrenees.	1764. 7. 8. H. ¼.	
94 erubescens. Tr.	(bh.) blush-flowered.	1825. — H. ¼.	
95 superbus. DC.	(re.) superb.	Europe.	1596. 7. 9. H. ¼.	Bot. mag. 297.
96 Fischeri. DC.	(re.) Fischer's.	Russia.	1826. 6. 8. H. ¼.	Swt. br. fl. gar. 245.
VACCARIA. Dod.	VACCARIA. Decandria Digynia.			Flor. græc. t. 380.
1 sessilifolia.	(re.) sessile-leaved.	Europe.	1596. 7. 8. H. ⊙.	
Saponaria Vaccaria.	B.M. 2290. Gypsophila Vaccaria. F.G.			
2 perfoliata.	(re.) perforiate.	E. Indies.	1820. — H. ⊙.	
Saponaria perforata. DC.				
SAPONARIA. DC.	SOAPWORT.	Decandria Digynia. L.		
1 officinalis. DC.	(bh.) common.	England. 7. 10. H. ¼.	Eng. bot. t. 1060.
β flòre pleno.	(bh.) double-flowered.	— H. ¼.	
2 ocyroides. DC.	(ro.) Basil-leaved.	Europe.	1768. 5. 7. H. ¼.	Bot. mag. 154.
3 glutinosa. DC.	(re.) small-flowered.	Tauria.	1823. — H. ¾.	— 2855.
4 pòrigens. DC.	(bh.) hairy.	Levant.	1680. 7. 8. H. ⊙.	Jacq. vind. 2. t. 109.
5 orientalis. DC.	(re.) small annual.	1732. 6. 8. H. ⊙.	Dill. elt. t. 167. f. 204.
6 deprissa. DC.	(ro.) depressed.	Sicily.	1828. — H. ¼.	Biv. Bon. t. 163. f. 1.
7 cæspitosa. DC.	(ro.) tufted.	Pyrenees.	1827. — H. ¼.	
8 Smithii. DC.	(re.) Smith's.	Levant.	1828. — F. ¼.	Flor. græc. t. 389.
cæspitosa. F.G. non DC.				
9 lutea. DC.	(ye.) yellow.	Switzerland.	1804. — H. ¼.	All. ped. t. 23. f. 1.
10 bellidifolia. DC.	(ye.) Daisy-leaved.	Italy.	— H. ¼.	Smith spic. t. 5.
CUCUBALUS. DC.	CAMPION.	Decandria Trigynia. L.		
bacciferus. DC.	(gr.) berry-bearing.	England. 6. 7. H. ¼.	Eng. bot. 1577.
SILENE. DC.	CATCHFLY.	Decandria Trigynia. L.		
1 acáulis. DC.	(re.) stemless.	Britain. 6. 8. H. ¼.	Eng. bot. 1081.
β alba. DC.	(wh.) white-flowered.	— H. ¼.	
2 pumilio. DC.	(pu.) dense.	Germany.	1823. — H. ¼.	Stur.deut.1.fas.22.t.11.
3 fimbriata. DC.	(wh.) fringe-flower'd.	Crete.	1803. 5. 8. H. ¼.	Bot. mag. 908.
4 laceria. DC.	(wh.) jagged-flower'd.	Caucasus.	1818. — H. ¼.	— 2255.
5stellata. DC.	(wh.) four-leaved.	N. America.	1696. 6. 8. H. ¼.	— 1107.
6 inflata. DC.	(wh.) bladder.	Britain. 5. 9. H. ¼.	Eng. bot. 164.
Cucubalus Béhen. E.B.				
7 linearis.	(wh.) linear-leaved.	Naples.	1820. — H. ¼.	Ten. flor. nap. t. 37.
Cucubalus angustifolius. T.N.				
8 Willdenowii.	(wh.) Willdenow's.	S. Europe.	— — H. ¼.	
Cucubalus glaucus. W.en. supp.				
9 marítima. E.B.	(wh.) sea.	Britain. 8. 9. H. ¼.	Eng. bot. 957.
10 fabaria. H.K.	(wh.) thick-leaved.	Sicily.	1731. 6. 8. F. ¼.	Flor. græc. t. 415.
11 Béhen. DC.	(bh.) Cretan.	Crete.	1713. 6. 7. H. ⊙.	— t. 416.
12 indica. DC.	(pu.) Indian.	Nepaul.	1824. — F. ¼.	
13 viscinoides. DC.	(wh.) Viscago-like.	Dahuria.	1826. — H. ¼.	
14 procumbens. DC.	(wh.) procumbent.	Siberia.	1820. — H. ¼.	Murr. c. gæt. p. 83. t.2.
15 rubella. DC.	(re.) small red.	Portugal.	1732. 5. 6. H. ⊙.	Flor. græc. t. 426.
16 apetala. DC.	(gr.) petalless.	Spain.	1801. 6. 7. H. ⊙.	
17 spergulifolia. DC.	(wh.) Spurry-like.	Armenia.	1824. — F. ¼.	Desf. cor. Tourn. t. 55.
18 Gypsophila. DC.	(wh.) Gypsophilà-like.	— H. ¼.	
19 angustifolia. DC.	(wh.) narrow-leaved.	Caucasus.	1824. — H. ¼.	
20 Otites. DC.	(st.) Spanish.	England. 7. 8. H. ¼.	Eng. bot. 85.
Cucubalus Otites. E.B.				
21 wolgensis. DC.	(wh.) Wolga.	Wolga.	1821. 6. 8. H. ¼.	
22 parviflora. DC.	(st.) small-flowered.	Hungary.	1796. — H. ¼.	
23 effusa. DC.	(st.) spreading.	Wolga.	1821. — H. ¼.	
24 sibirica. DC.	(wh.) Siberian.	Siberia.	1773. — H. ¼.	
25 multiflora. DC.	(wh.) many-flowered.	Hungary.	1794. — H. ¾.	Hall. g. 1. p. 150. ic.
26 ruthénica. DC.	(wh.) Russian.	Russia.	1827. — H. ¼.	W. et K. hung. 1. t. 56.

CARYOPHYLLEÆ.

27	staticifolia. DC.	(wh.)	Statice-leaved.	Levant.	1828.	6. 8.	H. ȝ.	Flor. græc. t. 434.
28	tatarica. DC.	(wh.)	Tartarian.	Tartary.	1769.	—	H. ȝ.	Flor. græc. t. 432.
29	gigantea. DC.	(wh.)	gigantic.	Africa.	1738.	—	G. ȝ.	Tourn. it. 2. p. 361. ic.
30	viscosa. DC.	(wh.)	clammy.	Levant.	1739.	7.	H. ♂.	Eng. bot. 922.
31	cónica. DC.	(fl.)	corn.	England.	6. 7.	H. Ⓛ.	Clus. his. I. p. 288. f. 2.
32	conóidea. DC.	(pu.)	conoid.	S. Europe.	1683.	—	H. Ⓛ.	Eng. bot. 1178.
33	undulata. DC.	(re.)	wave-leaved.	C. B. S.	1775.	8.	G. ♂.	Dill. elt. t. 311. f. 401.
34	ánglica. DC.	(wh.)	English.	England.	6. 7.	H. Ⓛ.	Vail. par. t. 16. f. 12.
35	lusitánica. DC.	(fl.)	Portugal.	Portugal.	1732.	—	H. Ⓛ.	Schrank. pl. rar. t. 39.
36	tridentata. DC.	(ro.)	three-toothed.	Spain.	1824.	—	H. Ⓛ.	Flor. græc. t. 412.
37	gállica. DC.	(fl.)	French.	France.	1683.	5. 6.	H. Ⓛ.	Eng. bot. 86.
38	ocymoïdes. DC.	(pu.)	Basil-like.	1819.	6. 7.	H. Ⓛ.	Flor. græc. t. 408.
39	dísticha. DC.	(ro.)	two-ranked.	1817.	—	H. Ⓛ.	Magn. h. mons. 171. ic.
40	cerastoïdes. DC.	(re.)	Cerastium-like.	S. Europe.	1732.	7. 8.	H. Ⓛ.	Ten. fl. nap. 1. t. 39.
41	quinquevulnera. DC.	(p.w.)	variegated.	England.	6. 8.	H. Ⓛ.	Flor. græc. t. 409.
42	nocturna. DC.	(w.re.)	spiked.	S. Europe.	1683.	—	H. Ⓛ.	Flor. græc. t. 414.
43	reflexa. H.K.	(wh.)	reflexed.	1726.	—	H. Ⓛ.	Flor. græc. t. 413.
44	mutabilis. H.K.	(bh.)	changeable.	1688.	—	H. Ⓛ.	Jacq. vind. 3. t. 81.
45	micropétala. DC.	(re.)	small-petaled.	Chili.	1822.	6. 7.	H. Ⓛ.	Swt. br. fl. gar. 58.
46	micránta. DC.	(re.)	small-blossomed.	Portugal.	1825.	—	H. Ⓛ.	Bot. mag. 114.
47	hirsuta. Lag.	(re.)	hairy.	1824.	—	H. Ⓛ.	Dill. elt. t. 313. f. 403.
	<i>sabuletorum</i> . Lk. <i>hirsutissimum</i> . DC.							Jacq. aust. 2. t. 120.
48	canescens. DC.	(re.)	canescent.	Naples.	1823.	6. 8.	H. ȝ.	W. et K. hung. 3. t. 212.
49	decumbens. DC.	(wh.)	decumbent.	S. Europe.	1828.	—	H. Ⓛ.	Jacq. aust. 1. t. 96.
50	diffusa. DC.	(fl.)	diffuse.	Italy.	1825.	—	H. ȝ.	Flor. dan. 4.
51	divaricata. F.G.	(wh.)	divaricate.	Levant.	1827.	—	H. ♂.	Flor. græc. t. 420.
	<i>racemosa</i> . DC.							Jacq. col. sup. t. 3. f. 3.
52	dichotoma. DC.	(wh.)	dichotomous.	Hungary.	1791.	—	H. Ⓛ.	W. et K. hung. 2. t. 175.
53	nyctantha. DC.	(st.)	various-leaved.	1815.	6. 8.	H. Ⓛ.	Dill. elt. t. 311. f. 403.
54	trinervia. DC.	(re.)	three-nerved.	Rome.	1821.	—	H. Ⓛ.	Flor. græc. t. 422.
55	bellidifolia. DC.	(re.)	Daisy-leaved.	1794.	—	H. Ⓛ.	Flor. græc. t. 425.
56	vespertina. DC.	(re.)	evening.	Barbary.	1796.	7. 8.	H. Ⓛ.	Lodd. bot. cab. 454.
	<i>bipartita</i> . Desf. atl. 1. t. 100.							W. et K. hung. 3. t. 164.
57	obtusifolia. DC.	(re.)	blunt-leaved.	1823.	6. 8.	H. Ⓛ.	All. auct. p. 28. t. 1. f. 3.
58	colorata. DC.	(fl.)	coloured.	1804.	—	H. Ⓛ.	Flor. græc. t. 419.
59	crassifolia. DC.	(br.)	thick-leaved.	C. B. S.	1774.	7. 8.	G. ♂.	Eng. bot. 465.
60	gracilis. DC.	(wh.)	slender.	1823.	6. 8.	H. Ⓛ.	W. et K. hung. 3. t. 213.
61	jeniseensis. DC.	(st.)	two-coloured.	Siberia.	1817.	6. 7.	H. ȝ.	Jacq. aust. 2. t. 120.
62	ciliata. DC.	(pu.)	ciliated.	Crete.	1804.	—	H. Ⓛ.	Flor. græc. t. 419.
63	pendula. DC.	(re.)	pendulous.	Sicily.	1731.	5. 7.	H. Ⓛ.	Eng. bot. 466.
64	longicaulis. DC.	(wh.)	long-stalked.	Spain.	1818.	6. 7.	H. Ⓛ.	Flor. græc. t. 420.
65	secundiflora. DC.	(wh.)	side-flowered.	—	—	H. Ⓛ.	Jacq. aust. 1. t. 96.
66	viscosissima. DC.	(wh.)	viscous.	Naples.	1822.	—	H. Ⓛ.	Flor. græc. t. 421.
67	quadridentata. DC.	(p.w.)	four-toothed.	Italy.	1783.	5. 6.	H. Ⓛ.	Flor. græc. t. 422.
	<i>Lychnis quadridentata</i> . w.							Flor. græc. t. 425.
68	pusilla. W.en.	(wh.)	dwarf.	Hungary.	1804.	6. 7.	H. ȝ.	Lodd. bot. cab. 455.
69	alpéstris. DC.	(wh.)	Austrian.	Austria.	1774.	5. 7.	H. ȝ.	W. et K. hung. 2. t. 164.
70	rupéstris. DC.	(re.)	rock.	Switzerland.	—	H. ȝ.	All. auct. p. 28. t. 1. f. 3.
71	glaucofolia. DC.	(bh.)	glaucous-leav'd.	Spain.	1823.	—	H. ȝ.	Flor. græc. t. 423.
72	inaperta. DC.	(st.)	unopen-flow'r'd.	S. Europe.	1732.	6. 7.	H. ȝ.	Dill. elt. t. 311. f. 403.
73	clandéstina. DC.	(w.p.)	hidden-flower'd.	C. B. S.	1801.	—	H. Ⓛ.	Flor. græc. t. 424.
74	portensis. W.	(pu.)	Oporto.	Portugal.	1759.	7. 8.	H. Ⓛ.	W. et K. hung. 3. t. 213.
75	antirrhina. DC.	(wh.)	Snap-dragon.	Virginia.	1732.	6. 7.	H. Ⓛ.	Jacq. aust. 2. t. 120.
76	geminiflora. DC.	(pu.)	twin-flowered.	1816.	—	H. Ⓛ.	Flor. græc. t. 425.
77	flavescens. DC.	(st.)	yellowish.	Hungary.	1804.	—	H. ȝ.	Flor. græc. t. 426.
78	linoides.	(st.)	Flax-like.	1817.	7. 8.	H. Ⓛ.	Flor. græc. t. 427.
	<i>Linaria</i> . DC. non Flor. græc.							Flor. græc. t. 428.
79	linifolia. F.G.	(fl.)	Flax-leaved.	Levant.	1828.	6. 8.	H. ȝ.	Flor. græc. t. 429.
80	crética. DC.	(re.)	Cadian.	Candia.	1732.	5. 8.	H. Ⓛ.	Flor. græc. t. 430.
81	sedoides. DC.	(re.)	Sedum-like.	Crete.	1804.	6. 8.	H. Ⓛ.	Flor. græc. t. 431.
82	sícula. Cyr.	(ro.)	Sicilian.	S. Europe.	1825.	—	H. Ⓛ.	Flor. græc. t. 432.
	<i>divaricata</i> . DC. non Flor. græc.							Flor. græc. t. 433.
83	Saxifraga. DC.	(w.br.)	Saxifrage.	S. Europe.	1640.	—	H. ȝ.	Flor. græc. t. 434.
84	petræa. DC.	(wh.)	fine-toothed.	Hungary.	1806.	—	H. ȝ.	Flor. græc. t. 435.
85	campánula. DC.	(wh.)	bell-flowered.	Piedmont.	1822.	—	H. ȝ.	Flor. græc. t. 436.
86	longipétala. DC.	(g.st.)	long-petaled.	Aleppo.	—	—	H. Ⓛ.	Flor. græc. t. 437.
87	nútans. E.B.	(wh.)	Nottingham.	England.	6. 7.	H. ȝ.	Flor. græc. t. 438.
88	infícta. W.K.	(wh.)	smooth.	Hungary.	1816.	—	H. ȝ.	Flor. græc. t. 439.
89	latifolia. H.	(wh.)	broad-leaved.	Europe.	1825.	—	H. ȝ.	Flor. græc. t. 440.

90 <i>saxátilis</i> . B.M. (st.) stone.	Siberia.	1800.	6. 7.	H. ¼.	Bot. mag. 689.
91 <i>lívida</i> . W. en.(w.br.) livid.	Carniola.	1816.	—	H. ¼.	
92 <i>tenuis</i> . DC. (st.) slender.	Siberia.	—	—	H. ¼.	
93 <i>quadrífida</i> . DC. (wh.) four-cleft.	Italy.	1820.	—	H. ¼.	
94 <i>viridiflóra</i> . DC. (gr.) green-flowered.	Portugal.	1739.	—	H. ♂.	Herm. parad. t. 199.
95 <i>chlorántha</i> . DC. (st.) pale-flowered.	Germany.	1732.	6. 8.	H. ¼.	Dill. elt. t. 316. f. 408.
96 <i>psammítis</i> . DC. (st.) sulphur-colour'd.	—	1820.	—	H. ¼.	All. ped. t. 44. f. 2.
97 <i>nicaéensis</i> . DC. (w.p.) villous-clammy.	S. Europe.	—	—	H. ⊙.	Jacq. vind. 1. t. 59.
98 <i>cathólica</i> . DC. (wh.) panicled.	Italy.	1711.	7. 9.	H. ¼.	
99 <i>Mussini</i> . H. (wh.) Mussin's.	Russia.	1824.	—	H. ¼.	
100 <i>diversifólia</i> . DC. (wh.) various-leaved.	—	1820.	—	H. ⊙.	
101 <i>répens</i> . DC. (wh.) creeping.	Siberia.	1806.	7. 8.	H. ¼.	
102 <i>glauca</i> . DC. (pu.) glaucous-leav'd.	—	1824.	—	H. ⊙.	
103 <i>tenuifólia</i> . DC. (pu.) slender-leaved.	Dahuria.	1820.	—	H. ⊙.	
104 <i>virgínica</i> . DC. (re.) Virginian.	Virginia.	1783.	5. 8.	H. ¼.	Pluk. alm. t. 203. f. 1.
105 <i>Catesbæi</i> . DC. (cr.) Catesby's.	Carolina.	—	—	H. ¼.	Catesb. car. 54. ic.
106 <i>strícta</i> . DC. (re.) upright.	Spain.	1802.	6. 7.	H. ⊙.	
107 <i>Muscípula</i> . DC. (re.) Spanish.	—	1596.	7. 8.	H. ⊙.	Clus. hist. I. p. 289. f. 1.
108 <i>noctiflóra</i> . DC. (fl.) night-flowering.	England.	—	7.	H. ⊙.	Eng. bot. 291.
109 <i>ornáta</i> . DC. (cr.) dark-coloured.	C. B. S.	1775.	5. 9.	G. ♂.	Bot. mag. 382.
110 <i>córsica</i> . DC. (st.) Corsican.	Corsica.	1825.	6. 8.	H. ¼.	Bocc. mus. t. 54.
111 <i>egyptiaca</i> . DC. (fl.) Egyptian.	Egypt.	1800.	7. 8.	G. ♂.	
112 <i>sericea</i> . DC. (fl.) silky.	Corsica.	1801.	6. 8.	H. ⊙.	All. ped. t. 79. f. 3.
113 <i>pícta</i> . DC. (w.p.) painted.	Mt. Carmel.	1818.	7. 9.	H. ⊙.	Swt. br. fl. gar. 92.
114 <i>bicolor</i> . DC. (w.p.) two-coloured.	France.	—	6. 8.	H. ⊙.	DC. ic. gall. rar. t. 42.
115 <i>lævigátâ</i> . F.G. (re.) smooth.	Greece.	1828.	7. 9.	H. ⊙.	Flor. græc. t. 418.
116 <i>reticulátâ</i> . DC. (re.) netted-flow'r'd.	Barbary.	1804.	7. 8.	H. ⊙.	Desf. atl. 1. t. 99.
117 <i>pensylvánica</i> . DC. (re.) Pensylvanian.	N. America.	1806.	6. 9.	H. ¼.	Bot. reg. 247.
<i>incarnátâ</i> . Bot. cab. 41.					
118 <i>vallésia</i> . DC. (pu.) hard-rooted.	Switzerland.	1765.	6. 8.	H. ¼.	All. ped. t. 23. f. 2.
119 <i>fruticósa</i> . DC. (fl.) shrubby.	Sicily.	1629.	6. 7.	F. ♂.	Flor. græc. t. 428.
120 <i>cáspica</i> . DC. (st.) Caspian.	Caucasus.	1823.	—	H. ♂.	
121 <i>am. cénâ</i> . DC. (wh.) Tartarian.	Tartary.	1779.	7.	H. ¼.	Bot. mag. 1997.
122 <i>supina</i> . DC. (wh.) trailing.	Caucasus.	1804.	6. 8.	H. ¼.	
123 <i>depréssâ</i> . DC. (wh.) depressed.	Iberia.	1823.	—	H. ¼.	Jacq. vind. 3. t. 84.
124 <i>paradoxa</i> . DC. (wh.) Dover.	England.	—	7.	H. ¼.	Bot. mag. 807.
125 <i>chloræfólia</i> . DC. (wh.) Armenian.	Armenia.	1796.	8. 9.	H. ¼.	Jacq. obs. 4. t. 79.
126 <i>italíca</i> . DC. (wh.) Italian.	Italy.	1759.	5. 6.	H. ♂.	
<i>Cucúbalus itálicus</i> . L.					
127 <i>polyphylla</i> . DC. (w.p.) many-leaved.	Austria.	6. 7.	H. ¼.	Clus. hist. I. t. 290.
128 <i>nemoralis</i> . DC. (wh.) wood.	Hungary.	1823.	—	H. ¼.	W. et K. hung. 3. t. 249.
129 <i>longiflóra</i> . DC. (wh.) long-flowered.	—	1793.	7. 9.	H. ¼.	— 1.p.7.t.8.
130 <i>bupleuroïdes</i> . DC. (fl.) spear-leaved.	Persia.	1801.	6. 7.	H. ¼.	Tourn. it. t. 154.
131 <i>pilosa</i> . s.s. (wh.) pilose.	Hungary.	1825.	6. 9.	H. ¼.	W. et K. hung. 3. t. 248.
<i>mollissima</i> . W.K.					
132 <i>mollissima</i> . s.s. (wh.) velvet.	Italy.	1739.	7. 9.	H. ¼.	
<i>Cucúbalus mollissimus</i> . L.					
133 <i>inclusa</i> . H. (wh.) included.	—	1828.	6. 9.	H. ⊙.	
134 <i>pínguis</i> . H. (wh.) fleshy-leaved.	C. Spartel.	1825.	—	H. ⊙.	
135 <i>táurica</i> . F. (wh.) Taurian.	Tauria.	1828.	—	H. ¼.	
136 <i>dahúrica</i> . F. (wh.) Dahurian.	Dahuria.	1827.	—	H. ¼.	
137 <i>scàbra</i> . F. (wh.) rough.	Siberia.	—	—	H. ¼.	
138 <i>marginátâ</i> . Sct. (wh.) margined.	—	1825.	—	H. ¼.	
139 <i>stenopetalâ</i> . Led. (re.) narrow-petaled.	Siberia.	1827.	—	H. ⊙.	
140 <i>saponariæfólia</i> . Sct. (re.) Soapwort-ld.	—	1822.	—	H. ⊙.	
141 <i>Persoñii</i> . Sct. (wh.) Persoon's.	Europe.	—	—	H. ¼.	
142 <i>vesicária</i> . (pu.) bladdered.	Siberia.	1822.	6. 9.	H. ⊙.	
143 <i>régia</i> . DC. (cr.) splendid.	N. America.	1811.	5. 8.	G. ♂.	Bot. mag. 1724.
144 <i>Atocion</i> . DC. (pu.) Orchis-flow'r'd.	Levant.	1781.	5. 7.	H. ⊙.	Jacq. vind. 3. t. 32.
<i>orchidéa</i> . H.K. Flor. græc. t. 427.					
145 <i>pseudo-Atocion</i> . DC. (pu.) Atlantic.	Atlantic.	1818.	—	H. ⊙.	
146 <i>Arméria</i> . DC. (re.) Lobel's.	England.	7. 9.	H. ⊙.	Eng. bot. 1398.
<i>a rúbra</i> .	(re.) red-flowered.	—	—	H. ⊙.	
β <i>álba</i> .	(wh.) white-flowered.	—	—	H. ⊙.	
147 <i>congestâ</i> . DC. (re.) close-flowered.	Greece.	1816.	—	H. ⊙.	
148 <i>compácta</i> . DC. (re.) compact.	Russia.	1816.	—	H. ♂.	
<i>VISC'ARIA</i> . Rechl.	<i>VISC'ARIA</i> . Decandria Pentagynia. L.				
1 <i>vulgáris</i> . Rechl. (re.) common.	Britain.	5. 7.	H. ¼.	Eng. bot. 788.
<i>Lychnis viscária</i> . L.					
β <i>flore-pléno</i> .	(re.) double-red.	—	—	H. ¼.	

CARYOPHYLLEÆ.

2	albiflora.	(wh.) white-flowered.	5. 7.	H. ȝ.	
LY'CHNIS. DC.		LY'CHNIS.	Decandria Pentagynia. L.			
1	chalcedonica. DC.	(sc.) scarlet.	Japan.	1596.	6. 7.	H. ȝ.
	β albiflora.	(wh.) white-flowered.	—	—	H. ȝ.
	γ flôre pleno.	(sc.) double-flowered.	—	—	H. ȝ.
2	Flós-jòvis. DC.	(re.) umbelled.	Piedmont.	1726.	7.	H. ȝ.
	Agrostémma Flós-jòvis. B.M.					Bot. mag. 398.
3	coronata. w.	(sc.) Chinese.	China.	1774.	6. 9.	H. ȝ.
	grandiflora. Jacq.	ic. rar. t. 84.				Bot. mag. 223.
4	fúlgens. DC.	(sc.) brilliant.	Siberia.	1819.	7. 9.	H. ȝ.
5	Cœ'lí-rosa. DC.	(fl.) smooth-leaved.	Levant.	1713.	7. 8.	H. ⊙.
	Agrostémma Cœ'lí-rosa. L.					Bot. mag. 295.
AGROSTE'MMA.		AGROSTE'MMA.	Decandria	Pentagynia. L.		
1	apétala. DC.	(wh.) petalless.	Lapland.	1819.	6. 8.	H. ȝ.
	Ly'chnis apétala. L.					Lin. fl. lap. t. 12. f. 1
2	brachypétala. L. en.	(wh.) short-petal'd.	—	—	H. ȝ.
3	sylvestris. DC.	(re.) red-flowered.	Britain.	6. 9.	H. ȝ.
	Ly'chnis dióica rùbra. E. B.					Eng. bot. 1579.
	β flôre-pleno.	double-flowered.	—	—	H. ȝ.
4	dióica. DC.	(wh.) white-flowered.	Britain.	—	H. ȝ.
	Ly'chnis dióica álba. E.B. vespertina. Sibthorp.					Eng. bot. 1580.
	β flôre-pleno.	(wh.) double-white.	—	—	H. ȝ.
	γ viridiflora.	(gr.) double-green.	—	—	H. ȝ.
5	dichlinis. DC.	(ro.) long-pedicled.	Spain.	1824.	—	H. ȝ.
6	læta. DC.	(re.) small.	Portugal.	1778.	7.	H. ⊙.
7	córsica. DC.	(re.) Corsican.	Corsica.	1824.	6. 8.	H. ⊙.
8	pyrenáica. DC.	(re.) Pyrenean.	Pyrenees.	1820.	4. 6.	H. ȝ.
9	suècica. L.C.	(re.) Swedish.	Sweden.	6. 9.	H. ȝ.
10	alpina. DC.	(re.) Alpine.	Scotland.	4. 5.	H. ȝ.
	Ly'chnis alpina. E.B.					Lodd. bot. cab. 881.
11	Flósculæ. DC.	(re.) Ragged Robin.	Britain.	6. 9.	H. ȝ.
	Ly'chnis Flósculæ. L.					Eng. bot. 573.
	β flôre-pleno.	(re.) double.	—	—	H. ȝ.
12	coronaria. DC.	(ro.) Rose Campion.	Italy.	1596.	—	H. ȝ.
	a rùbra.	(ro.) red-flowered.	—	—	H. ȝ.
	β álba.	(wh.) white-flowered.	—	—	H. ȝ.
	γ plena.	(ro.) double-flowered.	—	—	H. ȝ.
GITHAGO. Desf.		CORN-COCKLE.	Decandria	Pentagynia. L.		
1	segetum. Desf.	(re.) common.	Britain.	6. 8.	H. ⊙.
	Agrostémma Githago. L.					Eng. bot. 741.
2	nicaéensis.	(wh.) long-calyxed.	Italy.	1794.	—	H. ⊙.
VEL'EZIA. DC.		VEL'EZIA.	Pentandria Digynia. L.			
1	rígida. DC.	(re.) rigid.	Spain.	1683.	7.	H. ⊙.
2	quadridentata. F.G.(r.)	four-toothed.	Levant.	1826.	—	F. ȝ.
DR'YPI'S. DC.		DR'YPI'S.	Pentandria Trigynia. L.			
spinosa. DC.	(bh.) prickly.	Italy.	1775.	6. 7.	H. ȝ.	Bot. mag. 2216.

Tribus II. ALSINEÆ. DC.

ORT'EGIA. DC.	ORT'EGIA.	Triandria Monogynia. L.				
1 hispánica. DC.	(gr.) Spanish.	Spain.	1768.	6. 7.	H. ȝ.	
2 dichotoma. DC.	(gr.) forked.	Italy.	1781.	8. 9.	H. ȝ.	All ped. t. 4. f. 1.
	hispánica. Cav. ic. 1. t. 47.					
BUFFONIA. DC.	BUFFONIA.	Tetrandria Digynia. L.				
1 ánnua. DC.	(wh.) annual.	England.	6.	H. ⊙.	Eng. bot. 1313.
	tenuifolia. E.B.					
2 perénis. DC.	(wh.) perennial.	France.	1820.	6. 7.	H. ȝ.	Lam. ill. t. 87. f. 2.
SAG'INA. DC.	PEARL-WORT.	Tetrandria Tetragynia. L.				
1 procumbens. DC.	(gr.) procumbent.	Britain.	5. 9.	H. ⊙.	Eng. bot. 880.
	β multipléx.	(wh.) double-flowered.	—	—	H. ȝ.
2 cerastoïdes. DC.	(wh.) Cerastium like.	—	6. 7.	H. ⊙.	Eng. bot. 166.
	Cerástium tetrandrón H. K.					
3 filiformis. DC.	(wh.) filiform.	Pyrenees.	1824.	—	H. ⊙.	
4 marítima. E.B.	(wh.) sea-side.	Ireland.	5. 8.	H. ⊙.	Eng. bot. 2195.
5 strícta. DC.	(wh.) straight.	N. Europe.	1828.	—	H. ⊙.	
6 apétala. DC.	(wh.) small-flowered.	Britain.	5. 6.	H. ⊙.	Eng. bot. 881.

MŒ'NCHIA. P.S.	MŒ'NCHIA.	Tetrandria Tetragynia. L.				
gláucia. p.s.	(wh.) glaucous.	Britain.	4. 5. H. ⊙.	Eng. bot. 609.	
quaternélla. L. en.	Sagina erécta. E.B.					
MŒHRINGIA. DC.	MŒHRINGIA.	Octandria Digynia. L.				
1 muscôsa. DC.	(wh.) mossy.	S. Europe.	1775.	6. 7. H. ¼.	Lam. ill. t. 314.	
2 sedifolia. DC.	(wh.) Stonecrop-ld.	—	1823.	— H. ¼.	W.ber.m.1818.t.3.f.23.	
ELATINE. DC.	WATERWORT.	Octandria Tetragynia. L.				
1 Hydropiper. DC.	(wh.) small.	Britain.	7. 8. H. ¼.	Eng. bot. 955.	
2 hexândra. DC.	(wh.) hexandrous.	France.	— H. ¼.	DC. ic. rar. 1. t. 43. f. 1.	
BER'GIA. DC.	BE'RGIA.	Decandria Pentagynia. L.				
verticillâta. DC.	(wh.) whorl-leaved.	E. Indies.	1826.	7. 10. H. ⊙.	Delil. fl. æg. t. 26. f. 1.	
MOLLUGO. W.	MOLLUGO.	Triandria Trigynia. L.				
1 verticillâta. DC.	(wh.) whorled.	Virginia.	1748.	6. 8. H. ⊙.	Ehret pict. t. 6.	
2 Schránkii. DC.	(wh.) Schrank's.	Brazil.	1821.	7. 9. H. ⊙.	Schrank pl. rar. 64. ic.	
3 Línkii. DC.	(wh.) Link's.	—	—	— H. ⊙.		
	triphyl'la. L. en. non Lour.					
PHARN'ACEUM. W	PHARN'ACEUM.	Pentandria Trigynia. L.				
1 Mollùgo. R.S.	(wh.) LadiesBedstraw.	E. Indies.	1752.	7. 8. S. ⊙.	Burm. ind. t. 5. f. 4.	
	Mollugo Spérghula. DC.					
2 cordifoliūm. R.S.	(wh.) heart-leaved.	C. B. S.	1820.	— H. ⊙.	Jacq. h. scheen. t. 349.	
3 dichótomum. R.S.	(wh.) forked.	—	1783.	— H. ⊙.		
4 Cerviâna. R.S.	(wh.) umbelled.	Russia.	1771.	6. 7. H. ⊙.	Lam. ill. t. 214. f. 1.	
5 bellidifòium. R.S.	(w.) Daisy-leaved.	W. Indies.	1826.	— H. ⊙.	Plum. amer. t. 21. f. 1.	
HOLO'STEUM. DC.	HOLO'STEUM.	Triandria Trigynia. L.				
1 umbellatum. DC.	(wh.) umbelled.	England.	7. 8. H. ⊙.	Flor. dan. t. 1204.	
β röseum.	(ro.) rose-coloured.	—	— H. ⊙.	Eng. bot. 27.	
2 diandrum. DC.	(wh.) diandrous.	Jamaica	1822.	6. 8. S. ⊙.	Swartz ic. t. 7.	
SPERG'ULA. DC.	SPURREY.	Decandria Pentagynia. L.				
1 arvénis. DC.	(wh.) rough-seeded.	Britain.	7. 8. H. ⊙.	Eng. bot. 1535.	
2 pentândra. DC.	(wh.) smooth-seeded.	England.	6. 7. H. ⊙.	— 1536.	
3 pálida. DC.	(pa.) pale.	C. B. S.	7. 9. G. ¼.		
4 nodôsa. DC.	(wh.) knotted.	Britain.	7. 8. H. ¼.	Eng. bot. 694	
5 laricina. DC.	(wh.) Larch-leaved.	Siberia.	1823.	6. 8. H. ¼.	Sm. ic. ined. 1. t. 18.	
6 saginoides. DC.	(wh.) Sagina like	Scotland.	— H. ¼.	Eng. bot. 2105.	
7 subulâta. DC.	(wh.) awl-leaved.	Britain.	6. 7. H. ¼.	— 1082.	
8 glâbra. DC.	(wh.) smooth.	Europe.	1824.	— H. ¼.	All. ped. t. 64. f. 1.	
LA'RBEA. DC.	LA'RBEA.	Decandria Trigynia. L.				
aquática. DC.	(wh.) water.	Britain.	7. 8. H.w. ¼.	Eng. bot. t. 1074.	
Stellaria aquática. Poll. pal. n. 422.		uliginosa. E.B.	alsine Hoff. germ.	1. t. 5.		
DRYM'ARIA. DC.	DRYM'ARIA.	Pentandria Trigynia. L.				
1 cordâta. DC.	(wh.) heart-leaved.	W. Indies.	1814.	7. 9. H. ⊙.	Lam. ill. t. 51. f. 2.	
	Holosteum cordatum. L.					
2 divaricâta. DC.	(wh.) divaricate.	Lima.	1823.	— H. ⊙.		
STELL'ARIA. DC.	STICHWORT.	Decandria Trigynia. L.				
1 némorum. DC.	(wh.) wood.	Britain.	4. 6. H. ¼.	Eng. bot. 92.	
2 latifolia. DC.	(wh.) broad-leaved.	Germany.	1816.	6. 8. H. ⊙.		
3média. DC.	(wh.) Chickweed.	Britain.	1. 12. H. ⊙.	Eng. bot. 537.	
	Alsine media. L.					
4 dichótomâ. DC.	(wh.) forked.	Siberia.	1774.	6. 8. H. ¼.	Smith ic. ined. 1. t. 14.	
5 bulbôsa. DC.	(wh.) bulbiferous.	Carinthia.	1820.	— H. ¼.	Jacq. ic. rar. 3. t. 468.	
6 víscida. DC.	(wh.) viscous.	Tauria.	1804.	— H. ⊙.	W. et K. hung. 1. t. 22.	
	Cerastium anomálum. w.					
7 sabulôsa. DC.	(wh.) gravel.	Persia.	1828.	— H. ⊙.		
8 Holóstea. DC.	(wh.) greater.	Britain.	4. 6. H. ¼.	Eng. bot. 511.	
9 Laxmánni. DC.	(wh.) Laxmann's.	Siberia.	1827.	— H. ¼.		
10 velutina. DC.	(wh.) velvety.	Siberia.	1824.	6. 8. H. ¼.		
11 graminea. DC.	(wh.) grass-leaved.	Britain.	5. 8. H. ¼.	Eng. bot. 803.	
12 glâuca. DC.	(wh.) glaucous marsh.	—	— H. ¼.	— 825.	
13 cerastoides. DC.	(wh.) Alpine.	Scotland.	— H. ¼.	Eng. bot. 911.	
14 multicáulis. w.	(wh.) many-stalked.	Carinthia.	1815.	6. 8. H. ¼.	Jacq. coll. 1. t. 19.	
	cerastoides. Jacq. coll.					
15 arenâria. DC.	(wh.) sand.	Spain.	1799.	6. 8. H. ⊙.		
16 scapígera. DC.	(wh.) naked-stalked.	Scotland.	6. 7. H. ¼.	Eng. bot. 1269.	
17 Fischeriana. DC.	(wh.) Fischer's.	Siberia.	1825.	— H. ¼.		
18 Edwârdii. B.	(wh.) Edwards's.	N. America.	1827.	— H. ¼.		
	nítida. Hook.					
19 humifusa. DC.	(wh.) trailing.	N. Europe.	1825.	— H. ⊙.	Flor. dan. t. 978.	
20 læta. s.s.	(wh.) short-petaled.	N. America.	1827.	— H. ⊙.		

CARYOPHYLLEÆ.

21 gráclis. s.s.	(wh.) slender.	N. America.	1827.	6. 7.	H. ⊖.
22 murális. DC.	(wh.) wall.	Crete.	1821.	—	H. ⊖.
23 élégans. DC.	(wh.) elegant.	Siberia.	1823.	—	H. ȝ.
24 lóngipes. DC.	(wh.) long pedicled.	N. America.	1821.	—	H. ȝ.
25 mollugínea.	(wh.) Mollugo-like.	New Spain.	1825.	7. 9.	H. ⊖.
<i>Alsine molluginea. Lag. Arenaria? molluginea. DC.</i>					
AREN'ARIA. DC.	SANDWORT.	Decandria Trigynia. L.			
1 segetális. DC.	(wh.) hedge.	France.	1805.	6. 10.	H. ⊖.
<i>Alsine segetalis. L.</i>					
2 rúbra. DC.	(re.) purple.	Britain.	6. 8.	H. ⊖.
3 marína. L.	(re.) sea-side.	—	—	H. ȝ.
4 salina. DC.	(re.) salt marsh.	Bohemia.	1820.	—	H. ⊖.
5 rubéllea. F.L.	(re.) red-flowered.	England.	—	H. ⊖.
6 glutinósa. DC.	(pu.) glutinous.	Astracan.	1828.	—	H. ⊖.
7média. DC.	(re.) marine.	Britain.	—	H. ȝ.
<i>marina. E.B. non L.</i>					
8 glandulósa. Jac.	(wh.) glandular.	Europe.	1820.	6. 7.	H. ⊖.
9 canadénis. DC.	(re.) Canadian.	N. America.	1812.	—	H. ⊖.
10 graminifólia. DC.	(wh.) grass-leaved.	Caucasus.	1815.	—	H. ȝ.
<i>β glaberrima. DC. (w.) smooth.</i>					
11 longifólia. DC.	(wh.) long-leaved.	Volga.	1820.	—	H. ȝ.
12 dahúrica. DC.	(wh.) Dahurian.	Dahuria.	1824.	—	H. ȝ.
13 nardifólia. F.	(wh.) Nardus-leaved.	Siberia.	1827.	—	H. ȝ.
14 otitoídes. DC.	(wh.) dense-flowered.	—	—	—	H. ȝ.
15 rígida. DC.	(wh.) rigid.	—	—	—	H. ȝ.
16 formósa. DC.	(wh.) handsome.	Dahuria.	—	—	H. ȝ.
17 Gmelíni. DC.	(wh.) Gmelin's.	—	1827.	—	H. ȝ.
18 cephalótes. DC.	(wh.) close-headed.	Tauria.	1820.	—	H. ȝ.
19 imbricáta. DC.	(wh.) imbricate.	Caucasus.	1821.	—	H. ȝ.
20 juniperina. DC.	(wh.) Juniper-leaved.	Armenia.	1800.	—	H. ȝ.
21 stricta. DC.	(wh.) upright.	N. America.	1812.	5. 6.	H. ȝ.
22 pròcera. DC.	(wh.) tall.	—	1827.	—	H. ȝ.
23 laricifólia. L.	(wh.) Larch-leaved.	Switzerland.	8.	H. ȝ.
24 rostráta. W.en.	(wh.) beaked.	Hungary.	1816.	6. 7.	H. ȝ.
25 striáta. w.	(wh.) striated.	Switzerland.	1683.	—	H. ȝ.
26 ártica. DC.	(wh.) small.	Siberia.	1807.	—	H. ȝ.
27 austriaca. DC.	(wh.) Austrian.	Austria.	1793.	6. 9.	H. ȝ.
28 Villársii. P.S.	(wh.) Villars's.	Dauphiny.	1826.	—	H. ȝ.
29 grandifóra. DC.	(wh.) great-flowered.	Switzerland.	1783.	—	H. ȝ.
30 triflóra. L.	(wh.) three-flowered.	—	1816.	—	H. ȝ.
31 Hélmii. DC.	(wh.) long-peduncled.	Siberia.	1824.	—	H. ȝ.
32 macrocárpa. DC.	(wh.) large-capsuled.	N. America.	1828.	—	H. ȝ.
33 hírta. DC.	(re.) hairy.	Greenland.	1827.	—	H. ȝ.
34 vérra. DC.	(wh.) vernal.	Britain.	5. 8.	H. ȝ.
35 cæspítosa. DC.	(wh.) tufted.	S. Europe.	1824.	—	H. ȝ.
36 ramosissima. DC.	(wh.) branching.	Hungary.	1820.	—	H. ȝ.
37 saxátilis. DC.	(wh.) rock.	Germany.	1732.	7. 8.	H. ȝ.
38 péncola. DC.	(wh.) pendulous.	Hungary.	1816.	6. 7.	H. ȝ.
39 tenuifólia. DC.	(wh.) fine-leaved.	England.	—	H. ⊖.
40 villósa. DC.	(wh.) villous.	Siberia.	1828.	—	H. ȝ.
41 calycína. DC.	(wh.) large-calyxed.	Barbary.	1824.	—	H. ⊖.
42 triándra. DC.	(wh.) triandrous.	—	—	—	H. ⊖.
43 mediterránea. DC. (w.)	Mediterranean.	S. Europe.	1821.	—	H. ⊖.
44 recúrva. DC.	(wh.) recurved-leav'd.	—	—	—	H. ȝ.
45 hirsuta. DC.	(wh.) hairy.	Tauria.	1824.	—	H. ȝ.
46 setácea. DC.	(wh.) bristly.	France.	1820.	—	H. ȝ.
47 fasciácula. DC.	(wh.) dense-flowered.	S. Europe.	1815.	—	H. ⊖.
48 fastigiáta. E.B.	(wh.) fastigiate.	Scotland.	6. 7.	H. ⊖.
49 filifólia. DC.	(wh.) thread-leaved.	Arabia.	—	H. ȝ.
50 mucronáta. DC.	(wh.) mucronate.	—	1777.	—	H. ⊖.
51 uliginósa. DC.	(wh.) marsh.	Switzerland.	1824.	—	H. ȝ.
52 polygonóides. DC.	(wh.) procumbent.	—	—	—	H. ȝ.
53 margináta. DC.	(wh.) margined.	Siberia.	1824.	—	H. ȝ.
54 canéscens. DC.	(wh.) canescent.	—	1828.	—	H. ȝ.
<i>macrocárpa. H. non Ph.</i>					
55 tetraquétrá. DC.	(wh.) square-stalked.	S. France.	1731.	8.	H. ȝ.
56 Gieseckii. DC.	(wh.) Gieseke's.	Greenland.	1827.	—	H. ȝ.
57 lanceoláta. DC.	(wh.) lanceolate.	Piedmont.	1820.	—	H. ȝ.
58 cherlerioides. P.S.	(wh.) cherleria-leav'd.	S. France.	7. 8.	H. ȝ.

Vail. par. t. 3. f. 3.

Eng. bot. 852.

Flor. lond. t.

Eng. bot. 958.

Jacq. Schœnb. 3. t. 355.

Schrad. h. gøtt. t. 5.

Gmel. sib. 4. t. 61. f. 1.

Sm. ic. ined. t 35.

Jacq. aust. 3. t. 272.

Vill. dauph. 4. t. 47. f. 6.

Jacq. aust. 3. t. 270.

Vill. de ph. 3. t. 47.

All ped.n.1711.t.10.f.1.

Cav. ic. 3. t. 249. f. 2.

Flor. dan. t. 1646.

Eng. bot. 512.

Barr. ic. t. 580.

W. et K. hung. 1. t. 87.

Eng. bot. 219.

Schrank. h. mon. t. 30.

Jacq. coll. 2. t. 16. f. 1.

Flor. græc. t. 442.

Eng. bot. 1744.

Vahl. symb. 1. t. 12.

DC. ic. gall. t. 46.

Jacq. coll. 1. t. 15.

All. ped. 2. t. 89. f. 1.

Horn. fl. dan t. 1518.

All. fl. ped. t. 26. f. 5.

Vill. delph. 4. t. 47. f. 1.

59	montàna. DC.	(wh.) mountain.	Spain.	1800.	4. 7.	H. ȝ.	Bot. mag. 1118.
60	serpyllifòlia. DC.	(wh.) Thyme-leaved.	Britain.	6. 7.	H. ȝ.	Eng. bot. 923.
61	ciliàta. DC.	(wh.) fringed.	Ireland.	3. 8.	H. ȝ.	—— 1745.
62	multicàulis. W.	(wh.) many-stalked.	Europe.	1794.	7. 8.	H. ȝ.	Jacq. coll. 1. t. 17. f. 1.
63	trinèrvia. DC.	(wh.) three-nerved.	Britain.	5. 7.	H. ȝ.	Eng. bot. 1483.
64	norvègica. DC.	(wh.) Norway.	Norway.	1828.	6. 7.	H. ȝ.	Flor. dan. t. 1269.
65	baleàrica. DC.	(wh.) Majorca.	Majorca.	1787.	3. 8.	H. ȝ.	L. Her. stirp. 1. t. 15.
66	biflòra. DC.	(wh.) two-flowered.	S. Europe.	1823.	——	H. ȝ.	All. ped. t. 64. f. 3.
67	procùmbens. DC. (re)	procumbent.	Egypt.	1801.	6. 8.	F. ȝ.	Vahl. symb. 1. t. 33.
ADEN'ARIUM. DC. SEA CHICKWEED. Decandria Tri-Pentagynia.							
	peplòides. DC.	(wh.) common.	Britain.	6. 7.	H. ȝ.	Eng. bot. 189.
<i>Arenaria peploides.</i> E.B.							
CERA'STIUM. DC. MOUSE-EAR CHICKWEED.							
1	nemoràle. DC.	(wh.) wood.	Caucasus.	1823.	5. 6.	H. ȝ.	Gmel. sib. 4. t. 62. f. 2.
2	perfoliatum. DC.	(wh.) perfoliate.	Greece.	1725.	6. 7.	H. ȝ.	Dill. elt. t. 217. f. 284.
3	caucasicum. DC.	(wh.) Caucasean.	Caucasus.	1825.	——	H. ȝ.	Schrank. h. mon. t. 75.
4	dahuricum. DC.	(wh.) glaucous.	Dahuria.	1815.	5. 9.	H. ȝ.	
	amplexicáule. B.M. 1789.						
5	máximum. DC.	(wh.) greatest.	Siberia.	1792.	6. 7.	H. ȝ.	
6	dichotórum. DC.	(wh.) forked.	Spain.	1725.	——	H. ȝ.	
7	ruderàle. DC.	(wh.) pendulous-fruit.	Caucasus.	1824.	——	H. ȝ.	
8	táuricum. DC.	(wh.) Taurian.	Tauria.	——	——	H. ȝ.	
9	vulgátum. DC.	(wh.) common.	Britain.	4. 6.	H. ȝ.	Eng. bot. 789.
10	viscòsum. DC.	(wh.) clammy.	——	4. 9.	H. ȝ.	—— 790.
11	barbulátum. L.en.	(wh.) bearded.	S. Europe.	1821.	——	H. ȝ.	
12	holosteóides. DC.	(wh.) Holosteum like.	Sweden.	——	——	H. ȝ.	
13	pentándrum. DC.	(wh.) five-stamened.	Spain.	1820.	——	H. ȝ.	
14	semidecadrûm. DC.	(wh.) least.	Britain.	3. 4.	H. ȝ.	Eng. bot. 1630.
15	brachypétalum. DC.	(wh.) small-flowered.	France.	1816.	4. 5.	H. ȝ.	DC. ic. gall. t. 44.
16	diffúsom. DC.	(wh.) spreading.	4. 9.	H. ȝ.	
17	serpylliòlium. W.en.	(wh.) Thyme-ld.	S. Europe.	1824.	5. 6.	H. ȝ.	
18	mánticum. DC.	(wh.) long pendul'd.	Hungary.	1801.	6. 7.	H. ȝ.	W. et K. hung. 1. t. 96.
19	campanulátum. DC.	(wh.) early-flow ^d .	Italy.	1826.	4. 6.	H. ȝ.	Viv. ann. bot. 1. p.2.t.1.
	pr ^r æcox. T.N.						
20	grandiflòrum. DC.	(wh.) large-flowered.	——	1820.	6. 7.	H. ȝ.	W. et K. hung. 2. t. 168.
21	argénteum. M.B.	(wh.) silvery-leaved.	Iberia.	1821.	——	H. ȝ.	
22	tomentòsum. DC.	(wh.) white-leaved.	S. Europe.	1648.	——	H. ȝ.	Col.phyt.ed.1744.t.31.
23	Bieberstéini. DC.	(wh.) Bieberstein's.	Tauria.	1820.	——	H. ȝ.	DC. pl. rar. gen. t. 11.
24	lanátum. DC.	(wh.) woolly.	Pyrenees.	——	——	H. ȝ.	
25	alpinum. DC.	(wh.) Alpine.	Britain.	——	H. ȝ.	Eng. bot. 472.
26	ovátum. DC.	(wh.) oval-leaved.	Carinthia.	1816.	——	H. ȝ.	
27	latifòlium. DC.	(wh.) broad-leaved.	Britain.	——	H. ȝ.	Eng. bot. 473.
28	aquáticum. E.B.	(wh.) water.	——	——	H.w. ȝ.	—— 538.
29	sylváticum. DC.	(wh.) wood.	Hungary.	1826.	——	H. ȝ.	W. et K. hung. 1. t. 97.
30	arvénse. DC.	(wh.) field.	——	5. 8.	H. ȝ.	Eng. bot. 93.
31	repens. W.	(wh.) creeping.	S. Europe.	1759.	5. 7.	H. ȝ.	Lam. ill. t. 392. f. 2.
32	strictum. W.	(wh.) upright.	Austria.	1793.	——	H. ȝ.	Scop. carn. t. 19. f. 1.
33	suffruticòsum. W.	(wh.) suffruticose.	S. Europe.	1796.	——	H. ȝ.	
34	pensylvanicum. DC.	(wh.) Pensylvanian.	N. America.	1823.	——	H. ȝ.	
35	pubescens. DC.	(wh.) pubescent.	——	1822.	——	H. ȝ.	
36	matréns. L.en.	(wh.) Matram.	Hungary.	1823.	——	H. ȝ.	
37	diòicum. H.K.	(wh.) Spanish.	Spain.	1766.	6. 7.	H. ȝ.	
38	pilòsum. L.en.	(wh.) hairy-leaved.	1823.	——	H. ȝ.	
39	Sprengeli. DC.	(wh.) Sprengel's.	——	——	H. ♂.	
40	Tenoreánum. DC.	(wh.) Tenore's.	Italy.	1825.	——	H. ȝ.	
41	Samniànum. DC.	(wh.) Samnian.	——	——	——	H. ȝ.	
42	hirsütum. DC.	(wh.) hairy.	——	——	——	H. ȝ.	
43	glandulòsum. Lk.	(wh.) glandular.	1827.	——	H. ȝ.	
44	inflatùm. Lk.	(wh.) inflated.	——	——	H. ȝ.	
CHERL'ERIA. DC. CHERL'ERIA.							
	sedoides. DC.	(wh.) dwarf.	Decandria Trigynia. L.	7. 8.	H. ȝ.	Eng. bot. 1212.
			Scotland.			

ORDO XXVII.

LINEÆ. DC. prodr. 1. p. 423.

ORDO XXVIII.

MALVACEÆ. *DC. prodr.* 1, p. 429.

MA'LOPE. DC. **MA'LOPE.** *Monadelphia* *Polyandria*. L.
 1 *malacoïdes*. DC. (*pu.*) *Barbary.* *Barbary.* 1710. 6. 7. G. ♂. Cav. diss. 2. t. 27. f. 1.
 2 *trifida*. DC. (*pu.*) *trifid.* —————— 1808. 7. 8. H. ♂. Swt. br. fl. gar. t. 153.

MA'LVA. DC.	MALLOW. Monadelphia Polyandria. L.	W. Indies.	1726.	7. 8.	S. ♂.	Cav. diss. 2. t. 22. f. 2.
1 tricuspida. DC. (<i>ye.</i>) Jamaica. <i>coromandeliæna.</i> w. <i>americana.</i> Cav.			1756.	6. 7.	H. ♂.	
2 americana. DC. (<i>ye.</i>) American.	—		1798.	—	G. ♀.	Cav. diss. 5. t. 138. f. 1.
3 scabra. DC. (<i>ye.</i>) rough-stemmed. Peru. <i>scoparia.</i> Jacq. ic. 139.			1782.	8. 9.	G. ♀.	L. her. stirp. t. 27.
4 scoparia. DC. (<i>ye.</i>) Birch-leaved.	—	Mauritius.	1816.	7. 8.	S. ♀.	Cav. dis. 5. t. 138. f. 3.
5 borbonica. DC. (<i>ye.</i>) Bourbon.	—	Peru.	1798.	—	G. ♀.	— 2. t. 20. f. 4.
6 polystachya. DC. (<i>ye.</i>) many-spiked.	—	Jamaica.	1726.	9. 10.	S. ♀.	— 2. t. 20. f. 2.
7 spicata. DC. (<i>ye.</i>) simple-spiked.	—	Brazil.	—	—	S. ♀.	— 2. t. 21. f. 3.
8 ovata. L. en.	(<i>ye.</i>) oval-leaved.	S. America.	1821.	7. 10.	S. ♀.	Pluk. alm. t. 356. f. 1.?
9 subhastata. DC. (<i>ye.</i>) subhastate.	—	India.	1820.	—	S. ♀.	
10 tomentosa. DC. (<i>ye.</i>) tomentose.	—	Java.	1824.	—	S. ♀.	
11 waltherifolia. DC. (<i>ye.</i>) Waltheria-ld.	—	E. Indies.	—	7. 8.	H. ♂.	Pluk. t. 74. f. 6.
12 trachelifolia. DC. (<i>ye.</i>) Trachelium-ld.	—	Cuba.	1815.	5. 7.	S. ♀.	
13 gangética. DC. (<i>ye.</i>) Indian.	—	Bithynia.	—	—	H. ♂.	Cav. dis. 2. t. 26. f. 4.
14 leprosa. DC. (<i>wh.</i>) leprosus.	—	Crete.	1828.	—	H. ♂.	— 5. t. 138. f. 2.
15 Sherardiæna. DC. (<i>re.</i>) Sherard's.	—	Spain.	1710.	7.	H. ♂.	Desf. atl. 2. t. 170.
16 crética. DC. (<i>fl.</i>) Cretan.	—	Egypt.	1815.	6. 8.	H. ♂.	Cav. dis. 2. t. 15. f. 2.
17 hispánica. DC. (<i>ro.</i>) Spanish.	—	Spain.	1739.	6. 7.	H. ♂.	— 2. t. 17. f. 1.
18 stipulacea. DC. (<i>pu.</i>) long-stipuled.	—	Spain.	1815.	—	H. ♂.	— 5. t. 137. f. 2.
19 ægyptia. DC. (<i>bh.</i>) Egyptian.	—	Europe.	1759.	7. 8.	H. ♂.	— 2. t. 17. f. 3.
20 trifida. DC. (<i>bl.</i>) large-flowered.	—	Italy.	1597.	7. 10.	H. ♀.	Bot. mag. 2297.
21 Tournefortiana. DC. (<i>r.</i>) Tournefort's.	—	Italy.	1824.	—	H. ♀.	— 2793.
22 Alcæa. DC. (<i>pi.</i>) Vervain.	—	Britain.	—	7. 8.	H. ♀.	
23 Morènii. s.s. (<i>pi.</i>) Moreni's.	—	Sicily.	1825.	—	H. ♂.	Eng. bot. 754.
alceoides. Ten. fl. nap. t. 64.	—	Spain.	1828.	6. 8.	H. ♂.	Bot. mag. 2298.
24 moschata. DC. (<i>pi.</i>) musk.	—	S. Europe.	1768.	6. 7.	H. ♂.	Cav. dis. 2. t. 23. f. 2.
β laciniata. DC. (<i>wh.</i>) white-flowered.	—	S. Europe.	1828.	—	H. ♂.	Swt. br. fl. gar. 81.
25 hirsuta. T. (<i>pi.</i>) hairy.	—	Sicily.	1828.	6. 8.	H. ♂.	
26 fastigiata. DC. (<i>re.</i>) fastigiate.	—	Barbary.	1768.	6. 7.	H. ♂.	
27 mauritiæna. DC. (<i>pu.</i>) Ivy-leaved.	—	China.	1828.	—	H. ♂.	Eng. bot. 671.
28 excelsa. Pr. (<i>pu.</i>) tall.	—	Egypt.	1779.	6. 7.	H. ♀.	
29 sylvæstris. DC. (<i>pu.</i>) common.	—	Egypt.	1828.	6. 10.	H. ♀.	Eng. bot. 1092.
30 vulgàris. T. (<i>pu.</i>) prostrate.	—	Barbary.	1828.	—	H. ♂.	— 241.
31 Henníngii. DC. (<i>pu.</i>) Henning's.	—	China.	1823.	6. 8.	H. ♀.	
32 rotundifolia. DC. (<i>wh.</i>) round-leaved.	—	China.	1823.	—	H. ♂.	
β pusilla. E.B. (<i>wh.</i>) dwarf.	—	China.	1820.	—	S. ♀.	
33 brasiliensis. DC. (<i>pi.</i>) Brazilian.	—	China.	1823.	—	H. ♂.	Cav. dis. 2. t. 25. f. 1.
34 niceænus. DC. (<i>w.p.</i>) Italian.	—	Egypt.	1779.	—	H. ♂.	Cav. dis. 2. t. 26. f. 1.
35 microcarpa. DC. (<i>fl.</i>) small-fruited.	—	Barbary.	1833.	—	H. ♂.	Jacq. vind. 1. t. 40.
36 parviflora. DC. (<i>ro.</i>) small-flowered.	—	China.	1828.	6. 10.	H. ♂.	
37 verticillata. DC. (<i>w.p.</i>) whorl-flowered.	—	Egypt.	1573.	6. 8.	H. ♂.	Cav. dis. 2. t. 23. f. 1.
38 mareotica. DC. (<i>fl.</i>) headed.	—	Syria.	1823.	—	H. ♂.	
39 crispa. DC. (<i>bh.</i>) curled.	—	China.	1796.	4. 8.	G. ♀.	Bot. mag. 1998.
40 flexuosa. DC. (<i>re.</i>) flexuous.	—	Egypt.	1812.	—	G. ♀.	Cav. dis. 2. t. 23. f. 4.
41 amœna. DC. (<i>pi.</i>) lobe-leaved.	C. B. S.	—	1727.	5. 7.	G. ♀.	— 2. t. 18. f. 2.
calycina. B.R. 297.	—	—	1713.	1. 12.	G. ♀.	Bot. reg. 295.
42 calycina. DC. (<i>pi.</i>) large-calyxed.	—	—	1800.	5. 9.	G. ♀.	Jacq. ic. 1. t. 140.
43 virgata. DC. (<i>fl.</i>) twiggy.	—	—	1825.	—	G. ♀.	Cav. dis. 2. t. 21. f. 2.
44 capensis. DC. (<i>w.r.</i>) Cape.	—	—	1791.	6. 8.	G. ♀.	
45 balsamica. DC. (<i>re.</i>) balsamic.	—	—	1792.	6. 9.	G. ♀.	And. hot. rep. 182.
46 oxyacanthoides. DC. (<i>w.r.</i>) Hawthorn-ld.	—	—	1803.	3. 5.	G. ♀.	Cav. dis. 2. t. 21. f. 1.
47 tridactylites. DC. (<i>wh.</i>) reflex-flowered.	—	—	1759.	5. 7.	G. ♀.	Bot. reg. 296.
reflexa. A.R. t. 135.	—	—	1805.	5. 8.	G. ♀.	Jacq. schenb. 3. t. 294.
48 divaricata. DC. (<i>wh.</i>) straddling.	—	—	1731.	7. 8.	G. ♀.	Wend. h. her. 1. t. 4.
49 retusa. DC. (<i>y.bh.</i>) blunt-leaved.	—	—	1732.	5. 9.	G. ♀.	Bot. reg. 561.
50 frágans. DC. (<i>pu.</i>) fragrant.	—	—	1796.	—	G. ♀.	Jacq. schenb. 2. t. 139.
51 stricta. DC. (<i>w.r.</i>) upright.	—	—	1780.	1. 2.	G. ♀.	Cav. ic. 1. t. 30.
52 bryonifolia. DC. (<i>re.</i>) Bryony-leaved.	—	—	1798.	11. 12.	G. ♀.	Cav. dis. 5. t. 137. f. 1.
53 grossulariæfolia. DC. (<i>r.</i>) Gooseberry-ld.	—	—	1798.	—	G. ♀.	Cav. ic. 3. t. 278.
54 aspérima. DC. (<i>wh.</i>) roughest.	—	—	1795.	7. 8.	G. ♀.	Cav. dis. 2. t. 35. f. 1.
55 láctea. DC. (<i>wh.</i>) paniced white.	Mexico.	—	1759.	6. 8.	H. ♂.	Jacq. vind. 2. t. 156.
viticifolia. Cav.	—	—	1768.	—	H. ♂.	— 2. t. 140.
56 capitata. DC. (<i>pu.</i>) various-leaved.	Peru.					
57 miniata. DC. (<i>re.</i>) painted.	S. America.					
58 operculata. DC. (<i>pu.</i>) lid capsuled.	Peru.					
59 peruviana. DC. (<i>pu.</i>) Peruvian.						
60 líménais. DC. (<i>bl.</i>) blue-flowered.						

MALVACEÆ.

61 itálica. DC.	(re.) Italian.	Italy.*	1826.	6. 8. H. ¼.	
62 anómala. L. en.	(re.) anomalous.	C. B. S.	1814.	— G. ¼.	Pl. sel. hort. ber. t. 22.
63 umbelláta. DC.	(cr.) umbelled.	S. America.	1814.	1. 3. G. ℥.	Lod. Bot. cab. 222.
64 rósea. DC.	(ro.) rose-coloured.	Mexico.	1828.	— F. ℥.	Flor. mex. ic. ined.
65 abutilóides. DC.	(pu.) Bahama.	Bahama Isl.	1725.	6. 9. G. ℥.	Jacq. scheenb. 3. t. 293.
66 obtusilóbata. B.M.	(pu.) blunt-lobed.	Chili.	1824.	7. 8. F. ℥.	Bot. mag. 2787.
67 élegans. DC.	(y.re.) elegant.	C. B. S.	1791.	5. 8. G. ℥.	Cav. dis. 2. t. 16. f. 1.
68 angustifólia. DC.	(li.) narrow-leaved.	Mexico.	1780.	8. G. ℥.	Bot. mag. 2839.
69 caroliniána. DC.	(re.) Carolina.	Carolina.	1723.	6. 7. H. ○.	Cav. dis. 2. t. 15. f. 1.
70 prostráta. DC.	(sc.) trailing.	Brazil.	1806.	5. 7. G. ¼.	Bot. mag. 2515.
71 decúmbens. DC.	(re.) decumbent.	S. America.	1815.	6. 8. G. ¼.	
KITAIB'ELIA. DC.	KITAIE'ELIA.	Monadelphia Polyandria. L.			
vitífolia. DC.	(wh.) Vine-leaved.	Hungary.	1801.	7. 9. H. ¼.	Bot. mag. 821.
ALTH'EA. DC.	ALTH'EA.	Monadelphia Polyandria. L.			
1 officinalis. DC.	(fl.) Marsh-Mallow.	Britain.	7. 9. H. ¼.	Eng. bot. 147.
2 taurinénsis. DC.	(pu.) intermediate.	S. Europe.	1823.	— H. ¼.	
3 narbonénsis. DC.	(pu.) Narbonne.	—	1780.	8. 9. H. ¼.	Jacq. ic. 1. t. 138.
4 cannabina. DC.	(ro.) Hemp-leaved.	—	1597.	6. 7. H. ¼.	Jacq. aust. 2. t. 101.
5 hirsúta. DC.	(bh.) hairy.	England.	1683.	— H. ○.	— 2. t. 170.
6 Ludwígi. DC.	(wh.) Ludwig's.	Sicily.	1791.	— H. ○.	Cav. ic. 5. t. 423.
7 leucántha. OT.	(wh.) white-flowered.	Mt. Sinai.	1826.	6. 8. H. ♂.	
8 nudiflóra. F.	(ro.) naked-flowered.	Altay.	1824.	6. 10. H. ♂.	
9 Froloviána. F.	(pu.) Siberian.	Siberia.	1826.	6. 8. H. ♂.	
10 glabrifólia. Led.	(pu.) smooth-leaved.	—	1827.	— H. ♂.	
11 acáulis. DC.	(p.ye.) stemless.	Aleppo.	1680.	— H. ○.	Cav. dis. 2. t. 27. f. 3.
12 carib'aea. DC.	(ro.) West Indian.	W. Indies.	1816.	3. 4. S. ♂.	Bot. mag. 1916.
13 striáta. DC.	(li.) streaked.	—	1820.	6. 8. H. ♂.	
14 pálida. DC.	(li.) pale-flowered.	Hungary.	1805.	— H. ♂.	Cav. dis. 2. t. 28. f. 1.
15 rósea. DC.	(va.) Hollyhock.	China.	1573.	7. 9. H. ♂.	
β bilob'a. DC.	(vu.) two-lobed.	—	— H. ♂.	Cav. dis. 2. t. 29. f. 3.
16 sinénsis. DC.	(v.) China.	—	— H. ○.	Bot. mag. 892.
17 flexuósa. DC.	(pu.) flexuous.	E. Indies.	1803.	6. 8. S. ¼.	Cav. dis. 2. t. 28. f. 2.
18 ficifólia. DC.	(ye.) Fig leaved.	Levant.	1597.	6. 9. H. ♂.	
LAVATERA. DC.	LAVATERA.	Monadelphia Polyandria. L.			
1 triméstris. DC.	(li.w.) common annual.	S. Europe.	1633.	7. 9. H. ○.	Bot. mag. 109.
2 phoenicea. DC.	(cr.) scarlet.	Madeira.	1823.	6. 9. G. ℥.	Vent. malm. t. 120.
3 acerifólia. DC.	(li.) Maple-leaved.	Teneriffe.	—	— G. ℥.	Lois. h. amat. t. 322.
4 hispida. DC.	(ro.) hispid.	Algiers.	1804.	6. 7. G. ℥.	Desf. atl. 2. t. 171.
5 O'tibia. DC.	(pu.) downy-leaved.	France.	1570.	6. 10. G. ℥.	Cav. dis. 2. t. 32. f. 2.
6 unguiculáta. DC.	(pu.) clawed.	Greece.	1807.	7. 9. G. ℥.	Moris. ox. t. 17. f. 9.
7 micas. DC.	(pu.) glittering.	Spain.	1796.	6. 7. G. ℥.	Desf. atl. 2. t. 172.
8 lusitánica. DC.	(pu.) Portugal.	Portugal.	1731.	8. 9. G. ℥.	Bot. mag. 2269.
9 fláva. DC.	(ye.) yellow.	Sicily.	1823.	6. 8. H. ¼.	
10 plebéia. B.M.	(li.) vulgar.	N. S. W.	1820.	— G. ¼.	Bot. mag. 517.
Weinmanniána. Bes.					
11 thuringiaca. DC.	(pu.) large-flowered.	Germany.	1731.	7. 9. H. ¼.	Cav. dis. 2. t. 32. f. 3.
12 biénnis. DC.	(pu.) biennial.	Caucasus.	1823.	5. 8. H. ♂.	Bot. mag. 2226.
13 punctáta. DC.	(bh.) spotted-stalked.	Italy.	1800.	7. 9. H. ○.	Eng. bot. 1841.
14 marítima. DC.	(wh.) sea side.	S. Europe.	1597.	4. 6. G. ℥.	Cav. dis. 2. t. 32. f. 1.
15 trilóbata. DC.	(li.) three lobed.	Spain.	1759.	6. 7. G. ℥.	
16 arbórea. DC.	(li.) Tree Mallow.	Britain.	7. 10. H. ♂.	
17 neapolitána. DC.	(pu.) Neapolitan.	Naples.	1823.	— H. ¼.	
18 crética. DC.	(pu.) Cretan.	Crete.	1723.	7. 9. H. ○.	
19 lanceolátata. W.en.	(pu.) spear-leaved.	—	1823.	— H. ○.	
20 sylvestris. DC.	(pu.) wood.	Portugal.	—	— H. ○.	
21 ambígua. DC.	(pu.) ambiguous.	Naples.	1824.	— H. ○.	
22 móllis. Ehr.	(pu.) soft.	Egypt.	1825.	7. 10. H. ○.	
MALA'CHRA. DC.	MALA'CHRA.	Monadelphia Polyandria. L.			
1 rotundifólia. DC.	(ro.) round-leaved.	Brazil.	1823.	6. 9. S. ○.	Schrank. h. mon. t. 56
2 capitáta. DC.	(ye.) headed.	W. Indies.	1759.	8. 9. S. ○.	Cav. dis. 2. t. 33. f. 1. 2.
3 fasciáta. DC.	(ro.) fasciate.	Caracas.	1820.	— S. ○.	Jacq. ic. 3. t. 548.
4 trilóbata. DC.	(wh.) three-lobed.	—	—	— S. ○.	Cav. dis. 2. t. 33. f. 3.
5 radiáta. DC.	(pu.) rayed.	St. Domingo.	1794.	7. 9. S. ♂.	Jacq. ic. 3. t. 549.
6 alceafólia. DC.	(ye.) Hollyhock-ld.	Caracas.	1805.	8. 9. S. ○.	
7 heptaphylla. DC.	(ro.) many-flower'd.	Brazil.	1820.	6. 10. S. ¼.	
fasciáta. B. reg. 467. non DC.					
8 palmáta. DC.	(ye.) palmate.	—	1825.	— S. ○.	
UR'ENA. DC.	UR'ENA.	Monadelphia Polyandria. L.			
1 lobáta. DC.	(ro.) angular-leav'd.	China.	1731.	6. 7. S. ℥.	Dill. elt. t. 319. f. 412.

2	scabriúscula. DC.	(ro.) roughish.	India.	1820.	6. 7.	S. h.	Cav. dis. 6. t. 185. f. 1.
3	repánda. DC.	(ro.) netted-leaved.	—	—	—	S. h.	—
4	multifida. DC.	(ye.) multifid.	Mauritius.	1817.	10. 1.	S. h.	Cav. dis. 6. t. 184. f. 2.
5	tricúspis. DC.	(ye.) three-pointed.	—	1820.	6. 9.	S. h.	— 6. t. 183. f. 1.
6	reticulàta. DC.	(ye.) netted.	St. Domingo.	1816.	—	S. h.	— 6. t. 183. f. 2.
	americâna. L. non Sm.						
7	subtrilòba. DC.	(re.) various-leaved.	Brazil.	1823.	—	S. ⊙.	Schrank. h. mon. t. 79.
8	Swártzii. DC.	(re.) Swartz's.	Surinam.	—	S. h.	
	sinuata. Sw. americâna. Sm.						
9	sinuata. DC.	(re.) cut-leaved.	E. Indies.	1759.	7. 8.	S. h.	Cav. dis. 6. t. 185. f. 2.
10	muricâta. DC.	(re.) muricate-fruit'd	—	1823.	—	S. h.	
	PAVONIA. DC.	PAVONIA. Monadelphia Polyandria. L.					
1	spinifex. DC.	(ye.) prickly-seeded.	W. Indies.	1778.	7. 8.	S. h.	Bot. reg. 339.
2	Typhâlea. DC.	(wh.) globular-headed.	—	1823.	—	S. h.	Cav. dis. 2. t. 197.
3	ürens. DC.	(ro.) stinging.	Mauritius.	1801.	—	S. h.	Jacq. ic. 3. t. 522.
4	coccinea. DC.	(sc.) scarlet.	St. Domingo.	1816.	—	S. h.	Cav. dis. 3. t. 47. f. 1.
5	Columellâ. DC.	(pu.) angular-leaved.	Bourbon.	1807.	—	S. h.	— 3. t. 48. f. 3.
6	parviflôra. DC.	(st.) small-flowered.	—	1820.	—	S. h.	
7	racemosa. DC.	(ye.) naked-racemed.	Jamaica.	—	—	S. h.	Cav. dis. 3. t. 46. f. 1.
	spicata. Cav.						
8	præmorsa. DC.	(ye.) bitten-leaved.	C. B. S.	1774.	6. 8.	G. h.	Bot. mag. 436.
9	paniculâta. DC.	(ye.) paniced.	Caracas.	1823.	—	S. h.	Cav. dis. 3. t. 46. f. 2.
10	odorâta. DC.	(re.) fragrant.	E. Indies.	1807.	—	S. h.	
11	sidoídes. DC.	(re.) Sida like.	—	1820.	—	S. ♂.	
12	zeylânica. DC.	(bh.) Ceylon.	—	1790.	7. 9.	S. ⊙.	Burm. ind. t. 48. f. 2.
13	cancellâta. DC.	(re.) arrow-leaved.	Surinam.	1823.	—	S. ⊙.	Lin. sup. t. 311.
	MALVAVICUS. DC. ACHANIA. Monadelphia Polyandria. L.						
1	arboreus. DC.	(sc.) smooth-leaved.	Jamaica.	1714.	1. 12.	S. h.	Cav. dis. 3. t. 48. f. 1.
	Achania malvaviscus. W.						
2	mollis. DC.	(sc.) soft-leaved.	Mexico.	1780.	—	S. h.	Bot. reg. 11.
	Achania mollis. B.R.						
3	pilosa. DC.	(sc.) hairy.	Jamaica.	1780.	10. 11.	S. h.	Lodd. bot. cab. 829.
	LEBRETTONIA. DC. LEBRET'ONIA. Monadelphia Polyandria. L.						
	coccinea. DC.	(sc.) scarlet.	Brazil.	1823.	6. 8.	S. h.	Schrank. h. mon. t. 90.
	HIBISCUS. DC.	HIBÍSCUS. Monadelphia Polyandria. L.					
1	liliiflorus. DC.	(va.) Lily-flowered.	Bourbon.	1822.	8. 12.	S. h.	Cav. dis. 3. t. 57. f. 1.
2	Boryanus. DC.	(wh.) Bory St. Vincent's.	—	1826.	6. 10.	S. h.	
3	pedunculatus. DC.	(pi.) long-peduncled.	C. B. S.	1812.	5. 12.	S. h.	Bot. reg. 231.
4	Lámpas. DC.	(st.) three-pointed.	E. Indies.	1806.	10. 2.	S. h.	Cav. dis. 3. t. 56. f. 2.
5	membranaceus. DC.	(y.) leafy-calyxed.	—	1816.	—	S. h.	— 3. t. 57. f. 2.
6	spiralis. DC.	(y.fl.) spiral.	Mexico.	1826.	6. 9.	G. h.	Cavan. ic. 2. t. 162.
7	tubulosus. DC.	(y.p.) tubular.	E. Indies.	1796.	7. 10.	S. ♂.	Cav. dis. 3. t. 68. f. 2.
8	pentacarpus. DC.	(bh.) angular-fruited.	Venice.	1752.	—	G. ♀.	Jacq. ic. 1. t. 143.
9	virginicus. DC.	(ro.) Virginian.	Virginia.	1798.	—	H. ♀.	— 1. t. 142.
10	pentaspérminus. DC.	(ye.) five-seeded.	Jamaica.	1825.	—	S. h.	
11	Mánihot. DC.	(y.p.) palmated.	E. Indies.	1712.	7. 9.	S. h.	
12	palmatus. C.	(ye.p.) palmate-leav'd.	S. America.	—	S. h.	
13	fículneus. DC.	(wh.) Fig-leaved.	Ceylon.	1732.	6. 7.	S. h.	— 3. t. 52. f. 2.
	sinuatus. C.						
14	ficulneoides. B.R.	(st.) Fig-like.	S. America.	1824.	6. 8.	S. h.	Bot. reg. 938.
15	tetraphyllus. DC.	(ye.) four-leaved.	E. Indies.	1807.	7. 10.	S. ⊙.	
16	platanifolius.	(ye.) Plane-leaved.	W. Indies.	1820.	—	S. h.	Pl. sel. h. ber. t. 1.
	acerifolius. DC. non P.L. Pavonia platanifolia. L. en.						
17	Ròsa-sinensis. DC.	(sc.) China Rose.	E. Indies.	1731.	1. 12.	S. h.	Bot. mag. 158.
	β rùbro-plènus.	(sc.) double-red.	—	—	—	S. h.	Lod. bot. cab. 995.
	γ cárneo-plènus.	(sa.) double-salmon.	—	—	—	S. h.	
	δ variegátus.	(sc.w.) double-striped.	—	—	—	S. h.	Lod. bot. cab. 963.
	ε flávo-plènus.	(co.) double-buff.	—	—	—	S. h.	— 513.
	ζ lütœo-plènus.	(ye.) double-yellow.	—	1822.	—	S. h.	— 932.
18	acerifolius. P.L.	(va.) Maple-leaved.	China.	1798.	3. 6.	G. h.	Sal. par. lond. 22.
19	syriacus. DC.	(va.) Althaea frutex.	Syria.	1596.	8. 9.	H. h.	Bot. mag. 83.
	α purpureus.	(pu.) purple-flowered.	—	—	—	H. h.	
	β ruber.	(re.) red-flowered.	—	—	—	H. h.	
	γ álbus.	(wh.) white-flowered.	—	—	—	H. h.	
	δ variegátus.	(w.p.) striped-flowered.	—	—	—	H. h.	
	ε marginátus.	(pu.) variegated.	—	—	—	H. h.	
	ζ álbo-plènus.	(wh.) double-white.	—	—	—	H. h.	
	η purpureo-plènus.	(p.) double-purple.	—	—	—	H. h.	
20	prüiens. DC.	(y.pu.) stinging.	E. Indies.	1804.	7. 9.	S. ♂.	Andr. deposit. 498.

MALVACEÆ.

21 suratténis. DC. (<i>y.p.</i>)	prickly-stalk'd.	E. Indies.	1731. 7. 9. S. ⓧ.	Bot. mag. 1356.	
22 radiátus. DC. (<i>st.p.</i>)	rayed.	_____	1790. 6. 8. S. ⓧ.	_____ 1911.	
23 furcátus. DC. (<i>y.p.</i>)	forked.	Brazil.	1816. 7. 9. S. ⓧ.	Bot. reg. 878.	
24 unidéns. B.R. (<i>y.p.</i>)	one-toothed.	Carolina.	1824. 11. 4. S. ⓧ.	Bot. reg. 860.	
25 scáber. DC. (<i>st.</i>)	scabrous.	S. America.	1810. 7. 9. G.w. ⓣ.	Cav. dis. 3. t. 51. f. 1.	
26 strigósus. B.R. (<i>ro.</i>)	strigose.	Brazil.	1822. 1. 12. S. ⓧ.	_____ 3. t. 53. f. 2.	
27 bifúrcatus. DC. (<i>pu.</i>)	two-forked.	W. Indies.	1826. — S. ⓧ.	Jacq. ic. 3. t. 551.	
28 trilobus. DC. (<i>co.</i>)	three-lobed.	_____	1824. — S. ⓧ.	Bot. reg. 29.	
29 diversifolius. DC. (<i>y.p.</i>)	different-leav'd.	E. Indies.	1798. 6. 7. S. ⓧ.	Roxb. cor. 2. t. 190.	
30 heterophyllus. DC. (<i>w.p.</i>)	various-leaved.	N. S. W.	1808. 8. 9. G. ⓧ.	Rheed. mal. 6. t. 46.	
31 cannabinus. DC. (<i>y.p.</i>)	Hemp-leaved.	E. Indies.	1759. 6. 7. S. ⓧ.	Tozz. mus. d. fir. 2. t. 7.	
32 vitifolius. DC. (<i>y.p.</i>)	Vine-leaved.	_____	1690. 7. 10. S. ⓧ.	Swt. br. fl. gar. 286.	
33 obtusifolius. DC. (<i>y.p.</i>)	blunt-leaved.	_____	1820. — S. ⓧ.	Cav. dis. 3. t. 65. f. 2.	
34 Bámmia. L.en. (<i>y.p.</i>)	Egyptian.	Egypt. 6. 7. S. ⓧ.	Swt. br. fl. gar. 277.	
35 esculéntus. L.en. (<i>y.p.</i>)	eatable.	India.	1692. — S. ⓧ.	Wend. h. her. 4. t. 24.	
36 longifolius. DC. (<i>ye.</i>)	long-leaved.	E. Indies.	1820. 7. 9. S. ⓧ.	Bot. mag. 2385.	
37 moschéutus. DC. (<i>w.p.</i>)	swamp.	N. America. 8. 10. H.w. ⓣ.	_____ 360.	
38 palústris. DC. (<i>pu.</i>)	marsh.	_____	1759. 7. 9. H.w. ⓣ.	Cav. dis. 3. t. 60. f. 1.	
39 róseus. DC. (<i>ro.</i>)	rose-colour'd.	S. Europe.	1824. 8. 10. H.w. ⓣ.	Cav. dis. 3. t. 61. f. 1.	
40 aquáticus. DC. (<i>wh.</i>)	water.	Etruria.	1820. 7. 9. H.w. ⓣ.	Bot. reg. 589.	
41 incánu. DC. (<i>st.</i>)	hoary.	Carolina.	1806. 9. G.w. ⓣ.	Cav. dis. 3. t. 58. f. 1.	
42 militaris. DC. (<i>pu.</i>)	smooth.	Louisiana.	1804. 8. 9. G.w. ⓣ.	_____ 3. t. 62. f. 2.	
43 speciosus. DC. (<i>cr.</i>)	superb.	Carolina.	1778. 6. 9. G.w. ⓣ.	Bot. reg. 230.	
44 grandiflorus. DC. (<i>fl.</i>)	great-flower'd.	Georgia.	1823. — G.w. ⓣ.	Cav. dis. 3. t. 67. f. 3.	
45 ferrugineus. DC. (<i>re.</i>)	rusty.	Madagascar.	1820. — S. ⓧ.	Bot. reg. 337.	
46 lunariifolius. DC. (<i>st.p.</i>)	Lunaria-leav'd.	E. Indies. — S. ⓧ.	Cav. dis. 3. t. 69. f. 3.	
47 athiòpicus. DC. (<i>ye.</i>)	dwf. wedge-l'd.	C. B. S.	1774. 8. G. ⓣ.	Cav. dis. 3. t. 66. f. 1.	
48 mutabilis. DC. (<i>va.</i>)	changeable.	E. Indies.	1690. 10. 12. S. ⓧ.	Bot. mag. 209.	
β floré-pleno. (<i>va.</i>)	double-flowered.	_____ — S. ⓧ.	Bot. reg. 806.	
49 clypéatus. DC. (<i>r.y.</i>)	shield-capsuled.	Jamaica.	1759. 7. 8. S. ⓧ.	Cav. dis. 3. t. 62. f. 2.	
50 Abelmoschus. DC. (<i>ye.</i>)	Musk Okro.	India.	1640. 7. 9. S. ⓧ.	Bot. reg. 875.	
51 eriocárpus. DC. (<i>ye.</i>)	woolly-capsul'd.	E. Indies.	1820. — S. ⓧ.	Cav. dis. 3. t. 62. f. 2.	
52 phoeniceus. B.R. (<i>cr.</i>)	crimson-flow'd.	S. America.	1796. — S. ⓧ.	Bonp. nav. t. 29.	
53 hírtus. L. (<i>sc.</i>)	hairy.	E. Indies. 2. 10. S. ⓧ.	Cav. dis. 3. t. 70. f. 2.	
54 Rôsa malabárica. (<i>sc.</i>)	Malabar Rose.	_____	1818. 8. 10. S. ⓧ.	Bot. reg. 608.	
55 rhombifolius. DC. (<i>wh.</i>)	rhomb-leaved.	_____	1823. — S. ⓧ.	Cav. dis. 3. t. 55. f. 2.	
56 gossypinus. DC. (<i>re.</i>)	cottony.	C. B. S.	1818. — G. ⓧ.	Bot. reg. 917.	
57 micránthus. DC. (<i>ye.</i>)	small-flowered.	E. Indies.	1794. — S. ⓧ.	_____ 232.	
58 Trionum. DC. (<i>st.p.</i>)	BladderKetmia.	Italy.	1596. 6. 9. H. ⓧ.	Sloan. jam. 1. t. 234. f. 4.	
59 hispidus. B.R. (<i>y.p.</i>)	hispid.	C. B. S. — G. ⓧ.	Cav. dis. 3. t. 70. f. 3.	
60 africanus. R.B. (<i>st.p.</i>)	African.	Africa.	1826. 6. 10. H. ⓧ.	Bot. reg. 233.	
61 Richardsoni. (<i>st.</i>)	Richardson's.	N. S. W.	1822. — G. ⓧ.	Cav. dis. 3. t. 68. f. 1.	
62 vesicárius. DC. (<i>st.p.</i>)	African.	Africa.	1713. 7. 8. H. ⓧ.	Bonp. nav. t. 29.	
63 Sabdariffa. DC. (<i>y.ro.</i>)	Indian.	E. Indies.	1596. 6. 9. S. ⓧ.	Cav. dis. 3. t. 70. f. 2.	
64 digitátus. DC. (<i>ye.r.</i>)	fingered.	Brazil.	1818. — S. ⓧ.	Bot. reg. 608.	
β Kerriánus. DC. (<i>w.r.</i>)	Ker's.	Rio Janeiro.	— — — — —	Cav. dis. 3. t. 55. f. 2.	
65 tricúspis. DC. (<i>ye.</i>)	sharp-pointed.	Society Isl.	1820. — S. ⓧ.	Bot. reg. 917.	
66 racemosus. B.R. (<i>y.p.</i>)	racemed.	Nepaul.	1824. 8. 2. G. ⓧ.	_____ 232.	
67 tiliaceus. DC. (<i>ye.p.</i>)	Lime-tree-l'd.	E. Indies.	1739. 7. 8. S. ⓧ.	Sloan. jam. 1. t. 234. f. 4.	
68 elátus. DC. (<i>ye.p.</i>)	tall.	Jamaica.	1790. — S. ⓧ.	Cav. dis. 3. t. 70. f. 3.	
69 abutiloídes. DC. (<i>y.p.</i>)	smooth-leaved.	S. America.	1820. — S. ⓧ.	Bot. reg. 233.	
70 flavescens.	(<i>ye.</i>)	flavescient.	Brazil.	1822. 6. 9. S. ⓧ.	Cav. dis. 3. t. 68. f. 1.
71 borbónicus. DC. (<i>ye.</i>)	Bourbon.	I. Bourbon.	1826. — S. ⓧ.	Bot. reg. 233.	
72 Macleánius. BF. (<i>ye.</i>)	Macleay's.	W. Indies.	1827. 8. 9. S. ⓧ.	Cav. dis. 3. t. 68. f. 1.	
73 macrophyllus. DC. (<i>ye.</i>)	large-leaved.	E. Indies.	1820. — S. ⓧ.	Bot. reg. 233.	
74 collinus. H.B. (<i>ye.</i>)	hill.	_____	1822. — S. ⓧ.	Cav. dis. 3. t. 68. f. 1.	
75 cancellátus. H.B. (<i>ye.</i>)	netted.	Nepaul.	1823. 10. 2. S. ⓧ.	Bot. reg. 233.	
76 truncátus. H.B. (<i>ye.</i>)	truncate.	E. Indies.	— 8. 10. S. ⓧ.	Cav. dis. 3. t. 68. f. 1.	
77 scándens. H.B. (<i>ye.</i>)	climbing.	_____	1822. — S. ⓧ.	Bot. reg. 233.	
78 strictus. H.B. (<i>ye.</i>)	upright.	_____	1809. 8. 10. S. ⓧ.	Cav. dis. 3. t. 68. f. 1.	
79 gangéticus. L.en. (<i>ye.</i>)	dwarf.	_____	1804. 5. 10. S. ⓧ.	Bot. reg. 233.	
80 setosus. H.B. (<i>ye.</i>)	bristly.	_____	1820. — S. ⓧ.	Cav. dis. 3. t. 68. f. 1.	
81 tortuósus. H.B. (<i>ye.</i>)	twisted.	_____	1808. 6. 9. S. ⓧ.	Bot. reg. 233.	
82 pentaphylloides. H.B. (<i>ye.</i>)	five-leaved.	China.	1825. 7. 9. S. ⓧ.	Cav. dis. 3. t. 68. f. 1.	
83 pungens. H.B. (<i>ye.</i>)	pungent.	Nepaul.	— — — — —	Andr. rep. 286.	
84 fràgrans. H.B. (<i>ye.</i>)	fragrant.	E. Indies.	1826. — — — — —	Lagun'æa Patersònia. Bot. mag. 769. squâmea. Vent. malm. t. 42.	
85 Patersòni. DC. (<i>fl.</i>)	Norfolk Island.	Norfolk Isl.	1792. 6. 8. G. ⓧ.		

THESP'ESIA. DC.	THESP'ESIA. Monadelphia Polyandria. L.				
" populnea. DC. (y.p.) Poplar-leaved. E. Indies.	1770. S. h.	Cav. dis. 3. t. 56. f. 1.			
<i>Hibiscus populneus</i> . W.					
GOSSYPIUM. DC.	COTTON-TREE. Monadelphia Polyandria. L.				
1 herbaceum. DC. (y.p.) common.	E. Indies. 1594. 7. S. ♂.	Roxb. corom. t. 269.			
2 indicum. DC. (y.p.) Indian. 1694. 7. 8. S. ♀.	Cav. dis. 6. t. 169.			
3 arboreum. DC. (y.p.) tree. 1805. —— S. ♀.	—— 6. t. 195.			
4 vitifolium. DC. (y.p.) Vine-leaved. 1731. —— S. ♂.	—— 6. t. 166.			
5 hirsutum. DC. (ye.) hairy.	S. America. 1731. —— S. ♂.	—— 6. t. 167.			
6 eglandulosum. DC. (ye.) glandless. —— S. ♂.				
7 religiosum. DC. (bh.) spotted-bark'd.	E. Indies. 1777. —— S. ♀.	Cav. dis. 6. t. 164. f. 1.			
8 latifolium. DC. (ye.p.) broad-leaved. —— S. ♀.	Murr. c. gæt. p. 32. t. 1.			
9 barbadense. DC. (y.p.) Barbadoes.	Barbadoes. 1759. 8. 10. S. ♂.	Bot. reg. 84.			
10 obtusifolium. H.B. (y.p.) blunt-leaved.	E. Indies. 1822. 7. 8. S. ♀.				
REDO'UTEA. DC.	REDO'UTEA. Monadelphia Polyandria. L.				
heterophylla. DC. (st.p.) various-leaved.	St. Thomas. 1824. 7. 9. S. ⊖.	Vent. cels. t. 11.			
LOPI'MIA. DC.	LOPI'MIA. Monadelphia Polyandria. L.				
malacophylla. DC. (sc.) Mallow-leaved. Brazil.	1824. S. ♀.	Pl. sel. h. ber. t. 30.			
<i>Sida malacophylla</i> . L. en.					
PAL'AVIA. DC.	PAL'AVIA. Monadelphia Polyandria. L.				
1 malvæfolia. DC. (pu.) Mallow-leaved. Lima.	1794. 6. 8. H. ⊖.	Cav. dis. 1. t. 11. f. 4.			
2 moschata. DC. (pu.) musk-scented. 1822. —— H. ⊖.	—— 1. t. 11. f. 5.			
CRISTARIA. Ph.	CRISTARIA. Monadelphia Polyandria. L.				
? coccinea. Ph. (sc.) scarlet.	Missouri. 1811. 7. 9. H. ♀.	Bot. mag. 1673.			
<i>Mala coccinea</i> . Nut. <i>Sida?</i> coccinea. DC.					
AN'ODA. DC.	AN'ODA. Monadelphia Polyandria. L.				
1 hastata. DC. (bl.wh.) halberd-leav'd. Mexico.	1799. 6. 7. S. ⊖.	Bot. rep. 588.			
2 triloba. DC. (pu.) three-lobed. 1720. 7. 9. G. ⊖.	Cav. dis. 1. t. 10. f. 3.			
<i>Sida cristata</i> . W. non B.M.					
3 Dilleniâna. DC. (ro.) Dillenius's. 1725. 6. 11. G. ⊖.	Cav. dis. 1. t. 11. f. 1.			
<i>Sida cristata</i> . B.M. 330. non W.					
4 triangulâris. DC. (pu.) triangular-l'd. 1824. 7. 9. G. ⊖.				
5 incarnâta. DC. (fl.) flesh-color'd. 1825. —— G. ♂.				
6 acerifolia. DC. (bl.) Maple-leaved. 1809. 1. 12. S. ♂.	Bot. mag. 1541.			
<i>Sida hastata</i> . B.M. non W. centrota. s.s.					
7 parviflora. DC. (st.) small-flowered. New Spain.	1823. 6. 10. G. ⊖.	Cav. ic. 5. t. 431.			
PERI'PTERA. DC.	PERI'PTERA. Monadelphia Polyandria. L.				
punicea. DC. (cr.) crimson.	New Spain. 1813. 6. 9. S. ♀.	Bot. mag. 1644.			
<i>Sida periptera</i> . B.M. <i>Anoda punicea</i> . Lag.					
NUTTALLIA. B.A.	NUTTALLIA. Monadelphia Polyandria.				
1 digitata. B.A. (d.p.) digitate-leaved. N. America.	1824. 7. 10. H. ♀.	Swt. br. fl. gar. 129.			
2 pedata. B.A. (d.pu.) pedate-leaved. —— H. ♀.	Hook. ex. flor. 172.			
SIDA. Kth.	SIDA. Monadelphia Polyandria. L.				
1 linifolia. DC. (wh.) Flax-leaved.	S. America. 1822. 6. 9. S. ♀.	Cav. dis. 1. t. 2. f. 1.			
2 angustifolia. DC. (ye.) narrow-leaved.	Bourbon. 1726. 7. 9. S. ♀.	—— 1. t. 2. f. 2.			
3 linearis. DC. (ye.) linear-leaved.	New Spain. 1820. 6. 10. G. ♀.	Cav. ic. 4. t. 312. f. 1.			
4 viminea. L. en. (ye.) slender.	Brazil. 1823. —— S. ♀.				
5 spinosa. DC. (ye.) prickly.	E. Indies. 1880. —— S. ♂.	Cav. dis. 1. t. 1. f. 9.			
6 acuta. DC. (ye.) acute-leaved. 1827. —— S. ♀.	—— 1. t. 2. f. 3.			
7 ábla. DC. (wh.) white-flowered. 1732. 6. 7. S. ⊖.	Dill. elt. 2. t. 171. f. 210.			
8 stipulata. DC. (ye.) large-stipuled.	Mauritius. 1820. S. ♀.	Cav. dis. 1. t. 3. f. 10.			
9 jamaicensis. DC. (ye.) Jamaica.	Jamaica. 1818. 6. 9. S. ♀.	—— 1. t. 2. f. 5.			
10 bracteolata. DC. (ye.) bracteolate.	S. America. —— —— S. ♀.	Cav. dis. 5. t. 134. f. 1.			
11 carpinifolia. DC. (ye.) Hornbeam-l'd.	Canaries. 1774. 7. 9. G. ♀.				
12 carpinoides. DC. (ye.) Hornbeam-like. —— S. ♀.				
13 semicrenata. L.en.(y.) semicrenate.	Manila. 1826. —— S. ♀.				
14 acrântha. L.en. (ye.) close-flower'd.	Brazil. —— —— S. ♀.				
15 spiræifolia. L.en. (ye.) Spiræa-leaved. —— S. ♀.				
16 Schrankii. DC. (ye.) Schrank's.	Brazil. 1824. —— S. ♀.				
brasiliæ. L. en.					
17 betulina. H. (ye.) Birch-leaved.	S. America. 1823. —— S. ♀.				
18 erosa. DC. (ye.) jagged.	Brazil. —— —— S. ♀.				
19 ciliâris. DC. (cr.) ciliated.	St. Domingo. 1798. —— S. ♀.	Cav. dis. 1. t. 3. f. 9.			
20 alnifolia. DC. (ye.) Alder-leaved.	E. Indies. 1732. —— S. ♂.	Dill. elt. t. 172. f. 211.			
21 sessiliiflora. B.M. (ye.) sessile-flower'd.	S. America. 1826. 11. 2. S. ♀.	Bot. mag. 2857.			
22 frutescens. DC. (ye.) frutescent.	Mauritius. 1824. 6. 9. S. ♀.	Cav. dis. 1. t. 10. f. 1.			
23 bicolor. DC. (r.y.) two-colour'd.	New Spain. 1826. —— S. ♀.	Cav. ic. 4. t. 311.			
24 canariensis. DC. (wh.) Canary Tea.	Canaries. —— S. ♀.	Cav. dis. 1. t. 3. f. 8.			
ábla. Cav. non L.					

25 rhombifolia. DC. (<i>ye.</i>) rhomboid-l'd.	Brazil.	1732.	6. 8.	S. $\frac{1}{2}$.	Cav. dis. 1. t. 3. f. 12.
26 rhombóidea. DC. (<i>ye.</i>) rhomboid.	E. Indies.	1818.	—	S. $\frac{1}{2}$.	Cav. dis. 1. t. 3. f. 4.
27 retusa. DC. (<i>ye.</i>) retuse-leaved.	—	—	—	S. $\frac{1}{2}$.	
28 recisa. DC. (<i>ye.</i>) notched.	Brazil.	1823.	—	S. $\frac{1}{2}$.	
29 calyxhymènia. DC. (<i>st.</i>) New Holland.	N. Holland.	1820.	—	G. $\frac{1}{2}$.	
30 globiflora. B.M. (<i>st.</i>) globe-flowered.	Mauritius.	1825.	10. 12.	S. $\frac{1}{2}$.	Bot. mag. 2821.
31 maculatâ. DC. (<i>ye.r.</i>) corky. <i>suberosa</i> . L. her. stirp. t. 54.	St. Domingo.	1798.	S. $\frac{1}{2}$.	Cav. dis. 1. t. 3. f. 7.
32 acuminatâ. DC. (<i>ye.</i>) acuminate.	St. Domingo.	1805.	7. 9.	S. $\frac{1}{2}$.	
33 pilosa. DC. (<i>ye.</i>) pilose.	W. Indies.	1793.	—	S. $\frac{1}{2}$.	Cav. dis. 1. p. 9. t. 8.
34 hùmilis. DC. (<i>ye.</i>) dwarf. <i>unilocularis</i> . L'Herit. stirp. 1. t. 53. bis.	E. Indies.	1800.	7. 8.	S. $\frac{1}{2}$.	Cav. dis. 5. t. 134. f. 2.
35 rotundifolia. DC. (<i>pu.</i>) round-leaved.	Bourbon.	1826.	—	S. \odot .	Cav. dis. 1. t. 3. f. 6.
36 cordifolia. DC. (<i>ye.</i>) heart-leaved.	E. Indies.	1732.	6. 9.	S. \odot .	— 1. t. 3. f. 2.
37 multiflora. DC. (<i>ye.</i>) many-flowered.	Brazil.	1825.	—	S. $\frac{1}{2}$.	— 1. t. 3. f. 3.
38 arguta. DC. (<i>ye.</i>) sharp-leaved.	W. Indies.	1732.	7. 8.	S. $\frac{1}{2}$.	
39 althæfolia. DC. (<i>co.</i>) Althæa-leaved.	—	1818.	—	S. $\frac{1}{2}$.	Sloan. his. 1. t. 136. f. 2.
40 ûrens. DC. (<i>ye.</i>) stinging.	—	1781.	—	S. $\frac{1}{2}$.	Cav. dis. 1. t. 2. f. 7.
41 verticillatâ. DC. (<i>ye.</i>) whorl-flowered.	Brazil.	1823.	—	S. $\frac{1}{2}$.	— 1. t. 1. f. 12.
42 paniculatâ. DC. (<i>d.pu.</i>) dark purple. <i>atrosanguinea</i> . Jacq. ic. 1. t. 136.	S. America.	1795.	7. 9.	S. $\frac{1}{2}$.	— 1. t. 12. f. 5.
43 fruticosa. DC. (<i>wh.</i>) shrubby.	Jamaica.	6. 10.	S. $\frac{1}{2}$.	
44 villosa. DC. (<i>ye.</i>) villous.	S. America.	—	S. $\frac{1}{2}$.	
45 verruculatâ. DC. (<i>ye.</i>) wart-stem'd. arguta. F. in L. en. non Swartz.	Brazil.	1827.	5. 9.	S. $\frac{1}{2}$.	
46 graviéolens. DC. (<i>ye.</i>) strong-scented.	E. Indies.	1820.	6. 9.	S. $\frac{1}{2}$.	
47 purpurascens. DC. (<i>pu.</i>) purplish.	Brazil.	—	7. 9.	S. $\frac{1}{2}$.	
48 pâtens. DC. (<i>pu.</i>) spreading.	Abyssinia.	1806.	—	S. $\frac{1}{2}$.	
49 conférta. DC. (<i>st.</i>) close-flowered.	Brazil.	1826.	—	S. $\frac{1}{2}$.	
50 hirsuta. DC. (<i>wh.</i>) hairy.	—	S. $\frac{1}{2}$.	
51 lasiostega. DC. (<i>ye.</i>) woolly.	Brazil.	1823.	—	S. $\frac{1}{2}$.	
52 pentacárpos. DC. (<i>ye.</i>) five-capsuled.	E. Indies.	1823.	8. 10.	S. \odot .	
53 triloba. DC. (<i>wh.</i>) three-lobed.	C. B. S.	1794.	—	G. $\frac{1}{2}$.	Jacq. schœn. 2. t. 142.
54 ricinoïdes. DC. (<i>bh.</i>) Ricinus-like. <i>palmata</i> . Cav. dis. 1. t. 3. f. 3.	Peru.	1826.	7. 10.	H. \odot .	L'Her. stirp. 1. t. 56.
55 jatrophoides. DC. (<i>vi.</i>) Jatropha-like. <i>palmata</i> . Jacq. ic. rar. 3. t. 547.	Peru.	1787.	8.	S. $\frac{1}{2}$.	L. Her. stirp. 1. t. 56.
56 malvaeflora. B.R. (<i>bh.</i>) Mallow-flow'd.	Columbia.	1826.	9. 11.	H. $\frac{1}{2}$.	Bot. reg. 1036.
57 Napæa. DC. (<i>u.h.</i>) smooth. <i>Napæa laevis</i> . L.	Virginia.	1743.	8. 9.	H. $\frac{1}{2}$.	Bot. mag. 2193.
58 dioica. DC. (<i>wh.</i>) rough.	Virginia.	1759.	8. 9.	H. $\frac{1}{2}$.	Cav. dis. 5. t. 132. f. 2.
BASTA'R'DIA. Kth. BASTA'R'DIA. Monadelphia	Monadelphia	1822.	—	Polyandria. L.	
1 viscosa. K.S. (<i>ye.</i>) viscous. <i>Sida viscosa</i> . DC.	S. America.	1822.	7. 8.	S. $\frac{1}{2}$.	Sloan. his. 1. t. 139. f. 4.
2 foetida. (<i>ye.</i>) stinking. <i>Sida foetida</i> . DC. <i>Sida viscosa</i> . L.H.	Peru.	1795.	7. 8.	S. \odot .	L. Her. stirp. 1. t. 53.
GA'YA. Kth. GA'YA. Monadelphia Polyandria. L.					
1 occidentalis. H.S. (<i>vi.</i>) downy. <i>Sida occidentalis</i> . DC.	America.	1732.	7. 8.	S. \odot .	Cav. dis. 1. t. 4. f. 3.
2 hermannioïdes. K.S. (<i>ye.</i>) Hermannia-like.	Mexico.	1826.	4. 9.	G. $\frac{1}{2}$.	H. et B. n. gen. 5. t. 475.
3 disticha. (<i>ye.</i>) two-ranked. <i>Sida disticha</i> . Cav.	New Spain.	1820.	6. 9.	G. $\frac{1}{2}$.	Cav. ic. 5. t. 432.
4 nütans. (<i>ye.</i>) nodding. <i>Sida nütans</i> . DC. <i>calyptrata</i> . Cav.	Peru.	1825.	—	G. $\frac{1}{2}$.	L. Her. stirp. 1. t. 57.
AB'UTILON. Kth. AB'UTILON. Monadelphia Polyandria. L.					
1 periplocifolium. (<i>st.</i>) Periploca-l'd. <i>Sida periplocifolia</i> . DC.	India.	1691.	7. 8.	S. $\frac{1}{2}$.	Cav. dis. 1. t. 5. f. 2.
2 ferrugineum. K.S. (<i>ye.</i>) ferruginous.	Peru.	1822.	7. 8.	S. $\frac{1}{2}$.	
3 hernandioïdes. (<i>ye.</i>) Hernandia-l'd.	Hispaniola.	1798.	S. $\frac{1}{2}$.	L. Her. stirp. 1. t. 58.
4 lucianum. (<i>wh.</i>) St. Lucian. <i>Sida luciana</i> . DC.	St. Lucia.	1822.	S. $\frac{1}{2}$.	
5 Lechenaultianum. (<i>ye.</i>) Lechenault's. <i>Sida Lechenaultiana</i> . DC.	E. Indies.	1820.	S. $\frac{1}{2}$.	
6 nudiflorum. (<i>ye.</i>) naked-flower'd.	Peru.	1731.	5. 6.	S. $\frac{1}{2}$.	L. Her. stirp. 1. t. 59.
7 polyanthon. (<i>ye.</i>) many-flowered. <i>Sida polyantha</i> . DC.	1822.	—	S. $\frac{1}{2}$.	
8 contráctum. (<i>ye.</i>) contracted. <i>Sida contrácta</i> . L.en.	Madagascar.	1826.	5. 7.	S. $\frac{1}{2}$.	
9 áuritum. (<i>co.</i>) eared-stipuled.	Java.	1820.	12. 3.	S. $\frac{1}{2}$.	Bot.
	<i>Sida áurita</i> . B.M.				

10	triquètrum.	(ye.) triangular.	W. Indies.	1775.	7. 8.	S. h.	Jacq. vind. 2. t. 118.
11	incānum.	(ye.) hoary.	Sandwich Isl.	1821.	—	S. h.	
	<i>Sida incāna</i> . L. en.						
12	pulchellum.	(wh.) neat.	N. Holland.	—	—	S. h.	Bonpl. nav. t. 2.
13	umbellātum.	(ye.) umbelled.	Jamaica.	1788.	7. 9.	S. ♂.	Jacq. vind. t. 56.
14	gigāntēum.	(ye.) gigantic.	Caracas.	1823.	...	S. h.	Jacq. schœnb. 2. t. 141.
15	reflexūm.	(re.) reflex-flowered.	Peru.	1799.	7. 8.	S. h.	Cav. dis. 1. t. 7.
16	crispūm.	(wh.) curled.	America.	1726.	—	H. ⓧ.	— 1. t. 7. f. 1.
17	virgātūm.	(ye.) twiggy.	Peru.	1820.	—	S. h.	Cav. ic. 1. t. 73.
18	arbōreum.	(pu.) great-flowered.	—	1772.	—	S. h.	Lher. stirp. 1. t. 63.
19	Mauritiānum.	(ye.) hairy-capsuled.	Mauritius.	1789.	7. 9.	S. h.	Jacq. ic. 1. t. 137.
	<i>Sida planiflora</i> . Cav. dis. 1. t. 7. f. 4.						
20	tiliaefolium.	(ye.) Lime-tree-ld.	China.	1820.	—	S. ⓧ.	Jacq. f. ecl. 1. t. 35.
21	mūticūm.	awnless.	Egypt.	1825.	8. 10.	H. ⓧ.	
	<i>Sida mūtica</i> . DC.						
22	Avicénnae. g.	(ye.) broad-leaved.	S. France.	1596.	6. 8.	H. ⓧ.	Houtt. syst. 8. t. 61.
	<i>Sida Abūtilon</i> . DC.						
23	americānum.	(ye.) American.	Jamaica.	1730.	6. 8.	S. ⓧ.	
	<i>Sida americana</i> . DC.						
24	asiāticūm.	(ye.) small-flowered.	E. Indies.	1768.	7. 8.	S. ⓧ.	Cav. dis. 1. t. 7. f. 2.
25	populiſolium.	(ye.) Poplar-leaved.	—	1796.	—	S. ⓧ.	— 1. t. 7. f. 9.
26	hírtūm.	(ye.) hairy.	—	1820.	—	S. ⓧ.	Rumph. amb. 4. t. 10?
	<i>Sida hírta</i> . DC.						
27	mölle.	(ye.) large-leaved.	Peru.	1817.	10. 12.	S. h.	Bot. mag. 2759.
	<i>Sida móllis</i> . DC. <i>grandifolia</i> . Bot. reg. 360.						
28	mollissimum.	(st.) soft-leaved.	Peru.	1789.	6. 7.	S. ♂.	Cav. dis. 2. t. 14. f. 1.
29	permölle.	(ye.) woolly-leaved.	—	1818.	—	S. h.	
	<i>Sida permóllis</i> . DC.						
30	cornūtūm.	(ye.) horned.	S. America.	1827.	—	S. h.	
	<i>Sida cornūta</i> . W. enum.						
31	índicum.	(ye.) Indian.	E. Indies.	1731.	7. 8.	S. ⓧ.	Cav. dis. 1. t. 7. f. 10.
32	vesicarium.	(st.) bladdered.	Mexico.	1796.	—	S. h.	— 2. t. 14. f. 3.
33	gláucum.	(ye.) glaucous.	Senegal.	1822.	—	S. h.	Cav. ic. 1. t. 11.
34	álbidūm.	(ye.) white-leaved.	Canaries.	—	—	G. h.	
	<i>Sida albida</i> . DC.						
35	mollicómum.	(ye.) soft.	1804.	7. 9.	S. h.	
	<i>Sida sericea</i> . Cav? <i>Sida mollicòma</i> . DC.						
36	Somneriānum.	(ye.) Sonnerat's.	C. B. S.	1806.	6. 7.	G. ♂.	Cav. dis. 1. t. 6. f. 4.
	<i>LAGUN'EA</i> . DC.	LAGUN'EA. <i>Monadelphus Polyandria</i> . L.					
1	lobāta. DC.	(wh.) Maple-leaved.	Bourbon.	1787.	7. 8.	S. ⓧ.	Cav. dis. 5. t. 136. f. 1.
2	sinuāta. DC.	(wh.) sinuate.	—	S. ⓧ.	

ORDO XXIX.

BOMBACEÆ. DC. prodr. 1. p. 475.

HELI'CTERES. DC. SCREW-TREE. Monadelphia Dodecandria. L.

1	Isora. DC.	(re.) great-fruited.	E. Indies.	1733.	6. 7.	S. h.	Bot. mag. 2061.
2	baruénis. DC.	(wh.) small-fruited.	W. Indies.	1739.	9. 10.	S. h.	Jacq. amer. t. 149.
3	jamaicénis. DC.	(wh.) broad-leaved.	—	—	6. 7.	S. h.	Jacq. vind. 2. t. 143.
4	guazumæfölia. DC. (pu.)	Guazuma-ld.	S. America.	1820.	6. 8.	S. h.	
5	verbascifölia. DC.	(re.) Verbascum-ld.	Brazil.	—	—	S. h.	Lodd. bot. cab. 504.
6	ferruginātā. DC.	(ye.) yellow-flow'r'd.	—	—	—	S. h.	
7	lanceolāta. DC.	(pu.) spear-leaved.	E. Indies.	1823.	S. h.	
8	spicāta. H. B.	(pu.) spike-flowered.	—	—	S. h.	

MY'R'DIA. DC. MY'R'DIA. Monadelphia Dodecandria. L.

	turbināta. DC.	(wh.) short-flowered.	W. Indies.	1793.	S. h.	Fl. mex. ic. ined.
	PLAGIA'NTHUS. DC.	PLAGIA'NTHUS.	Monadelphia Dodecandria.				
	divaricātus. DC.	... small-leaved.	New Zealand.	1821.	G. h.	Forst. gen. t. 43.
	ADANSON'IA. DC.	SOUR-GOURD.	Monadelphia Polyandria. L.				
	discolor. DC.	(wh.) palmate.	Senegal.	1724.	S. 5.	Bot. mag. 2791-2792.

BOMBACEÆ.

CAROLI'NEA. DC.	CAROLI'NEA.	Monadelphia Polyandria. L.				
1 princéps. DC.	(re.) digitated.	W Indies.	1787.	S. h.	Aub. gui. 2. t. 291-2.
Páchira aquática. A.						
2 insignis. DC.	(sc.) great-flowered.	—	1796.	S. h.	Lodd. bot. cab. 1004.
3 robústa. L.C.	(re.) robust.	S. America.	1825.	S. h.	Hern. mex. 68. ic.
4 fastuosa. DC.	(cr.) crimson.	New Spain.	—	S. h.	Bot. mag. 1412.
5 minor. DC.	(re.) lesser.	Guiana.	1798.	7. 8.	S. h.	Lod. bot. cab. 752.
6 álba. B.C.	(wh.) white-flowered.	Brazil.	—	S. h.	
BO'MBAX. DC.	SILK-COTTON-TREE.	Monadelphia Polyandria.				
1 Céiba. DC.	(ro.) five-leaved.	S. America.	1692.	S. h.	Jacq. am. t. 176. f. 1.
quinatum. Jacq.						
2 septenátum. DC. seven-leaved.	—	1699.	S. h.	
heptaphyllum. L. non Cav.						
3 malabáricum. DC. (sc.) Malabar.	E. Indies.	S. h.	Roxb. cor. 3. t. 247.	
heptaphyllum. Roxb. non L.						
4 globosum. DC.	(pa.) globe-fruited.	Guiana.	1824.	S. h.	Aubl. guian. 2. t. 281.
5 buonopozense. DC. (cr.) Africán.	Africa.	1822.	S. h.	Beauv. fl. ow. 2. t. 83.	
ERIODE'NDRON. DC.	ERIODE'NDRON.	Monadelphia Dodecandra.				
1 leianthérum. DC. (wh.) Brazilian.	Brazil.	1822.	S. h.	Cav. dis. 5. t. 152. f. 1.	
Bómbax eriánthos. Cav.						
2 anfractuósom. DC. (st.) twisted-anther'd	—	1739.	S. h.	Cav. dis. 5. t. 151.	
Bómbax pentándrum. L.						
α indicum. DC.	(ye.) East Indian.	E. Indies.	—	S. h.	Rheed. mal. 3. t. 49. 51.
β caribicum. DC.	(ro.) West Indian.	W. Indies.	—	S. h.	Jacq. am. t. 176. f. 70.
γ africanum. DC. (cr.) African.	Africa.	1823.	S. h.		
OCHR'OMA. DC.	OCHR'OMA.	Monadelphia Pentandria. L.				
1 Lagópus. DC.	(wh.) downy-leaved.	Jamaica.	1802.	S. h.	Sw. fl. 2 t. 23.
2 tomentosa. DC.	(wh.) woolly-leaved.	S. America.	1816.	S. h.	
CHEIROST'EMON. DC.	CHEIROST'EMON.	Monadelphia Decandra.				
platanoides. DC.	(wh.) plane-leaved.	New Spain.	1820.	11. 1.	S. h.	H. et B. pl. æq. 1. t. 24.
Chiranthodéndron. Larr. dis. ic.						

ORDO XXX.

BUTTNERIACEÆ. Kunth Synops. 3. p. 266.

Tribus I. STERCULIEÆ. DC. prodr. 1. p. 481.

STERC'ULIA. DC.	STERC'ULIA.	Monocia Monadelphia. L.				
1 Balánghas. DC.	(st.) many-seeded.	E. Indies.	1787.	6. 9.	S. h.	Rheed. mal. 1. t. 49.
2 nòbilis. DC.	(st.) few-seeded.	China.	—	—	S. h.	Bot. reg. 185.
Balánghas. B.R. Southuélia nòbilis. Par. Lond. 69.						
3 acuminata. DC.	(ye.) taper-pointed.	Africa.	1822.	S. h.	Beau. fl. ow. 1. t. 24.
4 grandiflora. DC.	(ye.) large-flowered.	E. Indies.	1820.	S. h.	
5 coccinea. DC.	(re.) scarlet-fruited.	—	—	—	S. h.	
6 angustifolia. DC.	(ye.) narrow-leaved.	Nepaul.	1823.	S. h.	
7 guttata. DC.	(fl.) spotted.	E. Indies.	1819.	S. h.	Rheed. mal. 4. t. 61.
8 heterophylla. DC.	(ye.) various-leaved.	Africa.	1824.	S. h.	Beauv. fl. ow. 1. t. 40.
9 Ivira. DC.	(fl.) hairy capsule.	W. Indies.	1723.	S. h.	Aub. gui. 2. t. 279.
crinata. Cav. dis. 5. t. 162.						
10 macrophylla. DC.	(ye.) large-leaved.	E. Indies.	1822.	S. h.	Cav. dis. 5. t. 145.
11 plataniifolia. DC.	(gr.) Plane-leaved.	China.	1757.	7.	G. h.	Roxb. cor. 1. t. 25.
12 colorata. DC.	(re.) scarlet.	E. Indies.	1818.	S. h.	— 1. t. 24.
13 urens. DC.	(gr.) stinging.	—	1793.	6. 9.	S. h.	
14 villosa. DC. vilous-leaved.	—	1818.	S. h.	
15 foetida. DC.	(fl.) fetid.	—	1690.	S. h.	Cav. dis. 5. t. 145.
HERITIE'RA. DC.	LOOKING-GLASS PLANT.	Dioecia Monadelphia.				
1 littoralis. DC.	(re.) large-leaved.	E. Indies.	1780.	S. h.	Rheed. mal. 6. t. 21.
2 minor. DC. small-leaved.	—	1816.	S. h.	Gärt. fr. 2. t. 98. f. 2.

Tribus II. BUTTNERIEÆ.

THEOBR'OMA. DC.	CHOCOLATE-NUT. Polyadelphia Decandria. L.					
1 Cacão. DC.	(ye.) smooth-leaved. S. America.	1739.	6. 9.	S. h.	Lodd. bot. cab. 545.	
2 guianénsis. DC.	(ye.) woolly-leaved. Guiana.	1803.	S. h.	Aub. gui. 2. t. 275.	
3 carib'æa.	(ye.) Caribbean. W. Indies.	1821.	S. h.		
4 bicolor. DC.	(re.) two-coloured. S. America.	1820.	S. h.	H. et B. pl. eq. 1. t. 30.	
ABR'OMA. DC.	ABR'OMA. Polyadelphia Dodecadria. L.					
1 augusta. DC.	(d.pu.) smooth-stalked. E. Indies.	1770.	6. 9.	S. h.	Bot. reg. 518.	
2 fastuosa. DC.	(d.pu.) rough-stalked. N. Holland.	1800.	—	S. h.	Sal. par. lond. 102.	
3 mollis. DC.	(d.pu.) soft. Moluccas.	—	—	S. h.		
GUAZ'UMA. DC.	BASTARD CEDAR. Polyadelphia Dodecadria. L.					
1 ulmifolia. DC.	(ye.) Elm-leaved. W. Indies.	1739.	8. 9.	S. h.	Trew ehret. t. 76.	
<i>Bubròma Guazuma.</i> w.						
2 tomentosa. DC.	(ye.) tomentose. S. America.	—	—	S. h.	Cav. ic. 3. t. 299.	
3 polybòtrys. DC.	(ye.) velvety. W. Indies.	—	—	S. h.		
COMMERS'ONIA. DC.	COMMERS'ONIA. Pentandria Pentagynia. L.					
1 echinata. DC.	(wh.) oval-leaved. Moluccas.	1806.	6. 7.	S. h.	Rumph. amb. 3. t. 119.	
2 platiphylla. DC.	(wh.) broad-leaved.	—	—	S. h.	And. bot. rep. 519.	
RUL'INGIA. Br.	RUL'INGIA. Pentandria Pentagynia. L.					
1 dasphylla.	(wh.) hairy-leaved. N. Holland.	1808.	4. 7.	G. h.	Bot. rep. 603.	
<i>Commersonia dasphylla.</i> A.R. <i>Byttnèria dasphylla.</i> DC.						
2 pannosa. B.M.	(wh.) cloth-leaved. N. Holland.	1819.	4. 7.	G. h.	Bot. mag. 2191.	
3 hermanniæfolia. DC.	(w.) Hermannia-ld.	—	—	G. h.		
BUTTN'ERIA. L.	BUTTN'ERIA. Pentandria Monogynia. L.					
1 herbacea. DC.	(wh.) herbaceous. E. Indies.	1820.	4. 7.	S. ȝ.	Roxb. cor. 1. t. 29.	
2 grandifolia. DC.	(wh.) large-leaved.	—	—	S. h.		
3 catalpæfolia. DC.	(wh.) Catalpa-leav'd. Caracas.	1823.	S. h.	Jacq. schoenb. 1. t. 46.	
4 cordata. DC.	(wh.) heart-leaved. Peru.	—	—	S. h.	Cav. dis. 5. t. 150.	
5 microphylla. DC.	(va.) small-leaved. S. America.	1816.	S. h.	— 5. t. 148. f. 2.	
6 scabra. DC.	(wh.) rough.	1793.	7. 8.	S. h.	— 5. t. 148. f. 1.	
AY'ENIA. DC.	AY'ENIA. Pentandria Monogynia.					
1 pusilla. DC.	(pu.) small. Jamaica.	1756.	7. 9.	S. ♂.	Mill. ic. t. 118.	
2 lævigata. DC.	(cr.) smooth.	—	—	S. h.		
KLEINH'OVIA. DC.	KLEINH'OVIA. Monadelphia Dodecadria.					
Hóspita. DC.	(pu.) purple-flower'd. Moluccas.	1800.	7. 9.	S. h.	Cav. dis. 5. t. 146.	

Tribus III. LASIOPETALEÆ. DC.

SERI'NGIA. DC.	SERI'NGIA. Pentandria Monogynia.					
platiphylla. DC.	(wh.) broad-leaved. N. Holland.	1802.	4. 7.	G. h.	Gay dis. p. 13. t. 1. 2.	
<i>Lasiopétalum arboréscens.</i> H.K.						
LASIOPE'TALUM. DC.	LASIOPE'TALUM. Pentandria Monogynia.					
1 ferrugineum. DC.	(wh.) rusty.	N. Holland.	1791.	4. 7.	G. h.	Bot. mag. 1766.
2 parviflòrum. DC.	(wh.) small-flowered.	—	1810.	—	G. h.	Linn. trans. 10. t. 19. f. 2.
THOM'ASIA. DC.	THOM'ASIA. Pentandria Monogynia.					
1 purpurea. DC.	(pu.) purple.	N. Holland.	1803.	4. 7.	G. h.	Bot. mag. 1755.
<i>Lasiopétalum purpureum.</i> H.K. <i>coccineum.</i> Donn. cant.						
2 foliosa. DC.	(wh.) leafy.	N. Holland.	1822.	—	G. h.	Gay dis. p. 24. t. 7.
3 solanacea. DC.	(wh.) Solanum-like.	—	1803.	—	G. h.	— p. 26. t. 6.
<i>Lasiopétalum solanaceum.</i> B.M. 1486.						
4 querifolia. DC.	(pu.) Oak-leaved.	—	—	—	G. h.	Bot. mag. 1485.

Tribus IV. HERMANNIEÆ. DC.

MEL'OCHIA. DC.	MEL'OCHIA. Monadelphia Pentandria. DC.					
1 pyramidata. DC.	(vi.) pyramidal.	Brazil.	1768.	7. 8.	S. h.	Jacq. vind. 1. t. 30.
2 tomentosa. DC.	(vi.) downy.	W. Indies.	—	5. 6.	S. h.	Cav. dis. 6. t. 172. f. 2
3 parvifolia. DC.	(wh.) small-leaved.	Caracas.	1823.	S. h.	

RIEDLE'IA. DC.	RIEDLE'IA. Monadelphia Pentandria. L.
1 velutina. DC.	soft-leaved. E. Indies. 1820. 6. 8. S. h.
2 depréssâ. DC. (st.) depressed.	Havannah. — — S. h.
3 supina. DC.	procumbent. E. Indies. 1823. — — S. g.
4 corchorifolia. DC. (bh.) Corchorus-ld.	— 1732. 7. 8. S. ⊕.
<i>Melochia corchorifolia. L.</i>	Pluk. alm. t. 232. f. 4.
5 nodiflora. DC. (wh.) knot flowered.	S. America. 1823. — — S. h.
<i>Mougeotia nodiflora. Kth. Melochia nodiflora. Sw.</i>	Sloan. his. t. 135. f. 2.
6 caracasâna. DC. (wh.) Caracas.	Caracas. 1823. — — S. h.
<i>Mougeotia caracasâna. Kth. Melochia caracasâna. Jacq.</i>	Jacq. ic. 3. t. 507.
7 serrata. DC. (pu.) saw-leaved.	St. Domingo. 1827. 6. 8. S. 4.
WALTH'ERIA. DC.	WALTH'ERIA. Monadelphia Pentandria. L.
1 americâna. DC. (ye.) American.	S. America. 1691. 5. 10. S. h.
2 glâbra. DC. (ye.) smooth.	W. Indies. 1823. — — S. 4.
<i>lævis. L. en.</i>	Schrank. h. mon. t. 55.
3 ïndica. DC. (ye.) Indian.	E. Indies. 1759. 6. 8. S. h.
4 elliptica. DC. (ye.) woolly-leaved.	— 1812. — — S. h.
5 microphylla. DC. (ye.) small-leaved.	— 1823. — — S. h.
HERMANNIA. DC.	HERMANNIA. Monadelphia Pentandria. L.
1 multifida. DC. (ye.) multifid-leaved.	C. B. S. 1825. 5. 8. G. h.
2 halicâcaba. DC. (ye.) bladder-calyx'd.	— 1820. — — G. h.
3 comosa. DC. (ye.) tufted.	— 1. 12. G. h.
4 althaeifolia. DC. (ye.) Althaea-leaved.	— 1728. 3. 7. G. h.
5 althæoides. L. en. (ye.) Althaea-like	— ... G. h.
6 plicâta. DC. (ye.) plaited-leaved.	— 1774. 11. 12. G. h.
7 candidans. DC. (ye.) white-leaved.	— 4. 6. G. h.
8 mollis. DC. (ye.) soft-leaved.	— 1818. — — G. h.
9 decumbens. DC. (ye.) decumbent.	— — — G. h.
10 hyssopifolia. DC. (ye.) Hyssop-leaved.	— 1725. — — G. h.
11 trifoliata. DC. (ye.) three-leaved.	— 1752. 5. 8. G. h.
12 triphylla. DC. (st.) three-parted.	— 1820. — — G. h.
13 glandulosa. DC. (ye.) glandular.	— — — G. h.
14 frâgrans. DC. (ye.) fragrant.	— — — G. h.
15 disermafolia. DC. (ye.) simple-flow'r'd.	— 1795. 3. 4. G. h.
16 disticha. DC. (ye.) round-leaved.	— 1789. 5. 8. G. h.
<i>rotundifolia. Jacq. schœn. 3. t. 121.</i>	Jacq. schœnb. 1. t. 121 S. et Wend. sert.h.t. 10..
17 melochioïdes. DC. (ye.) Melochia-like.	— 1818. — — G. h.
18 bryoniifolia. DC. (ye.) Bryony-leaved.	— — — G. h.
19 salvifolia. DC. (ye.) Sage-leaved.	— 1795. 4. 6. G. h.
20 micans. DC. (y.or.) glittering.	— 1790. 5. 8. G. h.
<i>latifolia. Jacq. schœn. 3. t. 119.</i>	Cav. dis. 6. t. 180. f. 2. S. et Wend. s. han. t. 5.
21 involucrata. DC. (st.) involucred.	— 1794. 5. 6. G. h.
22 scordifolia. DC. (or.y.) Germanander-ld.	— 4. 11. G. h.
23 denudata. DC. (ye.) smooth.	— 1774. 5. 7. G. h.
24 alniifolia. DC. (ye.) Alder-leaved.	— 1728. 2. 5. G. h.
25 cuneifolia. DC. (ye.) wedge-leaved.	— 1791. 8. 9. G. h.
26 holosericea. DC. (ye.) velvet-leaved.	— 1792. 5. 6. G. h.
27 hirsuta. DC. (ye.) hairy-branch'd.	— 1790. — — G. h.
28 scâbra. DC. (ye.) rough-leaved.	— 1789. 3. 4. G. h.
29 multiflora. DC. (ye.) many-flowered.	— 1791. 3. 5. G. h.
30 flâumea. DC. (re.) flame-flowered.	— 1794. 1. 12. G. h.
31 angulâris. DC. (or.) angular-calyx'd.	— 1791. 4. 5. G. h.
32 trifurcata. DC. (re.) three-forked.	— 1789. 4. 7. G. h.
33 odorata. DC. (re.) sweet-scented.	— 1780. 2. 10. G. h.
34 lavandulæfolia. DC. (y.) Lavender-ld.	— 1732. 5. 9. G. h.
35 velutina. DC. (ye.) mucronate.	— 1818. — — G. h.
36 filifolia. DC. (or.) thread-leaved.	— 1816. 5. 8. G. h.
37 procumbens. DC. (st.) procumbent.	— 1792. 5. 6. G. h.
38 tenuifolia. DC. (ye.) slender-leaved.	— ... 6. 7. G. h.
39 incisa. DC. (ye.) cut-leaved.	— 1806. — — G. h.
40 pulverâta. DC. (or.) powdered.	— 1800. 5. 8. G. h.
41 argentea. DC. (or.) silvery-leaved.	— 1818. — — G. h.
42 coronopifolia. DC. (ye.) Buckshorn-ld.	— ... 6. 8. G. h.
43 hispidula. s.s. (ye.) bristly.	— 1823. 3. 8. G. h.
MAHERNIA. DC.	MAHERNIA. Monadelphia Pentandria.
1 verticillatâ. DC. (ye.) whorl-leaved.	C. B. S. 1820. 6. 8. G. h.
2 resedæfolia. DC. (re.) Reseda-leaved.	— 1818. — — G. h.
3 bipinnatâ. DC. (re.) winged-leaved.	— 1752. — — G. h.
<i>pinnatâ. B.M.</i>	Cav. dis. 6. t. 176. f. 1. Bot. mag. 277.

4 incisa. DC.	(st.) cut-leaved.	C. B. S.	1792.	7. 9.	G. $\frac{h}{2}$.	Bot. mag. 353.
5 diffusa. DC.	(ye.) diffuse.	_____	1774.	6. 8.	G. $\frac{h}{2}$.	Lodd. bot. cab. 187.
6 heterophylla. DC.	(ye.) various-leaved.	_____	1731.	4. 5.	G. $\frac{h}{2}$.	Cav. dis. 6. t. 178. f. 1.
<i>Hermánnia grossulariaefolia. L.</i>						
7 seselifolia. DC.	(ye.) Seseli-leaved.	_____	1818.	6. 8.	G. $\frac{h}{2}$.	Cav. dis. 6. t. 177. f. 3.
8 pulchella. DC.	(re.) neat.	_____	1792.	7. 9.	G. $\frac{h}{2}$.	Jacq. schoenb. 1. t. 53.
9 glabrata. DC.	(ye.) sweet-scented.	_____	1789.	4. 5.	G. $\frac{h}{2}$.	
<i>odorata. A.R. 85.</i>						
10 vesicaria. DC.	(ye.) inflated.	_____	1818.	—	G. $\frac{h}{2}$.	Cav. dis. 6. t. 181. f. 2.
11 grandiflora.	(sc.) great-flowered.	_____	1791.	5. 8.	G. $\frac{h}{2}$.	
<i>Hermánnia grandiflora. H.K.</i>						
12 Burchellii.	(sc.) Burchell's.	_____	1818.	—	G. $\frac{h}{2}$.	Bot. reg. 224.
<i>grandiflora. B.R. non H.K.</i>						
13 biserrata. DC.	(ye.) double-tooth'd.	_____	1825.	—	G. $\frac{h}{2}$.	Cav. dis. 6. t. 200. f. 2.

Tribus V. DOMBEYACEÆ. DC.

RUI'ZIA. DC.	RUI'ZIA. Monadelphia Polyandria. L.					
1 lobata. DC.	(ye.) lobed-leaved.	Bourbon.	1820.	S. $\frac{h}{2}$.	Cav. dis. 3. t. 36. f. 1.
2 variabilis. DC.	(bh.) various-leaved.	_____	1792.	5.	S. $\frac{h}{2}$.	Jacq. schoen. 3. t. 295.
PENTAP'ETES. DC.	PENTAP'ETES. Monadelphia Dodecandra. L.					
1 phenicea. DC.	(cr.) scarlet-flower'd. India.	1690.	7. 8.	S. $\frac{f}{2}$.	Bot. reg. 575.	
2 ovata. DC.	(sn.) oval-leaved.	New Spain.	1805.	6. 9.	S. $\frac{f}{2}$.	Cav. ic. 5. t. 433.
<i>Broteria ovata. Cav.</i>						
ASS'ONIA. DC.	ASS'ONIA. Monadelphia Dodecandra.					
1 populnea. DC.	(wh.) Poplar-leaved.	Bourbon.	1822.	S. $\frac{h}{2}$.	Cav. dis. 3. t. 42. f. 3.
2 viburnoides. DC.	(wh.) Viburnum-l'd.	_____	—	S. $\frac{h}{2}$.	
DOMBEYA. DC.	DOMBEYA. Monadelphia Dodecandra.					
1 cordifolia. DC.	(wh.) heart-leaved.	E. Indies.	1820.	S. $\frac{h}{2}$.	
2 ferruginea. DC.	(wh.) rusty.	Mauritius.	1815.	S. $\frac{h}{2}$.	Cav. dis. 3. t. 42. f. 2.
3 ovata. DC.	(wh.) oval-leaved.	Bourbon.	1822.	S. $\frac{h}{2}$.	— 3. t. 41. f. 2.
MELH'ANIA. DC.	MELH'ANIA. Monadelphia Pentandra.					
1 Erythroxylon. DC.	(wh.) red-wood.	St. Helena.	1722.	5. 8.	S. $\frac{h}{2}$.	Bot. mag. 1000.
2 Melanoxylon. DC.	(wh.) black-wood.	_____	1800.	7. 8	S. $\frac{h}{2}$.	Pluk. mant. 6. t. 333. f. 5.
3 Burchellii. DC.	(wh.) Burchell's.	C. B. S.	1818.	—	S. $\frac{h}{2}$.	
PTEROSPE'RUM. DC.	PTEROSPE'RUM. Monadelphia Dodecandra.					
1 acerifolium. DC.	(wh.) Maple-leaved.	E. Indies.	1790.	7. 9.	S. $\frac{h}{2}$.	Bot. mag. 620.
2 suberifolium. DC.	(wh.) various-leaved.	_____	1783.	9. 10.	S. $\frac{h}{2}$.	— 1526.
3 lanceafolium. DC.	(w.) lance-leaved.	_____	1818.	—	S. $\frac{h}{2}$.	
4 canescens. H.B.	(wh.) canescent.	Ceylon.	—	—	S. $\frac{h}{2}$.	
5 semisagittatum. DC.	(w.) fringe-bracted.	E. Indies.	—	—	S. $\frac{h}{2}$.	
ASTRAP'EA. DC.	ASTRAP'EA. Monadelphia Dodecandra.					
1 Wallichii. DC.	(sc.) Wallich's.	Mauritius.	1820.	12. 1.	S. $\frac{h}{2}$.	Bot. reg. 691.
2 viscosa.	viscous.	Madagascar.	1823.	S. $\frac{h}{2}$.	
3 tiliæfolia.	(wh.) Lime-leaved.	Bourbon.	1824.	S. $\frac{h}{2}$.	
KY'DIA. DC.	KY'DIA. Monadelphia Dodecandra.					
1 calycina. DC.	(wh.) calycine.	E. Indies.	1818.	S. $\frac{h}{2}$.	Roxb. cor. 3. t. 215.
2 fraterna. DC.	(wh.) long-stamen'd.	_____	1823.	S. $\frac{h}{2}$.	— 3. t. 216.

Tribus VI. WALLICHIEÆ. DC.

ERIOL'ÆNA. DC.	ERIOL'ÆNA. Monadelphia Polyandria.					
Wallichii. DC.	Wallich's.	E. Indies.	1823.	S. $\frac{h}{2}$.	DC.m.mus.10. p.102.ic.

ORDO XXXI.

TILIACEÆ. DC. prodr. 1. p. 503.

ENT'ELEA. B.M.	CORK-TREE. Polyantria Monogynia.			
arboréscens. B.M. (<i>wh.</i>)	New Zealand. N. Zealand. 1821. 5. 6. G. h.			Bot. mag. 2480.
SPARRMA'NNIA. DC.	SPARRMA'NNIA. Polyantria Monogynia. L.			
1 africana. DC. (<i>wh.</i>) African.	C. B. S. 1790. 3. 7. G. h.			Bot. mag. 516.
2 rugosa. Bhi. (<i>wh.</i>) rugged-leaved.	——— 1825. 3. 6. G. h.			
HELIOLA'RPU. DC.	HELIOLA'RPU. Dodecandria Monogynia.			
americana. DC. (<i>wh.</i>) American.	Vera Cruz. 1733. S. h.			Trew ehret. t. 45.
CO'RCHORUS. DC.	CO'RCHORUS. Polyandria Monogynia. L.			
1 siliquosus. DC. (<i>ye.</i>) Germanander-ld.	S. America. 1732. 6. 8. S. 4.			Jacq. vind. 3. t. 59.
2 foliosus. s.s. (<i>ye.</i>) leafy.	Cuba. 1827. —— S. 4.			
3 hirtus. DC. (<i>ye.</i>) hairy.	S. America. 1820. —— S. 4.			Plum. ed. B. t. 103. f. 2.
4 pilolobus. DC. (<i>ye.</i>) hairy-podded.	——— —— S. 4.			
5 americanus. Sck. (<i>ye.</i>) American.	S. America. 1825. —— S. h.			
6 trilocularis. DC. (<i>ye.</i>) three-celled.	Arabia. 1790. 7. 8. S. ♂.			Jacq. vind. 2. t. 173.
7 olitorius. DC. (<i>ye.</i>) bristly-leaved.	India. 1640. 6. 8. S. ♂.			Bot. mag. 2810.
8 aestuans. DC. (<i>ye.</i>) Hornbeam-ld.	S. America. 1731. 6. 7. S. ♂.			Jacq. vind. 1. t. 85.
9 acutangulus. DC. (<i>ye.</i>) acute-angled.	India. 1816. —— S. h.			Pluk. phyt. t. 44. f. 1.
10 capsularis. DC. (<i>ye.</i>) heart-leaved.	E. Indies. 1725. —— S. ♂.			Jacq. f. eclog. 2. t. 120.
11 hirsutus. DC. (<i>ye.</i>) woolly-capsul'd.	S. America. 1752. —— S. h.			Jacq. vind. 3. t. 59.
TRIUMFE'TTA. DC.	TRIUMFE'TTA. Dodecandria Monogynia. L.			
1 Láppula. DC. (<i>ye.</i>) prickly-seeded.	W. Indies. 1739. 7. 8. S. h.			Plum. ic. t. 255.
2 heterophylla. DC. (<i>ye.</i>) various-leaved.	——— 1823. —— S. h.			Pluk. t. 425. f. 3.
3 rubricaulis. DC. (<i>ye.</i>) red-stem'd.	Caracas. 1826. —— S. h.			
4 suborbiculata. DC. (<i>ye.</i>) subrotund.	E. Indies. —— S. h.			
5 glandulosa. DC. (<i>ye.</i>) gland-toothed.	Mauritius. —— S. h.			Lam. ill. t. 400. f. 1.
6 tricholàda. DC. (<i>ye.</i>) oval-leaved.	Nepaul. 1823. —— S. ♂.			
7 oblongata. DC. (<i>ye.</i>) oblong-leaved.	——— —— S. ♂.			
8 annua. DC. (<i>ye.</i>) annual.	Java. 1760. 8. 9. S. ♂.			Bot. mag. 2296.
9 angulata. DC. (<i>ye.</i>) angular-leaved.	E. Indies. 1739. 6. 7. S. h.			Lam. ill. t. 400. f. 2.
10 semitriloba. DC. (<i>ye.</i>) Mallow-leaved.	W. Indies. 1773. —— S. h.			Jacq. vind. 3. t. 76.
11 havanensis. DC. (<i>ye.</i>) Havannah.	Cuba. 1825. —— S. h.			
12 micropétala. B.R. (<i>ye.</i>) small-petal'd.	E. Indies. —— 4. 8. S. h.			Bot. reg. 1058.
13 petiolaris. B.R. (<i>ye.</i>) petiolate.	Peru. 1819. 5. 6. S. h.			Lindl. collect. t. 20.
rhomboidea. L. col. non Jacq.				
14 rhomboidea. DC. (<i>ye.</i>) rhomboid.	S. America. 1819. 5. 6. S. h.			Jacq. amer. t. 90.
15 althaeoides. DC. (<i>ye.</i>) Althaea-leaved.	Cayenne. —— —— S. h.			
macrophylla. Vahl. ecl. 2. p. 34.				
16 grandiflora. DC. (<i>ye.</i>) large-flowered.	W. Indies. 1810. —— S. h.			
17 trilocularis. DC. (<i>ye.</i>) three-celled.	E. Indies. 1817. 10. 4. S. h.			
GRE'WIA. DC.	GRE'WIA. Polyantria Monogynia. L.			
1 tomentosa. DC. tomentose.	Java. 1818. S. h.			Juss. an. 2. t. 49. f. 1.
2 hirsuta. DC. hairy.	——— 1816. 7. 9. S. h.			
3 bracteata. DC. (<i>gr.</i>) bracted.	E. Indies. 1818. —— S. h.			
4 Mallocócca. DC. rough-fruited.	Tongatabu. 1792. 8. 9. S. h.			Forst. gen. t. 39.
Mallocócca crenata. Forst.				
5 oppositifolia. DC. sharp-leaved.	Nepaul. 1818. S. h.			
6 umbellata. DC. umbel-flowered.	E. Indies. —— S. h.			
7 bicolor. DC. two-coloured.	Senegal. 1822. S. h.			Juss. ann. t. 50. f. 2.
8 Róthii. DC. Roth's.	E. Indies. 1819. S. h.			
9 salvifolia. DC. (<i>ye.</i>) Sage-leaved.	——— S. h.			
10 flava. DC. (<i>ye.</i>) yellow.	C. B. S. 1818. —— G. h.			
11 ovalifolia. DC. oval-leaved.	E. Indies. —— S. h.			
12 Micrócos. DC. (<i>st.</i>) panicled.	——— 1779. —— S. h.			Rheed. mal. 1. t. 56.
<i>Micrócos paniculata. L.</i>				
13 paniculata. DC. villous-leaved.	——— 1820. S. h.			
14 orientalis. DC. (<i>wh.</i>) oriental.	——— 1767. 7. 8. S. h.			Rheed. mal. 5. t. 46.
15 pilosa. DC. (<i>wh.</i>) pilose.	——— 1804. —— S. h.			
16 glandulosa. DC. glandular.	Mauritius. 1827. S. h.			Juss. an. 2. t. 48. f. 1.
17 serrulata. DC. serrulate.	E. Indies. 1818. S. h.			
18 occidentalis. DC. (<i>pu.</i>) Elm-leaved.	C. B. S. 1690. 7. 9. G. h.			Bot. mag. 422.
19 asiatica. (<i>li.</i>) Asiatic.	E. Indies. 1792. 7. 8. S. h.			Sonn. voy. 2. t. 138.

20	<i>subin'aqualis</i> . DC.	oblique-leaved.	E. Indies.	1816.	S. h.	
21	<i>villifera</i>	wool-bearing.	1826.	S. h.	
	<i>villosa</i> . Roth. <i>nec aliorum</i> .						
22	<i>áspéra</i> . DC.	rough-leaved.	1818.	S. h.	
23	<i>tiliaefolia</i> . DC.	Lime-leaved.	1812.	S. h.	
24	<i>villosa</i> . DC.	villous.	1816.	S. h.	
25	<i>therabinthinacea</i> . DC.	strong-scented.	S. h.	
26	<i>obtusifolia</i> . DC.	blunt-leaved.	C. B. S.	1818.	G. h.	
27	<i>ulmifolia</i> . H.B.	Chinese.	China.	G. h.	
28	<i>carpinifolia</i> . DC. (<i>pu.</i>)	Hornbeam-ld.	Africa.	1822.	S. h.	
29	<i>megalocarpa</i> . B.O.(<i>re</i>)	black-fruited.	S. h.	
							Beauv. fl. ow. 2. t.
							2. t. 102.
BROWNLOWIA. R.		BROWNLOWIA.	Polyandria Monogynia.				
	<i>elata</i> . R.	(ye.) tall.	E. Indies.	1812.	S. h.	Roxb. cor. 3. t. 265.
TILIA. DC.		LIME-TREE.	Polyandria Monogynia. L.				
1	<i>microphylla</i> . DC. (<i>wh.</i>)	small-leaved.	Britain.	8. 9.	H. 5.	Eng. bot. 1705.
	<i>parvifolia</i> . E.B.						
2	<i>intermédia</i> . DC. (<i>wh.</i>)	intermediate.	—	H. 5.	Hayn. et Sv. bot. t. 40.
	<i>europea</i> . Eng. bot. 610.						
	β <i>laciniatu</i> .	(<i>wh.</i>) jagged-leaved.	6. 8.	H. 5.	
3	<i>corallina</i> . E.F.	(<i>wh.</i>) coral-twigg'd.	Britain.	7.	H. 5.	
4	<i>rúbra</i> . DC.	(<i>wh.</i>) red-twigg'd.	Europe.	6. 8.	H. 5.	
5	<i>platyphylla</i> . DC. (<i>wh.</i>)	broad-leav'd.	Britain.	—	H. 5.	Vent. dis. p. 6. t. 1. f. 2.
	<i>grandifolia</i> . E.F.						
6	<i>glabra</i> . DC.	(<i>wh.</i>) broad-leaved.	N. America.	1752.	6. 7.	H. 5.	Vent. dis. p. 9. t. 2.
	<i>americana</i> . L.						
7	<i>laxiflora</i> . DC.	(<i>wh.</i>) loose-flowered.	1820.	—	H. 5.	
8	<i>pubescens</i> . DC.	(<i>wh.</i>) pubescent.	1726.	7. 8.	H. 5.	Vent. dis. p. 10. t. 3.
	β <i>leptophylla</i> . DC. (<i>w.</i>)	narrow-leaved.	—	H. 5.	
9	<i>heterophylla</i> . DC. (<i>wh.</i>)	white-leaved.	6. 8.	H. 5.	Vent. dis. p. 16. t. 5.
	<i>álba</i> . Mich. arb. 3. t. 2.						
10	<i>argéntea</i> . DC.	(<i>wh.</i>) silver-leaved.	Hungary.	1767.	—	H. 5.	Vent. dis. p. 13. t. 4.
	<i>álba</i> . W. et K. hung. 1. t. 3. <i>rotundifolia</i> . Vent.						
11	<i>petiolaris</i> . DC.	(<i>wh.</i>) long-petioled.	1824.	H. 5.	
MUNTINGIA. DC.		MUNTINGIA.	Polyandria Monogynia. L.				
	<i>Calabura</i> . DC.	(<i>wh.</i>) Jamaica.	Jamaica.	1690.	6. 7.	S. h.	Jacq. amer. t. 107.
APEIBA. DC.		APEIBA.	Polyandria Monogynia. L.				
1	<i>Tibórbou</i> . DC.	(ye.) hairy.	S. America.	1756.	S. h.	Aub. gui. 1. t. 213.
	<i>Aubletia Tibórbou</i> . w.						
2	<i>Petóumo</i> . DC.	(ye.) hoary.	1817.	S. h.	Aub. gui. 1. t. 215.
3	<i>áspera</i> . DC.	(ye.) prickly-capsul'd.	Cayenne.	1792.	S. h.	— 1. t. 216.
	<i>Aubletia áspora</i> . w.						
4	<i>glabra</i> . DC.	(gr.) smooth.	1817.	S. h.	Aub. gui. 1. t. 214.
	<i>Aubletia l'avis</i> . w.						
SL'ONEA. DC.		SL'ONEA.	Polyandria Monogynia. L.				
1	<i>dentata</i> . DC.	(ye.) Chestnut-leav'd.	S. America.	1752.	S. h.	Plum. ic. 244.
2	<i>sinemariensis</i> . DC.	(ye.) round-leaved.	W. Indies.	1820.	S. h.	Aub. gui. 1. t. 212.
GYROSTEMON. DC.		GYROSTEMON.	Dioecia Polyandria.				
	<i>ramulosum</i> . DC.	branching.	N. Holland.	1820.	G. h.	Desf. m. mus. 6. t. 6.
BE'RRYA. R.C.		BE'RRYA.	Polyandria Monogynia.				
	<i>ammonilla</i> . R.C. (<i>st.</i>)	Ceylon.	Ceylon.	1818.	S. h.	Roxb. corom. 3. t. 264.

ORDO XXXII.

ELÆOCARPEÆ. DC. *prodr.* 1. p. 519.

ELÆOCARPUS. DC. ELÆOCARPUS. Dodecandria Monogynia.

1	<i>serrátu</i> . DC.	(<i>pu.</i>) saw-leaved.	E. Indies.	1774.	S. h.	Burm. zeyl. t. 40.
2	<i>cyáneus</i> . DC.	(<i>wh.</i>) blue-fruited.	N. Holland.	1803.	6. 8.	G. h.	Bot. mag. 1737.
	<i>reticulatus</i> . B.R.	657.					

ACER'ATIUM. DC. ACER'ATIUM. Dodecandria Monogynia.

oppositifolium. DC.	opposite-leaved.	Amboyna.	1818.	S. h.	
---------------------	------------------	----------	-------	------	-------	--

DICERA. DC.	DICERA. Dodecandra Monogynia.
dentata. DC. (wh.) tooth-leaved.	N. Zealand. 1820. 6. 8. G. h. Forst. gen. t. 40.
FRIE'SIA. DC.	FRIE'SIA. Dodecandra Monogynia.
peduncularis. DC. (wh.) peduncled.	Van Diem. 1818. G. h. Lab. n. holl. 2. t. 155. <i>Elæocarpus peduncularis.</i> Lab.

ORDO XXXIII.

CHLENACEÆ. DC. prodr. 1. p. 521.

+ Genus *Chlenaceis* affine.

HUG'ONIA. DC.	HUG'ONIA. Monadelphia Decandria.
1 My'stax. DC. (ye.) entire-leaved.	E. Indies. 1818. 6. 9. S. h. Rheed. mal. 2. t. 29.
2 serrata. DC. (ye.) saw-leaved.	Mauritius. 1820. — S. h. Cav. dis. 3. t. 73. f. 1.

ORDO XXXIV.

TERNSTRŒMIACEÆ. DC. prodr. 1. p. 523.

Tribus I. TERNSTRŒMIEÆ. DC.

TERNSTRŒMIA. DC.	TERNSTRŒMIA. Polyandria Monogynia.
1 peduncularis. DC. (w.) long-peduncled.	W. Indies. 1818. 6. 8. S. h.
2 punctata. DC. (wh.) spotted-leaved.	Guiana. 1824. S. h.
3 dentata. DC. (ye.) toothed.	——— 1826. S. h.
4 venosa. DC. (wh.) veined-leaved.	Brazil. 1820. 6. 8. S. h.
5 serrata. H. B. (wh.) saw-leaved.	Silhet. — S. h.

Tribus II. FREZIEREÆ. DC.

CL'EYERA. DC.	CL'EYERA. Polyandria Monogynia.
japonica. DC. (wh.) Japan.	Japan. 1822. S. h. Kæmp. am. 5.p.774. ic.
FREZIE'R. DC.	FREZIE'R. Polyandria Monogynia.
thaœoides. DC. (wh.) Tea-like.	Jamaica. 1818. S. h.
EURYA. DC.	E'URYA. Polygamia Monœcia.
1 chinensis. DC. (wh.) China.	China. 1818. 1. 3. G. h. Lodd. bot. cab. 1213.
2 multiflora. DC. (wh.) many-flowered.	Nepaul. 1823. S. h.
LETTS'OMIA. DC.	LETTS'OMIA. Polyandria Monogynia.
tomentosa. DC. tomentose.	Peru. 1823. S. h. Ruiz et P. prod. t. 14.

Tribus III. SAURAUJEÆ DC.

SAURA'UJA. DC.	SAURA'UJA. Polyandria Pentagynia.
1 excelsa. DC. (wh.) tall.	Caracas. 1824. S. h. Willd. n. act. ber. 3.t.4.
2 napaulensis. DC. (wh.) Nepaul.	Nepaul. — S. h.

Tribus IV. LAPLACEÆ. DC.

WITTELSBA'CHIA. M.N. WITTELSBA'CHIA. Polyandria Monogynia.

- 1 *Gossy'pium*. M.N. (*ye.*) large-flowered. E. Indies. 1822. S. h. Sonn. voy. 2. t. 133.
Cochlospermum Gossy'pium. DC. *Bombax Gossy'pium*. Cav. dis. 5. t. 157.
- 2 *vitifolia*. M.N. (*ye.*) Vine-leaved. S. America. 1820. S. h. Fl. mex. ic. ined.
- 3 *Cochlospermum serratifolium*. DC. *hibiscoides*. K.S. *Bombax vitifolium*. W.en.
- 3 *insignis*. M.N. (*ye.*) spot-flowered. Brazil. 1821. S. h. Mart. nov. gen. 1. t. 55.

Tribus V? GORDONIEÆ. DC.

MALACHODE'NDRON. DC. MALACHODE'NDRON. Monadelphia Polyandria. L.

- ovatum*. DC. (*wh.*) oval-leaved. N. America. 1785. 8. 9. H. h. Bot. reg. 1104.
- Stuártia pentagynia*. Ex. bot. t. 101.

STUARTRIA. W. STUARTRIA. Monadelphia Polyandria. L.

- virginica*. DC. (*wh.*) Virginian. N. America. 1742. 5. 8. H. h. Cav. dis. 5. t. 159. f. 2.
- Malachodéndron*. L. *marylandicum*. A.R. t. 73.

GORD'ONIA. DC. GORD'ONIA. Monadelphia Polyandria. L.

- 1 *Lasiánthus*. DC. (*wh.*) smooth. N. America. 1739. 8. 11. F. h. Bot. mag. 668.
- 2 *haematóxylon*. DC. (*w.*) red-wooded. Jamaica. 1820. S. h.
- 3 *pubescens*. DC. (*wh.*) pubescent. N. America. 1774. 8. 11. F. h. Vent. malm. t. 1.
- 4 *Franklini*. L'Her. (*w.*) Franklin's. F. h.

POLYSP'ORA. N.L.F. POLYSP'ORA. Monadelphia Polyandria.

- axillaris*. N.L.F. (*wh.*) axillary-flow'r'd E. Indies. 1816. 11. 3. S. h. Bot. reg. 349.
- Camélia axillaris*. B.R. et B.M. 2047.

ORDO XXXV.

CAMELLIEÆ. DC. prodr. 1. p. 529.

CAMELLIA. Ker. CAMELLIA. Monadelphia Polyandria. L.

- | | | | | | |
|---|---------------------------------------|---------|---------------|-------|------------------------|
| 1 <i>Thea</i> . L. en. | (<i>wh.</i>) Bohea Tea. | China. | 1768. 11. 2. | G. h. | Lob. bot. cab. 226. |
| 2 <i>viridis</i> . L. en. | (<i>wh.</i>) green Tea. | _____ | _____ 10. 12. | G. h. | _____ 227. |
| <i>Thea chinensis</i> . B.M. 998. | | | | | |
| 3 <i>euryoides</i> . B.R. | (<i>wh.</i>) small-flower'd. | _____ | 1824. 1. 5. | G. h. | Bot. reg. 983. |
| 4 <i>oleifera</i> . B.R. | (<i>wh.</i>) oleiferous. | _____ | 1819. 1. 3. | G. h. | _____ 942. |
| 5 <i>Sesánqua</i> . DC. | (<i>wh.</i>) Lady Banks's. | _____ | 1811. 11. 1. | G. h. | Thunb. fl. jap. t. 30. |
| β <i>semiduplex</i> . | | | | | |
| γ <i>flôre-plêno</i> . (<i>wh.</i>) double-flowered. | | | | | |
| 6 <i>maliflora</i> . B.R. | (<i>ro.</i>) pink-flowered. | _____ | 1823. _____ | G. h. | Bot. reg. 12. 1091. |
| <i>Sesanqua</i> β <i>rosea</i> . Bot. reg. 547. | | | | | |
| 7 <i>Kissi</i> . DC. | (<i>wh.</i>) Nepaul. | Nepaul. | 1823. | G. h. | Bot. reg. 983. |
| 8 <i>reticulata</i> . B.R. | (<i>sc.</i>) Captain Rawes'. China. | _____ | 1822. 2. 4. | G. h. | Bot. reg. 1078. |
| 9 <i>japonica</i> . DC. | (<i>va.</i>) common. | _____ | 1739. 12. 5. | G. h. | Cav. dis. 6. t. 160. |
| 1 <i>râbra</i> . | (<i>sc.</i>) single-red. | _____ | _____ | G. h. | Bot. mag. 42. |
| 2 <i>álba</i> . | (<i>wh.</i>) single-white. | _____ | | G. h. | Bot. reg. 353. |
| 3 <i>semiduplex</i> . | (<i>re.</i>) semidouble red. | _____ | | G. h. | Andr. rep. 559. |
| 4 <i>rûbro-plêna</i> . | (<i>re.</i>) double red. | _____ | | G. h. | _____ 199. |
| 5 <i>cárnea</i> . | (<i>pi.</i>) Middlemist's. | _____ | | G. h. | Lob. bot. cab. 455. |
| 6 <i>myrtifolia</i> . | (<i>pi.</i>) Myrtle-leaved. | _____ | | G. h. | Bot. mag. 1670. |
| involuta. Bot. reg. 633. | | | | | |
| 7 <i>hexangularis</i> . | (<i>re.</i>) six sided. | _____ | | G. h. | Bot. mag. 1670. |
| 8 <i>atrorubens</i> . | (<i>sc.</i>) Loddiges' red. | _____ | | G. h. | Lob. bot. cab. 170. |
| 9 <i>anemoniflora</i> . | (<i>cr.</i>) red Waratah. | _____ | | G. h. | Bot. mag. 1654. |
| 10 <i>dianthiflora</i> . | (<i>v.</i>) Carnation War. Knights. | _____ | | G. h. | _____ 2577. |
| 11 <i>blanda</i> . | (<i>bh.</i>) blush Waratah. China. | _____ | | G. h. | |
| 12 <i>pompónia</i> . | (<i>bh.</i>) Kew-blush. | _____ | | G. h. | Bot. reg. 22. |

CAMELLIÆ.

13 <i>paeoniiflora.</i>	(pi.) <i>Paeony-flowered.</i> China.	12. 5. G. h.	Lod. bot. cab. 238. 1198.
14 <i>Welbánkii.</i>	(wh.) <i>Welbank's.</i>	G. h.	Bot. reg. 708.
15 <i>lúteo-álba.</i>	(st.) <i>pale yellow.</i>	G. h.	— 112.
16 <i>flavescens.</i>	(st.) <i>buff.</i>	G. h.	Andr. rep. 25.
17 <i>álbo-plena.</i>	(wh.) <i>double white.</i>	G. h.	
18 <i>flimbriáta.</i>	(wh.) <i>fringed white.</i>	1. 4. G. h.	Andr. rep. 91.
19 <i>imbricáta.</i>	(re.) <i>imbricate-petal'd.</i>	G. h.	
20 <i>variegátâ.</i>	(v.) <i>double striped.</i>	G. h.	
21 <i>crassínervis.</i>	(re.) <i>thick nerved.</i>	G. h.	
22 <i>expânsa.</i>	(re.) <i>expanded.</i>	G. h.	
23 <i>conchiflóra.</i>	(re.) <i>shell-flowered.</i>	G. h.	
24 <i>rubricáulis.</i>	(sc.) <i>Lady Ad'l. Campbell's</i>	G. h.	
25 <i>longifòlia.</i>	(re.) <i>long-leaved.</i> Malcolm's.	G. h.	
26 <i>Chandlèri.</i>	(v.) <i>striped Waratah.</i> Chandler's.	1822.	12. 5. G. h.	Chandl. camel. t. 1. 2.
	versicolor. Bot. reg. 887.			
27 <i>Aitonî.</i>	(sc.) <i>large single-red.</i>	—	G. h.	Chandl. camel. t. 3.
28 <i>altheæiflora.</i>	(re.) <i>Hollyhock-fl'd.</i>	—	G. h.	— t. 4.
29 <i>corallina.</i>	(re.) <i>coral-flowered.</i>	—	G. h.	— t. 5.
30 <i>insignis.</i>	(re.) <i>splendid.</i>	—	G. h.	— t. 6.
31 <i>flórida.</i>	(re.) <i>cluster-flowered.</i>	—	G. h.	— t. 7.
32 <i>anemonæflora. alba.</i> (w.) <i>white Anemone-fl'd.</i>	—	—	G. h.	— t. 8.
33 <i>Róssii.</i>	(re.) <i>Ross's.</i>	Ross's.	—	
34 <i>Wiltónia.</i>	(va.) <i>Mrs. Wilton's.</i>	Knight's.	—	
35 <i>Elphinstoniâna.</i>	(re.) <i>Miss Elphinstone's.</i>	—	G. h.	
36 <i>rosæflóra.</i>	(re.) <i>Rose-flowered.</i>	—	G. h.	
37 <i>Cliveâna.</i>	(re.) <i>Lady H. Clive's.</i>	—	G. h.	
38 <i>heterophyl'la.</i>	(re.) <i>various-leaved.</i>	—	G. h.	
39 <i>Egertóniæ.</i>	(re.) <i>Mrs. Egerton's.</i>	—	G. h.	
40 <i>Goussóniæ.</i>	(ro.) <i>Mrs. Goussoné's.</i>	—	G. h.	
41 <i>Barnabíæ.</i>	(pi.) <i>Mrs. Barnaby's.</i>	—	G. h.	
42 <i>radiâta.</i>	(re.) <i>rayed single.</i>	—	G. h.	
43 <i>supína.</i>	(re.) <i>supine.</i>	—	G. h.	
44 <i>Hibbertiâ.</i>	(re.) <i>Mr. Hibbert's.</i>	—	G. h.	
45 <i>Perc'ya.</i>	(re.) <i>D's. of Northumberland's</i>	—	G. h.	
46 <i>Woodsii.</i>	(ro.) <i>Mr. Woods's.</i>	Chandler's.	—	
47 <i>argéntea.</i>	(re.w.) <i>silvery-cup'd.</i>	—	G. h.	
48 <i>basílica.</i>	(re.) <i>princely.</i>	—	G. h.	
49 <i>princéps.</i>	(sc.) <i>double-carmine.</i>	—	G. h.	
50 <i>sanguínea.</i>	(cr.) <i>single-crimson.</i>	—	G. h.	
51 <i>serícea.</i>	(ro.) <i>double-silky.</i>	—	G. h.	
52 <i>Byrónii.</i>	(ro.) <i>Lord Byron's.</i>	—	G. h.	
53 <i>carnescens.</i>	(fl.) <i>single pale-red.</i>	—	G. h.	
54 <i>exímia.</i>	(re.) <i>fine double-red.</i>	—	G. h.	
55 <i>bicolor.</i>	(ro.w.) <i>single rosy & white.</i>	—	G. h.	
56 <i>purpurâscens.</i>	(re.) <i>single dark-red.</i>	—	G. h.	
57 <i>gigântea.</i>	(ro.) <i>large double.</i>	—	G. h.	
58 <i>lúcida.</i>	(re.) <i>double bright-red.</i>	—	G. h.	
59 <i>ornâta.</i>	(pi.) <i>pink Waratah.</i>	—	G. h.	
60 <i>papaverâcea.</i>	(re.) <i>Poppy-flower'd.</i> Loddiges.	—	G. h.	
61 <i>rotundifòlia.</i>	(re.) <i>round-leaved.</i>	—	G. h.	
62 <i>Spofforthiâna.</i>	(wh.) <i>red-lined white.</i> Spofforth.	—	G. h.	
63 <i>Herbérти.</i>	(ro.) <i>Mr. Herbert's.</i>	—	G. h.	
	Spofforthiâna. rosea. <i>Herbert MSS.</i>	—	G. h.	
64 <i>Haylöckii.</i>	(wh.) <i>Haylock's.</i>	—	G. h.	
65 <i>reflexa.</i>	(cr.) <i>single reflex-petaled.</i>	—	G. h.	

ORDO XXXVI.

OLACINEÆ. DC. prodr. 1. p. 531.

OLA'X. DC.

1 <i>scándens.</i> DC.	(wh.) <i>climbing.</i>	E. Indies.	1820.	1. 12. S. h. Roxb. cor. 2. t. 102.
2 <i>imbricáta.</i> DC.	(wh.) <i>imbricate.</i>	—	—	S. h. Roxb. cor. 2. t. 102.

SPERMA'XYRUM. DC.	SPERMA'XYRUM.	Triandria Monogynia.				
stríctum. DC.	(wh.) upright.	N. S. W.	1820.	G. h.	
XIM'ENIA. DC.	XIM'ENIA.	Octandria Monogynia. L.				
1 americàna. DC.	(wh.) spring.	W. Indies.	1759.	6. 9.	S. h.	Jacq. am. pict. t. 107.
2 inérmis. DC.	(wh.) unarmed.		1820.	S. h.	

ORDO XXXVII.

AURANTIACEÆ. DC. prodr. 1. p. 535.

ATALA'NTIA. DC.	ATALA'NTIA.	Monadelphia Octandria.				
monophylla. DC.	(wh.) simple-leaved.	E. Indies.	1777.	6. 8.	S. h.	Roxb. cor. 2. t. 83.
Limònìa monophylla. R.						
TRIPH'ASIA. DC.	TRIPH'ASIA.	Hexandria Monogynia.				
trifoliata. DC.	(wh.) three-leaved.	China.	1787.	7. 8.	S. h.	Jacq. ic. rar. t. 463.
Limònìa trifoliata. And. rep. 143.						
LIM'ONIA. DC.	LIM'ONIA.	Decandria Monogynia. L.				
1 acidissima. DC.	(wh.) sour-fruited.	E. Indies.	1824.	4. 8.	S. h.	Rumph. amb. 2. t. 43.
2 crenulata. DC.	(wh.) crenulate.		1808.	5. 7.	S. h.	Roxb. cor. 1. t. 86.
3 ambigùa. DC.	(wh.) ambiguous.			S. h.	
4 scàndens. H.B.	(wh.) climbing.	E. Indies.	1820.	S. h.	
CO'OKIA. DC.	CO'OKIA.	Decandria Monogynia. L.				
pùntatæ. DC.	(wh.) Wampee tree.	China.	1795.	5. 7.	S. h.	Jacq. schoenb. 1. t. 101.
MURRA'YA. DC.	MURRA'YA.	Decandria Monogynia. L.				
1 exòtica. DC.	(wh.) Ash-leaved.	E. Indies.	1771.	8. 9.	S. h.	Bot. reg. 434.
2 paniculata. DC.	(wh.) paniced.		1821.	4. 5.	S. h.	Hook. ex. flor. 134.
AGLA'IA. DC.	AGLA'IA.	Pentandria Monogynia.				
odoratæ. DC.	(st.) sweet-scented.	China.	1810.	2. 5.	S. h.	Rumph. amb. 5. t. 18.
BE'RGERA. DC.	BE'RGERA.	Decandria Monogynia. L.				
1 K'œnigii. DC.	(gr.) Kœnig's.	E. Indies.	1818.	3. 6.	S. h.	Lodd. bot. cab. 1019.
2 integrérima. DC.	(gr.) entire-leaved.		1820.	S. h.	
CLAUS'ENIA. DC.	CLAUS'ENIA.	Octandria Monogynia.				
pentaphylla. DC.	(wh.) five-leaved.	E. Indies.	1823.	S. h.	
GLYCO'SMIS. DC.	GLYCO'SMIS.	Decandria Monogynia.				
1 arbòrea. DC.	(wh.) tree.	E. Indies.	1796.	5. 8.	S. h.	Roxb. cor. 1. t. 85.
Limònìa arbòrea. R.C.						
2 citrifolia. H.T.	(wh.) Citron-leaved.	China.	1820.	2. 6.	G. h.	Bot. mag. 2416.
Limònìa citrifolia. DC.	parviflora. B.M.					
3 pentaphylla. DC.	(wh.) five-leaved.	E. Indies.	1790.	6. 7.	S. h.	Roxb. cor. 1. t. 84.
Limònìa pentaphylla. R.C.						
FER'ONIA. DC.	ELEPHANT APPLE.	Decandria Monogynia.				
elephántum. DC.	(bh.) Indian.	E. Indies.	1804.	S. h.	Roxb. cor. 2. t. 141.
'ÆGLE. DC.	BENGAL QUINCE.	Polyandria Monogynia.				
1 Mármelos. DC.	(wh.) thorny.	E. Indies.	1759.	S. h.	Roxb. cor. 2. t. 143.
CITRUS. DC.	ORANGE-TREE.	Polyadelphia Polyandria. L.				
1 Médicá. DC.	(wh.) Lemon-tree.	Asia.	1648.	5. 7.	G. h.	Ris. an. mus. 20. t. 2. f. 2.
2 Liméttæ. DC.	Bergamotte.		G. h.	20. t. 2. f. 1.
3 Limónum. DC.	(wh.) Citron.		G. h.	Ferr. hesp. t. 247. &c.
4 ácida. H.B.	(wh.) Lime.		G. h.	
5 Aurántium. DC.	(wh.) common.		1595.	G. h.	Ris. an. mus. 20. t. 1. f. 1. 2.
6 vulgàris. DC.	(wh.) Myrtle-leaved.		G. h.	Bot. reg. t. 346.
7 decumána. DC.	Shaddock.	India.	1724.	G. h.	Rumph. am. 2. t. 24. f. 2.
8 hy'strix. DC.	(wh.) very spiny.		G. h.	2. t. 28?
9 japónica. DC.	(wh.) small-fruited.	Japan.	1815.	G. h.	
10 nóbilis. DC.	(wh.) Mandarin.	China.	1805.	G. h.	Th. ic. jap. t. 15.
β minor.	(wh.) Tangerine.		G. h.	Andr. rep. 608.
11 Margarita. DC.	(wh.) sweet Lemon.		1820.	G. h.	Bot. reg. 211.
12 madurén sis. DC.	(wh.) spineless.	China.	1816.	G. h.	Rumph. amb. 2. t. 31.
13 angulátæ. DC.	(wh.) angular-fruited.	E. Indies.	G. h.	2. t. 32.
14 buxifolia. DC.	(wh.) refuse-leaved.	China.	G. h.	

ORDO XXXVIII.

HYPERICINEÆ. DC. prodr. 1. p. 541.

Subordo I. HYPERICINEÆ VERÆ. DC.

Tribus I. VISMIEÆ. DC.

HARO'NGA. DC.	HARO'NGA. Polyadelphus Dodecandria.					
1 madagascariensis. DC. (y.)	Madagascar. Madagascar. 1824.	S. h.	Lam. ill. t. 645.			
<i>Arungana paniculata</i> . P.S.						
VI'SMIA. DC.	VI'SMIA. Polyadelphus Polyandria.					
1 sessilifolia. DC. (y.)	sessile-leaved. Guiana. 1826.	S. h.	Aubl. guian. 2.t.312.f.2.			
2 guianensis. DC. (y.)	Guiana. 1828.	S. h.	2. t. 311.			
3 brasiliensis. DC. (y.)	Brasilian. Brazil. 1822. 5. 6.	S. h.	Chois. p. hyp. 35. t. 2.			
4 guineensis. DC. (y.)	African. Sierra Leon. 1823. —	S. h.	Lin. amoen. 8. t. 8. f. 1.			

Tribus II. HYPERICEÆ. DC.

ANDROSÆMUM. DC. TUTSAN. Monadelphus Polyandria.

officinale. DC. (y.) common.	Britain.	7. 9.	H. h.	Eng. bot. 1225.
HYPERICUM. DC.	ST. JOHN'S WORT. Polyadelphus Polyandria. L.			
1 elatum. DC. (y.) tall.	N. America. 1762. 7. 8.	H. h.		
2 frondosum. DC. (y.) green.	— 1806. —	H. h.		
3 amœnum. DC. (y.) pretty.	Carolina. 1812. —	H. h.		
4 grandifolium. DC. (y.) great-leaved.	Teneriffe. 1818. —	G. h.		
5 hircinum. DC. (y.) stinking.	S. Europe. 1640. 7. 9.	H. h.		
6 minus. W.D. (y.) smaller.	— —	H. h.		
7 foliosum. DC. (y.) shining.	Azores. 1778. 8.	G. h.		
7 floribundum. DC. (y.) many-flowered.	Madeira. 1779. —	G. h.		
8 olympicum. DC. (y.) Olympic.	Levant. 1706. 7. 9.	H. h.		
9 canariense. DC. (y.) Canary.	Canaries. 1699. —	G. h.		
10 chinense. DC. (y.) Chinese.	China. 1753. 3. 9.	G. h.		
<i>monogynum</i> . B.M.				
11 cordifolium. DC. (y.) heart-leaved.	Nepaul. 1822. 6. 9.	G. h.		
12 pyramidatum. DC. (y.) pyramidal.	Canada. 1759. 7. 8.	H. 4.		
13 A'scyron. DC. (y.) Siberian.	Siberia. 1774. 6. 9.	H. 4.		
14 ascyroides. DC. (y.) large-capsuled.	N. America. 1812. 6. 7.	H. 4.		
15 pátulum. DC. (y.) spreading.	Japan. 1820. 7. 9.	G. h.		
16 Urálum. D.P. (y.) Nepaul.	Nepaul. —	F. h.		
17 Kalmianum. DC. (y.) Kalmia-leaved.	Virginia. 1759. 6. 7.	H. h.		
18 calycinum. DC. (y.) large-flowered.	Ireland.	6. 9.	H. h.	
19 baleàricum. DC. (y.) warted.	Majorca. 1714. 3. 9.	G. h.		
20 virgínicum. DC. (re.) Virginian.	N. America. 1800. 7. 9.	H. 4.		
<i>Eldaea campanulata</i> . Ph.				
21 angulòsum. DC. (y.) tooth'd-flow'r'd.	N. America. 1812. 6. 7.	H. 4.		
22 punctatum. DC. (y.) spotted.	— 1789. 7. 9.	F. 4.		
<i>carymbosum</i> . W. maculatum. Walt.				
23 dolabriiforme. DC. (y.) hatchet-like.	— 1818. —	H. 4.		
24 densiflorum. DC. (y.) dense-flowered.	Virginia. 1812. 6. 8.	F. h.		
25 rosmarinifolium. DC. (y.) Rosemary-ld.	Carolina. —	F. h.		
26 virgatum. DC. (y.) twiggy.	N. America. 1823. 6. 8.	H. 4.		
27 prolificum. DC. (y.) prolific.	— 1758. —	H. h.		
<i>Kalmianum</i> . Maund, bot. gard. 194. t. 49. <i>nec aliorum. foliosum</i> . Jacq. schœnb. 3. t. 299. non Ait.				Wats. dend. brit. t. 88.
28 glaucum. DC. (y.) glaucous.	N. America. 1812. 7. 8.	H. 4.		
29 lœvigatum. H.K. (y.) smooth.	— 1772. —	H. 4.		
30 nudiflorum. DC. (y.) naked-pancled.	— 1811. 8. 10.	H. h.		
31 sphærocápon. DC. (y.) round-capsuled.	— 1827. 6. 8.	H. 4.		

HYPERICINEÆ.

32 quadrângulum. w. (ye.)	square-stalked.	Britain.	7. 8. H. ȝ.	Eng. bot. 370.	
33 dûbiūm. E.B.	(ye.) imperforate.	Europe.	— H. ȝ.	296.	
34 maculâtum. All.	(ye.) spot-flowered.	Barbary.	1802.	— F. ȝ.	All. ped. t. 83. f. 1.	
35 undulâtum. W.en.	(ye.) wave-leaved.	Dahuria.	1828.	— H. ȝ.	Chois. pr. hyp. p. 47.t.6.	
36 attenuâtum. DC.	(ye.) attenuated.	S. Europe.	1827.	— H. ȝ.		
37 rēpens. DC.	(ye.) creeping.	Japan.	1820.	— G. ȝ.	Thunb. jap. t. 31.	
38 japónicum. DC.	(ye.) Japan.	Greece.	1688.	— F. ȝ.	Bocc. mus. 2. t. 12.	
39 crispum. DC.	(ye.) curl-leaved.	N. America.	1818.	— H. ȝ.	Pluk. alm. t. 245. f. 6.	
40 pilosum. DC.	(ye.) pilose.	setòsum. H.K.				
41 simplex. DC.	(ye.) simple-stalked.	Persia.	1812.	— H. ⊖.	Pluk. alm. t. 421. f. 3.	
42 heterophyl'lum. DC.	(y.) various-leaved.	Egypt.	1787.	6. 7. G. ȝ.	Vent. cels. t. 68.	
43 ægyptiacum. DC.	(ye.) Egyptian.	Britain.	7. 8. H. ȝ.	Bot. reg. 196.	
44 humifusum. DC.	(ye.) trailing.	N. S. W.	1818.	— G. ȝ.	Eng. bot. 1226.	
45 pusillum. DC.	(ye.) small.	1822.	— G. ȝ.	Lab. nov. holl. 2. t. 175.		
46 involutum. DC.	(ye.) involute-fld.	Britain.	— H. ȝ.	2. t. 174.	
47 perforatum. DC.	(ye.) perforate.	Italy.	1820.	— H. ȝ.	Eng. bot. 295.	
48 veronése. Sck.	(ye.) Verona.	N. America.	1759.	6. 9. F. ȝ.		
49 quinquenérvium. DC.	(ye.) small-fld.	mùtilum. w.				
50 canadénse. DC.	(wh.) Canadian.	Canada.	1770.	7. 9. H. ȝ.		
51 perfoliánum. w.	(wh.) perfoliate.	Italy.	1785.	5. 6. F. ȝ.		
52 axillâre. DC.	(ye.) clustered.	fasciculâtum. w.	N. America.	1806.	7. F. ȝ.	
53 Elôdes. DC.	(ye.) marsh.	Britain.	7. 8. H. ȝ.	Eng. bot. 109.	
54 tomentosum. DC.	(ye.) woolly.	S. Europe.	1648.	7. 9. F. ȝ.	Moris.his.2.s.5.t.6.n.5.	
55 lanuginosum. DC.	(ye.) downy.	Levant.	1822.	— F. ȝ.		
56 hirsútum. DC.	(ye.) hairy.	Britain.	6. 7. H. ȝ.	Eng. bot. 1156.	
57 nummulârium. DC.	(ye.) Money-wort.	S. Europe.	1818.	— H. ȝ.	Lam. ill. t. 643.	
58 elegans. DC.	(ye.) elegant.	Siberia.	—	7. 9. H. ȝ.	Spr. fl. hal. n. 864. t. 9.	
<i>Kohliânum. Spr.</i>						
59 glandulósom. DC.	(ye.) glandular.	Madeira.	1777.	5. 8. G. ȝ.		
60 refléxum. DC.	(ye.) reflex-leaved.	Teneriffe.	1778.	6. 9. G. ȝ.		
61 púlchrum. DC.	(ye.) small upright.	Britain.	7. 8. H. ȝ.	Eng. bot. 1227.	
62 elodeoides. DC.	(ye.) close paniced.	Nepaul.	1823.	— G. ȝ.		
<i>nervosum. D.P.</i>						
63 barbâtum. DC.	(ye.) bearded.	Scotland.	6. 10. H. ȝ.	Eng. bot. 1985.	
64 montânum. DC.	(ye.) mountain.	Britain.	7. 8. H. ȝ.	371.	
65 fimbriátum. DC.	(ye.) fimbriate.	Pyrenees.	1818.	— H. ȝ.	Vill. delph. 3. t. 44.	
<i>Richeri. Vill.</i>						
66 aethiòpicum. DC.	(ye.) Ethiopian.	C. B. S.	1817.	— G. ȝ.		
67 ciliátum. DC.	(ye.) fringed.	S. Europe.	1739.	7. F. ȝ.	Boc. mus. 2. t. 127.	
68 serpylliòlum. DC.	(ye.) Thyme-leaved.	Levant.	1688.	7. 8. F. ȝ.	Mor.his.2 s.5.t.6.f.2.	
69 triplinérve. DC.	(ye.) triply-nerved.	N. America.	1823.	7. 9. H. ȝ.	Vent. cels. t. 58.	
70 empetrifòlium. DC.	(y.) Crowberry-lt.	S. Europe.	6. 8. H. ȝ.	Wats. dend. brit. 141.	
71 Còris. DC.	(ye.) Coris-leaved.	Levant.	1640.	5. 9. G. ȝ.	Bot. mag. 178.	
72 ericoïdes. DC.	(ye.) Heath-leaved.	S. Europe.	1818.	— G. ȝ.	Cav. ic. 2. t. 122.	
73 mexicanum. DC.	(ye.) Mexican.	Mexico.	1826.	— G. ȝ.	Linn. amœn. 8. t. 8. f. 2.	
74 fasciculâtum. DC.	(ye.) Aspalathus-like.	N. America.	1811.	6. 8. F. ȝ.		
<i>aspalathoides. w.</i>						
75 verticillâtum. DC.	(ye.) whorl-leaved.	C. B. S.	1784.	— G. ȝ.		
76 cochinchinense. DC.	(sc.) Cochinchina.	Cochinchina.	1822.	6. 8. G. ȝ.		
<i>A'SCYRUM. DC.</i>						
<i>A'SCYRUM. Monadelphia Polyandria.</i>						
1 pùlimum. DC.	(ye.) dwarf.	Georgia.	1806.	6. 8. G. ȝ.		
2 Crux A'ndreæ. DC.	(y.) St.Andrew's-cr.	N. America.	1759.	7. G. ȝ.	Burm. am. 146. t. 152.	
3 hypericoïdes. DC.	(ye.) Hypericum-like.	—	—	7. 9. G. ȝ.	— t. 152. f. 1.	
4 stâns. DC.	(ye.) large-flowered.	—	1806.	— G. ȝ.		
5 aplexicâule. DC.	(ye.) stem-clasping.	—	—	— G. ȝ.		
<i>SAROTHRA. W.</i>						
<i>SAROTHRA. Pentandria Trigynia.</i>						
gentianoïdes. w.	(ye.) Gentian-like.	N. America.	1768.	6. 10. H. ⊖.	Pluk. mant. t. 342. f. 2.	
hypericoïdes. n.	<i>Hypericum Sarothra.</i> M.					

Subordo II. HYPERICINEÆ ANOMALÆ. DC.

CARPODO'NTOS. DC. CARPODO'NTOS. Polyandria Pentagynia.

lúcida. DC. (wh.) shining. V. Diem. Is. 1820. G. ȝ. Lab. voy. 2. t. 18.

ORDO XXXIX.

GUTTIFERÆ. DC. prodr. 1. p. 557.

Tribus I. CLUSIEÆ. DC.

MAH'UREA. DC.	MAH'UREA. Polyandria Monogynia.				
palústris. DC. (pu.)	marsh.	Trinidad.	1820. S. h.	Aubl. guian. 1. t. 222.
M'ARILA. DC.	M'ARILA. Monadelphia Polyandria.				
racemosa. DC.	clustered.	W. Indies.	1827. S. h.	
GODO'YA. DC.	GODO'YA. Polyandria Monogynia.				
gemmiflora. M.N. (ye.)	gem-flowered.	Brazil.	1824. S. h.	Mart. nov. gen. 3. t. 74.
CL'USIA. DC.	BALSAM-TREE. Polygamia Monœcia. L.				
1 rosea. DC. (ro.)	Rose-coloured.	America.	1692.	7. 8. S. h.	Cates. car. 2. t. 99.
2 alba. DC. (wh.)	white-flowered.	S. America.	1752.	— S. h.	Jacq. amer. t. 166.
3 flava.	(ye.) yellow-flowered.	Jamaica.	1759.	9. S. h.	Bot. rep. 223.
4 venosa. DC. (wh.)	veiny-leaved.	S. America.	1733. S. h.	Plum. ic. 87. f. 2.
5 nemorosa. DC. (w.pu.)	wood.	W. Indies.	1826. S. h.	
HAV'ETIA. Kth.	HAV'ETIA. Dieccia Tetrandria.				
laurifolia. Kth. (wh.)	Laurel-leaved.	S. America.	1820. S. h.	H. et B. n. g. am. 5. t. 462.
Clusia tetrandra. W.					

Tribus II. GARCINIEÆ. DC.

MARIA'LVA. DC.	MARIA'LVA. Monadelphia Polyandria.				
guianensis. DC. (gr.)	Guiana.	Guiana.	1827. S. h.	Aubl. guian. 2. t. 364.
MICRA'NTHERA. DC.	MICRA'NTHERA. Dieccia Polyandria.				
clusiæfolia. DC.	Clusia-leaved.	Cayenne.	1823. S. h.	Chois. m. h. P. 1. t. 11. 12.
GARCI'NIA. DC.	GARCI'NIA. Polyandria Monogynia.				
1 Mangostana. DC. (ro.)	Mangosteen.	Java.	1789. S. h.	Lod. bot. cab. 845.
2 cornea. DC. (ye.)	nodding-flow'd.	E. Indies.	1817. S. h.	Rumph. amb. 3. t. 30.
3 Cambogia. DC. (ye.)	yellow-flower'd.	—	1822. S. h.	Rheed. mal. 1. t. 24.
4 Cowa. DC. (ye.)	polygamous.	—	— S. h.	

Tribus III. CALOPHYLLEÆ. DC.

MAMM'EA. DC.	MAMMEE-TREE. Polyandria Monogynia. L.				
1 americana. DC. (wh.)	America.	S. America.	1737. S. h.	Jacq. amer. t. 182. f. 32.
2 africana. G.D. (wh.)	African.	Sierra Leon.	1820. S. h.	
XANTHOCH'Ymus. DC.	XANTHOCH'Ymus. Polyadelphia Polyandria.				
1 pictoriæ. H.B. (wh.)	painters.	E. Indies.	1796. S. h.	Roxb. cor. 2. t. 196.
2 dulcis. H.B. (st.)	sweet.	—	1820. S. h.	— 3. t. 270.
3 ovalifolius. H.B.	oval-leaved.	—	— S. h.	
4 luteus. L.C. (ye.)	yellow.	E. Indies.	1823. S. h.	
5 macrophyllus. L.C.	large leaved.	—	— S. h.	
6 purpureus. L.C. (pu.)	purple.	—	— S. h.	
CALOPHY'LLUM. DC.	CALOPHY'LLUM. Polyandria Monogynia. L.				
1 Inophyllum. DC. (wh.)	sweet-scented.	E. Indies.	1793. S. h.	Rheed. mal. 4. t. 38.
2 Tacamaháca. DC. (wh.)	acute-leaved.	Bourbon.	1822. S. h.	Pluk. alm. t. 147. f. 3.
3 Cálabá. DC. (wh.)	Calaba-tree.	W. Indies.	1780. S. h.	Jacq. amer. t. 165.
4 spuriæ. DC. (wh.)	spurious.	E. Indies.	— S. h.	Rheed. mal. 4. t. 39.
Cálabá. L.					
PENTADE'SMA. H.T.	BUTTER-TREE. Polyandria Monogynia.				
butyracea. H.T. (ye.)	African.	Sierra Leon.	1821. S. h.	

Tribus IV. SYMPHONIEÆ. DC.

CANE'LLA. DC.	CANELLA. Dodecandra Monogynia.					
1 álba. DC. (pu.)	broad-leaved.	W. Indies.	1735.	S. h.	Linn. trans. 1. t. 8.
2 laurifolia. L.C. (pu.)	narrow-leaved.			S. h.	
MORONOB'EA. DC.	MORONOB'EA. Monadelphia Pentandria.					
coccinea. DC. (sc.)	scarlet.	Guiana.	1825.	S. h.	Aubl. guian. 2. t. 313.
<i>Sympônia globulifera</i> . L.						
SING'ANA. DC.	SING'ANA. Polyandria Monogynia.					
guianensis. DC. (wh.)	Guiana.	Guiana.	1827.	S. h.	Aubl. guian. sup. 2. t. 230.
<i>Sternbékia lateriflora</i> . W.						

ORDO XL.

MARCGRAVIACEÆ. DC. prodr. 1. p. 565.

Subordo I. MARCGRAVIEÆ. DC.

ANTHOL'OMA. DC.	ANTHOL'OMA. Polyandria Monogynia.					
montâna. DC. (ye.)	mountain.	N. Caledonia.	1820.	S. h.	Labill. voy. t. 41.
MARCGR'AVIA. DC.	MARCGR'AVIA. Polyandria Monogynia. L.					
1 umbellâta. DC. (gr.)	umbelled.	W. Indies.	1792.	6. 8.	S. h.	Hook. exot. flor. 160.
2 coriacea. DC. (gr.)	leathery-leav'd.	Guiana.	1820.	—	S. h.	
3 picta. DC. (gr.)	painted-leav'd.		1827.	S. h.	

Subordo II. NORANTEÆ. DC.

NORA'NTSEA. DC.	NORA'NTSEA. Polyandria Monogynia. L.					
1 guianensis. DC. (pu.)	Guiana.	Guiana.	1827.	S. h.	Aubl. guian. 1. t. 220.
2 brasiliensis. DC.	Brasilian.	Brazil.	1820.	S. h.	
3 índica.	Indian.	Mauritius.	1822.	S. h.	
RUYSCHIA. DC.	RUYSCHIA. Pentandria Monogynia.					
1 Souroubæa. DC. (st.)	Guiana.	Guiana.	1826.	S. h.	Aubl. guian. 1. t. 97.
2 clusiæföia. DC. (pu.)	Clusia-leaved.	W. Indies.	1823.	S. h.	Jacq. amer. t. 51. f. 2.

ORDO XLI.

HIPPOCRATEACEÆ. DC. prodr. 1. p. 567.

HIPPOCR'ATEA. DC.	HIPPOCR'ATEA. Triandria Monogynia. L.					
1 obcordata. DC. (st.)	obcordate.	W. Indies.	1819.	S. h.	Lam. ill. 1. t. 28. f. 1.
scândens. Jacq. amer. t. 9.						
2 ovata. DC. (ye.)	oval-leaved.	S. America.	1739.	S. h.	Lam. ill. 1. t. 28. f. 2.
volubilis. L.						
3 índica. DC. (ye.)	Indian.	E. Indies.	1818.	S. h.	Roxb. cor. 2. t. 130.
4 obtusifolia. DC. (st.)	blunt-leaved.		1820.	S. h.	
5 arborea. DC. (st.)	tree.		1822.	S. h.	Roxb. cor. 3. t. 205.
A'NTHODON. DC.	A'NTHODON. Triandria Monogynia.					
1 paniculatum. DC.	panicled.	Rio Janeiro.	1818.	S. h.	
2 ellipticum. DC.	elliptic-leaved.			S. h.	
TONSE'LLA. DC.	TONSE'LLA. Triandria Monogynia.					
1 pyrifòrmis. H.T. (gr.)	pear-fruited.	Sierra Leon.	1822.	S. h.	
2 scabra. R.S. (gr.)	rough.	Trinidad.	1824	S. h.	Aubl. guian. 1. t. 10.
Tontèlea scândens. Aub.						

HIPPOCRATEACEÆ.

JO'HNIA. F.I.	JO'HNIA. Triandria Monogynia. F. I.				
1 salacioides. F.I. (or.)	entire-leaved. E. Indies.	1822.	S. h.	—
2 coromandeliana. F.I.	saw-leaved.	—	—	S. h.	—
TRIG'ONIA. DC.	TRIG'ONIA. Monadelphia Pent-Decandra.				
1 villosa. DC. (y.re.)	villous. Cayenne.	1820.	S. h.	Aub. gui. 1. t. 149.
2 mollis. DC.	soft-leaved. Brazil.	1823.	S. h.	—
3 laevis. DC. (wh.)	smooth. Guiana.	1826.	S. h.	Aub. gui. 1. t. 150.

ORDO XLII.

ERYTHROXYLEÆ. DC. prodr. 1. p. 573.

ERYTHRO'XYLUM. DC. ERYTHRO'XYLUM. Decandria Trigynia. L.

1 hypericifolium. DC. (w.)	Hypericum-ld. Mauritius.	1818.	S. h.	Cav. dis. 8. t. 230.
2 havanense. DC. (wh.)	Havannah. W. Indies.	1822.	S. h.	Jacq. amer. t. 87. f. 2.
3 laurifolium. DC. (wh.)	Laurel-leaved. Mauritius.	—	—	S. h.	Cav. dis. 8. t. 226.
4 areolatum. DC. (wh.)	Carthagena. Carthagena.	1826.	S. h.	Jacq. amer. t. 187. f. 1. carthagénense. Jacq.

S'ETHIA. Kth.	S'ETHIA. Decandria Monogynia.				
índica. DC. (ye.)	Indian. E. Indies.	1823.	S. h.	Roxb. cor. 1. t. 88.
<i>Erythróxylon monogynum.</i> R.C.					

ORDO XLIII.

MALPIGHIACEÆ. DC. prodr. 1. p. 577.

Tribus I. MALPIGHIEÆ. DC.

MALPI'GHIA. DC. MALPI'GHIA. Decandria Trigynia. L.

1 fucata. DC. (li.)	painted. W. Indies.	1814.	1. 12.	S. h.	Bot. reg. 189.
macrophylla. Colla hort. ripul. t. 11.	non Juss.				
2 urens. DC. (li.)	stinging. S. America.	1737.	7. 10.	S. h.	Bot. reg. 96.
3 angustifolia. DC. (li.)	narrow-leaved. W. Indies.	—	7. 8.	S. h.	Lod. bot. cab. 321.
4 aquifolia. DC. (li.)	Holly-leaved. S. America.	1759.	8. 9.	S. h.	1079.
5 coccifera. DC. (li.)	kermes-oak-ld.	1733.	—	S. h.	Bot. reg. 568.
6 glabra. DC. (pu.)	smooth-leaved.	1757.	3. 7.	S. h.	Bot. mag. 813.
7 biflora. DC. (ye.)	two-flowered.	—	—	S. h.	Cav. dis. 8. t. 234. f. 2.
<i>punicifolia.</i> Cav.		—	S. h.	Plum. ic. t. 166. f. 2.
8 punicifolia. DC. (pu.)	Punica-leaved.	1690.	—	S. h.	
9 nitida. DC. (pu.)	glossy-leaved.	1733.	3. 8.	S. h.	
10 incana. DC. (ro.)	hoary.	Campeachy.	—	S. h.	
11? faginea. DC. (ye.)	Beech-leaved.	S. America.	1822.	Cav. dis. 8. t. 242.
12? dubia. DC. (ye.)	doubtful.	W. Indies.	1824.	Aub. gui. 1. t. 184.
BYRSON'IMA. DC.	BYRSON'IMA. Decandria Trigynia. L.				
1 verbascifolia. DC. (ye.)	Mullein-leaved.	Guiana.	1818.	3. 8.	Aub. gui. 1. t. 181.
2 nervosa. DC. (ye.)	prominent-n'd.	Brazil.	—	S. h.	
3 altissima. DC. (wh.)	tall.	Guiana.	1820.	Aub. gui. 1. t. 182.
4 crassifolia. DC. (ye.)	thick-leaved.	—	1793.	8.	
<i>Malpighia crassifolia.</i> w.				S. h.	
5 chrysophylla. DC. (ye.)	golden-leaved.	S. America.	1822.	Cav. dis. 8. t. 237.
6 spicata. DC. (ye.)	spiked.	Brazil.	1818.	5. 8.	Bot. mag. 2462.
7 lucida. DC. (re.)	wedge-leaved.	W. Indies.	1759.	—	
<i>Malpighia lucida.</i> w.				S. h.	
8 laevigata. DC. (st.)	smooth.	Cayenne.	1826.	S. h.
9 elegans. DC. (li.)	elegant.	Guiana.	1827.	S. h.
10 densa. DC. (re.)	dense-flower'd.	—	1825.	S. h.
11 coriacea. DC. (wh.)	leathery-leaved.	Jamaica.	1814.	—	S. h.

Sloan. his. 2. t. 163. f. 1.

12 pálida. DC.	(ye.) pale-leaved.	W. Indies.	1820.	5. 8.	S. h.	
13? volubilis. DC.	(ye.) twining.	—	1793.	8. 9.	S. h.	Bot. mag. 809.
	<i>Malpighia volubilis.</i> B.M.					
BUNCH'OSIA. DC.	BUNCH'OSIA.	Decandria Monogynia.				
1 glandulosa. DC.	(ye.) 2-glandled.	Antilles.	1804.	3. 5.	S. h.	Cav. dis. 8. t. 239. f. 2.
	<i>Malpighia glandulosa.</i> Cav.					
2 glandulifera. DC.	(ye.) 4-glandled.	Caracas.	1806.	—	S. h.	Jacq. coll. 5. f. 3.
	<i>Malpighia glandulifera.</i> Jacq. ic. 3. t. 469.					
3 cornifolia. DC.	(wh.) Cornus-leaved.	S. America.	1824.	S. h.	
4 polystachya. DC.	(ye.) many-spiked.	W. Indies.	1806.	3. 5.	S. h.	And. bot. rep. 604.
5 media. DC.	(ye.) intermediate.	—	1790.	—	S. h.	
	<i>Malpighia media.</i> H.K.					
6 tuberculata. DC.	(ye.) tuberculate.	Caracas.	1823.	S. h.	Jacq. schœn. 1. t. 104.
7 argentea. DC.	(ye.) silvery.	—	1820.	S. h.	Jacq. frag. t. 83.
8 nitida. DC.	(ye.) glossy-leaved.	W. Indies.	1733.	3. 8.	S. h.	Cav. dis. 8. t. 239. f. 1.
9 canescens. DC. downy-leaved.	—	1742.	S. h.	
10? paniculata. DC.	(pu.) paniced.	—	1733.	S. h.	
GALPHIMIA. DC.	GALPHIMIA.	Decandria Trigynia.				
1 hirsuta. DC.	(ye.) hairy.	Mexico.	1824.	3. 6.	S. h.	
2 glauca. DC.	(ye.) glaucous-leav'd.	—	1826.	S. h.	Cav. ic. 5. t. 489.
3 glandulosa. DC.	(ye.) glandular.	—	1824.	4. 6.	S. h.	— 6. t. 563.
THRYPALLIS. L.	THRYPALLIS.	Decandria Trigynia.				
? brachystachys. B.R. (y.) short-spiked.	Brazil.	1823.	9. 11.	S. h.	— Bot. reg. 1162.	

Tribus II. HYPTAGEÆ. DC.

HIPT'AGE. DC.	HIPT'AGE.	Decandria Monogynia.				
1 Madablotia. DC.	(wh.) clustered.	E. Indies.	1796.	3. 4.	S. h.	Roxb. cor. 1. t. 18.
	<i>Gärtnera racemosa.</i> A.R. 600.					
2 obtusifolia. DC.	(wh.) blunt-leaved.	China.	1822.	S. h.	
	<i>Gärtnera obtusifolia.</i> H.B.					
GAUDICHA'UDIA. DC.	GAUDICHA'UDIA.	Pentandria Monogynia.				
cynanchoides. DC. (ye.)	Cynanchum-like.	Mexico.	1824.	S. h.	H. et B. n. g. am. 5. t. 445.

Tribus III. BANISTERIEÆ. DC.

HIR'ÆA. DC.	HIR'ÆA.	Decandria Trigynia. L.				
1 odorata. DC. sweet-scented.	Guinea.	1822.	S. h.	
2 glaucescens. L.C. glaucescent.	—	—	S. h.	
3 nutans. DC. nodding.	E. Indies.	1818.	5. 8.	S. h.	
4 indica. DC.	(wh.) Indian.	—	—	—	S. h.	Roxb. cor. 2. t. 160.
TRIO'PTERIS. DC.	TRIO'PTERIS.	Decandria Trigynia. L.				
1 sericea. L.C. silky.	S. America.	1820.	S. h.	
2 jamaicensis. DC.	(ye.) Jamaica.	Jamaica.	1818.	5. 9.	S. h.	
3 lucida. K.S.	(pi.) glossy.	Cuba.	1822.	S. h.	H. et B. n. g. am. 5. t. 451.
TETRA'PTERIS. DC.	TETRA'PTERIS.	Decandria Trigynia. L.				
1 buxifolia. DC.	(ye.) Box-leaved.	St. Domingo.	5. 8.	S. h.	Cav. dis. 9. t. 262. f. 1.
2 acapulcensis. K.S.	(ye.) Acapulco.	Mexico.	1824.	S. h.	
3 acutifolia. DC.	(ye.) acute-leaved.	Cayenne.	1826.	S. h.	Cav. dis. 9. t. 261.
4 citrifolia. DC.	(ye.) Citron-leaved.	Jamaica.	1819.	—	S. h.	— 9. t. 260.
5 discolor. DC.	(ye.) two-coloured.	Guiana.	1827.	S. h.	
BANIST'ERIA. DC.	BANIST'ERIA.	Decandria Trigynia. L.				
1 auriculata. DC.	(ye.) eared.	Brazil.	1818.	S. h.	Cav. dis. 9. t. 255.
2 ciliata. DC.	(or.) ciliated.	—	1796.	S. h.	— 9. t. 254.
3 splendens. DC.	(ye.) shining.	S. America.	1812.	S. h.	— 9. t. 253.
	<i>heterophylla. w. fulgens.</i> Cav.					
4 dichotoma. DC.	(ye.) forked.	S. America.	S. h.	Cav. dis. 9. t. 256.
5 emarginata. DC.	(ye.) emarginate.	W. Indies.	1826.	S. h.	— 9. t. 249.
6 sericea. DC.	(ye.) silky.	Brazil.	1810.	S. h.	— 9. t. 258.
7 tomentosa. DC.	(ye.) woolly-leaved.	Antilles.	1818.	5. 8.	S. h.	
8 tiliæfolia. DC.	(pu.) Lime-leaved.	Java.	1820.	S. h.	Vent. choix, t. 50.

MALPIGHIACEÆ.

9 sinemariensis. DC. (<i>ye.</i>)	prickly-leaved.	Guiana.	1824.	S. h. Aubl. guian. 1. t. 185.
10 ferruginea. DC. (<i>bh.</i>)	ferruginous.	Brazil.	1818.	S. h. Cav. dis. 9. t. 248.
11 periplocæfolia. DC. (<i>ye.</i>)	Periploca-ld.	Porto Rico.	—	S. h.
12 laurifolia. DC. (<i>ye.</i>)	Laurel-leaved.	Jamaica.	1733.	7. 8.	S. h. Bet. reg. 937.
13 fulgens. DC.	(<i>ye.</i>) fulgent.	W. Indies.	1759.	S. h.
14 ovata. DC.	(<i>bh.</i>) oval-leaved.	St. Domingo.	1819.	7. 8.	S. h. Vent. choix. t. 51. A.
HETERO'PTERIS. DC. HETERO'PTERIS. Decandria Trigynia. L.					
1 purpurea. DC. (<i>pu.</i>)	purple.	W. Indies.	1759.	S. h. Cav. dis. 9. t. 246. f. 1.
2 parvifolia. DC. (<i>ye.</i>)	small-leaved.	St. Thomas.	1822.	S. h. Vent. choix. t. 51.
3 brachiata. DC. cross-branched.	S. America.	1759.	S. h.
4 chrysophylla. DC. (<i>or.</i>)	Star-apple-ld.	Brazil.	1793.	S. h. Jacq. schœn. 1. t. 105.
5 caerulea. DC. (<i>bl.</i>)	blue-flowered.	W. Indies.	1822.	S. h. Cav. dis. 9. t. 243.
6 floribunda. DC. (<i>bl.</i>)	many-flowered.	Mexico.	1826.	S. h.
7 nítida. DC. glossy.	Brazil.	1809.	S. h. Cav. dis. 9. t. 244.
8 appendiculata. DC.	... long-pointed.	W. Indies.	1823.	S. h.

ORDO XLIV.

ACERINEÆ. DC. prodr. 1. p. 593.

'ACER. DC.		MAPLE. Polygamia Monocia. L.			
1 oblóngum.	(<i>gr.</i>) oblong-leaved.	Nepaul.	1820.	H. 5.
2 tatáricum. DC.	(<i>gr.</i>) Tartarian.	Tartary.	1759.	5. 6.	H. 5.
3 striátum. DC.	(<i>st.</i>) striped-barked.	N. America.	1755.	—	H. 5.
	<i>pensylvanicum. L.</i>				Pall. fl. ros. t. 3.
4 Pseudo-Plátanus. (<i>st.</i>)	Sycamore.	Britain.	4. 5.	H. 5.
β variegátum.	(<i>st.</i>) variegated.	—	—	H. 5.
γ subobtúsum. DC. (<i>st.</i>)	blunt-leaved.	—	—	H. 5.
5 spicátum. DC.	(<i>st.</i>) mountain.	N. America.	1750.	—	H. 5.
	<i>montanum. H.K.</i>				Schm. arb. 1. t. 11.
6 hy'bridum. DC.	(<i>gr.</i>) hybrid.	Hybrid.	—	H. 5.
7 macrophyllum. DC. (<i>st.</i>)	large-leaved.	Columbia.	1827.	H. 5.
8 campéstre. DC.	(<i>gr.</i>) common.	Britain.	5. 6.	H. 5.
9 austriacum. L.en. (<i>gr.</i>)	Austrian.	Austria.	—	H. 5.
10 Opálus. w.	(<i>st.</i>) Italian.	Italy.	1752.	—	H. 5.
11 opalifólium. L.en. (<i>st.</i>)	Opalus-leaved.	S. Europe.	—	H. 5.
12 obtusátum. DC.	(<i>st.</i>) Hungarian.	Hungary.	1818.	—	H. 5.
13 neapolitanum. T.S. (<i>st.</i>)	Neapolitan.	Naples.	1820.	5. 6.	H. 5.
14 créticum. DC.	(<i>gr.</i>) Cretan.	Levant.	1752.	—	H. 5.
15 monspessulánum. DC. (<i>gr.</i>)	Montpelier.	France.	1739.	5.	H. 5.
16 ibéricum. DC.	(<i>gr.</i>) Iberian.	Iberia.	1826.	H. 5.
17 heterophyllum. DC. (<i>gr.</i>)	evergreen.	Levant.	1759.	5. 6.	H. 5.
	<i>sempervirens. L.</i>				Willd. arb. 10. t. 1. f. 1.
18 barbátum. DC.	(<i>gr.</i>) bearded.	N. America.	1812.	4. 5.	H. 5.
19 Lobélia. T.S.	(<i>st.</i>) Lobel's.	Italy.	1826.	H. 5.
20 platanoides. DC.	(<i>st.</i>) Norway.	Europe.	1683.	5. 7.	H. 5.
β laciniátum.	(<i>st.</i>) cut-leaved.	—	—	—	H. 5.
21 saccharinum. DC.	(<i>st.</i>) Sugar.	N. America.	1735.	4. 5.	H. 5.
22 nigrum. DC.	(<i>st.</i>) black-sugar.	—	1812.	—	H. 5.
23 circinnátum. DC. round-leaved.	Columbia.	1827.	H. 5.
24 eriocárpum. DC.	(<i>st.</i>) Sir C. Wager's.	N. America.	1725.	—	H. 5.
	<i>dasycárpum. W.</i>				Mich. f. arb. 2. t. 13.
25 rúbrum. DC.	(<i>re.</i>) red-flowered.	—	1656.	—	H. b.
26 palmátum. DC.	(<i>st.</i>) palmate.	China.	1820.	H. b.
NEGU'NDO. DC.	NEGU'NDO.	Dicecia Pentandria.			
	fraxinifólium. DC. (<i>st.</i>)	Ash-leaved.	N. America.	1688.	4.
	<i>Acer Negundo. L.</i>				Mich. f. arb. 2. t. 16.

ORDO XLV.

HIPPOCASTANEÆ. DC. prodr. 1. p. 597.

<i>ÆSCULUS.</i> DC.	HORSE-CHESTNUT.	Heptandria Monogynia.	L.		
1 <i>Hippocastanum.</i> DC. (<i>w.</i>) common.	Asia.	1629.	4. 5. H. 5.	Schm. arb.	1. t. 38.
β <i>variegata.</i> (<i>w.</i>) striped-leaved.	—	—	— H. 5.		
γ <i>flôre pleno.</i> (<i>w.h.</i>) double-flowered.	—	—	— H. 5.		
2 <i>rubicunda.</i> DC. (<i>re.</i>) red-flowered.	—	—	5. 6. H. 5.	Lod. bot. cab.	1242.
<i>cárnea.</i> W.D. 121 non B. reg.					
3 <i>cárnea.</i> B.R. (<i>fl.</i>) flesh-coloured.	—	—	5. 6. H. 5.	Bot. reg.	993.
4 <i>glabra.</i> DC. (<i>st.</i>) smooth-leaved.	N. America.	1812.	—	H. 5.	
5 <i>ohioensis.</i> DC. (<i>w.h.</i>) Ohio.	—	1818.	—	H. 5.	
6 <i>pallida.</i> DC. (<i>st.</i>) pale-flowered.	—	1812.	6. H. 5.		
<i>P'AVIA.</i>	BUCK'S-EYE-TREE.	Heptandria Monogynia.			
1 <i>macrostachya.</i> DC. (<i>w.</i>) long-spiked.	N. America.	1786.	7. 8. H. h.	Jacq. ecl.	1. t. 9.
<i>Æsculus parviflora.</i> H.K.					
2 <i>rûbra.</i> DC. (<i>re.</i>) red-flowered.	N. America.	1711.	5. 6. H. 5.	Duham. arb.	2. t. 19.
<i>Æsculus Pavia.</i> Lod. bot. cab. 1257.					
β <i>arguta.</i> B.R. (<i>fl.</i>) sharp-toothed.	—	—	— H. 5.	Bot. reg.	993.
γ <i>sublaciniata.</i> W.D. (<i>re.</i>) deep-toothed.	—	—	— H. 5.	Wats. dend. brit.	120.
3 <i>hybrida.</i> DC. (<i>fl.</i>) variegated-fld.	—	1812.	—	H. 5.	
4 <i>discolor.</i> (<i>fl.</i>) dwarf.	—	—	— H. b.	Bot. reg.	310.
<i>Æsculus discolor.</i> Ph. non <i>hybrida.</i> DC.					
5 <i>humilis.</i> B.R. (<i>re.</i>) dwarf.	—	—	— H. b.	Bot. reg.	1018.
6 <i>neglecta.</i> B.R. (<i>st.</i>) neglected.	—	—	— H. 5.		1009.
7 <i>flava.</i> DC. (<i>st.</i>) yellow-flowered.	—	1764.	— H. 5.	Lod. bot. cab.	1280.

ORDO XLVI.

RHIZOBOLEÆ. DC. prodr. 1. p. 599.

CAR'YOCAR. DC.	BUTTER-NUT.	Polyandria Tetragynia.			
1 <i>nuciferum.</i> DC. (<i>re.</i>) nut-bearing.	S. America.	1826.	S. h.	Bot. mag. 2728.
2 <i>glabrum.</i> DC. smooth-leaved.	Guiana.	1820.	S. h.	Aub. gui. 2. t. 240.
3 <i>tomentosum.</i> DC. (<i>w.h.</i>) woolly-leaved.	—	—	S. h.	1. t. 239.

ORDO XLVII.

SAPINDACEÆ. DC. prodr. 1. p. 601.

Tribus I. PAULLINIEÆ. DC.

CARDIOSPE'R'MUM. L.	HEART-SEED.	Octandria Trigynia.	L.		
1 <i>Halicácabum.</i> DC. (<i>w.</i>) smooth-leaved.	India.	1594.	6. 8.	S. ○. .	Bot. mag. 1049.
2 <i>grandiflorum.</i> DC. (<i>w.h.</i>) large-flowered.	Jamaica.	1823.	S. h. .	
3 <i>coluteoides.</i> DC. (<i>w.h.</i>) Colutea-like.	Caracas.	—	S. u. .	
4 <i>Coríndum.</i> L. (<i>w.h.</i>) Parsley-leaved.	Brazil.	1750.	7. 8.	S. ○. .	
5 <i>pubescens.</i> DC. (<i>w.h.</i>) pubescent.	N. Spain.	1820.	—	S. h. .	
<i>URVILL'EA.</i> Kth.	URVILL'EA.	Octandria Monogynia.			
ulmacea. Kth. (<i>w.h.</i>) Elm-leaved.	Caracas.	1823.	S. h. .	H. et B. n. gen. 5. t. 440.
SERI'ANA. Kth.	SERI'ANA.	Octandria Trigynia.	L.		
1 <i>sinuata.</i> DC. (<i>w.h.</i>) sinuate-leaved.	S. America.	7. 9.	S. h. .	Sch. ac. hafn. 3. t. 12. f. 1.

SAPINDACEÆ.

2 <i>divaricata</i> . DC. (wh.) <i>divaricata</i> .	W. Indies.	1823.	S. h. —	Sch. ac. hafn. 3.t.12.f.2.
3 <i>caracasana</i> . DC. (wh.) <i>tooth-leaved</i> .	Caracas.	1816. 7. 9.	S. h. —	Jacq. schœn. 1. t. 99.
4 <i>mexicana</i> . DC. (wh.) <i>Mexican</i> .	Mexico.	1825.	S. h. —	Sch. ac. hafn. 3.t.11.f.3.
5 <i>triternata</i> . DC. (wh.) <i>Supple-Jack</i> .	W. Indies.	1739.	S. h. —	Plum. ed. Burm. t. 112.
<i>Paulinia polyphylla</i> . L.				
PAULLINIA. DC. PAULLINIA.	Octandria Trigynia. L.			
1 <i>pinnata</i> . DC. (wh.) <i>winged-leaved</i> .	S. America.	1752. 7. 9.	S. h. —	Jacq. obs. 3. t. 62. f. 12.
2 <i>Vespertilio</i> . DC. (wh.) <i>Bat-winged</i> .	W. Indies.	1823. —	S. h. —	Sch. act. hafn. 3.t.11.f.1.
3 <i>meliæfolia</i> . DC. (wh.) <i>Melia-leaved</i> .	Brazil.	1818. —	S. h. —	Hook. ex. flor. 110.
4 <i>curassavica</i> . DC. (wh.) <i>shining-leaved</i> .	S. America.	1739. 6. 8.	S. h. —	Jacq. obs. 3. t. 61. f. 1.
5 <i>barbadensis</i> . DC. (wh.) <i>Barbadoes</i> .	W. Indies.	1786.	S. h. —	—— 3. t. 61. f. 9.
6 <i>polyphylla</i> . DC. (wh.) <i>many-leaved</i> .	S. America.	1739. 6. 8.	S. h. —	Pluk. alm. t. 168. f. 5?
7 <i>carthaginensis</i> . DC. (wh.) <i>binate</i> .	Trinidad.	1821. —	S. h. —	Jacq. obs. 3. t. 61. f. 6.
8 <i>caribaea</i> . DC. (wh.) <i>Caribbean</i> .	Caribees.	S. h. —	Jacq. obs. 3. t. 62. f. 7.
9 <i>hispida</i> . DC. (wh.) <i>hispid</i> .	Caracas.	1823.	S. h. —	Jacq. schœn. 3. t. 268.
10 <i>cauliflora</i> . DC. (wh.) <i>stem-flowering</i> .	——	1821. 6. 8.	S. h. —	Jacq. ic. rar. 3. t. 450.

Tribus II. SAPINDEÆ. DC.

SAPI'NDUS. DC.	SOAP BERRY.	Octandria Trigynia. L.		
1 <i>Saponaria</i> . DC. (wh.) <i>common</i>	W. Indies.	1697. 7. 9.	S. h. —	Comm. hort. 1. t. 94.
2 <i>marginatus</i> . DC. (wh.) <i>margined</i> .	Carolina. —	G. h. —	
3 <i>rigidus</i> . DC. (wh.) <i>Ash-leaved</i> .	S. America.	1759. —	S. h. —	Pluk. alm. t. 217. f. 7.
4 <i>senegalensis</i> . DC. (wh.) <i>Senegal</i> .	Senegal.	1823.	S. h. —	Rheed. mal. 4. t. 19.
5 <i>laurifolius</i> . DC. (wh.) <i>Laurel-leaved</i> .	E. Indies.	1818.	S. h. —	
6 <i>longifolius</i> . DC. (wh.) <i>long-leaved</i> .	——	——	S. h. —	
7 <i>emarginatus</i> . DC. (wh.) <i>emarginate</i> .	——	1820.	S. h. —	
8 <i>rubiginosus</i> . DC. (wh.) <i>rusty-leaved</i> .	——	——	S. h. —	Roxb. cor. 1. t. 62.
9? <i>indicus</i> . DC. (wh.) <i>various-leaved</i> .	——	——	S. h. —	
BLIGHIA. DC.	AKEE-TREE.	Octandria Monogynia.		
sápida. DC. (wh.) <i>Ash-leaved</i> .	Africa.	1723.	S. h. —	Kon. an. bot. 2. t. 16.17.
MATA'YBA. DC.	MATA'YBA.	Octandria Monogynia.		
guianensis. DC. (wh.) <i>Ash-leaved</i> .	Guiana.	1803. 7. 9.	S. h. —	Aub. gui. 1. t. 128.
<i>Ephèliis fraxinea</i> . W.				
SCHMI'DELIA. DC.	SCHMI'DELIA.	Octandria Monogynia. L.		
1 <i>serrata</i> . DC. (wh.) <i>saw-leaved</i> .	E. Indies.	1804. 6. 8.	S. h. —	Roxb. cor. 1. t. 61.
<i>Ornitrophe serrata</i> . W.				
2 <i>racemosa</i> . DC. (wh.) <i>clustered</i> .	——	1818.	S. h. —	Burm. ind. t. 32. f. 1.
3 <i>Cominia</i> . DC. (wh.) <i>yellow-berried</i> .	Jamaica.	1759. 6. 9.	S. h. —	Sloan. h. 2. t. 208 f. 1.
<i>Ornitrophe Cominia</i> . W. <i>Rhùs Cominia</i> . L.				
EUPH'ORIA. DC.	EUPH'ORIA.	Octandria Monogynia. L.		
1 <i>Litchi</i> . DC. (wh.) <i>Lee-Chee</i> .	China.	1786. 5. 6.	S. h. —	Lam. ill. t. 306.
<i>Dimocarpus Litchi</i> . W. <i>Scytalium chinensis</i> . G. fr. t. 42. f. 3.				
2 <i>Longana</i> . DC. (wh.) <i>loose-paniced</i> .	China.	—— —	S. h. —	Buchoz. ic. col. t. 99.
3 <i>verticillata</i> . B.R. (bh.) <i>whorl-flowered</i> .	Moluccas.	1822. 2. 3.	S. h. —	Bot. reg. 1059.
4 <i>Nephelium</i> . DC. (wh.) <i>Rambootan</i> .	E. Indies.	1809.	S. h. —	Lam. ill. t. 764.
<i>Nephelium lappaceum</i> . W. <i>Scytalium Rambootán</i> . H.B.				
CUP'ANIA. DC.	CUP'ANIA.	Octandria Monogynia.		
1 <i>tomentosa</i> . DC. (wh.) <i>woolly-leaved</i> .	W. Indies.	1818. 5. 7.	S. h. —	
<i>Trigoniis tomentosa</i> . Jacq. am. 102.				
2 <i>glabra</i> . DC. (wh.) <i>smooth-leaved</i> .	——	—— —	S. h. —	
3 <i>Saponaria</i> . DC. (wh.) <i>Soap-wort-l</i> .	——	1820.	S. h. —	
MOLIN'EA. J.	MOLIN'EA.	Octandria Monogynia.		
canescens. H.B. (wh.) <i>canescent</i> .	E. Indies.	1817. 5. 6.	S. h. —	
COSSI'GNIA. DC.	COSSI'GNIA.	Hexandria Monogynia.		
borbónica. DC. (wh.) <i>golden-leaved</i> .	Bourbon.	1811.	S. h. —	Lam. ill. t. 256.
<i>pinnata</i> . Lam. <i>Rúzia áurea</i> . Hort.				
MELIC'OCCA. DC.	HONEY-BERRY.	Octandria Monogynia. L.		
1 <i>bijuga</i> . DC. (wh.) <i>winged-leaved</i> .	Jamaica.	1778.	S. h. —	Jacq. amer. 103. t. 72.
2 <i>olijæformis</i> . DC. (wh.) <i>large-leaved</i> .	S. America.	1818.	S. h. —	

Tribus III. DODONÆACEÆ. DC.

KOELREUTERIA. DC.	KOELREUT'ERIA.	Octandria Monogynia. L.				
paniculata. DC.	(ye.) paniced.	China.	1763.	7. 8. H.	I ₂ .	Bot. reg. 320.
DODONÆA. DC.	Dodon'ÆA.	Octandria Monogynia. L.				
1 viscósa. DC.	(gr.) clammy.	S. America.	1690.	6. 7.	S. I ₂ .	Plum. ic. t. 247. f. 2.
2 jamaicensis. DC.	(gr.) Jamaica.	Jamaica.	—	—	S. I ₂ .	Cav. ic. t. 327.
viscosa. Cav.	angustifolia. Sw.					
3 Burmanniana. DC.	(gr.) Burman's.	E. Indies.	1758.	5. 8.	S. I ₂ .	Burm. zeyl. t. 23.
angustifolia. H.B.						
4 microcarpa. DC.	(gr.) small-fruited.	Bourbon.	1818.	—	S. I ₂ .	Lam. ill. t. 304. f. 2.
5 attenuata. F.T.	(gr.) attenuated.	N. S. W.	1824.	2. 6.	G. I ₂ .	Bot. mag. 2860.
6 salicifolia. DC.	(gr.) Willow-leaved.	C. B. S.	1758.	—	G. I ₂ .	
angustifolia. Lam.						
7 dioica. DC.	(gr.) diecious.	E. Indies.	—	S. I ₂ .	
8 triquetra. DC.	(gr.) three-sided.	N. S. W.	1790.	6. 8.	G. I ₂ .	Andr. rep. 230.
9 oblongifolia. L.en.	(gr.) oblong-leaved.	—	1816.	—	G. I ₂ .	Bot. reg. 1051.
10 cuneata. DC.	(gr.) wedge-leaved.	—	—	—	G. I ₂ .	Linn. trans. 11. t. 19.
11 asplenifolia. DC.	(gr.) Asplenium-ld.	—	—	—	G. I ₂ .	— 11. t. 20.
12 filiformis. DC.	(gr.) slender-leaved.	—	1820.	—	G. I ₂ .	
13 angustissima. DC.	(gr.) narrow-leaved.	—	—	—	G. I ₂ .	
14 elaeagnoides. DC.	(gr.) scaly leaved.	W. Indies.	1819.	S. I ₂ .	
15? pinnata. DC.	(gr.) winged-leaved.	N. S. W.	1810.	G. I ₂ .	

ORDO XLVIII.

MELIACEÆ. DC. prodr. 1. p. 619.

Tribus I. MELIEÆ. DC.

AIT'ONIA. W.	AIT'ONIA.	Monadelphia Octandria.				
capensis. w.	(pi.) Cape.	C. B. S.	1774.	5. 9.	G. I ₂ .	Bot. mag. 173.
TURRÆA. DC.	TURRÆA.	Decandria Monogynia. L.				
1 virens. DC.	(ye.) green.	E. Indies.	1822.	S. I ₂ .	Sm. ic. ined. 1. t. 10.
2 rigidula. DC.	(li.) rigid.	Mauritius.	1820.	S. I ₂ .	Vent. choix. t. 48.
QUIVISIA. DC.	QUIVISIA.	Octandria Monogynia.				
heterophylla. DC.	(bh.) various-leaved.	Mauritius.	1821.	S. I ₂ .	Cav. dis. 7. t. 213.
SAND'ORICUM. DC.	SAND'ORICUM.	Decandria Monogynia.				
indicum. DC.	(st.) Indian.	E. Indies.	1818.	S. I ₂ .	Roxb. corom. 3. t. 261.
M'ELIA. DC.	BEAD-TREE.	Decandria Monogynia. L.				
1 Azedarach. DC.	(li.) common.	Syria.	1656.	6. 8.	G. I ₂ .	Bot. mag. 1066.
2 supérbia. H.B.	(li.) superb.	E. Indies.	1821.	1. 2.	S. I ₂ .	
3 robusta. H.B.	(li.) robust.	—	—	—	S. I ₂ .	
4 sempervirens. DC.	(li.) evergreen.	Jamaica.	1656.	6. 9.	S. I ₂ .	Bot. reg. 643.
5 composita. DC.	(li.) compound-ld.	E. Indies.	1818.	1. 3.	S. I ₂ .	
6 Azadirachta. DC.	(li.) Ash-leaved.	—	1759.	6. 8.	S. I ₂ .	Cav. dis. 7. t. 208.
7 australis.	(li.) New Holland.	N. S. W.	1824.	G. I ₂ .	

Tribus II. TRICHILIEÆ. DC.

TRICHI'LIA. DC.	TRICHI'LIA.	Decandria Monogynia. L.				
1 hirta. DC.	(wh.) hairy.	Jamaica.	1820.	6. 7.	S. I ₂ .	Sloan. h. 2. t. 220. f. 1.
2 spondioides. DC.	(st.) Spondias-like.	—	—	—	S. I ₂ .	Jacq. schœn. 1. t. 102.
3 havanensis. DC.	(st.) smooth.	W. Indies.	1794.	—	S. I ₂ .	Jacq. amer. t. 175. f. 38.
glabra. L.						
4 odorata. A.R.	(gr.) sweet-scented.	—	1801.	—	S. I ₂ .	Andr. rep. 637.
5 terminalis. DC.	(wh.) terminal.	Jamaica.	1825.	—	S. I ₂ .	
6 glandulosa. DC.	(gr.) glandular.	N. S. W.	1821.	—	G. I ₂ .	
7 trifoliata. DC.	(wh.) three-leaved.	S. America.	1828.	—	S. I ₂ .	Jacq. amer. 129. t. 82.

EKEBERGIA. DC.	EKEBE'RGIA.	Decandria Monogynia.				
1 capensis. DC.	(wh.) Cape.	C. B. S.	1789.	7. 8.	G. h.	Lam. ill. t. 358.
2 indica. H.B.	(wh.) Indian.	E. Indies.	1826.	S. h.	
GUARÉA. DC.	GUARÉA.	Octandria Monogynia. L.				
1 trichilioïdes. L.	(wh.) Ash-leaved.	S. America.	1752.	5. 6.	S. h.	Jacq. am. t. 176. f. 37.
grandifolia. DC.	<i>Melia</i> Guára. Jacq.					
2 Swartzii.	(wh.) Swartz's.	W. Indies.	1822.	S. h.	Sloan. h. 2. t. 170. f. 1.
trichilioïdes. Sw.						
3 ramiflora. DC.	(li.) branch-flow'r'd. Porto Rico.	—	S. h.	Vent. choix. t. 41.	
HEYNEA. DC.	HEY'NEA.	Decandria Monogynia.				
1 trijuga. DC.	(wh.) Walnut-like.	Nepaul.	1812.	9. 10.	S. h.	Bot. mag. 1738.
2 quinquejuga. H.B. (w.)	five-paired.	Moluccas.	1821.	S. h.	
AMOO'RA. R.C.	AMOO'RA.	Hexandria Monogynia.				
cucullata. R.C.	(ye.) hooded.	E. Indies.	1824.	S. h.	Roxb. corom. 3. t. 258.

ORDO XLIX.

CEDRELEÆ. Brown gen. rem. 64.

CEDRELA. DC.	BASTARD CEDAR.	Pentandria Monogynia. L.				
1 odorata. DC.	(st.) fragrant.	W. Indies.	1739.	7. 9.	S. h.	Lam. ill. t. 137.
2 Toóna. DC.	(st.) Nepaul.	Nepaul.	1821.	S. h.	Roxb. cor. 3. t. 238.
3 velutina. DC. velvetty.	S. h.	
SWIETENIA. DC.	MAHOGANY-TREE.	Decandria Monogynia. L.				
1 Mahogoni. DC.	(wh.) common.	W. Indies.	1734.	S. h.	Cav. dis. 7. t. 209.
2 febrifuga. DC.	(wh.) Febrifuge.	E. Indies.	1796.	S. h.	Roxb. cor. 1. t. 17.
CHLORO'XYLON. DC.	CHLORO'XYLON.	Decandria Monogynia.				
Swieténia. DC.	(ye.) Satin-wood.	E. Indies.	1822.	S. h.	Roxb. cor. 1. t. 64.
Swieténia chloróxylon. R.C.						
FLINDE'RSIA. DC.	FLINDE'RSIA.	Decandria Monogynia.				
austrális. DC.	(wh.) New Holland.	N. Holland.	1821.	G. h.	Brown gen. rem. t. 1.
CARA'PA. DC.	CARA'PA.	Decandria Monogynia.				
1 prôcera. DC.	(st.) tall.	W. Indies.	S. h.	
2 guineáensis.	(st.) glossy-leaved.	Sierra Leon.	1793.	S. h.	
Afzélia spléndens. Hort.						
3 guianénsis. DC.	(st.) Guiana.	Guiana.	1820.	S. h.	Aubl. guian. 2. s. t. 387.

ORDO L.

AMPELIDEÆ. DC. prodr. 1. p. 627.

Tribus I. VINIFERÆ seu SARMENTACEÆ. DC.

CI'SSUS. DC.	CI'SSUS.	Tetrandria Monogynia. L.				
1 vitiginea. DC.	(re.) vine-leaved.	India.	1772.	7. 8.	S. h. —	Pluk. mant. t. 337. f. 4.
2 adnata. DC.	(gr.) pointed-leaved.	E. Indies.	1818.	—	S. h. —	
3 gláuca. DC.	(gr.) glaucous.	—	—	—	S. h. —	
4 c'æsia. DC.	(gr.) grey.	Guinea.	1822.	5. 6.	S. h. —	
5 r'epens. DC.	(pu.) creeping.	E. Indies.	1821.	—	S. h. —	Rheed. mal. 7. t. 48.
6 puncticulòsa DC.	(gr.) strumous.	Cayenne.	1818.	S. h. —	
7 uvífera. DC.	(gr.) berry-bearing.	Guinea.	1822.	5. 6.	S. h. —	
8 tiliacea. DC.	(gr.) Lime-leaved.	Mexico.	1825.	G. h. —	
9 quadranguläris. DC. (w.)	square-stalked.	E. Indies.	1790.	S. h. —	Rumph. amb. 5. t. 44. f. 2.
10 sicyoides. DC.	(gr.) naked-leaved.	Jamaica.	1768.	7. 8.	S. h. —	Jacq. amer. 22. t. 15.
11 smilacina. DC.	(gr.) Smilax-like.	S. America.	1826.	S. h. —	
12 ovata. DC.	(gr.) oval-leaved.	—	1822.	S. h. —	Brown. jam. t. 4. f. 1. 2.
13 antárctica. DC.	(gr.) Kanguru-Vine.	N. S. W.	1790.	6. 8.	G. h. —	Bot. mag. 2488.

14 glandulosâ. DC.	(gr.) glandular.	Arabia.	1824.	6. 7.	G. $\frac{1}{2}$. \sim .
15 capénsis. DC.	(gr.) Cape.	C. B. S.	1792.	—	G. $\frac{1}{2}$. \sim .
16 ácida. DC.	(gr.) acid.	Jamaica.	1692.	—	S. $\frac{1}{2}$. \sim . Jacq. schœn. 1. t. 33.
17 carnòsa. DC.	(wh.) flat-stemmed.	E. Indies.	1818.	S. $\frac{1}{2}$. \sim . Rheed. mal. 7. t. 9.
18 trifoliata. DC.	(re.) three-leaved.	Jamaica.	1739.	6. 8.	S. $\frac{1}{2}$. \sim . Sloan.jam.t.142.f.5.6.?
19 pacifîora. DC.	(gr.) few-flowered.	C. B. S.	1818.	5. 6.	G. $\frac{1}{2}$. \sim .
20 alâta. DC.	(re.) winged-stalked.	Jamaica.	—	—	S. $\frac{1}{2}$. \sim . Sloan.jam. 1. t. 144.
21 angustifòlia. DC.	(st.) narrow-leaved.	E. Indies.	1818.	S. $\frac{1}{2}$. \sim .
22 cirrhosa. DC.	(gr.) tendrilled.	C. B. S.	1822.	5. 6.	G. $\frac{1}{2}$. \sim .
23 quinâta. DC.	(gr.) wedge-leaved.	—	1790.	7.	G. $\frac{1}{2}$. \sim .
24 pentaphylla. DC.	(gr.) Japan.	Japan.	—	4. 9.	G. $\frac{1}{2}$. \sim .
25 quinquefòlia. B.M.	(gr.) five-leaved.	Brazil.	—	8. 9.	S. $\frac{1}{2}$. \sim . Bot. mag. 2443.
26 diversifòlia. DC.	(gr.) various-leaved.	—	—	S. $\frac{1}{2}$. \sim .
<i>heterophylla</i> . L.en.non DC.					
27 elongâta. DC.	(gr.) long-branched.	E. Indies.	1818.	S. $\frac{1}{2}$. \sim .
28 auriculâta. DC.	(gr.) ear-stipuled.	—	—	S. $\frac{1}{2}$. \sim .
29 orientalis. DC.	(gr.) oriental.	Levant.	—	6. 8.	G. $\frac{1}{2}$. \sim . Lam. ill. t. 84. f. 2.
AMPELO'PSIS. DC. AMPELO'PSIS. Pentandria Monogynia. L.					
1 cordâta. DC.	(gr.) heart-leaved.	N. America.	1803.	6. 8.	H. $\frac{1}{2}$. \sim .
<i>Cissus Ampelopsis</i> . P.S.					
2 hederacea. DC.	(gr.) Virginian creeper.	—	1629.	—	H. $\frac{1}{2}$. \sim . Corn. canad. t. 100.
<i>Hedera quinquefolia</i> . L. <i>Cissus hederacea</i> . P.S.					
3 hirsûta. DC.	(gr.) hairy.	—	1806.	—	H. $\frac{1}{2}$. \sim .
4 bipinnata. DC.	(gr.) Pepper Vine.	—	1700.	7. 9.	H. $\frac{1}{2}$. \sim . Pluk. mant. t. 412. f. 2.
<i>Vitis arborea</i> . W. <i>Cissus stans</i> . P.S.					
VITIS. DC. VINE. Pentandria Monogynia. L.					
1 vimifera. DC.	(gr.) common Grape. Asia.	—	6. 7.	H. $\frac{1}{2}$. \sim . Jacq. ic. 1. t. 50.
2 laciniosa. DC.	(gr.) Parsley-leaved.	1648.	—	H. $\frac{1}{2}$. \sim . Schm. ic. 34. t. 8.	
3 dentata. DC.	(gr.) toothed.	1818.	H. $\frac{1}{2}$. \sim .	
4 Wallichii. DC.	(gr.) Wallich's.	Nepaul.	—	H. $\frac{1}{2}$. \sim .
5 indica. DC.	(g.pu.) Indian.	E. Indies.	1692.	S. $\frac{1}{2}$. \sim . Rheed. mal. 7. t. 6.
6 lanâta. F.I.	(gr.) woolly.	—	1824.	S. $\frac{1}{2}$. \sim .
7 latifolia. F.I.	(br.) broad-leaved.	—	—	S. $\frac{1}{2}$. \sim . Rheed. mal. 7. t. 7.
8 caribæa. DC.	(st.) West Indian.	W. Indies.	—	S. $\frac{1}{2}$. \sim . Sloan. his. 2.t.210.f. 4.
9 Labrusca. DC.	(gr.) downy-leaved.	N. America.	1656.	5. 6.	H. $\frac{1}{2}$. \sim . Jacq. schœn. t. 426.
10 blanda.	(gr.) white-berried.	—	—	—	H. $\frac{1}{2}$. \sim .
11 æstivalis. DC.	(gr.) Fox Grape.	—	—	—	H. $\frac{1}{2}$. \sim . Jacq. schœn. t. 425.
<i>vulpina</i> . W.					
12 cordifolia. DC.	(gr.) Winter Grape.	—	1806.	H. $\frac{1}{2}$. \sim .
13 riparia. DC.	(gr.) sweet-scented.	—	—	5. 6.	H. $\frac{1}{2}$. \sim . Bot. mag. 2429.
14 rotundifolia. DC.	(gr.) Bullet Grape.	—	—	—	H. $\frac{1}{2}$. \sim .
15 palmata. DC.	(gr.) palmate-leaved.	—	1820.	—	H. $\frac{1}{2}$. \sim .

Tribus II. LEEACEÆ. DC.

LE'EA. DC.	LE'EA. Pentandria Monogynia. L.
1 sambucina. DC.	(st.) Elder-leaved.
2 críspa. DC.	(wh.) fringe-stalked.
3 pinnata. A.R.	(gr.) round-stalked.
4 hírta. DC.	(gr.) hairy.
5 robusta. F.I.	(gr.) robust.
6 macrophylla. DC.	(wh.) large-leaved.
E. Indies.	1790. S. $\frac{1}{2}$. Cav. dis. 7. t. 218.
C. B. S.	1676. 8. 10. S. $\frac{1}{2}$. Bot. rep. t. 355.
E. Indies.	1777. S. $\frac{1}{2}$.
—	1816. S. $\frac{1}{2}$.
—	1820. S. $\frac{1}{2}$.
—	1806. S. $\frac{1}{2}$.

ORDO LI.

GERANIACEÆ. DC. prodr. 1. p. 637.

Tribus I. GERANIEÆ. S.G. 1. p. vii.

SARCOCA'ULON. DC.	SARCOCA'ULON.	Monadelphia Dodecandria.
1 L'Heritiéri. (ye.)	L. Heritiér's.	C. B. S. 1790. 5. 6. G. ½. L. Herit. ger. t. 42.
<i>Monsónia</i> L'Heritiéri. DC.	<i>Monsónia spinosa</i> .	L. Her.
2 Burmánni. DC. (ye.)	Burman's.	C. B. S. 1790. 7. G. ½. Burm. afr. t. 31.
<i>Geránium spinósrum</i> . Cav. dis. 4. t. 75. f. 2.		
MONSONIA. S.G.	MONS'ONIA.	Monadelphia Dodecandria.
1 ováta. DC. (st.) oval-leaved.	C. B. S.	1774. 8. G. ½. Cav. dis. 4. t. 113. f. 1.
2 lobátâ. DC. (st. bh.) broad-leaved.	——	—— 4. 5. G. ¼. Sweet ger. 3. t. 273.
3 pilósa. W. en. (bh.) hairy.	——	1778. 7. 8. G. ¼. —————— 2. t. 199.
<i>filia</i> . A.R. 276. nec <i>aliorum</i> .		
4 speciosa. L. (va.) large-flowered.	——	1774. 4. 5. G. ¼. Bot. mag. 73.
<i>a rubra</i> . (ro.) red-flowered.	——	—— 12. 2. G. ¼. Sweet ger. 1. t. 77.
<i>β pálida</i> . (st.) pale-flowered.	——	—— 4. 5. G. ¼. ——————
GER'ANIUM. DC.	CRANES-BILL.	Monadelphia Decandria.
1 sibiricum. DC. (wh.) Siberian.	Siberia.	1758. 6. 7. H. ¼. Jacq. vind. 1. t. 19.
2 sanguíneum. DC. (cr.) bloody.	Britain. 6. 9. H. ¼. Eng. bot. 272.
3 prostrátum. L.en (std.) Lancashire.	—— —— H. ¼. Cav. dis. 4. t. 76. f. 3.
4 multífidum. s.g. (li.) multifid.	C. B. S.	1817. —— G. ¼. Sweet ger. 3. t. 245.
5 incánum. DC. (li.) hoary.	——	1701. 5. 7. G. ¼. Cav. dis. 4. t. 82. f. 2.
6 canéscens. DC. (li.) silky-leaved.	——	1787. —— G. ¼. L. Herit. ger. t. 38.
7 argénteum. DC. (bh.) silvery-leaved.	S. Europe.	1699. 6. 7. H. ¼. Sweet ger. 1. t. 59.
8 cinéreum. DC. (vi.) grey.	Pyrenees. 6. 8. H. ¼. Cav. dis. 4. t. 89. f. 1.
<i>várium</i> . L'Her. ger. t. 37.		
9 anemonefólium. DC. (p.) Anemone-l'd.	Canaries.	1778. 5. 8. G. ½. Sweet ger. 3. t. 244.
10 macrorhizon. DC. (re) long-rooted.	Italy.	1576. 5. 6. H. ½. —————— 3. t. 271.
11 tuberósrum. DC. (re) tuberous-rooted.	——	1596. 5. 8. H. ¼. —————— 2. t. 155.
12 ibéricum. DC. (bl.) Iberian.	Iberia.	1802. 6. 9. H. ¼. —————— 1. t. 84.
13 nodósum. DC. (std.) knotted.	England. 5. 10. H. ¼. Eng. bot. 1091.
14 angulátum. DC. (std.) angular-stalk'd.	——	1789. 5. 6. H. ¼. Bot. mag. 203.
15 Lambérti. s.g. (li.) Mr. Lambert's.	Nepaul.	1825. 8. 10. H. ¼. Sweet ger. 4. t. 338.
16 Wallichianum. s.g. (p.) Wallich's.	——	1820. 6. 10. H. ¼. —————— 1. t. 90.
17 Vlassovianum. DC. (st.) Vlassof's.	Siberia. 5. 8. H. ¼. —————— 3. t. 228.
18 stríatum. DC. (std.) streaked.	Italy.	1629. 5. 10. H. ¼. Bot. mag. 55.
19 refléxum. DC. (re) reflex-flowered.	——	1758. 5. 6. H. ¼. Cav. dis. 4. t. 81. f. 1.
20 ph'aëum. DC. (br.) dusky.	England. 4. 6. H. ¼. Eng. bot. 322.
21 fúscum. w. (da.) brown.	S. Europe.	1759. 7. H. ¼.
22 lívidum. w. (li.) wrinkle-leaved.	Switzerland.	1775. 6. 7. H. ¼.
23 erióstemon. DC. (bl.) woolly-stamen'd	Dahuria.	1822. 7. 9. H. ¼.
<i>a coréleum</i> . s.g. (bl.) bright blue.	——	—— H. ¼.
<i>β pállidum</i> . s.g. (pa.) pale blue.	Nepaul.	—— —— H. ¼.
24 sylváticum. DC. (pu.) wood.	Britain. 5. 6. H. ¼.
25 batracioídes. L.en. (b.) hairy.	Europe. —— H. ¼.
26 praténsé. DC. (bl.) Crowfoot-leav'd	Britain. 5. 7. H. ¼.
<i>a coréleum</i> .	(bl.) blue-flowered. —— H. ¼.
<i>β variegátum</i> .	(v.) variegated. —— H. ¼.
<i>γ álbum</i> .	(wh.) white-flowered. —— H. ¼.
<i>δ flóre pléno</i> .	(bl.) double-blue. —— H. ¼.
27 lóngipes. DC. (li.) long-peduncled.	——	1824. —— H. ¼.
<i>Londésii</i> . L.en.		
28 maculátum. DC. (bh.) spotted.	N. America.	1732. 5. 8. H. ¼. Sweet ger. 4. t. 332.
29 collinum. DC. (re.) hill.	Tauria.	1815. —— H. ¼.
30 palústre. DC. (re.) marsh.	Europe.	1732. —— H. ¼.
31 aconítifólium. DC. (wh.) Aconite-leaved.	Switzerland.	1775. 5. 6. H. ¼.
32 pilósrum. DC. (li.) pointed anth'l'd	N. Zealand.	1821. 5. 8. F. ¼.
33 parviflórum. DC. (li.) small-flowered.	Van Diem.	1816. —— G. ¼.
34 nepalénse. s.g. (li.) Nepaul.	Nepaul.	1819. —— H. ¼.
35 cristátum. DC. (li.) crested.	Iberia.	1823. —— H. ¼.
<i>albánum</i> . M.B.		

GERANIACEÆ.

36 <i>pyreniacum</i> . DC. (<i>li.</i>) mountain.	Britain.	5. 8. H. ȝ.	Eng. bot. 405.
37 <i>umbrosum</i> . P.S. (<i>li.</i>) naked-stalked.	Hungary. 1804.	6. 9. H. ȝ.	W.etKit.hung.2.t.124.
38 <i>mölle</i> . DC. (<i>li.</i>) Dove's-foot.	Britain.	4. 8. H. ȝ.	Eng. bot. 778.
39 <i>pusillum</i> . DC. (<i>li.</i>) small-flowered.	England.	6. 9. H. ȝ.	—— 385.
40 <i>rotundifolium</i> . DC. (<i>li.</i>) round-leaved.	——	6. 7. H. ȝ.	—— 157.
41 <i>columbinum</i> . DC. (<i>li.</i>) long-stalked.	Britain.	—— H. ȝ.	—— 259.
42 <i>dissectum</i> . DC. (<i>li.</i>) jagged-leaved.	——	5. 7. H. ȝ.	—— 753.
43 <i>carolinianum</i> . DC. (<i>st.</i>) veined petaled.	N. America. 1725.	7. 8. H. ȝ.	Cav. dis. 4. t. 84. f. 1.
44 <i>lanuginosum</i> . W.en. (<i>p.</i>) woolly.	——	—— H. ȝ.	Jacq. schoen. 2. t. 140.
45 <i>villosum</i> . DC. (<i>li.</i>) villous.	Naples. 1824.	—— H. ȝ.	Sweet ger. ser. 2. t.
46 <i>bohemicum</i> . DC. (<i>bl.</i>) Bohemian.	Bohemia. 1683.	6. 8. H. ȝ.	W. etKit.hung.2.t.123.
47 <i>divaricatum</i> . DC. (<i>li.</i>) straddling.	Hungary. 1799.	7. 8. H. ȝ.	Eng. bot. 75.
48 <i>lucidum</i> . DC. (<i>li.</i>) shining.	Britain.	5. 8. H. ȝ.	Eng. bot. 1486.
49 <i>Briceanum</i> . (<i>wh.</i>) Miss Brice's.	Bristol.	4. 10. H. ȝ.	Vill. dauph. 3. t. 40.
50 <i>Robertianum</i> . DC. (<i>pu.</i>) Herb-Robert.	Britain.	—— H. ȝ.	ERODIUM. DC. HERON'S-BILL. <i>Monadelphia Pentandria.</i>
51 <i>purpureum</i> . w. (<i>pu.</i>) purple.	——	—— H. ȝ.	S. Europe. 1640. 6. 7. H. ȝ.
			Spain. 1798. — H. ȝ.
			Dahuria. 1820. — H. ȝ.
			Italy. 1814. 5. 8. H. ȝ.
			Cyprus. 1788. 3. 8. G. ḥ.
			Crete. 1794. 5. 8. G. ȝ.
			Egypt. 1823. — G. ȝ.
			S. Europe. 1711. 6. 7. H. ȝ.
			Britain.
			Europe. 1823. — H. ȝ.
1 <i>petræum</i> . DC. (<i>pu.</i>) rock.	Numidia. 1803. 5. 6. H. ȝ.	Gouan. ill. t. 21. f. 1.	
2 <i>glandulosum</i> . DC. (<i>pu.</i>) glandulous.	Rome. 1724. — H. ȝ.	Cav. dis. 5. t. 125. f. 2.	
3 <i>Stephaniænum</i> . D.C. (<i>bl.</i>) multifid.	S. Europe. 1816. 6. 8. H. ȝ.	L'Herit. ger. t. 3.	
4 <i>alpinum</i> . DC. (<i>bl.</i>) Alpine.	England.	Sweet ger. 2. t. 111.	
5 <i>crassifolium</i> . DC. (<i>li.</i>) thick-leaved.	Italy. 1814. 5. 8. H. ȝ.	Cav. dis. 4. t. 113. f. 3.	
6 <i>laciniatum</i> . DC. (<i>li.</i>) laciniated.	Creta. 1794. 5. 8. G. ȝ.	Jacq. f. ecl. 1. t. 58.	
7 <i>hirtum</i> . DC. (<i>li.</i>) hairy.	Egypt. 1823. — G. ȝ.	Jacq. vind. 1. t. 18.	
8 <i>Ciconium</i> . DC. (<i>li.</i>) long-beaked.	S. Europe. 1711. 6. 7. H. ȝ.	Eng. bot. 1768.	
9 <i>cicutarium</i> . E.B. (<i>pi.</i>) Hemlock-leav'd	Britain.	Cav. dis. 4. t. 95. f. 2.	
10 <i>pimpinellæfolium</i> . (<i>pi.</i>) <i>Pimpinella</i> -ld.	Europe. 1823. — H. ȝ.	—— 5. t. 126. f. 3.	
11 <i>bipinnatum</i> . w. (<i>pi.</i>) Numidian.	Numidia. 1803. 5. 6. H. ȝ.	Bot. mag. 377.	
12 <i>românum</i> . w. (<i>pi.</i>) Roman.	Rome. 1724. — H. ȝ.	Sweet ger. t. 6.	
13 <i>caucalisæfolium</i> . S.G. (<i>pi.</i>) Caucalis-leaved.	S. Europe. 1816. 6. 8. H. ȝ.	Eng. bot. 902.	
14 <i>moschatum</i> . DC. (<i>li.</i>) musky.	England.	Cav. dis. 4. t. 90. f. 2.	
15 <i>bôtrys</i> . DC. (<i>bl.</i>) large-blue.	Italy. 1819. — H. ȝ.	—— 5. t. 126. f. 1.	
16 <i>mûrcicum</i> . DC. (<i>bl.</i>) small-blue.	Murcia. — — H. ȝ.	—— 4. t. 88. f. 2.	
17 <i>gruinum</i> . DC. (<i>bl.</i>) broad-leaved.	Crete. 1596. 6. 7. H. ȝ.	Stev.m.ac.P.3.t.15.f.2.	
18 <i>serotinum</i> . DC. (<i>bl.</i>) late-flowering.	Tyre. 1816. 6. 9. H. ȝ.		
			<i>multicaulé</i> . S.G. t. 137.
19 <i>chiuum</i> . DC. (<i>bl.</i>) Levant.	Levant. 1724. 6. 7. H. ȝ.	Cav. dis. 4. t. 92. f. 1.	
20 <i>hymenædes</i> . DC. (<i>li.</i>) three-leaved.	Barbary. 1789. 1. 12. H. ȝ.	Sweet ger. 1. t. 23.	
21 <i>Gussoniæ</i> . S.G. (<i>li.</i>) Gussone's.	Naples. 1822. 5. 8. H. ȝ.	—— 2. t. 200.	
22 <i>malachoides</i> . DC. (<i>pu.</i>) Mallow-leaved.	S. Europe. 1596. 5. 7. H. ȝ.	Cav. dis. 4. t. 91. f. 1.	
23 <i>ribifolium</i> . w. (<i>pu.</i>) Currant-leaved.	C. B. S.	Jacq. ic. rar. 3. t. 509.	
24 <i>incarnatum</i> . DC. (<i>fl.</i>) flesh-coloured.	1787. 5. 7. G. ḥ.	Sweet ger. 1. t. 94.	
25 <i>glaucophyllosum</i> . DC. (<i>li.</i>) glaucous-leav'd.	Egypt. 1732. 7. 8. H. ȝ.	—— 3. t. 283.	
26 <i>malopoides</i> . DC. (<i>li.</i>) velvetty-leaved.	Sicily. 1818. 5. 8. H. ȝ.	Cav. dis. 4. t. 90. f. 1.	
27 <i>littoreum</i> . DC. (<i>li.</i>) smooth.	Narbonne. — — H. ȝ.		
28 <i>maritimum</i> . DC. (<i>li.</i>) sea-side.	England.	Eng. bot. 646.	
29 <i>Reichardii</i> . DC. (<i>wh.</i>) dwarf.	Minorca. 1783. 4. 9. H. ȝ.	Bot. mag. 18.	
			<i>chamædryoides</i> , w. <i>Geranium Reichardii</i> , B.M.

Tribus II. *GRIELEÆ. S.G. p. VII.*

GRI'ELUM. W.	GRI'ELUM. Decandria Pentagynia.
1 <i>tenuifolium</i> . w. (<i>ye.</i>) large-flowered.	C. B. S. 1790. 4. 6. G. ḥ.
2 <i>laciniatum</i> . g. (<i>ye.</i>) jagged-leaved.	—— 1825. — G. ḥ.

Tribus III. *PELARGONIEÆ. S.G. p. VIII.*

PHYMATA'NTHUS. S.G. WART-FLOWER.	Monadelphia Pentandria.
1 <i>tricolor</i> . s.g. (<i>w.pu.</i>) three-coloured.	C. B. S. 1791. 6. 10. G. ḥ.
<i>Pelargónium tricolor</i> . B.M. 240.	
2 <i>elatius</i> . s.g. (<i>w.pu.</i>) tall.	—— 1795. — G. ḥ.

Sweet ger. 1. t. 43.
Sweet ger. 1. t. 96.

3	villòsus.	(w.pu.) villous.	Hybrid.	1823.	6. 10.	G. $\frac{1}{2}$.	
4	grandiflòrus.	(w.pu.) large-flowered.	_____	_____	_____	G. $\frac{1}{2}$.	
5	latifolius. c.c.	(w.pu.) broad-leaved.	_____	_____	_____	G. $\frac{1}{2}$.	
CAMP'YLIA.	S.G.	CAMP'YLIA.	Monadelphia Pentandria.				
1	carinàta. s.g.	(w.pu.) keeled-stipuled.	C. B. S.	1810.	4. 9.	G. $\frac{1}{2}$.	Sweet ger. I. t. 21.
2	variegàta. s.g.	(v.) variegated-fl'd.	Hybrid.	1823.	—	G. $\frac{1}{2}$.	3. t. 266.
3	holosericea. s.g.	(da.) silky-leaved.	_____	1820.	—	G. $\frac{1}{2}$.	1. t. 75.
4	blattaria. s.g.	(li.) downy-leaved.	C. B. S.	1790.	6. 9.	G. $\frac{1}{2}$.	1. t. 88.
5	élegans. s.g.	(bh.) elegant.	Hybrid.	1822.	—	G. $\frac{1}{2}$.	3. t. 222.
6	verbasciflora. s.g.	(li.) Mullein-fl'd.	C. B. S.	1811.	—	G. $\frac{1}{2}$.	2. t. 157.
7	càna. s.g.	(li.) hoary-leaved.	_____	1794.	7. 12.	G. $\frac{1}{2}$.	2. t. 114.
8	eristemon. s.g.	(wh.) velvet-leaved.	_____	_____	3. 6.	G. $\frac{1}{2}$.	Jacq. scheen. 2. t. 132.
Pelargònium eristemon.	Jacq.						
9	laciniatà. s.g.	(d.re.) jagged-leaved.	Hybrid.	1827.	4. 10.	G. $\frac{1}{2}$.	Swt. ger. ser. 2. t. 1.
10	coronopifolia.	(li.) Buck's-horn-ld.	C. B. S.	1791.	6. 10.	G. $\frac{1}{2}$.	Jacq. ic. rar. 3. t. 526.
11?	œnothéræ.	(li.) œnothera-like.	_____	1812.	4. 8.	G. $\frac{1}{2}$.	3. t. 525.
Pelargònium œnothéræ.	Jacq.						
OTI'DIA.	S.G.	OTI'DIA.	Monadelphia Pentandria.				
1	carnòsa. s.g.	(wh.) fleshy-stalked.	C. B. S.	1724.	6. 8.	G. $\frac{1}{2}$.	Swt. ger. 1. t. 98.
2	ceratophylla. s.g.	(wh.) horn-leaved.	Africa.	1786.	5. 7.	G. $\frac{1}{2}$.	Bot. mag. 315.
	Pelargònium ceratophyllum.	B.M.					
3	dasycaulón. s.g.	(wh.) thick-stemmed.	Africa.	1795.	7. 12.	G. $\frac{1}{2}$.	Bot. mag. 2029.
4	láxa. s.g.	(wh.) loose-panicled.	C. B. S.	1821.	6. 9.	G. $\frac{1}{2}$.	Sweet ger. 2. t. 196.
5	crithmifolia. s.g.	(wh.) Samphire-ld.	_____	1790.	4. 9.	G. $\frac{1}{2}$.	4. t. 354.
6	alternans. s.g.	(wh.) Parsley-leaved.	_____	1791.	5. 8.	G. $\frac{1}{2}$.	3. t. 286.
GRENVILLEA.	S.G.	GRENVILLEA.	Monadelphia Tetrandria.				
	conspicua. s.g.	conspicuous.	Africa.	1810.	7. 11.	G. $\frac{1}{2}$.	Sweetger. 3. t. 262. f. 2.
Geranium Grenevilleæ.	A.G. c. ic.						
H'OAERA.	S.G.	H'OAERA.	Monadelphia Pentandria.				
1	violàflora. s.g.	(wh.) white violet-fl'd.	C. B. S.	1821.	5. 8.	G. $\frac{1}{2}$.	Sweet ger. 2. t. 123.
2	nívea. s.g.	(wh.) snowy white.	_____	_____	—	G. $\frac{1}{2}$.	2. t. 182.
3	bubonifolia.†	(wh.) Bubon-leaved.	_____	1800.	3. 7.	G. $\frac{1}{2}$.	And. bot. rep. 328.
4	Leeána. s.g.	(wh.) Mr. Lee's.	_____	1823.	6. 8.	G. $\frac{1}{2}$.	Sweet ger. 3. t. 323.
5	laciniatà.†	(v.) jag-leaved.	_____	1800.	5. 6.	G. $\frac{1}{2}$.	And. bot. rep. 131.
6	pennifòrmis.†	(ye.) winged.	_____	_____	—	G. $\frac{1}{2}$.	269.
7	purpurascens.†	(pu.) purplish.	_____	_____	—	G. $\frac{1}{2}$.	204.
8	incrassata.†	(std.) fleshy-leaved.	_____	1801.	6. 7.	G. $\frac{1}{2}$.	Bot. mag. 761.
9	carinatà. s.g.	(ye.) boat-flowered.	_____	1812.	5. 8.	G. $\frac{1}{2}$.	Sweet ger. 2. t. 135.
10	corydaliflora. s.g.	(ye.) Fumitory-fl'd.	_____	_____	—	G. $\frac{1}{2}$.	1. t. 18.
11	rapacea.	(pi.) Turnep-rooted.	_____	1788.	4. 6.	G. $\frac{1}{2}$.	And. rep. 239.
	Geranium Selinum.	A.R.					
12	setosa. s.g.	(bh.) bristle-pointed.	_____	1818.	4. 8.	G. $\frac{1}{2}$.	Sweet ger. 1. t. 38.
13	pilosa.	(pu.) hairy.	_____	1801.	5. 7.	G. $\frac{1}{2}$.	And. bot. rep. 259.
	Geranium pilosum.	var. I. A.G. ic.					
14	bicolor.	two-coloured.	_____	_____	3. 8.	G. $\frac{1}{2}$.	Andr. geran. c. ic.
	Geranium pilosum.	var. 2. A.G. ic.					
15	blanda.	(bh.) bluish-flowered.	_____	_____	—	G. $\frac{1}{2}$.	Andr. geran. c. ic.
	Geranium pilosum.	var. 3. A.G. ic.					
16	congesta. s.g.	(sp.) crowded-fl'd.	_____	1820.	5. 10.	G. $\frac{1}{2}$.	Sweet ger. 3. t. 202.
17	retusa. s.g.	(da.) notched-petal'd.	Hybrid.	1824.	—	G. $\frac{1}{2}$.	4. t. 307.
18	selinifolia. s.g.	(d.pu.) Milk-parsley-ld.	_____	1821.	3. 8.	G. $\frac{1}{2}$.	2. t. 159.
19	radicata. s.g.	(ye.) fleshy-fringed-ld.	C. B. S.	1802.	5. 8.	G. $\frac{1}{2}$.	1. t. 174.
20	ciliata.	(wh.) ciliated.	_____	1795.	4. 6.	G. $\frac{1}{2}$.	L'Herit. ger. t. 7.
21	ovalifolia. s.g.	(wh.) oval-leaved.	_____	1820.	4. 8.	G. $\frac{1}{2}$.	Sweet ger. 2. t. 106.
22	reticulata. s.g.	(std.) netted-petaled.	_____	_____	—	G. $\frac{1}{2}$.	1. t. 91.
23	nervifolia.	(wh.) nerved-leaved.	_____	_____	5. 8.	G. $\frac{1}{2}$.	Jacq. ic. rar. 3. t. 517.
24	trifylla.	(fl.cr.) three-leaved.	_____	_____	—	G. $\frac{1}{2}$.	3. t. 515.
25	lutea.†	(ye.) small-yellow.	_____	1800.	—	G. $\frac{1}{2}$.	And. bot. rep. t. 123.
26	nummularifolia.	(ye.) money-wort-ld.	_____	1801.	—	G. $\frac{1}{2}$.	Sal. par. lond. t. 23.
27	integrifolia. c.c.	(w.r.) entire-leaved.	Hybrid.	1821.	4. 8.	G. $\frac{1}{2}$.	Sweet ger. 1. t. 72.
28	fusca. c.c.	(br.) brown-flowered.	_____	_____	—	G. $\frac{1}{2}$.	
29	nigricans. c.c.	(bl.) blackish.	_____	1822.	—	G. $\frac{1}{2}$.	
30	mágica. c.c.	(bl.) dark-flowered.	_____	_____	—	G. $\frac{1}{2}$.	
31	átra. s.g.	(da.) dark brown.	C. B. S.	1793.	—	G. $\frac{1}{2}$.	Sweet ger. 1. t. 72.
32	dioica. c.c.	(bl.) dioecious.	_____	1795.	6. 8.	G. $\frac{1}{2}$.	And. bot. rep. 209.
	Geranium melananthum.	A.R.					

33 undulæfödra. s.g. (da.)	waved-petaled.	C. B. S.	1821.	4. 8. G. ȝ.	Sweet ger. 3. t. 263.
34 melanátha. s.g. (bl.)	black-flowered.	_____	1790.	— G. ȝ.	— 1. t. 73.
35 vâria. s.g.	(da.) various-leaved.	Hybrid.	1821.	— G. ȝ.	— 2. t. 166.
36 Colvillii. s.g.	(cr.) Colvill's.	_____	1823.	— G. ȝ.	— 3. t. 260.
37 Jenkinsi. c.c.	(sc.) Jenkinson's.	_____	_____	— G. ȝ.	
38 diluta	(bh.) diluted.	_____	1821.	— G. ȝ.	
39 replicata.	(re.) replicate.	_____	_____	— G. ȝ.	
40 rubescens.	(re.) reddish.	_____	_____	— G. ȝ.	
41 intermixta.	(re.) intermixed.	_____	1822.	— G. ȝ.	
42 capitata.	(bh.) headed.	_____	_____	— G. ȝ.	
43 supérba.	(sc.) superb.	_____	_____	— G. ȝ.	
44 villôsa.	(sc.) villous.	_____	1821.	— G. ȝ.	
45 unguiculâta.	(re.) clawed.	_____	_____	5. 8. G. ȝ.	Sweet ger. 2. t. 151.
46 tenuifolia.	(bh.) slender-leaved.	_____	_____	— G. ȝ.	— 2. t. 132.
47 marginata.	(bh.) margined.	_____	1823.	— G. ȝ.	
48 atrosanguínea. s.g. (cr.)	dark-crimson.	_____	1822.	4. 8. G. ȝ.	
49 élégans. s.g.	(cr.) elegant.	_____	_____	— G. ȝ.	
<i>Pelargónium Sweetianum. DC.</i>					
50 coluteæfölia. s.g. (cr.)	BladderSenna-l'd.	_____	1823.	— G. ȝ.	Sweet ger. 4. t. 311.
51 orobifölia. s.g. (st. pu.)	Orobus-leav'd.	_____	_____	— G. ȝ.	— 4. t. 304.
52 coccinea. s.g.	(sc.) scarlet.	_____	_____	— G. ȝ.	— 4. t. 398.
53 recurviflora.	(sc.) recurv'd-petal'd.	_____	1821.	— G. ȝ.	
54 sanguinolénta.	(cr.) bloody-flower'd.	_____	1823.	— G. ȝ.	
55 pulchella.	(re.) neat.	_____	_____	— G. ȝ.	
56 amœna.	(re.) delightful.	_____	_____	— G. ȝ.	
57 pátent.	(re.) spreading.	_____	1822.	— G. ȝ.	
58 lilacina.	(li.) Lilac-coloured.	_____	_____	— G. ȝ.	
59 venosa. s.g.	(std.) veined-petaled.	_____	1823.	— G. ȝ.	Sweet ger. 3. t. 209.
60 literata.	(std.) lettered.	_____	_____	— G. ȝ.	
<i>labyrinthica. s.g. (std.) labyrinth-fl'd.</i>					
62 rósea. s.g.	(ro.) rose-coloured.	C. B. S.	1792.	— G. ȝ.	Sweet ger. 3. t. 276.
<i>Gerânium róseum. A.R. 173. Pelargónium condensátum. P.S.</i>					
63 hedsarifölia. s.g. (cr.)	Hedysarum-l'd.	Hybrid.	1823.	— G. ȝ.	Sweet ger. 4. t. 355.
64 galegofolia.	(cr.) Galega-leaved.	_____	_____	— G. ȝ.	
65 sisymbriifolia. s.g. (cr.)	Water-rocket-l'd.	_____	_____	— G. ȝ.	Sweet ger. 3. t. 292.
66 hirsuta.	(bh.) various-l'd. hairy.	C. B. S.	1788.	3. 5. G. ȝ.	Cav. dis. 4. t. 101. f. 2.
<i>DIMA'CRIA. S.G. DIMA'CRIA. Monadelphia Pentandria.</i>					
1 pinnâta. s.g.	(bh.) wing-leaved.	C. B. S.	1779.	4. 6. G. ȝ.	Sweet ger. 1. t. 46.
<i>Pelargónium viciæfölium. DC.</i>					
2 bipartita. s.g.	(st.) forked-leaved.	Hybrid.	1821.	4. 8. G. ȝ.	Sweet ger. 2. t. 142.
3 foliolosa. s.g.	(st.) leafy.	C. B. S.	1800.	— G. ȝ.	And. bot. rep. 311.
<i>Gerânium pinnatum. A.R.</i>					
4 astragalifolia. s.g. (st.)	Astragalus-l'd.	_____	1788.	6. 8. G. ȝ.	Sweet ger. 2. t. 103.
5 Smithiana. s.g.	(re.) Mr. E.D. Smith's.	Hybrid.	1823.	4. 8. G. ȝ.	— 4. t. 358.
6 tenellâ.†	(bh.) tender.	_____	1802.	— G. ȝ.	Andr. geran. c. ic.
<i>Gerânium tenellum. A.G.</i>					
7 imbûta.	(bh.) stained-petaled.	Hybrid.	1822.	4. 8. G. ȝ.	
8 fulgens. c.c.	(cr.) shining-scarlet.	_____	1821.	— G. ȝ.	
9 Andréwsi.†	(bh.p.) Andrew's's.	C. B. S.	1802.	— G. ȝ.	Andr. geran. c. ic.
<i>Gerânium heterophyllum. A.G. non Jacq.</i>					
10 élégans. s.g.	(sc.) elegant.	Hybrid.	1822.	— G. ȝ.	Sweet ger. 3. t. 202.
11 coronillæfölia.†	(co.) Coronilla-leav'd.	C. B. S.	1795.	6. 7. G. ȝ.	And. bot. rep. 305.
12 apifolia.†	(bh.) Parsley-leaved.	_____	1802.	4. 8. G. ȝ.	Andr. geran. c. ic.
<i>Gerânium apifölium. A.G. non Jacq.</i>					
13 auriculâta. s.g. (w.pu.)	eared-leaved.	C. B. S.	5. 8. G. ȝ.	Sweet ger. 4. t. 395.
<i>Pelargónium ciliatum. Jacq. ic. 3. t. 519.</i>					
14 sulphurea. s.g.	(st.) sulphur-color'd.	Hybrid.	1821.	— G. ȝ.	Sweet ger. 2. t. 163.
15 cárnea.	(fl.) flesh-coloured.	C. B. S.	1812.	4. 6. G. ȝ.	Jacq. ic. rar. 3. t. 512.
16 floribunda.†	(v.) many-flowered.	_____	1795.	3. 5. G. ȝ.	And. bot. rep. 420.
17 fissifolia.†	(bh.) cloven-leaved.	_____	_____	4. 8. G. ȝ.	— 378.
18 barbata. s.g.	(fl.) bearded.	_____	1790.	5. 8. G. ȝ.	Sweet ger. 4. t. 391.
19 recurvâta. s.g.	(wh.) recurved.	_____	_____	— G. ȝ.	And. rep. 323.
<i>Gerânium barbatum. v. minor. A.R.</i>					
20 aristata. s.g.	(wh.) awned.	_____	_____	— G. ȝ.	Andr. rep. 366.
<i>Gerânium barbatum. v. undulatum. A.R.</i>					
21 setigera. s.g.	(wh.) bristle-bearing.	_____	_____	— G. ȝ.	Andr. bot. rep. 303.
<i>Gerânium barbatum. A.R. nec aliorum.</i>					
22 revoluta.†	(pu.) revolute.	_____	1800.	7. 8. G. ȝ.	Andr. rep. 354.
23 oxalidifolia.†	(st.) Oxalis-leaved.	_____	1801.	5. 8. G. ȝ.	— 300.

GERANIACEÆ.

24 refléxa.†	(wh.) reflex-leaved.	C. B. S.	1800.	6. 7. G. ¼.	Andr. rep. 224.
25 rumicifòlia. s.g. (ye.)	Dock-leaved.	1825.	5. 8. G. ¼.	Sweet ger. 4. t. 318.	
26 depréssa. s.g. (st.)	depressed.	1812.	— G. ¼.	— 3. t. 290.	
27 longiflòra. s.g. (st.)	long-flowered.	—	— G. ¼.	Jacq. ic. rar. 3. t. 521.	
28 longiflòria. s.g. (ro.)	long-leaved.	—	— G. ¼.	— 3. t. 518.	
29 lanceolata. s.g. (bh.)	spear-leaved.	1826.	— G. ¼.	Sweet ger. 4. t. 387.	
30 undulatà.†	(wh.) wave-flowered.	1795.	5. 7. G. ¼.	Andr. rep. 292.	
31 spathulatà.†	(wh.) spatula-leaved.	—	4. 6. G. ¼.	— 152.	
32 affinis.†	(wh.) fring'd-spat.-ld.	—	— G. ¼.	— 282.	
33 virginea.†	(wh.) virgin.	—	5. 7. G. ¼.	— 317.	
34 radiata.†	(ye.) ray-leaved.	1801.	6. 8. G. ¼.	— 222.	
35 linearis.†	(st.) linear-petaled.	1800.	— G. ¼.	— 193.	
36 punctatà.†	(st.) dotted-flowered.	1794.	4. 6. G. ¼.	— 60.	
SEYMO'URIA. S.G.	SEYMO'URIA.	Monadelphia	Pentandria.		
1 asarifòlia. s.g. (da.)	Asarum-leaved.	C. B. S.	1821.	11. 2. G. ¼.	Sweet ger. 3. t. 206.
2 L'Heritiéri. s.g. (pu.)	L'Heritiér's.	—	1795.	4. 5. G. ¼.	L'Herit. ger. t. 43.
<i>Pelargónium dipétalum. L'Her.</i>					
PELARGONIUM. S.G. STORK'S-BILL.	Monadelphia	Heptandria.			
1 amœnum. s.g. (sa.)	delightful.	Hybrid.	1821.	4. 8. G. ¼.	Sweet ger. 2. t. 121
2 dimaciæflòrum. s.g. (sc.)	Dimacia-fld.	—	1822.	— G. ¼.	— 3. t. 220.
3 mündulum. s.g. (sc.)	spruce.	—	1823.	3. 10. G. ¼.	— 3. t. 288.
4 ringens. s.g. (sc.)	ringent-flow'rd.	—	—	— G. ¼.	— 3. t. 256.
5 concavum. s.g. (re.)	concave-petaled.	—	—	— G. ¼.	— 3. t. 237.
6 conclusum. s.g. (d.re.)	shut-petaled.	—	—	4. 10. G. ½.	— 4. t. 305.
7 hoareaeflòrum s.g. (sa.)	Hoarea-flow'r'd.	—	1821.	— G. ¼.	— 2. t. 133.
8 clásum. c.c. (re.)	closed petaled.	—	1822.	— G. ¼.	
9 patens. s.g. (sc.)	spreading-fld.	—	1820.	— G. ½.	
10 cruentum. s.g. (cr.)	blood-red.	—	1822.	— G. ¼.	
11 áulicum. (cr.)	courtly.	—	1823.	5. 10. G. ¼.	
12 Avroniànum. s.g. (sc.)	Avron's.	—	—	— G. ¼.	
13 ligulatum. s.g. (sc.)	ligulate-petaled.	—	1823.	4. 9. G. ½.	Sweet ger. 3. t. 394.
14 puniceum. c.c. (cr.)	puniceous.	—	—	— G. ¼.	— 4. t. 301.
15 pusillum. c.c. (sc.)	dwarf.	—	—	— G. ¼.	
16 pygmæum. c.c. (sc.)	small.	—	—	5. 8. G. ¼.	
17 venulosum. (sc.)	veined-petaled.	—	1821.	— G. ¼.	
18 intertèxtum. s.g. (sc.)	interwoven.	—	1822.	4. 8. G. ¼.	Sweet ger. 2. t. 185.
19 acídum. s.g. (cr.)	sour-leaved.	—	1823.	5. 9. G. ¼.	— 3. t. 261.
20 pallens. s.g. (st.)	cream coloured.	C. B. S.	1800.	— G. ½.	— 2. t. 148.
21 caryophyllaceum. (bh.)	Clove-scented.	Hybrid.	1822.	— G. ½.	— 4. t. 347.
22 paradoxum. (wh.)	paradoxical.	—	—	— G. ½.	
23 multiradiatum. s.g. (da.)	many-rayed.	C. B. S.	1818.	6. 10. G. ½.	Sweet ger. 2. t. 145.
24 rutaceum. s.g. (du.)	Rue-like.	Hybrid.	1823.	5. 10. G. ½.	— 3. t. 279.
25 pedunculatum. s.g. (da.)	long-peduncled.	—	—	— G. ½.	— 4. t. 346.
26 cenanthifòlium. s.g. (d.cr.)	Dropwort-ld.	—	—	6. 10. G. ½.	ser. 2. t. 13.
27 sphondyliifòlium. (d.cr.)	Sphondylium-ld	Hybrid.	1822.	4. 8. G. ¼.	— 3. t. 246.
28 chærophyllum. s.g. (cr.)	Cow-Parsley-ld	—	—	— G. ½.	— 3. t. 257.
29 nítidum. s.g. (sc.)	glossy.	—	—	— G. ½.	— 3. t. 298.
30 sanguineum. s.g. (cr.)	crimson.	C. B. S?	1819.	— G. ½.	— 1. t. 76.
31 árdens. s.g. (sc.)	glowing.	Hybrid.	1810.	3. 8. G. ½.	— 1. t. 45.
32 confertifòlium. (sc.)	close-leaved.	—	—	— G. ½.	— 3. t. 297.
<i>Gerânium árdens minor. A.G.</i>					
33 pulchellum. B.M. (v.)	nonesuch.	C. B. S.	1795.	3. 5. G. ½.	Sweet ger. 1. t. 31.
34 jonquillinum. s.g. (pi.)	Jonquil-scent'd.	Hybrid.	1822.	5. 10. G. ½.	— 3. t. 241.
35 sélectum. s.g. (d.pu.)	select.	—	—	— G. ¼.	— 2. t. 190.
36 imbricatum. s.g. (v.)	imbricate-pet'd.	C. B. S?	1800.	5. 8. G. ½.	— 1. t. 65.
37 bicolor. s.g. (da.)	two-coloured.	1778.	7. 8. G. ½.	Bot. mag. 201.
α átrum. s.g. (da.)	dark-edged.	—	— G. ½.	Sweet ger. 1. t. 97. f. 1.
β pállidum. s.g. (pu.)	pale-edged.	—	— G. ½.	— 2. t. 161.
38 quinquevúlerum. (da.)	dark-flower'd.	Hybrid.	1796.	5. 10. G. ½.	— 1. t. 97. f. 2.
39 blandfordianum. (bh.)	hoary-leaved.	—	1812.	1. 12. G. ½.	— 2. t. 101.
40 millefoliatum. s.g. (v.)	Milfoil-leaved.	C. B. S.	1819.	6. 10. G. ¼.	— 3. t. 230.
41 flavum. DC.	(st.) pale yellow.	—	1724.	7. 9. G. ¼.	— 3. t. 254.
42 scabridum. c.c. (br.)	rough-edged.	—	1822.	— G. ¼.	
43 triste. DC.	(da.) night-smelling.	—	1632.	5. 10. G. ¼.	Delaun. h. am. t. 27.
44 filipendulifòlium. (br.)	Dropwort-ld.	—	1812.	— G. ¼.	Sweet ger. 1. t. 85.
45 pentastictum.	(bh.) five-marked.	Hybrid.	1824.	— G. ¼.	— ser. 2. t. 25.
46 heracleifòlium. s.g. (br.)	Héraclum-ld.	C. B. S.	1820.	— G. ¼.	— 3. t. 211.
47 lobatum. DC.	(da.) Cow-parsnep-ld.	—	1710.	— G. ¼.	— 1. t. 51.
48 salebròsum. s.g. (d.pu.)	rugged-leaved.	Hybrid.	1824.	— G. ¼.	— 4. t. 309.

49 Lawranceanum.s.g.(d.pu.)	Mrs. Kearse's.	1824.	5. 10. G. ȝ.	Swt. ger. ser. 2. t. 23.
50 orbiculatum.	(da.) orbicular-ld. Hybrid	1823.	— G. ȝ.	
51 pulveruléntum. s.g.(sp.)	powdered-ld C. B. S.	1822.	— G. ȝ.	Sweet ger. 3. t. 218.
52 pedicellatum.s.g.(v.)	long-pedicled.	—	— G. ȝ.	— 3. t. 250.
53 lúridum. s.g.	(co.) lurid.	1811.	7. 12. G. ȝ.	— 3. t. 281.
<i>Geranium lúridum. A.G. ic.</i>				
54 rapæfólium.	(da) Turnep-leaved. Hybrid.	1821.	5. 10. G. ȝ.	Sweet ger. 2. t. 179.
55 glaucifólium.s.g.(da.)	Glaucium-ld.	—	— G. ȝ.	— 1. t. 61.
56 gibbósum. d.c.	(st.) knotted-stalk'd C. B. S.	1712.	— G. ȝ.	— 3. t. 213.
57 mutábilis. s.g.	(bh.) changeable-cold Hybrid.	1822.	7. 10. G. ȝ.	— 3. t. 239.
58 vespertinum. s.g.(pu.)	evening-scent'd	—	8. 12. G. ȝ.	
59 sinapifólium.	(pi.) Mustard-leav'd. Hybrid.	—	6. 10. G. ȝ.	
60 brassicáturn.	(st.) Cabbage-leav'd.	—	— G. ȝ.	
61 adscéndens.	(pa.) ascending.	—	— G. ȝ.	
62 circumfléxum.	(pa.) bending.	—	— G. ȝ.	
63 campyliaflórum. s.g.(li.)	Campylia-fld	1823.	— G. ȝ.	Sweet ger. 3. t. 251.
64 renifórmē. d.c.	(pu.) kidney-leaved. C. B. S.	1791.	1. 12. G. ȝ.	— 1. t. 48.
65 flexuósom. s.g.	(sc.) bent-stalked. Hybrid.	1821.	3. 10. G. ȝ.	— 2. t. 180.
66 sæpeflórens. s.g.(pi.)	frequent-flow'rg	1801.	— G. ȝ.	— 1. t. 58.
67 cómptum. s.g.	(pi.) decked.	1821.	— G. ȝ.	— 3. t. 255.
68 echinátum. d.c.(wh.)	prickly-stalked. C. B. S.	1789.	6. 10. G. ȝ.	— 1. t. 54.
69 armátum. s.g.(d.pu.)	long-spined.	—	3. 6. G. ȝ.	— 3. t. 214.
70 intermédianum.c.c.(li.)	intermediate. Hybrid.	1821.	6. 10. G. ȝ.	
71 particéps. s.g.	(li.) participant.	1806.	5. 10. G. ȝ.	Sweet ger. 1. t. 49.
72 cortusafólium. d.c.(wh.)	Cortusa-leav'd Africa.	1786.	6. 10. G. ȝ.	— 1. t. 14.
73 eréctum. s.g.	(pu.) upright. Hybrid.	1822.	— G. ȝ.	— 2. t. 187.
74 Stapletoni. s.g.(pu.)	Miss Stapleton's.	—	— G. ȝ.	— 2. t. 212.
75 crassicáule. s.g.(wh.)	Cowslip-scent'd. Africa.	1786.	— G. ȝ.	L'Herit. ger. t. 26.
a ábum. s.g.	(wh.) white-flowered.	—	— G. ȝ.	Sweetger. 2.t.192.f.a.
β maculátum.s.g.(sp.)	spotted-flowered	—	— G. ȝ.	— 2.t.192.f.b.
76 Victoriána. s.g.(sc.)	Princess Victoria's. Hybrid.	1826.	— G. ȝ.	— ser. 2. t. 7.
77 anómalam. s.g.(pi.)	anomalous.	1822.	4. 8. G. ȝ.	— 4. t. 315.
78 tenuifólium. d.c.(pi.)	slender-leaved. C. B. S.	1768.	5. 7. G. ȝ.	L'Herit. ger. t. 12.
79 hírtum. d.c.	(pi.) hairy.	—	3. 9. G. ȝ.	Sweet ger. 2. t. 113.
80 anthriscifólium.s.g.(sc.)	Anthriscus-ld. Hybrid.	1823.	7. 9. G. ȝ.	— 3. t. 233.
81 magnistipulátum.(sc.)	large-stipuled.	1825.	5. 10. G. ȝ.	— 4. t. 313.
82 fulgídum. d.c.	(sc.) Celandine-ld. C. B. S.	1723.	3. 10. G. ȝ.	— 1. t. 69.
83 aurantiacum.s.g.(or)	orange-colour'd. Hybrid.	1821.	4. 8. G. ȝ.	— 2. t. 198.
α undulátum.s.g.(or)	wave-leaved.	—	— G. ȝ.	— 2.t.198.f.a.
β planifólium.s.g.(or)	flat-leaved.	—	— G. ȝ.	— 2.t.198.f.b.
84 ardésrens. s.g.	(cr.) burnished.	1822.	— G. ȝ.	— 3. t. 231.
85 torrefáctum.s.g.(da.)	burnt-petaled.	—	— G. ȝ.	— 3. t. 243.
86 Pottéri. s.g.	(st.) Potter's.	—	6. 9. G. ȝ.	— 2. t. 147.
87 luculéntum.	(sc.) bright-flowered.	—	— G. ȝ.	
88 diléctum.	(sc.da.) chosen.	—	4. 9. G. ȝ.	
89 Loudoniánum.s.g.(d.sc.)	Mr. Loudon's.	1827.	5. 9. G. ȝ.	Sweet ger. ser. 2. t. 17.
90 eratinum. s.g.	(sc.) lovely.	—	— G. ȝ.	— ser. 2. t. 30.
91 nànum. s.g.	(sc.) pigmy.	1820.	— G. ȝ.	— 2. t. 102.
92 incúrvum. s.g.	(sc.) incurved-petal'd	1821.	— G. ȝ.	— 3. t. 249.
93 ignéscens. s.g.	(sc.) fiery-flowered.	1812.	— G. ȝ.	— 1. t. 2. & 55.
a mähr.	(sc.) large-flowered.	1818.	— G. ȝ.	— 1. t. 2.
β coccíneum.	(sc.) scarlet-flowered.	1822.	— G. ȝ.	
γ spléndens.	(sc.) splendid.	—	— G. ȝ.	
δ supérbum.	(sc.) superb.	—	— G. ȝ.	
ε stérile. s.g.	(sc.) barren-anthered.	1812.	— G. ȝ.	Sweet ger. 1. t. 55.
94 laxiflórum. s.g.	(sc.) spreading-umb'd.	1822.	— G. ȝ.	— 3. t. 216.
95 concrétem. s.g.	(sc.) compounded.	1827.	4. 9. G. ȝ.	— ser. 2. t. 24.
96 Leghkéckæ. s.g.(sc.)	Mrs. Legh Keck's	1826.	— G. ȝ.	— 4. t. 377.
97 Charlwoòdi. s.g.(d.sc.)	Charlwood's.	—	— G. ȝ.	— 4. t. 380.
98 incomparabile. (d.sc.)	incomparable.	1822.	— G. ȝ.	
99 eriocáulon. s.g. (sc.)	woolly-stalked.	1824.	— G. ȝ.	Sweet ger. 4. t. 357.
100 quadriflórum. s.g.(or.)	four-flowered.	—	— G. ȝ.	— 4. t. 321.
101 chelidoniítiflórum. s.g.(or.)	Chelidonium-ld	1825.	— G. ȝ.	— 4. t. 341.
102 chenopodiiflórum. s.g.(sc.)	Goosefoot-ld	—	— G. ȝ.	— 4. t. 328.
103 latídatum.s.g.(sc.)	broad-toothed.	1827.	— G. ȝ.	— ser. 2. t. 28.
104 variiflórum. s.g. (sc.)	various-leaved.	1823.	— G. ȝ.	— 3. t. 280.
105 diversilóbum.s.g.(re.)	various-lobed.	1825.	— G. ȝ.	— 4. t. 361.
106 Southcoteánum.s.g.(sc.)	Miss Southcote's	1826.	— G. ȝ.	— 4. t. 348.
107 insígnitum.s.g.(d.sc.)	marked-flower'd.	1823.	— G. ȝ.	— 3. t. 300.

GERANIACEÆ.

108 plectophyllosum. s.g. (d.sc.)	plaited-leaved.	Hybrid.	1826.	4. 9. G. h.	Sweet ger. ser. 2. t. 41.
109 floccosum. s.g. (cr.)	nappy.		1821.	— G. h.	— 2. t. 129.
110 rubicundum. (re.)	rubicund.		1822.	— G. h.	
111 hæmæthon.	(cr.)	blood-coloured.	1823.	— G. h.	
112 multiflorum. (sc.)	many-flowered.		—	— G. h.	
113 exile.	(sc.)	thin-branched.	1822.	— G. h.	
114 gracilescens.	(sc.)	slender-stalked.	—	— G. h.	
115 splendidum. (cr.)	splendid.		—	— G. h.	
116 consimile.	(sc.)	likened.	1822.	— G. h.	
117 Broughtoniæ.s.t. (sc.)	Lady Broughton's		—	— G. h.	Sweet ger. 2. t. 181.
118 Bakerianum.s.g. (sc.)	Mrs. Jenkinson's		—	— G. h.	— 3. t. 240.
119 áltum. s.g. (re.)	tall upright.		1827.	— G. h.	— ser. 2. t. 26.
120 Debúrghæ. s.g. (sc.)	Mrs. De Burgh's		—	— G. h.	— ser. 2. t. 21.
121 Cummingsæ.s.g. (re.)	Ly Gordon-Cumming's		—	— G. h.	— ser. 2. t. 35.
122 rubescens. s.g. (re.)	Lady Liverpool's		1819.	— G. h.	— 1. t. 30.
123 Boscawenæ.c.c. (re.)	Mrs. Boscawen's.		—	— G. h.	
124 pavoninum. s.g. (re.)	peacock-spott'd		—	— G. h.	
125 carnescens. s.g. (sa.)	pale red.		1822.	— G. h.	Sweet ger. 1. t. 40.
126 ramigerum. s.g. (sc.)	branching-veined		1826.	1. 12. G. h.	— 4. t. 388.
127 flaccidum. s.g. (d.re.)	flaccid-petal'd.		—	— 4. 9. G. h.	— 4. t. 352.
128 Smithii. s.g. (re.)	Smith's.		1819.	— G. h.	— 4. t. 337.
129 spectabilis. s.g. (sc.)	showy.		1821.	— G. h.	— 2. t. 110.
α maculatum. s.g. (sc.)	spot-flowered.		—	— G. h.	— 2. t. 136.
β atrorubens. s.g. (d.re.)	dark-red.		—	— G. h.	— 2. t. 136. f. 1.
γ striatum. s.g. (sc.)	streak-flowered.		—	— G. h.	
δ recurvum. s.g. (sc.)	recurved-petaled.		—	— G. h.	
ε purpurascens. (pu.)	purplish.		—	— G. h.	
130 Tibbitsianum. s.g. (sc.)	Mr. Tibbits'.		—	— G. h.	
131 flágans. s.g. (sc.)	burning.		1827.	— G. h.	Sweet ger. 2. t. 158.
132 malachræfölium. s.g. (re.)	Malachra-ld.		—	— G. h.	— ser. 2. t. 16.
133 magnifölium. s.g. (re.)	very large-ld.		—	— G. h.	— ser. 2. t. 2.
134 Lacóniæ. s.g. (d.sc.)	Lady Lacon's.		—	— G. h.	— ser. 2. t. 10.
135 Francisiæ. s.g. (re.)	Mr. Francis's.		1826.	— G. h.	— 4. t. 374.
136 cartilagineum. s.g. (re.)	horny-toothed.		1824.	5. 10. G. h.	— 4. t. 349.
137 dilütum. s.g. (re.)	stained-petal'd.		1824.	— G. h.	— 4. t. 382.
138 basilicum. s.g. (sc.)	princely.		1823.	5. 9. G. h.	— 3. t. 293.
139 Huemel. (d.re.)	Sir A. Hume's.		1824.	— G. h.	— 4. t. 360.
140 eriophorum. (sc.)	wool-bearing.		—	— 4. 10. G. h.	— 4. t. 324.
141 irregulare.	(sc.)	irregular-10.	1823.	— G. h.	
142 flamméum. c.c. (sc.)	flame-coloured.		1822.	— G. h.	
143 affine. s.g. (sc.)	related.		—	— G. h.	
144 Colvillii. s.g. (sc.)	Colvill's.		1820.	— G. h.	Sweet ger. 3. t. 277.
145 Vallèti. c.c. (sc.)	Vallet's.		1822.	— G. h.	— 1. t. 86.
146 heteromálum. s.g. (re.)	soft-leaved.		1823.	— G. h.	Sweet ger. sér. 2. t. 32.
147 hibiscifölium. (re.)	Hibiscus-ld.		—	— G. h.	
148 atrorubens. c.c. (re.)	dark-red.		1822.	— G. h.	
149 Allènii. s.g. (d.re.)	Allen's.		1823.	— G. h.	Sweet ger. 3. t. 229.
150 melanchòlicum. (d.re.)	melançholty.		1827.	— G. h.	
151 Daveyanum. s.g. (d.re.)	Davey's.		1819.	— G. h.	Sweet ger. 1. t. 32.
152 phœnicum. s.g. (d.re.)	reddish purple.		1824.	4. 12. G. h.	— 3. t. 207.
153 heterotrichum. s.g. (d.re.)	various-haired.		1823.	4. 10. G. h.	— 4. t. 303.
154 Burnettianum. s.g. (d.re.)	Miss Burnett's.		1826.	— G. h.	— 4. t. 369.
155 imperiale. s.g. (d.re.)	imperial.		—	— G. h.	— 4. t. 365.
156 saturatum. s.g. (d.re.)	saturated.		1827.	— G. h.	— sér. 2. t. 37.
157 régium. s.g. (sc.)	kingly.		1826.	— G. h.	— 4. t. 368.
158 Hesilrigæa. s.g. (pu.)	Lady Hesilrig'e's.		1824.	— G. h.	— 4. t. 319.
159 acetabulös: m. s.g. (sa.)	saucer-leaved.		1827.	— G. h.	— ser. 2. t. 44.
160 Knipeæ. s.g. (li.)	Mrs. Knipe's		1826.	— G. h.	— 4. t. 372.
161 calycinum. s.g. (li.)	large-calyxed.		1815.	— G. h.	— 1. t. 81.
α vèrum. s.g. (li.)	true.		—	— G. h.	— 1. t. 81. f. a.
β maculatum. s.g. (li.)	large-spotted.		1820.	— G. h.	— 1. t. 81. f. b.
γ angulatum. s.g. (li.)	angular-leaved.		—	— G. h.	— 1. t. 81. f. c.
162 planiflòrum. c.c. (li.)	flat-flowered.		—	— G. h.	
163 Dennisianum. s.g. (pu.)	Dennis's.		1819.	— G. h.	Sweet ger. 1. t. 20.
164 Barringtonii. d.c. (pu.)	Barrington's.	C. B. S.	—	— G. h.	
165 elatum. c.c. (pu.)	tall purple.	Hybrid.	1812.	— G. h.	
166 Tormánni. L.en. (pu.)	Torman's.		1821.	— G. h.	
167 Chandléri. c.c. (pu.)	Chandler's.		1814.	— G. h.	
168 crenulatum. s.g. (pu.)	crenulate-leav'd.		1821.	— G. h.	Sweet ger. 2. t. 162.

169	megalostictum. s.g. (d. pu.)	large-marked.	Hybrid.	1824.	4. 10. G. h.	Sweet ger. ser. 2. t. 8.
170	poculifolium. s.g. (pu.)	cup-leaved.		1826.	— G. h.	— ser. 2. t. 31.
171	Foljämbeæ. s.g. (r. pu.)	Mrs. Foljambe's		1825.	— G. h.	— 4. t. 312.
172	Robinsoni. s.g. (bh.)	Robinson's.		1822.	— G. h.	— 2. t. 150.
173	Most'yæ. s.g. (pu.)	Mrs. Mostyn's.		1816.	— G. h.	— 1. t. 10.
174	cucullatum. DC. (pu.)	hooded-leaved.	C. B. S.	1690.	— G. h.	Cav. dis. 4. t. 106. f. 1.
β	stratiiflorum. (pu.)	Prince Regent.		1810.	— G. h.	
γ	majör. c.c. (pu.)	Royal George.		1812.	— G. h.	
δ	grandiflorum. c.c. (pu.)	great-flowered.		1818.	— G. h.	
ε	parvifolium. c.c. (pu.)	smaller-leaved.		—	— G. h.	
175	cochlearium. W.en. (wh.)	concave-ld.		— G. h.	Wendl. col. 2. t. 64.
	concau'lium. Wendl.					
176	rübens. W.en. (re.)	red-flowered.		— G. h.	Wendl. col. 2. t. 54.
177	robustum. c.c. (pu.)	robust.	Hybrid.	1821.	— G. h.	
178	incarnatum. s.g. (fl.)	pale flesh-cold.		—	— G. h.	Sweet ger. 4. t. 308.
179	lasiocaulon. s.g. (bh.)	villous-stalked.		1826.	— G. h.	— 4. t. 364.
180	veniferum. s.g. (std.)	veined-petal'd.		1825.	— G. h.	— 4. t. 322.
181	foliosum. s.g. (r. pu.)	leafy.		1826.	— G. h.	— 4. t. 340.
182	anisodonton. s.g. (pu.)	unequal-toothed.		1825.	— G. h.	— 4. t. 399.
183	malacophyl'lium. s.g. (pu.)	soft-leaved.		—	— G. h.	— 4. t. 397.
184	angulosum. DC. (pu.)	Marshmallow-l.	C. B. S.	1724.	7. 9. G. h.	Cav. dis. 4. t. 112. f. 2.
185	cardiifolium. s.g. (pu.)	Cockle-shell-ld.		1816.	4. 9. G. h.	Sweet ger. 1. t. 15.
186	fasciculatum. c.c. (pu.)	Banbury.	Hybrid.	1803.	5. 10. G. h.	Andr. geran. c. ic.
	Gerânium speciosum var. flore purpureo. A.G.					
187	rugosum. c.c. (fl.)	wrinkled-leav'd.	C. B. S.	— G. h.	Andr. geran. c. ic.
188	speciosum. DC. (wh.)	specious.		1794.	— G. h.	
189	formosum. s.g. (li.)	variegated-fl'd.	Hybrid.	— G. h.	Sweet ger. 2. t. 120.
190	solubile. s.g. (pu.)	dissolvable-col'd.		1818.	— G. h.	— 1. t. 24.
191	multinervé. s.g. (pu.)	many-nerved.		—	— G. h.	— 1. t. 17.
192	tyrianthinum. s.g. (pu.)	royal purple.		1820.	— G. h.	— 2. t. 183.
193	albinotatum. s.g. (pu.)	white-marked.		1824.	— G. h.	— 4. t. 359.
194	'æmulum. s.g. (pu.)	rival.		—	— G. h.	— 2. t. 160.
195	accèdens. c.c. (pu.)	approached.		—	— G. h.	
196	villousum. s.g. (pu.)	villous.		—	— G. h.	Sweet ger. 1. t. 160.
197	floridum. s.g. (li.)	abundant-fl'd.		1816.	6. 10. G. h.	— 1. t. 41.
198	pulchrum. s.g. (pu.)	gay.		1820.	5. 10. G. h.	— 2. t. 107.
199	Watsoni. s.g. (pu.)	Watson's.		1812.	7. 11. G. h.	— 2. t. 130.
200	fusciflorum. s.g. (pu.)	brown-marked.		1822.	4. 10. G. h.	— 3. t. 210.
	fuscatum. s.g. errata.					
201	coarctatum. s.g. (pu.)	close-leaved.		1814.	6. 10. G. h.	— 1. t. 70.
202	paudentatum. s.g. (pu.)	distant-tooth'd.		1821.	5. 10. G. h.	— 2. t. 186.
203	nervosum. s.g. (pu.)	prominent-nerv'd.		1818.	— G. h.	— 1. t. 47.
204	exornatum. (pu.)	adorned.		1823.	— G. h.	— 4. t. 381.
205	Youngii. s.g. (wh.)	Young's.		1820.	— G. h.	— 2. t. 131.
206	Jenkinsi. s.g. (d. bh.)	Mr. Jenkinson's.		—	— G. h.	— 2. t. 154.
207	nùbilum. s.g. (d. bh.)	clouded.		1827.	— G. h.	— ser. 2. t. 20.
208	Brównii. s.g. (d. bh.)	Brown's.		—	— G. h.	— 2. t. 146.
209	diffòrme. s.g. (bh.)	various-leaved.		—	— G. h.	— 2. t. 105.
210	Obrériànum. s.g. (bh.)	Miss O'Brien's.		—	— G. h.	— ser. 2. t. 11.
211	melanostictum. s.g. (li.)	dark-spotted.		—	— G. h.	— ser. 2. t. 6.
212	vestiflùm. s.g. (wh.)	clothed.		—	— G. h.	— ser. 2. t. 27.
213	pùrum. s.g. (wh.)	pure white.		1824.	— G. h.	— 4. t. 334.
214	scitulum. s.g. (wh.)	neat white.		1826.	— G. h.	— 4. t. 390.
215	involucratum. s.g. (va.)	large-bracted.		— G. h.	
α	máximum. s.g. (w.)	largest-flowered.		1818.	— G. h.	Sweet ger. 1. t. 33.
β	álbidum. s.g. (wh.)	white-flowered.		1817.	— G. h.	
γ	incarnatum. s.g. (fl.)	commander in chief.		1815.	— G. h.	
δ	intermédia. s.g. (bh.)	high-admiral.		1817.	— G. h.	
ε	lilacinum. s.g. (li.)	Lilac-coloured.		—	— G. h.	
ζ	róseum. s.g. (ro.)	rosy-coloured.		—	— G. h.	
η	coriaceum. c.c. (w.)	leathery.		1820.	— G. h.	
ς	maculatum. c.c. (wh.)	coronation.		—	— G. h.	
216	polytrichum. s.g. (w.)	many-haired.		1823.	— G. h.	Sweet ger. 3. t. 274.
217	Baileyànum. s.g. (w.)	Bailey's.		1819.	— G. h.	— 1. t. 87.
218	Josephinæ. Lee. (wh.)	Josephine's.		1822.	— G. h.	Sweet ger. 3. t. 215.
219	formosissimum. s.g. (w.)	superb-white.	C. B. S.	1794.	— G. h.	— 3. t. 215. f. a.
α	ábum. s.g. (wh.)	white-flowered.		—	— G. h.	— 3. t. 215. f. b.
β	lineatum. s.g. (wh.)	lined-flowered.		—	— G. h.	— 3. t. 215. f. b.
220	Bóyleæ. s.g. (wh.)	Countess of Cork's.	Hybrid.	1818.	— G. h.	— 1. t. 50.

GERANIACEÆ.

221 pannifolium. s.g. (w.) cloth-leaved.	Hybrid.	1818. 5. 10. G. h.	Sweet ger. 1. t. 9.
222 biflorum. s.g. (wh.) two-flowered.	_____	1822. _____ G. h.	3. t. 287.
223 cándidum. s.g. (wh.) fair-flowered.	_____	1818. 4. 9. G. h.	2. t. 128.
224 blándum. s.g. (bh.) blush-flowered.	_____	1816. _____ G. h.	1. t. 4.
225 aceroídes. s.g. (fl.) Acer-leaved.	_____	1822. _____ G. h.	3. t. 242.
226 Gardnériæ. c.c. (wh.) Lady Gardner's.	_____	1817. _____ G. h.	
227 ramulòsum. s.g. (d.w.) small-branch'd.	_____	1820. _____ G. h.	Sweet ger. 2. t. 177.
228 maculátum. (d.bh.) spot-flowered. C. B. S.	_____	1805. _____ G. h.	Andr. ger. c. ic.
229 Hammerslèiæ. s.g. (b.) Mrs. Hammersley's. Hybrid.	_____	1822. _____ G. h.	Sweet ger. 3. t. 225.
230 pulchérrimum. s.g. (pu.) beautiful.	_____	1819. 5. 10. G. h.	2. t. 134.
231 Rollisòni. s.g. (d.ro.) Rollison's.	_____	1826. _____ G. h.	4. t. 371.
232 eriosépalon. s.g. (bh.) woolly-calyxed.	_____	_____ _____ G. h.	4. t. 375.
233 veniflòrum. s.g. (std.) veined-flowered.	_____	1822. _____ G. h.	3. t. 258.
234 sidæfólium. W.en. (li.) Sida-leaved.	_____	_____ 4. 8. G. h.	
235 acerifólium. DC. (li.) Maple-leaved. C. B. S.	_____	1784. 4. 6. G. h.	L'Herit. ger. t. 21.
236 cuneáutm. DC. (pu.) wedge-leaved. Hybrid.	_____	_____ 5. 8. G. h.	
237 Balbisianum. DC. (pu.) Balbis's.	_____	_____ _____ G. h.	
238 gloriósüm. L.en. (pu.) renowned.	_____	_____ _____ G. h.	
239 Beaufortiánum. s.g. (li.) Beaufort's. C. B. S.?	_____	_____ _____ G. h.	Sweet ger. 2. t. 138.
240 principissæ. s.g. (pu.) P's. Charlotte's.	_____	1810. 5. 10. G. h.	2. t. 139.
241 purpùreum. (pu.) purple-flowered.	_____	1800. _____ G. h.	Andr. geran. c. ic.
242 versicolor. s.g. (pu.) various-colour'd Hybrid.	_____	1819. _____ G. h.	Sweet ger. 1. t. 78.
243 pectinifólium. s.g. (li.) Scallop-shell-l'd.	_____	1820. _____ G. h.	1. t. 66.
244 Thynnèiæ. s.g. (pu.) Lady Bath's.	_____	1815. _____ G. h.	1. t. 74.
245 obtusifólium. s.g. (li.) blunt-leaved.	_____	_____ _____ G. h.	1. t. 25.
246 Beadoniæ. s.g. (pu.) Mrs. Beadon's.	_____	_____ _____ G. h.	2. t. 191.
247 adulterinum. DC. (pu.) hoary-trifid-l'd.	_____	1785. 4. 6. G. h.	1. t. 22.
248 dilatilòbium. s.g. (bh.) dilated-lobed.	_____	1826. _____ G. h.	4. t. 378.
249 gráficum. s.g. (pu.) written-petal'd.	_____	_____ _____ G. h.	ser. 2. t. 12.
250 obtusilòbium. s.g. (pu.) blunt-lobed.	_____	1806. 5. 10. G. h.	1. t. 8.
251 Comptoniæ. s.g. (li.) Ly. Northampton's	_____	1818. _____ G. h.	2. t. 122.
252 afflèus. s.g. (li.) numerous-fl'd.	_____	1821. _____ G. h.	2. t. 194.
253 Saundérsii. s.g. (pu.) Saunders's.	_____	_____ _____ G. h.	3. t. 205.
254 atropurpureum. s.g. (pu.) dark purple.	_____	1822. _____ G. h.	2. t. 152.
255 coruscans. s.g. (d.re.) glittering.	_____	1821. _____ G. h.	2. t. 173.
256 concínnum. s.g. (re.) comely.	_____	1819. _____ G. h.	2. t. 108.
257 Seymouriæ. s.g. (pu.) Mrs. Seymour's.	_____	_____ _____ G. h.	1. t. 37.
258 Lousadiánum. s.g. (bh.) Miss Lousada's.	_____	_____ _____ G. h.	1. t. 44.
259 lépidum. s.g. (li.) pretty.	_____	1822. _____ G. h.	2. t. 156.
260 inscriptum. s.g. (std.) scribbled.	_____	_____ _____ G. h.	2. t. 193.
261 labyrinthicum. (std.) labyrinth-mark'd.	_____	_____ _____ G. h.	
262 planifólium. s.g. (li.d.) flat-leaved.	_____	1823. _____ G. h.	Sweet ger. 3. t. 219.
263 incanescens. s.g. (pu.) whitish-leaved.	_____	_____ 6. 12. G. h.	3. t. 203.
264 notátum. s.g. (pu.) marked-petal'd.	_____	1821. 5. 10. G. h.	3. t. 208.
265 undulæflòrum. s.g. (bh.) waved-flower'd	_____	1824. _____ G. h.	4. t. 329.
266 lineáutm. s.g. (std.) striped-flower'd.	_____	1810. _____ G. h.	1. t. 16.
267 dumosùm. s.g. (pu.) bushy.	_____	1816. _____ G. h.	1. t. 19.
268 Scarboròviæ. s.g. (d.li.) L. Scarborough's	_____	1817. _____ G. h.	2. t. 117.
269 pustulòsum. s.g. (bh.) blistered-leav'd.	_____	1816. 5. 10. G. h.	1. t. 11.
270 bellulùm. s.g. (pu.) neat.	_____	_____ _____ G. h.	1. t. 60.
moschàtum. A.G. c. ic.	_____		
271 Fairliæ. s.g. (ro.) Mrs. Fairlie's.	_____	1821. _____ G. h.	Sweet ger. 2. t. 178.
272 scintillans. s.g. (sa.) sparkling.	_____	1812. _____ G. h.	1. t. 28.
273 platypetalon. s.g. (w.) broad-petaled.	_____	_____ _____ G. h.	2. t. 116.
a rigidum. s.g. (wh.) rigid.	_____	_____ _____ G. h.	2. t. 116. f. a.
β hirsütum. s.g. (w.) hairy.	_____	_____ _____ G. h.	2. t. 116. f. b.
274 optabile. s.g. (wh.) desirable.	_____	_____ _____ G. h.	1. t. 62.
275 ornátum. s.g. (wh.) ornate.	_____	1817. _____ G. h.	1. t. 39.
276 modéstum. s.g. (bh.) modest.	_____	1822. _____ G. h.	3. t. 204.
277 rhodoléntum. s.g. (bh.) rose-smelling.	_____	1824. _____ G. h.	3. t. 291.
278 picturátum. s.g. (d.bh.) pictured.	_____	1827. _____ G. h.	ser. 2. t. 34.
279 papyrâceum. s.g. (w.) paper-white.	_____	_____ _____ G. h.	ser. 2. t. 22.
280 Newshamianum. s.g. (bh.) Miss Newsham's.	_____	1821. _____ G. h.	2. t. 144.
281 Cosmiánum. s.g. (w.) perfumed.	_____	_____ _____ G. h.	2. t. 189.
282 Brightiánum. s.g. (w.) Miss Bright's.	_____	1823. _____ G. h.	3. t. 227.
283 dépendens. s.g. (wh.) pendent-petal'd.	_____	_____ _____ G. h.	2. t. 195.
284 Mattocksianum. s.g. (wh.) Mrs. Sweet's.	_____	_____ _____ G. h.	3. t. 234.
285 venústum. s.g. (bh.) comely-flower'd.	_____	1822. _____ G. h.	2. t. 167.
286 tinctum. s.g. (w.pu.) stained	_____	1826. _____ G. h.	ser. 2. t. 29.

287 obscūrum. s.g. (<i>wh.d.</i>) darkened-pet'd.	1818.	— G. h.	Sweet ger. 1. t. 89.
288 atrofūscum. s.g. (<i>d.pu.</i>) dark-brown.	—	— G. h.	— 1. t. 82.
289 acutilōbum. s.g. (<i>wh.</i>) acute-lobed.	1822.	— G. h.	— 2. t. 184.
290 compár. c.c. (<i>wh.p.</i>) partner.	1818.	— G. h.	
291 Hoareānum. s.g. (<i>bh.p.</i>) fair Rosamond.	—	— G. h.	Sweet ger. 1. t. 80.
292 Lambérti. s.g. (<i>pu.</i>) Lambert's.	—	— G. h.	— 2. t. 104.
293 recurvātum. s.g. (<i>li.</i>) recurved-petaled.	—	— G. h.	— 3. t. 223.
294 divérgens. s.g. (<i>li.</i>) spreading-petal'd.	1826.	— G. h.	Willd. h. ber. 1. t. 37.
295 penicillātum. W.en. (<i>wh.</i>) pencilled.	1794.	— G. h.	Andr. geran. c. ic.
296 dentātum. A.G. (<i>re.</i>) toothed.	...	— G. h.	
297 florēscens. H.G. (<i>wh.</i>) triangular-l'd.	1818.	— G. h.	
298 Sabīni. H.S. (<i>pu.</i>) Sabine's.	1821.	— G. h.	Bot. mag. 148.
299 betulīnum. DC. (<i>pu.</i>) Birch-leaved.	C. B. S.	1759.	Jacq. ic. rar. 3. t. 545.
300 alniōlīum. W.en. (<i>fl.</i>) Alder-leaved.	Hybrid.	...	Jacq. ic. rar. 3. t. 538.
301 betulæfōlium. Sch. (<i>bh.</i>) Birch-like.	...	— G. h.	Andr. geran. c. ic.
302 Willdenowī. s. (<i>bh.</i>) Willdenow's.	...	— G. h.	
303 hermanniōlīum. DC. (<i>bh.</i>) Hermannia-l'd.	C. B. S.	...	
304 rígidum. DC. (<i>wh.</i>) rigid.	1800.	4. 6. G. h.	
305 ribifōlium. DC. (<i>wh.</i>) Currant-leaved	1790.	5. 8. G. h.	
306 odoriferum. (<i>wh.</i>) odoriferous.	1798.	— G. h.	
Gerānium scábrum. A.G. nec aliorum.	...	— G. h.	
307 jocundūm. H.G. (<i>li.</i>) pleasant.	1818.	— G. h.	Andr. geran. c. ic.
308 Broadleæ. c.c. (<i>pu.</i>) Mrs. Broadley's.	—	— G. h.	
309 propinquum. c.c. (<i>li.</i>) neighbouring.	1818.	— G. h.	Sweet ger. ser. 2. t. 15.
310 décorum. s.g. (<i>li.</i>) neat-flowered.	—	— G. h.	— 3. t. 267.
311 delicātum. H.G. (<i>li.</i>) delicate.	—	— G. h.	Jacq. schoen. 2. t. 139.
312 semitrilobūm. DC. (<i>fl.</i>) three-lobed.	C. B. S.	1800.	Andr. geran. c. ic.
313 Irbyānum. c.c. (<i>ro.</i>) Irby's.	Hybrid.	...	Sweet ger. ser. 2. t. 42.
314 zingiberinūm. s.g. (<i>bh.</i>) Ginger-scented.	—	— G. h.	— 1. t. 71.
315 mixtūm. s.g. (<i>bh.</i>) mixed.	1826.	— G. h.	
316 myrtillūm. H.S. (<i>pu.</i>) Myrtle-scented.	1819.	5. 8. G. h.	
317 rūbidum. (<i>pu.</i>) red-purple.	C. B. S.	1820.	Andr. geran. c. ic.
Gerānium rūbens. A. G. non Wendl.	1805.	— G. h.	
318 spūrium. W.en. (<i>pu.</i>) spurious.	Hybrid.	...	
319 nōthum. W.en. (<i>pu.</i>) mungrel.	—	— G. h.	
320 crīspum. DC. (<i>li.</i>) curl-leaved.	C. B. S.	1774.	Sweet ger. 4. t. 383.
321 grātum. W.en. (<i>bh.</i>) grateful-scent'd.	Hybrid.	6. 11. G. h.	
322 consanguīneum. W.en. (<i>pu.</i>) kindred.	—	— 4. 8. G. h.	
323 pallidūm. W.en. (<i>pa.</i>) pale-flowered.	—	— G. h.	
324 citriodōrum. (<i>wh.</i>) Citron-scented.	C. B. S.	—	Andr. geran. c. ic.
325 uniflōrum. DC. (<i>fl.</i>) one-flowered.	—	— G. h.	
326 limōnum. s.g. (<i>pu.</i>) Lemon-scented.	Hybrid.	1823.	Sweet ger. 3. t. 278.
327 notābile. H.G. (<i>li.</i>) Hoare's Harlequin.	—	— G. h.	L'Herit. ger. t. 31.
328 scábrum. DC. (<i>li.</i>) rough wedge-l'd.	C. B. S.	1822.	— t. 30.
329 tricuspidātum. DC. (<i>wh.</i>) three-point'd.	—	— G. h.	
330 Perryānum. c.c. (<i>wh.</i>) Perry's.	Hybrid.	1775.	Wendl. coll. ic.
331 glaucoïdes. w.c. (<i>wh.</i>) glaucum-like.	—	— G. h.	Sweet ger. 1. t. 57.
332 glaucum. DC. (<i>wh.</i>) glaucous-leaved.	C. B. S.	1780.	Andr. geran. c. ic.
333 lanceolātum. (<i>wh.</i>) lance-leaved.	—	— G. h.	Wendl. coll. ic.
334 diversifōlium. w.c. (<i>wh.</i>) different-l'd.	—	— G. h.	
335 cuspidātum. DC. (<i>wh.</i>) sharp-pointed.	—	— G. h.	
336 sorōrium. DC. (<i>wh.</i>) sister.	Hybrid.	1816.	
337 Tankervilleæ. H.G. (<i>wh.</i>) Lady Tankerville's	—	— G. h.	
338 acutifōlium. (<i>wh.</i>) acute-leaved.	—	— G. h.	
339 trifurcatūm. c.c. (<i>wh.</i>) three-forked.	—	— G. h.	
340 lāvigātum. DC. (<i>wh.</i>) glaucous trifid-l.	C. B. S.	1821.	Cav. dis. 4. t. 121. f. 1.
341 trifoliātum. s.g. (<i>wh.</i>) trifoliate-leaved.	—	— G. h.	Sweet ger. 3. t. 294.
oxyphyllum. DC. Gerānium trifoliātum. A.G. ic.	—	— G. h.	
342 pátlum. DC. (<i>ro.</i>) spreading.	1812.	4. 7. G. h.	Jacq. ic. rar. 3. t. 541.
343 hepaticifōlium. DC. (<i>li.</i>) Hepatica-l'd.	1791.	5. 8. G. h.	
344 mitellafōlium. (<i>wh.</i>) Mitella-leaved.	1791.	5. 10. G. h.	Andr. geran. c. ic.
Gerānium hepaticifōlium. A.G. nec aliorum.	—	— G. h.	
345 fuscātum. DC. (<i>ro.</i>) dark-marked.	1812.	— G. h.	Jacq. ic. rar. 3. t. 540.
346 saniculæfōlium. DC. (<i>pu.</i>) Sanicle-leav'd	1806.	6. 9. G. h.	— 3. t. 539.
cortusafōlium. Jacq. non L. Her.	—	— G. h.	
347 crenæfōlium. c.c. (<i>w.</i>) notch'd-petaled.	1817.	— G. h.	
348 sclerātum. c.c. (<i>wh.</i>) Celery-leaved.	1819.	— G. h.	
349 viridifōlium. s.g. (<i>sa.</i>) bright-green-l'd.	1825.	5. 9. G. h.	
350 expánsum. (<i>re.</i>) expanded.	1823.	5. 10. G. h.	Sweet ger. 4. t. 331.
	H		

351	multidentatum. (re.) many-toothed.	Hybrid.	1823.	5. 10.	G. h.		Sweet ger. 3. t. 217.
352	grandidentatum. s.g. large-toothed.	_____	_____	_____	G. h.	_____	3. t. 270.
353	Belladonna. s.g.(bh.) painted-lady.	_____	_____	_____	G. h.		
354	triúmphans. c.c. (bh.) triumphant.	_____	1818.	_____	G. h.		
355	serratifólium. s.g.(bh.) saw-leaved.	_____	1822.	_____	G. h.	Sweet ger. 3. t. 221.	
356	variegátum. DC.(wh.) variegated-fl'd.	C. B. S.	1812.	_____	G. h.	Cav. dis. 4. t. 118. f. 3.	
357	amplissimum. w.c.(wh.) stately.	Hybrid.	1796.	_____	G. h.	Wendl. coll. ic.	
	<i>Geranium grandiflorum</i> . var. A.G. c. ic. <i>Formannii</i> .	Hort.					
358	grandiflorum. DC.(wh.) great-flower'd.	C. B. S.	1794.	_____	G. h.	Sweet ger. 1. t. 29.	
359	Parmentéri. c.c.(wh.) Parmentier's.	Hybrid.	1822.	_____	G. h.		
360	delphinifólium. DC. (vi.) Larkspur-ld.	_____	_____	G. h.		
361	gossypiifólium. (li.) Cotton-tree-ld.	_____	1805.	5. 10.	G. h.	Andr. ger. c. ic.	
	<i>Geranium grandiflorum</i> . var. <i>carnea</i> . A.G.						
362	électum. s.g. (wh.) elect.	Hybrid.	1804.	_____	G. h.	Sweet ger. 3. t. 238.	
363	asperifólium. s.g.(re.) rough-leaved.	_____	1807.	_____	G. h.	_____ 2. t. 169.	
364	Curtisiánum. DC.(wh.) Curtis's.	_____	_____	G. h.		
365	nóbile. DC. (li.) noble.	_____	1822.	_____	G. h.		
366	opulifólium. s.g. (sa.) Guelder-Rose-ld.	_____	1819.	_____	G. h.	Sweet ger. 1. t. 53.	
367	malváceum. (bh.) Mallow-like.	_____	_____	G. h.		
368	hæmastictum. s.g.(bh.) blood-spotted.	_____	1827.	_____	G. h.	Sweet ger. ser. 2. t. 18.	
369	psilophyllum. s.g.(re.) smooth-leaved.	_____	1825.	_____	G. h.	_____ 4. t. 356.	
370	lúcidum. s.g. (bh.) glossy-leaved.	_____	1826.	_____	G. h.	_____ 4. t. 373.	
371	Princeánum. s.g. (wh.) Prince's.	_____	1827.	_____	G. h.	_____ 4. t. 386.	
372	clárum. s.g. (wh.) clear white.	_____	1826.	_____	G. h.	_____ 4. t. 366.	
373	recurvifólium. s.g.(wh.) recurve-leav'd.	_____	1825.	_____	G. h.	_____ 4. t. 343.	
374	bryoníafólium. s.g.(bh.) Bryony-leav'd.	_____	1824.	_____	G. h.	_____ 4. t. 320.	
375	Richiánum. s.g.(bh.) Miss Rich's.	_____	1826.	_____	G. h.	_____ 4. t. 370.	
376	platanifólium. s.g.(li.) Plane-leaved.	_____	1824.	_____	G. h.	_____ 4. t. 326.	
377	verecúndom. s.g.(pi.) blushing.	_____	_____	_____	G. h.	_____ 4. t. 316.	
378	macránthon. s.g.(wh.) large-flowered.	_____	1821.	_____	G. h.	Sweet ger. 1. t. 83.	
379	Couátsiae. s.g. (sa.) Mrs. Coutts's.	_____	1822.	_____	G. h.	_____ 3. t. 269.	
380	rútulum. c.c. (sa.) light red.	_____	1821.	_____	G. h.		
381	schizophyllum. s.g.(sa.) jagged.	_____	1822.	_____	G. h.	Sweet ger. 3. t. 289.	
382	Barnardiánum. s.g.(sa.) Mr. Barnard's.	_____	1820.	_____	G. h.	_____ 2. t. 127.	
383	mucronatúm. s.g.(d.re.) mucronate.	_____	1823.	_____	G. h.	_____ 3. t. 275.	
384	Gurneyánum. s.g. (d.re.) Mr. Gurney's.	_____	1826.	_____	G. h.	_____ 4. t. 393.	
385	argútum. s.g. (sc.) sharp-toothed.	_____	1824.	_____	G. h.	_____ 4. t. 344.	
386	latifólium. s.g. (re.) broad-leaved.	_____	_____	_____	G. h.	_____ 4. t. 335.	
387	megalánthum. s.g.(re.) grand-flowered.	_____	1826.	_____	G. h.	ser. 2. t. 4.	
388	megálēion. s.g. (sc.) magnificent.	_____	_____	_____	G. h.	ser. 2. t. 5.	
389	cratégifólium. s.g.(re.) Hawthorn-ld.	_____	1824.	_____	G. h.	ser. 2. t. 19.	
390	cotylophylum. (re.) hollow-leaved.	_____	1827.	_____	G. h.	ser. 2. t. 33.	
391	Clintóniae. s.g. (re.) Ds.ofNewcastle's.	_____	_____	_____	G. h.	4. t. 392.	
392	campylosépalon. s.g.(pu.) reflex-calyxed.	_____	1826.	_____	G. h.	4. t. 379.	
393	Náirnii. s.g. (d.re.) Nairn's.	_____	1825.	_____	G. h.	4. t. 376.	
394	rhodánthum. s.g.(ro.) rose-colour'd.	_____	1824.	_____	G. h.	3. t. 282.	
395	rhodopétalon. s.g.(ro.) rosy-petal'd.	_____	1826.	_____	G. h.	ser. 2. t. 14.	
396	acutidentatum. s.g.(sa.) acute-toothed.	_____	1827.	_____	G. h.	ser. 2. t. 40.	
397	Milleri. (re.) Mr. Miller's.	_____	1824.	_____	G. h.		
398	conchiifólium. (li.) shell-leaved.	_____	1820.	_____	G. h.		
399	striátum. s.g. (bh.) streak-flowered.	_____	1818.	3. 8.	G. h.	Sweet ger. 1. t. 1.	
400	rigéscens. s.g. (li.) stiff-leaved.	_____	_____	4. 9.	G. h.	_____ 2. t. 112.	
401	eriophylum. s.g.(pu.) woolly round-ld.	_____	_____	5. 10.	G. h.	_____ 2. t. 141.	
402	interníxtum. (li.) intermixed.	_____	_____	4. 8.	G. h.		
403	abutiloídes. s.g. (pu.) Abutilon-like.	_____	1820.	4. 9.	G. h.	Sweet ger. ser. 2. t. 3.	
404	cordátum. DC. (pu.) heart-leaved.	C. B. S.	1774.	3. 8.	G. h.	1. t. 67.	
405	conduplicátum. DC.(pu.)curl'd-heart-ld.	Hybrid.	_____	G. h.		
406	rubro-cíncustum. DC.(pu.) red-edged.	_____	_____	G. h.		
407	Góweri. s.g. (pi.) Mr. LevisonGower's.	_____	1825.	5. 8.	G. h.	Sweet ger. 4. t. 333.	
408	papilionáceum. DC.(pu.)Butterfly.	C. B. S.	1724.	4. 7.	G. h.	1. t. 27.	
409	tomentosum.DC.(wh.) Pennyroyal.	_____	1790.	_____	G. h.	2. t. 168.	
410	vitifólium. DC. (li.) Vine-leaved.	_____	1724.	4. 8.	G. h.	Cav. dis. 4. t. 111. f. 2.	
411	capitátum. DC. (ti) Rose-scented.	_____	1690.	_____	G. h.	4. t. 105. f. 1.	
412	cónicolor. s.g. (re.) self-coloured.	Hybrid.	1820.	_____	G. h.	Sweet ger. 2. t. 140.	
	β major. s.g. (cr.) larger.	_____	_____	_____	G. h.	2. t. 140. f. b.	
413	congéstum. s.g. (pu.) close-headed.	_____	1825.	_____	G. h.	4. t. 325.	
414	calamistrátum. s.g. (pu.)curled-lobed.	_____	1828.	_____	G. h.	ser. 2. t. 36.	
415	gravéolens. DC. (li.) Odour of Rose.	C. B. S.	1774.	3. 7.	G. h.	L'Herit. ger. t. 17.	
	β variegátum. (li.) striped-leaved.	_____	_____	G. h.	Andr. geran. c. ic.	
	<i>Geranium capitatum</i> . var. A.G.						

416	áspérum. DC.	(fl.) rough multifid.	C. B. S.	1795.	5. 9.	G. h.	Roth. abh. p. 51. t. 10.
417	Rádula. DC.	(li.) rasp-leaved.	—	1774.	3. 7.	G. h.	L'Herit. ger. t. 16.
418	balsámeum. DC.	(fl.) balsamic.	—	1790.	5. 9.	G. h.	Jacq. ic. rar. 3. t. 53.
419	Vandésiae. s.g. (bh.)	Comtesse de Vandes'. Hybrid.	1818.	—	G. h.	Sweet ger. 1. t. 7.	
420	hispidum. DC.	(wh.) hispid.	C. B. S.	1790.	—	G. h.	Cav. dis. 4. t. 110. f. 1.
421	callistón. s.g. (sa.)	graceful.	—	1827.	—	G. h.	Sweet ger. ser. 2. t. 9.
422	anthemifólium. (re.)	Anthemis-ld.	Hybrid.	1822.	—	G. h.	Sweet ger. 2. t. 153.
423	pyrethrifólium. s.g. (sc.)	Feverfew-ld.	—	1821.	—	G. h.	— 4. t. 336.
424	tanacetifólium. s.g. (re.)	Tansy-leaved.	—	1824.	—	G. h.	— 4. t. 317.
425	sphærocéphalon. s.g. (sc.)	round-headed.	—	—	—	G. h.	— 3. t. 285.
426	Moreánum. s.g. (sc.)	More's victory.	—	1823.	—	G. h.	— 4. t. 310.
427	obcordátum. s.g. (sc.)	obcordate-petal'd.	—	—	—	G. h.	— 4. t. 330.
428	cuneíflorum. s.g. (sc.)	wedge-petal'd.	—	1825.	—	G. h.	— 2. t. 124.
429	chrysanthemifólium. (sc.)	Chrysanthm.-ld.	—	1821.	4. 10.	G. h.	— 3. t. 253.
430	Dobreeánum. s.g. (sc.)	Mrs. Dobree's.	—	1818.	—	G. h.	— 3. t. 272.
431	Bishóppae. s.g. (sc.)	Mrs. Bissopp's.	—	1822.	—	G. h.	— 3. t. 265.
432	signátum. s.g. (re.)	marked-leaved.	—	—	—	G. h.	— 4. t. 362.
433	Spini. C.R. (bh.d.)	Spin's.	—	—	—	G. h.	L'Herit. ger. t. 14.
434	querçifólium. DC. (bh.)	Oak-leaved.	C. B. S.	1774.	3. 8.	G. h.	— t. 15.
	bipinnatífidum. (bh.)	bipinnatifid.	—	—	—	G. h.	Sweet ger. ser. 2. t. 43.
435	tortuósom. s.g. (li.)	twisted-petal'd.	—	1826.	—	G. h.	— 1. t. 26.
436	exímium. s.g. (pi.)	select.	Hybrid.	1818.	4. 8.	G. h.	Sweet ger. 1. t. 35.
437	Esséxiæ. C.C. (pi.)	Lady Essex's.	—	—	—	G. h.	— 2. t. 164.
438	oblátum. s.g. (pi.)	oblade-leaved.	—	—	—	G. h.	— 3. t. 201.
439	Murryánum. s.g. (bh.)	Ld. Jas. Murray's.	—	—	—	G. h.	— 4. t. 307.
440	calocéphalon. s.g. (bh.)	pretty-headed.	—	1820.	—	G. h.	— 4. t. 314.
441	obovátum. s.g. (li.)	obovate-petal'd.	—	1825.	—	G. h.	Andr. geran. c. ic.
442	dilatátum. s.g. (li.)	dilated-leav'd.	—	1824.	—	G. h.	—
443	augústum. DC. (bh.)	august.	—	1809.	—	G. h.	Sweet ger. 1. t. 64.
	β Brúcti. C.C. (bh.)	Bruce's.	—	—	—	G. h.	— 1. t. 92.
444	Breesiánum. s.g. (pi.)	Brees'e's.	—	1818.	—	G. h.	— 1. t. 175.
445	Husseyánum. s.g. (pu.)	Lady Hussey's.	—	—	—	G. h.	— 3. t. 248.
446	Wellsiánum. s.g. (sc.)	Mr. Wells's.	—	1822.	3. 8.	G. h.	— 3. t. 264.
447	Kíngii. s.g. (sc.)	Mr. King's.	—	—	3. 12.	G. h.	— 3. t. 224.
448	Scóttii. s.g. (sc.)	Sir. C. Scott's.	—	—	3. 9.	G. h.	— 3. t. 252.
449	Pálkii. s.g. (cr.)	Mr. Palk's.	—	—	1. 12.	G. h.	— 3. t. 236.
450	rotundilóbum. s.g. (sc.)	round-lobed.	—	1823.	3. 9.	G. h.	— 4. t. 385.
451	latilóbum. s.g. (sc.)	broad-lobed.	—	—	—	G. h.	— 4. t. 400.
452	Russelliánum. s.g. (sc.)	Russell's.	—	1826.	4. 9.	G. h.	— 4. t. 384.
453	translúcens. s.g. (sc.)	transparent.	—	1824.	—	G. h.	— 4. t. 389.
454	Barclayánum. s.g. (sc.)	Mr. Barclay's.	—	—	—	G. h.	— 4. t. 353.
455	Harewoódiæ. s.g. (d.re.)	Lady Harewood's.	—	1826.	—	G. h.	— 3. t. 296.
456	Stewártii. s.g. (da.sc.)	Mr. Stewart's.	—	1825.	—	G. h.	Sweet ger. 2. t. 109.
457	lasiofíllum. s.g. (sa.)	woolly-divided-ld.	—	—	—	G. h.	— 2. t. 118.
458	Lechiánum. DC. (bh.)	acute-notched.	—	1820.	—	G. h.	Sweet ger. ser. 2. t. 38.
459	jatrophæfólium. DC. (fl.)	Jatropa-ld.	—	—	—	G. h.	L'Herit. ger. t. 20.
460	denticulátum. DC. (li.)	tooth-leaved.	C. B. S.	1789.	5. 8.	G. h.	Bot. mag. 143.
461	viscosissimum. s.g. (bh.)	viscous.	—	1820.	3. 8.	G. h.	Sweet ger. 3. t. 259.
	a ábum. (wh.)	white-flowered.	—	—	—	G. h.	—
	β cárneum. (fl.)	flesh-coloured.	—	—	—	G. h.	—
462	carbasinum. s.g. (su.)	linen-flower'd.	Hybrid.	1827.	—	G. h.	Sweet ger. ser. 2. t. 149.
463	glutinósom. DC.	(fl.) clammy.	C. B. S.	1777.	5. 6.	G. h.	—
	β nigréscens.	(fl.) black-marked.	—	—	—	G. h.	L'Herit. ger. t. 20.
464	cynosbatifólium. DC. (li.)	Cynosbat-ld.	Hybrid.	—	4. 8.	G. h.	Bot. mag. 143.
	Geraniúm oxoniéne. A.G. IC.	—	—	—	—	—	Sweet ger. 3. t. 259.
465	verbenaefólium. s.g. (li.)	Vervain-ld.	Hybrid.	1818.	—	G. h.	Sweet ger. 2. t. 149.
	piperátum. DC?	—	—	—	—	—	—
466	meliásum. s.g. (da.)	Balm-scented.	Hybrid.	—	—	G. h.	Sweet ger. 1. t. 5.
467	volatíflorum. s.g. (sc.)	flying-flower'd.	—	1823.	—	G. h.	— 3. t. 284.
468	brévipes.	(sc.) short-peduncled.	—	1824.	—	G. h.	—
469	ternatón. DC.	(li.) ternate-leaved.	C. B. S.	1789.	5. 8.	G. h.	Sweet ger. 2. t. 165.
470	patentissimum. w.c. (li.)	most-spreading.	—	1822.	—	G. h.	Wendl. coll. ic.
471	spinósom. DC.	(st.) thorny.	—	1795.	5. 6.	G. h.	Paters. it. t. ad. p. 67.
472	tripartitum. s.g. (st.)	brittle-stalked.	—	1789.	5. 8.	G. h.	Sweet ger. 2. t. 115.
	Geraniúm frágile. A.R. 37.	—	—	—	—	—	—
473	incisum. DC.	(st.) cut-leaved.	—	1791.	—	G. h.	Sweet ger. 1. t. 93.
474	canéscens. s.g. (bh.)	canescent.	—	—	—	G. h.	Jacq. schœn. 2. t. 136.
475	abrotanifólium. DC. (bh.)	Southernwood-ld.	—	1791.	—	G. h.	—
476	artemisiæfólium. DC.	Wormwood-ld.	—	1817.	—	G. h.	—

477	frægrans, DC. (wh.) Nutmeg-scent'd. Hybrid.	5. 10. G. $\frac{1}{2}$.	Sweet ger. 2. t. 172.
478	odoratissimum, DC. (wh.) sweet-scent'd. C. B. S.	1724.	— G. $\frac{1}{2}$.	3. t. 299.
479	exstipulatum, DC. (bh.) soft trifid-l'd. <i>Geranium suaveolens</i> , A.G. ic.	1779.	— G. $\frac{1}{2}$.	L'Herit. ger. t. 35.
480	dissectum, s.g. (wh.) dissected-l'd. Hybrid.	1822.	5. 11. G. $\frac{1}{2}$.	Sweet ger. 3. t. 247.
481	glomeratum, DC. (w.) close-headed. <i>australis</i> , s.g. 68. non nc.	1792.	5. 8. G. $\frac{1}{2}$.	Jacq. fl. ecl. 1. t. 98.
482	australe, DC. (wh.) New Holland.	—	— G. $\frac{1}{2}$.	Jacq. fl. ecl. 1. t. 100.
483	inodorum, DC. (w.pu.) scentless.	1796.	3. 8. G. $\frac{1}{2}$.	Sweet ger. 1. t. 56.
484	columbinum, DC. (pu.) Doves-foot.	C. B. S.	1795. 6. 10. G. $\frac{1}{2}$.	Jacq. schœn. 2. t. 133.
485	procumbens, DC. (bh.) procumbent.	—	1801. 4. 5. G. $\frac{1}{2}$.	Andr. bot. rep. 234.
486	humifusum, DC. (bh.) trailing.	Canaries.	— 4. 10. H. \odot .	Sweet ger. 1. t. 42.
487	chamaedryfolium, DC. (wh.) <i>Chamaedrys</i> -l'd. C. B. S.	1812.	5. 6. G. $\frac{1}{2}$.	Jacq. ic. rar. 3. t. 523.
488	altheoides, DC. (wh.) Althaea-leaved.	—	1724. 4. 6. G. $\frac{1}{2}$.	L'Herit. ger. t. 10.
489	grossularioides, DC. (fl.) Gooseberry-l'd.	—	1731. 4. 8. G. $\frac{1}{2}$.	Cav. dis. 4. t. 119. f. 2.
490	ancéps, DC. (pu.) flat-stalked.	—	1788. 5. 7. G. $\frac{1}{2}$.	Jacq. col. 4. t. 22. f. 3.
491	parviflorum, A.G. (pu.) small-flowered.	—	1800. — G. $\frac{1}{2}$.	Andr. geran. c. ic.
492	acgnaticum, L.T. (re.) <i>Tristan d'Acuna</i> , T.d'Acunna.	1818.	— G. $\frac{1}{2}$.	Sweet ger. 4. t. 327.
493	senecioïdes, DC. (wh.) Groundsel-l'd.	C. B. S.	1775. 6. 7. G. \odot .	Cav. dis. 4. t. 98. f. 1.
494	alchimilloides, DC. (wh.) mantle-leaved.	—	1693. 5. 10. G. $\frac{1}{2}$.	And. repos. 98.
495	Heritiæ, Jacq. (wh.) L'Heritiere's.	—	1820. — G. $\frac{1}{2}$.	Sweetger. 1. t. 36. f.a.
496	tabulare, DC. (wh.) rough-stalked.	—	1775. 5. 8. G. $\frac{1}{2}$.	— 1. t. 36. f. b.
497	ovale, DC. (ro.) oval-leaved.	—	1774. — G. $\frac{1}{2}$.	— 3. t. 232.
498	élégans, DC. (wh. pu.) elegant.	—	1795. — G. $\frac{1}{2}$.	L'Herit. ger. t. 9.
	α major, s.g. (wh.) smaller-flowered.	—	— G. $\frac{1}{2}$.	Sweet ger. 3. t. 235.
	β minor, s.g. (v.) larger-flowered.	—	— G. $\frac{1}{2}$.	And. repos. 98.
499	schizopetalum, s.g. (y. br.) divided-petal'd.	—	1821. 6. 10. G. $\frac{1}{2}$.	Sweetger. 1. t. 52.
500	pinguifolium, s.g. (bh.) greasy leaved. Hybrid? 5. 10. G. $\frac{1}{2}$.	— 1. t. 52.
501	latéripes, DC. (va.) Ivy-leaved.	C. B. S.	1787. — G. $\frac{1}{2}$.	L'Herit. ger. t. 24.
	α viridisfolium. (li.) green-leaved.	—	— G. $\frac{1}{2}$.	— Andr. geran. c. ic.
	β zonatum. (li.) Horseshoe-mark'd.	—	— G. $\frac{1}{2}$.	—
	γ rôseum. (ro.) rose-coloured.	—	— G. $\frac{1}{2}$.	—
	δ albumarginatum, DC. white-margined.	—	— G. $\frac{1}{2}$.	— Andr. geran. c. ic.
502	peltatum, DC. (li.) peltate-leaved.	—	1701. — G. $\frac{1}{2}$.	Bot. mag. 20.
	β variegatum, A.G. (li.) variegated-l'd.	—	— G. $\frac{1}{2}$.	Andr. geran. c. ic.
503	scutatum, s.g. (wh.) shield-leaved.	—	1819. — G. $\frac{1}{2}$.	Sweet ger. 1. t. 95.
JENKIN'SONIA. S.G. JENKINS'ONIA. <i>Monadelphus Heptandria</i> .				
1	tetragona, s.g. (li.) square-stalked.	C. B. S.	1774. 5. 8. G. $\frac{1}{2}$.	Sweet ger. 1. t. 99.
	β variegata. (li.) variegated.	—	— G. $\frac{1}{2}$.	— 1. t. 99.
2	quinata, s.g. (st.) quinate-leaved.	—	1793. — G. $\frac{1}{2}$.	— 1. t. 79.
	Pelargonium quinatum, B.M. 547. <i>Geranium præmorsum</i> , A.R. 150.	—	—	—
3	pendula, s.g. (pu.) pendulous-branch'd.	— 5. 8. G. $\frac{1}{2}$.	Sweet ger. 2. t. 188.
4	longicaulis, s.g. (bh.) long-stalked.	— — G. $\frac{1}{2}$.	Jacq. ic. rar. 3. t. 533.
	<i>Pelargonium longicáule</i> , Jacq.	—	—	—
5	anemonefolia, s.g. (re.) Anemone-l'd.	— — G. $\frac{1}{2}$.	Jacq. ic. rar. 3. t. 535.
6	caucalifolia, s.g. (bh.) Caucalus-leav'd.	—	1812. 3. 9. G. $\frac{1}{2}$.	— 3. t. 529.
7	multicaulis, s.g. (vi.) many-stalked.	—	1802. 6. 8. G. $\frac{1}{2}$.	— 3. t. 534.
8	láceræ, s.g. (bh.) torn-leaved.	—	1731. — G. $\frac{1}{2}$.	— 3. t. 532.
9	Synnoti, s.g. (pu.) Synnot's.	—	1825. — G. $\frac{1}{2}$.	Sweet ger. 4. t. 342.
10	coriandrifolia, s.g. (bh.) Coriander-l'.	—	1724. 3. 9. G. $\frac{1}{2}$.	— 1. t. 34.
	<i>Pelargonium coriandrifolium</i> , s.g.	—	—	—
11	myrrifolia, s.g. (ii.) Myrrh-leaved.	—	1696. 5. 8. G. $\frac{1}{2}$.	Jacq. ic. rar. 3. t. 531.
	<i>Pelargonium betonicum</i> , Jacq.	—	—	—
12	bullata, s.g. (wh.) blistered.	— — G. $\frac{1}{2}$.	Jacq. ic. rar. 3. t. 530.
13	canariensis, (wh.) Canary.	Canaries.	1802. 5. 9. G. $\frac{1}{2}$.	Willd. hort. ber. t. 17.
	<i>Pelargonium canariense</i> , w.	—	—	—
CIC'ONIUM. S.G. CIC'ONIUM. <i>Monadelphus Heptandria</i> .				
1	acetosum, s.g. (sa.) Sorrel-leaved.	C. B. S.	1710. 5. 9. G. $\frac{1}{2}$.	Andr. geran. c. ic.
2	scándens, s.g. (ro.) climbing.	—	1800. — G. $\frac{1}{2}$.	—
3	stenopetalum, s.g. (sc.) narrow-petal'd.	—	— G. $\frac{1}{2}$.	—
4	leptopetalum, (sc.) slender-petal'd.	—	— G. $\frac{1}{2}$.	Andr. geran. c. ic.
	<i>Geranium stenopetalum</i> , A.G. nec aliorum.	—	—	—
5	pùmilum, s.g. (sc.) dwarf.	C. B. S.	— G. $\frac{1}{2}$.	Andr. geran. c. ic.
	<i>Geranium zonale minima</i> , A.G.	—	—	—
6	lateritium, s.g. (sc.) brick-coloured.	—	— G. $\frac{1}{2}$.	Jacq. ecl. 1. t. 97. f. ult.
7	malvaefolium, (ii.) Mallow-leaved.	—	1812. — G. $\frac{1}{2}$.	— 1. t. 97.
	<i>Pelargonium malvaefolium</i> , Jacq.	—	—	—
8	heterogànum, s.g. (li.) six-stamened.	Hybrid.	1786. — G. $\frac{1}{2}$.	L'Herit. geran. t. 18.
9	álbidum, c.c. (wh.) white-flowered.	C. B. S. — G. $\frac{1}{2}$.	—

10	zonâle. s.g.	(va.) horse-shoe.	C. B. S.	1710.	4. 12.	G. h.	Cav. dis. 4. t. 98. f. 2.
	<i>a. lilacinum.</i>	(li.) lilac-coloured.	_____	_____	_____	G. h.	
	β coccineum.	(sc.) scarlet.	_____	_____	_____	G. h.	
	γ crystallinum.	(sc.) coral-stalked.	_____	_____	_____	G. h.	
	δ marginatum.	(sc.) white-edged.	_____	_____	_____	G. h.	
11	reticulatum. s.g.	(sc.) netted-veined.	Hybrid.	1810.	5. 10.	G. h.	Sweet ger. 2. t. 143.
12	Fothergillii. s.g.	(sc.) Fothergill's.	C. B. S.	_____	G. h.	3. t. 226.
	<i>a. coccineum.</i>	(sc.) scarlet.	_____	_____	_____	G. h.	3. t. 226. a.
	β purpureum.	(pu.) purple.	_____	_____	_____	G. h.	3. t. 226. b.
13	mónstrum. s.g.	(pu.) cluster-leaved.	Hybrid.	1784.	7. 8.	G. h.	1. t. 13.
14	aequaliflòrum. c.c.	(sc.) victory.	C. B. S.	1819.	5. 10.	G. h.	Sweet ger. 1. t. 63.
15	hybridum. s.g.	(sc.) hybrid.	C. B. S.	1732.	_____	G. h.	Andr. geran. c. ic.
16	oxyphyllum. c.c.	(wh.) sour-leaved.	Hybrid.	_____	G. h.	
	<i>Geranium miniatum v. album.</i> A.G.						
17	hùmile.	(sc.) low-spreading.	C. B. S.	_____	G. h.	
18	crenatum. s.g.	(sc.) scalloped-l'd.	_____	_____	G. h.	Sweet ger. 4. t. 345.
19	bracteosum.	(sc.) large-bracted.	Hybrid.	_____	G. h.	
	<i>Pelargonium bracteosum.</i> DC.						
20	Bentinckianum.	(sc.) Bentinck's.	C. B. S.	_____	G. h.	Sweet ger. 4. t. 350.
	<i>Pelargonium Bentinckianum.</i> DC.	<i>Geranium crenatum;</i> var. <i>mollifoliata.</i>	A.G. c. ic.				
21	fulgens. s.g.	(sc.) Basilisk.	Hybrid.	1823.	_____	G. h.	Sweet ger. 4. t. 339.
22	glabrifolium. s.g.	(sc.) smooth-leaved.	_____	1825.	_____	G. h.	4. t. 363.
23	cerinum. s.g.	(pi.) wax-flowered.	Hybrid?	_____	G. h.	2. t. 176.
24	inquinans. s.g.	(sc.) staining-gland'd.	C. B. S.	1714.	5. 9.	G. h.	Andr. Geran. c. ic.
25	micranthum. s.g.	(sc.) small-flowered.	_____	_____	G. h.	Sweet ger. 3. t. 295.
ISOPETALUM. S.G. ISOPETALUM. Monadelphia Pentandria.							
1	Cotyledonis. s.g. (wh.)	Hollyhock-l'd.	St. Helena.	1765.	5. 8.	G. h.	Sweet ger. 2. t. 126.
	<i>Pelargonium Cotyledonis.</i> W.						
2?	discipes. c.c. central-stalked.	_____	1808.	G. h.	

ORDO LII.

TROPÆOLEÆ. DC. prodr. 1. p. 683.

TROPÆOLUM. DC. INDIAN-CRESS. Octandria Monogynia. L.

1	minus. DC.	(ye.) small.	Peru.	1596.	6. 10.	H. ♂.	Bot. mag. 98.
	β flôre pleno.	(ye.) double-flowered.	_____	_____	_____	G. ♀.	
2	május. DC.	(ye.) great.	_____	1686.	_____	H. ♂.	Bot. mag. 23.
	β flôre pleno.	(ye.) double-flowered.	_____	_____	_____	G. ♀.	
3	hybridum. DC.	(ye.) hybrid.	Hybrid.	_____	G. ♀.	Berg. act. holm. 32. t. 1.
4	aduncum. DC.	(ye.) fringe-flower'd.	Peru.	1775.	_____	G. ♂.	Bot. reg. t. 718.
	<i>peregrinum.</i> B.R. non L.						
5	pinnatum. DC.	(ye.) pinnate-flower'd.	_____	_____	G. ♀.	Andr. reposit. t. 535.
6	tricolorum. B.F.G. (or.)	three-coloured.	Chile.	1828.	_____	H. ♀.	Sweet br. fl. gar. 3.t.270.

ORDO LIII.

BALSAMINEÆ. DC. prodr. 1. p. 685.

BALSAMINA. DC. BALSAM. Pentandria Pentagynia.

1	hortensis. DC.	(v.) garden.	E. Indies.	1596.	7. 10.	S. ♂.	Blackw. t. 583.
	<i>Impatiens Balsamina.</i> L.						
2	coccinea. DC.	(sc.) glandular-l'd.	_____	1808.	6. 9.	S. ♂.	Bot. mag. 1256.
3	mysorensis. DC.	(re.) Mysore.	_____	1823.	_____	S. ♂.	
4	bifida. DC.	(re.) bifid-spurred.	C. B. S.	1818.	7. 10.	H. ♂.	
5	capensis. DC.	(re.) Cape.	_____	_____	_____	H. ♂.	

6 cochleáta. DC.	(sc.) shell-spurred.	China.	1826.	— H. ⓧ.	
7 chinénsis. DC.	(pu.) Chinese.	China.	1823.	— H. ⓧ.	
IMPATIENS. DC.	TOUCH ME NOT.	Pentandria Monogynia. L.			
1 nátans. DC.	(ye.) floating.	E. Indies.	1823.	6. 10. S.w. ⓧ.	
2 biflóra. w.	(or.) two-flowered.	N. America.	— H. ⓧ.	Sweet br. fl. gar. t. 43.
<i>fúlva</i> . DC.					
3 pálida. DC.	(ye.) pale-flowered.		1812.	— H. ⓧ.	
4 Nòli-tángere. DC.	(ye.) common.	England.	— H. ⓧ.	Eng. bot. 937.
5 parviflóra. DC.	(ye.) small-flowered.	Russia.	1828.	— H. ⓧ.	
6 tripétala. DC.	(ye.) three-petaled.	E. Indies.	1823.	— S. ⓧ.	

ORDO LIV.

OXALIDEÆ. DC. prodr. 1. p. 689.

AVERRH'OA. DC.	AVERRH'OA.	Pent-Decandria Pentagynia. L.			
1 Carambòla. DC.	(st.) Carambola-tree.	E. Indies.	1793. S. ⓧ.	Rumph. amb. 1. t. 35.
2 Bilímbi. DC.	(st.) Bilimbi-tree.		1791.	5. 7. S. ⓧ.	— I. t. 36.
BIO'PHYTUM. DC.	BIO'PHYTUM.	Decandria Pentagynia.			
sensitivum. DC.	(ye.) sensitive.	E. Indies.	1824.	5. 7. S. 4.	Jacq. ox. n. 21. t. 78. f. 4.
<i>O'xalis sensitiva</i> . L.					
O'XALIS. DC.	WOOD-SORREL.	Decandria Pentagynia. L.			
1 Plumíeri. DC.	(ye.) Plumier's.	S. America.	1822.	4. 6. S. ⓧ.	Bot. reg. 810.
2 Barrelieri. DC.	(bh.) Barrelier's.		—	— S. ⓧ.	Jacq. ox. n. 4. t. 3.
3 fruticosa. DC.	(ye.) frutescent.	Rio Janeiro.	—	— S. ⓧ.	
4 perénmans. DC.	(br.y.) perennial.	N. S. W.	5. 9. G. 4.	
5 Dillénii. DC.	(ye.) Dillenius's.	America.	1798.	— H. ⓧ.	Dill. elth. 2. t. 221.
6 flórida. S.P.	(ye.) prostrate.		—	— H. ⓧ.	
7 strícta. DC.	(ye.) upright.	N. America.	1658.	6. 10. H. 4.	Jacq. ox. n. 9. t. 4.
8 corniculàta. DC.	(ye.) procumbent.	Britain.	5. 10. H. 4.	Eng. bot. 1726.
9 Lyònii. DC.	(ye.) Lyons'.	N. America.	1812.	— H. 4.	
10 microphylla. DC.	(ye.) small-leaved.	N. Holland.	— H. ⓧ.	
<i>rubens</i> . H.M.					
11 repens. DC.	(ye.) creeping-stalk'd.	C. B. S.	1793.	3. 4. G. 4.	Jacq. ox. n. 11. t. 78. f. 1.
12 rósea. DC.	(ro.) rose-coloured.	Chile.	1826.	3. 6. G. 4.	Bot. mag. 2830.
<i>floribunda</i> . B.R.	1123.				
13 Símsii.	(cr.) Sims's.		1822.	4. 6. G. 4.	Bot. mag. 2415.
<i>rósea</i> . Sims Bot. mag. nec aliorum.					
14 lateriflóra. DC.	(pu.) side-flowering.	C. B. S.	1826.	3. 6. G. 4.	Jacq. Schœnb. 2. t. 204.
15 macróstylis. DC.	(pu.) long-styled.		1793.	10. 11. G. b.	Jacq. ox. n. 22. t. 9.
16 tubiflóra. DC.	(vi.) tube-flowered.		1790.	— G. b.	— n. 23. t. 10.
17 canéscens. DC.	(li.) canescant.		1822.	— G. b.	— n. 24. t. 11.
18 secúnda. DC.	(li.) secund.		1790.	— G. b.	— n. 68. t. 12.
19 hírtia. DC.	(vi.) hairy-stalked.		1787.	1. 3. G. b.	— n. 26. t. 13
20 hirtélla. DC.	(li.) tooth-stamen'd.		1820.	— G. b.	— n. 27. t. 14.
21 multiflóra. DC.	(vi.) many-flowered.		1789.	2. 3. G. b.	— n. 28. t. 15.
22 fúlgida. B.R.	(cr.) crimson.		—	— G. b.	Bot. reg. 1073.
23 rubélla. DC.	(vi.) branching red.		1791.	9. 11. G. b.	Bot. mag. 1031.
24 rosácea. DC.	(vi.) rose-coloured.		1793.	10. 2. G. b.	— 1698.
25 vírginea. DC.	(wh.) virgin.		1822.	11. 12. G. b.	Jacq. Schœnb. 3. t. 275.
26 reptátrix. DC.	(li.) creeping-rooted.		1795.	— G. 4.	Jacq. ox. n. 33. t. 20.
27 incárnata. DC.	(bh.) flesh-coloured.		1739.	4. 6. G. b.	Jacq. vind. t. 71.
28 disticha. DC.	(ye.) distichous.		1822.	— G. b.	Jacq. ox. n. 31. t. 18.
29 venósa. DC.	(vi.) veined-flower'd.		1821.	— G. b.	
30 cùprea. B.C.	(co.) copper-colour'd.		1822.	— G. b.	Lodd. bot. cab. 824.
31 tetraphylla. DC.	(pu.) four-leaved.	Mexico.	—	— H. b.	— t. 790.
32 sericea. DC.	(ye.) silky-leaved.	C. B. S.	1794.	— G. b.	Jacq. ox. n. 13. t. 77. f. 1.
33 carnósa. B.R.	(ye.) fleshy.	Chile.	1825.	4. 9. H. 4.	Bot. reg. 1063.
34 bipunctatâ. Gr.	(li.) two-spotted.	Brazil.	—	4. 6. G. 4.	Bot. mag. 2781.
35 violacea. DC.	(vi.) violet-coloured.	N. America.	1772.	5. 6. H. b.	Jacq. vind. t. 180.
36 Déppii.	(li.) Depp's.	Mexico.	1827.	6. 8. G. b.	
37 Martiana. Zuc.	(li.) Martius's.	Brazil.	1828.	5. 9. S. 4.	
38 palústris. s.s.	(li.) marsh.		—	5. 12. S. 4.	
<i>papilionacea</i> . Zuc.					

39 caprīna. DC.	(vi.) goats-foot.	C. B. S.	1757.	3. 6. G. b.	Jacq. ox. t. 76. f. 1.
40 cérnua. DC.	(ye.) drooping-yellow.	—	—	2. 5. G. b.	n. 16. t. 6.
caprīna. B.M. 237. nea aliorum.					
β flōre plēno.	(ye.) double-flowered.	—		G. b.	
41 comprēssā. DC.	(ye.) compressed.	—	1794.	12. 1. G. b.	Jacq. ox. n. 19. t. 78. f. 3.
42 dentāta. DC.	(li.) toothed.	—	1793.	11. 12. G. b.	— n. 17. t. 7.
43 lívida. DC.	(fl.) livid.	—	—	10. 11. G. b.	— n. 18. t. 8.
44 purpurāta. DC.	(bh.) purplish-leav'd.	—	1820.	— G. b.	Jacq. schœnb. t. 356.
45 macropphylla. DC.	(vi.) large-leaved.	—		G. b.	
46 lobatā. DC.	(ye.) lobed.	Chile.	1821.	4. 6. H. b.	Bot. mag. 2386.
47 ténera. B.R.	(li.) tender.	Brazil.	1826.	5. 6. S. 4.	Bot. reg. 1046.
48 monophylla. DC.	(bh.) simple-leaved.	C. B. S.	1774.	10. 11. G. b.	Jacq. ox. n. 35. t. 79. f. 3.
49 lépida. DC.	(wh.) pretty.	—	1822.	10. 2. G. b.	— n. 34. t. 21.
50 rostrata. DC.	(bh.) beaked.	—	1795.	— G. b.	— n. 36. t. 22.
51 críspā. DC.	(wh.) curled.	—	1793.	— G. b.	— n. 37. t. 23.
52 leporīna. DC.	(std.) Hare's-eared.	—	1795.	10. 11. G. b.	— n. 30. t. 25.
53 asinīna. DC.	(ye.) asses-eared.	—	1792.	11. 12. G. b.	— n. 38. t. 24.
54 lanceafolia. DC.	(ye.) spear-leaved.	—	1794.	10. 11. G. b.	— n. 40. t. 26.
55 fabafolia. DC.	(ye.) Bean-leaved.	—	1794.	— G. b.	— n. 41. t. 27.
56 laburnifolia. DC.	(ye.) Laburnum. 1d.	—	1793.	9. 10. G. b.	— n. 42. t. 28.
57 sanguínea. DC.	(ye.) bloody-leaved.	—	1795.	10. 12. G. b.	— n. 43. t. 29.
58 tricolor. DC.	(std.) three-coloured.	—	1794.	— G. b.	— n. 63. t. 47.
59 ciliáris. DC.	(pu.) ciliated.	—	1793.	— G. b.	— n. 45. t. 30.
60 arcuata. DC.	(vi.) gland-covered.	—	1795.	— G. b.	— n. 46. t. 31.
61 fláccida. DC.	(re. wh.) flaccid.	—	1812.	— G. b.	— n. 66. t. 51.
62 ferruginata. DC.	(wh.) rust-spotted.	—	1822.	— G. b.	Jacq. schœnb. 3. t. 274.
63 ambigua. DC.	(v.) ambiguous.	—	1790.	9. 12. G. b.	Jacq. ox. n. 59. t. 43.
64 undulata. DC.	(wh.) wave-leaved.	—	1795.	10. 11. G. b.	— n. 60. t. 44.
65 glandulosa. DC.	(wh.) glandulous.	—	1822.	— G. b.	— n. 62. t. 46.
66 fuscata. DC.	(wh.) brown-spotted.	—	1795.	5. 6. G. b.	— n. 61. t. 45.
67 sulphurea. DC.	(st.) sulphur-colour'd.	—	—	10. 11. G. b.	— n. 77. t. 63.
68 breviscapa. DC.	(wh.) short-stalked.	—	1823.	9. 2. G. b.	— n. 72. t. 58.
69 speciosa. DC.	(pu.) specious.	—	1690.	9. 11. G. b.	— n. 74. t. 60.
70 rigidula. Jac.	(wh.) rigid.	—	1822.	— G. b.	— n. 73. t. 59.
71 variabilis. DC.*	(v.) variable.	—	1795.	10. 12. G. b.	— n. 67.
a longiscapa. DC.	(wh.) white-flowered.	—			— t. 52.
β rūbra. DC.	(re.) red-flowered.	—			— t. 53.
72 grandiflora. w.	(wh.) great-flowered.	—	1810.	— G. b.	— n. 68. t. 54.
β Simsii. DC.	(wh.) Sims's.	—		G. b.	Bot. mag. 1683.
73 láxula. Jac.	(wh.) loose.	—	1820.	— G. b.	Jacq. ox. n. 71. t. 57.
74 purpúrea. DC.	(pu.) purple.	—	1812.	— G. b.	— n. 70. t. 56.
75 convéxula. DC.	(or.) convex-leaved.	—	1789.	11. 1. G. b.	— n. 69. t. 55.
76 lèvigata. DC.	(pu.) smooth.	—	1818.	6. 8. H. ○.	
77 punctata. DC.	(bh.) spot-leaved.	C. B. S.	1823.	10. 2. G. b.	Jacq. ox. n. 82. t. 66.
78 strumosa. DC.	(wh.) swollen-styled.	—	1821.	— G. b.	— n. 79. t. 64.
79 marginata. DC.	(wh.) green-margin'd.	—	1812.	9. 12. G. b.	— n. 85. t. 68.
80 pulchella. DC.	(bh.) beautiful.	—	1795.	10. 11. G. b.	— n. 86. t. 69.
81 obtusa. DC.	(re.) blunt-leaved.	—	1812.	— G. b.	— n. 83. t. 79. f. 1.
82 lanata. DC.	(wh.) woolly-leaved.	—	1791.	— G. b.	— n. 81. t. 77. f. 2.
83 lutèola. DC.	(ye.) yellowish.	—	1822.	— G. b.	— n. 80. t. 65.
84 fállax. DC.	(ye.) related.	—	1825.	— G. b.	— n. 84. t. 67.
85 Acetosella. DC.	(wh.) common.	Britain.	4. 6. H. 4.	Eng. bot. 762.
86 americana. DC.	(wh.) American.	N. America.	— H. 4.	
87 tenella. DC.	(vi.) slender.	C. B. S.	1793.	— G. b.	Jacq. ox. n. 32. t. 19.
88 nátans. DC.	(wh.) floating.	—	1795.	9. 12. G. w.b.	— n. 78. t. 79. f. 2.
89 cruentata. DC.	(pu.) bloody.	—	1826.	— G. b.	Jacq. f. ecl. 1. t. 45.
90 filicaulis. DC.	(vi.) bilobed-leaved.	—	1815.	9. 10. G. b.	Jacq. schœnb. 2. t. 205.
91 bifida. DC.	(vi.) cloven-leaved.	—	1791.	— G. b.	Jacq. ox. n. 89. t. 79. f. 4.
92 bifurca. B.C.	(re.) forked-leaved.	—	1825.	6. 9. G. b.	Lodd. bot. cab. 1056.
93 cuneata. DC.	(wh.) wedge-leaved.	—	1818.	— G. b.	Jacq. ox. n. 55. t. 40.
94 cuneifolia. DC.	(wh.) wedge-shaped.	—	1793.	4. 5. G. b.	— n. 56. t. 41.
95 pusilla. DC.	(bh.) small.	—	1826.	— G. b.	— n. 57. t. 42.
96 linearis. DC.	(vi.) linear-leaved.	—	1795.	9. 11. G. b.	— n. 47. t. 32.
97 reclinata. DC.	(sc.) reclining.	—	—	— G. b.	— n. 49. t. 34.
98 grácilis. DC.	(re.) slender-stalked.	—	1818.	— G. b.	— n. 48. t. 33.
99 miniata. DC.	(re.) miniated.	—	1795.	5. 6. G. b.	— n. 50. t. 35.
100 glábra. DC.	(pu.) smooth.	—	1774.	1. 3. G. b.	— n. 58. t. 76. f. 3.
101 versicolor. DC.	(std.) striped-flower'd.	—	1791.	9. 10. G. b.	Bot. mag. 155.
102 elongata. DC.	(std.) elongated.	H 4			Jacq. ox. n. 52. t. 37.

OXALIDÆ.

103 am'ena. L.en. (ro.)	handsome.	C. B. S.	1791.	9. 10. G. b.	Jacq. schœnb. t. 206.
104 tenuifolia. DC. (std.)	fine-leaved.	_____	1790.	10. 11. G. b.	Lodd. bot. cab. t. 712.
105 polyphylla. DC. (ro.)	many-leaved.	_____	1791.	9. 1. G. b.	Jacq. ox. n. 59. t. 39.
106 filifolia. DC. (ro.)	slender-leaved.	_____	1818.	— G. b.	Jacq. schœn. 3. t. 273.
107 pentaphylla. DC. (li.)	five-leaved.	_____	1800.	11. 2. G. b.	Bot. mag. 1549.
108 lupinifolia. DC. (ye.)	Lupine-leaved.	_____	1791.	10. 11. G. b.	Jacq. ox. n. 92. t. 72.
109 flava. DC. (ye.)	narrow-leaved.	_____	1775.	3. 4. G. b.	Bot. reg. 117.
110 pectinata. DC. (ye.)	pectinated.	_____	1790.	9. 11. G. b.	Jacq. ox. n. 95. t. 75.
111 flabellifolia. DC. (ye.)	fan-leaved.	_____	1789.	— G. b.	— n. 94. t. 74.
112 tomentosa. DC. (wh.)	downy-leaved.	_____	1791.	4. 5. G. b.	— n. 96. t. 81.

ORDO LV.

ZYGOPHYLLEÆ. DC. prodr. 1. p. 703.

TRIBULUS. DC.	CALTROPS. Decandria Monogynia. L.				
1 cistoïdes. DC.	(ye.) Cistus-like.	S. America.	1752.	7. S. 4.	Bot. reg. 791.
2 terrestris. DC.	(ye.) small.	S. Europe.	1596.	6. 7. H. ♂.	Lam. ill. t. 346. f. 1.
3 subinermis. F.	(ye.) hairy.	Thibet.	1824.	6. 9. H. ♂.	
4 maximus. DC.	(st.) great.	Jamaica.	1728.	6. 7. S. ♂.	Jacq. ic. rar. t. 462.
5 trijugatus. DC.	(ye.) Nuttall's.	America.	1823.	— H. ♂.	
FAGONIA. DC.	FAGONIA. Decandria Monogynia. L.				
1 crética. DC.	(pu.) Cretan.	Candia.	1739.	6. 8. H. ♂.	Bot. mag. 241.
2 arábica. DC.	(pu.) Arabian.	Arabia.	1759.	— G. ♀.	
LA'RREA. DC.	LA'RREA. Decandria Monogynia.				
nítida. DC.	(ye.) glossy-leaved.	BuenosAyres.	1823.	6. 8. G. ♀.	Cav. ic. 6. t. 559.
ROEP'ERIA. J.R.	ROEP'ERIA. Decandria Monogynia.				
fabagifolia. J.R.	(ye.) Fabago-leaved.	N. Holland.	1822.	6. 8. G. ♀.	Juss. Rut. t. 2. f. 3.
ZYGOPHYLLUM. DC.	BEAN-CAPER. Decandria Monogynia. L.				
1 cordifolium. DC.	(ye.) heart-leaved.	C. B. S.	1774.	10. G. ♀.	
2 macrópterum. Led.	(ye.) large-winged.	Siberia.	1827.	7. 9. H. 4.	
3 Fabago. DC.	(st.) common.	Syria.	1596.	— H. 4.	Lam. ill. t. 345. f. 1.
4 fœtidum. DC.	(ye.) fœtid.	C. B. S.	1790.	6. 8. G. ♀.	Schr. s. han. p. 17. t. 9.
β insuáve. B.M.	(ye.) unpleasant.	_____	—	— G. ♀.	Bot. mag. 372.
5 maculatum. DC.	(ye.) spotted.	C. B. S.	1782.	10. 11. G. ♀.	
6 microphyllum. DC.	(ye.) small-leaved.	_____	1820.	— G. ♀.	Dill. elth. t. 116. f. 141.
7 Morgsana. DC.	(ye.) four-leaved.	_____	1732.	5. 9. G. ♀.	— t. 116. f. 142.
8 sessiliifolium. DC.	(ye.) sessile-leaved.	_____	1713.	7. 8. G. ♀.	
9 spinosum. DC.	(ye.) spiny.	_____	1818.	— G. ♀.	
10 coccineum. DC.	(sc.) scarlet.	Egypt.	—	— G. ♀.	Forsk. des. 87. ic. t. 11.
11 álbüm. DC.	(wh.) white.	_____	1770.	10. 11. G. ♀.	DC. pl. gras. t. 154.
12 prostratum. DC.	(ye.) hairy-jointed.	C. B. S.	1818.	— G. 4.	
13 tridentatum. DC.	(ye.) three-toothed.	Mexico.	1827.	9. 11. G. ♀.	Flor. mex. ic. ined.
GUAIACUM. DC.	LIGNUM VITÆ. Decandria Monogynia. L.				
1 officinale. DC.	(bl.) common.	W. Indies.	1694.	7. 9. S. ♀.	Lam. ill. t. 342.
2 verticale. DC.	(bl.) vertical-petal'd.	New Spain.	1823. S. ♀.	
3 arboreum. DC.	(ye.) tree.	S. America.	— S. ♀.	Jacq. amer. 130. t. 83.
Zygophyllum arboreum. Jac.					
PORLIE'RA. DC.	PORLIE'RA. Octandria Monogynia.				
hygrométrica. DC. (st.)	pinnate-leaved.	Peru.	1820. S. ♀.	Fl. per. v. 4. ined. t. 343.

§ 2. ZYGOPHYLLEÆ SPURIAE ALTERNIFOLIAE. DC.

MELIAN'THUS. DC.	HONEY-FLOWER. Didynamia Angiospermia. L.				
1 major. DC.	(br.) great.	C. B. S.	1688.	5. 7. G. ♀.	Bot. reg. 45.
2 minor. DC.	(br.) small.	_____	1696.	8. G. ♀.	Bot. mag. 301.
3 comosus. DC.	(br.) tufted.	_____	1818.	— G. ♀.	Comm. rar. t. 4.
BALANITES. DC.	BALANITES. Decandria Monogynia.				
ægyptiaca. DC.	(wh.) Egyptian.	Africa.	1822. S. ♀.	Del. fl. eg. t. 28. f. 1.
Ximènia ægyptiaca. L.					

ORDO LVI.

RUTACEÆ. DC. prodr. 1. p. 709.

Tribus I. DIOSMEÆ. DC.

RUTA. DC.	RUE. Decandria Monogynia. L.					
1 pinnata. DC.	(ye.) winged-leaved. Canaries.	1780.	3. 8.	G. h.	Bot. reg. 307.	
2 montana. DC.	(ye.) mountain. S. Europe.	1596.	8. 9.	H. Ȑ.	Jacq. ic. rar. 1. t. 76.	
3 graveolens. DC.	(ye.) common. —	1562.	6. 9.	H. h.	Duham. arb. 2. t. 61.	
4 bracteosa. DC.	(ye.) large-bracted. Sicily.	1823.	—	H. h.	Ten. fl. nap. 1. t. 36.	
5 divaricata. DC.	(ye.) spreading. Italy.	—	—	H. h.	Bot. mag. 2311.	
6 angustifolia. DC.	(ye.) narrow-leaved. S. France.	1722.	—	G. h.	2018.	
7 macrophylla. DC.	(ye.) long-leaved. Africa.	—	—	G. h.	—	
8 albidiflora. D.P.	(wh.) white-flowered. Nepal.	1820.	9. 10.	G. h.	Hook. ex. fl. t. 79.	
APLOPHYLLUM. J.R.	APLOPHYLLUM. Decandria Monogynia.					
1 patavina. J.R.	(ye.) Italian. Italy.	1820.	6. 9.	H. Ȑ.	Mich. gen. t. 19.	
2 pubescens. J.R.	(ye.) pubescent. Spain.	1816.	5. 8.	H. Ȑ.		
3 tuberculata. J.R.	(ye.) tubered. Egypt.	1824.	6. 8.	H. Ȑ.	Buxb. cent. 2. t. 28. f. 2.	
4 villosa. J.R.	(ye.) villous. Caucasus.	1828.	—	H. Ȑ.	Andr. rep. 565.	
5 linifolia. J.R.	(ye.) Flax leaved. Spain.	1752.	6. 9.	H. h.	Bot. mag. 2254.	
6 suaveolens. J.R.	(ye.) Primrose-scented. Tauria.	—	—	H. h.		
7 thesioides. F.	(ye.) Thesium-like. Caspia.	1828.	—	H. Ȑ.		
8 dahurica. J.R.	(st.) Milk-wort-ld. Dahuria.	1816.	7. 8.	H. Ȑ.	Gmel. sib. 4. t. 68. f. 2.	
Peganum dauricum. B. L.	Ruta dahurica. DC.					
PEGANUM. DC.	SYRIAN-RUE. Dodecadria Monogynia. L.					
Hármala. DC.	(wh.) common. Levant.	1570.	7. 8.	H. Ȑ.	Lam. ill. t. 401.	
DICTAMNUS. DC.	FRAXINELLA. Decandria Monogynia. L.					
1 Fraxinella. L.en. (re.) red-flowered. S Europe.	1596.	5. 7.	H. Ȑ.	Jacq. aust. 5. t. 428.		
2 álbus. L.en.	(wh.) white-flowered. —	—	—	H. Ȑ.		
CALODENDRON. DC.	CALODENDRON. Pentandria Monogynia.					
capense. DC.	(bh.) Cape. C. B. S.	1789.	—	G. h.	Lam. j. h. nat. 56. t. 3.	
ADENA'NDRA. W.en.	ADENA'NDRA. Pentandria Monogynia. L.					
1 uniflora. W.en. (wh.) one-flowered. C. B. S.	1775.	4. 7.	G. h.	Bot. mag. 273.		
2 acuminata.	(wh.) acute-leaved. —	1812.	—	G. h.	Lodd. bot. cab. 493.	
Diósma acuminata. B.C. non DC.						
3 am'ena.	(bh.) charming. —	1798.	—	G. h.	Bot. reg. 553.	
Diósma am'ena. B.C. 161.						
4 speciosa.	(wh.) umbel-flower'd. —	1790.	—	G. h.	Bot. mag. 1271.	
umbellata. W.en.	Diósma speciosa. B.M.					
5 frágans. R.S.	(re.) red-flowered. —	1812.	5. 7.	G. h.	Bot. mag. 1519.	
6 villosa. R.S.	(bh.) villous. —	—	—	G. h.		
Diósma villosa. DC. non Hort. tomentosa. Hort.						
7 marginata. R.S.	(bh.) marginate. —	1806.	4. 6.	G. h.		
8 linearis. B.D.	(wh.) linear-leaved. —	1800.	3. 7.	G. h.		
BAROSMA. W.en.	BAROSMA. Pentandria Monogynia. L.					
1 serratifolia. W.en. (wh.) saw-leaved. C. B. S.	1789.	3. 6.	G. h.	Bot. mag. 456.		
2 odorata. R.S.	(wh.) odoriferous. —	1790.	3. 5.	G. h.	Wend. coll. 1. t. 15.	
Diósma latifolia. B.C. 290.						
3 latifolia. R.S.	(wh.) broad-leaved. —	—	—	G. h.	Andr. rep. t. 33?	
4 crenata. B.D.	(wh.) crenated. —	1774.	1. 5.	G. h.	Lodd. bot. cab. t. 404.	
5 betulina. B.D.	(wh.) wedge-leaved. —	1790.	5. 7.	G. h.		
orbicularis. Hort.	Diósma betulina. DC.					
6 ovata. B.D.	(wh.) oval-leaved. —	—	—	2. 9. G. h.	Bot. mag. 1616.	
Diósma ovata. B.M.						
7 pulchella. B.D.	(pu.) blunt-leaved. —	1787.	2. 9.	G. h.	Bot. mag. 1357.	
Diósma pulchella. B.M.	Búcco pulchella. R.S.					
8 oblonga. B.D.	(wh.) oblong-leaved. —	1821.	4. 8.	G. h.		
9 dioica. B.D.	(pi.) dioecious. —	1816.	—	G. h.	Bot. reg. 502.	
Diósma dioica. B.R. linifolia. B.C. 400.						
10 angustifolia. B.D. (bh.) narrow-leaved. —	1820.	—	—	G. h.		
11 fetidissima. B.D. (bh.) strong-scented. —	1818.	—	—	G. h.		

MACRO'STYLIS. B.D.	MACRO'STYLIS.	Pentandria Monogynia.
1 lanceolata. B.D.	(li.) spear-leaved.	C. B. S. 1824. 4. 6. G. h.
Diósma barbata. s.s.		
2 barbigeria. B.D.	(li.) beard-bearing.	1826. — G. h.
3 obtusa. B.D.	(li.) blunt-leaved.	1824. — G. h.
4 squarroso. B.D.	(li.) squarrose.	1821. — G. h.
EUCH'ÆTIS. B.D.	EUCH'ÆTIS.	Pentandria Monogynia.
glomerata. B.D.	(wh.) close-flowered.	C. B. S. 1821. 4. 6. G. h.
ACMAD'ENIA. B.D.	ACMAD'ENIA.	Pentandria Monogynia.
1 juniperina. B.D.	(wh.) Juniper-leaved.	C. B. S. 1818. 4. 6. G. h.
2 obtusata. B.D.	(re.) blunt-leaved.	— G. h.
Diósma obtusata. R.S.		
3 laevigata. B.D.	(wh.) smooth.	— — G. h.
Diósma tetragona. Th. non L.		
4 pungens. B.D.	(wh.) pungent.	1823. — G. h.
5 tetragona. B.D.	(bh.) four-sided.	1720. 6. 8. G. h.
COLEON'EMA. B.D.	COLEON'EMA.	Pentandria Monogynia.
álba. B.D.	(wh.) white-flowered.	C. B. S. 1800. 3. 6. G. h.
Diósma álba. DC.	Adenándra álba. R.S.	
DIO'SMA. W.en.	DIO'SMA.	Pentandria Monogynia. L.
1 oppositifolia. DC.	(wh.) opposite-leav'd.	C. B. S. 1752. 3. 7. G. h.
2 succulenta. DC.	(wh.) succulent-l'd.	— 4. 6. G. h.
3 cupressina. DC.	(wh.) Cypress-like.	1790. 6. 7. G. h.
4 ramosissima. B.D.	(wh.) branching.	— 4. 8. G. h.
cypresina. Lam.		
5 Meyeriana. s.s.	(wh.) Meyer's.	1826. — G. h.
6 teretifolia. L.en.	(wh.) round-leaved.	1818. — G. h.
7 pectinata. DC.	(wh.) pectinated.	1812. 4. 6. G. h.
8 subulata. DC.	(wh.) awl-leaved.	— 3. 6. G. h.
9 rubra. DC.	(bh.) Heath-leaved.	1752. 2. 5. G. h.
ericifolia. Andr.	rep. t. 451.	
10 hirsuta. DC.	(wh.) hairy-leaved.	1731. — G. h.
11 ambigua. B.D.	(wh.) ambiguous.	1824. — G. h.
12 ericoides. DC.	(wh.) Heath-like.	1756. 3. 7. G. h.
13 longifolia. DC.	(wh.) long-leaved.	— — G. h.
AGATHO'SMA. W.en.	AGATHO'SMA.	Pentandria Monogynia.
1 hispida. s.s.	(wh.) rough-leaved.	C. B. S. 1786. 6. 8. G. h.
2 pátula. s.s.	(li.) spreading.	1821. — G. h.
3 brevifolia. s.s.	(li.) short-leaved.	1816. — G. h.
4 tenuissima. s.s.	(li.) slender-leaved.	— 2. 5. G. h.
5 virgata. s.s.	(wh.) twiggy.	1816. 4. 6. G. h.
6 erecta. B.D.	(wh.) upright.	— — G. h.
7 bruniades. s.s.	(li.) Brunia-like.	— 6. 8. G. h.
8 Thunbergiana. s.s. (li.)	Thunberg's.	— 4. 6. G. h.
Diósma ciliata. Th. non L.		
9 linifolia. s.s.	(li.) Flax-leaved.	1821. — G. h.
10 prolifera. s.s.	(wh.) proliferous.	— 4. 8. G. h.
11 ciliata. s.s.	(bh.) ciliated.	1774. — G. h.
12 Cerefolium. s.s.	(wh.) Chervil-scented.	1790. — G. h.
13 reflexa. L.en.	(li.) reflexed.	1816. 6. 8. G. h.
14 hirta. B.R.	(pu.) hairy.	1794. 5. 7. G. h.
Diósma Ventenatiána. s.s.	Búcco Ventenatiána. R.S.	
15 laxa. s.s.	(li.) loose-branched.	1816. — G. h.
16 villosa. W.en.	(li.) villous.	1786. 6. 8. G. h.
Búcco villosa. R.S.	Diósma Wendlandiána. DC.	
17 glabrata. B.D.	(li.) smooth.	1821. — G. h.
18 hybrida. s.s.	(li.) hybrid.	— — G. h.
19 rugosa. L.en.	(wh.) rugged-leaved.	1790. 4. 8. G. h.
20 mollis. B.D.	(wh.) soft.	1824. — G. h.
21 obtusa. DC.	(li.) blunt-leaved.	1774. — G. h.
Diósma ciliata. B.C. non L.		
α ovata.	(li.) oval-leaved.	— — G. h.
β oblóniga.	(li.) oblong-leaved.	— — G. h.
22 lanceolata.	(li.) lance-leaved.	— — G. h.
23 orbicularis. s.s.	(wh.) round-leaved.	1790. — G. h.
24 ambigua. DC.	(wh.) doubtful.	— — G. h.
25 imbricata. W.en.	(li.) imbricated.	1774. — G. h.
26 acuminata. W.en. (pa.)	taper-pointed.	1812. — G. h.
Diósma cordata. Hort.		
27 obtusata.	(li.) obtuse-leaved.	1816. — G. h.

B. et Wend. dios. t.A.f. 1.

Thunb. voy. 4. t. 5.

Wendl. coll. 1. t. 1.

Lodd. bot. cab. t. 303.

Wendl. coll. 1. t. 8.

Bot. reg. 563.

Wendl. coll. 1. t. 27.

Mill. ic. t. 124. f. 2.

Wendl. coll. 1. t. 19.

Wendl. coll. p. 17. t. 13.

Wendl. coll. 3. t. 77.

Bot. reg. 366.

Vent. malm. t. 93.

Bot. reg. 369.

Wendl. coll. 1. t. 2.

Lodd. bot. cab. 210.

Wendl. coll. 1. t. 13.

I. t. 14.

Bot. reg. 366.

Lodd. bot. cab. 461.

Wendl. coll. 1. t. 9.

I. t. 28.

Wendl. coll. 3. p. 7. t. 76.

EMPLÉ'URUM. DC.	EMPLÉ'URUM.	Monocia Tetrandria.				
serrulátum. DC.	(st.)	saw-leaved.	C. B. S.	1774.	6. 7.	G. h.
COR'RÉA. DC.	COR'RÉA.	Octandria Monogynia.				Sm. exot. bot. 2. t. 63.
1 álba. DC.	(wh.)	white.	N. S. W.	1793.	4. 7.	G. h.
2 rúfa. DC.	(wh.)	rusty-leaved.	_____	1819.	—	G. h.
3 pulchélla. s.F.A.	(pi.)	pretty.	N. Holland.	1824.	11. 5.	G. h.
4 speciòsa. DC.	(re.)	red-flowered.	N. S. W.	1806.	—	G. h.
5 virens. DC.	(gr.)	green-flowered.	_____	1800.	—	G. h.
PHEB'ALIUM. DC.	PHEB'ALIUM.	Decandria Monogynia.				
1 squammulósum. DC.	(wh.)	scaly.	N. S. W.	1820.	4. 6.	G. h.
2 elátum. F.T.	(wh.)	tall.	_____	1823.	4. 8.	G. h.
3 áureum. F.T.	(ye.)	golden.	_____	—	—	G. h.
4 lachnóides. F.T.	(wh.)	Lachnæa-like.	_____	—	—	G. h.
5 lineáre. ψ.	(wh.)	linear-leaved.	N. Holland.	1825.	—	G. h.
6 salicifolium. ψ.	(wh.)	Willow-leaved.	_____	—	—	G. h.
CR'OWEA. DC.	CR'OWEA.	Decandria Monogynia.				
saligna. DC.	(pi.)	Willow-leaved.	N. S. W.	1790.	7. 12.	G. h.
ERIO'STEMON. DC.	ERIO'STEMON.	Decandria Monogynia.				Bot. mag. 989.
1 buxifolium. DC.	(li.)	Box-leaved.	N. S. W.	1822.	4. 6.	G. h.
2 salicifolium. DC.	(re.)	Willow-leaved.	_____	—	—	G. h.
3 cuspidátum. F.T.(wh.)	sharp-pointed.	_____	1823.	4. 8.	G. h.	
4 obovále. F.T.	(li.)	oboval-leaved.	_____	—	—	G. h.
5 lanceolátum. DC.	(li.)	spear-leaved.	_____	—	—	G. h.
6 linearifolium. DC.	(wh.)	linear-leaved.	_____	—	—	G. h.
7 squámmum. DC.	(wh.)	scaly.	N. Holland.	1822.	—	G. h.
8 ericæfólum. M.C.	(li.)	Heath-leaved.	_____	1825.	—	G. h.
PHILOTH'ECA. L.T.	PHILOTH'ECA.	Monadadelphia Decandria.				
austrális. L.T.	(bh.)	Salsola-leaved.	N. S. W.	1822.	—	G. h.
BOR'ONIA. DC.	BOR'ONIA.	Octandria Monogynia.				Linn. trans. 11. t. 21.
1 pinnáta. DC.	(re.)	Hawthorn-scent'd.	N. S. W.	1794.	2. 5.	G. h.
2 tetrandría. DC.	(re.)	tetrandrous.	N. Holland.	1824.	—	G. h.
3 anemonæfólia. F.T.	(re.)	Anemone-leav'd.	N. S. W.	—	5. 8.	G. h.
4 aláta. DC.	(re.)	winged-leaved.	N. Holland.	—	4. 8.	G. h.
5 serrulátâ. DC.	(ro.)	Rose-scented.	N. S. W.	1816.	2. 5.	G. h.
6 denticulátâ. DC.	(re.)	tooth-leaved.	N. Holland.	1823.	4. 8.	G. h.
7 parviflóra. DC.	(re.)	small-flowered.	N. S. W.	1826.	—	G. h.
8 polygalæfólia. DC.	(re.)	Polygala-ld.	_____	1824.	—	G. h.
9 ledifólia. DC.	(re.)	Ledum-leaved.	_____	—	—	G. h.
CYMINO'SMA. DC.	CYMINO'SMA.	Octandria Monogynia.				
1 pedunculátâ. DC.	(wh.)	peduncled.	E. Indies.	1800.	—	G. h.
Jambolífera pedunculátâ. w.						Vahl. symb. 3. t. 61.
2 odorátâ. DC.	(wh.)	sweet-scented.	China.	1818.	—	G. h.
ZIE'RIA. DC.	ZIE'RIA.	Tetrandria Monogynia.				
1 lanceolátâ. DC.	(wh.)	spear-leaved.	N. S. W.	1808.	4. 7.	G. h.
Smithii. Bot. rep. 606.						Bot. mag. 1395.
2 macrophylla. DC.	(wh.)	large-leaved.	_____	1820.	—	G. h.
3 levigátâ. DC.	(wh.)	smooth-leaved.	_____	1822.	—	G. h.
4 octándra.	(gr.)	octandrous.	_____	1825.	4. 6.	G. h.
5 obcordata. F.T.	(wh.)	obcordate.	_____	1824.	—	G. h.
6 revolutâ. F.T.	(wh.)	revolute.	_____	—	3. 5.	G. h.
7 micropýlla. DC.	(wh.)	small-leaved.	_____	—	—	G. h.
8 pilósâ. DC.	(wh.)	hairy.	_____	—	—	G. h.
9 hirsútâ. DC.	(wh.)	hirsute.	_____	—	—	G. h.
MELIC'OPE. DC.	MELIC'OPE.	Octandria Monogynia.				
ternátâ. DC.	ternate-leaved.	N. Zealand.	1822.	—	G. h.
EV'ODIA. DC.	EV'ODIA.	Tetrandria Monogynia.				Lam. ill. t. 294.
tríphylla. DC.	(wh.)	three-leaved.	Pulo Penang.	1820.	—	S. h.
Fagàra triphylla. F.I.						Rumph. amb. 2. t. 62.
ZANTHO'XYLUM. DC.	ZANTHO'XYLUM.	Tetrandria Monogynia et Diœcia Pentandria.				
1 tragôdes. DC.	(wh.)	prickly-leaved.	St. Domingo.	1759.	8. 9.	S. h.
Fagàra tragôdes. w.						Jacq. amer. 21. t. 14.
2 affine. K.S.	(wh.)	related.	Mexico.	1826.	—	G. h.
3 Pteróta. DC.	(wh.)	Lentiscus-ld.	W. Indies.	1768.	—	S. h.
Fagàra lentiscifólia.	W. en.					Br. jam. 146. t. 5. f. 1.
4 piperitum. DC.	(wh.)	Ash-leaved.	Japan.	1773.	9.	G. h.
5 Ávicénnae. DC.	(wh.)	short-panicled.	China.	1823.	—	G. h.
6 heterophylum. DC.	(wh.)	various-leav'd.	Bourbon.	—	—	S. h.
7 emarginátum. DC.	(wh.)	notch-leaved.	Jamaica.	1739.	—	S. h.
8 acuminátum. DC.	(wh.)	taper-pointed.	_____	1818.	—	S. h.
						Sloan. j. 2. t. 168. f. 4.

RUTACEÆ.

9 spinòsum. DC.	(wh.) spiny.	Jamaica.	1824.	S. h.	
10 tricárpum. DC.	(gr.) Ash-leaved.	N. America.	1806.	H. h.	Cat. car. 1. t. 26.
11 fraxineum. DC.	(gr.) common.	—	1759.	3. 4.	H. h.	Duham. arb. 1. t. 97.
12 mítæ. DC.	(gr.) soft-leaved.	—	1812.	—	H. h.	
13 hermaphrodítum. DC.	(wh.) hermaphrodite.	Guiana.	1823.	S. h.	Aubl. gui. 1. t. 30.
14 juglandifólium. DC.	(wh.) Walnut-leaved.	W. Indies.	1822.	S. h.	Pluk. t. 239. f. 6.
15 ClavaHérculis. L.	(wh.) clubbed.	—	1739.	4. 5.	S. h.	— t. 239. f. 4.
16 aromáticum. DC.	(wh.) aromatic.	St. Domingo.	1823.	S. h.	Jacq. f. ecl. 1. t. 70.
17 armátum. DC.	(wh.) armed.	E. Indies.	1816.	S. h.	
18 nítidum. DC.	(wh.) shining-leaved.	China.	1822.	2. 3.	G. h.	Bot. mag. 2558.
19 sapindóides. DC.	(wh.) Licca-tree.	Jamaica.	G. h.	Br. jam. 207. t. 20. f. 2.
<i>Sapindus spinòsus. L.</i>						
SPIRANTH'ERA. DC. SPIRANTH'ERA. Pentandria Monogynia.						
odoratissima. DC.	(bh.) Jasmine-scent'd.	Brazil.	1823.	S. h.	N. et Mart. ac. b. xi. t. 31.
<i>Terpnánthus jasminiodorus. Nees.</i>						

Tribus II. CUSPARIEÆ. DC.

MONNIE'RA. DC.	MONNIE'RA. Monadelphia Diandria.					
trífolia. DC.	(wh.) three-leaved.	Guiana.	1792.	7. 8.	S. ⊖.	Aub. gui. 2. t. 293.
TIC'OREA. DC.	TICO'REA. Pentandria Monogynia.					
1 fe'tida. DC.	(wh.) strong-scented.	Guiana.	1825.	S. h.	Aubl. gui. 2. t. 277.
2 jasminiflòra. DC.	(wh.) Jasmine-fl'd.	Brazil.	1827.	S. h.	N. et Man. ac. b. 11. t. 18.
GALIP'EÀ. DC.	GALIP'EÀ. Diandria Monogynia.					
trifoliàta. DC.	(gr.) three-leaved.	Guiana.	1803.	S. h.	Aub. gui. 2. t. 269.

ORDO LVII.

SIMARUBEÆ. DC. prodr. 1. p. 733.

QUA'SSIA. DC.	QUA'SSIA. Decandria Monogynia.	L.				
amára. DC.	(re.) bitter.	Guiana.	1790.	6. 7.	S. h.	Bot. mag. 497.
SIMAR'UBA. DC.	SIMAR'UBA. Monœcia Decandria.					
1 officinális. DC.	(wh.) officinal.	Guiana.	1787.	S. h.	Aub. gui. 2. t. 331-2.
Quássia Simarùba. L.	Simarùba amára. Aub.					
2 gláuca. DC.	(wh.) glaucous-leav'd.	Cuba.	1824.	S. h.	
3 excélsa. DC.	(wh.) tall.	Jamaica.	1822.	S. h.	Sw. act. holm. 1788. t. 8.
Quássia excélsa. Sw.						
SIM'ABA. DC.	SIM'ABA. Decandria Monogynia.					
guianén sis. DC.	(wh.) Guiana.	Guiana.	1822.	S. h.	Aub. gui. 1. t. 153.
Zwingera amára. W.						

ORDO LVIII.

OCHNACEÆ. DC. prodr. 1. p. 735.

O'CHNA. DC.	O'CHNA. Polyandria Monogynia.	L.				
1 squarrósa. R.C.	(ye.) blunt-leaved.	E. Indies.	1790.	7. 8.	S. h.	Roxb. cor. 1. t. 89.
2 lúcida. DC.	(pu.) acute-leaved.	—	1818.	—	S. h.	Lam. ill. t. 472. f. 1.
3 nítida. DC.	(wh.) shining.	C. B. S.	1816.	S. h.	DC. ann. m. 17. n. 3. t. 2.
4 multiflòra. DC.	(ye.) many-flowered.	SierraLeon.	1822.	S. h.	— n. 4. t. 3.
5 mauritiána. DC.	(ye.) Mauritus.	Mauritus.	—	S. h.	— n. 8. t. 5.
6 atropurpurea. DC.	(ye.) purple-calyxed.	C. B. S.	1816.	S. h.	Pluk. alm. t. 263. f. 1. 2.

GO'MPHIA. DC.	Go'MPHIA.	Decandria Monogynia. W.				
1 obtusifolia. DC. (ye.)	blunt-leaved.	Madagascar. 1803.	S. h.	DC. ann. mus. n. 4. t. 8.	
<i>Ochna lavigata</i> . Vahl.						
2 guianensis. DC. (ye.)	Guiana.	Guiana. 1825.	S. h.	DC. ann. mus. n. 7. t. 9.	
3 Jabotapita. DC. (ye.)	spear-leaved.	W. Indies. 1818.	S. h.	Lam. ill. 472. f. 2.	
4 nítida. DC. (ye.)	glossy-leaved.	—	1803.	DC. an. mus. n. 13. t. 13.	
5 laurifolia. DC. (ye.)	laurel-leaved.	—	1822.	— n. 15. t. 15.	
6 mexicana. DC. (ye.)	Mexican.	Mexico. 1825.	G. h.	H. et B. pl. æq. 2. t. 74.	
WALK'ERA. DC.	WALK'ERA.	Pentandria Monogynia.				
serrata. DC. (ye.)	saw-leaved.	E. Indies. 1824.	S. h.	Rheed. mal. 5. t. 48.	

ORDO LIX.

CORIARIEÆ. DC. prodr. 1. p. 739.

CORIARIA. DC.	CORIARIA.	Mono-Dicecia Decandria.			
1 myrtifolia. DC. (gr.)	Myrtle-leaved.	S. Europe. 1629.	5. 8.	H. h.	Duham. arb. 1. t. 73.
2 sarmentosa. DC. (gr.)	sarmentose.	N. Zealand. 1820.	7. 9.	G. ¼.	Bot. mag. 2470.

** SUBCLASSIS II. CALYCIFLORÆ.

ORDO LX.

CELASTRINEÆ. Brown gen. rem. p. 22.

Tribus I. STAPHYLEACEÆ. DC. prodr. 2. p. 2.

STAPHYLEA. DC.	BLADDER-NUT.	Pentandria Trigynia. L.			
1 trifolia. DC. (wh.)	three-leaved.	N. America. 1640.	5. 6.	H. h.	Schmidt. arb. t. 80.
2 Bumálda. DC. (wh.)	Japan.	Japan. 1804.	6. 9.	G. h.	
<i>Bumálda trifolia</i> . Th.					
3 piñata. DC. (wh.)	five-leaved.	England.	4. 6.	H. h.	Eng. bot. 1560.
TURPINIA. DC.	TURPINIA.	Polygamia Dicecia. DC.			
paniculata. DC. (wh.)	panicled.	W. Indies. 1826.	S. h.	Vent. choix. t. 31.
<i>Dalrymplea domingensis</i> , s.s.					
DALRY'MPLEA. R.C. DALRY'MPLEA.	Pentandria Monogynia. F.I.				
pomifera. r.c. (wh.)	Apple-bearing.	E. Indies. 1823.	S. h.	Roxb. corom. 3. t. 279.
<i>Turpinia pomifera</i> . DC.					

Tribus II. EUONYMEÆ. DC.

EUONYMUS. DC.	SPINDLE-TREE.	Pentandria Monogynia. L.			
1 japonicus. DC. (wh.)	Japan.	Japan. 1804.	6. 8.	G. h.	Kämpf. ic. t. 8.
2 europaeus. DC. (wh.)	common.	Britain.	5. 7.	H. h.	Eng. bot. 362.
3 verrucosus. DC. (pu.)	warted.	Austria. 1763.	5. 6.	H. h.	Schmidt arb. t. 72.
4 latifolius. DC. (gr.)	broad-leaved.	S. Europe. 1730.	5. 7.	H. h.	— t. 74.
5 atropurpureus. DC. (d.pu.)	dark-purple.	N. America. 1756.	—	H. h.	Jacq. vind. 2. t. 120.
6 americanus. DC. (st.)	evergreen.	—	1683.	6. 7.	Schmidt arb. t. 75.
7 sarmentosus. N.	(st.) sarmentose.	—	1823.	—	
8 obovatus. N.	(gr.) obovate-leaved.	—	—	H. h.	

CELASTRINEÆ.

9 angustifolius. DC. (st.) narrow-leaved.	Georgia.	1806.	—	H. $\frac{1}{2}$.
10 lúcidus. D.P. (wh.) shining-leaved.	Nepaul.	1820.	6. 7.	H. $\frac{1}{2}$.
11 micránthus. D.P. (wh.) small-flowered.	—	—	—	H. $\frac{1}{2}$.
12 chinénsis. DC. (wh.) Chinese.	China.	1822.	5. 6.	H. $\frac{1}{2}$.
13 lácerus. D.P. (wh.) torn.	Nepaul.	1820.	6. 7.	H. $\frac{1}{2}$.
14 grandiflórus. F.I. (wh.) large-flowered.	—	1823.	—	H. $\frac{1}{2}$.
15 echinátus. F.I. (st.) spinous-fruited.	—	—	4. 5.	H. $\frac{1}{2}$. Bot. mag. 2767.
CELASTRUS. DC. STAFF-TREE.	Pentandria Monogynia. L.			
1 oleóides. DC. (wh.) Olive-leaved.	C. B. S.	1823.	5. 6.	G. $\frac{1}{2}$.
2 pterocárpus. DC. (wh.) wing-capsuled.	—	1816.	5. 8.	G. $\frac{1}{2}$. Burm. afr. t. 97. f. 1.
3 rostratus. DC. (wh.) beaked.	—	1821.	—	G. $\frac{1}{2}$.
4 tricuspidátus. DC. (wh.) three-pointed.	—	1816.	5. 6.	G. $\frac{1}{2}$. Buchoz. dec. 6. t. 3.
Cassine lævigatæ. Lam. 'Olea capensis'. Buchoz.				
5 lúcidus. DC. (wh.) shining.	C. B. S.	1722.	4. 9.	G. $\frac{1}{2}$. L'Her. stirp. 1. t. 25.
6 macrocárpus. DC. (wh.) large-capsuled.	Peru.	1826.	—	Flor. per. 3. t. 230. f. 16.
7 bullátus. DC. (wh.) scarlet-fruited.	Virginia.	1759.	7.	H. $\frac{1}{2}$. Pluk. alm. t. 28. f. 5.
8 scándens. DC. (wh.) climbing.	N. America.	1736.	5. 6.	H. $\frac{1}{2}$. Duham. arb. 1. t. 95.
9 punctátus. DC. (wh.) spotted-stalked.	Japan.	1816.	—	G. $\frac{1}{2}$.
10 nútans. F.I. (wh.) drooping-fld.	E. Indies.	1824.	—	S. $\frac{1}{2}$. L'Her. sert. ang. t. 10.
11 cassinoídes. DC. (wh.) crenated.	Canaries.	1779.	8. 9.	G. $\frac{1}{2}$.
12 lycioídes. DC. (wh.) Box-thorn-like.	—	1821.	—	G. $\frac{1}{2}$.
13 cérruus. DC. (wh.) drooping.	C. B. S.	1815.	5. 6.	G. $\frac{1}{2}$.
14 myrtifólius. DC. (wh.) Myrtle-leaved.	Jamaica.	—	6. 8.	S. $\frac{1}{2}$. Sloan. jam. 2. t. 193. f. 1.
15 tetragónus. DC. (wh.) four-sided.	C. B. S.	1816.	—	G. $\frac{1}{2}$.
16 ilicifólius. DC. (wh.) Holly-leaved.	Brazil.	1824.	—	S. $\frac{1}{2}$.
17 ilicinus. DC. (wh.) Ilex-like.	C. B. S.	1816.	5. 8.	G. $\frac{1}{2}$.
18 retúsus. DC. (wh.) notched.	Peru.	1823.	—	G. $\frac{1}{2}$. Flor. per. 3. t. 229. f. a.
emarginátus. F.P. non W.				
19 linearis. DC. (wh.) linear-leaved.	C. B. S.	1823.	5. 8.	G. $\frac{1}{2}$.
20 emarginátus. DC. (st.) emarginate.	E. Indies.	1820.	—	S. $\frac{1}{2}$. Bot. mag. 2114.
21 buxifólius. DC. (wh.) Box-leaved.	C. B. S.	1752.	5. 8.	G. $\frac{1}{2}$. — t. 2070.
22 cymósus. B.M. (wh.) cyme-flowered.	—	1815.	6. 8.	G. $\frac{1}{2}$.
23 multiflórus. DC. (wh.) many-flowered.	—	1816.	5. 8.	G. $\frac{1}{2}$.
24 pyracánthus. DC. (wh.) red-fruited.	—	1752.	—	G. $\frac{1}{2}$. Bot. mag. 1167.
25 flexuósus. DC. (wh.) flexuose.	—	1816.	—	G. $\frac{1}{2}$.
26 montánuus. DC. (wh.) mountain.	E. Indies.	1824.	—	S. $\frac{1}{2}$.
PLECTR'ONIA. W. PLECTR'ONIA.	Pentandria Monogynia.			
ventosa. w. (wh.) corymbed.	C. B. S.	1816.	—	G. $\frac{1}{2}$. Burm. afr. t. 94 ?
corymbosa. P.S. Celástrus? Plectrónia. DC.				
MA'YENUS. DC. MAYENUS.	Pentandria Monogynia.			
1 octogónus. DC. (gr.) angular-leaved.	Peru.	1786.	10. 11.	G. $\frac{1}{2}$.
Celástrus octogónus. w. Senácia octogóna. Lam.				
2 chilénsis. DC. (re.) Chile.	Chile.	1824.	—	G. $\frac{1}{2}$. Feuill. obs. 3. t. 27.
3 verticillátus. DC. (gr.) whorled.	Peru.	1823.	9. 11.	G. $\frac{1}{2}$. Flor. per. 3. t. 229. B.
ELÆODE'NDRON. DC. OLIVE-WOOD.	Pentandria Monogynia.			
1 orientális. DC. (wh.) taper-leaved.	Madagascar.	1771.	—	S. $\frac{1}{2}$. Jacq. ic. 1. t. 48.
índicum. Gært. fruct. 1. t. 57. Rubéntia olivina. J.				
2 gláucum. DC. (st.) glaucous.	Ceylon.	1823.	—	S. $\frac{1}{2}$. Rottb. n.a.hav. 2.t.4.f.1.
3 xylocárpum. DC. (wh.) leathery-leaved.	Antilles.	1816.	7. 8.	S. $\frac{1}{2}$. Vent. choix. t. 23.
Cassine xylocarpa. Vent.				
4 austrális. DC. (wh.) thick-leaved.	N. S. W.	1796.	6. 8.	G. $\frac{1}{2}$. Vent. malm. t. 117.
Portenschlägia australis. Tratt. arch. t. 250. Lamárckia dentata. Hort.				
5 integrifolium. (wh.) entire-leaved.	N. S. W.	—	—	G. $\frac{1}{2}$. Tratt. arch. t. 284.
Portenschlägia integrifolia. Tratt.				
6? cròceum. DC. (wh.) Cape Holly.	C. B. S.	1794.	5. 6.	G. $\frac{1}{2}$. Pet. Th. veg. afr. 1. t. 2.
Ilex cròcea. W.				
PTELI'DIUM. DC. PTELI'DIUM.	Tetrandria Monogynia.			
ovátum. DC. (gr.) oval-leaved.	Madagascar.	1825.	—	S. $\frac{1}{2}$. Pet. Th. veg. afr. 1. t. 2.

Tribus III. AQUIFOLIACEÆ. DC. prod. 2. p. 11.

CASSINE. DC.	CASSINE.	Pentandria Trigynia. L.		
1 Maurocénia. DC. (wh.) Hottentot-cherry.	C. B. S.	1690.	7. 8.	G. $\frac{1}{2}$. Dill. elt. t. 121. f. 147.
2 capénsis. DC. (wh.) Cape Phillyrea.	—	1629.	—	G. $\frac{1}{2}$. — t. 236. f. 303.
3 Colpoón. w. (wh.) Colpoon-tree.	—	1791.	—	G. $\frac{1}{2}$. Burm. afr. t. 86.

4	bárbara. R.S.	(wh.) quadrangular.	C. B. S.	1816.	—	G. h.	
5	æthiopica. R.S.	(wh.) Ethiopian.	_____	—	G. h.	
	oppositifolia. DC.	(wh.) opposite-leaved.	_____	—	S. h.	
7	excelsa. F.I.	(wh.) tall.	Nepaul.	1823.	G. h.	
	HARTOGIA. DC.	HARTOGIA.	Tetrandria Monogynia.				
	capensis. DC.	(wh.) Cape.	C. B. S.	1816.	G. h.	Thunb. n. gen. 5. p. 55. ic.
	Schrébera schinoides.	Th. prodr. t. 2.	<i>Elæodendron schinoides</i> .	s.s.			
	CURTISIA. DC.	CURTISIA.	Tetrandria Monogynia.				
	faginea. DC.	(wh.) Hassagay-tree.	C. B. S.	1775.	G. h.	Burm. afr. 235. t. 82.
	MYGINDA. DC.	MYGINDA.	Tetrandria Tetragynia.				
1	Uragoga. DC.	(pu.) saw-leaved.	S. America.	1790.	8. 9.	S. h.	Jacq. amer. 24. t. 16.
2	myrtifolia. DC.	(wh.) Myrtle-leaved.	N. America.	1818.	5. 8.	H. h.	
	<i>Ilex myrsinifolia</i> . Ph.						
3	Rhácoma. DC.	(re.) blunt-leaved.	Jamaica.	1798.	S. h.	Jacq. ic. 2. t. 311.
4	latifolia. DC.	(wh.) broad-leaved.	W. Indies.	1795.	4. 5.	S. h.	
	ILEX. DC.	HOLLY.	Tetrandria Tetragynia. L.				
1	Aquifoliūm. DC.	(wh.) common.	Britain.	4. 6.	H. h.	Eng. bot. 496.
β	heterophylla.	(wh.) various-leaved.	_____	—	H. h.	
γ	crassifolia.	(wh.) thick-leaved.	_____	—	H. h.	
δ	ferox.	(wh.) hedgehog.	_____	—	H. h.	
ε	echinata.	(wh.) striped hedgehog.	_____	—	H. h.	
ζ	flava.	(wh.) yellow-berried.	_____	—	H. h.	
η	senescent.	(wh.) spineless.	_____	—	H. h.	
δ	álbo-marginata.	(w.) white-edged.	_____	—	H. h.	
ι	áureo-marginata.	(w.) gold-edged.	_____	—	H. h.	
κ	picta.	(wh.) painted.	_____	—	H. h.	
2	recúrva. L.en.	(wh.) recurved-spined.	_____	—	H. h.	
3	baleàrica. DC.	(wh.) Minorca.	Minorca.	1815.	5. 6.	H. h.	Lodd. bot. cab. 549.
4	Perádo. DC.	(bh.) thick-leaved.	Madeira.	1760.	4. 5.	F. h.	Duham. ed. nov. 1. t. 2.
β	obtusa. DC.	(bh.) blunt-leaved.	_____	—	F. h.	
5	canariensis. DC.	(wh.) Canary.	Canaries.	1818.	5. 6.	F. h.	
6	opaca. DC.	(wh.) Carolina.	Carolina.	1744.	—	F. h.	Wats. dend. brit. t. 3.
7	laxiflora. DC.	(wh.) loose-flowered.	_____	1811.	—	F. h.	
8	chinensis. DC.	(wh.) Chinese.	China.	1810.	4. 5.	F. h.	Bot. mag. 2043.
9	Dahoón. DC.	(wh.) Dahoo.	Carolina.	1726.	5. 6.	F. h.	Willd. hort. ber. 1. t. 31.
β	laurifolia. DC.	(wh.) large-leaved.	_____	—	F. h.	Wats. dend. brit. t. 114.
10	Cassinæ. DC.	(wh.) Cassine-like.	_____	1726.	8.	F. h.	Catesb. car. 1. t. 31.
11	angustifolia. DC.	(wh.) narrow-leaved.	N. America.	1806.	5. 6.	F. h.	Wats. dend. brit. t. 4.
12	vomitaria. DC.	(wh.) South-sea Tea.	Florida.	1700.	6. 7.	F. h.	Jacq. ic. 2. t. 310.
13	latifolia. DC.	(wh.) broad-leaved.	Japan.	1824.	F. h.	
14	serrata. DC.	(wh.) saw-leaved.	_____	—	F. h.	
15	myrtifolia. DC.	(wh.) Myrtle-leaved.	W. Indies.	1806.	7. 8.	S. h.	
16	salicifolia. DC.	(wh.) Willow-leaved.	Mauritius.	1818.	4. 5.	S. h.	Jacq. col. 5. t. 2. f. 2.
	<i>Bulgaria lúcidula</i> . Wendl.						
17	Macocouá. DC.	(wh.) acuminate.	Trinidad.	1826.	S. h.	Aubl. gui. 1. t. 34.
18	paraguariensis. DC.	(w.) Paraguay Tea.	Brazil.	1823.	S. h.	Lamb. pin. ic.
	PRINOS. DC.	WINTER BERRY.	Hexandria Monogynia. L.				
1	deciduus. DC.	(wh.) deciduous.	N. America.	1760.	6. 7.	H. h.	Wats. dend. brit. 115.
	<i>Ilex prionoides</i> . R.S.						
2	ambiguus. DC.	(wh.) Carolina.	_____	1812.	7. 8.	H. h.	Wats. dend. t. 29.
3	verticillatus. DC.	(wh.) whorled.	_____	1736.	—	H. h.	_____ t. 30.
	<i>padifolius</i> . W. en.						
4	lævigatus. DC.	(wh.) smooth.	_____	1812.	—	H. h.	Wats. dend. t. 28.
5	lanceolatus. DC.	(wh.) scarlet berried.	_____	1811.	6. 7.	H. h.	
6	prunifolius. DF.	(wh.) Plum-leaved.	_____	1820.	—	H. h.	
7	gláber. DC.	(wh.) ever-green.	_____	1759.	7. 8.	H. h.	Wats. dend. brit. t. 27.
8	atomarius. DC.	(wh.) atomiferous.	Georgia.	1820.	6. 7.	H. h.	
9	coriaceus. DC.	(wh.) leathery-leaved.	_____	—	—	H. h.	
10	lúcidus. DC.	(wh.) shining-leaved.	_____	1778.	—	F. h.	
	NEMOPANTHES. DC.	NEMOPANTHES.	Polygamia Diœcia.				
	canadensis. DC.	(wh.) Canadian.	N. America.	1812.	6. 7.	H. h.	DC. pl. rar. gen. t. 3.
	<i>Ilex canadensis</i> . Mich. fl. amer. 2. t. 49.						

ORDO LXI.

RHAMNEÆ. DC. prodr. 2. p. 19.

ZI'ZYPHUS. R.S.	ZI'ZYPHUS. Pentandria Monogynia. R.S.						
1 Lòtus. R.S. (st.) Barbary.	Africa. 1731. G. h.						Desf. act. ac. 1788. t. 21.
2 mucronàta. W.en.(st.) mucronate.	C. B. S. 1820. G. h.						
<i>bubalina</i> . R.S.							
3 sinénsis. R.S. (wh.) Chinese.	China. 1818. 5. 6. H. h.						
4 Oenóplia. R.S. (st.) oblique-leaved.	Ceylon. S. h.						Burm. zeyl. 131. t. 61.
5 Jujùba. R.S. (st.) blunt-leaved.	E. Indies. 1759. 4. 5. S. h.						Rheed. mal. 4. t. 41.
6 soròria. R.S. (ye.) sister.	— — — 1823. S. h.						
7 álbens. F.I. (wh.) white-fruited.	China. 1822. G. h.						
8 Spina Christi. R.S.(st.) Christ's-thorn.	Egypt. 4. 5. G. h.						
9 Napéca. R.S. (wh.) spotted-fruited.	Ceylon. 1816. S. h.						Pluk. alm. t. 197. f. 3.
10 vulgáris. R.S. (ye.) common.	S. Europe. 1640. 8. 9. H. h.						—— t. 216. f. 6.
11 nítida. F.I. (ye.) shining.	China. 1822. G. h.						Pall. ross. 2. t. 59.
12 Xylop'yrus. R.S. (ye.) cork-barked.	E. Indies. — — — 1820. S. h.						
13 Caracúta. F.I. (ye.) villous-branch'd.	— — — 1820. S. h.						
14 microphylla. F.I. (ye.) small-leaved.	— — — 1824. S. h.						Pluk. alm. t. 197. f. 2.
15 incurva. F.I. (gr.) incurred.	Nepaul. 1820. 5. 6. F. h.						
16 flexuosa. F.I. (ye.) flexuose.	— — — 1820. F. h.						
PALITURUS. DC.	CHRIST'S-THORN. Pentandria Trigynia.						
1 aculeátus. DC. (ye.) common.	S. Europe. 1596. 6. 7. H. h.						Bot. mag. 1893.
<i>australis</i> . R.S. <i>vulgáris</i> . D.P. <i>Zizyphus Paliturus</i> . B.M.							
2 virgátus. DC. (ye.) twiggy.	Nepaul. 1819. 8. 9. H. h.						Bot. mag. t. 2535.
COND'ALIA.	COND'ALIA. Pentandria Monogynia.						
<i>microphylla</i> . DC. (gr.) small-leaved.	Chili. 1824. G. h.						Cav. ic. 6. t. 525.
BERCH'EMIA. DC.	BERCH'EMIA. Pentandria Monogynia.						
1 volubilis. DC. (gr.) climbing.	Carolina. 1714. 6. 7. H. h. — Jacq. ic. 2. t. 336.						
<i>Rhámnus volubilis</i> . L. <i>Zizyphus volubilis</i> . w. <i>Oenoplia volubilis</i> . R.S.							
2 lineáta. DC. (wh.) lined.	China. 1804. G. h.						Osb. it. 219. t. 7.
3 floribunda. Bro. (wh.) many-flowered.	Nepaul. 1827. G. h. — Brong. m. ram. pl. 2.f.1.						
SAGER'ETIA. Bro.	SAGER'ETIA. Pentandria Monogynia.						
1 theézans. Bro. (gr.) Tea-like.	China. 9. 12. G. h.						
<i>Rhámnus theézans</i> . L.							
2 hamòsa. Bro. (gr.) hook-thorned.	Nepaul. 1826. G. h. —						
<i>Zizyphus hamòsa</i> . F.I.							
RHAMNUS. DC.	BUCKTHORN. Pentandria Monogynia. L.						
1 Alatérnus. DC. (ye.) Alaternus.	S. Europe. 1629. 4. 6. H. h.						Duham. ed. nov. 3.t.14.
<i>a latifolius</i> .	(ye.) broad-leaved.	— —	H. h.			
<i>β balearicus</i> . DC.	(ye.) round-leaved.	— —	H. h.			
<i>γ hispánicus</i> . DC.	(ye.) Spanish.	— —	H. h.			
<i>δ vulgáris</i> . DC.	(ye.) oval saw-leaved.	— —	H. h.			
<i>ε glaber</i> .	(ye.) smooth-leaved.	— —	H. h.			
<i>ζ integrifolius</i> .	(ye.) entire-leaved.	— —	H. h.			
<i>η maculatus</i> .	(ye.) spot-leaved.	— —	H. h.			
<i>δ aureus</i> .	(ye.) gold-striped.	— —	H. h.			
<i>i argénteus</i> .	(ye.) silver-striped.	— —	H. h.			
2 Clusi. W.en.	(gr.) narrow-leaved.	1629.	— —	H. h.			
3 hybridus. DC.	(gr.) hybrid.	Hybrid.	5. 6.	H. h.			Clus. hist. 1. p. 50.
4 glandulòsa. DC.	(gr.) Madeira.	Canaries.	1785. 6. 7.	G. h.			L'Her. sert. t. 5.
5 integrifolius. DC.	(gr.) Teneriffe.		1820.	— —	G. h.		Vent. malm. t. 34.
<i>coriaceus</i> . Nees.							Nees hor. phys. t. 22.
6 longifolius. DC.	(gr.) long-leaved.	N. America?	1822. 5. 6.	H. h.			
7 primoídes. DC.	(gr.) Prinos-leaved.	C. B. S.	1778. 8. 9.	G. h.			L'Her. sert. ang. t. 9.
8 celtifolius. DC.	(gr.) Celtis-leaved.		— —	G. h.		Burm. afr. 242. t. 88.
9 cathárticus. DC.	(gr.) purging.	England.	5. 6.	H. h.		Eng. bot. 1629.
10 virgátus. DC.	(ye.) slender.	Nepaul.	1819. 4. 5.	H. h.			
11 tinctoriús. DC.	(st.) dying.	Hungary.	1823. 5. 6.	H. h.			Pl. rar. hung. 3. t. 255.
12 infectoriús. DC.	(ye.) yellow-berried.	S. Europe.	1683. 6. 7.	H. h.			Clus. hist. 1. p. 111. ic.
13 saxátilis. DC.	(gr.) rock.		1752. 5. 6.	H. h.			Jacq. aust. 1. t. 53.
14 oleoídes. DC.	(gr.) Olive-leaved.	Spain.	— —	6. 7.	H. h.		
15 buxifolius. DC.	(gr.) Box-leaved.	S. Europe.	1820. — —	H. h.			
16 pubéscens. DC.	(gr.) pubescent.	S. France.	1820. 6. 7.	H. h.			
17 crenulátus. DC.	(gr.) crenulate.	Teneriffe.	1778. 3. 4.	G. h.			

18	lycioïdes. DC.	(gr.) Box-thorn-like. Spain.	1752.	9. 12. H. h.	Cavan. ic. 2. t. 182.
	β latifolius.	(gr.) broader-leaved.	—	— H. h.	Pall. ross. 2. t. 62.
19	erythróxylon. DC.	(ye.) red-wooded.	Siberia.	1822. 4. 5. H. h.	H. et B. n. gen. a. 7. t. 616.
20	microphyllus. DC.	(ye.) small-leaved.	Mexico.	— G. h.	Scop. carn. ed. 2. t. 5.
21	pùmulus. L.	(bh.) dwarf.	Europe.	1752. 7. H. h.	
	β rupéstris. Vill.	(bh.) small.	Dauphiny.	— H. h.	
22	pusillus. T.P.	(gr.) little.	Naples.	1825. — H. h.	
23	valentinius. W.	(ye.) Spanish.	Spain.	— 6. 7. H. h.	Cavan. ic. 2. t. 181.
	pùmulus. c. non W.				
24	Wulfénii. s.s.	(ye.) Wulfen's.	Tergestum.	— — H. h.	Jacq. coll. 2. t. 11.
	pùmulus. Wulf. non L.				
25	dahuricus. DC.	(gr.) Dahurian.	Dahuria.	1826. H. h.	Pall. ross. 2. t. 61.
26	alnifòlius. DC.	(gr.) Alder-leaved.	N. America.	1778. 5. 6. H. h.	
27	franguloides. R.S.	(gr.) Frangula-like.	—	— H. h.	
28	Purshianus. DC.	(gr.) Pursh's.	—	1826. — H. h.	
	alnifòlius. Ph. nec aliorum.				
29	alpinus. DC.	(ye. pu.) Alpine.	Switzerland.	1752. — H. h.	Lodd. bot. cab. 1077.
30	carolinianus. DC.	(gr.) Carolina.	N. America.	1824. 5. 7. H. h.	Eng. bot. 250.
31	Frangula. DC.	(gr.) berry-bearing	Alder. Britain. 4. 5. H. h.	Wats. dend. brit. t. 11.
32	latifolius. DC.	(ye.) broad-leaved.	Azores.	1778. 6. 7. H. h.	
33	lanceolatus. Ph.	(gr.) spear-leaved.	N. America.	1812. 4. 5. H. h.	
34?	mystacinus. R.S.	(gr.) wiry.	Abyssinia.	1775. 11. S. h.	Th. act. gor. 1812. c. ic.
35?	tetragonus. DC.	(gr.) square-branch'd.	C. B. S.	1816. 6. 8. G. h.	
SC'UTIA. Bro.	SC'UTIA.	Pentandria Monogynia.			
1	índica. Bro.	(wh.) Indian.	E. Indies.	1824. S. h.	Gært. fruct. 2. t. 106.
	RhámnuS circumscissus. L.	Ceanòthus circumscissus. DC.			
RETANILLA. Bro.	RETANILLA.	Pentandria Monogynia.			
obcordata. Bro.	(ye.) obcordate.	ld.	Peru.	1822. 5. 7. G. h.	Vent. hort. cels. t. 92.
Colléttia obcordata. V.					
COLLE'TIA. Bro.	COLLE'TIA.	Pentandria Monogynia.			
1	spinosa. DC.	(ye.) spiny.	Peru.	1823. G. h.	Lam. ill. 2. t. 129.
2	serratifolia. DC.	(ye.) saw-leaved.	—	5. 8. G. h.	Vent. choix. t. 15.
3	pubescens. Bro.	(ye.) pubescent.	N. Holland.	1824. G. h.	
HOVENIA. DC.	HOVENIA.	Pentandria Monogynia.			
1	dúlcis. DC.	(gr.) smooth-leaved.	China.	1812. 4. 6. G. h.	Lam. ill. t. 131.
	acerba. Bot. reg. 501.				
2	inæquális. DC.	(gr.) pubescent-ld.	Nepaul.	1823. G. h.	
	dulcis. D.P. et F.I. nec aliòrum.				
COLUBRINA. Bro.	COLUBR'INA.	Pentandria Monogynia.			
1	ferrugínosa. Bro.	(st.) Bahama red-wood.	Bahama.	1726. 6. S. h.	Jacq. vind. 3. t. 50.
	RhámnuS colubrinus. L.	Ceanòthus colubrinus. DC.			
2	reclinata. Bro.	(wh.) oval-leaved.	Jamaica.	1758. 8. S. h.	Brown jam. t. 20. f. 2.
	Ceanòthus reclinatus. DC.	RhámnuS ellípticus. H.K.			
3	triflora. Bro.	(st.) three-flowered.	Mexico.	1826. G. h.	
4	cubensis. Bro.	(st.) Cuba.	Cuba.	1824. S. h.	Jacq. vind. 3. t. 49.
5	asiatica. Bro.	(st.) Asiatic.	Asia.	1691. 7. 8. S. h.	Cavan. ic. 5. t. 440. f. 1.
	Ceanòthus asiáticus. DC.				
CEANOTHUS. Bro.	CEAN'OTHUS.	Pentandria Monogynia. L.			
1	azáreus. DC.	(bl.) blue.	New Spain.	1818. 4. 5. G. h.	Bot. reg. 291.
	carvulus. Lod. bot. cab. 110.	bicolor.	R.S.		
2	tardiflorus. DC.	(wh.) late-flowering.	N. America.	1812. 9. 10. H. h.	
3	americanus. DC.	(wh.) New Jersey Tea.	—	1713. 7. 8. H. h.	Bot. mag. 1479.
4	ovatus. DC.	(wh.) smooth-leaved.	—	— H. h.	
5	perennis. DC.	(wh.) perennial.	—	1822. 6. 7. H. h.	
6	intermedius. DC.	(wh.) intermediate.	—	1812. — H. h.	
7	collinus. D.	(wh.) hill.	—	1827. — H. h.	
8	sanguineus. DC.	(wh.) bloody twigg'd.	—	1812. 5. 7. H. h.	
9	microphyllus. DC.	(wh.) small-leaved.	—	1806. 6. 7. H. h.	
10	buxifolius. DC.	(wh.) Box-leaved.	Mexico.	1824. 4. 6. G. h.	H. et B. n. gen. a. 7. t. 615.
11	macrocarpus. DC.	(st.) large-fruited.	New Spain.	1823. 6. 7. G. h.	Cavan. ic. 3. t. 276.
12	nepalensis. F.I.	(st.) Nepal.	Nepaul.	— H. h.	
WILLEM'ETIA. Bro.	WILLEM'ETIA.	Pentandria Monogynia.			
africana. Bro.	(st.) African.	C. B. S.	1712. 3. 4. G. h.	Pluk. phyt. 126. f. 1.	
	Ceanòthus africanaus. DC.				
POMADE'RRI'S. DC.	POMADE'RRI'S.	Pentandria Monogynia.			
1	globulosa. Bro.	(st.) round-headed.	N. Holland.	1803. 4. 5. G. h.	Lab. nov. hol. 1. t. 85.
	Ceanòthus globulus. Lab.				
2	spathulata. Bro.	(st.) spathulate-leav'd.	—	1821. — G. h.	Lab. nov. 1. t. 84.
3	discolor. DC.	(ye.) two-coloured.	—	1803. — G. h.	Swt. fl. aust. t. 41.
	Ceanòthus discolor. Vent. malm. t. 58.		I		

4 intermédia. DC. (st.) intermediate.	N. S. W.	1818.	4. 6.	G. h.	
5 acuminàta. L.en.(wh.) loose-flowered.	N. Holland.	1816.	4. 7.	G. h.	
6 ellíptica. DC. (ye.) elliptic-leaved.	_____	1805.	—	G. h.	Lab. nov. holl. 1. t. 86.
7 lanigera. (ye.) ferruginous. <i>Ceanothus laniger</i> . A.R.	_____	1806.	—	G. h.	Andr. rep. 569.
8 Símisia. (st.) Sims's. <i>lanigera</i> . Sims bot. mag. non Andr.	N. S. W.	1821.	—	G. h.	Bot. mag. 1823.
9 viridirùfa. Sieb. (st.) brown green.	N. S. W.	_____	—	G. h.	
10 ledifòlia. F.T. (ye.) Ledum-leaved.	_____	1824.	—	G. h.	
11 andromedæfòlia. F.T.(ye.) Andromeda-ld	_____	—	—	G. h.	
12 phylliræoïdes. DC.(ye.) Phillyrea-leaved.	_____	1803.	—	G. h.	Lab. nov. holl. 1. t. 87.
13 apétala. DC. (br.) petalless.	N. Holland.	_____	5. 6.	G. h.	
14 áspera. DC. (br.) rough.	N. S. W.	1818.	6. 7.	G. h.	
15 ligustrina. DC. (wh.) Privet-like.	_____	1826.	—	G. h.	Lodd. bot. cab. 120.
16 phylicifòlia. DC. (wh.) Phylica-leaved.	N. Holland.	1810.	3. 5.	G. h.	
CRYPTA'NDRA. DC. CRYPTAN'DRA. Pentandria Monogynia. R.S.					
1 ericifòlia. DC. (pa.) Heath-leaved.	N. S. W.	1823.	5. 7.	G. h.	Linn. trans. 10 t. 18. f. 1.
2 amára. DC. (pa.) bitter.	_____	—	4. 6.	G. h.	— 10. t. 18. f. 2.
3 spinescens. DC. (pa.) spiny.	_____	1824.	—	G. h.	
VENTIL'AGO. DC. VENTIL'AGO. Pentandria Monogynia. R.S.					
maderaspatâna. DC.(gr.) Madras.	E. Indies.	1822.	—	S. h. — Roxb. corom. 1. t. 76.	
GOUA'NIA. DC. GOUA'NIA. Polygamia Monœcia. L.					
1 integrifòlia. DC. entire-leaved.	S. America?	1824.	—	S. h. —	
2 domingénsis. DC. (gr.) Chaw-stick.	W. Indies.	1739.	—	S. h. — Jacq. amer. t. 179. f. 40.	
3 cordifòlia. DC. (ye.) heart-leaved.	Brazil.	1822.	—	S. h. —	
4 tomentòsa. DC. (ye.) woolly-leaved.	W. Indies.	1823.	—	S. h. — Gært.f. carp. 3.t.183.f.1	
5 mauritiâna. DC. (br.) rusty.	Mauritius.	—	—	S. h. —	
6 tiliaeñolia. DC. (gr.) Lime-leaved.	Bourbon.	1810.	4. 6.	S. h. —	
7 leptostàchya. DC. (gr.) slender-spiked.	E. Indies.	—	—	S. h. — Roxb. corom. 1. t. 98. <i>tiliaeñolia</i> . R.C. non Lam.	
8 Retinaria. DC. (gr.) oval-leaved.	Mauritius.	1824.	—	S. h. — Gært.fruct.2.t.120.f.4.	
TRICHOCE'PHALUS. Bro. TRICOCE'PHALUS. Pentandria Monogynia.					
1 stipularis. Bro. (wh.) stipuled.	C. B. S.	1786.	5. 9.	G. h.	Wendl. coll. t. 32.
<i>Phy'lica stipularis</i> . DC.					
2 spicata. Bro. (wh.) spiked.	—	—	1774. 11. 12.	G. h.	Lam. ill. t. 127. f. 3.
<i>Phy'lica spicata</i> . L. non B.M.					
PHY'LICA. Bro. PHY'LICA. Pentandria Monogynia. L.					
1 parvifòra. R.S. (wh.) small-flower'd.	C. B. S.	1790.	4. 7.	G. h.	
2 ericoides. R.S. (wh.) Heath-leaved.	—	—	9. 4.	G. h.	Bot. mag. 224.
3 glábrata. R.S. (wh.) smooth-leaved.	—	1820.	3. 5.	G. h.	
4 aceròsa. R.S. (wh.) needle-leaved.	—	1819.	4. 7.	G. h.	Spr.ber.mag.8.t.8 f.2.
5 lanceolâta. R.S. (wh.) spear-leaved.	—	1790.	4. 5.	G. h.	
6 eriophora. R.S. (wh.) pale-flowered.	—	1774.	11. 3.	G. h.	
7 calliosa. R.S. (wh.) callous-leaved.	—	—	3. 4.	G. h.	
8 imberbis. R.S. (wh.) beardless.	—	1823.	5. 9.	G. h.	Breyn. cent. 18. t. 7.
9 secúnda. R.S. (wh.) secund.	—	1812.	—	G. h.	
10? australis. L.en. (wh.) southern.	—	—	3. 5.	G. h.	
11 pùmila. DC. (wh.) dwarf.	—	1820.	4. 6.	G. h.	
12 bicolor. R.S. (wh.) two-coloured.	—	1823.	4. 5.	G. h.	
13 spicata. B.M. (wh.) close-spiked.	—	—	8. 12.	G. h.	Bot. mag. 2704.
14 hirsuta. R.S. (wh.) hairy.	—	1820.	5. 9.	G. h.	
15 pinea. R.S. (wh.) Pine-like.	—	1824.	—	G. h.	
16 villosa. R.S. (wh.) villous.	—	1790.	—	G. h.	
17 papillosa. R.S. (wh.) papillose.	—	1820.	3. 5.	G. h.	
18 rosmarinifòlia. R.S.(w.) Rosemary-ld.	—	1815.	5. 8.	G. h.	Lodd. bot. cab 849.
19 cylindrica. R.S. (wh.) cylindrical.	—	—	4. 8.	G. h.	Wendl. coll. 1. t. 7.
20 exélsa. R.S. (wh.) tall.	—	1820.	3. 5.	G. h.	— 3. t. 4.
21 horizontális. DC. (wh.) spreading. <i>plumosa</i> . Spr. non L.	—	1800.	6. 8.	G. h.	Spr. ber. mag. 8. f. 7.
22 squarròsa. DC. (wh.) squarrose.	—	1820.	3. 5.	G. h.	Lodd. bot. cab. 36.
23 Commelinì. R.S. (wh.) Commelin's.	—	1800.	5. 8.	G. h.	Comm. pral. 63. t. 13.
24 plumòsa. R.S. (wh.) feathered.	—	1752.	3. 5.	G. h.	Lam. ill. t. 127. f. 4.
25 capitâta. R.S. (wh.) headed.	—	1774.	—	G. h.	Bot. reg. 711.
26 reclinâta. R.S. (wh.) reclined.	—	1820.	—	G. h.	Wendl. coll. 2. t. 56.
27 pedicellâta. DC. (wh.) pedicled.	—	—	—	G. h.	
28 atráta. R.S. (wh.) dark-haired.	—	—	3. 5.	G. h.	
SOULANGIA. Bro. SOULANGIA. Pentandria Monogynia.					
1 axillaris. Bro. (wh.) axillary-flower'd.	C. B. S.	1812.	5. 6.	G. h.	Spr.ber. mag.8.t.8. f.4.
2 thymifòlia. Bro. (wh.) Thyme-leaved.	Antarctic.	1824.	—	S. h.	Vent. malm. t. 57.

3 oleæfòlia. Bro. (wh.) Olive-leaved.	C. B. S.	1. 4. G. h.	Lodd. bot. cab. t. 323.
<i>Phyllica spicata</i> . Lod. bot. cab. nec aliorum.			
4 paniculata. Bro. (wh.) panicled.	— — —	1812. 4. 6. G. h.	
<i>Phyllica paniculata</i> . DC. <i>myrtifolia</i> . Poir.			
5 buxiæfòlia. Bro. (wh.) Box-leaved.	— — —	1759. 5. 9. G. h.	Lodd. bot. cab. 848.
<i>Phyllica buxifolia</i> . Wend. coll. 1. t. 26.			
6 cordata. Bro. (wh.) heart-leaved.	— — —	1789. 5. 6. G. h.	Comm. præl. 62. t. 12.
7 orientalis. (wh.) oriental.	— — —	1820. — — G. h.	Herb. amat. t. 283.
<i>Phyllica orientalis</i> . Herb. amat. t. 283.			

† *Genera ad Rhamneas accedentia.***SCHÆFF'ERIA. DC. SCHÆFF'ERIA. Dioecia Tetrandria.**

frutescens. DC. (wh.) white-flowered.	W. Indies.	1793.	S. h.	Swartz.fl.ind.1.t.7.f.A.
β buxifolia. DC. (wh.) Box-leaved.	— — —	— — —	S. h.	Sloan.hist.2.t.209.f.1.
OL'NIA. DC. OL'NIA. Pentandria Monogynia.				
cymosa. DC. (wh.) sweet-scented.	C. B. S.	1812. 4. 6. G. h.		
<i>Sideroxylum cymosum</i> . L.				

ORDO LXII.

BRUNIACEÆ. DC. prodr. 2. p. 43.

BERZ'ELIA. Bro. BERZ'ELIA. Pentandria Monogynia.

1 abrotanoides. Bro. (w.) Southernwood-like.	C. B. S.	1787. 5. 7. G. h.	Wendl. coll. t. 45.
<i>Brunia abrotanoides</i> . Lodd. bot. cab. 355.			
2 lanuginosa. Bro. (wh.) woolly.	— — —	1774. 6. 8. G. h.	Wendl. coll. 1. t. 11.

BR'UNIA. R.S. BR'UNIA. Pentandria Monogynia. L.

1 l'ævis. R.S. (wh.) smooth.	C. B. S.	1823. 5. 7. G. h.	
2 nodiflora. R.S. (wh.) imbricated.	— — —	1786. 7. 8. G. h.	Breyn. cent. 22. t. 10.
3 deusta. R.S. (wh.) black-tipped.	— — —	1804. 6. 8. G. h.	
4 comosa. R.S. (wh.) tufted.	— — —	1823. — — G. h.	
5 verticillata. R.S. (wh.) whorled.	— — —	1794. — — G. h.	
6 virgata. Bro. (wh.) twiggy.	— — —	— — — G. h.	
7 squarrosa. R.S. (wh.) squarrose.	— — —	1818. — — G. h.	
8 alopecuroides. R.S. (w.) Fox-tail.	— — —	1816. 5. 7. G. h.	
9 laxa. R.S. (wh.) loose-branched.	— — —	— — — G. h.	
10 plumosa. R.S. (wh.) feathered.	— — —	1822. — — G. h.	
11 superba. R.S. (wh.) superb.	— — —	1791. — — G. h.	
12 fragarioides. R.S. (wh.) Strawberry-like.	— — —	1794. 5. 7. G. h.	
13 ciliata. R.S. (wh.) fringed.	— — —	1812. — — G. h.	
14 ericoides. R.S. (wh.) Heath-leaved.	— — —	— — — G. h.	Wendl. coll. II. t. 57.
15 arachnoidea. R.S. (wh.) cobweb.	— — —	1810. — — G. h.	t. 62.
16 formosa. R.S. (wh.) handsome.	— — —	1812. — — G. h.	
17 macrocephala. s.s. (w.) large-headed.	— — —	— — — G. h.	
18 racemosa. Bro. (wh.) clustered.	— — —	1790. 5. 9. G. h.	
<i>Phyllica racemosa</i> . L.			
19 pinifolia. Bro. (wh.) Pine-leaved.	— — —	1789. 5. 6. G. h.	Brong. brun. t. 1. f. 2.
<i>Phyllica pinifolia</i> . L.			
20 imbricata. (wh.) imbricated.	— — —	1801. 8. 11. G. h.	
<i>Phyllica imbricata</i> . Th.			

RASP'ALIA. Bro. RASP'ALIA. Pentandria Digynia.

microphylla. Bro. (w.) small-leaved.	C. B. S.	1804. 6. 8. G. h.	Brong. brun. t. 3. f. 1.
<i>Brunia microphylla</i> . Th.			

STA'AVIA. DC. STA'AVIA. Pentandria Monogynia.

1 radiata. DC. (wh.) rayed.	C. B. S.	1787. 5. 6. G. h.	Brong. brun. t. 2. f. 2.
2 glutinosa. DC. (wh.) clammy.	— — —	1793. 4. 5. G. h.	Lodd. bot. cab. 852.

3 nuda. Bro.	(wh.) naked.	C. B. S.	1821.	4. 5. G. h.	
4 ciliata. Bro.	(wh.) fringed.		1812.	5. 7. G. h.	
NEB'ELIA. Neck.	NEB'ELIA.	Pentandria Digynia.			
1 paleacea.	(wh.) chaffy.	C. B. S.	1791.	6. 8. G. h.	Wendl. coll. t. 21.
Berárdia. Bro. non p.s. <i>Brùnia paleacea</i> . R.S.					
2 affinis.	(wh.) related.	C. B. S.	1816.	— G. h.	
Berárdia affinis. Bro.					
3 phylloides.	(wh.) Phylica-leaved.		1822.	— G. h.	
Berárdia phylloides. Bro. <i>Brùnia phylloides</i> . Th.					
LINC'ONIA. L.	LINC'ONIA.	Pentandria Digynia.			
1 alopecuroidea. L. (wh.) Fox-tail.	C. B. S.	1824.	6. 8. G. h.	Swa.in.ber.m.1810.t.4.	
2 thymifolia. DC. (wh.) Thyme-leaved.		1825.	— G. h.	— p. 86. t. 4.	
3 cuspidata. DC. (wh.) cuspidate.			— G. h.	— 1811.p.284.t.7.f.1.	
AUDÓUINIA. Bro.	AUDÓUINIA.	Pentandria Monogynia.			
capitata. Fro. (wh.) headed.	C. B. S.	1790.	4. 6. G. h.		
Diósma capitata. DC.					
TITTMANNA. Bro.	TITTMANNA.	Pentandria Monogynia.			
lateriflora. Bro. (wh.) side-flowering.	C. B. S.	1812.	4. 6. G. h.	Brong. brun. t. 4. f. 2.	
THA'MNEA. Bro.	THA'MNEA.	Pentandria Monogynia.			
uniflora. Bro. (wh.) one-flowered.	C. B. S.	1810.	4. 6. G. h.	Brong. brun. t. 4. f. 3.	

ORDO LXIII.

SAMYDEÆ. DC. prodr. 2. p. 47.

SAM'YDA. DC.	SAM'YDA.	Decandria Monogynia. S.S.			
1 glábrata. DC.	(wh.) smooth.	Jamaica.	1818.	5. 8. S. h.	
2 villósa. DC.	(wh.) villous.		—	— S. h.	
3 decúrrens. s.s.	(wh.) decurrent.	Brazil.	1822.	— S. h.	
4 pubéscens. s.s.	(wh.) pubescent.	W. Indies.	1793.	— S. h.	
5 serrulata. DC.	(wh.) saw-leaved.		1723.	7. 8. S. h.	Jacq. col. 2. t. 17. f. 1.
6 spinulosa. DC.	(wh.) spiny.	St. Thomas.	1826.	... S. h.	Vent. choix. t. 43.
7 macrocarpa. DC.	(wh.) large-fruited.	Mexico.	—	... S. h.	
8 rósea. B.M.	(ró.) rose-coloured.	W. Indies.	1793.	6. 7. S. h.	Bot. mag. 550.
9 serrulata. A.R. 202. nec aliorum.					
9 nitida. DC.	(wh.) glossy-leaved.	Jamaica.	—	— S. h.	Br. jam. 217. t. 23. f. 3.
10 macrophylla. DC. (gr.)	large-leaved.	E. Indies.	1823.	... S. h.	
CASE'ARIA. DC.	CASE'ARIA.	Decandria Monogynia. S.S.			
1 glomerata. DC.	(wh.) cluster-flower'd. E. Indies.		1824.	5. 7. S. h.	
2 ovata. DC.	(gr.) oval-leaved.		—	— S. h.	Rheed. mal. 4. t. 49.
3 ramiflora. DC.	(wh.) branch-flower'd.	Guiana.	—	— S. h.	Aubl. guian. I. t. 127.
4 nitida. DC.	(wh.) shining.	Carthagena.	—	— S. h.	Act. helv. 8. p. 58. ic.
5 sylvétris. DC.	(wh.) wood.	Jamaica.	1823.	... S. h.	
6 parviflora. DC.	(wh.) small-flowered	W. Indies.	1818.	5. 7. S. h.	Sloan.hist.2.t.211.f.2.
7 serrulata. DC.	(wh.) serrulate.		—	— S. h.	
8 glábra. H.B.	(wh.) smooth.	E. Indies.	1820.	— S. h.	
9 hirsuta. DC.	(wh.) hairy.	W. Indies.	—	... S. h.	
10 zizyphoides. K.S. (wh.)	Zizyphus-like,	Caracas.	1824.	... S. h.	
11 arguta. K.S.	sharp-toothed.	Mexico.	—	... S. h.	

ORDO LXIV.

HOMALINEÆ. DC. prodr. 2. p. 53.

HOM'ALIUM. DC.	HOM'ALIUM.	Polyadelphia Polyandria.			
racemosum. DC. (wh.) racemed.		W. Indies.	1818.	5. 7. S. h.	Bot. reg. 519.
BLACKWELLIA. DC.	BLACKWELLIA.	Dodecandria Pentagynia.			
1 paniculata. DC. (wh.) paniced.		Bourbon.	1818.	5. 7. S. h.	

HOMALINEÆ.

117

2 glauca. DC.	(wh.) glaucous.	Mauritius.	1823.	S. h.	Vent. choix. t. 55.
3 nepalensis. DC.	(wh.) Nepaul.	Nepaul.	1824.	S. h.	
4 axillaris. DC.	(wh.) axillary-flower'd.	Madagascar.	1823.	S. h.	Lam. ill. t. 412. f. 1.
5 spiralis. DC.	(wh.) spiral.	Pegu.	1820.	S. h.	
6 fagifolia.	(wh.) Beech-leaved.	China.	1824.	5.	G. h.	
ASTRA'NTHUS. DC.	STAR-FLOWER.	Heptandria Tetragynia.				
cochinchinensis. (wh.)	Chinese.	China.	1821.	6. 7.	S. h.	Bot. mag. 2659.
ARISTOTELIA. DC.	ARISTOTE'LIA.	Dodecadandra Monogynia. L.				
Méqui. DC.	(wh.) shining-leaved.	Chile.	1773.	4. 5.	H. h.	Wats. dend. brit. t. 44.

ORDO LXV.

CHAILLETIACEÆ. *DC. prodr. 2. p. 57.*

CHAILL'ETIA. DC.	CHAILL'ETIA.	Pentandria Di-Trigynia.
1 toxicaria. D.D.	(wh.) poisonous.	Sierra Leone. 1822. S. h.
2 erecta. D.D.	(wh.) upright.	_____

ORDO LXVI.

AQUILARINEÆ. DC. prodr. 2: p. 59.

AQUIL'ARIA. DC. **AQUIL'ARIA.** Decandria Monogynia.
Agallòcha. H.B. Aloe-wood. E. Indies. 1824. S. L.

ORDO LXVIII.

CASSUVIEÆ. *R. Brown cong. p. 12.*

ANACARDIUM. DC.	CASHEW-NUT.	Enneandria Monogynia. L.	
1 occidentale. DC. (y.gr.)	common.	India.	1599. S. h. Catesb. car. 3. t. 9.
<i>Cassuvium pomiferum.</i> Lam. ill. 3. t. 322.			
2 dūbium. H.B.	(gr.) doubtful.	E. Indies.	1824. S. h.
SEMECA'RPUS. DC.	SEMECA'RPUS.	Pentandria Trigynia. L.	
Anacardium. DC. (y.gr.) common.		E. Indies.	1820. S. h.
α <i>angustifolium.</i> DC. (y.gr.) long-leaved.	_____	_____	S. h. Rumph. amb. 1. t. 70.
β <i>cuneifolium.</i> DC. (y.gr.) wedge-leaved.	_____	_____	S. h.
γ <i>obtusifolium.</i> DC. (y.gr.) broad-leaved.	_____	_____	S. h. Roxb. corom. 1. t. 12.
MANGIFERA. DC.	MANGO-TREE.	Pentandria Monogynia.	
1 indica. DC.	(ye.) common.	E. Indies.	1690. 6. 9. S. h. Andr. rep. 425.
2 oppositifolia. F.I.	(ye.) opposite-leaved.	_____	1823. S. h.
BUCHANANIA. R.C.	BUCHANANIA.	Decandria Monogynia.	
1 latifolia. DC.	(wh.) broad-leaved.	E. Indies.	1820. S. h.
2 angustifolia. DC.	(wh.) narrow-leaved.	_____ S. h. Roxb. corom. 3. t. 262
<i>Mangifera axillaris.</i> Lam. Cambessedea, K.S.			
RHUS. R.S.	SUMACH.	Pentandria Trigynia. L.	
1 Coriaria. R.S.	(gr.) Elm-leaved.	S. Europe.	1596. 7. H. h. Flor. græc. t. 290.
2 typhina. R.S.	(pu.) Virginian.	N. America.	1629. 7. 8. H. h. Wats. dend. t. 17. mas
α <i>arborescens.</i> R.S. (pu.) arborescent.	_____	_____	H. h. t. 18. fem
β <i>fruticosa.</i> R.S. (pu.) frutescent.	_____	_____	H. h.

CASSUVIEÆ.

3 javánica. R.S.	(wh.) Java.	Java.	1799.	7. 9.	S. h.	
4 semialata. R.S.	(wh.) Service-leaved.	Macao.	1780.	—	G. h.	Murr. com. got. vi. t. 3.
5 viridiflora. R.S.	(gr.) green-flowered.	N. America.	—	7.	H. h.	
6 glábra. R.S.	(gr.) smooth.	—	1726.	7. 9.	H. h.	Wats. dend. t. 15.
<i>a hermaphrodita.</i> DC. (gr.) <i>hermaphrodite.</i>	—	—	—	—	H. h.	Dill. elt. t. 243.
β dioica. DC.	(gr.) dioecious.	—	—	—	H. h.	Lam. ill. t. 207. f. 1.
7 élégans. R.S.	(sc.) scarlet-flowered.	—	—	—	H. h.	Wats. dend. t. 16.
8 Vérmix. R.S.	(gr.) varnish.	—	1713.	—	H. h.	— t. 19.
9 vernicifera. DC.	(gr.) Chinese varnish.	China.	—	—	G. h.	Kæmp. amoen. t. 792.
Rhus Vernix. Th. non Ph.						
10 A'mela. D.P.	(gr.) large-leaved.	Nepaul.	1823.	—	G. h.	
11 Wallichii.	(gr.) Wallich's.	—	—	—	G. h.	
<i>juglandifolia.</i> D.P. non Kth.						
12 fraxinifolia. D.P.	(gr.) Ash-leaved.	—	—	—	G. h.	
13 acuminata. DC.	(gr.) pointed-leaved.	—	—	—	G. h.	
14 succedanea. R.S.	(gr.) red Lac.	China.	1768.	6.	G. h.	Kæmp. amoen. t. 795.
15 juglandifolia. K.S.	(gr.) Walnut-leaved.	S. America.	1826.	—	G. h.	H. et B. n. ge. a. 7. t. 603-4.
16 copallina. R.S.	(gr.) Lentiscus-lea'd.	N. America.	1688.	8. 9.	H. h.	Jacq. schœn. 3. t. 341.
17 alata. R.S.	(gr.) winged-leaved.	C. B. S.	1818.	—	G. h.	
18 pauciflora. R.S.	(gr.) few-flowered.	—	—	—	G. h.	
19 pumila. R.S.	(gr.) dwarf poisonous.	N. America.	1806.	7.	H. h.	
20 digitata. R.S.	(gr.) fingered.	C. B. S.	1818.	—	G. h.	
21 obliqua. R.S.	(gr.) oblique-leaved.	—	—	—	G. h.	
22 schinoides. R.S.	(gr.) Schinus-like.	Brazil.	1825.	—	S. h.	
23 pentaphylla. R.S. (ye.)	various-leaved.	Barbary.	1816.	—	G. h.	Desf. atl. 1. t. 77.
24 chinensis. Mill.	(gr.) Chinese.	China.	1740.	—	G. h.	
25 râdicans. DC.	(wh.) rooting poison-oak.	N. America.	1724.	6. 7.	H. h.	Bot. mag. 1806.
26 Toxicodendron. DC. (w.) poison-oak.	—	—	1640.	—	H. h.	Schkuhr. hand. 1. t. 82.
27 cirrhiflora. R.S.	(gr.) tendril-flower'd.	C. B. S.	1825.	—	G. h.	
28 Thunbergiana. R.S. (gr.)	Thunberg's.	—	1697.	7. 8.	G. h.	
29 glauca. R.S.	(gr.) glaucous.	—	1818.	—	G. h.	
30 tridentata. R.S.	(gr.) three-toothed.	—	—	—	G. h.	
31 crenata. R.S.	(gr.) crenated.	—	—	6. 7.	G. h.	
32 dentata. R.S.	(gr.) rough-stalked.	—	1798.	—	G. h.	
33 sinuata. R.S.	(gr.) sinuated.	—	1818.	—	G. h.	
34 cuneifolia. R.S.	(gr.) wedge-leaved.	—	1816.	—	G. h.	
35 dimidiata. R.S.	(gr.) variable-leaved.	—	—	—	G. h.	
36 dissécta. R.S.	(gr.) dissected-leaved.	—	—	—	G. h.	
37 incisa. R.S.	(gr.) cut-leaved.	—	1789.	—	G. h.	
38 tomentosa. R.S.	(gr.) woolly-leaved.	—	1691.	—	G. h.	Comm. hort. ams. 1. t. 92.
39 albida. R.S.	(gr.) white-barked.	Barbary.	1824.	—	F. h.	
40 villosa. R.S.	(gr.) villous-leaved.	C. B. S.	1714.	7.	G. h.	Pluk. alm. t. 219. f. 8.
41 spicata. R.S.	(gr.) spiked.	—	1816.	—	G. h.	
42 pubescens. R.S.	(gr.) pubescent.	—	—	—	G. h.	
43 atomaria. DC.	(gr.) velvety.	—	1820.	6. 8.	G. h.	Jacq. schoenb. 3. t. 343.
44 parviflora. DC.	(gr.) small-flowered.	Nepaul.	1824.	—	G. h.	
45 mucronata. R.S.	(gr.) mucronate.	C. B. S.	1825.	—	G. h.	
46 viminalis. R.S.	(gr.) Willow-leaved.	—	1774.	6. 7.	G. h.	Jacq. schœn. 3. t. 344.
47 erosa. R.S.	(gr.) erose-leaved.	—	1816.	—	G. h.	
48 lancea. R.S.	(gr.) lance-leaved.	—	—	—	G. h.	
49 angustifolia. R.S. (wh.)	narrow-leaved.	—	1714.	—	G. h.	Pluk. alm. t. 219. f. 6.
50 rosmarinifolia. R.S. (gr.)	Rosemary-leaved.	—	1800.	—	G. h.	Burm. afr. t. 91. f. 1.
51 undulata. R.S.	(gr.) wave-leaved.	—	1818.	—	G. h.	Jacq. scheen. 3. t. 346.
52 laevigata. R.S.	(gr.) polished-leaved.	—	1758.	—	G. h.	— 3. t. 345.
<i>elongata.</i> Jacq.						
53 elliptica. R.S.	(gr.) elliptic-leaved.	—	1816.	—	G. h.	
54 pendulina. R.S.	(gr.) pendulous.	—	—	—	G. h.	
55 lúcida. DC.	(gr.) shining-leaved.	—	1697.	7. 8.	G. h.	Comm. hort. 1. t. 93.
56 Cavanillesii. DC. (wh.)	Cavanilles'.	Mexico.	—	—	G. h.	Cavan. ic. 2. t. 132.
<i>C. I. non DC.</i>						
57 Burmanni. DC.	(gr.) Burman's.	C. B. S.	1697.	7. 8.	G. h.	Burm. afr. t. 91. f. 2.
<i>lúcidum. a. H.K.</i>						
58 concinna. B.T.	(gr.) pretty.	—	1816.	—	G. h.	
59 nervosa. R.S.	(gr.) nerved-leaved.	—	—	—	G. h.	
60 pyrifides. B.T.	(gr.) Pyrus-like.	—	1816.	—	G. h.	
61 rígida. DC.	(gr.) rigid.	—	—	—	G. h.	
62 tridactylis. B.T.	(gr.) three-fingered.	—	1816.	—	G. h.	
63 serrafolia. B.T.	(gr.) saw-leaved.	Mogador	—	—	G. h.	
64 dioica. DC.	(ye.) dioecious.	—	1823.	—	F. h.	
<i>oxyacanthoides.</i> P.S.						

65 oxyacántha. DC. (ye.) Hawthorn-like.	Magador.	1823.	F. h.
66 lobata. DC. (gr.) lobed-leaved.	Teneriffe.	1824.	F. h.
67 heterophylla. DC. (gr.) various-leaved.	1816.	6. 8.	F. h.
68 zizyphina. R.S. (gr.) Zizyphus-like.	Sicily.	—	G. h.
69 Cotinus. R.S. (gr.) Venice.	S. Europe.	1656.	6. 7.	H. h.
LOB'ADIUM. Raf. LOB'ADIUM.	Pentandria Trigynia.			Jacq. aust. 3. t. 210.
1 aromáticum. (gr.) aromatic.	N. America.	1772.	5.	H. h.
Rhus aromática. R.S. Schmáltzia aromática. Desv.				Turp. ann. mus. 5. t. 30.
2 suavéolens. (gr.) sweet.	N. America.	1759.	—	H. h.
Schmáltzia suavéolens. Desv. Rhús suavéolens. R.S.				

ORDO LXVIII.

TEREBINTHACEÆ. Brown.

PIST'ACIA. DC.	PIST'ACHIA-TREE. Diœcia Pentandria. L.			
1 vèra. DC. (st.) true.	Levant.	1570.	4. 5	H. h.
Mas.P.trifolia.w. (st.) male-flowered.	—	—	—	H. h.
Fem. P. vera. w.(gr.) female.	—	—	—	H. h.
2 reticulata. w. (st.) netted-leaved.	—	1752.	—	H. h.
3 Terebinthus. DC. (st.) turpentine-tree.	S. Europe.	1656.	6. 7.	H. h.
β sphærocárpa. DC. (st.) round-fruited.	Levant.	—	—	H. h.
4 atlántica. DC. (st.) Atlantic.	Barbary.	1790.	G. h.
5 mexicana. DC. (st.) Mexican.	Mexico.	1825.	G. h.
6 fagaroïdes. DC. (st.) Fagara-like.	—	—	—	G. h.
7 Lentiscus. DC. (st.) Mastick-tree.	S. Europe.	1664.	5. 6.	H. h.
β massiliensis. Mill. (st.) narrow-leaved.	—	—	—	H. h.
ASTR'ONIUM. DC.	ASTR'ONIUM. Diœcia Pentandria.			
gravéolens. DC. (re.) strong-scented.	S. America.	1826.	S. h.
COMOCLADIA. DC.	MAIDEN-PLUM. Triandria Monogynia. L.			
1 ilicifolia. DC. (pu.) Holly-leaved.	Caribees.	1789.	7. 9.	S. h.
2 dentata. DC. (pu.) tooth-leaved.	W. Indies.	1790.	—	S. h.
3 propinqua. DC. (pu.) related.	Cuba.	1826.	—	S. h.
4 integrifolia. DC. (pu.) entire-leaved.	Jamaica.	1778.	—	S. h.
PICRA'MNIA. DC.	PICRA'MNIA. Diœcia Pentandria. L.			
1 Antidésma. DC. (st.) Ash-leaved.	Jamaica.	1793.	6. 8.	S. h.
2 pentandra. DC. (st.) pentandrous.	W. Indies.	1822.	—	S. h.
SCHINUS. DC.	SCHINUS. Diœcia Decandria. L.			
1 Mölle. DC. (wh.) Mastick-tree.	Peru.	1597.	7. 8.	G. h.
2 terebinthifolius. DC. (w.) Terebinthus-ld.	Brazil.	1822.	S. h.
DUVA'UA. DC.	DUVA'UA. Monœcia Octandria.			
1 dépendens. DC. (wh.) simple-leaved.	Chile.	1790.	6. 7.	H. h.
Δ nigris polygama. s.s. Schinus dépendens. H.K.	—	—	—	
2? dentata. DC. (wh.) tooth-leaved.	Owhyhee.	1795.	5. 7.	G. h.
Schinus dentata. A.R.				Andr. rep. 620.
MA'URIA. K.S.	MA'URIA. Octo-Decandria Monogynia.			
1 simplicifolia. K.S. (bh.) simple-leaved.	Peru.	1822.	G. h.
2 heterophylla. K.S. (bh.) various-leaved.	—	—	G. h.
				H. et B.n.gen.a.7.t.605.
				— 7. t. 606.

ORDO LXIX.

AMYRIDEÆ. Kth. gen. tereb. p. 21.

A'MYRIS. W.	A'MYRIS. Octandria Monogynia. L.			
1 n. ritima. DC. (wh.) sea-side.	Jamaica.	1824.	6. 7.	S. h.
2 floridana. DC. (wh.) Florida.	Florida.	1826.	—	G. h.
3 brasiliensis. s.s. (wh.) Brazilian.	Brazil.	1822.	—	S. h.

4 sylvática. DC.	(wh.) wood.	Carthagena.	1793.	6. 7.	S. h.	Jacq. amer. pict. t. 188.
5 Plumiéri. DC.	(wh.) Plumier's.	Antilles.	—	S. h.	Plum. ed. Burm. t. 100.
6 toxífera. DC.	(wh.) poisonous.	W. Indies.	1800.	—	S. h.	Catesb. car. I. t. 40.
7 pinnata. DC.	(wh.) pinnate.	S. America.	1824.	—	S. h.	H. et B. n.gen. 7.t.610.
8 Agallócha. H.B.	(wh.) hill.	E. Indies.	1820.	S. h.	
9 acuminata. H.B.	(wh.) acuminate.	—	1822.	S. h.	
10 heptaphylla. H.B.	(wh.) seven-leaved.	—	—	S. h.	
11 nána. H.B.	(wh.) dwarf.	—	1824.	S. h.	
12 punctata. H.B.	(wh.) dotted.	—	1818.	S. h.	

ORDO LXX.

SPONDIACEÆ. Kth. gen. tereb. p. 30.

SPO'NDIAS. DC. HOG-PLUM. Decandria Pentagynia. L.

1 Mómbin. L.	(re.) purple-fruited.	W. Indies.	1817.	6. 8.	S. h.	Jacq. amer. t. 88.
1 purpurea. DC.	<i>Myrobálanus</i> . Jacq. non L.					
2 Myrobálanus. L.	(wh.) yellow-fruited.	W. Indies.	1739.	—	S. h.	Sloan. hist. t. 219. f. 1. 2.
lútea. DC.	<i>Móbín</i> . Jacq. non L.					
3 mangifera. DC.	Mango-bearing.	E. Indies.	1818.	S. h.	
Mangifera pinnata. L.						
4 dulcis. DC.	Otaheite-apple.	Society Isles.	1793.	6. 8.	S. h.	Sonn. it. 2. t. 123.
5 axillaris. H.B.	axillary.	Nepaul.	1824.	S. h.	
6 acuminata. H.B.	taper-pointed.	Malabar.	—	S. h.	
POUPA'R'TIA. DC.	Poupártia.	Decandria Pentagynia.				
borbònica. DC.	(pu.) Bourbon.	Bourbon.	1824.	S. h.	

ORDO LXXI.

BURSERACEÆ. Kth. gen. tereb. p. 14.

BOSWELLIA. DC. BOSWE'LLIA. Decandria Monogynia.

1 glabra. DC.	(wh.) smooth.	Coromandel.	1824.	S. h.	Roxb. corom. 3. t. 207.
2 hirsuta. DC.	(wh.) hairy.	Amboyna.	—	S. h.	Rumph. amb. 2. t. 51.
3 serrata. DC.	(wh.) Olibanum-tree.	E. Indies.	1816.	S. h.	Asiat. res. 9.p 37.7.ic.

ELAPHRIUM. K.S. ELA'PHRIUM. Octandria Monogynia.

1 glábrum.	(st.) smooth.	Carthagea.	1818.	S. h.	Jacq. amer. t. 71. f. 4.
Fágara Eláphrium. w.						

2 copalliferum. DC.	(st.) pubescent.	Mexico.	1824.	S. h.	Hern. mex. 45. f. 1.
---------------------	------------------	---------	-------	------	-------	----------------------

BALSAMODE'NDRON. DC. BALSAM-TREE. Octandria Monogynia.

gileadense. DC.	(gr.) blunt-leaved.	Arabia.	1827.	S. h.	Vahl. symb. 1. t. 11.
-----------------	---------------------	---------	-------	------	-------	-----------------------

BURSE'A. DC. BUR'SERA. Polygamia Dicecia. L.

1 gummifera. DC.	(wh.) Jamaica Birch.	W. Indies.	1690.	S. h.	Jacq. amer. t. 65.
------------------	----------------------	------------	-------	------	-------	--------------------

2 serrata. Rox.	(wh.) saw-leaved.	E. Indies.	1818.	S. h.	
-----------------	-------------------	------------	-------	------	-------	--

3 simplicifolia. DC.	(wh.) simple-leaved.	Jamaica.	1824.	S. h.	
----------------------	----------------------	----------	-------	------	-------	--

COLOPH'ONIA. DC. COLOPH'ONIA. Hexandria Monogynia.

mauritianna. DC.	(pu.) panicled.	Mauritius.	1826.	S. h.	
------------------	-----------------	------------	-------	------	-------	--

CANARIUM. DC. CAN'ARIUM. Dicecia Hexandria.

1 commune. DC.	(wh.) common.	Moluccas.	1821.	S. h.	Rumph. amb. 2. t. 47.
----------------	---------------	-----------	-------	------	-------	-----------------------

2 microcarpum. DC.	(w.) small-fruited.	—	1826.	S. h.	— 2. t. 54.
--------------------	---------------------	---	-------	------	-------	-------------

3 Pimela. DC.	(wh.) Chinese.	China.	1823.	S. h.	Ann. bot. 1. t. 7. f. 2.
---------------	----------------	--------	-------	------	-------	--------------------------

GARUGA. DC. GARUGA. Decandria Monogynia.

1 pinnata. DC.	(ye.) winged-leaved.	E. Indies.	1808.	S. h.	Roxb. cor. 3. t. 208.
----------------	----------------------	------------	-------	------	-------	-----------------------

2 madagascariensis. DC.	(ye.) Madagascar.	Madagascar.	1824.	S. h.	
-------------------------	-------------------	-------------	-------	------	-------	--

ORDO LXXII.

PTELEACEÆ. *Kth. gen. tereb. p. 22.*

PTELEA. DC.	SHRUBBY TREFOIL. <i>Tetrandria Monogynia.</i> L.			
trifoliata. DC.	(wh.) three-leaved. N. America. 1704. 6. 7. H. h.	Schmidt arb. 2. t. 76.		
TODD'ALIA. DC.	TODD'ALIA. <i>Pentandria Monogynia.</i> L.			
1 aculeata. DC.	(wh.) prickly. E. Indies. 1790. 6. 8. S. h.	Rheed. mal. 5. t. 41.		
asiatica. R.S.	<i>Scopolia aculeata.</i> w.			
2 paniculata. DC.	(wh.) panicle. Mauritius. 1824. S. h.	Lam. ill. t. 139. f. 2.		
inermis. P.S.	<i>Scopolia paniculata.</i> s.s.			
CNE'ORUM. DC.	WIDOW-WAIL. <i>Triandria Monogynia.</i> L.			
1 tricoccum. DC.	(ye.) smooth. S. Europe. 1793. 4. 9. F. h.	Lam. ill. t. 27.		
2 pulverulentum. DC.	(y.) powdered. Teneriffe. 1820. — F. h.	Vent. cels. t. 77.		
SPATH'ELIA. DC.	SPATH'ELIA. <i>Pentandria Trigynia.</i> L.			
simplex. DC.	(re.) Sumach-leaved. Jamaica. 1778. 4. 6. S. h.	Bot. reg. 670.		

ORDO LXXIII.

CONNARACEÆ. *R. Brown cong. p. 12.*

CO'NNARUS. L.	CO'NNARUS. <i>Monadelphia Decandria.</i>			
1 paniculatus. H.B.	(wh.) panicle. Chittagong. 1824. S. h.			
2 nitidus. H.B.	(wh.) glossy. Silhet. — S. h.			
ROU'REA. Aub.	ROU'REA. <i>Monadelphia Decandria.</i>			
frutescens. Aub.	(wh.) pubescent. Guiana. 1820. S. h.	Aub. gui. 1. t. 187.		
OMPHAL'OBIVM. DC.	OMPHAL'OBIVM. <i>Monadelphia Decandria.</i>			
1 indicum. DC.	(wh.) Indian. E. Indies. S. h.	Gært. fruct. 1. t. 46.		
2 africanum. DC.	(wh.) African. Sierra Leon. 1822. S. h.	Cav. dis. 7. t. 221.		
EURYC'OMA. DC.	EURYC'OMA. <i>Pentandria Monogynia.</i>			
longifolia. DC.	(pu.) long-leaved. E. Indies. 1826. S. h.			
CNE'STIS. DC.	CNE'STIS. <i>Decandria Monogynia.</i>			
1 glabra. DC.	(pu.) smooth. Mauritius. 1824. S. h.	Lam. ill. t. 387. f. 1.		
2 polyphylla. DC.	(pu.) many-leaved. Madagascar. — S. h.	——— t. 387. f. 2.		
3 monadelpha. DC. monadelphous. E. Indies. 1826. S. h.			
BRUNE'LLIA. DC.	BRUNE'LLIA. <i>Dodecadria Monogynia.</i>			
inermis. DC. various-leaved. Peru. 1823. S. h.			
BR'UCEA. DC.	BR'UCEA. <i>Dioecia Tetrandria.</i>			
1 ferruginea. w.	(st.) ferruginous. Abyssinia. 1775. 4. 5. S. h.	Bruce's tr. 5. p. 69. ic.		
antidysentérica. DC.				
2 gracilis. DC.	(st.) slender. E. Indies. S. h.			
3 sumatrana. DC.	(pu.) Sumatra. Sumatra. 1822. 4. 6. S. h.			
TETRA'DIUM. DC.	TETRA'DIUM. <i>Tetrandria Monogynia.</i>			
trichotomum. DC. trichotomous. China. 1822. S. h.			
Bruea trichotoma. s.s.				
AILA'NTHUS. DC.	AILA'NTHUS. <i>Polygamia Monœcia.</i>			
1 glandulosa. DC.	(st.) Chinese. China. 1751. 8. H. 5.	Wats. dend. brit. t. 104.		
2 excelsa. DC.	(st.) Indian. E. Indies. 1800. S. 5.	Roxb. corom. 1. t. 23.		

ORDO LXXIV.

LEGUMINOSÆ. DC. prodr. 2. p. 93.

Subordo I. PAPILIONACEÆ. DC. l. c. p. 94.

Tribus I. SOPHOREÆ. DC. l. c. p. 94.

MYROSPERMUM. K.S. BALSAM-SEED. Decandria Monogynia.

frutescens. k.s. (bh.) shrubby.	Caracas.	1824.	S. $\frac{1}{2}$.	H. et B. n.g. 6. t. 570-71.
MYROXYLUM. K.S. BALSAM-WOOD. Decandria Monogynia.				
1 pubescens. k.s. (wh.) pubescent.	Carthagena.	1820.	S. $\frac{1}{2}$.	Lamb. cinch. 92. t. 1.
2 peruiferum. k.s. (wh.) Balsam of Peru. Peru.	1824.	S. $\frac{1}{2}$.	Lam. ill. t. 341. f. 1.	
3 toluferum. k.s. (wh.) Balsam of Tolu. S. America.	S. $\frac{1}{2}$.	Woodv. m. bot. 3. t. 193.	

SOPHORA. DC. S'OPHORA. Decandria Monogynia. L.

1 japonica. DC. (wh.) Japanese.	Japan.	1753.	8. 9. H. 5.	Andr. deposit. 585.
β pendula. L.C. (wh.) pendulous.	—	—	H. 5.	
γ variegata. L.C. (wh.) striped-leaved.	—	—	H. 5.	
2 chinensis. L.C. (wh.) Chinese.	China.	—	H. 5.	
3 glauca. DC. (pu.) glaucous.	E. Indies.	1818.	S. $\frac{1}{2}$.	Lam. ill. t. 325. f. 2.
4 tomentosa. DC. (ye.) downy.	—	1690.	S. $\frac{1}{2}$.	Trew. ehr. t. 59.
5 occidentalis. W. (ye.) West Indian.	W. Indies.	—	S. $\frac{1}{2}$.	
6 crassifolia. DC. (ye.) thick-leaved.	Senegal.	1824.	S. $\frac{1}{2}$.	
7 secundiflora. DC. (bl.) side-flowering.	New Spain.	1818.	G. $\frac{1}{2}$.	Cavan. ic. 5. t. 401.
<i>Virgilia secundiflora</i> . C.I. <i>Broussonetia secundiflora</i> . Ort. dec. 5. t. 7.				
8 macrocarpa. DC. (ye.) large-podded.	Chile.	1822.	4. 5. F. $\frac{1}{2}$.	Lodd. bot. cab. 1125.
9 littoralis. DC. (ye.) sea-side.	Brazil.	1824.	S. $\frac{1}{2}$.	Pluk. alnq. t. 104. f. 3.
10 robusta. H.B. robust.	Silhet.	1820.	S. $\frac{1}{2}$.	
11 velutina. B.R. velvetty.	Nepaul.	1824.	6. 7. G. $\frac{1}{2}$.	Bot. reg. 1185.
12 flavescens. DC. (ye.) pale-yellow.	Siberia.	1785.	5. 7. H. $\frac{1}{2}$.	Dill. elth. f. 136.
PSEUDOSOPHORA. PSEUDOSOPHORA. Diadelphia Decandria.				
1 alopecuroides. DC. (wh.) Fox-tail.	Levant.	1731.	7. 8. H. $\frac{1}{2}$.	
2 sericea. N. (wh.) silky.	Missouri.	1811.	6. 7. H. $\frac{1}{2}$.	

Astragalus carnosus. Ph.

EDWARDSIA. DC. EDWARDSIA. Decandria Monogynia.

1 grandiflora. DC. (ye.) large-flowered.	N. Zealand.	1772.	3. 6. F. $\frac{1}{2}$.	Bot. mag. 167.
<i>Sophora tetrapeta</i> . B.M.				
2 chrysophylla. DC. (ye.) golden-leaved.	—	—	F. $\frac{1}{2}$.	Bot. reg. 738.
3 microphylla. DC. (ye.) small-leaved.	—	1772.	F. $\frac{1}{2}$.	Bot. mag. 1442.
4 minima. L.C. (ye.) smallest.	—	1816.	F. $\frac{1}{2}$.	

ORMOSIA. DC. ORMOSIA. Decandria Monogynia.

1 dasycarpa. DC. (pu.) smooth-leaved.	W. Indies.	1793.	6. 7. S. $\frac{1}{2}$.	
<i>Sophora monosperma</i> . Swartz.				
2 coccinea. DC. (pu.) scarlet-seeded.	Guiana.	1826.	S. $\frac{1}{2}$.	Linn. trans. 10. t. 26.
VIRGILIA. DC. VIRGILIA. Decandria Monogynia.				Linn. trans. 10. t. 25.
1 capensis. DC. (bh.) Cape.	C. B. S.	1767.	7. 8. G. $\frac{1}{2}$.	Bot. mag. 1590.

2 intrusa. H.K. (ye.) small-flowered.	—	1790.	5. 8. G. $\frac{1}{2}$.	— t. 2617.
<i>Sophora sylvatica</i> . Burchell.				
3 aurea. DC. (ye.) golden.	Abyssinia.	1777.	7. G. $\frac{1}{2}$.	L'Herit. st. nov. 1. t. 75.
4 lutea. DC. (ye.) American.	N. America.	1812.	6. 7. G. $\frac{1}{2}$.	Mich. fil. arb. 3. t. 3.

MACROTR'OPIS. DC. MACROTR'OPIS. Decandria Monogynia.

1 foetida. DC. (wh.) strong-scented.	China.	1822.	4. 5. F. $\frac{1}{2}$.	
2 inodora. DC. (wh.) scentless.	—	1824.	F. $\frac{1}{2}$.	
ANAGYRIS. DC. BEAN TREFOIL. Decandria Monogynia. L.				
1 foetida. DC. (ye.) strong-scented.	Spain.	1570.	4. 5. F. $\frac{1}{2}$.	Lodd. bot. cab. t. 740.
β glauca. DC. (ye.) glaucous-leaved.	S. Europe.	1824.	F. $\frac{1}{2}$.	

2 latifolia. DC. (ye.) broad-leaved.	Teneriffe.	1815.	F. $\frac{1}{2}$.	
PIPTANTHUS. B.F.G. PIPTANTHUS. Decandria Monogynia.				
nepalensis. B.F.G. (ye.) Laburnum 1d.	Nepaul.	1819.	5. 6. H. $\frac{1}{2}$.	Swt. br. fl. gar. 264.
<i>Thermopsis napaulensis</i> . DC. laburnifolia. D.P. <i>Baptisia nepalensis</i> . Hook. ex. flor. 131.				

THERMO'PSIS. DC. THERMO'PSIS. Decandria Monogynia.						
1 rhombifolia. DC. (ye.) rhomb-leaved.	Louisiana.	1811.	6. 7. H. ȝ.			
Cytisus rhombifolius. Ph.						
2 fabacea. DC. (ye.) Bean-leaved.	Kamtschatka.	1824.	— H. ȝ.	Pall. astr. t. 90. f. 2.		
3 lanceolata. DC. (ye.) spear-leaved.	Siberia.	1776.	— H. ȝ.	Bot. mag. 1389.		
Sophora lupinoides. Pall. astr. t. 89. Podalyria lupinoides. B.M.						
4 corynensis. DC. (ye.) alpine.	Altay.	1824.	— H. ȝ.	Pall. astr. t. 90. f. 1.		
Sophora alpina. Pall. Podalyria alpina. W. enum.						
BAPTISIA. DC. BAPTISIA. Decandria Monogynia.						
1 perfoliata. DC. (ye.) perfoliate.	Carolina.	1732.	8. F. ȝ.	Lodd. bot. cab. 1104.		
Crotalaria perfoliata. L. Raphnia perfoliata. W.						
2 exaltata. B.F.G. (bl.) tall upright.	N. America.	1812.	6. 7. H. ȝ.	Swt. br. fl. gar. 97.		
3 auriculata. (bl.) eared.			— H. ȝ.			
4 australis. DC. (bl.) common.		1758.	— H. ȝ.	Bot. mag. 509.		
Sophora australis. B.M. Podalyria australis. Vent. cels. t. 56.						
5 confusa. (bl.) persistent-bract'd. N.America.	1812.	— H. ȝ.				
6 villosa. DC. (st.) villous.	1811.	— H. ȝ.				
7 versicolor. B.C. (va.) various-colour'd.	1824.	6. 8. H. ȝ.				
8 alba. DC. (wh.) white-flowered.	1724.	6. 7. H. ȝ.	Bot. mag. 1177.			
9 tinctoria. DC. (ye.) dyer's.	1759.	7. 8. H. ȝ.			t. 1099.	
Sophora tinctoria. L. Podalyria tinctoria. B.M.						
CYCLOPIA. DC. CYCLOPIA. Decandria Monogynia.						
1 genistoides. DC. (ye.) Genista-like.	C. B. S.	1787.	7. 8. G. h.	Bot. mag. 1259.		
Ibbetsonia genistoides. B.M. Gompholobium maculatum. A.R. 427.						
2 galioides. DC. (ye.) Galium-like.	C. B. S.	1818.	7. 8. G. h.	Pluk. alm. t. 413. f. 4.		
3 latifolia. DC. (ye.) broad-leaved.			— G. h.			
PODALYRIA. DC. PODALYRIA. Decandria Monogynia.						
1 Burchellii. DC. (pu.) Burchell's.	C. B. S.	1818.	5. 7. G. h.			
2 hirsuta. DC. (pu.) hairy.		1774.	7. 8. G. h.			
3 cuneifolia. DC. (wh.) wedge-leaved.		1804.	5. 8. G. h.	Vent. cels. t. 99.		
4 oleafolia. P.L. (pu.) Olive-leaved.			— G. h.	Sal. par. lond. 114.		
5 sericea. DC. (pu.) silky.		1778.	10. 1. G. h.	Bot. mag. 1923.		
6 myrtillifolia. DC. (pu.) Myrtle-leaved.		1795.	4. 7. G. h.			
7 buxifolia. DC. (pu.) Box-leaved.		1790.	5. 10. G. h.	Bot. reg. 869.		
8 styracifolia. DC. (pu.) Storax-leaved.		1792.	5. 7. G. h.	Bot. mag. 1580.		
9 calyptarata. W. (pu.) one-flowered.			4. 5. G. h.			
10 coruscans. Rch. (pu.) glittering.		1822.	— G. h.			
11 glauca. DC. (pu.) glaucous-leaved.			— G. h.	Lam. ill. t. 327. f. 4.		
12 argentea. DC. (li.) silvery-leaved.		1780.	2. 6. G. h.	Sal. par. lond. t. 7.		
biflora. Bot. mag. t. 753.						
13 liparioides. DC. (li.) Liparia-like.		1824.	4. 6. G. h.			
14 subbiflora. DC. (pu.) netted-leaved.			— G. h.			
15 cordata. DC. (pu.) heart-leaved.		1794.	5. 7. G. h.			
CHORIZEMA. DC. CHORIZEMA. Decandria Monogynia.						
1 ilicifolia. DC. (sc.) Holly-leaved.	N. Holland.	1803.	3. 10. G. h.	Labill. vey. 1. t. 21.		
2 nana. DC. (sc.) dwarf prickly.			— G. h.	Bot. mag. 1032.		
3 rhombaea. DC. (sc.) various-leaved.			4. 6. G. h.	Swt. flor. aust. t. 40.		
4 Henchmanni. B.R. (sc.) Henchmann's.		1825.	5. 9. G. h.	Bot. reg. 986.		
5 platylobiodes. DC. (ye.) Platylobiump-like.			— G. h.			
PODOL'OBIA. DC. PODOL'OBIA. Decandria Monogynia.						
1 trilobatum. DC. (ye.) three-lobed.	N. S. W.	1791.	4. 7. G. h.	Bot. mag. 1477.		
Chorozema trilobatum. L.T. Pultenæ ilicifolia A.R. 320.						
2 staurophyllum. DC. (ye.) spiny-leaved.	N. S. W.	1821.	3. 5. G. h.	Bot. reg. 959.		
3 scandens. DC. (ye.) climbing.		1825.	4. 6. G. h.			
Chorozema scandens. L.T.						
OXYL'OBIA. DC. OXYL'OBIA. Decandria Monogynia.						
1 obtusifolium. S.F.A. (sc.) blunt-leaved.	N. Holland.	1825.	4. 6. G. h.	Swt. flor. aust. t. 5.		
2 arborescens. DC. (ye.) arborescent.	V. Diem. Isl.	1805.	5. 9. G. h.	Bot. reg. 392.		
3 retusum. B.R. (or.) retuse-leaved.	N. Holland.	1822.	4. 6. G. h.		t. 913.	
4 ellipticum. DC. (ye.) elliptic-leaved.	V. Diem. Isl.	1805.	5. 9. G. h.	Lab. nov. hol. 1. t. 135.		
5 cordifolium. DC. (sc.) heart-leaved.	N. S. W.	1807.	4. 9. G. h.	Andr. rep. 492.		
6 spinosum. DC. (ye.) spiny.	N. Holland.	1823.	5. 8. G. h.			
7 Pulteneæ. DC. (ye.) wood.	N. S. W.		4. 6. G. h.			
CALLISTACHYS. DC. CALLISTACHYS. Decandria Monogynia.						
1 lanceolata. DC. (ye.) spear-leaved.	N. Holland.	1815.	5. 8. G. h.	Bot. reg. 216.		
2 ovata. DC. (ye.) oval-leaved.			— G. h.	Bot. mag. 1925.		
BRACHYS'EMA. DC. BRACHYS'EMA. Decandria Monogynia.						
1 latifolium. DC. (sc.) broad-leaved.	N. Holland.	1803.	4. 7. G. h.	Bot. reg. 118.		
2 undulatum. DC. (ye.) wave-leaved.		1820.	2. 7. G. h.		t. 642.	

GOMPHOL'OBIA. DC. GOMPHOL'OBIA. Decandria Monogynia.

1 barbigerum. DC. (ye.) bearded-flow'd. N. S. W.	1824.	5. 8. G. ♂.
2 grandiflòrum. DC. (ye.) large-flowered. ———	1803.	3. 9. G. ♂.
3 virgátum. DC. (ye.) twiggy. ———	1824.	5. 8. G. ♂.
4 latifòlium. DC. (ye.) broad-leaved. ———	1803.	3. 9. G. ♂.
5 fimbriatum. Sm. ex. bot. t. 58. <i>psoraleafolium</i> . Sal. par. lond. t. 6.		
5 glaucescens. F.T. (ye.) glaucous. N. S. W.	1821.	4. 6. G. ♂.
6 marginatum. DC. (ye.) small-flower'd. N. Holland.	1803.	3. 9. G. ♂.
7 pedunculare. DC. (ye.) long-stalk'd. N. S. W.	1825.	4. 7. G. ♂.
8 tetrathecoides. DC. (ye.) Tetraetheca-like. ———		G. ♂.
9 polymorphum. DC. (sc.) variable. N. Holland.	1803.	3. 8. G. ♂. — Bot. mag. 1533.
10 tomentosum. DC. (ye.) tomentose. ———	—	4. 7. G. ♂. Labill. nov. hol. t. 134.
11 glabratum. DC. (ye.) smooth. ———	1824.	— G. ♂.
12 venustum. DC. (pu.) purple-flowered. ———	1803.	— G. ♂.
13 pinnatum. DC. (ye.) wing-leaved. N. S. W.	1822.	5. 6. G. ♂.

BURTONIA. DC. BURTONIA. Decandria Monogynia.

1 scabra. DC. (pu.) rough-leaved. N. Holland.	1803.	5. 7. G. ♂.
2 minor. DC. (ye.) hairy-stalked. N. S. W.	1812.	3. 8. G. ♂.
3 sessilifolia. DC. (ye.) sessile-leaved. ———	1825.	— G. ♂.
4 furcellata. DC. (ye.) forked. N. S. W.	1822.	4. 6. G. ♂.
5 reticulata. DC. (ye.) netted. ———	—	— G. ♂.

Daviesia reticulata. L.T.

VIMIN'ARIA. DC. VIMIN'ARIA. Decandria Monogynia.

denudata. DC. (ye.) half-naked. N. S. W.	1789.	6. 9. G. ♂.
SPHÆROL'OBIA. DC. SPHÆROL'OBIA. Decandria Monogynia.		Bot. mag. 1190.

1 vimineum. DC. (ye.) yellow-flower'd. N. S. W.	1802.	5. 8. G. ♂.
2 medium. DC. (re.) red-flowered. N. Holland.	1803.	6. 8. G. ♂.

A'OTUS. DC. A'OTUS. Decandria Monogynia.

1 villosa. DC. (ye.) villous. V. Diem. Isl. 1790.	4. 6. G. ♂.	
2 virgata. DC. (ye.) slender. N. Holland.	1820.	4. 8. G. ♂.

DILLW'YNIA. DC. DILLW'YNIA. Decandria Monogynia.

1 floribunda. DC. (ye.) close-flowered. N. S. W.	1794.	4. 7. G. ♂.
2 ericifolia. DC. (ye.) Heath-leaved. ———	—	3. 7. G. ♂.

Sm. exot. bot. t. 26.
——— t. 25.
Lodd. bot. cab. 401.
Bot. mag. 944.
——— t. 1527.

3 juniperina. B.C. (ye.) Juniper-leaved. ———	1818.	6. 7. G. ♂.
4 glaberrima. DC. (ye.) smooth. ———	1800.	3. 7. G. ♂.

5 parvifolia. DC. (ye.) small-leaved. ———	—	G. ♂.
6 acicularis. DC. (ye.) needle-leaved. ———	1822.	5. 7. G. ♂.

7 rufis. DC. (ye.) villous-branched. ———	1812.	— G. ♂.
8 teretifolia. Sie. (ye.) round-leaved. ———	1825.	— G. ♂.

9 hispidula. Sie. (ye.) hispid. ———	—	G. ♂.
10 tenuifolia. DC. (ye.) slender-leaved. ———	1820.	— G. ♂.

11 phyllicoïdes. F.T. (ye.) Phyllica-leaved. ———	1824.	5. 8. G. ♂.
12 sericea. F.T. (ye.) silky. ———	—	G. ♂.

13 cinerascens. DC. (ye.) ash-coloured. V. Diem. Isl. 1818.	6. 7. G. ♂.
---	-------------

EUTAXIA. DC. EUTAXIA. Decandria Monogynia.		
1 myrtifolia. DC. (ye.) Myrtle-leaved. N. Holland.	1803.	3. 6. G. ♂.

Bot. mag. 2247.
Bot. mag. 1274.

Dillw'nia. Bot. mag. 1545. nec aliorum.		
2 pungens. S.F.A. (ye.) pungent-leaved. ———	1825.	6. 8. G. ♂.

SCLEROTHA'MNUS. DC. SCLEROTHA'MNUS. Decandria Monogynia.		
microphyllus. DC. (ye.) small-leaved. N. Holland.	1803.	5. 6. G. ♂.

GASTROL'OBIA. DC. GASTROL'OBIA. Decandria Monogynia.		
bilobum. DC. (ye.) two-lobed. N. Holland.	1803.	3. 5. G. ♂.

Bot. reg. 411.

EUCHILUS. DC. EUCHILUS. Decandria Monogynia.		
obcordatus. DC. (ye.) heart-leaved. N. Holland.	1803.	3. 6. G. ♂.

Bot. reg. 403.

PULTEN'EA. DC. PULTEN'EA. Decandria Monogynia.		
1 daphnoides. DC. (ye.) Daphne-leaved. N. S. W.	1792.	6. 7. G. ♂.

Bot. mag. 1394.
Andr. rep. t. 574.

2 obcordata. DC. (ye.) heart-leaved. V. Diem. Isl.	1808.	5. 7. G. ♂.
3 biloba. DC. (ye.) two-lobed. N. Holland.	1818.	— G. ♂.

Bot. mag. t. 2091.

4 scabra. H.K. (ye.) rough-leaved. N. S. W.	1803.	— G. ♂.
5 incurvata. F.T. (ye.) incurved-ld.	1825.	— G. ♂.

6 canescens. F.T. (ye.) canescent.	1824.	4. 7. G. ♂.
7 polifolia. F.T. (ye.) Poley-leaved.	—	— G. ♂.

8 argentea. F.T. (ye.) silvery.	1826.	— G. ♂.
9 ferruginea. DC. (ye.) ferruginous.	1818.	— G. ♂.

Linn. trans. 11. t. 23.

10 racemulòsa. DC.	(ye.) racemed.	N. S. W.	1821.	4. 7.	G. h.
11 parviflòra. DC.	(ye.) small-flowered.	— —	1824.	5. 7.	G. h.
12 procumbens. F.T.	(ye.) procumbent.	— —	— —	— —	G. h.
13 strícta. DC.	(ye.) upright.	— —	1803.	4. 6.	G. h.
14 ellíptica. DC.	(ye.) elliptic-leaved.	— —	1818.	4. 7.	G. h.
15 plumòsa. DC.	(ye.) feathered.	— —	1822.	5. 7.	G. h.
16 thymifòlia. DC.	(ye.) Thyme-leaved.	— —	— —	— —	G. h.
17 hypolámpra. DC.	(ye.) whorl-branched.	— —	1824.	— —	G. h.
18 villifera. DC.	(ye.) wool-bearing.	— —	— —	— —	G. h.
19 polygalifòlia. DC.	(ye.) Milkwort-ld.	— —	1818.	4. 7.	G. h.
20 fléxilis. DC.	(ye.) fragrant.	— —	1801.	4. 6.	G. h.
21 euchiña. DC.	(ye.) wedge-leaved.	— —	1824.	5. 7.	G. h.
22 retusa. DC.	(ye.) blunt.	— —	1789.	4. 5.	G. h.
23 capitellata. DC.	(ye.) small-headed.	— —	1825.	— —	G. h.
24 linophylla. DC.	(ye.) Flax-leaved.	— —	1789.	5. 7.	G. h.
25 microphylla. DC.	(ye.) small-leaved.	— —	1823.	— —	G. h.
26 stipularis. DC.	(ye.) scaly.	— —	1792.	4. 6.	G. h.
27 paleacea. DC.	(ye.) chaffy.	— —	1789.	5. 7.	G. h.
28 aristata. DC.	(ye.) awned-calyxed.	— —	1823.	— —	G. h.
29 echinula. DC.	(ye.) harsh-leaved.	— —	— —	— —	G. h.
30 vestita. DC.	(ye.) awned.	N. Holland.	1803.	4. 7.	G. h.
31 candida. B.C.	(ye.) white-leaved.	— —	1824.	— —	G. h.
32 tenuifolia. DC.	(ye.) slender-leaved.	— —	1818.	— —	G. h.
33 villosa. DC.	(ye.) villous.	N. S. W.	1790.	4. 5.	G. h.
34 pedunculata. B.M.	(ye.) pedunculated.	— —	1824.	5. 6.	G. h.
35 eriophora. L.C.	(ye.) wool-bearing.	— —	1794.	4. 7.	G. h.
36 incarnata. M.C.	(fl.) flesh-coloured.	N. Holland.	1824.	— —	G. h.
37 rosmarinifolia. ψ.	(ye.) Rosemary-leav'd.	— —	— —	— —	G. h.
38 áspéra. DC.	(ye.) rough.	N. S. W.	— —	5. 8.	G. h.
39 comosa. DC.	(ye.) tufted.	— —	— —	— —	G. h.
40 squarrosa. DC.	(ye.) squarrose.	— —	— —	— —	G. h.
41 phylloides. DC.	(ye.) Phyllica-like.	— —	— —	— —	G. h.
DAVIESIA. DC.	DAVIESIA. Decandria Monogynia.				
1 latifolia. DC.	(ye.) broad-leaved.	N. S. W.	1803.	5. 8.	G. h.
2 corymbosa. DC.	(ye.) corymb-flower'd.	— —	1804.	— —	G. h.
3 mimosoides. H.K.	(ye.) green-leaved.	— —	1800.	— —	G. h.
corymbosa. A.R. non Smith.					
4 glauca. B.C.	(ye.) glaucous-leaved.	— —	1812.	— —	G. h.
mimosoides. Bot. mag. 1957. non H.K.					
5 umbellulata. DC.	(ye.) small-umbelled.	— —	1822.	4. 8.	G. h.
6 incrassata. DC.	(ye.) thick-leaved.	— —	1820.	5. 8.	G. h.
7 acicularis. DC.	(ye.) needle-leaved.	— —	1804.	6. 7.	G. h.
8 ulicina. DC.	(ye.) Furze-like.	N. S. W.	1792.	4. 8.	G. h.
9 pungens. M.C.	(ye.) pungent.	N. Holland.	1825.	— —	G. h.
10 squarrosa. DC.	(ye.) squarrose.	N. S. W.	1822.	5. 7.	G. h.
11 cordata. DC.	(ye.) heart-leaved.	N. Holland.	1824.	— —	G. h.
12 alata. DC.	(ye.) winged.	— —	1820.	4. 6.	G. h.
MIRBELIA. DC.	MIRBELIA. Decandria Monogynia.				
1 reticulata. DC.	(pu.) netted-leaved.	N. S. W.	1792.	5. 8.	G. h.
2 pungens. M.C.	(pu.) pungent-leav'd.	N. Holland.	1824.	— —	G. h.
3 speciosa. DC.	(pu.) beautiful.	N. S. W.	— —	— —	G. h.
4 grandiflora. B.M.	(y.re.) large-flowered.	— —	1823.	— —	G. h.
5 Baxteri. M.C. Baxter's.	N. Holland.	1824.	G. h.
6 dilatata. DC.	(pu.) lobed-leaved.	— —	1803.	5. 8.	G. h.

Tribus II. LOTEÆ. DC. prodr. 2. p. 115.

Subtribus I. GENISTEÆ. DC. l. c. 115.

H'OVEA. DC.

H'OVEA. Monadelphia Decandria.

1 rosmarinifolia. F.T.	(bl.) Rosemary-ld.	N. S. W.	1824.	3. 6.	G. h.
2 longifolia. DC.	(bl.) long-leaved.	— —	1805.	3. 7.	G. h.
3 linearis. DC.	(bl.) linear-leaved.	— —	1796.	— —	G. h.
4 lanceolata. DC.	(bl.) spear-leaved.	— —	1805.	— —	G. h.

Bot. mag. 1588.

Linn. trans. 11. t. 24.

Swt. flor. aust. t. 37.
— — t. 35.

Bot. reg. 378.

Schrad. s. han. 3. t. 18.

Bot. mag. 475.

Lodd. bot. cab. t. 291.

Lod. bot. cab. 1236.

Bot. mag. 2086.

— — t. 967.

Bot. mag. 2859.

Bot. mag. 1757.

Andr. deposit. 526.

Lodd. bot. cab. 43.

Andr. deposit. 304.

Bot. mag. 1211.

Swt. flor. aust. t. 34.

Bot. mag. 2771.

Bot. reg. 1005.

— — t. 728.

Bot. reg. 1041.

5	elliptica. DC.	(bl.) elliptic-leaved.	N. S. W.	1817.	3. 7.	G. h.
	<i>Poirétia elliptica.</i> L.T.					
6	purpurea. S.F.A.	(pu.) purple-flowered.	_____	1824.	2. 5.	G. h.
7	latifolia. DC.	(bl.) broad-leaved.	_____	1817.	3. 6.	G. h.
8	Celsi. DC.	(bl.) Cels's.	N. Holland.	1818.	3. 7.	G. h.
	PLAGIOL'OBIIUM. S.F.A. PLAGIOL'OBIIUM. <i>Diadelphia Decandria.</i>					
1	chorizemæfolium. S.F.A. (bl.)	<i>Chorizema</i> -lt.	N. Holland.	1824.	2. 5.	G. h.
2	ilicifolium. S.F.A. (bl.)	Holly-leaved.	_____	1827.	—	G. h.
	PLATYL'OBIIUM. DC. FLAT-PEA. <i>Monadelphia Decandria.</i>					
1	formosum. DC.	(ye.) large-flowered.	N. S. W.	1790.	6. 8.	G. h.
2	parviflorum. DC.	(ye.) smaller-flower'd.	_____	1792.	5. 9.	G. h.
3	ovatum. DC.	(ye.) oval-leaved.	_____	1820.	5. 7.	G. h.
4	triangulare. DC.	(ye.) triangular-ld.	V. Diem. Isl.	1805.	5. 9.	G. h.
	BOSSI'EA. DC. BOSSI'EA. <i>Monadelphia Decandria.</i>					
1	scopoldriæ. DC. (ye.)	<i>Plank-plant.</i>	N. S. W.	1792.	5. 7.	G. h.
	<i>Platylöbium Scopoldriæ.</i> Andr. rep. 191.					
2	ensata. DC.	(ye.) sword-stemmed.	N. S. W.	1822.	—	G. h.
	<i>rufa.</i> Lod. bot. cab. 1119, non DC.					
3	rufa. DC.	(re.) red-flowered.	N. Holland.	1803.	6. 9.	G. h.
4	linophylla. DC.	(ye.) narrow-leaved.	_____	—	7. 9.	G. h.
5	heterophylla. DC. (ye.)	various-leaved.	N. S. W.	1792.	4. 12.	G. h.
	<i>lanceolata.</i> B.M. 1144. <i>Platylöbium lanceolatum et ovatum.</i> A.R. 205 et 276.					
6	rotundifolia. DC.	(ye.) round-leaved.	N. S. W.	1822.	5. 7.	G. h.
7	rhombifolia. DC.	(ye.) rhomb-leaved.	_____	—	4. 6.	G. h.
	<i>lenticularis.</i> Lod. bot. cab. 1238, non DC.					
8	microphylla. DC.	(ye.) small-leaved.	_____	1803.	5. 7.	G. h.
9	lenticularis. DC.	(ye.) orbicular-leaved.	_____	1822.	—	G. h.
10	foliosa. F.T.	(ye.) leafy.	_____	1823.	4. 7.	G. h.
11	buxifolia. F.T.	(ye.) Box-leaved.	_____	—	—	G. h.
12	prostrata. DC.	(ye.) prostrate.	_____	1803.	—	G. h.
13	cordifolia. S.F.A. (y.pu.)	heart-leaved.	N. Holland.	1824.	—	G. h.
14	cinerea. DC.	(y.pu.) sharp-leaved.	V. Diem. Isl.	1803.	5. 7.	G. h.
	GO'ODIA. DC. GO'ODIA. <i>Monadelphia Decandria.</i>					
1	lotifolia. DC.	(ye.) Lotus-leaved.	V. Diem. Isl.	1793.	4. 7.	G. h.
2	subpubescens. DC.	(ye.) hairy-leaved.	_____	1812.	—	G. h.
3	pubescens. DC.	(y.pu.) downy.	_____	1805.	—	G. h.
4	retusa. M.C.	(ye.) retuse-leaved.	N. Holland.	1824.	—	G. h.
	TEPHROTHA'MNUS. TEPHROTHA'MNUS. <i>Monadelphia Decandria.</i>					
	tomentosus.	(ye.) woolly.	C. B. S.	1798.	7. 8.	G. h.
	<i>Cytisus tomentosus.</i> A.R. <i>Goòdia? polysperma.</i> DC.					
	SCOTTIA. H.K. SCOTTIA. <i>Monadelphia Decandria.</i>					
1	dentata. DC.	(sc.) broad-leaved.	N. Holland.	1803.	1. 6.	G. h.
2	trapeziformis. M.C.	(sc.) trapeziform.	_____	1825.	—	G. h.
3	angustifolia. M.C. (br.)	narrow-leaved.	_____	—	1. 6.	G. h.
	TEMPLET'ONIA. DC. TEMPLET'ONIA. <i>Monadelphia Decandria.</i>					
1	retusa. DC.	(sc.) green-leaved.	N. Holland.	1803.	3. 7.	G. h.
2	glauca. DC.	(sc.) glaucous-leaved.	_____	—	—	G. h.
	RA'FNIA. DC. RA'FNIA. <i>Monadelphia Decandria.</i>					
1	cordata. DC.	(ye.) heart-leaved.	C. B. S.	1821.	5. 7.	G. h.
2	elliptica. DC.	(ye.) elliptic-leaved.	_____	1816.	—	G. h.
3	cuneifolia. DC.	(y.pu.) wedge-leaved.	_____	1820.	—	G. δ.
4	triflora. DC.	(ye.) three-flowered.	_____	1786.	—	G. δ.
	<i>Crotalária triflora.</i> L. <i>Borbónia cordata.</i> And. rep. t. 31. nec aliorum.					
5	lancea. DC.	(ye.) lance-leaved.	C. B. S.	1816.	5. 7.	G. δ.
	<i>Edmánnia lancea.</i> Th.					
6	opposita. DC.	(ye.) opposite-leaved.	_____	1825.	—	G. h.
7	angulata. DC.	(ye.) angular-stalked.	_____	1816.	—	G. h.
8	filifolia. DC.	(ye.) narrow-leaved.	_____	—	—	G. h.
	VASCO'A. DC. VASCO'A. <i>Monadelphia Decandria.</i>					
1	amplexicaulis. DC.	(ye.) stem-clasping.	C. B. S.	1812.	6. 7.	G. h.
	<i>Crotalária amplexicaulis.</i> L. <i>Ráfnia amplexicaulis.</i> Th.					
2	perfoliata. DC.	(ye.) perfoliate.	C. B. S.	1812.	7. 8.	G. h.
	<i>Borbónia perfoliata.</i> Th. <i>Crotalária amplexicaulis.</i> Lam.					
	BORB'ONIA. DC. BORB'ONIA. <i>Monadelphia Decandria.</i>					
1	barbata. DC.	(ye.) bearded.	C. B. S.	1816.	6. 8.	G. h.
2	trinervia. DC.	(ye.) three-nerved.	_____	1759.	—	G. h.
3	lanceolata. DC.	(ye.) many-nerved.	_____	1752.	—	G. h.
4	cordata. DC.	(ye.) heart-leaved.	_____	1759.	—	G. h.
5	ruscifolia. DC.	(ye.) Ruscus-leaved.	_____	1816.	—	G. h.
	Lam. ill. t. 610. f. 2.					
	Pluk. aim. t. 297. f. 4.					
	Jacq. schœn. b. 2. t. 217.					
	2. t. 218.					
	Bot. mag. 2128.					

Seba. thes. 1. t. 24. f. 5.

Lam. ill. t. 610. f. 2.

Pluk. aim. t. 297. f. 4.

Jacq. schœn. b. 2. t. 217.

2. t. 218.

Bot. mag. 2128.

6 parviflora. DC.	(ye.) small-flowered.	C. B. S.	1821.	6. 8. G. h.	
7 crenata. DC.	(ye.) notch-leaved.	——	1774.	—— G. h.	Bot. mag. 274.
8 ciliata. DC.	(ye.) fringed.	——	1816.	—— G. h.	Houtt. syst. 8. t. 62 f. 2.
ACHYR'ONIA. DC.	ACHYR'ONIA.	Diadelphia Decandria.			
villosa. DC.	(ye.) villous.	N. Holland.	1810.	5. 8. G. h.	Wendl. hort. her. 1.t.12.
LIP'ARIA. DC.	LIP'ARIA.	Diadelphia Decandria.			
sphaerica. DC.	(ye.) globe-headed.	C. B. S.	1794.	7. 8. G. h.	Bot. mag. 1241.
PRIESTLE'YA. DC.	PRIESTLE'YA.	Diadelphia Decandria.			
1 myrtifolia. DC.	(ye.) Myrtle-leaved.	C. B. S.	1825.	4. 7. G. h.	DC. Leg. mem. t. 29.
2 hirsuta. DC.	(ye.) shaggy-stemmed.	——	1792.	12. 4. G. h.	Bot. reg. 8.
<i>Liparia hirsuta</i> . B.R.					
3 laevigata. DC.	(ye.) polished.	——	1799.	7. 8. G. h.	DC. Leg. mem. t. 30.
<i>Borbonia laevigata</i> . Lod. bot. cab. 247.					
4 capitata. DC.	(ye.) headed.	——	1812.	3. 6. G. h.	
5 graminifolia. DC.	(ye.) narrow-leaved.	——	1800.	4. 7. G. h.	
6 teres. DC.	(ye.) smooth-stalked.	——	1816.	—— G. h.	
7 ericafolia. DC. (w.pu.)	Heath-leaved.	——	1812.	4. 8. G. h.	
8 sericea. DC.	(ye.) silky-leaved.	——	1794.	—— G. h.	
9 axillaris. DC.	(ye.) axillary-flower'd.	——	1824.	—— G. h.	
10 elliptica. DC.	(ye.) elliptic-leaved.	——	1816.	—— G. h.	
11 villosa. DC.	(ye.) villous-leaved.	——	1774.	—— G. h.	
12 vestita. DC.	(ye.) concave-leaved.	——	1800.	5. 6. G. h.	
<i>Liparia vestita</i> . B.M. <i>Liparia villosa</i> . A.R. 382. nec aliorum.					
13 tomentosa. DC.	(ye.) downy.	C. B. S.	1812.	5. 8. G. h.	
14 umbellifera. DC.	(ye.) umbel-flowered.	——	1825.	6. 7. G. h.	
HA'LLIA. DC.	HA'LLIA.	Monadelphia Decandria.			
1 alata. DC.	(pu.) wing-stalked.	C. B. S.	1818.	6. 8. G. h.	
2 flaccida. DC.	(pu.) long-leaved.	——	1789.	6. 9. G. ♂.	
3 angustifolia. DC.	(pu.) narrow-leaved.	——	1816.	—— G. ♂.	
4 cordata. DC.	(pu.) heart-leaved.	——	1787.	—— G. h.	
<i>Glycine monophylla</i> . Jacq. <i>Hedysarum cordatum</i> . Th. n. ac. ups. 6.t. 1.					Jacq. schœnb. 3. t. 296.
5 Asarina. DC.	(pu.) Asarum-like.	C. B. S.	1816.	6. 9. G. h.	
6 imbricata. DC.	(pu.) imbricated.	——	1812.	—— G. h.	
HEYLA'NDIA. DC.	HEYLA'NDIA.	Monadelphia Decandria.			
hebecarpa. DC.	(ye.) hairy-podded.	Ceylon.	1820.	6. 8. S. h.	
CROTAL'ARIA. DC.	CROTAL'ARIA.	Monadelphia Decandria.			
1 alata. DC.	(st.) winged-stalked.	Nepaul.	1818.	4. 8. S. h.	
2 platycarpa. DC.	(ye.) flat-podded.	N. America.	1824.	7. 9. H. ⓧ.	
3 Pürshii. DC.	(ye.) Pursh's.	——	1827.	—— H. ⓧ.	
<i>laevigata</i> . Ph. non Lam.					Pluk. alm. t. 277. f. 2.
4 parviflora. DC.	(ye.) small-flowered.	——	1823.	6. 9. H. ⓧ.	
5 sagittalis. DC.	(ye.) Virginian.	——	1731.	—— H. ⓧ.	Pluk. alm. t. 169. f. 6?
6 ovalis. DC.	(ye.) oval-leaved.	——	1827.	—— H. ⓧ.	Herm. lugd. p. 203. ic.
7 rubiginosa. DC.	(ye.) rusty.	E. Indies.	1807.	—— S. ⓧ.	
8 verrucosa. DC.	(bl.) blue-flowered.	——	1731.	—— S. ⓧ.	
<i>variega</i> . Jacq. ic. t. 144. <i>angulosa</i> . Cav. ic. 4. t. 321.					Andr. reposit. 308.
9 semperflorens. DC. (ye.) ever-blowing.	E. Indies.	1816.	3. 9. S. h.	Vent. hort. cels. t. 17.	
10 retusa. DC.	(ye.) wedge-leaved.	——	1731.	5. 9. S. ⓧ.	Bot. reg. 253.
11 Lechenaultii. DC.	(ye.) Lechenault's.	——	1826.	—— S. ⓧ.	
12 spectabilis. DC.	(pu.) showy.	——	1824.	—— S. ⓧ.	
13 pulcherrima. DC.	(ye.) Mysore.	——	1814.	—— S. ♀.	Bot. mag. 2027.
14 bengalensis. DC.	(ye.) Bengal.	——	1806.	—— S. ⓧ.	Pluk. alm. t. 169. f. 5.
15 junccea. DC.	(ye.) rush-stalked.	——	1700.	6. 9. S. ⓧ.	Bot. mag. 490.
16 tenuifolia. DC.	(ye.) slender-leaved.	——	1818.	—— S. ⓧ.	
17 fenestrata. DC.	(ye.) window-calyxed.	——	1816.	—— S. ⓧ.	
18 sericea. DC.	(ye.) silky.	——	1807.	—— S. ⓧ.	
19 tecta. DC.	(ye.) striped-flower'd.	——	1824.	6. 10. S. ⓧ.	
20 Burmanni. DC.	(ye.) Burmann's.	——	——	—— S. ⓧ.	
21 hirsuta. DC.	(ye.) hairy-podded.	——	1823.	—— S. ⓧ.	Burm. ind. t. 48. f. 1.
22 mysorensis. DC.	(ye.) Mysore.	——	1826.	—— S. ⓧ.	
23 montana. DC.	(ye.) mountain.	——	1816.	—— S. ♀.	
24 albida. DC.	(wh.) white-flowered.	——	1824.	—— S. ⓧ.	
25 paniculata. DC.	(ye.) paniced.	——	1807.	6. 8. S. h.	
26 pulchra. DC.	(ye.) short-podded.	——	1800.	3. 5. S. h.	
27 Növea-Holländia. DC. (pu.) New Holland.	N. Holland.	1820.	6. 8. S. h.		
28 paulina. DC.	(ye.) St. Paul.	Brazil.	——	7. 9. S. h.	Schr. pl. rar. h. mon. t. 88.
29 breviflora. DC.	(ye.) short-flowered.	——	1824.	—— S. h.	
30 bifaria. DC.	(ye.) two-ranked.	E. Indies.	——	—— S. h.	
31 dichotoma. DC.	(ye.) forked.	——	——	—— S. h.	

32 diffusa. DC.	(ye.) diffuse.	1818. 6. 8. S. ȝ.	Petiv. gaz. t. 30. f. 10.
33 biflora. DC.	(ye.) two-flowered. E. Indies. —	1790. — S. ⓧ.	Burm. ind. t. 48. f. 2.
34 nana. DC.	(ye.) dwarf. —	— S. ⓧ.	Andr. rep. t. 593.
35 bialata. DC.	(ye.) two-winged. —	1818. — S. ⓧ.	
36 tetragona. DC.	(ye.) square-stalked. Nepaul.	1806. 8. 9. S. ⓧ.	
37 linifolia. DC.	(ye.) Flax-leaved. E. Indies.	1823. 6. 7. S. ⓧ.	
38 pellita. DC.	(ye.) villous-podded. Jamaica.	1822. 7. 9. S. ȝ.	
39 virgultalis.	(ye.) twiggy. C. B. S.	1816. — G. ȝ.	
40 acuminata. DC.	(ye.) taper-pointed. —	— G. ȝ.	
41 spartoides. DC.	(ye.) Broom-like. —	— G. ȝ.	
42 thebaica. DC.	(ye.) stripe-flowered. Egypt.	1828. 7. 8. G. ȝ.	Delil. fl. eg. t. 37. f. 1.
43 nummularia. DC.	(ye.) money-wort-l'd. E. Indies.	— S. ȝ.	
44 tuberosa. DC.	(pu.) tuberous-rooted. Nepaul.	1816. 6. 7. S. ȝ.	
45 anthylloides. DC.	(ye.) large-calyxed. Java.	1789. 8. 9. S. ȝ.	Schr. pl. rar. mon. t. 12.
46 calycina. DC.	(st.) hispid-calyxed. E. Indies.	1818. 7. 9. S. ȝ.	
47 nepalensis. L.en.	(bl.) Nepaul.	1821. 6. 8. S. ⓧ.	Mart. acad. mun. 6.t.F.
48 Roxburghiana. DC.	(ye.) Roxburgh's. E. Indies.	1823. 7. 9. S. ȝ.	—— t. E.
49 chinensis. DC.	(ye.) Chinese. China.	1820. 6. 7. S. ȝ.	
50 hirta. DC.	(ye.) hairy. E. Indies.	1816. 6. 8. S. ⓧ.	Jacq. vind. 3. t. 64.
51 prostrata. DC.	(wh.) prostrate. —	1804. — S. ȝ.	
52 stipulacea. H.B.	(ye.) large-stipuled. —	1823. — S. ȝ.	Herb. amat. t. 238.
53 arborescens. DC.	(ye.) spreading. C. B. S.	1774. 6. 10. G. ȝ.	Rheed. mal. 9. t. 27.
<i>inanescens. w. capensis. Th.</i>			
54 turgida. DC.	(ye.) turgid-podded.	1820. — G. ȝ.	
55 laburnifolia. DC.	(ye.) Laburnum-l'd. E. Indies.	1739. 7. 9. S. ⓧ.	
56 pendula. DC.	(ye.) pendulous-pod'd. W. Indies.	1820. — S. ȝ.	
57 Brównei. DC.	(ye.) Browne's. Jamaica.	— S. ȝ.	
58 bracteata. DC.	(ye.) large-bracted. E. Indies.	1818. 5. 9. S. ȝ.	Mart. acad. mun. 6.t.H.
59 cytisoïdes. DC.	(ye.) Cytisus-like. Nepaul.	1823. 6. 8. S. ȝ.	6.t.G.
60 foliosa. DC.	(ye.) leafy. E. Indies.	1826. — S. ȝ.	
61 orixensis. DC.	(ye.) minute-flower'd. —	1816. 7. 8. S. ⓧ.	
62 virgata. DC.	(ye.) slender. —	1824. — S. ⓧ.	Bot. reg. 377.
63 purpurascens. DC.	(pu.) purplish. Mauritius.	1823. 8. 10. S. ⓧ.	
64 incána. DC.	(ye.) hoary. W. Indies.	1714. 6. 7. S. ⓧ.	
65 micans. DC.	(ye.) glittering. E. Indies.	1823. — S. ȝ.	
66 curtata. DC.	(ye.) short-keeled.	1820. — S. ⓧ.	
67 vitellina. DC.	(ye.) yellow-twigg'd. Brazil.	1819. 5. 7. S. ȝ.	Bot. reg. 447.
68 medicaginea. DC.	(ye.) Trefoil-leaved. E. Indies.	1818. 6. 7. S. ȝ.	Rott. n. act. nat. c.4.t.5.
69 Grahámi.	(ye.) Graham's. Mexico.	1824. 9. 12. S. ȝ.	Bot. mag. 2714.
<i>dichotoma B.M. non DC</i>			
70 Bojéri.	(ye.) Bojer's. Madagascar.	1828. S. ȝ.	
<i>dichotoma. Bojer. nec atiorum.</i>			
71 trichotoma. Boj.	(ye.) three-forked	— S. ȝ.	
72 stricta. DC.	(ye.) upright. E. Indies.	1826. 8. 10. S. ⓧ.	
73 purpurea. DC.	(pu.) purple-flowered. C. B. S.	1790. 3. 5. G. ȝ.	Bot. reg. 128.
74 microphylla. DC.	(ye.) small-leaved. Arabia.	1823. 8. 10. H. ⓧ.	
75 micrantha. DC.	(ye.) small-blossom'd. Ceylon.	1820. 6. 7. S. ⓧ.	
76 pallida. DC.	(st.) pale-flowered. Africa.	1775. — S. ⓧ.	
77 argentea. DC.	(ye.) silvery-leaved. C. B. S.	1823. — G. ȝ.	Jacq. schœnb. 2. t. 220.
78 pulchella. DC.	(ye.) large-flowered. —	1800. — G. ȝ.	Bot. mag. 1699.
79 Saltiana. A.R.	(ye.) Salt's. Abyssinia.	1810. 6. 7. S. ȝ.	Andr. deposit. 648.
80 angustifolia. DC.	(st.) narrow-leaved. C. B. S.	1815. 5. 9. G. ȝ.	Jacq. schœnb. 2. t. 219.
81 villosa. DC.	(ye.) villous. —	1823. — G. ȝ.	
82 latifolia.	(ye.) broad-leaved. E. Indies.	1825. — S. ȝ.	
83 elliptica. H.B.	(ye.) elliptic-leaved. China.	1823. 7. 9. G. ȝ.	
84 lanceolata. H.B.	(ye.) spear-leaved. W. Indies.	1806. — S. ȝ.	Dill. elth. t. 102. f. 131.
85 lotifolia. DC.	(ye.) Lotus-leaved. —	1732. 6. 7. S. ȝ.	
86 axillaris. DC.	(ye.) axillary-fl'd. Guinea.	1781. 7. 8. S. ⓧ.	Rheed. mal. 9. t. 28.
87 quinquefolia. DC.	(ye.) five-leaved. E. Indies.	1792. 6. 7. S. ⓧ.	Lodd. bot. cab. 1158.
<i>HYPOCALYPTUS. DC. HYPOCALYPTUS. Monadelphia Decandria.</i>			
obcordatus. DC.	(pu.) obcordate-leav'd. C. B. S.	1790. 4. 6. G. ȝ.	Lodd. bot. cab. 509.
<i>Crotalaria cordifolia. B.C.</i>			
VIBOR'GIA. DC.	VIBOR'GIA. Monadelphia Decandria.		
1 obcordata. DC.	(ye.) many-flower'd. C. B. S.	1812. 7. 9. G. ȝ.	
<i>Crotalaria floribunda. B.C. Anthyllis cuneata. Hort.</i>			
2 sericea. DC.	(ye.) silky. C. B. S.	1780. — G. ȝ.	
LODDIG'ESIA. DC.	LODDIG'ESIA. Monadelphia Decandria.		
oxalidifolia. DC.	(ro.) Oxalis-leaved. C. B. S.	1802. 5. 9. G. ȝ.	Bot. mag. 965.
DICH'ILUS. DC.	DICH'ILUS. Monadelphia Decandria.		
lebeckoïdes. DC.	(st.) Lebeckia-like. C. B. S.	1816. 6. 8. G. ȝ.	DC. leg. mem. 6. t. 35.

LEBE'CKIA. DC.	LEBE'CKIA.	Monadelphia Decandria.				
1 subnuda. DC.	(ye.) few-leaved.	C. B. S.	1823.	4. 6.	G. h.	
2 contaminata. DC.	(ye.) narrow-leaved.		1787.	—	G. h.	Pluk. alm. t. 424. f. 1.
3 sepiaria. DC.	(ye.) filiform-leaved.		1816.	—	G. h.	
4 subternata. DC.	(ye.) various-leaved.		—	—	G. h.	
5 sericea. DC.	(ye.) silky-leaved.		1774.	—	G. h.	Com. hort. amst. 2.t.107.
6 cytoides. DC.	(ye.) Cytisus-like.		—	—	G. h.	
SARCOPHY'LLUM. DC.	SARCOPHY'LLUM.	Monadelphia Decandria.				
carnosum. DC.	(ye.) fleshy-leaved.	C. B. S.	1812.	5. 8.	G. h.	
ASPA'LATHUS. DC.	ASPA'LATHUS.	Monadelphia Decandria.				
1 spinosa. DC.	(ye.) spiny.	C. B. S.	1823.	5. 8.	G. h.	Breyn. cent. t. 26.
2 aculeata. DC.	(ye.) yellow-spined.		—	—	G. h.	
3 acuminate. DC.	(ye.) taper-pointed.		1816.	—	G. h.	Lam. ill. t. 620. f. 4.
4 laricifolia. DC.	(ye.) Larch-leaved.		1825.	—	G. h.	
5 capitata. DC.	(ye.) headed.		1823.	6. 7.	G. h.	Lam. ill. t. 620. f. 2.
6 astroites. DC.	(ye.) spiny-bracted.		1825.	5. 8.	G. h.	Seba, thes. t. 24. f. 6.
7 chenopoda. DC.	(or) cluster-leaved.		1759.	7. 8.	G. h.	Bot. mag. 2225.
8 albens. DC.	(wh.) white-leaved.		1774.	—	G. h.	
9 argyræa. DC.	(wh.) silvery.		1818.	—	G. h.	
10 hystrix. DC.	(wh.) silky-spined.		1824.	—	G. h.	
11 thymifolia. DC.	(ye.) Thyme-leaved.		1825.	5. 8.	G. h.	
12 ericifolia. DC.	(ye.) Heath-leaved.		1780.	7. 8.	G. h.	
13 hispida. DC.	(ye.) hispid.		1823.	—	G. h.	
14 asparagoides. DC.	(ye.) Asparagus-like.		1812.	—	G. h.	
15 multiflora. DC.	(ye.) many-flowered.		1818.	—	G. h.	
16 uniflora. DC.	(ye.) one-flowered.		1812.	—	G. h.	
17 carnosa. DC.	(ye.) fleshy-leaved.		1795.	5. 8.	G. h.	
18 pinguis. DC.	(ye.) succulent-leaved.		1818.	7. 8.	G. h.	
19 crassifolia. DC.	(ye.) thick-leaved.		1800.	—	G. h.	
20 affinis. DC.	(ye.) likened.		1816.	—	G. h.	
21 genistoidea. DC.	(ye.) Genista-like.		1825.	6. 8.	G. h.	
22 squarrosa. DC.	(ye.) squarrose.		—	—	G. h.	
23 galoides. DC.	(ye.) Galium-like.		1816.	7. 8.	G. h.	
24 pinea. DC.	(ye.) Pine-leaved.		—	—	G. h.	
25 abietina. DC.	(ye.) Fir-leaved.		1826.	—	G. h.	
26 araneosa. DC.	(pa.) cobweb.		1795.	6. 7.	G. h.	Bot. mag. 829.
27 globosa. A.R.	(ye.) globe-headed.		1802.	—	G. h.	Andr. reposit. 302.
28 ciliaris. DC.	(ye.) ciliated.		1790.	7. 8.	G. h.	Bot. mag. 2233.
29 quinquefolia. DC.	(br.) five-leaved.		1816.	—	G. h.	
30 candicans. DC.	(ye.) white.		1774.	—	G. h.	
31 heterophylla. DC.	(ye.) various-leaved.		1828.	—	G. h.	
32 argentea. DC.	(ye.) silvery-leaved.		1759.	—	G. h.	
33 subulata. DC.	(ye.) awl-leaved.		1789.	—	G. h.	
34 sericea. DC.	(ye.) silky.		1816.	—	G. h.	
35 calliosa. DC.	(ye.) oval-spiked.		1812.	—	G. h.	Bot. mag. 2329.
36 fusca. DC.	(ye.) brown.		1816.	—	G. h.	
37 mucronata. DC.	(ye.) thorny-branch'd.		1796.	6. 7.	G. h.	
38 pedunculata. DC.	(ye.) small-leaved.		1775.	7. 8.	G. h.	Bot. mag. 344.
39 bracteata. DC.	(ye.) bracteate.		1816.	—	G. h.	
'ULEX. DC.	FURZE.	Monadelphia Decandria.				
1 europaeus. DC.	(ye.) common.	Britain.	3. 7.	H. h.	Eng. bot. 742.
2 nanus. DC.	(ye.) dwarf.		1. 12.	H. h.	— t. 743.
3 provincialis. DC.	(ye.) Provence.	S. Europe.	1815.	6. 8.	H. h.	Loisel. not. 105. t. 6.f. 2.
STAURACA'NTHUS. DC.	STAURACA'NTHUS.	Monadelphia Decandria.				
aphyllus. DC.	(ye.) leafless.	Portugal.	1815.	6. 8.	H. h.	
SPA'RTIUM. DC.	SPANISH BROOM.	Monadelphia Decandria.				
juncinum. DC.	(ye.) common.	S. Europe.	1548.	7. 9.	H. h.	Bot. mag. 85.
<i>Spartianthus juncineus</i> . L. en.						
<i>S. flôre pleno</i> .	(ye.) double-flowered.		—	H. h.	
GENISTA. DC.	GENI'STA.	Monadelphia Decandria.				
1 parviflora. DC.	(ye.) small-flowered.	Levant.	1739.	5. 7.	H. h.	Venten. cels. t. 87.
<i>Spartium parviflorum</i> . Vent.						
2 clavata. DC.	(ye.) clavate.	Barbary.	1812.	5. 8.	F. h.	Venten. hort. cels. t. 17.
<i>Spartium sericeum</i> . Vent. non Ait.						
3 canariensis. DC.	(ye.) Canary.	Canaries.	1656.	5. 9.	F. h.	Bot. reg. t. 217.
4 candicans. DC.	(ye.) hoary.	Spain.	1735.	4. 7.	H. h.	Wats. dend. brit. t. 80.
5 pâtens. DC.	(ye.) spreading.		1825.	5. 7.	H. h.	Cavan. ic. 2. t. 176.
6 linifolia. DC.	(ye.) Flax-leaved.		1739.	1. 6.	F. h.	Bot. mag. 442.
<i>Spartium linifolium</i> . Desf. atl. 2. t. 181.						

LEGUMINOSÆ.

7 triquètra. DC.	(ye.) triangular.	Corsica.	1770.	5. 6.	H. \ddot{h} .	Wats. dend. brit. t. 79'
8 bracteolàta. DC.	(ye.) bracteolate.	1826.	—	H. \ddot{h} .	
9 microphylla. DC.	(ye.) small-leaved.	Canaries.	1824.	—	F. \ddot{h} .	
10 umbellàta. DC.	(ye.) umbelled.	Barbary.	1799.	4. 6.	F. \ddot{h} .	Desf. atl. 2. t. 180.
<i>Spártium umbellátum</i> . Desf.						
11 radiàta. DC.	(ye.) rayed.	Italy.	1758.	5. 7.	H. \ddot{h} .	Bot. mag. 2260.
<i>Spártium radiátum</i> . B.M. Mill. ic. t. 249. f. 1.						
12 hórrida. DC.	(ye.) large-spined.	Pyrenees.	1826.	5. 7.	H. \ddot{h} .	Gilib. b. prat. 2.p. 239. ic.
13 lusitânica. DC.	(ye.) Portugal.	Portugal.	1771.	3. 5.	H. \ddot{h} .	Andr. deposit. t. 419.
14 Lobélia. DC.	(ye.) Lobel's.	S. Europe.	1826.	4. 6.	H. \ddot{h} .	Lob. adv. p. 409. icon.
15 ferox. DC.	(ye.) sharp-spined.	Barbary.	1800.	6. 7.	F. \ddot{h} .	Desf. atl. 2. t. 182.
16 Cupâni. DC.	(ye.) Cupani's.	Naples.	1826.	—	H. \ddot{h} .	Cup pan. ed. 1.v. 2.t. 233.
17 triacánthos. DC.	(ye.) three-spined.	Portugal.	1821.	5. 6.	H. \ddot{h} .	Brot. phyt 130. t. 54.
18 cuspidòs. DC.	(ye.) sharp-pointed.	C. B. S.	1816.	—	G. \ddot{h} .	
19 tricuspidàta. DC.	(ye.) three-pointed.	Algiers.	1826.	—	F. \ddot{h} .	Desf. alt. 2. t. 183.
20 gibraltárica. DC.	(ye.) Gibraltar.	Gibraltar.	—	—	F. \ddot{h} .	Jacq. ic. rar. t. 557.
21 sylvestris. DC.	(ye.) wood.	Hungary.	1824.	6. 8.	H. \ddot{h} .	
<i>hispanica</i> . Jacq. non L.						
22 aegyptiaca. DC.	(ye.) Egyptian.	Egypt.	1828.	—	H. \ddot{h} .	Brot. phyt. 133. t. 55.
23 falcata. DC.	(ye.) falcate-podded.	Portugal.	1821.	5. 7.	H. \ddot{h} .	
24 còrsica. DC.	(ye.) Corsican.	Corsica.	1826.	—	H. \ddot{h} .	Wats. dend. brit. t. 78.
25 Scórpius. DC.	(ye.) Scorpion.	S. Europe.	1570.	3. 4.	H. \ddot{h} .	Cavan. ic. 3. t. 211.
26 hispanica. DC.	(ye.) dwarf prickly.	Spain.	1759.	6. 8.	H. \ddot{h} .	Eng. bot. 132.
27 ánglica. DC.	(ye.) petty whin.	Britain.	5. 8.	H. \ddot{h} .	Fuchs. hist. 220. ic.
28 germánica. DC.	(ye.) German.	Germany.	1773.	6. 8.	H. \ddot{h} .	Bull. herb. t. 115.
29 púrgans. DC.	(ye.) purging.	S. France.	1768.	6. 7.	H. \ddot{h} .	Wats. dend. brit. t. 76.
30 cinérea. DC.	(ye.) ash-coloured.	—	1818.	5. 7.	H. \ddot{h} .	Bot. mag. 2265.
31 virgata. DC.	(ye.) long-twigg'd.	Madeira.	1777.	3. 6.	F. \ddot{h} .	
32 congesta. DC.	(ye.) close-branched.	Teneriffe.	4. 7.	F. \ddot{h} .	Jacq. ic. rar. 3. t. 556.
33 sericea. DC.	(ye.) silky-leaved.	Austria.	1812.	5. 6.	H. \ddot{h} .	Pall. it. 3. app. t. 5.f. 2.
34 aphylla. DC.	(vi.) violet-coloured.	Volga.	1818.	6. 7.	H. \ddot{h} .	
<i>Spártium aphyllum</i> . L. en.						
35 monospermá. DC. (wh.)	single-seeded.	S. Europe.	1690.	—	F. \ddot{h} .	Bot. mag. 683.
<i>Spártium monospermum</i> . B.M.						
36 sphærocárpa. DC.	(ye.) round-podded.	—	1731.	—	F. \ddot{h} .	Renealm. spec. t. 33.
37 aethnénis. DC.	(ye.) three-seeded.	Mt. Etna.	1822.	6. 8.	F. \ddot{h} .	Bot. mag. 2674.
<i>Partium aetnense</i> . B.M.						
38 tetragôna. DC.	(ye.) four-sided.	Poland.	1826.	—	H. \ddot{h} .	Viv. frag. ital. 1. t. 8.
39 depréssâ. DC.	(ye.) depressed.	Tauria.	1828.	—	H. \ddot{h} .	Eng. bot. 44.
40 scariòsa. DC.	(ye.) scariose.	Italy.	1820.	6. 8.	H. \ddot{h} .	
41 anxântica. DC.	(ye.) smooth.	—	—	—	H. \ddot{h} .	
42 tinctòria. DC.	(ye.) Dyer's green-weed.	Britain.	—	H. \ddot{h} .	
β latifolia. DC.	(ye.) broad-leaved.	Switzerland.	1824.	—	H. \ddot{h} .	
γ hirsuta. DC.	(ye.) villous.	—	—	H. \ddot{h} .	
δ praténis. DC.	(ye.) meadow.	Italy.	—	H. \ddot{h} .	
43 sibírica. DC.	(ye.) Siberian.	Siberia.	1785.	6. 8.	H. \ddot{h} .	Jacq. vind. 2. t. 190.
44 polygalæfólia. DC.	(ye.) Milkwort-leav'd.	Portugal.	1826.	—	H. \ddot{h} .	Poll. fl. veron. 2. t. 4.f. 7.
45 flòrida. DC.	(ye.) Spanish.	Spain.	1752.	—	H. \ddot{h} .	Wats. dend. brit. t. 77.
46 mántica. DC.	(ye.) prostrate.	Italy.	1824.	—	H. \ddot{h} .	
47 ovâta. DC.	(ye.) oval-leaved.	Hungary.	1816.	—	H. \ddot{h} .	
48 pátula. DC.	(ye.) spreading.	Siberia.	1828.	—	H. \ddot{h} .	
49 triangulàris. DC.	(ye.) three-sided.	Hungary.	1815.	5. 6.	H. \ddot{h} .	
50 sagittális. DC.	(ye.) jointed.	Germany.	1570.	—	H. \ddot{h} .	
51 diffusa. DC.	(ye.) diffuse.	Italy.	1816.	—	H. \ddot{h} .	
<i>Spártium decumbens</i> . Jacq. non Ait.						
52 prostrâta. DC.	(ye.) trailing.	Italy.	1826.	—	H. \ddot{h} .	Reyn. mem. 1.p. 211. ic.
53 procumbens. DC.	(ye.) procumbent.	Hungary.	1816.	6. 8.	H. \ddot{h} .	Bot. reg. 1150.
54 pilòsa. DC.	(ye.) hairy green-weed.	England.	5. 6.	H. \ddot{h} .	Eng. bot. 44.
55 micrânta. DC.	(ye.) small-blossom'd.	Spain.	1824.	—	H. \ddot{h} .	Orteg. dec. 6. t. 10. f. 1.
56 pilocárpa. DC.	(ye.) hairy-podded.	1820.	5. 7.	H. \ddot{h} .	
<i>CYTISUS</i> . DC.						
1 nubigènus. DC.	(wh.) cluster-flower'd.	Teneriffe.	1779.	5. 8.	H. \ddot{h} .	Duham. arb. t. 23.
2 álbüs. DC.	(wh.) white-flowered.	Portugal.	1752.	6. 7.	H. \ddot{h} .	
<i>Spártium multiflorum</i> . W.						
β incarnâtus B.C.	flesh-coloured.	—	1823.	—	H. \ddot{h} .	Lodd. bot. cab. 1052.
3 Laburnum. DC.	(ye.) Laburnum.	Europe.	1596.	5. 6.	H. \ddot{h} .	Bot. mag. 176.
4 alpinus. DC.	(ye.) Scotch Laburnum.	Hungary.	6. 7.	H. \ddot{h} .	W. et K. hung. 3. t. 260.
5 nigrans. DC.	(ye.) black-rooted.	Austria.	1730.	—	H. \ddot{h} .	Bot. reg. 802.
6 sessilifolius. DC.	(ye.) common.	Italy.	1029.	5. 6.	H. \ddot{h} .	Bot. mag. 255.

7	triflorus. DC.	(ye.) three-flowered. Spain.	1640.	6. 7.	H. h.	Clus. hist. 1. p. 94. f. 3.
8	mollis. DC.	(ye.) soft-leaved.	1820.	—	H. h.	
9	pátem. DC.	(ye.) woolly-podded. Portugal.	1752.	—	H. h.	
10	grandiflorus. DC.	(ye.) great-flowered.	1816.	—	H. h.	
11	scoparius. DC.	(ye.) common Broom. Britain.	4. 6.	H. h.	Eng. bot. 1330.
	<i>Spártium scopárium</i> . E.B.					
12	spinósus. DC.	(ye.) prickly. S. Europe.	1596.	6. 7.	F. h.	Lobel. ic. 2. t. 95.
13	lanigerus. DC.	(ye.) villous.	1822.	—	F. h.	
	<i>Spartium lanigerum</i> . Df. <i>Calyctome villosa</i> . L.en.					
14	proliferus. DC.	(wh.) silky. Canaries.	1779.	4. 5.	G. h.	Bot. reg. 121.
15	leucanthus. DC.	(wh.) cream-colour'd. Hungary.	1806.	6. 7.	H. h.	Bot. mag. 1438.
16	álbidus. DC.	(wh.) whitish. S. Europe.	1824.	—	F. h.	
17	purpureus. DC.	(pu.) purple-flower'd. Austria.	1792.	5. 8.	H. h.	Bot. mag. 1176.
	β álbis. DC.	(wh.) white-flowered.	—	—	H. h.	
18	biflorus. DC.	(ye.) two-flowered. Hungary.	1760.	5. 7.	H. h.	Bot. reg. 308.
19	elongátus. DC.	(ye.) long-branched.	1804.	—	H. h.	Wats. dend. brit. 82.
20	multiflorus. B.R.	(ye.) many-flowered.	1824.	—	H. h.	Bot. reg. 1191.
21	falcatus. DC.	(ye.) sickle-shaped.	1816.	6. 8.	H. h.	W. et K. hung. 3.t.238.
22	austriacus. DC.	(ye.) Austrian. Austria.	1741.	6. 9.	H. h.	Jacq. aust. 1. t. 21.
23	supinus. DC.	(ye.) trailing. S. Europe.	1755.	5. 8.	H. h.	— 1. t. 20.
24	hirsutus. DC.	(ye.) hairy.	1739.	6. 8.	H. h.	Jacq. obs. 4. t. 96.
25	serotinus. DC.	(ye.) late-flowering. Hungary.	1826.	—	H. h.	
26	capitátus. DC.	(ye.) cluster-flower'd. Austria.	1774.	6. 7.	H. h.	Jacq. aust. t. 33.
27	ciliátus. DC.	(ye.) fringed. Mt. Carpathian.	1824.	—	H. h.	
28	polytrichus. DC.	(ye.) many-haired. Tauria.	1822.	—	H. h.	Lobel. ic. 2. p. 41. f. 2.
29	argénteus. DC.	(ye.) silver-leaved. France.	1739.	7. 9.	H. h.	Lodd. bot. cab. t. 673.
30	calycinæ. DC.	(ye.) dwarf. Levant.	1816.	6. 8.	H. h.	
	nánus. Wats. dend. brit. t. 81.					
31	orientális. DC.	(ye.) oriental.	5. 7.	G. h.	
32	canéscens. DC.	(ye.) hoary.	1826.	—	H. h.	
33	prócerus. DC.	(ye.) tall. Portugal.	1816.	6. 7.	H. h.	
	ADENOCA'RPUS. DC. ADENOCA'RPUS. <i>Monadelphia Decandria</i> .					
1	hispanicus. DC.	(ye.) Spanish. Spain.	1824.	5. 7.	F. h.	Clus. hist. 1. p. 94. f. 1.
2	intermèdius. DC.	(ye.) intermediate. S. Europe.	7. 8.	H. h.	
3	parvifolius. DC.	(ye.) small-leaved. France.	4. 7.	H. h.	Du.arb.ed.n.5.t.47.f.2.
4	telonénensis. DC.	(ye.) loose-flowered. S. Europe.	1824.	5. 7.	H. h.	
	<i>Cytisus divaricatus</i> . β . Bot. mag. 1387?					
5	frankenioides. DC.	(ye.) Frankenia-like. Teneriffe.	1815.	—	G. h.	
	<i>Genista viscosa</i> . w. <i>Adenocarpus anagyrsus</i> . s.s.					
6	foliolosus. DC.	(ye.) leafy. Canaries.	1770.	7. 8.	F. h.	Bot. mag. 426.
	ON'ONIS. DC. REST-HARROW. <i>Monadelphia Decandria</i> .					
1	críspa. DC.	(ye.) curl-leaved. Spain.	1739.	6. 8.	F. h.	Rœm. arch. 1. p. 3. t. 1.
2	hispanica. DC.	(ye.) Spanish.	1799.	5. 9.	F. h.	Bot. mag. 2450.
3	vaginális. DC.	(ye.) sheathed. Egypt.	1815.	6. 8.	F. h.	Vent. cels. t. 32.
4	Nátrix. DC.	(ye.) yellow-shrubby. S. Europe.	1683.	5. 9.	F. h.	Bot. mag. 329.
5	picta. DC.	(std.) painted-flower'd. Barbary.	1825.	6. 8.	H. 2.	Desf. atl. 2. t. 137.
6	pínguis. w.	(ye.) greasy.	1739.	7. 8.	F. h.	
7	longifolia. DC.	(ye.) long-leaved. Canaries.	1816.	5. 6.	F. h.	
8	ramosíssima. DC.	(ye.) branching. Barbary.	1825.	7. 8.	H. h.	Desf. atl. 2. t. 186.
9	arenária. DC.	(ye.) sand. S. France.	—	H. h.	
10	biflora. DC.	(ye.) two-flowered. Barbary.	1824.	—	H. \odot .	
11	viscosa. DC.	(ye.) clammy. S. France.	1750.	—	H. \odot .	Barrel. ic. t. 1239.
12	brachycarpa. DC.	(ye.) short-podded. Spain.	1824.	—	H. \odot .	
13	brevifolia. DC.	(ye.) short-leaved. S. Europe.	—	H. \odot .	
14	sicula. DC.	(ye.) Sicilian. Sicily.	1822.	6. 8.	H. \odot .	
15	pubescens. DC.	(ye.) downy. S. Europe.	1680.	—	H. \odot .	Brot. phyt. 140. t. 58.
	<i>arthropoda</i> . Brot.					
16	ornithopodioides. DC.	(ye.) Bird's-foot. Sicily.	1713.	—	H. \odot .	Cavan. ic. 2. t. 192.
17	rotundifolia. DC.	(re.) round-leaved. Switzerland.	1570.	5. 7.	H. h.	Bot. mag. 335.
18	fruticosa. DC.	(re.) shrubby. S. Europe.	1680.	5. 7.	H. h.	— t. 317.
19	tridentata. DC.	(re.) three-toothed. Spain.	1752.	6. 8.	F. h.	Cavan. ic. 2. t. 152.
20	angustissima. DC.	(re.) narrow-leaved.	1826.	—	F. h.	
21	cenisia. DC.	(re.) crested. Italy.	1759.	—	H. \ddagger .	Barrel. ic. t. 1104.
22	fœtida. DC.	(re.) strong-scented. Barbary.	1804.	7. 10.	H. \odot .	
23	geminiflora. DC.	(re.) twin-flowered. Spain.	1825.	8. 10.	H. \odot .	
24	Schouwii. DC.	(re.) Schouw's. Italy.	1828.	7. 10.	H. \odot .	
25	reclinata. DC.	(re.) reclined. S. Europe.	1800.	6. 8.	H. \odot .	
26	mollis. DC.	(re.) soft. Spain.	1824.	—	H. \odot .	
27	Cherléri. DC.	(re.) dwarf. S. Europe.	1771.	6. 7.	H. \odot .	Bauh. hist. 2. p. 394.f.2

28 péninsula. DC.	(re.) pendulous.	Barbary.	1826.	7. 9. H. ⊖.	Desf. atl. 2. t. 191.
29 Siebéri. DC.	(re.) Sieber's.	Crete.	—	— H. ⊖.	
30 arboréscens. DC.	(re.) tree.	Barbary.	1825.	6. 8. F. ™.	Desf. atl. 2. t. 193.
31 altissima. DC.	(re.) stinking.	Italy.	1596.	5. 8. H. ¼.	All. ped. t. 41. f. 1.
	<i>hircina.</i> Jacq. vind. t. 93.				
32 procúrrens. DC.	(re.) rooting-branch'd.	Europe.	6. 8. H. ™.	Fuchs. hist. 60. ic.
33 répens. W.	(re.) creeping.	Britain.	— H. ¼.	Dill. elth. t. 25. f. 28.
34 spinosa. DC.	(re.) Cammock.	—	— H. ™.	Eng. bot. 682.
	<i>arvénensis.</i> E.B.				
35 antiquorum. W.	(re.) tall.	S. Europe.	1790.	— H. ¼.	
36 mitissima. DC.	(re.) cluster-flower'd.	Portugal.	1732.	6. 7. H. ⊖.	Dill. elth. t. 24. f. 27.
37 diffusa. DC.	(re.) diffuse.	Italy.	1824.	7. 8. H. ⊖.	
38 serrata. DC.	(pu.wh.) serrate.	Egypt.	1828.	6. 10. H. ⊖.	
39 hírta. DC.	(bl.) hairy.	Levant.	—	— H. ¼.	
40 alopecuroídes. DC.	(re.) Fox-tail.	Portugal.	1696.	7. 8. H. ⊖.	Schkuhr. handb. t. 194.
41 álba. DC.	(wh.) white-flowered.	Naples.	1826.	— H. ⊖.	
42 oligophylla. DC.	(re.w.) few-leaved.	—	1824.	— H. ⊖.	Ten. fl. nap. 2. t. 67.
43 capitata. DC.	(ye.) capitate.	Spain.	1818.	6. 7. H. ¼.	Cavan. ic. 2. t. 159. f. 2.
44 striata. DC.	(ye.) striated.	S. France.	1814.	— H. ¼.	
45 Colúmnæ. DC.	(ye.) small-flowered.	Europe.	1732.	— H. ¼.	All.p.1.n.1166.t.20.f.3.
	<i>minutissima.</i> Jacq. aust. t. 240. <i>parviflora.</i> P.S.				
46 minutissima. DC.	(ye.) pointed-calyx'd.	S. Europe.	1818.	6. 8. H. ⊖.	Cavan. ic. 2. t. 153.
	<i>barbata.</i> Cav. <i>saxatilis.</i> Lam.				
47 variegata. DC.	(ye.) variegated.	—	1784.	7. 8. H. ⊖.	Desf. atl. 2. t. 185.
48 pallida. DC.	(pa.) pale.	—	1824.	— H. ⊖.	
49 emarginata. Boj. emarginate.	Mauritius.	—	— S. ™.	
50 villosa. DC.	(ye.) villous.	C. B. S.	1825.	7. 9. G. ™.	
51 decumbens. DC.	(ye.) decumbent.	—	1822.	— G. ¼.	
52 geminata. DC.	(ye.) twin-podded.	—	1787.	— G. ¼.	
53 umbellata. DC.	(ye.) umbel-flowered.	—	1818.	— G. ¼.	
54 glabra. DC.	(ye.) smooth-leaved.	—	1816.	— G. ™.	
55 capensis. DC.	(pu.) Cape annual.	—	1823.	— H. ⊖.	
56 cernua. DC.	(ye.) hanging-podded.	—	1774.	— G. ™.	
REQUIE'NIA. DC.	REQUIE'NIA.	Monadelphia	Decandria.		
1 obcordata. DC.	(ye.) obcordate.	Senegal.	1824.	6. 8. S. ™.	DC. leg. mem. t. 37.
2 sphæropérma. DC.	(ye.) round-seeded.	C. B. S.	1816.	4. 7. G. ™.	— t. 38.
ANTHYLLIS. DC.	KIDNEY VETCH.	Monadelphia	Decandria.		
1 Gerárdi. DC.	(ro.) Gerard's.	S. Europe.	1806.	6. 8. H. ¼.	Ger. gallopr. t. 18.
2 onobrychioídes. DC.	(y.) Saintfoin-like.	Spain.	1818.	— H. ¼.	Cavan. ic. 2. t. 150.
3 cytisoides. DC.	(ye.) downy-leaved.	—	1731.	4. 6. F. ™.	Barrel. ic. 1182.
4 genistæ. DC.	(ye.) Genista-like.	—	1824.	— F. ™.	
5 Hermánniæ. DC.	(ye.) Lavender-ld.	Levant.	1739.	4. 7. F. ™.	Bot. mag. 2576.
6 aspálathi. DC.	(ye.) spiny.	—	—	— F. ™.	
7 erinácea. DC.	(bl.) prickly.	Spain.	1759.	4. 6. H. ™.	Bot. mag. 676.
8 hy'strix.	(wh.) porcupine.	S. Europe.	—	— H. ™.	
9 Bárba-Jòvis.	(st.) Jupiter's-beard.	—	1640.	3. 5. G. ™.	Bot. mag. 1927.
10 heterophylla. DC.	(v.) various-leaved.	—	1768.	6. 7. G. ™.	
11 sericea. DC.	(ye.) silky.	Spain.	1786.	— F. ™.	Swt. br. fl. gar. 1. t. 79.
12 montana. DC.	(ro.) mountain.	Europe.	1759.	— H. ¼.	Eng. bot. 104.
13 Vulneraria. DC.	(ye.) common.	Britain.	5. 8. H. ¼.	Dill. elth. t. 320. f. 413.
14 rústica. Mil.	(wh.) white-flower'd.	—	— H. ¼.	
15 Dilléni. Sct.	(sc.) red-flowered.	S. Europe.	— H. ¼.	Bot. mag. 108.
16 polypylla. Kit.	(ye.) many-leaved.	Hungary.	1827.	— H. ¼.	Cavan. ic. 1. t. 39. f. 2.
17 tetraphylla. DC.	(st.) four-leaved.	S. Europe.	1640.	7. 8. H. ⊖.	Cavan. ic. 1. t. 40.
18 cornicina. DC.	(st.) horned.	Spain.	1759.	— H. ⊖.	
19 hamosa. DC.	(st.) hooked.	Barbary.	1820.	— H. ⊖.	
20 lotoídes. DC.	(st.) Lotus-like.	Spain.	1739.	6. 7. H. ⊖.	
21 chilensis. DC.	(st.) Chile.	Chile.	1824.	— G. ¼.	
	<i>Lòtus subpinnatus.</i> Lag.				

Subtribus II. TRIFOLIEÆ. DC. prodr. 2. p. 171.

MEDICAGO. DC. MEDICK. Diadelphia Decandria.

1 circinnata. DC. (ye.) kidney-podded. Italy. 1640. 7. 8. H. ⊖.

Hymenocárpus circinnatus. L. en.

2 nummulària. DC. (ye.) money-wort. <i>circinnata</i> . G. <i>Hymenocárpos nummulárius</i> . L. en.	S. Europe.	— H. ⊙.	Gært. sem. 2. t. 155.f.7.
3 radiàta. DC. (ye.) ray-podded.	Italy. 1629.	6. 7. H. ⊙.	Moris. his. s. 2. t. 15. f. 3.
4 rupéstris. DC. (ye.) rock.	Tauria. 1827.	6. 8. H. 4.	
5 brachycárpa. DC. (ye.) short-podded.	Tiffin. 1818.	7. 8. H. ⊙.	
6 lupulina. DC. (ye.) nonesuch.	Britain.	5. 8. H. 3.	Eng. bot. 971.
7 cretacea. DC. (ye.) suffrutescent.	Tauria. 1805.	7. 8. H. 2.	Eng. bot. 1016.
8 falcata. DC. (ye.) sickle-podded.	Britain.	— H. 4.	
9 procumbens. DC. (ye.) procumbent.	Poland. 1827.	— H. 4.	
10 cancellata. DC. (ye.) netted-podded.	Caucasus. 1828.	— H. 4.	
11 suffruticosa. DC. (vi.ye.) frutescient.	Pyrenees. 1818.	6. 8. H. 2.	
12 arborea. DC. (ye.) moon trefoil.	Italy. 1596.	5. 11. H. 2.	Flor græc. 767.
13 sibírica. DC. (ye.) Siberian.	Siberia. 1824.	6. 9. H. ⊙.	
14média. DC. (bl.ye.) pale-flowered.	Europe.	6. 7. H. 4.	
15 sativa. DC. (bl.) Lucerne.	Britain.	— H. 4.	Eng. bot. 1749.
β versicolor. DC. (bl.ye.) various-colour'd.	—	— H. 2.	
16 mexicana. (bl.) Mexican.	Mexico. 1824.	5. 8. F. 2.	
17 prostrata. DC. (ye.) prostrate.	Hungary. 1793.	7. 8. H. 4.	Jacq. vind. 1. t. 89.
18 glomerata. DC. (ye.) clustered.	Italy. 1818.	— H. 4.	
19 glutinosa. DC. (vi.) glutinous.	Caucasus.	— H. 4.	
20 obscura. DC. (ye.) doubtful.	S. France. 1734.	— H. ⊙.	Retz. obs. fasc. 1. t. 1.
21 lœvis. DC. (ye.) smooth.	Tunis. 1816.	6. 8. H. ⊙.	
Hælix. w.			
22 orbicularis. DC. (ye.) flat-podded.	S. Europe. 1688.	7. 8. H. ⊙.	Gært. sem. 2. t. 155.
23 marginata. DC. (ye.) margined.	—	— H. ⊙.	Moris. his. s. 2. t. 15. f. 2.
24 applanata. DC. (ye.) obovate-leaved.	—	1824.	
25 scutellata. DC. (ye.) snail-podded.	S. Europe. 1562.	6. 8. H. ⊙.	Gært. sem. 2. t. 155.f.7.
26 rugosa. DC. (ye.) elegant.	—	1680.	Moris. his. s. 2. t. 15. f. 4.
élégans. w.			
27 saxatilis. DC. (ye.) stone.	Tauria. 1828.	6. 8. H. 4.	
28 tornata. DC. (ye.) smooth-podded.	S. Europe. 1658.	— H. ⊙.	
29 turbinata. DC. (ye.) Turban.	—	1680.	
30 tuberculata. DC. (ye.) wart-podded.	—	1658.	Moris. his.s. 2. t. 15. f. 5.
31 striata. DC. (std.) stripe-podded.	—	1828.	Bauh. hist. 2. f. 1.
32 apiculata. DC. (ye.) tufted.	Italy. 1800.	— H. ⊙.	
33 coronata. DC. (ye.) crowned.	S. Europe. 1660.	— H. ⊙.	
34 catalònica. DC. (ye.) Catalonian.	Catalonia. 1816.	— H. ⊙.	Schrank. h. mon. t. 28.
35 denticulata. DC. (ye.) toothed.	S. France. 1800.	— H. ⊙.	
36 flexuosa. DC. (ye.) flexuose.	Naples. 1824.	— H. ⊙.	
37 spinulosa. DC. (ye.) prickly-podded.	S. Europe. 1820.	— H. ⊙.	
38 pubescens. DC. (ye.) pubescent.	Montpelier.	1823.	
39 terebellum. DC. (ye.) short-spined.	S. Europe. 1798.	— H. ⊙.	
40 marina. DC. (ye.) sea-side.	—	1596.	Cavan. ic. 2. t. 130.
41 tentaculata. DC. (ye.) distichous-spin'd.	—	1824.	Gært. sem. 2. t. 155.f.7?
42 Hornemanniana. DC. (ye.) Hornemann's.	Morocco.	1823.	
43 littoralis. DC. (ye.) shore.	S. Europe. 1827.	— H. 4.	J. Bauh. his. 2.p. 385. ic.
44 lappacea. DC. (ye.) bur-podded.	—	— H. ⊙.	
45 pentacycla. DC. (ye.) five-whorled.	—	— H. ⊙.	
46 hístrix. DC. (ye.) porcupine.	Naples. 1824.	— H. ⊙.	
47 disciformis. DC. (ye.) hook-spined.	Montpelier.	1826.	
48 carstiensis. DC. (ye.) creeping-rooted.	Carinthia.	1789.	Bot. mag. 909.
49 nigra. DC. (ye.) black.	France.	—	Moris. his. s. 2. t. 15. f. 19.
50 triboloides. DC. (ye.) Caltrops-like.	S. Europe.	1739.	
51 cylindracea. DC. (ye.) cylinder-podded.	—	1824.	
52 græca. DC. (ye.) Grecian.	Greece. 1804.	— H. ⊙.	
53 minima. DC. (ye.) least.	England.	— H. ⊙.	Flor. dan. t. 211.
54 arenaria. DC. (ye.) sand.	Italy. 1824.	— H. ⊙.	
55 muricóleptis. DC. (ye.) arch-spined.	Sicily.	—	
56 uncinata. DC. (ye.) hooked.	S. Europe. 1820.	— H. ⊙.	
57 distans. DC. (ye.) distant.	—	— H. ⊙.	Moris. his.s. 2. t. 15. f. 21.
58 aculeata. DC. (ye.) prickly.	—	1802.	
59 maculata. DC. (ye.) heart.	England.	5. 6. H. ⊙.	Eng. bot. 1616.
polymórfha. E.B.			
60 Gerárdi. DC. (ye.) Gerard's.	Hungary. 1816.	6. 8. H. ⊙.	Moris. his. s. 2. t. 15. f. 18.
villlosa. DC. fl. fr. 4. p. 545.			
61 agréstis. DC. (ye.) field.	Italy. 1824.	— H. ⊙.	
62 rigidula. DC. (ye.) thorny-podded.	S. Europe. 1730.	— H. ⊙.	
63 muricata. DC. (ye.) prickly.	England.	5. 8. H. ⊙.	Vaill. par. t. 33. f. 7.
64 sphærocárpos. DC. (ye.) round-podded.	Italy. 1826.	6. 8. H. ⊙.	Sebast. pl. rom.p.15 t. 3.

65 mûrex. DC.	(ye.) Burdock-podded,	1802.	6. 8. H. ⊖.	Breyn. cent. 81. t. 34.
66 laciñata. DC.	(ye.) cut-leaved.	S. Europe.	1683.	— H. ⊖.	Jacq. coll.sup.t.15. f. 2.
67 Tenoreâna. DC.	(ye.) Tenore's.	Naples.	1827.	— H. ⊖.	Moris. his. s. 2.t.15. f. 8.
68 granaténsis. DC.	(ye.) Spanish.	Spain.	1816.	— H. ⊖.	— s. 2.t.15. f. 7.
69 intertéxta. DC.	(ye.) hedgehog.	S. Europe.	1629.	— H. ⊖.	— s. 2.t.15. f. 9.
70 ciliáris. DC.	(ye.) fringed.	France.	1686.	7. 8. H. ⊖.	
71 échinus. DC.	(ye.) Sea-egg.	S. Europe.	1820.	6. 8. H. ⊖.	
72 ægagróphila. DC.	(ye.) goats.	1824.	7. 9. H. ⊖.	
73 cásrica. L.en.	(ye.) Caspian.	Caucasus.	1820.	6. 8. H. ⊖.	
74 ovâta. H.E.	(ye.) oval-leaved.	S. Europe.	1818.	— H. ⊖.	
75 strumâria. H.E.	(ye.) strumous.	—	—	— H. ⊖.	
76 microdon. E.	(ye.) small-toothed.	Egypt.	1826.	— H. ⊖.	
77 pedunculâta. E.	(ye.) peduncled.	—	—	— H. ⊖.	
TRIGONE'LLA. DC.	FENUGREEK.	Diadelphia Decandria.			
1 cœrulea. DC.	(bl.) blue.	Switzerland.	1562.	7. 9. H. ⊖.	Bot. mag. 2283.
	<i>Trifolium cœruleum.</i> B.M. <i>Melilotus cœrulea.</i> Lam.				
2 Besseriana. DC.	(bl.) Besser's.	Poland.	—	— H. ⊖.	
3 marítima. DC.	(ye.) sea-side.	Egypt.	1827.	— H. ⊖.	
4 uncinâta. DC.	(st.) hooked.	Iberia.	1823.	6. 9. H. ⊖.	
	<i>Melilotus hamosum.</i> M.B. <i>uncinata.</i> Bess.				
5 littoralis. DC.	(ye.) shore.	Sicily.	1826.	— H. ⊖.	
6 calliceras. DC.	(ye.) rostrate.	Iberia.	1821.	— H. ⊖.	DC. hort. gen. 2. ic.
	<i>oxyrhynchus.</i> F.				
7 prostrâta. DC.	(ye.) prostrate.	S. France.	—	— H. ⊖.	J.Bauh.his.2.p.365.f.2.
8 gladiâta. DC.	(wh.) sword-podded.	Tauria.	1829.	— H. ⊖.	
9 Fœnum-græcum. DC.	(st.) common.	S. France.	1597.	— H. ⊖.	Schk. s. han. 2. t. 211.
10 spinôsa. DC.	(ye.) thorny.	Candia.	1710.	7. 9. H. ⊖.	Breyn.cent.79.t.33.f.1.
11 striâta. DC.	(ye.) streaked.	Abyssinia?	1824.	6. 9. H. ⊖.	Alp. ægypt. t. 124.
12 hamosa. DC.	(ye.) hook-podded.	Egypt.	1640.	7. 8. H. ⊖.	
13 flexuosa. DC.	(st.) flexuose.	—	1818.	6. 8. H. ⊖.	
14 Fischeriana. DC.	(ye.) Fischer's.	Tiflin.	—	— H. ⊖.	
	<i>flexuosa.</i> M.B. non Delil.				
15 anguina. DC.	(ye.) snake-podded.	Egypt.	1828.	— H. ⊖.	Delil. fl.egypt.t.110.f.2.
16 tenuis. DC.	(ye.) slender.	Tiflin.	1818.	— H. ⊖.	
17 elongâta. W. en.	(ye.) elongated.	—	1823.	— H. ⊖.	
18 cancellâta. DC.	(ye.) netted-podded.	—	1818.	— H. ⊖.	
19 monspeliaca. DC.	(ye.) Montpelier.	Montpelier.	1710.	— H. ⊖.	W. et K. hung. 2.t. 142.
20 parviflôra. DC.	(ye.) small-flowered.	Switzerland.	1824.	— H. ⊖.	
21 pinnatifida. DC.	(ye.) pinnatifid.	Spain.	1801.	— H. ⊖.	Cavan. ic. 1. t. 38.
	<i>mèdia.</i> Delil.				
22 polycerata. DC.	(ye.) broad-leaved.	S. Europe.	1640.	7. 9. H. ⊖.	
23 ægyptiaca. DC.	(ye.) Egyptian.	Egypt.	1823.	6. 8. H. ⊖.	
24 ornithory'ncchos. DC.	(ye.) beak-podded.	Russia.	1825.	6. 9. H. ⊖.	Lodd. bot. cab. 1391.
25 crassifolia. DC.	(ye.) thick-leaved.	Egypt.	1828.	6. 8. H. ⊖.	
26 ruthénica. DC.	(ye.) small.	Siberia.	1741.	5. 7. H. ♀.	Gmel. sib. 4. t. 9.
27 laciñata. DC.	(ye.) laciniate.	Egypt.	1823.	7. 9. H. ⊖.	DC. ic. rar. 1. t. 29.
28 platycápos. DC.	(ye.) round-leaved.	Siberia.	1741.	6. 9. H. ♀.	Eng. bot. t. 1047.
29 hy'brida. DC.	(ye.) hybrid.	France.	1806.	— H. ♀.	
30 ornithopodioides. DC.	(re.) Bird's-foot.	Britain.	—	— H. ⊖.	
	<i>Trifolium ornithopodioides.</i> E.B. <i>Melilotus ornithopodioides.</i> P.S.				
31 corniculâta. DC.	(ye.) horse-shoe.	S. Europe.	1597.	6. 7. H. ⊖.	Moris. his.s. 2.t.16.f.11.
32 esculenta. DC.	(ye.) esculent.	E. Indies.	1815.	6. 9. S. ⊖.	
POCO'CKIA. DC.	Poco'ckia.	Diadelphia Decandria.			
crética. DC.	(ye.) Cretan.	Candia.	1713.	6. 9. H. ⊖.	Moris. his. s. 2. t. 14.f.3.
	<i>Trifolium créticum.</i> L. <i>Melilotus crética.</i> P.S.				
MELIL'OTUS. DC.	MELLILOT.	Diadelphia Decandria.			
1 Kochiâna. DC.	(ye.) Koch's.	Germany.	1816.	6. 9. H. ♂.	
2 dentâta. DC.	(ye.) toothed.	Hungary.	1802.	6. 8. H. ♀.	W. et K. hung. 1. t. 42.
3 lineâris. DC.	(ye.) linear-leaved.	Spain.	1816.	— H. ⊖.	
4 ruthénica. DC.	(wh.) Russian.	Sarepta.	1828.	— H. ⊖.	
5 melanospérma. DC.	(ye.) black-seeded.	Tauria.	1825.	— H. ♂.	
6 officinalis. DC.	(ye.) common.	Britain.	—	6. 9. H. ⊖.	Eng. bot. 1340.
7 palústris. DC.	(ye.) marsh.	Hungary.	1824.	— H. ♂.	W. et K. hung. 3.t.266.
8 altíssima. DC.	(wh.) tallest.	S. Europe.	1818.	— H. ⊖.	
9 leucântha. DC.	(wh.) white-flowered.	Europe.	—	— H. ♂.	Sturm. deuts. 1. f. K.
	<i>vulgâris.</i> W.en.				
10 macrorhiza. DC.	(ye.) long-rooted.	Hungary.	1801.	7. 9. H. ♀.	W. et K. hung. 1. t. 26.
11 parviflôra. DC.	(ye.) small-flowered.	S. Europe.	1810.	6. 8. H. ⊖.	
12 rugulosa. W.en.	(wh.) rugged-podded.	E. Indies.	—	— H. ⊖.	

13	indica. W. en.	(ye.) Indian.	E. Indies.	1680.	6. 8. H. ⓧ.	Pluk. alm. t. 45. f. 4.
14	Bámeti. h.	(ye.) one-seeded.	1818.	— H. ⓧ.	
15	mínima. DC.	(wh.) small-flowered.	E. Indies.	1810.	8. 10. H. ⓧ.	
16	segetális. DC.	(ye.) hedge.	Spain.	1824.	6. 8. H. ⓧ.	
17	pólonica. DC.	(ye.) Polish.	Poland.	1778.	— H. ⓧ.	
18	táurica. DC.	(wh.) Taurian.	Tauria.	1823.	6. 9. H. ⓧ.	Camer. hort. t. 29.
19	ítalica. DC.	(ye.) Italian.	Italy.	1596.	— H. ⓧ.	
20	rotundifòlia. T.N.	(ye.) round-leaved.	1824.	— H. ⓧ.	
21	grácilis. DC.	(ye.) slender.	France.	—	6. 8. H. ⓧ.	
22	neapolitana. T.P.	(ye.) Neapolitan.	Naples.	1827.	— H. ⓧ.	
23	pállida. DC.	(ye.) pale-podded.	Poland.	1828.	— H. ⓧ.	
24	suavéolens. DC.	(ye.) sweet-scented.	Dahuria.	1826.	— H. 4.	
25	hirsuta. Bal.	(ye.) hairy.	—	— H. ⓧ.	
26	procumbens. H.Pr.	(ye.) procumbent.	1829.	— H. ⓧ.	
27	arvénis. DC.	(ye.) corn-field.	Germany.	1822.	6. 9. H. ⓧ.	
28	petitpierriana.w.	(wh.) rough-podded.	1816.	— H. ⓧ.	
29	Besseriana. DC.	(ye.) Besser's. imbricata. Bes.	Tauria.	1826.	6. 8. H. ⓧ.	
30	plicata. F.	(ye.) plaited.	1823.	— H. ⓧ.	
31	messanénis. DC.	(ye.) Sicilian.	Sicily.	1680.	— H. ⓧ.	
32	sulcata. DC.	(ye.) furrowed.	Barbary.	1798.	— H. ⓧ.	
	mauritánica. W. enum.					
TRIFOLIUM. DC.						
	TREFOIL. Diadelphia Decandria. L.					
1	angustifólium. DC. (ro.)	narrow-leaved.	S. Europe.	1640.	6. 8. H. ⓧ.	Barrel. ic. t. 698.
2	intermédia. DC. (wh.)	intermediate.	Italy.	1829.	— H. ⓧ.	
3	púrpureum. DC. (pu.)	purple.	Montpelier.	1826.	— H. ⓧ.	Loisel. fl gall. t. 14.
4	rúbens. DC.	(re.) long-spiked.	S. Europe.	1633.	6. 9. H. 4.	Jacq. aust. 4. t. 385.
5	cœrulescens. DC. (pu.)	bluish.	Tauria.	1827.	— H. ⓧ.	
6	incarnátum. DC. (sc.)	scarlet.	Italy.	1596.	7. 9. H. ⓧ.	Bot. mag. 328.
7	Molinieri. w.	(bh.) blush-flowered.	1823.	— H. ⓧ.	
8	arvénse. DC.	(wh.) Hare's-foot.	Britain.	7. 8. H. ⓧ.	Engl. bot. v. 14. t. 944.
9	grálice. P.S.	(re.) slender.	France.	1818.	— H. ⓧ.	Barrel. ic. 901.
10	ligósticum. DC.	(re.) awned.	Portugal.	1820.	6. 8. H. ⓧ.	Balb. act. ac. it. 1. f. 2.
	arrectisétum. Brot. phyt. t. 63. f. 1. aristátum. H.					
11	divaricátum. DC. (wh.)	spreading.	1827.	— H. ⓧ.	
12	gemélum. DC.	(wh.) double-headed.	Spain.	1825.	— H. ⓧ.	
13	phleódes. DC.	(re.) Phleum-spiked.	—	— H. ⓧ.	
14	lappáceum. DC. (st.)	burr-spiked.	S. Europe.	1787.	— H. ⓧ.	Barrel. ic. 871.
15	echinátum. DC. (wh.)	bristly.	Caucasus.	1823.	— H. ⓧ.	
16	erináceum. DC.	(wh.) villous-leaved.	Iberia.	1827.	— H. ⓧ.	
17	malacántum. DC. (wh.)	soft-headed.	1824.	— H. ⓧ.	
18	sylváticum. DC. (pu.)	wood.	S. France.	1826.	— H. 4.	
19	Boccóni. DC.	(re.) Boccone's.	S. Europe.	1825.	7. 8. H. ⓧ.	Brot. phyt. t. 63. f. 2.
	semiglábrum. Brot. collínium. Poir.					
20	striátum. DC.	(pu.) soft-knotted.	Britain.	6. 7. H. ⓧ.	Eng. bot. v. 26. t. 1843.
21	tenuifòrum. DC.	(pa.) slender-flower'd.	Naples.	1824.	6. 8. H. ⓧ.	
22	scábrum. DC.	(wh.) rough.	Britain.	5. 7. H. ⓧ.	Eng. bot. v. 13. t. 903.
23	marítimum. DC. (li.)	sea-shore.	6. 8. H. ⓧ.	v. 4. t. 220.
	rígidum. Savi. fl. pis. 2. p. 159. t. 1. f. 1.					
24	supinum. DC.	(bh.) spreading-branch'd.	S. Europe.	1827.	— H. ⓧ.	Savi. obs. trif. p. 46. f. 2.
25	constantinopolitánum. DC.	(ye.) Turkish.	Constantinople.	—	— H. ⓧ.	
26	cíncum. DC.	(st.) girded.	Montpelier.	1820.	— H. ⓧ.	
27	alexandrinum. DC. (st.)	Egyptian.	Egypt.	1798.	— H. ⓧ.	
28	ochroleúcum. DC. (st.)	sulphur-color'd.	Britain.	— H. 4.	Eng. bot. v. 17. t. 1224.
29	tricocéphalum. DC. (st.)	hairy-headed.	Caucasus.	1829.	— H. 4.	
30	canéscens. DC. (wh.)	gray.	1803.	5. 7. H. 4.	Bot. mag. 1168.
31	pannónicum. DC. (wh.)	Hungarian.	Hungary.	1752.	6. 8. H. 4.	Jacq. obs. 2. t. 42.
32	barbátum. DC.	(ye.) bearded.	Montpelier.	1827.	— H. 4.	
33	olympicum. B.M.	(st.) long-flowered.	Levant.	1810.	— H. 4.	Bot. mag. 2790.
34	arménium. DC.	(st.) Armenian.	Armenia.	1816.	5. 7. H. 4.	
35	squarrosum. DC. (wh.)	various-leaved.	Spain.	1640.	7. 8. H. ⓧ.	Moris. his. s. 2. t. 13. f. 1.
	β flavicans. DC. (st.)	yellowish.	— H. ⓧ.	Savi. obs. tri. p. 65. t. 3.
36	álbidum. DC.	(st.) whitish.	1796.	— H. ⓧ.	
37	cónicum. DC.	(st.) conical.	1823.	6. 8. H. ⓧ.	
38	Kitaibeliana. DC.	Kitaibel's.	Hungary.	1828.	— H. ⓧ.	
39	elongátum. DC.	(re.) long-flowered.	Galatia.	1810.	— H. 4.	
40	alpéstre. DC.	(pu.) oval-spiked.	Europe.	1789.	— H. 4.	Bot. mag. 2779.
41	médium. DC.	(pu.) Cow-grass.	Britain.	7. 8. H. 4.	Eng. bot. v. 3. t. 190.
42	bracteátum. DC.	(pu.) large-bracted.	Morocco.	1804.	6. 8. H. ⓧ.	
				K 4		

LEGUMINOSÆ.

43· expánsum. DC. (pu.) expanded.	1804.	6. 7.	H. ¼.	
44 pratense. DC. (pu.) common clover.	Britain.	5. 9.	H. ¼.	Eng. bot. v. 25. t. 1770.
a perenne.	(pu.) perennial.	—	H. ¼.	Sincl. gr. wob. 221.
β sativum.	(pu.) cultivated.	—	H. ½.	
45 vaginátum. Schl. (ye.) sheathed.	Switzerland.	1824.	—	H. ¼.	
pratense β flávicans. Ser. in DC. prodr. 2. p. 195. ochroleucum. γ. DC.					
46 microphyllum. DC. (p.) small-leaved.	France.	1824.	5. 9.	H. ¼.	Sturm deuts. 1. fas. 16.
47 nòricum. DC. (st.) shaggy.	Carinthia.	1823.	6. 7.	H. ¼.	
48 pensylvanicum. DC. (p.) Buffalo-clover.	N. America.	1811.	6. 9.	H. ¼.	W. et K. hung. 1. t. 36.
49 pálidum. DC. (wh.) pale.	Hungary.	1803.	6. 8.	H. ♂.	— 1. t. 50.
50 diffusum. DC. (pu.) diffuse.	1801.	—	H. ♂.	Desf. atl. 2. t. 209. f. 1.
51 hirtum. DC. (pu.) hairy.	S. Europe.	1806.	—	H. ♂.	
hispidum. Desf.					
52 pictum. W.	(pu.) painted-leaved.	—	—	H. ♂.	
53 pauciflórum. DC. (pu.) few-flowered.	Archipelago.	1828.	—	H. ♂.	Barrel. ic. 859.
54 Cherléri. DC. (st.) hispid-headed.	S. Europe.	1750.	5. 7.	H. ♂.	Alpin. exot. t. 306.
55 globosum. DC. (wh.) globular-headed.	Levant.	1713.	6. 8.	H. ♂.	Eng. bot. v. 22. t. 1545.
56 clypéatum. DC. (wh.) oriental.	1711.	—	H. ♂.	
57stellátum. DC. (ro.) star-headed.	England.	7. 8.	H. ♂.	W. et K. hung. 3. t. 269.
58 leucánthum. DC. (wh.) white-flowered.	Tauria.	1823.	6. 8.	H. ♂.	
59 Waldsteiniánum. (pu.) Waldstein's.	Hungary.	1826.	—	H. ♂.	All ped. t. 59. f. 3.
reflexum. W. K. non L.					Eng. bot. v. 15. t. 1049.
60 saxátile. DC. (wh.) rock.	Switzerland.	1816.	6. 8.	H. ♂.	— v. 15. t. 1063.
61 suffocátem. DC. (wh.) suffocated.	England.	—	H. ♂.	Sturm deuts. 1. fasc. 15.
62 glomerátum. DC. (ro.) round-headed.	6. 7.	H. ♂.	
63 parviflórum. DC. (li.) small-flowered.	Hungary.	1805.	7. 8.	H. ♂.	W. et K. hung. 1. t. 37.
strictum. Sturm.					Eng. bot. v. 25. t. 1769.
64 stríctum. DC. (bh.) upright.	S. Europe.	—	—	H. ♂.	
65 rēpens. DC. (wh.) white clover.	Britain.	5. 9.	H. ¼.	Schrank pl. rar. t. 47.
β pentaphyllum. P.S. (w.) purple 5-leaved.		—	H. ¼.	Sturm.deuts.fl.1.fas.15.
66 anómala. DC. (wh.) anomalous.	1827.	—	H. ¼.	Vill. delph. 3. t. 41.
67 palléscens. DC. (st.) pale.	Hungary.	1804.	6. 8.	H. ¼.	W. et K. hung. 1. t. 27.
68 cæpítosum. DC. (wh.) turfy.	Switzerland.	1815.	—	H. ¼.	W. hort. berol. t. 108.
69 angulátum. DC. (bh.) angular.	Hungary.	1803.	—	H. ♂.	Sturm deuts.fl.1.fas.15.
70 suavéolens. DC. (pu.) sweet-scented.	S. Europe.	1818.	—	H. ♂.	Mich. n. gen. t. 25. f. 25.
71 hybrídum. DC. (wh.) hybrid.	1805.	—	H. ♂.	Savi. fl. pi. 2. t. 1. f. 2.
72 Micheliana. DC. (st.) Michel's.	Italy.	1815.	—	H. ♂.	
73 élégans. DC. (li.) elegant.	S. Europe.	1818.	—	H. ¼.	
formosum. Savi. Vail. par. t. 22. f. 1.					Flor. dan. t. 1172.
74 reflexum. DC. (pu.) reflexed.	Virginia.	1794.	—	H. ¼.	
75 montanum. DC. (wh.) mountain.	Europe.	1786.	7. 8.	H. ¼.	Eng. bot. v. 15. t. 1048.
76 rupéstre. DC. (wh.) flat-headed.	Italy.	1824.	—	H. ¼.	W. et K. hung. 2. t. 165.
77 Prutetiánum. DC. (re.) soft hairy.	1828.	—	H. ¼.	
78 subterrâneum. DC. (w.) subterraneous.	England.	5. 6.	H. ♂.	
79 vesiculósom. DC. (re.) recurved.	Hungary.	1805.	6. 8.	H. ♂.	Cupan. panp. t. 69. f. 1.
recúrveum. W.K.					Sturm deuts.fl.1.fas.16.
80 spumósom. DC. (pu.) bladdered.	France.	1771.	—	H. ♂.	Eng. bot. v. 15. t. 1050.
81 resupinátum. DC. (pu.) resupinate.	Germany.	1713.	—	H. ♂.	Barrel. ic. 864.
82 fragíferum. DC. (li.) Strawberry.	England.	7. 8.	H. ¼.	Cupan.panp. t. 97. f. 1?
83 tomentósom. DC. (ye.) woolly.	S. Europe.	1640.	6. 8.	H. ♂.	Buxb. cent. 3. t. 31. f. 1.
84 phýsodes. DC. (li.) swollen-calyx'd.	Iberia.	1829.	—	H. ¼.	— 3. p. 18.t.31.f.2.
85 alátum. DC. (li.) netted-calyx'd.	Sicilý.	1820.	7. 8.	H. ♂.	
Cupáni. Tineo.					Sturm.deut.fl.1.fasc.15.
86 uníflórum. DC. (re.) one-flowered.	S. Europe.	1822.	6. 8.	H. ¼.	Kth. pl. legum. t. 53.
β Sternbergianum. DC. (wh.) Sternberg's.	—	—	—	H. ¼.	Bonan. sic. t. 245.
87 exímium. DC. (pu.) choice.	Dahuria.	1828.	—	H. ¼.	Bot. reg. 1070.
β albiflórum. DC. (wh.) white-flowered.	Altay.	—	—	H. ¼.	
88 alpínum. DC. (re.wh.) alpine.	Italy.	1775.	—	H. ¼.	Sturm deuts.fl.1.fasc.16.
89 involucrátum. DC. (re.) involucred.	Mexico.	1825.	—	H. ♂.	— fl. 1. fasc. 16.
90 Gussóni. DC. (pu.) Gussone's.	Sicily.	1822.	—	H. ¼.	Bot. mag. 557.
91 fimbriátum. B.R. (re.) fringed.	N. America.	1826.	9. 10.	H. ¼.	
92 tridentátum. B.R. (re.) three-toothed.	—	—	8. 10.	H. ¼.	Labill. dec. 5. t. 10.
93 cyathíferum. B.R. (re.) cup-like.	—	—	—	H. ¼.	
94 bádiúm. DC. (br.) villous-stalked.	Pyrenees.	6. 8.	H. ¼.	
95 agríarium. DC. (ye.) golden.	Europe.	1815.	—	H. ♂.	
96 spádiceum. DC. (ye.) bay-coloured.	—	1778.	—	H. ¼.	
97 decípiens. DC. (ye.) decipient.	—	1827.	6. 8.	H. ♂.	
98 speciósom. DC. (br.) large-flowered.	Candia.	1752.	—	H. ♂.	
comósom. Labil.					

99 procumbens. DC. (st.)	Hop.	Britain.	6. 7. H. ⊙.	Eng. bot. v. 14. t. 945.
100 parisiense. DC.	Paris.	France.	— H. ⊙.	Sturm deuts. 1. fas. 16.
pâtens. Sturm.					
101 Sebastiâni. DC.	(ye.) Sebastian's.	Italy.	1823.	— H. ⊙.	Sebas. pl. rom. fas. 2.t.4.
102 flavescens. DC.	(ye.) yellowish.	Sicily.	1828.	6. 8. H. ⊙.	Eng. bot. v. 18. t. 1256.
103 mînus. E.B.	(ye.) lesser yellow.	Britain.	— H. ⊙.	— v. 18. t. 1257.
104 filiforme. E.B.	(ye.) slender yellow.		— H. ⊙.	
105 comosum. W.	(wh.) tufted.	N. America.	1798.	— H. ¼.	
106 tenuifolium. DC.	(wh.) slender-leaved.	Naples.	1824.	— H. ⊙.	
107 capense. DC.	(st.) Cape.	C. B. S.	1825.	— G. ¼.	
108 congestum. DC.	(wh.) close-headed.	Sicily.	1829.	— H. ⊙.	
109 formosum. DC.	(pi.) handsome.	Isl. Melo.	1828.	— H. ¾.	
110 lasiocéphalum. DC.	(w.) woolly-headed.	C. B. S.	1825.	— H. ⊙.	
111 villosum. Pr.	(wh.) villous.	Sicily.	1818.	5. 7. H. ⊙.	
112 plicatum. Pr.	(br.) plaited.		1824.	6. 8. H. ⊙.	
113 Ehrenbergii. H. ber.	(w.) Ehrenberg's.	Egypt.	1826.	— H. ⊙.	
114 libanaticum. E.	(wh.) Lebanon.	Mt. Lebanon.	1827.	— H. ⊙.	
115 albicans. E.	(wh.) white-headed.	Egypt.	1829.	— H. ⊙.	
LUPINA STER. MH.	LUPINA STER.	Diadelphia Decandria.			
1 pentaphyllus. Mh. (re.)	five-leaved.	Siberia.	1741.	7. 8. H. ¼.	Bot. mag. 879.
Trifolium Lupinaster. B.M.	Péntaphyllo Lupináster. P.S.				
2 albens. F.	(wh.) white-flowered.	Volga.	1827.	— H. ¼.	
3 purpurascens. F.	(pu.) purple.		1821.	6. 8. H. ¼.	
4 cœrulescens. F.	(pu.) blue purple.		1826.	— H. ¼.	
DORY'CNIUM. DC.	DORY'CNIUM.	Diadelphia Decandria.			
1 réctum. DC.	(ro.) upright.	S. Europe.	1640.	6. 8. H. ½.	Barrel. ic. t. 544.
Lótus réctus. w.					
2 latifolium. DC.	(wh.) broad-leaved.	Iberia.	1818.	6. 9. H. ½.	
3 hirsutum. DC.	(ro.) hairy.	S. Europe.	1683.	— H. ½.	Bot. mag. 336.
Lótus hirsutus. B.M.					
4 sericeum.	(bh.) silky.		1820.	— H. ½.	
Lótus sericeus. DC. cat. h. monsp. p. 122.					
5 argenteum. DC.	(ye.) silvery.	Egypt.	1825.	— H. ½.	
6 parviflorum. DC.	(ye.) small-flowered.	S. Europe.	1824.	7. 8. H. ⊙.	Delil. fl. æg. 113. t. 40.
7 subbiflorum. DC.	(ye.) two-flowered.	Spain.	1820.	6. 8. H. ⊙.	Desf. atl. 2. t. 211.
8 herbaceum. DC.	(wh.) herbaceous.	S. Europe.	1802.	6. 9. H. ¼.	Vill. dauph. 3. t. 41.
9 suffruticosum. DC.	(wh.) suffrutescent.		1640.	7. 9. F. ½.	Lobel. ic. 2.p. 51. f. 1. 2.
10 microphyllum.	(ro.) small-leaved.	C. B. S.	1826.	— G. ½.	Bot. mag. 2808.
Lótus microphyllus. B.M.	2808.				
L'OTUS. DC.	BIRD'S-FOOT TREFOIL. Diadelphia Decandria.				
1 édulis. DC.	(ye.) esculent.	S. Europe.	1710.	7. 8. H. ⊙.	Cavan. ic. 2. t. 157.
2 ornithopodioïdes.	(ye.) claw-podded.		1683.	6. 8. H. ⊙.	— 2. t. 163.
3 peregrinus. DC.	(ye.) flat-podded.		1713.	7. 8. H. ⊙.	
4 indicus. DC.	(ye.) Indian.	E. Indies.	— H. ⊙.	
5 tetraphyllum. DC.	(ye.) four-leaved.	Archipelago.	1828.	— H. ⊙.	Pluk. phyt. t. 200. f. 7.
6 flexuosum. DC.	(ye.) flexuous.		1820.	6. 8. H. ⊙.	Viv. fl. lib. t. 17. f. 13.
7 glaucus. DC.	(ye.) glaucous.	Madeira.	1777.	— G. ¾.	
8 sessilifolius. DC.	(ye.) sessile leaved.	Canaries.	1816.	6. 9. G. ¼.	
9 anthylloides. DC.	(ye.) Anthyllis-like.	C. B. S.	1812.	— G. ½.	Venten. malm. t. 92.
10 atropurpureum. DC.	(da.) dark-purple.	Cap. Verd Isl.	1823.	3. 9. G. ½.	
11 jacobæus. DC.	(da.) dark-flowered.		1714.	1. 12. G. ½.	Bot. mag. 79.
12 arenarius. DC.	(ye.) sand.	Portugal.	1824.	6. 8. H. ⊙.	
13 créticus. DC.	(ye.) silver-leaved.	Levant.	1680.	6. 9. G. ½.	Cavan. ic. 2. t. 156.
β varians. DC.	(ye.) variable.		1829.	— G. ½.	
14 cytisoides. DC.	(ye.) downy.	S. Europe.	1752.	7. 8. H. ¼.	Barrel. ic. t. 1031.
15 Dioscòridis. DC.	(ye.) Dioscorides's.	Crete.	1658.	6. 8. H. ⊙.	All. ped. 1. t. 59. f. 1.
16 Gebélia. DC.	(ro.) Aleppo.	Aleppo.	1823.	5. 9. F. ¼.	Vent. h. cels. t. 57.
17 arábicus. DC.	(re.) red-flowered.	Arabia.	1773.	7. 9. F. ¼.	Jacq. vind. t. 155.
18 australis. DC.	(re.) New Holland.	N. S. W.	1803.	5. 9. G. ½.	Bot. mag. 1365.
19 Candolleanus. ed. 1. (re.) De Candolle's.			1812.	6. 9. G. ½.	
australis β angustifoliola. DC.					
20 albidus. B.C.	(wh.) white-flowered.		1823.	— G. ½.	Lodd. bot. cab. 1063
21 obovatus. ed. 1.	(ye.) glaucous obovate.		1803.	4. 9. G. ¼.	
22 decumbens. DC.	(ye.) decumbent.	France.	1820.	5. 7. H. ¼.	
23 Forstéri.	(ye.) Forster's.	Britain.	— H. ¼.	
decumbens. E.F. non DC.					
24 pilosissimus. DC.	(ye.) hairy.	S. France.	1826.	6. 8. H. ⊙.	
25 hispidus. DC.	(ye.) hispid.	S. Europe.	1820.	— H. ⊙.	Loisel. fl. gal. t. 16.
26 suavolens. DC.	(ye.) fragrant.		1804.	— H. ¼.	

LEGUMINOSÆ.

27 angustissimus. DC. (ye.) narrow-podded. S. Europe.	5. 7.	H. ♂.	Bauh. hist. 2. p. 356.f.2.
28 diffusus. DC. (ye.) slender-podded. England.	—	H. ♂.	Eng. bot. v. 13. t. 925.
29 ciliatus. DC. (ye.) ciliated. Italy. 1820.	6. 8.	H. ♂.	W. et K. hung. 3. t. 229.
30 gracilis. DC. (ye.) slender. Hungary. 1818.	5. 7.	H. ♂.	
31 lanuginosus. DC. (ro.) drooping-fld. Levant. 1818.	7. 8.	G. ♀.	
32 comibricensis. DC. (bh.) Portuguese. Portugal. 1800.	6. 8.	H. ♂.	Brot. phyt. t. 53.
33 glaberrimus. DC. (bh.) smooth.	1820.	—	Bot. mag. 1233.
34 odoratus. DC. (ye.) sweet-scented. Barbary. 1804.	—	H. ♀.	Eng. bot. v. 30. t. 2090.
35 corniculatus. E.B. (ye.) common. Britain.	—	H. ♀.	— v. 30. t. 2091.
36 major. E.B. (ye.) large hairy.	—	H. ♀.	
37 tenuis. W.K. (ye.) slender. Hungary. 1816.	—	H. ♀.	Cavan. ic. 2. t. 164.
depresso. W. enum. humifusus. W. en. ex. Link enum.			
38 pedunculatus. DC. (ye.) long-peduncled. Spain. 1814.	—	H. ♀.	
39 palustris. DC. (ye.) marsh. Levant. 1824.	—	H. ♀.	Bot. mag. 151.
TETRAGONO'LOBUS. DC. WINGED PEA. Diadelphia Decandria.			Desf. atl. t. 210.
1 purpureus. DC. (pu.) purple-flower'd. S. Europe. 1596.	6. 8.	H. ♂.	Jacq. aust. 4. t. 361.
Lotus tetragonolobus. B.M.			Flor. dan. 800.
2 biflorus. DC. (ye.) two-flowered. Sicily. 1754.	—	H. ♂.	
3 conjugatus. DC. (pu.) twin-podded. Montpelier. 1759.	—	H. ♂.	
4 siliquosus. L.en. (ye.) square-podded. S. Europe. 1683.	7. 8.	H. ♀.	
5 maritimus. L. en. (ye.) sea-side. —	—	H. ♀.	

Subtribus III. CLITORIÆ. DC. prodr. p. 216.

PSOR'ALEA. DC.	PSOR'ALEA.	Diadelphia Decandria. L.	
1 odoratissima. DC. (bl.) sweet-scented. C. B. S.	1795.	5. 10.	G. ♂.
pinnata. Andr. rep. 474. nec aliorum.			Jacq. schoenb. 2. t. 229.
2 arborea. B.M.	(bl.) tree.	1814.	—
3 pinnata. DC.	(bl.) winged-leaved.	1690.	5. 7.
4 laevigata. DC.	(pu.) smooth.	1824.	—
5 verrucosa. DC.	(bl.) warted.	1774.	5. 8.
angustifolia. Jacq.			Jacq. schoenb. 2. t. 226.
6 filiformis. DC.	(bl.) slender.	1821.	—
7 axillaris. DC.	(bl.) axillary-flower'd.	1816.	6. 7.
8 tenuifolia. DC.	(bl.) slender-leaved.	1793.	—
9 fascicularis. DC.	(bl.) fascicle-flowered.	1816.	—
10 multicaulis. DC.	(v.) many-stalked.	1793.	8. 10.
11 aphylla. DC.	(bl.) leafless.	1790.	6. 8.
12 lathyrifolia. DC.	(bl.) Lathyrus-leaved.	1823.	—
13 decumbens. DC.	(bl.) decumbent.	1774.	4. 6.
14 repens. DC.	(bl.) creeping	—	7. 8. G. ♂.
15 hirta. DC.	(bl.) hairy.	1713.	5. 9. G. ♂.
16 Jacquiniana.	(bl.) Jacquin's.	—	— G. ♂.
hirta. Jacq. nec aliorum.			Jacq. schoenb. 2. t. 228.
17 aculeata. DC.	(bl.) prickle-leaved.	1774.	—
18 bracteata. DC.	(bl.) oval-spiked.	1781.	6. 8. G. ♂.
19 involucrata. DC.	(bl.) large-bracted.	1824.	7. 8. G. ♂.
20 spicata. DC.	(bl.wh.) long-spiked.	1774.	—
21 Stachydis. DC.	(bl.) Stachy's-like.	1793.	4. 6. G. ♂.
22 striata. DC.	(bl.) streaked.	1816.	6. 8. G. ♂.
23 argentea. DC.	(bl.) silvery.	—	— G. ♂.
24 tomentosa. DC.	(bl.) woolly.	1825.	—
25 capitata. DC.	(bl.) headed.	1793.	—
26 corylifolia. DC.	(vi.) Hazel-leaved.	1739.	6. 7. S. ♂.
27 virgata. DC.	(bl.) twiggy.	1826.	6. 8. H. ♀.
28 arenaria. DC.	(bl.) sand.	1823.	—
lanceolata et elliptica. Ph. ex Nutt.			Bot. mag. 665.
29 palentina. DC.	(bl.) Palestine.	1771.	4. 9. F. ♀.
30 bituminosa. DC.	(pu.) bituminous.	1570.	—
31 sericea. DC.	(pu.) long-peduncled.	1815.	1. 12. G. ♂.
pedunculata. B.R.			Jacq. vind. 2. t. 184.
32 Mutisii. DC.	(pu) Mutis's.	1828.	6. 8. G. ♂.
33 esculenta. DC.	(bl.) Bread-root.	1811.	— H. ♀.
			Kth. pl. legum. t. 54.
			Pursh. amer. 2. t. 22.

34	cuspidatæ. DC.	(bl.) large-rooted. macrorhiza. Fraser.	Louisiana.	1811.	6. 8. H. 2.	
35	incanæ. DC.	(bl.) silky.	Missouri.	1813.	— H. 2.	
36	lupinellæ. DC.	(pu.) narrow-leaved.	Carolina.	1812.	— H. 2.	
37	tenuiflora. DC.	(pu.) slender-flower'd.	Missouri.	1826.	— H. 2.	
38	longifolia. DC.	(wh.) long-leaved.	Louisiana.	1811.	— H. 2.	
39	melilotoides. DC.	(bl.) Melilot-like.	Carolina.	1814.	— F. 2.	
40	Onobrychis. DC.	(pu.) rough-podded.	Louisiana.	1812.	— H. 2.	
41	glandulosa. DC.	(v.) glandular.	Peru.	1770.	5. 9. H. h.	
42	pubescens. DC.	(bl.) pubescent.	Lima.	1823.	— G. 2.	
43	canescens. DC.	(bl.) canescent.	Carolina.	—	6. 8. F. 2.	
44	divaricata. DC.	(bl.) divaricate.	Peru.	1820.	5. 9. G. h.	
45	dentata. DC.	(wh.pu.) tooth-leaved. americana. Jacq. schœnb. t. 227.	Madeira.	1816.	— F. h.	Lobel. ic. 2. t. 31. f. 1.
	CARMICHAELIA. B.R. CARMICHAELIA.	Diadelphia Decandria.				
	australis. B.R.	(li.) New Zealand.	New Zealand.	1822.	3. 5. G. h.	Bot. reg. 912.
	Lotus arboreus. Forster.					
	INDIGOFERA. DC.	INDIGO.	Diadelphia Decandria. L.			Roxb. cor. 2. t. 194.
1	linifolia. DC.	(wh.) Flax-leaved.	E. Indies.	1792.	7. 8. S. (○).	
2	paniculata. DC.	(re.) panicled.	Sierra Leon.	1824.	— S. h.	
3	filifolia. DC.	(re.) filiform-leaved.	C. B. S.	1812.	7. 10. G. h.	
4	echinata. DC.	(re.) prickly-podded.	E. Indies.	1824.	6. 9. S. 2.	
5	cordifolia. DC.	(pu.) heart-leaved.	—	1826.	— S. h.	
6	monophylla. DC.	(pu.) simple-leaved.	N. S. W.	—	— G. h.	
7	diphylla. DC.	(ro.) two-leaved.	Africa.	1824.	7. 10. S. (○).	
8	subulata. DC.	(li.) awl-stipuled.	W. Indies.	—	— S. (○).	
9	Lechenaultii. DC.	(pu.) Lechenault's.	Bengal.	1826.	7. 10. S. (○).	
10	trifoliata. DC.	(re.) trifoliate.	E. Indies.	1818.	7. 8. S. 2.	
11	multicaulis. DC.	(re.) many-stalked.	Nepaul.	1824.	7. 10. S. (○).	
12	glandulosa. DC.	(re.) glandular.	E. Indies.	1818.	5. 7. S. 2.	
13	biflora. DC.	(pu.) two-flowered.	—	1826.	— S. h.	
14	denudata. DC.	(pu.) smooth-leaved.	C. B. S.	1790.	— G. h.	Jacq. schœnb. 2. t. 233.
15	amœna. DC.	(sc.) scarlet-flowered.	—	1774.	3. 5. G. h.	Bot. reg. 300.
16	rígida. DC.	(re.) rigid.	E. Indies.	1820.	5. 8. S. h.	
17	virgata. DC.	(re.) twiggy.	Nepaul.	1823.	7. 8. S. h.	
18	argentea. DC.	(re.) silvery.	Egypt.	1776.	— S. h.	L'Herit. strip. t. 79.
19	tinctoria. DC.	(re.) East Indian.	E. Indies.	1731.	— S. h.	Rheed. mal. 1. t. 54.
20	brachycarpa. DC.	(re.) short-podded.	W. Indies.	—	— S. h.	Lam. ill. t. 626. f. 2.
21	A'nil. DC.	(re.) West Indian.	—	1731.	— S. h.	
22	cœrulea. H.B.	(pu.) blue-flowered.	E. Indies.	1820.	6. 7. S. h.	
23	leptostachya. DC.	(pu.) slender-spiked.	—	1827.	6. 8. S. h.	
24	atropurpurea. DC.	(pu.) dark-purple.	Nepaul.	1820.	7. 8. S. h.	
25	cylindrica. DC.	(ro.) cylindric-podd'd.	C. B. S.	1822.	6. 8. G. h.	
26	juncæ. DC.	(ro.) rushy.	—	—	7. 10. G. h.	Herb. amat. t. 227.
	aphylla. L. enum. Lebeckia contaminata. B.R.	104. nuda. B.M. 2214.				
27	australis. DC.	(li.) round-stem'm'd.	N. S. W.	1790.	3. 6. G. h.	Bot. reg. 386.
28	angulata. B.R.	(li.) angular-stalked.	—	1824.	— G. h.	— t. 991.
29	macrostachya. DC.	(ro.) long-spiked.	China.	1822.	4. 6. G. h.	Vent. malm. t. 44.
30	frutescens. DC.	(ro.) frutescent.	C. B. S.	1823.	7. 9. G. h.	Jacq. schœnb. t. 365.
31	divaricata. DC.	(ru.) spreading.	—	1824.	— G. h.	
32	mucronata. DC.	(re.) mucronate.	Jamaica.	—	— S. h.	
33	elliptica. H.B.	(re.) elliptic-leaved.	Bengal.	1820.	7. 8. S. h.	
34	pulchella. H.B.	(re.) pretty.	E. Indies.	1823.	— S. h.	
35	arborea. H.B.	(re.) tree.	—	—	... S. h.	
36	purpúascens. H.B.	(p.) purplish.	—	1827.	7. 8. S. h.	
37	violacea. H.B.	(vi.) violet-coloured.	—	1825.	— S. h.	
38	uncinata. H.B.	(re.) hooked.	—	—	— S. h.	
39	circinata. Rox.	(re.) rounded.	—	1822.	— S. h.	
40	lateritia. DC.	(re.) brick-coloured.	Guinea.	—	— S. (○).	Jacq. ic. t. 359.
	hirsuta. Jacq. nom Linn.					
41	viscosa. DC.	(re.) clammy.	E. Indies.	1806.	6. 8. S. (○).	Wend. sert. han. 2. t. 12.
42	dendroides. DC.	(ro.) tree-like.	Guinea.	1822.	— S. (○).	Jacq. ic. 3. t. 571.
43	polyphylla. DC.	(re.) many-flowered.	Nepaul.	1825.	4. 8. G. 2.	
44	heterotricha. DC.	(ro.) various-haired.	C. B. S.	1816.	— G. h.	
45	hirsuta. DC.	(pu.) hairy-leaved.	E. Indies.	1739.	7. 9. S. h.	Burm. zeyl. 37. t. 14.
46	endecaphylla. DC.	(re.) eleven-leaved.	Guinea.	1822.	— S. (○).	Bot. reg. 789.
47	alternans. DC.	(re.) alternate-leav'd.	C. B. S.	1816.	— G. h.	
48	fragrans. DC.	(re.) fragrant.	E. Indies.	1822.	— S. 2.	

LEGUMINOSÆ.

49 pusilla. DC.	(re.) small.	Madagascar.	1822.	7. 9.	S. h.	
50 glabra. DC.	(re.) smooth.	E. Indies.	1824.	—	S. ⓧ.	Pluk. alm. t. 166. f. 1.
51 emeaphylla. DC.	(pu.) trailing.		1776.	—	S. ⓧ.	— t. 166. f. 2.
52 cytisoides. DC.	(ro.) angular-stem'd.	C. B. S.	1774.	—	G. h.	Bot. mag. 742.
53 lotoides. DC.	(ro.) Lotus-like.		1816.	—	G. h.	Comm. hort. 2. t. 84?
54 angustifolia. DC.	(pu.) narrow-leaved.		1774.	6. 10.	G. h.	Bot. mag. 465.
55 aspalathoides. DC.	(re.) Aspalathus-like.	Ceylon.	1823.	4. 6.	S. h.	
56 filiformis. DC.	(ro.) filiform.	C. B. S.	1816.	—	G. h.	
57 alopecuroides. DC.	(ro.) Fox-tail.		1825.	—	G. h.	
58 coriacea. DC.	(ro.) leathery-leaved.		1774.	7. 8.	G. h.	
59 sarmentosa. DC.	(ro.) dwarf.		1786.	6. 8.	G. h.	
60 microphylla. P. S.	(ro.) small-leaved.		1812.	1. 12.	G. h.	
61 digitata. DC.	(ro.) finger-leaved.		1820.	4. 8.	G. h.	
62 sessilifolia. DC.	(ro.) sessile-leaved.		1816.	—	G. h.	
63 spinosa. DC.	(re.) spiny.	Arabia.	1822.	—	S. h.	
64 candicans. DC.	(ro.) white-leaved.	C. B. S.	1774.	5. 9.	G. h.	Bot. mag. 198.
65 psoraloides. DC.	(ro.) Psoralea-like.		1758.	7. 8.	G. h.	— t. 476.
66 stipularis. DC.	(ro.) stipuled.		1816.	6. 7.	G. h.	
67 incana. DC.	(ro.) hoary.		1812.	5. 7.	G. h.	Bot. reg. 957.
68 trita. DC.	(re.) oval-leaved.	E. Indies.	1802.	6. 8.	S. ⓧ.	
69 procumbens. DC.	(re.) procumbent.	C. B. S.	1816.	5. 7.	G. h.	
70 prostrata. DC.	(re.) prostrate.	E. Indies.	1828.	8. 10.	S. ⓧ.	
CLIT'ORIA. DC.	CLIT'ORIA.	Diadelphia Decandria.	L.			
1 lascivia. Boj.	Madagascar.	1826.	—	S. h.	
2 heterophylla. DC.	(bl.) various-leaved.	E. Indies.	1812.	7. 8.	S. ⓧ.	Bot. mag. 2111.
3 Ternatea. DC.	(bl. wh.) wing-leaved.		1739.	5. 9.	S. h.	Rheed. mal. 8. t. 38.
α cærulaea.	(bl.) blue-flowered.		—	—	S. h.	Bot. mag. 1542.
β alba.	(wh.) white-flowered.		—	—	S. h.	
4 marianna. DC.	(bl.) Maryland.	N. America.	1759.	8. 9.	H. 4.	
5 mexicana. DC.	(pu.) Mexican.	Mexico.	1823.	6. 8.	G. 4.	
6 virginiana. DC.	(pu.) Virginian.	America.	1732.	7. 8.	G. 4.	Dill. elth. 1. f. 76.
calcarigera. Sal. par. lond. t. 51.						
7 occidentalis.	(pu.) West Indian.	Trinidad.	1824.	6. 8.	S. h.	Bot. reg. 1047.
virginiana. B. R. 1047. non Linn.						
8 brasiliæna. DC.	(bl.) Brazilian.	Brazil.	1759.	—	S. 4.	Breyn. cent. t. 32.
9 Plumiéri. DC.	(pu.) Plumier's.	W. Indies.	1815.	8. 11.	S. h.	Bot. reg. 268.
10 erecta. H. B. erect.	S. America.	1822.	—	S. h.	
11 Berteriana. DC.	(ye.) Bertero's.	St. Domingo.	1824.	6. 8.	S. 4.	
12 arborescens. DC. tree.	Trinidad.	1804.	—	S. h.	
NEUROCA'RPU M. DC.	NEUROCA'RPU M.	Diadelphia Decandria.				
1 guianense. DC.	(pu.) Guiana.	Guiana.	1826.	6. 8.	S. 4.	Aubl. guian. 2. t. 305.
2 falcatum. DC.	(pu.) sickle-podded.	St. Domingo.	—	—	S. h.	Plum. spec. 8. t. 85.
COLOG'ANIA. DC.	COLOG'ANIA.	Diadelphia Decandria.				
1 intermédia. DC.	(fl.) intermediate.	Mexico.	1824.	6. 8.	G. 4.	
2 angustifolia. DC.	(vi.) narrow-leaved.		—	—	G. 4.	Kth. mim. 209. t. 58.
3 Broussonnetii. DC.	(vi.) Broussonnet's.		1820.	—	S. 4.	
GALA'CTIA. DC.	GALA'CTIA.	Diadelphia Decandria.				
1 pendula. DC.	(ro.) pendulous.	Jamaica.	1794.	7. 8.	S. h.	Böt. reg. 269.
2 sericea. DC.	(pu.) silky.	Isl. Bourbon.	1824.	—	S. h.	
3 cubensis. DC.	(ro.) Cuba.	Cuba.	1826.	—	S. 4.	
4 mollis. DC.	(pu.) soft.	Carolina.	—	—	G. 4.	Dill. elth. 1. f. 170.
5 glabælla. DC.	(re.) smooth.		1826.	6. 8.	G. 4.	
VILMORINIA. DC.	VILMORINIA.	Diadelphia Decandria.				
multiflora. DC.	(pu.) many-flowered.	W. Indies.	1823.	7. 8.	S. h.	
Clitòria multiflora. Swartz.						
COLL'EA. DC.	COLL'EA.	Monadelphia Pent-Decandria.				
1 speciosa. DC.	(pu.) pentandrous.	Peru.	1824.	6. 8.	G. h.	DC. leg. me. p. 245. t. 40.
2 trinervia. DC.	(pu.) decandrous.	E. Indies.	1826.	—	S. h.	— p. 247. t. 41.
OTO'PTERA. DC.	OTO'PTERA.	Monadelphia Decandria.				
Burchellii. DC.	(pu.) Burchell's.	C. B. S.	1816.	5. 7.	G. h.	DC. leg. mem. 250. t. 42.
PUER'ARIA. DC.	PUER'ARIA.	Monadelphia Decandria.				
1 tuberosa. DC.	(ye.) tuberous-root'd.	E. Indies.	1806.	6. 8.	S. h.	
Hedysarum tuberosum. W.						
2 Wallichii. DC.	(ye.) Wallich's.	Nepaul.	1826.	4. 6.	G. h.	DC. leg. mem. 254. t. 43.
DUM'ASIA. DC.	DUM'ASIA.	Diadelphia Decandria.				
1 villosa. DC.	(ye.) villous.	Nepaul.	1824.	8. 12.	G. h.	DC. leg. mem. 257. t. 44.
2 pubescens. DC.	(ye.) pubescent.		—	—	G. h.	Bot. reg. 962.
GLYCINE. DC.	GLYCINE.	Diadelphia Decandria.	L.			
1 clandestina. DC.	(bl.) hidden-flower'd.	N. S. W.	1810.	4. 8.	G. h.	

2 dèbilis. DC.	(ye.) hairy.	E. Indies.	1778.	6. 7.	S. ȝ. . .
3 mímina. DC.	(bl.) small.	N. S. W.	1812.	4. 10.	G. ȝ. . .
4 striàta. DC.	(li.) streaked.	S. America.	1823.	7. 9.	S. ȝ. . . Jacq. vind. 1. t. 76.
5 parviflòra. DC.	(re.) small-flowered.	E. Indies.	1812.	7. 8.	S. ȝ. . .
6 senegalénis. DC.	(ye.) Senegal.	Senegal.	1822.	—	S. ȝ. . .
7 heterophylla. DC.	(ye.) various-leaved.	C. B. S.	1816.	6. 8.	G. ȝ. . .
8 argéntea. DC.	(ye.) silvery-leaved.	—	—	—	G. ȝ. . .
9 secúnda. DC.	(ye.) side-flowering.	—	1825.	6. 9.	G. ȝ. . .

CHÆTOCALYX. DC. CHÆTOCALYX. Diadelphia Decandria.

vincentína. DC. (ye.) St. Vincent. W. Indies. 1823. 6. 8. S. ȝ. . . Bot. reg. 799.

Glycine vincentina. B.R. 799.

Subtribus IV. GALEGEÆ. DC. prodr. 2. p. 243.

PETALOST'EMUM. DC. PETALOST'EMUM. Monadelphia Pentandria.

1 cándidum. DC.	(wh.) white.	N. America.	1811.	7. 8.	H. ȝ. Mich. amer. 2. t. 37. f. 1.
2 cárneum. DC.	(li.) flesh-coloured.	Florida.	—	—	H. ȝ.
3 violáceum. DC.	(pu) purple-flower'd.	Missouri.	—	7. 9.	H. ȝ. Bot. mag. 1707.
4 villósum. DC.	(ro.) villous.	—	1826.	—	H. ȝ.
5 corymbósium. DC.	(wh.) corymbose.	Carolina.	1806.	—	H. ȝ.

Dâlea Kuhnistèra. w. Kuhnistéra carolinensis. Lam. Cylopôgon virgatum. Rafin.

D'ALEA. DC. D'ALEA. Monadelphia Decandria.

1 laxiflòra. DC.	(wh.) loose-flowered.	Louisiana.	1811.	7. 8.	H. ȝ.
2 áurea. DC.	(ye.) golden.	—	—	—	H. ȝ.
3 alopecuroïdes. DC.	(w.) Fox-tail.	—	1812.	—	H. ⓧ. Mich. amer. 2. t. 38.
Dâlea Linnæi. Mich.	—	—	—	—	—

4 Cliftiána. w.	(bl.) Vera Cruz.	Vera Cruz.	1737.	7. 8.	H. ⓧ. Linn. h. cliff. t. 22.
5 Lagòpus. DC.	(vi.) downy-spiked.	Mexico.	1780.	10. 11.	G. ⓧ. Cavan. ic. 1. t. 86.
6 sericea. DC.	(d.vi.) silky.	—	1825.	9. 11.	G. ȝ. Kth. mim. 166. t. 48.
gracilis. Kth.	—	—	—	—	—

7 tuberculáta. DC.	(pu.) tubercled.	—	—	—	G. ȝ.
8 citriodòra. DC.	(v.) leafy.	New Spain.	1780.	10. 11.	G. ⓧ. Cavan. ic. 3. t. 271.

Psoralea citriodora. Cav. foliolosa. H.K.

9 lutea. DC.	(ye.) yellow.	Mexico.	1823.	7. 9.	G. ȝ. Cavan. ic. 4. t. 325.
10 mutabilis. DC.	(v.) changeable.	—	1821.	9. 10.	G. ȝ. Bot. mag. 2486.

bicolor. Hook. ex. flor. 43.

11 nútans. DC.	(vi.) nodding.	—	1824.	7. 9.	G. ȝ. Cavan. ic. 3. t. 201.
12 reclinàta. DC.	(vi.) reclining.	—	—	5. 8.	G. ȝ. Cavan. ic. 3. t. 240.

13 tomentosa. DC.	(vi.) woolly.	—	1823.	7. 9.	G. ȝ. Cavan. ic. 3. t. 240.
14 prostràta. DC.	(ye.) prostrate.	—	1826.	—	G. ȝ. —

15 mucronàta. DC.	(wh.) mucronate.	—	—	—	G. ȝ. —
16 Thouíni. DC.	(vi.) Thouin's.	—	1824.	6. 9.	H. ⓧ. Schrank hort. mon. t. 9.

17 phymatòdes. DC.	(vi.) obovate-leaved.	Caracas.	—	7. 9.	G. ȝ. Jacq. ic. 3. t. 563.
18 enneaphylla. DC.	(re.) nine-leaved.	W. Indies.	1772.	—	S. ȝ. —

GLYCYRHIZA. DC. LIQUORICE. Diadelphia Decandria. L.

1 glábra. DC.	(pu.) common.	S. Europe.	1562.	7. 9.	H. ȝ. Lam. ill. t. 625. f. 2.
2 glandulífera. DC.	(pu.) glandular.	Hungary.	1805.	6. 8.	H. ȝ. W. et K. hung. 1. t. 21.

3 lepidóta. DC.	(wh.) scaly.	Missouri.	1811.	7. 8.	H. ȝ. Bot. mag. 2150.
4 foetida. DC.	(ye.) strong-scented.	Barbary.	1823.	6. 9.	H. ȝ. Desf. atl. 2. t. 199.

5 echináta. DC.	(pu.) prickly-headed.	Italy.	1596.	—	H. ȝ. Bot. mag. 2154.
6 aspérírrima. DC.	(li.) rough.	Siberia.	1795.	7. 8.	H. ȝ. Pall. it. app. t. M. f. 3.

7 uralénis. DC.	(li.) Siberian.	—	1826.	—	H. ȝ. —
8 hirsùta. DC.	(li.) hairy.	Levant.	1739.	—	H. ȝ. —

GAL'EGA. DC. GOAT'S-RUE. Monadelphia Decandria.

1 officinalis. DC.	(pu.) officinal.	Spain.	1568.	6. 9.	H. ȝ. Schk. hand. 2. t. 208.
α purpùrea.	(pu.) purple-flowered.	—	—	—	H. ȝ. —

β álba.	(wh.) white-flowered.	—	—	—	H. ȝ. —
2 pérsica. DC.	(wh.) Persian.	Persia.	1816.	—	H. ȝ. Swt. br. fl. gar. 3. t. 244.

3 bilòba. B.F.G.	(pu.) two-lobed.	1823.	—	H. ȝ. 2. t. 159.
4 orientális. DC.	(bl.) oriental.	Levant.	1801.	—	H. ȝ. Bot. reg. 326.

TEPHR'OSIA. DC.	TEPHR'OSIA.	Mono-Diadelphia Decandria.	—	—	—
1 suberòsa. DC.	(ro.) cork-barked.	E. Indies.	1818.	5. 8.	S. ȝ. —

Robinia suberòsa. H.B.	—	—	—	—	—
2 fruticòsa. DC.	(ro.) frutescent.	—	1820.	S. ȝ. —

LEGUMINOSÆ.

3 <i>sophoroïdes</i> . DC. (<i>wh.</i>)	Sophora-like.	Nepaul.	1826.	G. $\frac{1}{2}$.	
4 <i>brachystachya</i> . DC. (<i>ro.</i>)	short-spiked.			G. $\frac{1}{2}$.	
5 <i>candida</i> . DC. (<i>ro.</i>)	white-leaved.	E. Indies.	1810.	5. 8.	S. $\frac{1}{2}$.	
<i>Robinia candida</i> . H.B.						
6 <i>toxicaria</i> . DC. (<i>bh.</i>)	Fish poison.	S. America.	1791.	—	S. $\frac{1}{2}$.	Plum. ic. t. 135.
7 <i>virginiæna</i> . DC. (<i>re.</i>)	Virginian.	Virginia.	1765.	6. 8.	H. $\frac{1}{2}$.	Pluk. alm. t. 23. f. 2.
8 <i>hispidula</i> . DC. (<i>li.</i>)	bristle-podded.		1826.	—	H. $\frac{1}{2}$.	
9 <i>ochroleuca</i> . DC. (<i>st.</i>)	straw-colour'd.	W. Indies.	1799.	—	S. $\frac{1}{2}$.	Jacq. ic. t. 150.
10 <i>caribæa</i> . DC. (<i>re. wh.</i>)	various-colour'd.	Caribees.	1786.	—	S. $\frac{1}{2}$.	Jacq. amer. t. 125.
11 <i>grandiflora</i> . DC. (<i>re.</i>)	large-flowered.	C. B. S.	1774.	6. 9.	I. $\frac{1}{2}$.	Bot. reg. 769.
<i>Galèga grandiflora</i> . B.R. <i>Galèga ròsea</i> . Lam.						
12 <i>villosa</i> . DC. (<i>pu.</i>)	villous.	E. Indies.	1779.	6. 8.	S. $\frac{1}{2}$.	Burm. zeyl. t. 33.
13 <i>purpurea</i> . DC. (<i>pu.</i>)	purple.		1768.	7. 8.	S. $\frac{1}{2}$.	— t. 32.
<i>Galèga purpurea</i> . L.						
14 <i>lanceæfolia</i> . DC. (<i>st.</i>)	lance-leaved.		1824.	—	S. $\frac{1}{2}$.	
15 <i>argentea</i> . DC. (<i>pu.</i>)	silver.		1826.	S. $\frac{1}{2}$.	Pluk. alm. t. 52. f. 1.
16 <i>piscatória</i> . DC. (<i>pu.</i>)	woolly.		1778.	6. 8.	S. $\frac{1}{2}$.	
17 <i>leucantha</i> . DC. (<i>wh.</i>)	white-flowered.	Mexico.	1826.	6. 10.	G. $\frac{1}{2}$.	H. et B.n.gen.a.6.t.577.
18 <i>coronillæfolia</i> . DC. (<i>st.</i>)	Coronilla-leav'd.	Bourbon.	—	—	S. $\frac{1}{2}$.	
19 <i>capensis</i> . DC. (<i>re.</i>)	Cape.	C. B. S.	1823.	5. 9.	G. $\frac{1}{2}$.	Jacq. ic. t. 574.
20 <i>cineræa</i> . DC. (<i>vi.</i>)	cinerous.	W. Indies.	1824.	6. 9.	S. \odot .	— t. 575.
21 <i>hypargy'rea</i> . DC. (<i>pu.</i>)	various-leaved.	E. Indies.	1825.	—	S. $\frac{1}{2}$.	
<i>heterophylla</i> . F.						
22 <i>stricta</i> . DC. (<i>pu.</i>)	straight-podded.	C. B. S.	1774.	5. 6.	I. $\frac{1}{2}$.	Jacq. schœnb. t. 236.
23 <i>linearis</i> . DC. (<i>pu.</i>)	linear-leaved.	Guinea.	1822.	S. $\frac{1}{2}$.	
24 <i>pallens</i> . DC. (<i>li.</i>)	pale-flowered.	C. B. S.	1787.	6. 8.	I. $\frac{1}{2}$.	
25 <i>apollinea</i> . DC. (<i>bl.</i>)	Egyptian.	Egypt.	1822.	6. 7.	F. $\frac{1}{2}$.	Delil. fl. eg. t. 53. f. 5.
26 <i>tricolor</i> . DC. (<i>bl.</i>)	three-coloured.	N. Holland.	1824.	7. 9.	G. $\frac{1}{2}$.	Hook. ex. flor. t. 185.
<i>Galèga tricolor</i> . H.E.F.						
27 <i>Barclayana</i> . (<i>pu.</i>)	Mr. Barclay's.	Madagascar.	1823.	5. 10.	S. $\frac{1}{2}$.	Hook. ex. flor. t. 188.
<i>Dalbergia Barclayii</i> . H.E.F.						
28 <i>Telfairii</i> . (<i>pu.</i>)	Mr. Telfair's.			—	S. $\frac{1}{2}$.	
29 <i>Colonila</i> . L.T.	(<i>pu.</i>) Colonil.	E. Indies.	1818.	6. 8.	S. $\frac{1}{2}$.	
30 <i>racemosa</i> . (<i>pu.</i>)	racemed.		1822.	—	S. $\frac{1}{2}$.	
<i>Robinia racemosa</i> . H.B.						
31 <i>pentaphylla</i> . (<i>pu.</i>)	five-leaved.		1824.	—	S. $\frac{1}{2}$.	
<i>Galèga pentaphylla</i> . H.B.						
32 <i>biflora</i> . DC. (<i>pu.</i>)	two-flowered.	1826.	6. 9.	G. $\frac{1}{2}$.	
33 <i>capitulata</i> . DC. (<i>pu.</i>)	headed.	Owyhee.	—	—	S. $\frac{1}{2}$.	
34 <i>incana</i> . (<i>pu.</i>)	hoary.	E. Indies.	1820.	—	S. $\frac{1}{2}$.	
<i>Galèga incana</i> . H.B.						
35 <i>Heyniâna</i> . (<i>bl.</i>)	Heyne's.		1807.	7. 9.	S. $\frac{1}{2}$.	
<i>Galèga Heyniâna</i> . H.B.						
36 <i>sericea</i> . DC. (<i>pu.</i>)	silky.	C. B. S.	1824.	—	I. $\frac{1}{2}$.	
37 <i>filifolia</i> . DC. (<i>re.</i>)	filiform-leaved.			—	I. $\frac{1}{2}$.	
AMORPHA. DC.	BASTARD INDIGO.	Monadelphia Decandria.				
1 <i>fruticosa</i> . DC. (<i>bl.</i>)	shrubby.	Carolina.	1724.	6. 8.	H. $\frac{1}{2}$.	Bot. reg. 427.
2 <i>croceolanata</i> . W.D. (<i>pu.</i>)	yellow-haired.		1812.	—	H. $\frac{1}{2}$.	Wats. dend. brit. t. 139.
3 <i>frâgrans</i> . B.F.G. (<i>pu.</i>)	fragrant.			—	H. $\frac{1}{2}$.	Swt. br. fl. gar. t. 241.
<i>nâna</i> . Bot. mag. t. 2112. nec aliorum.						
4 <i>emarginata</i> . L.C. (<i>bl.</i>)	emarginate.			—	H. $\frac{1}{2}$.	
<i>fruticosa</i> β <i>emarginata</i> . Ph.						
5 <i>Lewisii</i> . L.C. (<i>bl.</i>)	Lewis's.			—	H. $\frac{1}{2}$.	
<i>fruticosa</i> γ <i>angustifolia</i> . Ph.						
6 <i>glabra</i> . DC. (<i>pu.</i>)	smooth.			—	H. $\frac{1}{2}$.	
7 <i>nâna</i> . DC. (<i>pu.</i>)	dwarf.	Missouri.	1811.	7. 8.	H. $\frac{1}{2}$.	
<i>microphylla</i> . Ph.						
8 <i>herbacea</i> . DC. (<i>pu.</i>)	pubescent.	Carolina.	1803.	6. 8.	H. $\frac{1}{2}$.	Lodd. bot. cab. 689
<i>pùbescens</i> . Ph.						
9 <i>canescens</i> . DC. (<i>bl.</i>)	canescent.	Missouri.	1812.	7. 8.	H. $\frac{1}{2}$.	
NISSOLIA. DC.	NISSOLIA.	Mono-Diadelphia Decandria.				
1 <i>fruticosa</i> . DC. (<i>st.</i>)	shrubby.	S. America.	1766.	7. 11.	S. $\frac{1}{2}$.	Jacq. vind. t. 167.
2 <i>racemosa</i> . DC. (<i>st.</i>)	racemed.	W. Indies.	6. 8.	S. $\frac{1}{2}$.	
3 <i>glabrata</i> . DC. (<i>wh.</i>)	smooth-leaved.	S. America.	1818.	7. 11.	S. $\frac{1}{2}$.	
4 <i>robiniaeæfolia</i> . DC.	Robinia-leaved.	St. Vincent.	1824.	—	S. $\frac{1}{2}$.	
5 <i>micróptera</i> . DC. (<i>ye.</i>)	small-winged.			5. 8.	S. $\frac{1}{2}$.	
6 <i>retusa</i> . DC. (<i>ye.</i>)	retuse-leaved.	S. America.	1823.	S. $\frac{1}{2}$.	
MULLERA. DC.	MULLERA.	Monadelphia Octo-Decandria.				
moniliformis. DC.	bracelet.	Guiana.	1792.	S. $\frac{1}{2}$.	Aubl. guian. 4. t. 356.

LONCHOCA'RPUS. DC. <i>LONCHOCA'RPUS.</i> Mono-Diadelpbia Decandria.						
1 <i>dominénsis.</i> DC. (<i>re.</i>) oval-leaved.	St. Domingo.	1759.	S. \natural .		
2 <i>violaceus.</i> DC. (<i>vi.</i>) violet.	W. Indies.	—	5. 8.	S. \natural .	Jacq. amer. t. 177. f.49.	
<i>Robinia violacea.</i> Jacq.						
3 <i>formosianus.</i> DC. (<i>vi.</i>) Formosa.	Africa.	1822.	S. \natural .	Beauv. fl. owar. 2 t. 76.	
4 <i>sericeus.</i> DC. (<i>pu.</i>) silky.	W. Indies.	1823.	S. \natural .		
5 <i>pyxidarius.</i> DC. (<i>pu.</i>) Cuba.	Cuba.	1822.	S. \natural .		
6 <i>latifolius.</i> DC. (<i>pu.</i>) broad-leaved.	S. America.	1814.	S. \natural .		
<i>Amerínum latifolium.</i> W.						
7 <i>pubescens.</i> DC. (<i>pu.</i>) pubescent.	Caracas.	1820.	S. \natural .		
8 <i>róseus.</i> DC. (<i>ro.</i>) rose-coloured.	S. America.	S. \natural .		
9 <i>sepium.</i> DC. (<i>ro.</i>) hedge.	W. Indies.	1820.	S. \natural .	Jacq. amer. t. 129. f.101.	
10 <i>Swartzii.</i> DC. (<i>ro.</i>) Swartz's.	—	—	S. \natural .		
<i>Robinia sépium.</i> Sw.						
11 <i>Nicou.</i> DC. (<i>pu.</i>) climbing.	Guiana.	1826.	S. \natural .	Aubl. guian. 2. t. 308.	
<i>Robinia Nicou.</i> Aubl. scándens. w.						
ROBI'NIA. DC.	ROBI'NIA. Diadelpbia Decandria. L.					
1 <i>Pseudacacia.</i> DC. (<i>wh.</i>) com.or Locust Tree. N. Amer.	1640.	5. 6.	H. 5.		Schmidt arb. 1. t. 32.	
2 <i>inérmis.</i> DC. (<i>wh.</i>) spineless.	—	H. 5.		
<i>spectabilis.</i> L.C.						
3 <i>macrophylla.</i> L.C. (<i>ro.</i>) long-leaved.	Siberia.	1797.	6. 7.	H. 5.		
4 <i>sophoræfolia.</i> L.C. (<i>wh.</i>) Sophora-leaved.	Hybrid.	1811.	—	H. 5.		
5 <i>amorphæfolia.</i> L.en. (<i>w.</i>) Amorpha-leav'd.	—	1820.	—	H. 5.		
6 <i>umbraculifera.</i> DC. (<i>w.</i>) Parasol.	—	1811.	H. 5.		
<i>Robinia inérmis.</i> Dum. Cours. non DC.						
7 <i>dúbia.</i> DC. (<i>bh.</i>) doubtful.	—	—	H. 5.		
8 <i>strícta.</i> L.C. (<i>wh.</i>) upright.	—	H. 5.		
9 <i>tortuosa.</i> DC. (<i>wh.</i>) twisting.	—	6. 7.	H. 5.		
10 <i>monstrósa.</i> L.C. (<i>wh.</i>) monstrous.	—	—	H. 5.		
11 <i>péndula.</i> L.en. (<i>wh.</i>) pendulous.	—	—	H. 5.		
12 <i>viscosa.</i> DC. (<i>bh.</i>) clammy.	N. America.	1797.	6. 8.	H. 5.	Venten. cels. t. 4.	
<i>glutinosa.</i> Bot. mag. 580.						
13 <i>hispida.</i> DC. (<i>ro.</i>) Rose Acacia.	Carolina.	1743.	5. 9.	H. \natural .	Bot. mag. 311.	
14 <i>rósea.</i> Ell. (<i>ro.</i>) smooth-branch'd.	—	1812.	—	H. \natural .		
15 ? <i>guineensis.</i> W.en. (<i>ro.</i>) Guinea.	Guinea.	1821.	S. \natural .		
<i>Cy'tisus hispidus.</i> w.						
16? <i>purpurea.</i> L.en. (<i>pu.</i>) purple.	—	1816.	4. 6.	S. \natural .		
POIT'EA. DC.	POIT'EA. Diadelpbia Decandria.					
galegóides. DC. (<i>ro.</i>) Galega-like.	St. Domingo.	1826.	S. \natural .	Venten. choix. t. 36.	
COURS'ETIA. DC.	COURS'ETIA. Diadelpbia Decandria.					
1 <i>tomentosa.</i> DC. (<i>ye.</i>) woolly.	Peru.	1824.	6. 8.	G. \natural .	Cavan. ic. t. 84.	
<i>Láthyrus fruticòsus.</i> Cav. <i>O'robos tomentòsus.</i> Df. <i>Vicia fruticòsa.</i> w.						
2 <i>virgata.</i> DC. (<i>ye.</i>) slender.	Trinidad.	1820.	—	S. \natural .	Cavan. ic. t. 293.	
<i>Æschynómene virgata.</i> Cav.						
SESBA'NIA. DC.	SESBA'NIA. Diadelpbia Decandria.					
1 <i>ægyptiaca.</i> DC. (<i>ye.</i>) Egyptian.	Egypt.	1680.	7. 8.	S. σ .	Alp. ægypt. 81. t. 82.	
2 <i>occidentalis.</i> DC. (<i>ye.</i>) West Indian.	W. Indies.	1823.	6. 8.	S. \natural .	Plum. ed Bur. t. 125. f.1.	
3 <i>aculeata.</i> DC. (<i>ye.</i>) prickly.	E. Indies.	1690.	7. 10.	S. \odot .	Jacq. ic. t. 564.	
4 <i>cannabina.</i> DC. (<i>ye.</i>) Hemp-like.	—	1800.	—	S. \odot .		
5 <i>affinis.</i> DC. (<i>ye.</i>) likened.	—	1822.	5. 8.	S. σ .		
6 <i>picta.</i> DC. (<i>std.</i>) painted.	W. Indies.	—	4. 9.	S. \natural .	Bot. reg. 873.	
7 <i>macrocárpa.</i> DC. (<i>st.</i>) long-podded.	Louisiana.	—	7. 10.	H. \odot .	Bart. fl. amer. 1. t. 28.	
8 <i>paludosa.</i> DC. (<i>ye.</i>) marsh.	E. Indies.	1810.	—	S. \odot .		
9 <i>gracilis.</i> DC. (<i>ye.</i>) slender.	—	1824.	—	S. \odot .		
10 <i>sericea.</i> DC. (<i>ye.</i>) silky.	E. Indies.	1818.	—	S. \odot .		
11 <i>procumbens.</i> H.B. (<i>ye.</i>) procumbent.	—	—	—	S. \odot .		
12 <i>uliginosa.</i> (<i>ye.</i>) swamp.	—	—	—	S. \odot .		
<i>Æschynómene uliginosa.</i> H.B.						
AGA'TI. DC.	AGA'TI. Diadelpbia Decandria.					
1 <i>grandiflora.</i> DC. (<i>bh.</i>) great-flowered.	E. Indies.	1768.	7. 8.	S. \natural .	Rheed. mal. 1. t. 51.	
<i>Sesbánia grandiflora.</i> P.S. <i>Æschynómene grandiflora.</i> L. <i>Coronilla grandiflora.</i> w.						
2 <i>coccinea.</i> DC. (<i>sc.</i>) red-flowered.	E. Indies.	1824.	7. 8.	S. \natural .	Rumph. amb. 1. t. 77.	
GLOTTIDIUM. DC.	GLOTTIDIUM. Diadelpbia Decandria.					
floridànum. DC. (<i>ye.</i>) flat-podded.	Florida.	1822.	7. 8.	S. \odot .	Jacq. ic. 1. t. 148.	
<i>Sesbánia platycárpa.</i> P.S. <i>Robínia vesicária.</i> Jacq. <i>Pháca floridâna.</i> w.						
PISCIDIA. DC.	JAMAICA DOGWOOD. Monadelphbia Decandria.					
1 <i>Erythrìna.</i> DC. (<i>v.</i>) oval-leaved.	W. Indies.	1690.	S. \natural .	Sloan. jam. 2. t. 176. f.4-5.	
2 <i>carthagénensis.</i> DC. (<i>v.</i>) obovate-leaved.	—	—	S. \natural .	Plum. ed Bur. t. 133. f.2.	

DAUBENT'ONIA. DC.	DAUBENT'ONIA. Diadelphia Decandria.						
1 punicea. DC. (pu.) red-flowered.	Mexico. 1824.	S. h.	Cavan. ic. 4. t. 316.				
2 longifolia. DC. (ye.) long-leaved.	New Spain. 1828.	S. h.	4. t. 315.				
CORYNE'LLA. DC.	CORYNE'LLA. Diadelphia Decandria.						
polyántha. DC. (pu.) many-flowered.	W. Indies 1824.	S. h.					
<i>Robinia polyántha.</i> Swartz.							
CARAG'ANA. DC.	CARAG'ANA. Diadelphia Decandria.						
1 Altagána. DC. (ye.) flat-podded.	Dahuria. 1789. 4. 6.	H. h.	L'Herit. stirp. t. 76.				
2 microphylla. DC. (ye.) small-leaved.	Siberia. 1816. —	H. h.	Pall. ross. t. 42. f. 1-2.				
3 arenaria. L.C. (ye.) sand.	—	1802. 6. 7.	H. h.	Bot. mag. 1886.			
<i>Robinia Caragana. β. arenaria.</i> B.M.							
4 arboréscens. DC. (ye.) common.	—	1752. 4. 6.	H. h.	Schmidt arb. I. t. 33.			
<i>Robinia Caragana.</i> L.							
5 Chamágua. DC. (ye.) shining.	China. 1773. 5. 6.	H. h.	L'Herit. stirp. t. 77.				
6 frutéscens. DC. (ye.) shrubby.	Siberia. 1752. 4. 6.	H. h.	Swt. br. fl. gar. 3.t.227.				
7 móllis. M.B. (ye.) soft-leaved.	Tauria. 1822. —	H. h.					
8 grandiflora. DC. (ye.) large-flowered.	Iberia. —	—	H. h.				
9 pygmæa. DC. (ye.) pigmy.	Siberia. 1751. —	H. h.	Bot. reg. 1021.				
10 mongólica. F. (ye.) Mongolian.	Tartary. 1826. —	H. h.	DC.leg.mem.t. 11. f. 45.				
11 Redówskii. F. (ye.) Redowski's.	Siberia. 1822. —	H. h.	Pall. ross. I. t. 44.				
12 spinosa. DC. (ye.) thorny.	—	1775. —	H. h.	Pall. astr. t. 86.			
13 tragacanthoídes. DC. (ye.) Goat's-thorn-like.	—	1816. —	H. h.	— p. 113. t. 85.			
14 jubata. DC. (bh.) bearded.	—	1796. —	H. h.				
HALIMODE'NDRON. DC.	SALT-TREE. Diadelphia Decandria.						
1 argéteum. DC. (pu.) silvery-leaved.	Iberia. 1779. 5. 6.	H. h.	Bot. mag. 1016.				
2 triflorum. (pu.) smooth-leaved.	—	H. h.	L'Herit. stirp. n. 162.			
<i>Robinia triflora.</i> L'Herit.							
CALOPHA'CA. DC.	CALOPHA'CA. Diadelphia Decandria.						
wolgárica. DC. (ye.) wing-leaved.	Siberia. 1786. 5. 6.	H. h.	Wats. dend. brit. 83.				
<i>Cytisus wolgáricus.</i> L. pinnátus. Pall. ross. I. t. 47.							
COLUTE'A. DC.	BLADDER SENNA. Diadelphia Decandria.						
1 arboréscens. DC. (ye.) common.	S. Europe. 1568. 6. 8.	H. h.	Bot. mag. 81.				
2 cruenta. DC. (re.) red-flowered.	Levant. 1710. —	H. h.	Schmidt arb. t. 119.				
3 Haléppica. DC. (ye.) Pocock's.	—	1752. 5. 10.	H. h.	— t. 120.			
<i>Pococcii. H.K. Istria.</i> Mill. ic. t. 100.							
4média. DC. (ye.) smaller.	6. 8. H. h.				
5 nepalénis. DC. (ye.) Nepaul.	Nepaul. 1822. 8. 9.	H. h.	Wats. dend. brit. t. 140.				
SPHÆROPH'YSA. DC.	SPHÆROPH'YSA. Diadelphia Decandria.						
1 Salsula. DC. (re.) salt-field.	Dahuria. 1829.	H. h.	Bot. mag. 2622.				
2 cásica. DC. (pu.) Caspian.	Caucasus. —	H. h.	Pall. itin. 4. t. 9. f. 1.			
<i>Colutea cásica.</i> M.B. <i>Phaca Sálssula.</i> Bieb. casp. 210.							
SWAINS'ONA. DC.	SWAINS'ONA. Diadelphia Decandria.						
1 galegífolia. DC. (sc.) red-flowered.	N. S. W. 1800. 7. 8.	G. h.	Bot. mag. 792.				
2 coronillæfolia. DC. (pu.) purple-flowered.	—	1802. —	G. h.	— t. 1725.			
3 lessertiæfolia. DC. (pu.) small-flowered.	N. Holland. 1822. —	G. h.					
4 astragalifolia.c.c.(wh.) white-flowered.	N. S. W. 1802. —	G. h.					
<i>galegífolia.</i> var. <i>albiflora.</i> B.R.							
LESSE'R'TIA. DC.	LESSE'R'TIA. Diadelphia Decandria.						
1 ánnua. DC. (pu.) annual.	C. B. S. 1731. 6. 7.	G. ○.	Hook. exot. flor. t. 84.				
2 diffusa. DC. (pu.) procumbent.	—	1792. 7. 8.	G. ○.	Jacq. ic. 3. t. 576.			
<i>Galéga dóbia.</i> Jacq.							
3 fruticòsa. B.R. (pu.) frutescent.	—	1826. —	G. ○.				
4 perénnans. DC. (ro.) perennial.	—	1753. —	G. ¼.	Bot. reg. 970.			
<i>Colutèa perénnans.</i> Jacq. <i>fstulosa.</i> Hort.				Jacq. vind. 3. t. 3.			
5 púlchra. DC. (ro.) pretty.	—	1817. 5. 8.	G. ○.				
6 falciformis. DC. (pu.) sickle-podded.	—	1825. —	G. ½.	Bot. mag. 2064.			
7 annulàris. DC. (pu.) ring-podded.	—	1816. —	G. ○.	DC. leg. mem. t. 46.			
8 brachystachya. DC. (pu.) short-spiked.	—	—	G. h.				
9 prostrata. DC. (pu.) prostrate.	—	1820. —	G. ○.				
10 virgáta. L.O. (pu.) slender.	—	1828. 6. 8.	G. ○.				
11 vesicária. DC. (pu.) bladder-podded.	—	1825. —	G. ½.				
12 tomentosa. DC. (pu.) woolly.	—	1822. —	G. ¼.				
13 procúmbens. DC. (pu.) procumbent.	—	—	G. h.				
SUTHERLA'NDIA. DC.	SUTHERLA'NDIA. Diadelphia Decandria.						
1 frutéscens. DC. (sc.) frutescent.	C. B. S. 1683. 6. 9.	F. h.	Bot. mag. 181.				
<i>Colutèa frutéscens.</i> B.M.							
β canescens. (sc.) hoary-l'd dwarf.	—	1816. —	F. h.				
2 microphylla. DC. (sc.) small-leaved.	—	—	F. h.				

Subtribus V. *ASTRAGALEÆ*. DC. prodr. 2. p. 273.

PHÀ'CA. DC.	BASTARD VETCH. Diadelphia Decandria.						
1 bætica. DC. (wh.) hairy.	Spain. 1640.	7. 8.	H. ¼.	Moris. his. s. 2. t. 8. f. 1.			
2 frigida. DC. (st.) small.	Austria. 1795.	—	H. ¼.	Jacq. aust. t. 166.			
3 alpina. DC. (st.) Alpine.	—	1759.	—	H. ¼.	Jacq. ic. 1. t. 151.		
4 membranacea. F. (st.) membranous.	Dahuria. 1823.	—	H. ¼.				
5 lutea. Led. (ye.) yellow.	Siberia. 1827.	—	H. ¼.				
6 caespitosa. DC. (st.) tufted.	Missouri. 1826.	—	H. ¼.				
<i>Astragalus triphyllus</i> . Ph.							
7 densifolia. DC. (re.) dense-leaved.	California. 1822.	—	H. ¼.				
8 glabra. DC. (v.) smooth.	S. Europe. —	—	H. ¼.	Lodd. bot. cab. 490.			
9 austriaca. DC. (pa.) trailing.	—	1779.	5. 6.	H. ¼.	—	t. 429.	
10 astragalina. DC. (v.) procumbent.	Europe. 1771.	6. 7.	H. ¼.	Flor. dan. t. 51.			
11 lapponica. DC. (v.) Lapland.	Lapland. 1824.	7. 8.	H. ¼.	—	—	8. t. 1396.	
12 oroboides. DC. (bl.) Orobus-like.	N. Europe. 1826.	—	H. ¼.				
13 triangularis. F. (bl.) triangular.	Siberia. 1824.	—	H. ¼.				
14 exaltata. F. tall.	Altay. 1828.	H. ¼.				
15 arenaria. DC. (pu.) sand.	Siberia. 1796.	7. 8.	H. ¼.	Pall. it. 4. t. 19. f. 3-4.			
OXYTROPIS. DC. OXYTROPIS. Diadelphia Decandria.				Bot. mag. 843.			
1 montana. DC. (bl.) mountain.	Europe. 1531.	7. 8.	H. ¼.				
<i>Astragalus montanus</i> . B.M.							
2 cyanea. DC. (bl.) bright blue.	Iberia. 1826.	—	H. ¼.				
3 sordida. DC. (pa.) sordid.	N. Europe.	5. 8.	H. ¼.	Flor. dan. t. 1041.			
4 argyre'a. DC. (pu.) silvery.	Altay. 1828.	—	H. ¼.				
5 uralensis. DC. (pu.) villous.	Siberia. 1800.	6. 8.	H. ¼.	Pall. ast. t. 42. f. A.			
6 sericea. DC. (pu.) hairy mountain.	Scotland.	5. 8.	H. ¼.	Eng. bot. t. 466.			
7 cœrulea. DC. (bl.) blue-flowered.	Siberia. 1827.	6. 8.	H. ¼.	Pall. ast. t. 52.			
8 arctica. DC. (pu.) northern.	Melvill. Isl.	—	—				
9 argentata. DC. (pa.) silver-leaved.	Siberia. —	—	H. ¼.	Pall. ast. t. 48.			
10 ambigua. DC. (pu.) ambiguous.	—	1824.	5. 8.	H. ¼.	—	t. 43.	
11 setosa. DC. (pu.) pointed-calyxed.	—	1828.	6. 8.	H. ¼.	—	t. 44.	
12 Lamberti. DC. (pu.) Mr. Lambert's.	Missouri. 1811.	8. 9.	H. ¼.	Bot. mag. 2147.			
13 grandiflora. DC. (ro.) large-flowered.	Siberia. 1824.	5. 8.	H. ¼.	Pall. astr. t. 46.			
14 cändigans. DC. (pa.) hoary.	—	1827.	6. 8.	H. ¼.	—	t. 49.	
15 songarica. DC. (vi.) spear-podded.	—	1824.	5. 8.	H. ¼.	—	t. 51.	
16 sylvatica. DC. (pu.) wood.	—	1820.	—	H. ¼.	—	t. 78.	
17 longirostra. DC. (pu.) long-beaked.	—	1824.	6. 8.	H. ¼.	DC. astr. n. 17. t. 5.		
<i>Gmelini</i> . F.							
18 brevirrostra. DC. (pu.) short-beaked.	Altay. 1802.	7. 9.	H. ¼.	DC. astr. n. 19. t. 6.			
<i>Astragalus alticus</i> . Pall. astr. t. 45.							
19 filiformis. DC. (bl.) filiform.	Dahuria. 1824.	7. 8.	H. ¼.	DC. astr. n. 16. t. 4.			
20 nigrescens. DC. (bl.) black-haired.	Siberia. 1828.	—	H. ¼.	Pall. astr. t. 53.			
21 campéstris. DC. (st.) field.	Scotland.	—	H. ¼.	Eng. bot. v. 36. t. 2522.			
<i>Astragalus campestrus</i> . E.B.							
22 sulphurea. F. (st.) sulphur-color'd.	Siberia. 1827.	7. 8.	H. ¼.				
23 microphylla. DC. (pu.) small-leaved.	—	1823.	—	H. ¼.	Pall. ast. t. 76.		
24 pumila. DC. (bl.) small.	—	1828.	—	H. ¼.			
25 verticillaris. DC. (pu.) whorl-leaved.	—	1824.	—	H. ¼.	Amm. ruth. t. 19. f. 1.		
26 myriophylla. DC. (va.) many-leaved.	—	—	—	H. ¼.	Pall. astr. t. 71.		
27 oxyphylla. DC. (pu.) sharp-leaved.	—	1823.	7. 9.	H. ¼.	—	t. 74.	
28 prostrata. DC. (pu.) prostrate.	—	1820.	6. 9.	H. ¼.	—	t. 72.	
<i>Astragalus dahuricus</i> . Pall. ast. <i>Phaea prostrata</i> . L.							
29 pilosa. DC. (st.) hairy.	Siberia. 1732.	6. 8.	H. ¼.	Bot. mag. 2483.			
30 floribunda. DC. (pu.) many-flowered.	—	1827.	5. 8.	H. ½.	Pall. astr. t. 37.		
31 glabra. DC. (vi.) smooth.	Dahuria. —	—	—	H. ¼.	DC. astr. n. 31. t. 8.		
32 deflexa. DC. (vi.) small-flowered.	Siberia. 1800.	6. 8.	H. ¼.	L'Herit. stirp. t. 80.			
<i>Astragalus retroflexus</i> . Pall. astr. t. 27. <i>hians</i> . Jacq. ic. t. 252. <i>parviflorus</i> . Lam.							
33 Fischeri. DC. (bl.) Fischer's.	Siberia. 1828.	—	H. ¼.				
34 vaginalis. DC. (bl.) sheathed.	—	—	—	H. ¼.			
35 tenella. DC. (bl.) slender.	—	—	—	H. ¼.			
36 dichotoma. DC. (bl.) pubescent.	—	1815.	—	H. ¼.			
37 longicuspis. Led. (pu.) long-pointed.	—	1827.	—	H. ¼.			
ASTRAGALUS. DC. MILK-VETCH. Diadelphia Decandria. L.							
1 hypoglottis. DC. (pu.) purple mountain.	Britain.	6. 7.	H. ¼.	Eng. bot. v. 4. t. 274.			
2 dasyglottis. DC. (pu.) creeping-root'd.	Siberia. 1828.	—	H. ¼.				

LEGUMINOSÆ.

3 purpureus. DC. (pu.) purple.	S. France.	1824.	6. 7. H. ¼.	DC. astr. n. 17. t. 12.
4 sínicus. L. (ro.) Chinese annual.	China.	1763.	7. 9. H. ⊖.	Bot. mag. 1350.
lotóides. DC.				
5 onobrychioídes. DC. (pu.) Saintfoin-like.	Persia.	1828.	— H. ½.	DC. astr. n. 39. t. 16.
canescens. DC. ast. cephalotes.	Pall. astr. t. 24.			
6 leontinus. DC. (pu.) Lion's-tail.	Austria.	1816.	5. 7. H. ¼.	Lodd. bot. cab. 432.
7 Doniánus. DC. (pu.) Mr. D. Don's.	Nepaul.	1821.	8. 9. H. ¼.	
macrorhizus. D.P. non Cav.				
8 arenárius. DC. (bl.) sand.	N. Europe.	1798.	6. 7. H. ¼.	Retz. obs. 3. t. 3.
9 bayonénsis. DC. (pu.) woolly-podded.	S. France.	1828.	— H. ¼.	
10 macrópterus. DC. (pu.) large-winged.	Siberia.	1827.	— H. ¼.	
11 taúricus. DC. (pu.) Taurian.	Tauria.	1824.	7. 8. H. ¼.	Pall. astr. t. 38.
12 austriacus. DC. (pu.) Austrian.	Austria.	1840.	6. 7. H. ¼.	Jacq. aust. 2. t. 195.
13 olópterus. DC. (pu.) long-racemed.	Altay.	1827.	— H. ¼.	
14 melilotoides. DC. (pu.) Melilot-like.	Siberia.	1785.	— H. ¼.	Pall. astr. t. 41.
15 sulcatus. DC. (pu.) furrowed.		—	7. 8. H. ¼.	Jacq. vind. 3. t. 40.
16 gráclis. DC. (li.) slender.	Missouri.	1812.	6. 7. H. ¼.	
17 subulátus. DC. (pu.) awl-podded.	Tauria.	1806.	5. 8. H. ¼.	Pall. astr. t. 20.
18 ceratoides. DC. (pu.) triangular-pod'd.	Altay.	1827.	— H. ¼.	— t. 20. A.
19 virgátus. DC. (pu.) twiggy.	Siberia.	1806.	— H. ½.	— t. 18.
várius. Gmel. itin. 2. t. 17.				
20 brachylóbüs. DC. (pu.) short-podded.		1827.	— H. ½.	
21 dahúricus. DC. (pu.) Dahurian.	Dahuria.	1824.	— H. ¼.	Pall. it. 3. ap. t. w. f. 1.
22 Steveniánus. DC. (pu.) Steven's.	Iberia.	1829. H. ¼.	
23 tenuifoliús. w. (pu.) fine-leaved.	Siberia.	1780.	7. 8. H. ¼.	Swt. br. fl. gar. t. 73.
linearifoliús. DC.				
24 Onob'ychis. DC. (pu.) purple-spiked.	Europe.	1640.	6. 8. H. ¼.	Jacq. aust. 1. t. 38.
25 flaccidus. DC. (wh.) flaccid.	Iberia	1826.	— H. ¼.	
26 viminéus. DC. (pu.) twiggy.	Siberia.	1824.	7. 8. H. ½.	Pall. astr. t. 21.
27 fruticosus. DC. (pu.) woody.		1804.	6. 8. H. ½.	— t. 19.
28 adsúrgens. DC. (bl.) assurgent.		1824.	— H. ¼.	— t. 31.
29 Laxmánni. DC. (pa.) Laxmann's.		1804.	— H. ¼.	Jacq. vind. 3. t. 37.
30 missourién sis. DC. (pu.) Missouri.	Missouri.	1826.	6. 8. H. ¼.	
31 caryocárpus. DC. (vi.) swelled-podded.	Louisiana.	1811.	— H. ¼.	Bot. reg. 176.
32 pentaglótis. DC. (pu.) rough Spanish.	Spain.	1739.	6. 7. H. ⊖.	Cavan. ic. 2. t. 188.
33 Gláux. DC. (pu.) small Spanish.		1596.	— H. ⊖.	Clus. hist. 2. p. 242. ic.
34 oxyglótis. DC. (pu.) sharp-podded.	Tauria.	1824.	6. 9. H. ⊖.	
35 psiloglótis. DC. (pu.) smooth-podded.	Astracan.	1829.	— H. ⊖.	
36 cruciátus. DC. (pu.) oblong-leaved.	Iberia.	1823.	— H. ⊖.	Lobel. ic. 2. t. 83.
37 Stélla. DC. (pu.) star-podded.	S. Europe.	1658.	— H. ⊖.	
38 tribuloiðes. DC. (pu.) Caltrops-like.	Egypt.	1828.	— H. ⊖.	
39 sesámeus. DC. (pu.) Sesama.	S. Europe.	1616.	— H. ⊖.	Garid. prov. t. 12.
40 vesicárius. DC. (pa.) white Italian.	Europe.	1737.	7. 8. H. ¼.	Vill. dauph. t. 42. f. 1.
álibdus. W. et K. hung. 1. t. 40. dealbátus. Pall. astr. t. 23. f. 1.				
41 albicáulis. DC. (wh.) white-stalked.	Caucasus.	1803.	7. 8. H. ¼.	DC. astr. n. 59. t. 21.
dealbátus Pall. astr. t. 23. f. 2 et 3. Oxytropis dealbátus. P.S.				
42 Nuttallianus. DC. (bl.) Nuttall's.	N. America.	1826.	6. 8. H. ¼.	
micránthus. N. non Desv.				
43 cymbæcarpos. DC. (pa.) boat-podded.	Portugal.	1800.	— H. ⊖.	Brot. phyt. t. 59.
44 striátellus. DC. (bl.) striated.	Tauria.	1829.	— H. ⊖.	Bieb. cent. 1. t. 20.
45 annulariás. DC. (ro.) ring-podded.	Egypt.	1800.	— H. ⊖.	DC. astr. app. n. 6. t. 7.
46 hispídulus. DC. (pa.) rough-podded.		1828.	— H. ⊖.	
47 scorpioides. DC. (bl.) blue-flowered.	Spain.	1816.	6. 9. H. ⊖.	
canaliculátus. W. enum. ex Link enum. 2. p. 254.				
48 mareóticus. DC. (li.) Alexandrian.	Egypt.	1828.	— H. ⊖.	Delil. fl. eg. t. 39. f. 3.
49 corrugátus. DC. (pu.) rugged-podded.		—	— H. ⊖.	
50 reticulátus. DC. (pa.) netted.	Volga.	—	— H. ⊖.	
51 contortuplicátus. DC. (st.) wave-podd'd.	Siberia.	1764.	7. 8. H. ⊖.	Pall. astr. t. 79.
52 triméstris. DC. (st.) Egyptian.	Egypt.	1739.	— H. ⊖.	Jacq. vind. 2. t. 174.
53 aégicéras. DC. (st.) Goat's-horn.		1816.	6. 8. H. ⊖.	
54 brachycéras. DC. (st.) short-podded.	Persia.	1828.	— H. ⊖.	
55 hamósus. DC. (st.) hook-podded.	Spain.	1633.	— H. ⊖.	Moris. ox. s. 2. t. 9.f.10.
bucéras. W. enum. ex Link enum.				
56 epiglótis. DC. (st.) heart-podded.	S. Europe.	1737.	— H. ⊖.	Herm. lugd. t. 77.
57 triangulárus. DC. (st.) triangular.	1828.	— H. ⊖.	Munt. phyt. t. 110.
58 báeticus. DC. (st.) slender-podded.	S. Europe.	1759.	— H. ⊖.	
59 capitátus. w. (st.) headed.	Levant.	—	7. 8. H. ¼.	
60 microphyllus. DC. (st.) small-leaved.	Siberia.	1773.	6. 7. H. ¼.	
61 semibiloculárus. DC. (st.) semibilocular.		1804.	6. 8. H. ¼.	DC. astr. n. 64. t. 23.

62 réptans. DC.	(wh.) creeping.	Mexico.	1829.	6. 8. F. ȝ.	Willd. hort. ber. 2. t. 88.
63 odoratùs. DC.	(st.) sweet-scented.	Levant.	1816.	— H. ȝ.	DC. astr. n. 67. t. 24.
64 stipulátus. DC.	(st.) large-stipuled.	Nepaul.	1821.	8. 9. H. ȝ.	Bot. mag. 2380.
65 glycyphyllos. DC.	(st.) sweet-leaved.	Britain.	6. 7. H. ȝ.	Eng. bot. 3. t 203.
66 depresso. DC.	(st.) depressed.	Europe.	1772.	5. 6. H. ȝ.	Lodd. bot. cab. 680.
67 leucophæus. DC.	(wh.) dwarf-white.	1776.	5. 8. H. ȝ. t. 111.
68 Cicer. DC.	(st.) bladder-pod'd.	Europe.	1570.	6. 7. H. ȝ.	Jacq. aust. t. 251.
69 uliginosus. DC.	(st.) marsh.	Siberia.	1752.	6. 8. H. ȝ.	Pall. astr. t. 26.
70 mieranthus. DC.	(st.) small-blossom'd.	1824.	— H. ȝ.	Lodd. bot. cab. 372.
71 canadensis. DC.	(st.) woolly.	N. America.	1732.	6. 7. H. ȝ.	Dill. elth. t. 39. f. 45.
72 carolinianus. DC.	(st.) Carolina.	— H. ȝ.	DC. astr. n. 70. t. 26.
73 falcatus. DC.	(st.) sickle-podded.	Siberia.	1737.	— H. ȝ.	
<i>virscens. H.K.</i>					
74 falciformis. DC.	(st.) sickle-like.	Barbary.	— H. ȝ.	Desf. atl. 2. t. 207.
75 asper. DC.	(st.) rough.	Astracan.	1796.	7. 8. H. ȝ.	Jacq. ic. 1. t. 33.
<i>chloranthus. Pall. astr. t. 25.</i>					
76 chinensis. DC.	(st.) upright.	Chinese.	1795.	6. 8. H. ȝ.	Linn. f. decad. 1. t. 3.
77 galegiformis. DC.	(st.) Goat's-rue-like.	Siberia.	1729.	— H. ȝ.	Pall. astr. t. 29.
78 racemosus. DC.	(wh.) racemed.	Louisiana.	1827.	6. 8. H. ȝ.	
<i>galegoidea. N.</i>					
79 alopecias. DC.	(ye.) axillary-headed.	Siberia.	1823.	— H. ȝ.	Pall. astr. p. 12. t. 9.
80 alopecuroides. DC.	(ye.) Foxtail-like.	1737.	— H. ȝ.	—— t. 8.
81 narbonensis. DC.	(ye.) French.	S. Europe.	1780.	— H. ȝ.	—— t. 10.
82 ponticus. DC.	(ye.) villous-calyxed.	Tauria.	1824.	— H. ȝ.	—— t. 11.
83 vulpinus. DC.	(ye.) Fox-tail.	Siberia.	1815.	— H. ȝ.	—— t. 7.
<i>alopecuroides. Pall. astr. t. 7.</i>					
84 macrocephalus. DC.	(ye.) large-headed.	Persia.	1829.	— H. ȝ.	
85 christianiæ. DC.	(st.) great-yellow.	Armenia.	1737.	7. 8. H. ȝ.	Tourn. it. 2. t. 254.
86 tomentosus. DC.	(st.) downy-leaved.	Egypt.	1800.	— H. ȝ.	DC. astr. n. 80. t. 29.
87 Siebéri. DC.	(st.) Sieber's.	1828.	— H. ȝ.	
88 Bárbá-Jóvís. DC.	(pu.) silvery-leaved.	Armenia.	— H. ȝ.	
<i>Tragacantha argentea. Mill.</i>					
89 caucasicus. DC.	(wh.) Caucasian.	Caucasus.	1822.	— H. ȝ.	Pall. astr. p. 2. t. 2.
90 breviflorus. DC.	(pu.) short-flowered.	Armenia.	1826.	6. 8. H. ȝ.	Lodd. bot. cab. 1388.
91 arnacantha. DC.	(li.) prickly-leaved.	Tauria.	1640.	— H. ȝ.	Pall. astr. p. 1. t. 1.
<i>Potérium. Pall. astr. nec aliorum.</i>					
92 siculus. DC.	(bh.) Sicilian.	Sicily.	1827.	— H. ȝ.	
93 Potérium. DC.	(wh.) small goat's-thorn.	S. Europe.	1640.	— H. ȝ.	Morris. ox. s. 2. t. 13.f. 2.
94 massiliensis. DC.	(wh.) sea-side.	1827.	— H. ȝ.	Duham. arb. 2. t. 100.
<i>Tragacantha. Pall. astr. t. 4. f. 1. 2.</i>					
95 aristatus. DC.	(wh.) awned.	Pyrenees.	1791.	5. 7. H. ȝ.	Garid. aix. t. 104.
96 lagurus. DC.	(ye.) hare's-tail'd.	Armenia.	1829.	— H. ȝ.	DC. astr. n. 104. t. 36.
97 armatus. W.	(ye.) armed.	Barbary.	6. 7. H. ȝ.	Desf. atl. 2. t. 194.
<i>Anthyllis tragacanthoides. Desf.</i>					
98 halicacabus. DC.	(ye.) bladdered.	Armenia.	1806.	— H. ȝ.	Schreb. dec. p. 5. t. 3.
<i>Phaca vesicaria. Schreb.</i>					
99 laguroides. DC.	(pu.) Hare's-tail-like.	Siberia.	1829.	— H. ȝ.	Pall. astr. t. 16.
100 ammodytes. DC.	(wh.) silky-leaved.	1823.	6. 7. H. ȝ.	—— p. 7. t. 5.
101 dasyanthus. DC.	(ye.) woolly-headed.	1828.	— H. ȝ.	—— p. 79. t. 65.
<i>eriocéphalus. W. et K. hung. t. 46.</i>					
102 caprinus. DC.	(ye.) scented.	Barbary.	1683.	— H. ȝ.	Moris. ox. s. 2. t. 24.f. 3.
103 longiflorus. DC.	(ye.) long-flowered.	Tartary.	— H. ȝ.	Pall. astr. t. 60.
104 utriger. DC.	(ye.) turgid-podded.	1825.	— H. ȝ.	—— t. 61.
105 exscapus. DC.	(ye.) hairy-podded.	Hungary.	1787.	5. 7. H. ȝ.	Jacq. ic. 3. t. 561.
106 diffusus. DC.	(st.) diffuse.	Tauria.	1824.	6. 7. H. ȝ.	Pall. astr. t. 68.
107 buchtorumensis. DC.	(ye.) pointed-pod'd.	Siberia.	1826.	— H. ȝ.	—— t. 62. f. A.
108 monspessulanus. DC.	(pu.) Montpelier.	S. Europe.	1710.	7. 8. H. ȝ.	Bot. mag. 375.
109 incanus. DC.	(pu.) hoary.	1759.	6. 7. H. ȝ.	Magn. monsp. p. 32. ic.
110 macrorhizus. DC.	(pu.) long-rooted.	Spain.	1821.	6. 8. H. ȝ.	Cavan. ic. 2. t. 133.
111 brachycarpus. DC.	(pu.) short-podded.	Siberia.	1820.	— H. ȝ.	Bot. mag. 2335.
112 physodes. DC.	(pu.) inflated	1759.	— H. ȝ.	DC. astr. n. 127. t. 48.
113 Pallasii. DC.	(pu.) Pallas's.	Caucasus.	1828.	— H. ȝ.	Pall. astr. t. 59.
114 testiculatus. DC.	(bh.) woolly-podded.	1824.	— H. ȝ.	—— t. 67.
115 rupifragus. DC.	(pu.) dry rock.	Tauria.	1828.	— H. ȝ.	—— t. 70.
116 reduncus. DC.	(pu.) crook-podded.	Volga.	— H. ȝ.	—— p. 109. t. 82.
117 chlorostachys. H.C.	(st.) straw-colour'd.	N. America.	1826.	9. 10. H. ȝ.	
118 uncatus. DC.	(wh.) hooked.	Aleppo.	1768.	7. 8. H. ȝ.	
119 ràdicans. DC.	(pu.) rooting-branch'd.	1828.	— H. ȝ.	

BISE'RRULA. DC. HATCHET-VETCH. Diadelphia Decandria. L.
Pelecinus. DC. (bl.) common. S. Europe. 1640. 7. 9. H. ⊙. Tourn. inst. 417. t. 234.

Tribus III. HEDYSAREÆ. DC. prodr. 2. p. 307.

Subtribus I. CORONILLEÆ. DC. loc. cit. p. 308.

SCORPIURUS. DC. CATERPILLAR. Diadelphia Decandria. L.

1 lœvigâta. DC.	(ye.) smooth.	Levant.	1820.	6. 8.	H. ⊙.	Flor. græc. t. 718.
2 muricâta. DC.	(ye.) two-flowered.	S. Europe.	1640.	—	H. ⊙.	Moris. ox. s. 2. t. 11. f. 4.
3 sulcâta. DC.	(ye.) three-flowered.	—	1596.	—	H. ⊙.	Desf. atl. 2. f. 4.
4 acutifolia. DC.	(ye.) acute-leaved.	Corsica.	1828.	—	H. ⊙.	Viv. fl. lyb. t. 19. f. 4.
5 vermiculâta. DC.	(ye.) common.	S. Europe.	1621.	—	H. ⊙.	Moris. ox. s. 2. t. 11. f. 3.

CORONILLA. DC. CORONILLA. Diadelphia Decandria. L.

1 E'merus. DC.	(ye.) Scorpion Senna. France.	1596.	4. 6.	H. ⊙.	Bot. mag. 445.	
2 júncea. DC.	(ye.) rush-like.	—	1656.	6. 8.	F. ⊙.	Bot. reg. 820.
3 valentina. L.	(ye.) nine-leaved.	S. Europe.	1596.	2. 6.	F. ⊙.	Bot. mag. 185.
stipulâris. Lam.						
4 pentaphylla. DC.	(ye.) five-leaved.	Algiers.	—	3. 6.	F. ⊙.	Mill. ic. t. 289. f. 2?
5 argéntea. DC.	(ye) silvery-leaved.	Crete.	1664.	5. 6.	F. ⊙.	— 2. t. 289. f. 3.
6 glânea. DC.	(ye.) seven-leaved.	France.	1722.	9. 5.	F. ⊙.	Bot. mag. t. 13.
7 viminális. DC.	(bh.) slender.	Mogador.	1798.	5. 11.	G. ⊙.	Sal. par. lond. t. 13.
8 mímina. DC.	(ye.) least.	S. Europe.	1658.	7. 8.	H. ⊙.	Bot. mag. 2179.
9 coronâta. DC.	(ye.) large-headed.	—	1776.	6. 8.	H. ⊙.	Clus. hist. 1. p. 98. f. 2.
10 montâna. DC.	(ye.) mountain.	Europe.	1816.	6. 9.	H. ⊙.	Scop. carn ed. 2. t. 44.
coronâta. B.M. 907. non L.						
11 ibérica. DC.	(ye.) Iberian.	Iberia.	1818.	7. 10.	H. ⊙.	Swt. br. fl. gar. t. 25.
12 parviflôra. DC.	(ye.) small-flowered.	Crete.	1828.	—	F. ⊙.	Jacq. vind. 1. t. 25.
13 crética. DC.	(v.) Cretan.	—	1731.	6. 8.	H. ⊙.	Bot. mag. 258.
14 vâria. DC.	(pu.) purple.	Europe.	1597.	6. 11.	H. ⊙.	

ASTROL'OBium. DC. ASTROL'OBium. Diadelphia Decandria.

1 ebracteátum. DC.	(ye.) round-podded.	Portugal.	—	5. 6.	H. ⊙.	Dalech. his. 1. 487. f. 1.
	<i>Ornithòpus lœvigatus</i> . Sm. <i>nudiflorus</i> . Lag. <i>ebracteatus</i> . Brot.					
2 dûrum. DC.	(ye.) perennial.	S. Europe.	1805.	6. 7.	H. ⊙.	Cavan. ic. 1. t. 41.
3 repândum. DC.	(ye.) repand.	Barbary.	—	—	H. ⊙.	Lam. ill. t. 631. f. 4.
4 scorpíoides. DC.	(ye.) Purslane-leav'd.	S. Europe.	1596.	—	H. ⊙.	Cavan. ic. 1. t. 37.
	<i>Ornithòpus scorpioides</i> . L. <i>trifoliatus</i> . Lam.					

ORNITH'OPUS. DC. BIRD'S-FOOT. Diadelphia Decandria. L.

1 compréssus. DC.	(ye.) hairy.	S. Europe.	1730.	6. 7.	H. ⊙.	Berg. phyt. t. 191.
2 intermèdius. RO. (ro.) intermediate.		Portugal.	1816.	—	H. ⊙.	
satívus. Brot. fl. lus. 2. p. 160.						

3 perpusillus. DC.	(v.) common.	Britain.	—	5. 6.	H. ⊙.	Eng. bot. v. 6. t. 369.
--------------------	--------------	----------	---	-------	-------	-------------------------

HIPPOCR'EPIs. DC. HORSESHOE-VETCH. Diadelphia Decandria. L.

1 baleàrica. DC.	(ye.) shabby.	Minorca.	1776.	5. 6.	F. ⊙.	Bot. mag. 427.
2 helvética.	(ye.) Swiss.	Switzerland.	1822.	5. 8.	H. ⊙.	Eng. bot. 1. t. 31.
3 comôsa. DC.	(ye.) tufted.	England.	—	4. 8.	H. ⊙.	Ten. fl. neap. 2. t. 69.
4 glânea. DC.	(ye.) glaucous.	Naples.	1825.	5. 8.	H. ⊙.	
5 syriaca. E.	(ye.) Syrian.	Syria.	1826.	6. 8.	H. ⊙.	
6 scâbra. DC.	(ye.) rough.	Spain.	—	—	H. ⊙.	
7 multisiliquosa. DC.	(ye.) many-podded.	S. Europe.	1683.	—	H. ⊙.	
8 monocárpa. DC.	(ye.) one-podded.	Caucasus.	1825.	—	H. ⊙.	
9 unisiliquosa. DC.	(ye.) single-podded.	Italy.	1570.	6. 7.	H. ⊙.	Lam. ill. t. 630. f. 3.
10 ciliâta. DC.	(ye.) ciliated.	Spain.	1823.	6. 8.	H. ⊙.	
11 biflôra. DC.	(ye.) two-flowered.	—	1824.	—	H. ⊙.	

SECUR'GERA. DC. HATCHET-VETCH. Diadelphia Decandria.

Coronilla. DC.	(ye.) common.	S. Europe.	1562.	7. 8.	H. ⊙.	Lam. ill. t. 629.
----------------	---------------	------------	-------	-------	-------	-------------------

Subtribus II. EUHEDISAREÆ. DC. prodr. 2. p. 313.

PICT'ETIA. DC. PICT'ETIA. Diadelphia Decandria.

aristâta. DC. (ye.) awned. St. Domingo. 1820. 4. 6. S. h. DC. leg. mem. t. 47. f. 5.

Æschynomene aristâta. Jacq. schoenb. 2. t. 237.

ORMOCA'RPUM. DC.	ORMOCA'RPUM.	Diadelphia Decandria.				
sennoides. DC.	(ye.) Senna-like.	E. Indies.	1818.	6.7.	S. $\frac{1}{2}$.	
<i>Hedy'sarum sennoides.</i> w.						
POIR'ETIA. DC.	POIR'ETIA.	Monadelphia Decandria.				
scándens. DC.	(ye.) climbing.	St. Domingo.	1826.	S. $\frac{1}{2}$.	Venten. choix. t. 42.
<i>Turpína punctatâ.</i> P.S.	<i>Gly'cine punctatâ.</i>	Lam. ill. t. 609.				
TETRAN'EMA.	TETRAN'EMA.	Monadelphia Decandria.				
nútans.	(li.) drooping-flđ.	E. Indies.	1823.	5.8.	S. $\frac{1}{2}$.	Bot. mag. 2867.
<i>Hedy'sarum nútans.</i> Gr.	<i>Desmodium nútans.</i> B.M.					
ZO'R'NIA. DC.	ZO'R'NIA.	Monadelphia Decandria.				
1 angustifòlia. DC.	(ye.) narrow-leaved.	E. Indies.	1733.	7.8.	S. \odot .	Rheed. mal. 9. t. 82.
<i>diphylla.</i> P.S.	<i>Hedy'sarum diphyllum.</i> L.					
2 reticulàta. DC.	(ye.) netted-bracted.	W. Indies.	—	S. \odot .	
3 thymifòlia. DC.	(ye.) Thyme-leaved.	Mexico.	1826.	—	G. $\frac{1}{2}$.	
4 zeylonénis. DC.	(ye.) Ceylon.	Ceylon.	1733.	—	S. \odot .	Burm. zeyl. t. 50. f. 1.
	<i>conjugatâ.</i> Sm.	<i>Hedy'sarum conjugatum.</i> w.				
5 dyctiocárpa. DC.	(ye.) netted-podded.	N. Holland.	1824.	—	H. \odot .	
6 tetraphylla. DC.	(ye.) four-leaved.	Carolina.	1823.	—	H. $\frac{1}{2}$.	
7 capénis. DC.	(ye.) Cape.	C. B. S.	1824.	5.6.	G. $\frac{1}{2}$.	Mich. amer. 2. t. 41.
						Thunb. n. ac. ups. 6.t.3.
STYLOSA'NTHES. DC.	STYLOSA'NTHES.	Monadelphia Decandria.				
1 procúmbens. DC.	(ye.) procumbent.	Jamaica.	1820.	6.8.	S. $\frac{1}{2}$.	Swar.ac.h.1789.t.11.f.1
2 viscósa. DC.	(ye.) viscous.	—	—	—	S. $\frac{1}{2}$.	— t. 9. f. 2.
3 mucronata. DC.	(ye.) mucronate.	E. Indies.	1818.	—	S. $\frac{1}{2}$.	Burm. zeyl. t. 106. f. 2.
	<i>Hedy'sarum hamatum.</i> Burm.	<i>A'rachis fruticosa.</i> Retz.				
4 elatior. DC.	(ye.) tall.	Carolina.	1823.	7.8.	H. $\frac{1}{2}$.	
5 guianénis. DC.	(ye.) Guiana.	Guiana.	1826.	—	S. \odot .	Pluk. alm. t. 447. f. 7.
						Aubl. guian. 2. t. 309.
ADE'SMIA. DC.	ADE'SMIA.	Decandria Monogynia.				
1 muricata. DC.	(ye.) prickly-podded.	Patagonia.	1793.	6.7.	G. $\frac{1}{2}$.	Jacq. ic. 3. t. 568.
	<i>Hedy'sarum muricatum.</i> Jacq.	<i>Patagónium Hedsaroídes.</i> Schk.				
2 hispídula. DC.	(ye.) hispid.	Peru.	1826.	6.8.	G. $\frac{1}{2}$.	DC. leg. mem. t. 48.
3 pappòs. DC.	(ye.) featherly-podd'd.	Chili.	1823.	—	G. $\frac{1}{2}$.	
ÆSCHYNO'MENE. DC.	ÆSCHYNO'MENE.	Diadelphia Decandria. L.				
1 áspéra. DC.	(ye.) rough-stem'm'd.	E. Indies.	1759.	6.7.	S. \odot .	Breyn. cent. 51. t. 52.
2 sensitiva. DC.	(wh.) sensitive.	W. Indies.	1733.	5.8.	S. $\frac{1}{2}$.	Plum. ic. t. 139.
3 índica. DC.	(ye.) Indian.	E. Indies.	1799.	6.7.	S. \odot .	Rheed. mal. 9. t. 18.
4 americana. DC.	(ye.) hairy.	Jamaica.	1732.	7.8.	S. \odot .	Sloan.jam.1.t.118.f.3.
5 pátila. DC.	(ye.) spreading.	Mauritius.	1826.	6.8.	S. $\frac{1}{2}$.	
6 subviscosa. DC.	(v.) glandular.	E. Indies.	1820.	9.10.	S. \odot .	
7 híspida. DC.	(ye.) hispid.	N. America.	1803.	8.9.	H. \odot .	Bart. fl. amer. t. 29.
8 micránta. DC.	(ye.) small-flowered.	Madagascar.	1826.	6.8.	S. $\frac{1}{2}$.	
9 viscidula. DC.	(ye.) viscous.	N. America.	1818.	7.8.	H. \odot .	
10 falcata. DC.	(ye.) sickle-podded.	Brazil.	1825.	—	S. \odot .	
II péndula. Boj.	(ye.) pendulous.	Madagascar.	—	—	S. \odot .	
SMI'THIA. DC.	SMI'THIA.	Diadelphia Decandria.				
1 sensitiva. DC.	(ye.) sensitive.	E. Indies.	1785.	7.10.	S. \odot .	Sal. par. lond. t. 92.
2 conférta. DC.	(ye.) close-flowered.	N. Holland.	1825.	—	S. \odot .	
3 áspera. H.B.	(ye.) rough.	E. Indies.	1812.	—	S. \odot .	
LO'UREA. DC.	LO'UREA.	Diadelphia Decandria.				
1 Vespertiliònìs. DC. (w.) bat-winged.	Cochinchina.	1780.	7.8.	S. δ .	Jacq. ic. 3. t. 566.	
	<i>Hedy'sarum Vespertiliònìs.</i> Jacq.	<i>Chrística lunata.</i> Mœnch.				
2 renifòrmis. DC.	(pu.) kidney-leaved.	China.	1826.	—	S. δ .	
UR'ARIA. DC.	UR'ARIA.	Diadelphia Decandria.				
1 picta. DC.	(pu.) painted-leaved.	Guinea.	1788.	6.8.	S. δ .	Jacq. ic. 3. t. 567.
	<i>Hedy'sarum pictum.</i> Jacq.					
2 crinita. DC.	(br.) crook-podded.	E. Indies.	1780.	7.9.	S. $\frac{1}{2}$.	Burm. ind. t. 56.
3 lagopòides. DC.	(wh.) Harefoot-like.	—	1790.	—	S. $\frac{1}{2}$.	— t. 53. f. 2.
	<i>Hedy'sarum lagopodioides.</i> L.	<i>Lespedéza lagopodioides.</i> P.S.				
4 Lágopus. DC.	(pu.) Hare's-foot.	Neapul.	1823.	6.8.	S. $\frac{1}{2}$.	
5 hamosa.	(wh.) hooked.	E. Indies.	—	6.7.	S. $\frac{1}{2}$.	
	<i>Doòdia hamòsa et Hedy'sarum hamòsum.</i> H.B.					
6 alopecuroïdes.	(wh.) Fox-tail.	E. Indies.	—	—	S. $\frac{1}{2}$.	
	<i>Doòdia alopecuroïdes et Hedy'sarum alopecuroïdes.</i> H.B.					
7 lagocéphala. DC.	(ye.) hairy.	Brazil.	—	7.9.	S. $\frac{1}{2}$.	
NICÓL'SONIA. DC.	NICÓL'SONIA.	Diadelphia Decandria.				
barbáta. DC.	(pu.) bearded.	W. Indies.	1826.	6.8.	S. $\frac{1}{2}$.	
DESM'ODIUM. DC.	DESM'ODIUM.	Diadelphia Decandria.				
1 dùbiūm. B.R.	doubtful.	Neapul.	1823.	7.8.	S. $\frac{1}{2}$.	Bot. reg. 967.
2 umbellátum. DC. (wh.) umbel-flower'd.	E. Indies.	1801.	6.7.	S. $\frac{1}{2}$.	Jacq. schoenb. 3, t. 297.	
3 auriculátum. DC. (pu.) eared.	Timor.	1825.	—	S. $\frac{1}{2}$.		

LEGUMINOSÆ.

4 triquètrum. DC. (<i>pu.</i>) triangular.	E. Indies.	1802.	7. 8.	S. $\frac{1}{2}$.	Burm. ind. t. 52. f. 2.
5 pseudotriquètrum. DC. (<i>pu.</i>) three-sided.	Nepaul.	1818.	—	S. $\frac{1}{2}$.	
6 alàtum. DC. (<i>pu.</i>) winged-stalked.	E. Indies.	1809.	—	S. $\frac{1}{2}$.	
7 g'yrans. DC. (<i>re.</i>) moving plant.	—	1775.	7. 9.	S. $\frac{1}{2}$.	Jacq. ic. t. 565.
8 gyroïdes. DC. (<i>li.</i>) gyrans-like.	—	1818.	—	S. $\frac{1}{2}$.	
9 ganglîcum. DC. (<i>li.</i>) oval-leaved.	—	1762.	—	S. $\frac{1}{2}$.	Burm. zeyl. t. 49.
10 maculâtum. DC. (<i>li.</i>) spotted.	—	1732.	6. 8.	S. \odot .	Dill. elth. t. 141. f. 168.
<i>Hedy'sarum maculatum.</i> L. <i>Aeschynomene maculata.</i> Poir.					
11 sagittatum. DC. (<i>re.</i>) arrow-leaved.	—	1807.	—	S. $\frac{1}{2}$.	
12 angustifòlium. DC. (<i>pu.</i>) narrow-leaved.	Mexico.	1825.	—	S. $\frac{1}{2}$.	
13 latifòlium. DC. (<i>bl.</i>) broad-leaved.	E. Indies.	1800.	7. 8.	S. $\frac{1}{2}$.	Bot. reg. t. 356.
<i>Hedy'sarum latifolium.</i> B.R.					
14 velutinum. DC. (<i>vi.</i>) velvety.	Trinidad.	1820.	—	S. $\frac{1}{2}$.	
15 lasiocarpum. DC. (<i>ro.</i>) woolly-podded.	Guinea.	1822.	7. 9.	S. $\frac{1}{2}$.	Beauv. fl. ów. 1. t. 18.
16 canadense. DC. (<i>pu.</i>) Canadian.	N. America.	1640.	—	H. $\frac{1}{2}$.	Corn. canad. t. 45.
17 canescens. DC. (<i>pu.</i>) canescent.	—	1733.	—	H. $\frac{1}{2}$.	
18 marylandicum. DC. (<i>p.</i>) Maryland.	—	1725.	7. 10.	H. $\frac{1}{2}$.	Dill. elth. t. 144. f. 171.
19 obtusum. DC. (<i>vi.</i>) obtuse.	—	1805.	7. 8.	H. $\frac{1}{2}$.	
20 ciliare. DC. (<i>vi.</i>) fringed.	—	1823.	—	H. $\frac{1}{2}$.	
21 viridiflorum. DC. (<i>pu.</i>) green-flower'd.	—	1787.	7. 9.	H. $\frac{1}{2}$.	Pluk. alm. t. 308. f. 5.
22 lavigatum. DC. (<i>pu.</i>) smooth-stalk'd.	—	1826.	—	H. $\frac{1}{2}$.	
23 bracteatum. DC. (<i>pu.</i>) bracted.	—	1806.	7. 8.	H. $\frac{1}{2}$.	
24 cuspidatum. DC. (<i>pu.</i>) sharp-pointed.	—	—	—	H. $\frac{1}{2}$.	
25 glabrum. DC. (<i>pu.</i>) smooth.	—	1824.	7. 9.	H. $\frac{1}{2}$.	
26 paniculatum. DC. (<i>pu.</i>) paniced.	—	1781.	7. 8.	H. $\frac{1}{2}$.	Pluk. alm. t. 432. f. 6.
27 acuminatum. DC. (<i>pu.</i>) glutinous.	—	1805.	—	H. $\frac{1}{2}$.	
<i>Hedy'sarum glutinosum.</i> W. <i>acuminatum.</i> Ph.					
28 nudiflorum. DC. (<i>pu.</i>) naked-flowered.	—	1823.	—	H. $\frac{1}{2}$.	
29 pauciflorum. DC. (<i>wh.</i>) small-flowered.	—	1828.	—	H. $\frac{1}{2}$.	
30 rotundifolium. DC. (<i>pu.</i>) round-leaved.	—	1733.	—	H. $\frac{1}{2}$.	
31 Aparines. DC. (<i>li.</i>) hairy-stalked.	Mexico.	1820.	7. 9.	S. $\frac{1}{2}$.	
32 cinereum. DC. (<i>pu.</i>) white-branched.	—	1826.	—	S. $\frac{1}{2}$.	H. et B.n.plam. 6.t. 599.
33 uncinatum. DC. (<i>vi.</i>) uncinate.	Caracas.	1823.	6. 7.	S. $\frac{1}{2}$.	Jacq. schoenb. 3. t. 208.
34 incanum. DC. (<i>pu.</i>) hoary.	W. Indies.	1820.	7. 8.	S. $\frac{1}{2}$.	Plum.ed.Bur.t.149.f.1.
35 erythrinæfòlium. DC. (<i>wh.</i>) <i>Erythrina</i> -ld.	Trinidad.	—	—	S. $\frac{1}{2}$.	
36 supinum. DC. (<i>pu.</i>) trailing.	Jamaica.	—	—	S. $\frac{1}{2}$.	Sloan. jam. 1. t. 118. f. 2.
37 tortuosum. DC. (<i>pu.</i>) twisted-podded.	—	1781.	—	S. $\frac{1}{2}$.	— 1. t. 116. f. 2.
38 adscendens. DC. (<i>pu.</i>) ascending.	—	1820.	7. 9.	S. $\frac{1}{2}$.	H. et B.n.plam. 6.t. 597.
39 mexicanum. (<i>bl.</i>) Mexican.	Mexico.	1820.	11. 5.	S. $\frac{1}{2}$.	Bot. reg. 815.
<i>Hedy'sarum adscendens</i> β <i>cærileum.</i> DC.					
40 trigonum. DC. (<i>li.</i>) trigonal.	Jamaica.	1733.	7. 9.	S. $\frac{1}{2}$.	
41 spirale. DC. (<i>wh.</i>) spiral-podded.	—	...	—	S. $\frac{1}{2}$.	
42 áspерум. DC. (<i>pa.</i>) harsh.	—	1820.	6. 8.	S. $\frac{1}{2}$.	
43 cæspitosum. DC. (<i>bh.</i>) tufted.	Mauritius.	—	—	S. $\frac{1}{2}$.	
44 mauritiánum. DC. (<i>wh.</i>) creeping-rooted.	—	1822.	7. 9.	S. $\frac{1}{2}$.	
45 Scálpe. DC. (<i>pu.</i>) repand-leaved.	Madagascar.	1823.	7. 8.	S. $\frac{1}{2}$.	
<i>Hedy'sarum repandum.</i> Poir. non Vahl.					
46 diversifolium. DC. (<i>li.</i>) different-leav'd.	—	1820.	—	S. $\frac{1}{2}$.	
47 spectabile. (<i>pu.</i>) showy.	Zanzibar.	1825.	—	S. $\frac{1}{2}$.	
<i>Hedy'sarum spectabile.</i> Boj.					
48 triflorum. DC. (<i>pu.</i>) three-flowered.	India.	1796.	6. 7.	S. $\frac{1}{2}$.	Burm. ind. t. 54. f. 2.
49 heterophyllo. DC. (<i>pu.</i>) various-leav'd.	Ceylon.	1820.	—	S. $\frac{1}{2}$.	Burm. zeyl. t. 54. f. 1.
50 parvifolium. DC. (<i>pu.</i>) slender.	Nepaul.	1823.	7. 9.	S. $\frac{1}{2}$.	
<i>Hedy'sarum tenellum.</i> D.P. non Kth.					
51 Roxburghii. (<i>pu.</i>) Roxburgh's.	E. Indies.	1825.	—	S. $\frac{1}{2}$.	
<i>diffusum.</i> DC. p. 335. non p. 336.					
52 floribundum. D.P. (<i>pu.</i>) abundant-fl'd.	Nepaul.	1823.	—	S. $\frac{1}{2}$.	
53 retusum. D.P. (<i>pu.</i>) retuse-leaved.	—	—	—	S. $\frac{1}{2}$.	
54 confertum. DC. (<i>li.</i>) close-flowered.	—	1826.	—	S. $\frac{1}{2}$.	
55 sambuénse. D.P. (<i>li.</i>) villous-leaved.	—	—	—	S. $\frac{1}{2}$.	
56 concinnum. DC. (<i>li.</i>) neat.	—	1823.	—	S. $\frac{1}{2}$.	
57 trichocáulim. DC. (<i>vi.</i>) hairy-stalked.	—	1827.	—	S. $\frac{1}{2}$.	
58 Leschenaultia. DC. (<i>li.</i>) Leschenault's.	E. Indies.	—	—	S. $\frac{1}{2}$.	
59 capitatum. DC. (<i>pu.</i>) headed.	Ceylon.	—	7. 8.	S. $\frac{1}{2}$.	Burm. ind. t. 54. f. 1.
<i>Hedy'sarum conicum.</i> Poir.					
60 laxum. DC. (<i>li.</i>) loose.	Nepaul.	1828.	—	S. $\frac{1}{2}$.	
61 tomentosum. DC. (<i>wh.</i>) woolly.	China.	1782.	6. 7.	G. $\frac{1}{2}$.	
62 rēpens. DC. (<i>pu.</i>) creeping.	W. Indies.	1820.	—	S. $\frac{1}{2}$.	
63 malacophyllum. DC. (<i>li.</i>) soft-leaved.	Manila.	1822.	—	S. $\frac{1}{2}$.	

64	grandifolium. DC. (<i>li.</i>) large-leaved.	E. Indies.	1822.	6. 7.	S. $\frac{1}{2}$.	
65	serotinum. DC. (<i>pu.</i>) late-flowering.	N. America.	1806.	8. 10.	H. $\frac{1}{2}$.	
66	cephalotes. H.B. (<i>li.</i>) close-headed.	E. Indies.	1825.	6. 8.	S. $\frac{1}{2}$.	
67	altissimum. (<i>pa.</i>) tallest.		—	...	S. $\frac{1}{2}$.	
68	heptaphyllum. (<i>li.</i>) seven-leaved.		1820.	6. 8.	S. $\frac{1}{2}$.	
69	abyssinicum. DC. (<i>li.</i>) Abyssinian.	Abyssinia.	1802.	7.	S. $\frac{1}{2}$.	
70	bracteatum. H.B. (<i>li.</i>) bracted.	E. Indies.	1820.	7. 8.	S. $\frac{1}{2}$.	
71	rhomboideum. (<i>pa.</i>) rhomboid.		—	7. 9.	S. $\frac{1}{2}$.	
	<i>Hedy' sarum rhombifolium.</i> H.B. non Elliot.					
72	orbiculatum. H.B. (<i>li.</i>) round-leaved.		1818.	—	S. $\frac{1}{2}$.	
73	recurvatum. H.B. (<i>li.</i>) recurved.		1824.	—	S. $\frac{1}{2}$.	
74	divergens. H.B. (<i>pa.</i>) divergent.		—	—	S. $\frac{1}{2}$.	
75	purpureum. H.B. (<i>pu.</i>) purple-flowered.		1818.	6. 8.	S. $\frac{1}{2}$.	
76	pátentis. H.B. (<i>li.</i>) spreading		1824.	—	S. $\frac{1}{2}$.	
77	arboreum. tree.		1818.	S. $\frac{1}{2}$.	
	<i>Hedy' sarum arboreum.</i> H.B.					
78	lacteum. Boj. (<i>st.</i>) milk-coloured.	Zanzibar.	1825.	10. 11.	S. $\frac{1}{2}$.	
79	Helsinbergii. Boj. Helsingberg's.	Madagascar.	—	S. $\frac{1}{2}$.	
DICE'RMA. DC.	DICE'RMA.	Diadelphus Decandria.				
1	pulchellum. DC. (<i>ye.</i>) neat.	E. Indies.	1799.	7. 8.	S. $\frac{1}{2}$.	Burm. zeyl. t. 52.
	<i>Hedy' sarum pulchellum.</i> w. Zórnia pulchella. P.S. <i>Phyllodium pulchellum.</i> Desv.					
2	élégans. DC. (<i>ye.</i>) elegant.	China.	1826.	7. 8.	G. $\frac{1}{2}$.	
3	biarticulatum. DC. (<i>ye.</i>) two-jointed.	E. Indies.	1808.	—	S. $\frac{1}{2}$.	Burm. zeyl. t. 50. f. 2.
	<i>Hedy' sarum biarticulatum.</i> L.					
HEDY'SARUM. DC.	HEDY'SARUM.	Diadelphus Decandria. L.				
1	grandiflorum. DC. (<i>st.</i>) large-flowered.	Iberia.	1820.	6. 8.	H. $\frac{1}{2}$.	Pall. itin. 2. p. 743. t. Y.
	<i>sericeum.</i> M.B. <i>Astragalus grandiflorus.</i> L.					
2	argenteum. DC. (<i>pu.</i>) silvery.	Siberia.	1827.	—	H. $\frac{1}{2}$.	Gmel. sib. 4. t. 13.
3	candidum. DC. (<i>pu.</i>) silver-leaved.		1796.	7. 8.	H. $\frac{1}{2}$.	
4	splendens. DC. (<i>pa.</i>) glossy-leaved.		1828.	—	H. $\frac{1}{2}$.	
5	pállidum. DC. (<i>pa.</i>) pale-flowered.	Barbary.	1811.	—	H. $\frac{1}{2}$.	
6	coronarium. DC. (<i>sc.</i>) F. Honeysuckle. Italy.		1596.	6. 8.	H. $\frac{1}{2}$.	
	<i>a coccineum.</i> (<i>sc.</i>) scarlet-flowered.		—	—	H. $\frac{1}{2}$.	Gært. fruct. 2. t. 155.
	<i>β ábum.</i> (<i>wh.</i>) white-flowered.		—	—	H. $\frac{1}{2}$.	
7	humble. DC. (<i>pu.</i>) dwarf.	Spain.	1640.	7. 8.	H. $\frac{1}{2}$.	J. Bauh. his. 2.p. 336. f. 1.
8	spinosissimum. DC. (<i>pu.</i>) thorny.		1731.	—	H. $\frac{1}{2}$.	Pluk. alm. t. 50. f. 2.
9	capitatum. DC. (<i>ro.</i>) headed.	Barbary.	...	—	H. $\frac{1}{2}$.	Bot. mag. 1251.
10	flexuosum. DC. (<i>pu.</i>) wave-podded.	Asia.	1680.	—	H. $\frac{1}{2}$.	Schk. hand. 2. t. 207.
11	cárneum. DC. (<i>fl.</i>) flesh-coloured.	Siberia.	1818.	—	H. $\frac{1}{2}$.	Lodd. bot. cab. 312.
12	elongatum. F. (<i>pu.</i>) long-stalked.	Russia.	1823.	6. 8.	H. $\frac{1}{2}$.	— t. 1401.
13	lasiocarpum. DC. (<i>pu.</i>) woolly-podded.	Altay.	1824.	—	H. $\frac{1}{2}$.	
14	taúricum. DC. (<i>ro.</i>) Taurian.	Tauria.	1804.	7. 8.	H. $\frac{1}{2}$.	Bieb. cent. 2. t. 85.
	<i>róseum.</i> Bot. mag. t. 996. non DC.					
15	róseum. DC. (<i>ro.</i>) rose-coloured.	Siberia.	1803.	—	H. $\frac{1}{2}$.	
16	altaicum. DC. (<i>pu.</i>) Altay.		1828.	—	H. $\frac{1}{2}$.	
17	venustum. DC. (<i>pu.</i>) handsome.	Altay.	—	—	H. $\frac{1}{2}$.	
18	fruticosum. DC. (<i>li.</i>) shrubby.	Siberia.	1782.	6. 7.	H. $\frac{1}{2}$.	Pall. it. 3. . 5. f. 1.
19	ibéricum. DC. (<i>li.</i>) Iberian.	Iberia.	1824.	6. 8.	H. $\frac{1}{2}$.	
20	Razoumowiánum. DC. (<i>li.</i>) net-podded.	Scythia.	1828.	—	H. $\frac{1}{2}$.	
21	cretaceum. DC. (<i>pu.</i>) slender-spiked.	Volga.	1818.	—	H. $\frac{1}{2}$.	
22	consanguíneum. DC. (<i>pu.</i>) related.	Altay.	—	—	H. $\frac{1}{2}$.	
23	obscürum. DC. (<i>pu.</i>) creeping-rooted.	Europe.	1640.	7. 8.	H. $\frac{1}{2}$.	
24	brachisèma. DC. (<i>pu.</i>) short-lipped.	Altay.	1820.	—	H. $\frac{1}{2}$.	
25	sibiricum. DC. (<i>pu.</i>) Siberian.	Siberia.	1798.	6. 7.	H. $\frac{1}{2}$.	
26	alpinum. W.	Alpine.	—	—	H. $\frac{1}{2}$.	
27	caucásicum. DC. (<i>pu.</i>) Caucasian.	Caucasus.	1824.	6. 8.	H. $\frac{1}{2}$.	
28	argyrophyllum. F. (<i>pu.</i>) silvery-leaved.	Altay.	1827.	—	H. $\frac{1}{2}$.	
ONOBRY'CHIS. DC.	SAINTFOIN.	Diadelphus Decandria.				
1	sativa. DC. (<i>ro.</i>) common.	Britain.	6. 7.	H. $\frac{1}{2}$.	Eng. bot. v. 2. t. 96.
	<i>Hedy' sarum Onobry'chis.</i> Jacq. aust. t. 352.					
2	tanaitica. F. (<i>ro.</i>) spreading.	Siberia.	1823.	—	H. $\frac{1}{2}$.	
3	carpáthica. DC. (<i>ro.</i>) Carpathian.	Carp. Mount.	1818.	—	H. $\frac{1}{2}$.	
4	incurvata. DC. (<i>ro.</i>) incurved.	1826.	—	H. $\frac{1}{2}$.	
5	montana. DC. (<i>pu.</i>) mountain.	Pyrenees.	1816.	—	H. $\frac{1}{2}$.	
6	conférta. DC. (<i>pu.</i>) close-flowered.	Caucasus.	1827.	6. 8.	H. $\frac{1}{2}$.	
7	procumbens. DC. (<i>ro.</i>) procumbent.	Iberia.	1823.	—	H. $\frac{1}{2}$.	
8	supina. DC. (<i>ro.</i>) trailing.	S. Europe.	1816.	—	H. $\frac{1}{2}$.	
9	glabra. DC. (<i>ro.</i>) smooth.	Tauria.	1829.	—	H. $\frac{1}{2}$.	
10	arenaria. DC. (<i>ro.</i>) sand.	Hungary.	1818.	—	H. $\frac{1}{2}$.	

LEGUMINOSÆ.

11	mareótica. DC.	(ro.) villous-calyxed.	Siberia.	1829.	6. 8. H. ȝ.		
12	álba. DC.	(wh.) white-flowered.	Hungary.	1804.	7. 8. H. ȝ.	W. et K. hung. 2. t. 111.	
13	petr'aea. DC.	(wh.) stone.	Caucasus.	1818.	6. 8. H. ȝ.		
14	grátilis. DC.	(li.) slender.	Poland.	1828.	— H. ȝ.		
15	saxátilis. DC.	(wh.) rock.	S. France.	1790.	— H. ȝ.		
16	Cáput-gálli. DC.	(vi.) Cock's-head.	S. Europe.	1731.	7. 8. H. ȝ.	All. ped. 1191. t. 19. f. 1.	
17	Crísta-gálli. DC.	(vi.) Cock's-comb.	—	1710.	6. 8. H. ȝ.	Lobel. ic. 2. p. 81. f. 1.	
18	foveoláta. DC.	(vi.) Sicilian.	Sicily.	1828.	— H. ȝ.	Gært. fruct. 2. t. 148.	
19	aéquidentáta. DC.	(re.) equal-toothed.	Levant.	1823.	— H. ȝ.	Desv. j. bot. 1. t. 23.	
20	Pallásii. DC.	(st.) Pallas's.	Tauria.	—	— H. ȝ.	Bieb. pl. ros. 1. t. 35.	
	Buxbaumiána. s.s.	<i>Hedy' sarum Pallásii.</i> w. Buxb.	cent. 2. t. 42.				
21	radiáta. DC.	(st.) rayed.	Caucasus.	1823.	7. 9. H. ȝ.	Desf. an. mus. 12. t. 13.	
	<i>Hedy' sarum Buxbaumii.</i> M.B.						
22	ptolemáica. DC.	(ye.) striped-flower'd.	Egypt.	1829.	— H. ȝ.	Delil. fl. eg. t. 39. f. 1.	
	ELE'OTIS. DC.	ELE'OTIS.	Diadelphia Decandria.				
1	monophyll'a. DC.	(li.) simple-leaved.	E. Indies.	1820.	4. 8. S. ȝ.	Burm. ind. t. 50. f. 2.	
2	sorória. DC.	(bh.) ternate-leaved.	—	—	— S. ȝ.	Desv. j. bot. 3. t. 6.f.31.	
	<i>Hedy' sarum soròrium.</i> L.						
	LESPED'EZA. DC.	LESPED'EZA.	Diadelphia Decandria.				
1	júncea. DC.	(wh.) slender-branch'd.	Siberia.	1776.	7. 8. G. ȝ.	Linn. f. dec. 1. t. 4.	
	<i>Hedy' sarum júncum.</i> L. <i>Anthyl'lis cuneàta.</i> Dum. Cours.						
2	reticuláta. DC.	(pu.) netted.	N. America.	1823.	— H. ȝ.		
3	sessiliflóra. DC.	(pu.) sessile-flowered.	—	—	— H. ȝ.		
4	eriocárpa. DC.	(vi.) woolly-podded.	Nepaul.	1812.	7. 9. G. ȝ.		
5	frutescens. DC.	(pu.) frutescent.	Carolina.	1739.	— G. ȝ.		
6	capitáta. DC.	(li.) headed.	N. America.	1789.	6. 8. H. ȝ.		
7	angustifólia. DC.	(li.) narrow-leaved.	Carolina.	1827.	— H. ȝ.		
8	Stüvei. DC.	(pu.) Stuve's.	New Jersey	1823.	7. 9. H. ȝ.		
9	polystachya. DC.	(wh.) hairy.	N. America.	1789.	6. 8. H. ȝ.		
	<i>Hedy' sarum hirtum.</i> W.						
10	villósa. DC.	(wh.) villous.	—	1812.	— H. ȝ.		
11	violacea. DC.	(vi.) violet-colour'd.	—	1789.	7. 8. H. ȝ.		
12	divergens. DC.	(vi.) divergent.	—	1827.	— H. ȝ.		
13	procumbens. DC.	(pu.) procumbent.	—	1812.	— H. ȝ.		
14	prostráta. DC.	(pu.) prostrate.	—	1818.	— H. ȝ.		
	E'BENUS. DC.	E'BENUS.	Monadelphia Decandria.				
1	crética. DC.	(bh.) Cretan.	Crete.	1737.	6. 7. F. ȝ.	Mich. amer. 2. t. 40.	
	<i>Anthyl'lis crética.</i> Lam.						
2	pinnáta. DC.	(pi.) wing-leaved.	Barbary.	1786.	7. 8. G. ȝ.	Bot. mag. 1092.	
	<i>Anthyl'lis sericea.</i> W.						
3	Sibthórii. DC.	(pi.) Sibthorp's.	Greece.	1826.	— G. ȝ.	Desf. ac. par. 1. t. 3.	
	FLEMÍ'NGIA. DC.	FLEMÍ'NGIA.	Diadelphia Decandria.				
1	strícta. DC.	(re.) straight.	E. Indies.	1798.	7. 9. S. ȝ.	DC. leg. mem. t. 53.	
2	semialátä. DC.	(ro.) winged.	Nepaul.	1805.	7. 8. S. ȝ.	Roxb. corom. 3. t. 248.	
3	congéstä. DC.	(re.) crowded spiked.	E. Indies.	1802.	7. 9. S. ȝ.	—	3. t. 249.
4	nána. DC.	(re.) dwarf.	—	1804.	8. 9. S. ȝ.		
5	lineáta. DC.	(re.) branch-spiked.	—	1793.	7. 8. S. ȝ.	Burm. ind. t. 53. f. 1.	
6	angustifólia. DC.	(re.) narrow-leaved.	—	1812.	7. 9. S. ȝ.		
7	procumbens. DC.	(re.) procumbent.	—	1816.	8. S. ȝ.		
8	prostráta. DC.	(re.) prostrate.	China.	1820.	7. 9. S. ȝ.		
9	strobilífera. DC.	(wh.) Beech-leaved.	E. Indies.	1787.	— S. ȝ.		
	<i>Hedy' sarum strobilíferum.</i> L. <i>Zórnia strobilífera.</i> P.S.						

Subtribus III. ALHAGEÆ. DC. prod. 2. p. 352.

ALH'AGI. DC.	ALH'AGI.	Diadelphia Decandria.			
1	mauròrum. DC.	(re.) Levant.	Levant.	1714.	7. 8. G. ȝ.
					Tourn. cor. 54. t. 489.
	<i>Hedy' sarum Alhagi.</i> L. <i>mannifera.</i> Dv. <i>Mánnia hebráica.</i> D.D.				
2	camelòrum. DC.	(re.) Caspian.	Caucasus.	1824.	— H. ȝ.
ALSICA'RPU'S. DC.	ALSICA'RPU'S.	Diadelphia Decandria.			
1	bupleurifòlius. DC.(li.)	Bupleurum-ld.	E. Indies.	1793.	7. 8. S. ȝ.
	<i>Hedy' sarum bupleurifòlium.</i> Roxb. corom. 2. t. 194.				Bot. mag. 1722.

2 <i>vaginális</i> . DC.	(<i>pu.</i>) sheathed.	E. Indies.	1790.	7. 8.	S. ⊙.	Burm. zeyl. t. 49. f. 1.
3 <i>nummularifólius</i> . DC.	(<i>pu.</i>) moneywort.	—	1777.	7. 9.	S. ⊙.	Petiv. gaz. t. 26 f. 1.
4 <i>monilífer</i> . DC.	(<i>pu.</i>) neck-lace.	—	1824.	—	S. Ȑ.	Burm. ind. t. 52. f. 3.
5 <i>glumáceus</i> . DC.	(<i>vi.</i>) rugged-podded.	Arabia.	—	—	G. Ȑ.	
6 <i>styracifólius</i> . DC.	(<i>pu.</i>) Storax-leaved.	E. Indies.	1796.	—	S. Ȕ.	
<i>Hedy'sarum styracifólium</i> , L.						

Tribus IV. VICIEÆ. DC. prodr. 2. p. 353.

CICER. DC.	CHICK-PEA.	Diadelphia Decandria. L.				
1 <i>arietinum</i> . DC.	common.	S. Europe.	1548.	7. 8.	H. ⊙.	Bot. mag. 2274.
2 <i>sóngaricum</i> . DC.	large-flowered.	Persia.	1828.	—	H. ⊙.	D.C. leg. mem. t. 54.
FABA. DC.	BEAN.	Diadelphia Decandria.				
ν ulgàris. DC.	(<i>wh.bk.</i>) common.	Egypt.	5. 8.	H. ⊙.	Tourn. inst. t. 222.
α equina.	(<i>wh.bk.</i>) Horse.	—	—	H. ⊙.	
β præcox.	(<i>wh.bk.</i>) Mazagon.	—	—	H. ⊙.	
γ ensiformis.	(<i>wh.bk.</i>) long-pod.	—	—	H. ⊙.	
δ albiflora.	(<i>wh.</i>) white-blossomed.	—	—	H. ⊙.	
ϵ purpurea.	(<i>pu.</i>) purple-flowered.	—	—	H. ⊙.	
ζ macrospérma.	(<i>wh.bk.</i>) broad Windsor.	—	—	H. ⊙.	
η chlorospermá.	(<i>wh.bk.</i>) green Windsor.	—	—	H. ⊙.	
VICIA. DC.	VETCH.	Diadelphia Decandria.				
1 <i>pisiformis</i> . DC.	(<i>st.</i>) Pea-shaped.	S. Europe.	1739.	7. 8.	H. Ȑ.	Jacq. aust. 4. t. 364.
2 <i>amœna</i> . DC.	(<i>pu.</i>) beautiful.	Siberia.	1829.	—	H. Ȑ.	Gmel. sib. 4. n. 11. t. 3?
3 <i>caroliniàna</i> . DC.	(<i>wh.</i>) Carolina.	Carolina.	—	—	H. Ȑ.	
4 <i>dumetòrum</i> . DC.	(<i>vi.</i>) great wood.	France.	1752.	5. 7.	H. Ȑ.	Sturm deuts. 1. fas. 31.
5 dentatâ. F.	(<i>vi.</i>) toothed.	Russia.	1818.	6. 7.	H. Ȑ.	
6 <i>sylvatica</i> . DC.	(<i>wh.vi.</i>) wood.	Britain.	7. 8.	H. Ȑ.	Eng. bot. 2. t. 79.
7 <i>americâna</i> . DC.	(<i>vi.</i>) American.	N. America.	1818.	—	H. Ȑ.	Desf. cor. p. 86. t. 66.
8 <i>variegâta</i> . DC.	(<i>v.</i>) variegated.	Caucasus.	1829.	—	H. Ȑ.	
9 <i>purpurea</i> . DC.	(<i>pu.</i>) purple.	Tauria.	1827.	—	H. Ȑ.	
10 <i>alpestrís</i> . DC.	(<i>pu.</i>) alpine.	—	1828.	—	H. Ȑ.	
11 <i>cassubica</i> . DC.	(<i>vi.</i>) Cassubian.	Germany.	1711.	6. 7.	H. Ȑ.	Sturm deuts. 1. fas. 31.
12 abbreviâta. DC.	(<i>vi.</i>) abbreviated.	Caucasus.	1818.	—	H. Ȑ.	
13 <i>cinérea</i> . DC.	(<i>pa.bl.</i>) ash-coloured.	Tauria.	1829.	—	H. Ȑ.	
14 Crâcca. DC.	(<i>vi.</i>) tufted.	Britain.	7. 9.	H. Ȑ.	Eng. bot. 17. t. 1168.
15 Bivônea. DC.	(<i>ro.</i>) Bivoni's.	Sicily.	1828.	—	H. Ȑ.	
16 Gerârdi. DC.	(<i>vi.</i>) Gerard's.	S. Europe.	1711.	6. 7.	H. Ȑ.	Jacq. aust. t. 229.
17 pseudocrâcca. DC.	(<i>vi.</i>) slender-leaved.	Italy.	1824.	7. 9.	H. ⊙.	
<i>tenuifòlia</i> . Tenore. non Roth.						
18 <i>microphylla</i> . DC.	(<i>li.</i>) small-leaved.	Archipelago.	1828.	6. 8.	H. Ȑ.	
19 <i>polyphylla</i> . DC.	(<i>li.</i>) many-leaved.	Algiers.	1823.	—	H. Ȑ.	
20 <i>tenuifòlia</i> . DC.	(<i>vi.</i>) slender-leaved.	Germany.	1799.	6. 7.	H. Ȑ.	Sturm deuts. 1. fas. 31.
21 <i>consentîna</i> . DC.	(<i>st.</i>) straw-coloured.	Naples.	1824.	—	H. Ȑ.	
22 <i>capénisia</i> . DC.	(<i>pu.</i>) Cape.	C. B. S.	1802.	6. 8.	G. Ȑ.	
23 <i>canéscens</i> . DC.	(<i>bl.</i>) hoary.	Libanus.	1800.	—	H. ⊙.	Labil. dec. syr. 1. t. 7.
24 <i>ochroleúca</i> . DC.	(<i>st.</i>) sulphur-color'd.	Italy.	1824.	—	H. Ȑ.	
25 punctatâ. DC.	(<i>pu.</i>) spot-leaved.	Switzerland.	1826.	—	H. ⊙.	
26 onobrychioïdes. DC.	(<i>pu.</i>) Saintfoin-like.	Europe.	1789.	—	H. ⊙.	Sturm deuts. 1. fas. 31.
27 atropurpurea. DC.	(<i>d.pu.</i>) dark-purple.	Algiers.	1815.	—	H. ⊙.	Bot. reg. t. 871.
28 benghalénsis. DC.	(<i>sc.</i>) Bengal.	E. Indies.	1792.	7. 9.	H. ⊙.	Herm. lugd. t. 625.
29 perénnisi. DC.	(<i>pu.</i>) perennial.	S. Europe.	1820.	6. 8.	H. Ȑ.	
30 longifòlia. DC.	(<i>st.</i>) long-leaved.	Syria.	1824.	—	H. Ȑ.	
31 argénteâ. DC.	(<i>ro.</i>) silvery.	Pyrenees.	1829.	—	H. Ȑ.	
32 villòsa. DC.	(<i>pu.</i>) villous.	Germany.	1815.	—	H. ⊙.	Sturm deuts. 1. fas. 31.
33 dispérma. DC.	(<i>li.</i>) two-seeded.	S. France.	1826.	—	H. ⊙.	
34 bienniâ. DC.	(<i>vi.</i>) biennial.	Siberia.	1753.	7. 9.	H. ♂.	Gmel. sib. 4. t. 2.
35 Nissoliâna. DC.	(<i>re.</i>) red-flowered.	Levant.	1773.	6. 7.	H. ⊙.	
36 pellùcida. DC.	(<i>pu.</i>) pellucid.	C. B. S.	1812.	6. 8.	G. Ȑ.	Jacq. schœnb. 2. t. 222.
37 biflòra. DC.	(<i>bl.</i>) two-flower'd.	Algiers.	1801.	—	H. ⊙.	Desf. atl. 2. t. 197.
38 calcarâta. DC.	(<i>pu.</i>) spurred.	—	1790.	7. 9.	H. ⊙.	
<i>monantha</i> . W.						
39 syriaca. DC.	(<i>vi.</i>) Syrian.	Syria.	1822.	—	H. ⊙.	
40 pimpinelloïdes. DC.	(<i>pu.</i>) Pimpinella-like.	Rome.	—	6. 9.	H. ⊙.	

41	sativa. DC.	(pu.) common.	Britain.	5. 7.	H. ⊙.	Eng. bot. 5. t. 334.
	α nemorális. P.S.	(pu.) wood.	—	—	—	H. ⊙.	Sturm deuts. 1. fas. 31.
	β segetális. DC.	(pu.) hedge.	—	—	—	H. ⊙.	Eng. bot. 5. t. 334.
42	leucospermá. L.en.(wh)	white-seeded.	—	—	—	H. ⊙.	Sturm deuts. 1. fas. 31.
43	angustifólia. W.en.(pu.)	narrow-leaved.	Europe.	—	—	H. ⊙.	
44	megalospérma. W.en.(pu.)	large-seeded.	Tauria.	1798.	6. 8.	H. ⊙.	
45	incisa. DC.	(pu.) cut-leaved.	—	1829.	—	H. ⊙.	
46	rèpens. DC.	(pu.) creeping.	Archipelago.	1828.	—	H. 4.	
47	canadén sis. DC.	(li.) Canadian.	Canada.	—	—	H. ⊙.	
48	globosa. DC.	(bl.) globular.	—	1804.	—	H. ⊙.	
49	cordát a. DC.	(pu.) heart-leaved.	Germany.	1828.	—	H. ⊙.	Sturm deuts. 1. fas. 32.
50	pilosa. DC.	(pu.) pilose.	Tauria.	1820.	—	H. ⊙.	
51	peregrina. DC.	(pu.) broad-podded.	S. Europe.	1779.	7. 8.	H. ⊙.	Sturm deuts. 1. fas. 32.
52	Michauxii. DC.	(wh.) Michaux's.	—	1803.	—	H. ⊙.	
53	amphicárpa. DC.	(pu.) subterraneous.	S. France.	1815.	—	H. ⊙.	Ger. m. an. 6.v.3.p.344.
54	lathróides. DC.	(pu.) spring.	Britain.	4. 6.	H. ⊙.	Eng. bot. 1. t. 30.
55	laevigáta. DC.	(br.) smooth-podded.	England.	7. 9.	H. 4.	— 7. t. 483.
56	dúbia. DC.	(pu.) doubtful.	—	1827.	—	H. 4.	
57	lútea. DC.	(ye.) yellow-flowered.	Britain.	6. 8.	H. 4.	Eng. bot. 7. t. 481.
58	hírta. P.S.	(wh.) hairy.	France.	1820.	—	H. 4.	
59	hy'brid a. DC.	(st.) hairy-flower'd.	England.	—	—	H. 4.	Eng. bot. 7. t. 482.
60	spuria. DC.	(st.) spurious.	Sicily.	1826.	—	H. 4.	
61	grandiflóra. DC.	(ye.) large-flowered.	Switzerland.	1823.	—	H. 3.	Scop. fl. carn. 2. t. 42.
62	sordida. DC.	(ye.) large-yellow.	Hungary.	1802.	—	H. ⊙.	W. et K. hung. 2. t. 134.
63	Biebersteini. DC.	(ye.) Bieberstein's.	Poland.	1823.	—	H. 4.	
64	tricolor. DC.	(ye.pu.) three-coloured.	Naples.	1827.	—	H. ⊙.	Seb. et Maur. pr. rom. t. 4.
65	sepium. DC.	(pu.) bush.	Britain.	5. 6.	H. 4.	Eng. bot. 22. t. 1515.
66	pannónica. DC.	(wh.) Hungarian.	Hungary.	1658.	6. 7.	H. ⊙.	Jacq. aust. 1. t. 34.
67	pyrenaica. DC.	(pu.) Pyrenean.	Pyrenees.	1828.	—	H. ⊙.	DC. ic. rar. 1. t. 33.
68	striáta. M.B.	(std.) striped.	Tauria.	1824.	—	H. ⊙.	
69	truncátula. DC.	(st.) truncate-leav'd.	Caucasus.	1828.	6. 8.	H. ⊙.	Roth. abhand. 10. t. 2.
70	arbonén sis. DC.(d.pu.)	broad-leaved.	France.	1596.	—	H. ⊙.	Sturm fl. germ. 1.fas. 32.
71	serratifólia. W.	(pu.) saw-leaved.	Hungary.	1723.	—	H. ⊙.	Roth. abhand. 10. t. 1.
72	platycárgos. DC.	(pu.) flat-podded.	Germany.	—	—	H. ⊙.	
73	bicolor. DC.	(v.) two-coloured.	—	1818.	—	H. 4.	
ER'VUM. DC.		TARE. Diadelphia Decandria. L.					
1	Léns. DC.	(pa.) Lentil.	France.	1548.	5. 7.	H. ⊙.	Sturm fl. germ. 1.fas. 32.
	Cicer Léns. w. Léns esculénta. Mœnch.						
2	nigricans. DC.	(pa.) dark-seeded.	Tauria.	1823.	6. 7.	H. ⊙.	
	lentoidés. T.P.						
3	lenticula. DC.	(bl.) one-flowered.	Carinthia.	1824.	6. 8.	H. ⊙.	Sturm fl. germ. 1.fas. 32.
	soloniense. Schrank pl. rar. t. 48. uniflòrum. Seb. rom. pl. fas. 2. t. 4.						
4	hirsútum. DC. (pu.wh.)	hairy.	Britain.	6. 7.	H. ⊙.	Eng. bot. 14. t. 970.
5	dispérnum. s.s.	(li.) two-seeded.	E. Indies.	1824.	6. 8.	H. ⊙.	
6	camelòrum. s.s.	(li.) broad-podded.	—	—	—	H. ⊙.	
7	péndulum. s.s.	(li.) pendulous-podd'd.	E. Indies.	1828.	—	H. ⊙.	Sturm fl. germ. 1.fas. 32.
8	Ervília. DC.	(pu.) officinal.	S. Europe.	1596.	6. 8.	H. ⊙.	
	Ervília sativa. L. en. Vicia Ervília. w.						
9	monánthos. DC.	(li.) single-flowered.	—	1790.	7. 8.	H. ⊙.	Sturm fl. ge. 1.fas. 32, ic.
	Vicia articulata. w. Láthyrus monánthos. W.en.						
10	agrientinum. DC.	(li.) Sicilian.	Sicily.	1827.	—	H. ⊙.	
11	tetraspérnum. DC.	(li.) four-seeded.	Britain.	6. 8.	H. ⊙.	Eng. bot. 17. t. 1223.
12	grácile. DC.	(li.) slender.	S. Europe.	1823.	—	H. ⊙.	Lois. fl. gal. t. 12.
	tenússimum. P.S. longifolium T.N. Vicia laxiflora. Brot. phyt. t. 54.						
13	pubéscens. DC.	(li.) pubescent.	S. Europe.	1828.	6. 8.	H. ⊙.	
14	Loiseleurii. DC.	(li.) Loiseleur's.	Tauria.	—	—	H. ⊙.	
15	diphy'lhum. DC.	(li.) two-leaved.	—	—	—	H. ⊙.	
PI'SUM. DC.		PEA. Diadelphia Decandria. L.					
1	sativum. DC.	(wh.) cultivated.	S. Europe.	5. 10.	H. ⊙.	Lam. ill. t. 633.
	α saccharátum. DC.(wh.) sugar.			—	H. ⊙.	Tourn. inst. t. 215.
	β macrocárpon. DC.(wh.) marrowfat.			—	H. ⊙.	
	γ víride.	(wh.) green marrowfat.		—	H. ⊙.	
	δ rugósrum.	(wh.) Knight's marrowfat.		—	H. ⊙.	
	ϵ chlorospermum.	(wh.) green rouncticval.		—	H. ⊙.	
	ζ cæruléscens.	(wh.) blue Prussian.		—	H. ⊙.	
	η prolificum.	(wh.) prolific.		—	H. ⊙.	
	θ præcox.	(wh.) early Charlton.		—	H. ⊙.	
	ι uniflòrum.	(wh.) early frame.		—	H. ⊙.	

<i>κ</i> <i>hamile.</i>	(<i>wh.</i>) dwarf.	S. Europe.	H. ⓧ.	Tabern. icon. t. 495.
<i>λ</i> <i>umbellatum.</i> DC. (<i>pi.</i>)	rose or crown.	H. ⓧ.	
<i>μ</i> <i>quadratum.</i> DC. (<i>pu.</i>)	grey or swine's.	H. ⓧ.	
2 <i>elatum.</i> DC.	(<i>pa.pu.</i>) upright.	Iberia.	1827.	6. 10. H. ⓧ.	Moris. his. 2. s. 2 t. 1.f.4.
3 <i>arvense.</i> DC.	(<i>pu.</i>) field.	Europe.	6. 9. H. ⓧ.	Eng. bot. 15. t. 1046.
4 <i>maritimum.</i> DC. (<i>pu.</i>)	sea.	England.	7. 8. H. ⓧ.	
5 <i>americanum.</i> DC. (<i>pu.</i>)	American.	America.	H. ⓧ.	
6 <i>Jomárdi.</i> DC.	(<i>wh.</i>) angular-stalk'd.	Egypt.	1818.	6. 9. H. ⓧ.	
7 <i>thebaicum.</i> DC.	(<i>wh.</i>) short-peduncled.	H. ⓧ.	
LA'THYRUS. DC.	LA'THYRUS.	Diadelphia Decandria.	L.		
1 <i>sylvestris.</i> DC.	(<i>pu.</i>) wood.	Britain.	7. 8. H. ⓧ.	Eng. bot. v. 12. t. 805.
2 <i>intermedius.</i> DC.	(<i>ro.</i>) intermediate.	N. Europe.	1818.	7. 8. H. ⓧ.	Flor. dan. t. 785.
3 <i>magellanicus.</i> DC. (<i>pu.</i>)	Lord Anson's Pea.	Cape Horn.	1744.	—	
4 <i>latifolius.</i> DC.	(<i>ro.</i>) everlasting Pea.	England.	7. 9. H. ⓧ.	Eng. bot. v. 16. t. 1108.
β <i>albiflorus.</i>	(<i>wh.</i>) white-flowered.	H. ⓧ.	Bieb. cent. 1. t. 22.
5 <i>rotundifolius.</i> DC.	(<i>ro.</i>) round-leaved.	Tauria.	1822.	—	Eng. bot. v. 10. t. 670.
6 <i>pratensis.</i> DC.	(<i>ye.</i>) meadow.	Britain.	6. 8. H. ⓧ.	Bot. mag. 111.
7 <i>tuberosus.</i> DC.	(<i>ro.</i>) tuberous-rooted.	Europe.	1596.	7. 8. H. ⓧ.	
8 <i>rōseus.</i> DC.	(<i>ro.</i>) rose-coloured.	Iberia.	1828.	—	Linn. fil. dec. t. 20.
9 <i>pisiformis.</i> DC.	(<i>std.</i>) Siberian.	Siberia.	1759.	6. 8. H. ⓧ.	Swt. br. fl. gar. t. 194.
10 <i>mutabilis.</i> B.F.G.	(<i>std.</i>) changeable.	1825.	—	
11 <i>venosus.</i> DC.	(<i>pu.</i>) veined.	N. America.	1823.	7. 8. H. ⓧ.	
12 <i>californicus.</i> B.R.	(<i>pu.</i>) Californian.	California.	1827.	6. 10. H. ⓧ.	Bot. reg. 1144.
13 <i>palustris.</i> DC.	(<i>pu.</i>) marsh.	Britain.	7. 8. H. ⓧ.	Eng. bot. v. 3. t. 169.
14 <i>myrtifolius.</i> DC.	(<i>pu.</i>) Myrtle-leaved.	N. America.	1823.	—	
15 <i>polymorphus.</i> DC. (<i>pu.</i>)	variable.	Missouri.	H. ⓧ.	
16 <i>heterophyllum.</i> DC. (<i>li.</i>)	various-leaved.	Europe.	1731.	7. 9. H. ⓧ.	J. Bauh. his. 2. p. 304.f.1.
17 <i>stipulaceus.</i> DC.	(<i>pu.</i>) large-stipuled.	N. America.	1827.	—	
18 <i>Apháca.</i> DC.	(<i>st.</i>) yellow vetchling.	England.	6. 7. H. ⓧ.	Eng. bot. v. 17. t. 1167.
19 <i>Nissolia.</i> DC.	(<i>cr.</i>) crimson.	5. 6. H. ⓧ.	— v. 2. t. 112.
20 <i>inconspicuus.</i> DC.	(<i>pa.</i>) small-flowered.	Levant.	1739.	7. 8. H. ⓧ.	DC. ic. rar. 1. t. 32.
21 <i>sphaericus.</i> DC.	(<i>sc.</i>) scarlet.	S. Europe.	1800.	6. 7. H. ⓧ.	
<i>coccineus.</i> P.S. <i>axillaris.</i> Lam.	<i>inconspicuus.</i> Jacq. vind. t. 86.	<i>Orobous coccineus.</i> Lodd. b. cab. t. 883.			
22 <i>micranthus.</i> DC.	(<i>pu.</i>) small-blossom'd.	S. France.	6. 7. H. ⓧ.	Buxb. cent. 3. t. 42. f.2.
23 <i>angulatus.</i> DC.	(<i>pu.</i>) angular-seeded.	S. Europe.	1683.	—	
24 <i>longepedunculatus.</i> DC. (<i>pu.</i>) long-stalked.	1824.	6. 9. H. ⓧ.	
25 <i>leptophyllum.</i> DC. (<i>pu.</i>)	slender-leaved.	Caucasus.	1828.	—	
26 <i>setifolius.</i> DC.	(<i>ro.</i>) bristle-leaved.	S. Europe.	1739.	6. 7. H. ⓧ.	J. Bauh. 2. p. 308. ic.
27 <i>amphicarpos.</i> DC.	(<i>li.</i>) Earth Pea.	Levant.	1680.	6. 9. H. ⓧ.	Swt. br. fl. gar. 3. t. 236.
28 <i>sativus.</i> DC.	(<i>bl.wh.</i>) chickling vetch.	S. Europe.	1640.	6. 8. H. ⓧ.	Jacq. f. eclog. t. 116.
α <i>cæruleus.</i>	(<i>bl.</i>) blue-flowered.	H. ⓧ.	Bot. mag. 115.
β <i>álbis.</i>	(<i>wh.</i>) white-flowered.	H. ⓧ.	
29 <i>Cícera.</i> DC.	(<i>pu.</i>) flat-podded.	1633.	—	Jacq. f. eclog. t. 115.
30 <i>annuus.</i> DC.	(<i>ye.</i>) yellow annual.	1621.	—	Buxb. cent. 3. t. 42. f.1.
31 <i>hirsutus.</i> DC.	(<i>pu.</i>) rough-podded.	England.	7. 8. H. ⓧ.	Eng. bot. v. 18. t. 1255.
32 <i>hirtus.</i> DC.	(<i>pu.</i>) hairy.	1824.	7. 9. H. ⓧ.	
33 <i>odoratus.</i> DC.	(<i>va.</i>) Sweet Pea.	Sicily.	1700.	6. 9. H. ⓧ.	Bot. mag. t. 60.
α <i>purpureus.</i> DC.	(<i>pu.</i>) dark-purple.	H. ⓧ.	— t. 60.
β <i>ruber.</i>	(<i>re.</i>) red-flowered.	H. ⓧ.	
γ <i>rōseus.</i> DC.	(<i>ro.</i>) painted lady.	Ceylon.	H. ⓧ.	
δ <i>álbis.</i>	(<i>wh.</i>) white-flowered.	H. ⓧ.	
34 <i>grandiflorus.</i> DC.	(<i>pu.</i>) large-flowered.	Italy.	1814.	—	Bot. mag. t. 1938.
35 <i>tingitanus.</i> DC.	(<i>pu.</i>) Tangier Pea.	Barbary.	1680.	—	— t. 100.
36 <i>bithynicus.</i> DC.	(<i>vi.</i>) tumid-podded.	S. Europe.	1816.	6. 8. H. ⓧ.	All. ped. 1. t. 26. f. 2.
<i>tumidus.</i> w. <i>túrgidus.</i> Lam.	<i>Vicia bithynica.</i> L.				
37 <i>incurvus.</i> DC.	(<i>pu.</i>) curve-podded.	Siberia.	1802.	7. 8. H. ⓧ.	Roth abh. 13. t. 4.
38 <i>alatus.</i> DC.	(<i>pu.</i>) wing-stalked.	Italy.	1829.	—	
39 <i>Cly'menum.</i> DC.	(<i>re.bl.</i>) various-flower'd.	Levant.	1713.	6. 7. H. ⓧ.	Pluk. alm. t. 114. f. 6.
40 <i>articulatus.</i> DC.	(<i>ro.wh.</i>) joint-podded.	S. Europe.	1640.	7. 9. H. ⓧ.	Bot. mag. 258.
41 <i>auriculatus.</i> Bert.	(<i>pu.</i>) eared.	Italy.	1820.	6. 8. H. ⓧ.	
42 <i>spurius.</i> DC.	(<i>pu.</i>) bastard.	1815.	—	
43 <i>tenuifolius.</i> DC.	(<i>bl.</i>) slender-leaved.	Algiers.	1824.	6. 7. H. ⓧ.	
44 <i>Ochrus.</i> DC.	(<i>ye.</i>) pale-yellow.	S. Europe.	1633.	—	Schk. hand. 2. t. 280.
<i>Ochrus pallida.</i> P.S. <i>Pisum Ochrus.</i> L.					
45 <i>purpureus.</i> DC.	(<i>pu.</i>) purple-flower'd.	Levant.	1826.	6. 8. H. ⓧ.	Desf. cor. p. 81. t. 61.
46 <i>cornutus.</i> DC.	(<i>pu.</i>) horned.	1824.	—	
47 <i>aphyllus.</i>	(<i>pu.</i>) leafless.	Smyrna.	1828.	7. 8. H. ⓧ.	
PLATYSTYLIS. B.F.G.	PLATYSTYLIS.	Diadelphia Decandria.			
1 <i>cyanea.</i> B.F.G.	(<i>bl.</i>) blue-flowered.	Iberia.	1826.	7. 8. H. ⓧ.	Swt. br. fl. gar. 3. t. 239.

2 sessilifolius. B.F.G. (pu.) sessile-leaved.	Levant.	1824.	6. 8. H. 4.	Buxb. cent. 2. t. 38.
O'robus sessilifolius. Bot. Mag. 2796. digitatus. M.B.				
O'ROBUS. DC.	BITTER VETCH. Diadelphia Decandria. L.			
1 hirsutus. DC. (li.) hairy.	Levant.	1818.	5. 6. H. 4.	Bot. mag. t. 2345.
2 formosus. DC. (pu.) handsome.	Caucasus.	1824.	— H. 4.	Bot. mag. t. 2098.
3 lathyroides. DC. (bl.) upright.	Siberia.	1758.	6. 7. H. 4.	— t. 521.
4 vénus. DC. (pu.) spring.	Europe.	1629.	3. 4. H. 4.	Gmel. fl. sib. 4. t. 5.
5 venosus. DC. (bl.) veined.	Siberia.	1818.	5. 6. H. 4.	W. et K. hung. 2. t. 126.
6 alpétris. DC. (pu.) alpine.	Hungary.	1818.	— H. 4.	Ten. fl. nap. t. 68.
7 multiflorus. DC. (li.) many-flowered.	Italy.	1824.	5. 7. H. 4.	W. et K. hung. 3. t. 242.
8 variegatus. DC. (pu.) variegated.	Naples.	1818.	5. 6. H. 4.	Vicia oroboides. W.K.
9 vicíofides. DC. (pu.) Vetch-like.	Hungary.	—	6. 8. H. 4.	
10 luteus. DC. (ye.) yellow.	Siberia.	1759.	6. 7. H. 4.	Lodd. bot. cab. 783.
11 laevigatus. DC. (ye.) smooth.	Hungary.	1818.	— H. 4.	W. et K. hung. 3. t. 243.
12 aurantius. DC. (ye.) orange-color'd.	Iberia.	1824.	6. 8. H. 4.	
13 sylvaticus. DC. (std.) wood.	Britain.	6. 7. H. 4.	Eng. bot. v. 8. t. 518.
14 ochroleucus. DC. (st.) straw-colour'd.	Hungary.	1816.	— H. 4.	W. et K. hung. 2. t. 118.
15 niger. DC. (pu.) black-rooted.	Europe.	1596.	5. 7. H. 4.	Bot. mag. t. 2261.
16 erectus. DC. (pu.) upright.	1820.	6. 7. H. 4.	Eng. bot. v. 17. t. 1153.
17 tuberosus. DC. (pu.) tuberous-rooted.	Britain.	5. 6. H. 4.	Pluk. alm. t. 210. f. 2.
18 pyrenaicus. L. (pu.) Pyrenean.	Pyrenees.	1699.	— H. 4.	
19 tenuifolius. Roth. (pu.) slender-leaved.	Germany.	1818.	5. 6. H. 4.	Bot. mag. t. 675.
20 varius. DC. (v.) parti-coloured.	Italy.	1759.	— H. 4.	Gmel. sib. 4. p. 14. t. 5.
21 angustifolius. DC. (v.) narrow-leaved.	Siberia.	1766.	— H. 4.	
22 canescens. DC. (bl. wh.) canescent.	S. Europe.	1823.	— H. 4.	
23 pallescens. M.B. (wh.) pale-flowered.	Tauria.	1824.	— H. 4.	
24 albus. DC. (wh.) white-flowered.	Austria.	1794.	— H. 4.	Swt. br. fl. gar. 1. t. 22.
pannonicus. Jacq. aust. 1. t. 39.				
25 atropurpureus. DC. (d. pu.) dark-purple.	Algiers.	1826.	6. 7. H. 4.	Desf. atl. 2. t. 196.
26 Fischeri. B.F.G. (d. pu.) Fischer's.	Siberia?	1827.	— H. 4.	Swt. br. fl. gar. 3. t. 289.
27 saxatilis. DC. (pu.) rock.	S. France.	1823.	6. 8. H. 4.	Venten. cels. t. 94.
28 longifolius. N. (li.) long-leaved.	Missouri.	1827.	— H. 4.	

Tribus V. PHASEOLEÆ. DC. prodr. 2. p. 381.

'ABRUS. DC.	WILD LIQUORICE. Monadelphia Enneandria.			
precatorius. DC. (ro.) common.	India.	1680.	3. 5. S. li. — Rheed. mal. 3. t. 39.	
α erythrosperma. (ro.) red-seeded.	—	—	— S. li. — Rumph. amb. 5. t. 32.	
β melanospérma. (ro.) black-seeded.	—	—	— S. li. —	
SWEETIA. DC.	SWEETIA. Diadelphia Decandria.			
1 longifolia. DC. (pu.) long-leaved.	S. America.	1824.	6. 7. S. li. — Jacq. ic. 3. t. 572.	
Gâlega longifolia. Jacq. Tephrosia longifolia. P.S.				
2 filiformis. DC. (pu.) slender.	—	1826.	— S. li. — Jacq. ic. 3. t. 573.	
3 lignosa. DC. (pu.) woody.	Trinidad.	1820. S. li. —	
Glycine lignosa. P.S.				
MACRA'NTHUS. DC.	MACRA'NTHUS. Diadelphia Decandria.			
cochinchinensis. DC. (wh.) eatable-podd'd. Cochinchina.	1826.	6. 9. H. 4.		
WESTONIA. S.S.	WESTONIA. Monadelphia Decandria.			
trifoliata. s.s. (st. re.) trifoliate.	1820.	6. 8. S. 4.	
Rôthia trifoliata. DC. Glycine humifusa. W. en.				
TERAMNUS. DC.	TERAMNUS. Monadelphia Pentandria.			
1 volubilis. DC. (re.) twining.	Jamaica.	1822.	7. 8. S. li. —	
2 uncinatus. DC. (re.) hooked.	—	—	— S. li. — Plum. ed. Burm. t. 221.	
AMPHICARPÆA. DC.	AMPHICARPÆA. Diadelphia Decandria.			
1 monoica. DC. (pa.) pale-flowered.	N. America.	1781.	8. 10. H. 4.	Rœm. arch. 1. p. 3. t. 2.
2 comosa. (bl.) tufted.	—	1812.	7. 9. H. 4.	
Glycine comosa. Ph.				
3 sarmentosa. DC. (pa.) sarmentose.	—	1805.	— H. 4.	Schk. bot. an. 12. t. 2.
Glycine sarmentosa. Ph. monoica. Schkuhr.				
KENNEDYA. DC.	KENNEDYA. Diadelphia Decandria.			
1 rubicunda. DC. (cr.) dingy-flowered.	N. S. W.	1788.	3. 8. G. li. — Bot. mag. 208.	
2 prostrata. DC. (sc.) scarlet one-fl'd.	—	1790.	— G. li. — t. 270.	
Glycine coccinea. B.M.				
3 coccinea. DC. (sc.) many-flowered.	N. Holland.	1803.	5. 8. G. li. — Swt. fl. aust. t. 23.	
4 Comptoniana. DC. (pu.) Comptonian.	—	—	3. 6. G. li. — Bot. reg. 298.	
Glycine Comptoniana. B.R.				

5 heterophylla. <i>(pu.)</i> various-leaved.	H. Holland.	1826.	G. \natural . \sim .
6 inophylla. <i>(pu.)</i> few-leaved.	1825.	G. \natural . \sim .
7 dilatata. <i>(pu.)</i> dilated-leaved.	G. \natural . \sim .
8 monophylla. <i>(pu.)</i> simple-leaved.	N. S. W.	1790.	3. 8.	G. \natural . \sim . Vent. malm. t. 106.
<i>Glycine bimaculata</i> . Bot. mag. t. 263.				
9 ovata. DC. <i>(pu.)</i> oval-leaved.	1820.	—	G. \natural . \sim . Bot. mag. 2169.
<i>cordata</i> . Bot. reg. 944.				
RHYNCH'OSIA. DC. RHYNCH'OSIA. Diadelphia Decandria.				
1 reniformis. DC. <i>(ye.)</i> kidney-leaved.	Carolina.	1806.	7. 8.	F. γ .
2 difformis. DC. <i>(ye.)</i> downy climbing.	1732.	6. 9.	F. γ . \sim . Dill. elth. t. 26. f. 29.
<i>Glycine tomentosa</i> β <i>volubilis</i> . M.				
3 erecta. DC. <i>(br.)</i> upright.	—	F. γ .
4 caribæa. DC. <i>(ye.)</i> trailing.	W. Indies.	1742.	9. 10.	S. \natural . \sim . Bot. reg. t. 275.
<i>Glycine caribæa</i> . B.R.				
5 mollis. DC. <i>(ye.)</i> soft-leaved.	Guinea.	1823.	6. 7.	S. \natural . \sim . Hook. ex. flor. 201.
<i>Glycine mollis</i> . H.E.F.				
6 minima. DC. <i>(ye.)</i> slender-stalked.	W. Indies.	—	—	S. \natural . \sim . Jacq. obs. 1. t. 22.
7 volubilis. DC. <i>(ye.)</i> black-podded.	China.	1826.	—	S. \natural . \sim .
8 phaseoloides. DC. <i>(ye.)</i> Abrus-seeded.	W. Indies.	1820.	—	S. \natural . \sim . Bot. mag. t. 2284.
9 precatoria. DC. <i>(ye.)</i> Abrus-like.	S. America.	—	—	S. \natural . \sim .
10 macrophylla. DC. <i>(ye.)</i> large-leaved.	Cuba.	1826.	—	S. \natural . \sim .
11 reticulata. DC. <i>(ye.)</i> netted.	Jamaica.	1779.	7. 9.	S. \natural . \sim .
12 rhombifolia. DC. <i>(ye.)</i> rhomb-leaved.	E. Indies.	1815.	5. 8.	S. \natural . \sim .
13 medicaginea. DC. <i>(ye.)</i> Medick-like.	Ceylon.	1826.	—	S. \natural . \sim . Burm. zeyl. 188. t. 88.
14 suavolens. DC. <i>(ye.)</i> sweet-scented.	E. Indies.	1816.	—	S. \natural .
15 viscosa. DC. <i>(ye.)</i> viscous.	1823.	6. 7.	S. \natural . \sim .
16 viscosa. DC. <i>(ye.)</i> clammy.	1824.	—	S. \natural . \sim .
17 Fridericiana. DC. <i>(ye.)</i> blunt-leaved.	Manilla.	1826.	—	S. \natural . \sim .
18 scarabœoides. DC. <i>(ye.)</i> rough-stemmed.	E. Indies.	—	—	S. \odot . \sim . Pluk. alm. t. 52. f. 3.
19 biflora. DC. <i>(ye.)</i> two-flowered.	E. Indies.	1827.	6. 8.	S. \odot . \sim .
20 angustifolia. DC. <i>(ye.)</i> narrow-leaved.	C. B. S.	1795.	—	G. \natural . \sim . Jacq. schoen. 2. t. 231.
21 Tötta. DC. <i>(ye.)</i> smooth-stemmed.	1819.	—	G. \natural . \sim .
FAG'ELIA. DC. FAG'ELIA. Diadelphia Decandria.				
bituminosa. DC. <i>(ye.)</i> clammy.	C. B. S.	1774.	4. 9.	G. \natural . \sim . Bot. reg. t. 261.
<i>Glycine bituminosa</i> . B.R.				
WISTERIA. DC. WIST'ERIA. Diadelphia Decandria.				
1 frutescens. DC. <i>(bl.)</i> handsome.	N. America.	1724.	6. 9.	H. \natural . \sim . Bot. mag. t. 2103.
<i>speciosa</i> . N. <i>Glycine frutescens</i> . B.M. <i>Apios frutescens</i> . Ph.				
2 chinensis. DC. <i>(p.bl.)</i> Chinese.	China.	1818.	5. 7.	H. \natural . \sim . Swt. br. fl. gar. 3. t. 211.
<i>Glycine chinensis</i> . Bot. mag. 2083. <i>sinensis</i> . Bot. reg. 650.				
APIOS. DC. 'APIOS. Diadelphia Decandria.				
tuberosa. DC. <i>(ro.)</i> tuberous-rooted.	N. America.	1640.	8. 9.	H. γ . \sim . Bot. mag. t. 1198.
<i>Glycine Apios</i> . B.M.				
AMPH'ODUS. B.R. AMPH'ODUS. Diadelphia Decandria.				
ovatus. B.R. <i>(pu.)</i> oval-leaved.	Trinidad.	1824.	4. 6.	S. \natural . \sim . Bot. reg. t. 1101.
PHASEOLUS. DC. KIDNEY BEAN. Diadelphia Decandria. L.				
1 Caracalla. DC. <i>(pu.)</i> Snail-flower.	E. Indies.	1690.	8. 9.	S. γ . \sim . Bot. reg. t. 341.
2 tuberosus. DC. <i>(ye.)</i> tuberous-rooted.	China.	1825.	—	S. γ . \sim .
3 perennis. DC. <i>(pu.)</i> perennial.	Carolina.	1823.	7. 9.	F. γ . \sim .
4 macrostachyus. DC. <i>(pu.)</i> long-spiked.	N. America.	1824.	—	H. γ . \sim .
5 alatus. DC. <i>(pu.)</i> winged.	Carolina.	1732.	7. 8.	H. \odot . Dill. elth. t. 235. f. 303.
6 multiflorus. DC. <i>(sc.)</i> scarlet-runner.	S. America.	1633.	6. 10.	H. γ . \sim . Schk. hand. 2. t. 199. a.
<i>a coccineus</i> . DC. <i>(sc.)</i> scarlet-flowered.	—	—	H. γ . \sim . Moris. his. 52. t. 5. f. 4.
<i>β albiflorus</i> . DC. <i>(wh.)</i> white-flowered.	—	—	H. γ . \sim .
7 vulgaris. DC. <i>(wh.li.)</i> common.	E. Indies.	1597.	5. 9.	H. \odot . \sim . Lobel. ic. 2. p. 59. ic.
8 compræssus. DC. <i>(wh.)</i> white-runner.	—	6. 9.	H. \odot . \sim . Savi. mem. 3. t. 10. f. 20.
9 oblóngus. DC. <i>(li.)</i> oblong-seeded.	—	—	H. \odot . \sim . — 3. p. 17. t. 10. f. 14.
10 saponaceus. DC. <i>(wh.)</i> spotted-white.	—	—	H. \odot . \sim . — 3. p. 19. t. 10. f. 15.
11 tumidus. DC. <i>(wh.)</i> thick-seeded.	—	—	H. \odot . \sim . — 3. p. 19. t. 10. f. 16.
12 haematócarpus. DC. <i>(li.)</i> red-podded.	—	—	H. \odot . \sim . — 3. p. 20. t. 10. f. 17.
13 sphæricus. DC. <i>(li.)</i> roundish-seeded.	—	—	H. \odot . \sim . — 3. p. 20. t. 10. f. 18.
14 gonospérmus. DC. <i>(wh.)</i> angular-seeded.	—	—	H. \odot . \sim . — 3. p. 21. t. 10. f. 19.
15 derásus. DC. <i>(gr.wh.)</i> narrow-podded.	Brazil.	1823.	—	H. \odot . \sim . Schk. h. mon. 1. t. 89.
16 lunátus. DC. <i>(gr.)</i> cimeter-podded.	E. Indies.	1779.	—	S. \odot . \sim . Hout. syst. 8. t. 63. f. 1.
17 inam'onus. DC. <i>(gr.wh.)</i> various-colour'd.	Africa.	1794.	7. 8.	H. \odot . \sim . Jacq. vind. t. 66.
18 Xuarésii. DC. <i>(gr.wh.)</i> pointed-leaved.	1823.	6. 9.	H. \odot . \sim .
19 adenánthus. DC. <i>(wh.bl.)</i> painted-flower'd.	Guiana.	1821.	—	S. \odot . \sim .
20 macrocarpus. DC. <i>(wh.)</i> long-podded.	—	—	H. \odot . \sim .
21 calcaratus. H.B. <i>(li.)</i> spurred.	E. Indies.	1820.	7. 8.	S. δ . \sim .

LEGUMINOSÆ.

22 dolichoïdes. H.B. (<i>pu.</i>)	Dolichos-like.	E. Indies.	1820.	7. 8. S. 4. ↗.
23 mínimus. H.B. (<i>li.</i>)	small.	China.	1823.	— H. ♂. ↗.
24 angustifòlius. H.B. (<i>li.</i>)	narrow-leaved.	—	1818.	6. 8. H. ♂. ↗.
25 microspérmos.DC. (<i>pu.</i>)	small-seeded.	Cuba.	1821.	6. 9. S. ♂.
26 heterophyllus.DC. (<i>re.</i>)	different-leaved.	Mexico.	1818.	7. 8. H. ♂. ↗.
27 acomitifòlius. DC. (<i>ye.</i>)	Aconite-leaved.	E. Indies.	1731.	6. 8. S. ♂. ↗. Jacq. obs. 3. t. 52.
28 sublobatus. H.B. (<i>li.</i>)	lobed-leaved.	—	1820.	— S. ♂. ↗.
29 subtrilobus. L.en. (<i>ye.</i>)	acute-lobed.	Brazil.	1818.	— S. ♂. ↗.
30 diversifòlius. DC. (<i>pu.</i>)	various-leaved.	N. America.	1806.	6. 8. H. ♂.
31 trilobus. DC. (<i>pu.</i>)	three-lobed.	E. Indies.	1777.	7. 8. S. ♂. Burm. ind. t. 50. f. 1.
32 angulosus. DC. (<i>bh.</i>)	angular-stalked.	S. America.	1818.	7. 8. S. ♂. ↗.
33 farinósus. DC. (<i>re.</i>)	mealy.	E. Indies.	1759.	S. 4. ↗. Niss. act. par. 1730. t. 42.
34 stipularis. DC. (<i>ye.</i>)	large-stipuled.	Peru.	1805.	— H. ♂.
35 mésoleucus.L.en. (<i>wh.</i>)	Brazilian.	Brazil.	1818.	— H. ♂. ↗.
36 hélylus. DC. (<i>ro.</i>)	pale red.	Carolina.	1732.	— H. ♂. ↗. Dill. elth. t. 233. f. 300.
37 vexillatus. DC. (<i>bh.</i>)	sweet-scented.	—	—	— H. ♂. ↗. Jacq. vind. 2. t. 102.
38 Múngo. DC. (<i>ye.</i>)	small-fruited.	E. Indies.	1790.	6. 7. H. ♂.
39 hirtus. w.	hairy.	C. B. S.	1818.	7. 8. H. ♂. ↗.
40 Hernandésii. DC. (<i>ye.</i>)	Hernandez.	Mexico.	1824.	— H. ♂. Hern. mex. 887. ic.
41 chrysanthos. DC. (<i>ye.</i>)	golden-flowered.	—	—	— H. ♂.
42 radiatus. DC. (<i>bh.</i>)	rayed.	China.	1732.	6. 7. H. ♂. Dill. elth. t. 235. f. 305.
43 scáber. DC. (<i>st.</i>)	rough-podded.	—	—	6. 9. H. ♂. Moris. ox. s. 2. t. 5. f. 8.
44 lathyróides. DC. (<i>re.</i>)	Lathyrus-like.	Jamaica.	1786.	7. 8. S. ♂. Sloan. hist. 1. t. 116. f. 1.
45 semerectus DC. (<i>re.</i>)	dark-red.	S. America.	1732.	— S. ♂. ↗. Bot. reg. t. 743.
46 violaceus. DC. (<i>vi. wh.</i>)	violet-flowered.	Africa?	—	6. 9. H. ♂.
47 Máx. DC. (<i>re.</i>)	hairy-podded.	E. Indies.	1758.	6. 7. S. ♂. Rumph. amb. 5. t. 140.
48 gláber. H.B. (<i>li.</i>)	smooth.	Mauritius.	1823.	7. 8. S. ♂.
49 torosus. H.B. (<i>li.</i>)	twisted-podded.	Nepaul.	—	— H. ♂.
S'JOJA. DC.	Soy.	Diadelphia Decandria.		
hispida. DC. (<i>vi.</i>)	hispid.	E. Indies.	1790.	7. 8. S. ♂. Jacq. ic. 1. t. 145.
Dólchos Soja. Jacq.	Sōja japonica. Savi.			
β pálida. DC. (<i>ye.</i>)	pale-flowered.	—	—	S. ♂.
DO'LICHOS. DC.	Do'LICHOS.	Diadelphia Decandria.	L.	
1 lignósus. DC. (<i>ro.</i>)	woody.	E. Indies.	1776.	7. 8. S. ♂. ↗. Smith spicil. t. 21.
β? falciſormis.	(<i>ro.</i>) falcate-podded.	C. B. S.	—	G. ♂. ↗. Bot. mag. t. 382.
2 Jacquiní. DC. (<i>wh.</i>)	Jacquin's.	W. Indies.	1822.	7. 9. S. ♂. ↗.
3 spurius. DC. (<i>pu.</i>)	spurious.	Guiana.	1826.	— S. ♂. ↗.
4 tuberosus. DC. (<i>li.</i>)	tuberous-rooted.	Martinica.	—	— S. ♂. ↗. Plum. ed. Burm. t. 220.
5 articulatus. DC. (<i>pu.</i>)	jointed.	St. Domingo.	1828.	— S. ♂. ↗.
6 hirsutus. DC. (<i>pu.</i>)	hirsute.	China.	1802.	6. 7. G. ♂. ↗. Kämpf. ic. t. 41.
7 pubescens. DC. (<i>st.</i>)	pubescent.	S. America.	1826.	7. 9. G. ♂. ↗.
8 pilosus. DC. (<i>li.</i>)	hairy-podded.	E. Indies.	1790.	7. 8. S. ♂. ↗.
9 tetraspermus. DC. (<i>li.</i>)	four-seeded.	—	1816.	5. 8. S. ♂. ↗.
10 capensis. DC. (<i>ye.</i>)	Cape.	C. B. S.	1818.	— G. 4. ↗.
11 falcatus. DC. (<i>li.</i>)	falcate-podded.	E. Indies.	1820.	7. 8. S. 4. ↗.
12 heterophyllus.DC. (<i>pu.</i>)	various-leaved.	Canaries.	1827.	6. 9. H. ♂. ↗.
13 lütens. DC. (<i>ye.</i>)	yellow-flowered.	Jamaica.	1812.	7. 8. S. 4. ↗.
14 biflorus. DC. (<i>li.</i>)	two-flowered.	E. Indies.	1776.	— S. ♂. Pluk. alm. t. 213. f. 4.
15 Catiáng. DC. (<i>pu.</i>)	small-podded.	—	1793.	— S. ♂.
16 monachalis. DC. (<i>pu.</i>)	hybrid.	Portugal.	1826.	— H. ♂.
17 siénensis. DC. (<i>fl.</i>)	Chinese.	China.	1776.	— H. ♂. Rumph. amb. 5. t. 134.
18 niloticus. DC. (<i>st.</i>)	Egyptian.	Egypt.	1828.	— H. ♂. Delil. fl. eg. t. 38. f. 1.
19 Lùbia. DC. (<i>bl. wh.</i>)	cultivated.	—	—	— H. ♂.
20 vexillatus. DC. (<i>st.</i>)	hispid.	S. America.	—	— H. ♂. ↗. Dill. elt. 2. f. 302.
21 lobatus. DC. (<i>li.</i>)	lobe-leaved.	C. B. S.	1818.	6. 7. G. 4. ↗. Hout. syst. 8. t. 64. f. 1.
22 virosus. H.B. (<i>li.</i>)	virose.	E. Indies.	—	5. 8. S. ♂. ↗. Rheed. mal. 8. t. 45.
23 gangéticus. H.B. (<i>pu.</i>)	Ganges.	Bengal.	1824.	6. 7. S. 4. ↗.
24 glutinosus. H.B. (<i>pu.</i>)	glutinous.	E. Indies.	1820.	— S. ♂. ↗.
25 unguiculatus. DC. (<i>li.</i>)	Bird's-foot.	W. Indies.	1780.	— S. ♂. ↗. Jacq. vind. 1. t. 23.
26 tranquebáricus.DC. (<i>vi.</i>)	Tranquebar.	E. Indies.	1801.	6. 7. S. ♂. ↗. Jacq. vind. 3. t. 70.
27 melanophthalmus.DC. (<i>vi.</i>)	black-circled.	—	1822.	7. 9. H. ♂. ↗.
28 sesquipedális. DC. (<i>wh.</i>)	long-podded.	W. Indies.	1781.	8. 9. S. ♂. ↗. Jacq. vind. 1. t. 67.
29 Brunélli. DC. (<i>wh.</i>)	branching.	—	1826.	6. 9. S. ♂.
30 sphæropérmos.DC. (<i>st.</i>)	round-seeded.	S. America.	1823.	7. 8. S. ♂. Sloan. jam. 1. t. 117.
31 reticulatus. DC. (<i>li.</i>)	netted-leaved.	N. S. W.	1781.	6. 7. G. ♂. ↗.
32 phaseoloïdes. H.B. (<i>li.</i>)	Kidney Bean-like.	China.	1823.	7. 8. G. 4. ↗.
33 frutescens. D.P. (<i>pu.</i>)	frutescent.	Nepaul.	1824.	— G. ♂.
34 incanus. Wal.	(<i>pu.</i>) hoary.	E. Indies.	1825.	— S. ♂.

VI'GNA. DC.	VI'GNA.	Diadelphia Decandria.						
1 glâbra. DC. (ye.) smooth.	Amerika.	1805.	7. 8.	H. ⊙.~.	Jacq. vind.	1. t. 90.		
Dôlichos lutèolus. Jacq.								
2 villòsa. DC. (ye.) villous.	Chile.	1826.	—	H. ⊙.~.				
LA'BLAB. DC.	LA'BLAB.	Diadelphia Decandria.						
1 vulgàris. DC. (pu.) common.	Egypt.	1694.	6. 7.	S. ⊙.	Bot. mag.	t. 896.		
Dôlichos Láblab. B.M.								
2 purpùreus. (pu.) purple.	E. Indies.	1790.	5. 10.	S. ȳ.~.	Exot. bot.	2. t. 74.		
Dôlichos purpùreus. Bot. reg. t. 830.								
3 benghalénus. (wh.) Bengal.	Bengal.	1820.	7. 8.	S. ȳ.~.	Jacq. vind.	2. t. 124.		
4 nankínicus. DC. (wh.) Chinese.	China.	1821.	—	H. ⊙.~.	Savi. dis.	1821. f. 8. d-g.		
5 leucocárpus. DC. (wh.) white-podded.	E. Indies.	—	—	S. ⊙.~.		f. 9. a-d.		
6 cultrátus. DC. (wh.) sharp-edged.	Japan.	1818.	6. 7.	H. ⊙.~.	Banks ic.	kämpf. t. 25.		
7 microcárpus. DC. (pu.) short-podded.	E. Indies.	1824.	—	S. ⊙.~.	Rumph. am.	5. t. 141. f. 1.		
8 perennans. DC. (wh.) perennial.	China.	1823.	7. 8.	H. ⊙.~.		5. t. 137.		
Dôlichos ábus. w.								
PACHYRH'IZUS. DC.	PACHYRH'IZUS.	Diadelphia Decandria.						
1 angulatus. DC. (vi.) angular-leaved.	E. Indies.	1781.	7. 8.	S. ȳ.~.	Rumph. amb.	5. t. 132.		
Dôlichos bulbosus. L.								
2 trilobus. DC. (pu.) three-lobed.	China.	1825.	6. 8.	S. ȳ.~.				
PAROCH'ETUS. D.P.	PAROCH'ETUS.	Diadelphia Decandria.						
1 communis. D.P. (ii.) common.	Nepaul.	1827.	6. 8.	G. ȳ.				
2 mäjor. D.P. (ii.) larger.	—	—	—	G. ȳ.				
DIOCLE'A. DC.	DIOCLE'A.	Diadelphia Decandria.						
Jacquiniana. DC. (re.) Jacquin's.	Martinica.	1826.	6. 9.	S. ȳ.~.	Jacq. amer.	t. 123.		
Dôlichos rüber. Jacq.								
PSOPHOCA'RPU'S. DC.	PSOPHOCA'RPU'S.	Diadelphia Decandria.						
tetragonólobus. DC. (bl.) square-podded.	Mauritius.	1816.	9. 11.	S. ⊙.	Rumph. amb.	5. t. 133.		
Dôlichos tetragonólobus. L.								
CANAVALIA. DC.	CANAVALIA.	Monadelphia Decandria.						
1 obtusifolia. DC. (pu.) blunt-leaved.	E. Indies.	1818.	7. 8.	S. ȳ.~.	Rheed. mal.	8. t. 43.		
Dôlichos rotundifolius. Vahl.								
β emarginata. DC. (pu.) emarginate.	—	—	—	S. ȳ.~.	Jacq. schoen.	2. t. 221.		
2 gladiata. DC. (bh.) sabre-podded.	—	1801.	6. 7.	S. ȳ.~.	Jacq. ic.	3. t. 560.		
Dôlichos gladiatus. w. Malóccchia gladiata. Savi.								
3 ensiformis. DC. (pu.) cimeter-podded.	E. Indies.	1778.	7. 8.	S. ⊙.	Jacq. ic.	3. t. 559.		
4 rôsea. DC. (ro.) rose-colour'd.	Jamaica.	1812.	—	S. ȳ.				
5 bonariensis. B.R. (pu.) Buenos Ayres.	BuenosAyres.	1826.	6. 10.	G. ȳ.~.	Bot. reg.	t. 1199.		
MUCUNA. DC.	Cow-ITCH.	Diadelphia Decandria.						
1 urens. DC. (st.) broad-podded.	W. Indies.	1691.	6. 7.	S. ȳ.~.	Jacq. amer.	t. 182. f. 84.		
Dôlichos urens. L. Stizolobium urens. P.S.								
2 altissima. DC. (ii.) tallest.	Martinica.	1779.	6. 8.	S. ȳ.~.	Jacq. amer.	t. 182. f. 85.		
3 pruriens. DC. (vi.) common.	India.	1680.	—	S. ȳ.~.	Rheed. mal.	8. t. 85.		
Dôlichos pruriens. L. Stizolobium pruriens. P.S. Carpopôgon pruriens. H.B.								
4 gigantea. DC. (pa.) gigantic.	E. Indies.	1815.	7. 9.	S. ȳ.~.	Rheed. mal.	8. t. 36.		
5 elliptica. DC. (ii.) elliptic-leaved.	Peru.	1828.	—	G. ȳ.~.				
6 mitis. DC. (ii.) oblique-leaved.	—	1823.	7. 9.	G. ȳ.~.				
Negrétia mitis. R.P.								
7 comosa. DC. (wh.) tufted.	Guiana.	1827.	—	S. ȳ.~.				
8 monosperma. DC. (ii.) one-seeded.	E. Indies.	1818.	—	S. ȳ.~.				
Carpopôgon monospermum. H.B.								
9 atropurpurea. DC. (d.pu.) dark-purple.	—	1820.	—	S. ȳ.~.				
10 anguinea. (pu.) snake-like.	—	1822.	—	S. ȳ.~.				
Carpopôgon anguineum. Roxb.								
11 nivea. DC. (wh.) snowy-white.	—	1820.	7. 9.	S. ⊙.~.				
12 imbricata. DC. (pu.) imbricated.	—	1815.	—	S. ȳ.~.				
13 bracteata. DC. (pu.) bracteate.	—	1822.	—	S. ȳ.~.				
14 capitata. (pu.) headed.	—	—	8. 11.	S. ⊙.~.				
Carpopôgon capitatum. H.B.								
PHYLLÓ'OBÍUM. DC.	PHYLLÓ'OBÍUM.	Diadelphia Decandria.	S.S.					
1 chinense. DC. (wh.) Chinese.	China.	1818.	6. 8.	G. ȳ.				
2 zanzibaréne. Boj. Zanzibar.	Zanzibar.	1825.	—	S. ȳ.				
CAJ'ANUS. DC.	PIGEON PEA.	Diadelphia Decandria.						
1 bicolor. DC. (ye.pu.) two-coloured.	E. Indies.	1687.	7. 8.	S. ȳ.	Jacq. vind.	2. t. 119.		
Cy'tisus pseudo-cájan. Jacq.								
2 flavus. DC. (ye.) yellow-flowered.	—	—	—	S. ȳ.	Jacq. obs.	1. t. 1.		
Cy'tisus Cájan. L.								
LUP'INUS. DC.	LUPINE.	Monadelphia Decandria.						
1 ábus. DC. (wh.) white.	Levant.	1596.	7. 8.	H. ⊙.	Blackw. herb.	t 282.		

LEGUMINOSÆ.

2 Térnis. DC.	(wh.bl.) Egyptian.	Egypt.	1802.	6. 7. H. ⊖.
3 várus. DC.	(bl.) variable.	S. Europe.	1596.	7. 9. H. ⊖.
4 hirsútus. DC.	(bl.) blue-flowered.	——	1629.	—— H. ⊖.
5 bracteoláris. DC.	(bl.) bracteolate.	Monte VIDEO.	1828.	—— H. ⊖.
6 pilósus. DC.	(ro.) rose.	S. Europe.	1710.	—— H. ⊖.
7 angustifólius. DC.	(bl.) narrow-leaved.	——	1686.	—— H. ⊖.
8 linífolius. DC.	(bl.) Flax-leaved.	——	1790.	—— H. ⊖.
9 lúteus. DC.	(ye.) yellow.	S. Europe.	1596.	—— H. ⊖.
β sulphureus.	(st.) straw-coloured.	——	——	—— H. ⊖.
10 microcárpus. DC.	(bl.) small-podded.	Chile.	1822.	4. 6. H. ⊖.
11 micránthus. B.R.	(pu.) small-flowered.	Columbia.	1827.	5. 7. H. ⊖.
12 áridus. B.R.	(bl.pu.) arid.	——	6. 9. H. ⊖.	
13 bicolor. B.R.	(bl.wh.) two-coloured.	——	5. 10. H. Ȑ.	
14 lépidus. B.R.	(bl.li.) lively.	——	7. 10. H. Ȑ.	
15 laxíflorus. B.R.	(bl.pi.) lax-flowered.	——	8. 10. H. Ȑ.	
16 littorális. B.R.	(pu.bl.) shore.	——	6. 10. H. Ȑ.	
17 ornátus. B.R.	(bl.) adorned.	——	5. 11. H. Ȑ.	
18 plumósus. B.R.	(bl.) feathered.	——	——	H. Ȑ.
19 Sabini. D.	(bl.) Sabine's.	——	6. 10.	H. Ȑ.
20 mexicanus. DC.	(bl.) Mexican.	Mexico.	1819.	7. 9. F. Ȑ.
21 perénnis. DC.	(bl.) smooth perennial.	N. America.	1658.	5. 8. F. Ȑ.
β albiflórus.	(wh.) white-flowered.	——	1817.	—— F. Ȑ.
22 polyphyllus. B.R.	(bl.) many-leaved.	Columbia.	1827.	—— H. Ȑ.
23 leucophyllus. B.R.	(pu.) white-leaved.	——	6. 8.	H. Ȑ.
24 nootkáensis. DC.	(bl.) hairy perennial.	Nootka sound.	1794.	—— H. Ȑ.
β variabilis.	(va.) variable coloured.	——	——	H. Ȑ.
25 variegátus. F.	(v.) variegated.	Silka.	1828.	—— H. Ȑ.
26 argyréus. DC.	(bl.) silvery-leaved.	——	——	F. Ȑ.
27 arbústus. B.R.	(pu.) half-shrubby.	California.	1827.	6. 9. F. Ȑ.
28 canaliculátus. B.F.G.	(bl.) channel-id.	Buenos Ayres.	1828.	6. 9. H. Ȑ.
29 versícolor. B.F.G.	(v.) various-colour'd.	Mexico.	1828.	—— H. Ȑ.
30 tomentósus. DC.	(va.) woolly.	Peru.	1826.	6. 10. H. Ȑ.
31 arbóreus. DC.	(ye.) tree.	S. America.	1793.	7. 9. H. Ȑ.
32 mutábilis. B.F.G.	(wh.y.) changeable.	Bogota.	1825.	—— F. Ȑ.
33 villósus. DC.	(pu.) simple-leaved.	Carolina.	1787.	—— F. Ȑ.
CYLÍSTA. DC.	CYLÍSTA. Diadelphia Decandria.			
1 scariósa. DC.	(bh.) rough.	E. Indies.	1806.	4. 6. S. Ȑ.
2 tomentósa. DC.	(ye.) tomentose.	——	1818.	—— S. Ȑ.
3 albiflóra. DC.	(wh.) white-flowered.	Mauritius.	4. 5. S. Ȑ.
4 villósa. DC.	(wh.) hairy.	C. B. S.	1776.	4. 6. G. Ȑ.
Dólichos hirtus. A.R.				
ERYTHR'INA. DC.	CORAL-TREE. Diadelphia Decandria. L.			
1 herbácea. DC.	(sc.) herbaceous.	Carolina.	1724.	6. 9. G. Ȑ.
2 cárnea. DC.	(fl.) flesh-coloured.	Vera Cruz.	1733.	5. 6. G. Ȑ.
β pubescens.	(fl.) pubescent-leaved.	——	——	G. Ȑ.
3 Corallodéndron. DC.	(sc.) common.	W. Indies.	1690.	—— S. Ȑ.
4 enneándra. DC.	(sc.) nine-anthered.	——	S. Ȑ.
velutina. Jacq. non W.				
5 macrophylla. DC.	(sc.) large-leaved.	——	1816.	—— S. Ȑ.
6 mítis. DC.	(sc.) smooth-stem'd.	Caracas.	1818.	6. 8. S. Ȑ.
7 poánthes. DC.	(sc.) crowded-flower'd.	——	1826.	4. 6. S. Ȑ.
8 secundiflóra. DC.	(sc.) side-flowering.	Brazil.	1818.	—— S. Ȑ.
9 speciosa. DC.	(sc.) large-flowered.	W. Indies.	1805.	5. 10. S. Ȑ.
10 umbrósia. DC.	(sc.) deltoid-leaved.	Caracas.	1824. S. Ȑ.
11 Humeána. s.s.	(sc.) Sir A. Hume's.	C. B. S.	1818.	5. 6. S. Ȑ.
cáffra. Bot. reg. t. 736. non Thunb.				
12 cáffra. Th.	(pu.) Cape-tree.	——	1816. S. Ȑ.
13 fúlgens. DC.	(sc.) fulgent.	E. Indies.	1801.	5. 10. S. Ȑ.
14 arboréscens. DC.	(cr.) arborescent.	Nepaul.	1818. S. Ȑ.
15 índica. DC.	(sc.) Indian.	E. Indies.	1814.	5. 6. S. Ȑ.
16 pícta. DC.	(cr.) prickly-leaved.	——	1696.	5. 10. S. Ȑ.
17 aculeatíssima. Df.	(sc.) very prickly.	——	1816.	—— S. Ȑ.
18 velutína. DC.	(sc.) velvety.	Caracas.	1824. S. Ȑ.
19 fúscia. DC.	(br.re.) red-brown.	China.	1800. S. Ȑ.
20 laurífolia. Jacq.	(sc.) Laurel-leaved.	Brazil.	5. 10. H. Ȑ.
21 crista-gálli. DC.	(sc.) Cockscomb.	——	1771.	6. 10. H. Ȑ.
22 nervosa. DC.	(sc.) nerved-leaved.	——	1824.	—— S. Ȑ.
23 glauca. DC.	(sn.) glaucous.	Caracas.	—— S. Ȑ.
24 piscidioídes. DC.	(sc.) Piscidia-leaved.	E. Indies.	1800. S. Ȑ.

J. Bauh. his. 2. p. 289. ic.

Knor. del. 2. t. L. 7.
Roth. abhand. 14. t. 5.
Bot. mag. t. 140.Bot. mag. t. 2413.
Bot. reg. t. 1251.
—— t. 1242.
—— t. 1109.
—— t. 1149.
—— t. 1140.
—— t. 1198.
—— t. 1216.
—— t. 1217.Bot. reg. t. 457.
Bot. mag. t. 202.
Bot. reg. t. 1096.
—— t. 1124.
—— t. 1311.
—— t. 2136.Bot. reg. t. 1230.
Swt. br. fl. gar. t. 283.
—— ser. 2. t. 12.
Swt. br. fl. gar. t. 261.
Bot. mag. t. 682.
Swt. br. fl. gar. t. 130.
Pursh. fl. amer. 2. t. 21.Roxb. corom. 1. t. 92.
—— t. 221.
Bot. mag. t. 1859.
Andr. rep. t. 446.Bot. mag. t. 877.
Trew. ehr. 2. t. 8.
Bot. reg. t. 389.
Comm. h. ams. 1. t. 108.
Jacq. schoen. 4. t. 466.
Jacq. schoen. 2. t. 216.
Bot. reg. t. 1246.
Linn. trans. 14. t. 12.
Andr. rep. t. 443.
Bot. mag. t. 2431.Roxb. corom. 3. t. 219.
Rheed. mal. 6. t. 7.
Rumph. amb. 2. t. 77.
Rumph. amb. 2. t. 78.
Swt. br. fl. gar. t. 142.
—— t. 2124.

25 incàna. w.	(sc.) hoary.	1820.	6. 8.	S. h.
26 lithosperma. W.H. stony-seeded.	N. Holland.	1824.	S. h.
27 arbórea. D.L. tree.	Trinidad.	1822.	S. h.
28 strícta. H.B.	(sc.) upright.	E. Indies.	1816.	4. 6.	S. h.
29 ovalifolia. H.B. oval-leaved.	S. h.
30 suberòsa. H.B. cork-barked.	1818.	S. h.
31 resupinata. H.B. resupinate.	1822.	S. h.
32 abyssinica. DC. Abyssinian.	Africa.	1820.	S. h.
RUDO'LPHIA. DC.	RUDO'LPHIA.	Diadelphia Decandria.			
1 volùbilis. DC.	(sc.) twining.	Portorico.	1820.	S. h. — Vahl ecl. 3. t. 30.
2 portoricensis. DC.	(sc.) Portorico.	1820.	S. h. —
<i>Erythrina portoricensis</i> . Df.					
3 dùbia. DC.	(sc.) doubtful.	S. America.	1815.	S. h. — H. et B. n. gen. 6. t. 591.
B'UTEA. DC.	B'UTEA.	Diadelphia Decandria.			
1 frondosa. DC.	(sc.) downy-branch'd.	E. Indies.	1796.	S. h. Roxb. corom. 1. t. 21.
2 supérba. DC.	(sc.) smooth-branch'd.	1798.	S. h. v. 1. t. 22.
3 parviflora. DC.	(sc.) small-flowered.	Coromandel.	1820.	S. h.

Tribus VI. DALBERGIEÆ. DC. prod. 2. p. 415.

DE'RRIS. DC.	DE'RRIS.	Monadelphia Decandria.			
trifoliata. DC.	(wh.) three-leaved.	China.	1826.	4. 6.	G. h. —
PONG'AMIA. DC.	PONG'AMIA.	Monadelphia Decandria.			
1 glabra. DC.	(wh.) smooth-leaved.	E. Indies.	1699.	6. 8.	S. h. Vent. malm. t. 28.
2 ellíptica.	(wh.) elliptic-leaved.	1820.	—	S. h. —
<i>Galedupa elliptica</i> . H.B.					
3 chinensis. DC.	(ye.) Chinese.	China.	1827.	—	S. h.
4 uliginosa. DC. marsh.	E. Indies.	—	S. h. —
5 piscidia. Piscidia-leaved.	1816.	S. h.
<i>Galedupa piscidia</i> . H.B.					
6 marginata. H.B. margined.	1824.	S. h. —
DALBERGIA. DC.	DALBERGIA.	Mono-Diadelphia Decandria.			
1 ougeinensis. DC.	(wh.) Hindoo.	E. Indies.	1828.	S. h.
2 latifolia. DC.	(wh.) broad-leaved.	1811.	S. h. Roxb. corom. 2. t. 113.
3 Sissoo. DC.	(wh.) large.	1823.	S. h.
4 rubiginosa. DC.	(wh.) rusty.	1811.	6. 8.	S. h. — Roxb. corom. 2. t. 115.
5 robusta. DC.	(wh.) robust.	—	S. h.
6 scandens. DC.	(ro.) scandent.	—	5. 7.	S. h. — Roxb. corom. 2. t. 192.
7 volubilis. DC.	(wh.) twining.	1820.	—	S. h. — v. 2. t. 191.
8 frondosa. DC.	(wh.) frondose.	1818.	S. h.
9 paniculata. DC.	(wh.) paniced.	1811.	S. h. Roxb. corom. 2. t. 114.
10 emarginata. H.B.	(wh.) emarginate.	1823.	S. h.
11 zeylanica. H.B.	(wh.) Ceylon.	Ceylon.	1816.	S. h.
12 marginata. H.B.	(wh.) margined.	E. Indies.	1820.	5. 7.	S. h. —
13 tamarindifolia. H.B.	(wh.) Tamarind-leav'd.	—	S. h. —
14 stipulacea. H.B.	(wh.) stipuled.	—	6. 8.	S. h.
15 rimosa. H.B.	(wh.) rimose.	1824.	S. h.
16 alata. H.B.	(wh.) winged.	1823.	S. h.
17 Crówei. DC.	(wh.) Silhet.	—	S. h.
PTEROCA'RPU'S. DC.	PTEROCA'RPU'S.	Mono-Diadelphia Decandria.			
1 Dráco. DC.	(ye.) officinal.	S. America.	1818.	S. h. — Jacq. amer. t. 183. f. 92.
<i>officinalis</i> . Jacq. hemiptera. Gært.					
2 suberosus. DC.	(ye.) corky.	Guiana.	1826.	S. h. Aubl. guian. 2. t. 299.
3 Marsipium. DC.	(st.) leathery-leaved.	E. Indies.	1811.	S. h. Roxb. corom. 2. t. 116.
4 dalbergioïdes. DC.	(ye.) Andaman red-wood.	1820.	S. h.
5 indicus. DC.	(ye.) Indian.	1818.	S. h.
6 santalinus. DC.	(ye.) red sandal wood.	1800.	S. h.
7 santalinoïdes. DC.	(ye.) African.	Sierra Leon.	1826.	S. h.
8 flávus. DC.	(ye.) yellow-barked.	China.	—	S. h. Rumph. amb. 3. t. 117.
9 scandens. DC.	(vi.) climbing.	Caracas.	1822.	6. 8.	S. h. —
<i>Amerinnum scandens</i> . W.					
DREPANOCA'RPU'S. DREPANOCA'RPU'S.	Monadelphia Decandria.				
lunatus. DC.	(wh.) crescent-podded.	S. America.	1792.	S. h. Plu.ed.Burm.t.201.f.2.
Pterocárpus lunatus. L.	Pterocárpus ápterus.	Gært. fruct. t. 156.			

- | | |
|--------------------------------------|--|
| ECASTAPHYLLUM. DC. | ECASTAPHYLLUM. Diadelphia Octo-Decandria. |
| Brównei. DC. | (wh.) Browne's. W. Indies. 1733. 6. 8. S. $\frac{1}{2}$. Brown.jam.t.32.f.1. |
| <i>Pterocarpus Ecastaphyllum.</i> L. | |
| AMERIMNUM. DC. | AMERIMNUM. Monadelphia Decandria. |
| Brównei. DC. | (wh.) Browne's. W. Indies. 1793. 6. 8. S. $\frac{1}{2}$. Brown.jam.288.t.32.f.3 |
| BRYA. DC. | BR'YA. Monadelphia Decandria. |
| E'benus. DC. | (ye.) Jamaica Ebony. W. Indies. 1713. 7. 8. S. $\frac{1}{2}$. Brown.jam.t.31.f.2. |

Subordo II.—*Tribus VII.* SWARTZIEÆ. DC. prodr. 2. p. 422.

- | | | | | | | |
|------------------------------|---|---------------|-------|-------|---|--------------------------------|
| SWA'RTZIA. DC. | SWA'RTZIA <i>Monadelphia</i> Dec.- <i>Polyandria</i> . | | | | | |
| 1 <i>simplicifolia</i> . DC. | (ye.) simple-leaved. | W. Indies. | 1822. | | S. h. | |
| | | | | | <i>Rittera simplex</i> . Vahl. <i>Possira simplex</i> . Swartz. | |
| 2 <i>grandiflora</i> . DC. | (ye.) great-flowered. | Trinidad. | 1822. | | S. h. | Vahl. pl. am. dec. 1. t. 9. |
| 3 <i>pinnata</i> . DC. | (ye.) wing-leaved. | — | 1820. | | S. h. | Corr. an. mus. 9. t. 24. f. 2. |
| 4 <i>tomentosa</i> . DC. | (ye.) woolly. | Cayenne. | 1826. | | S. h. | Aubl. guian. 2. t. 307. |
| BA'PHIA. DC. | BA'PHIA <i>Decandria</i> <i>Monogynia</i> . | | | | | |
| nítida. DC. | (wh.) glossy. | Sierra Leone. | | 5. 9. | S. h. | Lodd. bot. cab. t. 367. |

Subordo III.—*Tribus VIII. MIMOSÆ*. DC. prodr. 2. p. 424.

- | ENT'ADA. DC. | ENT'ADA. Polygamia Monœcia. | | | | | | |
|---|--|-------|-------|---------|------------------------------|--|--|
| 1 Gigalòbium. DC. (<i>wh.</i>) West Indian. | W. Indies. | 1780. | 4. 6. | S. h.~. | | | |
| 2 Pursætha. DC. (<i>wh.</i>) East Indian. | E. Indies. | — | — | S. h.~. | Rumph. amb. 5. t. 4. | | |
| Parrâna. s.s. <i>Acacia scândens</i> . W. <i>Mimosa scândens</i> . L. | | | | | | | |
| 3 monostachya. DC. (<i>wh.</i>) single-spiked. | — | — | — | S. h.~. | Rheed. mal. 9. t. 77. | | |
| Rheedii. s.s. <i>Mimosa Entada</i> . W. | | | | | | | |
| 4 polystachya. DC. (<i>wh.</i>) many-spiked. | W. Indies. | 1816. | 4. 9. | S. h.~. | DC. leg. mem. t. 61—62. | | |
| MIM'OSA. DC. | SENSITIVE-PLANT. Polygamia Monœcia. L. | | | | | | |
| 1 sensitiva. DC. (<i>ro.</i>) true. | Brazil. | 1648. | 4. 9. | S. h.~. | Botan. regist. t. 25. | | |
| 2 obtusifolia. DC. (<i>ro.</i>) blunt-leaved. | — | 1816. | — | S. h.~. | | | |
| 3 strigosa. DC. (<i>ro.</i>) strigose. | S. America. | 1822. | — | S. h.~. | | | |
| 4 floribunda. DC. (<i>ro.</i>) many-flowered. | Cumana. | 1824. | 6. 7. | S. h.~. | | | |
| 5 viva. DC. (<i>ro.</i>) lively. | Jamaica. | 1739. | 7. 9. | S. y.~. | Sloan. his. 2. t. 182. f. 7. | | |
| 6 polycarpa. DC. (<i>ro.</i>) many-podded. | Peru. | 1816. | — | S. h.~. | Kth. mim. p. 8. t. 3. | | |
| 7 pudica. DC. (<i>ro.</i>) humble-plant. | Brazil. | 1638. | 4. 9. | S. h.~. | Bot. reg. v. 11. t. 941. | | |
| 8 pudibunda. DC. (<i>ro.</i>) fringe-stipuled. | — | 1822. | — | S. h.~. | | | |
| 9 polydactyla. DC. (<i>ro.</i>) many-fingered. | Demerara. | 1824. | — | S. h.~. | Kth. mim. p. 14. t. 5. | | |
| 10 hamata. DC. (<i>ro.</i>) hooked. | E. Indies. | 1820. | 4. 9. | S. h.~. | | | |
| 11 intermedia. DC. (<i>ro.</i>) intermediate. | Caracas. | 1825. | — | S. h.~. | Kth. mim. p. 16. t. 6. | | |
| 12 viscosa. DC. (<i>ro.</i>) viscid. | Brazil. | — | — | S. h.~. | | | |
| 13 canescens. DC. (<i>wh.</i>) canescent. | Guinea. | 1826. | — | S. h.~. | | | |
| 14 hispida. DC. (<i>wh.</i>) hispid. | Caracas. | 1824. | 6. 9. | S. h.~. | | | |
| 15 ciliata. DC. (<i>wh.</i>) ciliated. | Brazil. | — | — | S. h.~. | | | |
| 16 polycantha. DC. (<i>wh.</i>) many-spined. | Africa. | 1828. | — | S. h.~. | Bruce itin. 5. t. 7. | | |
| 17 asperata. DC. (<i>wh.</i>) various spined. | W. Indies. | 1822. | 6. 7. | S. h.~. | DC. leg. mem. t. 63. | | |
| 18 pigra. L. (<i>wh.</i>) straight-spined. | Vera Cruz. | 1733. | — | S. h.~. | Mill. ic. 2. t. 182. f. 3. | | |
| 19 Sicaria. DC. (<i>wh.</i>) subulate-spined. | Brazil. | 1828. | — | S. h.~. | | | |
| 20 rubicaulis. DC. (<i>st.</i>) Bramble-stalk'd. E. Indies. | — | 1799. | 6. 7. | S. h.~. | Roxb. corom. 2. t. 200. | | |
| octandra. R.C. <i>spinosiliqua</i> . Rottler. <i>Rottleri</i> . S.S. | | | | | | | |
| 21 Ceratonia. DC. (<i>wh.</i>) round-leaved. | St. Domingo. | 1800. | | S. h.~. | Plum. ed. Burm. 17. t. 8. | | |
| Acacia Ceratonia. w. | | | | | | | |
| 22 casta. DC. (<i>wh.</i>) chaste. | E. Indies. | 1741. | 6. 8. | S. h.~. | Comm. hort. 1. t. 28. | | |
| 23 angulata. s.s. (<i>wh.</i>) angular-stalked. | Brazil. | 1826. | — | S. h.~. | | | |
| 24 latispinosa. s.s. (<i>ye.</i>) broad-spined. | Madagascar. | 1822. | — | S. h.~. | | | |
| 25 Barclayana. Mr. Barclay's. | — | — | | S. h.~. | | | |
| 26 brasiliensis. s.s. (<i>wh.</i>) Brazilian. | Brazil. | 1824. | 6. 8. | S. h.~. | | | |
| 27 abstergens. s.s. (<i>ro.</i>) spotted-stalked. | E. Indies. | 1818. | — | S. h.~. | | | |
| 28 ferruginea. s.s. (<i>ro.</i>) ferruginous. | — | — | — | S. h.~. | | | |

GAGNEB'INA. DC.	GAGNEB'INA.	Decandria Monogynia.				
1 tamariscina. DC. (st.)	Tamarisk-like.	Mauritius.	1824.	S. h.	DC. leg. mem. t. 64. f. B.	
Acâcia tamariscina. w.	Mimôsa tamariscina. Lam.	Pluk. t. 329. f. 3.				
2 axillâris. DC. (st.)	axillary-spiked.	Mauritius.	1822.	S. h.	DC. leg. mem. t. 64. f. A.	
ERYTHROPHL'EUM. B.	RED-WATER-TREE.	Decandria Monogynia.				
leonénse. G.D.	Sierra Leone.	Sierra Leone. 1822.	S. h.		
PARKIA. B.	PA'RKIA.	Monadelphia Decandria.				
africana. B. (cr.)	African.	Africa. 1822. 6. 8.	S. h.	Beauv. fl. ow. 2. t. 90.		
I'nga biglobôsa. DC.	Mimôsa biglobôsa. Jacq.	amer. t. 179. f. 87.				
INGA. DC.	INGA.	Polygamia Monœcia.				
1 quassiaefolia. DC. (wh.)	Quassia-leaved.	Brazil.	1820.	S. h.		
2 pilosiuscula. DC. (wh.)	glossy-leaved.	Cayenne.	1827.	S. h.	Vahl. ecl. 3. t. 24.	
Mimôsa lúcidâ. Vahl.						
3 setifera. DC. (wh.)	setiferous.	Guiana.	1824.	S. h.		
4 álba. DC. (wh.)	white.	Cayenne.	1804.	7. 8.	S. h.	
5 Feuillei. DC. (wh.)	Feuillee's.	Peru.	1824.	—	S. h.	Feuil. obs. 3. p. 2. t. 19.
6 fastuosa. DC. (re.)	villous-flower'd.	Caracas.	1820.	S. h.	Jacq. fragm. t. 10.
Mimôsa fastuosa. Jacq.						
7 sapindoides. DC. (wh.)	Soap-berry-like.	—	1826.	S. h.	
8 spuria. DC. (wh.)	spurious.	Cumana.	1823.	7. 8.	S. h.	Kth. mim. 39. t. 12.
9 vêra. DC. (wh.)	sweet-fruited.	S. America.	1739.	—	S. h.	Sloan. hist. t. 183. f. 1.
Mimôsa Inga. L.						
10 rhoifolia. DC. (wh.)	Rhus-leaved.	Brazil.	1815.	—	S. h.	
11 Bourgôni. DC. (st.)	Beech-leaved.	W. Indies.	1752.	—	S. h.	Aubl. guian. 2. t. 358.
marginatâ. w. Mimôsa sagifolia. L. non Jacq.						
12 marginata. DC.	margined.	Caracas.	—	—	S. h.	
13 laurina. DC. (wh.)	Laurel-leaved.	—	1768.	3. 5.	S. h.	
14 punctata. DC. (wh.)	white-spotted.	—	1818.	—	S. h.	Jacq. amer. t. 164.
Mimôsa sagifolia. Jacq. non L.						
15 splendens. DC. (wh.)	shining-leaved.	—	1825.	S. h.	
16 nodosa. DC. (wh.)	knobbed.	Ceylon.	1690.	4. 6.	S. h.	Pluk. alm. t. 211. f. 5.
17 microphylla. DC. (wh.)	small-leaved.	Cumana.	1815.	S. h.	
18 dulcis. DC. (wh.)	Sappan fruit.	E. Indies.	1800.	5. 7.	S. h.	Roxb. corom. 1. t. 99.
19 U'nguis-câti. DC. (wh.)	four-leaved.	W. Indies.	1690.	4. 6.	S. h.	Jacq. schœn. 2. t. 34.
20 mellifera. DC. (wh.)	honey-bearing.	Arabia.	1826.	S. h.	
21 ligustrina. DC. (wh.)	Privet-leaved.	Caracas.	—	S. h.	Jacq. frag. t. 32. f. 5.
22 fœtida. DC. (wh.)	strong-smelling.	W. Indies.	1818.	S. h.	Jacq. schœn. 3. t. 390.
23 tergemina. DC. (pu.)	trifoliolate.	Martinica.	—	4. 6.	S. h.	Jacq. amer. t. 177. f. 81.
24 emarginata. DC. (ro.)	emarginate.	Mexico.	1825.	S. h.	Kth. mim. 54. t. 17.
25 latifolia. DC. (ro.)	broad-leaved.	—	1768.	3. 5.	S. h.	Plum. ed. Burn. t. 9.
26 circinalis. DC. (sc.)	spiral-podded.	W. Indies.	1726.	4. 6.	S. h.	Catesb. car. 2. t. 97.
27 purpurea. DC. (pu.)	Soldier-wood.	—	1733.	3. 5.	S. h.	Bot. reg. t. 129.
Mimôsa purpurea. Andr. deposit. t. 372.						
28 xylocarpa. DC. (wh.)	woody-podded.	E. Indies.	1822.	4. 7.	S. h.	Roxb. corom. 1. t. 100.
29 bigemina. DC. (wh.)	two-paired.	—	1823.	—	S. h.	Rheed mal. 6. t. 12.
30 Jirínga. DC. (wh.)	twisted-podded:	—	1828.	S. h.	
31 comosa. DC. (wh.)	tufted.	Jamaica.	1820.	4. 6.	S. h.	
32 cyclocarpa. DC. (wh.)	shell-podded.	Caracas.	1826.	S. h.	Jacq. fragm. t. 34. f. 1.
33 Samân. DC.	linear-podded.	Jamaica.	—	S. h.	— t. 9.
34 corcondiana. DC. (wh.)	Coromandel.	E. Indies.	1818.	S. h.	
35 pulcherrima. Cerv. (pu.)	handsome.	Mexico.	1822.	S. h.	
ANNELE'IA. P.L.	ANNELE'IA.	Monadelphia Polyandria.				
1 grandiflora. (cr.)	large-flowered.	Mexico.	1769.	6. 9.	S. h.	Andr. deposit. t. 592.
Mimôsa grandiflora. A.R.	Acâcia grandiflora. w.	Inga anomala. Kth. mim. t. 22.				
2 Houstoni.	(cr.) Houston's.	Vera Cruz.	1729.	9. 11.	S. h.	Bot. reg. t. 98.
falcifolia. Sal. par. lond. 64.	Acâcia Houstoni. w.	Inga Houstoni. DC.				
SCHRA'NKIA. DC.	SCHRA'NKIA.	Polygamia Monœcia.				
1 aculeata. DC. (ro.)	prickly.	Vera Cruz.	1733.	7. 8.	S. Ȑ.	Mill. ic. 1. t. 182. f. 1.
2 leptocarpa. DC. (ro.)	slender-podd'd.	St. Domingo.	1827.	—	S. Ȑ.	
3 uncinata. DC. (ro.)	hooked.	N. America.	1789.	—	H. Ȑ.	Vent. choix. t. 28.
DARLINGT'ONIA. DC.	DARLINGT'ONIA.	Pentandria Monogynia.				
1 brachylôba. DC. (wh.)	short-podded.	N. America.	1803.	9. 10.	H. Ȑ.	DC. leg. mem. t. 66.
2 glandulosa. DC. (wh.)	glandular.	—	1806.	—	F. h.	Vent. choix. t. 27.
Mimôsa glandulosa. M.						
DESMA'NTHUS. DC.	DESMA'NTHUS.	Polygamia Monœcia.				
1 lacustris. DC. (st.)	marsh.	S. America.	1823.	6. 8.	S. w. ⊖.	H. et B. pl. aeq. 1. t. 16.
2 nâtans. DC. (ye.)	floating.	China.	1800.	7. 9.	S. w. ⊖.	Andr. deposit. t. 629.
3 polypyllus. DC. (wh.)	many-leaved.	—	1826.	—	S. w. ⊖.	
4 triquètrus. DC. (wh.)	triquetrous.	E. Indies.	1823.	—	S. w. ⊖.	

5 plenus. DC.	(ye.) double-yellow.	Vera Cruz.	1733.	7. 9. S.w. ⊕.	Mill. ic. t. 182. f. 2.
6 depresso. DC.	(wh.) depressed.	W. Indies.	1828. S. ℡.	Kth. mim. p. 115. t. 35.
7 diffusus. DC.	(wh.) prostrate.	1731.	7. 9. S. ℡.	Pluk. alm. t. 307. f. 3.
8 virgatus. DC.	(wh.) twiggy.	E. Indies.	1774.	— S. ℡.	Jacq. vind. 1. t. 80.
9 strictus. DC.	(wh.) upright.	Caribees.	1827.	— S. ℡.	
10 punctatus. DC.	(wh.) spotted-stalked.	Jamaica.	1636.	— S. ♂.	Comm. hort. 1. t. 31.
11 divergens. DC.	(ye.) prickly.	Abyssinia.	1816.	6. 7. S. ℡.	Bruce itin. 5. t. 6.
12 callistachys. DC. (ro.ye.)	pretty-spiked.	1820.	— S. ℡.	
13 cinereus. DC. (ro.ye.)	ash-coloured.	E. Indies.	1739.	— S. ℡.	Roxb. corom. 2. t. 174.
Mimosa cinerea. R.C.					
14 nūtans. DC. (ro.ye.)	nodding.	Senegal.	1826.	— S. ℡.	
ADENANTH'ERA. DC. BASTARD FLOWER-FENCE.		Decandria Monogynia.			
1 pavonina. DC.	(st.) pale-pellow.	E. Indies.	1759.	5. 8. S. ℡.	Rumph. amb. 3. t. 109.
2 falcatâ. DC.	(ye.) woolly-leaved.	1812.	— S. ℡.	— 3. t. 111.
PR'OSOPIS. DC.	Pr'osopis.	Polygamia Monœcia.			
1 spicigera. DC.	(ye.) eatable-podded.	E. Indies.	1812.	6. 7. S. ℡.	Roxb. corom. 1. t. 63.
2 Adenanthera. DC. (ye.)	prickly.	Coromandel.	1821. S. ℡.	
Adenanthera aculeata. H.B.					
3 hórrida. DC.	(wh.) long-spined.	S. America.	1826. S. ℡.	Kth. mim. p. 106. t. 33.
4 dulcis. DC.	(wh.) sweet-podded.	New Spain.	1824. S. ℡.	— p. 110. t. 34.
5 siliquastrum. DC. (wh.)	falcate-podded.	Chile.	1827. S. ℡.	
6 cumanensis. DC. (wh.)	Cumana.	Cumana.	1824. S. ℡.	
7 domingensis. DC. (wh.)	St. Domingo.	St. Domingo.	1826. S. ℡.	
8 juliflora. DC.	(ye.) long-spiked.	Jamaica.	1793.	5. 6. S. ℡.	
Acacia juliflora. W. Mimosa juliflora. Swartz.					
LAGONY'CHIUM. DC. LAGONY'CHIUM.		Decandria Monogynia.			
Stephaniænum. DC.	Stephan's.	Levant.	1821. H. ℡.	Breyn. cent. 1. t. 56. f. 4.
AC'ACIA. DC.	Ac'acia.	Polygamia Monœcia.			
1 alata. DC.	(ye.) wing-stalked.	N. Holland.	1803.	4. 7. G. ℡.	Bot. reg. 396.
2 dolabrifòrmis. DC. (ye.)	hatchet-leaved.	1818.	4. 6. G. ℡.	
3 decipiens. DC.	(ye.) paradoxical.	1803.	3. 6. G. ℡.	Bot. mag. t. 1745.
4 biflora. DC.	(ye.) two-flowered.	— G. ℡.	Wendl. dis. n. 3. t. 2.
5 hastulata. DC.	(ye.) hastulate.	1824.	4. 6. G. ℡.	
6 nervosa. DC.	(ye.) nervose.	— G. ℡.	
7 paradóxa. DC.	(ye.) wave-leaved.	1818.	3. 6. G. ℡.	Wendl. dis. n. 4. t. 3.
undulata. Bot. reg. t. 843.					
8 ornithophoræ. s. F.A. (ye.)	Bird-leaved.	1825.	4. 6. G. ℡.	Swt. fl. aust. t. 24.
9 armata. DC.	(ye.) armed.	1803.	— G. ℡.	Bot. mag. t. 1653.
10 hybrida.	(ye.) Whitley's mule.	Hybrid.	1822.	3. 6. G. ℡.	Lodd. bot. cab. 1342.
11 genistifolia. DC.	(ye.) Genista-leaved.	N. Holland.	1820.	— G. ℡.	
12 juniperina. DC. (st.)	Juniper-leaved.	N. S. W.	1790.	— G. ℡.	Vent. malm. t. 64.
ulicifolia. Wendl. coll. 2. t. 6. Mimosa ulicina. Sal. prodr.					
13 Brônnii. DC.	(st.) Brown's.	N. S. W.	1796.	— G. ℡.	
aciculäris. H.K. non w.					
14 echinula. DC.	(st.) pungent.	1824.	— G. ℡.	
15 pugioniformis. DC. (st.)	needle-leaved.	1803.	— G. ℡.	Wendl. dis. 26. t. 9.
16 asparagoides. F.T. (st.)	Asparagus-like.	1824.	— G. ℡.	
17 Cunninghamii. F.T. (ye.)	Cunningham's.	1825.	— G. ℡.	
taxifolia. F.T. non w.					
18 diffusa. DC.	(ye.) spreading.	1822.	3. 6. G. ℡.	Bot. reg. t. 634.
prostrata. Lodd. bot. cab. 631.					
19 sulcata. DC.	(ye.) furrowed-leaved.	N. Holland.	1803.	4. 8. G. ℡.	Bot. reg. v. 11. t. 928.
20 canaliculata.	(ye.) glaucous-furrow'd.	1823.	5. 8. I. ℡.	
21 saligna. DC.	(ye.) Willow-leaved.	1820.	3. 6. G. ℡.	Labil. n. hol. 2. t. 235.
22 emarginata. DC.	(st.) emarginate.	1816.	— G. ℡.	
23 strîcta. DC.	(ye.) upright.	N. S. W.	1790.	2. 5. G. ℡.	Bot. mag. t. 1121.
24 leprösa. DC.	(ye.) leprous.	1824.	3. 6. G. ℡.	
25 dodoneæfolia. DC. (ye.)	clammy.	1818.	4. 6. G. ℡.	Wendl. dis. n. 19. t. 7.
viscosa. Wendl.					
26 verniciflora. F.T.	(ye.) varnished.	1819.	3. 6. G. ℡.	
27 lanigera. F.T.	(ye.) woolly-stem'd.	1824.	— G. ℡.	Bot. mag. t. 2922.
virgata. Lodd. bot. cab. 1246.					
28 multinervia. DC.	(ye.) many-nerved.	N. Holland.	—	— G. ℡.	
eglandulosa. DC.	(ye.) glandless.	—	— G. ℡.	
30 hispida. DC.	(ye.) little harsh.	N. S. W.	1794.	4. 5. G. ℡.	Sm. nov. holl. t. 16.
31 cochlearis. DC.	(ye.) spoon-leaved.	N. Holland.	1820.	3. 6. G. ℡.	Labill. n. holl. 2. t. 234.
32 coriacea. DC.	(ye.) leathery.	1824.	— G. ℡.	
33 anceps. DC.	(ye.) flat-edged.	—	— G. ℡.	
34 lauriifolia. DC.	(ye.) Laurel-leaved.	S. Sea Isl.	1775.	— S. ℡.	

35 elongatæ. DC.	(ye.) elongated.	N. S. W.	1824.	3. 6.	G. h.	
36 trinervatæ. DC.	(ye.) three-nerved.	_____	1823.	4. 6.	G. h.	
37 calamifolia. DC.	(ye.) quill-leaved.	_____	1822.	1. 12.	G. h.	Bot. reg. t. 839.
38 quadrilateralis. DC.	(ye.) tetragonal-leav'd	_____	_____	4. 6.	G. h.	
39 falcata. DC.	(st.) sickle-leaved.	_____	1790.	5. 6.	G. h.	Lodd. bot. cab. 1115.
40 falciformis. DC.	(st.) sickle-shaped.	_____	1824.	4. 6.	G. h.	
41 penninervis. DC. (wh.)	feather-nerved. impressa. Bot. reg. v. 13. t. 1115!	Lodd. bot. cab. 1319.	1822.	—	G. h.	Bot. mag. t. 2754.
42 platyphylla. (st.)	broad-leaved.	N. Holland.	—	—	I. h.	
43 melanoxylon. DC. (wh.)	black-wooded.	V. Diem. Isl.	1808.	—	G. h.	Bot. mag. t. 1659.
44 heterophylla. DC. (ye.)	various-leaved.	Bourbon.	1820.	—	S. h.	
45 amena. DC.	(st.) handsome.	N. S. W.	1824.	4. 6.	G. h.	Wendl. dis. 16. t. 4.
46 myrtifolia. DC.	(st.) Myrtle-leaved.	_____	1789.	2. 5.	G. h.	Swt. flor. aust. t. 49.
47 vestita. DC.	(ye.) clothed.	N. Holland.	1820.	3. 6.	G. h.	Bot. reg. t. 698.
48 marginata. DC.	(ye.) thick margined.	N. S. W.	1803.	4. 6.	G. h.	Wendl. dis. 19. t. 5. A.
49 pyrifolia. DC.	(ye.) Pear-leaved.	_____	1824.	—	G. h.	
50 binervata. DC.	(ye.) two-nerved.	_____	—	—	G. h.	
51 bivenosa. DC.	(ye.) two-veined.	_____	—	—	G. h.	
52 buxifolia. F.T.	(ye.) Box-leaved.	_____	—	3. 5.	G. h.	
53 lunata. DC.	(ye.) lunate-leaved.	V. Diem. Isl.	1816.	3. 6.	G. h.	Swt. flor. aust. t. 42.
β brevifolia. B.C.	(ye.) short-leaved.	_____	—	—	G. h.	Lodd. bot. cab. t. 1235.
54 obtusata. DC.	(ye.) obtuse.	N. S. W.	1824.	4. 6.	G. h.	
55 crassiúscula. DC.	(st.) thick-leaved.	_____	1803.	2. 6.	G. h.	Wendl. dis. 20. t. 8.
56 suavolens. DC.	(st.) sweet-scented.	_____	1790.	—	G. h.	Lodd. bot. cab. t. 730.
β platycarpa. DC.	(st.) flat-podded.	_____	—	—	G. h.	
57 angustifolia. DC.	(st.) narrow-leaved.	_____	—	—	G. h.	Lodd. bot. cab. t. 768.
<i>Mimosa angustifolia</i> . Jacq. schœnb. 3. t. 391.						
58 linifolia. DC.	(wh.) Flax-leaved	_____	—	—	G. h.	
59 abietina. DC.	(wh.) Fir-leaved.	_____	—	3. 6.	G. h.	
60 subulata. DC.	(ye.) awl-leaved.	N. Holland.	1824.	4. 6.	G. h.	Bonpl. nav. t. 45.
61 spinulosa.	(ye.) small-spined.	_____	1822.	3. 6.	I. h.	
62 cordiòlia. ψ.	(ye.) heart-leaved.	_____	1824.	4. 6.	I. h.	
63 dillwyniaæfolia. ψ.	(st.) Dillwynia-leaved.	_____	—	—	G. h.	
64 thegonocarpa. ψ.	(ye.) angular-podded.	_____	—	—	G. h.	
65 rigens. ψ.	(ye.) stiff.	_____	—	—	G. h.	
66 Cyclopis. ψ.	(ye.) Cyclopia-like.	_____	—	—	G. h.	
67 mucronulata. ψ.	(ye.) sharp-pointed.	_____	—	—	G. h.	
68 oleifolia. ψ.	(ye.) Olive-leaved.	_____	—	—	G. h.	
69 pendula. ψ.	(wh.) pendulous.	_____	—	—	G. h.	
70 leucophylla. c.c.	(ye.) white-leaved.	_____	1820.	3. 6.	I. h.	
71 taxifolia. DC.	(ye.) Yew-leaved.	China.	—	—	G. h.	
72 Oxycedrus. DC.	(ye.) sharp-leaved.	N. S. W.	1824.	4. 6.	G. h.	Swt. flor. aust. t. 6.
<i>taxifolia</i> . Lodd. bot. cab. t. 1225. non w.						
73 verticillatæ. DC.	(ye.) whorl-leaved.	V. Diem. Isl.	1780.	3. 5.	G. h.	
α glabra. DC.	(ye.) smooth-leaved.	_____	—	—	G. h.	
β angusta. DC.	(ye.) narrow-leaved.	_____	—	—	G. h.	
γ latifolia. DC.	(ye.) broad-leaved.	_____	—	—	G. h.	
74 linearis. B.M.	(wh.) linear-leaved.	N. S. W.	1812.	8. 12.	G. h.	
75 longissima. W.A.	(ye.) long-linear-l'd.	_____	—	—	G. h.	
76 uncinata. C.C.	(ye.) hooked-leaved.	N. Holland.	1823.	9. 1.	G. h.	
77 mucronata. DC.	(wh.) mucronate.	N. S. W.	1812.	—	G. h.	
78 floribunda. DC.	(wh.) many-flowered.	_____	1796.	5. 6.	G. h.	
79 doratòxylon. F.T.	(ye.) spear-wood.	_____	1824.	3. 6.	G. h.	
80 obtusifolia. F.T.	(ye.) blunt-leaved.	_____	—	4. 6.	G. h.	
81 homomalla. DC.	(ye.) downy-leaved.	_____	—	—	G. h.	
82 longifolia. DC.	(ye.) long-leaved.	_____	1792.	3. 6.	G. h.	
β latifolia.	(ye.) broader-leaved.	_____	—	—	G. h.	
83 intertæcta. DC.	(ye.) netted-veined.	_____	1824.	4. 6.	G. h.	
84 glaucescens. DC.	(ye.) glaucescens.	_____	1790.	2. 6.	G. h.	
85 leucolobia.	(ye.) white-podded.	_____	1824.	3. 6.	G. h.	
<i>dealbata</i> . F.T. non Link.						
86 cinerascens. DC.	(ye.) greyish.	_____	—	—	G. h.	
87 subcorulea. B.R.	(ye.) bluish-leaved.	N. Holland.	—	4. 6.	I. h.	
88 rübida. F.T.	(st.) red-margined.	N. S. W.	1824.	—	G. h.	Bot. reg. t. 1075.
89 Sophoræ. DC.	(ye.) Sophora-podd'd.	V. Diem. Isl.	1805.	—	G. h.	
90 heteromalla. C.C.	(ye.) broad woolly-l'd.	N. Holland.	1822.	—	I. h.	Labil. n. holl. 2. t. 237.
91 coronillaefolia. DC.	(ye.) Coronilla-leaved.	Mogador.	1826.	4. 5.	G. h.	
92 strumbulifera. DC.	(ye.) twisted-podded.	Peru.	1823.	4. 7.	G. h.	
93 reticulata. DC.	(ye.) netted.	C. B. S.	1816.	4. 6.	G. h.	Pluk. alm. t. 123, f. 2.

94 pulchélla. DC. (ye.) zigzag spiny.	N. Holland.	1803.	4. 7.	G. h.	Lodd. bot. cab. t. 212.
95 hispidissima. DC. (ye.) hispid.	_____	1824.	4. 6.	G. h.	
96 detinens. DC. (ye.) detaining.	C. B. S.	1816.	6. 10.	I. h.	
97 viridirámis. DC. (ye.) green-branched.	_____	—	4. 6.	I. h.	
98 cassioïdes. DC. (ye.) Cassia-like.	S. America.	1822.	—	S. h.	
99 díptera. DC. (ye.) two-winged.	_____	—	—	S. h.	
100 lophanthóides. DC. (st.) crested.	Jamaica.	1826.	5. 6.	S. h.	
101 lophántha. DC. (st.) two-spiked.	N. Holland.	1803.	11. 4.	G. h.	Bot. reg. t. 361.
<i>Mimosa élégans</i> . Andr. deposit. 563.					
102 guianén sis. DC. (wh.) Guiana.	Guiana.	1803.	—	S. h.	Aubl. guian. 2. t. 357.
103 elephantina. B.T. (ye.) Elephant.	C. B. S.	1816.	—	I. h.	
104 pulchérrima. DC. (re.) beautiful.	Brazil.	1826.	—	S. h.	
105 aspidioïdes. DC. (wh.) Aspidium-leav'd. Guiana.	1827.	—	—	S. h.	
106 dealbáta. DC. (ye.) white-leaved.	V. Diem. Isl.	1818.	3. 6.	G. h.	Bot. reg. t. 371.
decúrrens β mollis. B.R.					
107 Wallichiána. DC. (ye.) Wallich's.	E. Indies.	1827.	—	S. h.	
108 adenanthéra. H.B. . . perennial.	_____	1825.	—	S. h.	
109 ferruginea. DC. (wh.) ferruginous.	_____	—	—	S. h.	
110 Sándra. DC. (ye.) hook-spined.	_____	1789.	5. 6.	S. h.	Roxb. corom. 3. t. 225.
111 Súma. H.B. (ye.) Shumee-tree.	_____	1822.	—	S. h.	
112 Catéchu. DC. (ye.) medicinal.	_____	1790.	—	S. h.	Roxb. corom. 2. t. 175.
113 fasciculáta. DC. (wh.) fascicled.	Mexico.	1828.	—	S. h.	Kth. mim. 75. t. 23.
114 cáffra. DC. (ye.) Hottentot.	C. B. S.	1816.	5. 6.	I. h.	
115 Senegal. DC. (wh.) Senegal.	Africa.	1823.	—	S. h.	
116 cornígera. DC. (ye.) Cuckold-tree.	S. America.	1692.	5. 6.	S. h.	
α americana. DC. (ye.) brown-spined.	_____	—	—	S. h.	Comm. hort. 1. t. 107.
β indica. DC. (ye.) yellow-spined.	E. Indies.	—	—	S. h.	Sebathes. 1. t. 70. f. 13.
117 latrótrum. DC. (wh.) white-spined.	_____	1824.	—	S. h.	
118 Séyal. DC. (ye.) Egyptian.	Egypt.	1828.	—	G. h.	Delil. fl. eg. t. 52. f. 2.
119 hórida. DC. (ye.) common-cape.	C. B. S.	1823.	3. 5.	I. h.	Pluk. t. 121. f. 4.
<i>Mimosa leucacántha</i> . Jacq. capén sis. B.T.					
120 ebúrnæa. DC. (ye.) ivory-thorned.	E. Indies.	1792.	4. 6.	S. h.	Roxb. corom. 2. t. 199.
121 tortuosa. DC. (st.) twisted.	Jamaica.	1822.	—	S. h.	
122 Burmanniána. DC. (ye.) Burmann's.	E. Indies.	1824.	—	S. h.	
123 véræ. DC. (ye.) Egyptian-thorn.	Egypt.	1596.	7. 8.	I. h.	Vest egypt. t. 8.
<i>Mimosa nilótica</i> . L.					
124 arábica. DC. (ye.) Gum Arabic.	Arabia.	—	6. 8.	S. h.	Roxb. corom. 2. t. 149.
125 hebecláda. DC. (ye.) stoloniferous.	C. B. S.	1816.	4. 6.	I. h.	
<i>stolonifera</i> . B.T.					
126 maurocéana. DC. (wh.) Marocca.	Marocca.	—	—	G. h.	
127 farnesiána. DC. (ye.) Sponge-tree.	St. Domingo.	1656.	6. 8.	I. h.	Duham. ar.ed.n.2.t.28.
128 álba. DC. (wh.) white.	E. Indies.	1828.	—	S. h.	
129 albicans. DC. (wh.) whitish.	Mexico.	1827.	—	S. h.	Kth. mim. 87. t. 27.
130 leucophl'æa. DC. (st.) panicled.	E. Indies.	1812.	4. 6.	S. h.	Roxb. corom. 2. t. 150.
131 hæmatóxylon. DC. (ye.) hoary.	C. B. S.	1816.	—	S. h.	
atomiphyl'a. B.T.					
132 tomentósa. DC. (st.) tomentose.	E. Indies.	—	5. 7.	S. h.	
133 índica. DC. (ye.) Indian.	_____	1824.	—	S. h.	
134 flexuosa. DC. (ye.) flexuose.	Cumana.	1822.	—	S. h.	
135 peruviana. DC. (ye.) Peruvian.	Peru.	—	—	S. h.	
136 brachyacántha. DC. . . short-spined.	S. America.	1816.	—	S. h.	
137 ciliàta. DC. . . fringed.	_____	1823.	—	S. h.	
138 aciculáris. DC. . . needle-spined.	_____	—	—	S. h.	
139 Giraffæ. DC. (ye.) Camelopard's.	C. B. S.	1816.	—	I. h.	
140 rhodacántha. DC. (ye.) red-spined.	—	—	—	S. h.	
141 heteracántha. DC. (ye.) various-spined.	C. B. S.	—	—	I. h.	
142 Bancroftiána. DC. (st.) Bancroft's.	Jamaica.	1826.	—	S. h.	
143 polycéphala. DC. (wh.) many-headed.	Mauritius.	—	—	S. h.	
144 subinérmis. DC. (ye.) scarcely spiny.	Jamaica.	—	—	S. h.	
145 édulis. W. en. . . eatable.	S. America.	1822.	—	S. h.	
146 acanthocárpa. DC. (bh.) prickly-podd'd.	New Spain.	1818.	4. 7.	F. h.	
<i>Mimosa aculeaticárpa</i> . P.S.					
147 acantholòba. DC. (st.) prickly-fruited.	S. America.	1822.	—	S. h.	
148 concinna. DC. (bh.) neat.	E. Indies.	1794.	6. 8.	S. h.	
149 tamarindifolia. DC. (wh.) Tamarind-l'd.	W. Indies.	1774.	—	S. h.	Jacq. schoenb. 3. t. 396.
150 I'ntsia. DC. (st.) angular-stalked.	E. Indies.	1778.	6. 8.	S. h.	Rheed. mal. 6. t. 4.
151 pennata. DC. (ye.) fine-leaved.	—	1773.	—	S. h.	Burm. zeyl. 2. t. 1.
152 c'asia. DC. (wh.) gray.	—	—	—	S. h.	Pluk. mant. 1. t. 330. f. 3.
153 centrophylla. DC. (wh.) prickly-leaved.	Jamaica.	1826.	—	S. h.	

154 Courantiâna. DC. (wh.)	Courant's.	1824.	6. 8.	S. h.	
155 arróphula. DC. (wh.)	Nepaul.	Nepaul.	1823.	G. h.	
156 scándens. DC.	climbing.	Brazil.	1826.	S. h. (—)	
157 sarmentosa. DC.	trailing.	—	S. h.	
158 trichòdes. DC. (wh.)	warted-stalked.	Caracas.	1824.	6. 8.	S. h.	Jacq. scheenb. 3. t. 394.
159 formôsa. DC. (wh.)	handsome.	Mexico.	1826.	S. h.	Kth. mim. t. 32.
160 nigrans. DC. (st.)	unequal-winged.	N. Holland.	1803.	5. 7.	G. h.	Bot. mag. t. 2188.
161 strigosa. DC. (ye.)	ciliate-winged.	—	3. 6.	G. h.	
	ciliata. H. K. nec aliorum.					
162 vâga. DC. (st.)	Brazilian.	Brazil.	1823.	S. h.	Marcg. bras. p. 111.f.1.
163 Lébbeck. DC. (pa.)	Egyptian.	Egypt.	4. 6.	G. h.	Pluk. mant. 2.t.331.f.1.
164 prôcera. DC. (pa.)	lofty.	Coromandel.	1824.	S. h.	Roxb. corom. 2. t. 121.
165 odoratissima. DC. (pa.)	fragrant.	1790.	4. 6.	S. h.	— 2. t. 120.
166 speciosa. DC. (wh.)	splendid.	E. Indies.	1742.	8. 9.	S. h.	Jacq. ic. 1. t. 198.
167 glâuca. DC. (wh.)	glaucous.	S. America.	1690.	6. 9.	S. h.	Trew ehret. t. 46.
168 leucocéphala. DC. (wh.)	white-headed.	1818.	4. 6.	S. h.	Plum. ed. Burm. t. 6.
169 latisiliqua. DC. (wh.)	broad-podded.	W. Indies.	1777.	3. 6.	S. h.	Jacq. ic. 3. t. 633.
170 portoricén sis. DC. (wh.)	Porto Rico.	Porto Rico.	1820.	S. h.	— 3. t. 632.
171 caracasâna. DC. (pu.)	Caracas.	Caracas.	1823.	S. h.	Bot. reg. t. 721.
172 Lambertiâna. DC. (pu.)	Mr. Lambert's.	Mexico.	1820.	7. 9.	G. h.	
173 tetragôna. DC. (wh.)	square-stalked.	Caracas.	1822.	S. h.	Bot. mag. t. 2651.
174 quadrangulâris. DC. (wh.)	quadrangular.	1816.	5. 7.	S. h.	— t. 1750.
175 discolor. DC. (ye.)	two-coloured.	N. S. W.	1788.	3. 6.	G. h.	
<i>Mimosa discolor.</i> Andr. deposit. t. 235.	<i>Mimosa botrycéphala.</i> Vent. cels. t. 1.					
176 angulâta. DC. (ye.)	angular.	N. Holland.	1824.	4. 6.	G. h.	
177 pubescens. DC. (ye.)	hairy-stemmed.	N. S. W.	1790.	3. 6.	G. h.	Bot. mag. t. 1263.
178 filicina. DC. (wh.)	fern-leaved.	Mexico.	1825.	4. 6.	G. h.	Cavan. ic. 1. t. 78.
179 villôsa. DC. (wh.)	villous.	Jamaica.	1826.	—	S. h.	
180 frondôsa. DC. (wh.)	frondose.	E. Indies.	1816.	S. h.	
181 venûsta. W. en. (ye.)	charming.	S. America.	—	S. h.	
182 stipulâta. DC. (wh.)	stipuled.	Bengal.	1820.	5. 6.	S. h.	
183 divaricata. DC. (wh.)	straddling.	E. Indies.	—	—	S. h.	Jacq. scheenb. t. 395.
184 umbellifera. DC. (wh.)	small-umbell'd.	Mexico.	1825.	—	G. h.	Kth. mim. t. 31.
185 arbôrea. DC. (bh.)	tree.	Jamaica.	1768.	—	S. h.	Sloan. his. 2.t.182.f.1.2.
186 amâra. DC. (wh.)	bitter.	Coromandel.	1823.	4. 6.	S. h.	Roxb. corom. 2. t. 122.
187 Nemû. DC. (bh.)	Japan tree.	Japan.	—	G. h.	Banks ic. Kämpf. t. 19.
188 Julibríssin. DC. (bh.)	smooth tree.	Levant.	1745.	8. 9.	H. h.	Scop. insub. 1. t. 8.
189 polyphylla. DC. (wh.)	many-leaved.	St. Martha.	1826.	6. 9.	S. h.	Vent. malm. t. 61.
190 decurrans. DC. (ye.)	decurrent.	N. S. W.	1790.	5. 7.	G. h.	Swt. flor. aust. t. 12.
191 mollissima. DC. (ye.)	soft-leaved.	—	1818.	4. 6.	G. h.	Flor. mexic. ic. ined.
192 esculentâ. DC. (wh.)	eatable.	New Spain.	1829.	S. h.	Kth. mim. t. 30.
193 peregrina. DC. (wh.)	white-flowered.	S. America.	1780.	7. 8.	S. h.	
194 grâta. DC. (wh.)	grateful.	Brazil.	1828.	S. h.	
195 microphylla. DC. (wh.)	small-leaved.	Caracas.	1826.	S. h.	
196 chrysostachys. (ye.)	golden-spiked.	Mauritius.	1823.	S. h.	
197 lúcida. H. B. glossy.	E. Indies.	1816.	S. h.	
198 Sirîssa. H. B.	(wh.) Shireesh.	—	—	S. h.	
199 elâta. H. B.	(wh.) tall.	—	1820.	S. h.	
200 Kalkôra. H. B.	(wh.) large.	—	—	S. h.	
201 Smithiâna. H. B. Smith's.	Silhet.	1823.	S. h.	
202 semicordâta. H. B. (wh.)	semicordate.	E. Indies.	—	S. h.	
203 fruticôsa. H. B.	(wh.) frutescent.	China.	1820.	5. 6.	G. h.	
204 pedunculâta. H. B. (wh.)	peduncled.	E. Indies.	1822.	S. h.	
205 triquètra. H. B.	(wh.) triquetrous.	Coromandel.	1824.	S. h.	
206 robûsta. B. T.	(ye.) robust.	C. B. S.	1816.	I. h.	
207 litakunénsis. B. T.	(ye.) Litakun.	—	—	I. h.	
208 guilandinæfolia. (gr.)	Guilandina-ld.	N. Holland.	1818.	7. 9.	I. h.	

Subordo IV. CÆSALPINEÆ.

Tribus V. GEOFFREÆ. DC. prodr. 2. p. 473.

ARACHIS. DC.

1 hypogæa. DC.

2 africana. w.

(ye.) American.

(ye.) African.

EARTH-NUT.

S. America.

Africa.

Diadelphia Decandria. L

1712. 5. 6. H. ⊕.

1824. — H. ⊕.

Trew ehret. t. 3. f. 3.

LEGUMINOSÆ.

VOANDZE'IA. DC.	VOANDZE'IA.	Diadelphia Decandria.			
subterrânea. dc. (ye.)	subterraneous.	Madagascar. 1820. 6. 8. S. ⊙.	DC. leg. me. t. 20. f. 106.		
Glycine subterrânea.	Linn. dec. 37. t. 17.				
PERA'LTEA. DC.	PERA'LTEA.	Diadelphia Decandria.			
lupinoïdes. dc. (pu.)	Lupine-like.	New Spain. 1827. 9. 11. F. ℥.	H. et B. n. gen. 6. t. 589.		
BRONGNIA'R'TIA. DC.	BRONGNIA'R'TIA.	Diadelphia Decandria.			
podalyrioïdes. dc.(fl.)	Podalyria-like.	New Spain. 1827. 9. 11. F. ℥.	H. et B. n. gen. 6. t. 588.		
AND'IRA. DC.	AND'IRA.	Diadelphia Decandria.			
1 racemosa. dc. (pu.)	racemed.	Brazil. 1824. S. ℥.	Aubl. gui. sup. t. 373.		
2 inermis. dc. (pu.)	bastard cabbage-tree.	Jamaica. 1778. S. ℥.	Phil. trans. 1777. t. 10.		
GEOFFRO'YA. DC.	GEOFFRO'YA.	Diadelphia Decandria.			
1 spinosa. dc. (st.)	spiny.	S. America. 1823. S. ℥.	Jacq. amer. t. 180. f. 62.		
2 Bredemeyeri. dc.(ye.)	striated.	Caracas. 1824. S. ℥.			
Robinia striata. w.					
3 violacea. dc. (vi.)	violet.	Guiana. 1827. S. ℥.	Aubl. guian. t. 301.		
BRO'WNEA. DC.	BRO'WNEA.	Monadelphia Dec-Dodecandria.			
1 Rosa. dc. (ro.)	mountain Rose.	Trinidad. 1816. S. ℥.	Ber. act. ang. 1771. t. 8-9.		
speciosa. Sieber.					
2 coccinea. dc. (sc.)	scarlet-flower'd.	W. Indies. 1793. 7. 9. S. ℥.	Jacq. amer. t. 121.		
3 latifolia. dc. (ro.)	broad-leaved.	Caracas. 1826. S. ℥.	Jacq. fragm. t. 17.		
4 racemosa. dc. (ro.)	racemed.	—— —— —— S. ℥.	—— t. 18-19.		
5 grandicéps. dc. (ro.)	close-headed.	—— —— 1828. S. ℥.	—— t. 22-23.		
DIPTERIX. DC.	TONQUIN BEAN.	Monadelphia Octo-Decandria.			
odorata. dc. (pu.)	sweet-scented.	Guiana. 1793. S. ℥.	Aubl. guian. 3. t. 296.		
Coumaroūna odorata.	Aubl. Baryosma Tonga. Gært. fruct. 2. t. 93.				

Tribus X. *CASSIEÆ.* DC. prodr. 2. p. 478.**MORI'NGA.** DC. HORSE-RADISH-TREE. Decandria Monogynia.

1 pterigospérma. DC. (st.)	wing-seeded.	E. Indies. 1759. S. ℥.	Jacq. ic. 3. t. 461.
oleífera. Lam.	Zeylánica. P.S.	Hyperanthèra Moriṅga. Vahl.	Guilindina Moriṅga. L.
2 polygona. dc.	(st.) many-sided.	—— —— 1822. S. ℥.	

GLEDITSCHIA. DC. **GLEDITSCHIA.** Polygamia Diœcia. L.

1 triacanthos. dc. (gr.)	three-thorned.	N. America. 1700. 6. 8. H. 5.	Wats. dend. brit. 2. t. 138.
β inermis. dc.	(gr.) spineless.	—— —— —— H. 5.	DC. leg. mem. t. 22. f. 109
2 brachycarpa. dc. (gr.)	curved-spined.	—— —— —— H. 5.	
3 monospermum. dc. (gr.)	single-seeded.	—— —— 1723. —— H. 5.	Mich. f. arb. 3. t. 10.
4 sinensis. dc. (gr.)	Chinese.	China. 1774. —— H. 5.	DC. leg. mem. 1. t. 1.
horrida. w.			
5 macracantha. dc. (gr.)	large-spined.	—— —— —— H. 5.	
6 ferox. dc.	(gr.) flat-spined.	—— —— —— H. 5.	
orientalis. Bosc.			

GYMNOCLADUS. DC. KENTUCKY COFFEE-TREE. Diœcia Decandria.

canadensis. dc. (wh.)	Canadian.	N. America. 1748. 5. 6. H. 5.	Mich. amer. 2. t. 51.
Guilindina dioica. L.			

GUILAND'INA. DC. NICKER-TREE. Decandria Monogynia. L.

1 Bónduc. L.T.	(ye.) Sumatra.	Sumatra. 1640. S. ℥.	Rumph. amb. 5. t. 48.
2 Bonducélla. L.T.	(ye.) common.	E. Indies. —— —— —— S. ℥.	Schrink. h. mon. t. 68.
3 microphylla. dc. small-leaved.	—— —— —— S. ℥.	Rumph. amb. 5. t. 49. f. 2?

COULTERIA. DC. **COULTERIA.** Decandria Monogynia.

1 tinctoria. dc.	(ye.) dyer's.	Peru. 1818. S. ℥.	H. et B. n. ge. am. 6. t. 569.
Poinciâna Târa. R.P.	Cesalpina pectinata. Cavan.		

2 mexicana. dc. (ye.) Mexican. New Spain. 1826. S. ℥. Flor. mexic. ic. ined.**CESALPINIA.** DC. **BRASILE'TTO.** Decandria Monogynia. L.

1 Nùga. dc.	(ye.) acute-leaved.	Amboyna. 1801. 4. 6. S. ℥.	Rumph. amb. 5. t. 50.
2 paniculata. dc.	(ye.) paniced.	E. Indies. 1812. S. ℥.	Rheed. mal. 6. t. 19.
3 brasiliensis. dc.	(ye.) Brazilian.	Brazil. 1818. 4. 6. S. ℥.	
4 Sappán. dc.	(ye.) narrow-leaved.	E. Indies. 1773. —— S. ℥.	Roxb. corom. 1. t. 16.

5	dig'yna. DC.	(ye.) digynous.	E. Indies.	1828.	S. h.	Will.n.ac.1803.v.4.t.3.
6	mimosoïdes. DC.	(ye.) Mimosa-leaved.	—	1806.	5. 7.	S. h.	Rheed. mal. 6. t. 8.
7	bijûga. DC.	(ye.) broad-leaved.	Jamaica.	1770.	6. 9.	S. h.	Sloan.j. 2. t. 181. f. 2. 3.
8	bahaménsis. DC.	(u.h.) emarginate.	Bahama.	1823.	S. h.	Catesb. car. 2. t. 51.
9	Crista. DC.	(ye.) oval-leaved.	Jamaica.	4. 6.	S. h.	Plum. gen. 28. t. 9.
10	glandulòsa. DC.	(ye.) glandular.	St. Domingo.	1828.	S. h.	
11	Cacaláco. DC.	(ye.) round-leaved.	Mexico.	1826.	S. h.	H. et B. pl. æq. 2. t. 137.
12	coriaria. DC.	(ye.) leathery.	W. Indies.	—	S. h.	Kth. mim. t. 45.
13	lebbekkoïdes. DC.	(ye.) Lebbek-podd'd.	China.	1812.	S. h.	
14	punctatâ. DC.	(ye.) dotted-leaved.	Brazil.	1818.	S. h.	
15	prôcerâ. s.s.	(ye.) tall.	Cuba.	1822.	S. h.	
16	cassioïdes. DC.	(ye.) Cassia-like.	S. America.	1823.	S. h.	
17	mucronatâ. DC.	(ye.) mucronate.	Brazil.	—	S. h.	
18	olæospérma. H.B.	(ye.) oil-seeded.	E. Indies.	1800.	5. 6.	S. h. ↗	
19	cucullatâ. H.B.	(ye.) hooded.	—	1822.	S. h. ↗	
20	sepiâria. H.B.	(ye.) Mysore-thorn.	—	1806.	6. 9.	S. h. ↗	
21	resupinatâ. H.B.	(ye.) resupinate.	—	1828.	S. h. ↗	
22	enneaphylla. H.B.	(ye.) nine-leaved.	—	1812.	S. h. ↗	
23	chinénsis. H.B.	(ye.) Chinese.	China.	—	S. h. ↗	
24	tortuosa. H.B.	(ye.) twisting.	E. Indies.	1822.	S. h. ↗	
25	sumatrâna. H.B.	(ye.) Sumatra.	—	—	S. h. ↗	
POINCIANA. DC. FLOWER FENCE.		Decandria Monogynia. L.					
1	règia. B.M.	(sc.) superb.	Madagascar.	1827.	S. h.	Bot. mag. t. 2884.
2	puschérmina. DC.	(or.) Barbadoes.	E. Indies.	1691.	6. 9.	S. h.	— t. 995.
3	elâtâ. DC.	(ye.) fringe-petal'd.	—	1778.	—	S. h.	
<i>Casalpínia elâtâ. s.s.</i>							
HOFFMANSE'GGIA. DC. HOFFMANSE'GGIA.		Decandria Monogynia.					
1	falcâria. DC.	(ye.) sickle-podded.	Chile.	1806.	7. 8.	F. ↗	Cavan. ic. t. 392.
2	prostrâta. DC.	(ye.) prostrate.	Lima.	1824.	—	F. ↗	
3	elâtâ. DC.	(ye.) fringe-petal'd.	—	—	—	S. h.	
MELANOSTI'CTA. DC. MELANOSTI'CTA.		Decandria Monogynia.					
Burchellii. DC.		(ye.) Burchell's.	C. B. S.	1816.	G. h.	DC. leg. mem. t. 69.
POM'ARIA. DC. POM'ARIA.		Decandria Monogynia.					
glandulòsa. DC.		(ye.) glandular.	New Spain.	1826.	5. 6.	G. h.	Cavan. ic. 5. t. 402.
HÉMATO'XYLON. D. LOGWOOD.		Decandria Monogynia. L.					
campechianum. DC.		(ye.) common.	S. America.	1724.	S. h.	Catesb. car. 2. t. 66.
PARKINS'ONIA. DC. PARKINS'ONIA.		Decandria Monogynia. L.					
aculeâtâ. DC.		(ye.) prickly.	W. Indies.	1739.	6. 9.	S. h.	Jacq. amer. t. 80.
CA'DIA. DC. CA'DIA.		Decandria Monogynia.					
vâria. DC.		(bh.) various-colour'd.	Arabia.	1755.	7. 10.	S. h.	
purpùrea. w. <i>Spendónea tamarindifolia</i> . Desf.							Picc. hort. pan. 9. ic.
CERAT'ONIA. DC. CAROB-TREE.		Polygamia Dicecia.					
Siliqua. DC.		(gr.) St.John's bread.	Levant.	1570.	9. 10.	G. h.	Andr. reposit. 567.
HARDWI'CKIA. DC. HARDWI'CKIA.		Oct-Decandria Monogynia.					
1	binâtâ. DC.	(ye.) binate-leaved.	E. Indies.	1824.	S. h.	Roxb. corom. t. 209.
2	pinnâtâ. DC.	(ye.) pinnate-leaved.	—	1826.	S. h.	
JON'ESIA. DC. JON'ESIA.		Octandria Monogynia.					
Asôca. DC.		(or.) winged-leaved.	E. Indies.	1796.	4. 6.	S. h.	Rheed. mal. 5. t. 59.
pinnâtâ. w. <i>Sarâca indica</i> . L.							
TACHIGALIA. DC. TACHIG'ALIA.		Decandria Monogynia.					
paniculâtâ. DC.		(ye.) panicled.	Guiana.	1827.	S. h.	Aubl. guian. 1.t.143.f 1.
MOLDENHAW'ERA. DC. MOLDENHAW'ERA.		Decandria Monogynia.					
floribûnda. DC.		(ye.) many-flowered.	Brazil.	1828.	S. h.	Neuw. reis. v. 2. ic.
TAMARI'NDUS. DC. TAMARIND-TREE.		Monadelphia Triandria.					
1	indica. DC.	(ye.) East Indian.	E. Indies.	1633.	6. 7.	S. h.	Rheed. mal. 1. t. 23.
2	occidentalis. DC.	(ye.) West Indian.	W. Indies.	—	—	S. h.	Jacq. amer. t. 10.
CA'SSIA. DC. CA'SSIA.		Decandria Monogynia. L.					
1	brasiliâna. DC.	(ye.) large.	Brazil.	1822.	S. h.	Breyn. cent. t. 21.
grândis. L. mollis. Vahl. <i>Cathartocárpus grândis</i> . P.S.							
2	ferruginea. DC.	(ye.) ferruginous.	Brazil.	1828.	S. h.	Rumph. amb. 2. t. 22.
3	Roxburghii. DC.	(ye.) Roxburgh's.	Coromandel.	1823.	6. 7.	S. h.	Woody. med. bot. t. 163.
marginatâ. H.B. non w.							
4	trinitatis. DC.	(ye.) Trinidad.	Trinidad.	1824.	S. h.	
5	javâica. DC.	(fl.) Java.	Java.	1779.	6. 7.	S. h.	
6	Fistula. DC.	(ye.) purging.	E. Indies.	1731.	—	S. h.	
7	rhombifòlia. H.B.	(ye.) rhomb-leaved.	—	1820.	—	S. h.	
8	nodôsa. H.B.	(ye.) knotted.	—	1824.	—	S. h.	
9	conspicua. G.D.	(ye.) drumstick-tree.	Sierra Leon.	1822.	S. h.	
10	bacillaris. DC.	(ye.) smooth.	Surinam.	1818.	—	S. h.	Houst. rel. t. 17.
<i>Cathartocárpus Bacillus</i> , Bot. Reg. t. 881.							

LEGUMINOSÆ.

11 speciōsa. DC.	(ye.) handsome.	Brazil.	1828.	6. 8.	S. h.
12 melanocarpa. DC.	(ye.) black-podded.	Jamaica.	1825.	—	S. h.
13 corymbosa. DC.	(ye.) corymbose.	Buenos Ayres.	1796.	7. 8.	S. h.
14 floribunda. DC.	(ye.) many-flowered.	New Spain.	1822.	5. 8.	S. h.
15 lavigata. DC.	(ye.) smooth.	—	1811.	—	S. h.
16 pendula. DC.	(ye.) pendulous.	S. America.	1820.	—	S. h.
17 torosa. DC.	(ye.) twisted.	China.	1807.	5. 7.	G. h.
<i>chinensis.</i> Jacq. non Lam.					
18 chinensis. DC.	(ye.) Chinese.	—	—	—	G. h.
19 gigantea. DC.	(ye.) gigantic.	Jamaica.	1825.	—	S. h.
20 coromandeliæna. DC. (or.)	Coromandel.	Coromandel.	1822.	5. 7.	S. h.
21 Sphæra. DC.	(ye.) round-podded.	E. Indies.	1653.	7. 9.	S. h.
22 alata. DC.	(ye.) winged-podded.	S. America.	1731.	—	S. h.
23 Rumphiana. DC.	(ye.) Rumphius's.	E. Indies.	1822.	—	S. h.
24 bracteata. DC.	(ye.) bracteate.	Surinam.	—	—	S. h.
25 obovata. DC.	(ye.) obovate-leaved.	Egypt.	1640.	7. 9.	G. (○)
<i>Senna.</i> Lam. ill. t. 332. f. 2.					
26 lanceolata. DC.	(ye.) spear-leaved.	—	1759.	6. 8.	G. (○)
<i>orientalis.</i> P.S. <i>acutifolia.</i> Delil. ill. eg. 75. t. 27. f. 1.					
27 ligustrinoïdes. DC. (ye.)	Privet-like.	Arabia.	1829.	—	G. h.
28 Browniana. DC.	(ye.) Brown's.	Mexico.	1828.	—	S. h.
29 cuspidata. W.en.	(ye.) sharp-pointed.	S. America.	1820.	5. 7.	S. h.
30 sericea. DC.	(ye.) silky-leaved.	—	1731.	5. 8.	S. (○)
<i>sensitiva.</i> Jacq.					
31 obtusifolia. DC.	(ye.) blunt-leaved.	Jamaica.	1732.	7. 8.	S. (○)
32 humilis. DC.	(ye.) dwarf.	S. America.	—	—	S. (○)
33 Tora. DC.	(st.) oval-leaved.	E. Indies.	1693.	—	S. (○)
34 toroïdes. H.B.	(st.) Mysore.	—	1812.	—	S. (○)
35 ciliaris. DC.	(ye.) fringe-stipuled.	—	—	—	S. (○)
36 Tagera. DC.	(ye.) long-podded.	—	1803.	—	S. (○)
37 venustula. DC.	(ye.) pretty.	Cumana.	1825.	—	S. h.
38 aversiflora. B.M.	(ye.) contrary-flow'd.	Brazil.	1823.	4. 12.	S. h.
39 viminea. DC.	(ye.) twiggy.	W. Indies.	1786.	6. 7.	S. h.
40 chrysotricha. DC.	(ye.) golden-podded.	Guiana.	1828.	—	S. h.
41 Apoucouita. DC.	(ye.) acuminate.	—	1823.	—	S. h.
<i>acuminata.</i> W. non Moench.					
42 sennoïdes. DC.	(ye.) Senna-leaved.	E. Indies.	1808.	7. 8.	S. h.
43 bicapsularis. DC.	(ye.) six-leaved.	W. Indies.	1730.	5. 6.	S. h.
44 mollissima. DC.	(ye.) soft-leaved.	S. America.	1816.	7. 8.	S. h.
45 artemisioides. DC. (ye.)	Wormwood-like.	N. S. W.	1823.	4. 7.	G. h.
46 glutinosa. DC.	(ye.) glutinous.	N. Holland.	—	—	G. h.
47 oxyadena. DC.	(ye.) acute-glanded.	Jamaica.	1820.	—	S. h.
48 dispar. DC.	(ye.) various-leaved.	S. America.	1822.	5. 8.	S. h.
49 acapulcensis. DC.	(ye.) Acapulco.	Mexico.	1824.	—	S. h.
50 sulfurea. DC.	(st.) straw-coloured.	E. Indies.	1800.	—	S. h.
<i>arborescens.</i> Vahl. non Miller.					
51 coluteoides. DC.	(ye.) Colutea-like.	—	—	—	S. h.
52 Barclayana. S.F.A.	(ye.) Mr. Barclay's.	N. Holland.	1824.	6. 10.	G. h.
53 biflora. DC.	(ye.) two-flowered.	W. Indies.	1766.	12. 4.	S. h.
54 tomentosa. DC.	(st.) tomentose.	S. America.	1822.	6. 8.	S. h.
<i>multiglandulosa.</i> Jacq.					
55 mexicana. DC.	(ye.) Mexican.	Mexico.	1824.	—	S. h.
56 stipulacea. DC.	(ye.) large-stipuled.	Chile.	1786.	—	S. h.
57 auriculata. DC.	(ye.) eared.	E. Indies.	1777.	7. 9.	S. h.
58 frondosa. DC.	(ye.) smooth shrubby.	W. Indies.	1769.	3. 4.	S. h.
59 australis. B.M.	(ye.) New Holland.	N. Holland.	1820.	5. 12.	I. h.
60 polypylla. DC.	(ye.) many-leaved.	Portorico.	1828.	6. 9.	S. h.
61 marginata. DC.	(ye.) white-edged.	Surinam.	1822.	—	S. h.
62 aegyptiaca. DC.	(ye.) Egyptian.	Egypt.	1823.	7. 9.	G. (○)
63 esculenta. H.B.	(ye.) esculent.	E. Indies.	—	—	S. (○)
64 occidentalis. DC.	(ye.) occidental.	W. Indies.	1759.	3. 8.	S. h.
65 purpurea. DC.	(ye.) purple-stalked.	E. Indies.	1823.	6. 11.	S. h.
66 hirsuta. DC.	(ye.) hairy.	S. America.	1778.	7. 9.	S. h.
<i>caracasana.</i> Jacq.					
67 linearis. DC.	(ye.) linear-podded.	Carolina.	1828.	5. 7.	F. 4.
68 pátula. DC.	(ye.) shining.	W. Indies.	1778.	8. 9.	S. h.
69 pubescens. DC.	(ye.) pubescent.	—	1818.	5. 8.	S. h.
70 sulcata. DC.	(ye.) furrowed.	S. America.	1826.	6. 8.	S. h.
71 ruscifolia. DC.	(ye.) Ruscus-leaved.	Caracas.	1816.	5. 7.	G. h.

Bot. mag. t. 633.

Jacq. ic. 1.t. 73.

Jacq. fragm. t. 100.

Bur. zeyl. t. 98.

Jacq. obs. 2. t. 45. f. 2.

Rumph. amb. 7. t. 18.

Jacq. f. ecl. 1. t. 87.

Tabern. ic. 507.

Kth. mim. t. 41.

Jacq. ic. 3. t. 459.

Dill. elth. t. 62. f. 72.

Plum. ed. Burm. t. 76. f. 2.

Dill. elth. t. 63. f. 73.

Rheed. mal. 2. t. 53.

Bot. mag. t. 2638.

Sloan. jam. 2.t.180.f.67.

Coll. mon. p. 99. t. 13.

Aubl. gui. 1. t. 146.

Jacq. ic. 1. t. 70.

Bot. mag. t. 633.

Rheed. mal. 6. t. 9-10.

Coll. mon. 102. t. 12.

Swt. flor. aust. t. 32.

Bot. mag. t. 810.

Jacq. ic. 1. t. 72.

Jacq. schœnb. t. 203.

Feuill. per. 3. t. 42.

Pluk. alm. t. 314. f. 4.

Jacq. ic. 1. t. 74.

Bot. mag. t. 2670.

Jacq. ic. t. 460.

Bot. reg. t. 83.

——— t. 856.

Jacq. schœnb. t. 270.

Jacq. fragm. t. 57.

Coll. mon. p. 110. t. 6.

Jacq. ic. 1. t. 71.

72	ligustrina. DC.	(ye.) Privet-leaved.	America.	1726.	6. 8.	S. h.	Bot. reg. t. 109.
73	robinioïdes. DC.	(ye.) Robinia-like.	S. America.	1820.	—	S. h.	Schkuhr hand. 1.t.113.
74	maryländica. DC.	(ye.) Maryland.	N. America.	1723.	8. 10.	H. 4.	
75	latifolia. DC.	(ye.) broad-leaved.	Trinidad.	1826.	6. 8.	S. h.	
76	arboréscens. DC.	(ye.) arborescent.	New Spain.	—	S. h.	
77	opáca. Gr.	(or.) dull-leaved.	S. America.	1825.	8. 12.	S. h.	
78	pulchella. Boj.	(ye.) pretty.	Zanzibar.	1825.	7. 9.	S. h.	
79	Milléri. DC.	(ye.) Miller's.	Jamaica.	6. 8.	S. h.	
80	Houstoniana. DC.	(ye.) Houstoun's.	7. 10.	S. ⊙.	
81	Plumiéri. DC.	(ye.) Plumier's.	Guadaloupe.	1822.	—	S. h.	Plum. ed. Burm. t. 77.
	planisiliqua. Lam.	non Linn.					
82	hírta. DC.	(ye.) hairy-branched.	S. America.	1822.	6. 9.	S. h.	
83	sumatrana. DC.	(ye.) Sumatra.	Sumatra.	1823.	—	S. h.	
84	formosa.	(ye.) beautiful.	E. Indies.	—	—	S. h.	
	speciosa. H.B. non Schrader.						
85	emarginata. DC.	(ye.) notch-leaved.	Jamaica.	1759.	5. 6.	S. h.	Sloan. his. 2.t.180.f.1-4.
86	atomaria. DC.	(ye.) woolly-leaved.	S. America.	1822.	3. 8.	S. h.	
87	reticulata. DC.	(ye.) netted-leaved.	Brazil.	—	4. 6.	S. h.	
88	spectabilis. DC.	(ye.) showy.	Caracas.	1823.	5. 7.	S. h.	Coll. mon. p. 115. t. 7.
89	sétigera. DC.	(ye.) seta-bearing.	E. Indies.	1826.	—	S. h.	
90	montana. DC.	(ye.) mountain.	1822.	—	S. h.	
91	A'bsus. DC.	(sn.) four-leaved.	Ceylon.	1777.	6. 7.	S. ⊙.	Burm. zeyl. t. 97.
92	hispida. DC.	(ye.) hispid.	Cayenne.	1826.	—	S. ⊙.	
93	diphylla. DC.	(ye.) two-leaved.	W. Indies.	1781.	5. 7.	S. ♂.	Cavan. ic. 5.t. 600. f. 1.
94	bifoliolata. DC.	(ye.) two-leafleted.	Brazil.	1825.	—	S. h.	Coll. mon. 120.t.9.f.B.
95	exigua. H.B.	(ye.) dwarf.	Bengal.	1820.	8. 10.	S. ⊙.	
96	gracilis. DC.	(ye.) slender.	S. America.	1816.	6. 8.	S. h.	
97	uniflora. DC.	(re.) one-flowered.	Brazil.	1824.	—	S. h.	
98	lineata. DC.	(ye.) lined.	Jamaica.	1820.	5. 7.	S. h.	
99	prostrata. DC.	(ye.) prostrate.	S. America.	1823.	7. 9.	S. h.	
100	triflora. DC.	(ye.) three-flowered.	1825.	—	S. ⊙.	Jacq. schoenb. t. 480.
101	Burmanni. DC.	(ye.) Burmanni's.	C. B. S.	1816.	—	G. h.	
102	virgata. DC.	(ye.) slender-twigg'd.	Jamaica.	1820.	5. 8.	S. h.	Linn. hort. cliff. t. 36.
103	nictitans. DC.	(ye.) Virginian.	N. America.	1800.	7.	G. ♂.	Bot. mag. t. 157.
104	chamacrista. DC.	(ye.) dwarf.	America.	1699.	6. 9.	G. ♂.	
105	chamaecristoides. DC.	(ye.) large-flowd.	Vera Cruz.	—	S. h.	
106	strícta. Sck.	(ye.) straight.	Jamaica.	—	—	S. ♂.	Schk. hort. mon. t. 34.
107	glandulosa. DC.	(ye.) glandular.	W. Indies.	1822.	—	S. ♂.	
108	mimosoides. DC.	(ye.) Mimosa-leaved.	E. Indies.	1806.	—	S. ⊙.	
109	capensis. DC.	(ye.) Cape.	C. B. S.	1816.	4. 6.	G. h.	Coll. mon. 124. t. 19.
110	pūmila. DC.	(ye.) small.	E. Indies.	1820.	7. 10.	S. ⊙.	Pluk. alm. t. 120.f. 1.
111	procumbens. DC.	(ye.) procumbent.	N. America.	1806.	6. 9.	H. ⊙.	Comm. petrop. t. 11.
112	Wallichiana. DC.	(ye.) fine-leaved.	Nepaul.	1812.	—	S. ⊙.	
	dimidiata. D.P.						
113	ténera.	(ye.) small-branched.	E. Indies.	1818.	—	S. ♂.	
	tenella. H.B. non Kth.						
114	microphylla. DC.	(ye.) small-leaved.	W. Indies.	1822.	—	S. ⊙.	
115	Hornemannia. DC.	(ye.) Hornemann's.	1827.	—	S. ⊙.	
	venosa. Horn. non Zucc.						
AFZ'ELIA. DC.	AFZ'ELIA.	Octandria Monogynia.					
africana. DC.	(cr.) African.	Sierra Leone.	1821.	S. h.		
SCHOT'TIA. DC.	SCHOT'TIA.	Decandria Monogynia.					
1	speciosa. DC.	(cr.) superb.	C. B. S.	1759.	7. 12.	I. h.	Andr. deposit. t. 348.
2	tamarindifolia. DC.	(cr.) Tamarind-leav'd.	—	1795.	5. 9.	I. h.	Bot. mag. t. 1153.
3	stipulata. DC.	(cr.) falcate-stipuled.	—	1794.	—	I. h.	
4	alaia. DC.	(re.) winged.	—	1816.	6. 7.	I. h.	
5	omplifolia. DC.	(re.) simple-leaved.	—	—	—	I. h.	
6	latifolia. DC.	(re.) broad-leaved.	—	—	5. 8.	I. h.	Hook. ex. flor. t. 159.
	Omphalobium Schötia	s.s. Jacq. ecl. ic.					
COPAI'FERA. DC.	BALSAM OF CAPEVI.	Decandria Monogynia. L.					
1	officinalis. DC.	(st.) officinal.	S. America.	1774.	S. h.	
2	guianensis. DC.	(st.) Guiana.	Guiana.	1826.	S. h.	
3	coriacea. DC.	(st.) leathery-leaved.	Brazil.	1824.	S. h.	
CYNOM'ETRA. DC.	CYNOM'ETRA.	Decandria Monogynia. L.					
1	cauliflora. DC.	(re.) stem-flowered.	E. Indies.	1804.	S. h.	Rumph. amb. 1. t. 62.
2	ramiflora. DC.	(re.) branch-flowered.	—	1826.	S. h.	— 1. t. 63.
3	polyandra. DC.	(re.) many stamen'd.	—	1822.	S. h.	Roxb. corom. t. 286.
PARIVO'A. DC.	PARIV'OA.	Diadelphia Decandria.					
grandiflora. DC.	(pu.) large-flowered.	Cayenne.	1816.	S. h.	Aubl. guian. 2. t. 303	
Dimorphia grandiflora. Linn.	trans. 9. t. 20.						

O'UTEA. DC.	O'UTEA. Tri-Tetrandria Monogynia.
1 guianensis. DC. (wh.) Guiana.	Guiana. 1827. S. h.
2 bijuga. DC. (wh.) two-paired.	E. Indies. 1823. S. h.
<i>Macrolobium bijugum</i> . Linn, trans.	
VOU'APA. DC.	Vou'APA. Triandria Monogynia.
bifolia. DC. two-leaved.	Guiana. 1823. S. h.
HYMEN'EA. DC.	LOCUST-TREE. Decandria Monogynia.
1 Courbaril. DC. (wh.) leathery-leaved.	S. America. 1688. 6. 8. S. h.
2 Candolleana. DC. (wh.) Decandolle's.	Mexico. 1825. S. h.
3 verrucosa. DC. (wh.) wart-podded.	Madagascar. 1808. 6. 7. S. h.
BAUHINIA. DC.	BAUH'NIA. Monadelphia v. Decandria Monogynia.
1 divaricata. DC. (wh.) dwarf.	W. Indies. 1742. 6. 9. S. h.
2 Lamarckiana. DC. (wh.) Lamarck's.	S. America. — S. h.
3 spathacea. DC. (wh.) spathaceous.	New Spain. 1825. S. h.
4 subtropidifolia. DC. (w.) round-leaved.	S. America. 1816. 4. 6. S. h.
5 aurita. DC. (wh.) long-eared.	Jamaica. 1756. 9. 10. S. h.
6 porræta. DC. (wh.) smooth-leaved.	W. Indies. 1737. 7. 8. S. h.
7 latifolia. DC. (wh.) broad-leaved.	New Spain. 1825. S. h.
8 candida. DC. (wh.) white-flowered.	E. Indies. 1777. 5. 6. S. h.
9 acuminata. DC. (wh.) acute-leaved.	1808. 7. 8. S. h.
10 grandiflora. DC. (wh.) large-flowered.	Peru. 1824. S. h.
11 aculeata. DC. (wh.) prickly-stalked.	W. Indies. 1737. 6. 8. S. h.
12 forficata. DC. (wh.) forked.	Brazil. 1824. S. h.
13 emarginata. DC. (wh.) emarginate.	Carthagena. — S. h.
14 rufescens. DC. rufous.	Africa. 1816. S. h.
15 parviflora. DC. (wh.) small-flowered.	E. Indies. 1808. S. h.
16 U'ngula. DC. (wh.) villous-leaved.	Caracas. 1827. S. h.
aculeata. Jacq. amer. t. 177. f. 2.	
17 variegata. DC. (v.) variegated-fl'd.	E. Indies. 1690. 6. 7. S. h.
18 tomentosa. DC. (st.) tomentose.	1808. S. h.
19 pubescens. DC. (wh.) pubescent.	W. Indies. 1824. S. h.
20 multinervia. DC. (wh.) many-nerved.	Caracas. 1825. S. h.
21 racemosa. DC. large-leaved.	E. Indies. 1790. S. h. — Vahl. symb. 3. t. 62.
22 purpurea. DC. (pu.) purple.	1778. S. h.
23 retusa. DC. (wh.) retuse-leaved.	1822. S. h.
24 corymbosa. DC. (wh.) corymbose.	China. 1816. S. h. — DC. leg. mem. t. 70.
25 anguina. DC. (wh.) snake.	E. Indies. 1818. S. h. — Roxb. corom. 3. t. 285.
scandens. L. excl. synon. Rumph. Rheed. mal. 8. t. 30-31.	
26 Língua. DC. (st.) tongue-leaved.	E. Indies. 1799. S. h. — Rumph. amb. 5. t. 1.
scandens. L. excl. syn. Rheed.	
27 coccinea. DC. (sc.) scarlet-flowered.	China. 1826. S. h. —
28 guianensis. DC. (wh.) Guiana.	Guiana. 1824. S. h. — Aubl. guian. 1. t. 145.
29 heterophylla. DC. (wh.) various-leaved.	Caracas. 1824. S. h. — Kth. mim. t. 46.
30 cumanensis. DC. (wh.) Cumaná.	Cumana. 7. 8. S. h. — Bot. reg. t. 1133.
31 anatómica. L. en. jagged-leaved.	S. America. 1816. S. h.
32 triándra. H.B. (wh.) triandrous.	E. Indies. 1823. S. h.
33 semibifida. H.B. (wh.) half-cleft.	1818. S. h. —
34 malabarica. H.B. Malabar.	Malabar. 1823. S. h.
35 ferruginea. H.B. ferruginous.	E. Indies. 1822. S. h. —
36 cordifolia. H.B. heart-leaved.	1828. S. h. —
37 spicata.	spiked.
38 speciosa. L.C.	handsome.
39 microphylla. L.C.	small-leaved.
40 esculentæ. B.T. (wh.) eatable-rooted.	C. B. S. 1816. — I. 4.
CE'RCIS. DC.	JUDAS-TREE. Decandria Monogynia. L.
1 Siliquastrum. DC. (ro.) European.	S. Europe. 1596. 5. 6. H. h.
2 canadensis. DC. (ro.) American.	N. America. 1730. H. h.
CR'UDYA. DC.	CR'UDYA. Octo-Decandria Monogynia.
1 spicata. DC. (wh.) spiked.	Guiana. 1825. S. h.
<i>Apalatða spicata</i> . Aubl.	
2 aromatica. DC. (gr.) aromatic.	1828. S. h.
<i>Touchirða aromatica</i> . Aubl.	
COD'ARIUM. DC.	VELVET TAMARIND. Diandria Monogynia.
1 acutifolium. DC. (re.) acute-leaved.	Sierra Leone. 1800. S. h.
<i>nítidum</i> . Vahl. <i>Diàlum guineense</i> . w.	
2 obtusifolium. DC. (re.) blunt-leaved.	— S. h.

Aubl. guian. 1. t. 9.
Linn. trans. 12. t. 17.

Aubl. guian. 1. t. 7.

Lam. ill. t. 330. f. 1.
H. et B. n. gen. 6. t. 566.
Lam. ill. t. 330. f. 2.

Linn. hort. cliff. t. 15.

Cavan. ic. 5. t. 406.
Miller ic. t. 61.
Bot. mag. t. 1708.
Cavan. ic. t. 405.

Rheed. mal. 1. t. 34.

Plum.ed. Bur.t. 44. f. 1.

Jacq. fragm. t. 15. f. 1.
Rheed. mal. 1. t. 32.
Burm. zeyl. t. 18.

Burm. zeyl. t. 18.

Rheed. mal. 1. t. 32.

Burm. zeyl. t. 18.

Vahl. symb. 3. t. 62.

Rheed. mal. 1. t. 33.

DC. leg. mem. t. 70.

Roxb. corom. 3. t. 285.

Kth. mim. t. 46.

Bot. reg. t. 1133.

Aubl. guian. 1. t. 145.

Aubl. guian. 1. t. 147.

Miller ic. t. 2.

Bot. mag. t. 1138.

Aubl. guian. 1. t. 147.

Aubl. guian. 1. t. 148.

Rœm. arch. 1. t. 6.

Tribus X. DETARIEÆ. DC. prodr. 2. p. 521.

DETARIUM. DC. DETARIUM. Decandria Monogynia.
senegalensis. DC. (gr.) Senegal. Senegal. 1822. S. h.

ORDO LXXV.

CHRYSOBALANEÆ. Brown cong. p. 14.

CHRYSOBA'LANUS.	DC.	COCOA PLUM.	Icosandria Monogynia.	L.			
1	Icáco.	DC.	(wh.) West Indian.	W. Indies.	1752.	4. 6.	S. h.
2	ellipticus.	DC.	(wh.) Pigeon plum.	Sierra Leone.	1822.	S. h.
3	oblongifolius.	DC.	(wh.) American.	Georgia.	1812.	4. 5.	G. h.
PARIN'ARIUM.	DC.	PARIN'ARIUM.	Dodecandria Monogynia.				
1	excésum.	G.D.	(wh.) rough plum.	Sierra Leone.	1822.	S. h.
2	macrophy'lum.	G.D.	(w.) gingerbread plum.	—	—	—	S. h.
GRANG'ERIA.	DC.	GRANG'ERIA.	Dodecandria Monogynia.				
borbónica.	DC.	(wh.) Bourbon.	Bourbon.	1822.	S. h.	Lam. ill. t. 427.
HIRTE'LLA.	DC.	HIRTE'LLA.	Tri-Dodecandria Monogynia.				
1	triándra.	DC.	(wh.) triandrous.	W. Indies.	1782.	S. h.
ameríicana.	Jacq.	non Aubl.	paniculata.	R.S.			Jacq. amer. t. 8.
2	glandulosa.	DC.	(wh.) glandular.	Brazil.	1827.	S. h.
3	racemosa.	DC.	(bl.) racemed.	Guiana.	1782.	S. h.
ameríicana.	Aubl.	non Jacq.					Aubl. guian. I. t. 98.
4	oblongifolia.	DC.	(ro.) oblong-leaved.	New Spain.	1826.	S. h.
5	nemorosa.	DC.	(ye.) wood.	Brazil.	1825.	S. h.

ORDO LXXVI.

AMYGDALINÆ. D. Don prodr. flor. nepal. p. 239.

AMY'GDALUS.	K.S.	ALMOND.	Icosandria Monogynia.	L.			
1	nàna. w.	(pi.)	common dwarf.	Russia.	1683.	3. 4.	H. h.
2	geórgica.	Df.	(pi.) Georgian.	Georgia.	1820.	—	H. h.
3	campéstris.	Bes.	(pi.) Besser's.	Poland.	1829.	—	H. h.
			Besseriana.	Schott.			
4	orientalis.	w.	(ro.) silvery-leaved.	Levant.	1756.	—	F. h.
		argéntea.	Lam.				
5	incána.	w.	(ro.) woolly.	Caucasus.	—	H. h.
6	prostrata.	K.S.	(pi.) prostrate.	Crete.	1802.	5.	H. h.
		Prúnus prostrata.	B.R.				
7	pùmila.	w.	(pi.) double dwarf.	Africa.	1683.	5. 6.	H. h.
		Pridius sinénisis.	P.S. Prúnus japonica,	B.R. is a very			Bot. mag. t. 2176.
				different plant, and a true Cérasus.			
8	sibírica.	L.C.	(pi.) Siberian.	Siberia.	1820.	3. 4.	H. h.
9	communis.	DC.	(pi.) common.	Barbary.	1548.	—	H. 5.
a	amara.	DC.	(pi.) bitter.	—	—	—	H. 5.
β	dulcis.	DC.	(bh.) sweet.	—	—	—	H. 5.
γ	frágilis.	DC.	(bh.) tender.	—	—	Lam. ill. t. 430. f. 2.
δ	macrocarpa.	DC.	(pi.) large-fruited.	—	—	Nois. jard. fr. t. 3. f. 2.
ε	persicoides.	DC.	(pi.) Almond Peach.	Hybrid.	—	—	Bot. reg. t. 1160.
10	cochinchinensis.	DC.	(pi.) Chinese.	Cochinchina.	1825.	—	H. 5.
11	Pérsica.	w.	(pi.) common Peach.	Persia.	1562.	4. 5.	Nois. j. fr. 1-16.22-35. ic.
	β	flóre pleno.	(pi.) double-blossomed.	—	—	—	H. 5.
	δ	nàna.	(pi.) dwarf.	China.	1826.	—	H. h.
	ε	nectarina.	(pi.) Nectarine.	Persia.	1562.	—	Nois. jard. fr. t. 20-21.

ARMENIACA. DC.	APRICOT.	Icosandria Monogynia.						
1 <i>vulgàris</i> . DC.	(wh.) common.	Levant.	1548.	2. 4.	H. 5.			Lam. ill. t. 431.
α <i>ovalifòlia</i> . DC.	(wh.) oval-leaved.				H. 5.			Nois. jar. fr. t. 1-2.f.1-2.
β <i>cordifòlia</i> . DC.	(wh.) heart-leaved.				H. 5.			— t. 1.f.3.t.2.f.3.
2 <i>dasyçarpa</i> . DC.	(wh.) black-fruited.		3. 4.	H. 5.			Bot. reg. t. 1243.
<i>Prunus dasycarpa</i> . B.R.					H. $\frac{1}{2}$.			
β <i>persicifòlia</i> . DC.(wh.)	Peach-leaved.				H. $\frac{1}{2}$.			Du. ar. ed.no.5.t.52.f.1.
3 <i>sibirica</i> . DC.	(pi.) Siberian.	Siberia.	1788.	4.	H. $\frac{1}{2}$.			Pall. ross. I. t. 8.
4 <i>brigantia</i> . DC.	(wh.) spiny.	S. Europe.			H. $\frac{1}{2}$.			Duh. arb.ed.n. 5. t. 59.
PRUNUS. DC.	PLUM-TREE.	Icosandria Monogynia. L.						
1 <i>spinosa</i> . DC.	(wh.) Sloe-tree.	Britain.		3. 4.	H. $\frac{1}{2}$.			Eng. bot. v. 12. t. 842.
α <i>vulgàris</i> . DC.	(wh.) common.				H. $\frac{1}{2}$.			Duham. ed.n.5.t.54.f.1.
β <i>microcarpa</i> . DC.(wh.)	small-fruited.				H. $\frac{1}{2}$.			
γ <i>macrocarpa</i> . DC.(wh.)	large-fruited.				H. $\frac{1}{2}$.			
δ <i>ovata</i> . DC.	(wh.) oval-leaved.				H. $\frac{1}{2}$.			
2 <i>insititia</i> . DC.	(wh.) Bullace-tree.	Britain.		4.	H. $\frac{1}{2}$.			
3 <i>candidans</i> . DC.	(wh.) snowy.		1818.	4. 5.	H. $\frac{1}{2}$.			Blackw. herb. t. 494.
4 <i>Cocomilia</i> . DC.	(wh.) twin-flowered.	Calabria.	1826.		H. $\frac{1}{2}$.			Eng. bot. v. 12. t. 841.
5 <i>maritima</i> . DC.	(wh.) sea-side.	N. America.	1800.		H. $\frac{1}{2}$.			Bot. reg. t. 1135.
6 <i>pubescens</i> . DC.	(wh.) pubescent.		1819.		H. $\frac{1}{2}$.			
7 <i>divaricata</i> . DC.	(wh.) spreading.	Caucasus.	1828.		H. $\frac{1}{2}$.			
8 <i>cerasifera</i> . s.s.	(wh.) Myrobalan.	N. America.	1629.	4. 5.	H. 5.			
9 <i>pyramidalis</i> . DC.	(wh.) pyramidal.	Europe.		4.	H. 5.			Duham. arb. 2.t.2.f.15.
10 <i>domestica</i> . DC.	(wh.) common.	England.			H. 5.			— 2. t. 15—20.
α <i>armenioides</i> . DC.(wh.)	Apricot-like.				H. 5.			Eng. bot. v. 25. t. 1783.
β <i>Claudiæna</i> . DC.	(wh.) Reine Claude.				H. 5.			Duham. arb. 2.t.13—14.
γ <i>damascena</i> . DC.(wh.)	Damson.				H. 5.			— 2. t. 11—12.
δ <i>turonensis</i> . DC.	(wh.) Monsieur hatif, &c.				H. 5.			— 2. n. 13. t. 5.
ϵ <i>Juliæna</i> . DC.	(wh.) Saint Julien.				H. 5.			— 2.n.16.t.20.f.1.
ζ <i>Cathartinea</i> . DC.(wh.)	Saint Catherine.				H. 5.			Duh. ed. nov.5.t.54.f.2.
η <i>Aubertiæna</i> . DC.(wh.)	Imperatrice.				H. 5.			Duham. arb. 2. f. 1.
CERASUS. DC.	CHERRY TREE.	Icosandria Monogynia.						Nois. jar. fr. t. 57-58.f.4.
1 <i>ávium</i> . DC.	(wh.) small-fruited.	Britain.		4. 5.	H. 5.			Blackw. t. 425.
α <i>sylvestris</i> .	(wh.) wood.				H. 5.			Du.ed.no.1.n.2.t.4.f.D.
β <i>macrocarpa</i> . DC.(wh.)	large-fruited.				H. 5.			— 5. n. 3.t.4.f.B.C.
γ <i>pálida</i> . DC.	(wh.) pale-fruited.				H. 5.			
δ <i>multiplæx</i> . DC.	(wh.) double-blossomed.				H. 5.			
2 <i>duracina</i> . DC.	(wh.) hard-fruited.	Europe.			H. 5.			
α <i>cordigera</i> . DC.	(wh.) Bigarreaux.				H. 5.			
β <i>obtusata</i> . DC.	(wh.) blunt-fruited.				H. 5.			
γ <i>mamillaris</i> . DC.	(wh.) large-leaved.				H. 5.			
3 <i>juliæna</i> . DC.	(wh.) early.	S. Europe.			H. 5.			Nois. jard. fr. t. 6. f. 1.
β <i>Heaumiana</i> . DC.(wh.)	long-leaved.				H. 5.			Duh.ed.no.5.t.18.f.B.
γ <i>pendula</i> . DC.	(wh.) pendulous-branch'd.				H. 5.			— 5. n. 12. t. 16. f.A.
4 <i>Caproniana</i> . DC.	(wh.) common.				H. 5.			— 5. n. 5. t. 15. f.A.
α <i>Montmorencyana</i> . DC.	May Duke, &c.				H. 5.			— n. 29. t. 19.
β <i>pallescens</i> . DC.	(wh.) pale-fruited.				H. 5.			
γ <i>gobéttæ</i> . DC.	(wh.) Kentish.				H. 5.			Duh.ed.no.5.n.14.t.7.
δ <i>polyg'yna</i> . DC.	(wh.) polygynous.				H. 5.			— n. 18. t. 11.
ϵ <i>multiplæx</i> . DC.	(wh.) semi-double.				H. 5.			— t. 12. f. A.
ζ <i>persiciflora</i> . DC.	(wh.) double-flowered.				H. 5.			Duham. arb. 4. t. 3.
κ <i>variegata</i> . DC.	(wh.) striped-leaved.				H. 5.			— p. 173. t. 5.
θ <i>griottæ</i> . DC.	(wh.) globular-fruited.				H. 5.			
ι <i>cordigera</i> . DC.	(wh.) red-heart.				H. 5.			Duh.ed.nov.n.21.t.14.
5 <i>sempervirens</i> . DC.	(wh.) late-fruited.	N. America.		5. 6.	H. $\frac{1}{2}$.			Duh.ed.n.5.t.16.f.1.
6 <i>chamæcérasus</i> . DC.	(wh.) bastard.	Germany.	1597.	5.	H. $\frac{1}{2}$.			Wats. dend. brit. 131.
7 <i>persicifolia</i> . DC.	(wh.) Peach-leaved.	N. America.	1818.	4. 5.	H. $\frac{1}{2}$.			Jacq. ic. 1. t. 90.
8 <i>pumila</i> . DC.	(wh.) dwarf.			1756.	5.	H. $\frac{1}{2}$.		
9 <i>pygmæa</i> . DC.	(wh.) small.			1818.		H. $\frac{1}{2}$.		
10 <i>nigra</i> . DC.	(wh.) black-fruited.	Canada.	1773.	5. 6.	H. $\frac{1}{2}$.			Mill. ic. t. 89. f. 2.
11 <i>borealis</i> . DC.	(wh.) northern.	N. America.	1818.		H. $\frac{1}{2}$.			Bot. mag. t. 1117.
12 <i>depressa</i> . DC.	(wh.) depressed.			1805.	5.	H. $\frac{1}{2}$.		
13 <i>Susquehannæa</i> . W.en.	(wh.) glaucous-ld.	N. America.	1800.		H. $\frac{1}{2}$.			
14 <i>hyemalis</i> . DC.	(wh.) mountain.			1805.		H. $\frac{1}{2}$.		
15 <i>Chicasa</i> . DC.	(wh.) Chicassaw.			1806.	4. 5.	H. $\frac{1}{2}$.		
16 <i>pubescens</i> . DC.	(wh.) pubescent.			1822.	5.	H. $\frac{1}{2}$.		
17 <i>japonica</i> . DC.	(pi.) Japan.	Japan.	1810.	3. 5.	H. $\frac{1}{2}$.			
18 <i>serrulata</i> . H.C.	(bh.) saw-leaved.	China.	1822.	4. 5.	H. $\frac{1}{2}$.			
19 <i>salicina</i> . H.C.	(bh.) Willow-like.				H. $\frac{1}{2}$.			Bot. reg. t. 27.

20	pseñudo-cerasus. <i>HC.</i> (<i>wh.</i>) Chinese. <i>paniculata.</i> B.R. <i>non</i> Thunb.	China.	1820.	4. 5.	F. $\frac{1}{2}$.	Bot. reg. t. 800.
21	pensylvánica. <i>DC.</i> (<i>wh.</i>) Pensylvanian.	N. America.	1773.	5.	H. $\frac{1}{2}$.	Willd. arb. t. 3. f. 3.
22	Maháleb. <i>DC.</i> (<i>wh.</i>) perfumed.	S. Europe.	1714.	4. 5.	H. $\frac{1}{2}$.	Jacq. aust. 3. t. 227.
23	Pádus. <i>DC.</i> (<i>wh.</i>) Bird.	Britain.	—	H. $\frac{1}{2}$.	Eng. bot. v. 20. t. 1383.
24	rúbra. <i>w.</i> (<i>wh.</i>) red-fruited.	England.	—	H. $\frac{1}{2}$.	Willd. arb. t. 4. f. 2.
25	virginiana. <i>DC.</i> (<i>wh.</i>) Virginian.	N. America.	1724.	5. 6.	H. $\frac{1}{2}$.	— t. 5. f. 1.
26	serotina. <i>DC.</i> (<i>wh.</i>) American Bird.	—	1629.	—	H. $\frac{1}{2}$.	Wats. dend. brit. 48.
27	occidentális. <i>DC.</i> (<i>wh.</i>) West Indian.	Jamaica.	1784.	12. 1.	S. $\frac{1}{2}$.	Catesb. car. 2. t. 94?
28	lusitánica. <i>DC.</i> (<i>wh.</i>) Portugal Laurel. Portugal.	—	1648.	6.	H. $\frac{1}{2}$.	Mill. ic. t. 196. f. 1.
29	Híxa. W. en. (<i>wh.</i>) Teneriffe.	Teneriffe.	1824.	—	F. $\frac{1}{2}$.	Blackw. herb. t. 512.
30	Laurocérasus. <i>DC.</i> (<i>wh.</i>) Laurel.	Levant.	1629.	4. 5.	H. $\frac{1}{2}$.	
31	caroliniána. <i>DC.</i> (<i>wh.</i>) evergreen.	Carolina.	1759.	5.	H. $\frac{1}{2}$.	

ORDO LXXVII.

POMACEÆ. D. *Don prodr. flor. nepal. p. 236.*

ME'SPILUS. K.S.		ME'SPILUS. Icosandria Di-Pentagynia.				
1	germánica. <i>DC.</i> (<i>wh.</i>) Medlar.	England.	5. 7.	H. 5.	Eng. bot. v. 22. t. 1523.
α	<i>sylvestris. DC.</i> (<i>wh.</i>) wild.	—	—	—	H. 5.	
β	<i>strícta. DC.</i> (<i>wh.</i>) upright.	—	—	—	H. 5.	Duham. arb. fr. 1. t. 3.
γ	<i>diffusa. DC.</i> (<i>wh.</i>) Dutch.	—	—	—	H. 5.	Exot. bot. 1. t. 18.
2	grandiflóra. s.s. (<i>wh.</i>) large-flow red. <i>Smithii. DC.</i>	S. Europe.	1800.	5. 6.	H. 5.	Schmidt arb. t. 90.
3	Pyracántha. s.s. (<i>wh.</i>) evergreen thorn.	—	1629.	5.	H. $\frac{1}{2}$.	
4	crenuláta. <i>D.P.</i> (<i>wh.</i>) crenulate.	Nepaul.	1818.	6. 7.	H. $\frac{1}{2}$.	
5	mexicána. <i>DC.</i> (<i>wh.</i>) Mexican.	Mexico.	1824.	5. 6.	F. $\frac{1}{2}$.	
6	Crús-gálli. <i>L.</i> (<i>wh.</i>) Cockspur-thorn. N. America.	—	1691.	5. 6.	H. 5.	Wats. dend. brit. t. 56.
7	pyracanthifolia. (<i>wh.</i>) Pyracantha-leaved.— <i>lúcida.</i> Dum-Cours. bot. cult. ed. 2. v. 5. p. 448. <i>non</i> Miller.	—	—	—	H. 5.	
8	salicifolia. <i>L.C.</i> (<i>wh.</i>) Willow-leaved.	—	—	—	H. 5.	
9	lineáris. <i>DF.</i> (<i>wh.</i>) linear-leaved.	—	—	—	H. 5.	
10	nána. Dum-C.	—	dwarf.	—	H. $\frac{1}{2}$.	
11	spinosissima. <i>L.C.</i> (<i>wh.</i>) long-spined.	S. Europe.	—	—	H. $\frac{1}{2}$.	Lodd. bot. cab. 1100.
12	punctátâ. <i>W. en.</i> (<i>wh.</i>) spotted-fruited. <i>N. America.</i>	1746.	—	—	H. 5.	Wats. dend. brit. t. 57.
α	<i>rúbra. Ph.</i> (<i>wh.</i>) red-fruited.	—	—	—	H. $\frac{1}{2}$.	
β	<i>aurea. Ph.</i> (<i>wh.</i>) yellow-fruited.	—	—	—	H. $\frac{1}{2}$.	
13	ellíptica. <i>W. en.</i> (<i>wh.</i>) oval-leaved.	—	1765.	5.	H. 5.	
14	prunifolia. s.s. (<i>wh.</i>) Plum-leaved.	—	1812.	5. 6.	H. 5.	
15	prunellifolia. <i>DC.</i> (<i>wh.</i>) Sloe-leaved.	—	1824.	—	H. $\frac{1}{2}$.	
16	latifolia. <i>Poir.</i> (<i>wh.</i>) broad-leaved.	N. America.	—	—	H. 5.	
17	parvifolia. <i>W. en.</i> (<i>wh.</i>) Gooseberry-leav'd.	—	1704.	—	H. $\frac{1}{2}$.	Wats. dend. brit. 65.
	<i>anthocárpos. s.s.</i> (<i>wh.</i>) <i>Cratægus parvifolia. DC. tomentosa. L.</i> Trew. ehr. t. 17.					
18	ovalifolia. <i>DC.</i> (<i>wh.</i>) oval-leaved.	—	—	—	H. $\frac{1}{2}$.	
19	víridis. <i>DC.</i> (<i>wh.</i>) smooth green.	—	—	—	H. $\frac{1}{2}$.	
20	pauciflora. <i>Poir.</i> (<i>wh.</i>) few-flowered.	Switzerland.	1818.	—	H. $\frac{1}{2}$.	
21	lúcida. <i>DC.</i> (<i>wh.</i>) glossy-leaved.	—	—	—	H. $\frac{1}{2}$.	
22	apiifolia. s.s. (<i>wh.</i>) Parsley-leaved.	—	1812.	—	H. $\frac{1}{2}$.	
23	pyrifolia. s.s. (<i>wh.</i>) Pear-leaved.	—	1765.	6.	H. 5.	
24	glandulosa. s.s. (<i>wh.</i>) hollow-leaved. <i>N. America.</i>	1750.	5. 6.	H. 5.		
	<i>Cratægus sanguinea. Pall. ross. 1. t. 11.</i>					
25	spatulata. s.s. (<i>wh.</i>) spatula-leaved.	—	1806.	—	H. 5.	Lodd. bot. cab. t. 1261.
26	coccinea. s.s. (<i>wh.</i>) scarlet-fruited.	—	1683.	4. 5.	H. 5.	Wats. dend. brit. t. 62.
27	purpúrea. <i>W.D.</i> (<i>wh.</i>) purple-branch'd.	—	—	—	H. 5.	— t. 60.
28	cordata. s.s. (<i>wh.</i>) Maple-leaved.	—	1738.	5. 6.	H. 5.	— t. 63.
	<i>Cratægus cordata. Bot. reg. t. 1151.</i>					
29	lobata. <i>Poir.</i> (<i>wh.</i>) lobed-leaved.	—	1818.	—	H. 5.	
30	florentina. <i>Be.</i> (<i>wh.</i>) Florentine.	Florence.	1824.	—	H. 5.	Lodd. bot. cab. t. 1021.
31	nígra. s.s. (<i>wh.</i>) black-fruited.	Hungary.	1818.	—	H. 5.	

POMACEÆ.

32 pentágyna. s.s. (<i>wh.</i>) five-styled.	Hungary.	1818.	5. 6.	H. 5.	
33 fláva. W. en. (<i>wh.</i>) yellow-berried.	N. America.	1724.	—	H. 5.	Wats. dend. brit. t. 59.
34 sphaer. Poir. (<i>wh.</i>) cloven-leaved.	—	1818.	—	H. 5.	
35 Oliveriana. DC. (<i>wh.</i>) Oliver's.	Levant.	—	—	H. 5.	
36 Poiretiána. DC. (<i>wh.</i>) Poiret's. <i>linearis</i> . Poir. <i>non</i> P.S.	N. America.	—	H. 5.	
37 dissécta. Dum. (<i>wh.</i>) cut-leaved.	Persia.	1828.	H. 5.	
38 Oxyacántha. s.s. (<i>wh.</i>) Hawthorn. <i>Cratægus Oxyacántha</i> . DC.	Britain.	5. 6.	H. 5.	Eng. bot. v. 35. t. 2504.
β major. (<i>wh.</i>) great-fruited.	—	—	—	H. 5.	
γ àurea. (<i>wh.</i>) yellow-berried.	—	—	—	H. 5.	
39 præcox.	(<i>wh.</i>) Glastonbury.	—	—	3. 5.	H. 5.
40 oxyacanthoïdes. B.R. Hawthorn-like.	S. Europe.	5. 6.	H. 5.	
β multiplex.	(<i>wh.</i>) double-flowered.	—	—	H. 5.	
41 élégans. Poir. (<i>ro.</i>) rose-coloured.	—	—	—	H. 5.	
β ruberrima.	(<i>re.</i>) dark-red.	—	1827.	—	H. 5.
42 eriocárpa. Lind. (<i>wh.</i>) woolly-fruited.	Britain.	—	—	H. 5.	
43 monógyna. Pal. (<i>wh.</i>) one-styled.	Siberia.	—	—	H. 5.	
44 heterophylla. DC. (<i>wh.</i>) various-leaved.	Levant?	1818.	—	H. 5.	
45 Azaròlus. s.s. (<i>wh.</i>) Azarole.	S. Europe.	1640.	—	H. 5.	
46 Arónia. W. en. (<i>wh.</i>) Aronian.	Levant.	—	—	H. 5.	
47 orientális. Poir. (<i>wh.</i>) oriental.	—	1824.	—	H. 5.	
48 tanacetifolia. A.R. (<i>wh.</i>) Tansy-leaved.	—	1789.	—	H. $\frac{1}{2}$.	
49 odoratissima. A.R. (<i>wh.</i>) sweet-scented.	Crimea.	—	—	H. $\frac{1}{2}$.	
50 Celsiana. W. en. (<i>wh.</i>) Cels's.	Levant.	1822.	—	H. $\frac{1}{2}$.	
51 laciniata. DC. (<i>wh.</i>) jagged.	Sicily.	—	—	H. $\frac{1}{2}$.	
52 melanocárpa. s.s. (<i>wh.</i>) black-berried.	Tauria.	1818.	—	H. $\frac{1}{2}$.	
53 pectinata. L. en. (<i>wh.</i>) pectinated.	Persia.	1824.	—	H. $\frac{1}{2}$.	
54 flabellata. L.C. (<i>wh.</i>) fan-leaved.	—	1823.	—	H. $\frac{1}{2}$.	
55 pruinosa. Lk. (<i>wh.</i>) frosted.	—	1826.	—	H. $\frac{1}{2}$.	
56 succulenta. Scr. (<i>wh.</i>) succulent.	—	—	—	H. $\frac{1}{2}$.	
57 subvillosa. Scr. (<i>wh.</i>) villous.	—	1827.	—	H. $\frac{1}{2}$.	
58 punicea. F. crimson.	Dahuria.	—	—	H. $\frac{1}{2}$.	
RAPHIOLEPIIS. DC. RAPHIOLEPIIS.	Icosandria	Digynia.			
1 indica. DC. (<i>wh.</i>) Indian.	China.	1806.	5. 8.	G. $\frac{1}{2}$.	Bot. mag. t. 1726.
<i>Cratægus indica</i> . B.M.					
2 phæostemon. DC. (<i>wh.</i>) red-stamen'd.	—	1818.	—	G. $\frac{1}{2}$.	Bot. reg. t. 468.
<i>indica</i> . Bot. reg. <i>non</i> Bot. mag.					
3 rúbra. DC. (<i>wh.</i>) red-stemmed.	—	1806.	—	G. $\frac{1}{2}$.	Lindl. coll. t. 3.
4 salicifolia. DC. (<i>wh.</i>) Willow-leaved.	—	1820.	—	G. $\frac{1}{2}$.	Bot. reg. t. 652.
PHOTINIA. DC.	PHOTINIA.	Icosandria	Digynia.		
1 serrulata. DC. (<i>wh.</i>) smooth-leaved.	China.	1804.	4. 7.	H. $\frac{1}{2}$.	Bot. mag. t. 2105.
<i>Cratægus glabra</i> . B.M.					
2 arbutifolia. DC. (<i>wh.</i>) Arbutus-leaved.	California.	1796.	6. 8.	H. $\frac{1}{2}$.	Bot. reg. t. 491.
3 integrifolia. DC. (<i>wh.</i>) entire-leaved.	Nepaul.	1820.	5. 6.	H. $\frac{1}{2}$.	
<i>Pyrus integréríma</i> . D.P.					
4 dúbia. DC. (<i>wh.</i>) doubtful.	—	1822.	—	H. $\frac{1}{2}$.	Linn. trans. 13. t. 10.
<i>Mespilus tinctoria</i> . D.P. <i>Mespilus bengalensis</i> . Roxb.					
ERIOB'OTRYA. DC. LOQUAT.	Icosandria	Pentagynia.			
1 japonica. DC. (<i>wh.</i>) Japan.	Japan.	1787.	9. 10.	H. $\frac{1}{2}$.	Bot. reg. t. 365.
<i>Mespilus japonica</i> . Vent. malm. t. 19.					
2 elliptica. DC. (<i>wh.</i>) elliptic-leaved.	Nepaul.	1822.	H. $\frac{1}{2}$.	
<i>Mespilus Cuila</i> . D.P.					
COTONEA'STER. DC. COTONEA'STER.	Icosandria	Di-Pentagynia.			
1 vulgáris. DC. (<i>wh.</i>) common.	Europe.	1656.	4. 5.	H. $\frac{1}{2}$.	Flor. dan. t. 112.
<i>Mespilus Cotoneaster</i> . L.					
2 affinis. DC. (<i>bh.</i>) likened.	Nepaul.	1820.	6. 7.	H. $\frac{1}{2}$.	
3 tomentosa. DC. (<i>wh.</i>) Quince-leaved.	Europe.	1759.	4. 5.	H. $\frac{1}{2}$.	Wats. dend. brit. t. 55.
<i>Mespilus eriocarpa</i> . W.D.					
4 denticulata. K.S. (<i>wh.</i>) tooth-leaved.	Mexico.	1826.	F. $\frac{1}{2}$.	
5 laxiflora. J.F. (<i>wh.</i>) loose-flowered.	Nepaul.	1820.	5. 7.	H. $\frac{1}{2}$.	H. et B. n. ge. am. 6. t. 556.
6 frígida. B.R. (<i>wh.</i>) mountain.	—	1823.	4. 5.	H. $\frac{1}{2}$.	Bot. reg. t. 1229.
7 acuminata. DC. (<i>bh.</i>) pointed-leaved.	—	1820.	4. 6.	H. $\frac{1}{2}$.	Linn. trans. v. 13. t. 9.
<i>Mespilus acuminata</i> . Lodd. bot. cab. t. 919.					
8 rotundifolia. B.R. (<i>wh.</i>) round-leaved.	—	—	—	H. $\frac{1}{2}$.	Bot. reg. t. 1187.
<i>microphylla</i> β <i>Uva úrsi</i> . B.R.					
9 microphylla. B.R. (<i>wh.</i>) small-leaved.	—	—	—	H. $\frac{1}{2}$.	Bot. reg. t. 1114.
10 buxifolia. B.R. (<i>wh.</i>) Box-leaved.	—	1824.	—	H. $\frac{1}{2}$.	

AMELA'NCHIER. DC. AMELA'NCHIER. Icosandria Pentagynia.

1 vulgàris. DC.	(wh.) common.	S. Europe.	1596.	4. 5.	H. h.
Mésphilus Amelánchier.	Jacq. aust. t. 300.	Pyrus Amelánchier. W.			
2 Botryàpium. DC.	(wh.) snowy.	N. America.	1746.	—	H. h.
Mésphilus canadénis. L.	Pyrus Botryàpium. w.	Arónia Botryàpium. P.S.			
3 ovalis. DC.	(wh.) oval-leaved.		1800.	—	H. h.

4 sanguinea. DC.	(wh.) red-branched.		1812.	—	H. h.
AR'ONIA. P.S.	AR'ONIA.	Icosandria. Di-Pentagynia.			

1 arbutifòlia. N.	(wh.) red-berried.	N. America.	1700.	5. 6.	H. h.
pyrifòlia. P.S.	Pyrus arbutifòlia. Ph.				

2 grandifòlia.	(wh.) large-leaved.		—	H. h.
Pyrus grandifòlia. B.R.					

3 floribunda.	(wh.) many-flowered.		—	H. h.
4 xanthocárpa. L.C.	(wh.) yellow-fruited.		1820.	—	H. h.

5 melanocárpa. N.	(wh.) black-fruited.		1700.	—	H. h.
6 chamæmélipilus. P.S. (wh.)	Bastard Quince. Pyrenees.	1683.	—	H. h.	

Mésphilus chamæmélipilus. L.	Cratagus chamæmélipilus. Jacq. aust. t. 231.				
P'YRUS. DC.	PEAR & APPLE. Icosandria Di-Pentagynia.				

1 communis. DC.	(wh.) common Pear.	England.	4.	H. 5.
α Achras. DC.	(wh.) pointed-leaved.		—	H. 5.

β pyraster. DC.	(wh.) round-leaved.		—	H. 5.
γ sativa. DC.	(wh.) cultivated.		—	H. 5.

2 Bollwylleriana. DC.	(wh.) woolly-leaved.	Germany.	1786.	4. 5.	H. 5.
3 salivifolia. DC.	(wh.) Sage-leaved.	France.	1824.	—	H. 5.

4 amygdaliformis. DC.	(wh.) Almond-like.		1820.	—	H. 5.
5 elæagnifolia. DC.	(wh.) Oleaster-leaved.	Tauria.	1818.	—	H. 5.

6 salicifolia. DC.	(wh.) Willow-leaved.	Siberia.	1780.	5. 6.	H. 5.
7 crenata. D.P.	(wh.) crenated.	Nepaul.	1820.	—	H. 5.

8 sináica. DC.	(wh.) Mount Sinai.	Levant.	—	5.	H. 5.
pérsica. P.S.					

9 nivalis. DC.	(wh.) white-leaved.	Austria.	—	H. 5.
10 Pashia. D.P.	(wh.) rough-fruited.	Nepaul.	1825.	5. 6.	H. 5.

11 Nüssia. D.P.	(wh.) small-fruited.		1823.	—	H. 5.
12 sinensis. B.R.	(wh.) Chinese.	China.	—	—	H. 5.

13 acerba. DC.	(bh.) sour.	Europe.	4. 5.	H. 5.
14 Málus. DC.	(bh.) common Apple.	Britain.	—	H. 5.

15 præcox.	(bh.) Paradise.	Russia.	1784.	4.	H. 5.
16 dioica. DC.	(st.) linear petal'd.	4. 5.	H. 5.

17 astracanica. DC.	(bh.) Astracan.	Astracan.	1824.	—	H. 5.
18 spectabilis. DC.	(pi.) Chinese Apple.	China.	1780.	5.	H. 5.

19 prunifolia. DC.	(bh.) Siberian Crab.	Siberia.	1758.	4. 5.	H. 5.
20 tomentosa. DC.	(bh.) woolly.		1824.	—	H. 5.

21 rubicunda. DC.	(bh.) red-fruited.		1820.	—	H. 5.
22 baccata. DC.	(bh.) small Crab.	Siberia.	1784.	—	H. 5.

23 coronaria. DC.	(pi.) sweet-scented.	Virginia.	1724.	5.	H. 5.
24 angustifolia. DC.	(bh.) narrow-leaved.	N. America.	1750.	—	H. h.

25 pumila. Mil.	(bh.) dwarf evergreen.	4. 5.	H. h.
26 Aria. DC.	(wh.) white-beam.	Britain.	5. 6.	H. 5.

27 alpina. W. en.	(wh.) Alpine.	Europe.	1820.	—	H. 5.
longifolia. Duham.					

28 intermédia. DC.	(wh.) Swedish.	Sweden.	1789.	4. 5.	H. 5.
29 latifolia. P.S.	(wh.) broad-leaved.	France.	5. 6.	H. 5.

30 edulis. W. en.	(wh.) eatable-fruited.	Europe.	1816.	—	H. 5.
31 lanata. D.P.	(wh.) woolly.	Nepaul.	1820.	—	H. 5.

32 torminalis. DC.	(wh.) Wild Service.	England.	4. 5.	H. 5.
33 auriculata. DC.	(wh.) eared-leaved.	Egypt.	1828.	—	H. 5.

34 pinnatifida. DC.	(wh.) pinnatifid.	England.	5. 6.	H. 5.
35 arbúscula. Poir.	(wh.) little tree.	Germany.	—	H. 5.

36 microcarpa. DC.	(wh.) small-berried.	N. America.	—	H. 5.
37 aucuparia. DC.	(wh.) Mountain Ash.	Britain.	—	H. 5.

38 americana. DC.	(wh.) purple-berried.	Canada.	1782.	—	H. 5.
39 sorbifolia. W.D.	(wh.) Service-leaved.	5.	H. 5.

40 Sórbus. DC.	(wh.) Service-tree.	England.	5. 6.	H. 5.
doméstica. E.B.	Sórbus doméstica. L.				

41 lanuginosa. DC.	(wh.) woolly-leaved.	Hungary.	1820.	—	H. 5.
spuria. DC.	(bh.) spurious.	Hybrid.	—	H. 5.

hybrida. Mœnch.	Weis. t. 6.	Sórbus spuria. P.S.			
			N		

Schmidt. arb. t. 85.

Schmidt. arb. t. 84.

Bot. reg. t. 1171.

Mill. ic. t. 109.

Bot. reg. t. 1154.

Bot. reg. t. 1006.

Schmidt. arb. t. 86.

— t. 87.

Eng. bot. v. 25. t. 1784.

Gärt. fr. t. 87. f. 2.

Lodd. bot. cab. t. 1009.

Jacq. aust. 2. t. 107.

Bot. reg. t. 1248.

Flor. dan. t. 1101.

Eng. bot. v. 3. t. 179.

Bot. mag. t. 267.

Mill. ic. 2. t. 269.

Wats. dend. brit. 132.

Pall. fl. ross. t. 10.

Bot. reg. t. 651.

Wats. dend. brit. t. 52.

Eng. bot. v. 5. t. 298.

Eng. bot. v. 33. t. 23-31.

Eng. bot. v. 5. t. 337.

Wats. dend. brit. t. 54.

— t. 53

Eng. bot. v. 5. t. 350.

CYD'ONIA. DC.	QUINCE. Icosandria Pentagynia.						
1 vulgàris. DC. (wh.) common.	S. Europe.	1573.	5. 6.	H. 5.		Jacq. aust. 4. t. 342.	
α malifòrmis. DC. (wh.) Apple-fruited.	—	—	—	H. 5.		Duha. arb. fr. 1. p. 195. ic.	
β lusitànica. DC. (wh.) large-fruited.	—	—	—	H. 5.		Blackw. herb. t. 137.	
γ oblonga. DC. (wh.) oblong-fruited.	—	—	—	H. 5.		Bot. reg. t. 905.	
2 sinénsis. DC. (wh.) Chinese.	China.	1815.	4. 5.	H. 5.		Bot. mag. t. 692.	
3 japónica. DC. (sc.) Japan.	Japan.	1796.	1. 12.	H. $\frac{1}{2}$.		Lodd. bot. cab. t. 541.	
α rùbra.	(sc.) red-flowered.	—	—	H. $\frac{1}{2}$.			
$P'yrus$ japonica. B.M. $Malus$ japonica. A.R. 462.	(wh.) white-flowered.	—	1812.	—	H. $\frac{1}{2}$.		
β álba.							

ORDO LXXVIII.

ROSACEÆ. D. Don prodr. fl. nepal. p. 229.

Tribus I. ROSEÆ. DC. prodr. 2. p. 596.

R'OSA. DC.	ROSE. Icosandria Polygynia. L.						
1 berberifòlia. W. (ye.)	Berry-ld. Persia.	1790.	6. 8.	H. $\frac{1}{2}$.		Redout. ros. 1. 27. t. 2.	
simplicifòlia. P.L. 101.	<i>Lo'wea berberifòlia</i> . Bot. reg. 1261.						
2 fèrox. L.R.	(pu.) hedgehog. Caucasus.	1796.	6. 8.	H. $\frac{1}{2}$.		Bot. reg. t. 420.	
3 kamchática. L.R. (pu.)	Kamtschatka. Kamtschatka.	1802.	7. 8.	H. $\frac{1}{2}$.	—	t. 419.	
β nitens. B.R.	(pu.) glossy.	1820.	—	H. $\frac{1}{2}$.	—	t. 824.	
4 involucrata. L.R. (wh.)	floral-leaved. China.	1814.	5. 10.	H. $\frac{1}{2}$.	—	t. 739.	
5 clinophylla. R.R. (wh.)	naked-flowered.	1820.	—	H. $\frac{1}{2}$.	Redout. ros. 1. t. 10.		
6 bracteata. L.R. (wh.)	Macartney.	1795.	6. 10.	H. $\frac{1}{2}$.	—	1. t. 6.	
β scabriúscula. L.R. (wh.)	rough Macartney.	—	—	H. $\frac{1}{2}$.	Bot. mag. 1377.		
7 nítida. W.en.	(ro.) dwarf Labrador. N. America.	1773.	7. 8.	H. $\frac{1}{2}$.	Lindl. ros. t. 2.		
8 ràpa. R.R.	(ro.) double Burnet.	1726.	—	H. $\frac{1}{2}$.	Redout. ros. 1. 7. t. 2.		
9 lúcida. W.	(re.) single Burnet.	1773.	—	H. $\frac{1}{2}$.	Miss Lawr. ros. t. 24.		
10 Lindlèyi. s.s.	(ro.) spreading.	1726.	—	H. $\frac{1}{2}$.	Lindl. ros. t. 3.		
láx'a. L.R. nom Retz.							
11 parviflora. w.	(bh.) small-flowered.	1724.	—	H. $\frac{1}{2}$.			
α Single Pensylvanian (bh.)	—	—	—	H. $\frac{1}{2}$.	Miss Lawr. ros. t. 68.		
β Double Pensylvanian (bh.)	—	—	—	H. $\frac{1}{2}$.	—	t. 66.	
12 Woodsii. L.R.	(pi.) Woods's.	1815.	6. 7.	H. $\frac{1}{2}$.			
13 gemella. s.s.	(re.) twin-flowering.	1800.	7. 8.	H. $\frac{1}{2}$.			
14 Lyònii. s.s.	(ro.) Lyon's.	1812.	—	H. $\frac{1}{2}$.			
15 florida. L.C.	(re.) flowering.	N. America.	6. 7.	H. $\frac{1}{2}$.			
16 carolina. L.R.	(re.) Carolina.	1726.	—	H. $\frac{1}{2}$.	Lindl. ros. t. 4.		
17 pensylvanica. L.en.	(re.) Pensylvanian.	—	—	H. $\frac{1}{2}$.			
18 blanda. L.R.	(re.) Hudson's Bay.	1773.	5. 8.	H. $\frac{1}{2}$.	Miss Lawr. ros. t. 27.		
19 fraxinifolia. L.R.	(re.) Ash-leaved.	Newfoundland.	4. 6.	H. $\frac{1}{2}$.	Bot. reg. 458.		
β Red Alpine Rose. (re.)	—	—	—	H. $\frac{1}{2}$.	Miss Lawr. ros. t. 75.		
20 cinnamònea. E.B. (ro.)	Single Cinnamon. England.	5.	H. $\frac{1}{2}$.	Eng. bot. v. 34. t. 2388.		
β flòre semipleno. (ro.)	semidouble.	—	H. $\frac{1}{2}$.	Lindl. ros. t. 5.		
γ fæcundissima. (ro.)	double Cinnamon. Europe.	4. 5.	H. $\frac{1}{2}$.	Flor. dan. t. 688.		
21 majális. L.R.	(ro.) dwarf Cinnamon. S. Europe.	—	H. $\frac{1}{2}$.			
22 frutetòrum. s.s.	(bh.) short pedunc'd. Volhynia.	1820.	5. 6.	H. $\frac{1}{2}$.			
23 ambigua. s.s.	(bh.) ambiguous.	—	H. $\frac{1}{2}$.			
24 macrophylla. L.R. (ro.)	long-leaved.	Nepaul.	1822.	Lindl. ros. t. 6.		
25 americana. L.en.	(ro.) American.	America.	—	—			
26 alpina. L.R.	(re.) Alpine.	Europe.	1683.	6. 7.	H. $\frac{1}{2}$.		
α Alpine Rose. (re.) common.	—	—	—	H. $\frac{1}{2}$.	Bot. reg. 424.		
β pyrenàica. L.R.	(re.) Pyrenean.	—	—	H. $\frac{1}{2}$.	Miss Lawr. ros. t. 30.		
γ pendulina. L.R.	(re.) pendulous.	—	1726.	—	Jacq. schenb. 4. t. 416.		
27 pygm'aea. s.s.	(ro.) pigmy.	Tauria.	—	Miss Lawr. ros. t. 9.		
28 rubella. E.B.	(re.) long-fruited.	England.	7.	Eng. bot. v. 36. t. 2521.		
β melanocarpa. L.R.	(re.) black-fruited.	—	—	H. $\frac{1}{2}$.			

29	muricata. L.en.	(ro.) muricate.	1823.	6. 7.	H. h.	
30	balsamica. s.s.	(ro.) balsamic.	—	—	H. h.	Lindl. ros. t. 7.
31	strícta. L.R.	(ro.) upright.	N. America.	1726.	—	H. h.	Miss Lawr. ros. t. 36.
	Upright Carolina. (ro.)				—	H. h.	
32	suavis. s.s.	(ro.) sweet.	1817.	—	H. h.	Lindl. ros. t. 8.
33	aciculáris. L.R.	(ro.) grey Siberian.	Siberia.	3. 4.	H. h.	
34	pauciflora. Ly.	(ro.) few-flowered.	—	—	H. h.	Bot. reg. 46.
35	sulphurea. w.	(st.) double yellow.	Levant.	1629.	7.	H. h.	Lindl. ros. t. 9.
36	lutescens. Ph.	(st.) yellowish. hispida. Bot. mag. t. 1570.	Siberia.	1780.	5. 6.	H. h.	
37	viminea. L.R.	(ro.) twiggy.	Russia.	1826.	5. 6.	H. h.	
38	spinossissima. E.B. (wh.) Scotch.	Scotch.	Britain.	6. 7.	H. h.	Eng. bot. v. 3. t. 187.
	β revérsa. L.R.	(wh.) reversed prickle.	—	—	H. h.	Bot. reg. 431.
	γ striped-flowered Scotch.	(std.)	—	—	H. h.	Miss Lawr. ros. t. 15.
	δ Red Scotch.	(re.)	—	—	H. h.	— t. 62.
	ε Double Scotch.	(wh.)	—	—	H. h.	— t. 63.
	ζ Tall Scotch.	(wh.)	—	—	H. h.	— t. 19.
	η Marbled Scotch.	(pu.) (wh.)	—	—	H. h.	— t. 78.
	♀ sanguisorbiolia. L.R. (wh.)	Sanguisorba-ld.	—	—	H. h.	
39	altáica. W.en.	(wh.) Pallas's.	Siberia.	—	H. h.	Pall. ros. 2. t. 75.
40	ochroleúca. L.en.	(st.) sulphur-color'd.	1814.	—	H. h.	
41	melanocárpa. s.s.	(wh.) small black fruited.	S.Europe.	1823.	—	H. h.	
42	grandiflora. L.R.	(wh.) large flowered.	Siberia.	5. 6.	H. h.	Bot. reg. 888.
43	myriacántha. DC.	(wh.) numerous-spin'd.	S.France.	1820.	—	H. h.	Lindl. ros. t. 10.
44	involúcta. E.B.	(bh.) Dr. Walker's.	Hebrides.	6. 7.	H. h.	Eng. bot. v. 29. t. 2068.
45	réversa. L.R.	(bh.) reversed.	Hungary.	1817.	—	H. h.	W. et K. hung. 3.t. 264.
46	Biebersteinii. Ly.	(ro.) Bieberstein's.	Caucasus.	1821.	5.	H. h.	
47	Doniána. L.T.	(wh.) Mr. Don's.	Britain.	6. 7.	H. h.	
48	gracilis. L.T.	(ro.) slender.	—	7. 8.	H. h.	Eng. bot. v. 9. t. 583.
	villósa. E.B. non L.						
49	Sabini. L.T.	(ro.) Mr. Sabine's.	—	—	—	H. h.	
50	marginata. L.R.	(ro.) margined.	Germany.	1823.	—	H. h.	
51	corúscans. L.en.	(ro.) glittering.	—	—	H. h.	
52	damascéna. L.R.	(re.) red damask.	Syria?	1575.	6. 7.	H. h.	Miss Lawr. ros. t. 38.
53	centifòlia. L.R.	(ro.) Provins.	Caucasus.	1596.	—	H. h.	— t. 8.
	β muscósula. L.R.	(ro.) single moss.	1724.	—	H. h.	Bot. reg. 53.
	γ multipléx.	(ro.) double moss.	—	—	H. h.	Bot. mag. t. 69.
	δ albiflóra.	(wh.) white moss.	—	—	H. h.	Bot. reg. t. 102.
	ε pomponia. L.R.	(pi.) Pompone.	—	—	H. h.	Redout. ros. 1. 65. t. 21.
	ζ bipinnata. L.R.	(pi.) bipinnate.	Caucasus.	—	H. h.	— 2. 11. t. 4.
54	gállica. L.R.	(re.) officinal.	Europe.	1596.	—	H. h.	Bot. reg. 448.
	β pùmila. L.R.	(re.) dwarf.	Austria.	1773.	—	H. h.	Jacq. aust 2. t. 198.
55	pulchéllea. s.s.	(re.) neat.	France.	—	H. h.	
56	parviflória. L.R.	(pu.) Burgundy.	Europe.	—	H. h.	Bot. reg. t. 452.
57	turbinata. L.R.	(ro.) Frankfort.	1629.	—	H. h.	Miss Lawr. ros. t. 69.
58	villósa. L.T.	(bh.) villous.	Britain.	6.	H. h.	Eng. bot. v. 35. t. 2459.
	móllis. E.B.						
	a single apple bearing.	(bh.) —	—	—	—	H. h.	Miss Lawr. ros. t. 33.
	β cærulea. L.T.	(bh.) blue-stalked.	—	—	—	H. h.	
	γ concavifòlia. L.T.	(bh.) concave-leaved.	Scotland.	—	—	H. h.	
	δ suberécta. L.T.	(bh.) suberect.	Yorkshire.	—	—	H. h.	
	ε double apple bearing.	(pi.)	—	—	—	H. h.	
59	scabriúscula. L.T. (bh.)	roughish.	Britain.	—	—	H. h.	
60	heterophylla. L.T.	(bh.) various-leaved.	Scotland.	—	—	H. h.	
61	tomentosa. L.T.	(bh.) downy-l ^o .	Britain.	—	—	H. h.	Eng. bot. v. 14. t. 990.
	β sylvestris. L.T.	wood.	—	—	—	H. h.	
	γ canescens. L.T.	(bh.) canescents.	—	—	—	H. h.	
	δ incánea. L.T.	(bh.) hoary.	—	—	—	H. h.	
	ε resinosa. L.R.	(re.) resinous.	Ireland.	—	—	H. h.	
62	nuda. L.T.	(bh.) naked.	Westmoreland.	—	—	H. h.	
63	álba. L.R.	(wh.) single white.	Europe.	1597.	6. 7.	H. h.	Miss Lawr. ros. t. 37.
64	montána. s.s.	(bh.) mountain.	Tauria.	1820.	—	H. h.	
65	hibérnica. L.T.	(pi.) Irish.	Ireland.	6. 11.	H. h.	Eng. bot. v. 31. t. 2196.
66	lutea. L.R.	(ye.) yellow.	Germany.	1596.	6.	H. h.	Bot. mag. 363.
67	punicea. Mil.	(re. ye.) red and yellow.	Persia.	—	—	H. h.	— t. 1077.
68	Sherárdi. Sm.	(bh.) Sherard's.	Britain.	6. 7.	H. h.	
69	Dicksoniàna. L.B.	(bh.) Dickson's.	Ireland.	—	—	H. h.	
70	rubiginosa. E.B.	(ro.) sweet briar.	Britain.	—	—	H. h.	Eng. bot. v. 14. t. 991.
71	micrántcha. L.T.	(ro.) small.	England.	—	—	H. h.	— v. 35. t. 2490.

ROSACEÆ.

72	sépium. DC.	(bh.) hedge.	France.	— H. h.		
73	Borréri. L.T.	(bh.) Borrer's.	Britain.	— H. h.	Eng. bot. v. 36. t. 2579.	
	<i>dumetorum</i> . E.B. nec aliorum.						
74	umbellata. DC.	(ro.) umbel-flowered.		— H. h.		
75	adenophylla. s.s.	(bh.) glandulous.		1816.	— H. h.		
76	pruinosa. L.C.	(ro.) frosted.		1817.	5. 6. H. h.		
	<i>pulverulenta</i> . L.R. non M.B.						
77	glutinosa. L.R.	(bh.) glutinous.	Levant.	1821.	6. 7. H. h.	Cupan. panp. ed. 1.t.61.	
	<i>rubiginosa crética</i> . Red. ros. 125. t. 47.						
78	Montezumæ. K.S.	(ro.) Mexican.	Mexico.	1823. H. h.		
79	rêpens. Scop.	(bh.) creeping.		1825.	6. 7. H. h.	Redout. ros. 1. t. 16.	
80	uncinata. Bes.	(bh.) hooked.	Poland.	1826.	— H. h.		
81	Boreykiána. Bes.	(bh.) Boreyki's.		1827.	— H. h.		
82	Fischeriana. Bes.	(bh.) Fischer's.	Russia.	1825.	— H. h.		
83	gorenkensis. Bes.	(bh.) Gorenki.		—	— H. h.		
84	guttesteiniæsis. Bes.	(bh.) Guttenstein.	Poland.	1822.	— H. h.		
85	Kosinskiæna. Bes.	(bh.) Koscinski.	Europe.	1825.	— H. h.		
86	polyphylla. Bes.	(bh.) many-leaved.		1827.	— H. h.		
87	terebinthinacea. Bes.	(bh.) Turpentine.	Germany.	1825.	— H. h.		
88	Redowskiæna. Bes.	(bh.) Redowski's.		1828.	— H. h.		
89	uncinella. Bes.	(bh.) small-hooked.	Europe.	1825.	— H. h.		
90	dimorpha. Bes.	(bh.) variable.	Volhynia.	1821.	— H. h.		
91	caryophyllacea. Bes.	(bh.) Clove-scent'd.	Poland.	1825.	— H. h.		
92	glaucæ. Bes.	(bh.) glaucous.		—	— H. h.		
93	nitidula. Bes.	(bh.) glossy.	Europe.	—	— H. h.		
94	floribunda. Bes.	(bh.) abundant-fl'd.		1828.	— H. h.		
95	solstitialis. Bes.	(bh.) summer.	Poland.	1821.	— H. h.		
96	acuta. F.	(bh.) acute-leaved.	Caucasus.	—	— H. h.		
97	purpurascens. F.	(pu.) purplish.	Tanaim.	—	— H. h.		
98	rupincola. F.	(bh.) rocky.	Altay.	—	— H. h.		
99	saxatilis. M.B.	(bh.) rock.	Tauria.	1825.	— H. h.		
100	caucásica. M.B.	(bh.) Caucasian.	Caucasus.	1793.	5. 6. H. h.	Lindl. ros. t. 11.	
101	cæsia. L.T.	(bh.) grey-leaved.	Scotland.	7. H. h.	Eng. bot. v. 33. t. 2367.	
102	sarmatæca. L.T.	(ro.) sarmentose.	Britain.	6. 7. H. h.		
	β nitens. L.T.	(ro.) glossy-leaved.		— H. h.		
103	bractescens. L.T.	(bh.) long-bracted.	Lancashire.	— H. h.		
104	dumetorum. L.T.	(bh.) bushy.	England.	6. H. h.		
105	Forstéri. Sm.	(bh.) Forster's.		— H. h.		
106	collina. L.T.	(bh.) hill.		— H. h.	Jacq. aust. 2. t. 197.	
107	camina. L.T.	(bh.) dog or hep.	Britain.	6. 7. H. h.	Eng. bot. v. 14. t. 902.	
	β cera. L.T.	(bh.) waxy.		— H. h.		
	γ glandulifera. L.T.	(bh.) glandular.		— H. h.		
	δ simpliciuscula. L.T.	(bh.) simple-calyxed.		— H. h.		
108	surculosa. L.T.	(bh.) surculose.		— H. h.		
109	leucanthæna. Len.	(wh.) white-flowered.	Siberia.	1824. H. h.		
110	rubrifolia. L.R.	(re.) reddish-leaved.	S. Europe.	5. 6. H. h.	Bot. reg. t. 430.	
111	sericea. L.R.	(bh.) silky-leaved.	Nepaul.	1823. H. h.	Lindl. ros. t. 12.	
112	microphylla. L.R.	(re.) small-leaved.	China.	1822.	6. 8. H. h.	Bot. reg. t. 919.	
113	longifolia. w.	(bh.) Willow-leaved.		2. 10. F. h.	Redout. ros. 2. 27. t. 12.	
114	índica. w.	(ro.) common China.		1789.	1. 12. H. h.	Miss Lawr. ros. t. 26.	
	β centifolia.	(ro.) large-double.		— H. h.		
	γ purpurea.	(pu.) purple China.		— H. h.		
	δ subalba.	(bh.) whitish China.		— H. h.		
	ϵ media.	(ro.) middle China.		— H. h.		
	ζ minor.	(ro.) small China.		— H. h.		
	η frâgrans.	(pu.) animating.		— H. h.		
115	odorata.	(pa.) sweet-scented.		1810.	2. 12. H. h.	Andr. ros. ic.	
	índica odorata. A.R.	índica frâgrans. R.R. 1. 115. t. 42.			Bot. reg. 804.	
	β flavescens.	(st.) yellow China.	China.	1823.	— H. h.		
116	semperflorens. L.R.	(cr.) red China.		1789.	1. 12. F. h.	Exot. bot. 2. t. 91.	
	α diversifolia.	v.(re.) single.		— F. h.	Vent. cels. t. 35.	
	β atrorubens.	(cr.) double red.		1789.	— F. h.	Bot. mag. 284.	
	γ resplendens.	(cr.) resplendent.		— F. h.		
	δ frâgrans.	(re.) fragrant.		1820.	— F. h.		
	ϵ Terneauxii.	(re.) Terneaux's.		— F. h.		
	ζ Bichonii.	(re.) Bichon's.		— F. h.		
117	Lawranceana. L.R.	(bh.) Miss Lawrence's.		1810.	— F. h.	Bot. reg. t. 538.	
	β flore pleno.	(bh.) double-flowered.		1824.	— F. h.		

118	<i>systyla</i> . L.T.	(<i>bh.</i>) clustered dog.	Britain.	6. 7.	H. h.	Eng. bot. v. 27. t. 1895.
	β <i>lanceolata</i> . L.R.	(<i>bh.</i>) spear-leaved.	Ireland.		H. h.	
	γ <i>Monsoniae</i> . L.R.	(<i>bh.</i>) Lady Monson's.	England.		H. h.	
119	<i>arvensis</i> . L.T.	(<i>wh.</i>) field.	Britain.		H. h.	Eng. bot. v. 3. t. 188.
	β <i>scandens</i> .	(<i>wh.</i>) <i>Ayrshire</i>		H. h.	Bot. mag. 2054.
120	<i>geminata</i> . Rau.	(<i>bh.</i>) Swiss.	Switzerland.		H. h.	
121	<i>semperfiriens</i> . w.	(<i>wh.</i>) evergreen.	S. Europe.	1629.	6. 8.	H. h.	Bot. reg. t. 465.
122	<i>Boursoultii</i> .	(<i>ro.</i>) <i>Boursoult's</i> .	Hybrid.	1821.	4. 7.	H. h.	
123	<i>hyacinthina</i> .	(<i>ro.</i>) Hyacinth-scented.				H. h.	
124	<i>Grevillii</i> .	(<i>ro. bh.</i>) Greville's.	China.		H. h.	
125	<i>Roxburghii</i> .	(<i>ro.</i>) Roxburgh's.			H. h.	
126	<i>multiflora</i> . w.	(<i>va.</i>) many-flowered.		1804.	6. 8.	H. h.	Bot. reg. 425.
	α <i>rosea</i> .	(<i>ro.</i>) common.				H. h.	Bot. mag. t. 1059.
	β <i>albiflora</i> .	(<i>wh.</i>) white-flowered.		1825.		H. h.	
127	<i>Brunoniæ</i> . L.R.	(<i>wh.</i>) Mr. Brown's.	Nepaul.	1820.		H. h.	Lindl. ros. t. 14.
128	<i>moschata</i> . L.R.	(<i>wh.</i>) musk-scented.	Africa.	1596.	7. 9.	H. h.	Miss Lawr. ros. t. 64.
	β <i>flôre plêno</i> .	(<i>wh.</i>) double musk.				H. h.	t. 53.
129	<i>nivea</i> . Dup.	(<i>wh.</i>) snow white.	Hybrid.	1823.		H. h.	Bot. reg. 861.
130	Noisettiana. R.R.	(<i>bh.</i>) Noisette's.		1816.	5. 11.	H. h.	
	β <i>grandiflora</i> .	(<i>bh.</i>) large-flowered.		1827.		H. h.	
	γ <i>Fraseri</i> .	(<i>bh.</i>) Fraser's.		1815.		H. h.	
	δ <i>purpurea</i> .	(<i>pu.</i>) purple.		1823.		H. h.	
131	<i>rufibolia</i> . L.R.	(<i>re.</i>) bramble-leav'd.	N. America.	1800.	7. 9.	H. h.	
	β <i>fenestrâlis</i> . L.R.	(<i>re.</i>) thin-leaved.				H. h.	Lindl. ros. t. 15.
132	<i>laevigata</i> . L.R.	(<i>wh.</i>) smooth.	Georgia.	1826.		H. h.	Bot. mag. t. 2847.
133	<i>sinica</i> . L.R.	(<i>wh.</i>) shining-leaved.	China.	1759.	6. 8.	F. h.	Lindl. ros. t. 16.
134	<i>Banksiae</i> . L.R.	(<i>wh.</i>) Lady Banks's.		1807.		H. h.	Bot. reg. t. 397.
	α <i>albiflora</i> .	(<i>wh.</i>) white-flowered.				H. h.	Bot. mag. t. 1954.
	β <i>lutea</i> .	(<i>ye.</i>) yellow-flowered.		1823.		H. h.	v. 13. t. 1105.

GARDEN VARIETIES.

I. *spinosisima*.

- 1 Aberdeen.
- 2 Aberfoil.
- 3 alba.
- 4 Alloa.
- 5 Ancram.
- 6 Anderson's double lady's [blush.]
- 7 Arbroath.
- 8 Argyll.
- 9 Arrochar.
- 10 Aurora borealis.
- 11 Balloch.
- 12 Banff.
- 13 Bannochburn.
- 14 Borisdale.
- 15 Bass.
- 16 Bengloe.
- 17 Ben Lomond.
- 18 Ben More.
- 19 Berwick.
- 20 bicolor.
- 21 Biggar.
- 22 Birnam.
- 23 Blair Athol.
- 24 Boharm.
- 25 Borthwick.
- 26 Buchan.
- 27 Bute.
- 28 Caithness.
- 29 Calder.
- 30 Callender.
- 31 Campsie.
- 32 carnescens.
- 33 Carron.
- 34 Cheviot.
- 35 ciphieri.

GARDEN VARIETIES.

- 36 Clydesdale.
- 37 Cromarty.
- 38 Dalkeith.
- 39 Dalrymple.
- 40 Dornock.
- 41 double blush Princess.
- 42 double blush Provinc.
- 43 double crimson.
- 44 double crimson mottled.
- 45 double dark marbled.
- 46 double dark red.
- 47 double lady's blush.
- 48 double light marbled.
- 49 double light purple.
- 50 double pink.
- 51 double purple.
- 52 double rose blush.
- 53 Douglas.
- 54 Dunbarton.
- 55 Dunblane.
- 56 Dumferline.
- 57 Dumfries.
- 58 Dunbar.
- 59 Duncrieff.
- 60 Dundee.
- 61 Dunglass.
- 62 Dunkeld.
- 63 Dunlop.
- 64 Dummore.
- 65 Dutch double blush.
- 66 dwarf bicolor.
- 67 Dysart.
- 68 Eden.
- 69 Elgin.
- 70 Etterick.
- 71 Fairy.
- 72 Falkirk.

GARDEN VARIETIES.

- 73 Falkland.
- 74 Falla.
- 75 Fife.
- 76 flavescens.
- 77 Forfar.
- 78 Forth.
- 79 Fort William.
- 80 fulgens.
- 81 Furness.
- 82 Galloway.
- 83 Glasgow.
- 84 Glenco.
- 85 Glenfallach.
- 86 Glengarry.
- 87 Gourrock.
- 88 Grahamston.
- 89 Grampian.
- 90 Greenock.
- 91 Haddington.
- 92 Halkirk.
- 93 Hamilton.
- 94 Hawick.
- 95 Hawthorndean.
- 96 Hector.
- 97 humiliis.
- 98 Huntly.
- 99 incarnata.
- 100 Inverary.
- 101 Inverness.
- 102 Invermay.
- 103 Janus.
- 104 Jedburgh.
- 105 Jura.
- 106 Keith.
- 107 Kelso.
- 108 Kilmarnock.
- 109 Kincardine.

GARDEN VARIETIES.

- 110 Kinnaird.
 111 Kinross.
 112 Kirkcaldy.
 113 Kirkwall.
 114 Lanark.
 115 Laxford.
 116 Leith.
 117 Leslie.
 118 Lismore.
 119 Lochaber.
 120 Lochard.
 121 Loch fine.
 122 Lochleven.
 123 Lochlomond.
 124 Lothian.
 125 Maidenkirk.
 126 Melrose.
 127 Moncrieff.
 128 Monteith.
 129 Montgomery.
 130 Montrose.
 131 Mount Stuart.
 132 Mull.
 133 Moray.
 134 mutabilis.
 135 Nevis.
 136 Northumberland.
 137 Paisley.
 138 penicillata.
 139 Pentland.
 140 perpetual.
 141 picta. L.B.C. 687.
 142 Proteus.
 143 red, double light.
 144 Roberton.
 145 Rosslyn.
 146 Rothsay.
 147 rubicunda.
 148 Selkirk.
 149 Shetland.
 150 Sky.
 151 small double light purple.
 152 small double white.
 153 small double yellow.
 154 Staffa.
 155 Stirling.
 156 Strathmore.
 157 striata.
 158 Stronsa.
 159 Sutherland.
 160 Tiviotdale.
 161 Tarbet.
 162 Thornhill.
 163 Thurso.
 164 Tranent.
 165 true double red.
 166 two-coloured large double.
 167 two-coloured small double.
 168 variegata.
 169 venulosa glabra.
 170 venulosa hispida.
 171 white, large double.
 172 white, large semidouble.
 173 white, small double.
 174 Whitley's double white.
 175 yellow, globe double.
 176 yellow, large double.
 177 yellow, pale double.

GARDEN VARIETIES.

- II. *gallica*.
- 178 admirable.
 179 aigle noir.
 180 aimable beaute.
 181 albanian.
 182 Amaranth.
 183 Antwerp.
 184 atlas.
 185 beaute rouge.
 186 beaute supreme.
 187 belle aurore.
 188 belle cramoisie.
 189 belle herminie.
 190 belle pourpre.
 191 belle violette.
 192 Bijou.
 193 Bishop. M.L.R. 20.
 194 black frizzled.
 195 blue.
 196 blue and purple.
 197 blush hundred-leaved.
 198 bouquet rouge royale.
 199 bright purple.
 200 brunette.
 201 Brussels.
 202 Buonaparte.
 203 burning coal.
 204 cardinal M.L.R. 59.
 205 carmine.
 206 carmine, brillante.
 207 carmine, proliferous.
 208 carnation.
 209 Catalonian.
 210 champion.
 211 chancellor.
 212 changeable.
 213 cherry.
 214 Clementine.
 215 coquette.
 216 couleur de feu.
 217 cramoisie, grand.
 218 crown.
 219 Cupid.
 220 damask black.
 221 dark marbled.
 222 dark violet.
 223 delicious.
 224 dingy.
 225 double velvet.
 226 double marbled. M.L.R. 57.
 227 Duc de Guiche.
 228 Duchess d'Orleans.
 229 dutch crimson.
 230 dutch tree.
 231 dutch 100 leaved.
 232 early Ranunculus.
 233 enchanter.
 234 enfant de France.
 235 Eucharis.
 236 Fanny Bias.
 237 favourite purple.
 238 fiery.
 239 Flanders.
 240 Flemish.
 241 formidable.
 242 fringed. M.L.R. 82.
 243 garnet.
 244 gay.

GARDEN VARIETIES.

- 245 giant. M.L.R. 49.
 246 gloria mundi.
 247 gloria mundi flore pallido.
 248 gloria mundi pourpre.
 249 granaat appel.
 250 grand monarque.
 251 grand purple.
 252 grand Sultan.
 253 Henry IV.
 254 Hervy.
 255 imperatrice.
 256 incomparable.
 257 infernal.
 258 invincible.
 259 Italian.
 260 Josephine.
 261 Junon.
 262 king.
 263 La Dauphine.
 264 la grand belle pourpre.
 265 Leyden.
 266 light purple.
 267 Lisbon. M.L.R. 67.
 268 lively.
 269 L'ombre agreable.
 270 L'ombre superbe.
 271 lurid.
 272 maiden.
 273 Majorca.
 274 Malabar.
 275 Malta.
 276 manteau royal.
 277 marbled.
 278 marbled, grand.
 279 Margaret.
 280 matchless.
 281 Mauve.
 282 mignonne.
 283 mignonne, blush.
 284 mignonne, dark.
 285 mignonne, favorite.
 286 mignonne, incised.
 287 mignonne, red.
 288 mignonne, semidouble.
 289 mignonne, striped.
 290 mirabelle.
 291 Mogul.
 292 Montaubon.
 293 Morocco.
 294 mottled, black.
 295 mundi. M.L.R. 13.
 296 natalie.
 297 negrette.
 298 negro.
 299 ninon de l'Enclos.
 300 nonpareil.
 301 nonsuch.
 302 noir de Hollande.
 303 Normandy.
 304 officinal.
 305 officinal blush.
 306 officinal carmine.
 307 Orleans.
 308 ornement de parade.
 309 Paestana.
 310 paradise.
 311 paragon.
 312 pavot.

GARDEN VARIETIES.

- 313 perruque.
 314 petite 100 leav'd. M.L.R. 55.
 315 petite panachée.
 316 phœnix.
 317 plicate. M.L.R. t. 87.
 318 Pluto. M.L.R. 32.
 319 Pomona.
 320 pompadour.
 321 pompone bizard.
 322 Poniatowsky.
 323 poppy.
 324 porcelaine.
 325 Portland.
 326 pourpre bouquet.
 327 pourpre charmante.
 328 pourpre de Tyr.
 329 pourprée, grand belle.
 330 pourprée, point.
 331 pourpre velours.
 332 prince.
 333 princess.
 334 prince William V.
 335 prolific.
 336 Pronville.
 337 Proserpine.
 338 Provins pulmonaire.
 339 purple blue.
 340 purple crimson.
 341 pyramidal.
 342 queen.
 343 Ranunculus.
 344 red and violet.
 345 Roi de France.
 346 Roi de pourprés.
 347 royal crimson.
 348 royal red.
 349 royal purple.
 350 royal virgin. M.L.R. 7.
 351 sable.
 352 Saint John's.
 353 Sanspareil.
 354 sceptre.
 355 semidouble velvet.
 356 shell. M.L.R. 89.
 357 single velvet.—51.
 358 Singleton's 100 leaved.—85.
 359 Spanish.
 360 stadtholder.
 361 Stepney. M.L.R. 46.
 362 striped nosegay.
 363 striped velvet.
 364 superb red.
 365 Sultan. M.L.R. 35.
 366 Trafalgar.
 367 triumphant.
 368 Tuscany. B.R. v. 6. t. 448.
 369 two-colored.
 370 Venetian.
 371 victory.
 372 violette and rouge.
- III. centifolia.**
- 373 aurora.
 374 Belle d'Aunay.
 375 Blandford, M.L.R. 21.
 376 black mottled.
 377 blush royal. M.L.R. 47.

GARDEN VARIETIES.

- 378 blush cabbage.
 379 Bourbon.
 380 bright crumpled.
 381 carmine.
 382 Centfeuilles Anemone.
 383 Centfeuilles de Bruxelles.
 384 Centfeuilles de Hesse.
 385 Centfeuilles gaufrière.
 386 chamois.
 387 cluster.
 388 Constance.
 389 Cumberland.
 390 dragon.
 391 Duchesse d'Angoulême.
 392 Duchesse de Berri.
 393 early 100 leaved.
 394 Elysian.
 395 emperor.
 396 grand chamois.
 397 grand marbled.
 398 grand Provins.
 399 Louis XVIII.
 400 Malta.
 401 mère gryone.
 402 mignon, scarlet.
 403 mignon, crimson.
 404 mignon, purple.
 405 mottled purple.
 406 Neapolitan.
 407 one-sided.
 408 Oillet.
 409 pencilled.
 410 petite Holland.
 411 Persian.
 412 pompone, gros.
 413 pourprée aimable.
 414 pourprée favorite.
 415 pourprée violette.
 416 Provins, single. R.R. 1. t. 1.
 417 Provins, common. M.L.R. 8.
 418 Provins, cabbage.
 419 Provins, blush cabbage.
 420 Provins, scarlet. M.L.R. 22.
 421 Provins, Childings.—43.
 422 Provins, blush.—1.
 423 Provins, white.—4.
 424 Provins, Shailler's.—76.
 425 Provins, Colvill's.
 426 Provins, damask.
 427 Provins, dwarf.
 428 Provins, invincible.
 429 Provins, Dutch.
 430 Provins, imperial.
 431 Provins, royal.
 432 Provins, early.
 433 Provins, semidouble.
 434 red Belgic.
 435 rouge superbe.
 436 sans petales.
 437 souchet.
 438 Spong's.
 439 superb carmine.
 440 surpassante.
 441 Syren.
 442 Trianon, Belle de
 443 Versailles.
 444 Vilnorin.
- * *centifolia muscosa.*
 445 single moss.
 446 common moss.
 447 blush moss.
 448 dark moss.
 449 mottled moss.
 450 prolific moss.
 451 royal moss.
 452 scarlet moss.
 453 semidouble moss.
 454 striped moss.
 455 Shailler's white moss.
 456 Bristol white moss.
 † *centifolia pomponia.*
 457 de Meaux. M.L.R. 31.
 458 de Rheims.—71.
 459 dwarf Bagshot.
 460 mignon charmante.
 461 mossy de Meaux.
 462 mossy Pompone.
 463 Pompone. M.L.R. 50.
 464 proliferous Pompone.
 465 St. Francis. M.L.R. 88.
 466 small provins.
- IV. damascena.**
- 467 argentea.
 468 Aurora.
 469 bifera carnea.
 470 bifera de Naples.
 471 bifera grandiflora.
 472 Belgique carnée.
 473 Belgique violette.
 474 belle couronnee.
 475 blush Belgic. M.L.R. 70.
 476 blush monthly.—18.
 477 blush damask.—52.
 478 Brunswick.
 479 couronnée petite.
 480 damas argente.
 481 damas pourpre.
 482 early blush.
 483 Egyptian.
 484 Emperor.
 485 favorite mignon.
 486 felicité.
 487 Goliath. M.L.R. t. 80.
 488 gracieuse.
 489 grande monarque.
 490 great royal. M.L.R. 58.
 491 imperial blush. M.L.R. 90.
 492 incomparable.
 493 lesser Belgic.
 494 pale cluster.
 495 paragon.
 496 Parnassus.
 497 Paëstana.
 498 petite 100-leaved.
 499 perpetual.
 500 prolific.
 501 quatre saisons.
 502 quatre saisons blanche.
 503 quatre saisons, flesh-colored.
 504 quatre saisons Francois.
 505 quatre saisons panaché.
 506 quatre saisons pompone.
 507 quatre saisons sans épines.
 508 quatre saisons, semidouble.

ROSACEÆ.

GARDEN VARIETIES.

- 509 red damask. M.L.R. 38.
 510 red monthly. M.L.R. 5.
 511 red Belgic.—83.
 512 rouge agathe.
 513 Swiss.
 514 valiant.
 515 Watson's blush.
 516 white damask.
 517 white monthly. M.L.R. 17.
 518 York and Lancaster.—10.
 519 Zealand.

V. alba.

- 520 agate.
 521 agate magnifique.
 522 agate Marie Louise.
 523 agate nouvelle.
 524 agate prolifere.
 525 agate royale.
 526 agate superb.
 527 agate thalie.
 528 belle aurore.
 529 belle Henriette.
 530 blanche à cœur vert.
 531 blanche de Belgique.
 532 bouquet blanc.
 533 celestial.
 534 cluster maiden's blush. [M.L.R. 23.]
 535 double white blush.
 536 Duc d'York.
 537 Eliza.
 538 feuille fermée.
 539 full double white. M.L.R. 25.
 540 grand cuisse de nymphe.
 541 great maiden's blush. [M.L.R. 230.]
 542 Joanne d'Arc.
 543 Moraga la favourite.
 544 muscat rouge.
 545 nova cœlestis.
 546 nova plena.
 547 petite cuisse de nymphe.
 548 rosea.
 549 semidouble white.
 550 Simonville.
 551 small maiden's blush.
 552 spineless virgin.
 553 thornless virgin.
 554 triangularis.

VI. rubiginosa.

- 555 Clementine sweet briar.
 556 cluster blush sweet briar.
 557 double mossy sweet briar. [M.L.R. 72.]
 558 double red sweet briar.—61.
 559 double rosy sweet briar.
 560 double tree sweet briar.
 561 eglantine sweet briar.
 562 eglanteria alba.
 563 evergreen sweet briar.
 564 large blush sweet briar.
 565 maiden sweet briar.
 566 Manning's sweet briar. [M.L.R. 41.]
 567 monstrous sweet briar.

GARDEN VARIETIES.

- 568 petite hessoise.
 569 royal sweet briar. [M.L.R. 74.]
 570 scarlet sweet briar.
 571 semidouble dwarf.
 572 semidouble white sweet briar. [M.L.R. 65.]
 573 Zabeth.

VII. indica et semperflorens.

- 574 atronigra.
 575 Bengale a bouquet.
 576 Bengale à floré panaché.
 577 Bengale blanche.
 578 carnescens.
 579 Chiffonnée.
 580 Cérisé éclatante.
 581 cucullata.
 582 elegant.
 583 florida.
 584 gigantea.
 585 lie de vin.
 586 lúcida.
 587 mäjor.
 588 monstrosa.
 589 moonshine.
 590 nigra.
 591 sanguinea.
 592 sans épines.
 593 Thisbe.
 594 Veloutée.

VIII. various garden Roses.

- 595 abundant.
 596 abyala.
 597 Achaia.
 598 Achilles.
 599 Adelina.
 600 admirable.
 601 Adonis.
 602 African black.
 603 agate Japanese.
 604 aglace noir.
 605 aimable violette.
 606 Ajax.
 607 aland.
 608 alba feuille fermee.
 609 alba bifera.
 610 Albanian.
 611 alba nova pleno.
 612 alba nova cœlestis.
 613 albo-rosea.
 614 alba triangularis.
 615 aline.
 616 Amazon.
 617 amour constant.
 618 Andalusian.
 619 anemoneflora.
 620 Arcadian.
 621 arcario.
 622 archbishop.
 623 ardoise.
 624 Armenian.
 625 armida.
 626 Assyrian.
 627 Athenian.
 628 augusta.
- 629 aurelia.
 630 Baden.
 631 Balbec.
 632 beauté frappante.
 633 beauté insurmontable.
 634 beauté panache.
 635 beauté renommée.
 636 beauté sanspareille.
 637 beauté tendre.
 638 belladonna.
 639 belle actrise.
 640 belle aimable.
 641 belle aurore descemet.
 642 belle brune.
 643 belle cataline.
 644 belle cerisie.
 645 belle d'aunay.
 646 belle du jour.
 647 belle flamande.
 648 belle forme.
 649 belle galathee.
 650 belle hative.
 651 belle incarnate.
 652 belle inconnue.
 653 belle matilde.
 654 belle mahæca.
 655 belle parade.
 656 belle Tigridia.
 657 belle zulme.
 658 belle rouge.
 659 belle sans flatterie.
 660 belle sanspareille.
 661 belle therese.
 662 belle vue.
 663 bellona.
 664 Berkshire.
 665 bizard royal.
 666 black purple.
 667 blanche superbe.
 668 blondine.
 669 blush Italian.
 670 blush velvet.
 671 bold.
 672 Bonifacius.
 673 bouquet blanc.
 674 bouquet panache.
 675 bouquet parfait.
 676 bouquet superb.
 677 Bourbon.
 678 Brabant.
 679 bracelet d'amour.
 680 brigitt.
 681 brilliant.
 682 brune brillante.
 683 brunette aimable.
 684 brunette superbe.
 685 bucephalus.
 686 bullata.
 687 cabinet.
 688 Calypso.
 689 camellote.
 690 cannabina.
 691 canopus.
 692 Carthaginian.
 693 Castile.
 694 Cecropia.
 695 Ceres.

GARDEN VARIETIES.

GARDEN VARIETIES.

- 696 cervalis.
 697 chance.
 698 charmante pourpre.
 699 chausse.
 700 chiron.
 701 cicris pale rose.
 702 Cleopatra.
 703 comtesse de Genlis.
 704 cornelian.
 705 cossack.
 706 couleur a la mode.
 707 couleur de cendre.
 708 couleur de cerise.
 709 couleur excellent.
 710 coupee.
 711 couronne.
 712 couronne des roses.
 713 couronne imperiale.
 714 cramoisie eblouissant.
 715 cramoisie elegant.
 716 cramoisie fonce.
 717 cramoisie imperial.
 718 cramoisie incomparable.
 719 cramoisie minor.
 720 cramoisie nouveau.
 721 cramoisie panache.
 722 cramoisie sanspareil.
 723 cramoisie triomphant.
 724 crimson.
 725 crimson perpetual.
 726 crown imperial.
 727 crown purple.
 728 cybele.
 729 danae.
 730 Danish.
 731 dark mignonne.
 732 dark mottled.
 733 dark shell.
 734 darling.
 735 de bengal gros violet
 736 Dejania.
 737 delicate.
 738 Derby.
 739 descemets.
 740 diadem.
 741 dianthiflora.
 742 Dickson's duchess.
 743 Dido.
 744 don des dames.
 745 Dione.
 746 discolor.
 747 Dorothee.
 748 double apple bearing.
 [M.L.R. 29.]
 749 double dog.—60.
 750 double mimms.
 751 double Pæony.
 752 double purple.
 753 double red thornless.
 754 double white thornless.
 755 due de Brabant.
 756 Duchess.
 757 duchess d'Oldenburgh.
 758 duke of Clarence.
 759 Durham.
 760 Dutch Cinnamon.
 761 Dutch velvet.

GARDEN VARIETIES.

- 762 dwarf burnet-leaved.
 763 dwarf Chester.
 764 dwarf prolific.
 765 early red.
 766 erato.
 767 Etna.
 768 euphrosyne.
 769 evratina.
 770 ex alba rosea.
 771 fair maid.
 772 favonius.
 773 favorite.
 774 favorite des dames.
 775 fausse unique.
 776 felicie.
 777 felicite.
 778 Felix.
 779 feu amoureux.
 780 feuille chiffonee.
 781 feuille crenulee.
 782 feuille de chene.
 783 feuille de frene.
 784 few-spined.
 785 fine panachee.
 786 fine crimson.
 787 fine purple.
 688 flavia pourpre.
 789 flesh-coloured.
 790 flesh-coloured four seasons.
 791 fleur de parade.
 792 fleur de pomme.
 793 fleur et feuille marbree.
 794 fleur panachee.
 795 flora.
 796 flore nigrante.
 797 florentine.
 798 flore rubra.
 799 floribunda.
 800 florio.
 801 flos puniceus.
 802 folie de corsé.
 803 four seasons semidouble.
 804 franconian.
 805 Frankfort.
 806 Frankfort agate.
 807 freckled.
 808 frilled.
 809 frizzled.
 810 gallica foliacea.
 811 garnett.
 812 Gascony.
 813 Genoese.
 814 Genoese velvet.
 815 gienvieve.
 816 globe.
 817 gloomy.
 818 gloria rubrorum.
 819 gloriosa.
 820 glowing.
 821 grand Alexandre.
 822 grande Angloise.
 823 grand Cæsar.
 824 grand centifolia.
 825 grande chausse.
 826 grand cramoisi.
 827 grande foncee.
 828 grand Mogul.

GARDEN VARIETIES.

- 829 grand pivoine.
 830 grande pompadour.
 831 grand pyramidal.
 832 grand rouge.
 833 grand triumphant.
 834 grandesse royale.
 835 grandeur royal.
 836 great Mogul.
 837 grey.
 838 grideline.
 839 grideline sans epines.
 840 Hainault.
 841 Haarlem.
 842 harpagus.
 843 Helena.
 844 Hebe.
 845 hero.
 846 Hertford.
 847 Hildesheim.
 848 Hollandia magniflora.
 849 hortulanus.
 850 hundred leaved mignonne.
 851 hyacinth.
 852 hybrida perfecta.
 853 hybride nouvelle.
 854 iconium.
 855 idumea.
 856 illustrious.
 857 imperatrice.
 858 imperatrice de France.
 859 Indian queen.
 860 inermis.
 861 iris noir.
 862 Isabelle.
 863 Isis.
 864 jaqueline.
 865 jeanne d'arc.
 866 Jersey.
 867 jessica.
 868 jolie.
 869 Julian.
 870 king Agrrippa.
 871 king of Mexico.
 872 la belle elise.
 873 la belle rosine.
 874 l'abondance.
 875 la constance.
 876 la coquette.
 877 la Diane.
 878 l'admirable.
 879 la fidele.
 880 la gracieuse.
 881 la grandesse.
 882 l'aimable de Stors.
 883 lake.
 884 la maculee.
 885 la magnifique.
 886 la majesteuse.
 887 la mignarde.
 888 l'amitie.
 889 la moderne.
 890 Lancaster.
 891 lanczman.
 892 la negresse.
 893 la parfaite.
 894 la Parisienne.
 895 la petite lisette.

ROSACEÆ.

GARDEN VARIETIES.

- 896 la plus belle.
 897 la plus rouge.
 898 la pomme grenade.
 899 la precieuse pourpre.
 900 la reine.
 901 la reine des roses.
 902 large platter.
 903 large perfect.
 904 la rose de Medicis.
 905 la rose d'Ispahan.
 906 la royale.
 907 la superbe.
 908 la superbe en brun.
 909 la tendresse.
 910 la victorieuse.
 911 la virginité.
 912 leafy.
 913 Leander.
 914 le dauphin.
 915 Lee's perpetual.
 916 le grandeur.
 917 le grand lowendael.
 918 le grand suarrow.
 919 le grand visir.
 920 Leipsick.
 921 les alliées.
 922 le violet triomphant.
 923 light 100 leaved.
 924 l'impayable.
 925 l'importeante.
 926 l'obscurité.
 927 Lodona.
 928 Lodoiska.
 929 l'ombre panaché.
 930 l'ombre sanspareille.
 931 London pride.
 932 Louis 18th.
 933 Lubeck.
 934 Ludolph.
 935 Ludovicus.
 936 Luxemburg.
 937 Lyonnaise.
 938 lystra.
 939 madder.
 940 magdalen.
 941 maiden blush.
 942 mantle.
 943 marble apple bearing.
 944 Margarettæ.
 945 Marie anne.
 946 marquis de la Romana.
 947 Marseilles.
 948 maxima.
 949 mere gigon.
 950 merveilleuse.
 951 Mexican.
 952 Milanese.
 953 Minerva.
 954 miroir des dames.
 955 mon bijou.
 956 monstrosa.
 957 monstrous cluster.
 958 monstrouis 100 ld. M.L.R. 73.

GARDEN VARIETIES.

- 963 mourning.
 964 multiplex.
 965 Napolean.
 966 Napoli di romana.
 967 Narbonne.
 968 Nassau.
 969 negre panaché.
 970 Neptune.
 971 new mottled.
 972 new Provinc.
 973 nigritienne.
 974 Niobe.
 975 noblesse.
 976 noir foncé.
 977 non plus ultra.
 978 nouvelle favorite.
 979 nouvelle de Provinc.
 980 nouvelle pivoine.
 981 Oldenburg.
 982 Oliver.
 983 Olympic.
 984 omphale.
 985 orient.
 986 orion.
 987 ornement de park.
 988 ornement de violet.
 989 oublie des François.
 990 painted.
 991 painted Belgic.
 992 Palestine.
 993 panachée admirable.
 994 panachée nouvelle.
 995 panachée sanspareille.
 996 panachée superbe.
 997 Pannonian.
 998 paradise.
 999 Paris 100-leaved.
 1000 Parnassus.
 1001 parson.
 1002 party coloured.
 1003 Patmos.
 1004 Palmyra.
 1005 Paulina.
 1006 Pearson's gigantic.
 1007 Pegasus.
 1008 pencilled mignonne.
 1009 pencilled Provinc.
 1010 Penelope.
 1011 perfecta.
 1012 perle de l'orient.
 1013 perpetuelle rouge.
 1014 Peruvian.
 1015 petit Caesar.
 1016 petit cramoisi.
 1017 petit favorite.
 1018 petit mignon.
 1019 pierian.
 1020 pilgrim.
 1021 pink velvet.
 1022 plicata purpurea.
 1023 Pomeranian.
 1024 pomegranate.
 1025 pomona.
 1026 pompone.
 2027 pompone blanc.
 1028 pompone bifera.
 1029 pompone quatre saisons.

GARDEN VARIETIES.

- 1030 pompone varin.
 1031 Pope's cluster.
 1032 portlandica grandiflora.
 1033 portlandica perpetua.
 1034 Portobella.
 1035 Portugal.
 1036 pourpre agreeable.
 1037 pourpre de Paris.
 1038 pourpre sans epines.
 1039 pourpre de Vienne.
 1040 pourpre imperial.
 1041 pourpre incomparable.
 1042 pourpre obscur.
 1043 pourpre sans defaut.
 1044 pourpre sanspareil.
 1045 pourpre superb.
 1046 predominant.
 1047 Presburg.
 1048 prince.
 1049 prince d'Aremberg.
 1050 princely.
 1051 prince regent.
 1052 princess Charlotte.
 1053 princesse noble.
 1054 professor.
 1055 purple crown.
 1056 purple imperial.
 1057 purple velvet.
 1058 pyramide pourpre.
 1059 quatre saisons blanche.
 1060 quatre saisons François.
 1061 quatre saisons panaché.
 1062 quatre saisons sans epines.
 1063 queen's Provinc.
 1064 ragged robin.
 1065 Raphael.
 1066 Ratisbon.
 1067 ravenna.
 1068 red cluster.
 1069 red crown.
 1070 red Provinc.
 1071 red velvet.
 1072 refulgent.
 1073 regina florum.
 1074 reine Caroline.
 1075 reine d'Hongrie.
 1076 renoncule nouvelle.
 1077 rex rubrorum.
 1078 riche en fleurs.
 1079 roi des negres.
 1080 roi de Rome.
 1081 roi de Maroc.
 1082 roi des pays bas.
 1083 Roman.
 1084 Rosabel.
 1085 Rosanna.
 1086 rose agreeable.
 1087 rose bouquet.
 1088 rose de Ceres.
 1089 rose des dames.
 1090 rose du roi.
 1091 rose d'Orleans.
 1092 rose du prince.
 1093 rose la mode.
 1094 rose lee.
 1095 rose pivoine.
 1096 rose prolifere.

GARDEN VARIETIES.

- 1097 rouge agreeable.
 1098 rouge bien vif.
 1099 rouge eclatant.
 1100 rouge formidable.
 1101 rouge frappant.
 1102 rouge luisant.
 1103 rouge panaché.
 1104 rouge sans epines.
 1105 rouge semidouble.
 1106 rouge vegetable.
 1107 royal bouquet.
 1108 royal carmine.
 1109 royal gabina.
 1110 royal mantle.
 1111 royal viscous.
 1112 rubrispina.
 1113 rubiginosa umbellata.
 1114 ruby.
 1115 sable.
 1116 St. Antoine.
 1117 St. Catharine.
 1118 St. George.
 1119 St. Germain.
 1120 St. Gothard.
 1121 St. Jago.
 1122 St. Mark.
 1123 St. Patrick.
 1124 sans defaut.
 1125 sans petales.
 1126 Sarmatian.
 1127 scarlet Brabant.
 1128 sceptre.
 1129 scone purple.
 1130 semidouble hep.
 1131 semidouble mignonette.
 1132 semidouble monthly.
 1133 semidouble new.
 1134 seigneur d'Ærtzelhaar.
 1135 septum cerise.
 1136 Seville.
 1137 Sheffield.
 1138 Shylock.
 1139 Silenus.
 1140 Silesian.
 1141 shining semidouble.
 1142 simplicité.
 1143 singuliere agate.

GARDEN VARIETIES.

- 1144 Sirius.
 1145 soliditas.
 1146 Southampton.
 1147 Spartan.
 1148 spineless red.
 1149 spiral.
 1150 stebon.
 1151 standard.
 1152 state.
 1153 Stephanus.
 1154 striped monthly.
 1155 striped Provins.
 1156 sugar loaf.
 1157 sulphurea.
 1158 sulphurea minor.
 1159 superb.
 1160 superb amaranth.
 1161 superbe brune.
 1162 superbe pyramide.
 1163 surpassé Singleton.
 1164 surpassé tout.
 1165 Swiss.
 1166 Syrian.
 1167 Tangiers.
 1168 ten-leaved.
 1169 temple d'Apollon.
 1170 tête de mort.
 1171 Theseus.
 1172 Tigris.
 1173 tricolor.
 1174 tomentosa alba.
 1175 toujours.
 1176 tous les mois gris.
 1177 transparent.
 1178 transparent nouvelle.
 1179 Trebonius.
 1180 tree burnet-leav'd.
 1181 tree paeony.
 1182 tresorier.
 1183 triomphe.
 1184 triomphe des dames.
 1185 triomphe royale.
 1186 Triton.
 1187 turban.
 1188 two-coloured 100-leaved.
 1189 two-coloured mignonette.
 1190 vacuna.

GARDEN VARIETIES.

- 1191 vagrant.
 1192 velours cramoisie.
 1193 veloute.
 1194 venerable.
 1195 venetatus.
 1196 ventome noir.
 1197 Venus.
 1198 venusta.
 1199 veritas.
 1200 vermillion.
 1201 vesta.
 1202 Vesuvius.
 1203 Victoria.
 1204 Vidua.
 1205 villosa nouvelle.
 1206 Vilimorin.
 1207 violette.
 1208 violette agreeable.
 1209 violette aimable.
 1210 violette brillante.
 1211 violette curieuse.
 1212 violette foncée.
 1213 violette maculée.
 1214 violette nouvelle.
 1215 violette sanspareille.
 1216 violette superbe.
 1217 violette superieure.
 1218 virgin's.
 1219 virginale.
 1220 virbilis.
 1221 virgo cramoisie.
 1222 ulterio.
 1223 ultramarine.
 1224 umbrella.
 1225 unique rouge.
 1226 Waterloo.
 1227 Watson's blush.
 1228 Watson's white.
 1229 Wellington.
 1230 woolly-leaved.
 1231 white damask.
 1232 white Pomponie.
 1233 York.
 1234 Yorkshire blush.
 1235 Yorkshire Provins.
 1236 zatre.
 1237 Zenobia.

Subordo II. POTENTILLÆ. Juss. gen. 337.

R'UBUS. L.

BRAMBLE. Icosandria Polygynia. L.

- 1 rosæfolius. DC. (*wh.*) Rose-leaved. Mauritius. 1811. 10. 4. G. h. Smith ic. 3. t. 60.
 2 eglantèria. DC. (*wh.*) Eglantine. N. Holland. 1824. 5. 6. G. h. Bot. mag. t. 1783.
 3 coronarius. (*wh.*) double-flowered. China. 1811. 10. 4. H. h. Bot. reg. t. 854.
rosæfolius β *coronarius*. B.M.
 4 pinnatus. DC. (*wh.*) pinnate. St. Helena. 1789. 6. 7. G. h.
 5 gracilis. DC. (*wh.*) slender. E. Indies. 1826. — S. h.
 6 micranthus. D.P. (*re.*) small-flowered. Nepaul. 1818. 6. 8. H. h. Bot. reg. t. 854.
pauciflorus B.R.
 7 distans. D.P. (*re.*) distant-leaved. — — H. h.
 8 apetalus. DC. (*gr.*) petalless. Mauritius. 1823. — S. h.
 9 macropodus. DC. (*wh.*) large-footed. N. Holland. 1822. — G. h.
 10 strigosus. DC. (*wh.*) strigose. N. America 6. 7. H. h.

ROSACEÆ.

11	pedunculosus. D.P. (<i>wh.</i>) long-peduncled.	Nepaul.	1820.	7. 8. H. h.	
12	occidentalis. DC. (<i>wh.</i>) American raspberry.	N.Amer.	1696.	5. 6. H. h.	Dill. elth. t. 247. f. 319.
13	asper. DC. (<i>wh.</i>) rough.	Nepaul.	1822.	6. 8. H. h.	Eng. bot. v. 34. t. 2442.
14	Idæus. DC. (<i>wh.</i>) Raspberry.	Britain.	5. 6. H. h.	Willd. hort. ber. t. 82.
15	laciñatus. DC. (<i>bh.</i>) jagged-leaved.	6. 9. H. h.	Eng. bot. v. 12. t. 826.
16	cæsius. DC. (<i>wh.</i>) Dewberry.	Britain.	6. 8. H. h.	
17	foliolosus. D.P. (<i>li.</i>) leafy.	Nepaul.	1823.	7. 8. H. h.	
18	flagellarris. DC. (<i>wh.</i>) shining-leaved.	N. America.	1789.	6. 7. H. h.	
19	inermis. W.en. (<i>wh.</i>) smooth.	1805.	— H. h.	
20	corylifolius. E.B. (<i>wh.</i>) Hazel-leaved.	Britain.	7. H. h.	Eng. bot. v. 12. t. 827.
21	agrætis. DC. (<i>wh.</i>) field.	Hungary.	1818.	7. 8. H. h.	W. et K. hung. 3. t. 268.
22	spectabilis. DC. (<i>pu.</i>) showy.	Columbia.	1827.	6. 8. H. h.	Ph. fl. amer. 1. t. 16.
23	plicatus. W.N. (<i>wh.</i>) plaited-leaved.	Britain.	7. 8. H. h.	Wei. et N. rub. germ. t. 1.
24	affinis. W.N. (<i>wh.</i>) related.	6. 9. H. h.	— p. 22. t. 3 et 36.
25	ulmifolius. DC. (<i>ro.</i>) Elm-leaved.	Gibraltar.	1824.	7. 8. H. h.	
26	Linkianus. DC. (<i>wh.</i>) double-flowered.	6. 9. H. h.	
27	albicans. DC. (<i>wh.</i>) white-leaved.	Hungary.	1818.	— H. h.	Eng. bot. v. 10. t. 715.
28	fruticosus. DC. (<i>fl.</i>) Blackberry.	Britain.	— H. h.	
	β pomponius. DC. (<i>bh.</i>) semidouble.	— H. h.	
	γ leucocarpus. DC. (<i>bh.</i>) white-berried.	— H. h.	
	δ inermis. DC. (<i>bh.</i>) smooth-stemmed.	— H. h.	
29	tomentosus. W.N. (<i>wh.</i>) woolly-leaved.	Germany.	1816.	6. 8. H. h.	Wei. et N. rub. germ. t. 8.
30	sancus. DC. (<i>pk.</i>) holy.	Levant.	1823.	— H. h.	Schreb. dec. t. 8.
31	canescens. DC. (<i>bh.</i>) canescent.	Piedmont.	1826.	— H. h.	
32	collinus. DC. (<i>bh.</i>) hill.	Montpelier.	1820.	7. 8. H. h.	Noc. et Balb. fl. tic. t. 9.
33	fastigiatus. W.N. (<i>wh.</i>) close-flowered.	Britain.	— H. h.	Wei. et N. rub. germ. t. 2.
34	suberectus. E.B. (<i>wh.</i>) upright.	6. 9. H. h.	Eng. bot. v. 36. t. 2572.
35	cordifolius. W.N. (<i>wh.</i>) heart-leaved.	7. 8. H. h.	Wei. et N. rub. germ. t. 5.
36	Ménkii. s.s. (<i>wh.</i>) velvety.	Germany.	1816.	— H. h.	
37	Schlechtendaliæ. s.s. (<i>w.</i>) Schlechtendale's.	— H. h.	
38	tiliaefolius. s.s. (<i>wh.</i>) Lime-tree-ld.	— H. h.	
39	Schleicheri. s.s. (<i>wh.</i>) Schleicher's.	1824.	— H. h.	
40	hóridus. s.s. (<i>wh.</i>) very prickly.	— H. h.	
41	leucostachys. s.s. (<i>wh.</i>) white-spiked.	Britain.	— H. h.	
42	rhamnifolius. W.N. (<i>wh.</i>) Rhamnus-leaved.	— H. h.	Wei. et N. rub. germ. t. 6.
43	abruptus. L.B. (<i>wh.</i>) abrupt.	— H. h.	Wei. et N. rub. germ. t. 4.
44	nítidus. W.N. (<i>bh.</i>) glossy-leaved.	Britain.	7. 8. H. h.	
45	argutus. DC. (<i>wh.</i>) sharp-toothed.	N. America.	1824.	6. 7. H. h.	
46	cuneifolius. DC. (<i>wh.</i>) wedge-leaved.	1811.	6. 8. H. h.	
47	sanguinolentus. DC. (<i>w.</i>) red-wooded.	Mauritius.	1823.	— S. h.	
48	villus. DC. (<i>wh.</i>) shaggy.	N. America.	1777.	7. 8. H. h.	
49	hirtus. W.K. (<i>wh.</i>) hairy.	Britain.	— H. h.	
50	vulpinus. Df. (<i>re.</i>) fox-tail.	Germany.	1816.	— H. h.	
	<i>Sprengeliæ.</i> W.N.				
51	jamaicensis. DC. (<i>wh.</i>) Jamaica.	Jamaica.	1820.	— S. h.	Sloan. jam. 2. t. 213. f. 1.
52	hispidus. DC. (<i>wh.</i>) bristly.	Canada.	1768.	— H. h.	
53	trivialis. Ph. (<i>bh.</i>) American dewberry.	N.Amer.	1789.	6. 7. H. h.	
54	macrophyllus. W.N. (<i>w.</i>) large-leaved.	Wales.	7. 8. H. h.	
55	vulgaris. W.N. (<i>bh.</i>) common.	Britain.	— H. h.	
56	diversifolius. L.B. (<i>bh.</i>) different-leaved.	— H. h.	
57	discolor. W.N. (<i>bh.</i>) two-coloured.	— H. h.	
58	fusco-ater. W.N. (<i>bh.</i>) dark-brown.	— H. h.	
59	pallidus. W.N. (<i>bh.</i>) pale-leaved.	— H. h.	
60	Kœhleri. W.N. (<i>bh.</i>) Kœhler's.	— H. h.	
61	rudis. W.N. (<i>bh.</i>) way-side.	— H. h.	
62	rubricaulis. W.N. (<i>bh.</i>) red-stalked.	Germany.	1816.	— H. h.	
63	echinatus. L.B. (<i>bh.</i>) hedgehog.	Britain.	— H. h.	
64	dumetorum. W.N. (<i>wh.</i>) bushy.	— H. h.	
65	tetraphyllus. W.en. (<i>w.</i>) four-leaved.	N. America.	1812.	6. 7. H. h.	
66	parvifolius. DC. (<i>pu.</i>) small-leaved.	China.	1818.	6. 8. H. h.	Bot. reg. t. 496.
67	canadensis. DC. (<i>wh.</i>) purple-stalked.	N. America.	1811.	6. 7. H. h.	Eng. bot. v. 32. t. 2233.
68	saxatilis. DC. (<i>wh.</i>) stone.	Britain.	— H. 4.	
69	triflorus. Rich. (<i>wh.</i>) three-flowered.	N. America.	— H. 4.	
70	arcticus. DC. (<i>cr.</i>) dwarf crimson.	Scotland.	5. 8. H. 4.	Eng. bot. v. 23. t. 1585.
71	pistillatus. Ph. (<i>ro.</i>) close-styled.	Labrador.	1802.	6. 7. H. 4.	Sm. exot. bot. 2. t. 86.
72	obovalis. DC. (<i>wh.</i>) oboval-leaved.	N. America.	1823.	5. 7. H. 4.	
73	parviflorus. DC. (<i>wh.</i>) Vine-leaved.	1824.	— H. 4.	
74	stellatus. DC. (<i>ro.</i>) star-flowered.	1823.	6. 7. H. 4.	Sm. ic. fasc. 3. t. 64.

75	<i>Chamæmorus</i> , DC. (<i>wh.</i>) Cloud-berry.	Britain.	6. 7. H. ȝ.	Eng. bot. v. 10. t. 716.
76	<i>floridus</i> . DC. (<i>wh.</i>) large-flowered.	N. America.	1827.	6. 8. H. ȝ.	
77	<i>odoratus</i> . DC. (<i>re.</i>) flowering raspberry.	—	1700.	6. 9. H. ȝ.	Bot. mag. t. 323.
78	<i>reflexus</i> . DC. (<i>wh.</i>) reflexed.	China.	1818.	6. 7. G. ȝ.	Bot. reg. t. 461.
79	<i>Hamiltonianus</i> , DC. (<i>wh.</i>) Hamilton's. <i>rugosus</i> . D.P. non Smith.	Nepaul.	—	— H. ȝ.	
80	<i>mollucanus</i> . DC. (<i>wh.</i>) Moluccas.	Amboyna.	1810.	— S. ȝ.	Rumph. amb. 5. t. 47. f. 2.
81	<i>rugosus</i> . DC. (<i>wh.</i>) rugose.	S. America.	1824.	— S. ȝ.	
82	<i>tiliaceus</i> . DC. (<i>wh.</i>) Lime-tree-ld.	Nepaul.	—	— H. ȝ.	
83	<i>paniculatus</i> . DC. (<i>wh.</i>) paniced.	—	1826.	7. 9. H. ȝ.	
84	<i>betulinus</i> . DC. (<i>wh.</i>) Birch-leaved.	—	1825.	— H. ȝ.	
DALIBAR'DA. DC.	DALIBAR'DA.	Icosandria Pent-Decagynia.			
1	<i>rêpens</i> . DC. (<i>wh.</i>) Violet-leaved. <i>violaceoides</i> . M.	N. America.	1768.	5. 6. H. ȝ.	Mich. amer. 1. t. 27.
FRAG'ARIA. DC.	STRAWBERRY.	Icosandria Polygynia.			
1	<i>monophylla</i> . s.s. simple-leaved.	—	1773.	5. 6. H. ȝ.	Bot. mag. t. 63.
2	<i>vésca</i> . DC. (<i>wh.</i>) wood.	Britain.	4. 6. H. ȝ.	Eng. bot. v. 22. t. 1524.
<i>a sylvestris</i> . DC. (<i>wh.</i>) wild.	—	—	—	— H. ȝ.	
β <i>semperflorens</i> . DC. (<i>wh.</i>) ever-bearing.	—	—	4. 12.	H. ȝ.	Nois. jard. fr. t. 11. f. 2.
γ <i>álba</i> . (<i>wh.</i>) white-fruited.	—	—	—	— H. ȝ.	
δ <i>minor</i> . DC. (<i>wh.</i>) small English.	Britain.	—	—	— H. ȝ.	
ϵ <i>hortensis</i> . DC. (<i>wh.</i>) garden.	—	—	—	— H. ȝ.	
ζ <i>effágellis</i> . DC. (<i>wh.</i>) short-shooted.	—	—	—	— H. ȝ.	
η <i>multiplex</i> . DC. (<i>wh.</i>) double-blossomed.	—	—	4. 6.	H. ȝ.	Blackw. herb. t. 77. f. 3.
θ <i>botryfórmis</i> . DC. (<i>wh.</i>) crown-flowered.	—	—	—	— H. ȝ.	
ι <i>muricata</i> . DC. (<i>gr.</i>) monstrous.	—	—	—	— H. ȝ.	
3	<i>calycina</i> . DC. (<i>wh.</i>) large-calyxed.	Britain.	—	— H. ȝ.	
4	<i>collina</i> . DC. (<i>wh.</i>) Alpine.	Germany.	1768.	4. 11. H. ȝ.	
<i>a rúbra</i> . (<i>wh.</i>) red Alpine.	—	—	—	— H. ȝ.	
β <i>álba</i> . (<i>wh.</i>) white Alpine.	—	—	—	— H. ȝ.	
5	<i>platanoides</i> . DC. (<i>bh.</i>) Plane-leaved.	N. America.	4. 6. H. ȝ.	
6	<i>Majauéfa</i> . DC. (<i>wh.</i>) plaited-leaved.	France.	—	— H. ȝ.	
<i>a bifera</i> . DC. (<i>wh.</i>) double-bearing.	—	—	—	— H. ȝ.	
β <i>dùbia</i> . DC. (<i>wh.</i>) flat-fruited.	—	—	—	— H. ȝ.	
7	<i>Breslingea</i> . DC. (<i>wh.</i>) leathery-leaved.	—	—	— H. ȝ.	
<i>a abortiva</i> . DC. (<i>wh.</i>) sterile.	—	—	—	— H. ȝ.	
β <i>nigra</i> . DC. (<i>wh.</i>) black-fruited.	—	—	—	— H. ȝ.	
γ <i>pendula</i> . DC. (<i>wh.</i>) pear-fruited.	—	—	—	— H. ȝ.	
δ <i>hispida</i> . DC. (<i>wh.</i>) Longchamp.	France.	—	—	— H. ȝ.	
ϵ <i>viridis</i> . DC. (<i>wh.</i>) green-fruited.	—	—	—	— H. ȝ.	Nois. jard. fr. t. 13. f. 2.
ζ <i>pratensis</i> . DC. (<i>wh.</i>) field.	France.	—	—	— H. ȝ.	
8	<i>elatior</i> . DC. (<i>wh.</i>) tall.	Europe.	—	— H. ȝ.	
9	<i>moschata</i> . Ehr. (<i>wh.</i>) Hautboy.	England.	—	5. 6. H. ȝ.	Eng. bot. v. 31. t. 2197.
<i>elatior</i> . E.B.					
β <i>dioica</i> . DC. (<i>wh.</i>) dioecious.	—	—	—	— H. ȝ.	
10	<i>virginiana</i> . DC. (<i>wh.</i>) scarlet.	N. America.	1829.	4. 6. H. ȝ.	Nois. jard. fr. t. 12. f. 2.
β <i>hybridia</i> . DC. (<i>wh.</i>) Keen's seedling.	Hybrid.	—	—	— H. ȝ.	
11	<i>chilensis</i> . DC. (<i>wh.</i>) Chile.	S. America.	1727.	5. 6. H. ȝ.	Dill. elth. t. 20. f. 140.
12	<i>grandiflora</i> . DC. (<i>wh.</i>) Pine.	Surinam.	1759.	— H. ȝ.	Mill. ic. t. 288.
β <i>anánassa</i> . DC. (<i>wh.</i>) Pine apple.	—	—	—	— H. ȝ.	Nois. jard. fr. t. 14. f. 2.
γ <i>calyculata</i> . DC. (<i>wh.</i>) Bath scarlet	—	—	—	— H. ȝ.	— t. 11. f. 1. t. 14. f. 1.
δ <i>tincta</i> . DC. (<i>wh.</i>) red-fleshed.	—	—	—	— H. ȝ.	
13	<i>canadénensis</i> . DC. (<i>wh.</i>) Canadian.	Canada.	—	— H. ȝ.	
14	<i>índica</i> . DC. (<i>ye.</i>) yellow-flower'd.	Nepaul.	1805.	5. 10. H. ȝ.	Bot. reg. t. 61.
	<i>Duchesnea fragarioides</i> . Smith Linn. trans. 10. p. 373.				
COMARO'PSIS. DC.	COMARO'PSIS.	Icosandria Pent-Octagynia.			
1	<i>fragarioides</i> . DC. (<i>ye.</i>) strawberry-like.	N. America.	1803.	5. 6. H. ȝ.	Mich. amer. 1. t. 28.
	<i>Dalibarda fragarioides</i> . M.				
2	<i>Doniana</i> . DC. (<i>ye.</i>) Don's.	—	—	— H. ȝ.	Bot. mag. t. 1567.
	<i>Dalibarda fragarioides</i> . B.M. non Mich.	<i>Waldsteinia Doniana</i> . Tratt.			
3	<i>pedata</i> . DC. (<i>wh.</i>) pedate.	—	1827.	— H. ȝ.	Smith. ic. ined. t. 63.
	<i>Rubus pedatus</i> . Sm. <i>Dalibarda pedata</i> . s.s.				
HORK'ELIA. B.M.	HORK'ELIA.	Decandria Polygynia.			
congesta. B.M.	(<i>wh.</i>) tufted-flowered.	N. America.	1827.	7. 9. H. ȝ.	Bot. mag. t. 2880.
WALDSTEINIA. DC.	WALDSTEINIA.	Icosandria Diginia.			
geoídes. DC.	(<i>ye.</i>) Avens-leaved.	Hungary.	1804.	6. 7. H. ȝ.	Lodd. bot. cab. 402.
COL'URIA. B.	COL'URIA.	Icosandria Polygynia.			
potentilloídes. B.	(<i>or.</i>) Siberian.	Siberia.	1780.	6. 7. H. ȝ.	Jacq. vind. 3. t. 68.
Géum potentilloídes. w.	<i>Drýas geoídes</i> . Pall.	<i>Sievérsia geoídes</i> , s.s.			

ROSACEÆ.

SIEVER'SIA. B.

SIEVER'SIA. B.	SIEVER'SIA.	Icosandria Polygynia.	N. America.	1827.	6. 7. H. 4.	Bot. mag. t. 2863.
1 Péckii. B.M.	(ye.) Peck's.		Austria.	1597.	5. 9. H. 4.	
2 radiata.	(ye.) radiate.			1822.	— H. 4.	
Gènum radiatum. Ph.						
3 triflora. B.M.	(ye., bh.) three-flowered.			1827.	— H. 4.	Bot. mag. t. 2858.
4 montana. s.s.	(ye.) mountain.	Austria.		1597.	5. 9. H. 4.	Jacq. aust. 4. t. 373.
β minor.	(ye.) smaller mountain.	Alps.			— H. 4.	Barrel. rar. t. 399.
5 glaciális. s.s.	(ye.) arctic.	Siberia.	1827.	— H. 4.		Brown ch. mel. p. 18. t. C.
6 Róssii. B.	(ye.) Ross's.	Melville Isl.			— H. 4.	
7 anemonoídes. s.s. (wh.)	Anemone-like.	Kamtschatka.	1824.		— H. 4.	Jacq. aust. 5. ap. t. 22.
8 réptans. s.s.	(ye.) creeping.	Switzerland.	1775.	6. 8. H. 4.		
G'EM. B.	A VENS.	Icosandria Polygynia. L.				
1 agrimonoides. Ph. (wh.)	Agrimony-leaved.	N. America.	1811.	6. 7. H. 4.	Murr. com. gæt. 5. t. 2?	
2 hispidum. s.s.	(wh.) hispid.	Sweden.	1828.	— H. 4.	Reich. ic. cent. t. 3.	
3 strictum. w.	(ye.) upright.	N. America.	1778.	5. 7. H. 4.	Jacq. ic. 1. t. 93.	
aléppicum. Jacq.						
4 intermédia. w.	(ye.) intermediate.	Europe.	1794.	5. 8. H. 4.	W. hort. ber. 1. t. 69.	
5 hy'bridum. Jacq.	(ye.) hybrid.		— H. 4.	Jacq. ic. 1. t. 94.	
6 rivale. w.	(ye., bh.) water.	Britain.	5. 9. H. 4.	Eng. bot. 2. t. 106.	
7 pyrenaicum. dc.	(ye.) Pyrenean.	Pyrenees.	1804.	6. 7. H. 4.	Lam. ill. t. 44.	
8 brachypétalum. dc. (ye., bh.)	short-petaled.	1826.	5. 8. H. 4.		
9 sylvaticum. dc.	(ye.) wood.	S. Europe.	1820.	6. 7. H. 4.		
altanticum. Desf. atl. 1. p. 401.	biflorum. Brot.					
10 Thomasianum. dc. (ye.)	Thomas's.	Pyrenees.	1828.	— H. 4.		
11 Quéllyon. B.F.G. (sc.)	scarlet Chile.	Chile.	1826.	5. 10. H. 4.	Swt. br. fl. gar. t. 292.	
coccineum. Bot. reg. t. 1088.	non Flor. græc.					
β grandiflorum. (sc.)	large-flowered.		1829.	— H. 4.		
12 macrophyllum. dc.	(ye.) large-leaved.	Kamtschatka.	1804.	6. 7. H. 4.	Murr. com. gæt. 5. t. 4.	
13 canadense. dc.	(ye.) Canadian.	Canada.	1778.	5. 8. H. 4.		
14 heterophyllum. dc. (wh.)	various-leaved.	— H. 4.		
15 Besserianum.	(ye.) Besser's.	Volhynia.	1828.	— H. 4.		
intermédia. Besser non Willd.						
16 hirsutum. L.en. (st.)	hairy.	N. America.	1818.	5. 7. H. 4.		
17 virginianum. dc. (wh.)	Virginian.		7. 8. H. 4.	Herm. parad. t. 111.	
18 álbum. dc.	(wh.) white-flowered.		1730.	— H. 4.	Jacq. vind. 2. t. 175.	
canadense. Jacq. non Murr.						
19 urbánum. dc.	(ye.) common.	Britain.	5. 8. H. 4.	Eng. bot. v. 20. t. 1400.	
20 ranunculoides. dc.	(ye.) Ranunculus-like.		1827.	— H. 4.		
DR'YAS. DC.	DR'YAS.	Icosandria Polygynia. L.				
1 octopétala. dc.	(wh.) mountain.	Britain.	7. 8. H. 4.	Eng. bot. v. 7. t. 451.	
2 integrifolia. dc.	(wh.) entire-leaved.	Greenland.	1824.	4. 8. H. 4.	Hook. ex. flor. t. 220.	
POTENTILLA. DC.	CINQUEFOIL.	Icosandria Polygynia. L.				
1 nivea. dc.	(ye.) snowy-leaved.	N. Europe.	1816.	6. 8. H. 4.	Lodd. bot. cab. 460.	
2 angustifolia. dc.	(ye.) narrow-leaved.	Siberia.	1826.	— H. 4.	Lehm. pot. t. 19.	
3 macrántha. dc.	(ye.) great-flowered.		1829.	5. 8. H. 4.		
4 uniflora. dc.	(ye.) single-flowered.	Dahuria.	1827.	— H. 4.		
5 mínima. dc.	(ye.) small.	Switzerland.	1816.	4. 6. H. 4.		
6 Brauneána. N.P.	(ye.) Braune's.			— H. 4.		
7 glaciális. dc.	(ye.) very hairy.			5. 7. H. 4.		
8 frígida. dc.	(ye.) frigid.	Dauphiny.		— H. 4.		
9 grandiflora. dc.	(ye.) large-flowered.	Siberia.	1640.	6. 7. H. 4.		
10 villosa. dc.	(ye.) villous.	N. America.	1827.	— H. 4.		
11 norvégica. dc.	(ye.) Norwegian.	N. Europe.	1764.	6. 8. H. 4.		
12 Morisóni. dc.	(wh.) Morison's.	N. America.	1680.	7. 8. H. 4.		
monsplériensis. L.	Moris. his. s. 2. t. 20. f. 2. hirsuta. M.					
13 hirsuta. dc.	(ye.) hairy-leaved.	Greenland.	1820.	6. 8. H. 4.	Flor. dan. t. 1390.	
14 subacaúlis. dc.	(ye.) short-stalked.	S. France.		5. 7. H. 4.	Scop. carn. 1. t. 22.	
velutina. L.P.						
15 Tormentilla dc.	(ye.) Septfoil.	Britain.	5. 10. H. 4.	Eng. bot. v. 12. t. 863.	
Tormentilla officinalis. E.B.	erecta. W.					
16 nemorális. N.P.	(ye.) creeping-wood.		6. 8. H. 4.	Eng. bot. v. 12. t. 864.	
Tormentilla réptans. E.B.						
17 umbròsa. dc.	(go.) shaded.	Tauria.	1821.	— H. 4.	Eng. bot. v. 12. t. 862.	
18 réptans. dc.	(ye.) common.	Britain.	— H. 4.		
19 bithy'nica. dc.	(ye.) Bithynian.	Bithynia.	1822.	— H. 4.		
20 simplex. dc.	(ye.) sarmentose.	N. America.	1804.	5. 8. H. 4.		
sarmen'tosa. W.en. caroliniána. Poiret.						
21 canadensis. dc.	(ye.) Canadian.	Canada.	1818.	— H. 4.	Nestl. pot. t. 10. f. 1.	
22 vérna. E.B.	(ye.) spring.	Britain.	3. 5. H. 4.	Eng. bot. v. 1. t. 37.	

23 <i>cinerea</i> . DC. (ye.) ash-coloured. <i>subacaulis</i> . Jacq. non L.	S. Europe. 1820.	4. 6. H. 2.	Jacq. ic. 3. t. 491.
24 <i>pátula</i> . DC. (ye.) spreading.	Hungary. 1806.	6. 8. H. 2.	W. et K. hung. 2. t. 199.
25 <i>opáca</i> . DC. (ye.) opaque-leaved.	Scotland.	— H. 2.	Ser. mus. helv. 1. t 4.f.1.
26 <i>alpéstris</i> . E.F. (ye.) alpine. <i>aúrea</i> . E.B. nec <i>aliorum</i> .	—	— H. 2.	— v. 8. t. 561.
27 <i>aúrea</i> . DC. (go.) golden.	Alps Europe.	— H. 2.	Ser. mus. helv. t. 8.f.a.b.
28 <i>salisburgénsis</i> . Jac. (ye.) broad-lobed.	Switzerland. 1825.	— H. 2.	Jacq. ic. 3. t. 490.
29 <i>Halléri</i> . DC. (ye.) Haller's.	—	— H. 2.	Hall. hist. t. 21. f. 4.
30 <i>Thomásii</i> . DC. (ye.) silky-haired.	Italy. 1825.	— H. 2.	Ten. fl. nap. 1. t. 44.
31 <i>calábria</i> . T.N. (ye.) Calabrian.	—	— H. 2.	— 1.p. 287.t.45.
32 <i>impólita</i> . Wah. (ye.) unpolished.	Carpathian. 1818.	— H. 2.	
33 <i>Guntheri</i> . L.P. (ye.) Gunther's.	Bohemia. 1823.	— H. 2.	Lehm. pot. 97. t. 10.
34 <i>argéntea</i> . E.B. (ye.) silvery-leaved.	Britain.	— H. 2.	Eng. bot. v. 2. t. 89.
35 <i>collina</i> . DC. (ye.) hill.	Germany. 1823.	— H. 2.	
36 <i>intermédia</i> . DC. (ye.) intermediate. <i>opáca</i> . E.B. non L.	Scotland.	— H. 2.	Eng. bot. v. 35. t. 2449.
37 <i>elongata</i> . Bes. (ye.) long-stalked.	Russia. 1827.	— H. 2.	
38 <i>incisa</i> . DC. (ye.) jagged-leaved.	—	5. 8. H. 2.	Tausch. h. can. 1. t. 10.
39 <i>canéscens</i> . DC. (ye.) canescent.	S. Europe. 1822.	6. 8. H. 2.	
40 <i>thuringiaca</i> . DC. (ye.) decumbent.	Germany. —	— H. 2.	Gmel. fl. bad. 3.n.38.f.2.
41 <i>stipuláris</i> . DC. (ye.) large-stipuled.	Siberia. 1807.	— H. 2.	Lehm. pot. t. 8.
42 <i>hírica</i> . DC. (ye.) hairy.	S. Europe. 1725.	5. 9. H. 2.	Nestl. pot. 44. t. 7.
43 <i>pedáta</i> . N.P. (ye.) pedate-leaved.	Italy. 1818.	— H. 2.	Jacq. ic. 1. t. 92.
44 <i>astracánica</i> . s.s. (ye.) Astracan.	Levant. 1787.	6. 8. H. 2.	Lehm. pot. 90. t. 9.
45 <i>taúrica</i> . s.s. (ye.) Taurian.	Tauria. 1824.	— H. 2.	Nestl. pot. 42. t. 6.
46 <i>récta</i> . s.s. (ye.) upright.	S. Europe. 1648.	— H. 2.	Noc. et Balb. fl. tic.t.10.
47 <i>obscura</i> . s.s. (ye.) obscure.	Siberia. 1816.	6. 7. H. 2.	Lehm. pot. 86. t. 9.
48 <i>laciniòsa</i> . s.s. (ye.) jagged.	Hungary. 1818.	— H. 2.	Swt. br. fl. gar. t. 124.
49 <i>atrosanguínea</i> . DC. (cr.) dark crimson.	Nepaul. 1820.	6. 9. H. 2.	— v. 3. t. 279.
50 <i>Russelliána</i> . B.F.G. (sc.) Russell's.	Hybrid. 1827.	— H. 2.	— v. 2. t. 136.
51 <i>formòsa</i> . D.P. (ro.) handsome. <i>nepalénensis</i> . Hook. ex. flor. t. 88.	Nepaul. 1820.	5. 9. H. 2.	
52 <i>ruthénica</i> . DC. (ye.) Russian.	Siberia. 1799.	5. 7. H. 2.	Moris. his. s. 2. t. 20. f. 2.
53 <i>diffusa</i> . DC. (ye.) diffuse.	—	— H. 2.	
54 <i>chrysántha</i> . L.en. (go.) golden-flowered.	—	— H. 2.	
55 <i>declinata</i> . H.H. (ye.) declining.	—	1827.	
56 <i>pùmila</i> . s.s. (ye.) dwarf.	N. America. 1818.	6. 8. H. 2.	Lodd. bot. cab. t. 706.
57 <i>Weinmannii</i> . L.C. (ye.) Weinmann's.	Thuringia. 1822.	— H. 2.	
58 <i>ornithopodioídes</i> . (ye.) Bird's-foot.	—	1827.	
59 <i>lanuginosa</i> . F. (ye.) woolly.	Caucasus. 1821.	— H. 2.	
60 <i>Sieversiána</i> . Ot. (ye.) Sievers's.	—	1829.	
61 <i>normális</i> . F. (ye.) normal.	Siberia. 1827.	— H. 2.	
62 <i>divaricáta</i> . s.s. (ye.) divaricate.	Corsica. 1824.	— H. 2.	
63 <i>fruticòsa</i> . DC. (ye.) shrubby.	England.	6. 8. H. 2.	Eng. bot. 2. t. 88.
64 <i>dahuríca</i> . N.P. (ye.) Dahurian.	Dahuria. 1824.	— H. 2.	Nestl. pot. 31. t. 1.
65 <i>floribúnida</i> . Ph. (ye.) cluster-flower'd.	N. America. 1811.	5. 9. H. 2.	Wats. dend. brit. t. 70.
66 <i>bifúrcea</i> . DC. (ye.) bifid-leaved.	Siberia. 1773.	6. 7. H. 2.	Gmel. it. 1. t. 37. f. 1.
67 <i>supina</i> . DC. (ye.) trailing.	Europe. 1696.	7. 8. H. 2.	Jacq. aust. 5. t. 406.
68 <i>pimpinelloídes</i> . DC. (ye.) Burnet-leaved.	Levant. 1758.	6. 8. H. 2.	Buxb. cent. 1. t. 48.
69 <i>geoides</i> . DC. (ye.) Avens-like.	Tauria. 1820.	— H. 2.	
70 <i>Sprengeliána</i> . DC. (ye.) Sprengel's.	Siberia. 1824.	— H. 2.	Lehm. pot. 49. t. 3.
71 <i>viscòsa</i> . DC. (ye.) clammy. <i>hispida</i> . N.P. non W.	—	1797.	Nestl. pot. t. 36.
72 <i>pensylvánica</i> . DC. (ye.) Pensylvanian.	N. America. 1725.	6. 8. H. 2.	Jacq. vind. 2. t. 189.
73 <i>pectinàta</i> . F. (ye.) pectinated.	Siberia. 1824.	— H. 2.	
74 <i>argùta</i> . Ph. (ye.) sharp-toothed.	N. America. 1818.	5. 7. H. 2.	
75 <i>multífida</i> . DC. (ye.) cut-leaved.	Siberia. 1759.	— H. 2.	Ser. mus. helv. 1. t. 8.
76 <i>agrimonióides</i> . s.s. (ye.) Agrimony-ld.	Caucasus. 1823.	6. 8. H. 2.	
77 <i>verticilláris</i> . DC. (ye.) whorled.	Dahuria. —	— H. 2.	
78 <i>sericea</i> . DC. (ye.) silky.	Siberia. 1780.	5. 6. H. 2.	Lehm. pot. 66. t. 6.
79 <i>pulchéllea</i> . DC. (ye.) neat.	Melville Isl. 1827.	5. 7. H. 2.	
80 <i>Egèdii</i> . DC. (ye.) smooth-leaved.	Greenland. —	— H. 2.	Flor. dan. t. 1578.
81 <i>ansérina</i> . DC. (ye.) wild Tansey. β <i>geminiflòra</i> . DC. (ye.) twin-flowered.	Britain.	5. 9. H. 2.	Eng. bot. 12. t. 861.
γ <i>cóncolor</i> . DC. (ye.) silky-leaved.	—	— H. 2.	Moris. his. s. 2. t. 20. f. 4.
82 <i>cánídicans</i> . DC. (ye.) hoary-leaved.	America. 1826.	— H. 2.	Nestl. pot. 34. t. 3. f. 2.
83 <i>spléndens</i> . DC. (ye.) glossy-leaved. <i>striata</i> . s.s.	Nepaul. 1818.	7. 9. H. 2.	Swt. br. fl. gar. 2. t. 150.

ROSACEÆ.

84 <i>Còmarum</i> . DC. (pu.) marsh.	Britain.	6. 8. H. ȝ.	Eng. bot. v. 3. t. 172.
<i>Còmarum palustre</i> . E.B.				
85 <i>Salesovii</i> . DC. (ye.) white shrubby.	Siberia.	1823.	— H. ȝ.	Lehm. pot. 35. t. 1.
86 <i>glábria</i> . DC. (wh.) smooth white.	—	1818.	6. 10. H. ȝ.	Lodd. bot. cab. t. 914.
87 <i>rupéstris</i> . DC. (wh.) rock.	England.	6. 8. H. ȝ.	Eng. bot. v. 29. t. 2058.
88 <i>fragarioídes</i> . DC. (wh.) Strawberry-like.	Siberia.	1773.	5. 7. H. ȝ.	Lehm. pot. 50. t. 4.
89 <i>álba</i> . DC. (wh.) white-flowered.	Wales.	6. 9. H. ȝ.	Eng. bot. v. 20. t. 1384.
90 <i>cauléscentis</i> . DC. (wh.) caulescent.	Austria.	1759.	5. 7. H. ȝ.	Jacq. aust. t. 220.
91 <i>Clausiana</i> . s.s. (wh.) <i>Clusius's</i> .	S. Europe.	1806.	7. 8. H. ȝ.	Bot. mag. t. 1327.
92 <i>lupinoídes</i> . s.s. (wh.) close-flowered.	Pyrenees.	1739.	6. 7. H. ȝ.	Lodd. bot. cab. t. 6.
93 <i>valderia</i> . DC. (st.) crowded-flower'd.	S. Europe.	1818.	6. 8. H. ȝ.	All. fl. ped. t. 24. f.
94 <i>nítida</i> . DC. (bh.) glossy.	—	1798.	— H. ȝ.	Jacq. aust. ap. t. 25.
95 <i>apennina</i> . DC. (wh.) Apennine.	Italy.	1825.	— H. ȝ.	Ten. fl. nap. 1. t. 46.
96 <i>tridentata</i> . DC. (wh.) trifid-leaved.	Scotland.	5. 7. H. ȝ.	Eng. bot. v. 34. t. 2389.
97 <i>mícrantha</i> . DC. (wh.) small-blossom'd.	S. Europe.	1816.	6. 8. H. ȝ.	Ser. mus. helv. 1. t. 5.
98 <i>Fragaria</i> . DC. (wh.) sterile strawberry.	Britain.	3. 6. H. ȝ.	Eng. bot. v. 25. t. 1785.
<i>Fragaria stérilis</i> . E.B. <i>Còmarum fragarioídes</i> . W. enum.				
99 <i>emarginata</i> . Ph. (wh.) emarginate.	Labrador.	1828.	— H. ȝ.	
100 <i>Jamesoniána</i> . DC. (ye.) Jameson's.	Greenland.	1822.	6. 7. H. ȝ.	
SIBBAL'DIA. DC. SIBBA'LDIA. Pentandria Pentagynia.				
1 <i>procumbens</i> . DC. (ye.) procumbent.	Britain.	6. 8. H. ȝ.	Eng. bot. v. 13. t. 897.
2 <i>parviflora</i> . DC. (ye.) small-flowered.	Levant.	— H. ȝ.	
3 <i>erécta</i> . DC. (ye.) upright.	Siberia.	1806.	— H. ȝ.	Amm. ruth. 112. t. 15.
4 <i>grandiflora</i> . DC. (ye.) large-flowered.	Dahuria.	1828.	— H. ȝ.	
5 <i>polygyna</i> . DC. (ye.) many-styled.	Siberia.	1824.	— H. ȝ.	
AGRIMONIA. DC. AGRIMONY. Dodecadenia Digynia.				
1 <i>Eupatòria</i> . DC. (ye.) common.	Britain.	6. 8. H. ȝ.	Eng. bot. v. 19. t. 1335.
2 <i>nepalénensis</i> . DC. (ye.) Nepaul.	Nepaul.	1820.	— H. ȝ.	
3 <i>odoráta</i> . DC. (ye.) sweet-scented.	Italy.	1640.	7. 8. H. ȝ.	
4 <i>dahúrica</i> . DC. (ye.) Dahurian.	Dahuria.	1824.	6. 8. H. ȝ.	
5 <i>pilósa</i> . DC. (ye.) hairy.	Siberia.	1820.	— H. ȝ.	
6 <i>parviflora</i> . DC. (ye.) small-flowered.	N. America.	1766.	6. 7. H. ȝ.	
7 <i>suavéolens</i> . DC. (st.) fragrant.	—	1820.	— H. ȝ.	
8 <i>striáta</i> . DC. (wh.) striped-flowered.	—	1812.	6. 8. H. ȝ.	
9 <i>rèpens</i> . DC. (ye.) creeping-rooted.	Levant.	1737.	7. 9. H. ȝ.	
AREMONTIA. DC. AREM'ONIA. Pent-Decandria Digynia.				
agrimonoídes. DC. (ye.) Agrimony-like.	Italy.	1739.	6. 8. H. ȝ.	Col. ecphr. 1. t. 144.
<i>Spallanzánia agrimonoides</i> . s.s. <i>Agrimónia agrimonoides</i> . L.				

Subordo III. SANGUISORBEÆ. DC. prodr. v. 2. p. 588.

ALCHIMILLA. DC. LADIES MANTLE. Di-Tetrandria Monogynia. L.				
1 <i>capénsis</i> . DC. (gr.) Cape.	C. B. S.	1818.	4. 6. G. ȝ.	Lam. ill. t. 86. f. 2.
2 <i>vulgáris</i> . DC. (gr.) common.	Britain.	6. 8. H. ȝ.	Eng. bot. v. 9. t. 597.
β <i>variegáta</i> .	(gr.) variegated-leav'd.	— H. ȝ.	
γ <i>glábria</i> . DC. (gr.) smooth.	Pyrenees.	— H. ȝ.	
3 <i>hybrida</i> . DC. (gr.) mountain.	Saxony.	— H. ȝ.	Pluk. t. 240. f. 2.
montánu. W. en.				
4 <i>pubéscens</i> . DC. (gr.) pubescent.	Caucasus.	1813.	— H. ȝ.	Willd. hort. ber. 2. t. 79.
5 <i>alpina</i> . DC. (gr.) silvery.	Britain.	7. H. ȝ.	Eng. bot. v. 4. t. 244.
6 <i>sericea</i> . DC. (gr.) silky.	Caucasus.	1813.	6. 8. H. ȝ.	Bocc. mus. p. 18. t. 1.
7 <i>pentaphylla</i> . DC. (gr.) five-leaved.	Switzerland.	1784.	7. H. ȝ.	H. et B. n. g. am. 6. t. 561.
8 <i>sibbaldiaefolia</i> . DC. (gr.) Sibbaldia-ld.	Mexico.	1823.	— F. ȝ.	Eng. bot. v. 15. t. 1011.
9 <i>arvénsis</i> . DC. (gr.) Parsley pier.	Britain.	4. 6. H. ⊕.	
<i>Aphanes</i> . w. <i>Aphanes arvénsis</i> . L.				
10 <i>cornucopioídes</i> . DC. (gr.) Spanish.	Spain.	1825.	4. 7. H. ⊕.	
CEPHAL'OTUS. DC. CEPHAL'OTUS. Dodecadria Hexagynia.				
folícularis. DC. (wh.) pitcher-plant.	N. Holland.	1822.	4. 6. G. w. ȝ.	Lab. nov. holl. 2. t. 145.
MARGYRICA'RPU'S. DC. MARGYRICA'RPU'S. Diandria Monogynia.				
setósus. DC. (gr.) bristle-pointed.	S. America.	1829. F. ȝ.	Flor. peruv. 1. t. 8. f. d.
AC'ENA. Kth.	Ac'ENA.	Tri-Pentandria Monogynia.		
1 <i>elongata</i> . DC. (gr.) long-spiked.	Mexico.	1827.	4. 8. F. ȝ.	
2 <i>latebròsa</i> . DC. (gr.) hairy.	C. B. S.	1774.	4. 6. G. ȝ.	Gärt. fruct. 1. t. 32.

3 pinnatifida. DC. (gr.) pinnatifid.	Chili.	1823.	4. 6.	G. 4.	Fl. per. 1. t. 104. f. 1. b.
4 sericea. DC. (gr.) silky.	Port Desire.	—	G. 4.	Jacq. f. ecl. 1. t. 55.
ANCI'STRUM. Kth. ANCI'STRUM.	Di-Pentandria Monogynia.				
1 sanguisorbae. L. (gr.) Burnet-leaved.	N. Zealand.	1796.	6.	F. 4.	Gært. fruct. t. 32.
2 ovalifolium. DC. (gr.) oval-leaved.	Peru.	1802.	5. 6.	F. 4.	Vent. h. cels. t. 5.
<i>rēpens</i> . Vent.					Fl. per. 1. t. 103. f. b.
3 argenteum. K.S. (gr.) silvery-leaved.	S. America.	1823.	—	F. ½.	Lam. ill. t. 22. f. 2.
4 adscendens. DC. (gr.) ascending.	Magellan.	—	—	H. 4.	— t. 22. f. 3.
5 lavigatum. DC. (gr.) smooth	—	1790.	6. 8.	H. 4.	
6 magellanicum. DC. (gr.) Magellan.	—	—	—	H. 4.	
7 lucidum. Lam. (gr.) shining.	Falkland Isl.	1777.	5. 6.	H. 4.	
ANGUISO'RBA. DC. GREAT-BURNET.	Tetrandria Monogynia. L.				
1 officinalis. DC. (pu.) officinal.	Britain.	6. 8.	H. 4.	Eng. bot. v. 19. t. 1312.
β auriculata. DC. (pu.) eared.	Italy.	—	H. 4.	Bocc. mus. 19. t. 9.
2 carnea. DC. (pi.) flesh-coloured.	—	1818.	—	H. 4.	Schrank. h. mon. t. 69.
3 rūbra. Sck. (re.) red-spiked.	Dahuria.	1824.	—	H. 4.	
4 præcox. Bes.	(pi.) early-flowered.	Siberia.	1827.	5. 7.	
5 angustifolia. F.	(pi.) narrow-leaved.	—	1820.	6. 8.	
6 tenuifolia. DC.	(re.) slender-leaved.	Dahuria.	—	—	
7 íngrica.	(wh.) glossy-leaved.	—	1829.	—	
8 mauritánica. DC. (gr.) Barbary.	Algiers.	—	H. 4.	Moris. ox. s. 8. t. 18. f. 4.
9média. DC. (wh.) short-spiked.	Canada.	1785.	—	H. 4.	— s. 8. t. 18. f. 8.
10 canadensis. DC. (wh.) Canadian.	—	1633.	—	H. 4.	— s. 8. t. 18. f. 12.
POTERIUM. DC.	BURNET. Monocotyledon Polyandria. L.				Moris. ox. s. 8. t. 18 f. 5.
1 spinosum. DC. (wh.) prickly shrubby.	Levant.	1595.	4. 8.	G. ½.	
β crispum. DC. (wh.) curled-leaved.	—	—	—	G. ½.	Bot. mag. 2341.
γ inerme. DC. (wh.) spineless.	—	—	—	G. ½.	Eng. bot. v. 12. t. 860.
2 caudatum. DC. (wh.) shrubby.	Canaries.	1779.	1. 4.	G. ½.	Barrel. ic. t. 632.
3 Sanguisorba. DC. (pu.) common.	England.	7.	H. 4.	W. et K. hung. 2. t. 198.
4 agrimonifolium. DC. (pu.) Agrimony-ld.	Spain.	1818.	6. 7.	H. 4.	Flor. græc. t. 942.
5 hy'bridum. W. (pu.) sweet.	France.	1683.	—	H. 4.	
6 polygamum. DC. (pu.) Hungarian.	Hungary.	1803.	7. 8.	H. 4.	
7 villosum. DC. (pu.) villous.	Levant.	1829.	—	H. 4.	
8 verrucosum. E. (pu.) warted.	Egypt.	1827.	—	H. 4.	
CERCOCAR'PUS. DC. CERCOCAR'PUS.	Icosandria Monogynia.				
fothergilloides. DC. (pu.) Fothergilla-like.	Mexico.	1828.	—	G. ½.	H. et B. n. gen. 6. t. 559.
CLIFFO'R'TIA. DC.	CLIFFO'R'TIA. Dicoccia Polyandria. L.				
1 ilicifolia. DC. (st.) Holly-leaved.	C. B. S.	1714.	5. 9.	G. ½.	Linn. hort. cliff. t. 30.
2 cordifolia. DC. (st.) heart-leaved.	—	1816.	—	G. ½.	
3 tridentata. DC. (st.) three-toothed.	—	1752.	6. 7.	G. ½.	Linn. hort. cliff. t. 31.
4 ruscifolia. DC. (st.) Ruscus-leaved.	—	—	—	G. ½.	
arachnoidea. B.C. 260.					
5 graminea. DC. (st.) grass-leaved.	—	1825.	—	G. ½.	
6 ferruginea. DC. (st.) ferruginous.	—	1823.	—	G. ½.	Pluk. alm. t. 275. f. 2.
7 cuneata. DC. (st.) wedge-leaved.	—	1787.	4.	G. ½.	
8 strobilifera. DC. (st.) cone-bearing.	—	1816.	5. 7.	G. ½.	
9 ericæfolia. DC. (st.) heath-leaved.	—	1799.	7. 9.	G. ½.	
10 sarmentosa. DC. (st.) twiggy.	—	1795.	6. 8.	G. ½.	
11 falcatæ. DC. (st.) falcate-leaved.	—	1816.	—	G. ½.	
12 ternata. DC. (st.) ternate-leaved.	—	—	—	G. ½.	Linn. hort. cliff. t. 32.
13 trifoliata. DC. (st.) three-leaved.	—	1752.	4. 7.	G. ½.	Pluk. alm. t. 319. f. 4.
14 obcordata. DC. (st.) obcordate.	—	1790.	6. 8.	G. ½.	
15 oblîqua. DC. (st.) oblique-leaved.	—	1816.	—	G. ½.	
16 dentata. DC. (st.) toothed.	—	—	—	G. ½.	
17 crenata. DC. (st.) notched-leaved.	—	1791.	7. 8.	G. ½.	
18 pulchella. DC. (st.) beautiful.	—	1795.	4. 5.	G. ½.	
19 cinerea. DC. (st.) cinereous.	—	1800.	6. 7.	G. ½.	

ORDO LXXIX.

SPIRÆACEÆ. D. Don prodr. flor. nepal. p. 227.

PURSHIA. DC. PU'RSHIA. Icosandria Monogynia.

tridentata. DC. (ye.) three-toothed. Columbia. 1827. H. ½. Pursh. fl. amer. 1. t. 15.

Tigæra tridentata. Ph.

KE'RRIA. DC.	KE'RRIA. Icosandria Pent-Octagynia.						
japónica. DC. (ye.) Japanese.	Japan.	1804.	4. 9.	H. h.			
β flôre plêno. (ye.) double-flowered.				H. h.			Andr. deposit. t. 587.
Côrchorus japónicus. Bot. mag. t. 1296.							
SPIRÆA. DC.	SPIRÆA. Icosandria Pentagynia. L.						
opulifolia. DC. (wh.) Guelder-rose-ld.	N. America.	1690.	6. 7.	H. h.			Schmidt arb. 1. t. 52.
capitata. DC. (wh.) headed.	Columbia.	1827.	—	H. h.			
ulmifolia. DC. (wh.) Elm-leaved.	Siberia.	1790.	—	H. h.			Jacq. vind. 2. t. 140.
flexuosa. DC. (wh.) flexuose.		1820.	—	H. h.			Camb. mon. spir. t. 26.
bella. DC. (ro.) red-flowered.	Nepaul.	1818.	6. 9.	H. h.			Bot. mag. t. 2426.
chamaedrifolia. DC. (wh.) Germanander-ld.	Siberia.	1780.	6. 7.	H. h.			Bot. reg. t. 1222.
média. Ph. (wh.) middle.	Canada.	1827.	—	H. h.			
oblongifolia. W.K. (wh.) oblong-leaved.	Hungary.	1815.	5. 7.	H. h.			W. et K. hung. 3. t. 235.
câna. DC. (wh.) hoary-leaved.		1824.	6. 7.	H. h.			3. t. 227.
trilobata. DC. (wh.) three-lobed.	Siberia.	1801.	5. 6.	H. h.			Pall. ross. 1. t. 17.
alpina. DC. (wh.) Alpine.		1806.	6. 7.	H. h.			1. t. 20.
thalictroides. DC. (wh.) meadow-rue-ld.		1790.	5. 6.	H. h.			I. t. 18.
pikowiensis. DC. (wh.) Pikow.	Poland.	1824.	—	H. h.			
hypericifolia. W. (wh.) Hypericum-ld.	N. America.	1640.	4. 5.	H. h.			Schmidt arb. 1. t. 26.
uralensis. (wh.) Fischer's.	Siberia.	1822.	—	H. h.			
crenata. F. non W. hypericifolia β uralensis. DC.							
acutifolia. W.en. (wh.) acute-leaved.	Siberia.		—	H. h.			
crenata. L. (wh.) crenated.		1739.	—	H. h.			
crataegifolia. L.en. (wh.) Hawthorn-leav'd.		1824.	5. 6.	H. h.			Lodd. bot. cab. t. 671.
obovata. W.K. (wh.) obovate leaved.	Hungary.	1816.	—	H. h.			t. 1403.
savrânicâ. Bes. (wh.) small-flowered.	Poland.	1826.	—	H. h.			Pall. ross. 1. t. 16.
Besseriana. (wh.) Besser's.			—	H. h.			Schmidt arb. 1. t. 49.
crenata. Bes. non L.							Eng. bot. v. 21. t. 1468.
canescens. DC. (wh.) canescent.	Nepaul.	1825.	6. 8.	H. h.			Wats. dend. brit. t. 133.
ceanothifolia. DC. (wh.) Ceanothus-ld.		1820.	5. 6.	H. h.			t. 66.
corymbosa. DC. (wh.) corymbose.	Virginia.	1812.	—	H. h.			Schmidt arb. 1. t. 51.
vacciniifolia. D.P. (wh.) Vaccinium-ld.	Nepaul.	1824.	6. 8.	H. h.			
betulaefolia. DC. (wh.) Birch-leaved.	Siberia.	1812.	6. 7.	H. h.			
lævigata. DC. (wh.) smooth-leaved.		1774.	4. 6.	H. h.			
salicifolia. DC. (ro.) Willow-leaved.	Britain.		—	H. h.			
álba. W.D. (wh.) white-flowered.			—	H. h.			
carpinifolia. W.en. (bh.) broad-leaved.	N. America.		7. 8.	H. h.			
tomentosa. DC. (ro.) tomentose.		1736.	7. 9.	H. h.			
ariæfolia. DC. (ro.) Aria-leaved		1827.	6. 8.	H. h.			
discolor. DC. (wh.) two-coloured.			—	H. h.			
sorbifolia. DC. (wh.) Service-leaved.	Siberia.	1759.	7. 9.	H. h.			Schmidt arb. 1. t. 58.
grandiflora. (wh.) large-flowered.	Kamtschatka.	1817.	7. 8.	H. h.			Pall. ros. 1. t. 25.
sorbifolia β alpina. DC.							
Aráncus. DC. (wh.) Goat's-beard.	Siberia.	1633.	6. 7.	H. Ȑ.			Pall. ros. 1. t. 26.
americana.	(wh.) glossy-leaved.	N. America.	1825.	—	H. Ȑ.		
Ulmâria. DC. (wh.) Meadow-sweet.	Britain.	6. 10.	H. Ȑ.			Eng. bot. v. 14. t. 960.
a tomentosa. DC. (wh.) tomentose.		—	H. Ȑ.			
β mûltiplex. DC. (wh.) double-flowered.		—	H. Ȑ.			
γ variegata. DC. (wh.) variegated-leav'd.		—	H. Ȑ.			
lobata. DC. (ro.) lobed-leaved.	N. America.	1765.	7. 8.	H. Ȑ.			Jacq. vind. 1. t. 88.
palmata. DC. (wh.) palmated.	China.	1823.	—	H. Ȑ.			
digitata. DC. (ro.) digitate.	Siberia.		—	H. Ȑ.			
Filipendula. DC. (wh.) Dropwort.	Britain.	6. 10.	H. Ȑ.			Pall. ros. 1. t. 27.
β mûltiplex. DC. (wh.) double-flower'd.		—	H. Ȑ.			Eng. bot. v. 4. t. 284.
GILLE'NIA. DC.	GILLE'NIA. Dodecadria Pentagynia.						
trifoliata. DC. (bh.) three-leaved.	N. America.	1713.	6. 7.	H. Ȑ.			Bot. mag. t. 489.
Spiraea trifoliata. B.M.							
stipulacea. DC. (bh.) large-stipuled.		1805.	—	H. Ȑ.			Bart. med. bot. t. 6.

ORDO LXXX.

CALYCANTHEÆ. DC. prodr. 3. p. 1.

CALYCA'NTHUS. DC. ALLSPICE-TREE. Icosandria Polygynia. L.

1 floridus. DC. (d. pu.) Carolina. Carolina. 1726. 5. 8. H. h.

Bot. mag. t. 503.

2 prunifolius.	(d.pu.) Plum-leaved.	Carolina.	1822.	5. 8. H. h.	
3 glaucus. DC.	(d.pu.) glaucous-leaved.	—————	1806.	— H. h.	Guimp. ab. holz. t. 5.
<i>fértilis.</i> Bot. reg. t. 404.					
4 oblongifolius. N.	(d.pu.) oblong-leaved.	—————	1812.	— H. h.	
5 lavigatus. DC.	(d.pu.) smooth-leaved.	—————	1806.	— H. h.	Bot. reg. t. 481.
CHIMONA'NTHUS. DC.	JAPAN ALLSPICE.	Icosandria Polygynia.			
frágans. DC.	(ye.) fragrant.	Japan.	1766.	12. 3. H. h.	Bot. mag. t. 466.
<i>Calycanthus pr'acox.</i> B.M.					
<i>a verus.</i>	(ye.pu.) common.	—————	—————	H. h.	Hort. Kew. v. 2. t. 10.
<i>β luteus.</i>	(ye.) yellow-flowered.	—————	1814.	— H. h.	
<i>γ grandiflorus.</i> B.R. (ye.) large-flowered.	—————	—————	1812.	— H. h.	Bot. reg. t. 451.

ORDO LXXXI.

GRANATEÆ. D. Don in Jameson Edinb. phil. journ. jul. 1826. p. 134.

P'UNICA. DC.	POMEGRANATE.	Icosandria Monogynia. L.			
1 Granátum. DC.	(or.) common.	S. Europe.	1548.	6. 9. H. h.	Bot. mag. t. 1832.
<i>α rùbrum.</i> DC.	(or.) red-flowered.	—————	—————	H. h.	Schkuhr handb. t. 131.
<i>β flòre plèno.</i> DC.	(or.) double red.	—————	6. 8. H. h.	Trew ehret. t. 71. f. 2.
<i>γ flavescens.</i>	(ye.) yellow-flowered.	China.	1810.	— F. h.	
<i>δ albescens.</i>	(wh.) white-flowered.	—————	—————	F. h.	Andr. deposit. t. 96.
<i>ε múltiplex.</i>	(wh.) double white.	—————	F. h.	
2 nàna. DC.	(or.) dwarf.	S. America.	1723.	7. 9. F. h.	Bot. mag. t. 634.

ORDO LXXXII.

MEMECYLEÆ. DC. prodr. 3. p. 5.

MEM'ECYLON. DC.	MEM'ECYLON.	Octandria Monogynia. L.			
1 angulatum. DC.	(pu.) square-stalked.	Mauritius.	1826.	5. 7. S. h.	
2 capitellatum. DC.	(pu.) headed.	Ceylon.	1796.	7. 8. S. h.	Burm. zeyl. t. 30.
3 èdule. DC.	(pu.) eatable-fruited.	E. Indies.	1823.	— S. h.	Roxb. corom. 1. t. 82.
4 grande. DC.	(bl.) tree.	—————	1824. S. h.	Rheed. mal. 2. t. 14.
MOURIRIA. DC.	MOURIRIA.	Decandria Monogynia.			
1 myrtloides. DC.	(wh.) sessile-leaved.	Jamaica.	1818.	6. 8. S. h.	Sloan. his. 2. t. 87. f. 3.
<i>Petaloma myrtilloides.</i> Swartz fl. ind. occ. 2. p. 833. t. 14.					
2 guianensis. DC.	(ye.) Guiana.	Guiana.	1825.	— S. h.	Aubl. guian. 1. t. 180.
3 grandiflora. DC.	(wh.) great flowered.	Brazil.	1822. S. h.	
4 alternifolia. H.B. alternate-leav'd.	E. Indies.	1823. S. h.	

ORDO LXXXIII.

COMBRETACEÆ. DC. prodr. 3. p. 9.

Tribus I. TERMINALIEÆ. DC. prodr. 3. p. 9.

BUCIDA. DC.	OLIVE-BARK-TREE.	Decandria Monogynia. L.			
1 Búceras. DC.	(wh.) Jamaica.	Jamaica.	1793.	7. 10. S. h.	Bot. reg. t. 907.
2 capitata. DC.	(wh.) headed.	W. Indies.	1826. S. h.	Vahl. ecl. 1. t. 8.
TERMIN'ALIA. DC.	TERMIN'ALIA.	Polygamia Momocia. L.			
1 Tanibouca. DC.	(wh.) Guiana.	Guiana.	1826. S. h.	Aubl. guian. 1. t. 178.
2 angustifolia. DC.	(wh.) narrow-leaved.	E. Indies.	1692.	5. 7. S. h.	Jacq. vind. 3. t. 100.
<i>Catappa Benzoin.</i> Gært. fruct. 2. p. 206. t. 127.					

COMBRETACEÆ.

3 mauritiâna. DC.	(wh.) Mauritia.	Mauritius.	1822. S. h.	Lam. ill. t. 848. f. 2.
4 Catáppa. DC.	(wh.) large-leaved.	E. Indies.	1778.	5. 7. S. h.	Jacq. ic. 1. t. 197.
5 subcordâta. K.S.	(wh.) heart-leaved.	W. Indies.	1816. S. h.	
6 moluccâna. DC.	(wh.) smooth-leaved.	Moluccas.	— S. h.	Rumph. amb. 1. t. 68.
7 glâbrata. DC.	(wh.) smooth-leaved.	Society Isles.	1822. S. h.	Spreng. antiqu. bot. t. 2.
8 latifolia. DC.	(wh.) broad-leaved.	W. Indies.	1778.	6. 8. S. h.	
9 bengalénsis. DC.	(wh.) Bengal.	E. Indies.	1826. S. h.	
10 Bellérica. DC.	(wh.) fetid-flowered.	—	1818. S. h.	Roxb. corom. 2. t. 198.
11 Chébula. DC.	(wh.) oval-leaved.	—	1796.	6. 8. S. h.	— 2. t. 197.
12 citrina. DC.	(st.) Citron-leaved.	—	1823. S. h.	Gært. fruct. 2. t. 97.
13 pròcera. DC.	(wh.) lofty.	—	1816. S. h.	Roxb. corom. 3. t. 224.
14 gangética. H.B.	(wh.) Ganges.	—	1823. S. h.	
15 biticaria. H.B.	(wh.) Silhet.	—	1824. S. h.	
16 arbùscula. DC.	(wh.) small tree.	Jamaica.	1822. S. h.	
17 sericea. DC.	(wh.) silky-leaved.	C. B. S.	1816. I. h.	
18 Fat'ræa. DC.	(wh.) Madagascar.	Madagascar.	1826. S. h.	
PENTA'PTERA. DC.	PENTA'PTERA.	Polygamia Monogynia.			
1 tomentosa. DC.	(wh.) woolly.	E. Indies.	1824. S. h.	DC. diss. t. 1.
2 paniculâta. DC.	(wh.) panicled.	—	1826. S. h.	
3 Arjûna. DC.	(wh.) glaucous-leaved.	—	1825. S. h.	DC. diss. t. 2.
4 bialâtâ. H.B.	(wh.) two-winged.	—	— S. h.	
GETONIA. DC.	GETONIA.	Decandria Monogynia.			
1 floribûnda. DC.	(wh.) many-flowered.	E. Indies.	1815.	5. 7. S. h.	Roxb. corom. 1. t. 87.
2 nûtans. DC.	(wh.) nodding.	—	1823. S. h.	
CONOCA'RPU. DC.	BUTTON-TREE.	Pent-Decandria Monogynia.			
1 erécta. w.	(wh.) upright.	Jamaica.	1752.	6. 8. S. h.	Jacq. amer. t. 52. f. 1.
2 procùmbens. w.	(wh.) procumbent.	Cuba.	1730.	— S. h.	— t. 52. f. 2.
3 acuminâta. DC.	(wh.) pointed-leaved.	E. Indies.	1822. S. h.	
4 latifòlia. DC.	(wh.) broad-leaved.	—	1816. S. h.	
LAGUNCUL'ARIA. DC.	LAGUNCUL'ARIA.	Decandria Monogynia.			
racemôsa. DC.	(gr.) racemed.	W. Indies.	1825. S. h.	Jacq. amer. t. 53.
POIV'REA. DC.	POIV'REA.	Decandria Monogynia.			
1 comôsa.	(cr.) tufted.	Sierra Leone.	1821.	5. S. S. h.	Bot. reg. t. 1165.
Combrètum comôsum. Bot. reg.					
2 intermédia. G.D.	(cr.) intermediate.	—	— S. h.	
3 alternifòlia. DC.	(wh.) alternate-leav'd.	S. America.	1826. S. h.	Jacq.am.pic.t.260.f.27.
Combrètum decândrum. Jacq.					
4 coccinea. DC.	(sc.) scarlet-flower'd.	Madagascar.	1818.	3. 10. S. h.	Lam. ill. t. 282. f. 2.
Combrètum purpureum. Bot. reg. 429.	Cristaria coccinea. Sonn. itin. 2. t. 140.				non Pursh.
5 Roxburghii. DC.	(wh.) Roxburgh's.	E. Indies.	1822. S. h.	Roxb. corom. 1. t. 59.
Combrètum decândrum. Roxb.					non Jacq.

Tribus II. COMBRETEÆ. DC. prodr. 3. p. 18.

COMBR'ETUM. DC.	COMBR'ETUM.	Octandria Monogynia. L.			
1 secûndum. G.D.(y.e.sc.)	side-flowering.	W. Indies.	1822. S. h.	Jacq.am.pic.t.260.f.26.
2 élégans. DC.	(re.) elegant.	S. America.	1823. S. h.	Aubl. gui. 1. t. 137.
3 formosum. G.D.(y.e.re.)	beautiful.	Brazil.	1824. S. h.	
4 grandiflòrum. G.D.(sc.)	large-flowered.	Sierra Leone.	1821. S. h.	
5 farinòsum. G.D.	(or.) mealy-leaved.	Mexico.	1823. G. h.	
6 laxum. G.D.	(wh.) loose-racemed.	S. America.	1822. S. h.	Lam. ill. t. 282. f. 1.
7 pulchéllum. G.D.	(sc.) pretty.	Brazil.	1823. S. h.	
8 nânum. G.D.	(wh.) dwarf.	Nepaul.	1826. G. h.	
9 micrânthum. G.D.(sc.)	small-flowered.	Sierra Leone.	1828. S. h.	
parviflòrum. Reichenb.	hort. bot. t. 62.				
CACO'UCIA. DC.	CACO'UCIA.	Dodecadria Monogynia.			
coccinea. DC.	(sc.) scarlet.	Guiana.	1815. S. h.	Aubl. guian. 1. t. 179.
Schousb'aea coccinea. w.					
QUISQU'ALIS. DC.	QUISQU'ALIS.	Decandria Monogynia. L.			
1 índica. DC.	(er.) Indian.	China.	1815.	5. 8. S. h.	Bot. reg. t. 492.
2 glâbra. DC.	(cr.) smooth.	E. Indies.	1825.	— S. h.	Burm. Ind. t. 28. f. 2.
3 pubescens. s.s. pubescent.	China.	1815. S. h.	
4 villôsa. s.s. villous.	Pegu.	1818. S. h.	

ORDO LXXXIV.

VOCHYSIEÆ. DC. prodr. 3. p. 25.

VOCHYSIA. DC.	VOCHYSIA. Monandria Monogynia.
1 guianensis. DC. (ye.) Guiana.	Guiana. 1822. S. h. Aubl. guian. 1. t. 6.
Cucularia excelsa. W.	—
2 tomentosa. DC. (ye.) woolly.	1825. S. h.

ORDO LXXXV.

RHIZOPHOREÆ. DC. prodr. 3. p. 31.

OLISBEA. DC.	OLISBEA. Decandria Monogynia.
rhizophoræfolia. DC. Rhizophora-ld. W. Indies.	1825. S. h.
RHIZOPHORA. DC.	RHIZOPHORA. Octandria Monogynia.
Mângle. DC. (ye.) common.	India. 1818. S. h. Jacq. amer. t. 89.
CARA'LLIA. DC.	CARA'LLIA. Dodecadria Monogynia.
lùcida. DC. (ye.) glossy.	E. Indies. 1826. S. h. Roxb. corom. 3. t. 211.

ORDO LXXXVI.

ONAGRARIEÆ. DC. prodr. v. 3. p. 35.

Tribus I. MONTINIEÆ. DC. prodr. loc. cit.

MONTI'NIA. DC.	MONTI'NIA. Diocia Tetrandria. L.
ácris. DC. (wh.) Sea purslane-ld. C. B. S.	1774. 7. 8. G. h. Smith spicil. t. 15.
caryophyllacea. Thunb.	

Tribus II. FUCHSIEÆ. DC. prod. v. 3. p. 36.

FUCHSIA. DC.	FUCHSIA. Octandria Monogynia. L.
1 microphylla. DC. (re.) small-leaved.	Mexico. 1827. 6. 9. F. h. Swt. br. fl. gar. s. 2. t. 16.
2 lycoides. DC. (re.) Box-thorn-ld.	Chile. 1796. 4. 10. G. h. Bot. mag. t. 1024.
3 parviflora. DC. (re.) small-flowered.	Mexico. 1824. 6. 9. F. h. Bot. reg. t. 1048.
4 arborescens. DC. (ro.) arborescent.	— 1823. 6. 10. G. h. — t. 943.
5 gráclis. DC. (sc.) slender.	— 1822. 5. 10. F. h. — t. 847.
β multiflora. DC. (sc.) many-flowered.	— 1824. — F. h. — t. 1052.
6 macrostémma. DC. (sc.) large-crowned.	Chile. 1823. — G. h. Flor. per. 3. t. 324. f. b.
tenella. DC. (sc.) weak.	— 1822. — G. h.
7 cónica. B.R. (sc.) conical-tubed.	— 1824. — F. h. Bot. reg. t. 1062.
8 virgata. (sc.) Colvill's.	Mexico. 1825. — F. h.
9 coccinea. DC. (sc.) scarlet.	Chile. 1788. — F. h. Bot. mag. t. 97.
10 excorticata. DC. (gr. re.) changeable.	N. Zealand. 1821. — G. h. Bot. reg. t. 857.

Tribus III. ONAGREÆ. DC. prodr. v. 3. p. 40.

CHAMÆNERION. Tourn.	CHAMÆNERION. Octandria Monogynia.
1 spicatum. G.B. (re.) spiked.	England. 7. 8. H. 4. Eng. bot. v. 28. t. 1947.
Epilobium angustifolium. E.B.	
β albiflorum. (wh.) white-flowered.	— — H. 4.

2	speciosum. L.C.	(re.) handsome.	Europe.	7. 8. H. 4.	
		<i>Epilobium angustifolium</i> β latum. DC.	<i>latifolium</i> . Schmidt.			
3	americanum.	(re.) American.	N. America.	1825.	— H. 4.	
4	rosmarinifolium. DC.	(re.) Rosemary-l'd.	Europe.	1775.	6. 8. H. 4.	Reich. ic. 4. t. 341.
		<i>Epilobium angustissimum</i> . Bot. mag. t. 76.				
5	Halleri. Retz.	(re.) Haller's.	Switzerland.	1798.	— H. 4.	— 4. t. 342.
		<i>Epilobium Dodon'æi</i> . Allioni, <i>rosmarinifolium</i> β <i>alpinum</i> . DC.				
6	latifolium.	(li.) broad-leaved.	Europe.	1779.	7. 8. H. 4.	Sal. par. lond. t. 58.
		<i>halimifolium</i> . P.L.				
EPIL'OBÍUM. DC. WILLOW-HERB. Octandria Monogynia. L.						
1	alpinum. DC.	(ro.) alpine.	Britain.	6. 7. H. 4.	Eng. bot. v. 28. t. 2001.
2	origanifolium. DC.	(ro.) Marjoram-leav'd.	—	7. 8. H. 4.	— v. 28. t. 2000.
		<i>alsinifolium</i> . E.B.				
3	róseum. E.B.	(ro.) rose-coloured.	—	— H. 4.	— v. 10. t. 693.
4	alpéstre. s.s.	(ro.) simple-stalked.	Switzerland.	— H. 4.	
5	montánum. DC.	(li.) mountain.	Britain.		— H. 4.	Eng. bot. v. 17. t. 1177.
6	nútans. DC.	(bh.) nodding.	Bohemia.	1827.	— H. 4.	Reich. ic. 2. t. 197.
7	Hornemánni. DC.	(li.) Hornemann's.	Finland.		— H. 4.	— p. 73. t. 180.
		<i>nútans</i> . Horn. nec <i>aliorum</i> .				
8	colorátum. DC.	(ro.) pink-flowered.	N. America.	1805.	— H. 4.	
9	dahúricum. DC.	(wh.) Dahurian.	Dahuria.	1816.	— H. ⊖.	
10	hirsútum. DC.	(pu.) Codlins & cream.	Britain.	— H. 4.	Eng. bot. v. 12. t. 838.
11	tomentósüm. DC.	(pu.) woolly.	Levant.	1821.	— H. 4.	Vent. hort. cels. t. 90.
12	villósüm. DC.	(pu.) villous.	C. B. S.	1799.	— G. 4.	
13	cylindricum. DC.	(ro.) round-stemmed.	Nepaul.	1827.	— H. 4.	
14	palústre. DC.	(ro.) round-stalked.	Britain.	— H. 4.	
15	rivuláre. DC.	(ro.) rivulet.	N. Europe.	1828.	— H. 4.	
16	parviflórum. DC.	(pu.) small-flowered.	Britain.	— H. 4.	
17	tetragónum. DC.	(pu.) square-stemmed.	—	— H. 4.	Eng. bot. v. 5. t. 346.
18	obscúrum. P.S.	(pu.) obscure.	Europe.	1820.	— H. 4.	Reich. ic. 2. t. 170.
19	canéscens. H.H.	(ro.) canescent.	—	1826.	— H. 4.	Eng. bot. v. 12. t. 795.
		<i>fruticosa</i> . Jacq. ex DC. prod'r.				— v. 28. t. 1948.
GA'URA. DC.	GA'URA. Octandria Monogynia. L.					Reich. ic. 2. t. 199.
1	bíennis. DC.	(bh.) biennial.	N. America.	1762.	8. 10. H. ♂.	Bot. mag. t. 389.
2	angustifólia. DC.	(fl.) narrow-leaved.	Carolina.	1816.	7. 9. F. ♂.	Jacq. ic. 3. t. 457.
		<i>fruticosa</i> . Jacq. ex DC. prod'r.				
3	sinuáta. DC.	(fl.) sinuate-leaved.	Arkansa.	1826.	— H. ♂.	Cavan. ic. 4. t. 396. f. 1.
4	tripétala. DC.	(bh.) three-petaled.	Mexico.	1804.	8. 10. H. ⊖.	
5	linisólia. DC.	(wh.) Flax-leaved.	Arkansa.	1826.	7. 9. H. ♂.	
6	coccínea. DC.	(ro.) pale-scarlet.	Louisiana.	1811.	8. 10. H. 4.	
7	mutábilis. DC.	(ye.) changeable-col'd.	N. Spain.	1795.	7. 9. F. ♂.	
8	enotheriflóra. DC.	(ye.) canescent.	—	1816.	— F. ♂.	Bot. mag. t. 388.
CENOTH'ERA. DC. EVENING PRIMROSE. Octandria Monogynia. L.						
1	dentata. DC.	(ye.) toothed.	Chile.	1818.	6. 9. H. ⊖.	Lindl. coll. t. 10.
2	micránta. s.s.	(ye.) small-blossom'd.	California.	1820.	— H. ⊖.	
		<i>hirta</i> . L. enum.				
3	cheiranthifólia. DC.	(ye.) Wallflower-l'd.	Chile.	—	— F. ♂.	
4	bíennis. DC.	(ye.) common.	N. America.	1629.	— H. ♂.	Bot. reg. t. 1040.
5	élata. DC.	(ye.) tall upright.	Mexico.	1827.	7. 10. F. ♂.	Eng. bot. v. 22. t. 1534.
6	grandiflóra. DC.	(ye.) large-flowered.	N. America.	1778.	6. 8. H. ⊖.	
7	suavéolens. DC.	(ye.) fragrant.	—	— H. ♂.	Bot. mag. t. 2068.
8	Simsiana. DC.	(ye.) Sims's.	Mexico.	1816.	7. 10. F. ♂.	
		<i>corymbosa</i> . B.M. non Lamarck.				Bot. mag. t. 1974.
9	macrocéarpa. DC.	(ye.) broad-leaved.	N. America.	1811.	6. 8. H. 4.	Swt. br. fl. gar. t. 5.
10	missouriensis. B.M.	(ye.) narrow-leaved.	—	—	— H. 4.	Bot. mag. t. 1592.
		<i>alata</i> . N. non Rafin.				
11	salicifolia. DC.	(ye.) Willow-leaved.	—	1818.	6. 9. H. ⊖.	Meerb. ic. 1. t. 34.
12	parviflóra. DC.	(ye.) small-flowered.	N. America.	1757.	6. 8. H. ♂.	
13	cruciáta. N.	(st.) cross-petaled.	—	1821.	6. 10. H. ♂.	Murr. com. goet. 6. t. 1.
14	muricáta. DC.	(ye.) prickly-stalked.	N. America.	1789.	7. 9. H. ♂.	Jacq. ic. 3. t. 455.
15	noctúrnsa. DC.	(ye.) night-smelling.	C. B. S.	1790.	4. 8. G. ♂.	Bot. mag. t. 365.
16	longiflóra. DC.	(ye.) long-flowered.	BuenosAyres.	1776.	7. 9. H. ♂.	Schkuhr hand. 1. t. 105.
17	villósä. DC.	(ye.) villous.	C. B. S.	1791.	— G. ♂.	Bot. mag. t. 2403.
18	mollíssima. DC.	(ye.) soft wave-l'd.	BuenosAyres.	1732.	6. 10. H. ♂.	Jacq. ic. 3. t. 456.
19	odoráta. DC.	(ye.) sweet-scented.	Patagonia.	1790.	4. 8. H. 4.	Bot. reg. t. 147.
		<i>a glaucescens</i> . DC. (ye.) glaucescent.		—	— H. 4.	
		<i>β viréscens</i> . DC. (ye.) purple-stalked.		—	— H. 4.	
20	striáta. DC.	(ye.) striped-seeded.	—	1816.	6. 9. H. ⊖.	Murr. com. get. 5. t. 9.
21	sinuáta. DC.	(ye.) scollop-leaved.	N. America.	1770.	7. 8. H. ⊖.	Swt. br. fl. gar. t. 167.
22	tenélla. DC.	(pu.) slender.	Chile.	1822.	6. 9. H. ⊖.	

23 <i>tenuifolia</i> . DC.	(<i>pu.</i>) slender-leaved. Chile.	1829.	6. 9. H. ⊙.	Cavan. ic. 4. t. 397.
24 <i>viminea</i> . B.M.	(<i>pu.</i>) twiggy. N. America.	1827.	— H. ⊙.	Bot. reg. t. 1220.
25 <i>decumbens</i> . D.	(<i>pu.</i>) decumbent. —	—	— H. ⊙.	— t. 1221.
26 <i>quadrivulnera</i> . D.	(<i>pu.</i>) four-spotted. —	—	— H. ⊙.	— t. 1119.
27 <i>Lindleyana</i> . D.	(<i>pu.</i>) Lindley's. —	—	— H. ⊙.	Swt. br. fl. gar. s. 2. t. 19.
28 <i>rōseo-álba</i> . DC. (<i>wh. ro.</i>)	rosy and white. America.	1826.	— H. ⊙.	— t. 268.
29 <i>purpurea</i> . DC.	(<i>pu.</i>) purple-flower'd. N. America.	1794.	5. 9. H. ⊙.	Bot. mag. t. 352.
30 <i>Romanzovii</i> . DC.	(<i>pu.</i>) Romanzoff's. —	1819.	— H. ⊙.	Bot. reg. t. 662. 562
31 <i>corymbosa</i> . DC.	(<i>ye.</i>) corymbose. —	1826.	6. 9. H. ♂.	
32 <i>spectabilis</i> . H.H.	(<i>ye.</i>) showy. Mexico.	1822.	— F. ♂.	
33 <i>gauroídes</i> . DC.	(<i>ye.</i>) Gaura-like. Baltimore.	1816.	7. 9. H. ♂.	
34 <i>média</i> . DC.	(<i>ye.</i>) intermediate. N. America.	1821.	6. 9. H. ♀.	
35 <i>serrulata</i> . DC.	(<i>ye.</i>) saw-leaved. —	—	6. 10. H. ♀.	Swt. br. fl. gar. t. 133.
36 <i>strícta</i> . Led.	(<i>ye.</i>) straight. —	1822.	— H. ⊙.	
37 <i>amœna</i> . Leh.	(<i>ye.</i>) delicate. —	1826.	— H. ⊙.	
38 <i>nervosa</i> . H.H.	(<i>ye.</i>) nerved-leaved. —	—	— H. ♂.	
39 <i>cæspitosa</i> . DC.	(<i>wh.</i>) tufted. N. America.	1811.	6. 9. H. ♀.	Bot. mag. t. 1593.
40 <i>acaúlis</i> . DC.	(<i>wh.</i>) short-stalked. Chile.	1822.	— H. ♀.	Bot. reg. t. 763.
41 <i>taraxacifolia</i> . B.F.G. (<i>w.</i>)	Dandelion-ld.	1823.	— H. ♀.	Swt. br. fl. gar. t. 294.
<i>grandiflora</i> . Flor. peruv. 3. t. 318. f. 6. <i>nec aliorum</i> .				
42 <i>triloba</i> . DC.	(<i>st.</i>) root-flowering. N. America.	1821.	6. 10. H. ⊙.	Bot. mag. t. 2566.
<i>rhizocarpa</i> . S.S.				
43 <i>virgata</i> . DC.	(<i>pu.</i>) slender-twigg'd. Peru.	1823.	6. 9. H. ⊙.	Flor. per. 29. t. 315.
44 <i>pinnatifida</i> . DC.	(<i>ye.</i>) pinnatifid. Mexico.	1825.	— F. ♀.	
45 <i>tetraptera</i> . DC.	(<i>wh.</i>) four-winged. —	1796.	— H. ⊙.	Bot. mag. t. 468.
46 <i>latiflora</i> . DC.	(<i>wh.</i>) broad-flowered. —	1826.	— H. ♀.	Flor. mex. ined. t. 376.
47 <i>glaúca</i> . DC.	(<i>ye.</i>) glaucous-leav'd. N. America.	1812.	5. 10. H. ♀.	Bot. mag. t. 1606.
48 <i>fruticosa</i> . DC.	(<i>ye.</i>) frutescent. —	1737.	6. 8. H. ♀.	— t. 332.
49 <i>ambigua</i> . DC.	(<i>ye.</i>) ambiguous. —	—	... H. ♀.	Swt. br. fl. gar. t. 184.
50 <i>serotina</i> . B.F.G.	(<i>ye.</i>) late-flowered. —	1820.	7. 10. H. ♂.	
51 <i>tetragona</i> . DC.	(<i>ye.</i>) blunt-leaved. —	—	6. 8. H. ♀.	
52 <i>Fraseri</i> . DC.	(<i>ye.</i>) Fraser's. —	1811.	5. 10. H. ♀.	Bot. mag. t. 1674.
53 <i>pūmila</i> . DC.	(<i>ye.</i>) dwarf. —	1757.	5. 9. H. ♀.	— t. 355.
54 <i>rōsea</i> . DC.	(<i>ro.</i>) rose-coloured. Peru.	1783.	— F. ♀.	— t. 347.
55 <i>linéaris</i> . DC.	(<i>ye.</i>) linear-leaved. N. America.	6. 8. H. ♀.	Swt. br. fl. gar. t. 253.
56 <i>speciosa</i> . DC.	(<i>wh.</i>) handsome. Louisiana.	1821.	— H. ♀.	Bot. reg. t. 1142.
57 <i>pallida</i> . B.R.	(<i>bh.</i>) pale. N. America.	1827.	— H. ♀.	
58 <i>albicalvis</i> . DC.	(<i>wh.</i>) white-stalked. —	1811.	5. 9. H. ⊙.	
<i>pinnatifida</i> . N. non Kth.				
59 <i>Nuttallii</i> .	(<i>wh.</i>) Nuttall's. N. America.	—	5. 8. H. ♀.	
<i>albicaulis</i> . N. non Pursh.				
60 <i>canadensis</i> . DC.	(<i>ye.</i>) Canadian. Canada.	1825.	6. 9. H. ⊙.	
CLA'RKIA. Ph.	CLA'RKIA. Octandria Monogynia.			
pulchella. Ph.	(<i>ro.</i>) beautiful. N. America.	1827.	5. 10. H. ⊙.	Bot. reg. t. 1100.
β <i>albiflora</i> .	(<i>wh.</i>) white-flowered. —	—	— H. ⊙.	
γ <i>integriloba</i> .	(<i>ro.</i>) entire-lobed. —	—	— H. ⊙.	

Tribus IV. JUSSIEÆ. DC. prodr. v. 3. p. 52.

JUSSI'ÆA. DC.

1 <i>variabilis</i> . DC.	(<i>ye.</i>) variable. Guiana.	1820.	7. 9. S.w. ♂.	
2 <i>leptocarpa</i> . DC.	(<i>ye.</i>) narrow-podded. Mississippi.	1824.	— H. ⊙.	H. et B.n. gen. 6. t. 532.
3 <i>pilosa</i> . DC.	(<i>ye.</i>) hairy. Caracas.	1829.	— S.w. ♀.	
4 <i>grandiflora</i> . DC.	(<i>ye.</i>) great-flowered. Carolina.	1812.	7. 10. G.w. ♀.	Bot. mag. t. 2122.
5 <i>repens</i> . DC.	(<i>ye.</i>) creeping. E. Indies.	1816.	— S.w. ♀.	Rheed. mal. 2. t. 51.
6 <i>Swartziana</i> . DC.	(<i>ye.</i>) Swartz's. W. Indies.	1826.	— S.w. ♀.	
7 <i>inclinata</i> . DC.	(<i>ye.</i>) inclined. Surinam.	1820.	— S.w. ⊙.	
8 <i>acuminata</i> . DC.	(<i>ye.</i>) acuminate. Jamaica.	—	— S. ⊙.	
9 <i>linifolia</i> . DC.	(<i>ye.</i>) Flax-leaved. W. Indies.	—	— S. ⊙.	
10 <i>erecta</i> . DC.	(<i>ye.</i>) upright. S. America.	1739.	— S.w. ♂.	Gärt. fruct. t. 31.
11 <i>frutescens</i> . DC.	(<i>ye.</i>) frutescent.	1829.	4. 6. S.w. ♀.	
12 <i>longifolia</i> . DC.	(<i>ye.</i>) long-leaved. Brazil.	1828.	6. 9. S.w. ♀.	DC. pl. rar. gen. 1. t. 4.
13 <i>decurrens</i> . DC.	(<i>ye.</i>) decurrent. N. America.	—	— H.w. ♀.	Abb. ins. t. 40.
14 <i>ovalifolia</i> . DC.	(<i>ye.</i>) oval-leaved. Madagascar.	1823.	7. 9. S.w. ♀.	Bot. mag. t. 2530.
15 <i>Burmanni</i> . DC.	(<i>ye.</i>) Burmann's. E. Indies.	1808.	8. 9. S.w. ♂.	Andr. deposit. 621.
<i>suffruticosa</i> , A.R.				

ORNAGRARIÆ.

16 villòsa. DC.	(ye.) villous.	E. Indies.	1826.	8. 9.	S.w. $\frac{1}{2}$.	Rheed. mal. 2. t. 50.
17 octonérvia. DC.	(ye.) eight-nerved.	S. America.	7. 9.	S.w. $\frac{1}{2}$.	Lam. ill. t. 280. f. 1.
<i>Oenothera octovalvis.</i> Jacq. amer. t. 70.						
18 octofíla. DC.	(ye.) eight-veined.	Caribees.	—	S.w. $\frac{1}{2}$.	Plum.edBur.t.175.f.1.
19 scábra. DC.	(ye.) rough.	Brazil.	1816.	—	S.w. $\frac{1}{2}$.	
20 hírta. DC.	(ye.) hairy.	S. America.	1823.	—	S.w. $\frac{1}{2}$.	Plum.edBur.t.174.f.2.
LUDWI'GIA. DC.	LUDWI'GIA.	Tetrandria Monogynia.	L.			
1 jussiaeóides. DC.	(ye.) Jussiaea-like.	Mauritius.	1826.	7. 9.	S.w. $\frac{1}{2}$.	
2 parviflòra. DC.	(ye.) small-flowered.	E. Indies.	1812.	—	S. \odot .	
ISNA'RDIÀ. DC.	ISNA'RDIÀ.	Tetrandria Monogynia.	L.			
1 alternifòlia. DC.	(ye.) alternate-leav'd.	N. America.	1752.	6. 7.	H.w. $\frac{1}{2}$.	Lam. ill. t. 77.
<i>Ludwigia alternifolia.</i> L. <i>macrocarpa.</i> M.						
2 hirsúta. DC.	(ye.) hairy.	—	1812.	6. 8.	H.w. $\frac{1}{2}$.	
3 palústris. DC.	(gr.) marsh.	Europe.	1776.	7. 8.	H.w. $\frac{1}{2}$.	Schk. handb. 1. t. 25.
4 nítida. (gr.) American.	America.	—	—	—	H.w. $\frac{1}{2}$.	
<i>Ludwigia nítida.</i> M. <i>apétala.</i> Walter. <i>palústris.</i> Torrey.						

Tribus V. CIRCAEÆ. DC. prodr. v. 3. p. 61.

LOP'EZIA. DC.	LOP'EZIA.	Monandria Monogynia.				
1 racemòsa. DC.	(pu.) smooth.	Mexico.	1792.	8. 10.	G. δ .	Bot. mag. t. 254.
2 cordàta. DC.	(pu.) heart-leaved.	—	1826.	7. 10.	H. \odot .	
3 coronàta. DC.	(pu.) coronet-flower'd.	—	1805.	—	H. \odot .	Swt. br. fl. gar. t. 108.
4 oppositifòlia. DC.	(pu.) opposite-leaved.	—	1825.	—	H. \odot .	
5 integrifòlia. DC.	(pu.) entire-leaved.	—	—	—	H. \odot .	
6 miniàta. DC.	(pu.) frutescent.	—	1824.	9. 11.	G. $\frac{1}{2}$.	Jacq. f. ecl. t. 109.
7 hirsúta. DC.	(pu.) hairy.	—	1796.	—	G. δ .	Jacq. coll. 5. t. 15. f. 4.
8 pùmila. DC.	(pu.) dwarf.	—	1825.	7. 9.	H. \odot .	
CIR'C'EA. DC.	ENCHANTER'S NIGHTSHADE.	Diandria Monogynia.				
1 lutetiána. DC.	(wh.) common.	Britain.	6. 8.	H. $\frac{1}{2}$.	Eng. bot. v. 15. t. 1056.
2 canadénsis. L.	(wh.) smooth-stalked.	N. America.	—	H. $\frac{1}{2}$.	
3 alpína. DC.	(wh.) mountain.	Britain.	6. 9.	H. $\frac{1}{2}$.	Eng. bot. v. 15. t. 1057.

Tribus VI. HYDROCARYEÆ. DC. prodr. v. 3. p. 63.

TR'APA. DC.	WATER-CALTROPS.	Tetrandria Monogynia.	L.			
1 nàtans. DC.	(wh.) floating.	S. Europe.	1781.	6. 8.	H.w. \odot .	Bot. reg. t. 259.
2 quadrispinòsa. DC.	(wh.) four-spined.	E. Indies.	1823.	—	S.w. \odot .	
3 bispinòsa. DC.	(wh.) two-spined.	—	1822.	6. 9.	S.w. \odot .	Roxb. corom. t. 234.
4 bicòrnis. DC.	(wh.) two-horned.	China.	1790.	—	G.w. $\frac{1}{2}$.	Braam ic. chin. t. 22.

ORDO LXXXVII.

HALORAGEÆ. DC. prodr. v. 3. p. 65.

Tribus I. CERCODIANÆ. DC. prodr. v. 3. p. 65.

SERPI'CULA. DC.	SERPI'CULA.	Monœcia Tetrandria.	L.			
rèpens. DC.	(gr.) creeping.	C. B. S.	1789.	7. 8.	G. $\frac{1}{2}$.	Berg. cap. t. 5. f. 10.
GONIOCA'RPUS. DC.	GONIOCA'RPUS.	Octandria Tetragynia.				
1 micránthus. DC.	(wh.) small-flowered.	Japan.	1805.	6. 9.	H. \odot .	Thunb. fl. jap. t. 15.
2 scáber. DC.	(wh.) rough-leaved.	China.	1820.	—	H. \odot .	Ann. bot. 1. t. 12. f. 6.
3 tecniórides. DC.	(wh.) Teucrium-like.	N. S. W.	1826.	—	H. \odot .	
HALOR'AGIS. DC.	HALOR'AGIS.	Octandria Di-Tetragynia.				
1 prostrata. DC.	(gr.) prostrate.	N. S. W.	1818.	4. 9.	G. $\frac{1}{2}$.	
2 digyna. DC.	(gr.) two-styled.	N. Holland.	1826.	—	G. $\frac{1}{2}$.	Labil. n. holl. 1. t. 129.

CERC'ODIA. DC.	CERC'ODIA. Octandria Tetragynia.					
1 erécta. DC. (gr.) upright.	N. Zealand. 1772.	4. 9. G. ½.	Jacq. ic. 1. t. 69.			
<i>Halorágis alata.</i> Jacq. <i>Cercòdia.</i> H.K. <i>Tetragónia ivafolia.</i> L.						
PROSERPIN'ACA. DC. PROSERPIN'ACA. Triandria Trigynia.						
1 palústris. DC. (wh.) marsh.	N. America. 1822.	7. 8. H.w.○.	Jus. an. mus. 3. t. 30. f. 1.			
2 pectinata. DC. (wh.) pectinated.	—	—	H.w.○.	Lam. ill. 1. t. 50. f. 1.		
MYRIOPHY'LLUM. WATER MILFOIL. Monoezia Tetr-Octandria.						
1 spicatum. DC. (ro.) spiked.	Britain.	7. 8. H.w. ¼.	Eng. bot. v. 2. t. 83.			
2 pectinatum. DC. (ro.) pectinated.	Europe. 1828.	— H.w. ¼.	Flor. dan. t. 1046.			
<i>verticillatum.</i> F.D.						
3 verticillatum. DC. (ro.) verticillate.	England.	— H.w. ¼.	Eng. bot. v. 4. t. 218.			

Tribus II. **CALLITRICHINEÆ.** DC. prodr. v. 3. p. 70.

CALLI'TRICE. DC.	WATER STARWORT. Monandria Digynia. L.					
1 vérrna. DC. (wh.) vernal.	Britain.	4. 10. H.w.○.	Eng. bot. v. 11. t. 722.			
2 autumnalis. DC. (wh.) autumnal.	— 6. 10. H.w.○.	Gmel. sib. 3. t. 1.			

Tribus III. **HIPPURIDEÆ.** DC. prodr. v. 3. p. 71.

HIPP'URIS. DC.	MARE'S-TAIL. Monandria Monogynia. L.					
vulgáris. DC. (gr.) common.	Britain.	5. H.w. ¼.	Eng. bot. v. 11. t. 763.			

ORDO LXXXVIII.**CERATOPHYLLEÆ.** Gray brit. pl. arr. 2. p. 554.

CERATOPHY'LLUM. DC.	HORNWORT. Monoezia Polyandria. L.					
1 demérsum. DC. (gr.) spiny-fruited.	Britain.	7. 9. H.w. ¼.	Eng. bot. v. 14. t. 947.			
2 submérsum. DC. (gr.) naked-fruited.	— 9. H.w. ¼.	— v. 10. t. 679.			

ORDO LXXXIX.**LYTHRARIEÆ.** DC. prodr. v. 3. p. 75.Tribus I. **SALICARIEÆ.** DC. prodr. v. 3. p. 75.

PE'PLIS. DC.	WATER PURSLANE. Hexandria Monogynia. L.					
Pórtula. DC. (re.) common.	Britain.	7. 9. H.○.	Eng. bot. v. 17. t. 1211.			
AMMA'NNIA. DC.	AMMA'NNIA. Tetr-Heptandria Monogynia.					
1 ægyptiaca. DC. (gr.) Egyptian.	Egypt. 1818.	7. 8. H.w.○.	Willd. hort. ber. t. 6.			
2 vesicatoria. DC. (ye.) bladdered.	E. Indies. 1825.	— S.w.○.	Pluk. alm. t. 136. f. 22.			
3 occidentális. DC. (gr.) occidental.	Portorico. 1829.	— S.w.○.				
4 cáspica. DC. (gr.) Caspian.	Astracan. 1818.	— H.w.○.				
5 ramosior. DC. (pu.) branching.	Virginia. 1759.	— H.w.○.	Bocc. mus. t. 104.			
6 latifolia. DC. (wh.) broad-leaved.	W. Indies. 1733.	— S.w.○.	Sloan. his. 1. t. 7. f. 4.			
7 hùmílis. DC. (ro.) small.	N. America. 1823.	— H.w.○.				
8 baccifera. R.S. (ye.) berry-bearing.	China. 1820.	— H.w.○.	Burm. ind. t. 15. f. 3.			
9 verticillata. DC. (pu.) whorled.	—	— H.w.○.	Lam. ill. t. 77. f. 3.			
10 multiflora. DC. (re.) many-flowered.	E. Indies. 1818.	— S.w.○.				
11 débilis. DC. (li.) cluster-flower'd.	—	— S.w.○.				
12 diffusa. DC. (wh.) spreading.	—	— S.w.○.				
13 rotundifolia. DC. (fl.) round-leaved.	Nepaul. 1818.	— H.w.○.				
14 pentandra. DC. (wh.) pentandrous.	E. Indies. 1823.	— S.w.○.				

15 sanguinolenta. DC. (cr.)	bloody.	Jamaica.	1803.	7. 8. S.w.○.	
16 octandra. DC.	(cr.) octandrous.	E. Indies.	1818.	— S.w.○.	Roxb. corom. 2. t. 133.
17 auriculata. DC.	(pu.) eared.	Egypt.	1820.	— H.w.○.	Willd. h. ber. 1. t. 7.
18 rübra. DC.	(re.) red-flowered.	Nepaul.	1822.	— H.w.○.	
19 pallida. DC.	(pa.) pale-flowered.	1828.	— H.w.○.	
20 viridis. DC.	(gr.) green.	1818.	— H.w.○.	
21 Wormskiöldii. F. Wormskiold's.	Siberia.	1828.	— H.w.○.	
L'YTHRUM. DC.	L'YTHRUM. Di-Dodecandria Monogynia.				
1 thymifolium. DC. (li.)	Thyme-leaved.	S. Europe.	1820.	7. 9. H.○.	
2 thesioides DC. (li.)	Thesum-like.	Caucasus.	1828.	— H.○.	
3 lineare. DC. (wh.)	linear-leaved.	N. America.	1812.	7. 8. H. ¼.	
4 alatum. DC. (pu.)	winged-stalked.	1812.	5. 11. F. ¼.	
5 hyssopifolium. DC. (li.)	Hyssop-leaved.	England.	8. 9. H.○.	
6 lanceolatum. DC. (pu.)	spear-leaved.	Carolina.	1822.	7. 9. H. ¼.	
7 álbum. DC. (wh.)	white-flowered.	Mexico.	1827.	— H. ½.	
8 Graefferi. DC. (pu.)	Græffær's.	S. Europe.	1825.	— H.○.	
9 Salicaria. DC. (pu.)	common.	Britain.	7. 9. H.w. ¼.	Eng. bot. v. 15. t. 1061.
10 pubescens. (pu.)	pubescent.	N. America.	7. 8. H. ¼.	
<i>Salicaria</i> β pubescens. Ph.					
11 diffusum. B.F.G. (pu.)	spreading.	1822.	7. 9. H. ¼.	Swt. br. fl. gar. t. 149.
12 virgatum. DC. (ro.)	slender-branch'd.	Europe.	1776.	6. 9. H. ¼.	Bot. mag. t. 1003.
C'UPHEA. DC.	C'UPHEA. Dodecandria Monogynia.				
1 ciræoïdes. DC. (vi.)	Ciræa-like.	S. America.	1819.	8. 10. H.○.	
2 spicata. DC. (ro.)	spiked.	Peru.	—	— H.○.	
3 Melvilla. DC. (sc.gr.)	scarlet & green.	Guiana.	1822.	6. 10. S. ½.	
4 viscosissima. DC. (li.)	clammy.	America.	1776.	7. 10. H.○.	
5 procumbens. DC. (li.)	procumbent.	Mexico.	1816.	6. 9. H.○.	
6 lanceolata. DC. (li.)	spear-leaved.	1796.	5. 8. G. ½.	
7 æquipétala. DC. (vi.)	equal-flowered.	1828.	6. 9. H.○.	Cavan. ic. 4. t. 382. f. 2.
8 virgata. DC. (vi.)	twiggy.	—	— H.○.	— t. 382. f. 1.
9 decandra. DC. (pu.)	ten-stamen'd.	Jamaica.	1789.	6. 10. S. ½.	
10 parviflora. DC. (ro.)	small-flowered.	Demerara.	1824.	10. 12. S. ½.	
11 Simsii. ed. 1. (li.)	Sims's.	Trinidad.	1823.	8. 10. S. ½.	
<i>serpyllifolia</i> . B.M. non Kth. Trinitatis. DC.					
12 multiflora. DC. (vi.)	many-flowered.	1822.	— S. ½.	Lodd. bot. cab. t. 808.
ACISANTH'ERA. DC.	ACISANTH'ERA. Decandria Monogynia.				
quadrata. DC. (pi.)	four-sided.	Jamaica.	1804. S. ¼.	Brown. jam. t. 22. f. 1.
HEIMIA. DC.	HEIMIA. Dodecandria Monogynia.				
1 salicifolia. DC. (ye.)	Willow-leaved.	Mexico.	1821.	6. 9. F. ½.	
2 myrtifolia. Ot. (ye.)	Myrtle-leaved.	1827.	— F. ½.	Swt. br. fl. gar. t. 281.
DEC'DON. DC.	DEC'DON. Decandria Monogynia.				
verticillatum. DC. (pu.)	whorl-flowered.	N. America.	1759.	7. 9. H.w. ¼.	
<i>Lythrum verticillatum</i> . L. <i>Nesaea verticillata</i> . Kth.					
NES'EA. DC.	NES'EA. Oct-Dodecandria Monogynia.				
triflora. DC. (bl.)	three-flowered.	Mauritius.	1802.	8. 9. S.w. ¼.	
<i>Lythrum triflorum</i> . L.					
LAWS'ONIA. DC.	LAWS'ONIA. Octandria Monogynia. L.				
1 spinosa. w. (wh.)	prickly.	E. Indies.	1759.	6. 9. S. ½.	Rheed. mal. 1. t. 40.
2 inermis. w. (wh.)	smooth.	Egypt.	—	— G. ½.	Rauw. ic. 60. t. 7.
GIN'ORIA. DC.	GIN'ORIA. Dodecandria Monogynia.				
americana. DC. (bl.)	American.	Cuba.	1827. S. ½.	Jacq. amer. t. 91.
GRISLEA. DC.	GRI'SLEA. Oct-Dodecandria Monogynia.				
1 secunda. DC. (re.)	secund.	Caracas.	1824.	6. 8. S. ½.	
2 tomentosa. DC. (sc.)	downy.	E. Indies.	1804.	5. 12. S. ½.	Bot. reg. t. 30.
<i>Woodfordia floribunda</i> . Sal. par. lond. t. 42. <i>Lythrum fruticosum</i> . L.					

Tribus II. LAGERSTRÆMIEÆ. DC. prodr. v. 3. p. 92.

LAGERSTRÆMIA. DC. LAGERSTRÆMIA. Icosandria Monogynia.

1 indica. DC.	(re.) Indian.	China.	1759.	8. 10. F. ½.	Bot. mag. t. 405.
β carnea.	(fl.) flesh-coloured.	1816.	— F. ½.	
γ álba.	(wh.) white-flowered.	—	— F. ½.	
2 parviflora. DC.	(wh.) small-flowered.	E. Indies.	—	— S. ½.	Roxb. corom. 1. t. 66.
3 speciosa. DC.	(ro.) beautiful.	China.	1826. S. ½.	Munch. haus. 5. t. 356.
4 grandiflora. DC.	(ro.) large-flowered.	E. Indies.	1809. S. ½.	
5 reginae. DC.	(ro.) oblong-leaved.	1792. S. ½.	Roxb. corom. 1. t. 651.

ORDO XC.

TAMARISCINEÆ. DC. prodr. v. 3. p. 95.

TA'MARIX. DC.	TAMARISK.	Tetr-Pentandria Trigynia.					
1 gállica. DC.	(fl.) red-wooded.	England.	7. 8.	H. ½.	Eng. bot. v. 19. t. 1318.	
2 Pallásii. DC.	(fl.) Pallas's.	Caucasus.	1827.	—	H. ½.	Pall. fl. ross. t. 77.	
pentándra. Pall.	<i>gállica</i> . M.B. nec aliorum.						
3 índica. DC.	(bh.) Indian.	E. Indies.	6. 8.	S. ½.		
4 dioíca. DC.	(bh.) diœcious.	—	1812.	—	S. ½.		
5 artículata. DC.	(pi.) jointed.	—	1816.	S. ½.	Vahl. symb. 2. t. 32.	
6 tetrándra. DC.	(wh.) tetrandrous.	Tauria.	1821.	7. 8.	H. ½.		
MYRIC'ARIA. DC.	MYRIC'ARIA.	Monadelphia Decandria.					
1 germánica. DC.	(pi.) green-wooded.	Germany.	1582.	6. 9.	H. ½.	Flor. dan. t. 234.	
Támarix germánica. L.							
2 dahúrica. DC.	(pi.) Dahirian.	Dahuria.	1827.	H. ½.		

ORDO XCI.

MELASTOMACEÆ. DC. prodr. v. 3. p. 99.

Subordo I. MELASTOMEÆ.—Tribus I. LAVOISIEREÆ. DC. loc. cit. 100.

MERI'ANIA. DC.	MERI'ANIA.	Decandria Monogynia.					
1 leucántha. DC.	(wh.) white-flowered.	Jamaica.	1825.	6. 8.	S. ½.		
2 rósea. DC.	(ro.) mountain rose.	—	—	—	S. ½.	Tussac fl. ant. t. 6.	
3 purpúrea. DC.	(pu.) purple.	—	—	—	S. ½.	Swart. fl. ind. oc. 2. t. 15.	

Tribus II. RHEXIEÆ. DC. prodr. 3. p. 114.

SPENN'ERA. DC.	SPENN'ERA.	Octo-Decandria Monogynia.					
1 paludosa. DC.	(ro.) marsh.	Brazil.	1825.	6. 8.	S. ¼.		
2 frágilis. DC.	(wh.) brittle.	Trinidad.	1822.	—	S. ½.		
3 aquática. DC.	(ro.) marsh.	S. America.	1793.	3. 9.	S. ½.	Aubl. guian. 1. t. 169.	
Rhéxia aquática. Bonpl. rhex. t. 40.	<i>Melastoma aquática</i> . Aubl.						
4 pendulifolia. DC.	(ro.) hanging-leaved.	Guiana.	1826.	—	S. ½.	Bonpl. nav. t. 26.	
5 glandulosá. DC.	(wh.) glandulous.	—	1824.	—	S. ½.	—	70. t. 27.
MICROLIC'CIA. DC.	MICROLIC'CIA.	Decandria Monogynia.					
1 brevífolia. DC.	(pu.) short-leaved.	Guiana.	1823.	7. 10.	S. ⊖.	Aubl. guian. 1.t.155.f.b.	
<i>Melastoma trivalvis</i> . Aubl.	<i>Rhéxia trivalvis</i> . Vahl.						
2 bivalvis. DC.	(pu.) two-valved.	Trinidad.	1822.	6. 8.	S. ⊖.	Aubl. guian. 1.t.155.f.a.	
3 recúrva. DC.	(pu.) recurved-leav'd.	—	1820.	—	S. ¼.		
RHE'XIA. DC.	RHE'XIA.	Octandria Monogynia. L.					
1 mariána. DC.	(li.) Maryland.	N. America.	1759.	7. 9.	H. ¼.	Pluk. mant. t. 428. f. 1.	
<i>a purpúrea</i> . DC.	(li.) purplish.	—	—	—	H. ¼.		
β rubélla. DC.	(re.) reddish.	—	—	—	H. ¼.	Swt. br. fl. gar. t. 41.	
2 virginica. DC.	(pu.) Virginian.	—	—	6. 8.	H. ¼.	Bot. mag. t. 968.	
3 ciliosa. DC.	(pu.) ciliated.	—	1812.	8. 10.	H. ¼.	Swt. br. fl. gar. t. 298.	
4 angustifolia. DC.	(wh.) narrow-leaved.	—	—	7. 9.	H. ¼.		
OXYSP'ORA. DC.	OXYSP'ORA.	Octandria Monogynia.					
paniculata. DC.	(wh.) panicled.	Nepaul.	1826.	6. 8.	S. ½.		
<i>Arthrostémma paniculatum</i> . D.P.							

Tribus III. OSBECKIÆ. DC. prodr. 3. p. 127.

LASIA'NDRA. DC. LASIA'NDRA. Decandria Monogynia.

- 1 Fontanesiâna. DC. (pu.) rough-leaved. Brazil. 1818. 4. 8. S. h. Bonpl. rhe. t. 36.
Melástoma granulosa. B. reg. 671. *Rhéxia Fontanésii*. Bonpl.
 2 Langsdorfiâna. DC. (pu.) Langsdorf's. Brazil. 1821. — S. h. Bonpl. rhe. t. 51.
 3 argéntea. DC. (bl.) silvery-leaved. — 1816. — S. h. — t. 12.
Rhéxia holosericea. Bot. reg. t. 323. *Pleròma holosericeum*. Don.

CHÆTOGA'STRA. DC. CHÆTOGÀ'STRA. Decandria Monogynia.

- 1 lanceolâta. DC. (wh.) spear-leaved. Trinidad. 1827. 1. 5. S. h. Bot. mag. t. 2836.
 2 hypericoïdes. DC. (wh.) Hypericum-like. — 1820. 8. 12. S. (O). Aubl. guian. 1. t. 168.
Rhéxia villosissima. Bonpl. rhe. t. 31. *Melástoma villosa*. Aubl.
 3 ciliâris. DC. (pu.) fringed. S. America. 1822. 4. 7. S. ¼. Vent. choix. t. 34.
Meriana ciliaris. Vent.
 4 grátilis. DC. (ro.) slender. Brazil. 1824. — S. ¼. Bonpl. rhe. t. 52.
 ARTHROSTE'MMA. DC. ARTHROSTÈ'MMA. Octandria Monogynia.

- 1 ladanoides. DC. (li.) bristly-stalked. Trinidad. 1820. 6. 9. S. (O). Bonpl. rhe. t. 27.
Rhéxia ladanoides. Bonpl. *Melástoma trichótoma*. w.
 2 villòsum. DC. (vi.) villous. Trinidad. — — S. h. Aubl. guian. 1. t. 129. f. 1.
 3 versicolor. DC. (bh.) changeable-col'd. Brazil. 1825. 5. 9. S. h. Bot. reg. t. 1066.
Rhéxia versicolor. B.R.

OSBECKIA. DC. OSBECKIA. Octo-Decandria Monogynia.

- 1 urceolâris. DC. (pu.) urceolate. Brazil. 1825. 4. 9. S. h. Bot. mag. t. 2235.
 2 Simsii. DC. (pu.) Sims's. Mauritius. 1820. — S. h.
Melástoma osbeckioides. B.M.
 3 glomerâta. DC. (ro.) cluster-flower'd. S. America. 1821. 12. 4. S. h. Bot. mag. t. 2838.
 β *albiflora* DC. (wh.) white-flowered. Brazil. — — S. h. Lodd. bot. cab. t. 334.
Rhéxia glomerata. B.C.
 4 chinénica. DC. (ro.) Chinese. China. 1816. 3. 10. S. h. Bot. reg. t. 542.
 5 zeylânica. DC. (wh.) Ceylon. Ceylon. 1749. 5. 9. S. h. — t. 565.
 6 angustifoliâ. DC. (ro.) narrow-leaved. Nepaul. 1826. — S. h.
 7 nepalénica. H.E.F. (ro.) handsome. — 1820. — S. h. Hook. ex. flor. t. 31.
speciosa. D.P.
 8stellâta. DC. (li.) starry-haired. — 1818. — S. h.
 9 ternifoliâ. DC. (li.) whorl-leaved. — 1825. — S. h.
 10 octândra. DC. (ro.) octandrous. Ceylon. 1815. 4. 8. S. h.
Melástoma octandra. L.

- 11 decumbens. DC. (ro.) decumbent. Africa. 1821. — S. ¼. Beauv. fl. ow. t. 41.

Melástoma decumbens. Beauv.

TIBOUCHINA. DC. TIBOUCHINA. Decandria Monogynia.

 α spera. DC. (pu.) rough. Guiana. 1820. 4. 6. S. h. Aubl. guian. 1. t. 177.

MELASTOMA. DC. MELA'STOMA. Decandria Monogynia. L.

1 affine. DC. (ro.) scaly-stalked. E. Indies. 1816. 1. 8. S. h.

2 malabáthricum. DC. (ro.) Malabar. — 1793. — S. h.

3 ásperum. DC. (ro.) rough. — 1815. 8. 11. S. h.

4 macrocárpus. DC. (ro.) large-fruited. — 1793. 1. 8. S. h.

malabáthrica. B.M. nec aliorum.

5 sanguíneum. DC. (li.) red-veined. Sunda. St. 1818. 4. 12. S. h.

6 rubrolimbatum. DC. (wh.) red-limbed. E. Indies. 1823. — S. h.

7 cándidum. DC. (li.) white-leaved. China. 1820. 4. 8. S. h.

8 normâle. DC. (li.) Nepaul. 1819. 2. 6. S. h.

nepalénica. B.C.

9 Wallíchii. DC. (li.) Wallich's. — 1824. — S. h.

10 decémfidum. DC. (pu.) ten-cleft. E. Indies. 1826. 1. 8. S. h.

11 cymosum. DC. (li.) cymose. Sierra Leone. 1792. 4. 8. S. h.

corymbosum. Bot. mag. t. 984.

12 elongatum. DC. (li.) tuberous-rooted. — 1822. S. ¼.

13 pulverulèntum. DC. (re.) powdered. Sumatra. — S. h.

PLEROMA. DC. PLEROMA. Decandria Monogynia.

1 vimineum. DC. (vi.) slender. Brazil. 1820. 4. 8. S. h.

Rhéxia viminea. B.R.

2 heteromállum. DC. (bl.) woolly-leaved. — — — S. h.

Melástoma heteromálla.

DIPLOSTEGIUM. DC. DIPLOSTEGIUM. Decandria Monogynia.

canescens. DC. (ro.) hoary-stem'm'd. Brazil. 1820. 4. 8. S. h.

ACIOTIS. DC. ACIOTIS. Octandria Monogynia.

discolor. DC. (bh.) two-coloured. W. Indies. 1793. 3. 9. S. ¼.

Tribus IV. MICONIEÆ. DC. prod. v. 3. p. 152.

CLID'EMIA. DC.	CLID'EMIA. Decandria Monogynia.
1 aggregata. DC. (wh.) clustered.	Peru. 1820. 7. 10. S. h.
2 hirta. DC. (wh.) hairy.	Jamaica. 1740. 3. 6. S. h.
<i>Melastoma hirta.</i> B.M.	
3 crenata. DC. (wh.) crenated.	W. Indies. 1820. — S. h.
4 elegans. DC. (wh.) elegant.	Trinidad. 1822. — S. h.
5 biserrata. DC. (pu.) doubly serrate.	Brazil. 1825. — S. h.
6 spicata. DC. (wh.) spiked.	Trinidad. 1820. — S. h.
7 strigillosa. DC. (wh.) strigose.	Jamaica. 1825. — S. h.
8 agréstis. DC. (wh.) field.	Cayenne. 1822. 4. 8. S. h.
9 lanata. DC. (wh.) sweet-scented.	Trinidad. — — S. h.
TOC'OCA. DC.	TOC'OCA. Decandria Monogynia.
guianensis. DC. (ro.) Guiana.	Guiana. 1822. 4. 7. S. h.
<i>Melastoma physiphora.</i> Vahl.	
MAI'ETA. DC.	MAI'ETA. Decandria Monogynia.
guianensis. DC. (ro.) Guiana.	Guiana. 1824. S. h.
OSS'EA. DC.	OSS'EA. Octandria Monogynia.
purpurascens. DC. (pu.) purplish.	Jamaica. 1822. 3. 6. S. h.
<i>Melastoma purpurascens.</i> Swartz.	
SAGR'EA. DC.	SAGR'EA. Octandria Monogynia.
sessiliflora. DC. (re.) sessile-flowered.	Guiana. 1793. 4. 8. S. h.
<i>Melastoma rúbra.</i> Bonpl. mel. t. 39.	Vahl. ic. am. t. 19.
2 hirtella. DC. (wh.) hispid-fruited.	Jamaica. 1824. — S. h.
3 umbrósa. DC. (wh.) shaded.	W. Indies. — 3. 7. S. h.
4 pilosa. DC. (wh.) pilose.	Jamaica. 1828. — S. h.
5 microphylla. DC. (wh.) small-leaved.	— 1820. 5. 6. S. h.
6 hirsuta. DC. (wh.) purple-haired.	— 1823. — S. h.
TETRAZY'GIA. DC.	TETRAZY'GIA. Tetr-Octandria Monogynia.
tetrandra. DC. (wh.) tetrandrous.	Jamaica. 1815. 3. 6. S. h.
<i>Melastoma tetrandra.</i> Swartz.	
2 discolor. DC. (wh.) two-coloured.	W. Indies. 1793. 5. 7. S. h.
<i>Melastoma discolor.</i> Jacq. amer. t. 84.	Bonpl. mel. t. 34.
3 angustifolia. DC. (wh.) narrow-leaved.	— 1823. — S. h.
HETEROTRÍCHUM. DC.	HETEROTRÍCHUM. Decandria Monogynia.
níveum. DC. (bh.) snowy-white.	St. Domingo. 1820. 5. 10. S. h.
<i>Melastoma nivea.</i> Lam.	
2 pátns. DC. (bh.) spreading.	— 1825. 4. 8. S. h.
CONOSTEGIA. DC.	CONOSTEGIA. Dec-Dodecandria Monogynia.
procera. DC. (wh.) tall.	Jamaica. 1822. S. h.
2 Balbiiana. DC. (wh.) Balbis's.	— 1825. S. h.
3 semicrenata. DC. (wh.) semicrenate.	W. Indies. 1823. 4. 6. S. h.
DIPLOCHITA. DC.	DIPLOCHITA. Octo-Decandria Monogynia.
Fothergilla. DC. (wh.) Fothergill's.	Trinidad. 1815. S. h.
<i>Melastoma Fothergilla.</i> s.s. <i>Fothergilla mirabilis.</i> Aubl. gui. 1. t. 175.	Bonpl. mel. t. 32.
2 Swartziana. DC. (ro.) Swartz's.	Jamaica. 1815. 4. 6. S. h.
<i>Melastoma Tamônea.</i> Swartz.	Bonpl. mel. t. 33.
3 macrophylla. DC. large-leaved.	Mexico. 1820. S. h.
L'OREYA. DC.	L'OREYA. Decandria Monogynia.
arborescens. DC. (wh.) arborescent.	Guiana. 1822. S. h.
<i>Melastoma arborescens.</i> Aubl.	Aubl. guian. 1. t. 163.
MIC'ONIA. DC.	MIC'ONIA. Decandria Monogynia.
racemosa. DC. (ro.) racemed.	Trinidad. 1822. 4. 8. S. h.
<i>Melastoma racemosa.</i> Aubl. guian. t. 156.	Bonpl. mel. t. 27.
2 purpurascens. DC. (wh.) purple-berried.	Guiana. 1804. 3. 6. S. h.
<i>Melastoma purpurea.</i> W.	Aubl. guian. t. 154.
3 ciliata. DC. (wh.) fringed.	— 1824. 4. 7. S. h.
4 holosericea. DC. (wh.) velvety.	W. Indies. 1816. — S. h.
5 argentea. DC. (wh.) silvery-leaved.	— 1827. — S. h.
6 impetiolaris. DC. (wh.) great-leaved.	— 1822. — S. h.
<i>Melastoma macrophylla.</i> Lam.	Bonpl. mel. t. 29.
7 Lambertiæna. DC. (wh.) Mr. Lambert's.	— 1824. — S. h.
8 caudata. DC. (ro.) tailed-leaved.	S. America. 1822. 5. 8. S. h.
9 pyramidalis. DC. (wh.) pyramidal.	Trinidad. — — S. h.
<i>Melastoma pyramidalis.</i> Lam.	Pluk. alm. t. 159. f. 1.

MELASTOMACEÆ.

10	Acinodéndron. (wh.) oval-leaved. <i>Melástoma Acinodéndron.</i> L.	Jamaica.	1804.	5. 8.	S. $\frac{1}{2}$.	Plum. ic. 142. f. 2.
11	lævigåta. DC. (wh.) smooth.	Caribees.	1815.	4. 8.	S. $\frac{1}{2}$.	Bot. reg. t. 363.
12	prasina. DC. (wh.) green.	Jamaica.	1824.	—	S. $\frac{1}{2}$.	
13?	ecostata. (pu.) ribless. <i>Melástoma ecostata.</i> H.K.	—	1793.	5. 6.	S. $\frac{1}{2}$.	
	BL'AKEA. DC.	BL'AKEA. Dodecadria Monogynia. L.				
1	trinérvia. DC. (ro.) three-ribbed.	Jamaica.	1789.	6. 7.	S. $\frac{1}{2}$.	Bot. mag. t. 451.
2	quinquénervis. DC. (fl.) five-nerved. <i>triplinérvia.</i> L.	Guiana.	1820.	—	S. $\frac{1}{2}$.	Aubl. guian. I. t. 210.

ORDO XCII.

ALANGIEÆ. DC. prodr. 3. p. 203.

ALA'NGIUM. DC.	ALA'NGIUM. Icosandria Monogynia.					
1	decapétalum. DC. (wh.) Sage-leaved. <i>Gre'wia salvifolia.</i> L.	E. Indies.	1779.	5. 8.	S. $\frac{1}{2}$.	Rheed. mal. 4. t. 17.
2	hexapétalum. DC. (wh.) six-petaled.	—	1822.	—	S. $\frac{1}{2}$.	Rheed. mal. 4. t. 26.

ORDO XCIII.

PHILADELPHEÆ. DC. prodr. v. 3. p. 205.

PHILA'DE'LPHUS. DC.	SYRINGA. Icosandria Monogynia. L.					
1	coronárius. DC. (wh.) common. β variegátus.	S. Europe.	1596.	5. 6.	H. $\frac{1}{2}$.	Bot. mag. t. 391.
	(wh.) striped-leaved.	—	—	—	H. $\frac{1}{2}$.	
	γ multipléz.	(wh.) double-flowered.	—	—	H. $\frac{1}{2}$.	
2	nánus. Mil.	(wh.) dwarf.	—	6. 7.	H. $\frac{1}{2}$.	
3	Zeyhéri. DC.	(wh.) Zeyher's.	N. America.	1812.	—	H. $\frac{1}{2}$.
4	floribún dus. DC.	(wh.) many-flowered.	—	1816.	—	H. $\frac{1}{2}$.
5	verrucósus. DC.	(wh.) warted. <i>grandiflórus.</i> Bot. reg. t. 570. non w.	—	1812.	—	H. $\frac{1}{2}$.
6	latifólius. DC.	(wh.) broad-leaved.	—	1820.	—	H. $\frac{1}{2}$.
7	grandiflórius. DC.	(wh.) large-flowered.	—	1816.	—	H. $\frac{1}{2}$.
8	speciosus. DC.	(wh.) beautiful.	—	1820.	—	H. $\frac{1}{2}$.
9	láxus. DC.	(wh.) loose-branched.	—	1816.	—	H. $\frac{1}{2}$.
10	hirsútus. DC.	(wh.) slender hairy.	—	—	—	
11	indórius. DC.	(wh.) entire-leaved.	Carolina.	1738.	—	H. $\frac{1}{2}$.
	DECUM'ARIA. DC.	Dodecadria Monogynia. L.				
1	bárbara. Ph.	(wh.) smaller.	Carolina.	1785.	7. 8.	H. $\frac{1}{2}$.
2	sarmentósa. Ph.	(wh.) larger.	—	1758.	—	H. $\frac{1}{2}$.
						Bosc. act. par. I. t. 13.

ORDO XCIV.

MYRTACEÆ. DC. prodr. v. 3. p. 207.

Tribus I. CHAMÆLAUCIEÆ. DC. prodr. 3. p. 208.

CA'LYTHRIX. DC.	CA'LYTHRIX. Icosandria Monogynia.					
1	glábria. DC.	(wh.) smooth.	N. S. W.	1818.	4. 6.	G. $\frac{1}{2}$.
2	ericoides. F.T.	(wh.) Heath-like.	—	1823.	—	G. $\frac{1}{2}$.
3	scábra. DC.	(wh.) rough-leaved.	—	1824.	—	G. $\frac{1}{2}$.

DARWI'NIA. L.T.	DARWI'NIA. Decandria Monogynia.					
1 fascicularis. L.T. (wh.) fascicled.	N. S. W.	1826.	4. 6.	G. h.	Linn. trans. v. 11. t. 22.	
2 taxifolia. F.T. (wh.) Yew-leaved.		1827.	—	G. h.		
VERTICO'R'DIA. DC.	VERTICO'R'DIA. Decandria Monogynia.					
1 Fontanésii. DC. (wh.) Desfontaines'. Chamaelæcum plumosum. Df.	N. Holland.	1826.	4. 6.	G. h.	Desf. mem. mus. 5. t. 4.	
2 Brownii. DC. (wh.) Brown's.		—	—	G. h.	Desf. mem. mus. 5. t. 19.	
CHAMÆLAUCIUM. DC.	CHAMÆLAUCIUM. Decandria Monogynia.					
ciliatum. DC. (wh.) fringed.	N. Holland.	1825.	4. 6.	G. h.	Df. mem. mus. 5. t. 3. f.B.	
GENETYL'LIS. DC.	GENETYL'LIS. Icosandria Monogynia.					
diosmóides. DC. (wh.) Diosma-like.	N. Holland.	1827.	4. 6.	G. h.	DC. dict. cl. v. 11. c. ic.	
PILE'ANTHUS. DC.	PILE'ANTHUS. Icosandria Monogynia.					
Limacis. DC. (wh.) clubbed-leaved.	N. Holland.	1824.	4. 7.	G. h.	Lab. n. holl. 2. t. 149.	

Tribus II. LEPTOSPERMEÆ. DC. prodr. 3. p. 209.

Subtribus I. MELALEUCEÆ. Stamina polyadelpa. DC. loc. cit. p. 210.

ASTA'RTEA. DC.	ASTA'RTEA. Polyadelphia Icosandria.						
fasciculáris. DC. (wh.) fascicled.	V. Diem. Isl.	1824.	4. 6.	G. h.	Lab. n. holl. 2. t. 170.		
Melaleuca fasciculáris. Labill.							
TRISTANIA. DC.	TRISTANIA. Polyadelphia Icosandria.						
1 nerifolia. DC. (ye.) Oleander-leav'd. N.S.W.		1804.	7. 9.	G. h.	Bot. mag. t. 1058.		
Melaleuca nerifolia. B.M. salicifolia. Andr. deposit. t. 485.							
2 persicafolia. F.T. (ye.) Peach-leaved.	N. S. W.	1824.	—	G. h.			
3 laurina. DC. (ye.) Laurel-leaved.		1798.	—	G. h.			
4 deprésssa. DC. (ye.) depressed.		1820.	—	G. h.			
5 álbens. DC. (ye.) white-leaved.		1822.	—	G. h.			
6 conférta. DC. (ye.) Pittosporum-ld.		1805.	—	G. h.			
7 suavéolens. DC. (ye.) sweet-scented.	N. Holland.	1826.	G. h.	Gärt. fruct. 1. t. 35.		
BEAUFO'R'TIA. DC.	BEAUFO'R'TIA. Polyadelphia Icosandria.						
1 decussáta. DC. (sc.) splendid.	N. Holland.	1803.	3. 8.	G. h.	Bot. reg. t. 18.		
2 spársa. DC. (sc.) alternate-leav'd.		—	—	G. h.			
CALOTHA'MNUS. DC.	CALOTHA'MNUS. Polyadelphia Icosandria.						
1 quadrífida. DC. (sc.) four-cleft.	N. Holland.	1803.	1. 12.	G. h.	Bot. mag. t. 1506.		
2 clavata. M.C.	(sc.) club-leaved.		1826.	—	G. h.	Lodd. bot. cab. t. 1447.	
3 villósa. DC.	(sc.) hairy 5-cleft.		1803.	—	G. h.		
4 grátilis. DC.	(sc.) long-leaved.		—	—	G. h.	Bot. reg. t. 1099.	
MELALEUCA. DC.	MELALEUCA. Polyadelphia Icosandria.						
1 Leucadéndron. DC. (wh.) aromatic.	E. Indies.	1796.	S. h.	Rumph. amb. 2. t. 16.		
2 minór. DC. (wh.) Cajuputi-tree.		1816.	S. h.	Rumph. amb. 2. t. 17. f. 1.		
3 viridiflóra. DC. (gr.) green-flowered.	N. S. W.	1798.	4. 6.	G. h.	Cavan. ic. t. 333.		
Metrosidéros quinquenervia. Cavan.							
4 paludósa. DC. (re.) long-leaved red.	N. Holland.	1803.	7. 9.	G. h.			
5 globifera. DC. (wh.) globe-fruited.		—	—	G. h.			
6 diosmifolia. DC. (gr.) Diosma-leaved.		1794.	6. 7.	G. h.	Andr. deposit. t. 476.		
7 styphelioídes. DC. (wh.) Stypelia-leav'd.	N. S. W.	1793.	4. 6.	G. h.			
8 genistifolia. DC. (wh.) Genista-leaved.		—	—	G. h.			
9 lanceolátâ. DC. (st.) spear-leaved.	N. Holland.	1812.	5. 6.	G. h.			
10 striâta. DC. (wh.) rigid-leaved.		1803.	—	G. h.			
11 thymoídes. DC. (ye.) yellow-spear leav'd.		—	—	G. h.			
12 squámea. DC. (pu.) scaly-branched.	V. Diem. Isl.	1805.	6. 8.	G. h.			
13 nodosa. DC.	(wh.) Whin-leaved.	N. S. W.	1790.	—	G. h.		
14 tenuifolia. DC.	(wh.) slender-leaved.		1824.	—	G. h.		
15 ericifolia. DC.	(wh.) Heath-leaved.		1788.	7. 9.	G. h.		
16 erubescens. DC.	(li.) Lilac-coloured.		1820.	—	G. h.		
17 armillâris. DC.	(wh.) pale-flowered.		1788.	6. 8.	G. h.		
ericifolia. Andr. rep. t. 175. non Sm.	Metrosidéros armillâris.	G.					
18 uncinâta. DC.	(re.) hook-leaved.	N. Holland.	1803.	—	G. h.		
19 scábra. DC.	(pu.) rough-leaved.		—	—	G. h.		
20 juniperoides. DC.	(st.) Juniper-like.	N. S. W.	1826.	—	G. h.		
21 pulchélla. DC.	(li.) pendulous.	N. Holland.	1803.	6. 9.	G. h.	Lodd. bot. cab. t. 200.	
22 thymifolia. DC.	(pu.) Thyme-leaved.	N. S. W.	1792.	—	G. h.	Sm. exot. bot. 1. t. 36.	
coronata. Andr. rep. t. 278.	gnidiefolia. Vent. malm. t. 7.						
23 decussáta. DC.	(li.) cross-leaved.	N. Holland.	1803.	6. 9.	G. h.	Bot. mag. t. 2268.	
24 tetragóna. DC.	(li.) four-sided.		1824.	—	G. h.		

25 <i>cuticulâris</i> . DC.	(<i>pu.</i>) solitary-flower'd. N. Holland.	1824.	6. 9. G. h.	Lab. n. holl. 2. t. 171.	
26 <i>fúlgens</i> . DC.	(<i>sc.</i>) splendid.	1803.	6. 8. G. h.	Bot. reg. t. 103.	
27 <i>linariifolia</i> . DC.	(<i>wh.</i>) Toad-flax-leav'd. N. S. W.	1793.	5. 7. G. h.	Sm. exot. bot. 1. t. 56.	
28 <i>abietina</i> . DC.	(<i>wh.</i>) Fir-leaved.	N. Holland.	1824.	G. h.	
29 <i>hypericifolia</i> . DC.	(<i>sc.</i>) Hypericum-ld.	N. S. W.	1792.	7. 9. G. h.	Andr. deposit. t. 200.
30 <i>elliptica</i> . DC.	(<i>cr.</i>) elliptic-leaved.	N. Holland.	1824.	G. h.	Lab. n. holl. 2. t. 173.
31 <i>squarrôsa</i> . DC.	(<i>wh.</i>) Myrtle-leaved.	N. S. W.	1792.	G. h.	Bot. mag. t. 1935.
	<i>myrtifolia</i> . Vent. malm. t. 47.				
32 <i>gibbosa</i> . DC.	(<i>li.</i>) gibbous.	V. Diem. Isl.	1822.	6. 8. G. h.	Lab. n. holl. 2. t. 172.
33 <i>imbricata</i> . L.en.	(<i>wh.</i>) imbricated.	N. Holland.	1803.	5. 6. G. h.	
34 <i>sprengelioïdes</i> . DC.	(<i>wh.</i>) Sprengelia-like.	N. S. W.	1824.	G. h.	
35 <i>calycina</i> . DC.	(<i>wh.</i>) permanent-cup'd.	N. Holland.	1803.	6. 8. G. h.	
36 <i>densa</i> . DC.	(<i>pu.</i>) dense-leaved.			G. h.	
37 <i>incana</i> . DC.	(<i>st.</i>) hoary.	N. S. W.	1812.	G. h.	Bot. reg. t. 410.
	<i>canescens</i> . Otto hort. ber. p. 87. Abbild. t. 81.				
EUDE'SMIA. DC.	EUDE'SMIA.	Polyadelphia Icosandria.			
tetragona. DC.	(<i>wh.</i>) square-stemmed.	N. Holland.	1824.	6. 8. G. h.	Swt. flor. aust. t. 21.

Subtribus II. EULEPTOSPERMEÆ. Stamina libera. DC. prod. 3. p. 216.

EUCALY'PTUS. DC.	EUCALY'PTUS.	Icosandria Monogynia.			
1 <i>cornuta</i> . DC.	(<i>wh.</i>) horned.	N. Holland.	1803.	3. 6. G. h.	Lab. voy. 1. t. 20.
2 <i>tereticornis</i> . DC.	(<i>wh.</i>) long-horned.	N. S. W.	1804.	4. 7. G. h.	Andr. deposit. t. 400.
3 <i>resinifera</i> . DC.	(<i>wh.</i>) red gum-tree.		1788.	G. h.	Link et Ot. abb. t. 45.
4 <i>longifolia</i> . DC.	(<i>wh.</i>) long-leaved.		1818.	G. h.	Smith nov. holl. t. 13.
5 <i>robusta</i> . DC.	(<i>wh.</i>) brown gum-tree.		1794.	8. 9. G. h.	Cavan. ic. 4. t. 342.
6 <i>rostrata</i> . Cav.	(<i>wh.</i>) beaked.		1804.	G. h.	
7 <i>marginata</i> . DC.	(<i>wh.</i>) thick-edged.		1794.	4. 7. G. h.	
8 <i>incrassata</i> . DC.	(<i>wh.</i>) thick-leaved.	N. Holland.	1820.	G. h.	Lab. nov. holl. 2. t. 150.
9 <i>persicifolia</i> . DC.	(<i>wh.</i>) Peach-leaved.	N. S. W.	1812.	G. h.	Lodd. bot. cab. t. 501.
10 <i>punctata</i> . DC.	(<i>wh.</i>) black-dotted.		1810.	G. h.	
11 <i>acervula</i> . DC.	(<i>wh.</i>) flat-peduncled.		1818.	G. h.	
12 <i>virgata</i> . DC.	(<i>wh.</i>) slender-twiggled.		1820.	5. 8. G. h.	
13 <i>micrantha</i> . DC.	(<i>wh.</i>) small-flowered.		1824.	G. h.	
14 <i>stellulata</i> . DC.	(<i>wh.</i>) starry.		1816.	G. h.	
15 <i>oblonga</i> . DC.	(<i>wh.</i>) oblong-leaved.		1819.	6. 8. G. h.	
16 <i>viminialis</i> . DC.	(<i>wh.</i>) slender.	N. Holland.	1816.	G. h.	Lab. nov. holl. 2. t. 151.
17 <i>capitellata</i> . DC.	(<i>wh.</i>) small-headed.	N. S. W.	1804.	4. 7. G. h.	White's voy. 226. ic.
18 <i>triangularis</i> . L.en.	(<i>wh.</i>) three-flowered.		1820.	4. 6. G. h.	
19 <i>saligna</i> . DC.	(<i>wh.</i>) Willow-like.		1804.	4. 7. G. h.	
20 <i>angustifolia</i> . L.en.	(<i>wh.</i>) narrow-leaved.		1816.	G. h.	
21 <i>microphylla</i> . DC.	(<i>wh.</i>) small-leaved.			G. h.	
22 <i>stenophylla</i> . DC.	(<i>wh.</i>) slender-leaved.			G. h.	
23 <i>myrtifolia</i> . DC.	(<i>wh.</i>) Myrtle-leaved.		1812.	4. 6. G. h.	
24 <i>elongata</i> . DC.	(<i>wh.</i>) taper-pointed.		1820.	G. h.	
25 <i>média</i> . DC.	(<i>wh.</i>) acuminate.			G. h.	
26 <i>reticulata</i> . DC.	(<i>wh.</i>) netted-leaved.		1816. G. h.	
27 <i>umbellata</i> . DC.	(<i>wh.</i>) umbel-flowered.		1820.	5. 8. G. h.	
28 <i>ovata</i> . DC.	(<i>wh.</i>) oval-leaved.	N. Holland.		G. h.	Lab. nov. holl. 2. t. 153.
	<i>muconata</i> . L. enum.				
29 <i>scabra</i> . DC.	(<i>wh.</i>) rough-leaved.	N. S. W.	1803.	G. h.	
30 <i>pilularis</i> . DC.	(<i>wh.</i>) small-fruited.		1804.	2. 5. G. h.	
31 <i>radiata</i> . DC.	(<i>wh.</i>) rayed.		1824.	3. 6. G. h.	
32 <i>stricta</i> . DC.	(<i>wh.</i>) straight.		 G. h.	
33 <i>haemastomma</i> . DC.	(<i>wh.</i>) red-mouthed.		1803.	4. 7. G. h.	White's voy. 226. c. ic.
	<i>racemosa</i> . Cavan. ic. 4. n. 377.				
34 <i>ligustrina</i> . DC.	(<i>wh.</i>) Privet-like.	N. S. W.	1824. G. h.	
35 <i>amygdalina</i> . DC.	(<i>wh.</i>) Almond-leaved.	N. Holland.	1822. G. h.	Lab. nov. holl. 2. t. 154.
36 <i>ambigua</i> . DC.	(<i>wh.</i>) ambiguous.		1824. G. h.	
37 <i>Lindleyana</i> . DC.	(<i>wh.</i>) Lindley's.	N. S. W.	1816.	4. 6. G. h.	Bot. reg. t. 947.
	<i>longifolia</i> . B.R. non Link.				
38 <i>botryoides</i> . DC.	(<i>wh.</i>) clustered.		1803.	4. 7. G. h.	Cavan. ic. 4. t. 341.
	<i>platypodus</i> . Cavan.				
39 <i>piperita</i> . DC.	(<i>wh.</i>) Peppermint-tree.		1788.	G. h.	Reich. gart. mag. t. 42.
40 <i>pallens</i> . DC.	(<i>wh.</i>) white-leaved.		1822.	G. h.	
41 <i>obliqua</i> . DC.	(<i>wh.</i>) oblique-leaved.	V. Diem. Isl.	1774.	4. 8. G. h.	Sal. par. lond. t. 15.

42	corymbosa. DC. (<i>wh.</i>) corymbose-flid.	N. S. W.	1788.	4. 8. G. h.	Cavan. ic. 4. t. 340.
43	paniculata. DC. (<i>wh.</i>) panicled.	_____	1801.	6. 8. G. h.	
44	cneorifolia. DC. (<i>wh.</i>) Willow-wail-ld.	_____	1824.	— G. h.	
45	obtusifolia. DC. (<i>wh.</i>) blunt-leaved.	_____	— G. h.	
46	Cunninghami. (<i>wh.</i>) Cunningham's.	_____	1825. G. h.	
	<i>microphylla. F.T. non Link.</i>				
47	gomphocéphala. DC. (<i>wh.</i>) persistent-headed.	_____	1824. G. h.	
48	Glòbulus. DC. (<i>wh.</i>) large tree.	V. Diem. Isl.	1804.	5. 6. G. h.	Lab. voy. 1. t. 13.
	<i>pulverulenta. L. enum. non B.M. glauca. DC.</i>				
49	diversifolia. DC. (<i>wh.</i>) various-leaved.	N. S. W.	1810.	5. 8. G. h.	Bonpl. nav. 1. t. 13.
50	perfoliata. C.C. (<i>wh.</i>) perfoliate glaucous.	N. Holland.	1803.	4. 6. G. h.	
51	cordata. DC. (<i>wh.</i>) heart-leaved.	V. Diem. Isl.	1816.	11. 3. G. h.	Lab. nov. holl. 2. t. 152.
52	pulverulenta. DC. (<i>wh.</i>) powdered.	N. S. W.	—	— G. h.	Bot. mag. t. 2087.
53	purpurascens. DC. (<i>wh.</i>) purple-branch'd.	_____	1804. G. h.	
54	tuberculata. DC. (<i>wh.</i>) warted-branch'd.	_____	1816. G. h.	
55	rigida. DC. (<i>wh.</i>) rigid-leaved.	_____	— G. h.	
56	hypericifolia. (<i>wh.</i>) Hypericum-leaved.	_____	1804. G. h.	
	ANGÓ'PHORA. DC. ANGÓ'PHORA. Icosandria Monogynia.				
1	cordifolia. DC. (<i>wh.</i>) hispid.	N. S. W.	1789.	5. 8. G. h.	Bot. mag. t. 1960.
	<i>Metrosideros hispida. Sm. exot. bot. 1. t. 42. hirsuta. Andr. rep. t. 281. anomala. Vent. malm. t. 2.</i>				
2	floribunda. (<i>wh.</i>) many-flowered.	N. S. W.	1788.	— G. h.	
	<i>Metrosideros floribunda. L.T. non Vent.</i>				
3	intermédia. DC. (<i>wh.</i>) intermediate.	_____	1820.	— G. h.	
4	lanceolata. DC. (<i>wh.</i>) ribbed.	_____	1816.	— G. h.	Cavan. ic. 4. t. 337.
	<i>Metrosideros costata. Gaert. fruct. 1. t. 34. f. 2.</i>				
	CALLISTEMON. DC. CALLIST'EMON. Icosandria Monogynia.				
1	pinifolium. DC. (<i>gr.</i>) Pine-tree-leav'd.	N.S.W.	1806.	6. 8. G. h.	Wendl. coll. 1. t. 16.
2	viridiflorum. DC. (<i>gr.</i>) green-flowered.	_____	1820.	— G. h.	Bot. mag. t. 2602.
	<i>ruscifolium. C.C. Metrosideros viridiflora. B.M.</i>				
3	lophanthum. S.F.A. (<i>st.</i>) crested.	_____	1816.	— G. h.	Swt. flor. aust. t. 29.
4	leptoştachyum. S.F.A. (<i>st.</i>) slender-spik'd.	_____	—	— G. h.	
5	salignum. DC. (<i>st.</i>) Willow-leaved.	_____	1788.	5. 7. G. h.	Bonpl. nav. t. 4.
	<i>Metrosideros saligna. Vent. cels. t 70. non Bot. mag.</i>				
6	pállidum. DC. (<i>st.</i>) pale-leaved.	N. S. W.	1816.	— G. h.	Bonpl. nav. p. 101. t. 41.
7	Siebéri. DC. (<i>st.</i>) Sieber's.	_____	1820.	— G. h.	
8	rígidum. DC. (<i>sc.</i>) rigid-leaved.	_____	1800.	3. 8. G. h.	Bot. reg. t. 393.
9	lineare. DC. (<i>sc.</i>) linear.	_____	1788.	6. 7. G. h.	Sert. hann. t. 11.
10	rugulésum. DC. (<i>sc.</i>) tubercle-leaved.	_____	1820.	— G. h.	
11	scábrum. B.C. (<i>sc.</i>) rough-leaved.	N. Holland.	1824.	— G. h.	Lodd. bot. cab. t. 1288.
12	linearifolium. DC. (<i>sc.</i>) linear-leaved.	_____	1820.	— G. h.	
13	microphyllum. M.C. (<i>sc.</i>) small-leaved.	_____	1825.	— G. h.	
14	formosum. M.C. (<i>sc.</i>) handsome.	_____	—	— G. h.	
15	lanceolatum. DC. (<i>sc.</i>) spear-leaved.	N. S. W.	1788.	11. 6. G. h.	Bot. mag. t. 260.
	<i>Metrosideros citrina. B.M. lanceolata. L.T.</i>				
	β semperflorens. B.C. (<i>sc.</i>) ever-blowing.	_____	—	— G. h.	Lodd. bot. cab. t. 523.
16	marginatum. DC. (<i>sc.</i>) thick-margined.	_____	1816.	6. 8. G. h.	Cavan. ic. 4. t. 332.
17	gláicum. (<i>sc.</i>) splendid.	N. Holland.	1803.	3. 6. G. h.	Bonpl. nav. t. 34.
	<i>Metrosideros glauca. Bonp. speciosa. Bot. mag. t. 1761.</i>				
18	hybridum. DC. (<i>sc.</i>) hybrid.	N. S. W.	1824.	4. 6. G. h.	
	METROSID'EROS. DC. METROSID'EROS. Icosandria Monogynia.				
1	véra. DC. (<i>gr.</i>) Ironwood.	E. Indies.	1815.	5. 8. S. h.	Lindl. coll. t. 18.
2	flórida. DC. (<i>st.</i>) many-flowered.	N. Zealand.	1820.	— G. h.	
3	glomulifera. DC. (<i>st.</i>) cluster-flower'd.	N. S. W.	1805.	— G. h.	
4	umbellata. DC. (<i>re.</i>) umbel-flowered.	_____	1822.	— G. h.	Cavan. ic. 4. t. 337.
5	angustifolia. DC. (<i>wh.</i>) narrow-leaved.	C. B. S.	1787.	— G. h.	Hout. pfly. sys. 3. t. 25. f. 2.
	STENOSPERMUM. STENOSPERMUM. Icosandria Monogynia.				
1	capitatum. (<i>li.</i>) headed.	N. S. W.	1824.	4. 6. G. h.	Reich. hort. bot. 1. t. 84.
	<i>Metrosideros capitata. DC.</i>				
2	corifolium. (<i>wh.</i>) Coris-leaved.	_____	1791.	6. 7. G. h.	Vent. malm. t. 46.
	<i>Leptospermum ambiguum. Sm. exot. bot. t. 59. Metrosideros corifolia. Vent.</i>				
	AG'ONIS. DC. AG'ONIS. Icosandria Monogynia.				
1	marginata. DC. (<i>wh.</i>) white margined.	N. Holland.	1816.	6. 7. G. h.	Lab. nov. holl. 2. t. 148.
	<i>Leptospermum marginatum. Labill.</i>				
2	flexuosa. DC. (<i>wh.</i>) flexuose.	_____	1812.	— G. h.	Colla h. rip. ap. 1. t. 2.
	<i>Leptospermum flexuosum. L. enum. Metrosideros flexuosa. W. enum.</i>				
3	salígra. (<i>wh.</i>) Willow-like.	N. S. W.	1812.	6. 7. G. h.	
4	ovalifolia. (<i>wh.</i>) oval-leaved.	N. Holland.	—	— G. h.	
	<i>Fabricia myrtifolia. Hort. nec aliorum.</i>				
5	linearifolia. DC. linear-leaved.	_____	1824. G. h.	

6 elliptica.	(wh.) elliptic-leaved.	N. Holland.	1824. G. h.
<i>Billottia elliptica</i> . M.C.				
LEPTOSPERMUM. DC. SOUTH SEA MYRTLE. Icosandria Monogynia.				
1 emarginatum. DC.	emarginate.	N. S. W.	1816.	6. 7. G. h.
2 sericeum. DC.	(wh.) silky-leaved.	V. Diem. Isl.	1820.	— G. h.
3 tuberculatum. DC. (wh.) tuberculate.	stellatum. Cavan.	N. S. W.	1816.	— G. h.
4 grandifolium. DC. (wh.) large-leaved.	—	—	1803.	— G. h.
5 lanigerum. DC. (wh.) woolly-leaved.	—	—	1774.	— G. h.
6 pubescens. w.	(wh.) hoary-leaved.	—	—	— G. h.
7 trinervé. DC.	(wh.) three-nerved.	—	1812.	— G. h.
8 grandiflorum. B.C. (wh.) large-flowered.	—	—	1816.	— G. h.
9 scoparium. DC. (wh.) N. Zealand Tea.	N. Zealand.	—	1772.	— G. h.
10 squarrosum. G.	(wh.) squarrose.	N. S. W.	—	— G. h.
11 floribundum.	(wh.) many-flowered.	—	—	— G. h.
12 rubricaulis. L.en.	(wh.) red-stemmed.	—	1812.	— G. h.
13 flavescens. DC.	(wh.) yellowish.	—	1787.	5. 7. G. h.
<i>Melaleuca Thæa</i> . Wendl. sert. han. 1. t. 14.				
14 gnidiæfolium. DC. (wh.) Gnidiæ-leaved.	—	—	1824.	— G. h.
15 thymifolium. F.T. (wh.) Thyme-leaved.	—	—	1823.	— G. h.
16 multicaule. F.T. (wh.) many-branched.	—	—	—	— G. h.
17 porphyllum. DC. (wh.) dotted-leaved.	—	—	—	— G. h.
18 parvifolium. DC. (wh.) small-leaved.	—	—	1795.	— G. h.
19 myrtifolium. DC. (wh.) Myrtle-leaved.	—	—	1824.	6. 8. G. h.
20 obovatum. S.F.A. (wh.) obovate-leaved.	—	—	1823.	— G. h.
21 attenuatum. DC. (wh.) fine-branched.	—	—	1795.	— G. h.
22 multiflorum. DC. (wh.) abundant-flower'd.	—	—	1820.	— G. h.
23 juniperinum. DC. (wh.) Juniper-leaved.	—	—	1790.	— G. h.
24 baccatum. DC.	(wh.) berry-fruited.	—	1790.	6. 7. G. h.
25 arachnoideum. DC. (w.) cobweb.	—	—	1795.	5. 7. G. h.
26 trilocularis. DC.	(wh.) trilocular.	—	1816.	— G. h.
27 dubium. DC.	(wh.) doubtful.	—	1824.	6. 7. G. h.
28 pendulum. DC.	(wh.) pendulous.	—	—	— G. h.
FABRICIA. DC. FABRICIA. Icosandria Monogynia.				
1 myrtifolia. DC.	(wh.) Myrtle-leaved.	N. S. W.	1803.	5. 6. G. h.
2 levigata. DC.	(wh.) smooth-leaved.	—	1788.	— G. h.
3 sericea.	(wh.) silky-leaved.	N. Holland.	1803. G. h.
4 stricta. B.C.	(wh.) upright.	N. S. W.	1822.	5. 7. G. h.
BÆCKEÆA. DC. BÆCKEÆA. Pent-Dodecandra Monogynia.				
1 frutescens. DC.	(wh.) Chinese.	China.	1806.	9. 1. G. h.
2 gracilis. F.T.	(wh.) slender-branch'd.	N.S.W.	1824.	6. 9. G. h.
3 virgata. DC.	(wh.) slender-twiggly.	—	1806.	8. 10. G. h.
4 linifolia. DC.	(wh.) Flax-leaved.	—	1820.	6. 8. G. h.
5 diffusa. DC.	(wh.) spreading.	—	1824.	— G. h.
6 densifolia. DC.	(wh.) close-leaved.	—	1812.	— G. h.
7 pulchella. DC.	(wh.) neat.	—	1825.	— G. h.
8 diosmifolia. DC.	(wh.) Diosma-leaved.	—	1823.	— G. h.
9 camphorata. DC.	(wh.) Camphor-scented.	—	1803.	8. 10. G. h.
10 microphylla. DC.	(wh.) small-leaved.	—	1812.	— G. h.
11 brevifolia. DC.	(wh.) short-leaved.	—	1824.	— G. h.
12 ramosissima. DC.	(wh.) branching.	—	1825.	7. 9. G. h.
13 crenulata. DC.	(wh.) notch leaved.	—	1822.	— G. h.
14 diosmoides. DC.	(wh.) Diosma-like.	—	1825.	— G. h.

Tribus III. MYRTEÆ. DC. prodr. v. 3. p. 230.

SONNERATIA. DC. SONNERATIA. Icosandria Monogynia. L.

1 acida. DC.	(re.) acid-fruited.	E. Indies.	1822. S. h.
2 alba. DC.	(wh.) white-flowered.	—	1824. S. h.
3 apetala. DC.	(gr.) petalless.	—	1826. S. h.

NELITRIS. DC. NELITRIS. Icosandria Monogynia.

paniculata. DC.	(wh.) paniced.	Moluccas.	1820. S. h.
-----------------	----------------	-----------	-------	------------

CAMPOMANESIA. DC. CAMPOMANESIA. Icosandria Monogynia.

lineatifolia. DC.	(wh.) yellow-fruited.	Peru.	1824.	4. 6. G. h.
-------------------	-----------------------	-------	-------	-------------

PSIDIUM. DC. GUAVA. Icosandria Monogynia. L.

1 pumilum. DC.	(wh.) dwarf.	E. Indies.	1824.	— S. h.
2 aromaticum. DC.	(wh.) aromatic.	W. Indies.	1779.	6. 8. S. h.

Rheed. mal. 3. t. 40.

Rumph. amb. 3. t. 73,

Sym.emb.to Ava. 3.t.25.

Flor. per. prod. t. 13.

Rumph. amb. 1. t. 49.

Aubl. guian. 1. t. 191.

Lab. nov. holl. 2. t. 147.

Cavan. ic. 4. t. 330.

Bot. mag. t. 1810.

Lodd. bot. cab. t. 1192.

White itin. 229. c. ic.

Lodd. bot. cab. t. 514.

Andr. rep. t. 622.

Forst. itin. 1. t. 22.

Rœm. et Ust. mag. 7.t.2.

Bot. mag. t. 2695.

Cav. ic. 4. t. 330. f. 2.

Swt. flor. aust. t. 36.

Cav. ic. 4. t. 331. f. 1.

Vent. malm. 89.

Cavan. ic. 4. t. 331. f. 2.

Gært. fruct. 1. t. 35. f. 3.

Vent. malm. t. 84.

Gært. fruct. 1. t. 35. f. 4.

Bot. mag. t. 1304.

Lodd. bot. cab. t. 1219.

Bot. mag. t. 2802.

Bot. mag. t. 2127.

Linn. trans. 8. t. 12.

Bot. mag. t. 2694.

3 montanum. DC. (wh.) mountain.	W. Indies.	1779.	5. 7.	S. h.	Trew. ehr. t. 43.
4 pyrifera. DC. (wh.) Pear-fruited.	—	1656.	6. 8.	S. h.	Rumph. amb. 1. t. 48.
5 pomiferum. DC. (wh.) Apple-fruited.	—	1692.	—	S. h.	Jacq. scheenb. 3.t.366.
6 sapidissimum. s.s. (wh.) pleasant.	—	1823.	—	S. h.	Linn. trans. 11. t. 17.
7 guineense. DC. (wh.) Guinea.	Guinea.	—	—	S. h.	Raddi mem. 1821. t. 1.
8 polycarpon. DC. (wh.) many-fruited.	Trinidad.	1809.	4. 8.	S. h.	
9 Araca. DC. (wh.) egg-fruited.	Brazil.	1820.	5. 7.	S. h.	
10 fluviatile. DC. (wh.) Guiana. guianense. P.S.	Guiana.	1823.	—	S. h.	
11 oligospermum. DC. (wh.) few-seeded.	Brazil.	—	—	S. h.	Lindl. coll. t. 16.
12 Cattleyanum. DC. (wh.) Mr. Cattley's.	—	1816.	1. 12.	S. h.	Bot. mag. t. 1779.
13 cordatum. DC. (wh.) spice.	W. Indies.	1811.	5. 7.	S. h.	
14 nigrum. DC. (wh.) black-fruited.	China.	—	—	S. h.	
15 rubrum. DC. (wh.) red-fruited.	—	1820.	—	S. h.	
16 chinense. L.C. (wh.) yellow Chinese.	—	1818.	—	S. h.	
17 fragrans. L.C. (wh.) fragrant.	—	—	4. 8.	S. h.	
18 myrtifolia. L.C. (wh.) Myrtle-leaved.	—	—	—	S. h.	
19 latifolium. DC. (wh.) broad-leaved.	—	1820.	6. 8.	S. h.	
20 indicum. DC. (wh.) Indian.	E. Indies.	1824.	—	S. h.	
JOSSI'NIA. DC. Jossi'NIA. Icosandria Monogynia.	—	—	—	—	
1 mespiloides. DC. (wh.) Mespilus-like.	Isl. Bourbon.	1826.	—	S. h.	
2 elliptica. DC. (wh.) elliptic-leaved.	Mauritius.	1824.	—	S. h.	
Eugénia elliptica. Lam. non Smith.	—	—	—	—	
3 buxifolia. DC. (wh.) Box-leaved.	Isl. Bourbon.	1822.	—	S. h.	
MYRTUS. DC. MYRTLE. Icosandria Monogynia. L.	—	—	—	—	
1 communis. DC. (wh.) common.	S. Europe.	1597.	7. 8.	F. h.	Lam. ill. t. 410.
A. melanocarpa. DC. (wh.) black-berried.	—	—	—	F. h.	
α romana. DC. (wh.) broad-leaved.	—	—	—	F. h.	Mill. ic. t. 184. f. 1.
β multipléx. (wh.) double-blossomed.	—	—	—	F. h.	
B. leucocarpa. DC. (wh.) white-fruited.	Levant.	—	—	F. h.	
2 tarentina. (wh.) Box-leaved.	S. Europe.	1597.	—	F. h.	
3 italica. (wh.) upright Italian.	—	—	—	F. h.	
β variegata. (wh.) striped-leaved.	—	—	—	F. h.	
γ maculata. (wh.) blotch-leaved.	—	—	—	F. h.	
4 bætica. (wh.) Orange-leaved.	—	1597.	—	F. h.	Blackw. t. 114.
5 lusitánica. (wh.) Portugal.	Portugal.	—	—	F. h.	Clus. hist. 1. p. 66. f. 1.
6 bélga. (wh.) sharp-leaved.	S. Europe.	—	—	F. h.	
7 mucronata. (wh.) small-leaved.	—	—	—	F. h.	
8 tomentosa. DC. (ro.) woolly-leaved.	China.	1776.	5. 9.	S. h.	Bot. mag. t. 250.
9 tenuifolia. DC. (wh.) slender-leaved.	N. S. W.	1824.	—	G. h.	
10 lúcida. DC. (wh.) shining.	Surinam.	1793.	4. 8.	S. h.	
MYRCIA. DC. MYRCIA. Icosandria Monogynia.	—	—	—	—	
1 punctata. DC. (wh.) punctated.	Santa Cruz.	1825.	—	S. h.	Plum. ic. t. 208. f. 2.
2 coriacea. DC. (wh.) Sumach-leaved.	Hispaniola.	1759.	5. 7.	S. h.	Pluk. alm. t. 155. f. 3.
3 acris. DC. (wh.) broad-lid.	Allspice. W. Indies.	—	—	S. h.	Lodd. bot. cab. t. 178.
4 pimentooides. DC. (wh.) Allspice-like.	—	—	—	S. h.	
Myrtus Piménta. β latifolia. Lodd. bot. cab. t. 178.	—	—	—	—	
5 divaricata. DC. (wh.) spreading-flow'd.	W. Indies.	1820.	—	S. h.	Plum. ic. t. 208. f. 1.
6 sororia. DC. (wh.) sister.	Trinidad.	1822.	—	S. h.	
7 spléndens. DC. (wh.) glossy-leaved.	Hispaniola.	—	—	S. h.	Jacq. coll. 2. t. 4.
Eugénia periplocafolia. Jacq.	—	—	—	—	
8 Mini. (wh.) small-fruited.	Guiana.	1803.	—	S. h.	Aubl. guian. 1. t. 197.
Eugénia Mini. Aubl.	—	—	—	—	
9 crassinervia. DC. (wh.) thick-nerved.	—	1780.	—	S. h.	
10 bracteata. DC. (wh.) leafy-bracted.	Brazil.	1824.	—	S. h.	
11 pseudo-Mini. DC. (wh.) Brazilian.	—	1822.	—	S. h.	
CALYPTRA'NTHES. DC. CALYPTRA'NTHES. Icosandria Monogynia.	—	—	—	—	
1 rígida. DC. (wh.) rigid-leaved.	Jamaica.	1822.	—	S. h.	Brown. jam. t. 37. f. 2.
2 Chytracilia. DC. (wh.) forked-paniced.	—	1778.	3. 5.	S. h.	—
3 Zuzy'gium. DC. (wh.) oval-leaved.	—	—	5. 7.	S. h.	240. t. 7. f. 2.
SYZYGIUM. DC. SYZYGIUM. Icosandria Monogynia.	—	—	—	—	
1 obovatum. DC. (wh.) obovate-leaved.	Mauritius.	1822.	—	S. h.	
2 paniculatum. DC. (wh.) paniced.	—	1821.	4. 6.	S. h.	
Eugénia paniculata. Lam.	—	—	—	—	
3 glomeratum. DC. (wh.) cluster-flowered.	—	1824.	—	S. h.	
4 Jambolana. DC. (wh.) Jambolana-tree.	E. Indies.	1796.	5. 7.	S. h.	Rumph. amb. 1. t. 42.
Calyptránthes Jambolana. w. Eugénia Jambolana.	Lam.	—	—	—	
5 inophyllum. DC. (wh.) equal-leaved.	E. Indies.	1826.	—	S. h.	

6	caryophyllifolium. DC. (wh.)	Clove-leav'd. E. Indies.	1822.	5. 7.	S. h.	Rumph. amb. 1. t. 41.
	Eugénia caryophyllia. Lam.	Calyptráneas caryophyllifolia. w.				
7	venosum. DC.	(wh.) veined-leaved. Nepaul.	1824.	S. h.	
8	fruticosum. DC.	(wh.) shrubby. E. Indies.	—	S. h.	
9	zeylanicum. DC.	(wh.) Ceylon.	1798.	6. 7.	S. h.	Herm. zeyl. 435. ic.
CARYOPHYLLUS. DC.	CLOVE-TREE.	Icosandria Monogynia. L.				
aromaticus. DC.	(wh.) aromatic.	Moluccas.	1797.	S. h.	Bot. mag. t. 2749-2750.
ACM'ENA. DC.	ACM'EMA.	Icosandria Mónogynia.				
elliptica. DC.	(wh.) elliptic-leaved.	N. S. W.	1790.	5. 9.	G. h.	Bot. mag. t. 1872.
Eugénia elliptica. B.M.	Myrtus Smithii. s.s.	Metrosidéros floribunda. Vent.	malm. t. 75.			
EUGÉNIA. DC.	EUG'ENIA.	Icosandria Monogynia.				
1	Michelii. DC.	(wh.) one-flowered. Brazil.	1759.	1. 3.	S. h.	Bot. mag. t. 473.
	uniflora. w. Plinìa pedunculata. B.M.					
2	ligustrina. DC.	(wh.) Privet-leaved. Caribees.	1798.	8. 9.	S. h.	
3	balsámica. DC.	(wh.) balsamic. W. Indies.	1816.	6. 7.	S. h.	Jacq. frag. t. 45. f. 2.
4	angustifolia. DC.	(wh.) narrow-leaved. St. Domingo.	1825.	S. h.	Plum.ed Bur.t.207.f. 2.
5	obscura. DC.	(wh.) obscure. Brazil.	1824.	10. 11.	S. h.	Bot. reg. t. 1044.
	Myrtus obscura. B.R.					
6	alpina. DC.	(wh.) alpine. Jamaica.	1827.	6. 8.	S. h.	
7	latifolia. DC.	(wh.) broad-leaved. Guiana.	1793.	—	S. h.	Aubl. guian. 1. t. 199.
8	Lambertiána. DC.	(wh.) Mr. Lambert's. W. Indies.	1794.	S. h.	
9	Parkeriana. DC.	(wh.) Parker's. Trinidad.	—	S. h.	
10	Roxburghii. DC.	(wh.) Roxburgh's. E. Indies.	1798.	6. 7.	S. h.	Andr. deposit. t. 619?
	zeylanica. Roxb. non w.					
11	dísticha. DC.	(wh.) globe-berried. Jamaica.	1793.	4. 7.	S. h.	Lindl. coll. t. 19.
	Myrtus disticha. L. coll.					
12	glabrata. DC.	(wh.) smooth. W. Indies.	1825.	—	S. h.	
13	axillaris. DC.	(wh.) axillary. Jamaica.	1793.	9. 10.	S. h.	
14	floribunda. DC.	(wh.) many-flowered. Santa Cruz.	1822.	4. 6.	S. h.	
15	buxifolia. DC.	(wh.) Box-leaved. W. Indies.	—	—	S. h.	
16	baruensis. DC.	(wh.) many-flowered. —	—	—	S. h.	Jacq. ic. 3. t. 486.
17	dumosa. DC.	(wh.) bushy. —	1793.	6. 7.	S. h.	
18	biflora. DC.	(wh.) two-flowered. Jamaica.	1759.	4. 5.	S. h.	Brown jam. t. 25. f. 3.
	Myrtus biflora. w.					
19	Sieberiana. DC.	(wh.) Sieber's. Trinidad.	1824.	—	S. h.	
20	trinervia. DC.	(wh.) three-nerved. N. S. W.	1823.	G. h.	
	Myrtus trinervia. L.T.					
21	trinitatis. DC.	(wh.) Trinidad. Trinidad.	—	S. h.	
22	Grégi. DC.	(wh.) Greg's. Dominica.	1776.	6. 7.	S. h.	Gärt. fruct. 1. t. 33.
	Grégia aromática. G.	Myrtus Grégi. w.				
23	fragrans. DC.	(wh.) sweet-scented. Jamaica.	1790.	4. 5.	S. h.	Bot. mag. t. 1242.
24	virgultosa. DC.	(wh.) twiggy. W. Indies.	1787.	7. 8.	S. h.	
	Myrtus virgultosa. w.					
25	corymbosa. DC.	(wh.) corymbose. E. Indies.	1824.	S. h.	Rheed. mal. 5. t. 27.
26	polchella. H.B.	(wh.) pretty. —	1822.	S. h.	
27	glandulifera. H.B.	(wh.) glandular. —	1823.	S. h.	
28	obtusifolia. H.B.	(wh.) blunt-leaved. —	—	S. h.	
29	cerasoides. H.B.	(wh.) Cherry-fruited. —	—	S. h.	
30	Piménta. DC.	(wh.) Allspice. W. Indies.	1723.	5. 7.	S. h.	Bot. mag. t. 1236.
	Myrtus Piménta. w.					
JAMB'OSA. DC.	JAMB'OSA.	Icosandria Monogynia.				
1	vulgaris. DC.	(st.) narrow-leaved. E. Indies.	1768.	2. 7.	S. h.	Bot. mag. t. 1696.
	Eugenia Jámbo. B.M.					
2	venosa. DC.	(bh.) veined-leaved. Madagascar.	1824.	S. h.	
3	macrophylla. DC.	(wh.) large-leaved. E. Indies.	1822.	S. h.	
4	macrocarpa.	(wh.) large-fruited. —	—	S. h.	
	Eugenia macrocarpa. H.B.					
5	purpurascens. DC.	(pu.) purple-flower'd. W. Indies.	1768.	5. 8.	S. h.	Andr. deposit. t. 458.
	Eugenia malaccensis. Smith exot. bot. t. 61.					
6	malaccensis. DC.	(wh.) Malay Apple. E. Indies.	—	—	S. h.	Rheed. mal. 1. t. 18.
7	amplexicaulis. DC.	(gr.) stem-clasping. Sumatra.	1820.	3. 6.	S. h.	Bot. reg. t. 1033.
8	áquea. DC.	(ro.) watery. Moluccas.	1819.	—	S. h.	Rumph. amb. 1.t.38.f.2.
	Eugenia áquea. H.B.					
9	acuminata.	(gr.) taper-pointed. Amboyna.	1816.	5. 6.	S. h.	
	Eugenia acuminata. H.B.					
10	laurifolia. DC.	(wh.) Laurel-leaved. E. Indies.	1824.	S. h.	
11	ternifolia.	(wh.) three-leaved. —	1822.	—	S. h.	
	Eugenia ternifolia. H.B.					

- 12 australis. DC. (*wh.*) Myrtle-leaved. N. Holland. 1812. 5. 9. G. h. Bot. mag. t. 2230.
Eugènia australis. Colla hort. rip. app. t. 8. *Eugènia myrtifolia*. Bot. reg. t. 627.

Tribus IV. *BARRINGTONIEÆ*. DC. prodr. v. 3. p. 288.BARRINGTONIA. DC. BARRINGTONIA. *Monadelphia Polyandria*. L.

- 1 speciosa. DC. (*pu.wh.*) beautiful. E. Indies. 1786. S. h. Rumph. amb. 3. t. 114.
 2 racemosa. DC. (*wh.*) racemed. Malabar. 1822. S. h. Rheed. mal. 4. t. 6.
Eugènia racemosa. L.

STRAVADIUM. DC. STRAVADIUM. *Monadelphia Polyandria*.

- 1 álbium. DC. (*wh.*) white-flowered. Moluccas. 1822. S. h. Rumph. amb. 3. t. 116.
 2 rúbrum. DC. (*re.*) red-flowered. — — — S. h. — — — 3. t. 115.

Barringtònia acutángula, G. *Eugènia acutángula*. L.

- GUSTAVIA. DC. GUSTAVIA. *Monadelphia Polyandria*. L.
 1 augusta. DC. (*wh.*) large-flowered. Guiana. 1794. S. h. Aubl. guian. 1. t. 192.
 2 fastuosa. DC. (*wh.*) six-petaled. — — — 1824. S. h. — — — 1. t. 193.

Tribus V. *LECYTHIDEÆ*. DC. prodr. v. 3. p. 290.LECYTHIS. DC. LECYTHIS. *Monadelphia Polyandria*.

- 1 Ollaria. DC. (*wh.*) sessile-leaved. Brazil. 1820. S. h. Jacq. amer. t. 109.
 2 minor. DC. (*wh.*) smaller. Carthagena. 1823. S. h. Au.gu.2.t.283.284.285.
 3 grandiflora. DC. (*ro.*) large-flowered. Cayenne. 1822. S. h. — — 2. t. 289.
 4 Idatimon. DC. (*fl.*) flesh-coloured. Guiana. — — — S. h. — — 2.t.286et.t.285.f.1.
 5 amara. DC. (*ye.*) bitter-seeded. — — — S. h. — — 2. t. 287.
 6 parviflora. DC. (*st.*) small-flowered. — — — S. h. — — 2. t. 287.

BERTHOLLETIA. DC. BERTHOLLETIA. *Monadelphia Polyandria*.

- excelsa. DC. (*st.*) lofty. Brazil. 1824. S. h. H. et B. pl. equ. 1. t. 36.

COUROUPITA. DC. COUROUPITA. *Monadelphia Polyandria*.
guianensis. DC. (*bh.*) Guiana. Guiana. 1820. S. h. Aubl. guian. 2. t. 282.
Lecythis bracteata. W. Pèkea *Couroupita*. Juss.

MYRTACEÆ DUBIAE. DC. prodr. v. 3. p. 394.

CAREYA. R.C. CAREYA. *Monadelphia Polyandria*.

- 1 sphærica. H.B. (*wh.*) spherical. E. Indies. 1823. S. h. Roxb. corom. 3. t. 218.
 2 arborea. R.C. (*gr.bh.*) tree. — — — S. h. — — — 3. t. 217.

GRIAS. DC. ANCHOVY PEAR. *Polyandria Monogynia*.

- cauliflora. DC. (*wh.*) stem-flowering. Jamaica. S. h. Sloan.his.2.t.217.f.1.2.

MARLEA. R.C. MARLEA. *Icosandria Monogynia*.
begoniifolia. R.C. (*wh.*) Begonia-leaved. E. Indies. 1824. S. h. Roxb. corom. 3. t. 283.

ORDO XCV.

CUCURBITACEÆ. DC. prodr. v. 3. p. 297.

Tribus II. *CUCURBITEÆ*. DC. prod. 3. p. 299.LAGENARIA. DC. GOULD. *Monœcia Monadelphia*.

- 1 vulgaria. DC. (*wh.*) bottle. India. 1597. 7. 9. H. ⊕. — — — H. ⊕. Moris. his. s. 1. t. 5. f. 1.
 a *Gourd*. DC. (*wh.*) common. — — — — — H. ⊕. — — — H. ⊕. Moris. his. s. 1. t. 5. f. 1.
 β *Cougourdy*. DC. (*wh.*) Cow-gourd. China. — — — — — H. ⊕. — — — H. ⊕. Rumph. amb. 5. t. 144.
 γ *depressa*. DC. (*wh.*) flat-fruited. India. — — — — — H. ⊕. — — — H. ⊕. — — — H. ⊕. Rumph. amb. 5. t. 144.
 δ *turbinata*. DC. (*wh.*) top-fruited. — — — — — H. ⊕. — — — H. ⊕. — — — H. ⊕. Moris. his. s. 1. t. 5. f. 2.
 ε *clarata*. DC. (*wh.*) club-fruited. — — — — — H. ⊕. — — — H. ⊕. — — — sect. 1. t. 5. f. 3.

CUCURBITACEÆ.

2 vittata, DC.	(wh.) ribbon.	E. Indies.	7. 9. H. ⊙.~.	
3 idolátrica, DC.	(wh.) Pear-fruited.	H. ⊙.~.	
CUCUMIS. DC.	CUCUMBER.	Monœcia Monadelphia. L.			
1 Mélö. DC.	(st.) Melon.	Tartary.	1570.	5. 9. F. ⊙.	Blackw. herb. t. 329.
2 deliciosus. DC.	(st.) delicious.	E. Indies.	1818.	— F. ⊙.	
3 sativus. DC.	(ye.) common.	1573.	7. 9. H. ⊙.	Blackw. herb. t. 4.
α viridis. DC.	(ye.) green-fruited.	—	— H. ⊙.	
β flavus. DC.	(ye.) pale-fruited.	—	— H. ⊙.	
γ álbis. DC.	(ye.) white Turkey.	—	— H. ⊙.	
δ variegatus. DC.	(ye.) striped-fruited.	—	— H. ⊙.	
ε fastigiatus. DC.	(ye.) clustered.	—	— H. ⊙.	
4 flexuösus. DC.	(ye.) flexuous snake.	1597.	5. 9. F. ⊙.	Lob. stirp. p. 363. f. 2.
β reflexus. DC.	(ye.) angular-leaved.	—	— F. ⊙.	
5 anguinus. W.	(ye.) snake-fruited.	1820.	7. 9. F. ⊙.	
6 jamaicensis. DC.	(ye.) round-fruited.	Jamaica.	1818.	— F. ⊙.	
7 Châte. DC.	(ye.) hairy.	Levant.	1759.	6. 7. H. ⊙.	Alp. ægypt. t. 117.
8 Dudaim. DC.	(ye.) Apple-fruited.	1705.	7. 8. H. ⊙.	Andr. reposit. t. 548.
9 prophetarum. DC.	(ye.) globe-fruited.	1777.	6. 9. F. ⊙.	Jacq. vind. 1. t. 9.
10 africanus. DC.	(ye.) African.	C. B. S.	1816.	— G. 4.~.	Herm. parad. t. 134.
11 Anguria. DC.	(ye.) round prickly.	Jamaica.	1692.	7. 9. H. ⊙.	Mill. ic. 1. t. 33.
12 Citrullus. DC.	(ye.) Water Melon.	Africa.	1597.	5. 9. F. ⊙.	Rumph. amb. 5. t. 146.
α Pásteca. DC.	(ye.) firm-fruited.	—	— F. ⊙.	
β Jace. DC.	(ye.) juicy-fruited.	—	— F. ⊙.	
13 pubescens. DC.	(ye.) pubescent.	1815.	6. 9. F. ⊙.	Ser. dis. t. 3.
14 maculatus. DC.	(ye.) spotted-fruited.	—	— F. ⊙.	Blackw. herb. t. 441.
15 Cocoly'nthis. DC.	(ye.) bitter.	Japan.	1551.	5. 8. F. ⊙.	
16 perennis. DC.	(ye.) perennial.	N. America.	1824.	6. 8. F. ⊙.	
17 coronarius.	(ye.) crowned.	1827.	— F. ⊙.	
18 Momordica. H.B.	(ye.) Momordica-like.	E. Indies.	1812.	7. 9. F. ⊙.	
19 utilissimus. H.B.	(ye.) useful.	1820.	— F. ⊙.	
20 turbinatus. H.B.	(ye.) top-shaped.	—	— F. ⊙.	
21 maderaspatanus. W. (ye.) Madras.	1805.	7. 8. F. ⊙.	Pluk. alm. t. 170. f. 2.
Bryonia maderaspatana. DC.					
22 integrifolius. H.B. (ye.) entire-leaved.	1825.	— F. ⊙.~.	
LUFFA. DC.	LUFFA.	Pentandria Monogynia.			
1 foetida. DC.	(ye.) strong-scented.	E. Indies.	1812.	6. 10. F. ⊙.~.	Bot. mag. t. 1638.
2 acutangula. DC.	(ye.) acute-angled.	1692.	6. 9. F. ⊙.~.	Jacq. vind. 3. t. 73-74.
Cucumis acutangulus. Jacq.					
3 Plukenetiæna. DC.	(ye.) Plukenet's.	—	— F. ⊙.~.	Pluk. alm. t. 172. f. 1.
4 clavata. H.B.	(ye.) erect.	1820.	6. 10. F. ⊙.	
5 amara. H.B.	(ye.) bitter.	—	— F. ⊙.~.	
6 graveolens. H.B.	(ye.) strong-smelling.	1823.	— F. ⊙.~.	
7 racemosa. H.B.	(ye.) racemed.	—	— F. ⊙.~.	
8 egyptiaca. DC.	(ye.) Egyptian.	Egypt.	1739.	7. 8. F. ⊙.	Alp. ægyp. t. 58.
Momordica Luffa. L.					
BENINCASA. DC.	BENINCASA.	Polygamia Monœcia.			
cerifera. DC.	(ye.) musky.	E. Indies.	7. 9. F. ⊙.	Savi. mem. c. 1818. p. 6.
BRYONIA. DC.	BRYONY.	Monœcia Monadelphia. W.			
1 rostrata. DC.	(st.) rostrate.	E. Indies.	1825.	7. 9. S. ⊙.~.	
2 scábra. DC.	(st.) globe-fruited.	C. B. S.	1774.	9. 10. G. 4.~.	
3 verrucosa. DC.	(st.) warted.	Canaries.	1779.	6. 10. G. 4.~.	
4 scabratæ. DC.	(st.) roughish.	E. Indies.	1826.	— S. 4.~.	
5 punctatæ. DC.	(st.) dotted.	C. B. S.	1816.	— G. 4.~.	
6 cordatæ. DC.	(st.) cordate.	—	— G. 4.~.	
7 grändis. DC.	(re.) great-flowered.	E. Indies.	1783.	5. 8. S. 4.~.	Rump. amb. 5. t. 166. f. 1.
8 umbellata. DC.	(st.) umbel-flowered.	1825.	— S. 4.~.	Rheed. mal. 8. t. 26?
9 hederifolia. DC.	(st.) Ivy-leaved.	Teneriffe.	—	— G. 4.~.	Jacq. fragm. 73. t. 113.
10 epigæa. DC.	(st.) leathery-leaved.	E. Indies.	1815.	— G. 4.~.	
11 scabrélla. DC.	(st.) rough-seeded.	1781.	6. 8. S. ⊙.~.	
12 latebrósa. DC.	(st.) hairy.	Canaries.	1779.	— G. 4.~.	
13 trilobata. DC.	(st.) three-lobed.	C. B. S.	1816.	— G. ⊙.~.	
14 álba. DC.	(wh.) black-fruited.	Europe.	1807.	— H. 4.~.	Lam. ill. t. 796.
15 dioica. DC.	(st.) red-fruited.	Britain.	5. 9. H. 4.~.	Eng. bot. v. 7. t. 439.
16 nítida. DC.	(st.) glossy.	1824.	— S. 4.~.	
17 crética. DC.	(st.) Cretan.	Candia.	1759.	7. 9. G. 4.~.	Desf. an. mus. 12. t. 17.
18 quinqueloba. DC.	(st.) five-lobed.	C. B. S.	6. 10. G. 4.~.	Bot. reg. t. 82.
19 ficifolia. DC.	(st.) Fig-leaved.	Buenos Ayres.	1726.	— G. 4.~.	Dill. elth. t. 50. f. 58.
20 variegata. DC.	(st.) variegated.	America.	— S. 4.~.	
21 racemosa. DC.	(st.) racemed.	Trinidad.	1820.	— S. 4.~.	Plum. amer. 83. t. 97.

22 pinnatífida. DC.	(st.) pinnatifid.	C. B. S.	1816.	7. 8. G. ¼.~.
23 divisa. W. en.	(st.) divided-leaved.	1818.	6. 10. G. ¼.~.
24 palmata. DC.	(st.) palmated.	Ceylon.	1778.	7. 8. S. ¼.~.
25 Garcini. DC.	(st.) round-lobed.	1818.	— S. ¼.~. Burm. ind. t. 57. f. 3.
26 laciñiosa. DC.	(st.) laciniated.	E. Indies.	1710.	— S. ¼.~. Herm. lugd. 95. t. 97.
27 africana. DC.	(st.) African.	C. B. S.	1759.	7. 8. G. ¼.~. Herm. parad. t. 708.
28 dissecta. DC.	(st.) smooth-leaved.	1710.	— G. ¼.~.
29 tenella. H.B.	(st.) slender.	E. Indies.	1820.	— S. ¼.~.
30 Wallichiana. DC.	(st.) filiform.	Nepaul.	—	— S. ¼.~.

filiformis. H.B.

SI'CYOS. DC.	SINGLE-SEEDED CUCUMBER.	Monocotia Monadelphia. W.
1 angulatus. DC.	(ye.) angular-leaved.	N. America. 1710. 7. 9. H. ¼.~. Dill. elth. t. 51. f. 59.
2 parviflòrs. DC.	(ye.) small-flowered.	Quito. 1823. — H. ¼.~.
3 microphyllus. DC.	(ye.) small-leaved.	Mexico. — — H. ¼.~.
4 vitifolius. DC.	(ye.) Vine-leaved. — H. ¼.~.
5 laciniatus. DC.	(ye.) jagged-leaved.	S. America. 1827. — F. ¼.~. Plum. ed Burm. t. 243.

ELAT'ERIUM. DC.	ELAT'ERIUM.	Monocotia Monadelphia.
1 gemellum. DC.	(wh.) twin-tendrilled.	Mexico. 1826. 6. 9. H. ¼.~.
2 carthaginense. DC.	(wh.) Carthaginian.	Caracas. 1824. — F. ¼.~. Jacq. am. pict. t. 232.
3 tamnoïdes. DC.	(st.) Tamus-like.	Mexico. 1818. — H. ¼.~.

MOMO'R'DICA. DC.	MOMO'R'DICA.	Monocotia Monadelphia. W.
1 balsamina. DC.	(ye.) Balsam Apple.	E. Indies. 1568. 6. 7. F. ¼.~. Blackw. herb. 6. t. 539.
2 muricata. DC.	(ye.) tubercled. — F. ¼.~. Rheed. mal. 8. t. 10.
3 Charantia. DC.	(ye.) hairy. 1710. — F. ¼.~. Bot. mag. t. 2455.
β abbreviata. DC.	(ye.) short-fruited. — F. ¼.~.

4 cylindrica. DC.	(ye.) cylindrical.	China. 1816. — F. ¼.~.
5 operculata. DC.	(ye.) rough-fruited.	W. Indies. 1731. 6. 9. F. ¼.~. Comm. rar. t. 22.
6 Elaterium. DC.	(st.) squirting cucumber.	S. Europe. 1548. — H. ¼.~. Bot. mag. t. 1914.
7 dioica. DC.	(ye.) dioecious.	E. Indies. 1820. 7. 8. S. ¼.~. Rheed. mal. 8. t. 18.
8 monadelpha. H.B.	(ye.) monadelphous. 1812. — S. ¼.~.
9 mixta. H.B.	(ye.) mixed. 1820. — S. ¼.~.
10 umbellata. H.B.	(ye.) umbel-flowered. 1818. — S. ¼.~.
11 tubiflora. H.B.	(ye.) tube-flowered. 1824. 7. 10. F. ¼.~.

S'ECHIUM. DC.	СНОЧО.	Monocotia Monadelphia.
1 édule. DC.	(ye.) eatable.	W. Indies. 1823. 7. 9. F. ¼.~. Jacq. amer. t. 163.
2 americanum. DC.	(ye.) American.	Jamaica. 1825. — F. ¼.~.
3 palmatum. DC.	(st.) palmated.	Mexico. 1827. 9. 11. F. ¼.~. Moc. et Ses. fl. mex. ic.

MELO'THRIA. DC.	MELO'THRIA.	Monocotia Triandria.
1 pénula. DC.	(ye.) pendulous.	America. 1752. 6. 7. H. ¼.~. Plum. ic. t. 66. f. 2.
2 foetida. DC.	(ye.) strong-scented.	Guinea. 1825. — S. ¼.~. Jacq. ic. 3. t. 624.

Trichosanthes sativissima. Jacq.

TRICHOSA'NTHES. DC.	SNAKE-GOURD.	Monocotia Monadelphia. W.
1 anguina. DC.	(wh.) common.	China. 1755. 5. 8. F. ¼.~. Bot. mag. t. 722.
2 colubrina. DC.	(wh.) panicled. 1820. — F. ¼.~. Jacq. f. eclog. t. 128.
3 dioica. H.B.	(wh.) dioecious.	E. Indies. — 5. 6. S. ¼.~.
4 cucumerina. DC.	(wh.) Cucumber-like. 1804. 6. 8. F. ¼.~. Rheed. mal. 8. t. 15.
5 lobata. H.B.	(wh.) lobed-leaved. 1812. — F. ¼.~.
6 cordata. H.B.	(wh.) heart-leaved. 1823. — S. ¼.~.
7 palmata. H.B.	(wh.) palmated. 1825. — S. ½.~.
8 corniculata. DC.	(wh.) horn-gourd.	W. Indies. 1810. 6. 7. S. ¼.~. Bot. mag. t. 2703.
<i>tuberosa. B.M.</i>	<i>Ceratosanthes tuberosa. s.s.</i>	

JOLI'FFIA. DC.	JOLI'FFIA.	Dicecia Monadelphia.
africana. DC.	(pu.) African.	Zanzibar. 1825. 6. 8. S. ½.~. Bot. mag. t. 2681.
♀ fa'mina.	(pu.) female.	— — — S. ½.~. — t. 2681.

*Feuillæa pedata. B.M.**♂ máscula. (pu.) male.**Telfaria pedata. Hook. bot. mag. t. 2751.*

CUCU'RBITA. DC.	PUMPKIN.	Monocotia Monadelphia. W.
1 máxima. DC.	(ye.) Elephant. 7. 10. H. ¼.~. Tourn. inst. n. 2. t. 34.
α Potiro. DC.	(ye.) yellow-fruited. — H. ¼.~.
β viridis. DC.	(ye.) green-fruited. — H. ¼.~.
γ Courgero. DC.	(ye.) small-fruited. — H. ¼.~.
2 Melopepo. DC.	(ye.) Squash. 1597. 5. 9. H. ¼.~. Moris. hist. 1. s. 1. t. 8.
3 moschata. DC.	(ye.) musk Melon.	W. Indies. — 6. 9. F. ¼.~. Dalech. hist. 616. f. 3.
4 villosa. DC.	(ye.) woolly-fruited.	E. Indies. 1818. — F. ¼.~.
5 Pèpo. DC.	(ye.) common.	Levant. 1570. — H. ¼.~.
α subrotunda. DC.	(ye.) round-fruited.	— — — H. ¼.~.
β oblonga. DC.	(ye.) oblong-fruited.	— — — H. ¼.~.
6 farinosa. DC.	(ye.) mealy-fruited.	E. Indies. 1822. — F. ¼.~.

7 verrucosa. DC. (ye.) warted.	1658.	6. 9. H. ○.~.	Bauh. hist. 2. p. 222. ic.
8 subverrucosa. DC. (ye.) oblong-fruited.	— H. ○.~.	
9 aurantia. DC. (ye.) orange-gourd.	— H. ○.~.	
α orangina. DC. (ye.) true orange.	— H. ○.~.	
β colocynthoides. DC. (ye.) variegated-fruit'd.	— H. ○.~.	
10 ovifera. DC. (ye.) egg-shaped. Astracan.	— H. ○.~.	Dod. pempt. 670. f. 1.
α pyriformis. DC. (ye.) pear-fruited.	— H. ○.~.	Ser. diss. t. 1.
β subglobosa. DC. (ye.) white-striped.	— H. ○.~.	
γ grisea. DC. (ye.) white-spotted.	— H. ○.~.	
11 foetidissima. DC. (ye.) strong-scented. Mexico. 1825.	— H. ○.~.	
12 marsupialis. Jacq. (ye.) bag-bearing.	1821.	— H. ○.~.	
13 dipsacea. E. (ye.) Teasel-like. Egypt. 1827.	— H. ○.~.	
ANG'URIA. DC. ANG'URIA. Monoclea Diandria. W.			
1 trilobata. DC. (re.) three-lobed. Martinico. 1793.	6. 7. S. ¼.~.	Jacq. am. pict. t. 234.	
2 pedatisecta. DC. (re.) cut-leaved. Peru. 1827.	7. 10. H. ○.~.	Feuil per. 1. t. 41.	
3 pedata. DC. (re.) pedate-leaved. St. Domingo. —	6. 8. S. ¼.~.	Jacq. am. pict. t. 155.	

GENERA CUCURBITACEIS AFFINIA.

GRON'OVIA. DC.	GRON'OVIA. Pentandria Monogynia. L.
scándens. DC.	(st.) climbing. Jamaica. 1731. 6. 7. S. ¼.~.
CA'RICA. S.S.	PAPAW-TREE. Diocia Decandria. L.
1 Papáya. s.s.	(st.) common. India. 1690. 7. 8. S. h. Bot. reg. t. 459.
2 pyrifórmis. s.s.	(ro.) Pear-fruited. Peru. 1825. S. h. Feuil. journ. 3. t. 39.
3 citrifórmis. s.s.	(st.) Citron-fruited. Guinéa. — S. h.
4 caulinflóra. s.s.	(st.) stem-flowering. Caracas. 1806. 6. 8. S. h. Jacq. schoenb. 3. t. 311.
5 microcarpa. s.s.	(st.) small-fruited. — — — S. h. — 3. t. 309. 310.
6 monoíca. Df.	(st.) monœcious. — 1816. — S. h.
7 digitáta. s.s.	(st.) spiny. Guiana. 1822. S. h. Aubl. guian. 2. t. 246.
spínosa. W.	

ORDO XCVI.

PASSIFLOREÆ. DC. prodr. v. 3. p. 321.

Tribus I. PAROPSIEÆ. DC. prodr. v. 3. p. 322.

SMEA'THMA'NNIA. L.T.	SMEA'THMA'NNIA. Polyandria Pentagynia.
lævigàta. L.T. (wh.) smooth.	Sierra Leone. 1822. S. h.

Tribus II. PASSIFLOREÆ VERÆ. DC. prodr. v. 3. p. 322.

PASSIFL'ORA. DC.	PASSION-FLOWER. Monadelphia Pentandria.
1 holosericea. DC.	(st.) silky-leaved. Vera Cruz. 1733. 5. 8. S. h.~. Bot. reg. t. 59.
2 pálida. DC.	(gr.) pale-flowered. W. Indies. 1820. 5. 10. S. h.~. — t. 660.
3 cùprea. DC.	(co.) copper-colour'd. Bahama Isl. 1724. 7. 8. S. h.~. Jacq. ic. 3. t. 606.
4 multiflóra. DC.	(wh.) many-flowered. Vera Cruz. 1731. 6. 9. S. h.~. Cavan. diss. 10. t. 272.
5 pubescens. DC.	(wh.) pubescent. Caracas. 1825. — S. h.~.
6 mexicana. DC.	(wh.) Mexican. Mexico. 1812. — S. h.~. Jus. an. mus. 6. t. 38. f. 2.
7 normális. DC.	(cr.) linear-lobed. Jamaica. 1771. 5. 7. S. h.~.
8 angustifòlia. DC.	(gr.) narrow-leaved. W. Indies. 1773. 6. 9. S. h.~. Bot. reg. t. 188.
heterophylla. Jacq. schen. 2. t. 181.	
9 maculata. DC.	(wh.) painted-leaved. N. America. 1812. — G. h.~. Pluk. t. 210. f. 3.
10 grácidis. DC.	(wh.) slender. — 1823. 6. 10. S. ○.~. Bot. reg. t. 870.
11 lutéa. DC.	(st.) pale yellow. — 1714. 5. 8. F. ¼.~. — t. 79.
12 mímina. DC.	(gr.) small. S. America. 1690. 7. 9. S. h.~. — t. 144.
13 hirsúta. DC.	(gr.) hairy-stalked. W. Indies. 1778. 8. 10. S. h.~. Plum. amer. t. 88.
14 suberósa. DC.	(wh.) cork-barked. — 1759. 6. 9. S. h.~. Sm. exot. bot. 1. t. 28.

15	peltáta. DC.	(st.) peltate.	W. Indies.	1778.	8. 10.	S. h.~, Bot. reg. t. 507.
16	hederácea. DC.	(wh.) Ivy-leaved.	Antilles.	6. 7.	S. h.~, Plum. amer. t. 84.
17	pannósa. DC.	(fl.) cloth-leaved.	W. Indies.	1824.	6. 9.	S. h.~.
18	cuneáta. W.en.	(wh.) wedge-leaved.	S. America.	1820.	6. 10.	S. h.~.
19	multifórmis. DC.	(wh.) multiform.	Caracas.	1825.	—	S. h.~, Jacq. frag. t. 67. f. 1. caracásina. W.en.
20	heterophylla. DC.	(wh.) various-leaved.	St. Domingo.	1817.	—	S. h.~, Plum.ed.Bur.t.139.f.1.
21	rubricaulis. Jac.	(re) red-stalked.	S. America.	1821.	6. 8.	S. h.~.
22	perfoliáta. DC.	(pu.) perfoliate.	W. Indies.	1806.	7. 9.	S. h.~, Bot. reg. t. 78.
23	rúbra. DC.	(fl.) red-fruited.	—	1731.	4. 9.	S. h.~, — t. 95.
24	capsuláris. DC.	(wh.) angular-fruited.	—	1826.	—	S. h.~, Bot. mag. t. 2868?
25	biflóra. DC.	(wh.) crescent-leaved.	—	1733.	—	S. h.~, Sm. ic. pict. 1. t. 1. lunata. Sm. non Juss. <i>Vespertilio</i> . Miss Lawr. pass. t. 8. nec aliorum.
26	β? acutilobá. (wh.) acute-lobed.	W. Indies.	—	—	—	S. h.~, Bot. reg. t. 577.
27	Vespertilio. DC.	(wh.) bat-winged.	—	1732.	5. 6.	S. h.~, Dill. elth. t. 137. f. 164.
28	Maximiliána. DC.	(wh.) two-coloured.	Brazil.	1820.	6. 10.	S. h.~, Bot. reg. t. 597.
29	Vespertilio. B.R. nec aliorum. discolor.	Lodd. bot. cab. t. 565.	—	—	—	—
30	oblongáta. DC.	(wh.) oblong-leaved.	Jamaica.	1822.	—	S. h.~.
31	tuberósa. DC.	(pa.) tuberous-root'd.	S. America.	1784.	—	S. h.~, Bot. reg. t. 432. punctáta. Lodd. bot. cab. t. 101.
32	rotundifolia. DC.	(wh.) round-leaved.	W. Indies.	1779.	5. 8.	S. h.~, Cavan. dis. t. 290.
33	punctáta. DC.	(st.) dotted-leaved.	Peru.	1784.	5. 6.	S. h.~, — 6. t. 269.
34	serratifolia. DC.	(pu.) notch-leaved.	W. Indies.	1731.	5. 10.	S. h.~, Bot. mag. t. 651.
35	coccínea. DC.	(sc.) scarlet-flower'd.	Guiana.	1823.	—	S. h.~, Aubl. guian. 3. t. 324.
36	malucináta. DC.	(v.) mucronate.	Brazil.	1822.	—	S. h.~, Cavan. diss. t. 282.
37	malifórmis. DC.	(v.) Apple-fruited.	W. Indies.	1731.	7. 11.	S. h.~, Bot. reg. t. 94.
38	tiliácea. DC.	(wh.) Lime-tree-ld.	Peru.	1822.	—	S. h.~, Feuill. obs. 2. t. 12.
39	ligularis. DC.	(v.) taper-pointed.	—	—	4. 8.	S. h.~, Juss. an. mus. 6. t. 40.
40	quadranguláris. DC.	(v.) square-stalked.	Jamaica.	1768.	8. 10.	S. h.~, Bot. reg. t. 14.
41	alátia. DC.	(v.) wing-stalked.	W. Indies.	1772.	4. 8.	S. h.~, Bot. mag. t. 66.
42	álbita. DC.	(wh.) white-flowered.	Brazil.	1816.	7. 11.	S. h.~, Bot. reg. t. 677.
43	anguláta. Cerv.	(wh.) angular-stalked.	Mexico.	1823.	—	G. h.~.
44	laurifolia. DC.	(v.) Laurel-leaved.	W. Indies.	1690.	6. 8.	S. h.~, Bot. reg. t. 13.
45	tinifolia. DC.	(v.) Tinus-leaved.	—	1822.	—	S. h.~, Juss. an. mus. 6.t.41.f.2.
46	racemosa. DC.	(sc.) racemed.	Brazil.	1816.	1. 12.	S. h.~, Bot. reg. t. 285.
47	sanguínea. DC.	(cr.) crimson.	Hybrid.	1820.	—	G. h.~.
48	cæruleo-racemosa. (pu.) Whitley's mule.	—	—	—	—	G. h.~, Lodd. bot. cab. t. 573.
49	α quinquelobata. H.T. (pu.) five-lobed.	—	—	—	—	G. h.~, Hort. trans. 4. t. 9.
50	β trilobata. H.T. (pu.) three-lobed.	—	—	—	—	G. h.~.
51	γ racemifera. (pu.) raceme-bearing.	—	—	—	—	G. h.~.
52	alátio-cærulea. DC.	(ro.) Masters's mule.	Hybrid.	1823.	4. 8.	G. h.~, Bot. reg. t. 848.
53	stipuláta. DC.	(wh.) glaucous-leaved.	Cayenne.	1779.	8. 9.	S. h.~, Aubl. guian. 2. t. 325. glaúca. Bot. reg. t. 88. non Humb.
54	incarnáta. DC.	(bh.) flesh-coloured.	N. America.	1629.	7. 9.	H. h.~, Miss Lawr. pass. ic.
55	β? integrilobata. DC.	(bh.) entire-lobed.	—	1812.	—	H. h.~, Bot. reg. t. 332.
56	culis. DC.	(bh.) eatable-fruited.	Brazil.	1816.	6. 10.	G. h.~, Bot. mag. t. 1989. incarnáta. β. Bot. reg. t. 152.
57	vitifolia. DC.	(wh.) Vine-leaved.	S. America.	1824.	—	S. h.~.
58	filamentosa. DC.	(v.) filamentose.	—	1819.	7. 10.	G. h.~, Cavan. diss. 10. t. 294.
59	palmáta. DC.	(v.) palmate-leaved.	Brazil.	—	—	F. h.~, Lodd. bot. cab. t. 97.
60	Colvillia. DC.	(v.) Colvill's mule.	Hybrid.	1823.	—	H. h.~, Swt. br. fl. gar. t. 126.
61	cærulea. DC.	(wh. bl.) common.	Brazil.	1699.	—	H. h.~, Bot. reg. t. 488. β glaucocephala. (wh. bl.) glaucous-leaved.
62	γ angustifolia. (wh. bl.) narrow long-l'd.	—	—	—	—	H. h.~.
63	chinénsis. (wh. bl.) Chinese.	China.	1819.	6. 10.	G. h.~.	
64	cærulea Lour. nec aliorum.	—	—	—	—	—
65	serrata. DC.	(pa.) saw-leaved.	Martinica.	1800.	—	S. h.~, Cavan. diss. 10. t. 296.
66	digitáta. R.P.	(pa.) digitate.	Peru.	1824.	—	S. h.~.
67	pedáta. DC.	(pa.) curl-flowered.	W. Indies.	1781.	—	S. h.~, Plum. am. t. 81.
68	picturáta. DC.	(v.) pictured.	Brazil.	1823.	6. 9.	S. h.~, Bot. reg. t. 673.
69	hibiscifolia. (wh. bl.) Hibiscus-leav'd.	Caribees.	1825.	—	S. h.~, Plum. am. t. 86.	
70	fœtida.	(wh. bl.) stinking.	W. Indies.	1731.	7. 9.	S. h.~, Bot. mag. t. 2619.
71	hircina. ed. 1. (wh. bl.) strong-scented.	Brazil.	—	5. 10.	S. h.~, Bot. reg. t. 321. fætida. Bot. reg. nec aliorum. hirsuta. Bot. cab. 138. nec aliorum.	
72	ciliáta. DC.	(wh. ro.) ciliated.	Jamaica.	1783.	7. 9.	S. h.~, Bot. mag. t. 288. DISE'MMA. DC.
73	DISE'MMA. Monadelphia Pentandria.	—	—	1822.	6. 10.	G. h.~, Bot. reg. t. 737. Herbertiána. DC. (st.) L'd.Caernarvon's.N.Holland.
74	Passiflóra Herbertiána. B.R.	—	—	1792.	—	G. h.~, Bot. reg. t. 233.
75	adianthifolia. DC. (or.) Adiantum-l'd.	Norfolk Isl.	—	—	—	Passiflóra aurántia. Andr. reposit. t. 295. Passiflóra glábra. Wendl. coll. 1. t. 17.

PASSIFLOREÆ.

MURUC'UIA. DC.	MURUC'UIA. Monadelphia Pentandria.
ocellata. DC.	(cr.) ocellated. W. Indies. 1739. 7. 8. S. h. — Bot. reg. t. 574.
<i>Passiflora Murucuja.</i> B.R. Lin. am. 1. t. 10. f. 10.	
TAC'SONIA. DC.	TAC'SONIA. Monadelphia Pentandria.
1 pinnatistipula. DC. (ro.) feather-stipuled. Chile. 1826. S. h. — Cavan. ic. 5. t. 428.	
2 peduncularis. DC. (ro.) long-peduncled. Peru. 1815. S. h. — — — 5. t. 426.	
<i>Passiflora peduncularis.</i> Cav.	
MODECCA. DC.	MODECCA. Dicecia Monadelphia.
1 lobata. DC.	(gr.) lobe-leaved. Sierra Leone. 1816. 6. 9. S. h. — Bot. reg. t. 433.
2 trilobata. H.B.	(gr.) three-lobed. E. Indies. 1823. — S. h. — Roxb. corom. 3. ic.
3 tuberosa. H.B.	(gr.) tuberous-rooted. 1822. — S. h. — Rumph. amb. 8. t. 20.

ORDO XCVII.

LOASEÆ. DC. prodr. v. 3. p. 339.

BARTONIA. DC.	BART'ONIA. Polyandria Monogynia.
1 ornata. DC.	(wh.) naked-seeded. Missouri. 1811. 7. 9. F. ♂. Bot. mag. t. 1487.
decapetala. B.M.	
2 nuda. DC.	(wh.) winged-seeded. — — — F. ♀.
BLUMENBA'CHIA. DC.	BLUMENBA'CHIA. Polyandria Monogynia.
insignis. DC.	(wh.) elegant. Chile. 1826. 7. 10. H. Ⓢ. Swt. br. fl. gar. t. 171.
<i>Loasa palmata.</i> Treviranus. pátula. Gr.	
SCYPHANTHUS. B.F.G.	CUP-FLOWER. Polyandria Monogynia.
élégans. B.F.G.	(ye.) elegant. Chile. 1824. 5. 9. F. h. — Swt. br. fl. gar. t. 238.
LOASA. DC.	LO ASA. Polyandria Monogynia.
1 bryoniaefolia. DC.	(ye.) Bryony-leaved. Chile. 1828. 6. 9. H. Ⓢ. Schrad. pl. rar. græt. ic.
2 tricolor. DC.	(ye.) three-coloured. 1822. 8. 10. H. Ⓢ. Bot. reg. t. 667.
3 nítida. DC.	(ye.) glossy-leaved. Peru. — 7. 10. H. Ⓢ. Hook. exot. bot. t. 83.
4 Placei. H.T.	(ye.) Place's. Chile. — — H. Ⓢ. Bot. reg. t. 785.
acanthifolia. B.R. non Lamarck.	
5 grandiflora. DC.	(ye.) large-flowered. Peru. 1825. — H. Ⓢ. Juss. an. mus. 5. t. 4. f. 2.
MENTZ'ELIA. DC.	MENTZ'ELIA. Polyandria Monogynia.
1 áspera. DC.	(or.) rough. W. Indies. 1773. 7. 9. S. Ⓢ. Plum. ed. Bur. t. 174. f. 1.
2 oligosperma. DC.	(or.) few-seeded. Louisiana. 1812. 5. 7. F. ♀. Bot. mag. t. 1760.
3 hispida. DC.	(or.) hispid. Mexico. 1824. 6. 9. F. ♀. Cavan. ic. 1. t. 70.

ORDO XCVIII.

TURNERACEÆ. DC. prodr. v. 3. p. 345.

TURN'ERA. DC.	TURN'ERA. Pentandria Trigynia. L.
1 ulmifolia. DC.	(ye.) Elm-leaved. Jamaica. 1733. 6. 9. S. h. Linn. h. cliff. t. 10.
2 angustifolia. B.M.	(ye.) narrow-leaved. — — 4. 9. S. h. Bot. mag. t. 281.
3 cuneiformis. DC.	(ye.) wedge-leaved. Brazil. 1821. — S. h. Bot. mag. t. 2106.
4 trioniflora. DC.	(ye.) Ketmia-flow'd. Trinidad. 1812. — S. h.
élegans. R.S.	
5 acuta. DC.	(ye.) acute-leaved. Jamaica. 1826. — S. h.
6 Pumilea. DC.	(ye.) nettle-leaved. 1796. 8. 9. S. Ⓢ. Sloan. hist. t. 127. f. 6.
7 rupéstris. DC.	(ye.) rock. Guiana. 1826. 6. 9. S. h. Aubl. guian. 1. t. 113. f. 1.
8 guianensis. DC.	(ye.) Guiana. — — — S. Ⓢ. — 1. t. 114.
9 cistoides. DC.	(ye.) Cistus-like. America. 1774. 6. 10. H. Ⓢ. Plum. ic. t. 150. f. 1.
10 racemosa. DC.	(ye.) raceme-flower'd. St. Domingo. 1789. 7. 10. S. Ⓢ. Jacq. vind. 3. t. 94.
PIRIQUETA. DC.	PIRIQU'ETA. Pentandria Trigynia.
villosa. DC.	(ye.) villous. Guiana. 1826. 7. 10. S. Ⓢ. Aubl. guian. 1. t. 117.

ORDO XCIX.

PORTULACEÆ. DC. prodr. v. 3. p. 351.

TRIA'NTHEMA. DC.		TRIA'NTHEMA. Pent-Decandria Mono-Trigynia.						
1	decândra. DC.	(gr.) trailing.	E. Indies.	1762.	7. 8. S. ♂.	Burm. ind. t. 31. f. 3.		
2	triquêtra. DC.	(gr.) triquetrous.	—	1826.	— S. ♂.			
3	obcordâta. H.B.	(gr.) obcordate.	—	1820.	— S. ♂.			
4	monogyna. DC.	(gr.) Purslane-leav'd. Jamaica.	1716.	— S. ♂.	DC. plant. gras. t. 109.			
PORTULACA. DC.		PURSLANE. Oct-Dodecadria Monogynia.						
1	oleracea. H.S.	(ye.) small.	Europe.	1562.	6. 8. H. ♂.	DC. plant. gras. t. 123.		
2	sativa. H.S.	(ye.) garden.	America.	1652.	6. 9. H. ♂.			
	β aurea. H.S.	(ye.) golden.	—	—	— H. ♂.			
3	parvifolia. H.S.	(ye.) small-leaved.	Jamaica.	1799.	8. 10. H. ♂.			
4	foliosa. DC.	(ye.) leafy.	Guinea.	1822.	6. 8. S. ♀.	Bot. reg. t. 793.		
5	involucrata. H.R.	(ye.) leafy-flowered.	—	1819.	8. 9. S. ♂.	Sloan. hist. 1. t. 129. f. 3.		
6	marginata. DC.	(ye.) red-margined.	Caracas.	1822.	6. 10. H. ♂.			
7	halimoïdes. DC.	(ye.) Halimus-leaved.	Jamaica.	1823.	8. 9. S. ♂.			
8	mucronata. DC.	(ye.) mucronate.	—	1827.	6. 9. S. ♂.			
9	quadrifida. DC.	(ye.) quadrifid.	Asia.	1773.	8. 9. S. ♂.	Jacq. coll. 2. t. 17. f. 4.		
10	meridiâna. H.S.	(ye.) noon-day.	E. Indies.	1791.	5. 6. S. ♂.	Rheed. mal. 10. t. 31.		
11	grandiflora. B.M.	(ea.) large-flowered.	Chile.	1828.	6. 8. F. ♀.	Bot. mag. t. 2885.		
	α purpurea.	(pu.) purple-flowered.	—	—	— F. ♀.	— t. 2885. f. 2.		
	β aurantiaca.	(or.) orange-coloured.	—	—	— F. ♀.	— t. 2885. f. 1.		
12	pilosâ. DC.	(pu.) hairy.	W. Indies.	1690.	— S. ♂.	Bot. reg. t. 792.		
13	setacea. H.S.	(pu.) bristly.	—	—	— S. ♂.	Pluk. phyt. t. 246. f. 6.		
ANACAMPSEROS. DC.		ANACAMPSEROS. Dodecadria Monogynia.						
1	telephiâstrum. DC.	(re.) round-leaved.	C. B. S.	1732.	7. 9. D.I. ♂.	DC. plant. gras. t. 3.		
	Portulaca Anacampseros. L.	Talinum Anacampseros. W.	Rulîngia Anacampseros. H.S.					
2	arachnoides. DC.	(bh.) cobweb.	C. B. S.	1790.	7. 9. D.I. ♂.	Bot. mag. t. 1368.		
3	varians.	(re.) various-leaved.	—	1812.	— D.I. ♂.			
	Rulîngia varians. H.S.s.							
4	rûbens. DC.	(re.) red-leaved.	—	1796.	— D.I. ♂.			
5	filamentosa. DC.	(ro.) thready.	—	1795.	— D.I. ♂.	Bot. mag. t. 1367.		
6	lanceolâta. DC.	(ro.) spear-leaved.	—	1796.	— D.I. ♀.			
7	angustifolia. DC.	(ro.) narrow-leaved.	—	1819.	— D.I. ♂.			
8	rufescens. DC.	(bh.) rufescent.	—	1818.	— D.I. ♂.			
9	trigona. DC.	(bh.) trigonal-leaved.	—	—	— D.I. ♂.			
10	lanigera. DC.	(ro.) woolly.	—	—	— D.I. ♂.	Burm. afr. t. 30. f. 2.		
11	intermédia. H.P.	(ro.) intermediate.	—	1823.	— D.I. ♂.			
12	polyphylla. DC.	(ro.) many-leaved.	—	—	— D.I. ♂.			
	Rulîngia polyphylla. H.S.s.							
TALINUM. DC.		TALINUM. Dodecadria Monogynia.						
1	teretifolium. DC.	(ro.) round-leaved.	N. America.	1820.	7. 8. H. ♀.	Lodd. bot. cab. t. 819.		
2	crassifolium. DC.	(ro.) thick-leaved.	Caribees.	1800.	8. 9. S. ♂.	Jacq. vind. 3. t. 52.		
3	fruticosum. W.	(wh.) white-flowered.	S. America.	—	— S. ♂.	Comm. hort. 1. t. 4.		
4	triangulare. DC.	(ye.) triangular.	W. Indies.	1739.	— S. ♂.	Jacq. obs. 1. t. 23.		
5	pátent. DC.	(re.) spreading-fl'd.	S. America.	1776.	8. 10. S. ♂.	Jacq. vind. 2. t. 151.		
	paniculatum. Gært. fruct. 2. t. 128. Portulaca paniculatum. Jacq. amer. non R.P.							
6	reflexum. DC.	(ye.) reflex-flowered.	S. America.	1800.	8. 10. S. ♂.	Bot. mag. t. 1543.		
7	cuneifolium. DC.	(re.) wedge-leaved.	Arabia.	1823.	— S. ♂.			
8	lineâre. K.S.	(ye.) linear-leaved.	Mexico.	1827.	— H. ♂.			
9	purpureum. F.	(pu.) purple.	—	—	— H. ♂.			
CALANDRINA. DC.		CALANDRINA. Dodecadria Monogynia.						
1	paniculata. DC.	(ro.) panicled.	S. America.	1812.	8. 10. H. ♂.			
2	pilosúscula. DC.	(ro.) ciliated.	Chile.	1823.	— H. ♂.	Hook. ex. flor. 1. t. 82.		
	Talinum ciliatum. Hooker non R.P. ex DC.							
3	Phacospérma. DC.	(ro.) Peruvian.	Peru.	1827.	— G. ♂.			
4	caulescens. DC.	(ro.) caulescent.	Mexico.	—	— H. ♂.	H. et B. n. ge. am. 6. t. 526.		
5	compréssâ. DC.	(ro.) compressed.	Chile.	1826.	— H. ♂.			
	β adscendens. Otto.	(ro.) ascending	—	—	— H. ♂.			
6	glaucâ. DC.	(ro.) glaucous-leaved.	—	1827.	— D.G. ♂.			
7	grandiflora. B.R.	(ro.) large-flowered.	—	1826.	6. 10. D.G. ♂.	Bot. reg. t. 1194.		
8	Andre'wsiâ.	(ro.) Andrews's.	W. Indies.	1812.	8. 10. D.G. ♂.	Andr. rep. t. 253.		
	Talinum pátent. A.R. nec aliorum.							
9	Lockhartiâ.	(ro.) Lockhart's.	Trinidad.	1825.	6. 9. D.G. ♂.			
	Talinum grandiflorum. Lockhart.							

PORTULACEÆ.

POR TULAC'ARIA. DC.	PURSLANE-TREE. Pentandria Trigynia.
A'fra. DC.	(ro.) African. Africa. 1732. 6. 9. D.G. h.
CLAYTONIA. DC.	CLAY'TONIA. Pentandria Monogynia. L.
1 perfoliata. DC.	(wh.) small-flowered. N. America. 1794. 5. 8. H. ⊖.
<i>cubensis</i> of Bonap.an.mus.7.t.6.	is a very different species, according to G.Don, who saw it growing wild.
2 alsinoïdes. DC.	(wh.) chickweed-like. Nootka-sound. 1794. 3. 6. H. ⊖.
3 lanceolata. DC.	(wh.) spear-leaved. N. America. 1812. 3. 5. H. ♀.
4 sibirica. DC.	(ro.) Siberian. Siberia. 1768. 5. 8. H. ⊖.
5 acutifolia. DC.	(wh.) acute-leaved. — 1827. H. ♀.
6 Vestiana. F.	(ro.) Vest's. Altay. H. ♀.
7 unalaschkensis. R. S. (wh.) Unalaschka.	Unalaschka. 1824. 3. 6. H. ⊖.
8 virginiana. L.	(wh.) notch-petal'd. N. America. 1748. 3. 8. H. ♀.
9 acutiflora.	(wh.) acute-petal'd. — H. ♀.
<i>virginica.</i> B.M. non L. <i>virginiana a. acutiflora.</i> DC.	—
10 grandiflora. B.F.G. (pi.)	large-flowered. N. America. H. ♀.
11 caroliniana. M.	(pi.) spatula-leaved. — 1789. — H. ♀.
MO'NTIA. DC.	WATER CHICKWEED. Triandria Trigynia. L.
fontana. DC.	(wh.) common. Britain. 4. 6. H. ⊖.
GINGINSIA. DC.	GING'NSIA. Pentandria Trigynia.
1 brevicaulis. DC.	(pi.) short-stalked. C. B. S. 1816. 5. 6. G. h.
2 elongata. DC.	(pi.) long-peduncled. — 1795. — G. h.
<i>Pharnaceum incanum.</i> L. <i>lineare.</i> Andr. <i>non alior.</i>	
3 aurantiaca. DC.	(or.) orange-flowered. C. B. S. — — G. h.
4 albicans. DC.	(wh.) white-flowered. — — — G. h.
5 conferta. DC.	(wh.) close-leaved. — 1782. 5. 10. G. h.
<i>Pharnaceum incanum.</i> B.M. non L.	
LIMEUM. W.	L'IMEUM. Heptandria Digynia. L.
africatum, w.	(wh.) African. C. B. S. 1774. 6. 7. G. ♀.

ORDO C.

PARONYCHIEÆ. DC. prodr. v. 3. p. 365.

Tribus I. TELEPHIEÆ. DC. prodr. v. 3. p. 366.

TEL'EPHIUM. DC.	ORPINE. Pentandria Trigynia. L.
Imperati. DC.	(wh.) true. S. Europe. 1658. 6. 8. H. ♀.
CORRIGIOLA. DC.	STRAPWORT. Pentandria Trigynia. L.
1 telephiifolia. DC.	(wh.) Orpine-leaved. Spain. 1822. 6. 8. H. ♀.
2 littoralis. DC.	(wh.) sand. England. 7. 8. H. ♀.

Tribus II. ILLECEBREÆ. DC. prodr. v. 3. p. 367.

HERNITARIA. DC.	RUPTURE-WORT. Pentandria Digynia. L.
1 cinerea. DC.	(gr.) cinereous. Montpellier. 1816. 7. 8. H. ⊖.
2 glabra. DC.	(gr.) smooth. England. — H. ♀.
3 hirsuta. DC.	(gr.) hairy. — — — H. ♀.
4 incana. DC.	(gr.wh.) hoary. S. Europe. 1816. 5. 8. H. ♀.
5 Besseri. DC.	(gr.) Besser's. Tauria. 1824. — — H. ♀.
6 alpina. DC.	(gr.) alpine. S. Europe. 1816. 7. 8. H. ♀.
7 fruticosa. DC.	(gr.) frutescent. Spain. 1814. 5. 8. H. h.
8 polygonoides. DC. (gr.)	Knot-grass-like. S. Europe. 1752. — F. h.
ANY'CHIA. DC.	ANY'CHIA. Pentandria Monogynia.
1 dichotoma. DC.	(gr.) forked. N. America. 1806. 5. 8. H. ♂.
<i>Quericia canadensis.</i> L.	
2 capillacea. DC.	smooth-stalked. — 1824. — H. ♂.
ILLECEBRUM. DC.	KNOT-GRASS. Pentandria Monogynia. L.
verticillatum. DC.	(wh.) whorled. England. 7. 8. H. ♀.
CHÆTONY'CHIA. DC.	CHÆTONY'CHIA. Pentandria Monogynia.
cymosa. DC.	(wh.) cymose. S. Europe. 1817. 6. 7. H. ♀.
<i>Illecebrum cymosum.</i> L.	
PARONY'CHIA. DC.	PARONY'CHIA. Pentandria Monogynia.
1 echinata. DC.	(wh.) bristly. S. Europe. 1817. 6. 7. H. ♀.

Schrad. journ. 1801. t. 4.

2 bonariensis. DC.	(wh.) Buenos Ayres.	BuenosAyres. 1826.	8. 10.	H. ⊙.	
brasiliæna.	Lam. dict.	5. p. 23.			
3 arabica. DC.	(wh.) Arabian.	Arabia.	1828.	7. 10.	H. ⊙.
4 polygonifolia. DC.	(wh.) Polygonum-ld.	Dauphiny.	1816.	5. 8.	H. ♀.
5 argentea. DC.	(wh.) silvery.	S. Europe.	1640.	—	H. ♀.
hispanica. s.s.	<i>Illécebrum Paronychia</i> . L.				Descr. de l'Egypt. t. 18.
6 italica. R.S.	(wh.) Italian.	Italy.	1816.	—	H. ♀.
7 capitata. DC.	(wh.) capitate.	Spain.	1683.	6. 8.	H. ♀.
8 serpylliælia. DC.	(wh.) Thyme-leaved.	S. Europe.	1820.	—	H. ♀.
9 nivea. DC.	(wh.) villous-leaved.	—	1812.	—	H. ♀.
10 tenuifolia. L.en.	(wh.) slender-leaved.	—	1822.	—	H. ⊙.
11 alsinefolia. R.S.	(wh.) Chickweed-leav'd.	Spain.	—	—	H. ⊙.
14 pubescens. R.S.	(wh.) pubescent.	S. France.	1818.	—	H. ♀.
					Vill. delph. 2. p. 410.ic.
					Flor. græc. t. 247.
					Schrad. journ. 1801.t.4.
					Barrel. ic. t. 687.
					Scop. del. ins. t. 13.
					Schra. jour. 1801.p.412.

Tribus III. POLYCARPÆÆ. DC. prodr. v. 3. p. 373.

POLYCARPÆA. DC. POLYCARPÆA. Pentandria Monogynia.

1 gnaphalodes. DC.	(wh.) woolly-leaved.	Canaries.	1818.	6. 8.	F. ♂.
<i>Hægea gnaphalodes</i> .	p.s.	<i>Möllia gnaphalodes</i> . s.s.	<i>Laháya gnaphalodes</i> . R.S.		
2 latifolia. DC.	(wh.) broad-leaved.	Teneriffe.	1816.	7. 8.	F. ♂.
3 Teneriffæ. DC.	(wh.) forked.	—	1779.	—	H. ⊙.
<i>Möllia diffusa</i> . W.	<i>Laháya diffusa</i> . Schult. syst.				Schrank pl. rar. t. 29.
4 aristata. DC.	(wh.) bearded.	Canaries.	1780.	6. 7.	F. ♂.
5 memphitica. DC.	(wh.) Egyptian.	Egypt.	1828.	7. 9.	H. ⊙.
6 carnosa. DC.	(wh.) fleshy-leaved.	Canaries.	1816.	6. 8.	F. ♂.
7 Smithii. DC.	(wh.) Smith's.	—	—	—	F. ♂.

ORT'EGIA. DC. ORT'EGIA. Triandria Monogynia. L.

1 hispanica. DC.	(gr.) Spanish.	Spain.	1768.	6. 7.	H. ♀.
2 dichotoma. DC.	(gr.) forked.	Italy.	1781.	8. 9.	H. ♀.
POLYCA'RPON. DC. ALL-SEED. Tri-Pentandria Trigynia.					

1 tetraphyllum. DC.	(wh.) four-leaved.	England	—	5. 8.	H. ⊙.
2 alsinefolum. DC.	(wh.) Chickweed-ld.	S. Europe.	—	—	H. ⊙.
<i>Laháya alsinefolia</i> . s.					Eng. bot. v. 15. t. 1031.
3 peploides. DC.	(wh.) Peplis-like.	Sicily.	1818.	6. 8.	H. ♀.
<i>Laháya polycarpoides</i> . s. <i>Möllia polycarpon</i> . s.s.					Bocc. sic. p. 71. t. 38.

Tribus IV. POLLICHIEÆ. DC. prodr. v. 3. p. 377.

POLLI'CHIA. DC. POLLICHIA. Monandria Monogynia.

campéstris. DC. (wh.) whorl-leaved. C. B. S. 1780. 8. 10. G. ♂. Smith spicil. 1. t. 1.

Tribus V. SCLERANTHEÆ. DC. prodr. v. 3. p. 377.

MNI'TARUM. DC. MNI'TARUM. Monandria Digynia.

1 biflorum. DC.	(gr.) two-flowered.	N. Holland.	1818.	5. 7.	G. ♀.
SCLER'ANTHUS. DC.	KNAWEL.	Decandria Digynia. L.			Forst. gen. 1. t. 1.
1 ánnuus. DC.	(gr.) annual.	Britain.	—	7. 8.	H. ⊙.
2 perennius. DC.	(gr.) perennial.	—	—	8. 9.	H. ♀.

Eng. bot. v. 5. t. 351.

v. 5. t. 352.

Tribus VI. QUERIACEÆ. DC. prodr. v. 3. p. 379.

QU'ERIA. DC. QU'ERIA. Triandria Trigynia. L.

hispanica. DC. (gr.) Spanish. Spain. 1800. 5. 9. H. ⊙. Quer. fl. esp. 6. t. 15. f. 2.

Tribus VII. MINUARTIEÆ. DC. prodr. v. 3. p. 379.

MINUA'R'TIA. DC. MINUA'R'TIA. Tri-Decandria Trigynia.

1 dichotoma. DC.	(wh.) forked.	Spain.	1771.	6. 7.	H. ⊙.
					Loefl. it. p. 121. t. 1. f. 3.

2 montàna. DC.	(wh.) mountain.	Spain.	1806.	6. 7. H. ⊖.	Lœfl. it. p. 122.t.1.f. 4.
3 campéstris. DC.	(wh.) field.	—	—	— H. ⊖.	Act. stock. 1758.t.1.f.3.
LŒFLI'NGIA. DC.	LŒFLI'NGIA.	Triandria Monogynia. L.			
hispánica. DC.	(wh.) Spanish.	Spain.	1770.	6. 7. H. ⊖.	Cavan. ic. 1. t. 94.

ORDO CI.

CRASSULACEÆ. DC. prodr. v. 3. p. 381.

Tribus I. CRASSULEÆ. DC. prodr. v. 3. p. 381.

TILL'EA. DC.	TILL'EA.	Tetrandria Tetragynia. L.			
1 muscòsa. DC.	(wh.) mossy.	England.	5. 7. H. ⊖.	Eng. bot. v. 2. t. 116.
2 moschàta. DC.	(wh.) musk-scented.	Magellan.	1794.	5. 11. H. ⊖.	
Crássula moschàta. Forster.					
3 verticillàris. DC.	(wh.) whorled.	N. S. W.	1823.	6. 9. H. ⊖.	
BULLIA'RDA. DC.	BULLIA'RDA.	Tetrandria Tetragynia.			
1 Vaillanti. DC.	(wh.) Vaillant's.	France.	1825.	6. 8. H.w. ⊖.	DC. plant. gras. t. 74.
2 aquática. DC.	(wh.) water.	N. Europe.	1816.	— H.w. ⊖.	
DASYST'EMON. DC.	DASYST'EMON.	Pentandria Pentagynia.			
calycinum. DC.	(wh.) leafy-calyxed.	N. Holland.	1823.	6. 9. H. ⊖.	
SE'PTAS. DC.	SE'PTAS.	Heptandria Heptagynia. L.			
1 capénisia. DC.	(wh.) Cape.	C. B. S.	1774.	8. 9. D.G. Ȑ.	Andr. rep. t. 98.
2 globifera. B.M.	(wh.) globe-flowering.	—	1809.	3. 5. D.G. Ȑ.	Bot. mag. t. 1472.
3 Umbélla. DC.	(wh.) skreen.	—	1800.	7. 8. D.G. Ȑ.	Jacq. ic. t. 352.
Crássulu Umbélla. Jacq.					
CRA'SSULA. H.R.	CRA'SSULA.	Pentandria Pentagynia. L.			
1 arboréscens. DC.	(pi.) arborescent.	C. B. S.	1730.	5. 6. D.G. Ȑ.	Bot. mag. t. 384.
Cotylédon. B.M.					
2 portulacea. DC.	(ro.) oblique-leaved.	—	1759.	4. 5. D.G. Ȑ.	DC. plant. gras. t. 79.
obliqua. H.S.					
3 láctea. H.S.	(wh.) milk white.	—	1774.	9. 2. D.G. Ȑ.	Bot. mag. t. 1771.
4 telephioídes. H.R.	(ro.) Orpine-like.	—	1820.	7. 8. D.G. Ȑ.	
5 ramosa. H.S.	(ro.) branching.	—	1774.	— D.G. Ȑ.	
6 revolvéns. H.P.	(wh.) recurved-leaved.	—	1820.	8. 10. D.G. Ȑ.	DC. plant. gras. t. 19.
7 tetragóna. H.S.	(wh.) square-leaved.	—	1711.	— D.G. Ȑ.	
8 biplanáta. H.P.	(wh.) flat-leaved.	—	1820.	— D.G. Ȑ.	DC. plant. gras. t. 2.
9 acutifòlia. H.S.	(wh.) acute-leaved.	—	1795.	9. 11. D.G. Ȑ.	
10 bibracteata. H.P.	(wh.) double-bracted.	—	1820.	8. 10. D.G. Ȑ.	
α minor. H.P.	(wh.) smaller.	—	—	— D.G. Ȑ.	
β major. H.P.	(wh.) larger.	—	—	— D.G. Ȑ.	
11 filicaulis. H.P.	(wh.) thready-stem'd.	—	—	— D.G. Ȑ.	
12 scábra. H.R.	(wh.) rough-leaved.	—	1730.	6. 7. D.G. Ȑ.	Dill. elth. t. 99. f. 117.
13 scarélla. H.R.	(wh.) patent-leaved.	—	1820.	6. 9. D.G. Ȑ.	
14 squamułosa. H.R.	(wh.) small-scaly.	—	1817.	— D.G. Ȑ.	Mart. cent. t. 24.
15 bullulàta. H.S.	(ye.) rough yellow.	—	1800.	6. 7. D.G. Ȑ.	
16 pruinòsa. DC.	(wh.) frosted.	—	1821.	— D.G. Ȑ.	Burm. afr. t. 9. f. 2.
17 columnàris. H.S.	(wh.) columnar.	—	1789.	7. 10. D.G. Ȑ.	
18 lycopodioides. DC.	(wh. pu.) whip-cord.	—	1760.	6. 7. D.G. Ȑ.	
imbricata. H.K.					
19 ericoïdes. H.P.	(wh.) large whip-cord.	—	1820.	9. 10. D.G. Ȑ.	
20 vestita. DC.	(ye.) powdered-leav'd.	—	6. 10. D.G. Ȑ.	
21 argéntea. H.S.	(wh.) silvery-leaved.	—	1820.	7. 9. D.G. Ȑ.	
22 perfóssa. DC.	(wh.) threaded.	—	1785.	8. 9. D.G. Ȑ.	DC. plant. gras. t. 25.
perfíllata. H.S.	Scop. del. ins. 3. t. 6. c.	—		— D.G. Ȑ.	
23 perforàta. DC.	(wh.) perfoliate.	—			
24 marginàlis. DC.	(wh.) red-margined.	—	1774.	7. 8. D.G. Ȑ.	Jacq. schœnb. 4. t. 471.
25 pellúcida. H.S.	(bh.) pellucid.	—	1732.	6. 9. D.G. Ȑ.	Dill. elth. t. 100. f. 119.
26 centauroídes. DC.	(ro.) Centaury-flow'd.	—	1774.	5. 7. D.G. Ȑ.	Bot. mag. t. 1765.
27 cordàta. DC.	(ro.) heart-leaved.	—	—	5. 8. D.G. Ȑ.	DC. plant. gras. 2.t.121.
28 spathulàta. DC.	(ro.) notch-leaved.	—	—	7. 9. D.G. Ȑ.	— 1. t. 49.
29 orbicùlaris. H.S.	(bh.) starry.	—	1731.	— D.G. Ȑ.	— 1. t. 43.
30 rosulàris. H.R.	(wh.) great starry.	—	1819.	— D.G. Ȑ.	
31 glomeràta. H.S.	(wh.) rough clustered.	—	1774.	8. 10. H. ⊖.	DC. plant. gras. 1. t. 57.

32 glábra. H.R.	(wh.) smooth-cluster'd. C. B. S.	1774.	6. 10. H. ⊙.	
33 expánsa. H.S.	(wh.) awl-leaved.	—	6. 8. H. ⊙.	
34 diffusa. H.S.	(wh.) diffuse.	—	— H. ⊙.	
35 subuláta. H.S.	(wh.) awl-leaved.	1820.	— H. ⊙.	Petiv. gaz. t. 89. f. 8.
36 lineoláta. H.S.	(wh.) channelled.	1774.	— D.G. ♂.	
37 biconvexa. H.S.	(wh.) doubly convex.	1800.	8. 9. D.G. ♂.	
38 pulchella. H.R.	(ye.) pretty.	1778.	7. 8. H. ⊙.	
39 spársa. H.S.	(st.) alternate-leaved.	1774.	6. 9. H. ⊙.	
PURG'OSEA. H.P.		Pentandria Pentagynia.		
1 linguæfólia. H.R.	(wh.) tongue-leaved.	C. B. S.	1803.	8. 9. D.G. ♀.
2 ciliáta. DC.	(st.) ciliated.	—	1732.	7. 8. D.G. ♀.
3 concínna.	(wh.) small neat.	—	1820.	D.G. ♂.
<i>Crásula concínna. H.R.</i>				
4 ligulifólia.	(wh.) latchet-leaved.	—	—	D.G. ♂.
<i>Crásula ligulifólia. H.R.</i>				
5 concinnéla. DC.	elegant.	—	1822.	D.G. ♂.
6 tomentosa. H.R.	(wh.) tomentose.	—	1790.	4. 5. D.G. ♂.
7 cotyledónis. DC.	(wh.) Cotyledon-like.	—	1800.	7. 9. D.G. ♀.
8 conspícuua.	(wh.) conspicuous.	—	1820.	D.G. ♀.
<i>Crásula conspícuua. H.R.</i>				
9 corymbulósa. DC.	(wh.) impress-dotted.	—	1818.	9. 11. D.G. ♀.
<i>pertísa. H.R. Crásula corymbulósa. L.en.</i>		Ot. et Lk. abb. t. 16.		
10 pertísa. H.P.	(wh.) small impress-dot'd.	—	1822.	D.G. ♀.
11 capitelláta. H.R.	(wh.) square-spiked.	—	1774.	7. 8. D.G. ♂.
12 thyrsiflóra. H.R.	(wh.) thyrs-flowered.	—	1824.	D.G. ♂.
13 obováta. H.R.	(wh.) obovate-leaved.	—	1818.	D.G. ♂.
14 aloídes. H.R.	(bh.) Aloe-like.	—	1774.	6. 8. D.G. ♂.
15 punctátæ. DC.	(wh.) dotted.	—	1759.	4. 8. D.G. ♂.
16 ramuliflóra. DC.	(wh.) branch-flowered.	—	1821.	6. 7. D.G. ♂.
17 turritá. H.R.	(bh.) tower-like.	—	1818.	9. 11. D.G. ♂.
<i>a álba.</i>	(wh.) white-flowered.	—	—	Jacq. schœnb. 1.t. 52.
<i>β rôsea. H.S.S.</i>	rose-coloured.	—	—	D.G. ♂.
GLOB'ULEA. DC.		Pentandria Pentagynia.		
1 cultráta. DC.	(wh.) sharp-leaved.	C. B. S.	1732.	7. 8. D.G. ♂.
2 rádicans. DC.	(wh.) rooting-branch'd.	—	1823.	12. 3. D.G. ♂.
3 atropurpurea. DC.	(wh.) purple-leaved.	—	1823.	7. 8. D.G. ♂.
4 Lingúa. DC.	(wh.) tongue-leaved.	—	—	5. 6. D.G. ♀.
5 língula. DC.	(wh.) lesser-tongue.	—	—	D.G. ♀.
6 capitátæ. DC.	(wh.) large grey-leav'd.	—	1819.	7. 8. D.G. ♀.
7 obvalláta. DC.	(wh.) Houseleek-leav'd.	—	1795.	D.G. ♂.
<i>Crásula obvalláta. L. Trew. pl. rar. 1. t. 11.</i>		DC. plant. gras. t. 61.		
8 canescens. DC.	(wh.) canescent.	—	1800.	D.G. ♀.
9 nudicáulis. DC.	(wh.) naked-stalked.	—	1732.	5. 9. D.G. ♀.
10 sulcátæ. DC.	(wh.) sulcate.	—	1820.	D.G. ♀.
11 impréssæ. DC.	(wh.) impressed.	—	1822.	7. 8. D.G. ♀.
<i>β minor. H.P.</i>	(wh.) lesser.	—	—	Comm. præl. t. 29.
12 paniculáta. DC.	(wh.) paniced.	—	1823.	D.G. ♀.
13 hispida. DC.	(wh.) hispid.	—	—	10. 11. D.G. ♂.
14 mesembryanthoïdes. DC.	(wh.) hedgehog.	—	1822.	7. 9. D.G. ♂.
<i>β áltior.</i>	(wh.) taller.	—	—	D.G. ♂.
15 móllis. DC.	(wh.) soft.	—	1774.	D.G. ♂.
16 subincána. DC.	(wh.) hoary frutescent.	—	1822.	D.G. ♂.
<i>a decumbens. H.P.</i>	(wh.) decumbent.	—	—	D.G. ♂.
<i>β erécta. H.P.</i>	(wh.) erect.	—	—	D.G. ♂.
CURTO'GYNE. H.R.		Pentandria Pentagynia.		
1 dejécta. DC.	(wh.) twisted-branch'd. C. B. S.	1821.	6. 7. D.G. ♂.	Jacq. schœnb. t. 423.
<i>Crásula dejécta. Jacq.</i>				
2 unduláta. H.R.	(wh.) lesser-waved.	—	1797.	8. 11. D.G. ♂.
3 undátæ. H.R.	(wh.) greater-waved.	—	1818.	D.G. ♂.
GRAMMA'NTHES. DC.		Pentandria Pentagynia.		
1 chloræflóra. DC.	(ye.) Chlora-flower'd. C. B. S.	1774.	6. 8. H. ⊙.	Herm. lugd. t. 553.
<i>Crásula dichótoma. L. Vauánthes chloræflóra. H.R.</i>				
2 retrofléxa.	(or.) orange-flower'd.	—	1788.	H. ⊙.
<i>Crásula retrofléxa. H.S.</i>				
3 gentianóides. DC.	(ye.) Gentian-flower'd.	—	1816.	H. ⊙.
<i>R'OCHEA. DC.</i>		Pentandria Pentagynia.		
1 falcata. DC.	(or.) falcated.	C. B. S.	1785.	6. 9. D.G. ♂.
<i>Crásula falcata. Bot. mag. t. 2035. obliqua. Andr. deposit. t. 414.</i>		DC. plant. gras. t. 103.		
<i>β minor. H.P.</i>	(sc.) smaller.	C. B. S.	1821.	D.G. ♂.

CRASSULACEÆ.

2 perfoliata. DC.	(vu.) great perfoliate. C. B. S.	1725.	7. 8. D.G. ½.	Mill. ic. t. 108.	
α coccinea. DC.	(sc.) scarlet.	—	— D.G. ½.		
β álba. DC.	(wh.) white-flowered.	—	— D.G. ½.	DC. plant. gras. t. 13.	
3 albiflòra. DC.	(wh.) white-flowered.	1820.	7. 9. D.G. ½.	Bot. mag. t. 2391.	
Crassula albiflòra. B.M.					
KALOSA'NTHES. H.R. KALOSA'NTHES.	Pentandria Pentagynia.				
1 coccinea. H.R.	(sc.) scarlet.	C. B. S.	1710.	6. 8. D.G. ½.	
Crassula coccinea. B.M. Röchea coccinea. DC.	plant. gras. t. 1.			Bot. mag. t. 495.	
β álba. H.R.	(wh.) white-flowered.	C. B. S.	1710.	6. 8. D.G. ½.	
2média. H.R.	(std.) great changeable.	—	1820.	— D.G. ½.	
3 versicolor. H.R.	(std.) various-coloured.	—	1816.	— D.G. ½.	
Crassula versicolor. B.R.					
4 bicolor. H.R.	(ye.sc.) two-coloured.	—	1820.	— D.G. ½.	
5 capitata.	(wh.) headed.	—	1822.	— D.G. ½.	
Crassula capitata. L.C. nec aliorum.				Lodd. bot. cab. t. 1029.	
6 odoratissima. H.R. (st.) sweet-scented.	—	1793.	6. 7. D.G. ½.	Andr. rep. t. 26.	
Crassula odoratissima. Jacq. schoenb. t. 434. Dietrichia odoratissima. Tratt.					
7 flava. H.S.	(ye.) yellow-flower'd. C. B. S.	1802.	6. 9. D.G. ½.	Burm. afr. t. 23. f. 3.	
8 cymosa. H.R.	(wh.) cymose.	—	1800.	— D.G. ½.	
9 biconvexa. H.P.	(wh.) convex-leaved.	—	1822.	— D.G. ½.	
10 jasmínea. H.R.	(wh.) Jasmine-flower'd.	—	1812.	6. 8. D.G. ½.	
Crassula jasmínea. B.M. Röchea jasmínea. DC.				Bot. mag. t. 2178.	
KALANCH'OE. DC. KALANCH'OE.	Octandria Tetracynia.				
1 spathulata. DC.	(ye.) spathulate.	China.	1820.	7. 9. D.G. ½.	
2 aegyptiaca. DC.	(or.) Egyptian.	Egypt.	1820.	7. 9. D.G. ½.	
3 rotundifolia. H.P. (ye.) round-leaved.	C. B. S.	—	— D.G. ½.	DC. plant. gras. t. 64.	
4 acutiflora. H.P. (wh.) white-flowered.	E. Indies.	1806.	— D.S. ½.	Andr. deposit. t. 560.	
Veria acutiflora. A.R.					
5 crenata. DC.	(ye.) great notched.	Sierra Leone.	1793.	— D.S. ½.	Andr. deposit. t. 21.
Cotyledon crenatum. Bot. mag. t. 1436.	Veria crenata. A.R.				
6 laciñata. DC.	(ye.) cut-leaved.	E. Indies.	1731.	— D.S. ½.	DC. plant. gras. t. 100.
7 ceratophylla. DC. (st.) buck's-horn-l'd.	China.	1820.	9. 1. D.G. ½.		
BRYOPHYLLUM. DC. BRYOPHYLLUM.	Octandria Tetracynia.				
calycinum. DC.	(br.) large-cupped.	Mauritius.	1800.	4. 7. D.G. ½.	Bot. mag. t. 1409.
COTYLEDON. DC. NAVELWORT.	Decandria Pentagynia. L.				
1 undulata. H.S.S.	(ye.) wave-leaved.	C. B. S.	1818.	7. 8. D.G. ½.	
2 orbiculata. H.S. (ye.bh.)	round-leaved.	—	1789.	— D.G. ½.	
3 ovata. H.S.	(ye.bh.) ovate-leaved.	—	—	— D.G. ½.	Bot. mag. t. 321.
orbiculata. B.M. orbiculata β obovata. DC.	pl. gras. t. 76.				
4 oblónga. H.S.	(ye.bh.) oblong-leaved.	C. B. S.	1690.	7. 10. D.G. ½.	
5 elata. H.S.S.	(ye.bl.) tall powdered.	—	1816.	6. 8. D.G. ½.	
6 crassifolia. H.P. thick-leaved.	—	1823. D.G. ½.	
7 viridis. H.P. green-leaved.	—	—	— D.G. ½.	
8 ramosa. H.S.S. (ye.bh.)	branching.	—	1768.	7. 10. D.G. ½.	Moris.his.s.12.t.7.f.39.
9 ramosissima. H.S.S. (ye.)	cluster-branch'd.	—	1816.	— D.G. ½.	
10 corúscans. H.S.S. (or.)	sparkling.	—	1818.	3. 5. D.G. ½.	Bot. mag. t. 2601.
canalifolia. H.P.					
11 ungulata. DC.	(pu.) ungulate.	—	1822.	5. 8. D.G. ½.	Bur.afr.dec.3.t.22.f.1.
12 papillaris. DC.	(or.) cross-leaved.	—	1813.	— D.G. ½.	Bot. reg. t. 915.
decessata.	Bot. mag. t. 2518.				
13 tricuspidata. H.P. (or.)	trident-leaved.	—	1820.	— D.G. ½.	
14 purpurea. DC.	(pu.) purple.	—	1813.	7. 8. D.G. ½.	Burm. afr. t. 20. f. 1.
15 rotundifolia. H.P. (ye.bh.)	round-leav'd.	—	1826.	— D.G. ½.	
16 cristata. H.P.	(pu.) Coxcomb-leaved.	—	1825.	8. 10. D.G. ½.	
17 clavifolia. H.P.	(pu.) club-leaved.	—	—	— D.G. ½.	
18 curviflora. DC.	(ye.bh.) curve-flower'd.	—	—	— D.G. ½.	Bot. mag. t. 2044.
19 tuberculosa. DC.	(or.) tubercled.	—	1822.	— D.G. ½.	DC. plant. gras. t. 1. t. 86.
20 ecalloides. DC.	(ye.) Cacalia-like.	—	1820.	5. 6. D.G. ½.	
21 ventricosa. DC.	(gr.) bellied-tubed.	—	1825.	6. 8. D.G. ½.	Burm. afr. d. 3.t.21.f.1.
22 spuriæ. DC.	(ye.) narrow-leaved.	—	1731.	7. 8. D.G. ½.	— t. 19. f. 1.
23 fascicularis. DC. (gr.re.)	clustered.	—	1759.	7. 9. D.G. ½.	— t. 18.
tardiflorum. Bonpl. nav. t. 37.					
24 triflora. DC.	(wh.pu.) three-flowered.	—	1821.	7. 19. D.G. ½.	
25 maculata. DC.	(v.) spotted-leaved.	—	1818.	5. 7. D.G. ½.	
26 rhombifolia. H.P.	(ye.) rhomb-leaved.	—	1823.	— D.G. ½.	
27 jasminiflora. DC.	(v.) Jasmine-flower'd.	—	1818.	6. 8. D.G. ½.	
28 hemisphærica. DC. (v.)	thick-leaved.	—	1731.	— D.G. ½.	DC. plant. gras. 1. t. 87.
29 mamillaris. DC.	(v.) mammillary.	—	1818.	5. 7. D.G. ½.	
30 cuneata. DC.	(ye.) large pubescent.	—	1813.	6. 8. D.G. ½.	

31	caryophyllacea. DC. (re.)	Clove-flowered.	C. B. S.	1826.	6. 8. D.G. ȏ.	Burm. dec. 2. t. 17.
32	mucronata. DC. (ye.)	mucronate.	—	1818.	— D.G. ȏ.	Burm. afr. t. 19. f. 2.
33	canaliculata. H.S.S. (v.)	channel-leav'd.	—	1817.	— D.G. ȏ.	
34	gracilis. H.S.S. (ye.)	slender.	—	1800.	5. 8. D.G. ȏ.	
35	dichotoma. H.R. (gr.)	fork-spined.	—	1818.	— D.G. ȏ.	
36	cuneiformis. H.P. (w.)	wedge-leaved.	C. B. S.	1823. D.G. ȏ.	
37	interjecta. H.P. (w.)	fleshy-stemmed.	—	— D.G. ȏ.	
PISTOR'NIA. DC.	PISTOR'NIA.	Decandria Pentagynia.				
hispanica. DC. (re.)	Spanish.	Spain.	1796.	6. 7. H. ♂.	DC. plant. grass. t. 122.	
Cotyledon hispanica.	Lefc. it. p. 77. t. 1.					
UMBILICUS. DC.	PENNYWORT.	Decandria Pentagynia.				
1 pendulinus. DC. (gr.)	drooping-flower'd.	Britain.	6. 9. H. ȏ.	Eng. bot. v. 5. t. 325.	
Cotyledon Umbilicus. E.B.						
2 erectus. DC. (ye.)	upright-flower'd.	England.	— H. ȏ.	Eng. bot. v. 22. t. 1522.	
Cotyledon luteum. E.B.						
3 horizontalis. DC. (st.)	horizontal.	Sicily.	1828.	— H. ȏ.		
OROSTACHYS. F.	OROSTACHYS.	Decandria Pentagynia.				
1 serrata. (pu.)	notch-leaved.	Crete.	1732.	6. 7. F. ȏ.	Dill. elth. t. 95. f. 112.	
Cotyledon serrata. H.S.						
2 spinosa. (ye.)	spiny.	Siberia.	— F. ȏ.	Murr. com. gæt. 7. t. 5.	
chlorantha. F. Cotyledon spinosum. L.	Gmel. sib. 4. t. 67. f. 2.					
3 malachophylla. F. (wh.)	soft-leaved.	Dahuria.	1815.	— H. Ⓣ.	Pall. it. 3. ap. t. G. f. 1.	
4 thyrsiflora. F. (bh.)	thyrsse-flowered.	—	1820.	— H. Ⓣ.		
ECHEVERIA. DC.	ECHEVERIA.	Decandria Pentagynia.				
1 coccinea. DC. (sc.)	scarlet.	Mexico.	1816.	8. 10. D.G. ȏ.	Bot. mag. t. 2572.	
Cotyledon coccinea. B.M.						
2 grandifolia. H.P. (or.)	large-leaved.	—	1826.	— D.G. ȏ.	Swt. br. fl. gar. t. 275.	
3 gibbiiflora. DC. (or.)	gibbous-flower'd.	—	—	— D.G. ȏ.	Bot. reg. t. 1247.	
4 caspiosa. DC. (ye.)	tongue-leaved.	California.	1796.	6. 8. D.F. ȏ.	Jacq. f. ecl. 1. t. 17.	
Cotyledon linguisiformis. H.K. Sédum Cotyledon. Jacq.						
RHODIOLA. L.	ROSE-ROOT.	Dicecia Octandria. L.				
1 rosea. L. (ye.)	common.	Britain.	5. 7. H. ȏ.	Eng. bot. v. 8. t. 508.	
Sédum Rhodiola.	DC. plant. gras. t. 143.					
2 sibirica. F. (ye.)	Siberian.	Siberia.	1826.	— H. ȏ.		
S'EDUM. DC.	STONECROP.	Decandria Pentagynia.				
1 Aizòon. DC. (ye.)	yellow Orpine.	Siberia.	1757.	7. 9. H. ȏ.	DC. plant. gras. t. 101.	
2 hybridum. DC. (st.)	Germander-leaved.	—	1766.	5. 7. H. ȏ.	Murr. com. gæt. 6. t. 5.	
3 involucratum. DC. (wh.)	involucrated.	Caucasus.	1822.	— H. ȏ.		
4 verticillatum. DC. (pu.)	whorl-leaved.	Kamtschatka.	7. 9. H. ȏ.	Linn. amoëna. 2. t. 4. f. 14.	
5 latifolium. DC. (st.)	broad-leaved.	Europe.	1794.	— H. ȏ.	Clus. hist. 2. p. 66. f. 1.	
6 maximum. H.S. (pu.)	great purple.	Spain.	—	— H. ȏ.		
7 albicans. H.S. (wh.)	great white.	Europe.	—	— H. ȏ.		
8 argutum. H.R. (pu.)	lance-toothed.	M.Carpathian.	1820.	— H. ȏ.		
9 paucidens. H.R. (wh.)	few-toothed.	Europe.	—	— H. ȏ.		
10 Teléphium. E.B. (pu.)	common Orpine.	Britain.	—	— H. ȏ.	Eng. bot. v. 19. t. 1319.	
11 triphyllum. (pu.)	three-leaved.	Portugal.	1794.	— H. ȏ.		
Anacampseros triphylla. H.S.						
12 purpureum. H.S. (pu.)	purple-stalked.	—	—	— H. ȏ.	DC. plant. gras. t. 92.	
13 telephioïdes. DC. (pu.)	Rhodiola-like.	N. America.	1810.	7. 8. H. ȏ.		
14 spuriun. DC. (pu.)	greater-fringed.	Caucasus.	—	7. 9. H. ȏ.	Bot. mag. t. 2370.	
15 roseum. DC. (ro.)	rose-coloured.	—	1827.	— H. ȏ.		
16 dentatum. DC. (li.)	large-toothed.	—	1810.	— H. ȏ.		
17 pectinatum. DC. (wh.)	pectinated.	—	1819.	— H. ȏ.		
18 lividum. DC. (wh.)	livid-leaved.	—	1820.	— H. ȏ.		
19 cruciatum. DC. (wh.)	cross leaved.	S. Europe.	1816.	6. 7. H. ȏ.	Lodd. bot. cab. t. 464.	
monregalense. Balb. misc. p. 23. t. 6.						
20 ciliare. H.S. (wh.)	fringed.	Caucasus.	1810.	7. 9. H. ȏ.	Bot. mag. t. 1807.	
oppositifolium. B.M. Anacampseros ciliaris. H.S.						
21 ternatum. DC. (bh.)	dwarf 3-leaved.	N. America.	1789.	7. 8. H. ȏ.	Bot. reg. t. 142.	
pulchrum. DC. (pu.)	pretty.	—	1824.	— H. ȏ.		
populifolium. DC. (wh.)	Poplar-leaved.	Siberia.	1780.	— H. ȏ.	Bot. mag. t. 211.	
Notarjanni. DC. (wh.)	few-flowered.	Naples.	1824.	— H. ȏ.	Ten. flor. nap. 1. t. 40.	
Anacampseros. DC. (pu.)	evergreen.	France.	1596.	7. 8. H. ȏ.	Bot. mag. t. 118.	
Anacampseros sempervirens. H.S.						
26 divaricatum. W. (li.)	spreading.	Madeira.	1777.	6. 7. G. ȏ.		
Sempervivum. DC. (re.)	houseleek-leav'd.	Iberia.	1823.	6. 8. H. ȏ.	Bot. mag. t. 2474.	
28 deltoides. DC. (pu.)	deltoid-leaved.	Naples.	1826.	— H. Ⓣ.		
stellatum. DC. (bh.)	starry.	S. Europe.	1640.	6. 7. H. Ⓣ.	Camer. hort. 7. t. 2.	
Cepaea. DC. (wh.)	panicked.	France.	—	7. 8. H. ♂.	Clus. hist. 2. p. 68. ic.	
spathulatum. W.K. (wh.)	spathulate.	Hungary.	1820.	— H. ♂.	W. et K. hung. 2. t. 104.	

CRASSULACEÆ.

32 tetraphyllum. F.G. (<i>wh.</i>) four-leaved.	Levant.	7. 8. H. ♂.	
33 cœruleum. DC. (<i>pu.</i>) blue pubescent.	Tunis.	1821.	7. 9. H. ♀.	Bot. reg. t. 520.
34 ánglicum. DC. (<i>wh.re.</i>) English.	England.	7. 8. H. ♀.	Eng. bot. v. 3. t. 171.
β microphyllum. H.R. (<i>wh.</i>) small-leaved.	—	—	— H. ♀.	
35 oblògnum. H.R. (<i>wh.</i>) oblong-leaved.	—	—	— H. ♀.	
36 atrátum. DC. (<i>wh.</i>) dark annual.	Italy.	1795.	7. 9. H. ⊖.	DC. plant. gras. t. 120.
37 glaucum. DC. (<i>wh.</i>) glaucous.	Hungary.	1816.	— H. ♂.	W. et K. hung. 2. t. 181.
38 sexfidum. DC. (<i>wh.</i>) six-cleft.	Caucasus.	1818.	— H. ⊖.	
39 pálens. H.R. (<i>wh.</i>) annual glaucous.	—	7. 9. H. ⊖.	Eng. bot. v. 6. t. 394.
40 villòsum. DC. (<i>ro.</i>) hairy.	Britain.	6. 7. H. ♀.	
41 pálidum. M.B. (<i>ro.</i>) pale red.	Tauria.	1820.	7. 9. H. ⊖.	DC. plant. gras. t. 55.
42 rübens. DC. (<i>re.</i>) annual red.	S. Europe.	1759.	5. 8. H. ⊖.	
Crássula rübens. DC.				Cavan. ic. t. 69. f. 2.
43 cæspitòsum. DC. (<i>re.</i>) tufted.	—	1788.	7. 8. H. ⊖.	
Crássula cæspitosa. Cav. Crássula Magnoliæ. DC. verticillàris.	L?			
44 dasyphyllum. DC. (<i>wh.</i>) thick-leaved.	England.	6. 7. H. ♀.	Eng. bot. v. 10. t. 656.
45 hispánicum. DC. (<i>wh.</i>) Spanish.	Spain.	1732.	— H. ♀.	Jacq. aust. 5. ap. t. 47.
46 ábum. DC. (<i>wh.</i>) white.	England.	6. 8. H. ♀.	Eng. bot. v. 22. t. 1578.
47 altàicum. F. Altay.	Altay.	1827.	— H. ♀.	
48 quinquéfidum. H.R. (<i>gr.wh.</i>) five-cleft.	—	6. 8. H. ♀.	
49 hexapétalum. H.R. (<i>ye.</i>) six-petal'd.	—	— H. ♀.	
50 quadrifidum. DC. (<i>ye.</i>) four-cleft.	Dahuria.	1800.	7. 8. H. ♀.	Pall. it. 3. a. t. P. f. 1.
51 nudum. DC. (<i>ye.</i>) naked-branch'd.	Madeira.	1777.	— G. ♀.	
52 ochroleùcum. H.P. (<i>st.</i>) tall sharp-cup'd.	Greece.	— F. ♀.	
53 altissimum. DC. (<i>st.</i>) tall blunt-cup'd.	Italy.	1769.	— H. ♀.	DC. plant. gras. t. 116.
Jacquini. H.P. Sempervivum sedifòrme. Jacq. vind. 1. t. 81.				
54 cæruléscens. H.P. (<i>st.</i>) great-straw-color'd.	—	— H. ♀.	
55 réflexum. H.P. (<i>ye.</i>) reflex-leaved.	Britain.	6. 7. H. ♀.	Eng. bot. v. 10. t. 695.
56 septangulare. H.P. (<i>st.</i>) seven-angled.	Europe.	7. 8. H. ♀.	Eng. bot. v. 35. t. 2477.
57 albescens. H.R. (<i>ye.</i>) white-leaved.	England.	6. 7. H. ♀.	
glaucum. E.B. non W.K.				DC. plant. gras. t. 116.
58 minus. H.P. (<i>ye.</i>) lesser short-leaved.	—	— H. ♀.	
refléxum. DC.				
59 spiràle. H.P. (<i>ye.</i>) spiral-leaved.	Europe.	— H. ♀.	
60 sexangulare. DC. (<i>ye.</i>) insipid.	England.	— H. ♀.	Eng. bot. v. 28. t. 1946.
61 acre. DC. (<i>ye.</i>) biting.	Britain.	— H. ♀.	— v. 12. t. 839.
62 recurvàtum. H.P. (<i>ye.</i>) slender recurv'd.	—	1820.	— H. ♀.	Bauh. icon. 3. 428.
63 anopétalum. DC. (<i>st.</i>) broader-leaved.	S. France.	1815.	— H. ♀.	
β aurantiacum. H.P. (<i>or.</i>) orange-colour'd.	—	—	— H. ♀.	
64 vires. H.P. (<i>ye.</i>) deep green.	Portugal.	1774.	— H. ♀.	
65 viridulum. H.P. (<i>ye.</i>) small-green.	—	1826.	6. 8. H. ♀.	Eng. bot. v. 3. t. 170.
66 rupéstre. H.P. (<i>ye.</i>) expanding glaucous.	England.	— H. ♀.	— v. 26. t. 1802.
67 Forsterianum. H.P. (<i>ye.</i>) Forster's.	Wales.	7. 8. H. ♀.	DC. plant. gras. t. 119.
68 saxatilé. DC. (<i>ye.</i>) rock.	S. Europe.	1739.	7. 9. H. ⊖.	Vill. dauph. 3. t. 45.
69 répens. DC. (<i>ye.</i>) creeping.	Pyrenees.	1829.	— H. ⊖.	
Guettardi. Villars.				
70 magellénse. DC. (<i>ye.</i>) Neapolitan.	Naples.	1826.	— H. ⊖.	
MONA'NTHES. H.R. MONA'NTHES.	Dodecandria Heptagynia.			
polyphylla. H.R. (<i>pu.</i>) many-leaved.	Canaries.	1777.	7. 9. D.G. ♀.	Bot. mag. t. 93.
Sempervivum monanthes. B.M.				
SEMPERVIVUM. DC. HOUSELEEK.	Dodecandria Dodecagynia. L.			
1 tortuòsum. DC. (<i>ye.</i>) twisted.	Canaries.	1779.	7. 8. D.G. ♀.	Bot. mag. t. 296.
2 villòsum. DC. (<i>ye.</i>) small villous.	Madeira.	1777.	6. 7. D.G. ♀.	
3 ciliátum. DC. (<i>wh.</i>) ciliated.	Teneriffe.	1815.	— D.G. ♀.	Bot. reg. t. 278.
4 glandulòsum. DC. (<i>ye.</i>) gland-edged.	Madeira.	1777.	3. 5. D.G. ♀.	
5 glutinòsum. DC. (<i>ye.</i>) glutinous.	—	—	7. 8. D.G. ♀.	
6 úrbicum. DC. (<i>ye.</i>) wedge-leaved.	Canaries.	1824.	— D.G. ♀.	
7 retisum. H.P. (<i>ye.</i>) retuse-leaved.	Teneriffe.	—	— D.G. ♀.	
8 arbóreum. DC. (<i>ye.</i>) tree.	Levant.	1640.	12. 3. D.G. ♀.	Bot. reg. t. 99.
β variegátum. H.R. (<i>ye.</i>) variegated.	—	—	— D.G. ♀.	
γ atropurpureum. H.R. (<i>ye.</i>) dark purple-leav'd.	—	—	— D.G. ♀.	
9 frutescens. H.P. (<i>ye.</i>) frutescent.	Teneriffe.	1824.	3. 5. D.G. ♀.	Lodd. bot. cab. t. 1328.
10 tabulæfòrme. DC. (<i>ye.</i>) table-shaped.	Madeira.	1815.	7. 9. D.G. ♀.	DC. plant. gras. t. 141.
11 canariénse. DC. (<i>wh.</i>) Canary Island.	Canaries.	1699.	6. 7. D.G. ♀.	Bot. mag. t. 1980.
12 Smithii. DC. (<i>ye.</i>) Smith's.	—	1815.	8. 10. D.G. ♀.	
13 barbátum. DC. (<i>ye.</i>) bearded.	Madeira.	—	6. 7. D.G. ♀.	
lineolàre. H.S.S. spathulátum. Horn.				
14 hybrídum. H.R. (<i>ye.</i>) hybrid.	Hybrid.	— D.G. ♀.	
15 cæspitòsum. DC. (<i>ye.</i>) latchet-leaved.	Canaries.	1815.	9. 4. D.G. ♀.	Bot. mag. t. 1978.
ciliátum. B.M. non W. barbátum. H.R. non Smith. ciliare. H.S.S.				

16 láxum. H.R.	(ye.) loose-flowered. Canaries.	1820. 7. 10. D.G. ♂.	
17 dodrantále. DC.	(st.) smooth-leaved. ———	1815. 7. 8. D.G. ♂.	
18 àureum. DC.	(ye.) chalice-form. Teneriffe.	——— 6. 7. D.G. ♀.	Bot. reg. t. 892.
	<i>calyciforme.</i> B.R. Lodd. bot. cab. 1368.		
19 punctatūm. DC.	(ye.) dotted-leaved. Canaries.	1824. 7. 9. D.G. ♂.	
20 stellatūm. DC.	(ye.) annual villous. Madeira.	1790. 7. 10. H. ⓧ.	Bot. mag. t. 1809.
	<i>villōsum.</i> B.M. non H.S.		
21 dichótomum. DC.	(ye.) forked. Canaries.	1824. 6. 8. D.G. ♂.	DC. jard. gen. t. 21.
22 africānum. DC.	(ye.) African. C. B. S.	1768. D.G. ♀.	
23 hirtum. DC.	(st.) hairy. Italy. 6. 7. H. ♀.	DC. plant. gras. t. 107.
	<i>soboliferum.</i> B.M. t. 1457 ex DC. <i>globiferum.</i> H.R. non DC.		
24 globiferum. DC.	(ye.) great-flowered. Germany.	1731. —— H. ♀.	Bot. mag. t. 507 et 2115.
25 tectòrum. DC.	(pu.) common. Britain. 6. 9. H. ♀.	Eng. bot. v. 19. t. 1320.
26 flagellifòrme. DC.	(re.) whip-lash. Siberia.	1818. 6. 8. H. ♀.	
27 montanūm. DC.	(re.) mountain. Switzerland.	1752. —— H. ♀.	DC. plant. gras. t. 105.
28 arachnoídeum. DC.	(pu.) Cobweb. Italy.	1699. —— F. ♀.	Bot. mag. t. 68.

Tribus II. PENTHOREÆ.

PENTH'ORUM. DC.	PENTH'ORUM. Decandria Pentagynia. L.		
sedoides. DC.	(st.) American. Virginia.	1768. 7. 8. H. ♀.	Act. ups. 1744. t. 2.
LEWI'SIA. Ph.	LEWI'SIA. Dodecandra Trigynia.		
rediviva. Ph.	(wh.) white-flowered. N. America.	1827. 7. 8. H. ♀.	

ORDO CII.

FICOIDEÆ. DC. *prodr. v. 3. p. 415.*

Tribus I. MESEMBRYANTHEMEÆ.

MESEMBRYA'NTHEMUM. FIG-MARYGOLD. Icosandria Pentagynia. L.			
1 minūtum. H.R.	(re.) small-white. C. B. S.	1795. 9. 11. D.I. ♀.	Bot. mag. t. 1376.
2 mīnimum. H.R.	(st.) small chequer'd. ———	1766. 9. 12. D.I. ♀.	Pet. gaz. t. 39. f. 3.
3 perpusillum. H.R.	(st.) small exerted. ———	1819. —— D.I. ♀.	
4 obcordellum. H.R.	(wh.) glaucous chequer'd. ———	1776. 10. 2. D.I. ♀.	Bot. mag. t. 1647.
5 obconéllum. H.R.	(wh.) green chequer'd. ———	1794. —— D.I. ♀.	
6 ficifòrme. H.R. fig-shaped. ———	1819. D.I. ♀.	
7 truncatéllum. H.R.	(st.) large dotted. ———	1795. 10. 2. D.I. ♀.	
8 fibulifòrme. H.R. cloth button. ——— 11. D.I. ♀.	
9 uvafòrme. H.R. Grape-like. ———	1820. D.I. ♀.	Burm. dec. t. 10. f. 2?
10 nucifòrme. H.R. nut-shaped. ———	1790. D.I. ♀.	
11 testiculare. H.R.	(wh.) broad short white. ———	1774. 11. D.I. ♀.	
12 octophyllum. H.R.	(ye.) narrow short white. ———	——— D.I. ♀.	Bot. mag. t. 1573.
	<i>testiculare.</i> B.M.		
	<i>β longiúsculum.</i> (ye.) longer-leaved. ———	——— D.I. ♀.	
13 testiculatum. DC.	(ro.) rosy. ———	——— D.I. ♀.	Jacq. frag. 73. t. 12. f. 2.
14 obtùsum. H.R.	(re.) obtuse cloven. ———	1792. 3. 4. D.I. ♀.	
15 fissum. H.R. cloven white. ———	1776. D.I. ♀.	
16 digitifòrme. H.R.	(wh.) finger-formed. ———	1775. 4. 6. D.I. ♀.	
	<i>digitatum.</i> H.R?		
17 magnipúnctum. H.R.	(ye.) great dotted club. ———	1816. 9. 11. D.I. ♀.	
	<i>a sesquiunciàle.</i> H.R. (ye.) thickest-leaved. ———	——— D.I. ♀.	
	<i>β unciale.</i> H.R. (ye.) thinner-leaved. ———	——— D.I. ♀.	
18 nobile. H.P.	(ye.) noble. ——— 11. D.I. ♀.	
19 cànūm. H.R.	(ye.) hoary club-leav'd. ———	1795. D.I. ♀.	
20 aloïdes. H.R.	(ye.) Aloe-like. ———	1816. 9. 11. D.I. ♀.	
21 caninūm. H.R.	(ye.) dog-chop. ———	1717. 8. 10. D.G. ♂.	DC. plant. gras. t. 95.
22 vulpinūm. H.R.	(ye.) fox-chop. ———	1795. —— D.G. ♂.	
23 hy'bridum. H.R.	(ye.) hybrid. Hybrid. —— D.G. ♀.	
24 álibidum. H.R.	(ye.) white. C. B. S.	1714. 7. 8. D.G. ♀.	Bot. mag. t. 1824.

25	albinotum.	H.P.	(ye.) keeled white dott'd.	C.B.S.	1824.	7. 9.	D.G. 4.	
26	albipunctum.	H.P.	(ye.) white-spotted.	_____	_____	8. 9.	D.G. 4.	
27	tigrinum.	H.P.	(ye.) tiger-chop.	_____	1790.	9. 11.	D.G. 4.	Bot. reg. t. 260.
28	felinum.	H.P.	(ye.) cat-chop.	_____	1730.	8. 11.	D.G. 4.	DC. plant. gras. t. 152.
29	lupinum.	H.P.	(ye.) wolf's chop.	_____	1822.	6. 8.	D.G. 4.	
30	agninum.	H.P.	(ye.) lamb's chop.	_____	_____	5. 6.	D.G. 4.	
31	mustelinum.	H.P.	(ye.) weasel-chop.	_____	1818.	8. 11.	D.G. 4.	
32	musculinum.	H.P.	(ye.) small trailing.	_____	_____	6. 8.	D.G. 4.	
33	ermuinum.	H.P.	(ye.) Ermine-chop.	_____	1823.	5. 6.	D.G. 4.	
	β mäjus.	H.P.	(ye.) red-tipped.	_____	_____	—	D.G. 4.	
34	murinum.	H.P.	(ye.) mouse-chop.	_____	1790.	9.	D.G. 4.	
35	dolabriiforme.	H.R.	(ye.) hatchet-leaved.	_____	1705.	5. 11.	D.G. 4.	Bot. mag. t. 32.
36	cárinans.	H.R.	(ye.) scymetar-leav'd.	_____	1820.	—	D.G. 4.	
37	scapíger.	H.P.	(ye.) great-scaped.	_____	1822.	8. 9.	D.G. 4.	
38	denticulatum.	H.R.	(st.) toothed-beak'd.	_____	1793.	4.	D.G. 4.	
	α cänum.	H.R.	(st.) white-leaved.	_____	_____	—	D.G. 4.	
	β glaucum.	H.R.	(st.) glaucous-leaved.	_____	_____	—	D.G. 4.	
	γ candidissimum.	H.R.	(st.) whitest.	_____	_____	—	D.G. 4.	
39	robustum.	H.R.	(ye.) robust-beaked.	_____	1795.	—	D.G. 4.	
40	compáctum.	H.R.	(ye.) compact.	_____	1780.	11.	D.G. 4.	
41	quadrifidum.	H.R.	(ye.) quadrifid.	_____	1795.	—	D.G. 4.	
42	bifidum.	H.R.	(ye.) bifid beaked.	_____	_____	—	D.G. 4.	
43	bibracteatum.	H.R.	(ye.) double-bracted.	_____	1803.	4. 11.	D.G. 4.	
44	purpurascens.	DC.	(ye.) purple-bracted.	_____	1826.	—	D.G. 4.	
45	rostratum.	H.R.	(ye.) heron-beaked.	_____	1732.	4. 5.	D.G. 4.	Dill. elth. f. 229.
	α tuberculatum.	H.R.	(ye.) tuberculate.	_____	—	—	D.G. 4.	
46	ramulosum.	H.R.	(ye.) small-beaked.	_____	1803.	3. 11.	D.G. 4.	
47	pisiforme.	H.R. pea-shaped.	_____	1796.	—	D.I. 4.	
48	moniliforme.	H.R.	(wh.) necklace.	_____	1791.	3. 4.	D.I. 4.	
49	scalpratum.	H.R.	(ye.) great tongue.	_____	1714.	8. 10.	D.G. 4.	Dill. elth. t. 183. f. 224.
50	frägrans.	H.R.	(ye.) sweet-scented.	_____	1820.	—	D.G. 4.	Ott. et Lk. abb. t. 43.
51	grandiflorum.	H.P.	(ye.) great-blistered.	_____	1824.	—	D.G. 4.	
52	præpingue.	H.R.	(ye.) soft-tongue.	_____	1792.	—	D.G. 4.	
53	médium.	H.R.	(ye.) large-tongue.	_____	1816.	—	D.G. 4.	
54	cultratum.	H.R.	(ye.) long cultrated.	_____	1820.	—	D.G. 4.	
55	lúcidum.	H.R.	(ye.) high-polished.	_____	1768.	—	D.G. 4.	Dill. elth. f. 227.
56	adscéndens.	H.R.	(ye.) long-peduncled.	_____	1805.	8. 11.	D.G. 4.	
57	pustulatum.	H.R.	(ye.) blistered tongue.	_____	1816.	—	D.G. 4.	
58	lóngum.	H.R.	(ye.) long-thin-leaved.	_____	1725.	8. 10.	D.G. 4.	Bot. mag. t. 1866.
	α flaccidum.	H.R.	(ye.) long-flacid.	_____	—	—	D.G. 4.	
	depresso.	B.M.	nec aliorum.	_____	—	—	D.G. 4.	
	β declive.	H.R.	(ye.) lesser erect-edged.	_____	—	—	D.G. 4.	
	γ angústius.	H.R.	(ye.) narrow-leaved.	_____	—	—	D.G. 4.	
	δ purpurascens.	H.R.	(ye.) purple green.	_____	—	—	D.G. 4.	
	ϵ uncatum.	H.R.	(ye.) leaden green.	_____	1819.	—	D.G. 4.	
	ζ attollens.	H.R.	(ye.) narrow dark.	_____	—	—	D.G. 4.	
59	linguiforme.	H.R.	(ye.) common tongue.	_____	1732.	11. 3.	D.G. 4.	Dill. elth. f. 226.
	β rufescens.	H.R.	(ye.) reddish green.	_____	1804.	—	D.G. 4.	
	γ subruciatum.	H.R.	(ye.) subruciate.	_____	1820.	8. 11.	D.G. 4.	
	δ prostratum.	H.R.	(ye.) prostrate.	_____	—	—	D.G. 4.	
	ϵ assurgens.	H.R.	(ye.) assurgent.	_____	1819.	—	D.G. 4.	
60	látum.	H.R.	(ye.) broad tongue.	_____	1732.	—	D.G. 4.	Dill. elth. f. 225.
	β brève.	H.R.	(ye.) short flat.	_____	1802.	—	D.G. 4.	
61	depresso.	H.R.	(ye.) small recurring-l'd.	_____	1795.	9. 11.	D.G. 4.	Dill. elth. f. 226?
	α pálens.	H.R.	(ye.) light green.	_____	—	—	D.G. 4.	
	β lívidum.	H.R.	(ye.) rufous green.	_____	1819.	—	D.G. 4.	
62	cruciatum.	H.R.	(ye.) soft cruciate.	_____	1792.	5. 11.	D.G. 4.	
63	taurinum.	H.R.	(ye.) Bull's-horn.	_____	1795.	9. 11.	D.G. 4.	
64	Sálmii.	H.R.	(ye.) Prince de Salm's.	_____	1818.	—	D.G. 4.	
	α decussatum.	H.R.	(ye.) decussate.	_____	—	—	D.G. 4.	
	β semicruciatum.	H.R.	(ye.) semicruciate.	_____	1820.	—	D.G. 4.	
	γ angustifolium.	H.R.	(ye.) narrow-leaved.	_____	—	—	D.G. 4.	
65	surréctum.	H.R.	(ye.) thick acute tongue.	_____	1819.	—	D.G. 4.	
66	heterophyllum.	H.R.	(ye.) various-leaved.	_____	1795.	—	D.G. 4.	
67	angustum.	H.R.	(ye.) narrow tongue.	_____	1790.	3. 10.	D.G. 4.	Andr. deposit. 540.
	β pálidum.	H.R.	(ye.) paler-leaved.	_____	1819.	—	D.G. 4.	Dill. elth. f. 242.
	γ heterophyllum.	H.K.	(ye.) semicylindric-l'd.	_____	—	8. 10.	D.G. 4.	
68	diffórmē.	H.R.	(ye.) deformed.	_____	1732.	—	D.G. 4.	
	β brevicaule.	H.R.	(ye.) short-stalked.	_____	—	—	D.G. 4.	

69	bidentatum. H.R. (<i>ye.</i>) double-toothed. C. B. S.	1816. 8. 10. D.G. ȝ.	
	α minus. H.R. (<i>ye.</i>) smaller.	— — — D.G. ȝ.	
	β majus. H.R. (<i>ye.</i>) larger.	— — — D.G. ȝ.	
70	bigibberatum. H.P. (<i>ye.</i>) lesser difformed.	1824. — — D.G. ḥ.	
71	semicylindricum. H.R. (<i>ye.</i>) small difformed.	1732. 3. 11. D.G. ḥ.	Dill. elth. t. 194. f. 241.
72	gibbosum. H.R. (<i>re.</i>) great pale gibbous.	1780. 1. 4. D.G. ḥ.	
73	luteo-viride. H.R. (<i>re.</i>) long pale gibbous.	1795. — — D.G. ḥ.	
74	perviride. H.R. (<i>re.</i>) dark green gibbous.	1792. 1. 5. D.G. ḥ.	
75	pubescens. H.R. (<i>re.</i>) hoary gibbous.	— — — D.G. ḥ.	
76	calamiforme. H.R. (<i>wh.</i>) white fl̄d. quill.	1717. 7. 9. D.G. ḥ.	DC. plant. gras. t. 5.
77	obsubulatum. H.R. ... reversed quill.	1796. — — D.G. ḥ.	
78	cylindricum. H.R. (<i>re.</i>) cylindric quill.	1792. 9. 2. D.G. ḥ.	
79	teretifolium. H.R. (<i>re.</i>) round quill.	1794. — — D.G. ḥ.	
80	teretisulcum. H.R. (<i>re.</i>) roundish quill.	— — — D.G. ḥ.	
81	bellidiflorum. H.R. (<i>re.</i>) daisy-flowered.	1717. 6. 8. D.G. ḥ.	Dill. elth. t. 224. f. 233.
	α glaucescens. H.R. (<i>re.</i>) small glaucous.	— — — D.G. ḥ.	
	β viride. H.R. (<i>ro.</i>) pea green.	1822. — — D.G. ḥ.	
82	subulatum. DC. (<i>std.</i>) great green.	1768. 6. 8. D.G. ḥ.	DC. plant. gras. t. 41.
83	acutum. H.R. (<i>pu.</i>) great awl-leaved.	1793. 11. 4. D.G. ȝ.	
84	punctatum. H.R. (<i>pu.</i>) middle-spotted.	— ... D.G. ȝ.	
85	diminutum. H.R. (<i>pu.</i>) small awl-leaved.	1789. 4. D.G. ȝ.	
	β caulinatum. H.R. (<i>pu.</i>) stalked.	1817. — — D.G. ȝ.	
86	löreum. H.R. (<i>st.</i>) leathery-stalk'd.	1732. 9. D.G. ḥ.	Dill. elth. t. 200. f. 255.
	β congestum. H.R. (<i>st.</i>) crowded-leaved.	1805. — — D.G. ḥ.	
87	diversifolium. H.R. (<i>st.</i>) shorthorn-leav'd.	1726. 3. 10. D.G. ḥ.	Dill. élth. t. 198. f. 252.
	α majus. H.R. (<i>std.</i>) larger.	1819. — — D.G. ḥ.	
	β glaucius. H.R. (<i>st.</i>) glaucescent.	1726. — — D.G. ḥ.	
	γ brevifolium. H.R. (<i>st.</i>) short-leaved.	1819. 8. D.G. ḥ.	
	δ l'ate-virens. H.R. (<i>st.</i>) pale green.	— — — D.G. ḥ.	
	ε atro-virens. H.R. (<i>st.</i>) dark green.	— — 8. 9. D.G. ḥ.	
88	decipiens. H.R. (<i>st.</i>) middle horn-ld.	1820. ... D.G. ḥ.	
89	dùbium. H.R. (<i>st. or.</i>) round stalked.	1800. 11. 5. D.G. ḥ.	Bradl. succ. d. 4. t. 40.
90	corniculatum. H.R. (<i>std.</i>) long-horn-leav'd.	1732. 3. 5. D.G. ḥ.	DC. plant. gras. t. 108.
91	purpureo-álbum. H.P. (<i>std.</i>) purple & white.	1824. 7. 10. D.G. ḥ.	
92	procumbens. H.R. (<i>v.</i>) purple-jointed.	1820. ... D.G. ḥ.	
93	tricolorum. H.R. (<i>st. re.</i>) three-coloured.	1794. 10. D.G. ḥ.	
94	pugioniforme. H.R. (<i>ye.</i>) great dagger-ld.	1714. 7. 9. D.G. ḥ.	DC. plant. gras. t. 82.
	β carnatum. H.R. (<i>ye.</i>) flesh-coloured.	... — — D.G. ḥ.	
	γ purpureum. H.R. (<i>pu.</i>) purple.	... — — D.G. ḥ.	
	δ biennne. H.R. (<i>ye.</i>) biennial.	... — — D.G. ḡ.	
95	capitatum. H.R. (<i>ye.</i>) lesser dagger-ld.	1717. 7. 9. D.G. ḥ.	Bot. reg. t. 494.
	β ramigerum. H.R. (<i>ye.</i>) branching.	1816. — — D.G. ḥ.	
96	brevicaulé. H.R. (<i>ye.</i>) dwarf dagger-ld.	— — — D.G. ḥ.	
97	corúscans. H.R. (<i>ye.</i>) glittering dagger.	1813. — — D.G. ḥ.	
98	elongatum. H.R. (<i>ye.</i>) tuberous long-ld.	1793. 5. D.G. ḥ.	DC. plant. gras. t. 72.
	α grandiflorum. (<i>ye.</i>) large-flowered.	— — — D.G. ḥ.	
	β minus. H.R. (<i>ye.</i>) lesser.	— — — D.G. ḥ.	Bot. reg. t. 493.
	γ fusiforme. H.R. (<i>ye.</i>) fusiform.	— — — D.G. ḥ.	Breyen. cent. t. 80.
99	válidum. H.P. (<i>ro.</i>) robust light green.	1823. — — D.G. ḥ.	
100	geminiflorum. H.R. (<i>ro.</i>) small pale.	1819. — — D.G. ḥ.	
101	simile. H.R. (<i>ro.</i>) short-jointed.	— — 5. 7. D.G. ḥ.	
102	láxum. H.R. (<i>ro.</i>) long-jointed.	— — — D.G. ḥ.	
103	sarmentosum. H.R. (<i>ro.</i>) sarmentose.	1805. 4. 5. D.G. ḥ.	
104	rigidicáule. H.R. (<i>ro.</i>) stiff-stemmed.	1819. — — D.G. ḥ.	
105	Schóllii. H.R. (<i>ro.</i>) Scholl's.	1818. — — D.G. ḥ.	Jacq. frag. t. 51. f. 2.
	aduncum. Jacq. non H. recárvum. H.S.s.		
106	filamentosum. H.R. (<i>ro.</i>) thready.	1732. 3. 4. D.G. ḥ.	Dill. elth. t. 212. f. 273.
107	serrulatum. H.R. (<i>ro.</i>) rosy-stalked.	1795. 11. 12. D.G. ḥ.	
	α glaucum. H.R. (<i>ro.</i>) glaucous.	— — — D.G. ḥ.	
	β viridius. H.R. (<i>ro.</i>) greenish.	— — — D.G. ḥ.	
108	rubricaule. H.R. (<i>ro.</i>) deep red-stalk'd.	1802. 12. 2. D.G. ḥ.	
	α gracilius. H.R. (<i>ro.</i>) slenderest.	— — — D.G. ḥ.	
	β densius. H.R. (<i>ro.</i>) crowded.	— ... — D.G. ḥ.	
	γ subvirens. H.R. (<i>ro.</i>) tall-green.	— ... — D.G. ḥ.	
109	acinaciforme. H.R. (<i>ro.</i>) great scymtar-ld.	1714. 8. 9. D.G. ḥ.	Dill. elth. f. 270. 271.
	α brève. H.R. (<i>ro.</i>) short-branched.	— — — D.G. ḥ.	Andr. deposit. 508.
	β longum. H.R. (<i>ro.</i>) longer-branched.	— — — D.G. ḥ.	
110	lævigatum. H.R. (<i>ro.</i>) smooth scymtar.	1802. — — D.G. ḥ.	

FICOIDEÆ.

	C. B. S.	1811.	8. 9.	D.G. ™.
111 rùbro-cíncutum. H.R. (ro.) red-edged. α cràssum. H.R. (ro.) thick-leaved. β compréssum. H.R. (ro.) compressed. γ ténérum. H.R. (ro.) tender.	—	—	D.G. ™.
112 subalátum. H.R. (ro.) wing-stalked.	—	—	D.G. ™.
113 édule. H.R. (ye.) Hottentot's fig.	—	1802.	D.G. ™.
114 dimidiáatum. H.R. (ro.) lesser eatable.	—	1690.	7. 8.	D.G. ™.
115 glaucésens. H.R. (ro.) glaucescent.	N. Holland.	1804.	7.	D.G. ™.
116 abbreviátum. H.P. short-jointed.	N. S. W.	1825.	D.G. ™.
117 Róssi. H.R. (ro.) Ross's.	V. Diem. Isl.	1821.	D.G. ™.
118 virésens. H.R. (ro.) virescent.	N. Holland.	1804.	6. 8.	D.G. ™.
119 æquilaterálè. H.R. (ro.) equal angled. β decag'num. DC. (ro.) ten-styled.	—	1791.	—	D.G. ™.
120 vires. H.R. (ro.) smooth-edged green.	C. B. S.	1821.	7. 8.	D.G. ™.
121 réptans. H.R. (ro.) creeping.	—	1774.	6. 8.	D.G. ™.
122 débile. H.P. (ro.) smooth weak.	—	1821.	—	D.G. ™.
123 austrálè. H.R. (bh.) New Zealand.	N. Zealand.	1773.	—	D.G. ™.
124 crassifolium. H.R. (ro.) thick-leaved.	C. B. S.	1727.	5. 8.	D.G. ™.
125 clavellátum. H.R. (re.) club-l'd. creeping.	N. Holland.	1803.	6. 7.	D.G. ™.
β aggregátum. H.R. (re.) cluster branched.	—	—	D.G. ™.	Dill. elth. t. 212. f. 272.
γ mìnus. H.R. (re.) shorter-branched.	—	—	D.G. ™.	DC. plant. gras. t. 89.
126 forcícatum. H.R. (re.) scissors-leaved.	C. B. S.	1758.	9. 10.	D.G. ™.
127 geminátum. H.R. (re.) white twin-shoot'd.	—	1792.	—	D.G. ™.
128 marginátum. H.R. (re.) white-edged.	—	1793.	—	D.G. ™.
129 rostélum. H.R. (wh. re.) little prostrate beak.	—	1820.	8. 10.	D.G. ™.
130 perfoliátum. H.R. (ro.) great-hooked. a tricánthum. H.R. (ro.) three-thorned. β monacánthum. H.R. (ro.) one-thorned.	—	1714.	6. 8.	D.G. ™.
131 uncinélum. H.R. (li.) small-hooked.	—	—	D.G. ™.	Jacq. vind. I. t. 26.
132 uncinátum. H.R. (ro.) greenish-hooked.	—	1819.	—	D.G. ™.
133 semidentátum. H.R. (ro.) slender-hooked.	—	1725.	8.	D.G. ™.
134 ûnidens. H.P. (li.) dwarf uncinate.	—	1818.	6. 8.	D.G. ™.
135 víride. H.R. (li.) deep green.	—	1824.	—	D.G. ™.
136 acutángulum. H.P. (wh.) sharp-angled.	—	1792.	7.	D.G. ™.
137 cútum. H.R. (wh.) short-sheathed. β mágus. H.R. (wh.) larger.	—	1820.	6. 8.	D.G. ™.
138 hamátum. w. (wh.) hook'd sheathing.	—	1811.	—	D.G. ™.
139 vaginátum. H.R. (wh.) rough sheathing. β parviflòrum. H.R. (wh.) smaller-flowered.	—	—	D.G. ™.	Dill. elth. t. 192. f. 240.
140 parviflòrum. H.R. (wh.) small-flowered.	—	1802.	7. 8.	D.G. ™.
141 rígidum. H.R. (wh.) rigid.	—	—	D.G. ™.	Bradl. suc. 5. t. 46.
142 tenélum. H.R. (wh.) least perfoliate.	—	1800.	8.	D.G. ™.
143 imbricátum. H.R. (wh.) imbricated. a mìnus. H.R. (wh.) smallest. β mágus. H.R. (wh.) blunt crowded. γ víride. H.R. (wh.) dull green.	—	1793.	—	D.G. ™.
144 multiflòrum. H.R. (wh.) many-flowered. a mágus. H.R. (wh.) larger white. β mìnus. H.R. (wh.) lesser white. γ rùbrum. H.R. (re.) red-flowered. δ pátens. H.R. (wh.) single flowered. ε nitens. H.R. (wh.) bright-green.	—	1792.	—	D.G. ™.
145 umbellátum. H.R. (wh.) umbel-flowered. β apétalum. H.R. (wh.) petalless.	—	—	D.G. ™.	Pluk. phyt. t. 117. f. 1.
146 tumídulum. H.R. (re.) tumid.	—	1816.	—	D.G. ™.
β mìnus. H.R. (re.) lesser.	—	—	D.G. ™.	Dill. elth. t. 208. f. 266.
147 foliosum. H.R. (re.) many-leaved.	—	—	9.	D.G. ™.
148 lineolátum. H.R. (re.) tumid sheathed. β mìnus. H.R. (re.) lesser. γ nitens. H.R. (re.) glittering.	—	1819.	6. 8.	D.G. ™.
149 serrátum. H.R. (ye.) saw-kneed.	—	—	D.G. ™.	Dill. elth. t. 192. f. 238.
150 lácerum. H.R. (re.) ragged-keeled.	—	1707.	6. 7.	D.G. ™.
151 heteropétalum. H.R. (ye.) various-petal'd. a albiflòrum. (wh.) white-flowered. β rôseum. (ro.) rose-coloured.	—	1792.	6.	D.G. ™.
152 glaucinum. H.R. (li.) glaucine. a angústum. H.R. (li.) slenderest-leaved. β cràssum. H.R. (li.) thicker-leaved.	—	1794.	5. 8.	D.G. ™.
	—	1818.	—	D.G. ™.
	—	—	D.G. ™.	Dill. elth. t. 192. f. 238.
	—	—	D.G. ™.	DC. plant. gras. t. 54.

153 mutáibile. H.R. (<i>wh. ro.</i>) changeable. <i>filamentosum. DC.</i>	C. B. S.	1792.	7. 9. D.G. ḥ.	DC. plant. gras. t. 60.
154 incláudens. H.R. (<i>pu.</i>) open-flowered.	—	1805.	6. 9. D.G. ḥ.	Andr. deposit. 388.
155 caulésrens. H.R. (<i>li.</i>) smooth delta-l'd.	—	1731.	5. 7. D.G. ḥ.	Dill. elth. f. 243-4.
156 deltoídes. H.R. (<i>li.</i>) great delta-leaved.	—	—	5. D.G. ḥ.	DC. plant. gras. 53.
157 muricátum. H.R. (<i>li.</i>) small delta-leaved. β minus. H.R. (<i>li.</i>) smaller-leaved.	—	—	— D.G. ḥ.	Dill. elth. f. 195. f. 27.
158 microphyllum. H.R. (<i>ro.</i>) small-green-leav'd.	—	1795.	— D.G. ḥ.	
159 mucronatum. H.R. (<i>ro.</i>) bristle-pointed.	—	1794.	6. 7. D.G. ḥ.	
160 pygmæum. H.R. (<i>ro.</i>) pigmy connate.	—	1820.	— D.G. ḥ.	
161 pulchéllum. DC. (<i>ro.</i>) small grey-leaved. <i>canescens. H.R.</i>	—	1793.	4. D.G. ḥ.	
β revolutum. H.R. (<i>ro.</i>) revolving-branched.	—	1819.	— D.G. ḥ.	
162 máximum. H.R. (<i>ro.</i>) great crescent-l'd.	—	1787.	3. 8. D.G. ḥ.	
163 lunátum. H.R. (<i>ro.</i>) small crescent-l'd.	—	—	4. 6. D.G. ḥ.	
164 falcátum. H.R. (<i>ro.</i>) small sickle-leav'd.	—	1727.	6. 8. D.G. ḥ.	Dill. elth. f. 275-6.
165 decúmbens. H.R. (<i>li.</i>) dwarf decumbent.	—	1759.	5. 10. D.G. ḥ.	
166 incárvatum. H.R. (<i>li.</i>) incurved. β dilatans. H.R. (<i>li.</i>) dilated.	—	1802.	6. D.G. ḥ.	
γ pallidius. H.R. (<i>li.</i>) pale flowering.	—	1805.	9. 10. D.G. ḥ.	
δ densifolium. H.R. (<i>li.</i>) dense lesser-leav'd.	—	1819.	— D.G. ḥ.	
ε róseum. H.R. (<i>ro.</i>) rosy.	—	—	— D.G. ḥ.	
167 confértum. H.R. (<i>li.</i>) crowded-leaved.	—	1805.	— D.G. ḥ.	
168 falciforme. H.R. (<i>li.</i>) sickle-shaped.	—	—	7. 8. D.G. ḥ.	
169 glomerátum. H.R. (<i>re.</i>) clustered.	—	1732.	6. 8. D.G. ḥ.	Dill. elth. t. 213. f. 274.
170 infléxum. H.R. (<i>re.</i>) inflex-leaved.	—	1818.	— D.G. ḥ.	
171 scábrum. H.R. (<i>re.</i>) rough. α pálidum. H.R. (<i>li.</i>) pale-bloomed.	—	1731.	7. D.G. ḥ.	Dill. elth. t. 197. f. 251.
β purpúreum. H.R. (<i>pu.</i>) purple.	—	—	— D.G. ḥ.	
172 versicolor. H.R. (<i>wh. re.</i>) changeable rough.	—	1795.	5. 8. D.G. ḥ.	
173 retrofléxum. H.R. (<i>re.</i>) reflex-flowered.	—	1794.	5. 10. D.G. ḥ.	
174 imbricáns. H.R. (<i>re.</i>) imbricate-leaved.	—	1818.	— D.G. ḥ.	
175 defléxum. H.R. (<i>re.</i>) smooth deflexed.	—	1774.	7. 10. D.G. ḥ.	
176 polyánthom. H.R. (<i>pu.</i>) copious-flowered.	—	1803.	8. D.G. ḥ.	
177 polypyllum. H.R. (<i>re.</i>) crowded-leaved.	—	1819.	6. 10. D.G. ḥ.	
178 violáceum. H.R. (<i>vi.</i>) deep violaceous.	—	1820.	10. D.G. ḥ.	
179 emarginátum. H.R. (<i>vi.</i>) notch-petaled.	—	1732.	6. 8. D.G. ḥ.	DC. plant. gras. t. 84.
180 dilatátum. H.R. (<i>wh.</i>) gibbous-keeled.	—	1795.	7. D.G. ḥ.	Dill. elth. t. 197. f. 250.
181 virgátum. H.R. (<i>li.</i>) slender-twigged.	—	1793.	2. 4. D.G. ḥ.	
182 bracteátum. H.R. (<i>re.</i>) double-bracted.	—	1774.	7. 10. D.G. ḥ.	
183 ancéps. H.R. (<i>li.</i>) broad-bracted. β pálidum. H.R. (<i>li.</i>) paler-flowered.	—	1811.	9. 10. D.G. ḥ.	
184 gracile. H.R. (<i>li.</i>) slender bracteate. <i>stellátum. H.S. non DC.</i>	—	1794.	8. 11. D.G. ḥ.	
β gracilius. H.R. (<i>li.</i>) pale flat-bracted.	—	—	— D.G. ḥ.	
185 radiátum. H.R. (<i>re.</i>) radiate.	—	1732.	— D.G. ḥ.	Dill. elth. t. 197. f. 249.
186 compréssum. H.R. (<i>re.</i>) rough compressed.	—	1792.	7. 9. D.G. ḥ.	
187 páturnum. H.R. (<i>li.</i>) paler-flowered.	—	1811.	10. 11. D.G. ḥ.	
188 ásperum. H.R. (<i>ro.</i>) large rough-leav'd. α mágus. H.R. (<i>ro.</i>) broader-leaved.	—	1818.	7. 9. D.G. ḥ.	
β carulésrens. H.R. (<i>ro.</i>) bluish-leaved.	—	—	— D.G. ḥ.	
189 formósium. H.R. (<i>re.</i>) white-eyed red.	—	1820.	— D.G. ḥ.	
190 am'ena. DC. (<i>re.</i>) beautiful.	—	—	8. 9. D.G. ḥ.	
191 spectábole. H.R. (<i>pu.</i>) glaucous showy.	—	1825.	— D.G. ḥ.	
192 conspicuum. H.R. (<i>re.</i>) dark-red showy.	—	1787.	5. 8. D.G. ḥ.	Bot. mag. t. 396.
193 turbinátum. DC. (<i>re.</i>) top-shaped.	—	1806.	9. 10. D.G. ḥ.	
194 blándum. H.R. (<i>bh.</i>) fair-flowered.	—	1826.	— D.G. ḥ.	Jacq. vind. t. 476.
195 curviflórum. H.R. (<i>bh.</i>) incurving white.	—	1816.	6. 12. D.G. ḥ.	Bot. reg. t. 582.
196 àureum. H.R. (<i>or.</i>) great orange.	—	1818.	— D.G. ḥ.	
197 aurántium. H.R. (<i>or.</i>) glaucous orange.	—	1750.	3. 10. D.G. ḥ.	Bot. mag. t. 262.
198 glaécum. H.R. (<i>ye.</i>) great gold-flower'd.	—	1793.	6. 8. D.G. ḥ.	
199 cymbifolium. H.P. (<i>ye.</i>) greater boat-l'd.	—	1696.	6. 7. D.G. ḥ.	DC. plant. gras. t. 146.
200 stríctum. H.R. (<i>ye.</i>) upright boat-l'd.	—	1823. D.G. ḥ.	
201 mucroniferum. H.R. (<i>ye.</i>) mucronate boat-l'd.	—	1795. D.G. ḥ.	
202 cymbifórmæ. H.R. (<i>re.</i>) small boat-leav'd.	—	1793. D.G. ḥ.	
203 granifórme. H.R. (<i>ye.</i>) grain-leaved.	—	1727.	9. 10. D.G. ḥ.	Bradl. succ. 2. t. 20.
204 mólle. H.R. (<i>re.</i>) soft.	—	1774.	10. D.G. ḥ.	

205	coccineum. H.R. (<i>sc.</i>) scarlet-flower'd. C. B. S.	1696.	5. 9.	D.G. ḥ.	Bot. mag. t. 59.
α	obtusum. H.R. (<i>sc.</i>) blunt-calyxed.	—	—	D.G. ḥ.	DC. plant. grass. t. 83.
β	acutius. H.R. (<i>sc. pu.</i>) acute-calyxed.	—	—	D.G. ḥ.	
γ	minus. H.R. (<i>sc.</i>) lesser.	—	—	D.G. ḥ.	
206	bicolòrum. H.R. (<i>sc. ye.</i>) two-coloured.	1732.	—	D.G. ḥ.	Dill. elth. t. 202. f. 258.
α	erectum. H.R. (<i>sc. ye.</i>) tall upright.	—	—	D.G. ḥ.	
β	patulum. H.R. (<i>sc. ye.</i>) spreading.	—	—	D.G. ḥ.	
γ	minus. H.R. (<i>sc. ye.</i>) dwarf spreading.	—	—	D.G. ḥ.	
207	inæquale. H.R. (<i>ye.</i>) unequal-cupped.	1716.	—	D.G. ḥ.	Bradl. succ. 1. t. 7.
208	tenuifolium. H.R. (<i>sc.</i>) slender-leaved.	1700.	6. 9.	D.G. ḥ.	Dill. elth. t. 201. f. 256.
β	erectum. H.R. (<i>sc.</i>) upright slender-ld.	—	—	D.G. ḥ.	DC. plant. gras. t. 82.
209	variabile. H.R. (<i>ye.</i>) changeable-fld.	1796.	6. 8.	D.G. ḥ.	
α	densius. H.R. (<i>ye.</i>) rough close-leaved.	—	—	D.G. ḥ.	
β	lævius. H.R. (<i>ye.</i>) smooth distant-ld.	—	—	D.G. ḥ.	
210	spiniforme. H.R. (<i>li.</i>) thorn-shaped.	1793.	9. 10.	D.G. ḥ.	
β	subaduncum. H.R. (<i>li.</i>) greater hook-ld.	1810.	10. 11.	D.G. ḥ.	
211	curvifolium. H.R. (<i>li.</i>) great curve-ld.	1799.	10.	D.G. ḥ.	
212	flexifolium. H.R. (<i>li.</i>) lesser curve-ld.	1818.	—	D.G. ḥ.	
α	perviride. H.R. (<i>li.</i>) dark-green.	—	—	D.G. ḥ.	
β	late virens. H.R. (<i>li.</i>) apple-green.	—	—	D.G. ḥ.	
213	inconspicuum. H.P. (<i>li.</i>) inconspicuous.	1824.	6. 9.	D.G. ḥ.	
214	aduncum. H.R. (<i>li.</i>) small curve-leav'd.	1795.	2. 3.	D.G. ḥ.	
215	filicâle. H.R. (<i>li.</i>) ready-stemmed.	1800.	9.	D.G. ḥ.	
216	spinosum. H.R. (<i>pu.</i>) thorny.	1714.	6. 9.	D.G. ḥ.	Dill. elth. t. 208. f. 265.
217	prodûctum. H.P. (<i>ro.</i>) great-cupped.	1820.	6. 8.	D.G. ḥ.	
218	lépidum. H.P. (<i>wh.</i>) pretty.	1824.	7. 9.	D.G. ḥ.	
219	stipulaceum. H.R. (<i>re.</i>) tall, stipuled.	1723.	5. 8.	D.G. ḥ.	Dill. elth. f. 267 & 268.
220	corallinum. H.R. (<i>ro.</i>) coral-stemmed.	1819.	—	D.G. ḥ.	
221	Haworthii. H.R. (<i>ro.</i>) Haworth's.	1793.	6. 1.	D.G. ḥ.	
222	læve. H.R. (<i>pu.</i>) blunt white-leav'd.	1774.	7. 9.	D.G. ḥ.	
223	veruculatum. H.R. (<i>ye.</i>) long spit-leaved.	1731.	5. 6.	D.G. ḥ.	
α	Dillenii. H.R. (<i>ye.</i>) Dillenius's.	—	—	D.G. ḥ.	Dill. elth. f. 259.
β	Candollii. H.R. (<i>ye.</i>) cluster spit-leaved.	—	—	D.G. ḥ.	DC. plant. gras. t. 36.
224	purpuro-crôceum. H.R. purple saffron-fld.	1780.	8. 10.	D.G. ḥ.	
225	flavò-crôceum. H.P. yellow and saffron.	1816.	5. 9.	D.G. ḥ.	
226	luteum. H.P. (<i>ye.</i>) upright yellow.	1822.	6. 8.	D.G. ḥ.	
227	lutèolum. H.P. (<i>ye.</i>) small yellow.	—	—	D.G. ḥ.	
228	crôceum. H.R. (<i>sn.</i>) saffron-flowered.	1816.	10. 11.	D.G. ḥ.	Jacq. frag. p. 17. t. 11. f. 2.
229	crystallinum. H.R. (<i>wh.</i>) diamond.	1727.	5. 8.	D.G. ♂.	DC. plant. gras. 128.
230	glaciâle. H.R. (<i>wh.</i>) Ice-plant. Greece.	1787.	—	G.○.	
231	pinnatifidum. H.R. (<i>ye.</i>) jagged-leaved. C. B. S.	1774.	5. 10.	G.○.	Bot. mag. t. 67.
232	sessiliñorum. H.R. (<i>ye.</i>) small ice-plant.	—	7. 8.	G.○.	
α	luteum. H.R. (<i>ye.</i>) hairy-yellow.	1816.	—	G.○.	
β	álbum. H.R. (<i>wh.</i>) white-flowered.	1774.	—	D.G. ḥ.	
233	humifusum. H.R. (<i>wh.</i>) narrow-ld. ice-plant.	1823.	5. 8.	D.G. ḥ.	
234	clandestinum. H.P. (<i>wh.</i>) minute-flower'd.	—	6. 10.	D.G. ♂.	Jacq. vind. 3. t. 7.
235	Aitonii. H.R. (<i>ro.</i>) Aiton's.	1795.	5. 8.	G.○.	
236	lanceolatum. H.R. (<i>wh.</i>) spear-leaved.	—	—	G.○.	
α	álbum. H.R. (<i>wh.</i>) white-flowered.	1812.	—	G.○.	
β	rôseum. H.R. (<i>ro.</i>) rosy.	1774.	5. 9.	D.G. ♂.	DC. plant. gras. t. 102.
237	cordifolium. H.R. (<i>re.</i>) heart-leaved.	—	7. 9.	G.○.	Bot. mag. t. 540.
238	pomeridiânum. H.R. (<i>ye.</i>) great yellow.	—	—	G.○.	Andr. deposit. t. 57.
β	Andre'wsii. H.R. (<i>ye.</i>) Andrews's.	—	G.○.	DC. plant. gras. t. 135.
239	Candollii. H.R. (<i>ye.</i>) Decandolle's. helianthoides. DC. nec aliorum.	—	G.○.	
240	pilosum. H.R. (<i>ye.</i>) hairy-yellow.	1800.	6. 8.	G.○.	Breyn. cent. t. 79.
241	calendulaceum. H.R. (<i>y.</i>) hispid narrow-cup'd.	1819.	8.	G.○.	
242	flácidum. DC. (<i>ye.</i>) flaccid.	1826.	—	D.G. ♂.	
243	helianthoides. H.R. (<i>ye.</i>) spatula-leaved.	1774.	8. 10.	G.○.	
244	cuneifolium. DC. (<i>pu.</i>) transparent. límpidum. H.R.	—	7.	G.○.	Jacq. ic. 3. t. 288.
245	pyrop'æum. H.S.S. (<i>v.</i>) annual 3-colour'd.	1820.	4. 7.	G.○.	
α	rubidum. (<i>pu.</i>) the ruby.	—	—	G.○.	Bot. mag. t. 2144.
β	tricolor. W. hort. ber. t. 22.	—	—	G.○.	
γ	rôseum. H.R. (<i>ro.</i>) the rosy.	—	—	G.○.	
γ	álbum. H.R. (<i>bh.</i>) blush white.	1819.	—	G.○.	
246	villosum. H.R. (<i>bh.</i>) villous.	1759.	7.	D.G. ḥ.	

247	caducum. H.R. (<i>wh.</i>) deciduous.	C. B. S.	1774.	7. 8.	G. \circ .	
248	apétalum. H.R. (<i>wh.</i>) dwarf spreading.	_____	1774.	—	G. \circ .	Jacq. vind. 3. t. 6.
249	nodiflòrum. H.R. (<i>wh.</i>) knot-flowered.	Egypt.	1739.	8. 10.	G. \circ .	DC. plant. gras. t. 88.
250	ciliatum. H.R. (<i>wh.</i>) ciliated.	C. B. S.	1774.	—	D.G. $\frac{1}{2}$.	
251	geniculiflòrum. H.R. (<i>ye.</i>) joint-flower'd.	Egypt.	1727.	7. 9.	D.G. $\frac{1}{2}$.	DC. plant. gras. t. 17.
252	Triplònum. H.R. (<i>wh.</i>) Aster-leaved.	C. B. S.	1700.	6. 10.	D.G. $\frac{1}{2}$.	Dill. elth. t. 179. f. 220.
253	expánsum. H.R. (<i>st.</i>) houseleek-leav'd.	_____	1705.	7. 8.	D.G. $\frac{1}{2}$.	_____ f. 223.
	<i>tortuðsum.</i> DC. plant. gras. t. 94. non L.					
254	várians. H.R. (<i>wh. st.</i>) varying.	_____	1706.	7. 10.	D.G. $\frac{1}{2}$.	Pet. gaz. t. 78. f. 10.
255	tortuðsum. H.R. (<i>st.</i>) twisting-leaved.	_____	1705.	6. 10.	D.G. $\frac{1}{2}$.	Dill. elth. t. 181. f. 222.
256	pállens. H.R. (<i>wh.</i>) pale-flowered.	_____	1774.	7. 8.	D.G. $\frac{1}{2}$.	DC. plant. gras. t. 47.
	<i>expánsum.</i> DC. non L.					
257	crassicaûle. H.P. (<i>ye.</i>) thick-stalked.	_____	1822.	5.	D.G. $\frac{1}{2}$.	
258	lorátum. H.R. (<i>wh.</i>) glaucous strap-l'd.	_____	1819.	—	D.G. $\frac{1}{2}$.	
	<i>relaxátum.</i> H.R. (<i>re.</i>) livid strap-leav'd.	_____	—		D.G. $\frac{1}{2}$.	
260	anatómicum. H.R. (<i>wh.</i>) skeleton-leav'd.	_____	1803.	7. 8.	D.G. $\frac{1}{2}$.	
	β frágile. H.R. (<i>wh.</i>) brittle.	_____	—		D.G. $\frac{1}{2}$.	
261	réctum. H.R. (<i>wh.</i>) upright egg-leav'd.	_____	1819.	—	D.G. $\frac{1}{2}$.	
262	crassuloides. H.R. (<i>bh.</i>) Crassula-like.	_____	—	7. 8.	D.G. $\frac{1}{2}$.	
263	incómpactum. H.R. (<i>wh.</i>) persistent cup'd.	_____	1816.	—	D.G. $\frac{1}{2}$.	
264	spléndens. H.R. (<i>wh.</i>) shining white fld.	_____	1716.	6. 8.	D.G. $\frac{1}{2}$.	DC. plant. gras t. 35.
265	flexuòsum. H.R. (<i>bh.</i>) flexuose-leaved.	_____	1795.	7. 8.	D.G. $\frac{1}{2}$.	
266	acuminátum. H.P. (<i>wh.</i>) long-cupped.	_____	1820.	8. 9.	D.G. $\frac{1}{2}$.	
267	albicauile. H.P. (<i>wh.</i>) white-twigg'd.	_____	1823.	8. 10.	D.G. $\frac{1}{2}$.	
268	sulcátum. H.R. (<i>wh.</i>) acute-cupped.	_____	1819.	—	D.G. $\frac{1}{2}$.	
269	fastigiátum. H.R. (<i>bi.</i>) level-topped.	_____	1794.	7. 9.	D.G. $\frac{1}{2}$.	
	β reflexum. H.R. (<i>br.</i>) reflex-leaved.	_____	1792.	8. 9.	D.G. $\frac{1}{2}$.	
270	umbelliflòrum. H.R. . umbel-flowered.	_____	1825.	—	D.G. $\frac{1}{2}$.	
271	longist'ylum. DC. (<i>bh.</i>) long-styled.	_____	1822.	—	D.G. $\frac{1}{2}$.	
	<i>pállens.</i> Jacq. schœnb. 3. t. 279. non Ait. pallescens. H.R.	C. B. S.	1804.	—	D.G. $\frac{1}{2}$.	Jacq. schœnb. 3. t. 278.
	β purpurascens. DC. (<i>pu.</i>) purpleish.	_____				
272	micráthon. H.R. (<i>wh.</i>) small-blossom'd.	_____	1800.	8. 10.	D.G. $\frac{1}{2}$.	
	<i>parviflòrum.</i> Jacq. non H.R.		1821.	6. 8.	D.G. $\frac{1}{2}$.	
273	júceum. H.R. (<i>ro.</i>) rushy.	_____	1819.	—	D.G. $\frac{1}{2}$.	
274	granulicauile. H.P. (<i>ro.</i>) shagreen-stem'd.	_____	1794.	6. 10.	D.G. $\frac{1}{2}$.	
275	ténue. H.R. (<i>wh.</i>) slender rushy.	_____	1774.	8. 10.	D.G. $\frac{1}{2}$.	
276	spinulíferum. H.R. (<i>st.</i>) spinulescent.	_____	1819.	—	D.G. $\frac{1}{2}$.	
277	gróssum. H.R. (<i>bh.</i>) gouty.	_____	1794.	7. 10.	D.G. $\frac{1}{2}$.	
278	salmónéum. H.R. (<i>ro.</i>) salmon-colour'd.	_____	1774.	7. 11.	D.G. $\frac{1}{2}$.	Bot. mag. t. 326.
279	canaliculátum. H.R. (<i>ye.</i>) channel-leav'd.	_____	1824.	—	D.G. $\frac{1}{2}$.	Jacq. frag. 52. f. 3.
280	viridiflòrum. H.R. (<i>gr.</i>) green-flowered.	_____	1821.	8. 11.	D.G. $\frac{1}{2}$.	
281	tenuiflòrum. H.R. (<i>re.</i>) slender-flower'd.	_____	1790.	7. 10.	D.G. $\frac{1}{2}$.	DC. plant. gras. t. 129.
282	longispinul. H.P. (<i>st.</i>) ros'y-cupp'd yellow.	_____	1821.	8. 9.	D.G. $\frac{1}{2}$.	
283	nítidum. H.R. (<i>ye.</i>) shrubby crystalline.	_____	1774.	6. 8.	D.G. $\frac{1}{2}$.	
	<i>brachiátum.</i> DC. nec aliorum.		1816.	—	D.G. $\frac{1}{2}$.	
284	subincánum. H.P. (<i>wh.</i>) soft-leav'd white.	_____	1823.	6. 9.	D.G. $\frac{1}{2}$.	
285	brachiátum. H.R. (<i>ye.</i>) three-forked.	_____	1714.	6. 10.	D.G. $\frac{1}{2}$.	DC. plant. gras. t. 78.
286	testáceum. H.R. (<i>sn.</i>) tile-coloured.	_____	1821.	—	D.G. $\frac{1}{2}$.	
287	macrorhízum. H.P. (<i>wh.</i>) large-rooted.	_____	1823.	6. 9.	D.G. $\frac{1}{2}$.	
288	tuberòsum. H.R. (<i>co.</i>) tuberous shrubby.	_____	1714.	6. 10.	D.G. $\frac{1}{2}$.	
	β minus. H.R. (<i>bl.</i>) lesser-leaved.	_____	—		D.G. $\frac{1}{2}$.	
289	noctiflòrum. H.R. (<i>ro. wh.</i>) night-flowering.	_____	—	6. 8.	D.G. $\frac{1}{2}$.	DC. plant. gras. t. 10.
	<i>a phænicéum.</i> H.R. (<i>ro.</i>) deep ros'y.	_____	—		D.G. $\frac{1}{2}$.	Dill. elth. f. 262.
	β stramíneum. H.R. (<i>st.</i>) straw-coloured.	_____	—		D.G. $\frac{1}{2}$.	_____ f. 263.
	γ elátum. H.R. (<i>wh. ro.</i>) fall-rosy.	_____	1819.	—	D.G. $\frac{1}{2}$.	
290	fúlvum. H.R. (<i>br. wh.</i>) grey-barked.	_____	1732.	—	D.G. $\frac{1}{2}$.	
291	clavátum. DC. (<i>wh.</i>) clubbed.	_____	—		D.G. $\frac{1}{2}$.	Jacq. schœnb. t. 108.
	<i>defoliátum.</i> H.R.					
292	horizontálæ. H.R. (<i>st.</i>) spreading-leav'd.	_____	1795.	—	D.G. $\frac{1}{2}$.	
293	speciosum. H.R. (<i>cr.</i>) glittering orange.	_____	1793.	5. 10.	D.G. $\frac{1}{2}$.	
294	micans. H.R. (<i>sc.</i>) dark orange.	_____	1704.	5. 8.	D.G. $\frac{1}{2}$.	Bot. mag. t. 448.
295	maculátum. H.R. (<i>sc.</i>) spotted-stalked.	_____	1792.	—	D.G. $\frac{1}{2}$.	
296	flávum. H.R. (<i>go.</i>) small-glittering.	_____	1820.	8.	D.G. $\frac{1}{2}$.	
297	obliquum. H.R. (<i>re.</i>) bright afternoon fig —	_____	1819.	—	D.G. $\frac{1}{2}$.	
298	parvifolium. H.R. (<i>pu.</i>) small-triquetrous.	_____	—	6. 10.	D.G. $\frac{1}{2}$.	
299	brevifolium. H.R. (<i>re.</i>) short-leaved.	_____	1777.	—	D.G. $\frac{1}{2}$.	Jacq. vind. t. 477.
	<i>erigerifolium.</i> Jacq.					
300	subglobosum. H.R. (<i>re.</i>) globular.	_____	1795.	—	D.G. $\frac{1}{2}$.	
301	puveruléntum. H.R. (<i>re.</i>) powdered.	_____	1792.	5.	D.G. $\frac{1}{2}$.	

302 hispidum. H.R. (<i>pu.</i>) hispid-stalked. α <i>Dilleniæ</i> . H.R. (<i>pu.</i>) the bushy. β <i>platypetalum</i> . H.R. tall.	C. B. S.	1704. 5. 10. D.G. ḥ. 1820. — D.G. ḥ. 1792. 5. 11. D.G. ḥ. 1820. — D.G. ḥ. 1824. 6. 8. D.G. ḥ. — D.G. ḥ.	DC. plant. gras. t. 66. Dill. elth. f. 278.
303 hirtellum. H.R. (<i>re.</i>) dwarf bristly.	—	1820. — D.G. ḥ.	
304 candens. H.R. (<i>wh.</i>) trailing hispid.	—	1820. — D.G. ḥ.	
305 subcompræsum. H.P. (<i>vi.</i>) upright twiggy. β <i>minus</i> . H.P. (<i>vi.</i>) lesser.	—	1824. 6. 8. D.G. ḥ. — D.G. ḥ.	
306 floribundum. H.R. (<i>fl.</i>) crowded-flower'd.	—	1704. 5. 10. D.G. ḥ.	
307 calycinum. H.R. (<i>wh.</i>) hispid long-cup'd.	—	1819. — D.G. ḥ.	
308 striatum. H.R. (<i>std.</i>) striped bristly. α <i>rōseum</i> . H.R. (<i>std.</i>) rosy. β <i>pallens</i> . H.R. (<i>std.</i>) pale-flowered.	—	1727. — D.G. ḥ. — D.G. ḥ. — D.G. ḥ.	Dill. elth. t. 215. f. 281. — t. 215. f. 281. DC. plant. gras. t. 132.
309 attenuatum. H.R. (<i>v.</i>) slender hispid.	—	1820. — G. ḥ.	
310 tuberculatum. DC. (<i>v.</i>) hispid-leaved. <i>hispidolum</i> . H.R. α <i>fastigiatum</i> . H.R. (<i>v.</i>) level-topped. β <i>rōseum</i> . H.R. (<i>ro.</i>) rosy-flowered.	—	1814. — G. ḥ. — G. ḥ. — G. ḥ. — G. ḥ.	
311 echinatum. H.R. (<i>ye.</i>) hedge-hog. α <i>luteum</i> . (<i>ye.</i>) yellow. β <i>album</i> . (<i>wh.</i>) white-flowered.	—	1774. 7. 10. G. ḥ. — G. ḥ. — G. ḥ.	DC. plant. gras. t. 24.
312 strumosum. H.R. (<i>st.</i>) tuberous hedge-hog.	—	1820. — G. ḥ.	
313 barbatum. H.P. (<i>pu.</i>) bearded bushy.	—	1793. 5. 10. G. ḥ.	
314 stelligerum. H.P. (<i>pu.</i>) procumbent bearded.	—	1705. 6. 8. G. ḥ.	
315 bulbosum. H.P. (<i>re.</i>) bearded tuberous.	—	1820. — G. ḥ.	
316 intonsum. H.P. (<i>re.</i>) black bearded. α <i>rubicundum</i> . H.P. (<i>re.</i>) red-flowered. β <i>album</i> . H.P. (<i>bh.</i>) white-flowered.	—	7. — G. ḥ. — G. ḥ. — G. ḥ.	
317stellatum. H.P. (<i>pu.</i>) small-bearded.	—	1716. 9. 10. G. ḥ.	DC. plant. gras. t. 29.
318 densum. H.P. (<i>pu.</i>) dwarf-bearded.	—	1732. 5. 8. G. ḥ.	Bot. mag. t. 1220.
HYMENO'GYNE. H.R. HYMENO'GYNE. Icosandria Dodecagynia. glabra. H.R. (<i>st.</i>) smooth. C. B. S. 1787. 7. 8. G. ⊕. <i>Mesembryanthemum glabrum</i> . H.S. non A.R.	—		
TETRAG'ONIA. W. TETRAG'ONIA. Icosandria Pentagynia. L.	—		
1 expansa. H.S. (<i>gr.</i>) horned.	N. Zealand.	1772. 8. 9. H. ⊕.	DC. plant. gras. t. 114.
2 crystallina. H.S. (<i>gr.</i>) diamond.	Peru.	1788. 6. 12. H. ⊕.	— t. 34.
3 echinata. H.S. (<i>gr.</i>) hedgehog.	C. B. S.	1774. 5. 8. G. ♂.	— t. 113.
4 hirsuta. DC. (<i>gr.</i>) hairy.	—	1823. — G. ♂.	
5 fruticosa. H.S. (<i>gr.</i>) shrubby.	—	1712. 7. 9. G. ḥ.	
6 linearis. H.R. (<i>gr.</i>) linear-leaved.	—	1819. 9. — G. ḥ.	
7 obovata. H.R. (<i>gr.</i>) obovate-leaved.	—	— G. ḥ.	
8 decumbens. H.S. (<i>gr.</i>) trailing.	—	1758. 7. 9. G. ḥ.	DC. plant. gras. t. 23.
9 Tetrapteris. H.S. (<i>gr.</i>) wing-seeded.	—	1795. — G. ḥ.	
10 spicata. H.S. (<i>gr.</i>) spiked.	—	— G. ♀.	Comm. hort. 2. t. 102.
11 herbacea. H.S. (<i>gr.</i>) tuberous-rooted.	—	1752. 6. 7. G. ♀.	
12 purpurea. Ot. (<i>pu.</i>) purple.	—	1825. — H. ⊕.	
SESUVIUM. DC. SESUVIUM. Icosandria Tri-Pentagynia.	—		
1 pedunculatum. P.S. (<i>pu.</i>) peduncled.	W. Indies.	1692. 6. 8. D.S. ♀.	Pluk. phyt. t. 216. f. 1.
2 sessile. P.S. (<i>pu.</i>) sessile-flowered.	—	— D.S. ♀.	DC. plant. gras. t. 9.
<i>Aizoön canariense</i> . Andr. rep. t. 201. nec <i>aliorum</i> .	—		
3 longifolium. DC. (<i>pu.</i>) long-leaved.	S. America.	1816. 7. 8. D.S. ♀.	Bot. mag. t. 1701.
4 revolutifolium. DC. (<i>pu.</i>) revolute-leaved.	Cuba.	— D.S. ♀.	Rumph. amb. 5.t.72.f.1.
5 repens. DC. (<i>pu.</i>) creeping.	E. Indies.	1816. — S. ⊕.	
6 microphyllum. DC. (<i>pu.</i>) small-leaved.	Havannah.	1823. — D.S. ♀.	
AIZO'ON. DC. AIZO'ON. Icosandria Pentagynia. L.	—		
1 canariense. DC. (<i>ye.</i>) Purslane-leav'd. Canaries.	—	1731. 7. 8. D.G. ♀.	DC. plant. gras. t. 136.
2 glineoides. DC. (<i>ro.</i>) hairy.	C. B. S.	1774. 6. 8. D.G. ♀.	DC. plant. gras. t. 30.
3 hispanicum. DC. (<i>wh.</i>) Spanish.	Spain.	1728. 7. 8. H. ⊕.	
4 tomentosum. DC. (<i>ye.</i>) woolly.	C. B. S.	1824. — D.G. ḥ.	
5 stellatum. DC. (<i>ye.</i>) starry.	—	— D.G. ḥ.	
6 perfoliatum. DC. (<i>ye.</i>) perfoliate.	—	1816. 6. 8. D.G. ḥ.	Burm. afr. t. 26. f. 2.
7 sarmentosum. DC. (<i>ye.</i>) trailing.	—	— D.G. ḥ.	
8 lanceolatum. w. (<i>ye.</i>) spear-leaved.	—	1752. 8. D.G. ♂.	
GL'INUS. DC. GL'INUS. Dodecandria Pentagynia. L.	—		
lotoides. DC. (<i>ye.</i>) hairy.	S. Europe.	1788. 7. 8. H. ⊕.	Bocc. sic. 21. t. 11.

Tribus II. REAUMURIEÆ.

REAUM'URIA. DC. REAUM'URIA. Icosandria Pentagynia.

1 vermiculata. DC. (*wh.*) white-flowered. Persia. 1829. G. ḥ.

Lob. ic. t. 380.

2 hypericoïdes. DC. (pu.) purple-flowered. Syria.	1800.	7. 10.	G. ♂.	Bot. mag. t. 2057.
<i>linifolia</i> . Sal. par. lond. t. 18.				
NITR'ARIA. DC.	NITR'ARIA.	Dodecandra Trigynia.		
1 Schobèri. L. (wh.) thick-leaved.	Siberia.	1778.	5. 8. H. ♂.	Andr. rep. t. 529.
2 sibírica. F. (wh.) Siberian.		1826. H. ♂.	Gmel. sib. 2. t. 98.
3 cáspera. W.en. (wh.) Caspian.	Caspia.	1822.	5. 8. H. ♂.	Pall. ross. t. 50. f. B.
4 tridentata. DC. (wh.) tridentate.	Africa.	—	— H. ♂.	

ORDO III.

CACTEÆ. DC. prodr. v. 3. p. 457.

Tribus I. OPUNTIACEÆ. DC. prodr. v. 3. p. 458.

MAMMILL'ARIA. H.S. MAMMILL'ARIA. Icosandria Monogynia.

1 magnimámma. DC. (re.) large-beaded.	Mexico.	1823.	6. 8. D.S. ♂.	
2 coronaria. DC. (sc.) the great.	—	1820.	— D.S. ♂.	Orteg. dec. t. 16.
3 discolor. DC. (wh.re.) two-coloured.	S. America.	1800.	— D.S. ♂.	DC. diss. t. 2. f. 2.
<i>Cáctus Spinii</i> . Colla. <i>depréssus</i> . DC. cat. h. monsp. p. 84. <i>pseudo-mammillaris</i> . Salm Dyck.				
4 lanífera. DC. (re.) wool-bearing.	Mexico.	1823.	6. 8. D.S. ♂.	DC. diss. t. 4.
5 flavescens. DC. (re.) yellow-spined.	S. America.	1811.	— D.S. ♂.	
6 strámnea. H.S.s. (re.) straw-coloured.	—	—	— D.S. ♂.	
7 prolífera. DC. (wh.) white-spined.	—	1688.	7. 8. D.S. ♂.	
8 simplex. DC. (wh.) small red-spin'd. W. Indies.	—	—	D.S. ♂.	DC. plant. gras. t. 3.
<i>Cáctus mammillaris</i> . L.				
9 missouriensis. L.C. (wh.) Missouri.	Missouri.	1813.	6. 8. D.G. ♂.	
<i>Cáctus mammillaris</i> . Nuttall. <i>nec aliorum</i> .				
10 vivipara. DC. (re.) viviparous.	—	—	— D.G. ♂.	
11 parvímámma. DC. small-beaded.	S. America.	1817. D.S. ♂.	
<i>Cáctus microthélè</i> , s.s.				
12 geminispina. DC. (re.) twin-spined.	Mexico.	1823.	6. 8. D.S. ♂.	DC. diss. t. 3.
13 glomerata. DC. (re.) tufted.	St. Domingo.	1825. D.S. ♂.	Plum. ed Bur. t. 201. f. 1.
14 pusilla. DC. (bh.) hoary starry.	S. America.	1816.	6. 9. D.S. ♂.	DC. diss. t. 2. f. 1.
<i>Cáctus stellatus</i> . Lodd. bot. cab. t. 79.				
15 Helicteres. DC. (ro.) simple obovate.	Mexico.	1827.	— D.S. ♂.	DC. diss. t. 5.
16 cómica. H.S.s. cone-beaded.	—	1808.	— D.S. ♂.	
17 coccinea. Gil. (sc.) scarlet-flower'd. Chile.	—	1827.	— D.S. ♂.	
18 speciosa. Gil. (re.) handsome.	—	—	— D.S. ♂.	
19 fusca. Ot. brown.	S. America.	—	— D.S. ♂.	
20 canescens. Ot. canescent.	—	—	— D.S. ♂.	
21 caspítosa. Ot. tufted.	—	—	— D.S. ♂.	
22 chrysántha. Ot. (ye.) yellow-flowered.	—	—	— D.S. ♂.	

MELOCA'CTUS. DC. MELO THISTLE. Icosandria Monogynia.

1 communis. DC. (re.) Turk's-cap.	W. Indies.	1688.	7. 8. D.S. ♂.	DC. plant. gras. t. 112.
β oblongus. DC. (re.) oblong.	—	—	— D.S. ♂.	
γ macrocéphalus. large-beaded.	—	—	— D.S. ♂.	
2 macracáthus. DC. (re.) large-spined.	S. America.	1820.	— D.S. ♂.	Lk. et Ot. dis. t. 12.
3 pyramidális. DC. pyramidal.	Curassao.	1824. D.S. ♂.	— p. 10. t. 25.
4 placentiformis. DC. (re.) black-spined.	Brazil.	—	7. 9. D.S. ♂.	Besl. h. eyst. 4. ord. f. 1.
<i>Cáctus Melocactus</i> . Besl. non L. <i>Melocactus Besleri</i> . Lk. et Ot. dis. p. 11. t. 21.				
5 Langsdórfii. DC. (ye.) Langsdorf's.	Brazil.	1826.	7. 9. D.S. ♂.	
6 Sellowii. DC. Sellow's.	Monte Video.	— D.S. ♂.	Lk. et Ot. dis. t. 22.
7 polyacanthus. DC. many-spined.	Brazil.	— D.S. ♂.	— p. 13. t. 16. f. 1.
ECHINOCA'CTUS. DC. ECHINOCACTUS. Icosandria Monogynia.				
1 gibbosus. DC. (wh.) gibbous.	Jamaica.	1808.	6. 8. D.S. ♂.	Bot. reg. t. 137.
<i>Cáctus gibbosus</i> . B.R.				
2 latispinus. H.P. (pu.) flat-spined.	Mexico.	1823. D.S. ♂.	Lk. et Ot. dis. t. 14.
<i>cornigerus</i> . DC. dis. t. 7?				
3 crispátus. DC. (pu.) curl-ribbed.	—	1826. D.S. ♂.	DC. diss. t. 8.
4 obvallátus. DC. (pu.) guarded.	—	— D.S. ♂.	— t. 9.
5 tenuispinus. DC. (ye.) slender-spined.	Brazil.	—	6. 8. D.S. ♂.	Lk. et Ot. dis. t. 19. f. 1.
6 melocactiformis. DC. (bh.) Melocactus-like.	Mexico.	— D.S. ♂.	DC. diss. t. 10.
7 tephracáthus. DC. ashy-spined.	Brazil.	1825. D.S. ♂.	Lk. et Ot. dis. t. 14. f. 2.
8 recúrvus. DC. recurve-spined.	Mexico.	1768. D.S. ♂.	
<i>Cáctus nóbilis</i> . L.				

CACTEÆ.

9 intortus. DC. twisted.	Antigua.	1768. D.S. \ddagger .	
10 Salmiânius. DC. Salm-Dyck's.	Curassao.	1824. D.S. \ddagger .	Lk. et Ot. dis. t. 26.
11 tuberculatus. DC. tubercled.	Mexico.	1826. D.S. \ddagger .	p. 17. t. 17.
12 gladiatus. DC. sword-spined.	1824. D.S. \ddagger p. 16. t. 27.
13 subuliferus. DC. awl-spined.	S. America.	1780. D.S. \ddagger p. 15. t. 23.
14 depresso. DC. depressed.	Monte Video.	1828. D.G. \ddagger .	Lk. et Ot. dis. t. 18.
15 orthacanthus. DC. straight-spined.	S. America.	1815. D.S. \ddagger p. 15. t. 15.
16 acutatus. DC. sharp-ribbed. D.G. \ddagger .	Lk. et Ot. dis. t. 24.
17 parvispinus. DC. small-spined.	S. America.	1815. D.S. \ddagger
18 intricatus. DC. intricate.	Monte Video.	1828. D.S. \ddagger
19 meonacanthus. DC. oblong.	Jamaica. D.S. \ddagger
20 inflatus. Gil. inflated.	Chile.	1828. D.S. \ddagger
21 erinaceus. Hedgehog.	1818. D.S. \ddagger
<i>Cactus erinaceus.</i> H.S.S.					
<i>CEREUS.</i> DC. <i>CEREUS.</i> Icosandria Monogynia.					
1 multangulâris. DC. many angled.	S. America.	1815. D.S. \ddagger .	
2 redactus. DC. removed.	Mexico.	1796. D.S. \ddagger .	
<i>Cactus nobilis.</i> H.S.					
3 Scopa. DC. setaceous.	Brazil.	1826. D.S. \ddagger .	
4 sénilis. DC. grey-headed.	Mexico.	1823. D.S. \ddagger .	
<i>Cactus senilis.</i> H.P.					
5 peruviana. DC. (re.wh.)	great Peruvian.	Peru.	1728.	8. 10. D.S. \ddagger .	DC. plant. gras. t. 58.
6 monströsus. DC. (re.wh.)	irregular.	S. America.	1816.	— D.S. \ddagger .	DC. diss. t. 11.
<i>Cactus abnormis.</i> W. enum. sup. 31.					
7 monoclonos. DC. (ro.wh.)	obcordate-petal'd.	Caribees.	6. 8. D.S. \ddagger .	Plum. ed. Burm. t. 191.
8 fimbriatus. DC.	(ro.) fringe-petal'd.	St. Domingo.	1826. D.S. \ddagger t. 195. f. 1.
9 hexagonus. H.S. (ro.wh.)	great 6-angled.	Surinam.	1690.	7. 8. D.S. \ddagger .	Bradl. succ. t. 1.
10 heptagonus. H.S. (wh.)	great 7-angled.	W. Indies.	1728.	— D.S. \ddagger .	
11 Hístrix. DC. Porcupine.	1818. D.S. \ddagger .	
12 ebûneus. DC. ivory-thorned.	Curassao.	— D.S. \ddagger .	
β monströsus. DC. monströus.	Hybrid.	— D.S. \ddagger .	Bradl. succ. t. 10.
13 griseus. H.S.	(wh.) grey-spined.	S. America.	1809.	9. 10. D.S. \ddagger .	
14 crenulatus. DC. crenulate.	1822. D.S. \ddagger .	
15 albispinus. DC. white-spined.	— D.S. \ddagger .	
16 regalis. DC. the royal.	1816. D.S. \ddagger .	
17 strictus. DC. straight.	1822. D.S. \ddagger .	
18 Haworthii. DC. Haworth's.	Caribees.	1811. D.S. \ddagger .	
<i>Cactus Haworthii.</i> s.s. <i>Cereus nobilis.</i> H.S.					
19 aureus. DC. gold-spined.	S. America.	1825. D.S. \ddagger .	
20 niger. DC.	(wh.) black.	1820. D.S. \ddagger .	
21 fulvispinosus. DC. tawny-spined.	1795. D.S. \ddagger .	
22 flavispinus. DC. yellow-spined.	1822. D.S. \ddagger .	
23 chiloensis. DC. ten-angled.	Chile.	1825. D.S. \ddagger .	
24 euphorbioides. DC. Euphorbia-like.	S. America.	1815. D.S. \ddagger .	
25 Royeni. DC.	(wh.) Royen's 9-angled.	W. Indies.	1728.	6. 9. D.S. \ddagger .	DC. plant. gras. t. 143.
26 lanuginosus. DC.	(wh.) woolly.	1690.	7. 8. D.S. \ddagger .	Herm. parad. t. 115.
27 repandus. DC.	(wh.) waved short-spin'd.	1728.	8. 10. D.S. \ddagger .	Bot. reg. t. 336.
28 subrepandus. DC.	(wh.) waved long-spin'd.	1817.	— D.S. \ddagger .	
29 polygonus. DC.	(wh.) many-sided.	St. Domingo.	1826. D.S. \ddagger .	Plum. ed. Burm. t. 196.
30 rosaceus. Ot.	(ro.) rosy.	1826. D.S. \ddagger .	
31 Déppii. Ot. Deppi's.	— D.S. \ddagger .	
32 incrustatus. Ot. incrusted.	— D.S. \ddagger .	
33 grändis. DC. straddling-spin'd.	1816. D.S. \ddagger .	
34 tetragonus. DC.	(wh.) great 4-angled.	S. America.	1710.	7. 8. D.S. \ddagger .	
35 paniculatus. DC.	(wh.re.) paniced.	St. Domingo.	1827. D.S. \ddagger .	Plum. ed. Burm. t. 192.
36 obtusus. DC. blunt 3-angled.	Brazil.	1819. D.S. \ddagger .	
37 serpentinus. DC.	(pu.wh.) white-spined.	S. America.	1817.	7. 8. D.S. \ddagger .	Lk. et Ott. abb. t. 91.
38 ambiguum. DC.	(pu.wh.) ambiguous.	1827.	— D.S. \ddagger .	Bot. mag. t. 17.
39 flagelliformis. DC.	(ro.) creeping.	Peru.	1690.	3. 6. D.G. \ddagger .	
40 caudatus. Gil.tailed.	Chile.	1828. D.G. \ddagger .	
41 horizontalis. Gil. horizontal.	— D.G. \ddagger .	
42 grandiflorus. DC.	(ye.wh.) night-flowered.	Jamaica.	1700.	6. 8. D.S. \ddagger .	Andr. deposit. t. 508.
43 Colvillii. Colvill's hybrid.	Hybrid.	1825. D.S. \ddagger .	
44 pentagonus. DC.	(wh.) five-angled.	S. America.	1769.	7. 8. D.S. \ddagger .	
45 réptans. H.S.S. creeping.	1813. D.S. \ddagger .	
46 alisetosus. H.S.S. white-spin'd trails.	St. Domingo.	1816. D.S. \ddagger .	
47 rádicans. DC. rooting-branch'd.	S. America.	1827. D.S. \ddagger .	
48 humilius. DC. slender dwarf.	— D.S. \ddagger .	
49 quadrangularis. DC.	(wh.) quadrangular.	S. America.	1809.	7. 9. D.S. \ddagger .	Plum. ed. Bur. t. 199. f. 1.

50	trípteris. DC. broad-lobed.	1827.	D.S. $\frac{1}{2}$.	
51	bifróns. H.S.S rooting 4-angular.	St. Domingo. 1818.	D.S. $\frac{1}{2}$.	
52	speciosissimus. DC. (sc. pu.)	beautiful.	Mexico.	1816.	6. 9. D.G. $\frac{1}{2}$.	Bot. reg. t. 486.
	Cáctus speciosissimus.	B.R. speciosus.	Colla hort. rip. t. 10.			
53	triangularis. DC. (wh.)	great triangular.	W. Indies.	1690.	7. 8. D.S. $\frac{1}{2}$.	Bot. mag. t. 1884.
54	trigonous. DC. (wh.)	small triangular.	—	1809.	—	D.S. $\frac{1}{2}$.
55	triqueter. DC. least triangular.	S. America.	1794.	D.S. $\frac{1}{2}$.
56	coccineus. DC. (sc.)	scarlet-flowered.	Brazil.	1828.	D.S. $\frac{1}{2}$.
57	Smithianus.	(sc.) Smith's hybrid.	Hybrid.	1824.	6. 9. D.G. $\frac{1}{2}$.	
58	extensus. DC. extended.	—	1826.	D.S. $\frac{1}{2}$.
59	squamulosus. DC. (ye.)	scaly.	Brazil.	—	D.S. $\frac{1}{2}$.
60	prismáticus. DC. prismatic.	—	—	D.S. $\frac{1}{2}$.
61	setáceus. DC. bristly.	Brazil.	1828.	D.S. $\frac{1}{2}$.
62	myosurus. DC. mouse-tail.	—	—	D.S. $\frac{1}{2}$.
63	tenuis. DC.	(ro.) slender-branched.	—	1826.	D.S. $\frac{1}{2}$.
64	tenuispinus. H.P. slender-spined.	Mexico.	—	D.S. $\frac{1}{2}$.
65	phyllanthoides. DC. (pi.)	rosy-flowered.	—	1816.	5. 8. D.G. $\frac{1}{2}$.	Bot. mag. t. 2092.
	Cáctus speciosus.	Bot. reg. t. 304.	Epiphyllum phyllanthoides.	H.R.		
66	Jenkinsoni.	(sc.) Mr. Jenkinson's.	Hybrid.	1824.	6. 9. D.G. $\frac{1}{2}$.	
67	Ackermannii. H.R. (sc.)	Ackermann's.	Mexico.	1824.	5. 7. D.G. $\frac{1}{2}$.	
68	truncatus. DC.	(cr.) truncated.	Brazil.	1818.	7. 11. D.S. $\frac{1}{2}$.	Bot. reg. t. 696.
	Cáctus truncatus.	Bot. mag. t. 2092.	Epiphyllum truncatum.	H.P.		
69	alatus. DC.	(gr.wh.) wing-stemmed.	Jamaica.	1818.	8. 11. D.S. $\frac{1}{2}$.	Bot. mag. t. 2820.
70	Hookeri. H.P.	(wh.) Hooker's.	S. America.	—	7. 8. D.S. $\frac{1}{2}$.	— t. 2692.
	Cáctus Phyllanthus.	B.M. nec aliorum.	Epiphyllum Hookeri.	H.P.		
71	Phyllanthus. DC. (wh.)	long-tubed.	S. America.	1710.	6. 7. D.S. $\frac{1}{2}$.	DC. plant. gras. t. 145.
72	gracilis. DC. slender.	—	1826.	D.S. $\frac{1}{2}$.
73	imbricatus. H.R. imbricated.	—	1820.	D.S. $\frac{1}{2}$.
74	cylindricus. H.S. cylindrical.	Peru.	1799.	D.G. $\frac{1}{2}$.
OPUNTIA. DC.		INDIAN-FIG. Icosandria Monogynia.				
1	foliosa. DC.	(ye.) greater Pin-pillow.	S. America.	1805.	6. 7. D.S. $\frac{1}{2}$.	
	pusilla. H.S. non DC.					
2	curassavica. DC.	(ye.) Pin-pillow.	Curassao.	1690.	—	D.S. $\frac{1}{2}$.
	α major. H.S.	(ye.) larger.	—	—	—	D.S. $\frac{1}{2}$.
	β media. H.S.	(ye.) middle.	—	—	—	D.S. $\frac{1}{2}$.
	γ minor. H.S.	(ye.) smaller.	—	—	—	D.S. $\frac{1}{2}$.
	δ longa. H.R.	(ye.) longer.	Brazil.	1820.	—	D.S. $\frac{1}{2}$.
3	frágilis. DC.	(ye.) brittle Pin-pillow.	Missouri.	1814.	D.F. $\frac{1}{2}$.
4	pusilla. DC.	(ye.) small Pin-pillow.	S. America.	1826.	D.S. $\frac{1}{2}$.
5	polyacantha. H.S.S. (st.)	many-spined.	Missouri.	1814.	7. 9. D.F. $\frac{1}{2}$.	
	missouriensis. DC.	Cáctus ferrox. N.				
6	média. H.S.S.	(ye.) lesser many-spined.	—	—	—	D.F. $\frac{1}{2}$.
7	ferox. DC.	(ye.) ferocious.	S. America.	1817.	D.S. $\frac{1}{2}$.
8	spinossissima. DC. cluster-spined.	Jamaica.	1732.	7. D.S. $\frac{1}{2}$.	
9	húmilia. H.S.	(ye.) humble.	W. Indies.	1795.	7. 8. D.S. $\frac{1}{2}$.	
	hórrida. DC.					
10	Dilleniæ. DC.	(ye.) Dillenius's.	S. America.	1731.	9. 10. D.S. $\frac{1}{2}$.	Bot. reg. t. 255.
	Cáctus Dilleniæ. B.R.	Dill. elth. 2. t. 296. f. 382.				
11	Tuna. DC.	(ye.) yellow-spined.	—	—	—	D.S. $\frac{1}{2}$.
12	polyántha. DC.	(ye.) many-flowered.	—	1811.	7. 9. D.S. $\frac{1}{2}$.	Dill. elth. t. 295. f. 380.
13	monacantha. DC.	(ye.) one-spined.	—	1816.	—	D.S. $\frac{1}{2}$.
14	elatior. DC.	(ye.) great black-spin'd.	—	1731.	7. 8. D.S. $\frac{1}{2}$.	Dill. elth. t. 204. f. 379.
15	nigricans. DC.	(ye.) lesser black-spined.	—	1795.	—	D.S. $\frac{1}{2}$.
	Cáctus Tuna y nigricans. B.M.					Bot. mag. t. 1557.
16	máxima. DC.	(ye.) great oblong.	S. America.	1768.	7. 9. D.S. $\frac{1}{2}$.	
17	triacantha. DC.	(ye.) three-spined.	—	1796.	—	D.S. $\frac{1}{2}$.
18	decumâna. H.R. great.	—	1820.	D.S. $\frac{1}{2}$.
19	elongata. H.S.S. various-spined.	—	1816.	D.S. $\frac{1}{2}$.
20	cochinillifera. DC.	(re.) Cochineal-fig.	—	1688.	7. 9. D.S. $\frac{1}{2}$.	Andr. rep. t. 523.
	Cáctus cochinillifer.	Bot. mag. t. 2741-2742.				
21	Ficus-Indica. DC.	(ye.) Indian Fig.	—	1731.	7. 8. D.S. $\frac{1}{2}$.	
22	pseudo-Tuna. DC.	(ye.) bastard.	—	—	—	D.S. $\frac{1}{2}$.
23	tomentosa. DC. woolly-branch'd.	—	1824.	D.S. $\frac{1}{2}$.
24	sericea. Gil. silky.	Chile.	1827.	D.S. $\frac{1}{2}$.
25	lanceolata. DC.	(ye.) spear-shaped.	—	1796.	7. 9. D.S. $\frac{1}{2}$.	
26	tuberculata. DC.	(ye.) tubercled.	—	1818.	—	D.S. $\frac{1}{2}$.
27	inérmis. DC.	(ye.) upright spineless.	—	1796.	—	D.S. $\frac{1}{2}$.
	stricta. H.S.					DC. plant. gras. n. 138. ic.
28	Amyclæa. DC.	(ye.) Neapolitan.	Naples.	1825.	—	D.F. $\frac{1}{2}$.

CACTEÆ.

29 subinermis. L.en. (<i>ye.</i>) nearly smooth	S. America.	1824.	7. 9. D.S. h.	
30 rubescens. DC. red-stemmed.	Brazil.	1828. D.S. h.	
31 crassa. DC. (<i>ye.</i>) thick-lobed.	Mexico.	1817.	6. 9. D.G. h.	
32 longispina. Ot. (<i>ye.</i>) long-spined.	S. America.	1824. D.S. h.	
33 leucantha. Ot. (<i>wh.</i>) white-flowered.	—	— D.S. h.	
34 pulvinata. DC. pulvinate.	Mexico.	1827. D.S. h.	
35 Hernándezii. DC. (<i>re.</i>) Hernandez's.	—	— D.G. h.	Hern. mex. 78. ic.
36 vulgaris. DC. (<i>ye.</i>) common.	America.	1596.	7. 9. D.F. h.	Mill. dict. t. 191.
Cactus Opuntia. Bot. mag. t. 2393.				
β minor. (<i>ye.</i>) lesser.	N. America.	1814.	— D.F. h.	
Cactus Opuntia. N.				
37 brasiliensis. DC. (<i>gr.</i>) round-stemm'd.	Brazil.	1817. D.S. h.	Pis. bras. p. 100. f. 2.
PERE'SKIA. DC. PERE'SKIA. Icosandria Monogynia.				
1 aculeata. DC. (<i>wh.</i>) Barbadoes Gooseby. W. Indies.	1696.	10. 11. S. h.		Dill. elth. t. 227. f. 294.
Cactus Pereskia. L.				
2 longispina. H.s.s. (<i>wh.</i>) long-spined.	S. America.	1808. S. h.	
3 portulacæfolia. DC. (<i>pu.</i>) Purslane-leav'd. W. Indies.	1820. S. h.		Plum. ed Bur. t. 197. f. 1.
4 grandifolia. DC. great-leaved.	Brazil.	1818. S. h.	

Tribus II. RHIPSALIDEÆ. DC. prodr. v. 3. p. 475.

RHIPS'ALIS. DC.	RHIPS'ALIS. Dodecandra Monogynia.			
1 Cass'ytha. DC. (<i>wh.</i>) pendulous.	W. Indies.	1758.	7. 10. S. h.	Gært. sem. 1. t. 28. f. 1.
2 Hookeriana. DC. (<i>wh.</i>) four-petal'd.	—	—	— S. h.	Hook. ex. flor. t. 2.
3 fasciculata. DC. (<i>wh.</i>) cluster-branch'd. S. America.	1800.	—	— S. h.	
4 parasitica. DC. (<i>cr.</i>) parasitical.	W. Indies.	—	— S. h.	Plum. ed Bur. t. 197. f. 2.
5 funalis. DC. (<i>wh.</i>) great-flowered.	S. America.	1816.	7. 8. S. h.	Bot. mag. t. 2740.
grandiflorus. B.M. Cactus funalis. s.s.				
6 salicornoides. DC. (<i>ye.</i>) Glass-wort-like. W. Indies.	1818.	5. 8. D.S. h.		Bot. mag. t. 2461.
7 mesembryanthoides. DC. Fig Marygold-like.	—	—	— D.S. h.	

ORDO CIV.

GROSSULARIEÆ. DC. prodr. v. 3. p. 477.

RIBES. DC.	CURRENT. Pentandria Monogynia. L.			
1 oxyacanthoides. DC. (<i>st.</i>) Hawthorn-leav'd. N. America.	1705.	4. 5. H. h.		Dill. elth. t. 139. f. 166.
2 ferox. DC. (<i>st.</i>) ferocious.	California.	1827.	— H. h.	Lodd. bot. cab. t. 884.
3 lacustre. DC. (<i>br.</i>) swamp.	N. America.	1812.	— H. h.	
4 Menziesii. DC. (<i>cr.</i>) Menzies'.	—	1827.	— H. h.	
5 speciosum. DC. (<i>pu.</i>) beautiful.	—	—	— H. h.	
6 uva-crispa. R.S. (<i>st.</i>) smooth Gooseberry. England.	—	—	— H. h.	
7 reclinatum. R.S. (<i>st.</i>) procumbent.	Germany.	1683.	— H. h.	Eng. bot. v. 29. t. 2057.
8 Grossularia. R.S. (<i>st.</i>) rough Gooseberry. England.	—	—	— H. h.	Eng. bot. v. 12. t. 1292.
9 caucasicum. DC. (<i>st.</i>) Caucasian.	Caucasus.	1820.	— H. h.	
10 setosum. B.R. (<i>st.</i>) bristly.	Missouri.	1827.	5. H. h.	Bot. reg. t. 1237.
11 hirtellum. DC. (<i>st.</i>) hairy.	N. America.	1812.	4. 5. H. h.	
12 gracie. DC. (<i>st.</i>) slender.	—	—	— H. h.	
13 Cynosbati. DC. (<i>gr.</i>) prickly-fruited.	Canada.	1759.	4. H. h.	Schmidt arb. t. 98.
14 triflorum. DC. (<i>br.</i>) three-flowered.	N. America.	1812.	4. 5. H. h.	Lodd. bot. cab. t. 1094.
15 Dikuscha. F. (<i>st.</i>) Fischer's.	Siberia.	1821.	— H. h.	
16 Stakori. F. (<i>st.</i>) Siberian.	—	—	— H. h.	
17 orientale. DC. (<i>st.</i>) oriental.	Syria.	1820.	— H. h.	
18 saxatile. DC. (<i>br.</i>) rock.	Siberia.	1828.	— H. h.	
19 diacantha. DC. (<i>st.</i>) two-spined.	—	1781.	5. 6. H. h.	Pall. ross. 2. t. 66.
20 alpinum. DC. (<i>st.</i>) Alpine.	Britain.	—	4. 5. H. h.	Eng. bot. v. 10. t. 704.
α sterile. DC. (<i>st.</i>) barren.	—	—	— H. h.	
β baciferum. DC. (<i>st.</i>) berry-bearing.	—	—	— H. h.	
21 resinosum. DC. (<i>st.</i>) clammy.	N. America.	1800.	— H. h.	Bot. mag. t. 1583.
22 procumbens. DC. (<i>br.</i>) trailing.	Dahuria.	1804.	5. 6. H. h.	Pall. ross. 2. t. 65.
23 multiflorum. DC. (<i>st.</i>) many-flowered.	Hungary.	1818.	5. H. h.	Bot. mag. t. 2368.
24 spicatum. DC. (<i>br.</i>) acid.	England.	—	— H. h.	Eng. bot. v. 18. t. 1290.

25 <i>rùbrum</i> . DC.	(<i>st.</i>) common red.	Britain.	4. 5. H. $\frac{1}{2}$.	Eng. bot. v. 18. t. 1289.
α <i>sylvestre</i> . DC.	(<i>st.</i>) wild.	—	— H. $\frac{1}{2}$.	
β <i>hortense</i> . DC.	(<i>st.</i>) cultivated.	— H. $\frac{1}{2}$.	
γ <i>cárneum</i> . DC.	(<i>st.</i>) pale-fruited.	— H. $\frac{1}{2}$.	
δ <i>variegátum</i> . DC.	(<i>st.</i>) stripe-fruited.	— H. $\frac{1}{2}$.	
ϵ <i>álbum</i> . DC.	(<i>st.</i>) white-fruited.	— H. $\frac{1}{2}$.	Berl. mem. t. 2. f. 15.
26 <i>pétř æum</i> . DC.	(<i>re.</i>) rock.	England.	5. 6. H. $\frac{1}{2}$.	Eng. bot. v. 10. t. 705.
27 <i>cérèum</i> . B.R.	(<i>wh.</i>) waxy.	Columbia.	1827.	4. 5. H. $\frac{1}{2}$.	Bot. reg. t. 1263.
28 <i>atropurpureum</i> . Led.	dark-purple.	Siberia.	— H. $\frac{1}{2}$.	
29 <i>carpáthicum</i> . R.S. (<i>st.</i>)	Carpathian.	Carpathian.	1816.	4. 5. H. $\frac{1}{2}$.	
30 <i>triste</i> . DC.	(<i>re.ye.</i>) dark-coloured.	Siberia.	1818.	— H. $\frac{1}{2}$.	
31 <i>nigrum</i> . DC.	(<i>wh.</i>) black-fruited.	Britain.	— H. $\frac{1}{2}$.	Eng. bot. v. 18. t. 1291.
32 <i>trífidum</i> . DC.	(<i>pu.</i>) trifid-calyxed.	N. America.	1828.	— H. $\frac{1}{2}$.	
33 <i>glandulòsum</i> . DC.	(<i>st.</i>) glandular.	Chile.	1820.	5. 6. H. $\frac{1}{2}$.	Flor. per. t. 232. f. b.
34 <i>prostrátum</i> . DC.	(<i>pu.</i>) prostrate.	N. America.	1777.	4. 5. H. $\frac{1}{2}$.	L'Herit. stirp. 1. t. 2.
<i>glandulòsum</i> . H.K. non R.P.					
35 <i>Biebersteini</i> . DC. (<i>st.</i>)	Bieberstein's.	Caucasus.	1820.	— H. $\frac{1}{2}$.	
<i>caucásicum</i> . M.B. nec aliorum.					
36 <i>rigens</i> . DC.	(<i>st.</i>) rigid.	N. America.	1812.	— H. $\frac{1}{2}$.	
37 <i>albinérvium</i> . DC.	(<i>st.</i>) white-nerved.	Canada.	1828.	— H. $\frac{1}{2}$.	
38 <i>laxiflòrum</i> . DC.	(<i>st.</i>) loose-flowered.	N. America.	1812.	— H. $\frac{1}{2}$.	
39 <i>viscosissimum</i> . DC. (<i>ye.</i>) viscous.		—	1827.	— H. $\frac{1}{2}$.	
40 <i>sanguíneum</i> . DC.	(<i>cr.</i>) crimson.	—	1820.	5. 6. H. $\frac{1}{2}$.	
41 <i>flòridum</i> . DC.	(<i>st.</i>) Pensylvanian.	—	1729.	4. 5. H. $\frac{1}{2}$.	Dill. elth. t. 244. f. 315.
42 <i>aùreum</i> . DC.	(<i>ye.</i>) golden.	Missouri.	1812.	4. 6. H. $\frac{1}{2}$.	Bot. reg. t. 125.
43 <i>longiflòrum</i> . L.C.	(<i>ye.</i>) long-flowered.	N. America.	1818.	— H. $\frac{1}{2}$.	
44 <i>tenuiflòrum</i> . B.R. (<i>ye.</i>)	slender-flower'd.	—	1824.	— H. $\frac{1}{2}$.	Bot. reg. t. 1274.
α <i>melanocárpa</i> .	(<i>ye.</i>) black-fruited.	—	—	— H. $\frac{1}{2}$.	
β <i>chrysocárpa</i> .	(<i>ye.</i>) yellow-fruited.	—	—	— H. $\frac{1}{2}$.	

ORDO CV.

ESCALLONEÆ. Kth. *synops.* v. 3. p. 364.

ESCALLONIA. L. ESCALLONIA. Pentandria Monogynia. L.

1 <i>rùbra</i> . R.S.	(<i>re.</i>) red-flowered.	Chile.	1826.	8. 10. G. $\frac{1}{2}$.	Bot. mag. t. 2890.
<i>Stereóxylon rùbrum</i> . Flor. per. v. 3. t. 236. f. b.					
2 <i>glandulòsa</i> . B.C. (<i>wh.</i>)	glandular.	Chile.	—	— G. $\frac{1}{2}$.	Lodd. bot. cab. t. 1291.
3 <i>floribúnda</i> . K.S. (<i>wh.</i>)	many-flowered.	New Granada.	1827. G. $\frac{1}{2}$.	
4 <i>caracasâna</i> . K.S. (<i>wh.</i>)	Caracas.	Caracas.	— S. $\frac{1}{2}$.	
ANO'PTERUS. B.P.	ANO'PTERUS.	Hexandria Monogynia.			
<i>glandulòsa</i> . B.P. (<i>wh.</i>) glandular.					
V. Diem. Isl. 1823.					
.... G. $\frac{1}{2}$.					
Lab. nov. holl. 1. t. 112.					

ORDO CVI.

SAXIFRAGEÆ. Juss. *gen.* 308.

SAXI'FRAGA. L. SAXIFRAGE. Decandria Dignynia. L.

1 <i>crassifòlia</i> . L.T.	(<i>ro.</i>) thick-leaved.	Siberia.	1765.	3. 5. H. $\frac{1}{4}$.	Bot. mag. t. 126.
<i>Bergénia crassifòlia</i> . Mön. <i>Mégæsea crassifòlia</i> . H.R.					
2 <i>cordifòlia</i> . L.T.	(<i>ro.</i>) heart-leaved.	—	1779.	— H. $\frac{1}{4}$.	
3 <i>média</i> .	(<i>ro.</i>) intermediate.	—	1819.	— H. $\frac{1}{4}$.	
<i>Mégæsea média</i> . H.R.					
4 <i>ligulàta</i> . L.T.	(<i>wh.</i>) fringed-leaved.	Nepaul.	1818.	— H. $\frac{1}{4}$.	Swt. br. fl. gar. 59.
<i>Mégæsea ciliàta</i> . H.R.					
5 <i>Géum</i> . L.T.	(<i>sp.</i>) kidney-leaved.	Ireland.	6. 7. H. $\frac{1}{4}$.	Eng. bot. v. 22. t. 1561.
6 <i>hirsùta</i> . L.T.	(<i>sp.</i>) hirsute.	—	5. 6. H. $\frac{1}{4}$.	— v. 33. t. 2322.
7 <i>polita</i> . L.en.	(<i>sp.</i>) polished.	—	— H. $\frac{1}{4}$.	
<i>Robertsónia polita</i> . H.S.					

8 dentata. L.en. (sp.) toothed. <i>Robertsónia dentata</i> . H.S.	Ireland.	5. 6. H. ȝ.	
9 crenata. (sp.) crenated. <i>Robertsónia crenata</i> . H.S. 1790.	— H. ȝ.	
10 sphæroïdea. (sp.) sphæroid. <i>Robertsónia sphæroïdea</i> . H.S.	Pyrenees? 1800.	— H. ȝ.	
11 punctata. L. (sp.) dotted-flower'd. Siberia.	1699.	— H. ȝ.	
12 serrata. L.en. (sp.) sawed-leaved. <i>Robertsónia serrata</i> . H.S.	Ireland.	— H. ȝ.	
13 umbrosa. L.T. (sp.) London pride.	Britain.	4. 6. H. ȝ.	Eng. bot. v. 10. t. 633.
14 cuneifolia. L.T. (sp.) wedge-leaved.	Switzerland. 1768.	5. 6. H. ȝ.	W. et K. hung. 1. t. 44.
15 hybrida. L.en. (sp.) hybrid. 1820.	— H. ȝ.	Sternb. sax. 17. t. 8. f. 3.
16 stellaris. L.T. (wh.) starry. δ <i>Schleicheria</i> . L.T. (wh.) <i>Schleicher's.</i> <i>Spatularia stellaris</i> . γ <i>depilata</i> . H.R.	Britain.	6. 7. H. ȝ.	Eng. bot. v. 3. t. 167.
17 leucanthemifolia. (wh.) Stock-leav'd.	N. America. 1812.	— H. ȝ.	Lap. pyr. sax. t. 25.
18 sarmentosa. L.T. (wh.) sarmentose.	China. 1771.	— F. ȝ.	Bot. mag. t. 92.
Diptera sarmentosa. Bork.	<i>Ligularia sarmentosa</i> . H.R.		
19 cuscuteæformis. B.C. (wh.) dodder-like.	China. 1815.	2. 7. F. ȝ.	Lodd. bot. cab. t. 186.
<i>Ligularia minor</i> . H.R.			
20 erosa. L.T. (wh.) jagged-leaved.	N. America. 1812.	5. 6. H. ȝ.	
Aulaxis nuda. H.R.			
β hirsuta. L.T. (wh.) hairy.	— — —	— H. ȝ.	
Aulaxis micranthifolia. H.R.			
21 rotundifolia. L.T. (wh.) round-leaved.	Austria. 1596.	— H. ȝ.	Bot. mag. t. 424.
<i>Miscopetalum rotundifolium</i> . H.R.			
β repanda. L.T. (wh.) repand.	— — —	— H. ȝ.	
22 granulata. L.T. (wh.) grain-rooted.	Britain.	5. H. ȝ.	Sternb. saxif. t. 5.
β multiplex.	(wh.) double-flowered.	— — —	Eng. bot. v. 7. t. 500.
23 bulbifera. L.T. (wh.) bulbiferous.	S. Europe.	6. 7. H. ȝ.	Sternb. sax. t. 12. f. 1.
24 cérnua. L.T. (wh.) drooping.	Scotland.	7. H. ȝ.	Eng. bot. v. 10. t. 66
<i>Lobelia cernua</i> . H.R.			
25 sibirica. L.T. (wh.) Siberian.	Siberia. 1802.	5. 6. H. ȝ.	Sternb. sax. p. 23. t. 25.
26 rivularia. L.T. (wh.) Alpine brook.	Scotland.	6. 7. H. ⊖.	Eng. bot. v. 32. t. 2275.
<i>Lobelia rivularia</i> . H.R.			
27 hederaea. L.T. (wh.) Ivy-leaved.	Levant. 1752.	7. H. ⊖.	Flor. græc. t. 379.
28 Hirculus. L.T. (ye.) yellow marsh.	England.	8. H. ȝ.	Eng. bot. v. 15. t. 1009.
<i>Hirculus ranunculoides</i> . H.R.			
29 flagellaria. L.T. (ye.) flagelliferous.	Baffin's Bay. 1820.	— H. ȝ.	Sternb. sax. p. 25. t. 6.
30 aizoïdes. L.T. (ye.) smaller mountain.	Britain.	7. 8. H. ȝ.	— t. 8. f. 1.
β autumnalis. (ye.) larger mountain.	— — —	— H. ȝ.	Eng. bot. v. 1. t. 39.
<i>Leptasea aizoïdes</i> . H.R.			
31 bronchiialis. L.T. (st.) spiny-fringed.	Siberia. 1819.	— H. ȝ.	Gmel. sib. 4. t. 65. f. 2.
32 tenella. L.T. (wh.) slender.	Carinthia. 1824.	— H. ȝ.	Jacq. coll. 3. t. 17.
33 juniperina. L.T. (ye.) Juniper-like.	Caucasus. 1826.	— H. ȝ.	Sternb. sax. p. 31. t. 10.
34 áspera. L.T. (st.) rough.	Switzerland. 1752.	8. H. ȝ.	Jacq. aust. 5. t. 31.
<i>Ciliaria aspera</i> . H.R.			
35 bryoides. L.T. (st.) thread-moss like.	— — —	6. 7. H. ȝ.	Jacq. misc. 2. t. 5. f. 1.
36 hieracifolia. L.T. (br.) Hawkweed-lid.	Hungary. 1789.	5. 6. H. ȝ.	W. et K. hung. 1. t. 18.
37 pensylvánica. L.T. (gr.) Pennsylvanian.	N. America. 1732.	— H. ȝ.	Dill. elt. t. 253. f. 328.
β glabra. (gr.) smoother-leaved.	— — —	— H. ȝ.	
38 semipubescens. L.T. (st.) semipubescent.	— — —	— H. ȝ.	
<i>Micranthes semipubescens</i> . H.S.			
39 virginianæ. L.T. (wh.) Virginian.	— — —	1790. 5. 7. H. ȝ.	Bot. mag. t. 1664.
<i>Dermasea pilosa</i> . H.R.			
40 congesta. H.S.L. (wh.) close-headed.	— — —	1812. — H. ȝ.	
nivalis. Ph. nec aliorum.			
41 nivalis. L.T. (wh.) clustered Alpine.	Britain.	6. 7. H. ȝ.	Eng. bot. v. 7. t. 440.
<i>Dermasea nivalis</i> . H.R.			
42 melaleuca. s.s. (wh.) Fischer's.	Siberia. 1826.	— H. ȝ.	
43 longiscapa. L.T. (wh.) long-scaped.	— — —	1824. 5. 7. H. ȝ.	
44 davurica. L.T. (wh.) Davurian.	Davuria. — — —	— H. ȝ.	
45 Cotyledon. L.T. (wh.) pyramidal.	Alps Europe. 1596.	5. 7. H. ȝ.	
<i>Chondrosea pyramidalis</i> . H.R.			
46 intermedia. (wh.) related. 1808.	7. H. ȝ.	
<i>Chondrosea intermedia</i> . H.R.			
47 lingulata. L.T. (sp.) tongue-leaved.	Pyrenees.	5. 7. H. ȝ.	
longifolia. s.s.			
β media. L.T. (sp.) middle.	— — —	— H. ȝ.	Sternb. sax. p. 1. t. 1. a.

48 crustatæ. L.en.	(sp.) crusted.	Pyrenees.	5. 7.	H. ¼.	Sternb. sax. t. 1. b.	
49 Aizooñ. L.T.	(sp.) large margined.	1731.	—	H. ¼.	Jacq. aust. t. 438.	
50 recta. P.S.	(sp.) straight-leaved.	Alps Europe.	—	H. ¼.	Lap. pyr. sax. t. 15.	
intacta. L.T. W. hort. ber. 2. t. 75.	<i>Chondrosea Aizoñ.</i> H.R.					
β minor. H.R.	(sp.) lesser.	Alps Europe.	—	H. ¼.		
γ pareiflōs. H.R.	(sp.) smaller-flowered.	—	—	H. ¼.		
51 mutata. L.T.	(ye.) saffron-colour'd.	Switzerland.	1779.	6. 7.	H. ¼.	
52 aretioidæ. L.T.	(ye.) Aretia-like.	—	1825.	—	H. ¼.	
53 Burseriâna. L.T.	(st.) Burser's.	Italy.	—	5. 6.	H. ¼.	
54 retusa. L.T.	(ro.) purple-flower'd.	Pyrenees.	—	3. 4.	H. ¼.	
55 oppositifolia. L.T. (re.) opposite-leaved.	Britain.	—	—	H. ¼.	Eng. bot. v. 1. t. 9.	
Antiphyllea cærulea. H.R.						
56 biflora. L.T.	(re.) two-fl'd purple.	Switzerland.	1825.	—	H. ¼.	
57 cæsia. L.T.	(st.) grey.	—	1752.	5. 6.	H. ¼.	
58 diapensioïdes. L.T. (wh.) Diapensia-like.	—	—	1828.	—	H. ¼.	
59 compacta. s.s.	(wh.) compact.	—	—	—	H. ¼.	
60 androsacea. L.T. (wh.) Androsace-leav'd.	—	—	1792.	—	H. ¼.	
61 sedoides. L.T.	(st.) Stonecrop-like.	Europe.	1825.	—	H. ¼.	
62 ténera. L.T.	(st.) flat-leaved.	Switzerland.	—	5. 6.	H. ¼.	
planifolia. s.s.						
63 geranioidæ. L.S. (wh.) cranesbill-leav'd.	Pyrenees.	1770.	4. 5.	H. ¼.	Lap. pyr. sax. t. 43.	
64 irrigua. L.T.	(wh.) crowfoot-leav'd.	Siberia.	1820.	—	H. ¼.	
65 pedatifida. L.T. (wh.) pedatifid.	Scotland.	—	5. 6.	H. ¼.	M.B. cent. pl. rar. t. 73.	
β minor. H.R.	(wh.) smaller villous.	—	—	H. ¼.	Eng. bot. v. 32. t. 2278.	
γ ladanifera. L.T. (wh.) ladaniferous.	Pyrenees.	—	—	H. ¼.	Lap. sax. pyr. t. 42.	
66 quinquefida. H.R. (wh.) quinquefid.	Scotland.	—	—	H. ¼.	Bot. mag. t. 1651.	
67 ceratophyllea. L.T. (wh.) shining-calyx'd.	Spain.	1804.	—	H. ¼.	Lap. pyr. sax. t. 31.	
68 ajugifolia. L.T.	(wh.) Bugle-leaved.	Pyrenees.	1770.	6. 7.	H. ¼.	
69 affinis. L.T.	(wh.) affined.	Ireland.	—	—	H. ¼.	
70 pentadactyla. L.T. (wh.) five-fingered.	Pyrenees.	1815.	5. 6.	H. ¼.	Lap. pyr. sax. t. 40.	
71 latifida. L.T.	(wh.) broad-cleft.	Spain.	1826.	—	H. ¼.	
72 decipiens. L.T. (wh.) palmate.	Wales.	—	—	H. ¼.	Sternb. sax. p. 55. t. 23.	
palmata. Eng. bot. v. 7. t. 455. cæspitosa. Flor. dan. t. 71.						
73 hirta. L.T.	(wh.) hairy.	Ireland.	—	—	H. ¼.	
74 platypétala. L.T. (wh.) broad-petaled.	Scotland.	—	6.	H. ¼.	Eng. bot. v. 32. t. 2291.	
75 Schradéri. s.s.	(wh.) Schrader's.	Siberia.	1826.	—	H. ¼.	— v. 32. t. 2276.
76 incurvifolia. L.T. (wh.) incurved-leav'd.	Ireland.	—	—	H. ¼.		
77 denudata. L.T. (wh.) half-naked.	Scotland.	—	—	H. ¼.		
78 Sternbergii. L.T. (wh.) Sternberg's.	Germany.	—	5. 6.	H. ¼.	Sternb. sax. t. 24.	
79 pulchélla. L.T.	(st.) pretty.	—	1818.	—	H. ¼.	
80 tridentata. L.T. (wh.) three-toothed.	—	—	—	—		
81 cæspitosa. L.T.	(st.) tufted.	Wales.	—	—	H. ¼.	
82 exarata. L.T.	(wh.) furrowed.	Pyrenees.	1818.	—	H. ¼.	
nervosa. Lap. pyr. sax. t. 39.					Eng. bot. v. 12. t. 794.	
83 pedemontana. L.T. (wh.) Piedmont.	Piedmont.	1824.	—	H. ¼.	Vill. delph. 4. t. 45.	
cymosa. Pl. rar. hung. t. 88. heterophyllea. Sternb. sax. t. 20. f. 1-2.					All. ped. t. 21. f. 6.	
84 moschata. L.T.	(ye.) musky.	Piedmont.	—	5. 6.	H. ¼.	
85 muscoïdes. L.T.	(st.) Moss-like.	England.	—	—	H. ¼.	
<i>Muscaria muscoïdes.</i> H.R.					Jacq. misc. 2. t. 21. f. 2.	
86 pygmæa. L.T.	(st.) pigmy.	Scotland.	—	—	Lap. pyr. sax. t. 35.	
moschata. E.B. non Jacq. <i>Muscaria cæspitosa.</i> H.R.					Eng. bot. v. 33. t. 2214.	
87 tridactylites. E.B. (bh.) Rue-leaved.	Britain.	—	3. 5.	H. ⊖.	Eng. bot. v. 7. t. 501.	
<i>Tridactylites annua.</i> H.R.						
88 controvérsa. s.s. (wh.) confused.	N. Europe.	—	—	H. ⊖.	Sternb. sax. t. 16.	
89 petræa. L.T.	(wh.) rock.	Carinthia.	1752.	4. 5.	H. ⊖.	
90 adscendens. L.T. (wh.) ascending.	Pyrenees.	—	5. 6.	H. ¼.	Jacq. ic. 1. t. 81.	
aquatica. Lap. pyr. sax. t. 28.					Sternb. sax. t. 19. f. 1.2.	
91 hypnoïdes. L.T. (wh.) mossy.	Britain.	—	4. 6.	H. ¼.	Eng. bot. v. 7. t. 454.	
β viscosa. L.T. (wh.) viscous.	—	—	—	H. ¼.		
γ angustifolia. L.T. (wh.) narrow-leaved.	—	—	—	H. ¼.		
δ ruscosa. L.T. (wh.) tufted.	—	—	—	H. ¼.		
ε pulchella. L.T. (wh.) larger.	—	—	—	H. ¼.		
ζ spatulata. H.R. (wh.) spatulate.	—	—	—	H. ¼.		
η latifida. H.R. (wh.) broad-cleft.	—	—	—	H. ¼.		
θ rhodopétala. H.R. (bh.) rosy-petaled.	—	—	—	H. ¼.		
ι recurvula. H.R. (wh.) recurring-cup'd.	—	—	—	H. ¼.		
κ aristata. H.R. (wh.) awned.	—	—	—	H. ¼.		
λ septifida. H.R. (wh.) seven-cleft.	—	—	—	H. ¼.		
92 condensata. L.T. (wh.) close.	Scotland.	—	—	H. ¼.	Gmel. fl. bad. t. 3.	

93 elongélla. L.T. (wh.) long peduncled. Scotland.	5. 6. H. ȝ.	Eng. bot. v. 32. t. 2277.
94 leptophylla. L.T. (wh.) narrow-leaved. Wales.	— H. ȝ.	
β angustifida. L.T. (wh.) narrow-lobed. —	— H. ȝ.	
95 lætevirens. L.T. (wh.) light-green. Scotland.	— H. ȝ.	
96 lanceolata. H.R. (wh.) spear-leaved. Europe. 1808.	— H. ȝ.	
obtusa. H.R. (wh.) blunt-leaved. —	1820. — H. ȝ.	
97 altifida. H.R. (wh.) deep-cleft. —	— H. ȝ.	
98 viscósa. H.R. (wh.) viscous. —	— H. ȝ.	
99 quinquedéns. H.R. (wh.) five-toothed. —	— H. ȝ.	
100 Haworthii. H.S.L. (wh.) Haworth's. —	— H. ȝ.	
hirta. H.R. non E.B.	— H. ȝ.	
101 æstivalis. f. (wh.) summer. Altay. 1821. 5. 7. H. ȝ.		
HEUCHERA. L. ALLUM-ROOT. Pentandria Digynia. L.		
1 americana. L. (gr.) viscid. N. America. 1656. 5. 7. H. ȝ.		Schkuhr hand. t. 58.
viscidia. Ph.		
2 pubescens. Ph. (bh.) pubescent. — 1812. — H. ȝ.		
3 hispida. Ph. (gr.) hispid. — 1820. — H. ȝ.		
4 villosa. M. (gr.) villous. — 1812. — H. ȝ.		
5 caulescens. Ph. (wh.) caulescent. — — H. ȝ.		
6 glabra. R.S. (gr.) smooth. Altay. 1827. — H. ȝ.		
TIARELLA. L. TIARELLA. Decandria Digynia. L.		
1 cordifolia. Ph. (wh.) heart-leaved. N. America. 1731. 4. 5. H. ȝ.		
2 Menziesii. Ph. (wh.) leafy-stemmed. — 1812. — H. ȝ.		
3 trifoliata. Ph. (wh.) three-leaved. N. America. 1826. — H. ȝ.		
4 polyphylla. D.P. (wh.) many-leaved. Nepaul. 1818. — H. ȝ.		
ASTILBE. D.P. ASTILBE. Oct-Decandria Digynia.		
1 decandra. D.P. (st.) decandrous. Carolina. 1812. 5. 6. F. ȝ.		Vent. malm. t. 54.
Tiarëlla bibernata. v.		
2 rivularis. D.P. (st.) rivulet. Nepaul. 1825. — H. ȝ.		
TELLIMA. B. TELLIMA. Decandria Digynia.		
grandiflora. B.R. (st. re.) large-flowered. N. America. 1826. 4. 5. H. ȝ.		Bot. reg. t. 1178.
Mitella grandiflora. Ph.		
MITELLA. L. MITE'LLA. Decandria Digynia. L.		
1 diphylla. w. (wh.) two-leaved. N. America. 1731. 4. 5. H. ȝ.		
2 cordifolia. w. (wh.) heart-leaved. — 1812. — H. ȝ.		
3 reniformis. Ph. (wh.) kidney-leaved. — 1758. 6. 8. H. ȝ.		
nuda. w.		
4 prostrata. M. (wh.) prostrate. — 1812. 5. 6. H. ȝ.		
5 pentandra. B.M. (wh.) pentandrous. — 1827. 6. 7. H. ȝ.		
CHRYSOSPLENIUM. L. GOLDEN SAXIFRAGE. Oct-Decandria Digynia.		
1 alternifolium. w. (ye.) alternate-leav'd. Britain. 4. 5. H. ȝ.		
2 oppositifolium. w. (ye.) opposite-leaved. — — H. ȝ.		
3 nepalense. D.P. (ye.) Nepaul. Nepaul. 1824. — H. ȝ.		
ADOXA. L. MOSCHATEL. Oct-Decandria Tetragynia.		
Moschatellina. E.B. (st.) tuberous. Britain. 3. 5. H. ȝ.		
HYDRA'NGEA. L. HYDRA'NGEA. Decandria Digynia. L.		
1 arboréscens. w. (wh.) shrubby. Virginia. 1736. 7. 8. H. ȝ.		Eng. bot. v. 7. t. 453.
vulgaris. Ph.		
2 cordata. Ph. (wh.) heart-leaved. Carolina. 1806. — H. ȝ.		Bot. mag. t. 437.
3 radiata. w. (wh.) white-leaved. — 1786. — H. ȝ.		
nivea. Wats. dend. brit. t. 43.		
4 heteromalla. D.P. (wh.) woolly-leaved. Nepaul. 1823. — H. ȝ.		
5 querifolia. w. (wh.) Oak-leaved. Florida. 1803. 6. 9. H. ȝ.		
6 hortensis. w. (pi.) changeable. China. 1788. 4. 9. H. ȝ.		
β cærulea. (bl.) blue-flowered. — — H. ȝ.		

ORDO CVII.

CUNONIACEÆ. Brown.

WEINMA'NNIA. L. WEINMA'NNIA. Octandria Digynia. L.		
1 glabra. L. (wh.) smooth. W. Indies. 1818. 5. 7. S. ȝ.		Lam. ill. t. 313. f. 1.
2 hirta. w. (wh.) hairy. — — S. ȝ.		
3 elliptica. K.S. (wh.) elliptic-leaved. Peru. 1823. G. ȝ.		
4 ovata. K.S. (wh.) oval-leaved. — — G. ȝ.		Cav. ic. 6. t. 566.

CUN'ONIA. L.	CUN'ONIA. Decandria Digynia. L.				
capénsis. w. (wh.) Cape.	C. B. S. 1816.	5. 7. G. b.			Bot. reg. t. 828.
CALLIC'OMA. A.R.	CALLIC'OMA. Dodecandria Digynia.				
serratifolia. A.R. (st.) saw-leaved.	N. S. W. 1793.	5. 8. G. b.			Andr. deposit. t. 566.
CERATOP'ETALUM. Sm.	CERATOP'ETALUM. Decandria Monogynia.				
gummiferum, Sm. (ye.) gum-bearing.	N. Holland. 1820.	G. b.		Sm. n. holl. 1. t. 3.
BAU'ERA. Sal.	BAU'ERA. Polyandria Digynia. H.K.				
1 rubiæfolia. Sal. (re.) Madder-leaved. N. S. W. 1793.	7. 12. G. b.				Sal. ann. bot. 1. t. 10.
rubioïdes. A.R. 198. B.M. 715.					
2 hùmilib. L.en. (re.) dwarf.	—	1804.	—	G. b.	Lodd. bot. cab. t. 1197.

ORDO CVIII.

UMBELLIFERÆ. Juss. gen. 218.

Subordo I. ORTHOSPERMEÆ. DC.

Tribus I. HYDROCOTYLINEÆ.—Subtribus I. *Hydrocotyleæ*. DC.

HYDROCO'TYLE. S.S. MARSH-PENNYWORT. Pentandria Digynia. L.

1 vulgàris. s.s. (bh.) common.	Britain.	5. 6. H.w. Ȑ.	Eng. bot. v. 11. t. 751.
2 umbellata. s.s. (wh.) umbelled.	N. America. 1795.	7. 8. H.w. Ȑ.	Spreng. umb. 1. t. 1.
3 bonariensis. s.s. (ye.) Buenos Ayres.	Buenos Ayres. 1828.	— F.w. Ȑ.	Cavan. icon. 5. t. 488.
4 americana. s.s. (gr.) American.	N. America. 1790.	5. 8. H.w. Ȑ.	Spreng. um.p.3.t.2.f.3.
5 villosa. s.s. (gr.) villous.	C. B. S. 1795.	8. G.w. Ȑ.	
6 nepalensis. H.E.F. (gr.) Nepaul.	Nepaul. 1818.	2. 6. G.w. Ȑ.	Hook. ex. flor. t. 30.

hispida. D.P.

7 nitidula. s.s. (gr.) glossy.	Java. 1821.	— S.w. Ȑ.	Hook. ex. flor. t. 29.
8 canariensis. (gr.) Canary.	Canaries. 1828.	4. 6. G.w. Ȑ.	
9 prolifera. Ot. (gr.) proliferous.	Brazil. 1824.	— S.w. Ȑ.	
10 repanda. s.s. (wh.) Pilewort-leav'd. N. America.	1806.	6. 7. G.w. Ȑ.	Spreng. umb. t. 2. f. 4.
11 asiatica. s.s. (wh.) thick-leaved. Asia.	1690.	7. 8. G.w. Ȑ.	Rheed. mal. 10. t. 46.
12 sibthorpioides. s.s. (wh.) Sibthorpioides. Mauritius.	1818.	— S.w. Ȑ.	

ERIG'ENIA. N. ERIG'ENIA. Pentandria Digynia.

bulbosa. N. (wh.) bulbous.	N. America. 1825.	3. 5. H. Ȑ.	
----------------------------	-------------------	-------------	--

DIDI'SCUS. DC. DIDI'SCUS. Pentandria Digynia.

1 incisa. DC. (wh.) cut-leaved.	N. S. W. 1820.	6. 8. H. ⊖.	Linn. trans. 10. t. 21.f.2.
Trachymène incisa. L.T.			

2 azurea. DC. (bl.) blue-flowered.	—	1827.	7. 9. H. ⊖.	Bot. mag. t. 2875.
cærulea. B.M. Trachymène cærulea.	Bot. reg. t. 1225.			

TRACHYM'ENE. DC. TRACHYM'ENE. Pentandria Digynia.

1 compressa. s.s. (st.) flat-stalked.	N. Holland. 1824.	5. 6. G. Ȑ.	Lab. n. holl. 1. t. 101.
2 linearis. s.s. (ye.) linear-leaved.	N. S. W. —	— G. b.	Cavan. icon. 5. t. 485.
Azorella linearifolia. Cav.			

3 ovata. s.s. (st.) oval-leaved.	—	1827.	— G. b.	Lab. n. holl. 1. t. 100.
Azorella ovata. Labill.				

ASTROTRICHA. DC. ASTROTRICHA. Pentandria Digynia.

floccosa. DC. (wh.) woolly.	N. S. W. 1825.	5. 7. G. b.	
-----------------------------	----------------	-------------	--

XANTH'OSIA. DC. XANTH'OSIA. Pentandria Digynia.

1 pilosa. L.T. (wh.) hairy.	N. S. W. 1826.	— G. Ȑ.	Linn. trans. 10. t. 22.f.1.
-----------------------------	----------------	---------	-----------------------------

BOWL'ESIA. S.S. BOWL'ESIA. Pentandria Digynia.

ténera. s.s. (wh.) slender.	MonteVideo. 1827.	6. 8. H. ⊖.	
-----------------------------	-------------------	-------------	--

Subtribus II. *Mulineæ*. DC.

SPANA'NTHE. S.S. SPANANTHE. Pentandria Digynia. S.S.

paniculata. s.s. (wh.) panicled.	Caracas. 1795.	7. 8. S. ♂.	Jacq. ic. 2. t. 350.
----------------------------------	----------------	-------------	----------------------

Hydrocotyle Spananthe. w.

Poz'oA. Poz'oA. Pentandria Digynia.			
-------------------------------------	--	--	--

coriacea. s.s. (wh.) leathery-leaved. Peru.	1827.	7. 9. F. Ȑ.	
---	-------	-------------	--

Tribus II. SANICULEÆ. DC.

ACTIN'OTUS. Lab. ACTIN'OTUS. Pentandria Digynia.

Heliánthi. Lab. (wh.) larger.	N. S. W. 1821.	6. 12. G. b.	Bot. reg. t. 654.
-------------------------------	----------------	--------------	-------------------

Eriocalia mäjor. Ex. bot. t. 80.

SANICULA. S.S.	SANICLE.	Pentandria Digynia. L.				
1 europæa. s.s. (wh.) wood.	Britain.	6. 7. H. ȝ.	Eng. bot. v. 2. t. 98.		
2 canadensis. s.s. (wh.) Canadian.	Canada.	1800.	— H. ȝ.			
3 marilandica. s.s.(wh.) Maryland.	N. America.	1765.	— H. ȝ.	Jacq. ic. 2. t. 348.		
HACQU'ETIA. DC.	HACQU'ETIA.	Pentandria Digynia.				
Epipactis. DC. (ye.) dwarf.	S. Europe.	1818.	5. 7. H. ȝ.	Jacq. aust. 5. t. 11.		
Astrantia Epipactis. w. Dondia Epipactis. s.s.						
ASTRA'NTIA. S.S.	ASTRA'NTIA.	Pentandria Digynia. L.				
1 major. w. (bh.) great black.	Alps Europe.	1596.	5. 9. H. ȝ.	Exot. bot. 2. t. 76.		
2 carnolica. w. (bh.) Carnolian.	Carniola.	1812.	5. 6. H. ȝ.	Jacq. aust. 5. app.t.10.		
3 caucasica. R.S. (bh.) Caucasian.	Caucasus.	1824.	— H. ȝ.			
intermedia. M.B.						
4 maxima. Pall. (bh.) Hellebore-leav'd. —	1804.	6. 7. H. ȝ.	Bot. mag. t. 1553.			
helleborifolia. P.L. 60. heterophylla. M.B.						
5 minor. w. (wh.) small.	Switzerland.	1686.	5. 6. H. ȝ.	Sm. exot. bot. 2. t. 77.		
ERY'NGIUM. S.S.	ERY'NGO.	Pentandria Digynia. L.				
1 aquatum. s.s. (wh.) marsh.	N. America.	1699.	7. 9. H. ȝ.	Bot. reg. t. 372.		
2 longitolum. s.s. (wh.) long-leaved.	Mexico.	1812.	— F. ȝ.	Cav. ic. 6. t. 555.		
3 gramineum. s.s. (bh.) grass-leaved.	New Spain.	1825.	7. 10. F. ȝ.	Laroch. eryng. t. 26.		
4 bromelifolium.s.s.(st.) Bromelia-leav'd.	—	1822.	— F. ȝ.	t. 28.		
5 monocéphalum. s.s.(pu.) one-headed.	Mexico.	1824.	— F. ȝ.	Cavan. icon. 6. t. 553.		
6 ebracteatum. s.s.(wh.) bractless.	Monte Video.	1810.	7. 9. F. ȝ.	Laroch. eryng. t. 32.		
7 ternatum. s.s. (pu.) ternate-leaved.	Crete.	— F. ȝ.	Moris. his. s. 7.t.36.f.24.		
8 planum. s.s. (bh.) flat-leaved.	Europe.	1596.	— H. ȝ.	Jacq. aust. 4. t. 391.		
9 tricuspidatum. s.s.(pu.) trifid.	Spain.	1699.	9. H. ♂.	Laroch. eryng. t. 9.		
10 asperifolium. s.s.(wh.) rough-leaved.	1824.	6. 8. H. ȝ.	t. 11.		
11 dichotomum. s.s. (bh.) forked.	S. Europe.	—	7. 9. H. ȝ.	Desf. atl. 1. t. 55.		
12 cœruleum. M.B. (bh.) blue.	Caucasus.	1816.	— H. ȝ.			
13 giganteum. s.s. (bh.) gigantic.	Armenia.	1823.	— H. ȝ.			
14 alpinum. s.s. (bh.) Alpine.	Switzerland.	1597.	— H. ȝ.			
15 spina-alba. DC. (wh.) white-spined.	Europe.	1816.	— H. ȝ.			
16 virgatum. s.s. (bh.) oval-leaved.	N. America.	1810.	6. 7. H. ȝ.			
17 gracie. s.s. (ye.) slender.	New Spain.	1818.	7. 9. F. ȝ.			
18 Baldwini. s.s. (bh.) small-headed.	Carolina.	1824.	— H. ȝ.			
19 aquifolium. s.s. (bh.) holly-leaved.	Spain.	1827.	— H. ȝ.			
20 corniculatum.s.s.(bh.) horned.	Portugal.	1803.	6. 8. H. ȝ.			
21 tenui. s.s. (bh.) slender annual.	—	— H. ♂.			
22 pusillum. s.s. (bh.) dwarf.	Spain.	1640.	— H. ȝ.			
23 virginianum. s.s. (pa.) Virginian.	Virginia.	7. 9. H. ȝ.			
24 serratum. s.s. (wh.) serrated.	New Spain.	1823.	8. 10. F. ȝ.			
25 Carlinæ. s.s. (bh.) Carline-like.	—	1827.	— F. ȝ.			
26 fœtidum. s.s. (wh.) stinking.	W. Indies.	1714.	— S. ♂.			
27 maritimum. s.s. (bh.) sea Holly.	Britain.	7. 10. H. ȝ.			
28 triquetrum. s.s. (bh.) triquetrous.	Sicily.	1827.	— H. ȝ.			
29 campestre. s.s. (wh.) field.	Britain.	7. 8. H. ȝ.			
30 Bourgatii. s.s. (bh.) cut-leaved.	S. France.	1731.	6. 8. H. ȝ.			
31 amethystinum.s.s.(bh.)amethystine.	Syria.	1648.	7. 8. H. ȝ.			
32 dilatatum. s.s. (bh.) dilated.	Portugal.	— H. ȝ.			
33 crinitum. s.s. (bh.) fringe-leaved.	Spain.	1826.	— H. ȝ.			
34 comosum. s.s. (pa.) tufted.	New Spain.	1818.	— F. ȝ.			
35 azureum. R.S. (bh.) azure blue.	—	— H. ȝ.	Laroch. eryng. t. 7.		

Tribus III. AMMINEÆ. DC.

R'UMIA. H.U.	R'UMIA.	Pentandria Digynia.				
1 taurica. H.U. (wh.) Taurian.	Tauria.	1820.	7. 8. H. ȝ.	Hoff. umb. p.173. f. 17.		
Câchrys taûrica. s.s.						
2 sibirica. F. (wh.) Siberian.	Siberia.	1826.	— H. ȝ.			
3 microcárpa. H.U.(wh.) small-fruited.	Caucasus.	1827.	— H. ȝ.	Moris. his. s. 9. t. 1.f. 3.		
4 athamanthoïdes.H.U.(wh.)Athamanta-like.	Siberia.	—	— H. ȝ.			
5 seseloides. H.U. (wh.) Seseli-like.	Tauria.	—	— H. ȝ.	Hoff.umb.ed.2.t.tit.f.4.		
CICUTA. S.S.	COWBANE.	Pentandria Digynia. L.				
1 virôsa. s.s. (wh.) long-leaved.	Britain.	7. H. ȝ.	Eng. bot. v. 7. t. 479.		
2 davûrica. s.s. (wh.) Daurian.	Siberia.	1824.	— H. ȝ.	Gmel. sib. 1. t. 47.		
3 maculâta. s.s. (wh.) spotted.	N. America.	1759.	7. 8. H. ȝ.	Pluk. alm. t. 76. f. 1.		
4 bulbifera. s.s. (wh.) bulb-bearing.	—	1823.	— H. ȝ.			
5 sinensis. R.S. (wh.) Chinese.	China.	1826.	— H. ȝ.			

ZIZIA. DC.	ZIZIA. Pentandria Digynia.						
aurea. DC.	(ye.) golden.	N. America.	1699.	5. 6.	H. 4.		
<i>Smyrnium aureum</i> . w.	<i>Thaspium aureum</i> .	N.	<i>Sison aureus</i> . s.s.				
'APIUM. DC.	CELERY. Pentandria Digynia. L.						
1 gravéolens. s.s.	(wh.) common.	Britain.	6. 8.	H. ♂.		
α fistulosum.	(wh.) hollow-stalked.	——	——	H. ♂.		
β sólidum.	(wh.) solid-stalked.	——	——	H. ♂.		
γ rúbrum.	(wh.) red-stalked.	——	——	H. ♂.		
δ napáceum. Mill.	(wh.) Turnep-rooted.	——	H. ♂.		
PETROSEL'INUM. DC.	PASLEY. Pentandria Digynia. L.						
1 sativum. K.U.	(gr.) common.	Sardinia.	1548.	6. 7.	H. ♂.		
β críspum.	(gr.) curled-leaved.	——	——	H. ♂.		
2 románum. Zuc.	(gr.) large.	Greece.	——	H. ♂.		
3 fractophyllum. Lag. (st.)	broken-leaved.	1825.	——	H. ♂.		
4 prostratum. s.s.	(gr.) prostrate.	V. Diem. Isl.	1818.	——	G. ♂.		
5 peregrinum. K.U. (wh.)	Lovage-like.	Portugal.	1633.	6. 7.	H. ♂.		
<i>Ligusticum peregrinum</i> . Jacq.	<i>Sison peregrinum</i> , R.S.						
6 ségetum. K.U.	(bh.) corn.	England.	8.	H. ⊙.		
<i>Sison ségetum</i> . E.B.							
TR'INIA. H.U.	TR'INIA. Pentandria Digynia. L.en.						
1 Hoffmánni. M.B. (wh.)	Hoffman's.	England.	5. 6.	H. 4.		
<i>Pimpinélla dioica</i> . E.B.							
2 Henníngii. M.B. (wh.)	Henning's.	Hungary.	1803.	6. 8.	H. 4.		
<i>Pimpinélla glauca</i> . w.							
3 Kitaibeli. M.B. (wh.)	Kitaibel's.	——	1827.	——	H. 4.		
4 ramosíssima. F.	(wh.) branching.	Siberia.	——	——	H. 4.		
HELOSCIADIUM. K.U.	HELOSCIADIUM. Pentandria Digynia.						
1 lateriflórum. K.U. (wh.)	narrow-leaved.	S. Europe.	1824.	7. 8.	H. ⊙.		
<i>Sison Ammi</i> . Jacq.	<i>Pimpinélla lateriflóra</i> . L. enum.						
2 inundátum. K.U. (wh.)	inundated.	Britain.	5. 6.	H.w. ⊙.		
<i>Sison inundátum</i> . E.B.	<i>Siùm inundátum</i> . E.F.	<i>Mèum inundátum</i> . s.s.					
3 nodiflórum. K.U. (wh.)	procumbent.	Britain.	7. 8.	H.w. 4.		
<i>Siùm nodiflórum</i> . E.B.							
4 repens. K.U.	(wh.) creeping.	——	6.	H.w. 4.		
5 bulbósom. K.U. (wh.)	bulb-bearing.	Switzerland.	1826.	6. 8.	H.w. 4.		
DISCOPLE'URA. DC.	DISCOPLE'URA. Pentandria Digynia.						
Nuttállii. DC.	(wh.) slender-leaved.	N. America.	1812.	7. 8.	H. ⊙.		
<i>A'mmi capilláceum</i> . M.	<i>Sison capilláceus</i> . s.s.						
LEPTOCA'ULIS. DC.	LEPTOCA'ULIS. Pentandria Digynia.						
divaricátus. DC. (ro.wh.)	spreading.	N. America.	1812.	6. 8.	H. ⊙.		
<i>Æthusa divaricata</i> . N.	<i>A'mmi divaricatum</i> . P.S.						
PTYCH'OTIS. K.U.	PTYCH'OTIS. Pentandria Digynia.						
1 heterophylla. K.U. (wh.)	Coriander-leav'd.	Pyrenees.	1778.	7. 8.	H. ♂.		
<i>Mèum heterophyllum</i> . s.s.	<i>Æthusa Bünius</i> . w.						
2 ammóides. K.U.	(wh.) Milfoil-leaved.	S. Europe.	1759.	6. 7.	H. ⊙.		
<i>Séseli ammóides</i> . s.s.							
3 verticillátæ. DC.	(wh.) whorled.	——	1825.	——	H. ⊙.		
4 cótica. DC.	(wh.) Egyptian.	Egypt.	1773.	——	H. ⊙.		
<i>Trachyspérmum cóticum</i> . L. enum.	<i>A'mmi cóticum</i> , L.	<i>Bünium cóticum</i> . R.S.					
FALC'ARIA. DC.	FALC'ARIA. Pentandria Digynia.						
agréstis.	(wh.) decurrent.	Europe.	1726.	7. 8.	H. 4.		
<i>Siùm Falcària</i> . R.S.	<i>Drepanophyllum agréstis</i> . K.U.						
SISON. S.S.	SISON. Pentandria Pentagynia. L.						
1 Amómum. E.B.	(wh.) hedge.	Britain.	7. 8.	H. ⊙.		
SCHU'L'TZIA. S.S.	SCHU'L'TZIA. Pentandria Digynia.						
crinita. s.s.	(wh.) slender-leaved.	Altay.	1828.	6. 7.	H. ♂.		
A'MMI. S.S.	BISHOP'S WEED. Pentandria Digynia. L.						
1 mágus. s.s.	(wh.) great.	S. Europe.	1551.	6. 7.	H. ⊙.		
2 Bœberi. L.en.	(wh.) Bœber's.	Crete.	1818.	7. 8.	H. 4.		
<i>cicutae'folum</i> . R.S.							
3 Visnágæ. s.s.	(wh.) Toothpick.	S. Europe.	1596.	6. 8.	H. ⊙.		
<i>Visnágæ daucoides</i> . G.							
4 glucifolium. s.s. (wh.)	glaucous-leav'd.	France.	1816.	——	H. ♂.		
ÆGOP'ODIUM. L.	GOUT-WEED. Pentandria Digynia. L.						
Podagrària. E.B. (wh.)	common.	Britain.	5. 7.	H. 4.		
<i>Sison Podagrària</i> . s.s.							
CARUM. K.U.	CARAWAY. Pentandria Digynia. L.						
1 Cárví. s.s.	(wh.) common.	Britain.	6.	H. ♂.		
<i>Féniculum Cárví</i> . L.en.							

UMBELLIFERÆ.

2	Bulbocastanum. k.u. (wh.)	Earth-nut.	Europe.	1797.	6. 7. H. 4.	Flor. dan. t. 220.
	<i>Bùniūm Bulbocastanum.</i> w. <i>My'rrhis Bùniūm.</i> R.S.					
3	flexuosum.	(wh.) common.	Britain.	5. 6. H. 4.	Eng. bot. v. 14. t. 988.
	<i>Bùniūm flexuosum.</i> E.B.					
4	verticillatum. k.u.	(wh.) whorl-leaved.	—	7. 8. H. 4.	Eng. bot. v. 6. t. 395.
	<i>Sium verticillatum.</i> E.B.					
B'UNIUM. DC.	B'UNIUM.	Pentandria Digynia. L.				
1	peucedanoïdes. M.B. (st.)	Sulphur-wort.	Caucasus.	1823.	7. 8. H. 4.	
2	luteum. M.B.	(ye.) yellow.	Volga.	1827.	— H. 4.	
3	virésens. DC.	(gr.) greenish-flower'd.	Europe.	1826.	— H. 4.	
	<i>Sium virésens.</i> R.S.					
4	acaulē. M.B.	(ye.) stalkless.	Caucasus.	1821.	6. 8. H. 5.	
	<i>A'mmi acaule.</i> s.s.					
CRYPTOT'ENIA. DC.	CRYPTOT'ENIA.	Pentandria Digynia.				
1	canadénsē. DC.	(wh.) Canadian.	N. America.	1699.	7. 8. H. 4.	Moris. his. s. 9. t. 11.
	<i>Sison canadénsē.</i> w. <i>My'rrhis canadénsis.</i> s.s.					
2	Thomasiil. DC.	(wh.) Thomas's.	Italy.	1827.	— H. 4.	
PIMPINE'LLA. K.U.	BURNET SAXIFRAGE.	Pentandria Digynia. L.				
1	mágnā. s.s.	(wh.) great.	England.	6. 8. H. 4.	Eng. bot. v. 6. t. 408.
2	média. P.S.	(wh.) middle.	Germany.	— H. 4.	Hoff. deut. fl. t. 397.
	<i>orientális.</i> Jacq. aust. t. 397.					
3	dissécta. w.	(wh.) cut-leaved.	France.	— H. 4.	Retz. obs. 3. t. 2.
4	rotundifólia. M.B. (wh.)	round-leaved.	Caucasus.	1816.	— H. 5.	Spr. Ani.ed. 2.t.4.f.1.4.
	<i>Sison rotundifolius.</i> s.s.					
5	Saxifraga. w.	(wh.) common.	Britain.	— H. 4.	Eng. bot. v. 6. t. 407.
6	Trágium. DC.	(wh.) Columna's.	S. Europe.	1824.	6. 8. H. 4.	Column. phyt. t. 17.
	<i>Trágium Colámnæ.</i> s.s.					
7	peregrina. w.	(wh.) nodding.	Italy.	1640.	— H. 4.	Jacq. vind. 2. t. 131.
8	aromática. M.B.	(wh.) aromatic.	Caucasus.	1824.	— H. 5.	
9	bubonóides. P.S. (wh.)	Bubon-like.	Portugal.	1814.	6. 8. H. 4.	
	<i>Trágium Brotéri.</i> R.S.					
10	villōsa. Scho.	(wh.) villous.	Barbary.	1824.	— H. 4.	
	<i>Trágium Schousb'ei.</i> R.S.					
11	taúrica. DC.	(wh.) Taurian.	Tauria.	1824.	6. 8. H. 4.	
	<i>Ledebüria taúrica.</i> L.en. <i>Trágium taúricum.</i> s.s.					
12	nigra. W.en.	(wh.) black-rooted.	Germany.	1683.	— H. 4.	Moris. sect. 9. t. 5.
13	lateriflóra. L.en.	(wh.) side-flowering.	Hungary.	1816.	— H. 4.	
14	dichótoma. w.	(wh.) dichotomous.	Spain.	1798.	— H. 4.	
15	A'nisum. L.	(wh.) Anise.	Egypt.	1551.	6. 8. H. ⊖.	Moris. his. s. 9. t. 9.
	<i>Sison A'nisum.</i> R.S. <i>A'nisum vulgare.</i> Gaert.					
S'IM. K.U.	WATER-PARSNEP.	Pentandria Digynia. L.				
1	latifólium. s.s.	(wh.) broad-leaved.	Britain.	7. 8. H. 4.	Eng. bot. v. 3. t. 204.
2	lancifólium. s.s.	(wh.) spear-leaved.	Siberia.	1823.	— H. 4.	Gmel. sib. 1. t. 47.
3	angustifólium. s.s. (wh.)	narrow-leaved.	Britain.	— H. 4.	Eng. bot. v. 2. t. 139.
4	Sísum. s.s.	(wh.) Skirret.	China.	1548.	— H. 4.	Schkuhr. hand. 1. t. 69.
BUPLE'URUM. S.S.	HARE'S-EAR.	Pentandria Digynia. L.				
1	coriaceum. s.s.	(st.) thick-leaved.	Gibraltar.	1784.	7. 8. G. 4.	L'Herit. st. nov. t. 67.
2	fruticosum. s.s.	(st.) shrubby.	S. Europe.	1596.	— H. 4.	Flor. græc. t. 263.
3	fruticéscens. s.s.	(st.) frutescent.	Spain.	1752.	8. 9. H. 4.	Cav. ic. 2. t. 106.
4	plantagineum. s.s. (st.)	plantain-leaved.	Barbary.	1809.	— G. 4.	Desf. atl. 1. t. 57.
	β canescens. P.S. (st.) hoary.				— G. 4.	
5	spinòsum. s.s.	(st.) thorny.	Spain.	1752.	— F. 4.	Gouan. ill. 8. t. 2. f. 3.
6	diffórme. s.s.	(st.) various-leaved.	C. B. S.	—	— G. 4.	Burm. afr. t. 71. f. 1.
7	rotundifólium. s.s. (ye.)	round-leaved.	England.	6. 7. H. ⊖.	Eng. bot. v. 2. t. 99.
8	protractum. s.s.	(st.) protracted.	Portugal.	1824.	— H. ⊖.	
9	heterophyllum. s.s. (st.)	various-leaved.	Egypt.	—	— H. ⊖.	
10	aureum. s.s.	(ye.) golden.	Siberia.	—	— H. 4.	
11	longifólium. s.s.	(st.) long-leaved.	Switzerland.	1713.	5. 7. H. 4.	Moris. sect. 9. t. 11.
12	pyrenaicum. s.s.	(st.) Pyrenean.	Pyrenees.	1814.	— H. 4.	Gouan. ill. t. 4.
13	angulosum. w.	(st.) angular-leaved.	Switzerland.	1759.	— H. 4.	Moris. sect. 9. t. 12.
14	ranunculoides. w. (st.)	Crowfoot-like.	Pyrenees.	1790.	7. 8. H. 4.	Park. theatr. f. 7.
15	caricifólium. w.	(st.) Carex-leaved.	S. France.	1824.	— H. 4.	
16	Burseriánū. W.en. (st.)	Burser's.	—	1820.	— H. 4.	
17	petr'aum. s.s.	(st.) alpine.	S. Europe.	1823.	— H. 4.	Rich. de Bellev. t. 206.
18	sibiricum. M.B.	(st.) Siberian.	Siberia.	1826.	— H. 4.	
19	graminifólium. s.s. (st.)	Grass-leaved.	Switzerland.	1768.	5. 7. H. 4.	Jacq. ic. 1. t. 56.
20	stellátum. s.s.	(st.) starry.	—	1775.	— H. 4.	Hall. helv. t. 18.
21	falcátum. s.s.	(st.) twisted-stalk'd.	Germany.	1739.	5. 9. H. 4.	Jacq. aust. 2. t. 158.

22	scorzoneraefolium. W.en.(st.)	Scorzonera-l ^d . Germany.	1818.	6. 8. H. ȝ.		
23	rígidum. R.S.	(st.) rigid.	S. France.	1739.	5. 9. H. ȝ.	Moris. sect. 9. t. 12.
24	núdum. s.s.	(st.) naked-stalked.	C. B. S.	1778.	10. G. ȝ.	
25	gibraltáricum. Lag.(st.)	Gibraltar.	Gibraltar.	1825.	— G. ȝ.	
26	glaúicum. DC.	(ye.) glaucous.	S. France.	1828.	— H. ⓧ.	
27	Odontites. s.s.	(bh.) narrow-leaved.	England.	— H. ⓧ.	Eng. bot. v. 35. t. 2468.
	<i>Odontites luteola</i> . R.S.					
28	júnceum. s.s.	(ye.) linear-leaved.	S. Europe.	1722.	7. 8. H. ⓧ.	Moris. sect. 9. t. 12.
29	Gerárdi. R.S.	(ye.) Gerard's.	—	1804.	— H. ⓧ.	Jacq. aust. t. 256.
30	baldénsē. R.S.	(ye.) channel-leav'd.	M. Baldo.	1818.	— H. ⓧ.	W. et K. hung. 3.t. 257.
	<i>exaltatum</i> . P.S.					
31	lancifólium. H.H.	(ye.) lance-leaved.	—	1824.	6. 9. H. ⓧ.	
32	semicompósitum. s.s. (ye.) dwarf.	Spain.	—	1778.	— H. ⓧ.	Flor. græc. t. 261.
	<i>Odontites semicompósita</i> . R.S.					
33	gráile. M.B.	(ye.) slender-twigg'd.	Tauria.	1827	— H. ⓧ.	
34	tenuíssimum. s.s. (ye.) slender.	England.	—	7. 8. H. ⓧ.		
35	trífidum. s.s.	(ye.) trifid-umbel'd.	Italy.	1824.	5. 9. H. ȝ.	Eng. bot. v. 7. t. 478.
	HETEROMORPHA. DC. HETEROMORPHA.	Pentandria Dignynia.				
	arboréscens. DC. (st.) arborescent.	C. B. S.	1816.	7. 8. G. ȝ.		
	<i>Bupleurum arboréscens</i> . s.s. <i>Tenòria arboréscens</i> . R.S.					

Tribus IV. SESELINEÆ. DC.

LICHTENSTEINIA. DC. LICHTENSTEINIA. Pentandria Dignynia.
inébrians. DC. (wh.) various-leaved. C. B. S. 1816. 8. 9. G. ȝ.
Cénâthe inébrians. s.s.

ŒNA'NTHE. S.S.	WATER-DROPWORT.	Pentandria Dignynia. L.				
1	fstulósa. s.s.	(bh.) common.	Britain.	6. 8. H. ȝ.	Eng. bot. v. 6. t. 363.
2	tenuifólia. s.s.	(wh.) slender-leaved.	C. B. S.	1816.	6. 8. G. ȝ.	
3	peucedanifólia. s.s.	(wh.) round-rooted.	S. Europe.	— H.w. ȝ.	Pollich. palat. t. 3.
4	silaifólia. s.s.	(wh.) oblong-rooted.	England.	— H.w. ȝ.	Eng. bot. v. 5. t. 348.
	<i>peucedanifolia</i> . E.B. non. s.s.					
5	gymnorhiza. s.s. (wh.) naked rooted.	N. Europe.	1824.	— H.w. ȝ.		
6	pimpinelloídes. s.s. (bh.) BurnetSaxifrage.	England.	—	— H.w. ȝ.	Eng. bot. v. 5. t. 347.	
7	prolífera. s.s.	(wh.) proliferous.	S. Europe.	1739.	— H.w. ȝ.	Jacq. vind. 3. t. 62.
8	globulósa. s.s.	(wh.) globe-headed.	Portugal.	1710.	— H. ⓧ.	Gouan. ill. 18. t. 9.
9	crocátâ. s.s.	(wh.) poisonous.	Britain.	—	— H.w. ȝ.	Eng. bot. v. 33. t. 2313.
10	Phellandrium. s.s. (wh.) waterHemlock.	—	—	6. 7. H.w. ⓧ.	—	v. 10. t. 684.
	<i>Phellandrium aquáticum</i> . E.B.					
11	apiifólia. s.s.	(wh.) Parsley-leaved.	Portugal.	1806.	6. 8. H. ȝ.	
12	austrális. R.S.	(wh.) southern.	S. Europe.	1824.	— H. ȝ.	
13	involucrâ. Cav.	(wh.) involucred.	Spain.	1820.	— H. ȝ.	
14	am'ena. Leh.	(wh.) handsome.	—	1827.	— H. ȝ.	

ANESORHIZA. DC. ANESORHIZA. Pentandria Dignynia.
interruptâ. (wh.) interrupted.

C. B. S.	1816.	6. 8. G. ȝ.				
<i>Anénthe interrupta</i> . s.s.						
SCLEROSCIADIUM. DC. SCLEROSCIADIUM.	Pentandria Dignynia.					
rígens. DC.	(wh.) rigid-stalked.	C. B. S.	1787.	6. 8. G. ȝ.		
<i>Cónium rígens</i> . w. <i>Bùntium rígens</i> . s.s.						

ÆTH'USA. R.S. FOOLS-PARSLEY. Pentandria Dignynia. L.

1	Cynápium. w.	(wh.) common.	Britain.	7. 9. H. ⓧ.	
2	cynapioides. M.B.	(wh.) Caucasian.	Caucasus.	1820.	— H. ȝ.	Hoff.umb.ed.2. t.tit.f.9.
3	élata. L.en.	(wh.) tall.	Siberia.	—	— H. ⓧ.	
4	fátua. R.S.	(wh.) fine-leaved.	—	1781.	— H. ȝ.	

FœniculUM. G. FENNEL. Pentandria Dignynia.

1	vulgáre. G.	(ye.) common.	Britain.	7. 8. H. ȝ.	
	<i>Anethum Fœniculum</i> . E.B. <i>Méum Fœniculum</i> . s.s.					
2	dúlcê. DC.	(ye.) sweet.	S. Europe.	1824.	— H. ⓧ.	
3	piperítum.	(ye.) peppery.	Italy.	—	— H. ⓧ.	DC. hort. m. ined. t. 96.
	<i>Méum piperítum</i> . R.S. <i>Anethum piperítum</i> . Bert.					

4 látuum. F. (ye.) yellow.

	Persia.	1829.	— H. ⓧ.			
KUNDMA'NNIA. DC. KUNDMA'NNIA.	Pentandria Dignynia.					
pastinacæfólia. (ye.) Parsnp-leaved.	Sicily.	1686.	7. 8. H. ȝ.	Jacq. vind. 2. t. 133.		

Brignòlia pastinacæfólia. K.U. *Sium sículum*. R.S.

DEVERRA. DC.	DEVERRA.	Pentandria Dignynia.				
1 tortuosa. DC.	(wh.) twisting.	Levant.	1828. G. ȝ.	Desf. atl. 1. t. 73.	
	<i>Bùbon tortuosum</i> . Df.					

2	aphy'lla. DC.	(wh.) leafless.	C. B. S.	1816.	G. $\ddot{\alpha}$.
	<i>Bûbon aphy'llus.</i>	Schlecht.				
SE'SELI. S.S.	MEADOW SAXIFRAGE.	Pentandria	Digynia.	S.S.		
1	élâtum. s.s.	(wh.) tall.	Austria.	1710.	7. 8. H. $\ddot{\alpha}$.	Gouan. ill. 16. t. 8.
2	vârium. s.s.	(wh.) variable.	1818.	— H. $\ddot{\alpha}$.	Jacq. aust. 2. t. 144.
3	glaucum. s.s. (pu.wh.)	glaucous.	France.	1759.	— H. $\ddot{\alpha}$.	Jacq. vind. 2. t. 129.
4	taûricum. s.s.	(wh.) Taurian.	Tauria.	1818.	6. 8. H. $\ddot{\alpha}$.	Jacq. aust. t. 55.
5	montânus. s.s.	(wh.) mountain.	Italy.	1658.	6. 7. H. $\ddot{\alpha}$.	W. et K. hung. t. 89.
6	ânnuum. s.s.	(bh.) annual.	Europe.	1816.	7. 8. H. $\ddot{\alpha}$.	
	<i>Cârum simplex. W.</i>					
7	campêstre. s.s.	(wh.) field.	Poland.	1827.	— H. $\ddot{\alpha}$.	
8	pimpinelloïdes. s.s. (wh.)	nodding-flower'd.	S. Europe.	1796.	— H. $\ddot{\alpha}$.	
9	Hippomârathrum. s.s. (bh.)	various-leav'd.	Austria.	1656.	— H. $\ddot{\alpha}$.	Jacq. aust. 2. t. 143.
10	tortuosum. s.s.	(wh.) crooked.	S. Europe.	1597.	10. H. $\ddot{\alpha}$.	Moris. sect. 9. t. 6.
11	leucospérnum. s.s. (wh.)	white-seeded.	Hungary.	1805.	7. H. $\ddot{\alpha}$.	W. et K. hung. t. 89.
12	proliferum. s.s.	(wh.) prolific.	1823.	5. 8. H. $\ddot{\alpha}$.	
13	chaerophylloïdes. s.s. (wh.)	Chervil-like.	C. B. S.	1816.	— G. $\ddot{\alpha}$.	
14	Boccòni. s.s.	(wh.) Boccone's.	Sicily.	1826.	— H. $\ddot{\alpha}$.	Bocc. sic. s. 27. 28.
15	divaricatum. s.s. (ye.)	shining-leaved.	N. America.	1812.	6. 7. H. $\ddot{\alpha}$.	Bot. mag. t. 1742.
16	striatum. s.s.	(wh.) striped-stalked.	C. B. S.	1816.	7. 8. G. $\ddot{\alpha}$.	
17	grâclie. s.s.	(ye.) slender.	Hungary.	1805.	6. 7. H. $\ddot{\alpha}$.	W. et K. hung. 2. t. 117.
18	peucedanifoliūm. K.U.	(wh.) smooth-stalk'd.	Hungary.	1826.	7. 9. H. $\ddot{\alpha}$.	t. 146.
19	petræum. K.U.	(wh.) glaucous.	Tauria.	1824.	7. 8. H. $\ddot{\alpha}$.	
	<i>Bûbon glaucus. s.s.</i>					
20	cuneifoliūm. M.B. (wh.)	wedge-leaved.	Sicily.	1823.	7. 9. H. $\ddot{\alpha}$.	Moris. his. s. 9. t. 7.
21	gummiñerum. K.U. (bh.)	gum-bearing.	Crimea.	1710.	— H. $\ddot{\alpha}$.	Sm. exot. bot. t. 20.
22	dichotomum. M.B. (wh.)	forked.	Tauria.	1816.	— H. $\ddot{\alpha}$.	Bot. mag. t. 2073.
23	ericocéphalum.	(wh.) woolly-headed.	Siberia.	1827.	— H. $\ddot{\alpha}$.	
24	côrsicum. L.en.	(wh.) dwarf annual.	Corsica.	1825.	— H. \odot .	
25	Pallâsii. Bess.	(wh.) Pallas's.	Siberia.	1823.	— H. $\ddot{\alpha}$.	
26	Lobelianum. Bess. (wh.)	Lobel's.	—	— H. $\ddot{\alpha}$.	
LIBAN'OTIS. DC.	LIBAN'OTIS.	Pentandria	Digynia.			
1	buchtorménsis. DC. (wh.)	Siberian.	Siberia.	1822.	7. 8. H. $\ddot{\alpha}$.	
	<i>Bûbon buchtorménsis.</i>	R.S. <i>Athamânta rigida.</i>	Horn.			
2	vulgâris. DC.	(wh.) mountain.	England.	— H. $\ddot{\alpha}$.	Eng. bot. v. 2. t. 138.
	<i>Athamânta Libanotis.</i>	E.B.				
CENOL'OPHIUM. DC.	CENOL'OPHIUM.	Pentandria	Digynia.			
Fischéri. DC.	(wh.) Fischer's.	Siberia.	1820.	6. 8. H. $\ddot{\alpha}$.		
	<i>Cnidium Fischéri.</i>	s.s. <i>Angélica Fischéri.</i>	R.S. <i>Ligústicum Fischéri.</i>	L. enum.		
CNI'DIUM. S.S.	CNI'DIUM.	Pentandria	Digynia.	S.S.		
1	Monniéri. s.s.	(wh.) annual.	S. Europe.	1771.	7. 8. H. \odot .	Jacq. vind. 1. t. 62.
	<i>Selinum Monniéri.</i>	Jacq.				
2	apioides. H.U.	(wh.) Parsley-leaved.	Europe.	1791.	6. 7. H. $\ddot{\alpha}$.	Jacq. aust. 5. ap. t. 44.
	<i>Ligústicum apioides.</i>	R.S. <i>silafolium.</i>	Jacq.			
3	venôsum. DC.	(wh.) veined.	Europe.	1818.	7. 8. H. $\ddot{\alpha}$.	
	<i>Séseli dûbium.</i>	R.S.				
THA'SPIUM. N.	THA'SPIUM.	Pentandria	Digynia.			
1	actaeifoliūm. N.	(ye.) Actaea-leaved.	N. America.	1824.	7. 8. H. $\ddot{\alpha}$.	
	<i>Ligústicum actaeifoliūm.</i>	M.				
2	barbinode. N.	(ye.) bearded-jointed.	—	1759.	6. 7. H. $\ddot{\alpha}$.	
	<i>Thâpsia trifoliata.</i>	R.S.				
TROCHISCA'NTHES. K.U.	TROCHISCA'NTHES.	Pentandria	Digynia.			
	<i>nodiflorus.</i>	K.U. (bh.) knot-flowered.	APennines.	1818.	6. 7. H. $\ddot{\alpha}$.	All. ped. t. 72.
	<i>Ligústicum nodiflorum.</i>	s.s. <i>verticillatum.</i>	W. et K. hung. 2. t. 171.			
ATHAMA'NTA. S.S.	SPIGNEL.	Pentandria	Digynia.	L.		
1	macedônica. s.s. (wh.)	Macedonian.	Greece.	1596.	6. 8. G. $\ddot{\alpha}$.	Lobel. hist. 406. ic. 708.
	<i>Bûbon macedônicum.</i>	L.				
2	condensata. w.	(wh.) close-headed.	Siberia.	1773.	7. 9. H. $\ddot{\alpha}$.	Gmel. sib. 1. t. 44.
3	sibîrica. w.	(st.) Siberian.	—	1771.	7. 8. H. $\ddot{\alpha}$.	Lam. ill. t. 194. f. 1.
4	sícula. s.s.	(wh.) Flixweed-leav'd.	Sicily.	1686.	6. 7. H. $\ddot{\alpha}$.	Ten. fl. nap. 1. t. 25.
5	Matthiolí. s.s.	(wh.) fine-leaved.	Carniola.	1802.	— H. $\ddot{\alpha}$.	Jacq. ic. rar. 1. t. 57.
6	stricta. Led.	(wh.) upright.	Siberia.	1827.	— H. $\ddot{\alpha}$.	
7	creténica. s.s.	(wh.) candy-carrot.	Austria.	1596.	— H. $\ddot{\alpha}$.	Jacq. aust. 1. t. 62.
8	incâna. s.s.	(wh.) hoary.	Siberia.	1802.	7. 8. H. $\ddot{\alpha}$.	
9	ânnua. w.	(wh.) annual.	Candia.	1731.	6. 7. H. \odot .	
LIGUSTICUM. S.S.	LIGUSTICUM.	Pentandria	Digynia.	L.		
1	scóticum. s.s.	(bh.) Scotch.	Britain.	— H. $\ddot{\alpha}$.	Eng. bot. v. 17. t. 1207.
2	obtusifoliūm. s.s. (wh.)	blunt-leaved.	Tangiers.	1824.	7. 8. H. $\ddot{\alpha}$.	
3	athamantoïdes. s.s. (st.)	Carrot-leaved.	Pyrenees.	1734.	— H. $\ddot{\alpha}$.	Scop. carn. t. 10.
	<i>Ammi daucifoliūm.</i>	w. <i>Athamânta pyrenaica.</i>	Jacq. vind. 2. t. 197.			

4 alatum. s.s. (ro.wh.) winged-stalked. Caucasus.	1823.	7. 8. H. 2.	
5 simplex. s.s. (wh.pu.) shining. Switzerland.	1775.	7. H. 3.	Jacq. misc. 2. t. 2.
Laserpitium simplex. Jacq. lúcidum. H.K.			
6 ferulaceum. s.s. (wh.) Fennel-leaved. Levant.	1752.	6. H. 2.	Allion. ped. t. 60.
Laserpitium dauricum. Jacq. vind. 3. t. 38.			
7 canticans. s.s. (wh.) pale.	1780.	7. 8. H. 2.	
8 pyrenaicum. K.U. (wh.) Pyrenean. Pyrenees.	1739.	6. 7. H. 2.	Gouan. ill. t. 7. f. 2.
Cnídium pyrenæum. s.s. Séseli aristatum. H.K.			
9 Seguierii. K.U. (wh.) Fennel-leaved. Italy.	1774.	7. 8. H. 2.	Jacq. vind. 1. t. 61.
Selinum Seguierii. Jacq. Imperatoria Seguierii. s.s.			
10 Fontanæsii. s.s. (wh.) fine-leaved. Algiers.	6. 7. H. 2.	Desf. atl. 1. t. 71.
Laserpitium peucedanoïdes. Desf.			
11 cuneifolium. Led. (wh.) wedge-leaved. Siberia.	1827.	— H. 2.	
SIL'AUS. K.U. SIL'AUS. Pentandria Digynia.			
1 pratensis. K.U. (st.) meadow. England.	6. 8. H. 2.	Eng. bot. v. 30. t. 2142.
Peucedanum Silaus. E.B. Cnídium Silaus. s.s.			
2 Matthioli. K.U. (ye.) late-flowering. Italy.	1824.	7. 8. H. 2.	
Peucedanum Matthioli. s.s.			
3 alpestris. K.U. (st.) Alpine. France.	1739.	6. 7. H. 2.	
WALLRO'THIA. DC. WALLRO'THIA. Pentandria Digynia.			
1 alpina. DC. (wh.) alpine. Hungary.	1827.	6. 8. H. 2.	W. et K. hung. 2. t. 182.
tuberosa. R.S. Búnium alpinum. W.K.			
2 tenuifolia. DC. (wh.) slender-leaved. Pyrenees.	1826.	— H. 2.	DC. mem. umb. t. 1.f.A.
M'EUM. S.S. BAWD-MONEY. Pentandria Digynia. S.S.			
1 athamanticum. s.s. (wh.) common. Britain.	4. 6. H. 2.	Eng. bot. v. 32. t. 2249.
2 Mutellina. s.s. (re.) Alpine. Austria.	1774.	7. 8. H. 2.	Jacq. aust. t. 56.
Phellandrium Mutellina. L.			
3 sibiricum. s.s. (ye.) Siberian. Siberia.	1820.	— H. 2.	
CONIOSEL'INUM. H.U. CONIOSEL'INUM. Pentandria Digynia.			
1 íngricum. F. (wh.) Ingrian. Russia.	1827.	7. 8. H. 2.	
2 tatáricum. F. (wh.) Tartarian. Tartary.	1825.	— H. 2.	
Angélica Fischéri. R.S.			
CRITHMUM. L.en. SAMPHIRE. Pentandria Digynia. L.			
1 Libanótis. L.en. (ye.) smooth-seeded. Sicily.	1570.	7. 8. H. 2.	Schk. handb. 1. t. 65.
Cáchrys Libanótis. s.s.			
2 marítimum. w. (st.) sea-side. Britain.	7. 9. H. 2.	Eng. bot. v. 12. t. 819.

Tribus V. ANGELICEÆ. DC.

LEVI'STICUM. K.U. LOVAGE. Pentandria Digynia.			
officinale. K.U. (ye.) common. Italy.	1596.	6. 7. H. 2.	Schk. handb. t. 68.
Ligósticum Levisticum. s.s.			
SELINUM. L. MILK-PARSLEY. Pentandria Digynia. L.			
1 Carvifolium. w. (bh.) Caraway-leav'd. Austria.	1774.	7. 8. H. 2.	Jacq. aust. 1. t. 16.
OSTE'RICUM. H.U. OSTE'RICUM. Pentandria Digynia.			
pratense. H.U. (wh.) field. Galicia.	1824.	7. 8. H. 2.	
Angélica pratensis. M.B.			
ANGÉLICA. K.U. ANGE'LICA. Pentandria Digynia. L.			
1 sylvestris. s.s. (wh.) wild. Britain.	6. 8. H. 2.	Eng. bot. v. 16. t. 1128.
2 Razoulli. s.s. (bh.) decurrent-l'd. Pyrenees.	1816.	— H. 2.	Gouan. ill. 13. t. 6.
3 montana. K.U. (wh.) mountain. Switzerland.	—	— H. 2.	
4 flavescens. Bess. (st.) yellowish. Pyrenees.	—	— H. 2.	
5 pyrenæa. s.s. (wh.) Pyrenean.	1818.	— H. 2.	Gouan. ill. p. 11. t. 5.
6 lúcida. s.s. (br.) shining. Canada.	1640.	— H. 2.	Jacq. vind. 3. t. 24.
7 chærophyllea. s.s. (wh.) Chervil-leaved. Livonia.	1827.	— H. 2.	
8 Wormskioldii. F. (wh.) Wormskold's. Kamtschatka.	—	— H. 2.	
9 lancifolia. s.s. (wh.) lance-leaved.	1818.	— H. 2.	
Selinum angelicastrum. L. enum.			
ARCHANGE'LICA. H.U. ARCHANGE'LICA. Pentandria Digynia.			
1 officinalis. H.U. (wh.) garden. England.	6. 8. H. 2.	Eng. bot. v. 36. t. 2561.
Angélica Archangélica. E.B.			
2 atropurpurea. H.U. (pu.) dark purple. Canada.	1759.	7. 8. H. 2.	Cornut. canad. t. 199.

Tribus VI. PEUCEDANEÆ. DC.

OP'OPANAX. K.U. OP'OPANAX. Pentandria Digynia.			
Chirónium. K.U. rough. S. Europe.	1640.	6. 7. H. 2.	W. et K. hung. 3.t.211.
Pastinaca Opópanax. w. Ferula Opópanax. s.s.			

FER'ULA. S.S.

GIANT-FENNEL. Pentandria Digynia. L.

1	lavigata, s.s.	(ye.) smooth.	C. B. S.	1774.	12. 3.	G. ḥ.
	Būbon lavigatum, w.					
2	nudicaulis, s.s.	(ye.) naked-stalked.	Sicily.	1825.	—	H. ȝ.
3	A'sa f'œtida, s.s.	(st.) Asa fœtida.	Persia.	1818.	7. 8.	H. ȝ.
4	communis, s.s.	(ye.) common.	S. Europe.	1597.	6. 7.	H. ȝ.
5	glauca, s.s.	(ye.) glaucous.	Italy.	1596.	—	H. ȝ.
6	sylvatica, s.s.	(ye.) wood.	Poland.	1827.	—	H. ȝ.
7	campéstris, s.s.	(ye.) field.	Tauria.	—	—	H. ȝ.
8	nodiflora, s.s.	(ye.) knot-flowered.	S. Europe.	1596.	—	H. ȝ.
	sulcata, Desf.					
9	Ferulago, s.s.	(ye.) Fennel-leaved.	—	—	—	H. ȝ.
	nodiflora, Jacq. non s.s.					

10	thyrsiflora, s.s.	(st.) thyrs-flower'd.	Crete.	1823.	—	F. ȝ.
11	tingitana, s.s.	(ye.) Tangier.	Barbary.	1680.	—	H. ḡ.
12	stricta, s.s.	(ye.) upright.	C. B. S.	1816.	—	G. ḡ.
13	nuda, s.s.	(ye.) naked.	Siberia.	1827.	—	H. ȝ.
14	caspica, s.s.	(ye.) Caspian.	Caspia.	1824.	—	H. ȝ.
15	orientalis, s.s.	(ye.) eastern.	Levant.	1759.	7. 8.	H. ȝ.
16	persica, s.s.	(ye.) stinking.	Persia.	1782.	—	H. ȝ.
17	tatarica, F.	(ye.) Tartarian.	Tartary.	1828.	—	H. ȝ.
18	capillaris, s.s.	(ye.) capillary.	Portugal.	—	—	H. ȝ.
19	desertorum, Led.	(ye.) desert.	Siberia.	1827.	—	H. ȝ.
20	songarica, R.S.	(ye.) Songarian.	—	—	—	H. ȝ.

ERIOSYNA'PHE. DC. ERIOSYNA'PHE. Pentandria Digynia.

longifolia, dc.	(ye.) long-leaved.	Wolga.	1823.	7. 8.	H. ȝ.
Frēula longifolia, F.					

PÆLIMBIA. DC. PÆLIMBIA. Pentandria Digynia.

Chabr'æi, dc.	(bh.) fine-leaved.	Europe.	1791.	7. 8.	H. ȝ.
Selinum Chabr'æi, Jacq.	Oreoselinum Chabr'æi, M.B.	Imperatòria Chabr'æi, R.S.			

PEUCED'ANUM. S.S. SULPHURWORT. Pentandria Digynia. L.

1	officinale, s.s.	(ye.) officinal.	England.	5. 7.	H. ȝ.
2	paniculatum, R.S.	(ye.) paniced.	Corsica.	1827.	—	H. ȝ.

3	alsaticum, w.	(wh.) small-headed.	Austria.	1774.	6. 8.	H. ȝ.
4	gálico, s.s.	(wh.) French.	France.	1816.	—	H. ȝ.
5	ítalicum, P.S.	(ye.) yellow-flower'd.	Italy.	5. 7.	H. ȝ.
6	ruthenicum, R.S.	(ye.) Russian.	Russia.	1824.	6. 8.	H. ȝ.
7	tauricum, R.S.	(ye.) Taurian.	Tauria.	1828.	—	H. ȝ.
8	longifolium, R.S.	(ye.) long-leaved.	Hungary.	1826.	—	H. ȝ.
9	arenarium, s.s.	(ye.) sand.	—	1816.	6. 7.	H. ȝ.
10	Morisóni, R.S.	(ye.) Morison's.	—	—	—	H. ȝ.
11	nodosum, s.s.	(wh.) knotted-stalked.	Crete.	—	H. ȝ.
12	aureum, s.s.	(ye.) golden.	Canaries.	1779.	6.	G. ḡ.
13	tenuifolium, s.s.	(ye.) slender-leaved.	C. B. S.	1816.	6. 8.	G. ḡ.
14	isetense, s.s.	(st.) straw-coloured.	Siberia.	1828.	—	H. ḡ.
15	sibiricum, w.	(ye.) Siberian.	—	1804.	6. 7.	H. ȝ.
16	palustre, K.U.	(wh.) marsh.	Britain.	7. 8.	H. ȝ.

	Selinum palustre, E.B.	Thyselinum palustre, H.U.				
17	baicalense, K.U.	(wh.) Baical.	Siberia.	1824.	—	H. ȝ.

18	Cervaria, K.U.	(wh. pu.) large-leaved.	Europe.	1597.	—	H. ȝ.
	Athamánta Cervaria, Jacq.	Ligósticum Cervaria, s.s.				

19	Oreoselinum, K.U.	(re.wh.) bending.	Europe.	1780.	7. 9.	H. ȝ.
	Athamánta Oreoselinum, Jacq.	Oreoselinum legitimum, M.B.				

20	montanum, K.U.	(wh.) mountain.	Switzerland.	1816.	7. 8.	H. ȝ.
21	involucratum, K.U.	(wh.) involucred.	S. France.	1824.	—	H. ȝ.

	Selinum peucedanoïdes, R.S.	Oreoselinum peucedanoïdes, H.U.				
22	austriacum, K.U.	(wh.) Austrian.	Austria.	1804.	—	H. ȝ.

	Selinum austriacum, Jacq.	Oreoselinum austriacum, H.U.				
23	rablense, K.U.	(wh.) linear-leaved.	N. Europe.	1816.	7. 8.	H. ȝ.

24	latifolium, (wh.)	broad-leaved.	Caucasus.	—	—	H. ȝ.
	Oreoselinum latifolium, M.B.					

25	polymorphum, Lag.	(wh.) variable.	Russia.	1824.	—	H. ȝ.
26	Bellárdi, Lag.	(wh.) Bellardii's.	N. Europe.	—	—	H. ȝ.

27	élégans,	(wh.) elegant.	—	—	—	H. ȝ.
	Oreoselinum élégans, H.U.					

28	podòlicum, H.U.	(wh.) Polish.	Poland.	1819.	—	H. ȝ.
29	nígricans, H.U.	(wh.) black-rooted.	Switzerland.	1823.	—	H. ȝ.

30	caucasicum,	(st.) Caucasian.	Caucasus.	1824.	—	H. ȝ.
31	Ostrúthium, K.U.	(wh.) Masterwort.	Scotland.	5. 7.	H. ȝ.

	Imperatòria Ostrúthium, Eng. bot.					

Moris. s. 9. t. 15 f. 3.

Spreng. umb. t. 7. f. 14.
Tourn. it. 3. t. 239.

Andr. rep. t. 558.

Jacq. aust. 5. ap. t. 5.

Flor. græc. t. 280.

Herm. parad. t. 165.

DC.mem.umb. pl. 1 f. E.

Eng. bot. v. 25. t. 1767.

Jacq. aust. 1. t. 72.

Gmel. sib. 1. t. 41.

W. et K. hung. t. 251.

Bot. reg. t. 20.

Moris. his. 3. s. 9. t. 15. f. 1.

Moris. his. s. 9. t. 17.

Eng. bot. v. 4. t. 229.

Gmel. sib. 1. t. 42-43.

W. et K. hung. 1. t. 20.

Jacq. aust. 1. t. 69.

Jacq. aust. 1. t. 68.

Jacq. aust. 1. t. 71.

Flor. dan. t. 1330.

Eng. bot. v. 20. t. 1380.

32 imperatorioïdes. k.u. (wh.) narrow-leav'd. Alps, Europe. 1818.	— H. 4.	Moris. his. s. 9. t. 4.
<i>Imperatoria angustifolia</i> . s.s. <i>Selinum imperatorioïdes</i> . L.en.		
IMPERATORIA. DC. MASTERWORT. Pentandria Digynia. L.		
verticillaris. s.s. (wh.) whorl-flowered. Italy. 1683.	7. H. 4.	Jacq. vind. 2. t. 130.
<i>Angelica verticillaris</i> . W.		
CALLIS'ACE. F. CALLIS'ACE. Pentandria Digynia.		
davurica. F. (wh.) Daurian. Dauria. 1820.	7. 8. H. 4.	
<i>Thysselinum davuricum</i> . s.s.		
B'UBON. K.U. B'UBON. Pentandria Digynia.		
Galbânum. k.u. (ye.) Lovage-leaved. C. B. S. 1696.	6. 8. G. ½.	Jacq. vind. 3. t. 36.
<i>Selinum Galbânum</i> . R.S.		
AN'ETHUM. Len. DILL. Pentandria Digynia. L.		
1 graveolens. w. (ye.) common. Spain. 1570.	6. 7. H. ½.	Hoff. umb. gen. t. 1.f. 13.
<i>Pastinaca Anethum</i> . s.s.		
2 ségetum. L.en. (ye.) Portugal. Portugal. 1796.	— H. ¼.	Jacq. vind. t. 132.
3 Sôwa. H.B. (st.) Indian. E. Indies. 1810.	— H. ¼.	
CAPNOPHYLLUM. S.S. CAPNOPHYLLUM. Pentandria Digynia. S.S.		
africânum. s.s. (wh.) Rue-leaved. C. B. S. 1759.	6. 9. H. ¼.	Jacq. vind. t. 194.
<i>Conium africânum</i> . w. <i>Rùmia capensis</i> . L.en.		
ARCHEMORA. DC. ARCHEMORA. Pentandria Digynia.		
rígida. DC. (wh.) rigid. Virginia. 1774.	7. 8. H. 4.	Moris. dis. s. 9. t. 7.f. 1.
<i>Sium rígidius</i> . w. <i>Cénâthe rígida</i> . n. <i>Pastinaca rígida</i> . s.s.		
PASTINACA. S.S. PARSNEP. Pentandria Digynia. L.		
1 lúcida. s.s. (ye.) shining-leaved. S. Europe. 1771.	6. 7. H. ½.	Jacq. vind. 2. t. 199.
2 sativa. W.en. (ye.) cultivated.	7. H. ¾.	Flor. dan. t. 1206.
3 opâca. W.en. (ye.) wild. Britain.	— H. ½.	Eng. bot. v. 8. t. 556.
<i>sativa β arvensis</i> . P.S.		
4 taraxacifolia. F. (ye.) Dandelion-ld. Russia. 1822.	— H. ¼.	
5 graveolens. s.s. (ye.) strong-scented. Tauria. 1819.	7. 8. H. 4.	Hoff. umb. t. 1.B.f. 6.c.
<i>Malabaíla graveolens</i> . H.U.		
6 panacifolia. F. (ye.) Panax-leaved. Tauria. 1826.	7. 8. H. 4.	
7 pimpinellifolia.s.s.(ye.) Burnet-Saxifrage. Caucasus. 1824.	— H. ½.	— p.209.t.1.B.f.6.a.b.
8 umbrôsa. F. (ye.) shady. 1821.	— H. ¼.	
9 dissécta. s.s. (ye.) cut-leaved. Levant. 1816.	6. 7. H. ½.	Venten. cels. t. 78.
10 nudicaulis. s.s. (ye.) naked-stalked. N. America. 1816.	— H. ¼.	
<i>Smyrnium nudicaule</i> . Ph.		
11 divaricata. Desf. (ye.) spreading.	1824.	— H. ¼.
ASTYD'AMIA. DC. ASTYD'AMIA. Pentandria Digynia.		
canariensis. DC. (ye.) cut-leaved. Canaries. 1780.	7. G. ½.	DC. dis. umb. t. 1. f. D.
<i>Bupleürum canariense</i> . s.s. <i>Crithmum latifolium</i> . L. <i>Tendria canariensis</i> . R.S.		
HERA'CLEUM. S.S. COW-PARSNEP. Pentandria Digynia. L.		
1 Sphondyl'ium. w.(wh.) common. Britain.	5. 6. H. ¼.	Eng. bot. v. 14. t. 939.
2 flavescens. w. (wh.) yellowish. Austria. 1789.	6. 7. H. ¼.	Jacq. aust. 2. t. 173.
3 élégans. w. (wh.) rough-leaved.	1800.	— 2. t. 175.
4 angustifolium.w.(wh.) narrow-leaved.	—	— 2. t. 174.
5 longifolium.Jacq.(wh.) long-leaved.	—	— 2. t. 374.
6 sibiricum. s.s. (st.) Siberian. Siberia. 1768.	— H. ¼.	Gmel. sib. 1. t. 50.
7 Panáces. s.s. (wh.) Fig-leaved.	1596.	Lobel. ic. 701.
8 pyrenäicum. w. (wh.) Pyrenean. Pyrenees. 1804.	— H. ¼.	
9 amplifolium.Lap.(wh.) large-leaved.	—	Lap. fl. pyr. t. 79.
10 gummiferum.w. (wh.) gum-bearing.	1816.	Wild.hort.ber. t. 53. 54.
11 speciosum. H.U. (wh.) handsome.	—	
12 ásperum. M.B. (wh.) rough. Caucasus.	—	
13 cordátum. Pr. (wh.) heart-leaved. Sicily. 1825.	— H. ¼.	
14 villosum. s.s. (wh.) villous. Siberia. 1816.	— H. ¼.	
15 grandiflorum.Stev.(w.)large-flowered. Caucasus. 1818.	— H. ¼.	
16 decipiens. H.U. (wh.) deceiving.	—	
17 marginatum. F. (wh.) margined. Siberia. 1816.	— H. ¼.	
18 pubescens. M.B. (wh.) pubescent. Caucasus. 1824.	— H. ¼.	
19 caucasicum. R.S. (wh.) Caucasian.	—	
20 alpinum. s.s. (wh.) Alpine. Switzerland. 1739.	6. 7. H. ¼.	Barrel. ic. 55.
21 austriacum. s.s. (wh.) Austrian. Austria. 1816.	— H. ¼.	Jacq. aust. t. 61.
<i>Tordyl'ium siifolium</i> . w.		
22 ligusticifolium. s.s.(wh.) Lovage-leaved. Tauria. 1824.	— H. ½.	
23 subvillous. F. (wh.) subvillous. Siberia. 1820.	— H. ¼.	
24 giganteum. F. (wh.) gigantic.	—	
25 básicum. F. (wh.) stately.	—	
26 laciniatum. F. (wh.) jagged.	1824.	
27 tauricum. F. (wh.) Taurian. Tauria. 1823.	— H. ¼.	

28 cásicum. f.	(wh.) Caspian.	Caucasus.	1828.	6. 7. H. ¼.	
29 lanatum. M.	(wh.) woolly.	N. America.	—	— H. ¼.	
30 pùmilum. DC.	(wh.) dwarf.	Dauphiny.	1800.	5. 7. H. ¼.	Vill. delph. 2. t. 14.
	<i>Peucedanum Vocontiorum</i> . s.s.				
ZOSÍMIA. H.U.	ZOSÍMIA.	Pentandria Digynia.			
orientális. H.U.	(ye.) Wormwood-ld.	Persia.	1816.	7. 8. H. ⊖.	Vent. choix. t. 22.
	<i>Herácleum absinthifolium</i> . s.s. <i>Tordy'lium absinthifolium</i> . P.S.				

Tribus VII. TORDYLINEÆ. DC.

HASSELQUI'STIA. S.S.	HASSELQUI'STIA.	Pentandria Digynia. L.			
1 ægyptiaca. s.s.	(wh.) Egyptian.	Egypt.	1768.	7. H. ⊖.	Jacq. vind. 1. t. 37.
2 cordátæ. s.s.	(wh.) heart-leaved.	—	1787.	— H. ⊖.	— 2. t. 193.
TORDYLIUM. K.U.	HARTWORT.	Pentandria Digynia. L.			
1 syriacum. s.s.	(wh.) Syrian.	Syria.	1597.	7. H. ⊖.	Jacq. vind. 1. t. 54.
2 cordátum. E.	(wh.) heart-leaved.	—	1826.	— H. ⊖.	Eng. bot. v. 17. t. 1173.
3 máximum. s.s.	(wh.) great.	England.	—	6. 7. H. ⊖.	Eng. bot. v. 34. t. 2440.
4 lusitánicum. W.en.(wh.) Portugal.	Portugal.	Portugal.	1820.	— H. ⊖.	Jacq. vind. 1. t. 53.
CONDYLOCA'RPU'S. K.U.	CONDYLOCA'RPU'S.	Pentandria Monogynia.			
1 officinális. k.u.	(bh.) officinal.	England.	—	7. H. ⊖.	
2 ápulus. k.u.	(wh.) small.	Italy.	1739.	— H. ⊖.	

Tribus VIII. SILERINEÆ. DC.

KRUB'ERA. H.U.	KRUB'ERA.	Pentandria Digynia.			
leptophylla. k.u.	slender-leaved.	Barbary.	1800.	6. 7. H. ⊖.	Desf. atl. 1. t. 66,
<i>Cónium dichótومum</i> . Df.	<i>Cáchrys dichótoma</i> . r.s.	<i>Tordy'lium peregrinum</i> . L.			
AGASY'LLIS. S.U.	AGASY'LLIS.	Pentandria Digynia.			
1 caucásica. s.u.	(wh.) Caucasean.	Caucasus.	1823.	5. 7. H. ¼.	
<i>Síler caucásicum</i> . s.s.	<i>Cáchrys latifolia</i> . M.B.				
2 salsa. s.u.	(st.) fine-leaved.	Siberia.	1804.	7. 8. H. ¼.	Pall. act. pet. 1779. t. 8.
<i>Síler sálsum</i> . r.s.	<i>Sison sálsum</i> . L.				
SILE'R. K.U.	SILE'R.	Pentandria Digynia.			
trilóbum. k.u.	(wh.) Columbine-leav'd.	Austria.	1796.	5. 7. H. ¼.	Jacq. aust. 2. t. 147.
<i>aquilegífolium</i> . r.s.	<i>Laserpítium aquilegífolium</i> . w.				

Tribus IX. CUMINEÆ. DC.

CU'MINUM. S.S.	CUMIN.	Pentandria Digynia. L.			
Cýminum. s.s.	(wh.) officinal.	Egypt.	1594.	6. 7. H. ⊖.	Cav. ic. 4. t. 360.

Tribus X. THAPSIEÆ. DC.

THA'PSIA. S.S.	DEADLY CARROT.	Pentandria Digynia. L.			
1 villósa. s.s.	(ye.) villous.	S. Europe.	1710.	6. 7. H. ¼.	Moris. s. 9. t. 18. f. 3.
2 f'etida. s.s.	(ye.) stinking.	Spain.	1596.	7. 8. H. ¼.	— t. 18. f. 7.
3 gummífera. s.s.	(wh.) gum-bearing.	S. Europe.	1824.	— H. ¼.	Desf. atl. 1. t. 72.
	<i>Laserpitium gummiferum</i> . Desf.				
4 Asclépium. s.s.	(ye.) oriental.	Levant.	—	— H. ¼.	Flor. græc. t. 286.
5 gargánica. s.s.	(ye.) Garganian.	Barbary.	1683.	— H. ¼.	Gouan. ill. 18. t. 10.
6 altíssima. r.s.	(ye.) tall.	—	—	— H. ¼.	
CYMO'PTERUS. DC.	CYMO'PTERUS.	Pentandria Digynia.			
glomeràta.	(wh.) clustered.	N. America.	1824.	5. 6. H. ¼.	
	<i>Selnum acáule</i> . Ph.				
LASERPI'TIUM. DC.	LASERWORT.	Pentandria Digynia. L.			
1 latífolium. s.s.	(wh.) broad-leaved.	Europe.	1640.	6. 7. H. ¼.	Jacq. aust t. 146.
2 Libanòtis. s.s.	(pu.) pubescent.	S. Europe.	1824.	— H. ¼.	Lob. ic. 705.
3 marginátum. s.s.	(st.) purple-margin'd.	Hungary.	1818.	— H. ¼.	W. et K. hung. 2. t 192.
4 Síler. s.s.	(wh.) mountain.	—	1640.	— H. ¼.	— 2. t. 145.

5 <i>alpinum</i> . K.U. (wh.) alpine. <i>trilobum</i> . s.s.	S. Europe. 1640.	5. 7. H. ȝ.	W. et K. hung. 3. t. 253.
6 <i>angustifolium</i> . w. (wh.) narrow-leaved.	S. Europe. 1738.	6. 7. H. ȝ.	Moris. s. 9. t. 19. f. 9.
7 <i>peucedanoides</i> . w. (wh.) fine-leaved.	Italy.	— H. ȝ.	Jacq. ic. 2. t. 350.
8 <i>resinosa</i> . s.s. (ye.) resinous.	Sicily. 1825.	— H. ȝ.	Hall. stirp. t. 19.
9 <i>hirsutum</i> . s.s. (wh.) hairy.	Switzerland. 1759.	— H. ȝ.	Jacq. ic. 1. t. 58.
10 <i>Archangelica</i> . s.s. (wh.) spotted-stalked.	Hungary. 1818.	— H. ȝ.	Jacq. aust. 2. t. 153.
11 <i>pilosum</i> . s.s. (st.) sulphur-color'd.	Caucasus. 1759.	— H. ȝ.	Pluk. phyt. t. 198. f. 6.
12 <i>pruthenicum</i> . s.s. (wh.) Prussian.	Germany. —	7. 8. H. ȝ.	Tourn. it. 2. t. 121.
13 <i>hispidum</i> . M.B. (bh.) hispid.	Caucasus. 1818.	— H. ȝ.	
14 <i>Athamantæ</i> . s.s. (wh.) branching.	Siberia. —	— H. ♂.	
15 <i>gallicum</i> . s.s. (wh.) French.	S. Europe. 1683.	6. 7. H. ȝ.	DC. mem. umb. t. 2.f.P.
16 <i>ferulaceum</i> . w. (wh.) Fennel-leaved.	Levant. 1752.	— H. ȝ.	
17 <i>pratense</i> . Led. (wh.) field.	Siberia. 1827.	— H. ȝ.	
18 <i>laxiflorum</i> . f. (wh.) loose-flowered.	—	1828. — H. ȝ.	
LOPHOSCIADIUM. DC. LOPHOSCIADIUM. Pentandria Digynia.			
millefolium. DC. (ye.) milfoil-leaved.	Levant. 1823.	7. 8. H. ȝ.	
Ferula meoides. s.s.	Madeira. 1785.	6. 7. G. ȝ.	Wend. s. han. 3. t. 13.

Tribus XI. DAUCINEÆ. DC.

ART'EDIA. S.S. ART'EDIA. Pentandria Digynia. L.			
squamata. s.s. (wh.) Fennel-leaved.	Levant. 1740.	7. H. ⊖.	Lam. ill. t. 193.
ORLA'YA. H.U. ORLA'YA. Pentandria Digynia.			
1 <i>grandiflora</i> . H.U. (wh.) great-flowered.	S. Europe. 1648.	7. 8. H. ⊖.	Jacq. aust. 1. t. 54.
<i>Caucalis grandiflora</i> . s.s.			
2 <i>maritima</i> . K.U. (bh.) sea-side.	—	1825. — H. ⊖.	Cavan. icon. 2. t. 101.
<i>Caucalis maritimus</i> . s.s.			
3 <i>platycarpus</i> . K.U. (wh.) broad-seeded.	—	1800. — H. ⊖.	Jacq. vind. 3. t. 10.
DAUCUS. S.S. CARROT. Pentandria Digynia. L.			
1 <i>Carota</i> . w. (wh.) wild.	Britain.	6. 7. H. ♂.	Eng. bot. v. 17. t. 1174.
β <i>lutea</i> .	(wh.) yellow-rooted, garden.—	— H. ♂.	
γ <i>aurantia</i> .	(wh.) long-orange.	— H. ♂.	
δ <i>præcox</i> .	(wh.) early horn.	— H. ♂.	
2 <i>mauritanicus</i> . w. (wh.) fine-leaved.	Spain. 1768.	— H. ♂.	All. ped. 2. t. 61. f. 1.
3 <i>polygamus</i> . L.en.(wh.) polygamous.	—	— H. ♂.	Jacq. vind. 3. t. 78.
4 <i>pissulus</i> . R.S. (wh.) small.	N. America. 1812.	— H. ♂.	
5 <i>maritimus</i> . E.B. (ro.) sea-side.	England.	— H. ♂.	Eng. bot. v. 36. t. 2560.
6 <i>gummifer</i> . s.s. (ro.) gum-bearing.	Italy. 1722.	— H. ♂.	Moris. sect. 9. t. 13.
7 <i>prolifer</i> . s.s. (wh.) proliferous.	Sicily. 1828.	— H. ♂.	Desf. atl. t. 63.
8 <i>hispidus</i> . s.s. (bh.) hispid.	Barbary. 1804.	— H. ♂.	
9 <i>siculus</i> . s.s. (bh.) Sicilian.	—	— H. ♂.	
10 <i>Gingidium</i> . s.s. (ro.) shining-leaved.	France. 1722.	— H. ♂.	Moris. s. 9. t. 13. f. 4.
<i>lucidus</i> . w.			
11 <i>muricatum</i> . s.s. (bh.) prickly-seeded	Barbary. 1683.	— H. ⊖.	Moris. s. 9. t. 14. f. 4.
12 <i>litoralis</i> . s.s. (wh.) sea-shore.	Greece. 1823.	— H. ⊖.	Flor. græc. t. 271.
13 <i>montevideensis</i> . Ot. (wh.) Monte Video.	Monte Video. 1826.	— H. ⊖.	
14 <i>parviflorus</i> . s.s. (wh.) small-flowered.	Barbary. 1823.	— H. ♂.	Desf. atl. 1. t. 68.
15 <i>crinitus</i> . K.U. (wh.) whorl-leaved.	—	1804. 6. 7. H. ȝ.	— 1. t. 62.
<i>Torilis crinita</i> . s.s.			
16 <i>pulcherrimus</i> . DC. (wh.) beautiful.	Caucasus. 1816.	— H. ♂.	
<i>Caucalis pulcherrima</i> . M.B.			
17 <i>orientalis</i> . DC. (wh.) oriental.	—	1699. — H. ♂.	Buxb. cent. 3. t. 28.
<i>Caucalis orientalis</i> . R.S.			
18 <i>Marschallianus</i> . DC. (wh.) Marschall's.	Odessa. 1824.	— H. ♂.	
<i>Caucalis littoralis</i> . M.B. <i>Cachrys littoralis</i> . s.s.			

Subordo II. CAMPYLOSPERMEEÆ. DC.

Tribus XII. CAUCALINEÆ. DC.

CA'UCALIS. S.S. BUR-PARSLEY. Pentandria Digynia. L.			
1 <i>daucoides</i> . s.s. (ro.) small.	England.	7. 8. H. ⊖.	Eng. bot. v. 3. t. 197.

2 mauritànica. s.s. (wh.)	Barbary.	Barbary.	1823.	7. 8. H. ⊖.	
3 leptophylla. s.s. (bh.)	fine-leaved.	Europe.	1739.	— H. ⊖.	Jacq. vind. t. 195.
4 pùmila. w.	(bh.) dwarf.	—	1640.	— H. ⊖.	Ger. gallop. n. 4. t. 10.
5 hispida. s.s. (st.)	hispid.	S. Europe.	1825.	— H. ⊖.	
6 humiliis. Lag.	(wh.) low.	—	—	— H. ⊖.	
TURG'ENIA. DC.	TURG'ENIA.	Pentandria Digynia.			
latifòlia. DC.	(bh.) broad-leaved.	England.	7. 8. H. ⊖.	Eng. bot. v. 3. t. 198.
Caucalis latifòlia. E.B.					
TOR'ILIS. S.S.	TOR'ILIS.	Pentandria Digynia. S.S.			
1 Anthriscus. s.s. (bh.)	upright.	Britain.	7. 8. H. ⊖.	Eng. bot. v. 14. t. 987.
2 infésta. E.F.	(wh.) spreading.	—	— H. ⊖.	— v. 19. t. 1314.
helvética. s.s. arvénica. L.en.	Le. Caucalis infésta. E.B.				
3 negligéta. R.S.	(wh.) confused.	Austria.	— H. ⊖.	Jacq. aust. t. 29.
4 ucrànica. s.s. (wh.)	Ukraine.	Ukraine.	1828.	— H. ⊖.	
5 nodòsa. s.s. (wh.)	knotted.	Britain.	5. 7. H. ⊖.	Eng. bot. v. 3. t. 199.
6 trichospérma. s.s. (fl.)	hairy-seeded.	Egypt.	1818.	— H. ⊖.	

Tribus XIII. SCANDICINEÆ. DC.

SCA'NDIX. DC.	CICELY.	Pentandria Digynia. L.			
1 Pécten. R.S.	(wh.) Venus's-comb.	Britain.	6. 7. H. ⊖.	Eng. bot. v. 20. t. 1397.
2 australis. R.S.	(wh.) southern.	S. Europe.	1713.	5. 7. H. ⊖.	Flor. græc. t. 285.
W'ylia australis. H.U.					
3 falcata. R.S.	(wh.) radiated.	Tauria.	—	— H. ⊖.	Hoff. umb. t. 2. f. 2.
W'ylia radians. H.U.					
4 grandiflora. R.S. (wh.)	large-flowered.	—	1827.	6. 7. H. ⊖.	Hoff. umb. p. 15. t. 2. f. 3.
5 brachycarpa. s.s. (wh.)	short-seeded.	Sicily.	1829.	— H. ⊖.	
6 ibérica. R.S.	(wh.) Iberian.	Iberia.	1828.	— H. ⊖.	Hoff. umb. t. 2. f. 4.
7 pinnatifida. R.S. (wh.)	cut-leaved.	Persia.	1805.	5. 7. H. ⊖.	Venten. cels. t. 14.
8 parviflora. R.S. (wh.)	small-flowered.	Tauria.	1827.	— H. ⊖.	
ANTHRI'SCUS. S.S.	ROUGH CHERVIL.	Pentandria Digynia. S.S.			
1 vulgàris. s.s. (wh.)	common.	Britain.	5. 6. H. ⊖.	Eng. bot. v. 12. t. 818.
Scàndix anthriscus. E.B.					
2 trichospérma. R.S. (wh.)	hairy-seeded.	Persia.	1827.	— H. ⊖.	Hoff. umb. 2.t.1.B.f.19.
3 nemorosa. s.s. (wh.)	wood.	Caucasus.	1823.	7. 8. H. ⊖.	W. et K. hung. 3.t. 224.
4 fumarioïdes. s.s. (wh.)	Fumitory-like.	Hungary.	1818.	— H. ♂.	Jacq. aust. 1. t. 63.
5 bulbösa. DC.	(wh.) bulbous-rooted.	Europe.	1726.	6. 7. H. ♂.	
Myrrhis bulbösa. s.s.	Charophyllum bulbosum. Jacq.				
6 sylvestris. K.U.	(wh.) smooth.	Britain.	4. 5. H. ⊖.	Eng. bot. v. 11. t. 752.
Chærophyllum sylvestre. E.B.					
7 Cerefolium. K.U. (wh.)	true Chervil.	—	6. 7. H. ⊖.	Eng. bot. v. 18. t. 1268.
Scàndix cerefolium. E.B.					
8 capensis. s.s. (wh.)	Cape.	C. B. S.	1816.	— G. ⊖.	
CHÆROPHYLLUM. S.S.	CHERVIL.	Pentandria Digynia. L.			
1 aromáticum. K.U. (wh.)	aromatic.	Scotland.	6. 8. H. ⊖.	Jacq. aust. 2. t. 150.
Myrrhis aromática. s.s.					
2 angelicæfölium. M.B. (wh.)	Angelica-leav'd.	Iberia.	1828.	6. 8. H. ⊖.	Jacq. aust. 2. t. 148.
3 hirsutum. W.	(wh.) hairy-leaved.	Switzerland.	1759.	6. 7. H. ⊖.	
4 cicutariöm. L.en. (bh.)	Cicuta-like.	S. Europe.	1816.	— H. ⊖.	Eng. bot. v. 30. t. 2103.
5 aüratum. L.	(st.) golden.	Scotland.	6. H. ⊖.	— v. 22. t. 1521.
6 temulémentum. E.B. (wh.)	rough.	Britain.	7. 8. H. ♂.	
7 monogònum. L.en. (wh.)	one-angled.	Hungary.	1820.	4. 6. H. ⊖.	
8 angulatum. Kit.	(wh.) angular.	—	—	— H. ⊖.	
9 humili. M.B.	(wh.) dwarf.	—	1825.	— H. ⊖.	
Anthriscus humili. Besser.	Myrrhis humili. R.S.				
10 nitidum. Wahl.	(wh.) glossy-seeded.	Carpathian-m.	1820.	— H. ⊖.	
11 Biebersteiniä. Lag.	(bh.) Bieberstein's.	Caucasus.	1824.	5. 6. H. ⊖.	Jacq. vind. 3. t. 25.
12 nodòsum. K.U.	(wh.) knotted.	Sicily.	1656.	5. 6. H. ⊖.	
Scàndix nodòsa. Jacq.	Anthriscus nodòsa. s.s.				
13 procumbens. Ph.	(wh.) procumbent.	N. America.	1699.	7. 8. H. ⊖.	Moris. his. s. 9.t. 11.f. ult.
14 ròseum. M.B.	(ro.) rose-coloured.	Iberia.	1824.	6. 7. H. ♂.	Hoff. umb. t. 1.B. f. 33.
15 coloratüm. W.	(st.) color'd-bract'd.	Illyria.	1806.	7. 8. H. ♂.	
16 tenuisölüm. Poir.	(wh.) slender-leaved.	Morocco.	—	— F. ♂.	Jacq. vind. t. 51.

SPHALLEROCA'RPUS. DC. SPHALLEROCA'RPUS. Pentandria Digynia. gracilis. DC. (wh.) slender. Siberia. 1827. 6. 8. H. ♂.	DC. dis. umb. t. 2.f. N.
<i>Myrrhis gracilis.</i> s.s. <i>Chærophyllum Cy'minum.</i> F.	
MOLOPOS'PERMUM. K.U. MOLOPOS'PERMUM. Pentandria Digynia. peloponnesiacum. K.U. (pa.) Hemlock-l'd. Switzerland. 1596. 5. 7. H. ♀.	Jacq. aust. app. t. 13.
<i>Ligusticum peloponnesiacum.</i> s.s.	
MYRRHIS. S.S. MYRRHIS. Pentandria Digynia. S.S. odorata. s.s. (wh.) sweet-scented. Britain. 5. 6. H. ♀.	Eng. bot. v. 10. t. 697.
OSMORH'IZA. DC. OSMORH'IZA. Pentandria Digynia. Claytöni. DC. (wh.) sweet-rooted. N. America. 1806. 7. 8. H. ♀.	Spreng. umb. t. 3. f. 6.
<i>Myrrhis Claytöni.</i> s.s. <i>Uraspermum hirsutum.</i> Bigelow. <i>Claytöni.</i> n.	

Tribus XIV. SMYRNEÆ. DC.

LAG'ŒCIA. L. WILD CUMIN. Pentandria Monogynia. L. cuminoïdes. s.s. (gr.ye.) common. Levant. 1640. 6. 7. H. ♂.	Flor. græc. t. 243.
OLIV'ERIA. V. OLIV'ERIA. Pentandria Digynia. S.S. decumbens. v. (bh.) Thyme-scented. Bagdad. 1816. 5. 7. H. ♂.	Venten. cels. t. 21.
ECHINO'PHORA. S.S. SEA PARSNEP. Pentandria Digynia. L. 1 spinosa. s.s. (bh.) prickly. England. 7. H. ♀.	Eng. bot. v. 34. t. 2413.
2 tenuifolia. s.s. (bh.) fine-leaved. Apulia. 1731. 7. 8. H. ♀.	Flor. græc. t. 266.
3 trichophylla. s.s. (bh.) slender-leaved. Armenia. 1829. — H. ♀.	
ARCT'OPUS. S.S. ARCT'OPUS. Polygamia Dicecia. L. echinatus. s.s. (wh.) rough. C. B. S. 1774. 6. 9. G. ♀.	Bot. reg. t. 705.
CA'CHRYS. Len. CA'CHRYS. Pentandria Digynia. L. 1 odontalgica. s.s. (ye.) Pallas's. Siberia. 1824. 7. 8. H. ♀.	Pall. reis. ap. 3. t. 9.
2 Morisoni. s.s. (ye.) Morison's. S. Europe. 1710. — H. ♀.	Moris. umb. t. 3. f. 1.
PERLEBIA. DC. PERLEBIA. Pentandria Digynia. triquête. DC. (wh.) winged. Levant. 1816. 6. 7. H. ♀.	Vent. cels. t. 79.
<i>Laserpitium triquetrum.</i> v. <i>Cächrys triquête.</i> s.s.	
HIPPOMAR'ATHRUM. Hippomar'athrum. Pentandria Digynia. siculum. L.en. (ye.) hairy-seeded. Sicily. 1640. 7. 8. H. ♀.	Moris. umb. t. 3. f. 2.
<i>Cächrys sicula.</i> s.s.	
MAGYD'ARIS. DC. MAGYD'ARIS. Pentandria Digynia. panacina. DC. (wh.) Parsnep-leav'd. Sicily. 1752. 7. 8. H. ♀.	Bocc. sic. 1. t. 1.
<i>Cächrys panacifolia.</i> v. <i>Athamanta panacifolia.</i> R.S.	
HE'RMAS. S.S. HE'RMAS. Polygamia Monocelia. W. 1 gigantea. s.s. (pu.) gigantic. C. B. S. 1794. 6. 7. G. ♀.	Thunb. act. pet. 14.t.11.
2 villosa. s.s. (bh.) villous. — 1795. — G. ♀.	Burm. afr. t. 71. f. 2
depauperata. I..	
1 ciliata. s.s. (wh.) ciliated. — 1816. — G. ♀.	Burm. t. 72. f. 1.
capitata. s.s. (wh.) headed. — 6. 8. G. ♀.	Thun. n. act. pet. 14.t.12.
5 quinquedentata. s.s. (wh.) toothed. — 1821. — G. ♀.	— 14. p. 533. t. 12.
CONIUM. H.U. HEMLOCK. Pentandria Digynia. L. 1 maculatum. s.s. (wh.) common. Britain. 6. 7. H. ♂.	Eng. bot. v. 17. t. 1191.
2 croaticum. w.k. (wh.) Hungarian. Hungary. — H. ♂.	
ARRAC'ACIA. DC. ARRAC'ACIA. Pentandria Digynia. esculent. DC. (wh.) esculent. S. America. 1822. 4. 5. F. ♀.	Hook. ex. flor. t. 152.
<i>Conium Arracacha.</i> H.E.F.	
PLEUROSPERMUM. S.S. PLEUROSPERMUM. Pentandria Digynia. S.S. austriacum. s.s. (wh.) Austrian. Austria. 1596. 6. 8. H. ♀.	Jacq. aust. 2. t. 151.
<i>Ligusticum austriacum.</i> w.	
SMYRNIUM. S.S. ALEXANDERS. Pentandria Digynia. L. 1 Olus'atum. s.s. (st.) common. " Britain. 5. 6. H. ♂.	Eng. bot. v. 4. t. 230.
2 perfoliatum. w. (st.) perfoliate. Italy. 1596. 5. H. ♀.	Lobel. ic. 709.
<i>Dodon'e.</i> s.s.	
3 Dioscoridis. s.s. (ye.) Dioscorides. S. Europe. — H. ♂.	Flor. græc. t. 289.
4 apuifolium. s.s. (ye.) Smallage-leav'd. Candia. 1731. 5. 7. H. ♀.	
5 nudicaulis. s.s. (ye.) naked-stalked. Caucasus. 1820. — H. ♀.	Spreng. umb. t. 4. f. 7.
6 cicutarium. s.s. (ye.) Cicuta-like. — 1829. — H. ♀.	
7 cordatum. s.s. (ye.) heart-leaved. N. America. 1759. 6. 7. H. ♂.	
8 integrerrimum. Ph. (ye.) entire-leaved. N. America. 1759. — H. ♀.	
PHYSOSPERMUM. S.S. PHYSOSPERMUM. Pentandria Digynia. S.S. commutatum. s.s. (wh.) Cornish. England. 7. 8. H. ♀.	Eng. bot. v. 10. t. 683.
<i>Ligusticum cornubiense.</i> E.B.	

Subordo III. CÆLOSPERMEÆ. DC.

Tribus XV. CORIANDREÆ. DC.

BIF'ORIS. S.S.	BIF'ORIS. Pentandria Digynia. S.S.				
1 flosculosa. s.s. (wh.) flosculose.	S. Europe. 1640.	6. 7.	H. ⊖.	Moris. sect. 9. t. 11.	
Còrion testiculatum. L.en. Coriàndrum testiculatum. L.					
2 ràdians. s.s. (wh.) rayed.	Tauria.		H. ⊖.	Hoff. umb. ed. 2.t. tit. f. 2.	
ATR'EMA. DC.	ATR'EMA. Pentandria Digynia.				
americana. dc. (wh.) American.	N. America. 1827.	6. 7.	H. ⊖.	DC. mem. umb. t. 18.	
Coriàndrum americanum. n.					
CORIA'NDRUM. S.S.	CORIANDER. Pentandria Digynia. L.				
sativum. s.s. (bh.) common.	England.	6.	H. ⊖.	Eng. bot. v. 1. t. 67.	

ORDO CIX.

ARALIACEÆ. Juss. gen. p. 217.

P'ANAX. L.	P'ANAX. Polygamia Dicecia. L.				
1 quinquefòlium. w.(gr.)	Ginseng.	N. America. 1740.	6.	H. Ȑ.	Bot. mag. t. 1333.
2 trifòlium. w.	(wh.) lesser.	1759.	5. 6.	H. Ȑ.	_____ t. 1334.
pusilla. a. B.M.					
3 fruticòsum. w.	(gr.) shrubby.	Ternate.	1800.	8. 9.	S. Ȑ.
4 digitatùm. H.B.	(gr.) fingered.	E. Indies.	1818.	S. Ȑ.
5 arboreùm. w.	(gr.) tree.	N. Zealand.	1822.	G. Ȑ.
6 chrysophyllum. R.S.	(gr.) golden-leaved.	Guiana.	—	S. Ȑ.
7 conchifòlium. H.B.	(gr.) shell-leaved.	—	1820.	S. Ȑ.
8 attenuatùm. w.	(gr.) attenuated.	W. Indies.	1825.	S. Ȑ.
9 aculeatùm. w.	(gr.) prickly.	China.	1773.	11.	S. Ȑ.
CUSS'ONIA. W.	Cuss'ONIA. Pentandria Digynia. L.				
1 thysiflòra. w.	(gr.) thyrsse-flower'd.	C. B. S.	1795.	8. 10.	G. Ȑ.
2 triptera. C.R.	(gr.) tripterous.	—	1816.	G. Ȑ.
3 spicatà. w.	(gr.) spike-flowered.	—	1789.	8. 10.	G. Ȑ.
SCIODAPHY'LLUM. S.S.	SCIODAPHY'LLUM. Pentandria Pentagynia. S.S.				
1 pentàndrum. s.s. (bh.)	pentandrous.	Peru.	1828.	G. Ȑ.
2 acuminatùm. s.s. (ye.)	acuminate-petal'd.	—	1818.	S. Ȑ.
Actinophy'llum acuminatùm. R.P.					_____ t. 310.
3 cònicum. s.s.	(bh.) conical-flower'd.	—	—	S. Ȑ.
4 pedicellatùm. s.s. (bh.)	pedicelled.	—	—	S. Ȑ.
Actinophy'llum pedicellatùm. R.P.					_____ t. 308.
5 anglatùm. s.s. (ye.)	tree.	—	1824.	G. Ȑ.
6 Bròwnii. s.s.	(gr.) upright.	W. Indies.	1793.	6. 8.	S. Ȑ.
Aralia Sciodaphy'llum. w.					Brown jam. t. 19. f. 1. 2.
AR'ALIA. D.P.	AR'ALIA. Pentandria Pentagynia. L.				
1 pentaphylla. s.s. (gr.)	five-l'd prickly.	China.	1822.	G. Ȑ.
2 spinosa. s.s.	(wh.) Angelica-tree.	Virginia.	1688.	9.	H. Ȑ.
3 racemosa. s.s.	(gr.) berry-bearing.	N. America.	1658.	6. 9.	H. Ȑ.
4 nudicaulis. s.s.	(wh.) naked-stalked.	—	1731.	6. 7.	H. Ȑ.
5 hispida. s.s.	(wh.) hispid.	—	1799.	—	H. Ȑ.
6 pubescens. dc.	(wh.) pubescent.	New Spain.	1819.	G. Ȑ.
GAST'ONIA. J.	GAST'ONIA. Dodecandria Monogynia.				
palmata. H.B.	(wh.) palmate-leaved.	E. Indies.	1818.	3. 4.	f.
H'EDERA. D.P.	Ivy. Pentandria Mono-Pentagynia.				
1 umbraculifera. (wh.)	large.	E. Indies.	1818.	3. 4.	S. Ȑ.
Aralia umbraculifera. H.B.					
2 digitata. H.B.	(wh.) fingered.	—	—	—	S. Ȑ.

3 aculeata. D.P.	(wh.) prickly Nepaul.	Nepaul.	1816.	F. h.	
4 frâgrans. D.P.	(wh.) fragrant.	—	—	F. h.	
5 cochleata.	(wh.) shell-leaved.	E. Indies.	1820.	S. h.	Rumph. amb. 4. t. 31.
Aralia cochleata. R.S.						
arborea. R.S.	(wh.) tree.	Jamaica.	—	S. h.	Jacq. schœnb. t. 51.
7 capitata. R.S.	(gr.) cluster-flower'd.	W. Indies.	1777.	8.	S. h.	Smith ic. pict. 4.
canariensis. R.S.	(gr.) Irish.	Canaries.	1807.	9. 10.	H. h.	Wild. ber. m. 2.t.5.f.1.
9 Hélix. R.S.	(gr.) common.	Britain.	—	—	H. h.	Eng. bot. v. 18. t. 1267.
β poética. R.S.	(gr.) Poet's.	—	—	—	H. h.	
γ álbo-variegata. (gr.) white-variegated.	—	—	—	—	H. h.	
δ flávo-variegata. (gr.) yellow-variegated.	—	—	—	—	H. h.	

ORDO CX.

CAPRIFOLIACEÆ. Juss. gen. p. 210.

CORNUS. L.

DOGWOOD. Tetrandria Monogynia. L.

1 alternifolia. R.S.	(wh.) alternate-leav'd.	N. America.	1760.	9.	H. h.	Schm. arb. 2. t. 70.
2 paniculata. R.S.	(wh.) panicle'd.	—	1758.	6. 7.	H. h.	— 2. t. 68.
3 stricta. R.S.	(wh.) upright.	—	—	—	H. h.	— 2. t. 67.
β variegata.	(wh.) variegated-leav'd.	—	—	—	H. h.	
4 oblonga. F.I.	(bh.) oblong-leaved.	Nepaul.	1818.	7. 8.	H. h.	
5 macrophylla. F.I.	(wh.) large-leaved.	—	1827.	—	H. h.	
6 circinata. R.S.	(wh.) Pensylvanian.	N. America.	1784.	—	H. h.	Schm. arb. 2. t. 69.
7 alba. R.S.	(wh.) white berried.	Siberia.	1741.	6. 9.	H. h.	— 2. t. 65.
8 sibirica. L.C.	(wh.) Siberian.	—	—	—	H. h.	
9 sevicia. R.S.	(wh.) blue-berried.	N. America.	1683.	8.	H. h.	Schm. arb. 2. t. 64.
10 sanguinea. R.S.	(wh.) common.	Britain.	—	6. 7.	H. h.	Eng. bot. v. 4. t. 249.
β variegata.	(wh.) variegated.	—	—	—	H. h.	
11 máscula. R.S.	(ye.) CornelianCherry.	Europe.	1596.	2. 4.	H. h.	Flor. græc. t. 151.
β variegata.	(ye.) variegated.	—	—	—	H. h.	
12 florida. R.S.	(ye.) great-flowered.	N. America.	1731.	4. 5.	H. h.	Bot. mag. t. 526.
13 capitata. F.I.	(ye.) headed.	Nepaul.	1820.	—	H. h.	
14 canadensis. R.S.	(wh.) Canadian.	Canada.	1774.	6. 8.	H. 4.	Bot. mag. t. 880.
15 suécica. R.S.	(wh.) dwarf.	Britain.	—	4.	H. 4.	Eng. bot. v. 5. t. 310.

SAMBUCUS. L.

ELDER. Pentandria Trigynia. L.

1 Ebulus. R.S.	(fl.) herbaceous.	Britain.	—	6. 7.	H. 4.	Eng. bot. v. 7. t. 475.
2 hùmilis. Mill.	(fl.) dwarf.	—	—	—	H. 4.	
3 chinensis. H.T.	(fl.) Chinese.	China.	1825.	—	H. 4.	
4 nigra. R.S.	(wh.) common.	Britain.	—	5. 7.	H. h.	Eng. bot. v. 7. t. 476.
β variegata.	(wh.) variegated.	—	—	—	H. h.	
γ aurea.	(wh.) golden-leaved.	—	—	—	H. h.	
δ pulvérulenta.	(wh.) powdered-leaved.	—	—	—	H. h.	
ε monstrosa.	(wh.) deformed.	—	—	—	H. h.	
ζ virescens.	(wh.) green-fruited.	—	—	—	H. h.	
η álbida.	(wh.) white-fruited.	—	—	—	H. h.	
5 ciniata. L.en.	(wh.) Parsley-leaved.	—	—	—	H. h.	Schm. arb. t. 144.
6 hybrida. Jac.	(wh.) hybrid.	—	1826.	—	H. h.	
7 rotundifolia. L.C.	(wh.) round-leaved.	Wiltshire.	—	—	H. h.	
8 dénisia. R.S.	(wh.) Canadian.	N. America.	1761.	6. 8.	H. h.	Schm. arb. t. 142.
demosa. R.S.	(st.) red-berried.	S. Europe.	1596.	5. 6.	H. h.	Jacq. ic. 1. t. 59.
β purpurea.	(br.) purple-flowered.	—	—	—	H. h.	
10 pubescens.	(st.) pubescent.	N. America.	1812.	—	H. h.	

VI

VIBURNUM. Pentandria Trigynia. L.

1 mucronatum. R.S.	(wh.) soft.	N. America.	1812.	6. 7.	H. h.	
2 acerifolium. R.S.	(wh.) Maple-leaved.	—	1736.	—	H. h.	
3 édoule. R.S.	(wh.) eatable-fruited.	—	1812.	7.	H. h.	
4 Oxycoccos. R.S.	(bh.) Cranberry-like.	—	—	—	H. h.	
5 O'pulus. R.S.	(wh.) Guelder Rose.	Britain.	—	5. 6.	H. h.	Eng. bot. v. 5. t. 332.
β roseum. R.S.	(wh.) Snowball tree.	—	—	—	H. h.	Schm. arb. t. 174.
γ variegatum.	(wh.) variegated-leav'd.	—	—	—	H. h.	
6 davuricum. R.S.	(wh.) Siótan.	Dahuria.	1785.	6. 7.	H. h.	Pall. ross. 1. t. 38.
7 lantanoides. R.S.	(wh.) Lantana-like.	N. America.	—	5. 6.	H. h.	
8 Lantana. R.S.	(wh.) wayfaring-tree.	Britain.	—	—	H. h.	Eng. bot. v. 5. t. 331.
9 nervosum. D.P.	(wh.) nerved-leaved.	Nepaul.	1826.	—	H. h.	

CAPRIFOLIACEÆ.

10 Lentago. R.S. (<i>wh.</i>) tree.	N. America.	1761.	7.	H. $\frac{1}{2}$.	Wats. dend. brit. t. 21.
11 squamatum. R.S. (<i>wh.</i>) scaly.	1812.	6. 7.	H. $\frac{1}{2}$.	_____ t. 24.	
12 pyrifolium. R.S. (<i>wh.</i>) pear-leaved.	_____	5. 6.	H. $\frac{1}{2}$.	_____ t. 22.	
13 prunifolium. R.S. (<i>wh.</i>) plum-leaved.	1731.	—	H. $\frac{1}{2}$.	_____ t. 23.	
14 pubescens. R.S. (<i>wh.</i>) pubescent.	1736.	6. 7.	H. $\frac{1}{2}$.		
15 dentatum. R.S. (<i>wh.</i>) tooth-leaved.	—	—	H. $\frac{1}{2}$.		
16 laevigatum. R.S. (<i>wh.</i>) Cassioberry bush.	1724.	7. 8.	H. $\frac{1}{2}$.		
17 nitidum. R.S. (<i>wh.</i>) shining-leaved.	1758.	5. 6.	H. $\frac{1}{2}$.		
18 cassinoïdes. R.S. (<i>wh.</i>) thick-leaved.	1761.	6. 7.	H. $\frac{1}{2}$.		
19 obovatum. R.S. (<i>wh.</i>) obovate-leaved.	1812.	5. 6.	H. $\frac{1}{2}$.		
20 odoratissimum. B.R. (<i>w.</i>) sweet-scented. China.	1810.	—	G. $\frac{1}{2}$.		
chinense. Zeyh.					
21 nudum. R.S. (<i>wh.</i>) oval-leaved.	N. America.	1752.	—	H. $\frac{1}{2}$.	
22 cylindricum. D.P. (<i>wh.</i>) tube-flowered.	Nepaul.	1826.	6. 8.	H. $\frac{1}{2}$.	
23 villosum. R.S. (<i>wh.</i>) villous.	Jamaica.	1820.	4. 6.	S. $\frac{1}{2}$.	
24 rugosum. R.S. (<i>wh.</i>) rugged-leaved.	Canaries.	1796.	4. 5.	H. $\frac{1}{2}$.	
rigidum. Vent. malm. t. 98.					
25 strictum. L.en. (<i>wh.</i>) upright Laurestine. S. Europe.	1596.	3. 5.	H. $\frac{1}{2}$.		
β virgatum. (<i>wh.</i>) twiggy.	—	—	H. $\frac{1}{2}$.		
26 Tinus. R.S. (<i>wh.</i>) common Laurestine.	—	—	12. 4.	H. $\frac{1}{2}$.	
27 lucidum. R.S. (<i>bh.</i>) shining Laurestine.	—	—	2. 5.	H. $\frac{1}{2}$.	
SYMPH'ORIA. S.S. St. PETER'S-WORT. Pentandria Monogynia.					
1 glomerata. s.s. (<i>bh.</i>) common.	N. America.	1730.	8. 9.	H. $\frac{1}{2}$.	
2 racemosa. s.s. (<i>bh.</i>) Snowberry.	—	—	1812.	7. 9.	H. $\frac{1}{2}$.
DIERVILLA. J. DIERVILLA. Pentandria Monogynia.					
humilis. p.s. (<i>ye.</i>) yellow-flower'd. N. America.	1739.	6. 7.	H. $\frac{1}{2}$.		
lutea Ph. Tournefortii. R.S. canadensis. s.s. Lonicera Diervilla. L.					
LONICERA. R.S. FLY-HONEYSUCKLE. Pentandria Monogynia. L.					
1 alpigena. s.s. (<i>re.</i>) red-berried.	Switzerland.	1596.	4. 5.	H. $\frac{1}{2}$.	
2 microphylla. s.s. (<i>re.</i>) small-leaved.	Siberia.	1818.	5. 6.	H. $\frac{1}{2}$.	
3 sibirica. R.S. (<i>re.</i>) Siberian.	—	—	—	H. $\frac{1}{2}$.	
4 orientalis. R.S. (<i>wh.</i>) oriental.	Levant.	1820.	—	H. $\frac{1}{2}$.	
caucasica. Pall.					
5 punicea. B.M. (<i>cr.</i>) crimson-flower'd.	—	—	4. 5.	H. $\frac{1}{2}$.	
6 coerulea. R.S. (<i>st.</i>) blue-berried.	Switzerland.	1629.	3. 4.	H. $\frac{1}{2}$.	
β altaica. F. (<i>st.</i>) Altay.	Altay.	1827.	—	H. $\frac{1}{2}$.	
7 canadensis. R.S. (<i>st.</i>) Canadian.	Canada.	1812.	4. 5.	H. $\frac{1}{2}$.	
8 involucrata. s.s. (<i>ye.</i>) involucred.	N. America.	1824.	5. 6.	H. $\frac{1}{2}$.	
9 ibérica. R.S. (<i>br.</i>) Iberian.	Iberia.	1818.	5. 6.	H. $\frac{1}{2}$.	
10 villosa. R.S. (<i>st.</i>) villous.	Canada.	—	4. 5.	H. $\frac{1}{2}$.	
Xylósteum villosum. M.					
11 Xylósteum. R.S. (<i>wh.</i>) fly.	England	—	6. 7.	H. $\frac{1}{2}$.	
12 canescens. R.S. (<i>wh.</i>) canescent.	Barbary.	1820.	—	H. $\frac{1}{2}$.	
biflora. Desf.					
13 ciliata. R.S. (<i>bh.</i>) ciliated.	N. America.	1824.	5. 6.	H. $\frac{1}{2}$.	
β alba. R.S. (<i>wh.</i>) white berried.	—	—	—	H. $\frac{1}{2}$.	
14 tatárica. R.S. (<i>wh.</i>) Tartarian.	Tartary.	1752.	4. 5.	H. $\frac{1}{2}$.	
α ábla.					
β rúbra.					
15 nigra. R.S. (<i>pu.</i>) black-berried.	Switzerland.	1597.	3. 4.	H. $\frac{1}{2}$.	
16 flexuosa. R.S. (<i>wh.</i>) flexuose.	China.	1806.	7. 9.	H. $\frac{1}{2}$.	
17 quadrifolia. R.S. (<i>wh.</i>) four-leaved.	Siberia.	1824.	—	H. $\frac{1}{2}$.	
18 pyrenaea. R.S. (<i>wh.</i>) Pyrenean.	Pyrenees.	1739.	5.	H. $\frac{1}{2}$.	
19 hispida. R.S. (<i>wh.</i>) hispid-branch'd.	Dahuria.	1821.	—	H. $\frac{1}{2}$.	
20 ligustrina. D.P. (<i>st.</i>) Privet-leaved.	Nepaul.	1825.	4. 5.	H. $\frac{1}{2}$.	
NINTO'OA. NINTO'OA. Pentandria Monogynia.					
1 longiflora. (<i>st.</i>) long-flowered.	China.	1816.	5. 10.	F. $\frac{1}{2}$.	Bot. reg. t. 1233.
Caprifolium longiflorum. B.R. japonicum. D.P.					
2 confusa. (<i>st.</i>) pale-flowered.	China.	1806.	7. 10.	F. $\frac{1}{2}$.	Andr. deposit. t. 583.
<i>Nin-foo</i> . Kämpf. amoen. 5. p. 785. Lonicera japonica. Bot. reg. t. 70. non Thunb.					
3 japonica. Th. (<i>re.</i>) red-flowered.	China.	1814.	4. 11.	H. $\frac{1}{2}$.	Wats. dend. brit. t. 117.
<i>Lonicera chinensis</i> . W.D. Lonicera flexuosa. Bot. reg. v. 9. t. 712. nec aliorum.					
CAPRIFOLIUM. R.S. HONEYSUCKLE. Pentandria Monogynia.					
1 italicum. R.S. (<i>bh.</i>) early.	England.	—	4. 6.	H. $\frac{1}{2}$.	Eng. bot. v. 12. t. 799.
ábum.					
<i>Lonicera Caprifolium</i> . E.B.					
β rúbrum. (<i>re.</i>) early red.	Italy.	—	—	H. $\frac{1}{2}$.	Schmidt arb. t. 106.
2 etruscum. R.S. (<i>wh. pu.</i>) Italian.	—	—	1824.	—	San. Viag. al. m. t. 1.
3 baleàricum. R.S. (<i>st.</i>) red-barked.	S. Europe.	—	—	H. $\frac{1}{2}$.	

4 dioicum. R.S. (re.) small-flowered. N. America. 1766.	6. 7. H. h. Bot. reg. t. 138.
<i>parviflorum</i> . Ph? <i>bracteosum</i> . M? <i>Lonicera media</i> . Murr. com. got. 1776. t. 3.	
5 gratum. R.S. (re.) evergreen. N. America. 1730.	6. 8. H. h. Hort. angl. n. 10. t. 8.
6 flavum. B.M. (ye.) golden-flowered. ———	1810. 5. 7. H. h. Bot. mag. t. 1318.
<i>Fraseri</i> . Ph. <i>Lonicera flava</i> . B.M.	
7 Douglassii. H.T. (ye.) Douglas's. Canada. 1824.	5. 7. H. h. Bot. mag. t. 781.
8 pubescens. H.E.F. (ye.) pubescent. N. America. 1820.	——— H. h. Hook. ex. flor. t. 27.
<i>Lonicera Goldii</i> . S.S.	
9 sempervirens. R.S. (sc.) Trumpet. ———	1656. 5. 8. H. h. Bot. mag. t. 781.
10 oblongifolium. (sc.) oblong-leaved. Carolina. ———	——— F. h. ——— t. 1753.
<i>Lonicera sempervirens</i> . β minor. B.M.	
11 ciliatum. R.S. (ye.) fringed. N. America. 1827.	——— H. h. ———
12 impléxum. R.S. (pa.) glaucous-leaved. Minorca. 1772.	6. 9. H. h. Bot. mag. t. 640.
13 Pericly'menum. R.S. (st.) Woodbine. Britain. ———	5. 7. H. h. Eng. bot. v. 12. t. 800.
α vulgáris. H.K. (st.) common. ———	——— H. h. Schmidt arb. t. 107.
β serotinum. H.K. (re.) late red. ———	——— 6. 8. H. h. ——— t. 108.
γ belgicum. H.K. (bh.) dutch. ———	——— 5. 7. H. h. Hort. angl. n. 5. t. 6.
δ quercifolium. H.K. (st.) Oak-leaved. Britain. ———	——— H. h. ———
ϵ variegátum. H.K. (st.) variegated-leav'd. ———	——— H. h. ———
TRIO'STEUM. R.S. FEVERWORT. Pentandria Monogynia. L.	
1 perfoliatum. R.S. (pu.) perfoliate. N. America. 1730.	6. 7. H. $\frac{1}{2}$.
2 angustifolium. R.S. (ye.) narrow-leaved. ———	1699. ——— H. $\frac{1}{2}$. Pluk. alm. t. 104. f. 2.
LINN'EA. W. LINN'EA. Didynamia Angiospermia. L.	
borealis. w. (bh.) two-flowered. Britain. ———	5. 8. H. $\frac{1}{2}$. Eng. bot. v. 7. t. 433.

ORDO CXI.

LORANTHEÆ. *Juss. ann. mus.* 12. p. 292.

LORA'NTHUS. L. LORA'NTHUS. Hexandria Monogynia.	
europæus. s.s. (st.) European. Europe. 1824. H. h. Jacq. aust. t. 30.
VI'SCUM. L. MISSELTOE. Diocia Tetrandria. L.	
álbum. E.B. (gr.) common. England. 5. H. h. Eng. bot. v. 21. t. 1470.	

ORDO CXII.

CHLORANTHEÆ. *R. Brown.*

CHLORA'NTHUS. R.S. CHLORA'NTHUS. Tetrandria Monogynia. R.S.	
1 in conspícuus. R.S. (st.) trailing. China. 1781. 5. 9. G. $\frac{1}{2}$. L'Her. sert. angl. t. 2.	
2 eréctus. H.S.L. (wh.) upright. ——— 1809. 6. 8. G. $\frac{1}{2}$.	
<i>elatiōr</i> . L.enum.	
3 odoriferus. (ye.) sweet-scented. ——— 1820. ——— G. $\frac{1}{2}$.	
<i>Creodus odorifer</i> . Loureiro.	
4 monostachys. s.s. (st.) single-spiked. ——— 1819. 2. 6. G. $\frac{1}{2}$. Bot. mag. t. 2190.	
5 monánder. B. (ye.) single-stamen'd. ——— 1817. 1. 12. G. $\frac{1}{2}$.	

ORDO CXIII.

LOPHIREÆ. *G. Don.*

LO'PHIRA. S.S. SCRUBBY OAK. Icosandria Monogynia. S.S.	
alata. s.s. (gr.) winged. SierraLeone. 1822. S. h.	

ORDO CXIV.

RUBIACEÆ. Juss. gen. 196.

Sect. I. GUETTARDEÆ. Kunth synops. 3. p. 67.

GUETTA'RDA. S.S.	GUETTA'RDA.	Monœcia Tetr-Pent-Hexandria.				
1 argéntea. s.s.	(wh.) silvery-leaved.	Jamaica.	1826.	S. h.	Brown. jam. t. 20. f. 1.
2 hirsüta. s.s.	(wh.) hairy.	Peru.	1822.	S. h.	Flor. per. 2. t. 145. f.a.
<i>Laugèria hirsüta.</i> R.P.						
3 tomentösa. s.s.	(wh.) tomentose.	Jamaica.	—	S. h.	Vahl. symb. 3. t. 57.
4 lúcida. s.s.	(wh.) glossy.	—	1818.	S. h.	
<i>Laugèria lúcida.</i> Sw.						
5 odoráta. R.S.	(br.) sweet-scented.	Havannah.	1827.	S. h.	Jacq.am.pic.t.259.f.16.
6 scábra. s.s.	(wh.) rough.	W. Indies.	—	S. h.	Vent. choix. 1. t. 1.
<i>Matthiola scábra.</i> L.						
7 rugösa. s.s.	(wh.) rough-leaved.	—	1793.	S. h.	
8 ellíptica. R.S. elliptic-leaved.	Jamaica.	1825.	S. h.	Rheed. mal. 4. t. 47.48.
9 speciosa. s.s.	(wh.) great-flowered.	E. Indies.	1771.	S. h.	
NONAT'ELIA. R.S.	NONAT'ELIA.	Pentandria Monogynia.				
1 racemösa. R.S.	(wh.) racemed.	Trinidad.	1825.	S. h.	Aubl. guian. 1. t. 72.
2 officinalis. R.S.	(wh.) officinal.	W. Indies.	1824.	S. h.	— 1. t. 70. f. 1.
<i>Psychotria involucrata.</i> W.						
VANGU'ERIA. J.	VANGU'ERIA.	Pentandria Monogynia. S.S.				
1 édulis. s.s.	(gr.) eatable.	E. Indies.	1809.	S. h.	Lam. ill. t. 159.
2 spinösa. F.I.	(gr.) spiny.	—	1818.	S. h.	
ERITHALIS. S.S.	ERITHALIS.	Pentandria Monogynia. L.				
fruticösa. s.s.	(wh.) shrubby.	Jamaica.	1793.	7. 8.	S. h.	Jacq.am.p.t. 260.f.20.
<i>odorifera.</i> R.S.						

Sect. II. HAMELIACEÆ. Kth. synops. 3. p. 63.

HAMELIA. S.S.	HAMELIA.	Pentandria Monogynia. L.				
1 axillär. s.s.	(ye.) axillary-flower'd.	Jamaica.	1820.	7. 8.	S. h.	
2 ventricösa. s.s.	(ye.) large-flowered.	W. Indies.	1778.	9. 11.	S. h.	Bot. reg. 1195.
<i>grandiflora.</i> par. lond. t. 55.						
3 chrysántha. s.s.	(ye.) yellow-flower'd.	—	1820.	—	S. h.	Jacq. ic. 2. t. 335.
4 pátns. s.s.	(sc.) spreading.	—	1752.	7. 8.	S. h.	Exot. bot. 1. t. 24.
5 sphærocárpa. P.S.(or.)	round-fruited.	Peru.	1811.	—	S. h.	Flor. per. t. 221. f. b.
ISE'R'TIA. S.S.	ISE'R'TIA.	Hexandria Monogynia. S.S.				
coccínea. s.s.	(sc.) scarlet-flower'd.	S. America.	1820.	S. h.	Aubl. guian. 1. t. 123.
SABI'CEA. R.S.	SABI'CEA.	Pentandria Monogynia.				
1 áspéra. R.S.	(wh.) rough.	Guiana.	1824.	S. h.	Aubl. guian. 1. t. 123.
2 hírta. R.S.	(wh.) hairy.	Jamaica.	1825.	S. h.	

Sect. III. GARDENIEÆ. Sal. par. lond. 46.

MUSS'ÆNDA. R.S.	MUSS'ÆNDA.	Pentandria Monogynia. L.				
1 corymbösa. F.I.	(or.) corymbose.	E. Indies.	1827.	5. 8.	S. h.	
2 frondösa. R.S.	(ye.) frondose.	—	1815.	5. 10.	S. h.	Bot. reg. t. 517.
3 glábra. R.S.	(ye.) smooth.	—	1820.	—	S. h.	
4 macrophylla. F.I.	(or.) large-leaved.	Nepaul.	1827.	S. h.	
5 pubéscent. R.S.	(ye.) pubescent.	China.	1805.	5. 9.	G. h.	Bot. mag. t. 2099.
6 arcuäta. R.S.	(ye.) arched.	Isle France.	1822.	S. h.	
LUCULIA. B.F.G.	LUCULIA.	Pentandria Monogynia.				
gratissima. B.F.G.(ro.)	sweet-scented.	Nepaul.	1818.	10. 2.	G. h.	Swt br. fl. gar. t. 145.
<i>Cinchona gratissima.</i> F.I.	<i>Mussænda Luculia.</i> D.P.					
RA'NDIA. R.S.	RA'NDIA.	Pentandria Monogynia. R.S.				
1 aculeäta. R.S.	(wh.) prickly.	W. Indies.	1733.	5. 6.	S. h.	Bot. mag. t. 1841.
<i>Gardénia Rändia.</i> W.						

2	obovata. R.S.	(wh.) obovate-leaved.	Peru.	1818.	5. 6.	S. h.	Flor. per. 2. t. 220. f. b.
3	rotundifolia. R.S.	(wh.) round-leaved.	_____	_____	_____	S. h.	_____
4	parvifolia. R.S.	(wh.) small-leaved.	Jamaica.	1825.	5. 7.	S. h.	Sloan. hist. 2. t. 207.f.1.
5	macrantha. R.S.	(wh.) large-flowered.	Sierra Leon.	1796.	8. 9.	S. h.	Sal. par. lond. t. 93.
	<i>longiflora</i> . P.L. non R.S.						
6	racemosa. H.B.	(wh.) racemed.	E. Indies.	1818.	S. h.	_____
7	<i>longiflora</i> . R.S.	(wh.) long-flowered.	_____	_____	S. h.	Lam. ill. t. 156. f. 3.
	<i>Gardenia multiflora</i> . W.						
8	stricta. F.I.	(wh.) upright.	_____	1824.	5. 7.	S. h.	_____
	<i>Macrocneum strictum</i> . R.S.						
9	horrida. R.S.	(wh.) large-spined.	China.	1825.	—	S. h.	_____
10	sinensis. R.S.	(wh.) Chinese.	_____	_____	—	S. h.	_____
POSOQUERIA. F.I. POSOQU'ERIA. Pentandria Monogynia.							
1	rígida. F.I.	(wh.) rigid.	Nepaul.	1825.	S. h.	_____
2	floribunda. F.I.	(wh.) many-flowered.	E. Indies.	_____	S. h.	_____
3	fasciculata. F.I.	(wh.) fascicle-flower'd.	_____	1824.	5. 6.	S. h.	_____
4	fragrans. F.I.	(wh.) fragrant.	_____	1822.	4. 6.	S. h.	_____
5	longispina. F.I.	(wh.) long-spined.	_____	1812.	—	S. h.	_____
	<i>Gardenia longispina</i> . s.s.						
6	nudans. F.I.	(wh.) drooping.	_____	1818.	7. 9.	S. h.	_____
7	dumetorum. F.I.	(wh.) bushy.	_____	1779.	—	S. h.	Roxb. corom. 2. t. 136.
	<i>Gardenia dumetorum</i> . w. <i>Canthium coronatum</i> . Lam.						
8	uliginosa. F.I.	(wh.) marsh.	E. Indies.	1802.	—	S. h.	Roxb. corom. 2. t. 135.
	<i>spinosa</i> .	(wh.) spiny.	China.	1804.	—	S. h.	Thunb. gar. n. 7.t.2.f.4.
	<i>Gardenia spinosa</i> . Th. <i>Canthium chinense</i> . P.S.						
10	longiflora. R.S.	(wh.) long-flowered.	Guiana.	1820.	S. h.	Aubl. guian. 1. t. 51.
GARD'ENIA. R.S. GARD'ENIA. Pentandria Monogynia. L.							
1	amœna. s.s.	(re.wh.) crimson-tipped.	China.	1810.	6. 8.	S. h.	Bot. mag. t. 1904.
2	armata. s.s.	(wh.) armed.	W. Indies.	1813.	S. h.	Jacq. amer. pict. t. 71.
	<i>Mussenda spinosa</i> . Jacq.						
3	tetracantha. s.s.	(wh.) four-thorned.	New Spain.	1818.	S. h.	_____
4	propinqua. B.R.	(wh.) strong-spined.	_____	1816.	5. 8.	S. h.	Bot. reg. t. 975.
5	montana. F.I.	(wh.) mountain.	E. Indies.	1818.	6. 8.	S. h.	_____
6	campanulata. F.I. (st.)	bell-flowered.	_____	1812.	—	S. h.	_____
7	micrantha. R.S.	(wh.) small-flowered.	China.	1806.	—	S. h.	Thunb. gar. n. 8.t. 1.f.2.
8	florida. s.s.	(wh.) Cape Jasmine.	_____	1754.	7. 10.	S. h.	Bot. reg. t. 449.
	β <i>flore-pleno</i> .	(wh.) double-flowered.	_____	—	—	S. h.	Ehret. pict. t. 15.
9	angustifolia. B.C.	(wh.) narrow-leaved.	_____	1820.	—	S. h.	Lodd. bot. cab. 512.
10	radicans. s.s.	(wh.) rooting.	_____	1804.	3. 6.	G. h.	Bot. reg. t. 73.
11	grandiflora. R.S.	(wh.) large-flowered.	_____	1816.	—	S. h.	Bot. mag. t. 2627.
	<i>florida</i> var. <i>ovalifolia</i> . Bot. mag. t. 2627.						
12	costata. F.I.	(wh.) ribbed.	E. Indies.	1825.	S. h.	_____
13	latifolia. s.s.	(wh.) broad-leaved.	_____	1787.	S. h.	Roxb. cor. 2. t. 134.
14	lucida. F.I.	(wh.) glossy-leaved.	_____	1820.	S. h.	_____
15	pubescens. R.S.	(wh.) pubescent.	_____	1825.	5. 7.	S. h.	_____
16	Thunbergia. s.s.	(wh.) starry.	C. B. S.	1773.	1. 3.	S. h.	Bot. mag. t. 1004.
17	Rothmannia. s.s. (sp.)	spotted-flower'd.	_____	1774.	7.	G. h.	— t. 690.
OXYA'NTHUS. DC. OXYA'NTHUS. Pentandria Monogynia.							
	speciosus. DC.	(wh.) tube-flowered.	Sierra Leon.	1789.	7. 12.	S. h.	Bot. mag. t. 1992.
	<i>Gardenia tubiflora</i> . A.R. 183.						
GENIPA. R.S. GENIP-TREE. Pentandria Monogynia.							
1	oblongifolia. R.S.	(st.) oblong-leaved.	Peru.	1823.	4. 6.	S. h.	Flor. per. 2. t. 220. f. a.
2	americana. R.S.	(wh.) American.	S. America.	1779.	5. 7.	S. h.	Plum. ic. t. 136.
3	Merianæ. R.S.	(wh.) hairy.	Cayenne.	1800.	—	S. h.	Merian. sur. t. 43.
	<i>Duroia eriopila</i> . L.						
4	édulis. R.S.	(wh.) eatable.	_____	1825.	—	S. h.	_____
5	escalenta. R.S.	(wh.) Cherry-fruited.	China.	_____	—	S. h.	_____
CA'NTHIUM. F.I. CA'NTHIUM. Pentandria Monogynia.							
1	angustifolium. F.I. (st.)	narrow-leaved.	E. Indies.	1826.	5. 7.	S. h.	_____
2	parvifolium. F.I.	(wh.) small-leaved.	_____	1824.	—	S. h.	_____
3	parviflorum. F.I. (ye.)	small-flowered.	_____	—	—	S. h.	_____
	<i>Wébera tetrandra</i> . W.						
4	didymum. F.I.	(wh.) cyme-flowered.	_____	1811.	—	S. h.	Gært. carp. 3. t. 196.
	<i>Wébera cymosa</i> . W.						
AZIMA. Lam. AZ'IMA. Tetrandria Monogynia.							
	tetracantha. P.S.	(wh.) four-spined.	E. Indies.	1758.	7.	S. h.	Seb. thes. 1. t. 13. f. 1.
	<i>Monettea barleroides</i> . L'Herit. stirp. nov. 1. t. 1.						
WE'BERIA. S.S. WE'BERIA. Pentandria Monogynia.							
1	corymbosa. s.s.	(wh.) corymbose.	E. Indies.	1759.	4. 6.	S. h.	Bot. reg. t. 119.

2 macrophylla. F.I. (wh.) large-leaved.	E. Indies.	1824.	5. 7.	S. h.	
3 odorata. F.I. (wh.) sweet-scented.	—	1826.	—	S. h.	
BURCHELLIA. B.R. BURCHELLIA. Pentandria Monogynia.					
1 capensis. B.R. (sc.) broad-leaved.	C. B. S.	1818.	3. 6.	G. h.	Bot. reg. t. 466.
2 parviflora. B.R. (or.) smaller-flower'd.	—	—	—	G. h.	t. 891.
bubalina. B.M. 2339.					
CATESBEA. S.S. LILY-THORN. Tetrandria Monogynia. L.					
1 spinosa. s.s. (wh.) spiny.	Providence I.	1726.	5. 9.	S. h.	Bot. mag. t. 131.
2 latifolia. B.R. (wh.) broad-leaved.	W. Indies.	1818.	9. 11.	S. h.	Bot. reg. t. 858.
3 parviflora. s.s. (wh.) small-flowered.	—	1810.	7. 10.	S. h.	Sloan. h. 2. t. 207. f. 1.
FERNELIA. J. FERNELIA. Tetrandria Monogynia.					
1 obovata. R.S. (wh.) obovate-leaved.	Isl. France.	1816.	—	S. h.	Lam. ill. t. 67. f. 1.
2 biflora. R.S. (wh.) two-flowered.	Bourbon.	—	—	S. h.	
Coccocypselum buxifolium. s.s. biflorum. w.					
3 uniflora. R.S. (wh.) one-flowered.	Mauritius.	—	—	S. h.	
buxifolia. β. Lamarck.					
COCCOCYPSOLUM. S.S. COCCOCYPSOLUM. Tetrandria Monogynia. S.S.					
rēpens. s.s. (li.) creeping.	W. Indies.	1793.	5.	S. 4.	Brown. jam. t. 6. f. 1.

Sect. IV. CINCHONEÆ. *Kth. synops. 3. p. 47.*

PORTLANDIA. S.S. PORTLANDIA. Pentandria Monogynia. L.					
1 grandiflora. R.S. (wh.) great-flowered.	Jamaica.	1775.	6. 8.	S. h.	Sm. ic. pict. 1. t. 6.
2 coccinea. R.S. (sc.) scarlet-flowered.	—	1812.	—	S. h.	
COUTAREA. S.S. COUTAREA. Hexandria Monogynia. S.S.					
speciosa. s.s. (sc.) Laurel-leaved.	Guiana.	1803.	—	S. h.	Aubl. guian. 1. t. 122.
Portlandia hexandra. Jacq. am. pict. t. 65.					
EXOSTEMMA. R.S. EXOSTEMMA. Pentandria Monogynia. R.S.					
1 caribæum. R.S. (wh.) Caribbean.	W. Indies.	1780.	6. 9.	S. h.	Andr. reposit. t. 481.
Cinchona caribæa. ad. w.					
2 floribundum. R.S. (bh.) many-flowered.	—	1794.	—	S. h.	Lamb. cinch. 27. t. 7.
PINCKNEYA. M. PINCKNEYA. Pentandria Monogynia.					
pūbens. m. (bh.) pubescent.	Georgia.	1786.	6. 7.	F. h.	Mich. amer. t. 13.
HYMENODICTYON. F.I. HYMENODICTYON. Pentandria Monogynia.					
1 thyrsiflorum. Wall. (st.) thyrs-flower'd. E. Indies.	1816.	—	—	S. h.	
2 excelsum. Wall. (wh.) tall.	—	—	—	S. h.	Roxb. cor. 2. t. 106.
Cinchona excelsa. R.S.					
HILLIA. S.S. HILLIA. Tetr-Hexandria Monogynia. S.S.					
1 longiflora. s.s. (wh.) long-flowered.	W. Indies.	1789.	2. 3.	S. h.	Bot. mag. t. 721.
2 tetrandra. s.s. (wh.) mountain.	Jamaica.	1793.	6. 7.	S. h.	
MACROCNEMUM. L. MACROCNEMUM. Pentandria Monogynia. L.					
1 jamaicense. R.S. (wh.) Jamaica.	Jamaica.	1806.	—	S. h.	Swartz obs. t. 3. f. 1.
2 speciosum. s.s. (li.) handsome.	Caracas.	1828.	—	S. h.	Jacq. schœnb. 1. t. 43.
Mussanda speciosa. R.S.					
3 coccineum. s.s. (sc.) scarlet.	Trinidad.	1825.	—	S. h.	
Mussanda coccinea. R.S.					
RONDEL'ETIA. R.S. RONDEL'ETIA. Pentandria Monogynia. L.					
1 americana. R.S. (wh.) American.	W. Indies.	1752.	8.	S. h.	Plum. ic. t. 242. f. 1.
2 laevigata. R.S. (wh.) smooth-leaved.	—	1790.	7. 8.	S. h.	Jacq. amer. pict. t. 62.
3 trifoliata. R.S. (re.) three-leaved.	Jamaica.	—	—	S. h.	
4 paniculata. F.I. (wh.) panicle'd.	E. Indies.	1820.	—	S. h.	
5 tinctoria. F.I. (wh.) dyer's.	—	1825.	—	S. h.	
6 racemosa. R.S. (wh.) racemed.	W. Indies.	1818.	—	S. h.	Brown. jam. 143. t. 2. f. 3.
7 tomentosa. R.S. (wh.) downy-leaved.	—	—	—	S. h.	
8 hirta. R.S. (br.ye.) hairy.	—	1776.	6. 8.	S. h.	Lodd. bot. cab. t. 350.

Sect. V. HEDYOTIDEÆ. *Kth. synops. 3. p. 40.*

HEDYOTIS. R.S. HEDYOTIS. Tetrandria Monogynia. L.					
1 diffusa. R.S. (wh.) spreading.	E. Indies.	1818.	7. 9.	S. ⊖.	
2 herbacea. R.S. (wh.) herbaceous.	—	1816.	6. 9.	S. ⊖.	Rheed. mal. 10. t. 35.
3 capensis. R.S. (wh.) Cape.	C. B. S.	1824.	7. 10.	H. ⊖.	
4 umbellata. R.S. (wh.) Indian Madder.	E. Indies.	1792.	7. 8.	S. 4.	Roxb. cor. 1. t. 3.
Oldenlandia corymbosa L.					

5 cymosa. R.S. (wh.)	Hyssop leaved. Jamaica.	1739.	6. 10.	S. $\textcircled{1}$.	Ehret pic. t.2. f.1. et.4.
Oldenländia corymbosa. w.					
6 ramosissima. R.S. (wh.)	branching.	1816.	—	S. $\textcircled{1}$.	
7 biflora. R.S. (pu.)	two-flowered. E. Indies.	—	—	S. $\textcircled{1}$.	
8 láctea. R.S. (st.)	milk-coloured. —	—	—	S. $\textcircled{1}$.	
9 campanuliflora. B.M. (pu.)	bell-flow'r'd. Brazil.	1825.	2. 10.	S. $\textcircled{1}$.	
MANETTIA. R.S.	MANETTIA. Tetrandria Monogynia. L.				
1 Lygustum. R.S. (re.)	veined-leaved. W. Indies.	1822.	2. 4.	S. $\textcircled{1}$. —. Brown. jam. t. 3. f. 2.	
2 coccinea. R.S. (sc.)	scarlet. Trinidad.	1806.	12. 4.	S. $\textcircled{1}$. —. Bot. reg. t. 693.	
BOUVA'RDIA. R.S.	BOUVA'RDIA. Tetrandria Monogynia.				
1 triphylla. R.S. (sc.)	blunt-leaved. Mexico.	1794.	4. 11.	G. $\textcircled{1}$.	Sal. par. lond. t. 88.
2 Jacquinii. k.s.	sharp-leaved. —	—	—	G. $\textcircled{1}$.	Jacq. scheob. 1. t. 257.
Ixòra americana. Jacq.	Bouvardia triphylla. Bot. reg. t. 107.				
versicolor. s.s. (re.ye.)	various-color'd. S. America.	1814.	7. 1.	S. $\textcircled{1}$.	Bot. reg. t. 245.
DENTE'LLA. W.	DENTE'LLA. Pentandria Monogynia. W.				
rēpens. w. (wh.)	creeping. N. Holland.	1802.	7.	H. $\textcircled{1}$.	Rumph.am.6.t.170.f.4.
SPERMADI'CTYON. B.R.	SPERMADI'CTYON. Pentandria Monogynia.				
1 suavolens. B.R. (wh.)	sweet-scented. E. Indies.	1816.	8. 12.	S. $\textcircled{1}$.	Bot. reg. t. 348.
Hamiltonia suavolens. F.I.					
2 azureum. B.R. (bl.)	blue-flowered. Nepaul.	1827.	1. 3.	S. $\textcircled{1}$.	Bot. reg. t. 1235.

Sect. VI. CEPHALANTHEÆ. *Kth. synops. 3. p. 37.*

NA'UCLEA. R.S.	NA'UCLEA. Pentandria Monogynia. L.				
1 Adina. Sm. (wh.)	Myrtle-leaved. China.	1804.	7. 8.	G. $\textcircled{1}$.	Bot. reg. t. 895.
Adina globiflora. Sal.	par. lond. t. 115.				
2 adinoïdes. s.s. (wh.)	Adina-like. —	1825.	—	G. $\textcircled{1}$.	
3 orientalis. R.S. (ye.)	oriental. E. Indies.	1800.	S. $\textcircled{1}$.	
4 Cadámba. F.I. (or.)	broad-leaved. —	1823.	S. $\textcircled{1}$.	Rheed. mal. 3. t. 33.
5 undulata. F.I. (ye.)	wave-leaved. —	—	S. $\textcircled{1}$.	
6 parvifolia. F.I. (ye.)	small-leaved. —	1825.	—	S. $\textcircled{1}$.	Roxb. corm. 1. t. 52.
7 cordifolia. R.S. (st.)	heart-leaved. —	—	S. $\textcircled{1}$.	— 1. t. 53.
8 purpurea. F.I. (pu.)	purple-flowered. —	1827.	S. $\textcircled{1}$.	— 1. t. 54.
UNCARIA. F.I.	UNCARIA. Pentandria Monogynia.				
Gámbier. F.I. (pi.)	Gambier-tree. E. Indies.	1826.	S. $\textcircled{1}$. —. Linn. trans. v. 9. t. 22.	
SARCOCE'PHALUS. Afz.	SARCOCE'PHALUS. Pentandria Monogynia.				
esculentus. Afz. (wh.)	eatable-fruited. Sierra Leone.	1822.	6. 8.	S. $\textcircled{1}$.	Hort. trans. 5. t. 18.
CEPHALA'NTHUS. L.	BUTTON-WOOD. Tetrandria Monogynia. L.				
occidentalis. L. (wh.)	American. N. America.	1735.	8.	S. $\textcircled{1}$.	Schmidt arb. 1. t. 45.
MORINDA. L.	MORINDA. Pentandria Monogynia. L.				
1 umbellata. R.S. (wh.)	umbelled. E. Indies.	1809.	6. 8.	S. $\textcircled{1}$.	
2 citrifolia. R.S. (wh.)	broad-leaved. Otaheite.	1793.	—	S. $\textcircled{1}$.	Rheed. mal. 1. t. 52.
3 tinctoria. F.I. (wh.)	dyer's.	—	1826.	S. $\textcircled{1}$.
4 bracteata. F.I. (wh.)	bracted.	E. Indies.	1816.	5. 7.	S. $\textcircled{1}$.
5 angustifolia. F.I. (wh.)	narrow-leaved.	—	—	—	S. $\textcircled{1}$.
6 Róyoc. R.S.	(wh.) Laurel-leaved. W. Indies.	1793.	7. 10.	S. $\textcircled{1}$.	Roxb. corom. 3. t. 237.
					Jacq vind. 1. t. 16.

Sect. VII. COFFEACEÆ. *Kth. synops. 3. p. 20.*

MITCHELLA. L.	MITCHELLA. Tetrandria Monogynia. L.				
rēpens. L.	(wh.) creeping. N. America.	1761.	6.	H. $\textcircled{1}$.	Cat. car. 1. t. 20.
PAVETTA. R.S.	PAVETTA. Tetrandria Monogynia. L.				
1 índica. R.S. (wh.)	Indian.	E. Indies.	1791.	8. 10.	S. $\textcircled{1}$.
2 tomentosa. R.S. (wh.)	woolly.	—	1824.	—	S. $\textcircled{1}$.
3 cáffra. R.S. (wh.)	Cape.	C. B. S.	1816.	—	G. $\textcircled{1}$.
4 pentandra. R.S. (wh.)	pentandrous.	W. Indies.	1823.	—	S. $\textcircled{1}$.
5 arenosa. R.S. (wh.)	tubercléd.	China.	3. 10.	S. $\textcircled{1}$.
índica. B.R. nec aliorum.					Plum. ic. t. 156. f. 1.
IX'ORA. L.	IX'ORA. Tetrandria Monogynia. L.				
1 grandiflora. B.R. (sc.)	great-flowered. E. Indies.	1814.	7. 9.	S. $\textcircled{1}$.	Bot. reg. t. 154.
coccinea. L. non B.M.					

2	Bandhūca. F.I.	(sc.) stem-clasping.	E. Indies.	1814.	1. 4.	S. h.	Bot. reg. t. 513.
3	fúlgens. F.I.	(sc.) glossy.	—	1820.	S. h.	Rumph. amb. 4. t. 46.
	longifolia. Sm.	<i>lanceolata</i> . Lam.					
4	flámmea. s.p.	(or.) flame-coloured.	China.	1690.	4. 8.	S. h.	Bot. mag. t. 169.
	coccinea. B.M.	<i>strícta</i> . F.I. <i>speciosa</i> . W. enum.	—	—			
5	crocátā. B.R.	(sn.) saffron-colour'd.	—	1820.	—	S. h.	Bot. reg. t. 782.
6	incarnátā. B.C.	(fl.) flesh-coloured.	—	—	—	S. h.	Lodd. bot. cab. 1048.
7	blánda. B.R.	(bh.) charming.	—	1812.	—	S. h.	Bot. reg. t. 100.
8	álba. w.	(wh.) white.	E. Indies.	1768.	6.	S. h.	Rheed. mal. 2. t. 14.
9	cuneifólia. F.I.	(wh.) wedge-leaved.	—	1820.	—	S. h.	Bot. reg. t. 648.
10	brachiáta. F.I.	(wh.) brachiate-paniced.	—	—	—	S. h.	Andrews's reposit. t. 78.
11	Pavéttā. A.R.	(wh.) fragrant.	—	—	—	S. h.	Vahl symb. 3. t. 52.
12	acuminátā. F.I.	(wh.) tapering-leaved.	—	1824.	3. 5.	S. h.	Bot. mag. t. 2505.
13	parviflóra. F.I.	(wh.) small-flowered.	—	1800.	8. 10.	S. h.	
14	barbáta. F.I.	(wh.) bearded.	—	1822.	6. 8.	S. h.	
15	undulátā. F.I.	(wh.) wave-leaved.	—	1818.	—	S. h.	
16	rósea. F.I.	(ro.) rose-coloured.	—	1820.	4. 8.	S. h.	Bot. reg. t. 540.
17	obováta. R.S.	(cr.) obovate-leaved.	—	1810.	—	S. h.	
	purpùrea. F?						
CHOM'ELIA. R.S.	CHOM'ELIA.	Tetrandria Monogynia. W.					
1	spinósa. R.S.	(wh.) spiny.	W. Indies.	1793.	S. h.	Jacq. amer. 18. t. 13.
2	fasciculátā. R.S.	(wh.) fascicled.	Jamaica.	1824.	4. 6.	S. h.	
SIDERODE'NDRON. S.S.	IRON-TREE.	Tetrandria Monogynia. W.					
	triflorum. s.s.	(ro.wh.) three-flowered.	W. Indies.	1793.	S. h.	Jacq. amer. 19. t. 175. f. 9.
TETRAM'ERIUM. G.	TETRAM'ERIUM.	Tetrandria Monogynia. S.S.					
1	odoratissimum. s.s. (wh.) sweet-scented.	W. Indies.	1793.	4. 6.	S. h.	Jacq. amer. t. 47.	
	Coffea occidentalis. w.	<i>Ixora americana</i> . Sw.					
2	paniculátum. s.s. (wh.) paniced.	Guiana.	1822.	S. h.	Aub. guian. t. 58.	
COFFEEA. L.	COFFEE-TREE.	Pentandria Monogynia. L.					
	árabica. w.	(wh.) Arabian.	Yemen.	1696.	8. 11.	S. h.	Bot. mag. t. 1303.
CEPHAE'LIS. S.S.	CEPHAE'LIS.	Pentandria Monogynia. L.					
1	eláta. s.s.	(pu.) tall.	Jamaica.	1793.	S. h.	
2	tomentósa. R.S.	(pu.) tomentose.	Trinidad.	1820.	4. 6.	S. h.	
3	púnicea. R.S.	(pu.) puniceous.	Jamaica.	1825.	—	S. h.	
4	violácea. R.S.	(vi.) violet-coloured.	W. Indies.	1824.	—	S. h.	Aubl. guian. 1. t. 60.
	<i>Tapogòmea violacea</i> . Aubl.						
5	axillaris. s.s.	(pu.) axillary-flow'r'd.	Brazil.	1816.	4. 5.	S. h.	Lind. coll. t. 21.
	calycina. L. col.						
6	glábra. R.S.	(bl.) smooth.	Trinidad.	1820.	5. 8.	S. h.	Aubl. guian. 1. t. 63.
7	álba. R.S.	(bh.) white.	Guiana.	1824.	—	S. h.	— 1. t. 62. f. 1.
8	involucrátā. R.S. (wh.) involucred.	—	—	—	S. h.	— 1. t. 64.
	<i>Carapichea guianensis</i> . Aubl.						
9	pedunculátā. P.L. (wh.) long-peduncled.	Sierra Leon.	2. 4.	S. h.	Sal. par. lond. t. 103. 29	
PSYCH'OTRIA. R.S.	PSYCH'OTRIA.	Pentandria Monogynia. L.					
1	asiática. R.S.	(wh.) Indian.	E. Indies.	1806.	4. 5.	S. h.	Jacq. amer. p. t. 260. f. 19.
2	glabréta. R.S.	(wh.) smooth.	Jamaica.	1822.	—	S. h.	
3	laurifólia. R.S.	(wh.) Laurel-leaved.	W. Indies.	1825.	5. 8.	S. h.	
4	f'œtens. R.S.	(wh.) strong-scented.	Jamaica.	—	—	S. h.	
5	citrifólia. R.S.	(wh.) Citron-leaved.	W. Indies.	1793.	1. 12.	S. h.	
6	ellíptica. B.R.	(wh.) elliptic-leaved.	Brazil.	1820.	2. 6.	S. h.	Bot. reg. t. 607.
7	myrtiphyl'um. R.S. (wh.) Myrtle-leaved.	Jamaica.	1823.	2. 5.	S. h.	Sloan. his. 2. t. 209. f. 2.	
8	parasítica. R.S.	(wh.) parasitic.	W. Indies.	1802.	5. 8.	S. h.	Jacq. amer. t. 51. f. 1.
9	brachiáta. R.S.	(wh.) cross-branched.	—	1793.	—	S. h.	
10	pubéscens. R.S.	(wh.) pubescent.	Jamaica.	1812.	S. h.	
11	corymbosa. R.S.	(pu.) corymbose.	—	—	3. 4.	S. h.	
12	pedunculátā. R.S. (st.) pedunculate.	—	1824.	5. 6.	S. h.	Jacq. schoenb. 3. t. 260.	
13	undátā. R.S.	(wh.) wave-leaved.	Bahama.	1818.	4. 6.	S. h.	
14	Reevesi. F.I.	(gr.) Reeves's.	China.	1823.	—	G. h.	
GEO'PHILA. D.P.	GEO'PHILA.	Pentandria Monogynia.					
	reniformis. D.P. (wh.) kidney-leaved.	W. Indies.	1793.	4. 6.	S. 4.	Jacq. amer. t. 46.	
	<i>Psychotria herbacea</i> . W.						
CHIOC'C'A. W.	SNOWBERRY.	Pentandria Monogynia. L.					
	racemosa. w.	(wh.) cluster-flower'd.	Jamaica.	1729.	2.	S. h.	Andr. deposit. t. 284.
ERN'ODEA. S.S.	ERN'ODEA.	Tetrandria Monogynia. W.					
	montána. s.s.	(pu.) mountain.	Sicily.	1820.	6. 7.	H. h.	Flor. græc. t. 143.
	<i>Aspérula calabrica</i> . L.						
SERISSA. S.S.	SERISSA.	Pentandria Monogynia. W.					
	f'œtida. s.s.	(wh.) Japanese.	Japan.	1787.	5. 9.	G. h.	Bot. mag. t. 361.
	β multiplex.	(wh.) double-flowered.	—	—	—	G. h.	

PLOC'AMA. W.	PLOC'AMA. Pentandria Monogynia. W.
péndula. w.	(wh.) pendulous. Canaries. 1779. G. h.
PÆ'D'ERIA. W.	PÆ'D'ERIA. Pentandria Monogynia. L.
1 fœtida. w.	(pu.) stinking. China. 1806. G. h. —. Icon. Kæmpf. 9.
2 Língun. Boj. tongue. Madagascar. 1823. S. h. —.

Sect. VIII. SPERMACOCEÆ. *Kth. synops. 3. p. 14.*RICHARDS'ONIA. *Kth. RICHARDS'ONIA.* Hexandria Monogynia. L.

1 scábra. B.F.G.	(u.h.) rough. S. America. 1812. 9. 10. H. ⊖.	Swt. br. fl. gar. t. 91.
2 pilosa. K.S.	(wh.) hairy. Lima. 1824. 7. 9. H. ⊖.	Flor. peruv. 3.t. 279.f.b.

PHYLLIS. W.	BASTARD HARE'S-EAR. Pentandria Dignynia. L.
Nòbla. w.	(br.) Canary. Canaries. 1699. 6. 7. G. h.

DI'ODIA. S.S.	DI'ODIA. Tetrandria Monogynia. W.
---------------	-----------------------------------

1 virginica. s.s.	(wh.) Virginian. Virginia. 1820. 6. 8. H. ȳ.	Jacq. ic. 1. t. 29.
2 sarmentosa. s.s.	(wh.) sarmentose. W. Indies. 1821. — S. h.	
3 verticillata. s.s.	(wh.) whorl-leaved. Santa Cruz. — S. ȳ.	

BIGEL'OVIA. S.S.	BIGEL'OVIA. Tetrandria Monogynia. S.S.
------------------	--

1 verticillata. s.s.	(wh.) whorl-flowered. Africa. 1732. 6. 8. S. h.	Dill. elth. t. 277. f. 358.
----------------------	---	-----------------------------

<i>Spermacoce verticillata.</i> L.		
------------------------------------	--	--

2 commutata. s.s.	(wh.) starry-headed. W. Indies. 1818. — S. ȳ.
3 stricta. s.s.	(wh.) upright. E. Indies. 1821. — S. ⊖.

<i>Spermacoce stricta.</i> L.		
-------------------------------	--	--

SPERMACOCE. S.S. BUTTON-WEED.	Tetrandria Monogynia. L.
-------------------------------	--------------------------

1 glabra. s.s.	(wh.) smooth. N. America. 1823. 7. 8. H. ⊖.
2 diodina. s.s.	(wh.) Diodia-like. 1820. — H. ⊖.

3 hispida. s.s.	(vi.) bristly. E. Indies. 1781. 8. 9. S. ⊖.
-----------------	---

4 spinosa. s.s.	(wh.) spiny. W. Indies. 1825. — S. ⊖.
-----------------	---------------------------------------

5 latifolia. s.s.	(wh.) broad-leaved. Guiana. 1803. 7. S. ♂.
-------------------	--

6 cornifolia. s.s.	(wh.) Dogwood-leav'd. Brazil. 1820. 8. 9. S. ⊖.
--------------------	---

7 stylosa. s.s.	(wh.) long-styled. Manila. — — S. ⊖.
-----------------	--------------------------------------

8 suffruticosa. s.s.	(bh.) suffruticose. 1823. 7. 8. S. h.
----------------------	--

9 radicans. s.s.	(wh.) rooting. Guiana. 1803. — S. ȳ.
------------------	--------------------------------------

10 áspera. s.s.	(wh.) rough. W. Indies. 1822. — S. ⊖.
-----------------	---------------------------------------

11 tenuior. s.s.	(bh.) slender. — 1732. 6. 8. H. ⊖.
------------------	------------------------------------

12 lanceolata. L.en.	(wh.) spear-leaved. — 1818. — H. ⊖.
----------------------	-------------------------------------

13 hirta. s.s.	(wh.) hairy. Jamaica. 1822. 7. 9. H. ⊖.
----------------	---

14 villosa. s.s.	(wh.) villous. — 1823. — S. ⊖.
------------------	--------------------------------

15 scábra. s.s.	(wh.) rough. E. Indies. 1816. — S. ⊖.
-----------------	---------------------------------------

16 capitata. s.s.	(wh.) headed. Peru. 1826. 6. 8. G. ȳ.
-------------------	---------------------------------------

17 rúbra. s.s.	(pi.) Cross-wort. S. America. 1760. 7. 8. S. ♂.
----------------	---

strigosa. B.M. 1558.		
----------------------	--	--

18 Fischéri. s.s.	(wh.) Fischer's. Jamaica. 1820. — S. ⊖.
-------------------	---

19 sexangulär. s.s.	(bl.) six-angled. W. Indies. 1822. — S. ⊖.
---------------------	--

20 involucrata. R.S.	(wh.) involucr. Carolina. 1812. — H. ⊖.
----------------------	---

21 Roxburi. hii. s.s.	(wh.) Roxburgh's. E. Indies. 1818. — S. ♂.
-----------------------	--

<i>lævis. F.I. non Lam.</i>		
-----------------------------	--	--

KNO'XIA. S.S.	KNO'XIA. Tetrandria Monogynia. W.
---------------	-----------------------------------

1 corymbosa. s.s.	(wh.) Sumatra. E. Indies. 1818. 4. 6. S. h.
-------------------	---

<i>Spermacoce sumatrénis. F.I.</i>		
------------------------------------	--	--

2 umbellata. s.s.	(wh.) umbel-flower'd. Nepaul. 1820. — S. h.
-------------------	---

<i>Spermacoce teres. F.I.</i>		
-------------------------------	--	--

Sect. IX. GALEÆ. *Kth. synops. 3. p. 10.*

ANTHOSPE'RUM. W. AMBER-TREE. Diœcia Tetrandria. L.

æthiopicum. s.s. (gr.)	Ethiopian. C. B. S. 1692. 6. 7. G. h.	Pluk. alm. t. 183. f. 1.
------------------------	---------------------------------------	--------------------------

CRUCIANELLA. R.S. CROSS-WORT.	Tetrandria Monogynia. L.
-------------------------------	--------------------------

1 angustifolia. R.S. (st.)	narrow-leaved. S. Europe. 1658. 6. 7. H. ⊖.	Sm. exot. bot. 2. t. 109.
----------------------------	---	---------------------------

2 ciliata. R.S. (st.)	ciliated. Levant. 1805. 7. 8. H. ⊖.
-----------------------	-------------------------------------

3 pátula. R.S. (ye.)	spreading. Spain. 1798. 6. 7. H. ⊖.
----------------------	-------------------------------------

4 latifolia. R.S. (gr.)	broad-leaved. S. Europe. 1633. — H. ⊖.
-------------------------	--

5 ægyptiaca. R.S. (st.)	Egyptian. Egypt. 1800. — H. ⊖.
-------------------------	--------------------------------

6 monspeliaca. R.S. (st.)	Montpellier. S. France. 1791. 7. 8. H. ⊖.
---------------------------	---

<i>Flor. græc. t. 140.</i>		
----------------------------	--	--

RUBIACEÆ.

7 marítima. R.S. (ye.) sea-side.	S. France. 1640.	7. 8. G. $\frac{1}{2}$.	Sabb. hort. 2. t. 13.
8 americana. R.S. (bl.) American.	Vera Cruz.	— S. $\frac{1}{2}$.	
9 pubescens. R.S. (pu.) pubescent.	Candia. 1799.	7. 8. G. $\frac{1}{2}$.	
10 áspera. M.B. (st.) rough.	Iberia. 1827.	6. 8. H. $\frac{1}{2}$.	
11 glomerata. M.B. (st.) clustered.	— 1828.	— H. $\frac{1}{2}$.	
12 molluginoïdes. R.S. (wh.) Mollugo-like.	Caucasus. 1800.	— H. $\frac{1}{2}$.	Buxb. cent. 2. t. 30. f. 1.
13 hispida. R.S. (pu.) hispid.	Vera Cruz.	— S. $\frac{1}{2}$.	
SHERARDIA. R.S. SHERARDIA.	Tetrandria Monogynia. L.		
arvénensis. R.S. (li.) little.	Britain.	4. 9. H. \odot .	Eng. bot. v. 13. t. 891.
R'UBIA. R.S. MADDER. Tetrandria Monogynia. L.	S. Europe. 1596.	6. H. $\frac{1}{2}$.	
1 tintorium. R.S. (gr.) dyer's.	Portugal. 1812.	6. 8. H. $\frac{1}{2}$.	
2 spléndens. R.S. (ye.) glossy.	Canaries. 1816.	— F. $\frac{1}{2}$.	
3 galioïdes. R.S. (gr.) Galium-like.	England.	7. H. $\frac{1}{2}$.	Eng. bot. v. 12. t. 851.
4 peregrina. R.S. (ye.) wild.	Majorca. 1762.	— F. $\frac{1}{2}$.	Flor. græc. t. 142.
5 lúcida. R.S. (ye.) shining.	Canaries. 1779.	9. F. $\frac{1}{2}$.	Jacq. ic. 1. t. 25.
6 fruticosa. R.S. (ye.) prickly-leaved.	Italy. 1823.	— H. $\frac{1}{2}$.	Bocc. mus. t. 75. f. 2-3.
7 Bocconi. R.S. (bh.) Boccone's.	S. Europe. 1772.	7. 8. F. $\frac{1}{2}$.	Lam. ill. t. 60. f. 2.
8 angustifolia. R.S. (st.) narrow-leaved.	Siberia. 1783.	7. H. $\frac{1}{2}$.	Pall. it. 3. ap. t. L. f. 1.
9 cordifolia. R.S. (wh.) heart-leaved.	Tetrandria Monogynia. L.		
ASPERULA. R.S. WOODRUFF. S. Europe. 1775.	6. H. $\frac{1}{2}$.	Moris. his. 3. t. 21. f. 4.	
1 laevigata. R.S. (wh.) shining.	Italy. 1739.	4. 6. H. $\frac{1}{2}$.	Lob. ic. 800. f. 1.
2 taurina. R.S. (li.) broad-leaved.	Hungary. 1818.	— H. $\frac{1}{2}$.	W. et K. hung. 2 t. 150.
3 longiflora. R.S. (li.) long-flowered.	S. Europe. 1823.	— H. $\frac{1}{2}$.	
4 aristata. R.S. (st.) awned.	Caucasus. —	6. 7. H. $\frac{1}{2}$.	
5 supina. R.S. (bh.) trailing.	Levant. 1818.	— H. $\frac{1}{2}$.	Eng. bot. v. 1. t. 33.
6 brevifolia. R.S. (li.) short-leaved.	England.	7. H. $\frac{1}{2}$.	
7 cynanchica. R.S. (bh.) small.	Caucasus. 1825.	— H. $\frac{1}{2}$.	
8 alpina. s.s. (bh.) Alpine.	Levant. 1775.	6. H. $\frac{1}{2}$.	Tabern. ic. 151.
9 crassifolia. R.S. (wh.) thick-leaved.	Italy. 1824.	— H. $\frac{1}{2}$.	Lobel. ic. t. 801. f. 2.
10 scabra. s.s. (wh.) rough.	Europe. 1764.	6. 7. H. $\frac{1}{2}$.	Bot. mag. t. 2146.
11 tinctoria. R.S. (bh.) narrow-leaved.	Hungary. 1801.	— H. $\frac{1}{2}$.	Ten. fl. nap. 1. t. 9.
12 montana. R.S. (bh.) mountain.	Europe. 1596.	7. H. \odot .	Ram. phil. t. 9. f. 1. 2.
13 arvensis. R.S. (bh.) field.	Arcadia. 1819.	5. H. $\frac{1}{2}$.	
14 arcadiensis. s.s. (ro.) Arcadian.	Italy. 1826.	— H. $\frac{1}{2}$.	
15 tomentosa. T.N. (pu.) woolly.	Pyrenees. —	5. 6. H. $\frac{1}{2}$.	
16 hirta. R.S. (pu.) thick-rooted.	S. Europe. 1820.	— H. $\frac{1}{2}$.	
17 hirsuta. R.S. (ro.) hairy creeping.	Crete. 1823.	6. 7. H. $\frac{1}{2}$.	Flor. græc. t. 119.
repens. Brot.	S. Europe. 1710.	6. 9. H. $\frac{1}{2}$.	Bocc. mus. 2. t. 116.
18 incana. R.S. (pu.) hoary.	Caucasus. 1825.	— H. $\frac{1}{2}$.	
19 galioïdes. s.s. (wh.) glaucous.	Greece. 1823.	— H. $\frac{1}{2}$.	Flor. græc. t. 118.
Gálium glaucum. L.	Crete. —	— H. $\frac{1}{2}$.	t. 117.
20 humifusa. s.s. (wh.) spreading.	Britain.	5. 6. H. $\frac{1}{2}$.	Eng. bot. 755.
21 longifolia. R.S. (wh.) long-leaved.	Levant. 1826.	— H. $\frac{1}{2}$.	
22 Aparine. R.S. (wh.) marsh.	Tetrandria Monogynia. S.S.		
rivalis. Sm. Gálium uliginosum. Pall.	S. Europe. 1739.	5. 7. H. \odot .	Colum. Ecphr. t. 297.
odorata. R.S. (wh.) sweet-scented.	— 1768.	— H. \odot .	Gært. fr. C. 2. t. 24. f. 1.
24 tyraicum. Bes. (wh.) Tyraican.	Canaries. 1780.	7. 9. H. \odot .	
VALANTIA. S.S. VALANTIA.	Tetrandria Monogynia. L.		
1 muralis. s.s. (st.) wall.	Britain.	5. 6. H. $\frac{1}{2}$.	Eng. bot. v. 2. t. 143.
2 hispida. s.s. (wh.) bristly.	Tauria. 1829.	5. 7. H. $\frac{1}{2}$.	
3 filiformis. s.s. (wh.) least.	— 1820.	— H. $\frac{1}{2}$.	Bot. mag. t. 2670.
GA'LUM. S.S. BEDSTRAW.	Hungary. 1799.	7. H. \odot .	W. et K. hung. 1. t. 33.
1 cruciatum. s.s. (ye.) Crosswort.	Europe. 1731.	7. 8. H. $\frac{1}{2}$.	1. t. 32.
Valantia cruciata. w.	—	— H. $\frac{1}{2}$.	
2 chersonense. R.S. (ye.) fringed.	Switzerland. 1823.	5. 6. H. $\frac{1}{2}$.	Scop. c.e. 2.1.n.144.t.2.
3 taūricum. R.S. (ye.) fragrant.	Egypt. 1752.	7. 8. H. \odot .	Lam. ill. t. 843. f. 3.
Valantia taūrica. B.M.	S. Europe. 1775.	7. H. $\frac{1}{2}$.	Buxb. cent. 2. t. 29.
4 pedemontanum. s.s. (ye.) Piedmont.	— 1820.	— H. $\frac{1}{2}$.	
5 Bauhinii. s.s. (ye.) Bauhin's.	Tartary. —	6. 8. H. $\frac{1}{2}$.	
Valantia glabra. w.	Caucasus. 1823.	— H. $\frac{1}{2}$.	
6 Scopolianum. R.S. (ye.) Scopoli's.	Britain.	— H. $\frac{1}{2}$.	Eng. bot. v. 26. t. 1857.
vérnum. Scop.	—	— H. $\frac{1}{2}$.	
7 Halleri. R.S. (st.) Haller's.	—	— H. $\frac{1}{2}$.	
8 articulatum. s.s. (ye.) jointed.	—	— H. $\frac{1}{2}$.	
9 rubioïdes. s.s. (st.) Madder-leaved.	—	— H. $\frac{1}{2}$.	
10 diffusum. s.s. (st.) spreading.	—	— H. $\frac{1}{2}$.	
11 tatáricum. s.s. (wh.) Tartarian.	—	— H. $\frac{1}{2}$.	
12 valantioïdes. s.s. (wh.) Valantia-like.	—	— H. $\frac{1}{2}$.	
13 palustre. s.s. (wh.) water.	—	— H. $\frac{1}{2}$.	

14	satureiaefolium.s.s. (<i>wh.</i>) Savory-leaved.	Caucasus.	1818.	7.	H.w. 4.	
15	trifidum. s.s. (<i>wh.</i>) 3-fid-flowered.	N. Europe.	—	—	H.w. 4.	Flor. dan. t. 48.
16	fruticosum. s.s. (<i>wh.</i>) frutescent.	Crete.	1823.	7. 8.	F. 4.	
17	helodes. R.S. (<i>wh.</i>) weak-branched.	Portugal.	1827.	—	H.w. 4.	
18	Claytoni. R.S. (<i>wh.</i>) Clayton's.	N. America.	—	—	H.w. 4.	
19	Witheringii. s.s. (<i>wh.</i>) rough.	England.	—	6. 7.	H. 4.	Eng. bot. v. 31. t. 2206.
20	glabrum. s.s. (<i>wh.</i>) smooth.	C. B. S.	1816.	—	G. 4.	
21	uliginosum. s.s. (<i>wh.</i>) marsh.	Britain.	—	7. 8.	H.w. 4.	Eng. bot. v. 28. t. 1972.
22	divaricatum. s.s. (<i>wh.</i>) divaricate.	S. France.	1820.	—	H. 4.	DC. ic. gal. rar. t. 24.
23	campestre. s.s. (<i>st.</i>) field.	Barbary.	1818.	8. 9.	H. 4.	
24	austriacum. R.S. (<i>wh.</i>) Austrian.	Europe.	1804.	6. 7.	H. 4.	Jacq. aust. t. 80.
25	Boccóni. R.S. (<i>bh.</i>) Boccone's.	—	1801.	5. 6.	H. 4.	Bocc. mus. t. 101.
26	obliquum. R.S. (<i>wh.</i>) oblique-leaved.	S. France.	—	—	H. 4.	Vill. delph. 2.n.9. t. 8.
27	pusillum. E.B. (<i>wh.</i>) least.	England.	—	7. 8.	H. 4.	Eng. bot. v. 2. t. 74.
28	alpestre. R.S. (<i>wh.</i>) smooth-leaved.	S. Europe.	1824.	—	H. 4.	Vill. delph. 2. t. 7. bis.
29	tenuifolium. R.S. (<i>wh.</i>) slender-leaved.	—	1820.	6. 7.	H. 4.	
30	ruberum. s.s. (<i>re.</i>) red.	Italy.	1597.	—	H. 4.	Ger. herb. 967. f. 3.
31	tenuissimum. s.s. (<i>br.</i>) slenderest.	Caucasus.	1818.	7. 9.	H. 4.	
32	vérum. s.s. (<i>ye.</i>) cheese-rennet.	Britain.	—	—	H. 4.	Eng. bot. v. 10. t. 660.
33	mutabile. R.S. (<i>st.</i>) changeable-color'd.	Poland.	1828.	7. 10.	H. 4.	
34	véro-simile. R.S. (<i>ye.</i>) large yellow.	—	1825.	7. 9.	H. 4.	
35	caucasicum. R.S. (<i>ye.</i>) Caucasean.	Caucasus.	1820.	—	H. 4.	
36	tyrolense. R.S. (<i>wh.</i>) Tyrolese.	Tyrol.	1801.	7.	H. 4.	Eng. bot. v. 24. t. 1673.
37	Mollugo. R.S. (<i>wh.</i>) great hedge.	Britain.	—	7. 8.	H. 4.	
38	elatum. P.S. (<i>wh.</i>) tall.	France.	1827.	—	H. 4.	
39	campanulatum.s.s. (<i>wh.</i>) bell-flowered. <i>glaucum.</i> Jacq. non L.	Europe.	1710.	6. 9.	H. 4.	Jacq. aust. t. 1. t. 81.
40	rectum. s.s. (<i>wh.</i>) upright.	Britain.	—	6. 7.	H. 4.	Eng. bot. v. 29. t. 2067.
41	lucidum. R.S. (<i>wh.</i>) glossy.	S. Europe.	1820.	—	H. 4.	All. ped. 1. t. 77. f. 2.
42	linifolium. s.s. (<i>wh.</i>) Flax-leaved.	—	1759.	—	H. 4.	Barr. ic. t. 356 et 583.
43	rigidum. s.s. (<i>wh.</i>) rigid.	Azores.	1778.	—	H. 4.	
44	sylvaticum. s.s. (<i>wh.</i>) wood.	S. Europe.	1658.	7. 8.	H. 4.	
45	cinerium. s.s. (<i>wh.</i>) cinereous.	Scotland.	—	8.	H. 4.	All. ped. t. 77. f. 4.
46	aristatum. s.s. (<i>wh.</i>) awned.	—	—	7. 8.	H. 4.	Barr. ic. t. 356.
47	purpureum. s.s. (<i>pu.</i>) purple.	Switzerland.	1731.	6. 7.	H. 4.	Noct. et B. fl. tic. 1. t. 3.
48	saxatile. s.s. (<i>bh.</i>) smooth heath.	Britain.	—	7. 8.	H. 4.	Eng. bot. v. 12. t. 815.
49	helveticum. s.s. (<i>wh.</i>) Swiss.	Switzerland.	1823.	—	H. 4.	J.m.ac.p.1714.t.15.f.1.
50	baldense. s.s. (<i>wh.</i>) tufted.	Mount Baldo.	1825.	—	H. 4.	
51	tricorne. E.B. (<i>st.</i>) three-flowered.	Britain.	—	7.	H. 4.	Eng. bot. v. 23. t. 1641.
52	spuriūm. E.B. (<i>gr.</i>) spurious.	—	—	6. 7.	H. 4.	— v. 26. t. 1871.
53	verrucosum. E.B. (<i>st.</i>) warted. <i>saccharatum.</i> R.S. <i>Valantia Aparine.</i> w.	—	—	6. 8.	H. 4.	— v. 31. t. 2173.
54	Cucullaria. s.s. (<i>ye.</i>) hooded.	Levant.	1780.	5. 6.	H. 4.	Buxb. cent. 1. t. 19. f. 2.
55	boreale. E.B. (<i>wh.</i>) cross-leaved.	Britain.	—	7.	H. 4.	Eng. bot. v. 2. t. 105.
56	septentrionale. R.S. (<i>wh.</i>) northern.	N. America.	1827.	—	H. 4.	
57	brachyphyllum. R.S. (<i>wh.</i>) short-leaved.	Caucasus.	—	—	H. 4.	
58	rotundifolium. s.s. (<i>wh.</i>) round-leaved.	Europe.	—	6. 7.	H. 4.	
59	ellipticum. s.s. (<i>wh.</i>) elliptic-leaved.	Sicily.	1820.	—	H. 4.	Jacq. aust. t. 94.
60	pilosum. R.S. (<i>wh.</i>) hairy.	N. America.	1778.	—	H. 4.	Barr. ic. t. 21. f. 4.
61	suavolens. R.S. (<i>wh.</i>) sweet-scented.	N. Europe.	1827.	—	H. 4.	
62	reflexum. R.S. (<i>wh.</i>) reflexed.	Tauria.	1818.	—	H. 4.	
63	humifusum. R.S. (<i>wh.</i>) trailing.	—	1820.	—	H. 4.	
64	maritimum. s.s. (<i>pu.</i>) sea-side.	S. Europe.	—	7. 8.	H. 4.	
65	triflorum. R.S. (<i>wh.</i>) three-flowered.	N. America.	1827.	—	H. 4.	Willd. hort. ber. t. 66.
66	anglicum. E.B. (<i>st.</i>) English.	England.	—	6. 7.	H. 4.	Eng. bot. v. 6. t. 384.
67	parvifolium. R.S. (<i>wh.</i>) small-leaved.	Switzerland.	1825.	—	H. 4.	
68	parisiense. R.S. (<i>st.</i>) Paris.	France.	1818.	—	H. 4.	
69	murale. s.s. (<i>st.</i>) wall.	Italy.	1805.	6. 8.	H. 4.	All. ped. 34. t. 77. f. 1.
	<i>Sherardia muralis.</i> w.					
70	verticillatum.R.S. (<i>wh.</i>) whorled.	France.	1829.	—	H. 4.	Flor. græc. t. 116.
71	græcum. s.s. (<i>pu.</i>) Candian.	Candia.	1798.	6. 7.	H. 4.	— t. 136.
72	capillare. R.S. (<i>wh.</i>) woolly-fruited.	Spain.	1828.	—	H. 4.	Cavan.icon.2.t.191.f.1.
73	Aparine. E.B. (<i>wh.</i>) Cleavers.	Britain.	—	5. 9.	H. 4.	Eng. bot. v. 12. t. 316.
74	infestum. R.S. (<i>st.</i>) infesting.	Europe.	1818.	—	H. 4.	W. et K. hung. 3.t.202.
75	scaberrimum. R.S. (<i>st.</i>) roughest.	Egypt.	—	6. 9.	H. 4.	

ORDO CXVI.

OPERCULARIEÆ. *Juss. ann. mus. 4. p. 418.*

OPERCUL'ARIA. G. OPERCUL'ARIA. Tetrandria Monogynia. W.

1 umbellàta. g. (wh.) umbel-flower'd. N. Holland.	1818.	6. 7. G. ȝ.	Gært. fr. 1. t. 24. f. 4.
2 Lamarckiàna. R.S. (wh.) Lamarck's.	N. S. W.	1825. 7. 9. H. ⊖.	Lam. ill. t. 58. f. 1.
3 áspéra. s.s. (wh.) rough.	N. Zealand.	1790. 6. 7. G. ȝ.	Gært. fr. 1. t. 24. f. 4.
4 hispida. s.s. (wh.) hispid.	N. S. W.	— — G. ȝ.	Juss. an. mus. 4. t. 70. f. 1.
aspera. J.			

5 sessiliflòra. R.S. (wh.) sessile-flow'r'd.	— —	1816. — — G. ȝ.	Juss. an. mus. 4. t. 70. f. 2.
6 ocympidòlia. R.S. (wh.) Basil-leaved.	— —	1820. — — G. ȝ.	

CRYPTOSPE'RUM. P.S. CRYPTOSPERMUM. Tetrandria Monogynia. P.S.

Yoñgii. P.S. (gr. re.) Young's.	N. S. W.	1793. 7. 8. G. ȝ.	Linn. trans. 3. t. 5.
Opercularia paleata. R.S.			

ORDO CXVII.

VALERIANEÆ. DC. fl. fr. ed. 3. v. 4. p. 418.

CENTRA'NTHUS. DC. CENTRA'NTHUS. Monandria Monogynia. R.S.

1 rùber. DC. (re.) common.	England. 6. 10. H. ȝ.	Eng. bot. v. 22. t. 1531.
Valeriana rùbra. E.B.			
a coccineus.	(sc.) scarlet.	— —	H. ȝ.
β cárneus.	(fl.) flesh-coloured.	— —	H. ȝ.
γ álbus.	(wh.) white-flowered.	— —	H. ȝ.

2 intermèdius.	(re.) spear-leaved.	Europe.	— — H. ȝ.
3 angustitòlius. D.C. (re.) narrow-leaved.	— —	1731. 5. 7. H. ȝ.	
4 Calcítrapa. R.S. (pi.) cut-leaved.	Portugal.	1570. — — H. ⊖.	

VALERIANA. DC. VALERIAN. Triandria Monogynia. L.

1 dioica. E.B.	(bh.) diocious.	Britain. 5. 7. H. ȝ.
2 Phù. R.S.	(wh.) garden.	Germany.	1597. — — H. ȝ.
3 officinális. E.B.	(bh.) great wild.	Britain. 6. 7. H. ȝ.
4 tenuífolia. Vahl.	(bh.) linear-leaved.	Germany. — — H. ȝ.
5 sambucifòlia. R.S. (bh.)	Elder-leaved.	Sweden.	1818. — — H. ȝ.

6 cardámines. M.B. (bh.) Ladies smock-l'd.	Caucasus.	1823. 6. 7. H. ȝ.	Flor. græc. 1. t. 29.
7 sisymbriifòlia. R.S. (pi.) water-rocket-l'd.	Levant.	1820. — — H. ȝ.	— — 1. t. 30.
8 montâna. R.S. (bh.) mountain.	Switzerland.	1748. — — H. ȝ.	
9 trípteris. R.S. (bh.) three-leaved.	— —	1752. 3. 5. H. ȝ.	
10 pyrénaïca. E.B. (ro.) heart-leaved.	Scotland. 5. 6. H. ȝ.	Eng. bot. v. 9. t. 628.

11 alliarizéfòlia. R.S. (wh.)	Alliaria-leaved.	Levant.	1825. 5. 8. H. ȝ.
12 tubérösa. R.S.	(ro.) tuberous-root'd.	S. Europe.	1629. — — H. ȝ.
13 elongàta. R.S.	(st.) elongated.	Austria.	1812. 6. 7. H. ȝ.
14 supina. R.S.	(ro.) creeping-root'd.	Germany.	1818. — — H. ȝ.
15 saxátilis. R.S.	(wh.) rock.	— —	1748. 7. H. ȝ.

16 saliúncia. R.S.	(bh.) headed-flower'd.	Italy.	1824. 6. 7. H. ȝ.
17 céltica. R.S.	(st.) raceme-flower'd.	Switzerland.	1748. — — H. ȝ.
18 capénisia. R.S.	(bh.) Cape.	C. B. S.	1816. — — F. ȝ.
PATRÍ'NIA. J.	PATRÍ'NIA.	Tetrandria Monogynia.	R.S.

1 sibírica. J.	(ye.) Siberian.	Siberia.	1759. 5. 6. H. ♂.
Valeriana sibírica. L. ruthénica. B.M.			
2 coronàta. s.s.	(ye.) crowned.	Altay.	1826. 5. 7. H. ♂.
3 rupéstris. J.	(re.) rock.	Siberia.	1801. — — H. ♂.
Valeriana sibírica. B.M.			

4 intermédia. R.S.	(ye.) intermediate.	— —	1823. — — H. ♂.
5 nudiuscula. s.s.	(ye.) naked-stem'm'd.	Altay.	1826. 6. 8. H. ♂.
6 scabiosæfòlia. s.s.	(ye.) Scabious-leav'd.	Dahuria.	— — H. ȝ.
7 serratulæfòlia. F.	(ye.) Serratula-leav'd.	— —	1826. — — H. ȝ.
F'EDIA. DC.	F'EDIA.	Diandria Monogynia.	

Cornucòpiae. R.S. (pu.) purple.	S. Europe.	1596. 5. 8. H. ⊖.	Bot. reg. t. 155.
Valeriana cornucòpiae. B.R.			

Bot. mag. t. 2325.
Jacq. aust. 3. t. 268.
Eng. bot. v. 23. t. 1591.
Buxb. cent. 2. t. 11.
Lapeyr. pyr. t. 53.

Jacq. ic. 3. t. 219.
Jacq. misc. 2. t. 17. f. 2.
Jacq. aust. 3. t. 267.
All. ped. 1. t. 70. f. 1.
Jacq. coll. 1. t. 1.

Swt. br. fl. gar. t. 154.

VALERIANELLA. DC. VALERIANELLA. *Triandria Monogynia.*

1 echinata. DC.	(<i>li.</i>) prickly-capsul'd. S. Europe.	1807.	7. 8. H. ⊙.	Reich. ic. C. 1. t. 68.
2 olitòria. DC.	(<i>bh.</i>) Lamb's Lettuce. Britain.	4. 8. H. ⊙.	Eng. bot. v. 12. t. 811.
	<i>Valeriana locusta.</i> E. B. <i>Fèdia olitòria.</i> R.S. <i>Fèdia locusta.</i> Reich. ic. t. 60.			
3 carinata. DC.	(<i>bh.</i>) keeled.	France.	H. ⊙.
4 dentata. DC.	(<i>bh.</i>) oval-fruited.	Britain.	H. ⊙.
	<i>Valeriana dentata.</i> E.B. <i>Fèdia dentata.</i> Reich. ic. 1. t. 62.			
5 discoidea. DF.	(<i>bh.</i>) discoid.	Italy.	1731.	4. 7. H. ⊙.
6 rotata. R.S.	(<i>bh.</i>) rotate.	—	1828.	H. ⊙.
7 vesicaria. DF.	(<i>bh.</i>) bladdered.	Candia.	1739.	H. ⊙.
8 hamata. DC.	(<i>bh.</i>) hooked.	France.	1825.	H. ⊙.
9 coronata. DC.	(<i>bh.</i>) prickly seeded.	Portugal.	1731.	H. ⊙.
10 radiata. DF.	(<i>bh.</i>) rayed.	N. America.	1818.	H. ⊙.
11 auricula. DC.	(<i>bh.</i>) eared.	Montpelier.	—	H. ⊙.
12 pumila. DC.	(<i>re.</i>) small.	S. Europe.	1823.	H. ⊙.
13 mixta. DF.	(<i>bh.</i>) mixed.	—	1828.	H. ⊙.
14 campanulata. Biv. (<i>bh.</i>) bell-flowered.	Italy.	1829.	H. ⊙.	
15 eriocarpa. DC.	(<i>bh.</i>) woolly-capsul'd.	France.	1823.	H. ⊙.
16 truncata. s.s.	(<i>bh.</i>) truncate.	Crete.	1829.	H. ⊙.
17 microcarpa. s.s.	(<i>bh.</i>) small-capsuled.	S. Europe.	1828.	H. ⊙.
18 chenopodiifolia. R.S. (<i>bh.</i>) Goosefoot-l'd.	N. America.	H. ⊙.	
19 uncinata. DF.	(<i>bh.</i>) uncinate.	Tauria.	1818.	H. ⊙.
20 dasycarpa. L.en.	(<i>bh.</i>) thick-capsuled.	—	—	H. ⊙.
21 muricata. M.B.	(<i>bh.</i>) muricate.	Caucasus.	1828.	H. ⊙.
22 tridentata. L.en.	(<i>bh.</i>) three-toothed.	Tauria.	1823.	H. ⊙.
23 angulosa. DC.	(<i>bh.</i>) angular.	—	1826.	H. ⊙.
24 congesta. B.R.	(<i>ro.</i>) close-headed.	N. America.	—	5. 6. H. ⊙.

ORDO CXVIII.

DIPSACEÆ. *Juss. gen. 194.*

MOR'INA. R.S.

pérsica. R.S.

DI'PSACUS. S.S.

fullonum. s.s.

sylvestris. s.s.

laciñatus. s.s.

divaricatus. s.s.

Gmelini. s.s.

férox. s.s.

pilosus. s.s.

inermis. F.I.

SCABITOSA. S.S.

arvénensis. s.s.

Trichera arvénensis. R.S.

campéstris. Bes.

sylvatica. s.s.

montána. s.s.

longifolia. s.s.

pubescens. s.s.

révera. s.s.

ciliata. s.s.

hy'brida. s.s.

integrifolia. s.s.

Hornemanni. s.s.

austrális. s.s.

SUCC'ISA. S.S.

pratensis. s.s.

Scabiosa Succisa. L.

glabrata.

Scabiosa glabrata. R.S.

syriaca. s.s.

Vaillanti. R.S.

transylvanica. s.s.

MOR'INA. *Diandria Monogynia.* L.

(re.) Persian. Persia. 1740. 7. 8. G. 4.

TEASEL. *Tetrandria Monogynia.* L.

(li.) manured. Britain. 7. H. ♂.

(li.) wild. — — H. ♂.

(li.) cut-leaved. Germany. 1683. 7. 8. H. ♂.

(li.) spreading-leav'd. Sicily. 1827. — — H. ♂.

(bl.) Gmelin's. Caucasus. 1820. — — H. ♂.

(bh.) fierce. Corsica. — — H. ♂.

(wh.) small. Britain. 8. H. ♂.

(st.) spineless. Nepal. 1820. 8. 10. H. ♂.

SCABIOSA. *Tetrandria Monogynia.* L.

(li.) field. Britain. 7. 10. H. 4.

Poland. 1824. — — H. 4.

(re.) broad-leaved. Austria. 1633. 7. H. 4.

(bl.) mountain. Caucasus. 1823. — — H. 4.

(li.) long-leaved. Hungary. 1802. 7. 8. H. 4.

(st.) pubescent. — 1820. — — H. 4.

(li.) reversed. Siberia. 1829. — — H. 4.

(st.) ciliated. Hungary. 1820. — — H. 4.

(li.) hybrid. S. Europe. — 6. 9. H. ⊙.

(pu.) entire-leaved. — 1748. 6. 8. H. ⊙.

(li.) Horneman's. 1825. — — H. 4.

(ye.) creeping-root'd. Italy. 1823. — — H. 4.

SUCC'ISA. S.S.

(bl.) Devil's-bit. Britain. 8. 10. H. 4.

Scabiosa Succisa. L.

(bl.) smooth-leaved. Austria. 1823. 7. 9. H. 4.

Syria. 1633. 7. H. 4.

Levant. 1826. — — H. ⊙.

Hungary. 1699. — — H. ⊙.

Flor. græc. 1. t. 28.

Eng. bot. v. 29. t. 2080.

—— v. 15. t. 1032.

Jacq. aust. 5. t. 403.

Jacq. f. ecl. t. 106.

—— t. 107.

Eng. bot. v. 13. t. 877.

Eng. bot. v. 10. t. 659.

Jacq. aust. 4. t. 362.

Jacq. f. eclog. t. 60.

W. et K. hung. 1. t. 5.

Eng. bot. v. 13. t. 878.

Flor. græc. t. 105.

Jacq. vind. 2. t. 111.

6 joppica. s.s.	(st.) Joppa.	Joppa.	1828.	7.	H. ⊖.	Reich. H. bot. t. 15.
7 uralensis. s.s.	(st.) Uralian.	Siberia.	1789.	7. 8.	H. ⊖.	Murr. c. got. 1782. t. 4.
8 tatárica. s.s.	(ye.) Tartarian.	Russia.	1759.	6. 8.	H. ♀.	Lin. act. ups. 1744. t. 1.
19 alpina. s.s.	(st.) Alpine.	Switzerland.	1570.	6. 7.	H. ♀.	Moris. s. 6. t. 13. f. 10.
Cephalaria alpina. R.S.	Scabiōsa alpīna. w.					
10 leucantha. s.s.	(wh.) white flowered.	S. Europe.	1633.	9. 10.	H. ♀.	Ger. emac. f. 8.
11 centauroides. s.s.	(st.) Centaury-like.	——	1816.	7. 8.	H. ♀.	W. et K. hung. 3. t. 230.
12 levigata. s.s.	(st.) smooth.	Hungary.	1823.	——	H. ♀.	—— 1. t. 13.
13 corniculata.	(st.) horned.	——	1801.	——	H. ♀.	
Cephalaria corniculata. R.S.	Scabiōsa corniculata. W.K.					
14 cretacea. M.B.	(st.) leathery-leaved.	Tauria.	1828.	——	H. ♀.	
15 rigidia. s.s.	(wh.) rough-leaved.	C. B. S.	1731.	7.	G. ḥ.	
16 attenuata.	(wh.) narrow-leaved.	——	1774.	7. 9.	G. ḥ.	
trifida. s.s.	Scabiōsa attenuata. L.					
ASTEROCE'PHALUS. S.S.	ASTEROC'PHALUS. Tetrandria Monogynia. S.S.					
1 prolifer. s.s.	(st.) prolific.	Egypt.	1683.	7. 9.	H. ⊖.	Flor. græc. t. 107.
Scabiōsa prolifera. L.						
2 graminifolius. s.s. (bl.)	grass-leaved.	Switzerland.	——	6.	H. ♀.	Bot. reg. t. 835.
Scabiōsa graminifolia. B.R.						
3 crēticus. s.s.	(li.) Cretan.	Crete.	1596.	6. 10.	G. ḥ.	Moris. s. 6. t. 15. f. 31.
4 limoniifolius. s.s.	(li.) Limonium-leav'd. Sicily.	——	6. 8.	H. ḥ.	
5 Massōni.	(li.) Masson's.	Azores.	1779.	——	H. ♀.	
lūcidus. s.s.	Scabiōsa lūcida. H.K.	Massōni. H.S.L. nītens. R.S.				
6 dichotomus. s.s.	(re.) forked.	Sicily.	1804.	6. 8.	H. ⊖.	Bocc. mus. t. 120.
7 africanus. s.s.	(wh.) African.	Africa.	1690.	7. 10.	G. ḥ.	Herm. parad. t. 219.
8 palæstinus. s.s.	(st.) Palestine.	Palestine.	1771.	7. 8.	H. ⊖.	Jacq. vind. 1. t. 96.
9 lūcidus.	(li.) shining.	France.	1800.	6. 9.	H. ♀.	
Scabiōsa lūcida. R.S.						
10 nōricus. R.S.	(li.) simple-stalked.	Caucasus.	1822.	——	H. ♀.	
11 strīctus. s.s.	(li.) upright.	Hungary.	1827.	——	H. ♀.	
12 silenifolius. s.s.	(li.) Silene-leaved.	——	1823.	7. 8.	H. ♀.	
13 canescens. s.s.	(li.) canescent.	——	1802.	——	H. ♀.	
14 suavolens. s.	(li.) sweet-scented.	France.	1827.	——	H. ♀.	
15 setifer. s.s.	(wh.) setiferous.	——	1820.	——	H. ♀.	
16 lyrātus. s.s.	(li.) lyrate-leaved.	Turkey.	1799.	——	G. ♂.	
17 micrānthus. s.s.	(ro.) small-flowered.	Armenia.	1823.	——	H. ♀.	
18 Biebersteini.	(li.) Bieberstein's.	Caucasus.	——	——	H. ⊖.	
Scabiōsa Biebersteini. R.S. sīcula. M.B.						
19 sīculus. s.s.	(li.) Sicilian.	Sicily.	1783.	——	H. ⊖.	Desf. an. mus. 11. t. 25.
20 am'ēnus. s.s.	(ro.) glossy.	——	1823.	——	H. ♀.	Jacq. vind. 1. t. 15.
21 stellātus.	(li.) starry.	Spain.	1596.	——	H. ⊖.	Moris. s. 6. t. 15. f. 39.
22 ceratophyllus. R.S.	(li.) stag's-horn-leav'd. Naples.	Naples.	1825.	——	H. ♂.	Ten. fl. nap. 1. t. 8.
23 rotātus. s.s.	(li.) rotate.	Iberia.	1827.	——	H. ⊖.	Reich. hort. t. 16.
24 diffūsus. s.s.	(li.) diffuse.	Teneriffe.	1825.	6. 8.	F. ♀.	Bot. mag. t. 247.
25 atropūrēus. s.s. (pu.) sweet.	E. Indies.	1629.	7. 9.	H. ♂.		
Scabiōsa atropūrēa. B.M.						
β cárneus.	(fl.) flesh-coloured.	——	——	——	H. ♂.	
γ rōseus.	(ro.) rose-coloured.	——	——	——	H. ♂.	
δ variegātus.	(v.) variegated.	——	——	——	H. ♂.	
ε álbis.	(wh.) white-flowered.	——	——	——	H. ♂.	
ζ prolifer.	(d.pu.) large-flowered.	——	1814.	——	H. ♂.	
26 caucásicus. s.s.	(wh.) Caucasean.	Caucasus.	1803.	7. 8.	H. ♀.	Moris. s. 6. t. 14. f. 2.
27 connātus. H.H.	(bh.) connate-leaved.	——	1823.	——	H. ♀.	Jacq. fragm. t. 125. f. 1.
28 élégans. R.S.	(vi.) elegant.	——	1803.	7. 8.	H. ♀.	Bot. mag. t. 886.
Scabiōsa caucásica. B.M. élégans. R.S.						
29 holosericeus. s.s.	(bl.) silky.	Pyrenees.	——	——	H. ♀.	
30 pyrenaicus. R.S.	(li.) Pyrenean.	——	1823.	——	H. ♀.	Cavan. icon. 2. t. 183.
31 tomentōsus. s.s.	(vi.) woolly.	Spain.	1826.	——	H. ♀.	Bot. reg. t. 717.
32 Webbianus. s.s.	(wh.) Webb's.	Levant.	1820.	——	H. ♀.	
Scabiōsa Webbiana. B.R.						
33 crenātus. s.s.	(fl.) crenated.	Sicily.	1824.	7. 8.	H. ♀.	Cyril. pl. rar. 1. t. 3.
34 ruteafolius. R.S.	(fl.) Rue-leaved.	——	1804.	6. 8.	H. ♀.	Bocc. sic. t. 40. f. 3.
35 urceolātus. s.s.	(st.) jagged.	Barbary.	——	7. 8.	H. ♀.	—— t. 52.
36 argénteus. s.s.	(wh.li.) silvery.	Levant.	1713.	6. 10.	H. ♀.	Flor. græc. t. 108.
37 marítimus. s.s.	(li.) sea-side.	S. Europe.	1818.	7. 9.	H. ⊖.	Bauh. hist. 3. p. 7. f. 2.
38 ucrānicus. s.s.	(wh.) Ukraine.	Ukraine.	1795.	9.	H. ♀.	Gmel. sib. 2. t. 87.
39 Columbāria. s.s.	(li.) fine-leaved.	Britain.	7. 8.	H. ♀.	Eng. bot. v. 19. t. 1311.
40 grandiflōrus. R.S.	(st.) great-flowered.	Barbary.	1804.	6. 9.	H. ♀.	Scop. del. ins 3. t. 14.
41 mólis. L.en.	(li.) soft.	——	1827.	——	H. ♀.	
Scabiōsa capillāta. R.S.						

42 altissimus. s.s.	(<i>li.</i>) tall.	C. B. S.	1816.	6. 9. G. Ȑ.	Jacq. vind. 2. t. 185.
43 mollissimus. s.s.	(<i>bl.</i>) soft.	S. Europe.	—	7. 8. H. Ȑ.	
44 pilōsus. s.s.	(<i>bh.</i>) hairy.	Spain.	1826.	— H. Ȑ.	
45 ochroleucus. s.s.	(<i>st.</i>) pale-flowered.	Germany.	1597.	— H. Ȑ.	Jacq. aust. 5. t. 439.
46 gramántius. s.s.	(<i>wh. li.</i>) cut-leaved.	S. Europe.	—	— H. Ȑ.	Gerard herb. 582, f. 2.
47 iseténis. s.s.	(<i>wh. li.</i>) Siberian.	Siberia.	1801.	— H. Ȑ.	Gmel. sib. 2. t. 88. f. 1.
48 bannáiticus. s.s.	(<i>fl.</i>) Hungarian.	Hungary.	1800.	— H. Ȑ.	W. et K. hung. 1. t. 12.
49 Colúmnae. Ten.	(<i>li.</i>) Columna's.	Italy.	1823.	— H. Ȑ.	Ten. fl. nap. 1. t. 7.
50 Scopólii. R.S.	(<i>st.</i>) Scopoli's.	S. Europe.	1822.	— H. Ȑ.	
51 comosus. R.S.	(<i>vi.</i>) tufted.	Dahuria.	1828.	— H. Ȑ.	
52 agréstis. s.s.	(<i>li.</i>) field.	Hungary.	—	— H. Ȑ.	W. et K. hung. 3. t. 204.
53 simplex. S.S.	(<i>wh.</i>) simple-stalked.	Barbary.	—	— H. Ȑ.	Desf. atl. 1. t. 39. f. 1.
54 incánus. R.S.	(<i>li.</i>) hoary.	—	—	— H. Ȑ.	
55 dahúricus. F.	(<i>li.</i>) Dahirian.	Dahuria.	1824.	— H. Ȑ.	
PTEROCE'PHALUS. S.S. PTEROCE'PHALUS.		Tetrandria Monogynia. S.S.			
1 plumósus. s.s.	(<i>li.</i>) feathered.	Greece.	1824.	7. 8. H. ⊖.	Flor. græc. 1. t. 3.
Scabiòsa plumòsa. Sm.	Knaútia plumòsa. L.				
2 Vaillántii. s.s.	(<i>wh.</i>) Vaillant's.	S. Europe.	1739.	— H. ⊖.	
Scabiòsa pappòsa. L.					
KNA'UTIA. R.S.	KNA'UTIA.	Tetrandria Monogynia. L.			
1 orientális. R.S.	(<i>re.</i>) red-flowered.	Levant.	1713.	6. 9. H. ⊖.	Schk. handb. 1. t. 22.
2 propóntica. R.S.	(<i>pu.</i>) purple-flowered.	—	1768.	— H. ♂.	Lam. ill. t. 58.

ORDO CXIX.

CALYCEREÆ. Brown in Linn. trans. v. 12. p. 132.

ACICA'RPHA. L.T.	ACICA'RPHA.	Syngenesia Necessaria. P.S.
spathulàta. L.T.	(<i>ye.</i>) spathulate-leav'd. Brazil.	1816. 6. 9. G. Ȑ.

ORDO CXX.

COMPOSITÆ. Adanson fam. 2. 103.

Subordo I. CICHORACEÆ. Kth. synops. 2. p. 352.

Tribus I. HIERACEÆ. D. Don in Edinb. philos. journ.

HIER'ACIUM. W.	HAWKWEED.	Syngenesia Polygamia Æqualis.			
1 alpéstre. w.	(<i>ye.</i>) smooth-leaved.	Austria.	1823.	5. 7. H. Ȑ.	Jacq. aust. 2. t. 191.
2 rupéstre. w.	(<i>ye.</i>) rock.	Switzerland.	—	— H. Ȑ.	All. auct. t. 1. f. 2.
3 alpínum. w.	(<i>ye.</i>) Alpine.	Britain.	—	7. 8. H. Ȑ.	Eng. bot. v. 16. t. 1110.
4 pùmilum. w.	(<i>ye.</i>) small.	Germany.	1804.	— H. Ȑ.	
5 glabrátum. w.	(<i>ye.</i>) smooth.	Switzerland.	1823.	6. 8. H. Ȑ.	
6 bulbósrum. w.	(<i>ye.</i>) bulbous-rooted.	S. Europe.	—	— H. Ȑ.	
7 Pilosélla. w.	(<i>st.</i>) Mouse-ear.	Britain.	—	5. 7. H. Ȑ.	Eng. bot. v. 16. t. 1093.
8 dùbium. w.	(<i>st.</i>) branching.	—	—	7. 8. H. Ȑ.	— v. 33. t. 2332.
9 Aurícula. w.	(<i>go.</i>) umbell'd mouse-ear.	England.	—	— H. Ȑ.	— v. 33. t. 2368.
10 fállax. W.en.	(<i>go.</i>) hairy spear-leav'd.	—	1816.	— H. Ȑ.	
11 angustifólium. w.	(<i>ye.</i>) narrow-leaved.	Mt. Cenis.	1822.	— H. Ȑ.	
12 florentinum. w.	(<i>ye.</i>) Florentine.	Germany.	1796.	— H. Ȑ.	Bauhin pin. t. 67.
13 præáltum. s.s.	(<i>ye.</i>) tall.	Europe.	1824.	— H. Ȑ.	
14 cymosum. w.	(<i>ye.</i>) small-flowered.	—	1739.	5. 6. H. Ȑ.	Moris. s. 7. t. 8. f. 10.
15 piloselloïdes. P.S.	(<i>ye.</i>) Mouse-ear like.	France.	1823.	5. 7. H. Ȑ.	Vill. delph. 3.p.100.t.27.
16 staticifólium. w.	(<i>ye.</i>) thrift-leaved.	Europe.	1804.	6. 7. H. Ȑ.	— 3. p. 116. t. 27.
17 præmórsrum. w.	(<i>ye.</i>) præmorse.	Siberia.	1820.	— H. Ȑ.	Gmel. sib. 2. t. 13.
18 succisaefólium. P.S.	(<i>ye.</i>) Succisa-leaved.	Piedmont.	1822.	— H. Ȑ.	
19 flagellare. W.en.	(<i>ye.</i>) creeping.	—	1816.	5. 7. H. Ȑ.	
20 bifúrcum. M.B.	(<i>ye.</i>) two-forked.	S. Europe.	1822.	— H. Ȑ.	
21 bifidum. W.en.	(<i>ye.</i>) bifid.	Hungary.	—	6. 7. H. Ȑ.	

22	incarnatūm. w.	(fl.) flesh-coloured.	Carniola.	1815.	6. 7. H. 4.	Jacq. ic. t. 578.
23	aurantiacum. w.	(or.) orange-colour'd.	Scotland.	— H. 4.	Eng. bot. v. 21. t. 1469.
24	venosum. w.	(ye.) red-veined.	N. America.	1790.	— H. 4.	
25	Gronovii. w.	(ye.) Gronovius's.		1798.	— H. 4.	
26	Gmelini. w.	(ye.) Gmelin's.	Siberia.		— H. 4.	
27	Bauhinii. s.s.	(ye.) Bauhin's.	Switzerland.	1820.	— H. 4.	Gmel. sib. 2. t. 8. f. 2.
28	Lawsoni. w.	(st.) Lawson's.	Britain.	— H. 4.	Eng. bot. v. 29. t. 2083.
29	collinum. w.	(ye.) hill.	Europe.	1824.	6. 8. H. 4.	
30	Gochnati. s.s.	(ye.) Gochnat's.			— H. 4.	
31	paniculatum. w.	(ye.) panicle.	Canada.	1800.	— H. 4.	
32	chondrilloides. w.	(ye.) Gum-succory.	Austria.	1640.	— H. 4.	
33	humile. w.	(ye.) low.	S. Europe.	1816.	— H. 4.	Jacq. aust. 5. t. 429.
	pumilum.	Jacq. Jacquinii. Vill. delph. 3. t. 28.				— t. 189.
34	calcareum. H.H.	(ye.) spurred.	S. Europe.	1823.	7. 8. H. 4.	
35	sonchifolium. M.B.	(ye.) Sonchus-leaved.	Caucasus.	1828.	— H. 4.	Willd. hort. ber. t. 10.
36	pictum. P.S.	(ye.) painted-leaved.	Switzerland.	1823.	— H. 4.	All. ped. t. 15. f. 2.
37	nigriscens. w.	(ye.) dark-coloured.	1801.	7. 8. H. 4.	Jacq. aust. 2. t. 190.
38	prunellæfolium. w.	(ye.) Prunella-leav'd.	Switzerland.	1823.	6. 7. H. 4.	
39	montanum. w.	(ye.) mountain.	S. Europe.	1775.	— H. 4.	Jacq. aust. 3. t. 286.
40	hybridum. s.s.	(ye.) hybrid.	Switzerland.	1825.	— H. 4.	Jacq. ic. 1. t. 163.
41	porrifolium. w.	(ye.) Leek-leaved.	Austria.	1640.	7. 8. H. 4.	All. ped. t. 81. f. 1.
42	saxatile. w.	(ye.) rock.	1801.	— H. 4.	Eng. bot. v. 31. t. 2210.
43	glaucum. w.	(ye.) glaucous.	S. Europe.	1807.	6. 7. H. 4.	
44	mollis. w.	(go.) soft-leaved.	Scotland.	7. 8. H. 4.	
45	croaticum. W.K.	(ye.) Hungarian.	Hungary.	1820.	— H. 4.	
46	Sternbergii. L.en.	(ye.) Sternberg's.			— H. 4.	
47	pallescens. s.s.	(st.) pale.	1815.	6. 7. H. 4.	
48	murorum. E.B.	(go.) wall.	Britain.	7. H. 4.	
49	maculatum. E.B.	(ye.) stained-leaved.	England.	7. 8. H. 4.	
50	ramosum. w.	(ye.) branching.	Hungary.	1805.	— H. 4.	
51	incisum. s.s.	(ye.) cut-leaved.	1823.	— H. 4.	
52	parvifolium. w.	(ye.) small-leaved.	Austria.		— H. 4.	
53	cycloniæfolium. s.s.	(ye.) Quince-leaved.	France.	1816.	— H. 4.	
54	denudatum. s.s.	(ye.) half-naked.	Pyrenees.	1828.	— H. 4.	Eng. bot. v. 29. t. 2031.
55	sylvaticum. E.B.	(ye.) wood.	Britain.	8. H. 4.	— v. 33. t. 2307.
56	pulmonarium. E.B.	(go.) lungwort.	Scotland.	7. 8. H. 4.	Gouan. ill. t. 21. f. 3.
57	lapsanoïdes. w.	(ye.) Lapsana-like.	Pyrenees.	1812.	— H. 4.	Gmel. sib. 2. t. 9.
58	lyratum. w.	(ye.) lyre-leaved.	Siberia.	1777.	— H. 4.	Eng. bot. v. 34. t. 2378.
59	cerinthoides. E.B.	(ye.) honeywort.	Scotland.	8. H. 4.	
60	flexuosum. w.	(ye.) bending-stalk'd.	Hungary.	1804.	7. 8. H. 4.	All. ped. 1. t. 15. f. 1.
61	amplexicaule. w.	(ye.) heart-leaved.	Pyrenees.	1739.	— H. 4.	W. et K. hung. 1. t. 99.
62	grandiflorum. w.	(ye.) great-flowered.	Europe.	1791.	— H. 4.	Jacq. aust. ap. t. 43.
63	intybaceum. w.	(ye.) endive-leaved.	1794.	— H. 4.	Vill. dauph. 3. t. 26.
64	Halleri. w.	(ye.) Haller's.	1802.	— H. 4.	Eng. bot. v. 34. t. 2379.
65	eriophyllum. L.en.	(ye.) villous.	1816.	— H. 4.	
66	villosum. E.B.	(st.) shaggy alpine.	Scotland.	— H. 4.	
67	Schraderi. DC.	(ye.) Schrader's.	Switzerland.	1823.	— H. 4.	
68	elongatum. Lap.	(ye.) elongated.	1824.	— H. 4.	
69	pilocéphalum. L.en.	(ye.) hairy-headed.	Europe.	1816.	— H. 4.	
70	trichocéphalum. w.	(ye.) bristly-headed.	1823.	— H. 4.	
71	prostratum. s.s.	(ye.) prostrate.	S. Europe.		— H. 4.	
72	calyculatum. H.H.	(ye.) long-scaled.	1826.	— H. 4.	
73	longifolium. H.H.	(ye.) long-leaved.	1823.	— H. 4.	
74	speciosissimum. s.s.	(ye.) handsome.	S. Europe.		— H. 4.	
	speciosum. H.H.					
75	ciliatum. w.	(ye.) fringed.	Crete.	1823.	— H. 4.	
76	Milléri. L.en.	(ye.) Millier's.	— H. 4.	
77	lanatum. w.	(ye.) woolly.	Hungary.	—	— H. 4.	
78	andryalaoides. P.S.	(ye.) Andryala-like.	France.	1818.	— H. 4.	
79	echioïdes. w.	(ye.) viper's-bugloss.	Hungary.	1802.	— H. 4.	
80	verruculatum. L.en.	(ye.) warted.	1823.	— H. 4.	
81	undulatum. w.	(ye.) wave-leaved.	Spain.	1778.	— H. 4.	
82	glutinosum. w.	(ye.) clammy.	S. Europe.	1796.	— H. 4.	
83	Kalmii. w.	(ye.) Kalm's.	Pensylvania.	1794.	8. H. 4.	W. et K. hung. 2. t. 127.
84	fasciculatum. Ph.	(ye.) bundled.	Canada.	7. 8. H. 4.	Vill. delph. 3. t. 29.
85	racemosum. w.	(ye.) racemed.	Hungary.	1818.	— H. 4.	W. et K. hung. 1. t. 85.
86	foliosum. w.	(ye.) leafy.	1805.	— H. 4.	
87	corymbosum. P.S.	(ye.) corymbose.	1820.	— H. 4.	
88	sabaudum. w.	(go.) frutescent.	Britain.	— H. 4.	
						Eng. bot. v. 5. t. 349.

89	umbellatum, w. (ye.) umbel-flowered.	Britain.	8. 9. H. ȝ.	Eng. bot. v. 25. t. 1771.
90	prenanthoides. E.B. (ye.) rough-bordered.	Scotland.	6. 9. H. ȝ.	— v. 32. t. 2235.
91	latifolium. s.s. (ye.) broad-leaved.	Pyrenees.	1823.	— H. ȝ.	Eng. bot. v. 30. t. 2122.
92	denticulatum. E.B. (ye.) small-toothed.	Scotland.	7. 8. H. ȝ.	Willd. hort. ber. t. 16.
93	lævigatum. w. (ye.) smooth.	1804.	8. 9. H. ȝ.	
94	canescens. L.en. (ye.) canescent.	1820.	7. 8. H. ȝ.	
95	auriculatum. s.s. (ye.) ear-leaved.	1826.	— H. ȝ.	
96	boreale. s.s. (ye.) northern.	N. Europe.	1828.	— H. ȝ.	
97	bracteolatum. L.en. (ye.) bracted.	S. Europe.	1823.	— H. ȝ.	
98	coronopifolium. L.en. (ye.) Buckshorn-ld.	—	— H. ȝ.	
99	parviflorum. Schl. (ye.) small-flowered.	Switzerland.	—	— H. ȝ.	
100	ochroleucum. Schl. (st.) sulphur-color'd.	1821.	— H. ȝ.	
101	ovatum. Schl. (ye.) oval-leaved.	1820.	— H. ȝ.	
102	compositum. s.s. (ye.) compound.	Pyrenees.	1823.	— H. ȝ.	
103	crassifolium. s. (ye.) thick-leaved.	—	— H. ȝ.	
104	rotundatum. s.s. (ye.) round-leaved.	Hungary.	—	— H. ȝ.	
105	sudeticum. St. (ye.) Swedish.	Sweden.	—	— H. ȝ.	
106	virgatum. Ph. (ye.) twiggy.	N. America.	1816.	— H. ȝ.	
107	virescens. Schl. (gr.) greenish.	Switzerland.	1820.	— H. ȝ.	
108	stoloniferum. Viv. (ye.) stoloniferous.	Levant.	1823.	— H. ȝ.	
109	verbascifolium. P.S. (ye.) Mullein-leav'd.	S. Europe.	1732.	5. 6. H. ȝ.	Mill. ic. 1. t. 146. f. 1.
<i>Andryala lanata</i> , s.s.					
HAPALOST'EPHIUM. D.P. HAPALOST'EPHIUM. Syngenesia Polygamia Äqualis.					
1	paludosum. D.P. (ye.) Succory-leaved.	Britain.	7. 8. H. ȝ.	Eng. bot. t. 1094.
<i>Hieracium paluddosum</i> . E.B.					
2	pyrenaicum. D.P. (ye.) Pyrenean.	Pyrenees.	1723.	— H. ȝ.	
<i>Hieracium blattarioides</i> . P.S.					
3	pilosum. D.P. (ye.) hairy.	—	—	— H. ȝ.	Herm. parad. t. 184.
4	austriacum. (ye.) Austrian.	Austria.	—	— H. ȝ.	Jacq. aust. t. 441.
5	sibiricum. D.P. (ye.) Siberian.	Siberia.	1755.	— H. ȝ.	Gmel. sib. 2. t. 10.
<i>Crépis sibirica</i> . L. <i>Hieracium sibiricum</i> . w.					
6	grandiflorum. (ye.) large-flowered.	Idria.	1824.	6. 8. H. ȝ.	
<i>Crépis grandiflora</i> . L.					
LAGO'SERIS. L.en. LAGO'SERIS. Syngenesia Polygamia Äequalis.					
1	bursifolia. L.en. (ye.) pinnatifid.	Sicily.	1823.	7. 8. H. ȝ.	Bocc. m. 2. t. 106 et 112.
<i>Crépis bursifolia</i> . L.					
2	versicolor. L.en. (ye.) various-color'd.	Dauria.	—	— H. ȝ.	
3	cernua. L.en. (ye.) nodding.	Calabria.	—	6. 8. H. ⊖.	
4	nemausensis. (ye.) Palestine.	S. Europe.	1794.	6. 7. H. ⊖.	Vill. delph. 3. t. 26.
<i>Crepis nemausensis</i> . w.					
5	leontodontoides. L.en. (ye.) Dandelion-like.	Italy.	1804.	7. 8. H. ♂.	
6	raphanifolia. L.en. (ye.) Radish-leaved.	1816.	6. 7. H. ȝ.	
7	tauriniensis. L.en. (ye.) bladdered.	S. Europe.	1823.	5. 7. H. ⊖.	
8	intybacea. L.en. (ye.) Succory-leaved.	Portugal.	1816.	6. 7. H. ♂.	
CR'EPI. S.S. CR'EPI. Syngenesia Polygamia Äequalis. L.					
1	nudicaulis. s.s. (ye.) naked-stalked.	Crete.	1828.	6. 8. H. ȝ.	
2	rigens. s.s. (ye.) bristle-leaved.	Azores.	1778.	7. 8. F. ȝ.	
3	ramosissima. s.s. (ye.) branched.	Archipelago.	1828.	— F. ȝ.	
4	virens. w. (ye.) green.	Switzerland.	1796.	6. 7. H. ⊖.	Lobel. ic. 229. f. 2.
5	bannatica. s.s. (ye.) Hungarian.	Hungary.	1823.	7. 8. H. ⊖.	
6	parviflora. s.s. (ye.) small-flowered.	Levant.	1829.	— H. ⊖.	
7	rhagadioloïdes. s.s. (ye.) Spanish.	Spain.	1818.	6. 7. H. ⊖.	Jacq. schoenb. 2. t. 144.
<i>Picris Rhagadiolus</i> . P.S.					
8	aspera. s.s. (ye.) rough.	Sicily.	1797.	7. 8. H. ⊖.	Flor. græc. t. 804.
9	Sprengeriana. s.s. (ye.) Portugal.	Portugal.	1783.	6. 7. H. ⊖.	Moris. his. 7. t. 5. f. 17.
<i>Picris Sprengeriana</i> . P.S. <i>Hieracium Sprengerianum</i> . L.					
10	hieracioïdes. s.s. (ye.) hawkweed-like.	Hungary.	1816.	7. 8. H. ȝ.	W. et K. hung. 1. t. 17.
11	rígida. s.s. (ye.) rigid.	—	1805.	5. 7. H. ȝ.	— 1. t. 19.
12	púlchra. s.s. (ye.) hawkweed-leav'd.	Scotland.	6. 9. H. ⊖.	Eng. bot. v. 33. t. 2325.
13	nána. s.s. (ye.) dwarf.	N. America.	1827.	— H. ȝ.	
14	heterosperma. s.s. (ye.) various-seeded.	Germany.	1824.	— H. ⊖.	
15	Dioscòridis. s.s. (ye.) Dioscorides'.	France.	1772.	6. 7. H. ⊖.	Flor. græc. t. 808.
16	Iodomiriensis. s.s. (ye.) rough-headed.	Germany.	1827.	6. 9. H. ⊖.	
<i>Gmeini</i> . s.					
17	neglecta. s.s. (ye.) neglected.	—	— H. ⊖.	
18	scábra. s.s. (ye.) rough.	S. Europe.	1829.	— H. ♂.	
19	agréstis. s.s. (ye.) field.	Hungary.	1801.	7. 8. H. ⊖.	
20	corymbosa. s.s. (ye.) corymbose.	Italy.	1828.	— H. ♂.	
21	cérnua. s.s. (ye.) nodding-flower'd.	Naples.	1824.	— H. ♂.	

COMPOSITÆ.

22 tectorum. E.B.	(ye.) smooth.	Britain.	6. 9. H. ⊖.	Eng. bot. v. 16. t. 1111.
23 Lachenali. DC.	(ye.) entire-leaved.	Europe.	6. 9. H. ⊖.	Balb. misc. p. 37. t. 9.
24 cinerea. P.S.	(ye.) red-stalked.	Britain.	— H. ⊖.	Eng. bot. v. 3. t. 149.
25 taurinensis. W.	(ye.) early-flowered.	Europe.	5. 6. H. ⊖.	— v. 6. t. 406.
26 biennis. s.s.	(ye.) biennial.	Britain.	6. 7. H. ♂.	
27 foetida. E.B.	(st.) strong-scented.	—	—	— H. ♂.	
28 graveolens.	(ye.) strong-smelling.	1824.	7. 9. H. ⊖.		
29 pulcherrima. F.	(ye.) elegant.	Siberia.	1823.	6. 8. H. ⊖.	
30 latifolia. P.S.	(ye.) broad-leaved.	Italy.	1816.	— H. ⊖.	
31 pinnatifida. s.s.	(ye.) pinnatifid.	S. Europe.	1820.	4. 6. H. ⊖.	
32 lacerata. s.s.	(ye.) torn-leaved.	Naples.	1826.	6. 7. H. ⊖.	
BEHRI'NIA. Sieb.	BEHRI'NIA. Syngenesia Polygamia Æqualis.				
chondrilloides. Sieb.	(ye.) Chondrilla-like. Carniola.	1829.	6. 7. H. ⊖.		
Crépis chondrilloides. s.s.	Wibellia chondrilloides. Hoppe.				
PRENA'NTHES. D.D.	PRENA'NTHES. Syngenesia Polygamia Æqualis. L.				Jacq. aust. 4. t. 317.
1 purpurea. w.	(pu.) purple-flower'd.	Germany.	1658.	7. 9. H. ¼.	
2 hispida. s.s.	(ye.) hispid.	Siberia.	1823.	6. 8. H. ¼.	
3 diversifolia. s.s.	(ye.) various-leaved.	—	1827.	— H. ♂.	
HARPAL'YCE. D.D.	HARPAL'YCE. Syngenesia Polygamia Æqualis.				
1 altissima. D.D.	(ye.) tallest.	N. America.	1696.	7. 8. H. ¼.	
Prenánthes altissima. W.					Pluk. alm. t. 317. f. 2.
2 cordata. D.D.	(ye.) heart-leaved.	—	1816.	— H. ¼.	
3 virgata. D.D.	(li.) tall slender.	—	1823.	8. 9. H. ¼.	
4 álba. D.D.	(wh.) white-flowered.	N. America.	1762.	7. 8. H. ¼.	
Prenánthes álba. B.M. suavis. Sal. par. lond. t. 85.					Bot. mag. t. 1079.
5 rubicunda. D.D.	(fl.) rubicund.	—	1823.	8. 10. H. ¼.	
6 serpentaria. D.D.	(pu.) Lion's-foot.	—	—	— H. ¼.	
7 racemosa. D.D.	(pu.) racemed.	Canada.	1822.	— H. ¼.	
CHORI'SMA. D.D.	CHORI'SMA. Syngenesia Polygamia Æqualis.				Pursh am. sept. 2. t. 24.
rēpens. D.D.	(ye.) creeping.	Siberia.	1828.	7. 9. H. ¼.	
Prenánthes rēpens. L.					Linn. am. ac. 2. t. 4. f. 23.
ANDRYALA. W.	ANDRYALA. Syngenesia Polygamia Æqualis.				
1 cheiranthifolia. w.	(ye.) various-leaved.	Madeira.	1777.	5. 10. F. ¼.	
2 nigricans. w.	(ye.) dark-headed.	Barbary.	1804.	6. 8. H. ⊖.	
3 pinnatifida. w.	(ye.) wing-leaved.	Madeira.	1778.	7. 8. F. ♂.	
4 crithmifolia. w.	(ye.) Samphire-leav'd.	—	—	6. 8. F. ♂.	
5 rugosina. w.	(ye.) downy.	Archipelago.	1753.	— F. ¼.	
6 incana. DC.	(ye.) hoary.	Pyrenees.	1826.	— H. ¼.	
RO'THIA. W.	RO'THIA. Syngenesia Polygamia Æqualis.				
1 runcinata. w.	(ye.) hoary.	S. Europe.	1711.	7. 8. H. ♂.	
2 cheiranthifolia. w.	(ye.) Stock-leaved.	Spain.	1768.	— H. ⊖.	
3 andryaloïdes. w.	(ye.) Spanish.	—	1810.	8. H. ⊖.	
Andryala Róthia. P.S.					Gärt. sem. 2. t. 174.
LAPS'AÑA. D.D.	NIPPLE-WORT. Syngenesia Polygamia Æqualis.				
1 communis. w.	(ye.) common.	Britain.	6. 9. H. ⊖.	Eng. bot. v. 12. t. 844.
2 lyrata. W.en.	(ye.) lyrate.	Caucasus.	1816.	7. 8. H. ¼.	
3 grandiflora. M.B.					
4 intermedia. M.B.	(ye.) intermediate.	—	1823.	— H. ⊖.	
4 pubescens. Zey.	(ye.) pubescent.	Europe.	1820.	— H. ⊖.	
5 crispa. W.en.	(ye.) curled.	1799.	— H. ⊖.	
<hr/>					
Tribus II. TARAXACEÆ. D.Don.					
LEO'NTODON. W.	DANDELION. Syngenesia Polygamia Æqualis. L.				
1 Taráxacum. w.	(ye.) common.	Britain.	4. 7. H. ¼.	
2 corniculatum. H.H.	(ye.) horned.	Europe.	1827.	— H. ¼.	Eng. bot. v. 8. t. 510.
3 bessarabicus. F.	(ye.) small-flowered.	S. Europe.	1823.	6. 8. H. ¼.	
4 alpinus. s.s.	(ye.) Alpine.	—	1827.	— H. ¼.	
5 caucasicus. M.B.	(ye.) Caucasean.	Caucasus.	—	— H. ¼.	
6 serotinus. w.	(ye.) late-flowered.	Hungary.	1816.	7. 9. H. ¼.	
7 glaucescens. s.s.	(ye.) glaucous.	Caucasus.	1820.	4. 7. H. ¼.	W. et K. hung. 2. t. 114.
8 obovatus. s.s.	(ye.) obovate-leaved.	Spain.	1805.	7. H. ¼.	
9 palustris. E.B.	(ye.) marsh.	Britain.	6. 7. H. ¼.	Eng. bot. v. 8. t. 553.
10 nigricans. w.k.	(ye.) dark-leaved.	Hungary.	1823.	7. H. ¼.	
11 lavigatus. s.s.	(ye.) smooth.	Spain.	—	— H. ¼.	W. et K. hung. 2. t. 110.
APA'RGINIA. W.	APA'RGINIA. Syngenesia Polygamia Æqualis.				
1 áspera. s.s.	(ye.) hairy.	Hungary.	1805.	6. 7. H. ¼.	
2 saxatilis. s.s.	(ye.) rock.	Naples.	1827.	— H. ¼.	

3	hispida, s.s.	(ye.) rough.	Britain.	7.	H. ȝ.	Eng. bot. v. 8. t. 554.
<i>Hedy'pnoidis hispida</i> , E. B. <i>Leontodon hispidum</i> . L.							
4	dùbia, w.	(ye.) tooth-leaved.	Germany.	—	—	H. ȝ.	Vill. dauph. 3. t. 25.
5	crispa, s.s.	(ye.) curled.	France.	1803.	7. 8.	H. ȝ.	Cavan. icon. t. 149.
6	fasciculata, s.s.	(ye.) bunch-rooted.	Naples.	1828.	—	H. ȝ.	
7	hispanica, s.s.	(ye.) Spanish.	Spain.	1825.	—	H. ȝ.	
8	strigosa, M.B.	(ye.) strigose.	Caucasus.	1828.	—	H. ȝ.	
9	caucásica, M.B.	(ye.) Caucasean.		1823.	—	H. ȝ.	
10	coronopifolia, s.s.	(ye.) buck's-horn-ld.	Barbary.	1820.	—	H. ȝ.	
11	incàna, s.s.	(ye.) hoary.	Europe.	1784.	5. 6.	H. ȝ.	
12	alpina, s.s.	(ye.) alpine.	Austria.	1816.	6. 8.	H. ȝ.	
13	Villársii, s.s.	(ye.) Villars's.	France.	1818.	—	H. ȝ.	
14	Taráxaci, s.s.	(ye.) Dandelion-like.	Britain.	7. 9.	H. ȝ.	
<i>Hedy'pnoidis Taráxaci</i> , E. B. <i>Hieracium Taráxaci</i> . L. <i>Picris Taráxaci</i> . All.							
15	aurantiaca, w.	(or.) orange-colored.	Hungary.	1816.	5. 6.	H. ȝ.	
16	crócea, w.	(sn.) saffron-colored.		1820.	—	H. ȝ.	
17	hástilis, s.s.	(ye.) shining-leaved.	S. Europe.	1796.	7. 8.	H. ȝ.	
OPORI'NIA. D.D.	OPORI'NIA.	Syngenesia Polygamia Æqualis.					Jacq. aust. 2. t. 164.
autumnális, D.D.	(ye.) autumnal.	Britain.		8. 9.	H. ȝ.	Eng. bot. v. 12. t. 830.
<i>Hedy'pnoidis autumnalis</i> , E. B. <i>Apárgia autumnalis</i> , w.							
THRI'NCIA. W.	THRI'NCIA.	Syngenesia Polygamia Æqualis.					
1	tuberòsa, DC.	(ye.) tuberous-rooted.	France.	1683.	5. 7.	H. ȝ.	Flor. græc. t. 797.
<i>Apárgia tuberòsa</i> , w. <i>Leontodon tuberòsum</i> . L.							
2	glabra, Schl.	(ye.) smooth.	Switzerland.	1823.	7. 8.	H. ȝ.	
3	hírta, w.	(ye.) simple-haired.	Britain.	—	H. ȝ.	Eng. bot. v. 8. t. 555.
<i>pilocályx</i> , Lag? <i>Hedy'pnoidis hírta</i> , E. B.							
4	nudicályx, Lag.	(ye.) naked-calyxed.	Spain.	1823.	—	H. ȝ.	
5	hispida, s.s.	(ye.) hispid.	S. Europe.	1815.	6. 8.	H. ⓧ.	
6	marocána, P.S.	(ye.) Morocco.	Morocco.	1799.	7. 8.	H. ⓧ.	
7	pygmæa, P.S.	(ye.) small.	Madeira.	—	H. ⓧ.	
<i>Hyoseris pygmæa</i> , w.							
CALLI'OPEA. D.D.	CALLI'OPEA.	Syngenesia Polygamia Æqualis.					
aurea, D.D.	(go.) golden-flower'd.	Italy.		1769.	5. 7.	H. ȝ.	Jacq. aust. 3. t. 297.
<i>Leontodon aureum</i> , Jacq. <i>Hieracium aureum</i> , w.							
ÆTH'ONIA. D.D.	ÆTH'ONIA.	Syngenesia Polygamia Æqualis.					
1	fruticosa, D.D.	(ye.) shrubby.	Madeira.	1785.	6. 7.	G. ȝ.	
<i>Hieracium fruticosum</i> , H.K.							
2	filiformis, D.D.	(ye.) fine-leaved.		1777.	—	G. ♂.	
<i>Crépis filiformis</i> , w.							
3	lagopoda, D.D.	hare's-foot.	Canaries.	1817.	—	G. ♂.	
TO'LPI'S. S.S.	TO'LPI'S.	Syngenesia Polygamia Æqualis.					
1	barbata, s.s.	(ye.) purple-eyed.	France.	1620.	6. 9.	H. ⓧ.	Bot. mag. t. 35.
<i>Crépis barbata</i> , B.M. <i>Drepània barbata</i> , DC.							
2	umbellata, s.s.	(ye.) umbelled.	Italy.	1818.	6. 7.	H. ⓧ.	
3	quadriaristata, Biv.	(ye.) four-awned.		1824.	—	H. ⓧ.	
4	virgata, s.s.	(ye.) twiggy.	S. Europe.	1826.	—	H. ⓧ.	
5	altissima, s.s.	(ye.) tall.		1818.	—	H. ⓧ.	
<i>Drepània ambigua</i> , DC. <i>Crépis ambigua</i> , Balbis.							
6	coronopifolia, s.s.	(ye.) fleshy-leaved.	Madeira.	1777.	8. 9.	H. ⓧ.	Desf. act. par. 38. t. 9.
<i>Crépis coronopifolia</i> , w. <i>succulenta</i> , H.K.							
HEDY'PNOIS. W.	HEDY'PNOIS.	Syngenesia Polygamia Æqualis.					
1	monspeliensis, w.	(ye.) branching.	S. Europe.	1683.	6. 7.	H. ⓧ.	
2	crética, w.	(ye.) Cretan.	Crete.	1731.	—	H. ⓧ.	Flor. græc. t. 813.
3	coronopifolia, T.C.	(ye.) buck's-horn-ld.	Italy.	1823.	—	H. ⓧ.	
4	tubæformis, T.C.	(ye.) thick-tubed.		—	—	H. ⓧ.	
5	mauritánica, w.	(ye.) Barbary.	Barbary.		—	H. ⓧ.	
6	rhagadioloïdes, w.	(ye.) Nipple-wort.	S. Europe.	1773.	7. 8.	H. ⓧ.	
7	pérsica, M.B.	(ye.) Persian.		1822.	6. 9.	H. ⓧ.	
8	péndula, w.	(ye.) pendulous.	Barbary.	1818.	—	H. ⓧ.	
9	aculeata, Jac.	(ye.) prickly.		1829.	—	H. ⓧ.	
RHAGAD'IOULS. W.	RHAGAD'IOULS.	Syngenesia Polygamia Æqualis.					
1	stellatus, w.	(ye.) starry.	S. Europe.	1633.	6. 7.	H. ⓧ.	Flor. græc. t. 817.
2	èdulis, w.	(ye.) heart-leaved.	Levant.	—	—	H. ⓧ.	— t. 818.
<i>Lapsana Rhagadiolus</i> , L.							
KÆLP'I'NIA. Pall.	KÆLP'I'NIA.	Syngenesia Polygamia Æqualis.					
linearis, Pall.	(ye.) linear-leaved.	Levant.		1788.	7. 8.	H. ⓧ.	Flor. græc. t. 819.
<i>Lapsana Kælpinia</i> , L. <i>Rhagadiolus Kælpinia</i> , w.							
ARNO'SERIS. G.	ARNO'SERIS.	Syngenesia Polygamia Æqualis.					
1	fa'tida, L.en.	(ye.) stinking.	Italy.	1722.	7. 8.	H. ȝ.	W. et K. hung. I. t. 49.
<i>Lapsana fa'tida</i> , w. <i>Hyoseris fa'tida</i> , P.S.							

COMPOSITÆ.

2 mínima. L.en. (st.) least.	Britain.	5. 7. H. ⊙.	Eng. bot. v. 2. t. 95.
<i>Hyoséris mínima</i> . E.B. <i>Lapsána pusilla</i> . w.	<i>Hyoséris</i> . Syngenesia Polygamia Æqualis.		
1 arenaria. w. (ye.) sand.	Morocco. 1800.	7. 8. H. ⊙.	
2 hispida. w. (ye.) hispid.	——— 1779.	——— H. ⊙.	
3 scabra. w. (ye.) rugged.	Sicily. 1789.	——— H. ⊙.	
4 lúcida. w. (ye.) shining.	Levant. 1770.	6. 8. H. ¼.	
5 radiata. w. (ye.) starry.	S. Europe. 1640.	——— H. ¼.	
ZACI'NTHA. W. <i>Zaci'ntha</i> . Syngenesia Polygamia Æqualis.			
verrucosa. w. (ye.) warted.	S. Europe. 1633.	6. 7. H. ⊙.	
<i>Lapsána Zacy'ntha</i> . L.			
TRO'XIMON. D.D. <i>Tro'ximon</i> . Syngenesia Polygamia Æqualis.			
1 glaucum. Ph. (ye.) glaucous-leaved. N. America.	1811.	7. 8. H. ¼.	
2 cuspidatum. Ph. (ye.) cuspidate. —————	———	4. 6. H. ¼.	
marginatum. N.			
CY'NTHIA. D.D. <i>Cy'nthia</i> . Syngenesia Polygamia Æqualis.			
amplexicaulis. D.D. (go.) stem-clasping. N. America.	1799.	7. 8. H. ¼.	
<i>Hyoséris amplexicaulis</i> . M. <i>Krigia amplexicaulis</i> . N. <i>Tróximon virginicum</i> . Ph.			
KRI'GIA. W. <i>Kri'gia</i> . Syngenesia Polygamia Æqualis.			
virginica. w. (ye.) Virginian.	N. America. 1811.	5. 7. H. ⊙.	
PICRIS. W. <i>Picris</i> . Syngenesia Polygamia Æqualis.			
1 hieracoides. w. (ye.) hawkweed-like.	England.	7. 8. H. ⊙.	
2 strigosa. M.B. (ye.) strigose.	Caucasus. 1823.	——— H. ¼.	
3 rígida. s.s.	Tauria. 1829.	——— H. ¼.	
4 hispida. H.K. (ye.) hispid.	Levant. 1789.	——— H. ¼.	
5 altissima. s.s. (ye.) tall.	Egypt. 1825.	——— H. ⊙.	
6 pauciflora. w. (ye.) few-flowered	S. France. 1818.	6. 8. H. ⊙.	
7 pilosa. s.s. (ye.) hairy.	Egypt. 1829.	——— H. ⊙.	
8 asplenioides. w. (ye.) Spleenwort-ld.	Barbary. 1803.	7. 8. H. ⊙.	
9 dahurica. f. (ye.) Dahurian.	Dahuria. 1820.	——— H. ⊙.	
10 kamtschatica. Led. (ye.) Kamtschata. Kamtschata. 1823.	——— H. ⊙.		
HELM'INTHIA. J. OX-TONGUE. Syngenesia Polygamia Æqualis.			
echoïdes. w. (ye.) bristly.	Britain.	6. 7. H. ⊙.	
<i>Picris echoïdes</i> . E.B.			
<hr/>			
Tribus III. HYPOCHÆRIDÆ. D. Don in Edinb. phil. journ.			
HYPOCHÆRIS. S.S. CAT'S-EAR. Syngenesia Polygamia Æqualis. L.			
1 helvética. s.s. (ye.) one-flowered.	Switzerland. 1799.	6. 7. H. ¼.	Jacq. ic. 1. t. 165.
2 hispida. s.s. (ye.) bristly.	S. Europe. 1804.	7. 8. H. ¼.	Willd. hort. ber. 1. t. 46.
3 arachnites. Biv. (ye.) Cobweb.	Italy.	——— H. ⊙.	
4 mínima. s.s. (ye.) least.	Barbary. 1797.	——— H. ⊙.	Flor. græc. t. 816.
5 glabra. s.s. (st.) smooth.	Britain.	——— H. ⊙.	Eng. bot. v. 8. t. 575.
ACHYRO'PHORUS. G. ACHYRO'PHORUS. Syngenesia Polygamia Æqualis.			
1 maculatus. D.D. (ye.) spotted.	England.	6. 7. H. ¼.	Eng. bot. v. 4. t. 225.
2 radicatus. D.D. (ye.) long-rooted.	Britain.	6. 9. H. ¼.	——— v. 12. t. 831.
<i>Hypoch'aris radicata</i> . E.B.			
3 Balbisii. W.en. (ye.) Balbis's.	S. Europe. 1822.	——— H. ¼.	
ROBE'R'TIA. DC. ROBE'R'TIA. Syngenesia Polygamia Æqualis.			
1 taraxacoides. DC. (ye.) Dandelion-like. Corsica.	1824.	7. 8. H. ¼.	Lois. gal. 530. t. 28.
2 pinnatifida. s.s. (ye.) pinnatifid.	Naples. 1827.	——— H. ¼.	
<i>Hypoch'aris pinnatifida</i> . Tenore.			
SER'IOLA. S.S. SER'IOLA. Syngenesia Polygamia Æqualis. L.			
1 ürens. s.s. (ye.) stinging.	S. Europe. 1773.	7. 8. H. ⊙.	Schmid. ic. t. 32.
2 æthnensis. w. (ye.) rough.	Italy. 1763.	——— H. ⊙.	Jacq. obs. 4. t. 79.
3 glauca. s.s. (ye.) glaucous.	Sicily. 1829.	——— H. ⊙.	
4 rubescens. s.s. (li.) reddish.	———	——— H. ⊙.	
5 álbicans. s.s. (ye.) whitish.	———	——— H. ⊙.	
6 cretensis. s.s. (ye.) Cretan.	Crete. 1820.	——— H. ⊙.	
7 Alliatae. Len. (ye.) wood.	Sicily. 1823.	——— H. ⊙.	
8 apargioïdes. Trev. (ye.) Apargia-like.	S. Europe. 1829.	——— H. ⊙.	
AGE'NORA. D.D. AGE'NORA. Syngenesia Polygamia Æqualis.			
levigata. D.D. (ye.) smooth.	Candia. 1772.	7. 8. H. ⊙.	Desf. atl. 2. t. 216.
<i>Seriola levigata</i> . w. <i>Rodigia levigata</i> . s.s.			
RODI'GIA. S.S. RODI'GIA. Syngenesia Polygamia Æqualis.			
commutata. s.s. (ye.) Poppy-leaved.	Caucasus. 1823.	6. 7. H. ⊙.	
<i>Crépis rhæadifolia</i> . M.B.			
SOLDEVILLA. S.S. SOLDEVILLA. Syngenesia Polygamia Æqualis.			
setosa. s.s. (ye.) bristly.	Spain. 1826.	7. 8. H. ⊙.	

Tribus IV. LACTUCEÆ. D. Don in Edinb. phil. journ.

LACTUCA. S.S.	LETTUCE.	Syngenesia Polygamia Æqualis. L.				
1 sativa. w.	(ye.) garden.	1562.	6. 9. H. ♂.	Rob. ic. pl. c. ic.	
2 crista. w.	(ye.) curled.	1570.	6. 8. H. ♂.		
3 palmata. w.	(ye.) jagged-leaved.	1683.	— H. ♂.		
laciniata. Roth.						
4 intybacea. w.	(ye.) Endive-leaved.	S. America.	1781.	— H. ♂.	Jacq. ic. 1. t. 162.	
5 quercina. w.	(ye.) Oak-leaved.	Europe.	1686.	5. 7. H. ♂.		
6 stricta. w.	(ye.) upright.	Hungary.	1805.	6. 7. H. ♂.	W. et K. hung. 1. t. 48.	
7 elongata. w.	(st.) elongated.	Pennsylvania.	—	— H. ♂.		
8 cichoriifolia. DC.	(ye.) Succory-leaved.	France.	1818.	— H. ♀.		
9 Scariola. w.	(st.) prickly.	Britain.	—	7. 8. H. ♂.	Eng. bot. v. 4. t. 268.	
10 maculata. L.en.	(st.) spot-leaved.	Europe.	—	— H. ♂.		
11 viriosa. E.B.	(st.) strong-scented.	Britain.	—	7. 9. H. ♂.	Eng. bot. v. 28. t. 1957.	
12 augustana. w.	(st.) entire-leaved.	Italy.	1791.	7. 8. H. ♂.	All. ped. 1. t. 52. f. 1.	
13 altissima. M.B.	(ye.) tall.	Caucasus.	1818.	— H. ♀.		
14 sagittata. W.K.	(ye.) arrow-leaved.	Hungary.	1805.	— H. ♂.	W. et K. hung. 1. t. 1.	
15 saligna. w.	(ye.) least.	England.	—	7. 8. H. ♂.	Eng. bot. v. 10. t. 707.	
16 muralis. D.D.	(st.) wall.	Britain.	—	— H. ♀.	— v. 7. t. 457.	
Prenanthes muralis. E.B.						
17 viminea. L.en.	(st.) rushy-twigg'd.	Austria.	1789.	— H. ♂.	Jacq. aust. 1. t. 9.	
Prenanthes viminea. Jacq.						
18 villosa. s.s.	(bl.) villous.	—	1825.	— H. ♂.	Jacq. schoenb. 3. t. 367.	
19 tuberosa. s.s.	(ye.) tuberous.	—	1820.	— H. ♀.	Jacq. vind. 1. t. 57.	
20 indica. s.s.	(ye.) Indian.	E. Indies.	1784.	8. 10. H. ♂.		
21? spectabilis. F.	... showy.	Persia.	1829.	— H. ♂?		
22 pallida. D.P.	(st.) pale-flowered.	Canada.	1704.	7. 9. H. ♀.	Rob. ic. 148 et 151.	
Sónchus pallidus. w.						
23 segusiana. s.s.	(pu.) various-leaved.	Italy.	1823.	7. 8. H. ♂.		
24 tenerrima. s.s.	(pu.) purple-flower'd.	S. Europe.	1815.	— H. ♀.		
25 perennis. s.s.	(bl.) perennial.	Germany.	1596.	6. 8. H. ♀.	Mill. ic. 2. t. 157.	
CHONDRILLA. W.	GUM SUCCORY.	Syngenesia Polygamia Æqualis. L.				
1 junccea. w.	(ye.) common.	France.	1633.	9. 10. H. ♀.	Jacq aust. 5. t. 427.	
2 graminea. M.B.	(ye.) grass-leaved.	Caucasus.	1824.	7. 9. H. ♀.		
3 latifolia. M.B.	(ye.) broad-leaved.	—	—	— H. ♂.		
AGATHYRSUS. D.D. AGATHYRSUS.	Syngenesia Polygamia Æqualis.					
1 alpinus. D.D.	(bl.) Lapland.	Lapland.	1804.	7. 8. H. ♀.	Smith ic. ined. t. 21.	
Sónchus lappónicus. w.						
2 ceruleus. D.D.	(bl.) blue-flowered.	Scotland.	—	— H. ♀.	Eng. bot. v. 34. t. 2425.	
Sónchus alpinus. w. non L.						
3 Plumieri. D.D.	(bl.) Plumier's.	Pyrenees.	1794.	— H. ♀.		
4 floridana. D.D.	(bl.) small-flowered.	N. America.	1713.	— H. ♂.		
5 acuminatus.	(bl.) taper-pointed.	—	1812.	8. 9. H. ♂.		
Sónchus acuminatus. Ph.						
6 macrophyllus.	(bl.) large-leaved.	N. America.	1820.	7. 8. H. ♀.		
Sónchus macrophyllus. Ph.						
7 spicatus. D.D.	(pa.) whitish-flower'd.	—	—	— H. ♂.		
Sónchus leucophæus. w.	Prenanthes autumnalis. Walter.					
8 prenanthoides. D.D. (bl.)	Prenanthes-like.	Caucasus.	1827.	— H. ♀.		
9 cacaliæfolius. D.D. (bl.)	Cacalia-leaved.	Iberia.	1826.	— H. ♀.		
10 sibiricus. D.D. (bl.)	Siberian.	Siberia.	1759.	7. 9. H. ♀.	Gmel. sib. 2. f. 3.	
Sónchus sibiricus. w.	Lactuca sibirica. D.P.					
11 tatáricus. D.D.	(bl.) Tártarian.	Tartary.	1784.	6. 7. H. ♀.		
12 pulchellus. D.D.	(bl.) entire-leaved.	Missouri.	1812.	— H. ♀.		
Sónchus pulchellus. Ph.	Lactuca integrifolia. N.					
13 sonchifolia.	(bl.) Sonchus-leaved.	Levant.	1824.	— H. ♂.		
Lactuca sonchifolia. w.						
LYGODESMIA. D.D. LYGOODESMIA.	Syngenesia Polygamia Æqualis.					
junccea. D.D.	(ro.) rush-like.	Missouri.	1827.	5. 6. H. ♀.		
Prenanthes junccea. Ph.						
ATALANTHUS. D.D. ATALA'NTHUS.	Syngenesia Polygamia Æqualis.					
1 pinnatus. D.D.	(ye.) pinnated.	Canaries.	1823.	6. 7. F. h.		
Prenanthes pinnata. w.						
2 arboreus.	(ye.) arborescent.	—	—	6. 8. F. h.		
Prenanthes arborea. W. énum.						
3 spinosus. D.D.	(ye.) prickly.	Barbary.	1640.	3. 5. F. h.	Park. theatr. 804. f. 7.	
		T 3				

COMPOSITÆ.

SO'NCHUS. W.	SOW THISTLE.	Syngenesia Polygamia Æ qualis, L.				
1 squarrösus. DC.	(ye.) shrubby.	Madeira.	1777.	4. 7.	G. $\ddot{\text{z}}$.	Jacq. ic. 1. t. 161.
fruticösus. L. non Jacq.						
2 Jacquinii. NC.	(ye.) Jacquin's.	—	—	—	G. $\ddot{\text{z}}$.	
fruticösus. Jacq. non L.						
3 lœvigatus. L.en.	(ye.) smooth.	—	1816.	—	G. $\ddot{\text{z}}$.	
4 lyratus. L.en.	(ye.) lyre-leaved.	—	—	—	G. $\ddot{\text{z}}$.	
5 hyoserifolius. H.H.	(ye.) Hyoseris-leaved.	—	1818.	—	G. $\ddot{\text{z}}$.	
6 radicatis. s.s.	(ye.) long-rooted.	Canaries.	1780.	7. 8.	G. $\ddot{\text{z}}$.	
7 gummifer. s.s.	(ye.) gummy.	—	1816.	—	G. $\ddot{\text{z}}$.	
8 divaricatus. L.en.	(ye.) spiny-leaved.	—	1828.	4. 6.	G. $\ddot{\text{z}}$.	
9 pinnatus. s.s.	(ye.) wing-leaved.	Madeira.	1777.	4. 7.	G. $\ddot{\text{z}}$.	All. ped. 1. t. 16. f. 2.
10 maritimus. s.s.	(ye.) sea-side.	Canaries.	1748.	7. 9.	H. $\ddot{\text{u}}$.	Eng. bot. v. 12. t. 843.
11 oleraceus. s.s.	(ye.) common-smooth.	Britain.	6. 12.	H. \odot .	
<i>l'ævis.</i> L.enum.	<i>oleraceus a. lavis.</i> W.enum.					
12 lacerus. W.en.	(ye.) torn-leaved.	Britain.	—	H. \odot .	
13 ásper. s.s.	(ye.) rough-leaved.	—	—	H. \odot .	Blackw. ic. t. 30.
<i>oleraceus γ. asper.</i> L.						
14 uliginosus. s.s.	(ye.) marsh.	Caucasus.	1823.	7. 9.	H. $\ddot{\text{u}}$.	
15 palustris. s.s.	(ye.) tall marsh.	Britain.	—	H. $\ddot{\text{u}}$.	Eng. bot. v. 13. t. 935.
16 arvensis. s.s.	(ye.) corn-field.	—	—	H. $\ddot{\text{u}}$.	— v. 10. t. 674.
17 hantoniensis.	(ye.) smooth-headed.	Hampshire.	—	H. $\ddot{\text{u}}$.	
18 tenerrimus. s.s.	(ye.) clammy.	S. Europe.	1691.	7. 8.	H. \odot .	Flor. græc. t. 790.
19 chondrilloides. s.s.	(ye.) Chondrilla-like.	Sicily.	1826.	—	H. $\ddot{\text{u}}$.	
20 caucasicus. F.	(ye.) Caucasean.	Caucasus.	1820.	—	H. $\ddot{\text{u}}$.	
21 flexuosus. Led.	(ye.) flexuose.	—	1822.	—	H. $\ddot{\text{u}}$.	
22 chinensis. F.	(ye.) Chinese.	China.	1820.	—	H. $\ddot{\text{u}}$.	
23 tenuax. F.	(ye.) tough.	Siberia.	1824.	—	H. $\ddot{\text{u}}$.	
24 longifolius. Trev.	(ye.) long-leaved.	—	—	—	H. $\ddot{\text{u}}$.	
BARKHA'USIA. DC.	BARKHA'USIA.	Syngenesia Polygamia Æ qualis.				
1 muricata. s.s.	(ye.) bristly-headed.	S. Europe.	1825.	6. 8.	H. $\ddot{\text{u}}$.	
2 bellidifolia. DC.	(ye.) Daisy leaved.	Corsica.	1826.	—	H. \odot .	
3 Suffreniana. DC.	(ye.) mealy-headed.	France.	1825.	—	H. \odot .	
4 spathulata. s.s.	(ye.) spathulate.	Sicily.	1829.	—	H. \odot .	
5 alpina. s.s.	(ye.) alpine.	Italy.	1739.	7. 8.	H. \odot .	Gmel. sib. 2. t. 5.
<i>Crèpis alpina.</i> W.						
6 apargioïdes. s.s.	(ye.) Apargia-like.	Germany.	1826.	—	H. $\ddot{\text{u}}$.	Jacq. aust. 3. t. 293.
7 vesicaria. s.s.	(ye.) bladdered.	Crete.	1828.	—	H. $\ddot{\text{u}}$.	
8 hyemalis. s.s.	(ye.) winter.	Sicily.	1829.	8. 11.	H. $\ddot{\text{u}}$.	
9 purpurea. s.s.	(pu.) purple.	—	1827.	7. 9.	H. $\ddot{\text{u}}$.	
10 rübra. s.s.	(li.) lilac.	Italy.	1632.	6. 8.	H. \odot .	Flor. græc. t. 801.
11 nicaensis. s.s.	(ye.) gland-headed.	S. Europe.	1824.	—	H. \odot .	
12 hispida. s.s.	(ye.) hispid.	—	1798.	—	H. \odot .	
13 setosa. DC.	(ye.) setose.	Switzerland.	1824.	—	H. \odot .	W. et K. hung. 1. t. 43.
14 pungens. P.S.	(ye.) pungent.	—	1826.	—	H. \odot .	
MYO'SERIS. L.en.	MYO'SERIS.	Syngenesia Polygamia Æ qualis.				
purpurea. L.en. (pu.) purple.	Tauria.	1823.	6. 7.	H. \odot .		
<i>Lugosericis taurica.</i> M.B. <i>Crèpis purpurea.</i> M.B.						

Tribus V. SCORZONERÆ. D. Don in Edinb. phil. journ.

SCORZON'ERA. W.	VIPER'S-GRASS.	Syngenesia Polygamia Æ qualis.				
1 tomentosa. w.	(ye.) white.	Armenia.	1789.	6. 7.	H. $\ddot{\text{u}}$.	
2 villosa. s.s.	(ye.) villous.	Levant.	1821.	6. 8.	H. $\ddot{\text{u}}$.	Scop. carn. t. 46.
3 hirsuta. s.s.	(ye.) hairy.	S. Europe.	1823.	—	H. $\ddot{\text{u}}$.	All. ped. t. 31. f. 33.
4 humili. w.	(ye.) dwarf.	Europe.	1597.	8.	H. $\ddot{\text{u}}$.	
5 austriaca. W.en.	(ye.) Austrian.	Austria.	—	H. $\ddot{\text{u}}$.	Jacq. aust. 1. t. 36.
humilis. Jacq. non w.						
6 plantaginifolia. Sch.	(ye.) Plantain-leav'd.	Switzerland.	1816.	—	H. $\ddot{\text{u}}$.	
7 hispanica. w.	(ye.) garden.	Spain.	1576.	6. 9.	H. $\ddot{\text{u}}$.	Lam. ill. t. 647. f. 5.
8 glastifolia. w.	(ye.) Woad-leaved.	Germany.	1816.	—	H. $\ddot{\text{u}}$.	
9 taurica. M.B.	(ye.) Taurian.	Tauria.	1820.	—	H. $\ddot{\text{u}}$.	
10 rumicifolia. L.en.	(ye.) Döck-leaved.	—	1822.	—	H. $\ddot{\text{u}}$.	
11 caricifolia. w.	(ye.) Carex-leaved.	Siberia.	1805.	—	H. $\ddot{\text{u}}$.	Pall. it. 3. ap. t. J. 1.f.1.
12 parviflora. w.	(ye.) small-flowered.	Austria.	1816.	—	H. $\ddot{\text{u}}$.	Jacq. aust. 4. t. 305.
13 mollis. M.B.	(ye.) soft.	Volga.	1829.	—	H. $\ddot{\text{u}}$.	
14 stricta. s.s.	(ye.) upright.	—	1821.	—	H. $\ddot{\text{u}}$.	
villosa. M.B. nec Scop.						

15 lanâta. M.B.	(ye.) woolly.	Iberia.	1823.	6. 7. H. ȝ.	
16 purpurea. w.	(pu.) purple-flower'd. Austria.	1759.	5. 6. H. ȝ.	Jacq. aust. 1. t. 35.	
17 angustifolia. w.	(ye.) narrow-leaved. S. Europe.	—	6. 8. H. ȝ.	W. et K. hung. 2. t. 122.	
18 ensifolia. M.B.	(ye.) sword-leaved.	Caucasus.	1823.	— H. ȝ.	
19 pusilla. s.s.	(st.) small.	Volga.	1827.	— H. ȝ.	Pall. it. 2. ap. t. L.
20 tuberosa. s.s.	(ye.) tuberous-rooted.	—	1825.	— H. ȝ.	Pall. it. 3. ap. t. Y. f. 3.
21 crispa. s.s.	(ye.) curled.	Tauria.	1821.	— H. ȝ.	
PODOSPERMUM. DC. PODOSPERMUM. Syngenesia Polygamia Æqualis.					
1 calcitrapifolium. DC.	(ye.) Calcitrapa-leav'd. Levant.	1823.	6. 8. H. ȝ.	Barr. ic. 800.	
2 resedifolium. DC.	(ye.) spreading.	Spain.	1729.	6. 7. H. ♂.	Bocc. sic. 13. t. 7.
3 laciniatum. DC.	(ye.) cut-leaved.	S. Europe.	1640.	— H. ♂.	Jacq. aust. 4. t. 356.
4 octangulare. DC.	(ye.) 8-angled.	—	1820.	— H. ♂.	
5 pumilum.	(ye.) small.	Spain.	1822.	7. 9. H. ♂.	Cav. ic. 2. t. 121. f. 2.
<i>Scorzonera pumila. w.</i>					
6 asperimum.	(ye.) roughest.	Galatia.	—	8. 9. H. ȝ.	
<i>Scorzonera asperrima. w.</i>					
7 áspérum.	(ye.) rough.	Levant.	—	7. 8. H. ȝ.	Desf. an. mus. 1. t. 9.
<i>Scorzonera áspéra. Desf. stylös. P.S.</i>					
8 subulatum. DC.	(ye.) grass-leaved.	Portugal.	1759.	— H. ȝ.	Jacq. obs. 4. t. 100.
<i>Scorzonera graminifolia. F.G. 784.</i>					
9 muricatum. DC.	(ye.) bristly.	Piedmont.	1818.	— H. ȝ.	
10 taraxacifolium.	(ye.) Dandelion-leav'd. Bohemia.	1801.	6. 8. H. ȝ.	Jacq. ic. 1. t. 106.	
<i>Scorzonera taraxacifolia. w.</i>					
LASIOSPERMUM. F. LASIOSPERMUM. Syngenesia Polygamia Æqualis.					
angustifolium. F.	(ye.) narrow-leaved.	Caucasus.	1825.	6. 8. H. ȝ.	
<i>Scorzonera eriosperma. M.B.</i>					
PICRIDIUM. P.S. PICRIDIUM. Syngenesia Polygamia Æqualis.					
1 vulgare. P.S.	(ye.) various-leaved.	France.	1773.	6. 8. H. ♂.	All. ped. 1. t. 16. f. 1.
<i>Scorzonera picroides. L. Sónchus picroides. w.</i>					
2 tingitanum. P.S.	(ye.) Tangier.	Barbary.	1713.	6. 9. H. ♂.	Bot. mag. t. 142.
<i>Scorzonera tingitana. B.M. Sónchus tingitanus. w.</i>					
3 hispánicum. V.	(ye.) Spanish.	Spain.	1816.	— H. ♂.	Jacq. schœnb. 2. t. 143.
4 albídum. P.S.	(ye.) pale-flowered.	France.	1781.	7. 10. H. ȝ.	Jacq. ic. 1. t. 164.
<i>Crépis albida. w.</i>					
ARNOPOGON. W. SHEEP'S-BEARD. Syngenesia Polygamia Æqualis.					
1 Dalechampii. w.	(ye.) great-flowered.	S. Europe.	1739.	6. 10. H. ȝ.	Bot. mag. t. 1623.
2 picroides. w.	(ye.) prickly-cupp'd.	—	1683.	7. 8. H. ♂.	Flor. græc. t. 781.
3 áspér. w.	(ye.) rough.	Montpellier.	1774.	— H. ♂.	— t. 782.
4 capense. w.	(ye.) Cape.	C. B. S.	1824.	6. 8. G. ♂.	Jacq. ic. 3. t. 577.
TRAGOPOGON. W. GOAT'S-BEARD. Syngenesia Polygamia Æqualis.					
1 pratensis. w.	(ye.) yellow.	Britain.	5. 6. H. ♂.	Eng. bot. v. 7. t. 434.
2 mutabilis. w.	(va.) changeable.	Siberia.	1816.	— H. ♂.	Jacq. ic. 1. t. 157.
3 undulatus. w.	(st.) wave-leaved.	Crimea.	1790.	— H. ♂.	— t. 158.
4 parviflorus. L.en.	(ye.) small-flowered.	—	1822.	— H. ♂.	
5 orientalis. w.	(ye.) oriental.	Levant.	1787.	6. 7. H. ♂.	
6 canus. w.k.	(ye.) hoary.	Hungary.	1822.	— H. ♂.	
7 major. w.	(ye.) great.	Austria.	1788.	5. 6. H. ♂.	Jacq. aust. 1. t. 29.
8 campéstris. s.s.	(ye.) field.	Poland.	1828.	— H. ♂.	
9 rósea. s.s.	(ro.) red-flowered.	Volga.	1827.	— H. ♂.	
10 porrifolius. w.	(pu.) purple.	England.	— H. ♂.	Eng. bot. v. 9. p. 638.
11 floccosus. w.k.	(ye.) woolly.	Hungary.	1816.	— H. ♂.	W. et K. hung. 2. t. 112.
12 dûbius. L.en.	(ye.) doubtful.	—	1822.	— H. ♂.	Flor. græc. t. 779.
13 crocifolius. w.	(vi.) Crocus-leaved.	Italy.	1739.	6. 7. H. ♂.	
14 angustifolius. w.	(vi.) narrow-leaved.	—	1816.	— H. ♂.	
15 villósus. w.	(ye.) hairy.	Spain.	1794.	5. 6. H. ♂.	
16 rhodánthus.	(ro.) rose-coloured.	Hungary.	1807.	7. H. ȝ.	W. et K. hung. 2. t. 121.
<i>Scorzonera rósea. w.k.</i>					
17 húmilia. F. dwarf.	Caucasus.	1829. H. ♂.	
18 phæopáppus. F. glossy-chaff'd.	Persia.	— H. ♂.	
GEROPOGON. W. OLD MAN'S-BEARD. Syngenesia Polygamia Æqualis.					
1 gláber. w.	(pi.) smooth.	Italy.	1704.	7. 8. H. ♂.	Bot. mag. t. 479.
2 hirsútus. w.	(pi.) hirsute.	—	1759.	— H. ♂.	Col. ecphr. 1. t. 231.
3 calyculatus. w.	(pi.) perennial.	—	1774.	6. 8. H. ȝ.	Jacq. vind. 2. t. 106.

Tribus VI. CICHOREÆ. D. Don in Edinb. phil. journ.

CICHORIUM. S.S.	SUCCORY. Syngenesia Polygamia Æqualis. L.
1 I'ntybus. s.s.	(bl.) common.

2	divaricatum. s.s. (bl.)	branching.	Barbary.	1798.	7. 9. H. ♂.		
3	spinosum. s.s.	(bl.) prickly.	Candia.	1633.	7. 8. F. ♀.	Flor. græc. t. 823.	
4	pūmilum. s.s.	(bl.) dwarf.	Levant.	1790.	— H. ♂.		t. 821.
5	Endivia. s.s.	(bl.) Endive.	E. Indies.	1548.	— H. ♂.		
	a latifolia.	(bl.) broad-leaved.		—	— H. ♂.		
	β crīspa.	(bl.) curled-leaved.		—	— H. ♂.		
	γ leucophylla.	(bl.) white-leaved.		—	— H. ♂.		
<hr/>							
SCOLYMYUS. S.S.							
	GOLDEN THISTLE. Syngenesia Polygamia Æqualis.						
1	maculatus. s.s.	(ye.) annual.	S. Europe.	1633.	7. 8. H. ♂.	Flor. græc. t. 824.	
2	hispanicus. s.s.	(ye.) Spanish.		1658.	7. 9. H. ♀.		t. 825.
3	grandiflorus. s.s.	(ye.) great-flowered.	Barbary.	1820.	— H. ♀.	Desf. atl. 2. t. 218.	
CATANA'NCHE. S.S.							
	CATANA'NCHE. Syngenesia Polygamia Æqualis. L.						
1	lutea. s.s.	(ye.) yellow.	Candia.	1640.	6. 7. H. ♂.	Flor. græc. t. 821.	
2	cœrulea. s.s.	(bl.) blue-flowered.	S. Europe.	1596.	7. 10. H. ♀.	Bot. mag. t. 293.	

SUBORDO II. LABIATIFLORÆ. *Lagasca et DC. in ann. mus.*

L'ERIA. DC.							
1	nūtans. DC.	(wh.) drooping-flower'd. W. Indies.	1793.	6. 7. S. ♂.	Plum. ic. t. 41. f. 1.		
	Tussilago nūtans. w.						
2	albicans. DC.	(wh.) woolly-leaved.	Jamaica.	1821.	— S. ♂.		
CHAP'TALIA. V.							
	CHAP'TALIA. Syngenesia Polygamia Necessaria.						
	tomentosa. v.	(bh.) woolly.	N. America.	1806.	5. 6. H. ♀.	Bot. mag. t. 2257.	
	Tussilago integrifolia. M.						
ONO'SERIS. DC.							
	ONO'SERIS. Syngenesia Polygamia Æqualis.						
1	pūpurea. k.s.	(pu.) purple.	S. America.	1827.	7. 9. F. ♀.	Smith. ic. ined. t. 65.	
	Atrāctylis purpurat̄a. Smith.						
2	mexicana. k.s.	(wh.) Mexican.	Mexico.	1828.	— F. ♂.	Smith. ic. ined. t. 66.	
CHÆTANTH'ERA. R.P. CHÆTANTH'ERA.							
	Syngenesia Polygamia Superflua.						
	ciliata. r.p.	(ye.) ciliated.	Chile.	1822.	7. 8. F. ♀.		
MUTI'SIA. L.							
	MUTI'SIA. Syngenesia Polygamia Æqualis. L.						
	speciosa. b.m.	(pu.) handsome.	Brazil.	1825.	9. 11. G. ♂. —	Bot. mag. t. 2705.	
	arachnoidea. Martius.						
CHUQUIR'AGA. J.							
	CHUQUIR'AGA. Syngenesia Polygamia Æqualis.						
	insignis. k.s.	(ye.) large-flowered.	S. America.	1824.	— G. ♂.	Lam. ill. t. 691.	
	Johannia insignis. w.						
DUMERILIA. DC.							
	DUMERILIA. Syngenesia Polygamia Æqualis.						
	paniculata. dc.	(pu.) panicled.	Chile.	1825.	7. 9. H. ♀.	DC. ann. mus. 19. t. 7.	
BAC'AZIA. R.P.							
	BAC'AZIA. Syngenesia Polygamia Æqualis.						
	spinosa. r.p.	(pi.) spiny.	Peru.	1825.	4. 5. F. ♂.		
TRIPT'LION. R.P.							
	TRIPT'LION. Syngenesia Polygamia Æqualis.						
	cordifolium, b.r.(wh.)	heart-leaved.	Peru.	1823.	6. 8. H. ♂.	Bot. reg. t. 853.	
LEUC'ERIA. Lag.							
	LEUC'ERIA. Syngenesia Polygamia Superflua.						
	senecioïdes. d.d. (wh.)	Ragwort-like.	Chile.	1822.	8. 10. F. ♀.	Hook. ex. flor. t. 101.	
	Trixis senecioïdes. Hooker.						
GASTROCA'RPHA. D.D. GASTROCA'RPHA.							
	Syngenesia Polygamia Æqualis.						
	runcinata. d.d.	(wh.) Turnep-leaved.	Chile.	1826.	8. 10. F. ♂.	Swt. br. fl. gar. t. 229.	
PERDICIUM. DC.							
	PERDICIUM. Syngenesia Polygamia Superflua.						
1	Anándria. dc.	(wh.) Siberian.	Siberia.	1759.	8. H. ♀.	Gmel. sib. 2. t. 68. f. 1.	
2	lyrātum.	(wh.) lyrate.		1825.	6. 8. H. ♀.		t. 67. f. 2.
	Tussilago lyrātā. w. Chaptalia lyrātā. s.s.						
TRIXIS. DC.							
	TRIXIS. Syngenesia Polygamia Superflua.						
1	frutēscens. s.s.	(wh.) frutescent.	Jamaica.	1826.	2. 4. S. ♂.	Lam. ill. t. 677. f. 2.	
	Perdicum radiale. L.						
2	divaricata. d.d.	(wh.) divaricate.	Brazil.	1824.	4. 6. S. ♂.	Bot. mag. t. 2765.	
	auriculata. b.m.	Perdicum brasiliense. Gr.					

SUBORDO III. CARDUACEÆ. *Don prodr. p. 166.*Div. I. CARDUACEÆ VERÆ. *Kth. synops. 2. p. 363.*

A'RCTIUM. L.	BURDOCK. Syngenesia Polygamia Æqualis.		
1 Lappa. w.	(pu.) smooth-headed. Britain.	7. 8. H. ♂.
			Eng. bot. v. 18. t. 1228.

2	mimus. M.B.	(pu.) smaller.	Europe.	7. 8. H. ♂.	
	Láppa minor. DC.					Eng. bot. v. 35. t. 2478.
3	Bardana. w.	(pu.) woolly-headed.	Britain.	— H. ♂.	
	tomentosum. P.S.					Bot. mag. t. 2589.
SERRA'TULA. D.P.	SAWWORT.	Syngenesia Polygamia Æqualis.				Linn. trans. v. 11. t. 38.
1	pulchella. B.M.	(pu.) purple-scaled.	Siberia.	1823.	6. 7. H. 4.	—
2	depréssâ. L.T.	(pu.wh.) depressed.	Caucasus.	1820.	— H. 4.	—
3	élégans. L.T.	(ro.) elegant.		—	— H. 4.	v. 11. t. 37.
4	alpina. E.B.	(pu.) Alpine.	Britain.	7. 8. H. 4.	Eng. bot. v. 9. t. 599.
	Saussurea alpina. DC.					
5	angustifolia. w.	(pu.) narrow-leaved.	Siberia.	1816.	— H. 4.	Gmel. sib. 2. t. 33.
6	discolor. w.	(pu.wh.) two-coloured.	Europe.	—	— H. 4.	
	Saussurea discolor. DC.					Moris. s. 7. t. 22. f. 1.
7	serrata.	(pu.) saw-leaved.		—	— H. 4.	
	Saussurea serrata. DC.					
8	salicifolia. w.	(re.) Willow-leaved.	Siberia.	1796.	— H. 4.	Gmel. sib. 2. t. 27.
9	alâta.	(pu.) winged.		1823.	— H. 4.	
	Saussurea alata. DC. non Serrátula alata. W.en.					
10	sálsa. M.B.	(pu.) salt.	Caucasus.	1824.	— H. 4.	
	Heterotrichum sálsum. M.B.	Saussurea salsa. s.s.				
11	lyrâta. F.	(pu.) lyrate-leaved.	Dahuria.	1827.	— H. 4.	
12	Gmelini. Led.	(pu.) Gmelin's.	Siberia.	—	— H. 4.	
13	elongâta. DC.	(pu.) elongated.		1820.	— H. 4.	
14	nitidula. Led.	(pu.) glossy.		1827.	— H. 4.	
15	pycnocéphala. Led.	(pu.) large-headed.		—	— H. 4.	
16	rigida. Led.	(pu.) rigid.		—	— H. 4.	
17	pygmæa. Jac.	(pu.) pigmy.	Austria.	1816.	— H. 4.	Jacq. aust. t. 440.
SYNCA'RPHA. DC.	SYNCA'RPHA.	Syngenesia Polygamia Æqualis.				
	gnaphaloides. DC.	(or.) squarrose.	C. B. S.	1815.	7. 9. G. ♀.	Pluk. alm. t. 302. f. 3.
	Stæhelina gnaphaloides. L.	Leysèra squarròsa. Th.				
CARDUUS. D.P.	THISTLE.	Syngenesia Polygamia Æqualis.				
1	alatus. B.F.G.	(pu.) winged-stalked.		1812.	6. 8. H. ♂.	Swt. br. fl. gar. t. 103.
2	tinctorius. D.P.	(pu.) common.	Britain.	7. 10. H. 4.	Eng. bot. v. 1. t. 38.
	Serrátula tintoriæ. E.B.					
3	quinquefolius. D.P.	(pu.) five-leaved.	Persia.	1804.	7. 8. H. 4.	Bot. mag. t. 1871.
	Serrátula quinquefolia. B.M.					
4	coronatus. D.P.	(pu.) lyre-leaved.	Italy.	1739.	— H. 4.	Gmel. sib. 2. t. 20.
	Serrátula coronata. w.					
5	nítidus.	(pu.) glossy.	Siberia.	1821.	— H. 4.	
	Serrátula nítida. F.					
6	multiflorus. L.	(pu.) Volga.		—	— H. 4.	Gmel. sib. 2. t. 28.
7	linearifolius. DC.	(pu.) linear-leaved.		1827.	— H. 4.	
8	Pícris.	(pu.) scariose.	Caucasus.	1822.	— H. 4.	
	Centaurea Picris. w.	Serrátula Picris. s.s.				
9	centauroïdes. D.P.	(pu.) Centaury-like.	Siberia.	1804.	— H. 4.	Gmel. sib. 2. t. 40.
10	radiatus. W.K.	(pu.) rayed.	Hungary.	1800.	— H. ♂.	W. et K. hung. 1. t. 11.
11	polyclónos. w.	(pu.) Siberian.	Siberia.	1823.	— H. 4.	
12	argútus.	(pu.) sharply-tooth'd.		—	— H. 4.	
	Serrátula argùta. L.en.					
13	cyanoides. w.	(pu.) blue-bottle-leav'd.		1778.	— H. 4.	Gmel. sib. 2. t. 15.
14	simplex.	(pu.) one-flowered.	Caucasus.	1817.	— H. 4.	Bot. mag. t. 2482.
	móllis. w.	Serrátula simplex. DC.				
15	xeranthemoides.	(pu.) everlasting-fl'd.		1804.	— H. 4.	Gmel. sib. 2. t. 47. f. 1.
	Serrátula xeranthemoides. M.B.	Centaurea radiata. w.				
16	lycopiifolius. w.	(pu.) horehound-leaved.	France.	1823.	— H. 4.	Vill. delph. 3. t. 19.
17	heterophyllum. s.s.	(pu.) various-leaved.	France.	1824.	— H. 4.	
18	stæchadifolius.	(pu.) woolly-headed.	Tauria.	1820.	— H. 4.	
	Serrátula stæchadifolia. s.s.					
19	leucographus. w.	(pu.) white-spotted.	Italy.	1752.	6. 7. H. ⓧ.	Jacq. vind. 3. t. 23.
20	árabicus. w.	(pu.) Arabian.	Arabia.	1789.	7. 8. H. ⓧ.	Jacq. ic. 1. t. 166.
	nútans. w.	(pu.) musk.	Britain.	— H. ⓧ.	Eng. bot. v. 16. t. 1112.
22	carlinoides. W.en.	(pu.) Pyrenean.	Pyrenees.	1784.	— H. ♂.	Gouan. ill. t. 23.
	Carlina pyrenæica. L.					
23	carlinæfolius. w.	(pu.) Carline-leaved.		1804.	— H. ♂.	Eng. bot. v. 14. t. 973.
24	acanthoïdes. w.	(li.) welted.	Britain.	6. 7. H. ⓧ.	
25	hamulösus. w.	(pu.) hooked.	Hungary.	1802.	— H. ⓧ.	Eng. bot. v. 6. t. 412.
26	tenuiflorus. E.B.	(li.) slender-flower'd.	Britain.	— H. ⓧ.	Flor. dan. t. 621.
27	crispus. w.	(li.) curled.	Europe.	1804.	7. 8. H. ♂.	

28	cándicans. w.	(pu.) hoary.	Hungary.	1805.	7. 8. H. ♂.	W. et K. hung. 1. t. 83.
29	personátus. w.	(pu.) cut-leaved.	Austria.	1776.	— H. ♂.	Jacq. aust. 4. t. 348.
30	polyánthemus. w.	(pu.) many-flowered.	Rome.	1739.	6. 7. H. ♂.	Triunf. obs. t. 103.
31	paniculátus. w.	(pu.) panicled.	S. Europe.	1781.	— H. ♀.	
32	pycnocéphalus. w.	(pu.) Italian.		1739.	7. 9. H. ♀.	Jacq. vind. 1. t. 24.
33	onopordioídes. M.B.	(pu.) Onopordum-like.	Iberia.	1824.	7. 8. H. ♀.	Jacq. vind. 2. t. 192.
34	argentátus. w.	(pu.) white Egyptian.	Egypt.	1789.	6. 8. H. ♂.	
35	arctioídes. w.	(pu.) pinnated.	Hungary.	1804.	7. 8. H. ♀.	
36	dùbius. s.s.	(pu.) doubtful.		1827.	— H. ♀.	
37	seminudus. s.s.	(pu.) half-naked.	Caucasus.	—	— H. ♀.	
38	uncinátus. s.s.	(pu.) crook-pointed.	Volga.	—	— H. ♀.	
39	alpéstris. w.	(pu.) alpine.	Hungary.	1805.	7. 8. H. ♀.	
40	deflorátus. w.	(pu.) different leav'd.	Austria.	1570.	7. 9. H. ♀.	
41	corymbósus. s.s.	(pu.) corymb-flower'd.	Naples.	1827.	— H. ♀.	
42	cinéreus. s.s.	(pu.) gray.	Caucasus.	—	— H. ♀.	
43	collinus. s.s.	(pu.) hill.	Hungary.	1828.	— H. ♀.	
44	Argemón. DC.	(pu.) Argemone-leav'd.	Pyrenees.	1816.	— H. ♀.	
45	parviflórus. w.	(pu.) small-flowered.	S. Europe.	1781.	6. 7. H. ♀.	
46	crassifólius. W.en.	(pu.) thick-leaved.		1805.	7. H. ♀.	
47	sumánus. L.en.	(pu.) Pollin's.	Italy.	1823.	7. 8. H. ♀.	
48	nítidus. w.	(pu.) glossy.	Hungary.	1806.	— H. ♀.	
49	nudicáulis. P.S.	(pu.) naked-stalked.	S. Europe.	1739.	— H. ♀.	
	cerinthoídes. w. gláucus.	Cav. <i>Centaurea nudicaulis</i> .	L. <i>Serrátula nudicaulis</i> . DC.			W. et K. hung. 1. t. 52.
50	clavulátus. s.s.	(pu.) clubbed.	Canaries.	1827.	— F. ♂.	Cav. ic. 3. t. 226.
51	álbidus. s.s.	(pu.) white-leaved.	Caucasus.	1829.	— H. ♀.	
52	argyróa. s.s.	(pu.) silvery.	Sicily.	1827.	— H. ♀.	
53	orientális. M.B.	(pu.) oriental.	Iberia.	1804.	7. H. ♀.	
54	salinus. M.B.	(pu.) narrow-leaved.	Siberia.	—	— H. ♀.	
55	völgénis. s.s.	(pu.) Volga.	Volga.	—	— H. ♀.	
56	leucánthus. s.s. (wh.)	white-flowered.	Spain.	1825.	— H. ♂.	
57	atriplicifólius. s.s. (pu.)	Orach-leaved.	Russia.	1823.	— H. ♀.	
SI'LYBUM. G.		SI'LYBUM. Syngenesia Polygamia Æqualis.				Cavan. icon. 2. t. 165.
1	mariánum. g.	(pu.) milk-thistle.	Britain.	7. H. ♂.	Eng. bot. v. 14. t. 976.
	Cárduuus mariánum. E.B.					Mill. ic. 2. t. 248.
2	cérnum. g.	(wh.) nodding.	Siberia.	1755.	6. 7. H. ♀.	
	Cnicus cérnus. w.					
HALOCHA'RIS. F.	HALOCHA'RIS.	Syngenesia Polygamia Æqualis.				
1	altáica. f.	(pu.) Altay.	Altay.	1827.	7. 8. H. ♀.	
2	carthamoídes. f.	(pu.) Carthamus-like.	Siberia.	1823.	— H. ♀.	
	Cnicus carthamoídes. w.	<i>Leuzea? carthamoídes</i> . DC.	<i>Cnicus salinus</i> . M.B.			
CIR'SIUM. G.		HORSE-THISTLE. Syngenesia Polygamia Æqualis.				
1	palústre. DC.	(li.wh.) marsh.	Britain.	7. 8. H. ♂.	Eng. bot. v. 14. t. 974.
	Cárduuus palústris. E.B.	<i>Cnicus palústris</i> . w.				
2	elátum. L.en.	(pu.) tall.		1823.	— H. ♀.	
3	uliginòsum. M.B.	(pu.) swamp.	Caucasus.	1820.	— H. ♀.	
4	cánum. M.B.	(pu.) hoary.	Austria.	1633.	— H. ♀.	
5	Acárna. DC.	(pu.) yellow.	Spain.	1683.	7. 9. H. ♂.	
6	pyrenáicum. DC.	(pu.) Pyrenean.	Pyrenees.	1818.	7. 8. H. ♀.	
7	monspessulánum. DC.	(pu.) Montpelier.	Montpelier.	1596.	6. 7. H. ♀.	
8	pannónicum. L.en.	(pu.) Hungarian.	Hungary.	1816.	7. 9. H. ♀.	
9	stríctum. s.s.	(pu.) straight.	Naples.	1820.	— H. ♀.	
10	lanceolátum. DC.	(pu.) common.	Britain.	6. 9. H. ♀.	
	Cárduuus lanceolátus. E.B.	<i>Cnicus lanceolatus</i> . w.				
11	desertórum. L.en.	(pu.) desert.	Siberia.	1823.	7. 8. H. ♀.	
12	cichoráceum. s.s.	(pu.) Succory-like.	Naples.	1824.	— H. ♀.	
13	ítalicum. s.s.	(pu.) Italian.	Italy.	1826.	— H. ♀.	
14	sículum. s.s.	(pu.) Sicily.	Sicily.	1828.	— H. ♀.	
15	neglécum. s.s.	(pu.) neglected.	Siberia.	1829.	— H. ♀.	
16	pungens. s.s.	(pu.) pungent.	Levant.	—	— H. ♀.	
17	paniculátum. s.s.	(pu.) panicled.	Pyrenees.	1827.	— H. ♀.	
18	créticum. s.s.	(wh.) Cretan.	Crete.	—	— H. ♀.	
19	ciliátum. L.en.	(pu.) fringed.	Siberia.	1787.	8. H. ♀.	
20	serrolátum. M.B.	(pu.) serrulate.	Tauria.	1820.	7. 8. H. ♀.	
21	laníflórum. M.B.	(pu.) woolly-flowered.		1823.	— H. ♀.	
22	arachnoídeum. M.B.	(pu.) cobweb.		—	— H. ♀.	
23	strigósium. M.B.	(pu.) strigose.	Caucasus.	1818.	— H. ♂.	
24	hórridum. M.B.	(pu.) very spiny.		1820.	— H. ♀.	
25	múnitus. M.B.	(pu.) armed.		1816.	— H. ♀.	
26	obvallátum. M.B.	(pu.) crowded-flower'd.		—	— H. ♀.	

Murr. com. got. 6. t. 5.

Jacq. aust. 1. t. 42-43.

Cavan. ic. 1. t. 53.

Jacq. obs. 4. t. 95.

Jacq. aust. 3. t. 5.

Eng. bot. v. 2. t. 107.

27 scleránthon. M.B. (wh.) hard-flowered. Caucasus.	1824.	7. 8. H. ♂.	
carlinoides. F. <i>Carlina Echinus</i> . M.B.		— H. ♀.	
28 echinátum. s.s. (pu.) hedgehog.	Barbary.	— H. ♂.	
29 eriophorum. DC. (pu.) woolly-headed.	Britain.	— H. ♂.	Eng. bot. v. 6. t. 386.
5 <i>Cárdus eriophorus</i> . E.B. <i>Cnicus eriophorus</i> . w.			
30 ferox. DC. (wh.) fierce.	S. Europe.	1683.	— H. ♂.
31 discolor. s.s. (pu.) two-coloured.	N. America.	1803.	— H. ♂.
32 inérme. L.en. (pu.) unarmed.	Pyrenees.	1818.	— H. ♀.
33 cynaroídes. s.s. (pu.) Artichoke-like.	Crete.	1828.	— H. ♀.
34 altíssimum. s.s. (pu.) giant.	N. America.	1726.	8. 9. H. ♀.
35 prátense. (pu.) meadow.	Britain.	6. 7. H. ♀.
ánglicum. DC. <i>Cárdus pratensis</i> . E.B. <i>Cnicus pratensis</i> . w.			Dill. elth. t. 69. f. 80
36 heterophyl'um. DC. (pu.) melancholy.		— v. 10. t. 675.
37 helenioídes. L.en. (pu.) Elecampane-ld.	Siberia.	1804.	— H. ♀.
38 serratuloides. s.s. (pu.) sawwort-like.		1752.	6. 10. H. ♀.
39 heteromálum. (pu.) Nepaul.	Nepaul.	1820.	— H. ♂.
<i>Cnicus heteromálus</i> . D.P. <i>Cárdus cándicans</i> . Wal.			Jacq. aust. 2. t. 127.
40 arvénse. DC. (li.) corn or way.	Britain.	Eng. bot. v. 14. t. 975.
<i>Serrátula arvénsis</i> . w. <i>Cárdus arvénsis</i> . E.B. <i>Cnicus arvénsis</i> . Ph.			
41 rivulare. s.s. (pu.) river.	Hungary.	1804.	7. 8. H. ♀.
tricephalòdes. DC. <i>Cnicus rivularis</i> . w.			Jacq. aust. 1. t. 91.
β salisburgénsē. smaller.			
42 ambíguum. DC. (pu.) ambiguous.	S. Europe.	1823.	— H. ♀.
43 orgyale. s.s. (pu.) tall.		— H. ♀.
44 carníolicum. s.s. (wh.) Carniola.	Carniola.	1792.	— H. ♀.
45 pauciflórum. s.s. (pu.) few-flowered.	Hungary.	1816.	— H. ♂.
46 tatáricum. s.s. (wh.) Tartarian.	Siberia.	1775.	7. 8. H. ♀.
47 rigens. s.s. (wh.) upright alpine.	Switzerland.		— H. ♀.
48 oléraceum. DC. (wh.) pale-flowered.	Europe.	1570.	— H. ♀.
49 Erisíthales. L.en. (st.) clammy.	France.	1752.	6. 8. H. ♀.
glutinòsum. DC. <i>Cnicus Erisíthales</i> . w.			Jacq. aust. 4. t. 310.
50 ochroleúcum. DC. (st.) pale yellow.	Switzerland.	1801.	7. H. ♀.
51 mítæ. s.s. (pu.) soft.	Siberia.	1818.	7. 8. H. ♂.
52 fimbriátum. s.s. (pu.) fringed.	Caucasus.	1829.	— H. ♀.
53 médium. s.s. (pu.) middle.	Italy.		— H. ♀.
54 níveum. s.s. (pu.) snowy-leaved.	Sicily.	1828.	— H. ♀.
55 dealbátum. s.s. (pu.) mealy-leaved.	Caucasus.		— H. ♀.
56 echinocéphalum. s.s. (pu.) spiny-headed.		1823.	— H. ♀.
57 lappáceum. M.B. (pu.) Burdock-like.		1826.	— H. ♀.
58 pinnatifidum. s.s. (pu.) pinnatifid.	Spain.		— H. ♀.
59 bulbósom. DC. (pu.) tuberos.	England.	8. 10. H. ♀.
<i>Cnicus tuberosus</i> . E.B. <i>Cárdus tuberosus</i> . L.			Eng. bot. v. 36. t. 2562.
60 setòsum. M.B. (pu.) setose.	Tauria.	1823.	7. 8. H. ♀.
61 acaule. DC. (pu.) dwarf.	Britain.	— H. ♀.
<i>Cárdus acaulis</i> . E.B. <i>Cnicus acaulis</i> . w.			Eng. bot. v. 3. t. 161.
β caulescens. P.S. (pu.) stalked.		— H. ♀.
62 Casabònæ. DC. (pu.) fish-bone.	S. Europe.	1714.	6. 8. F. ♂.
63stellátum. DC. (pu.) starry.	Italy.	1665.	6. 7. H. ♂.
64 Diacánthon. s.s. (pu.) two-spined.	Barbary.	1800.	— F. ♂.
65 áfrum. s.s. (pu.) African.		1802.	— F. ♂.
66 squarrosum. s.s. (pu.) squarrose.	Siberia.	1829.	— H. ♀.
67 incánum. s.s. (pu.) hoary.	Caucasus.		— H. ♀.
68 syriacum. L.en. (wh.) Syrian.	Levant.	1771.	7. 8. H. ♂.
69 spinosissimum. s.s. (wh.) feathery-headed.	Switzerland.	1759.	6. 8. H. ♀.
70 tatáricum. s.s. (wh.) Tartarian.	Tartary.	1829.	— H. ♀.
71 Bertolónii. s.s. (wh.) Bertoloni's.	Italy.	1828.	— H. ♀.
72 b'aeticum. s.s. (wh.) Baetian.	Spain.	1825.	— H. ♀.
73 semipectinátum. s.s. (w.) semipectinate.	Siberia.	1828.	— H. ♀.
74 Gmelini. F. (pu.) Gmelin's.	Baical.	1829.	— H. ♀.
75 arábicum. F. (pu.) Arabian.	Arabia.		— H. ♂.
76 pygmæum. DC. (pu.) pigmy.	Austria.	1816.	— H. ♀.
<i>Serrátula pygmæa</i> . w.			Jacq. aust. t. 440.
ERYTHRÓLENA. ERYTHRÓLENA. Syngenesia Polygamia Áequalis.			
conspicua. B.F.G. (sc.) conspicuous.	Mexico.	1824.	9. 10. F. ♂.
ONOPÓRDUM. DC. COTTON-THISTLE. Syngenesia Polygamia Áequalis.			
1 Acánthum. E.B. (pu.) woolly.	Britain.	7. 8. H. ♂.
2 taúricum. w. (pu.) Taurian.	Tauria.	1800.	— H. ♂.
3 vires. DC. (pu.) green.	S. Europe.	1822.	— H. ♂.
4 macrocánthum. w. (pu.) long-spined.	Barbary.	1798.	— H. ♂.
			Swt. br. fl. gar. t. 134.
			Eng. bot. v. 14. t. 977.
			Schousb. maroc. t. 5.

COMPOSITÆ.

5 illy'ricum. w.	(pu.) Illyrian.	S. Europe.	1648.	7. 8. H. ♂.	Jacq. vind. 2. t. 148.
6 arábicum. w.	(pu.) Arabian.	—	1686.	7. H. ♂.	— 2. t. 149.
7 elátum. Sm.	(pu.) tall.	Greece.	1818.	7. 8. H. ♂.	
8 deltoídes. w.	(pu.) Siberian.	Siberia.	1784.	8. H. ♀.	
9 gr'acum. w.	(pu.) Grecian.	Greece.	1799.	6. 7. H. ♀.	
10 cynaroídes. L.en.	(pu.) Cynara-like.	Caucasus.	1823.	7. 8. H. ♀.	
<i>Cártamus cynaroídes. M.B.</i>					
11 viscosum. Schr.	(pu.) viscous.	Europe.	1821.	— H. ♂.	
12 pyrenáicum. DC.	(pu.) Pyrenean.	Pyrenees.	1818.	— H. ♀.	
13 acaúlon. w.	(pu.) dwarf.	—	1739.	— H. ♂.	
BERA'DIA. P.S.	BERA'DIA.	Syngenesia Polygamia Æqualis.			
subacaúlis. p.s.	(pu.) round-leaved.	Italy.	1791.	7. 8. H. ♀.	
<i>Arctium lanuginosum. DC. Onopordum rotundifolium. w.</i>					
ACIL'EPIS. D.P.	ACIL'EPIS.	Syngenesia Polygamia Æqualis.			
squarrósa. D.P.	(pu.) squarrose.	Nepaul.	1821.	7. 8. H. ♀.	
CYN'ARA. W.	ARTICHOKE.	Syngenesia Polygamia Æqualis.			
1 Cardúneculus. w.	(bl.) Cardoon.	Candia.	1658.	8. 9. H. ♀.	
2 Scólymus. w.	(bl.) common.	S. Europe.	1548.	— H. ♀.	
3 spinosissima. s.s.	(bl.) very spiny.	Sicily.	1824.	— H. ♀.	
4 hórrida. w.	(bl.) very prickly.	Madeira.	1778.	— G. ♀.	
5 húmiliis. w.	(bl.) dwarf.	Spain.	1613.	7. 8. H. ♀.	
6 cósrica. Viv.	(bl.) Corsican.	Corsica.	1825.	— H. ♀.	
7 py'gmaea. w.	(bl.) small.	Spain.	1824.	— H. ♀.	
CESTR'INUS. Cass.	CESTR'INUS.	Syngenesia Polygamia Æqualis.			
carthamoídes. Cass.	(bl.) Carthamus-like.	Barbary.	1799.	7. H. ♀.	
<i>Cynara acaúlis. w. Serrátula acaúlis. DC.</i>					
CARLINA. W.	CARLINE-THISTLE.	Syngenesia Polygamia Æqualis.			
1 acaúlis. w.	(wh.) dwarf.	Italy.	1640.	6. H. ♀.	
β caule'scens. P.S.	(wh.) stalked.	—	—	— H. ♀.	
2 acanthifólia. w.	(wh.) Acanthus-leav'd.	Pyrenees.	1818.	6. 7. H. ♀.	
3 símplex. P.S.	(wh.) single-flowered.	Hungary.	1816.	— H. ♀.	
4 aggregáta. w.	(wh.) clustered.	—	1804.	6. 9. H. ♀.	
5 lyráta. w.	(wh.) lyre-leaved.	C. B. S.	1816.	— G. ♂.	
6 lanáta. w.	(ro.) woolly.	S. Europe.	1683.	6. 7. H. ♂.	
7 corymbósa. w.	(ye.) corymbed.	—	1640.	7. 8. H. ♀.	
8 vulgáris. w.	(bh.) common.	Britain.	—	5. 9. H. ♂.	
9 racemósa. w.	(ye.) racemed.	Spain.	1658.	6. 8. H. ♂.	
10 sícula. s.s.	(wh.) Sicilian.	Sicily.	1828.	— H. ♀.	
11 Biebersteinii. Ot.	(wh.) Bieberstein's.	Caucasus.	1821.	— H. ♀.	
STOB'EA. W.	STOB'EA.	Syngenesia Polygamia Æqualis.			
1 pinnáta. w.	(ye.) Carthamus-like.	C. B. S.	1812.	1. 12. G. ♂.	
2 glomeráta. s.s.	(ye.) clustered.	—	1816.	7. 8. G. ♀.	
<i>Cynara glomeráta. Thunberg.</i>					
3 atractyloides. s.s. (ye.)	Atractylis-like.	—	1823.	— G. ♀.	
CA'RTHAMUS. S.S.	BASTARD SAFFRON.	Syngenesia Polygamia Æqualis.			
1 tinctorius. w.	(or.) common.	Egypt.	1551.	6. 7. H. ♂.	
2 oxyacántha. s.s.	(ye.) sharp-spined.	Caucasus.	1829.	— H. ♂.	
HERACA'NTHA. L.en.	HERACA'NTHA.	Syngenesia Polygamia Æqualis.			
1 lanáta. L.en.	(ye.) woolly.	S. Europe.	1596.	7. 8. H. ♂.	
<i>Cártamus lanátus. B.M. Centaurea lanátu. DC.</i>					
2 crética. L.en.	(wh.) Cretan.	Candia.	1781.	6. 7. H. ♂.	
<i>Cártamus créticus. w.</i>					
3 taúrica. L.en.	(ye.) Taurian.	Tauria.	1818.	— H. ♂.	
4 glauca. (pu.)	glauca.	Iberia.	—	— H. ♂.	
<i>Cártamus glauca. M.B.</i>					
ONOBR'OMA. G.	ONOBR'OMA.	Syngenesia Polygamia Æqualis.			
1 tingitána.	(bl.) Tangier.	Barbary.	1759.	6. 7. H. ♀.	
<i>Cártamus tingitánu. w.</i>					
2 cerúlea. L.en.	(bl.) blue-flowered.	Spain.	1640.	— H. ♀.	
<i>Cártamus cerúleus. w.</i>					
3 arboréscens. s.s.	(ye.) tree.	Spain.	1731.	7. 8. F. ♂.	
<i>Cártamus arboréscens. w.</i>					
CARDUNCE'LLUS. DC.	CARDUNCE'LLUS.	Syngenesia Polygamia Æqualis.			
1 vulgáris. DC.	(bl.) common.	France.	1776.	5. 6. H. ♀.	
<i>Cártamus Carduncellus. w.</i>					
2 mitissimus. DC.	(bl.) small.	—	1734.	6. 7. H. ♀.	
CARLOW'IZIA. DC.	CARLOW'IZIA.	Syngenesia Polygamia Æqualis.			
salicifólia. DC.	(st.) Willow-leaved.	Madeira.	1784.	8. G. ♂.	
<i>Cártamus salicifolius. w. Onobroma salicifolium. L.en.</i>					

Jacq. vind. 2. t. 148.
— 2. t. 149.

Gouan. ill. t. 25.

Jacq. ic. 1. t. 167.

Vill. dauph. 3. t. 22.

Tabern. ic. 696.

Blackw. t. 458.

Flor. græc. t. 834.

Pluk. alm. t. 81. f. 2.

Desf. atl. 2. t. 223.

Knorr. thes. 2. t. c. 1.

Allion. ped. t. 51.

W. et K. hung. 2.t. 152.

Garid. aix. t. 21.

Flor. græc. t. 837.

Eng. bot. v. 16. t. 1144.

Bot. mag. t. 1788.

Bot. reg. t. 170.

Bot. mag. t. 2143.

Cavan. ic. 2. t. 128.

Bot. mag. t. 2293.

Lob. ic. 2. p. 20.

ATRA'CTYLIS. W.	ATRA'CTYLIS.	Syngenesia Polygamia Æqualis.			
humilis. w.	(wh.) dwarf.	Spain.	1759.	5. 7. H. 4.	Cav. ic. 1. t. 54.
ACA'RNA. W.	ACA'RNA.	Syngenesia Polygamia Æqualis.			
1 gummifera. w.	(pu.) gummy-rooted.	S. Europe.	1640.	6. 8. H. 4.	Cavan. ic. 3. t. 228.
2 serratoloides. s.s.(pu.)	Sawwort-like.	Palestine.	1827.	— H. 4.	
3 comosa. s.s.	(pu.) tufted.		—	— H. 4.	
4 cancellata. w.	(bl.) netted.	S. Europe.	—	6. 7. H. 4.	Lam. ill. t. 662. f. 1.
<i>Atracytis cancellata. P.S.</i>					
STOK'ESIA. W.	STOK'ESIA.	Syngenesia Polygamia Æequalis.			
cyanea. w.	(bl.) blue-flowered.	Carolina.	1766.	8. G. 4.	L'Herit. sert. 27. t. 38.
STÆHEL'INA. W.	STÆHEL'INA.	Syngenesia Polygamia Æequalis.			
1 dñbia. w.	(pu.) Rosemary-leav'd.	S. Europe.	1640.	6. 7. H. 4.	Lam. ill. t. 666. f. 4.
2 arborescens. w.	(re.) Storax-leaved.	Candia.	1739.	7. 9. F. 4.	Flor. græc. t. 345.
3 chamæpeuce. w.	(re.) Pine-leaved.		1640.	7. 11. F. 4.	— t. 347.
PTER'ONIA. W.	PTER'ONIA.	Syngenesia Polygamia Æequalis.			
1 camphorata. w.	(ye.) aromatic.	C. B. S.	1774.	6. 7. G. 4.	Pluk. mant. t. 345. f. 2.
2 stricta. w.	(ye.) cluster-flowered.	—	—	4. 6. G. 4.	
3 echinata. w.	(ye.) prickly-cupped.		1820.	— G. 4.	
4 flexicaulis. w.	(ye.) bending-stalk'd.		1812.	6. 8. G. 4.	
5 fasciculata. w.	(ye.) cluster-flowered.		1816.	— G. 4.	
6 pallens. w.	(pa.) pale.		—	— G. 4.	
7 oppositifolia. w.	(ye.) opposite-leaved.		1774.	7. G. 4.	Lam. ill. t. 667. f. 2.
8 glomerata. w.	(ye.) crowded.		1823.	6. 8. G. 4.	
9 scariosa. w.	(ye.) window-cupped.		1815.	— G. 4.	
Z'ŒGEA. W.	Z'ŒGEA.	Syngenesia Polygamia Frustranea.			
Leptairea. w.	(ye.) yellow-flower'd. Levant.		1779.	7. 8. H. 4.	Jacq. ic. 1. t. 177.
LE'UZEA. DC.	LE'UZEA.	Syngenesia Polygamia Frustranea.			DC. ann. mus. 16. t. 14.
1 conifera. DC.	(pu.) cone.	S. Europe.	1683.	6. 9. H. 4.	
<i>Centaura conifera. w.</i>					
2 altäica. L.en.	(pu.) large-flowered.	Altay.	1818.	— H. 4.	
3 dahurica. F.	(pu.) Daurian.	Dahuria.	1821.	— H. 4.	
4 australis. D.D.	(pu.) New Holland.	N. Holland.	1821.	8. 9. H. 4.	
GALA'CITITES. DC.	GALA'CITITES.	Syngenesia Polygamia Frustranea.			
tomentosa. DC.	woolly.	S. Europe.	1738.	7. H. 4.	DC. ann. mus. 16. t. 16.
<i>Centaura galactites. w.</i>					
CRUP'INA. Cas.	STARRY SCABIOUS.	Syngenesia Polygamia Frustranea.			
1 vulgaris. Cas.	(re.) black-seeded.	Italy.	1596.	6. 7. H. 4.	Colum. eph. 1. t. 34.
<i>Centaura Crupina. w.</i>					
2 crupinoides. P.S. (sn.)	saffron-colour'd.	Barbary.	1826.	— H. 4.	
3 Lippi. P.S.	Lippi's.	Egypt.	1739.	— H. 4.	Isn. act. par. 1719. t. 10.
<i>Centaura Lippi. w.</i>					
4 arenaria. P.S.	(pu.) sand.	Volga.	1821.	— H. 4.	
CENTA'UREA. W.	CENTAURY.	Syngenesia Polygamia Frustranea.			
1 moschata. w.	(li.) sweet sultan.	Persia.	1629.	7. 10. H. 4.	Knor. thes. 2. t. C. 4.
2 glauca. w.	(pu.) glaucous.	Caucasus.	1805.	6. 7. H. 4.	Swt. brit. fl. gar. 1. t. 51.
3 suavolens. w.	(ye.) yellow sultan.	Levant.	1683.	7. 10. H. 4.	Cornut. can. 69. t. 70.
4 alpina. w.	(ye.) Alpine.	Italy.	1640.	7. 8. H. 4.	
5 Centaurium. w.	(bl.) great.		1596.	— H. 4.	
6 ruthenica. w.	(st.) Russian.	Russia.	1806.	— H. 4.	Gmel. sib. 2. t. 41.
7 phrygia. w.	(pu.) feathery-cupp'd.	Switzerland.	1633.	6. 10. H. 4.	Flor. dan. t. 520.
8 rivularis. P.S.	(br.) river-side.	Portugal.	1812.	7. 9. H. 4.	
9 salicifolia. w.	(pu.) Willow-leaved.	Caucasus.	1816.	— H. 4.	
10 austriaca. w.	(pu.) Austrian.	Austria.	1815.	6. 10. H. 4.	
11 pectinata. w.	(pu.) pectinated.	France.	1727.	7. 10. H. 4.	Bocc. mus. 2. t. 2.
12 uniflora. w.	(pu.) one-flowered.	S. Europe.	1824.	6. 8. H. 4.	
13 flosculosa. w.	(pu.) flosculose.	Italy.	1817.	— H. 4.	
14 nervosa. w.	(pu.) nerved.	S. Europe.	1815.	6. 9. H. 4.	
15 linifolia. P.S.	(pu.) Flax-leaved.		1826.	— H. 4.	
16 capillata. P.S.	(pu.) hairy-headed.	Volga.	1827.	— H. 4.	
17 tricocéphala. w.	(pu.) downy-cupped.	Siberia.	1805.	7. 8. H. 4.	Gmel. sib. 2. t. 45. f. 1.2.
18 hyssopifolia. w.	(pu.) Hyssop-leaved.	Spain.	1812.	— F. 4.	Barrel. ic. 306.
19 coronopifolia. w.	(ye.) Buck's-horn-ld.	Levant.	1739.	6. 7. H. 4.	
20 procumbens. P.S.	(pu.) procumbent.	Italy.	1821.	— H. 4.	
21 nigra. w.	(pu.) black knapweed.	Britain.	...	5. 8. H. 4.	
22 pratensis. P.S.	(pu.) field.	France.	1824.	— H. 4.	Eng. bot. v. 4. t. 278.
23 nigræscens. w.	(pu.) dark.	Hungary.	1805.	6. 8. H. 4.	
24 decipiens. P.S.	(pu.) deceived.	France.	1826.	— H. 4.	
25 Triumfetti. w.	(re.) Cenian.	Mt. Cenix.	—	— H. 4.	
26 montana. w.	(bl.) mountain.	Austria.	1596.	— H. 4.	Bot. mag. t. 77.

27 <i>seusâna</i> . p.s.	silvery-fringed.	S. Europe.	1816.	6. 8. H. 4.	Barrel. ic. 389.
<i>axillaris</i> , w. <i>variegata</i> . Lam.					
28 <i>strícta</i> . w.k.	(<i>bl.</i>) upright.	Hungary.	1826.	— H. 4.	
29 <i>depræssa</i> . m.b.	(<i>bl.</i>) depressed.	Caucasus.	—	— H. 4.	
30 <i>lingulâta</i> . s.s.	(<i>bl.</i>) tongue-leaved.	Spain.	1825.	— H. 4.	
31 <i>spathulâta</i> . s.s.	(<i>bl.</i>) spathulate-leav'd.	Naples.	1827.	— H. 4.	
32 <i>Cyânus</i> . w.	(<i>bl. va.</i>) blue-bottle.	Britain.	— H. 4.	
<i>Cyânus ségetum</i> . Mench.					
33 <i>americâna</i> . n.	(<i>li.</i>) American.	N. America.	1823.	7. 10. H. 4.	
34 <i>ochroleúca</i> . w.	(<i>st.</i>) Caucasian.	Caucasus.	1801.	7. 8. H. 4.	
35 <i>africana</i> . w.	(<i>ye.</i>) African.	Africa.	1825.	— F. 4.	
36 <i>cheiranthifolia</i> . w. (<i>ye.</i>)	Stock-leaved.	Caucasus.	1829.	— H. 4.	
37 <i>ovina</i> . w.	(<i>ye.</i>) small-flowered.	—	1802.	7. 11. H. 4.	
38 <i>paniculâta</i> . w.	(<i>pu.</i>) paniced.	Europe.	1640.	7. 8. H. 4.	
39 <i>intybâcea</i> . p.s.	(<i>re.</i>) Succory-leaved.	S. Europe.	1778.	7. 9. H. 4.	
40 <i>leucânta</i> . Lam.	(<i>wh.</i>) white-blossom'd.	—	1826.	— H. 4.	
41 <i>spinôsa</i> . w.	(<i>wh.</i>) prickly-branch'd.	Candia.	1640.	— F. 4.	
42 <i>ragusina</i> . w.	(<i>ye.</i>) white-leaved.	—	1710.	6. 7. G. 4.	
43 <i>argéntea</i> . w.	(<i>ye.</i>) silver-leaved.	—	1739.	7. 8. F. 4.	
44 <i>Cinerâria</i> . w.	(<i>pu.</i>) hoary-leaved.	Italy.	1710.	— H. 4.	
45 <i>cinérea</i> . w.	(<i>pu.</i>) gray.	—	—	6. 7. H. 4.	
46 <i>dealbâta</i> . m.b.	(<i>pu.</i>) mealy.	Caucasus.	1804.	7. 8. H. 4.	
47 <i>cicutæfolia</i> . H.H.	(<i>pu.</i>) Cicutæ-leaved.	Europe.	1824.	— H. 4.	
48 <i>maculôsa</i> . p.s.	(<i>pu.</i>) spotted-cupped.	Siberia.	1816.	— H. 4.	
49 <i>Marschalliana</i> . s.s. (<i>pu.</i>)	Marschall's.	Caucasus.	1829.	— H. 4.	
50 <i>declinâta</i> . m.b.	(<i>pu.</i>) declining.	—	1827.	— H. 4.	
51 <i>leucophylla</i> . M.B. (<i>pu.</i>)	white-leaved.	—	1824.	— H. 4.	
52 <i>trinérvia</i> . w.	(<i>pu.</i>) three-nerved.	—	1828.	— H. 4.	
53 <i>dissécta</i> . T.N.	(<i>pu.</i>) cut-leaved.	Italy.	1824.	— H. 4.	
54 <i>semperfîrens</i> . w.	(<i>ye.</i>) evergreen.	Spain.	1683.	— H. 4.	
55 <i>coriâcea</i> . w.	(<i>re.</i>) leathery-leav'd.	Hungary.	1804.	6. 7. H. 4.	
56 <i>Scabiôsa</i> . w.	(<i>pu.</i>) greater-knapweed.	Britain.	6. 8. H. 4.	
57 <i>tatârica</i> . w.	(<i>ye.</i>) Tartarian.	Tartary.	1801.	7. 8. H. 4.	
58 <i>St'œbe</i> . w.	(<i>pu.</i>) wing-leaved.	Austria?	1759.	6. 7. H. 4.	
59 <i>Fischèri</i> . W.en.	(<i>pu.</i>) Fischer's.	Siberia.	1816.	— H. 4.	
60 <i>macrocéphala</i> . w.	(<i>st.</i>) large-headed.	Caucasus.	1805.	6. 8. H. 4.	
61 <i>atropurpùrea</i> . w.	(<i>pu.</i>) dark purple.	Hungary.	1802.	— H. 4.	
62 <i>calocéphala</i> . W.en. (<i>ye.</i>)	smooth-stalked.	Llevant.	1816.	— H. 4.	
63 <i>canariensis</i> . W.en. (<i>wh.</i>)	Canary.	Teneriffe.	1815.	— G. 4.	
64 <i>orientâlis</i> . w.	(<i>ye.</i>) oriental.	Levant.	1759.	7. 8. H. 4.	
65 <i>pulcherrima</i> . w.	(<i>pu.</i>) beautiful.	Caucasus.	1823.	— H. 4.	
66 <i>arachnoïdea</i> . P.S.	(<i>ye.</i>) cobweb.	Italy.	—	— H. 4.	
67 <i>sibírica</i> . w.	(<i>pu.</i>) Siberian.	Siberia.	1782.	— H. 4.	
68 <i>dilâta</i> . w.	(<i>li.</i>) pale-flowered.	S. Europe.	1781.	— H. 4.	
69 <i>alâta</i> . w.	(<i>ye.</i>) winged-stalked.	Tartary.	—	8. 9. H. 4.	
70 <i>elongâta</i> . s.s.	(<i>li.</i>) elongated.	Barbary.	1822.	— H. 4.	
71 <i>Béhen</i> . w.	(<i>ye.</i>) saw-leaved.	Levant.	1797.	7. 8. H. 4.	
72 <i>répens</i> . w.	(<i>ye.</i>) creeping.	—	1739.	6. 8. H. 4.	
73 <i>Jacea</i> . w.	(<i>pu.</i>) brown knapweed.	England.	7. 9. H. 4.	
74 <i>decumbens</i> . p.s.	(<i>pu.</i>) decumbent.	France.	1815.	— H. 4.	
75 <i>amâra</i> . w.	(<i>pu.</i>) bitter.	Italy.	7. 8. H. 4.	
76 <i>álba</i> . w.	(<i>pu.</i>) white.	Spain.	1597.	6. 9. H. 4.	
77 <i>spléndens</i> . w.	(<i>pu.</i>) shining.	—	—	7. 8. H. 4.	
78 <i>nítens</i> . w.	(<i>pu.</i>) glittering.	Caucasus.	1823.	— H. 4.	
79 <i>tagâna</i> . w.	(<i>pu.</i>) Portugal.	Portugal.	1640.	— H. 4.	
80 <i>babylônia</i> . w.	(<i>pu.</i>) Babylonian.	Levant.	1710.	6. 9. H. 4.	
81 <i>glastifolia</i> . w.	(<i>ye.</i>) woad-leaved.	Siberia.	1731.	— H. 4.	
82 <i>parviflôra</i> . s.s.	(<i>pu.</i>) small-flowered.	S. Europe.	1823.	— H. 4.	
83 <i>multifida</i> . p.s.	(<i>st.</i>) multifid.	—	1824.	— H. 4.	
84 <i>sordida</i> . s.s.	(<i>pu. ye.</i>) sordid.	—	1817.	— H. 4.	
85 <i>rígida</i> . s.s.	(<i>pu.</i>) rigid.	—	1824.	— H. 4.	
86 <i>Barrelieri</i> . s.s.	(<i>pu.</i>) Barrelier's.	Hungary.	1826.	— H. 4.	
87 <i>stereophylla</i> . s.s.	(<i>pu.</i>) variable-leaved.	Poland.	1828.	— H. 4.	
88 <i>spinulôsa</i> . s.s.	(<i>pu.</i>) small-spined.	Hungary.	1825.	— H. 4.	
89 <i>tenuifolia</i> . Leh.	(<i>pu.</i>) slender-leaved.	Siberia.	1829.	— H. 4.	
90 <i>subspinôsa</i> . Leh.	(<i>pu.</i>) somewhat spiny.	—	—	— H. 4.	
91 <i>jacobæfolia</i> . s.s.	(<i>ye.</i>) Jacobæa-leaved.	—	—	— H. 4.	
92 <i>reflexa</i> . s.s.	(<i>ye.</i>) crook-spined.	Iberia.	1801.	7. 8. H. 4.	
93 <i>centauroïdes</i> . s.s.	(<i>ye.</i>) lyre-leaved.	S. Europe.	1739.	5. 7. H. 4.	

Column. ephr. 1. t. 35.

Eng. bot. v. 4. t. 277.

Swt. br. fl. gar. ic.

Bot. mag. t. 1175.

Jacq. aust. 4. t. 320.

Barrel. ic. 359.

Bot. mag. t. 2493.

— t. 494.

Barrel. ic. t. 218.

Moris. s. 7. t. 26. f. 20.

Jacq. vind. 1. t. 92.

Gmel. sib. 2. t. 44.f.1.2.

Bocc. sic. t. 39. f. 3.

W. et K. hung. 2. t. 195.

Eng. bot. v. 1. t. 56.

Bot. mag. t. 1248.

W. et K. hung. 2. t. 116.

Viv. fl. it. fr. t. 8.

Gmel. sib. 2. t. 42. f. 2.

Venten. cels. t. 80.

Eng. bot. v. 24. t. 1678.

Bocc. mus. 31. t. 17.

Buxb. cent. 2. t. 15. f. 1.

Brot. phyt. lus. t. 3.

Alp. exot. t. 282.

Bot. mag. t. 62.

Barrel. ic. 310.

94 collina. s.s. (ye.) hill.	S. Europe.	1596.	6. 7. H. ȝ.	
95 pubescens. s.s. (ye.) downy.	1804.	7. 8. H. ȝ.	
96 nicaensis. s.s. (ye.) Italian.	Italy.	1827.	— H. ȝ.	Column. ecp. 1.t. et. f. 2.
97 rupéstris. s.s. (ye.) rock.	—	1804.	— H. ȝ.	
98 hybrida. s.s. (pu. ye.) hybrid.	—	1821.	— H. ȝ.	
99 sabulosa. s.s. (wh.) gravel.	Siberia.	1828.	— H. ȝ.	
100 Crocodylium. s.s. (bh.) blush-flowered.	Levant.	1777.	— H. ȝ.	Barrel. rar. t. 503.
101 aurea. s.s. (go.) great golden.	S. Europe.	1758.	7. 9. H. ȝ.	Bot. mag. t. 421.
102 scoparia. s.s. (pu.) broom.	Levant.	1829.	— H. ȝ.	Jacq. ic. t. 178.
103 Verutum. s.s. (ye.) dwarf.	—	1780.	8. 9. H. ȝ.	
104 muricata. s.s. (pu.) muricated.	Spain.	1621.	7. 8. H. ȝ.	
105 pubigera. P.S. (pu.) pubescent.	S. Europe.	1828.	— H. ȝ.	
106 peregrina. s.s. (pu.) soft-leaved.	—	1749.	— H. ȝ.	
107 cichoracea. s.s. (pu.) Succory-like.	Italy.	7. 9. H. ȝ.	Till. pis. p. 84. t. 27.
108 salmantica. s.s. (wh. pu.) ragwort-leaved.	S. Europe.	1596.	7. 8. H. ȝ.	Jacq. vind. 1. t. 64.
109 sonchifolia. s.s. (pu.) Sow-thistle-ld.	—	1780.	8. 10. H. ȝ.	Pluk. phyt. 39. f. 1.
110 Séridis. s.s. (pu.) purple-flower'd.	Spain.	1686.	6. 8. H. ȝ.	Pluk. alm. t. 38. f. 1.
111 napifolia. s.s. (pu.) Turnep-leaved.	Candia.	1691.	7. 9. H. ȝ.	Herm. parad. t. 189.
112 Zanónii. s.s. (pu.) Zanon's.	—	1822.	— H. ȝ.	
113 Isnárdi. s.s. (li.) Jersey.	Jersey.	7. 8. H. ȝ.	Eng. bot. v. 32. t. 2256.
114 áspéra. s.s. (pu.) rough.	S. Europe.	1772.	6. 10. H. ȝ.	Bocc. mus. 35. t. 26.
115 sphærocéphala. s.s. (pu.) globe-headed.	—	1683.	6. 8. H. ȝ.	Bot. mag. t. 2551.
116 auriculata. P.S. (pu.) eared.	—	1816.	7. 8. H. ȝ.	
117 polyacantha. s.s. (pu.) many-spined.	Portugal.	1804.	— H. ȝ.	
118 prolifera. P.S. (st.) proliferous.	Egypt.	1816.	— H. ȝ.	Vent. hort. cels. t. 16.
straminea. W. hort. ber. t. 26.				
119 ibérica. s.s. (pu.) Iberian.	Iberia.	1821.	— H. ȝ.	
120 ferox. s.s. (ye.) hedgehog.	Barbary.	1790.	7. 9. F. ȝ.	Desf. atl. 2. t. 242.
121 solstitialis. s.s. (ye.) St. Barnaby's.	England.	7. 8. H. ȝ.	Eng. bot. v. 4. t. 243.
122 melitensis. s.s. (ye.) cluster-headed.	Malta.	1710.	— H. ȝ.	Bocc. sic. t. 35.
123 A'damii. s.s. (ye.) Adam's.	Siberia.	1804.	— H. ȝ.	
124 sulphurea. W.en. (st.) straw-coloured.	—	1815.	— H. ȝ.	
125 sícula. s.s. (st.) Sicilian.	Sicily.	1710.	— H. ȝ.	Bocc. sic. t. 8. f. 1.
126 Stevénii. s.s. (ye.) Steven's.	Caucasus.	1826.	— H. ȝ.	
127 cruenta. W.en. (pu.) obovate-leaved.	S. Europe.	1816.	6. 8. H. ȝ.	
128 romana. L. (pu.) Roman.	Italy.	1739.	7. 9. H. ȝ.	Barrel. rar. t. 504.
129 benedicta. s.s. (st.) blessed thistle.	Spain.	1548.	6. 10. H. ȝ.	Zorn. ic. t. 122.
130 pullata. s.s. (br.) sad-coloured.	S. Europe.	1714.	6. 7. H. ȝ.	Mill. ic. 2. t. 152. f. 2.
131 eriophora. s.s. (ye.) woolly-headed.	Portugal.	1714.	6. 10. H. ȝ.	
132 ápula. s.s. (ye.) Apulian.	Sicily.	1829.	6. 8. H. ȝ.	Column. ecp. 1. t. 31.
133 Calcítrapa. s.s. (li.) Star-thistle.	Britain.	— H. ȝ.	Eng. bot. v. 2. t. 125.
134 calcitrapoides. s.s. (li.) Phœnician.	Levant.	1683.	6. 7. H. ȝ.	
135 ægyptiaca. s.s. (wh.) Egyptian.	Egypt.	1790.	6. 9. H. ȝ.	
136 balsamita. w. (ye.) Balsam-scented.	Armenia.	1824.	7. 8. H. ȝ.	
137 lanata. DC. (ye.) hoary.	Switzerland.	1825.	— H. ȝ.	
138 crética. s.s. (wh.) Cretan.	Candia.	1827.	— H. ȝ.	
139 bullata. Leh. blistered.	—	1829. H. ȝ?	
140 ucránica. Ot. Ukraine.	Ukraine.	— H. ȝ?	
RHAPO'NTICA. DC. RHAPO'NTICA.	Syngenesia Polygamia Frustranea.			
1 scariosa. DC. (pu.) common.	Switzerland.	1640.	7. 8. H. ȝ.	Bot. mag. t. 1752.
Centaurea Rhapontica. B.M.				
β lyrata. DC. (pu.) lyrate.	—	— H. ȝ.	
2 uniflora. DC. one-flowered.	Siberia.	1796.	— H. ȝ.	Gmel. sib. 2. t. 38.
Cnicus uniflorus. w.				
3 doronicæfolia. F. (pu.) Doronicum-leav'd.	—	1825.	6. 8. H. ȝ.	
4 dahurica. F. (pu.) Dahurian.	Dahuria.	1829.	— H. ȝ.	
CARDOP'ATUM. J. CARDOP'ATUM.	Syngenesia Polygamia Segregata.			
corymbosum. P.S. (bl.) umbelled.	S. Europe.	1640.	6. 7. H. ȝ.	Moris. s. 7. t. 33. f. 17.
Brótera corymbosa. w.				

Div. II. ECHINOPSIDEÆ. Kth. synops. 2. p. 364.

ROLA'NDRA. W.	ROLA'NDRA.	Syngenesia Polygamia Segregata. L.	
argentea. w. (wh.) silver-leaved.	W. Indies.	1714.	7. S. h.
ECHINO'PS. W.	GLOBE-THISTLE.	Syngenesia Polygamia Segregata. L.	Sloan. jam. 1. t. 7. f. 3.
1 sphærocéphalus. w. (wh.) great.	Austria.	1596.	7. 8. H. ȝ.
β exaltatus. Schr. (wh.) tall.	—	1820. H. ȝ.
			Flor. græc. t. 923.

2 spinosus. w. (wh.)	thorny-headed.	Egypt.	1597.	7. 8. H. ȝ.	Flo. græc. t. 924.
3 paniculatus. B.R. (bl.)	panicled.	Spain.	1815.	— H. ♂.	P. reg. t. 356.
4 hórridus. P.S. (wh.)	horrid.	Persia.	1817.	— H. ȝ.	
5 viscosus. M.B. (wh.)	viscous.	Caucasus.	1823.	— H. ♂.	
6 taúricus. W.en. (bl.)	Taurian.	Tauria.	1816.	— H. ȝ.	
7 dahúricus. F. (bl.)	Dahurian.	Dahuria.	1827.	— H. ȝ.	
8 ruthénicus. M.B. (bl.)	Russian.	Russia.	1819.	— H. ȝ.	
9 Rítro. w. (bl.)	small.	Europe.	1570.	7. 9. H. ȝ.	
10 tenuifòlius. F. (bl.)	slender-leaved.	Russia.	1823.	— H. ȝ.	
11 strictus. B.M. (bl.)	upright.	—	1822.	8. 10. H. ♂?	Bot. mag. t. 932.
12 pérsicus. F. (wh.)	Persian.	Persia.	1821.	— H. ȝ.	Bot. mag. t. 2457.
13 bannáticus. w.k. (wh.)	Hungarian.	Hungary.	1828.	— H. ȝ.	Flor. græc. t. 926.
14 lanuginósus. w. (bl.)	woolly.	Levant.	1736.	6. 7. H. ȝ.	
15 hùmilis. M.B. (bl.)	low.	Siberia.	1826.	6. 8. H. ȝ.	Bot. mag. t. 2109.
16 strigósus. w. (wh.)	annual.	Spain.	1729.	7. 9. H. ⓧ.	Mill. ic. t. 287.
GUND'ELIA. W.	GUND'ELIA.	Syngenesia Polygamia Segregata. L.			L.
Tournefôrtii. w. (pu.)	Tournefort's.	Levant.	1739.	6. 8. H. ȝ.	
SPHÆRA'NTHUS. W.	SPHÆRA'NTHUS.	Syngenesia Polygamia Segregata.			
1 móllis. H.B. (bl.)	soft.	E. Indies.	1818.	7. 9. S. ⓧ.	Burm. zeyl. t. 94. f. 3.
2 índicus. w. (pu.)	Indian.	—	1699.	8. 12. S. ȝ.	
3 cochinchinénsis.w.(bl.)	Cochinchina.	Cochinchina.	1826.	7. 8. H. ⓧ.	Lam. ill. t. 718. f. 1.
4 hirtus. w. (bl.)	hairy.	Manila.	1823.	— G. ⓧ.	Pluk. mant. t. 108. f. 7.
5 africáñus. w. (bl.)	African.	C. B. S.	1759.	7. 8. G. ⓧ.	
ANGIA'NTHUS. W.C. ANGIA'NTHUS.	Syngenesia Polygamia Segregata.				
tomentósus. w.c. (ye.)	woolly.	N. Holland.	1803.	7. 8. G. ȝ.	
Cassinia aurea. H.K. non L.T.					
BROTERA. P.S.	BROTERA.	Syngenesia Polygamia Segregata. P.S.			
trinerváta. p.s. (ye.)	three-nerved.	S. America.	1799.	7. 8. H. ⓧ.	
Contrayérala. Spr.	Nauenburgia trinerváta. w.				
CED'ERA. W.	CED'ERA.	Syngenesia Polygamia Segregata. L.			
1 prolífera. w. (ye.)	proliferous.	C. B. S.	1789.	5. 6. G. ȝ.	Bot. mag. t. 1637.
2 aliena. w. (ye.)	linear-leaved.	—	1821.	5. 7. G. ȝ.	Jacq. schoenb. 2. t. 154.
ELEPHHA'NTOPUS. W.	ELEPHANT'S-FOOT.	Syngenesia Polygamia Segregata.			
1 scâber. w. (re.)	rough-leaved.	E. Indies.	1695.	6. 9. S. ȝ.	
2 caroliniánus. w. (re.)	Carolina.	America.	1732.	7. 9. G. ȝ.	Rheed. mal. 10. t. 7.
3 tomentósus. w. (re.)	woolly.	W. Indies.	1733.	7. 8. S. ȝ.	Dill. elth. t. 106. f. 126.
4 móllis. k.s. (wh.)	soft-leaved.	Caracas.	1825.	— S. ȝ.	
5 nudiflòrus. w. (re.)	naked-flower'd.	W. Indies.	1822.	— S. ȝ.	
6 spicátus. w. (re.)	spiked.	—	1821.	— S. ȝ.	
7 angustifòlius. w. (re.)	narrow-leaved.	—	1823.	— S. ȝ.	
ST'CEBE. W.	ST'CEBE.	Syngenesia Polygamia Segregata. L.			
1 áethiòpica. w. (wh.)	Juniper-leaved.	C. B. S.	1759.	8. — G. ȝ.	
Seriphium áethiòpicum. P.S.					
2 ericóides. w. (wh.)	Heath-like.	—	1816.	7. 9. G. ȝ.	Moris. s. 7. t. 18. f. 12.
3 gomphrenoídes.w.(w.)	Gomphrena-like.	—	—	— G. ȝ.	Houtt. syst. 4. t. 34. f. 1.
4 gnaphaloídes. w. (wh.)	Gnaphalium-like.	—	1824.	— G. ȝ.	Burm. afr. t. 77. f. 1.
5 cinérea. w. (wh.)	Heath-leaved.	—	1774.	— G. ȝ.	Pluk. mant. t. 297. f. 1.
Seriphium cinéreum. P.S.					
6 alopecuroídes.w.(wh.)	Fox-tail-like.	—	1824.	— G. ȝ.	Pluk. alm. t. 445. f. 2.
7 reflexa. w. (wh.)	reflex-leaved.	—	1816.	— G. ȝ.	
CÆSULIA. Roxb.	CÆSULIA.	Syngenesia Polygamia Segregata. L.T.			
1 axilláris. w. (wh.)	axillary-flower'd.	E. Indies.	1804.	7. 9. S. ȝ.	Andr. reposit. t.
2 lancifòlia.	(wh.) spear-leaved.	—	1820.	— S. ȝ.	
LAGA'SCA. Cav.	LAGA'SCA.	Syngenesia Polygamia Segregata. L.T.			
móllis. w. (wh.)	soft.	S. America.	1815.	6. 9. G. ⓧ.	Bot. mag. t. 1804.
NO'CCA. W.	No'CCA.	Syngenesia Polygamia Segregata.			
1 rígida. c.i. (re.)	red-flowered.	Mexico.	1823.	8. 10. F. ȝ.	Swt. br. fl. gar. s. 2. t. 26.
Lagáscea rubra. H. et B. pl. amer. 4. t. 311.					
2 suaveolens. (wh.)	sweet-scented.	Mexico.	—	... F. ȝ.	
Lagáscea suavèolens. K.S.					
3 latifòlia. B.F.G. (wh.)	broad-leaved.	—	—	... F. ȝ.	Swt. br. fl. gar. t. 215.

Div. III. VERNONIACEÆ. Kth. synops. 2. p. 367. et ASTEREÆ. l. c. p. 99.

AMPHE'REPHIS. K.S. AMPHE'REPHIS. Syngenesia Polygamia Æqualis. intermédia. L.en. (bl.) intermediate. Brazil. 1822. 7. 9. H. ⓧ.

Swt. br. fl. gar. t. 225.

ASCARI'CIDA. Cass.	ASCARI'CIDA.	Syngenesia Polygamia $\ddot{\text{E}}$ qualis.		
anthelm. <i>sc.</i> (u.) Indian.	E. Indies.	1770. 8. 9. S. ♂.		Rheed. mal. 2. t. 24.
<i>indica.</i> Cass.	<i>rnònia anthelmintica.</i> W.	<i>Conyzza anthelmintica.</i> L.		
VERN'ONIA. W.	VERN'ONIA.	Syngenesia Polygamia $\ddot{\text{E}}$ qualis. W.		
1 noveboracénsis. (pu.) long-leav'd.	N. America.	1710. 9. 11. H. ♀.		Dill. elt. t. 263. f. 342.
2 præalta. w. (pu.) tall.		1732. — H. ♀.		— t. 264. f. 343.
3 angustifolia. Ph. (pu.) narrow-lea d.		1817. — H. ♀.		
4 glauca. w. (pu.) glaucous-lea d.		1710. — H. ♀.		Dill. elt. t. 262. f. 341.
<i>oligophylla.</i> M.				
5 altissima. N. (pu.) tallest.		1823. — H. ♀.		
6 pandurata. L.en. (pu.) fiddle form.		— H. ♀.		
7 serratuloides. K.S. (pu.) Sawwort-like.	Mexico.	1824. — H. ♀.		
8 axilliflora. Ot. (pi.) axillary-flower'd.		1829. 9. 10. F. ♀.		
9 centriflora. Lk. (pu.) centre-flower'd. Brazil.		1826. — S. ♂.		
10 sericea. B.R. (li.) silky.		1818. 9. 1. S. ♂.		
11 arborescens. P.S. (pu.) tree.	Jamaica.	1733. 11. 12. S. ♂.		
<i>Conyzza arborescens.</i> W.				
12 fruticosa. P.S. (pu.) shrubby.	S. America.	1818. — S. ♂.		Plum. ic. 95. f. 1.
<i>Conyzza fruticosa.</i> L.				
13 flexuosa. B.M. (pu.) flexuose-branch'd. Brazil.		1823. 9. 10. S. ♂.		Bot. reg. t. 522.
LI'ATRIS. W.	LI'ATRIS.	Syngenesia Polygamia $\ddot{\text{E}}$ qualis. W.		Plum. sp. 10. t. 130. f. 2.
1 spicata. w. (pu.) long-spiked.	N. America.	1732. 8. 10. H. ♀.		
β <i>pumila.</i> B.C. (pu.) dwarf.		— H. ♀.		
2 pycnostachya. M. (pu.) pubescent.		1732. — H. ♀.		
3 heterophylla. Ph. (pu.) various-leaved.		1790. 7. 8. H. ♀.		
4 intermedia. B.R. (pu.) intermediate.		1823. 8. 9. H. ♀.		
5 cylindracea. M. (pu.) cylindrical-cup'd.		1811. 10. H. ♀.		
6 pilosa. Ph. (pu.) hairy-leaved.		1783. — H. ♀.		
7 gracilis. Ph. (pu.) smooth-leaved.		1812. — H. ♀.		
8 turbinata. B.F.G. (pu.) turbinate-cup'd.		1823. — H. ♀.		
9 elegans. w. (pu.) hairy-cup'd.		1787. 9. 10. H. ♀.		
10 sphæroidea. M. (pu.) globular-cup'd.		1817. 8. 10. H. ♀.		
11 scariosa. w. (pu.) scarioso-cup'd.		1739. 9. 10. H. ♀.		
12 squarrosa. w. (pu.) rough-cup'd.		1732. — H. ♀.		
13 paniculata. w. (pu.) paniced.		1826. 8. 10. H. ♀.		
14 odoratissima. w. (pu.) sweet-scented.	Carolina.	1786. — F. ♀.		
15 corymbosa. w. (pu.) corymbose.	N. America.	1825. — F. ♀.		
BRACHYL'ENA. L.T. BRACHYL'ENA.	Syngenesia Polygamia Superflua.			
neriifolia. L.T. (st.) Oleander-leav'd. C. B. S.		1752. 8. 11. G. ♂.		
<i>Baccharis neriifolia.</i> w.				
BA'CCHARIS. W.	PLOWMAN'S-SPIKENARD.	Syngenesia Polygamia Superflua.		
1 ivafolia. w. (wh.) Peruvian.	America.	1696. 7. 8. G. ♂.		Schk. handb. 3. t. 244.
2 angustifolia. M. (wh.) narrow-leaved.	N. America.	1812. 7. 9. G. ♂.		
3 glomeruliflora. M. (wh.) close-flowered.		1817. 8. 10. H. ♂.		
4 halimifolia. w. (wh.) Groundsel-tree.		1683. 10. 11. H. ♂.		
5 Dioscoridis. w. (wh.) Dioscorides'.	Levant.	1822. 7. 9. G. ♂.		
6 scoparia. P.S. (wh.) broom.	Jamaica.	— — S. ♂.		
<i>Calea scoparia.</i> L.				
7 mollis. K.S. (wh.) soft.	Quito.	1824. 5. 6. F. ♂.		
8 genistelloides. K.S. (w.) wing-stemmed.	Brazil.	1829. 7. 10. G. ♂.		
alata. Hort. <i>Pterocladis genistelloides.</i> Lamb. herb.				
9 cerviflora. P.S. (wh.) small-flowered.	Peru.	1822. — S. ♂.		
10 lycopodioides. P.S. (wh.) Club-moss-like.	Mauritius.	1828. — S. ♂.		
<i>Conyzza lycopodioides.</i> Lam.				
N'YZA. W.	FLEA-BANE.	Syngenesia Polygamia Superflua. L.		
squarrosa. w. (ye.) great.	Britain.	— 7. 8. H. ♂.		
marylandica. Ph. (pu.) Maryland.	N. America.	— 8. 10. H. ♂.		Eng. bot. v. 17. t. 1195.
uphorata. Ph. (pu.) Camphor-scent'd.		1704. — H. ♀.		Dill. elt. t. 88. f. 104.
<i>echaris f. atida.</i> w.				— t. 89. f. 105.
1 tula. w. (pu. ye.) spreading.	China.	1758. 7. 9. H. ♂.		
4 p. <i>an-</i> w. (re.) axillary.	Mauritius.	1824. — H. ♂.		
5 a. <i>ifrons.</i> w. (st.) oval-leaved.	N. America.	1739. 8. 9. H. ♀.		
6 purpureascens. w. (li.) purplish.	Jamaica.	1822. — S. ♂.		
7 pindji. a. L.	Candia.	1714. 6. 7. F. ♂.		
8 ct. <i>cifolia.</i> w.		— — F. ♂.		
9 nifolia. F.G. (ye.) wh.				
e dida. w. non L.				
10 balsamifera. w. (re.) balsamiferus.	China.	1818. — S. ♀.		
11 chinensis. w. (bl.) Chinese.		1796. 7. 8. S. ♂.		

Mill. ic. 2. t. 247.
Pluk. alm. t. 87. f. 4.
Sloan. hist. 1. t. 159. f. 1.
Flor. græc. t. 865.
Barr. ic. t. 217.
Rumph. amb. 6. t. 24. f. 1.
— 6. t. 14. f. 2.

COMPOSITE.

12	cinérea. w.	(<i>pu.</i>) grey.	E. Indies.	7. 8.	S. ○.	Rumph. amb. 6.t.14.f.1.
13	fastigiata. W.en.	(<i>pu.</i>) clustered.	Senegal.	1820.	—	H. ○.	
14	prolífera. w.	(<i>wh.</i>) proliferous.	Java.	1819.	7. 9.	S. ½.	Rumph. am. 5.t.104.f.1.
15	hirsúta. w.	(<i>ye.</i>) shaggy.	China.	1767.	8. 9.	G. ♂.	
16	ægyptiaca. w.	(<i>ye.</i>) Egyptian.	Egypt.	1778.	7. 9.	H. ○.	Jacq. vind. 3. t. 19.
17	chilénica. s.s.	(<i>ye.</i>) Chile.	Chile.	1816.	8. 10.	H. ○.	
18	mauritiana. Boj.	(<i>pu.</i>) Mauritius.	Mauritius.	1821.	—	S. ○.	
19	Hubértii. Boj.	(<i>pu.</i>) Hubert's.	Madagascar.	1824.	7. 10.	S. ○.	
20	viscosa. Boj.	(<i>pu.</i>) viscous.	Mauritius.	—	8. 10.	S. ○.	
21	púmila. s.s.	(<i>ye.</i>) dwarf.	Crete.	1826.	7. 9.	H. ○.	
22	pinnatifida. R.	(<i>pu.</i>) pinnatifid.	E. Indies.	1816.	—	S. ½.	
23	Gouaniá. w.	(<i>ye.</i>) Gouan's.	Canaries.	1772.	7. 8.	G. ♂.	Jacq. vind. 3. t. 79.
24	sícula. w.	(<i>ye.</i>) red-stalked.	Sicily.	1779.	8. 9.	H. ○.	Bocc. sic. t. 31. f. 4.
25	aúrtia. w.	(<i>wh.</i>) eared.	E. Indies.	1818.	—	S. ○.	
26	fœtidia. w.	(<i>wh.re.</i>) stinking.	Africa.	1724.	—	G. ¼.	Mill. ic. 2. t. 233.
27	amœna. L.en.	(<i>pu.</i>) pretty.	—	1816.	—	S. ¼.	
28	glomerata. L.en.	(<i>pu.</i>) crowded.	—	1824.	—	S. ¼.	
29	spatulata. L.en.	(<i>bl.</i>) spatulate.	—	—	7. 9.	S. ¼.	
30	adnáta. k.s.	(<i>pu.</i>) decurrent.	Mexico.	—	—	S. ½.	
	Bácccharis adnáta. W.en.						
31	thapsoides. M.B.	(<i>pu.</i>) Thapsus-leav'd.	Caspian Sea.	1806.	—	H. ¼.	
32	sordida. w.	(<i>br.</i>) small-flowered.	S. Europe.	1570.	—	F. ½.	Barr. ic. t. 368.
33	saxatilis. w.	(<i>br.</i>) stone.	—	1640.	7. 8.	F. ½.	Schk. handb. 3. t. 241.
34	rupéstris. w.	(<i>ye.</i>) rock.	Arabia.	1790.	—	G. ½.	Schmidel ic. t. 36.
35	Tenórii. L.en.	(<i>br.</i>) Tenore's.	S. Europe.	1824.	—	F. ½.	
36	sericea. w.	(<i>ye.</i>) snowy.	Canaries.	1779.	—	G. ½.	
37	geminiflora. L.en.	(<i>br.</i>) twin-flowered.	S. Europe.	1824.	—	F. ½.	
38	inuloides. w.	(<i>pu.</i>) cluster-flower'd.	Teneriffe.	1780.	7. 8.	G. ½.	Jacq. ic. 1. t. 171.
39	odorata. w.	(<i>pu.</i>) sweet-scented.	India.	1759.	6. 8.	S. ½.	Plum. ic. t. 97.
40	incisa. w.	(<i>pu.</i>) cut-leaved.	C. B. S.	1774.	—	G. ½.	
41	carolinensis. w.	(<i>pu.</i>) Carolina.	Carolina.	1823.	—	G. ½.	Jacq. ic. 3. t. 185.
42	virgata. w.	(<i>pu.</i>) wing-stalked.	America.	1783.	8. 9.	G. ¼.	Sloan.hist.1.t.152.f.5.
43	rugosa. H.K.	(<i>br.</i>) St. Helena.	St. Helena.	1772.	11.	G. ½.	
	CARP'ESIUM. W.	CARE'ESIUM.	Syngenesia	Polygamia	Superflua. L.		
1	cérnum. w.	(<i>ye.</i>) drooping.	Austria.	1739.	7. 8.	H. ¼.	Jacq. aust. 3. t. 204.
2	abrotanoïdes. w.	(<i>ye.</i>) Southernwood	like. China.	1768.	—	H. ¼.	Osbeck it. t. 10.
3	torulosum. F.	(<i>ye.</i>) twisted.	Siberia.	1823.	6. 8.	H. ¼.	
4	Wulfeniànum. Schreb.	(<i>ye.</i>) Wulfen's.	—	1829.	—	H. ¾.	
	I'NULA. W.	I'NULA.	Syngenesia	Polygamia	Superflua. L.		
1	Helénium. w.	(<i>ye.</i>) Elecampane.	Britain.	7. 8.	H. ¼.	
2	O'culus-Chrísti.w.	(<i>ye.</i>) hoary.	Austria.	1759.	7. 9.	H. ¼.	Eng. bot. v. 22. t. 1546.
3	británica. w.	(<i>ye.</i>) creeping-rooted.	Germany.	—	—	H. ¼.	Jacq. aust. 3. t. 223.
4	undulata. w.	(<i>ye.</i>) wave-leaved.	Egypt.	1739.	7. 10.	H. ○.	Flor. dan. t. 413.
5	índica. w.	(<i>ye.</i>) Indian.	E. Indies.	—	—	S. ○.	Burm. zeyl. t. 55. f. 2.
6	paludosa. s.s.	(<i>ye.</i>) marsh.	Portugal.	1828.	—	H. ¼.	
7	áspera. s.s.	(<i>ye.</i>) rough.	—	—	—	H. ¼.	
8	campéstris. s.s.	(<i>ye.</i>) field.	Poland.	1827.	—	H. ¼.	
9	squarrosa. w.	(<i>ye.</i>) net-leaved.	S. Europe.	1768.	7. 9.	H. ¼.	Flor. græc. t. 875.
10	viscosa. w.	(<i>ye.</i>) clammy.	Barbary.	1633.	7. 8.	G. ¼.	Jacq. vind. 2. t. 165.
	Erigeron viscidum. Jacq.						
11	tuberosa. P.S.	(<i>ye.</i>) tuberous-root'd.	S. Europe.	1640.	—	H. ¼.	Moris. s. 7. t. 19. f. 20.
	Erigeron tuberosum. W.						
12	salicina. w.	(<i>ye.</i>) willow-leaved.	N. Europe.	1648.	8. 9.	H. ¼.	Flor. dan. t. 786.
13	Bubónium. w.	(<i>ye.</i>) Anstrian.	Austria.	1801.	7. 9.	H. ¼.	Jacq. aust. 5. ap. 4. t. 378.
14	hírta. w.	(<i>ye.</i>) hairy.	—	1759.	6. 9.	H. ¼.	Bot. mag. t. 1907.
15	glandulosa. w.	(<i>ye.</i>) glandular.	Georgia.	1804.	7. 8.	H. ¼.	
16	grandiflora. w.	(<i>ye.</i>) large-flowered.	Caucasus.	1825.	—	H. ¼.	
17	móllis. L.en.	(<i>ye.</i>) soft.	—	1823.	—	H. ¼.	
18	suavéolens. w.	(<i>ye.</i>) woolly-leaved.	S. Europe.	1758.	6. 8.	H. ¼.	Jacq. vind. 3. t. 31.
19	Vaillantii. w.	(<i>ye.</i>) Vaillant's.	France.	1739.	—	H. ¼.	Hall. helv. 1. t. 13.
20	germánica. w.	(<i>ye.</i>) German.	Germany.	1759.	6. 7.	H. ¼.	Jacq. aust. 3. t. 184.
21	ensifolia. w.	(<i>ye.</i>) sword-leaved.	Austria.	1793.	7. 9.	H. ¼.	—
22	saxátilis. s.s.	(<i>ye.</i>) rock.	S. Europe.	1825.	—	H. ¼.	—
	Erigeron glutinosum. L.						
23	critmoides. L.	(<i>ye.</i>) Samphire-leav'd.	England.	—	8. 9.	H. ¼.	F. 1. t. 1546.
24	provincialis. w.	(<i>ye.</i>) oval-leaved.	France.	1778.	7. 8.	H. ¼.	
25	montaña. w.	(<i>ye.</i>) mountain.	S. Europe.	1759.	—	H. ¼.	
26	bifróns. w.	(<i>ye.</i>) Italian.	—	1713.	6. 8.	H. ¼.	
27	thapsoides. L.en.	(<i>ye.</i>) Mullein-like.	—	1823.	—	H. ¼.	

28	verbascifolia. s.s. (ye.)	Mullein-leaved.	Caucasus.	1828.	6. 8. H. 4.		
29	calycina. s.s. (ye.)	reflex-scaled.	Italy.	1829.	— H. 4.		
30	saturejoides. w. (ye.)	Savory-leaved.	Vera Cruz.	1733.	... S. 4.	Reliq. houst. 8. t. 19.	
31	fœtida. w. (ye.)	stinking.	Malta.	1688.	6. 8. H. ⊖.	Bocc. sic. 26. t. 13.	
32	odora. w. (ye.)	sweet-scented.	S. Europe.	1824.	— H. 4.	Moris. s. 7. t. 21. f. 6.	
33	hybrida. s.s. (ye.)	hybrid.	Poland.	1828.	— H. 4.		
34	glabra. M.B. (ye.)	smooth.	Caucasus.	1826.	— H. 4.		
35	caspica. F. (ye.)	Caspian.	Volga.	—	— H. 4.		
PULICARIA. G.							
	FLEAWORT.		Syngenesia Polygamia Superflua.				
1	vulgaria. g. (ye.)	small.	England.	8. 9. H. ⊖.	Gært. sem. 2. t. 173. f. 7.	
	<i>Inula Pulicaria.</i>		Eng. bot. v. 17. t. 1196.				
2	villosa. Len. (ye.)	villous.	—	1821.	— H. ⊖.		
3	arábica. L.en. (ye.)	Arabian.	Arabia.	1823.	— F. 4.	Pluk. alm. t. 149. f. 4.	
4	dysentérica. L.en. (ye.)	meadow.	England.	— H. 4.	Eng. bot. v. 16. t. 1115.	
	<i>Inula dysentérica. E.B.</i>						
CHRYSO'PSIS. N.							
1	gossypina. n. (ye.)	cottony.	N. America.	1823.	6. 8. H. 4.		
	<i>Inula gossypina. M.</i>						
2	trichophylla. n. (ye.)	hairy-leaved.	—	1827.	— H. 4.		
3	mariána. n. (ye.)	American.	—	1742.	7. 8. H. 4.	Mill. ic. 1. t. 57. f. 1.	
IX'ODIA. H.K.							
	IX'ODIA.		Syngenesia Polygamia Äqualis.				
	achillæoides. H.K. (wh.)	Milfoil-like.	N. Holland.	1803.	3. 9. G. ȳ.	Bot. mag. t. 1534.	
H'UMEA. Sm.							
	H'UMEA.		Syngenesia Polygamia Äqualis.				
	élegans. H.K. (re.)	elegant.	N. S. W.	1800.	6. 10. G. ȳ.	Exot. bot. 1. t. 1.	
CASSINIA. L.T.							
	CASSINIA.		Syngenesia Polygamia Äqualis.				
1	spectabilis. L.T. (st.)	Humea-like.	N. Holland.	1821.	5. 10. G. ȳ.	Bot. reg. t. 678.	
2	affinis. L.T. (st.)	related.	N. S. W.	1823.	4. 6. G. ȳ.	Bot. reg. t. 764.	
3	aurea. L.T. (ye.)	golden.	—	1821.	— G. ȳ.		
4	longifolia. L.T. (st.)	long-leaved.	—	—	— G. ȳ.		
5	denticulata. L.T. (st.)	tooth-leaved.	—	1826.	6. 8. G. ȳ.		
6	leptophylla. L.T. (wh.)	small-leaved.	N. Zealand.	1821.	4. 6. G. ȳ.		
OZOTHAMNUS. L.T. OzOTHAMNUS.							
1	cinereus. L.T. (ye.)	rusty.	V. Diem. Isl.	1820.	4. 8. G. ȳ.	Lab. nov. hol. v. 2. t. 182.	
	<i>Chrysócoma cinerea. Lab.</i>						
2	rosmarinifolius. L.T. (ye.)	Rosemary-leav'd.	—	1822.	— G. ȳ.	Lab. nov. hol. v. 2. t. 181.	
	<i>Eupatorium rosmarinifolium. Lab.</i>						
3	ferrugineus. L.T. (ye.)	ferruginous.	—	—	— G. ȳ.	Lab. nov. hol. v. 2. t. 180.	
	<i>Eupatorium ferrugineum. Lab.</i>						
AMMOBIUM. Br.							
	AMM'OBIA.		Syngenesia Polygamia Äqualis.				
	alatum. B.M. (wh.)	winged-stalk'd.	N. S. W.	1823.	8. 9. F. 4.	Swt. br. fl. gar. t. 48.	
ANTENNARIA. L.T. ANTENN'ARIA.							
1	dioica. L.T. (ro.)	red-flowered.	Britain.	5. 7. H. 4.	Eng. bot. v. 4. t. 267.	
	<i>Gnaphalium dioicum. E.B.</i>						
2	alpina. L.T. (wh.)	Alpine.	Alps Europe.	1775.	6. 7. H. 4.		
3	carpatica. L.T. (wh.)	Carpathian.	Carp. mount.	—	— H. 4.	Flor. dan. t. 332.	
	<i>Gnaphalium alpinum. w. non L.</i>						
4	plantaginifolia. L.T. (wh.)	Plantain-leav'd.	Virginia.	1759.	6. 7. H. 4.	Pluk. alm. t. 348. f. 9.	
	<i>Gnaphalium plantaginifolium. w.</i>						
5	margaritacea. L.T. (wh.)	pearly.	England.	7. 9. H. 4.	Eng. bot. v. 29. t. 2018.	
6	tripinnervis. D.P. (wh.)	triple nerved.	Nepaul.	1823.	6. 9. F. 4.	Bot. mag. t. 2468.	
7	contorta. D.P. (wh.)	twisted-leaved.	—	1818.	— F. 4.	Bot. reg. t. 605.	
LEONTOP'ODIUM. L.T. LION'S-FOOT.							
1	alpinum. D.D. (wh.)	Swiss.	Switzerland.	1776.	6. 7. H. 4.	Bot. mag. t. 1958.	
	<i>Gnaphalium Leontopodium. B.M.</i>						
2	sibiricum. D.D. (wh.)	Siberian.	Siberia.	1829.	— H. 4.		
GNAPH'ALIUM. L.T. EVERLASTING.							
	GNAPH'ALIUM.		Syngenesia Polygamia Superflua.	L.			
1	luteo-álbum. w. (st.)	Jersey.	Jersey.	7. 8. H. ⊖.	Eng. bot. v. 14. t. 1002.	
2	albescens. w. (wh.)	white Jamaica.	Jamaica.	1793.	... S. ȳ.		
3	sanguineum. w. (re.)	bloody.	Egypt.	1768.	5. 7. H. 4.	Rauw. it. 285. t. 37.	
4	undulatum. w. (wh.)	wave-leaved.	Africa.	1732.	6. 9. H. ⊖.	Dill. elth. t. 108. f. 130.	
5	obtusifolium. w. (wh.)	blunt-leaved.	N. America.	1699.	7. 9. H. ⊖.	— t. 108. f. 131.	
6	purpureum. w. (pu.)	purple-flower'd.	S. America.	1732.	6. 9. G. ȳ.	— t. 109. f. 132.	
7	involucratum. w. (gr.)	involucrated.	N. Zealand.	1823.	7. 8. G. 4.	Bot. mag. t. 2582.	
8	americánum. w. (br.)	Jamaica.	Jamaica.	1815.	... S. ȳ.		
9	pensylvanicum. w. (st.)	American.	N. America.	—	7. 9. H. ⊖.		
10	sylvaticum. w. (st.)	highland.	Britannia.	8. 9. H. 4.	Eng. bot. v. 13. t. 913.	
11	récum. w. (br.)	upright wood.	—	—	— H. 4.	— v. 2. t. 124.	
12	coarctatum. w. (gr.)	crowded.	Chili.	1820.	— H. 4.		
13	polycéphalum. m. (st.)	many-headed.	N. America.	1826.	— H. ⊖.		

14	stachydisfölium.s.s.(br.)	Stachy's-leaved.	MonteVideo.	1826.	8. 9. H. ♂.	
15	supinum. w.	(br.) dwarf.	Scotland.	6. 7. H. ♀.	Eng. bot. v. 17. t. 1193.
16	pūsillum. w.	(br.) small.	Sweden.	1818.	— H. ♀.	Krock. siles. t. 41.
17	sphæricum.W.en.(br.)	spherical.	S. Europe.	—	— H. 2.	
18	cephaloideum.w.(br.)	headed.	—	—	— H. ♂.	
19	uliginosum. w.	(st.) marsh.	Britain.	8. 9. H. ♂.	Eng. bot. v. 17. t. 1194.
20	germánicum. w.	(st.) common.	—	—	6. 8. H. ♂.	— v. 14. t. 948.
21	gállum. w.	(st.) narrow-leaved.	England.	— H. ♂.	— v. 33. t. 2369.
22	montanum. w.	(st.) mountain.	Europe.	1820.	— H. ♂.	
23	pyramidalatum. w.(st.)	pyramidal.	S. Europe.	1779.	— H. ♂.	
24	mínimum. w.	(st.) least.	Britain.	7. 8. H. ♂.	Eng. bot. v. 17. t. 1157.
25	arvénse. w.	(st.) corn.	Europe.	1804.	— H. ♂.	
26	austrâle. f.	(st.) southern.	S. Europe.	1823.	— H. ♂.	
27	Sellówii. Lk.	(st.) Sellow's.	BuenosAyres.	1828.	— H. ♂.	
28	alátum. k.s.	(st.) wing-stalked.	Quito.	1829.	— G. ♀.	
29	pompeyânun. E.	(st.) Egyptian.	Egypt.	—	— H. ♂.	
30	Lágopus. s.s.	(wh.) Hare's-foot.	Siberia.	1820.	— H. ♂.	
METAL'ASIA. L.T. METAL'ASIA.						
			Syngenesia	Polygamia	Superflua.	
1	umbellata. D.D.	(ro.) umbelled.	C. B. S.	1816.	5. 7. G. ♂.	
2	muricata. D.D.	(ro.) muricate.	—	—	6. 9. G. ♂.	Burm. afr. t. 72. f. 2.
3	divergens. D.D.	(wh.) spreading.	—	—	— G. ♂.	
	<i>Gnaphálium divergens. w.</i>					
4	fastigiata.	(wh.) close-flowered.	—	1812.	5. 8. G. ♂.	
	<i>Gnaphálium fastigiátum. w.</i>					
5	aurea. D.D.	(ye.) golden.	—	1816.	— G. ♂.	
6	seriphioïdes.D.D.(wh.)	Seriphium-like.	—	1825.	— G. ♂.	
7	muconatá. L.T.	(ye.) muconate.	—	1824.	— G. ♂.	Burm. afr. t. 66. f. 3.
8	pungens. D.D.	(wh.) pungent.	—	1815.	— G. ♂.	
9	uniflora. D.D.	(wh.) single-flowered.	—	1816.	— G. ♂.	
10	phylicoïdes. D.D.(wh.)	Phyllica-leaved.	—	—	— G. ♂.	
SPIRAL'EPIS. D.D. SPIRAL'EPIS.						
			Syngenesia	Polygamia	Superflua.	
1	squarrösa.D.D.(pu.wh.)	squarrose.	C. B. S.	1816.	7. 10. G. ♀.	
	<i>Gnaphálium squarrösum. Jacq.</i>					
2	glomerata. D.D.	(wh.) cluster-flowered.	—	1774.	3. 9. G. ♀.	
3	tincta. D.D.	(bh.) stained.	—	1816.	— G. ♀.	
4?	declinata.	(wh.) creeping.	—	1787.	7. 9. G. ♀.	
	<i>Gnaphálium declinátum. w.</i>					
PETALA'CTE. D.D. PETALA'CTE.						
			Syngenesia	Polygamia	Superflua.	
1	coronata. D.D.	(wh.) crowned.	C. B. S.	1816.	5. 7. G. ♂.	
	<i>Gnaphálium coronátum. w.</i>					
2	bicolor. D.D.	(pu.wh.) two-coloured.	—	—	— G. ♂.	Burm. afr. t. 69. f. 3.
PENTATA'XIS. D.D. PENTATA'XIS.						
			Syngenesia	Polygamia	Superflua.	
	micrantha. D.D.	(wh.) small-flowered.	C. B. S.	1821.	5. 7. G. ♂.	
ASTEL'MA. Br. ASTEL'MA.						
			Syngenesia	Polygamia	Superflua.	
1	eximum. B.R.	(re.) giant.	C. B. S.	1793.	6. 9. G. ♂.	
	<i>Gnaphálium exínum. A.R. 654.</i>					
2	milleflörum.D.D.(wh.)	many-flowered.	—	1802.	6. 9. G. ♂.	
	<i>Gnaphálium milleflòrum. w.</i>					
3	Staelhelina. D.D.	(ye.) keel-leaved.	—	1801.	12. 3. G. ♂.	
4	crassifolium. D.D.(ye.)	thick-leaved.	—	1816.	3. 5. G. ♂.	Andr. deposit. t. 428.
5	protoïdes. D.D.(wh.)	Protea-like.	—	1821.	— G. ♂.	
6	speciosissimum. D.D.(wh.)	showy.	—	1691.	7. 9. G. ♂.	Andr. deposit. t. 51.
	<i>Xeránthemum speciosissimum. A.R. Elichr'ysum speciosissimum. w.</i>					
7	lineare. D.D.	(wh.) linear-leaved.	C. B. S.	1816.	7. 9. G. ♂.	
8	spiralis. D.D.	(re.wh.) spiral-leaved.	—	1801.	7. 10. G. ♂.	Andr. deposit. t. 262.
	<i>Elichr'ysum spirale. w.</i>					
9	variegatum. D.D.	(v.) large globular-fld.	—	—	5. 6. G. ♂.	
10	imbricatum. D.D.(ro.)	imbricated.	—	1816.	— G. ♂.	
	<i>Elichr'ysum imbricatum. w. Xeránthemum imbricatum. L.</i>					
11	candescens. D.D.	(ro.) elegant.	C. B. S.	1794.	8. 4. G. ♂.	
12	modestum. Sie.	(fl.) modest.	—	1824.	6. 7. G. ♂.	Bot. mag. t. 420.
	<i>Gnaphálium modéstum. B.M.</i>					
HELICHR'YSUM. P.S. HELICHR'YSUM.						
			Syngenesia	Polygamia	Superflua.	
1	grandiflorum.D.D.(w.)	great-flowered.	C. B. S.	1731.	6. 8. G. ♂.	
	<i>Gnaphálium grandiflorum. w.</i>					
2	fruticans. D.D.	(wh.) frutescent.	—	1779.	— G. ♂.	Bot. mag. t. 1802.
	<i>Astélma frúticans. B.R. 726. Gnaphálium frúticans. B.M. grandiflorum. A.R. 489.</i>					
3	crispum. D.D.	(re.) curled.	C. B. S.	1809.	— G. ♂.	
4	arboreum. D.D.	(re.) shrubby.	—	1770.	2. 8. G. ♂.	Bot. mag. t. 2710.

5	congéstum. D.D. (<i>fl.</i>) close-headed.	C. B. S.	1791.	5. 6. G. ḥ.	Bot. reg. t. 243.
	<i>Gnaphálium congéstum.</i> B.R.				
6	cárneum. P.S. (<i>fl.</i>) flesh-coloured.	——	1816.	4. 6. G. ḥ.	
7	pátulum. D.D. (<i>wh.</i>) spreading.	——	1771.	8. 1. G. ḥ.	
8	discolor. D.D. (<i>wh.fl.</i>) two-coloured.	——	1815.	5. 8. G. ḥ.	Burm. afr. t. 97. f. 4.
9	cephalòtes. D.D. (<i>wh.re.</i>) large-headed.	——	1789.	11. 1. G. ḥ.	Pluk. phyt. t. 410. f. 2.
10?	diosmæfólium. (<i>wh.</i>) Diosma-leaved.	——	1812.	3. 8. G. ḥ.	Vent. malm. t. 74.
	<i>Gnaphálium diosmæfólium.</i> v.				
11	ericoides. D.D. (<i>pa.</i>) Heath-leaved.	——	1774.	— G. ḥ.	Bot. mag. t. 435.
	<i>Gnaphálium ericoídes.</i> B.M. non <i>Helichrysum ericoídes.</i> P.S.				
12	teretifólium. D.D. (<i>wh.</i>) round-leaved.	C. B. S.	1812.	— G. ḥ.	Burm. afr. t. 77. f. 2.
13?	tephródes. (<i>wh.</i>) slender-leaved.	——	1816.	— G. ḥ.	
	<i>Gnaphálium tephródes.</i> L.en.				
14	acuminátum. (<i>wh.</i>) taper-pointed.	N. Holland.	—	— G. ḥ.	
	<i>Gnaphálium acuminátum.</i> L.en.				
15?	lasiocáulon. (<i>wh.</i>) woolly-stalked.	C. B. S.	1823.	6. 9. G. ḥ.	
	<i>Gnaphálium lasiocáulon.</i> L.en.				
16	St'echas. D.D. (<i>ye.</i>) common shrubby.	S. Europe.	1629.	6. 10. H. ḥ.	Barrel. ic. 410.
17	angustifólium. (<i>ye.</i>) narrow-leaved.	——	1818.	— H. ḥ.	
	<i>Gnaphálium angustifólium.</i> P.S.				
18	cónicum. (<i>ye.</i>) conical.	——	1824.	— H. ḥ.	
	<i>Gnaphálium cónicum.</i> L.en.				
19	ignéscens. D.D. (<i>re.</i>) red-flowered.	C. B. S.	1731.	6. 10. G. ḥ.	
20	microphyllum. D.D. (<i>ye.</i>) small-leaved.	Crete.	1823.	— G. ḥ.	
21?	dasyánthum. (<i>ye.</i>) hairy-flowered.	C. B. S.	1812.	— G. ḥ.	
	<i>Gnaphálium dasyánthum.</i> W.en.				
22	crassifólium. D.D. (<i>ye.</i>) thick-leaved.	S. Europe.	1774.	7. 9. G. ḥ.	
23	marítimum. D.D. (<i>ye.</i>) sea-side.	C. B. S.	1772.	6. 8. G. ḥ.	
24	apiculátum. D.D. (<i>ye.</i>) New Holland.	V. Diem. Isl.	1804.	1. 12. G. ḥ.	
	<i>Gnaphálium apiculátum.</i> B.R.				
25	f'atidum. D.D. (<i>ye.</i>) strong-scented.	C. B. S.	1692.	6. 9. G. ḡ.	Bot. mag. t. 1987.
	<i>Gnaphálium f'atidum.</i> B.M.				
26	cylindricum. D.D. (<i>ye.</i>) cylindrical.	——	1780.	6. 8. G. ȝ.	Pluk. phyt. 298. f. 4.
27	orientále. D.D. (<i>ye.</i>) eastern.	Africa.	1629.	4. 8. G. ḥ.	Comm. hort. 2. t. 55.
28	arenárium. D.D. (<i>ye.</i>) sand.	Europe.	1739.	7. 9. H. ȝ.	
29	gravéolens. (<i>ye.</i>) strong-smelling.	Tauria.	1819.	— H. ȝ.	Bot. mag. t. 2159.
	<i>Gnaphálium gravéolens.</i> M.B.				
30	odoratissimum. (<i>ye.</i>) sweet-scented.	C. B. S.	1691.	4. 8. G. ḥ.	Mill. ic. 1. t. 131. f. 2.
31	candidissimum. D.D. (<i>ye.</i>) white-leaved.	Caucasus.	1819.	7. 9. H. ȝ.	Dill. elt. t. 107. f. 127.
32	rútilans. D.D. (<i>re.</i>) shining-flower'd.	C. B. S.	1731.	6. 7. G. ȝ.	—— t. 107. f. 128.
33	cymosum. D.D. (<i>ye.wh.</i>) branching.	Africa.	——	4. 8. G. ḥ.	Volck. norib. t. 194.
34	helianthemifólium. D. (<i>wh.</i>) Sun-rose-leav'd.	C. B. S.	1774.	7. 10. G. ḥ.	Andr. deposit. 375
35	bracteátem. D.D. (<i>ye.</i>) wave-leaved.	N. Holland.	1799.	— H. ⓧ.	
	<i>chrysánthum.</i> P.S. <i>Xeránthemum bracteátem.</i> v. M. t. 2. <i>lúcidum.</i> Maunde.				
36	fulgidum. P.S. (<i>ye.</i>) great yellow.	C. B. S.	1774.	2. 10. G. ȝ.	
37	dealbátum. P.S. (<i>bh.</i>) herbaceous.	V. Diem. Isl.	1812.	1. 12. G. ȝ.	
38	paniculátum. P.S. (<i>wh.</i>) corymb-flower'd.	C. B. S.	1800.	6. 9. G. ḥ.	
	<i>argénteum.</i> A.R. 552. nec <i>alivorum.</i>				
39	argénteum. P.S. (<i>wh.</i>) silvery.	——	—	— G. ḥ.	
40	frágrans. A.R. (<i>ro.</i>) fragrant.	——	1803.	7. 9. G. ḥ.	
	<i>Elichrysum frágrans.</i> A.R.				
41	retortum. w. (<i>wh.</i>) trailing.	——	1732.	7. 8. G. ḥ.	Dill. elth. t. 322. f. 415.
42?	incánum. B.M. (<i>ro.wh.</i>) hoary-leaved.	V. Diem. Isl.	1828.	5. 6. G. ȝ.	Bot. mag. t. 2881.
43	rígidum. A.R. (<i>wh.</i>) rigid-leaved.	C. B. S.	1801.	— G. ḥ.	Andr. deposit. t. 387.
44	herbáceum. A.R. (<i>ye.</i>) shining-flowered.	——	1802.	7. 9. G. ȝ.	—— t. 487.
	<i>spléndens.</i> B.M. 1773.				
LEUCOSTE'MMA. D.D. LEUCOSTE'MMA. Syngenesia Polygamia Superflua.					
1	vestitum. D.D. (<i>wh.</i>) upright.	C. B. S.	1774.	7. 9. G. ḥ.	Burm. afr. t. 66. f. 1.
	<i>Elichrysum vestitum.</i> w.				
2	lingulátum. D.D. (<i>wh.</i>) tongue-leaved.	——	1812.	— G. ḥ.	
PHÉNO'COMA. D.D. PHÉNO'COMA. Syngenesia Polygamia Superflua.					
	<i>prolífera.</i> D.D. (<i>pu.</i>) proliferous.	C. B. S.	1789.	5. 11. G. ḥ.	
	<i>Elichrysum prolífera.</i> B.R. <i>Xeránthemum prolífera.</i> L.				
APHELE'XIS. D.D. APHELE'XIS. Syngenesia Polygamia Superflua.					
1	hùmilis. D.D. (<i>pu.</i>) dwarf.	C. B. S.	1812.	6. 9. G. ḥ.	Andr. deposit. t. 664.
	<i>Elichrysum spectáabile.</i> B.C. 59. <i>pinifólium.</i> P.S.				
2	sesamoídes. D.D. (<i>pu.</i>) superb.	C. B. S.	1739.	4. 6. G. ḥ.	Bot. mag. t. 425..
	<i>Elichrysum sesamoídes.</i> w. <i>Xeránthemum sesamoídes.</i> B.M.				

COMPOSITÆ.

3 fasciculata. D.D. (<i>vu.</i>) bundle-leaved.	C. B. S.	1799.	3. 9. G. ḥ.	Andr. deposit. t. 242.
<i>a lutescens.</i> (<i>st.</i>) yellowish.	—	G. ḥ.	
<i>β versicolor.</i> (<i>pu.wh.</i>) variegated-flower'd.	—	G. ḥ.	Breyne. ic. t. 16. f. 2.
<i>γ alba.</i> (<i>wh.</i>) white-flowered.	—	G. ḥ.	Andr. deposit. t. 279.
<i>δ rubra.</i> (<i>sc.</i>) red-flowered.	—	G. ḥ.	— t. 650.
4 ericoïdes. (<i>bh.</i>) filiform.	1796.	— G. ḥ.	Lam. ill. t. 693. f. 2.
<i>filiformis.</i> D.D. <i>Helichrysum ericoïdes.</i> P.S.				
XERA'NTHEMUM. D.D. XERA'NTHEMUM.	Syngenesia Polygamia Superflua.	L.		
1 ánnuum. D.D. (<i>pu.wh.</i>) common annual.	S. Europe.	1570.	7. 8. H. ○.	Jacq. aust. 4. t. 388.
<i>a rösem.</i> (<i>ro.</i>) rose-coloured.	—	H. ○.	
<i>β album.</i> (<i>wh.</i>) white-flowered.	—	H. ○.	
2 inapertum. D.D. (<i>pu.</i>) small-flowered.	1620.	— — H. ○.	Moris. his. s. 6. t. 12. f. 1.
CHARDI'NIA. D.D. CHARDI'NIA.	Syngenesia Polygamia Superflua.			
orientalis. D.D. (<i>wh.</i>) oriental.	Levant.	1713.	7. 8. H. ○.	
<i>xeranthemoïdes.</i> Desf. <i>Xeranthemum orientale.</i> w.				
FILAGO. W.	COTTON-ROSE. Syngenesia Polygamia Necessaria. W.			
1 pygmæa. w. (<i>br.</i>) pigmy.	S. Europe.	1629.	7. 8. H. ○.	Cavan. ic. 1. t. 36.
' <i>Eva</i> x pygmæa. G.				
2 asterisciflora. (<i>br.</i>) simple-stalked.	Spain.	1824.	— — H. ○.	Lam. ill. t. 694. f. 2.
' <i>Eva</i> x asterisciflora. P.S.				
MICR'OPUS. W.	MICR'OPUS. Syngenesia Polygamia Necessaria. W.			
1 supinus. w. (<i>pa.</i>) trailing.	S. Europe.	1710.	6. 9. H. ○.	Schk. hand. 3. t. 267.
2 erectus. w. (<i>pa.</i>) upright.	1683.	— — H. ○.	Lœfl. hisp. t 1. f. 3.
ERI'GERON. W.	ERI'GERON. Syngenesia Polygamia Superflua. L.			
1 graveolens. w. (<i>ye.</i>) strong-smelling.	S. Europe.	1633.	7. 8. H. ○.	
2 compòsitum. Ph. (<i>li.</i>) daisy-flowered.	N. America.	1811.	— — H. 4.	
3 caroliniànum. w. (<i>ye.</i>) Hyssop-leaved.	1727.	— — H. 4.	
4 canadense. w. (<i>wh.</i>) Canada.	England.	8. 9. H. ○.	Dill. elt. t. 306. f. 394.
5 bonariense. w. (<i>wh.</i>) buck's-horn.	S. America.	1732.	7. 8. H. ○.	Eng. bot. v. 29. t. 2019.
6 linifolium. w. (<i>wh.</i>) flax-leaved.	— — H. ○.	Dill. elt. t. 257. f. 334.
7 lavigatūm. L.en. (<i>wh.</i>) smooth.	Cayenne.	1823.	— — S. ♂.	
8 delphinifolium. w. (<i>wh.</i>) Larkspur-leav'd.	S. America.	1816.	— — S. ♂.	
9 strigosum. w. (<i>wh.</i>) strigose.	N. America.	1818.	— — H. ♂.	
10 heterophyllosum. w. (<i>wh.</i>) various-leaved.	1640.	7. 9. H. ♂.	Flor. græc. t. 486.
<i>Aster annuus.</i> L. <i>Diplopappus dubius.</i> Cas.				
11 chinense. w. (<i>wh.</i>) Chinese.	China.	1818.	— — H. ○.	Jacq. schœnb. 3. t. 303.
12 jamaicense. w. (<i>li.</i>) Jamaica.	Jamaica.	1820.	— — S. ○.	Swartz obs. t. 8. f. 2.
13 pubescens. k.s. (<i>wh.</i>) pubescent.	Mexico.	1827.	7. 11. H. 4.	
14 involucratum. Lk. involucred.	Brazil.	1828. S. 4.	
15 armerifolium. F. (<i>pu.</i>) Thrift-leaved.	Baical.	1829.	6. 8. H. 4.	
16 nudicaulé. M. (<i>wh.</i>) naked-stalked.	N. America.	1812.	7. H. 4.	
17 philadelphicum. w. (<i>pu.</i>) spreading.	1778.	7. 8. H. 4.	
18 purpureum. w. (<i>pu.</i>) purple.	Hudson's Bay.	1776.	— — H. 4.	
19 bellidifolium. w. (<i>li.</i>) Plantain-leav'd.	N. America.	1790.	— — H. 4.	
<i>serpentarium.</i> Hort.				
20 glabellum. N. (<i>pu.</i>) smoothish.	1827.	8. 12. H. 4.	Bot. mag. t. 2923.
21 podolicum. s.s. (<i>pu.</i>) Polish.	Poland.	—	7. 9. H. 4.	
22 Lehmanni. s.s. (<i>li.</i>) Lehman's.	1828.	— — H. 4.	
23 glaucum. B.R. (<i>li.</i>) glaucous-leav'd.	S. America.	1812.	1. 12. G. ḥ.	Bot. reg. t. 10.
24 chilense. D.P. (<i>wh.</i>) Chili.	Chili.	1816.	7. 9. H. ○.	
<i>Conyzza chilensis.</i> L.en.				
25 rivulare. s.s. (<i>bh.</i>) rivulet.	Trinidad.	1821.	— — S. ○.	
26 Villarsii. w. (<i>pu.</i>) Villars's.	Piedmont.	1804.	7. 8. H. 4.	Bot. reg. t. 583.
<i>atticum.</i> Vil.				
27 rupestre. Schl. (<i>li.</i>) rock.	Switzerland.	1818.	— — H. 4.	
28 acré. w. (<i>li.</i>) common.	Britain.	— — H. ♂.	Eng. bot. v. 17. t. 1158.
29 alpinum. w. (<i>li.</i>) Alpine.	Scotland.	7. H. 4.	— v. 7. t. 464.
30 uniflorum. w. (<i>li.</i>) dwarf.	8. 9. H. 4.	— v. 34. t. 2416.
31 pumilum. N. (<i>wh.</i>) dwarf.	N. America.	1827.	— — H. 4.	
32 humile. Gr. (<i>wh.</i>) low.	—	9. 10. H. ḥ.	
33 asperum. N. (<i>wh.</i>) rough.	1828.	8. 9. H. 4.	
34 gramineum. w. (<i>pu.</i>) grass-leaved.	Siberia.	1824.	— — H. 4.	Gmel. sib. 2. t. 76. f. 2.
35 caucasicum. M.B. (<i>pu.</i>) Caucasian.	Caucasus.	—	7. 8. H. 4.	
36 asteroïdes. L.en. (<i>wh.</i>) Starwort-like.	1812.	— — H. 4.	
JAS'ONIA. Cas.	JAS'ONIA.	Syngenesia Polygamia Superflua.		
longifolia. Cas. (<i>ye.</i>) long-leaved.	N. America.	1818.	7. 8. H. 4.	
<i>Erigeron longifolium.</i> P.S.				
CAL'OTIS. B.R.	CAL'OTIS.	Syngenesia Polygamia Superflua.		
cuneifolia. B.R. (<i>bl.</i>) wedge-leaved.	N. Holland.	1819.	7. 8. G. ḥ.	Bot. reg. t. 504.

DIPLOSTE'PHIUM. K.S. DIPLOSTE'PHIUM. Syngenesia Polygamia Superflua.

1	linariifolium. (pu.)	toad-flax-leav'd. N. America.	1699.	9. 10. H. $\frac{1}{2}$.	Lodd. bot. cab. t. 6.
2	A'ster pulcherrimus. L.B.C. linariifolius. w.	<i>Chrysópsis linariifolia</i> . n.	1759.	8. 10. H. $\frac{1}{2}$.	Pluk. alm. t. 14. f. 7.
2	rígido. (vi.) stiff-leaved.	1759.	8. 10. H. $\frac{1}{2}$.	Pluk. alm. t. 14. f. 7.	
3	A'ster rígido. Ph. nemoralis. w. non H.K.	1759.	8. 10. H. $\frac{1}{2}$.	Pluk. alm. t. 14. f. 7.	
3	linifolium. (wh.) Flax-leaved.	N. America.	1739.	7. 8. H. $\frac{1}{2}$.	
	<i>Chrysópsis linifolia</i> . n. A'ster linifolius. w.				
4	ábum. N. (wh.) white-flowered. Missouri.	1824.	8. 9. H. $\frac{1}{2}$.		
5	húmile. (wh.) low.	N. America.	1699.	8. 10. H. $\frac{1}{2}$.	Willd. hort. ber. t. 67.
	<i>A'ster humilius</i> . w. <i>Chrysópsis humilius</i> . n.				
6	cornifolium. w. (wh.) Dogwood-leaved.	1811.	— H. $\frac{1}{2}$.		
7	amygdalinum. (wh.) Almond-leaved.	1759.	7. 9. H. $\frac{1}{2}$.	Hoff. phyt. blat. t. 8. f. 2.	
	<i>A'ster umbellatus</i> . w. <i>Chrysópsis umbellata</i> . n.				
A'STER. W.	STARWORT.	Syngenesia Polygamia Superflua. L.			
1	reflexus. w. (re.wh.) reflexed-leav'd. C. B. S.	1759.	2. 9. G. $\frac{1}{2}$.	Bot. mag. t. 884.	
2	tomentosus. w. (wh.) tooth-leaved. N. S. W.	1793.	5. 7. G. $\frac{1}{2}$.	Wend. s.han. p. 8. t. 24.	
	<i>dentatus</i> . Andr. rep. t. 61.				
3	stellulatus. P.S.	starry-haired. V. Diem. Isl.	1823.	— G. $\frac{1}{2}$.	Lab. nov. hol. 2. t. 196.
4	phlogopappus. P.S. (vi.)	glossy-haired.	1827.	— G. $\frac{1}{2}$.	— 2. t. 195.
5	myrsinoides. P.S. (wh.)	Myrsine-leaved.	1824.	— G. $\frac{1}{2}$.	— p. 53. t. 202.
6	erubescens. s.s. (bh.)	blush-flowered. N. S. W.	1826.	— G. $\frac{1}{2}$.	
7	argophyllus. P.S. (wh.)	musk-scented. V. Diem. Isl.	1804.	— G. $\frac{1}{2}$.	Bot. mag. t. 1563.
8	liratus. B.M.	(wh.) fluted-stemm'd. N. S. W.	1812.	— G. $\frac{1}{2}$.	— t. 1509.
9	sericeus. w. (bl.)	silky-leaved. Missouri.	1802.	5. 11. G. $\frac{1}{2}$.	Venten. cels. t. 33.
10	Cymbalariae. w. (wh.)	Ivy-leaved. C. B. S.	1786.	— G. $\frac{1}{2}$.	Venten. malm. t. 95.
11	papposus. s.s.	(vi.) cluster-leaved.	1812.	— G. $\frac{1}{2}$.	
12	taxifolius. s.s.	(li.) Yew-leaved.	1824.	— G. $\frac{1}{2}$.	Jacq. schoenb. 3. t. 370.
13	angustifolius. w.	(bl.) narrow-leaved.	1804.	5. 7. G. $\frac{1}{2}$.	
14	villus. w.	(li.) villous.	1812.	— G. $\frac{1}{2}$.	
15	obtusatus. w.	(li.) obtuse-leaved.	1793.	— G. $\frac{1}{2}$.	
16	fruticulosus. w.	(li.) shrubby.	1759.	3. 7. G. $\frac{1}{2}$.	Jacq. frag. 9. t. 5. f. 4.
17	filiolius. v.	(wh.) thread-leaved.	1812.	— G. $\frac{1}{2}$.	Venten. malm. t. 82.
18	aculeatus. P.S.	(wh.) prickly-leaved. exasperatus. L. enum.	1818.	— G. $\frac{1}{2}$.	Lab. nov. hol. 2. t. 200.
19	tenellus. w.	(vi.) slender.	C. B. S.	1769. 4. 10. H. \odot .	Bot. mag. t. 33.
20	alpinus. w.	(pu.) Alpine.	Alps Europe.	1658. 5. 8. H. $\frac{1}{2}$.	— t. 199.
21	lusitanus. s.s.	(pu.) Portugal.	Portugal.	1820. 8. 10. H. $\frac{1}{2}$.	
22	salsuginosus. s.s.	(vi.) salt plain.	N. America.	1827. 5. 6. H. $\frac{1}{2}$.	Bot. mag. t. 2942.
23	pulchellus. w.	(re.) narrow-rayed.	Armenia.	1818. — H. $\frac{1}{2}$.	
24	caucasicus. w.	(pu.) Caucasian.	Caucasus.	1804. 7. 8. H. $\frac{1}{2}$.	Bot. mag. t. 2321.
25	alwartensis. B.M.	(bh.) large-flowered.		... — H. $\frac{1}{2}$.	
26	diversifolius. M.	(li.) various-leaved.	N. America.	1811. 7. 9. H. $\frac{1}{2}$.	
27	macrophyllus. w. (bl.)	large-leaved.		1739. — H. $\frac{1}{2}$.	
28	corymbosus. w.	(wh.) corymbed.		1765. 9. H. $\frac{1}{2}$.	
29	cordifolius. w.	(bh.) heart-leaved.		1759. 7. 8. H. $\frac{1}{2}$.	
30	paniculatus. w.	(li.) panicled.		1640. 9. 10. H. $\frac{1}{2}$.	
31	sagittifolius. w.	(bl.) arrow-leaved.		1818. 7. 9. H. $\frac{1}{2}$.	Corn. canad. t. 65.
32	undulatus. w.	(pu.) wave-leaved.		1699. 8. 10. H. $\frac{1}{2}$.	Herm. parad. t. 96.
33	præcox. W.en.	(li.) early-flowering.		1800. 7. 8. H. $\frac{1}{2}$.	
34	pallens. W.en.	(li.) pale-flowered.		... 9. 10. H. $\frac{1}{2}$.	
35	conyzoides. w.	(wh.) Conzya-like.		1778. 8. 9. H. $\frac{1}{2}$.	
36	altaicus. W.en.	(pu.) dwarf. pùmilus. F.	Siberia.	1804. 5. 8. H. $\frac{1}{2}$.	
37	nemoralis. H.K.	(li.) wood. ledijolius. Ph.	N. America.	1778. 8. 9. H. $\frac{1}{2}$.	
38	hyssopifolius. w.	(bh.) Hyssop-leaved.		1683. 9. 10. H. $\frac{1}{2}$.	Dodart. mem. t. 60.
39	punctatus. w.	(vi.) dotted.	Hungary.	1815. 8. 9. H. $\frac{1}{2}$.	W. et K. hung. 2. t. 109.
40	acrìs. w.	(bl.) acrid.	S. Europe.	1731. — H. $\frac{1}{2}$.	Pluk. alm. t. 271. f. 3.
41	cánus. w.	(bl.) hoary-leaved.	Hungary.	1816. — H. $\frac{1}{2}$.	W. et K. hung. 1. t. 30.
42	solidaginoides. w.	(wh.) Solidago-like.	N. America.	1699. — H. $\frac{1}{2}$.	Pluk. alm. t. 79. f. 2.
43	subulatus. N.	(li.) awl-leaved.		1823. 8. 10. H. \odot .	
44	pilosus. w.	(wh.) hairy.		1812. — H. $\frac{1}{2}$.	
45	foliolosus. w.	(wh.) leafy.		1732. 10. H. $\frac{1}{2}$.	Dill. elth. t. 35. f. 39.
46	tenuifolius. w.	(wh.) slender-leaved.		1725. 9. 10. H. $\frac{1}{2}$.	Pluk. alm. t. 78. f. 5.
47	dumosus. w.	(wh.) bushy.		1734. — H. $\frac{1}{2}$.	Herm. parad. 95.
	α violaceus. H.K.	(vi.) violet-flowered.		— — H. $\frac{1}{2}$.	
	β ábus. H.K.	(wh.) white-flowered.		— — H. $\frac{1}{2}$.	
48	ericoides. w.	(wh.) heath-leaved.		1758. 9. H. $\frac{1}{2}$.	Dill. elth. t. 36. f. 40.
49	multiflorus. w.	(wh.) many-flowered.		1732. 9. 10. H. $\frac{1}{2}$.	
50	ciliatus. w.	(wh.) ciliated.		... — H. $\frac{1}{2}$.	

51 canéscens, Ph. biénnis, N.	(li.) canescent.	N. America.	1812.	— H. ȝ.
52 pauciflórus, N.	(wh.) few-flowered.	N. America.	1825.	8. 9. H. ȝ.
53 sparsiflórus, w.	(wh.) scattered-flower'd.	—	1798.	9. 11. H. ȝ.
54 coridifolius, w.	(bl.) Coris leaved.	—	—	8. 11. H. ȝ.
55 squarrósus, w.	(bl.) ragged.	—	1801.	— H. ȝ.
56 argénteus, H.K.	(bl.) silver-leaved.	—	—	7. 9. H. ȝ.
57 cóncolor, w.	(pu.) self-coloured.	—	1759.	8. 11. H. ȝ.
58 reticulátus, Ph.	(wh.) netted-leaved.	—	1812.	— H. ȝ.
59 salicifolius, w.	(pu.) Willow-leaved.	—	1760.	9. 10. H. ȝ.
60 astivus, w.	(bl.) summer.	—	1776.	7. 8. H. ȝ.
61 pannónicus, w.	(vi.) Hungarian.	Hungary.	1815.	— H. ȝ.
62 Amélus, w.	(bl.) Italian.	Italy.	1596.	7. 9. H. ȝ.
a latifolius.	(bl.) broad-leaved.	—	—	H. ȝ.
β angustifolius.	(bl.) narrow-leaved.	—	—	H. ȝ.
63 ibéricus, M.B.	(bl.) Iberian.	Iberia.	1827.	7. 9. H. ȝ.
64 nòvæ ángliæ, w.	(bl.) New England.	N. America.	1710.	9. 10. H. ȝ.
β rúbra.	(re.) red-flowered.	—	1812.	— H. ȝ.
65 cyáneus, N.	(bl.) bright-blue.	—	1789.	— H. ȝ.
66 gláucus, N.A.	(bl.) glaucous-leaved.	—	1823.	8. 10. H. ȝ.
67 paludósus, w.	(bl.) marsh.	N. America.	1784.	— H. ȝ.
68 grandiflorus, w.	(bl.) great-flowered.	—	1720.	10. 11. H. ȝ.
69 caroliniánus, w.	(pu.) tall.	Carolina.	—	8. 9. H. ȝ.
70 oblongifolius, N.	(li.) oblong-leaved.	Missouri.	1823.	— H. ȝ.
71 phlogifolius, w.	(vi.) Phlox-leaved.	N. America.	1797.	7. 10. H. ȝ.
72 pátem, w.	(bl.) spreading.	—	1773.	9. 11. H. ȝ.
73 Tripoliúm, w.	(bl.) sea-side.	Britain.	—	8. 9. H. ȝ.
74 Schrebèri, N.A.	(wh.) Schreber's.	—	1827.	8. 10. H. ȝ.
75 salígnus, w.	(wh.) Sallow-leaved.	Germany.	1815.	— H. ȝ.
76 gravéolens, N.	(bl.) strong-scented.	Arkansa.	1825.	10. 11. H. ȝ.
77 púniceus, w.	(re.) red-stalked.	N. America.	1710.	7. 10. H. ȝ.
78 híspidus, w.	(wh.) rough-stalked.	China.	1804.	9. 10. H. ȝ.
79 dentátus, Th.	(bl.) tooth-leaved.	C. B. S.	1825.	7. 9. G. ȝ.
80 serrátus, Th.	(bl.) saw-leaved.	—	—	G. ȝ.
81 tatáricus, w.	(bl.) Tartarian.	Siberia.	1818.	— H. ȝ.
82 pyren'æus, s.s.	(vi.) Pyrenean.	Pyrenees.	1768.	6. 9. H. ȝ.
83 sibíricus, w.	(pu.) Siberian.	Siberia.	1768.	— H. ȝ.
84 incísus, f.	(li.) cut-leaved.	—	1816.	6. 9. H. ȝ.
85 amelloïdes, Bes.	(vi.) Amellus-like.	Poland.	1827.	7. 9. H. ȝ.
86 élegans, w.	(bl.) elegant.	—	1790.	8. 10. H. ȝ.
87 amplexicaúlis, w.	(bl.) stem-clasping.	N. America.	—	9. 11. H. ȝ.
88 adulterínus, w.	(re.) bastard.	—	—	8. 10. H. ȝ.
89 lèvigátus, w.	(li.) smooth-stem'd.	—	1794.	9. 11. H. ȝ.
90 longifolius, P.S.	(bl.) long-leaved.	—	1798.	10. H. ȝ.
91 versícolor, w. (wh.re.)	various-color'd.	—	1790.	8. 9. H. ȝ.
92 mutábilis, w.	(pu.) changeable.	—	1710.	9. 10. H. ȝ.
93 l'ævis, w.	(li.) smooth.	—	1753.	— H. ȝ.
94 concínnus, W.en.	(li.) neat.	—	1800.	— H. ȝ.
95 floribundus, w.	(li.) abundant-flower'd.	—	—	— H. ȝ.
96 nòvi bélgi, w.	(li.) New-York.	—	1710.	— H. ȝ.
97 fir'mus, L.en.	(li.) firm.	—	1818.	— H. ȝ.
98 bellidiflórus, W.en.	(li.) daisy-flowered.	—	—	— H. ȝ.
99 spectabilis, w.	(bl.) showy.	—	1777.	8. 9. H. ȝ.
100 serotínus, w.	(bl.) late-flowering.	—	—	9. 11. H. ȝ.
101 tardiflórus, w.	(li.) spear-leaved.	—	1775.	7. 9. H. ȝ.
102 blándus, Ph.	(bh.) charming.	—	1800.	10. 11. H. ȝ.
103 acuminatus, Ph. (wh.)	acuminata.	—	1806.	8. 10. H. ȝ.
104 éminens, W.en.	(re.) eminent.	—	—	9. 11. H. ȝ.
105 pubescens, N.A.	(re.) pubescent.	—	1827.	— H. ȝ.
106 Rádula, w.	(wh.) rasp-leaved.	—	1785.	— H. ȝ.
107 prenanthroídes, w.	(bl.) hairy-branch'd.	—	1817.	— H. ȝ.
108 strictus, Ph.	(vi.) upright dwarf.	—	1806.	— H. ȝ.
109 Tradescánti, w.	(wh.) Michaelmas daisy.	—	1633.	7. 9. H. ȝ.
110 vimineus, w.	(bl.) slender.	—	1818.	— H. ȝ.
111 recurvátus, w.	(li.) recurved.	—	1800.	8. 9. H. ȝ.
112 expánsus, N.A.	(bl.) expanded.	—	1827.	— H. ȝ.
113 fastigiátus, Led.	(bl.) clustered.	Siberia.	—	— H. ȝ.
114 láxus, W.en.	(wh.) loose-stalked.	N. America.	—	9. 11. H. ȝ.
115 simplex, W.en.	(wh.) simple-stalked.	—	—	8. 10. H. ȝ.

Rob. ic. 307.

Jacq. vind. 1. t. 8.
Bot. reg. t. 340.

Herm. parad. 98.

Bot. reg. 183. f. inf.
Hoff. phyt. 1. t. B. f. 1.

Bot. reg. t. 273.

Swt. brit. fl. gar. t. 234.
Eng. bot. v. 2. t. 87.Herm. lugd. t. 651.
Kämpf. ic. t. 29.

Gmel. sib. 2. t. 80. f. 1.

Moris. s. 7. t. 22. f. 26.

Herm. lugd. t. 67.

Herm. lugd. t. 69.

Bot. mag. t. 2707.

Moris. s. 7. t. 21. f. 42.

116	virgíneus. N.A. (wh.)	virgin.	N. America.	1826.	8. 10. H. 4.
117	polyphyllus. W.en.	(w.) many-leaved.	—	—	H. 4.
118	júnceus. w.	(fl.) slender-stalked.	—	1758.	9. 10. H. 4.
119	lanceolátus. w.	(wh.) lanceolate.	—	1811.	8. 11. H. 4.
120	dracunculoídes. w.	(w.) Tarragon-like.	—	—	9. 11. H. 4.
121	frágilis. w.	(wh.) brittle.	—	1800.	9. H. 4.
122	cárneus. N.A.	(fl.) flesh-coloured.	—	1827.	— H. 4.
123	pinifolius. L.en.	(wh.) Pine-leaved.	—	1823.	— H. 4.
124	laxiflorus. L.en.	(li.) loose-flowered.	—	—	8. 10. H. 4.
125	miser. w.	(wh.) meagre-flower'd.	—	1759.	9. 10. H. 4.
126	dívèrgens. w.	(wh.) spreading downy.	—	1758.	— H. 4.
127	dívèssus. w.	(wh.) diffuse.	—	1777.	— H. 4.
128	scopàrius. N.A.	(wh.) broom-like.	—	1827.	— H. 4.
129	péndulos. w.	(wh.) pendulous.	—	1758.	— H. 4.
130	parviflorus. L.en.	(w.) small-flowered.	—	1822.	— H. 4.
131	rigidulus. L.en.	(re.) stiffish.	—	—	H. 4.
132	myrtifolius. L.en.	(w.) Myrtle-leaved.	—	1819.	8. 9. H. 4.
133	abbreviátus. N.A. (bl.)	short-rayed.	—	1827.	— H. 4.
134	robustus. N.A.	(bl.) robust.	—	—	H. 4.

CALLISTEMA Cas. CHINA ASTER. Syngenesia Polygamia Superflua.

hortén sis. Cas.	(va.) garden.	China.	1731.	7. 9. H. ④.
α cærulea.	(bl.) blue-flowered.	—	—	H. ④.
β rúbra.	(re.) red-flowered.	—	—	H. ④.
γ álba.	(wh.) white-flowered.	—	—	H. ④.
δ variegáta.	(bl.wh.) variegated.	—	—	H. ④.
ϵ versicolor.	(re.wh.) red and white.	—	—	H. ④.
ζ multiplicéa.	(va.) double-flowered.	—	—	H. ④.
η brachyántha.	(va.) bonnet.	—	—	H. ④.

SOLIDAGO W. GOLDEN-ROD. Syngenesia Polygamia Superflua. L.

1	canadénsis. w.	(ye.) Canadian.	N. America.	1648.	7. 9. H. 4.
2	pròcera. w.	(ye.) great.	—	1758.	9. 10. H. 4.
3	serotína. w.	(ye.) upright smooth.	—	—	7. 8. H. 4.
4	gigántea. w.	(ye.) gigantic.	—	—	8. 9. H. 4.
5	ciliáris. w.	(ye.) ciliated.	—	1811.	— H. 4.
6	refléxa. w.	(ye.) hanging-leaved.	—	1758.	8. 9. H. 4.
7	frágans. W.en.	(ye.) fragrant.	—	—	7. 9. H. 4.
8	lateriflòra. w.	(ye.) lateral-flowered.	—	1758.	8. 9. H. 4.
9	glábra. s.s.	(ye.) smooth.	—	—	H. 4.
10	áspera. w.	(ye.) rough-leaved.	—	1732.	9. H. 4.
11	altíssima. w.	(ye.) tall.	—	1686.	8. 9. H. 4.
12	recurváta. W.en.	(ye.) recurved.	—	1814.	— H. 4.
13	rugósa. Ph.	(ye.) wrinkle-leaved.	—	1732.	— H. 4.
14	villósa. Ph.	(ye.) villous.	—	—	H. 4.
15	scábra. w.	(ye.) rough.	—	1811.	— H. 4.
16	pyramidáta. Ph.	(ye.) pyramidal.	—	1812.	8. 9. H. 4.
17	nemorális. w.	(ye.) woolly-stalked.	—	1769.	9. H. 4.
18	pátula. w.	(ye.) spreading.	—	1805.	9. 10. H. 4.
19	ulmifolia. w.	(ye.) Elm-leaved.	—	—	8. 10. H. 4.
20	argútâ. w.	(ye.) sharp-notched.	—	1758.	7. 8. H. 4.
21	júncea. w.	(ye.) rush-stalked.	—	1769.	8. 9. H. 4.
22	ellíptica. w.	(ye.) oval-leaved.	—	1759.	— H. 4.
23	asperáta. Ph.	(ye.) roughish-leaved.	Canada.	1825.	— H. 4.
24	odóra. w.	(ye.) sweet-smelling.	N. America.	1699.	8. 10. H. 4.
25	retrórsâ. M.	(ye.) reflex-leaved.	—	1824.	— H. 4.
26	lævigáta. N.	(ye.) fleshy-leaved.	—	—	10. 11. H. 4.
27	mexicâna. w.	(ye.) Mexican.	America.	1683.	7. 10. H. 4.
28	sempervírens. N.	(ye.) evergreen.	N. America.	1699.	9. 10. H. 4.
29	speciosa. N.	(ye.) handsome.	—	1812.	— H. 4.
30	multiflòra. P.S.	(ye.) many-flowered.	—	—	H. 4.
31	pauciflosculosa. M.	(ye.) slender-flower'd.	—	1811.	8. 10. H. 4.
32	bicolor. w.	(wh.ye.) two-coloured.	—	1759.	8. 9. H. 4.

A'ster bicolor. Nees.

33	petioláris. w.	(ye.) late-flowered.	—	1758.	10. 12. H. 4.
34	stricta. w.	(ye.) Willow-leaved.	—	—	9. H. 4.
35	hùmilis. Ph.	(ye.) dwarf.	—	1811.	7. 8. H. 4.
36	virgáta. M.	(ye.) slender.	—	1827.	— H. 4.
37	lívida. W.en.	(ye.) purple-stemmed.	—	1812.	— H. 4.
38	hírta. W.en.	(ye.) hairy.	—	—	9. 10. H. 4.
39	litheospermifolia. Ph.	(ye.) Gromwell-leav'd.	—	1811.	8. 10. H. 4.

Dill. elth. t. 34. f. 38.

Schk. hand. 3. f. 246.

Dill. elth. t. 305. f. 392.
Mart. cent. 14.

Dill. elth. t. 308. f. 396.

Pluk. alm. t. 116. f. 6.

Dodart. act 4. t. 219.
Corn. canad. t. 169.

Pluk. alm. t. 114. f. 8.

COMPOSITÆ.

40 hispida. w.	(ye.) hispid.	N. America.	1820.	8. 10. H. ȝ.
41 cæsia. w.	(ye.) Maryland.	——	1732.	9. 10. H. ȝ.
42 flexicaulis. w.	(ye.) crook-stalked.	——	1725.	9. H. ȝ.
43 latifolia. L.	(ye.) broad-leaved.	——	——	H. ȝ.
44 ambigua. w.	(ye.) angular-stalked.	1759.	7. 8. H. ȝ.	
45 macrophylla. Ph.	(ye.) large-leaved.	N. America.	——	8. 10. H. ȝ.
46 glomerata. M.	(ye.) close-flowered.	——	1820.	—— H. ȝ.
47 squarrosa. N.	(ye.) squarrose-cup'd.	——	1827.	—— H. ȝ.
48 alpestris. w.k.	(ye.) Alpine.	Europe.	——	7. 9. H. ȝ.
49 Virgaurea. w.	(ye.) common.	Britain.	——	—— H. ȝ.
50 cæmbrica. w.	(ye.) Welsh.	Wales.	——	7. 8. H. ȝ.
51 asiatica. F.	(ye.) Asiatic.	N. Asia.	1824.	—— H. ȝ.
52 axillaris. Ph.	(ye.) axillary.	N. America.	1811.	8. 10. H. ȝ.
53 erecta. Ph.	(ye.) upright.	——	——	H. ȝ.
54 viminea. w.	(ye.) twiggy.	——	1759.	9. H. ȝ.
55 angustifolia. N.	(ye.) narrow-leaved.	——	1824.	—— H. ȝ.
56 littoralis. s.s.	(ye.) sea-side.	Italy.	1827.	7. 9. H. ȝ.
57 decurrentis. s.s.	(ye.) decurrent.	China.	1823.	8. 10. H. ȝ.
58 puberula. N.	(ye.) pubescent.	N. America.	——	—— H. ȝ.
59 minuta. w.	(ye.) least.	Pyrenees.	1772.	7. 8. H. ȝ.
60 multiradiata. w.	(ye.) Labrador.	Labrador.	1776.	—— H. ȝ.
61 elata. Ph.	(ye.) tall hairy.	N. America.	1811.	8. 10. H. ȝ.
62 rigida. w.	(ye.) hard-leaved.	——	1710.	9. H. ȝ.
EUTHAMIA. N.	EUTHAMIA.	Syngenesia Polygamia Superflua.		
1 graminifolia. n.	(ye.) grass-leaved.	N. America.	1758.	8. 10. H. ȝ.
	<i>Chrysócoma graminifolia.</i> L. <i>Solidago lanceolata.</i> w.			
2 tenuifolia. N.	(ye.) slender-leaved.	N. America.	——	9. 10. H. ȝ.
BRACHYRIS. N.	BRACHYRIS.	Syngenesia Polygamia Superflua.		
	<i>Euthamia.</i> n. (ye.) strong-scented.	Missouri.	1827.	8. 10. H. ȝ.
	<i>Solidago Sarothrae.</i> Ph. <i>Brachyachyris Euthamiae.</i> s.s.			
CHRYSOCOMA. W.	GOLDY-LOCKS.	Syngenesia Polygamia Aequalis. L.		
1 comæra. w.	(ye.) greatshubby.	C. B. S.	1731.	6. 8. G. ȝ.
2 pátula. w.	(ye.) spreading.	——	1816.	—— G. ȝ.
3 cérrna. w.	(ye.) small shrubby.	——	1712.	5. 9. G. ȝ.
4 ciliaris. w.	(ye.) Heath-leaved.	——	1759.	7. 10. G. ȝ.
5 nivea. w.	(ye.) woolly-leaved.	——	1820.	5. 9. G. ȝ.
	<i>tomentosa.</i> Jacq. non L.			
6 scabra. w.	(ye.) rugged.	——	1732.	8. 9. G. ȝ.
7 denticulata. w.	(ye.) toothed.	——	——	—— G. ȝ.
8 Linos'rys. w.	(ye.) flax-leaved.	Europe.	——	9. 10. H. ȝ.
9 dracunculoides. w.	(ye.) Siberian.	Siberia.	——	—— H. ȝ.
10 biflora. w.	(ye.) two-flowered.	——	1741.	8. 9. H. ȝ.
11 graveolens. N.	(ye.) Rue-scented.	Missouri.	1822.	—— H. ȝ.
12 nudata. M.	(ye.) naked.	N. America.	1819.	—— H. ȝ.
13 virgata. N.	(ye.) slender.	——	——	—— H. ȝ.
14 villosa. w.	(ye.) villous.	Hungary.	1799.	—— H. ȝ.
BOLTONIA. W.	BOLTONIA.	Syngenesia Polygamia Superflua. L.		
1 asteroides. w.	(wh.) Starwort-flower'd. N. America.	1758.	8. 10. H. ȝ.	
2 glastifolia. w.	(pu.) glaucous-leaved.	——	——	9. H. ȝ.
AMELLUS. W.	AMELLUS.	Syngenesia Polygamia Superflua.		
Lychnitis. w.	(bl.) Cape.	C. B. S.	1768.	6. 7. G. ȝ.
SIDERANTHUS. Fras.	SIDERANTHUS.	Syngenesia Polygamia Superflua.		
1 villösus.	(ye.) villous.	Missouri.	1811.	8. 9. H. ȝ.
	<i>Sideranthus integrifolius.</i> Fras. <i>Amellus villosus.</i> Ph.			
2 spinulosus.	(ye.) spiny.	——	——	H. ȝ.
	<i>Amellus spinulosus.</i> Ph. <i>Sideranthus pinnatifidus.</i> Fras.			
STA'RKEA. W.	STA'RKEA.	Syngenesia Polygamia Superflua.		
1 umbellata. w.	(ye.) umbel-flower'd. Jamaica.	1768.	6. 7. S. ȝ.	
2 igniaria. D.D.	(ye.) close-flowered. S. America.	1823.	6. 8. F. ȝ.	
	<i>Hierba de Santa María.</i> incolarum. <i>Andromachia igniaria.</i> K.S.			
GRIND'ELIA. W.en.	GRIND'ELIA.	Syngenesia Polygamia Superflua. W.		
1 glutinosa. Dun.	(ye.) glutinous.	Mexico.	1803.	1. 12. F. ȝ.
	<i>Dónia glutinosa.</i> B.R. <i>Dorónicum glutinosum.</i> w.			
2 inuloides. W.en.	(ye.) Inula-like.	Mexico.	1815.	6. 9. F. ȝ.
3 pulchella. s.s.	(ye.) pretty.	Chile.	1827.	—— F. ȝ.
4 nítida. Bern.	(ye.) glossy.	Mexico.	1825.	—— F. ȝ.
5 coronopifolia. Leh.	(ye.) Buckshorn-leav'd.	——	1826.	—— F. ȝ.
6 angustifolia. B.R.	(ye.) narrow-leaved.	——	1822.	—— F. ȝ.
7 spatulata. L.en.	(ye.) spatulate-leav'd.	——	——	F. ȝ.

Pluk. alm. t. 235. f. 3.
—— t. 235. f. 4.

Eng. bot. v. 5. t. 301.
Dill. elth. t. 306. f. 393.

Lodd. bot. cab. t. 189.

Herm. parad. 243.

Bot. mag. t. 2546.

Bot. mag. t. 1972.

Comm. hort. 2. t. 45.
—— 2. t. 48.

Jacq. schœn. 2. t. 147.

Dill. elth. t. 88. f. 103.
Jacq. schœn. 3. t. 368.

Eng. bot. v. 35. t. 2505.

Gmel. sib. 2. t. 82. f. 1.

W. et K. hung. 1. t. 58.

Bot. mag. t. 2554.
—— t. 2381.

Bot. reg. t. 586.

Lam. ill. t. 682. f. 2.
H. et B. pl. æq. 2.t.112.

Bot. reg. t. 187.

Bot. reg. t. 248.

Bot. reg. t. 781.

COMPOSITÆ.

299

- | | | | | | |
|------------------------------|-------------------------------|----------------------------------|-------|--------------|------------------------------|
| 8 squarròsa. Dun. | (ye.) Snake-headed. | Missouri. | 1811. | 7. 9. F. ȝ. | Bot. mag. t. 1706. |
| 9 ciliàta. s.s. | (ye.) ciliated. | Arkansa. | 1821. | 9. 10. H. ȝ. | Hook. ex. flor. t. 45. |
| Dònia ciliàta. H.E.F. | | | | | |
| 10 villòsa. d. | (ye.) villous. | Columbia. | 1827. | 8. 10. H. ȝ. | |
| PODOL'EPIS. Lab. | PODOL'EPIS. | Syngenesia Polygamia Superflua. | | | |
| 1 rugàta. h.k. | (ye.) wrinkle-scaled. | N. Holland. | 1803. | 7. 8. G. ȝ. | Lab. nov. holl. 2. t. 208. |
| 2 acuminàta. h.k. | (ye.) sharp-scaled. | N. S. W. | — | 5. 8. G. ȝ. | Bot. mag. t. 956. |
| Scàlia jaceoïdes. B.M. | | | | | |
| 3 gràcilis. B.F.G. | (li.) slender. | N. Holland. | 1827. | 8. 10. H. ȝ. | Swt. br. fl. gar. t. 285. |
| GERB'ERIA. Cass. | GERB'ERIA. | Syngenesia Polygamia Superflua. | | | |
| 1 crenàta. B.R. | (li.) crenated. | C. B. S. | 1820. | 4. 8. G. ȝ. | Bot. reg. t. 855. |
| 2 asplenioïda.s.s.(ye.pu.) | Spleenwort-leaved. | — | 1822. | 4. 6. G. ȝ. | Burm. afr. 155. t. 26. f. 1. |
| Arnica Gerbera. L. | Dorónicum asplenioïdium. Lam. | | | | |
| A'RNICÀ. W. | A'RNICÀ. | Syngenesia Polygamia Superflua. | | | |
| 1 montàna. w. | (ye.) mountain. | Europe. | 1731. | 7. 8. H. ȝ. | Bot. mag. t. 1749. |
| 2 angustifòlia. s.s. | (ye.) narrow-leaved. | Greenland. | 1827. | 6. 8. H. ȝ. | |
| 3 fúlgens. Ph. | (ye.) glossy-flowered. | Missouri. | 1826. | — H. ȝ. | |
| 4 scorpioides. w. | (ye.) alternate-leaved. | Austria. | 1710. | — H. ȝ. | Jacq. aust. 4. t. 349. |
| 5 glaciàlis. w. | (ye.) alpine. | Germany. | 1816. | 7. 8. H. ȝ. | Jacq. ic. 3. t. 586. |
| 6 Dorónicum. w. | (ye.) Clusius's. | Austria. | — | — H. ȝ. | Jacq. aust. 1. t. 92. |
| Clüstí. All. t. 17. f. 1. 2. | | | | | |
| 7 lanígera. T.N. | (ye.) woolly. | Italy. | 1827. | — H. ȝ. | |
| 8 marítima. L. | (ye.) sea-side. | Kamtschatka. | 1821. | — H. ȝ. | |
| BELLIDIA'STRUM. Cas. | BELLIDIA'STRUM. | Syngenesia Polygamia Superflua. | | | |
| Michèllii. Cas. (wh.ro.) | Micheli's. | Austria. | 1570. | 6. 8. H. ȝ. | Bot. mag. t. 1196. |
| A'rnicà Bellidiástrum. B.M. | Dorónicum Bellidiástrum. L. | | | | |
| DIPLO'COMA. D.D. | DIPLO'COMA. | Syngenesia Polygamia Superflua. | | | |
| villòsa. D.D. | (ye.) villous. | Mexico. | 1827. | 8. 10. F. ȝ. | Swt. br. fl. gar. t. 246. |
| NEJA. D.D. | N'EJA. | Syngenesia Polygamia Superflua. | | | |
| gràcilis. D.D. | (ye.) slender. | Mexico. | 1828. | 9. 10. F. ȝ. | Swt. br. fl. gar. ser. 2. |
| DOR'ONICUM. W. | LEOPARD'S-BANE. | Syngenesia Polygamia Superflua. | | | |
| 1 Pardaliàncches.w. | (ye.) great. | Britain. | | 5. H. ȝ. | W. |
| 2 macrophyllum.s.s. | (ye.) large-leaved. | Europe. | 1821. | — H. ȝ. | Eng. bot. v. 9. t. 630. |
| 3 scorpioides. w. | (ye.) mountain. | Germany. | — | 4. 6. H. ȝ. | |
| 4 Colùmnae. T.S. | (ye.) Columna's. | Naples. | 1826. | — H. ȝ. | Column. ecphr. 2. t. 46. |
| 5 caucásicum. M.B. | (ye.) Caucasian. | Caucasus. | 1815. | — H. ȝ. | |
| orientàle. W.en. | | | | | |
| 6 altaicum. w. | (ye.) Siberian. | Siberia. | 1783. | 6. 8. H. ȝ. | Pall. a.pet. 1779. t. 16. |
| 7 dentatum. L.en. | (ye.) toothed. | | 1825. | 5. 6. H. ȝ. | |
| 8 plantaginum. w. | (ye.) Plantain-leav'd. | S. Europe. | 1570. | — H. ȝ. | |
| 9 austriacum. w. | (ye.) Austrian. | Austria. | 1816. | — H. ȝ. | Jacq. aust. t. 130. |
| LEPTOSTE'LMA. D.D. | LEPTOSTE'LMA. | Syngenesia Polygamia Superflua. | | | |
| máxima. D.D. | (wh.) gigantic. | Mexico. | 1828. | 8. 9. H. ȝ. | Swt. br. fl. gar. s.2.t.38. |
| BE'LLIS. W. | DAISY. | Syngenesia Polygamia Superflua. | L. | | |
| 1 sylvestris. w. | (wh.) large Portugal. | Portugal. | 1797. | 5. 7. H. ȝ. | Bot. mag. t. 2511. |
| 2 perénvis. w. | (wh.) common. | Britain. | | 3. 8. H. ȝ. | Eng. bot. v. 6. t. 424. |
| β horténisis. | (wh.) large double. | | | — H. ȝ. | Bot. mag. t. 228. |
| γ fistulòsa. | (re.) red quilled. | | | — H. ȝ. | |
| δ tubulòsa. | (wh.) white quilled. | | | — H. ȝ. | |
| ε prolíféra. | (wh.re.) Hen & Chicken. | | | — H. ȝ. | |
| 3 hy'bida. Ten. | (wh.) Italian. | Italy. | 1824. | 4. 8. H. ȝ. | |
| 4 ánnua. w. | (wh.) annual. | S. Europe. | 1759. | 3. 7. H. ⊙. | Bot. mag. t. 2174 |
| BE'LLIUM. W. | BE'LLIUM. | Syngenesia Polygamia Superflua. | L. | | |
| 1 bellidioïdes. w. | (wh.) small. | Italy. | 1796. | 6. 9. H. ⊙. | Lam. ill. t. 684. |
| 2 minutum. w. | (wh.) dwarf. | Levant. | 1772. | 6. 10. H. ȝ. | Schreb.a.ups.1.t.5. |
| PSI'ADIA. W. | PSI'ADIA. | Syngenesia Polygamia Necessaria. | W. | | |
| glutinòsa. w. | (ye.) glutinous. | Mauritius. | 1796. | 6. 8. S. ȝ. | Jacq. schoen. 2. t. 152. |

Subordo IV. EUPATOREÆ. *Kth. synops.* 2, p. 408.

- | | | | |
|-----------------------------|---|-------------------|-------------------------|
| K'UHNIA. W. | K'UHNIA. Syngenesia Polygamia Aequalis. | | |
| 1 eupatorioïdes. w.(wh.) | Eupatorium-like. N.America. 1812. | 6. 8. H. 24. | Linn. fil. dec. t. 11. |
| | <i>Critonia eupatorioïdes</i> . G. | | |
| 2 Critonia. w. | (st.) glandular-leaved. —— | 1816. —— H. 24. | Gært. carp.2.t.174.f.7. |
| | <i>Critonia Kühnia</i> , G. | | |
| 3 rosmarinifolia. s.s. | Rosemary-leaved. Cuba. | 1827. S. 24. | |

COMPOSITÆ.

EUPATORIUM. W.	EUPATORIUM.	Syngenesia	Polygamia	Aequalis.	L.			
1 Dálea. w.	(wh.) shrubby.	Jamaica.	1773.	8.	S. h.	Jacq. schen. 2. t. 146.		
2 parviflórum. w.	(wh.) small-flowered.	—	1823.	—	S. h.	Aub. gni. 2. t. 314.		
3 triflórum. w.	(wh.) three-flowered.	Trinidad.	1821.	7. 8.	S. h.	Plum. spec. 10. t. 129.		
4 macrophyllum. w.	(li.) large-leaved.	S. America.	1824.	—	S. 2.	Vahl symb. 3. t. 73.		
5 populiflórum. k.s.	(wh.) Poplar-leaved.	Mexico.	—	—	S. h.	H. et B. v. 4. t. 341.		
6 canéscens. w.	(wh.) canescent.	S. America.	1822.	6. 8.	S. h.	Dill. elth. t. 115. f. 140.		
7 veronicafólium. k.s.	(w.) Speedwell-leav'd.	Mexico.	1825.	—	S. h.	Willd. hort. ber. t. 32.		
8 fœniculáceum. w.	(st.) Fennel-leaved.	N. America.	1807.	6. 9.	H. 2.	Pluk. alm. t. 88. f. 4.		
9 coronopifólium. w.	(wh.) buckshorn-leaved.	—	1819.	8. 9.	H. 2.	Jacq. vind. 2. t. 164.		
10 hyssopifólium. w.	(wh.) Hyssop-leaved.	—	1699.	—	H. 2.	Eng. bot. v. 6. t. 428.		
11 sessiliflórum. w.	(wh.) sessile-leaved.	—	1777.	9. 10.	H. 2.	Jacq. ic. 1. t. 170.		
12 truncátum. w.	(wh.) truncate-leaved.	—	1822.	8. 10.	H. 2.	Corn. canad. t. 72.		
13 álbum. w.	(wh.) white-flowered.	—	—	—	H. 2.	Herm. parad. t. 158.		
14 lanceolátum. w.	(wh.) spear-leaved.	—	1819.	8. 11.	H. 2.	Pluk. alm. t. 87. f. 6.		
15 teucrifólium. w.	(wh.) Teucrium-leav'd.	—	1816.	—	H. 2.	Bot. mag. t. 2010.		
16 rotundiflórum. w.	(wh.) round-leaved.	—	1699.	7. 8.	H. 2.	Pluk. alm. t. 88. f. 3.		
17 pubéscens. w.	(wh.) pubescent.	—	1816.	8. 10.	H. 2.	Smith ined. t. 68.		
18 ceanothifólium. w.	(w.) Ceanothus-like.	—	1812.	8. 11.	H. 2.	Corn. canad. t. 21.		
19 altissimum. w.	(wh.) tall.	—	1699.	9. 10.	H. 2.	Jacq. schen. 3. t. 369.		
20 trifoliátum. w.	(wh.) three-leaved.	—	1768.	8. 10.	H. 2.	Cavan. ic. 1. t. 98.		
21 cannabinum. w.	(pu.) Hemp agrimony.	Britain.	—	7. 10.	H. 2.	Plum. sp. 10. ic. 128. f. 1.		
22 syriacum. w.	(ro.) Syrian.	Syria.	1807.	7. 9.	H. 2.	Pluk. alm. t. 177. f. 3.		
23 diffusum. s.s.	(wh.) spreading.	S. America.	1821.	6. 8.	S. 2.	Vent. malm. p. 3. t. 3.		
24 purpúreum. w.	(pu.) purple-stalked.	N. America.	1640.	9. 10.	H. 2.	Sm. ic. ined. t. 69.		
25 maculátum. w.	(pu.) spotted-stalked.	—	1656.	8. 9.	H. 2.	Jacq. ic. 3. t. 282.		
26 punctátum. Ph.	(pu.) punctated.	—	1815.	—	H. 2.	H. et B. v. 4. t. 340.		
27 verticillátum. w.	(pu.) whorl-leaved.	—	1811.	—	H. 2.	— v. 4. t. 345.		
28 perfoliátum. w.	(wh.) Feverwort.	—	1699.	8. 10.	H. 2.	Plum. ic. 130. f. 1.		
29 salviafólium. B.M.	(wh.) Sage-leaved.	—	1812.	—	H. 2.	ST'EVIA. W.		
30 lamífiolum. L.en.	(wh.) Lamium-leaved.	—	1818.	—	H. 2.	ST'EVIA. Syngenesia Polygamia Aequalis. W.		
31 paniculátum. Mil.	(wh.) paniced.	—	—	—	H. 2.	1 connáta. L.G.		
32 melissoídes. w.	(wh.) Balm-leaved.	—	1821.	8. 10.	H. 2.	(wh.) connate-leaved.		
33 aromáticum. w.	(wh.) aromatic.	—	1739.	7. 8.	H. 2.	New Spain.		
34 urticafólium. w.	(wh.) Nettle-leaved.	S. America.	1803.	—	G. 2.	1825.	8. 10.	F. h.
35 ageratoides. w.	(wh.) Ageratum-like.	N. America.	1640.	8. 10.	H. 2.	1826.	—	F. h.
36 deltoides. w.	(pi.) deltoid-leaved.	—	1818.	—	S. 2.	1827.	—	F. h.
37 scándens. L.en.	(ye.) climbing.	S. America.	1822.	6. 8.	S. h.	1828.	—	F. h.
38 squarrósom. w.	(st.) squarrose.	Mexico.	1824.	—	S. 2.	1829.	—	S. h.
39 atriplicifólium. w.	(li.) Orache-leaved.	W. Indies.	1820.	—	S. h.	1830.	—	S. h.
40 sinuátum. P.S.	(wh.) sinuate.	—	1824.	6. 8.	S. h.	1831.	—	S. h.
41 odorátum. w.	(wh.) sweet-scented.	Jamaica.	1752.	8. 10.	S. h.	1832.	—	S. h.
42 Ayapána. v.	(pu.) medicinal.	Brazil.	1821.	—	S. h.	1833.	—	S. h.
43 triplinérve. P.S.	... triply-nerved.	Santa Cruz.	—	—	S. h.	1834.	—	S. h.
44 ivæfolium. w.	(wh.) Iva-leaved.	Jamaica.	1794.	6. 7.	S. 2.	1835.	—	S. h.
45 stœchadifólium. L.	(vi.) Steechas-leav'd.	New Granada.	1826.	—	S. h.	1836.	—	S. h.
46 myosotifólium. w.	(bl.) Myosotis-leav'd.	S. America.	1824.	6. 8.	S. 2.	1837.	—	S. h.
47 iresinoïdes. k.s.	(wh.) Iresine-like.	—	—	—	S. 2.	1838.	—	S. h.
48 pulchellum. k.s.	(vi.) pretty.	Mexico.	—	—	S. h.	1839.	—	S. h.
49 xalapénse. k.s.	(wh.) six-angled.	—	—	—	S. h.	1840.	—	S. h.
50 repándum. w.	(wh.) repand-leaved.	Antilles.	1822.	—	S. h.	1841.	—	S. h.
51 ásperum. H.B.	(wh.) rough.	E. Indies.	1820.	6. 8.	S. h.	1842.	—	S. h.
MIK'ANIA. W.	MIK'ANIA.	Syngenesia Polygamia Aequalis. W.						
1 Houstóni. w.	(wh.) Houstoun's.	Jamaica.	1733.	7. 8.	S. h.			
2 hastáta. w.	(wh.) halbert-leaved.	—	—	—	S. h.	Brown jam. t. 34. f. 3.		
3 scándens. w.	(wh.) climbing.	N. America.	1714.	8. 9.	H. 2.	Jacq. ic. 1. t. 169.		
4 Guáco. k.s.	(wh.) medicinal.	S. America.	1823.	—	S. 2.	H. et B. pl. 2. t. 105.		
5 amára. w.	(wh.) bitter.	Trinidad.	1824.	—	S. h.	Aubl. gui. 2. t. 315.		
6 opífera. Mart.	(wh.) snakewort.	Brazil.	1822.	—	S. h.			
7 nummulária. D.D.	(wh.) moneywort.	Mexico.	1826.	6. 8.	S. 2.			
ST'EVIA. W.	ST'EVIA.	Syngenesia Polygamia Aequalis. W.						
1 connáta. L.G.	(wh.) connate-leaved.	New Spain.	1825.	8. 10.	F. h.			
2 subpubéscens. L.G.	(w.) downy-branch'd.	—	—	—	F. h.			
3 coriacea. L.G.	(wh.) leather-leaved.	Peru.	1826.	—	F. h.			
4 lúcida. L.G.	(wh.) glossy-leaved.	New Spain.	1822.	—	F. h.			
5 salicifólia. w.	(wh.) Willow-leaved.	Mexico.	1803.	—	F. h.			
6 angustifólia. k.s.	(wh.) narrow-leaved.	—	—	—	F. h.			
7 trífida. L.G.	(wh.) trifid-leaved.	New Spain.	1828.	—	F. 2.			
8 micránta. L.G.	(wh.) small-flowered.	—	1825.	8. 10.	F. 2.	Cavan. ic. 4. t. 354.		

9 incanescens. L.G. (<i>wh.</i>) villous-stalked.	New Spain.	1825.	8. 10.	F. 24.
10 pubescens. L.G. (<i>pu.</i>) pubescent.	Mexico.	1823.	—	F. 24.
11 tomentosa. K.S. (<i>ro.</i>) tomentose.	—	1825.	—	F. 24.
12 monardæfolia. K.S. (<i>ro.</i>) Monarda-leav'd.	—	—	8. 9.	F. 24.
13 fastigiata. K.S. (<i>wh.</i>) crowded-flower'd.	New Spain.	1828.	—	F. 24.
14 adenophora. L.G. (<i>wh.</i>) glandular.	Chile.	1822.	—	F. 24.
15 enarthrotricha. L.G. (<i>w.</i>) jointed haired.	New Spain.	1828.	—	F. 24.
16 ovata. L.G. (<i>wh.</i>) oval-leaved.	Mexico.	1816.	8. 9.	F. 24.
17 subtoctaristata. L.G. (<i>w.</i>) 8-awned.	Peru.	1824.	—	F. 24.
18 paniculata. L.G. (<i>wh.</i>) paniced.	New Spain.	—	—	F. 24.
19 suavolens. L.G. (<i>wh.</i>) sweet-scented.	—	1827.	—	F. 24.
20 rhombifolia. K.S. (<i>wh.</i>) rhomb-leaved.	Mexico.	1826.	8. 10.	F. 24.
21 nepetaefolia. K.S. (<i>wh.</i>) catmint-leaved.	—	—	—	F. 24.
22 Eupatoria. w. (<i>ro.wh.</i>) entire-leaved.	—	1798.	7. 9.	F. 24.
23 lanceolata. L.G. (<i>hh.</i>) spear-leaved.	—	1824.	—	F. 24.
24 hyssopifolia. B.M. (<i>wh.</i>) Hyssop-leaved.	—	—	—	F. 24.
25 ivæfolia. W.en. (<i>wh.</i>) Iva-leaved.	—	1816.	7. 9.	F. 24.
26 serrata. w. (<i>wh.</i>) saw-leaved.	—	1799.	—	F. 24.
27 virgata. K.S. (<i>wh.</i>) slender.	—	1824.	—	F. 24.
28 purpurea. W.en. (<i>pu.</i>) purple-flowered.	—	1812.	8. 9.	H. 24.
<i>Eupatoria. B.R. nec aliorum.</i>				
29 viscosa. K.S. (<i>ro.</i>) rose-coloured.	—	1824.	9. 10.	F. 24.
30 pilosa. L.G. (<i>ro.</i>) hairy.	New Spain.	1825.	8. 10.	F. 24.
31 microphylla. K.S. (<i>bh.</i>) small-leaved.	Mexico.	1828.	—	F. 24.
PALAFÖ'XIA. Lag. PALAFÖ'XIA.	Syngenesia Polygamia Æqualis.			
1 linearis. L.G. (<i>wh.</i>) linear-leaved.	Mexico.	1816.	8. 10.	F. 24.
<i>Stèvia linearis. w. Ageratum lineare.</i> Cav. ic. 3. t. 205.				Bot. mag. t. 2132.
2 callous. D.D. (<i>wh.</i>) callous.	Arkansa.	1824.	6. 8.	H. ⊖.
<i>Stèvia callosa. N.</i>				
AGERATUM. W. AGERATUM.	Syngenesia Polygamia Æqualis. L.			
1 stríctum. B.M. (<i>wh.</i>) upright.	Nepaul.	1820.	9. 11.	H. ⊖.
2 conyzoides. w. (<i>bl.</i>) hairy.	America.	1714.	7. 8.	H. ⊖.
3 cœruleum. P.S. (<i>bl.</i>) blue-flowered.	S. America.	1820.	—	H. ⊖.
4 mexicanum. B.M. (<i>bl.</i>) Mexican.	Mexico.	1823.	8. 10.	H. ⊖.
5 latifolium. w. (<i>wh.</i>) broad-leaved.	Peru.	1800.	7. 8.	H. ⊖.
6 Houstonianum. Mil. (<i>bl.</i>) Houstoun's.	S. America.	6. 8.	S. ⊖.
7 paniculatum. Scr. (<i>bl.</i>) paniced.	—	1827.	9. 10.	H. ⊖.
8 álbium. w. (<i>wh.</i>) white-flowered.	—	1822.	6. 9.	H. ⊖.
<i>hirtum. Lam.</i>				
9 maritimum. K.S. (<i>bl.</i>) sea-side.	Cuba.	1826.	—	S. 24.
10 cordifolium. H.B. (<i>wh.</i>) heart-leaved.	Bengal.	1820.	8. 10.	S. ⊖.
11 aquáticum. H.B. (<i>wh.</i>) water.	E. Indies.	—	—	S. ⊖.
CELESTINA. Cas. CŒLEST'INA.	Syngenesia Polygamia Æqualis.			
1 suffruticosa. (<i>bl.</i>) suffruticose.	S. America.	7. 10.	G. ½.
<i>Ageratum caelestinum. B.M.</i>				Bot. mag. t. 1730.
2 ageratoïdes. K.S. (<i>bl.</i>) Ageratum-like.	New Spain.	1824.	—	G. 24.
3 cœrulea. Cas. (<i>bl.</i>) blue-flowered.	N. America.	1732.	7. 11.	H. 24.
<i>Eupatorium caelestinum. w.</i>				Dill. elth. t. 114. f. 139.
AL'OMIA. K.S. AL'OMIA.	Syngenesia Polygamia Æqualis.			
ageratoïdes. K.S. (<i>wh.</i>) white-flowered.	New Spain.	1823.	7. 9.	G. 24.
PIQU'ERIA. W. PIQU'ERIA.	Syngenesia Polygamia Æqualis. W.			
1 trinervia. w. (<i>wh.</i>) three-nerved.	Mexico.	1798.	7. 8.	G. 24.
2 pilosa. K.S. (<i>wh.</i>) hairy.	—	1824.	—	G. 24.
3 artemisioides. K.S. (<i>wh.</i>) Wormwood-like.	Quito.	—	8. 12.	G. 24.
SELL'OA. Spr. SELL'OA.	Syngenesia Polygamia Æqualis.			
glutinosa. B.R. (<i>ye.</i>) clammy.	Brazil.	1815.	1. 5.	G. ½.
LEY'SERA. W. LEYS'ERA.	Syngenesia Polygamia Superflua. W.			
1 ciliata. w. (<i>ye.</i>) ciliated.	C. B. S.	1816.	7. 8.	G. 24.
2 gnaphalodes. w. (<i>or.</i>) woolly.	—	1774.	7. 9.	G. ½.
				Jacq. ic. 3. t. 588.

SUBORDO V. JACOBÆ. *Kth. synops. 2. p. 440.*

KLEI'NIA. W.	KLEI'NIA.	Syngenesia Polygamia Æqualis. W.
1 ruderalis. w. (<i>gr.ye.</i>) glaucous-leav'd.	S. America.	1824. 7. 10. S. ⊖.
2 Porophyllum. w. (<i>gr.ye.</i>) perforated.	—	1699. 6. 10. S. ⊖.
3 viridiflora. K.S. (<i>gr.</i>) green-flowered.	Mexico.	1823. — S. ½.

Jacq. amer. t. 127.
Cavan. ic. 3. t. 222.

4 tagetoídes. k.s. (gr.)	Tagetes-like.	Mexico.	1823.	6. 10. H. ⊖.	
5 suffruticosa. w. (pu.)	suffruticose.	S. America.	1825.	— G. ℡.	Cavan. ic. 3. t. 257.
6 colorata. k.s. (vi.)	violet-colour'd.	Mexico.	—	— H. ⊖.	
CAC'ALIA. W.	CAC'ALIA.	Syngenesia Polygamia Äqualis. L.			
1 papillaris. w. (wh.)	rough-stalked.	C. B. S.	1727.	... G. ℡.	Dill. elth. t. 55. f. 63.
2 Anteuphorbium. w. (st.)	oval-leaved.	—	1596.	2. 3. G. ℡.	— t. 55. f. 2. 3.
3 Kleínia. w. (wh.)	Oleander-leaved.	Canaries.	1732.	9. 10. G. ℡.	DC. plant. gras. t. 12.
4 Ficoides. w. (st.)	flat-leaved.	C. B. S.	1710.	6. 11. G. ℡.	— t. 90.
5 rēpens. w. (st.)	glaucous-leaved.	—	1759.	6. 10. G. ℡.	— t. 42.
6 cylindrica. dc.	(ye.) cylindrical.	Africa.	1818.	— G. ℡.	— t. ic.
7 carnosa. w. (st.)	narrow-leaved.	C. B. S.	1757.	6. G. ℡.	
8 canescens. w. (st.)	woolly-leaved.	—	1795.	... G. ℡.	
Haworthii. H.S.L.	Kleínia tomentosa. H.S. non C. tomentosa. L.				
9 longifolia. H.R.	long-leaved.	1820.	... S. ℡.	DC. plant. gras. t. 18.
pugioniformis. L.en.					
10 articulata. w. (wh.)	jointed.	C. B. S.	1775.	9. 11. G. ℡.	
laciniatá. Jacq. ic. 1. t. 168.					
11 tomentosa. w. (ye.)	tomentose.	—	1795.	— G. ℡.	Bot. reg. t. 923.
12 reticulata. w. (ye.)	netted.	Bourbon.	1823.	... G. ℡.	— t. 110.
13 appendiculata. w. (ye.)	appendaged.	Teneriffe.	1815.	6. 9. G. ℡.	— t. 102.
14 salicina. Lab.	(ye.) willow-leaved.	V. Diem. Isl.	1820.	4. 6. G. ℡.	Rheed. mal. 10. t. 68.
15 bicolor. w. (or.)	two-coloured.	E. Indies.	1804.	9. 5. S. ℡.	Bot. mag. t. 564.
16 ovalis. B.R.	(ye.) oval-leaved.	—	—	— S. ℡.	
17 sonchifolia. w. (re.)	Sonchus-leaved.	—	1768.	7. S. ⊖.	Gmel. sib. 2. t. 66.
18 coccinea. B.M.	(sc.) scarlet-flower'd.	S. America.	1799.	6. 7. H. ⊖.	
19 sarracénica. w. (st.)	creeping-rooted.	France.	1772.	8. 10. H. 2.	Pluk. alm. t. 101. f. 2.
20 runcinata. k.s. (ro.)	runcinate.	Mexico.	1825.	... H. 2.	
21 hastata. w. (wh.)	spear-leaved.	Siberia.	1780.	8. 10. H. 2.	
22 rhombifolia. W.en. (st.)	rhomb-leaved.	—	1816.	— H. 2.	
23 macrophylla. M.B. (st.)	large-leaved.	Caucasus.	1828.	— H. 2.	
24 suavolens. w. (wh.)	sweet-scented.	N. America.	1752.	— H. 2.	
25 atriplicifolia. w. (wh.)	Orache-leaved.	—	1699.	8. H. 2.	
26 cordifolia. k.s. (ye.)	heart-leaved.	Mexico.	1825.	7. 9. F. 2.	
27 reniformis. w. (wh.)	kidney-leaved.	N. America.	1801.	7. 8. H. 2.	
28 tuberosa. N.	(wh.) tuberous-rooted.	—	1812.	— H. 2.	
29 pinnata. W.en.	(ye.) pinnate-leaved.	Iberia.	1816.	— H. 2.	
30 peltata. k.s. (gr.)	peltated.	Mexico.	1825.	— G. 2.	
31 scandens. w. (pi.)	climbing.	C. B. S.	1774.	4. G. 2.	
EMI'LEA. Cas.	EMI'LEA.	Syngenesia Polygamia Äqualis.			H. et Bonpl. 4. t. 361.
flámmea. Cas. (or. pu.)	flame-flowered.	Java.	1823.	6. 9. S. ⊖.	
Cacália sagittata. w.					
ADENO'STYLES. Cas.	ADENO'STYLES.	Syngenesia Polygamia Äqualis.			
1 candidissima. Cas. (pu.)	hoary.	S. France.	1828.	7. 9. H. 2.	
Cacália tomentosa. Vill.	leucophylla. w.				Jacq. aust. 3. t. 235.
2 álbida. Cas.	(pi.) white-leaved.	Austria.	1739.	7. 8. H. 2.	
Cacália álbifrons. Jacq.					Jacq. aust. 3. t. 234.
3 virdis. Cas.	(li.) alpine.	—	—	— H. 2.	
Cacália alpina. Jacq.					
TUSSILAGO. W.	COLTSFOOT.	Syngenesia Polygamia Superflua. L.			
1 alpina. w. (pu.)	Alpine.	Austria.	1710.	3. 5. H. 2.	
2 discolor. w. (pu.)	two-coloured.	—	1633.	4. 5. H. 2.	Bot. mag. t. 84.
3 sylvestris. w. (pu.)	wood.	—	1816.	— H. 2.	Jacq. aust. 3. t. 247.
4 Fársara. w. (ye.)	common.	Britain.	3. 4. H. 2.	— 5. app. t. 12.
β variegata.	(ye.) variegated-leav'd.	—	...	— H. 2.	Eng. bot. v. 6. t. 429.
5 frigida. w. (pa.)	Lapland.	Lapland.	1710.	5. H. 2.	Flor. dan. t. 61.
6 frágans. w. (bh.)	fragrant.	Italy.	1806.	1. 3. H. 2.	Bot. mag. t. 1388.
7 álba. w. (wh.)	white butter-bur.	Europe.	1683.	1. 4. H. 2.	Flor. dan. t. 524.
8 corymbosa. s.s. (wh.)	corymbose.	Arctic regions.	1827.	— H. 2.	
9 nívea. w. (wh.)	downy-leaved.	Switzerland.	1713.	4. H. 2.	Retz. obs. 2. t. 3.
paradóxa. Retz.					
10 Petasites. E.B.	(bh.) common butter bur.	Britain.	3. 4. H. 2.	Eng. bot. v. 6. t. 431.
11 hybrida. E.B.	(bh.) large-leaved.	—	— H. 2.	— v. 6. t. 430.
12 spuria. w. (wh.)	lobe-leaved.	Germany.	1790.	— H. 2.	Retz. obs. 1. t. 2.
paradóxa. Roth.					
13 palmata. w. (wh.)	cut-leaved.	Labrador.	1778.	4. H. 2.	Hort. Kew. v. 3. t. 11.
SEN'ECIO. W.	GROUNDSEL.	Syngenesia Polygamia Superflua. W.			
1 reclinatus. w. (ye.)	grass-leaved.	C. B. S.	1774.	6. 8. G. ♂.	Jacq. ic. 1. t. 174.
2 hieracifolius. w. (ye.)	hawkweed-leav'd.	N. America.	1699.	8. H. ⊖.	Herm. parad. t. 226.
3 purpureus. w. (pu.)	purple.	C. B. S.	1774.	7. 9. G. 2.	Jacq. ic. 3. t. 580.

4 cérnuus. w.	(pu.) drooping.	E. Indies.	1780.	7. 8. H. ⊖.	Jacq. vind. 3. t. 98.
5 erubescens. w.	(bh.) blush-coloured.	C. B. S.	1774.	6. 10. G. ♂.	
6 hæmatophyllus. W.en.	(ye.) purple leaved.—		1789.	— G. ♀.	
7 divaricatus. w.	(ye.) straddling.	China.	1801.	7. G. ♂.	
8 croaticus. w.	(ye.) Croatian.	Hungary.	1805.	7. 8. H. ♀.	W. et K. hung. 2. t. 143.
9 Pseudo-China.w.(or.)	Chinese.	China.	1732.	6. 8. G. ♀.	Dill. elth. t. 258. f. 335.
10 japonicus. w.	(ye.) jagged-leaved.	Japan.	1774.	8. H. ♀.	
11 vulgäris. w.	(ye.) common.	Britain.	1. 12. H. ♀.	Eng. bot. v. 11. t. 747.
12 lanuginosus. s.s.	(ye.) woolly.	1827.	4. 8. H. ⊖.	
13 valerianæfolius.s.s.(ye.)	Valerian-leav'd.	Brazil.	1826.	6. 10. F. ⊖.	
14 peucedanifolius.w.(w.)	Peucedanum-leav'd.	C. B. S.	1816.	6. 8. G. ♀.	Jacq. ic. 3. t. 581.
<i>Calatia peucedanifolia. Jacq.</i>					
15 arabicus. w.	(ye.) Arabian.	Egypt.	1804.	7. 8. H. ♂.	
16 verbenæfolius.w.(ye.)	Vervain-leav'd.	1803.	6. 7. H. ⊖.	Jacq. vind. 1. t. 3.
17 glomeratus. L.en.	(ye.) close-flowered.	N. Holland.	1816.	8. 10. H. ⊖.	
18 cacalioïdes. L.en.	(ye.) Cacalia-like.	Brazil.	1820.	— H. ⊖.	
19 triflorus. w.	(ye.) three-flowered.	Egypt.	1776.	7. 9. H. ⊖.	
20 ægyptius. w.	(ye.) Egyptian.	1771.	7. 8. H. ⊖.	
21 crassifolius. w.	(ye.) thick-leaved.	S. Europe.	1815.	— H. ⊖.	
22 lividus. w.	(ye.) livid.	Britain.	— H. ⊖.	Eng. bot. v. 35. t. 2515.
23 telephifolius. w.	(ye.) Telephium-leav'd.	C. B. S.	1823.	— H. ⊖.	Jacq. frag. 1. t. 1. f. 3.
24 trilobus. w.	(ye.) three-lobed.	Spain.	1728.	6. 8. H. ⊖.	
25 cinerascens. w.	(ye.) gray.	C. B. S.	1774.	5. 7. G. ♂.	Jacq. schoen. 2. t. 150.
26 lilacinus. s.s.	(ti.) lilac-coloured.	1826.	6. 12. G. ♀.	Link et Ott. abb. ic.
27 viscosus. w.	(ye.) stinking.	Britain.	6. 10. H. ⊖.	Eng. bot. v. 1. t. 32.
28 sylvaticus. w.	(st.) mountain.	7. 8. H. ⊖.	— v. 11. t. 748.
29 coronopifolius. w.(ye.)	Buck'shorn-leav'd.	Spain.	1821.	— H. ⊖.	
30 nebrodensis. w.	(ye.) Nebrodes.	S. Europe.	1704.	6. 8. H. ⊖.	Barrel. rar. 401.
31 glaucus. w.	(ye.) sea-green.	Egypt.	1739.	— H. ⊖.	
32 Marmoræ. s.s.	(ye.) Sardinian.	Sardinia.	1829.	4. 5. H. ♂.	
33 hastatus. w.	(ye.) halbert-leaved.	C. B. S.	1722.	5. 8. G. ♀.	Dill. elth. t. 152. f. 184.
34 pubigerus. w.	(ye.) runcinate-leav'd.	1816.	— G. ♀.	Moris. s. 7. t. 18. f. 32.
35 vernalis. w.	(ye.) spring.	Hungary.	1803.	4. 6. H. ⊖.	W. et K. hung. 1. t. 24.
36 montanus. w.	(ye.) smooth mountain.	Germany.	1819.	— H. ⊖.	
37 rupestris. w.	(ye.) rock.	Hungary.	1805.	6. 7. H. ♀.	W. et K. hung. 2. t. 128.
38 lacinata. Bért.	(ye.) jagged.	Italy.	1822.	5. 8. H. ♂.	
39 dentatus. w.	(ye.) toothed-leaved.	C. B. S.	1823.	— G. ♀.	
40 squamosus. w.	(ye.) scaly.	1816.	— G. ♀.	
41 venustus. w.	(re.wh.) wing-leaved.	1774.	5. 11. G. ♀.	Bot. reg. t. 901.
42 élégans. w.	(re.wh.) elegant.	1700.	6. 8. H. ⊖.	
α rüber.	(re.) red-flowered.	—	— H. ⊖.	
β álbis.	(wh.) white-flowered.	—	— H. ⊖.	
γ pleno-álbis.	(wh.) double-white.	—	4. 11. G. ♀.	
δ pleno-rüber.	(re.) double red.	—	— G. ♀.	
43 speciosus. w.	(re.) red-flowered.	China.	1789.	7. 8. G. ♀.	Bot. mag. t. 238.
44 squálidus. w.	(ye.) inelegant.	England.	6. 10. H. ⊖.	Bot. reg. t. 41.
45 erucæfolius. w.	(ye.) Eruca-leaved.	Europe.	1816.	7. 8. H. ♀.	Eng. bot. v. 9. t. 600.
46 chrysanthemifolius.(ye.)	Chrysanthemum-ld.	Sicily.	1823.	7. 9. H. ♀.	
47 uniflorus. w.	(ye.) one-flowered.	S. Europe.	1799.	7. 8. H. ♀.	Bocc. sic. t. 67.
48 incanus. w.	(ye.) downy.	1759.	— H. ♀.	All. ped. t. 17. f. 8.
49 abrotanifolius. w.(ye.)	Southernwood-leav'd.	1640.	7. 10. H. ♀.	Pluk. alm. t. 39. f. 6.
50 artemisiæfolius.P.s.(ye.)	Wormwood-leav'd.	France.	1820.	— H. ♀.	Jacq. aust. 1. t. 79.
51 præaltus. s.s.	(ye.) lofty.	Italy.	1827.	7. 8. H. ♀.	
52 Othónnae. M.B.	(ye.) Othonna-like.	Caucasus.	1822.	— H. ♀.	
53 Marschallianus.s.s.(ye.)	Bieberstein's.	Ukraine.	1826.	— H. ♀.	
arenarius. M.B. non Thunb.					
54 grandiflorus. s.s.	(ye.) large-flowered.	C. B. S.	1824.	6. 8. G. ♀.	Pluk. mant. t. 422. f. 5.
55 glaucescens. s.s.	(ye.) glaucescient.	1826.	6. 10. H. ⊖.	
56 carniolicus. s.s.	(ye.) Carniolian.	Carniola.	1827.	7. 8. H. ♀.	
57 adonisifolius.Lois.(ye.)	Adonis-leaved.	1826.	7. 10. H. ♀.	
58 dahuricus. F.	(ye.) Dahurian.	Dahuria.	1821.	— H. ♀.	
59 umbellátus.s.s.(ye.pu.)	umbel-flowered.	C. B. S.	1822.	6. 8. G. ♀.	
60 canadensis. w.	(ye.) Canadian.	N. America.	1818.	— H. ♀.	
61 delphinifolius. w.	(ye.) Larkspur-leav'd.	Sicily.	1823.	— H. ♀.	Vahl symb. 2. t. 43.
62 tenuifolius. w.	(ye.) slender-leaved.	Britain.	7. 8. H. ♀.	Eng. bot. v. 8. t. 574.
63 Jacob'æa. w.	(ye.) common-ragwort.	— H. ♀.	— v. 16. t. 1130.
64 erraticus. s.s.	(ye.) water.	S. Europe.	1826.	5. 7. H. ♀.	
65 aquáticus. w.	(ye.) marsh.	— H. ♀.	
66 aureus. w.	(ye.) golden.	N. America.	1758.	7. 8. H. ♀.	Eng. bot. v. 16. t. 1131.

67	lyratifolius. s.s.	(ye.) lyrate-leaved.	S. Europe.	1826.	5. 6. H. 24.
68	fragrans. s.s.	(ye.) fragrant.	Siberia.	—	6. 8. H. 24.
69	Balsamitæ. w.	(ye.) various-leaved.	N. America.	1819.	— H. 24.
70	obovatæ. w.	(ye.) obovate-leaved.	—	—	H. 24.
71	leucophyllus. DC.	(ye.) white-leaved.	S. Europe.	1821.	— H. 24.
72	rosmarinifolius. w.	(ye.) Rosemary-leav'd.	C. B. S.	...	7. 8. G. h.
73	asper. w.	(ye.) rough.	—	1774.	— G. h.
74	linifolius. w.	(ye.) Flax-leaved.	Spain.	1817.	— H. 24.
75	rigescens. w.	(ye.) stiff-leaved.	C. B. S.	1815.	— G. h.
76	nervatæ. s.s.	(pu.) nerved-leaved.	—	1824.	7. 8. G. h.
77	paludosus. w.	(ye.) Bird's-tongue.	England.	...	6. 8. H. 24.
78	periscæfolius. DC.	(ye.) peach-leaved.	Pyrenees.	1817.	— H. 24.
79	nemorénsis. w.	(ye.) branching.	Austria.	1785.	7. 8. H. 24.
80	sarracénicus. w.	(ye.) creeping-root'd.	Britain.	...	7. 10. H. 24.
81	ovatæ. w.	(ye.) oval-leaved.	Germany.	1817.	— H. 24.
82	coriáceus. w.	(ye.) leathery-leaved.	Levant.	1728.	7. 8. H. 24.
83	Dòria. w.	(ye.) broad-leaved.	Austria.	1570.	7. 9. H. 24.
84	macrophyllus. M.B.	(ye.) large-leaved.	Caucasus.	1817.	— H. 24.
85	Dorónicum. w.	(ye.) Leopard's-bane.	S. Europe.	1705.	— H. 24.
86	umbrosus. w.K.	(ye.) ear-leaved.	Hungary.	1815.	— H. 24.
87	Barrelieri. w.	(ye.) Barrelier's.	Pyrenees.	1824.	— H. 24.
88	lanceus. w.	(ye.) spear-leaved.	C. B. S.	1774.	7. 10. G. h.
89	oporus. w.	(ye.) Jacquin's.	—	—	G. 24.
90	longifolius. w.	(ye.) long-leaved.	—	1775.	8. 11. G. h.
91	halimifolius. w.	(ye.) succulent-leaved.	—	1723.	7. G. h.
92	ilicifolius. w.	(ye.) Ilex-leaved.	—	1731.	6. 7. G. h.
93	rígidus. w.	(ye.) hard-leaved.	—	1704.	6. 9. G. h.
94	solidaginoides. w.	(ye.) Golden-rod-like.	—	1819.	— G. h.
95	vérnus. Biv.	(ye.) vernal.	Italy.	—	— H. 24.
96	cordifolius. s.s.	(ye.) heart-leaved.	C. B. S.	1827.	7. 9. G. 24.
97	arenárius. s.s.	(ye.) sand.	—	1826.	— G. 24.
98	spicátus.	(ye.) spike-flowered.	—	1828.	— H. 24.
ADENOTRI'CHA. B.R.	ADENOTRI'CHA.	Syngenesia	Polygamia Superflua.		
amplexicaulis. B.R.	(ye.) stem-clasping.	Chile.	1826.	5. 7. F. 24.	
SENECI'LLIS. G.	SENECI'LLIS.	Syngenesia	Polygamia Superflua.		
glauca. G.	(ye.) glaucous.	Siberia.	1790.	6. 7. H. ♂.	
Cineraria glauca. w.					
CINER'ARIA. W.	CINER'ARIA.	Syngenesia	Polygamia Superflua. L.		
1	americana. w.	(ye.) American.	S. America.	1823.	... S. h.
2	incána. P.S.	(ye.) hoary.	Jamaica.	—	6. 8. S. h.
3	discolor. P.S.	(ye.) two-coloured.	—	1822.	— S. h.
4	lúcida. P.S.	(ye.) glossy.	W. Indies.	—	— S. h.
5	glabrata. s.s.	(ye.) smooth.	—	1827.	4. 7. S. h.
6	geifolia. w.	(ye.) kidney-leaved.	C. B. S.	1710.	4. 8. G. h.
7	canescens. L.en.	(ye.) small-flowered.	—	1790.	— G. h.
8	aúrita. w.	(pu.) purple-flower'd.	Madeira.	1777.	6. 7. G. h.
9	láctea. W.en.	(wh.) milk-coloured.	—	1816.	— G. h.
10	lanáta. w.	(pu.) woolly.	Canaries.	1780.	5. 9. G. h.
11	hy'brid. W.en.	(ye.) hybrid.	Hybrid.	—	2. 5. G. h.
12	éláti. L.en.	(wh.) tall.	—	—	— G. h.
13	cruénta. w.	(pu.) purple-leaved.	Canaries.	1777.	— G. 24.
	aúrita. A.R. 24.	nec aliorum.			
14	pulchélla.	(re.) bright red-flower'd.	—	1818.	— G. 24.
15	populifolia. H.K.	(re.) Poplar-leaved.	—	1780.	6. 9. G. h.
16	hypoleúca. s.s.	(pu.) snowy-leaved.	C. B. S.	1828.	4. 7. G. h.
17	venústa. s.s.	(re.) charming.	—	—	G. 24.
18	lobatæ. w.	(ye.) lobed.	—	1774.	6. 8. G. h.
19	malvæfolia. w.	(li.) Mallow-leaved.	Azores.	1777.	8. G. 24.
20	petasites. B.M.	(ye.) butter-bur-leav'd.	Mexico.	1812.	12. 2. G. h.
21	præcox. w.	(ye.) early-flowering.	—	1824.	2. 4. G. h.
22	sibírica. w.	(ye.) Siberian.	Siberia.	1784.	6. 8. H. 24.
23	speciosa. L.en.	(ye.) large-flowered.	—	1818.	— H. 24.
24	caucásica. M.B.	(ye.) Caucasian.	Caucasus.	1820.	— H. 24.
25	gigántea. H.K.	(bh.) gigantic.	Cape Horn.	1801.	7. 8. H. 24.
26	lævis. s.s.	(ye.) smooth.	C. B. S.	1828.	4. 7. G. 24.
27	crenata. s.s.	(st.) crenate.	—	—	— G. 24.
28	palústris. w.	(ye.) marsh.	England.	—	— H. 24.
29	aurantiaca. w.	(or.) orange-colour'd.	Switzerland.	1818.	5. 7. H. 24.
30	campéstris. w.	(ye.) mountain.	N. Europe.	—	H. ♂.

Jacq. ic. 3. t. 587.

Barrel. rar. t. 802.

Jacq. coll. 5. t. 6. f. 1.

Eng. bot. v. 10. t. 650.

Ram. bul. phil. t. 11. f. 3.

Jacq. aust. 2. t. 184.

Eng. bot. v. 31. t. 2211.

Dill. elth. t. 105. f. 125.

Jacq. aust. 2. t. 185.

Jacq. aust. 5. t. ap. 45.

Barrel. ic. 146.

Jacq. schæn. 3. t. 304.

Comm. hort. 2. t. 71.

Dill. elth. t. 104. f. 124.

Comm. rar. t. 42.

Comm. hort. 2. t. 75.

Hout. syst. 4. t. 32. f. 1.

Bot reg. t. 1190.

Gmel. sib. 2. t. 74.

Plum. 2. ic. 154.

Comm. hort. 2. t. 73.

Bot. mag. t. 1990.

— t. 1786.

Bot. mag. t. 53.

Bot. mag. t. 406.

Venten. malm. t. 100.

Bot. mag. t. 1536.

DC. pl. rar. gen. t. 7.

Bot. mag. t. 1869.

Bot. reg. t. 812.

Sm. exot. bot. 2. t. 65.

Eng. bot. v. 3. t. 151.

Swt. br. fl. gar. t. 256.

Jacq. aust. t. 180.

31 integrifolia. E.B. (<i>ye.</i>) entire-leaved.	England.	5. 7.	H. 4.	Eng. bot. v. 3. t. 152.	
32 longifolia. w. (<i>ye.</i>) long-leaved.	Austria.	1792.	6. 8.	H. 4.	Jacq. aust. 2. t. 181.	
33 crista. w. (<i>ye.</i>) curled.	1818.	H. 4. 2. t. 178.	
34 cordifolia. w. (<i>ye.</i>) heart-leaved.	1739.	7. 8.	H. 4. 2. t. 176.	
35 media. W.K. (<i>ye.</i>) intermediate.	Hungary.	1818.	H. 4.	Jacq. aust. 2. t. 177.	
36 alpina. w. (<i>ye.</i>) Alpine.	Austria.	1683.	6. 8.	H. 4.		
37 aurea. w. (<i>ye.</i>) golden.	Siberia.	1825.	H. 4.		
38 rivularis. s.s. (<i>ye.</i>) rivulet.	Hungary.	1822.	H. 4.		
39 crassifolia. s.s. (<i>ye.</i>) thick-leaved.	Carinthia.	1825.	H. 4.		
40 alpestris. s.s. (<i>ye.</i>) rough-leaved.	N. Europe.	1826.	H. 4.		
41 spathulaefolia.s.s. (<i>ye.</i>) spatulate-leav'd.	Germany.	1820.	H. 4.		
42 papposa. s.s. (<i>ye.</i>) long-chaffed.	Galicia.	1828.	H. 4.		
43 parviflora. s.s. (<i>ye.</i>) small-flowered.	Caucasus.	1827.	H. 4.		
44 congesta. s.s. (<i>ye.</i>) close-headed.	Melville Isl.	5. 6.	H. 4.		
45 taraxacifolia.s.s. (<i>ye.</i>) Dandelion-leav'd.	Caucasus.	1824.	6. 8.	H. 4.		
46 exsquamata. (<i>ye.</i>) scaleless.	Portugal.	1827.	H. ⊙.		
<i>Senecio exsquameus.</i> Brot.						
47 frigida. s.s. (<i>ye.</i>) frigid.	N. America.	1827.	5. 6.	H. 4.		
48 scapiflora. s.s. (<i>ye.</i>) long-peduncled.	C. B. S.	1828.	G. 4.		
49 gibbosa. s.s. (<i>ye.</i>) gibbous-headed.	Sicily.	6. 8.	H. ½.		
50 Aitoniana. s.s. (<i>ye.</i>) Aiton's.	G. ½.		
51 maritima. w. (<i>ye.</i>) sea ragwort.	S. Europe.	1633.	7. 9.	F. ½.	Flor. græc. t. 871.	
52 bicolor. w. (<i>ye.</i>) shining-leaved.	1820.	F. ½.		
53 acanthifolia. s.s. (<i>ye.</i>) Acanthus-leav'd.	Levant.	1826.	G. ½.		
54 thyrsoidea. F. (<i>ye.</i>) thyrs-flower'd.	Armenia.	1827.	H. 4.		
55 angulata.	(<i>ye.</i>) angular-leaved.	Mexico.	12. 1.	G. 4.	
56 canadensis.	(<i>ye.</i>) Canadian.	Canada.	1739.	6. 8.	H. 4.	Jacq. schœn. 3. t. 308.
57 linifolia. w. (<i>ye.</i>) flax-leaved.	C. B. S.	G. ½.		
58 angustifolia. K.S. (<i>ye.</i>) narrow-leaved.	Mexico.	1825.	G. ½.		
59 humifusa. w. (<i>ye.</i>) trailing.	C. B. S.	1704.	7. 8.	G. 4.	Jacq. frag. 12. t. 7. f. 2	
60 viscosa. w. (<i>wh.</i>) clammy.	1774.	6. 8.	G. ½.		
61 vestita. H.R. (<i>ye.</i>) clothed.	1824.	5. 6.	G. 4.		
AGATHÆA. Cas.	AGATHÆA.	Syngenesia Polygamia Superflua.				
cœlestis. Cas.	(bl.) Cape-Aster.	C. B. S.	1753.	2. 9.	G. ½.	Bot. mag. t. 249.
<i>Cineraria amelloides.</i> w.						
KAULFUSSIA. Nees.	KAULFUSSIA.	Syngenesia Polygamia Superflua.				
1 amelloides. Nees. (bl.) blue-flowered.	C. B. S.	1819.	6. 7.	H. ⊙.	Bot. reg. t. 490.	
2 ciliata. s.s. (st.) fringed.	1822.	6. 9.	H. ⊙.		
TAGETES. W.	TAGETES.	Syngenesia Polygamia Superflua. L.				
1 lúcida. w. (<i>ye.</i>) sweet-scented.	Mexico.	1798.	7. 11.	F. 4.		
2 flórida. B.F.G. (<i>ye.</i>) gay-flowering.	1827.	8. 11.	F. 4.		
3 pátula. w. (<i>pu. ye.</i>) French Marygold.	1573.	7. 10.	H. ⊙.		
4 corymbosa. B.F.G. (va. corymbus-flower'd.	1825.	9. 11.	H. ⊙.		
<i>a purpurea.</i> (<i>pu.</i>) purple-flowered.	H. ⊙.		
<i>β lutea.</i> (<i>ye.</i>) yellow-flowered.	H. ⊙.		
5 erécta. w. (<i>ye.</i>) African Marygold.	1596.	6. 9.	H. ⊙.		
6 tenuifolia. w. (<i>ye.</i>) fine-leaved.	1797.	7. 10.	H. ⊙.	Lam. ill. t. 684.	
7 subvillösa. LG.	(<i>ye.</i>) villous-leaved.	1825.	8. 11.	H. ⊙.	Swt. br. fl. gar. t. 141.
8 minuta. w. (<i>ye.</i>) small-flowered.	Chili.	1728.	8. 10.	H. ⊙.	Dill. elth. t. 280. f. 362.	
9 clandestina. LG. (<i>ye.</i>) small-rayed.	Mexico.	1822.	H. ⊙.		
<i>coronopifolia.</i> W.en.					Cavan. ic. 4. t. 352.	
10 micrantha. w. (<i>ye.</i>) solitary-flower'd.	H. ⊙.		
11 glandulosa. L.en. (<i>ye.</i>) glandular.	1825.	H. ⊙.		
12 glandulifera. Sck. (<i>ye.</i>) gland-bearing.	1826.	9. 11.	H. ⊙.		
BŒB'ERA. W.	BŒB'ERA.	Syngenesia Polygamia Superflua.			Venten. cels. t. 36.	
1 glandulosa. n. (<i>ye.</i>) glandular.	N. America.	1823.	8. 9.	H. ⊙.		
<i>chrysanthemoïdes.</i> W.en. <i>Tagetes papposa.</i> v. <i>Dyssodia glandulosa.</i> Cav.						
2 Porophyllum. K.S. (<i>ye.</i>) Mexican.	Mexico.	H. ⊙.		
<i>Dyssodia Porophyllum.</i> W.en.						
3 fastigiata. K.S. (<i>ye.</i>) clustered.	1827.	9. 11.	H. ⊙.		
4 divaricata. (<i>ye.</i>) spreading.	1828.	H. ⊙.		
<i>Dyssodia divaricata.</i> Rich.						
5 pubescens. s.s. (<i>ye.</i>) pubescent.	New Spain.	1826.	8. 10.	H. ⊙.	Cavan. icon. 3. t. 212.	
<i>Dyssodia pubescens.</i> LG. <i>A'ster pinnatus.</i> Cavan.						
OTHO'NNA. W.	RAGWORT.	Syngenesia Polygamia Necessaria. L.				
1 flabellifolia. B.C. (<i>ye.</i>) fan-leaved.	C. B. S.	1822.	7. 9.	G. ½.	Lodd. bot. cab. t. 728.	
2 pinnatifida. P.S. (<i>ye.</i>) pinnatifid.	1823.	G. 4.	Bot. mag. t. 768.	
3 pinnata. w. (<i>ye.</i>) wing-leaved.	1759.	4. 6.	G. 4.		
4 ciliata. w. (<i>ye.</i>) fringed-leaved.	1824.	5. 6.	G. 4.		
5 Tagetes. w. (<i>ye.</i>) slender-stalked.	1823.	7. 9.	H. ⊙.		

6 pectinata. w. (ye.)	Wormwood-leav'd.C. B. S.	1731.	4. 6.	G. $\frac{1}{2}$.	Bot. mag. t. 306.
7 Athanasiæ. w. (ye.)	Athanasia-like.	1795.	11. 12.	G. $\frac{1}{2}$.	Jacq. schoen. 2. t. 242.
8 abrotanifolia. w. (ye.)	Southernwood-leav'd.	1692.	1. 5.	G. $\frac{1}{2}$.	Bot. reg. t. 108.
9 digitata. w. (ye.)	finger-leaved.	1822.	6. 9.	G. $\frac{1}{2}$.	
10 trifurcata. w. (ye.)	three-forked.			G. $\frac{1}{2}$.	
11 trifida. w. (ye.)	trifid-leaved.	1823.	—	G. $\frac{1}{2}$.	
12 retrofracta. w. (ye.)	axillary-flower'd.	1812.	—	G. $\frac{1}{2}$.	Jacq. schoen. 3. t. 376.
13 coronopifolia. w. (ye.)	Buckshorn-leav'd.	1731.	7. 9.	G. $\frac{1}{2}$.	Comm. hort. 2. t. 70.
14 cheirifolia. w. (ye.)	Stock-leaved.	1752.	4. 6.	H. $\frac{1}{2}$.	Bot. reg. t. 266.
15 crassifolia. w. (ye.)	thick-leaved.	C. B. S.	1710.	9. 10.	Mill. ic. 2. t. 245. f. 2.
16 denticulata. w. (ye.)	tooth-leaved.		1774.	4. 7.	Bot. mag. t. 1979.
17 heterophylla. w. (ye.)	various-leaved.		1812.	—	
18 Lingua. w. (ye.)	tongue-leaved.		1787.	5. 9.	G. $\frac{1}{2}$.
19 filicaulis. w. (ye.)	Yam-rooted.		1791.	4. 5.	G. $\frac{1}{2}$.
20 bulbosa. w. (ye.)	bulbous.		1774.	5. 6.	G. $\frac{1}{2}$.
21 capillaris. w. (ye.)	capillary-stalked.		1822.	6. 9.	H. \odot .
22 parviflora. w. (ye.)	small-flowered.		1704.	7. 8.	G. $\frac{1}{2}$.
23 perfoliata. H.K. (ye.)	perfoliolate.		1789.	5. 7.	G. $\frac{1}{2}$.
amplexicaulis. B.M. (ye.)	amplexicaulis.	1312. non Thunb.			Volk. norib. t. 226.
24 ericoides. w. (ye.)	Heath-leaved.		1815.	7. 8.	Jacq. schoen. 2. t. 238.
25 linifolia. w. (ye.)	Flax-leaved.		1825.	—	— 2. t. 241.
26 tenuissima. w. (ye.)	fine-leaved.		1759.	4. 7.	Breyn. cent. t. 66.
27 frutescens. w. (ye.)	frutescent.		1816.	7. 8.	
28 arborescens. w. (ye.)	tree.		1723.	—	Dill. elth. t. 103. f. 123.
29 cacaioides. w. (ye.)	tuberous.		1774.	5. 9.	
30 amplexicaulis. w. (ye.)	stem-clasping.		1826.	—	

Subordo VI. HELIANTHEÆ. Kth. synops. 2. p. 463.

HYMENOPAPPUS. J. HYMENOPAPPUS. Syngenesia Polygamia Äqualis.

1 pedatus. L.G. (wh.)	pedate-leaved.	Mexico.	1803.	7. 9.	G. \odot .	Cavan. ic. 4. t. 356.
Stévia pedata. B.M. 2040.	Florestina pedata. Cas.					
2 scabiösæs. Ph. (wh.)	Scabious-like.	Missouri.	1812.	6. 8.	H. \odot .	Jour. de hist. nat. n. 1. ic.
3 tenuifolius. Ph. (wh.)	slender-leaved.		1811.	—	F. δ .	
SCHK'UHRIA. W.	SCHK'UHRIA.	Syngenesia Polygamia Superflua. L.				Schk. hand. 3. t. 250. b.
abrotanoïdes. w. (wh.)	Wormwood-leav'd.	Mexico.	1798.	7. 9.	H. \odot .	
PE'CTIS. W.	PE'CTIS.	Syngenesia Polygamia Superflua. L.				
1 humifusa. w. (ye.)	trailing.	W. Indies.	1822.	7. 9.	S. \odot .	Plum. ic. t. 95. f. 2.
2 prostrata. w. (ye.)	prostrate.	New Spain.	1824.	—	H. \odot .	Cavan. ic. 4. t. 324.
3 ciliaris. w. (ye.)	ciliated.	Hispaniola.	1793.	7.	S. \odot .	Plum. ic. 151. f. 2.
4 linifolia. w. (ye.)	Flax-leaved.	Jamaica.	1732.	7. 8.	S. \odot .	Sloan. hist. I. t. 149. f. 3.
5 canescens. k.s. (ye.)	canescent.	Mexico.	1827.	8. 9.	H. \odot .	H. et B. n. gen. 4. t. 393.
6 punctata. w. (ye.)	dotted.	Hispaniola.	1824.	6. 8.	S. \odot .	Jacq. amer. t. 126.

HETEROSPERMUM. HETEROSPERMUM. Syngenesia Polygamia Superflua. W.

pinnatum. w. (ye.)	wing-leaved.	New Spain.	1799.	8. 9.	H. \odot .	Cavan. ic. 3. t. 267.
MELANANTHERA. M.	MELANANTHERA.	Syngenesia Polygamia Äequalis. W.				P.S.
1 linnæi. k.s. (li.)	Linnæus's.	S. America.	1799.	6. 8.	S. δ .	Jacq. ic. 3. t. 583.

BIDENS. L. Càlea áspéra. Jacq.

2 hastata. M. (wh.)	hastate-leaved.	N. America.	1732.	—	G. $\frac{1}{2}$.	Dill. elth. t. 47. f. 55.
β pandurata. Ph. (wh.)	fiddle-leaved.			—	G. $\frac{1}{2}$.	— t. 46. f. 54.

MARSHALLIA. W.

MARSHALLIA. W.	Syngenesia Polygamia Äequalis. W.					
1 lanceolata. Ph. (li.)	spear-leaved.	Carolina.	1812.	6. 7.	F. $\frac{1}{2}$.	
2 latifolia. Ph. (li.)	broad-leaved.		1806.	—	F. $\frac{1}{2}$.	

PERSOONIA. latifolia. M. Trattenickia latifolia. P.S.

PLATYPTERIS. K.S.	PLATYPTERIS.	Syngenesia Polygamia Äequalis.				
crocatæ. k.s. (sn.)	saffron-colour'd.	Mexico.	1812.	1. 3.	S. $\frac{1}{2}$.	Bot. mag. t. 1627.
Spilanthæ crocæta. B.M.						

SPLA'NTHES. W.

SPLA'NTHES. W.	SPLA'NTHES.	Syngenesia Polygamia Äequalis. L.				
1 tenella. k.s. (st.)	slender.	Caracas.	1823.	8. 10.	G. \odot .	Pluk. alm. t. 159. f. 4.
2 Pseudo-Acmella. w. (ye.)	spear-leaved.	Ceylon.	1768.	7.	S. \odot .	Jacq. ic. 3. t. 584.
3 uliginosa. w. (ye.)	marsh.	Jamaica.	1820.	7. 9.	S. \odot .	L'Herit. stirp. 7. t. 4.
4 exasperata. w. (wh.)	rough-stalked.	Caracas.	1825.	—	S. \odot .	Jacq. vind. 2. t. 135.
5 alba. w. (wh.)	white-flowered.	Peru.	1783.	6. 7.	H. \odot .	
6 oleracea. w. (ye.)	esculent.	E. Indies.	1770.	7. 9.	S. δ .	
7 pallida. r. (st.)	pale-flowered.	Jamaica.	1821.	—	S. \odot .	
8 multiflora. Ot. (ye.)	many-flowered.	1829.	—	S. \odot .	

ACME'LLA. P.S.	ACME'LLA. Syngenesia Polygamia Superflua. P.S.			
1 mauritiâna. p.s. (ye.)	Balm-leaved. Mauritus. 1768. 7. 8. S. ⊖.	Rumph. amb. 6. t. 65.		
<i>Spilánthes Acméllea. w.</i>				
2 répens. p.s.	(ye.) creeping. N. America. 1819. — H. ⊖.			
3 occidentális. p.s. (ye.)	oval-leaved. S. America. 1825. — H. ⊖.			
4 bupthalmoides. p.s. (ye.)	Ox-eye-leaved. — 1798. 7. 9. H. ⊖.	Jacq. schœnb. 2. t. 151.		
<i>A'nthemis bupthalmoides. Jacq.</i>				
LAV'ENIA. W.	LAV'ENIA. Syngenesia Polygamia Æqualis.			
erécta. w.	(ye.) upright. E. Indies. 1739. 7. 9. S. ⊖.	Burm. zeyl. t. 42.		
SA'LMEA. DC.	SA'LMEA. Syngenesia Polygamia Æqualis.			
1 scándens. dc.	(wh.) climbing. S. America. 1815. 4. 6. S. ḥ. ⊖.	Bot. mag. t. 2062.		
<i>Câlea Améllea. L.</i>				
2 hirsuta. dc.	(wh.) hairy. — — — — S. ḥ. ⊖.			
PETR'OB'UM. L.T.	PETR'OB'UM. Syngenesia Polygamia Æequalis.			
arbóreum.	(st.) tree. St. Helena. 1816. S. h.			
<i>Spilánthus arbòreus. Forst.</i>				
CALEA. L.T.	CALEA. Syngenesia Polygamia Æequalis. L.			
1 jamaicénis. l.l.	(li.) purple-flower'd. W. Indies. 1739. 6. 7. S. ȳ.	Sloan. jam. 1.t. 151.f.3.		
2 martinicénis.	(li.) Martinico. Martinico. 1821. S. ȳ.			
3 cordifòlia. l.t.	(ye.li.) heart-leaved. Jamaica. 1822. 6. 7. S. ȳ.			
CALEA'CTE. L.T.	CALEA'CTE. Syngenesia Polygamia Æequalis.			
1 urticifòlia. l.t.	(ye.) Nettle-leaved. Vera Cruz. S. h.			
<i>Solidago urticafòlia. Mill.</i>				
2 pinnatifida. l.t.	(ye.) pinnatifid. Brazil. 1816. 6. 8. S. h.			
ISOCA'RPHA. L.T.	ISOCA'RPHA. Syngenesia Polygamia Æequalis.			
oppositifòlia.	(wh.) opposite-leaved. W. Indies. 1739. 7. 8. S. ♂.			
<i>Câlea oppositifòlia. w.</i>				
NEUROL'ENA. L.T.	NEUROL'ENA. Syngenesia Polygamia Æequalis.			
lobâta.	(ye.) halberd-weed. W. Indies. 1733. 6. 7. S. h.	Bot. mag. t. 1734.		
<i>Câlea lobâta. B.M.</i>				
M'ADIA. W.	M'ADIA. Syngenesia Polygamia Superflua. W.			
1 viscòsa. w.	(ye.) viscous. Chili. 1794. 7. 8. H. ⊖.	Bot. mag. t. 2574.		
2 mellòsa. w.	(ye.) stem-clasping. — 1824. — H. ⊖.			
TETRAGONOTH'ECA. TETRAGONOTH'ECA.	Syngenesia Polygamia Superflua. W.			
helianthoides. w.	(ye.) Sunflower-like. Virginia. 1726. 8. 10. H. ȳ.	Schk. hand. 3. t. 263.		
XIMEN'ESIA. W.	XIMEN'ESIA. Syngenesia Polygamia Superflua. W.			
1 heterophylla. k.s.	(ye.) various-leaved. Mexico. 1827. 8. 10. F. ȳ.			
2 encelioïdes. w.	(ye.) oval-leaved. — 1795. 6. 11. G. ♂.	Cavan. ic. 2. t. 178.		
3 fœtida. k.s.	(ye.) fetid. — 1799. 7. 8. H. ⊖.	——— 1.t. 77.		
<i>Coreopsis fœtida. Cav. Simsia fœtidula. p.s.</i>				
4 cordatâ. k.s.	(ye.) heart-leaved. Mexico. — — F. ȳ.			
HEL'ENIUM. W.	HEL'ENIUM. Syngenesia Polygamia Superflua. W.			
1 autumnâle. w.	(ye.) smooth. N. America. 1729. 8. 10. H. ȳ.	Schk. hand. 3. t. 250.		
2 pubescens. w.	(ye.) downy. — 1776. 8. 9. H. ȳ.			
3 quadripartitum. L.en.	(ye.) four-cleft. — 1825. — G. ♂.			
4 quadridentatum. w.	(ye.) four-toothed. Louisiana. 1790. 5. 10. H. ȳ.	Bot. reg. t. 598.		
<i>Rudbeckia alata. Jacq. ic. t. 193.</i>				
5 pùmulum. W.en.	(ye.) dwarf. N. America. 1819. — H. ȳ.			
6 mexicanum. k.s.	(ye.) Mexican. Mexico. 1823. 8. 10. H. ȳ.			
ZI'NNIA. W.	ZI'NNIA. Syngenesia Polygamia Superflua. W.			
1 pauciflòra. w.	(ye.) yellow-flower'd. Peru. 1753. 7. 8. H. ⊖.	Mill. ic. 1. t. 64.		
2 multiflòra. w.	(re.) red-flowered. N. America. 1770. 6. 10. H. ⊖.	Bot. mag. t. 149.		
3 hybrida. B.M.	(sc.) hybrid. Mexico. 1818. 6. 9. H. ⊖.	——— t. 2123.		
4 verticillâta. w.	(re.) whorl-leaved. — 1789. 7. 8. H. ⊖.	Andr. repos. t. 189.		
5 violacea. c.l.	(pu.) purple-flowered. — 1796. 7. 10. H. ⊖.	——— t. 55.		
<i>élegans. Bot. mag. t. 527.</i>				
β coccinea. B.R.	(sc.) scarlet-flowered. — 1829. — H. ⊖.	Bot. reg. t. 1294.		
6 tenuiflòra. w.	(re.) slender-flower'd. — 1799. 7. 8. H. ⊖.	Bot. mag. t. 555.		
7 angustifolia. k.s.	(re.) narrow-leaved. — 1825. — H. ⊖.			
RELH'ANIA. W.	RELH'ANIA. Syngenesia Polygamia Superflua. W.			
1 squarrosa. w.	(ye.) cross-leaved. C. B. S. 1774. 5. 6. G. ḥ.			
2 genistæfolia. w.	(ye.) Genista-leaved. — 1823. 5. 7. G. ḥ.			
3 lateriflòra. w.	(ye.) side-flowering. — — — G. ḥ.			
4 pungens. w.	(ye.) pungent-leaved. — 1820. 6. 8. G. ḥ.	Bot. reg. t. 587.		
5 pedunculâta. w.	(ye.) dwarf. — 1824. 5. 8. G. ḥ.			
6 paleacea. w.	(ye.) chaffy. — 1825. — G. ḥ.	Gært. fruct. 2.t. 173. f.9.		
<i>Leyséra paleacea. Gært.</i>				
ATHRI'XIA. B.R.	ATHRI'XIA. Syngenesia Polygamia Superflua.			
capensis. B.R.	(li.) Cape. C. B. S. 1823. 6. 8. G. ȳ.	Bot. reg. t. 681.		

- | | | |
|---|--------------------------------------|--------------------------------------|
| MATA'XA. S.S. | MATA'XA. | Syngenesia Polygamia Superflua. |
| capénis, s.s. (ye.) fleshy-leaved. | C. B. S. | 1824. 6. 8. G. ᲃ. |
| <i>Lasiospérnum radiatum</i> . Treviranus. | | |
| LONGCHA'MPIA. W. | LONGCHA'MPIA. | Syngenesia Polygamia Superflua. W. |
| capillifolia, w. (st.) Leysera-like. | Africa. | 1824. 6. 8. H. ○. |
| <i>Gnaphálium leyseroides</i> . Df. <i>Pectis discoidea</i> . H.H. <i>Leyséra capillifolia</i> . Spr. | | |
| SANVIT'ALIA. W. | SANVIT'ALIA. | Syngenesia Polygamia Superflua. W. |
| procumbens, w. (ye.) trailing. | Mexico. | 1798. 7. 8. H. ○. |
| TR'IDAX. L. | TR'DAX. | Syngenesia Polygamia Superflua. L. |
| procumbens, l. (st.) procumbent. | Mexico. | 1804. 7. 8. H. ○. |
| <i>Balbisia elongata</i> . w. | | |
| COLUMEL'lia. W. | COLUME'LLIA. | Syngenesia Polygamia Superflua. W. |
| biennis, w. (ye.) biennial. | C. B. S. | 1820. 7. 9. G. ♂. |
| <i>Nestlèra biennis</i> , s.s. | | Jacq. schœn. 3. t. 301. |
| MEY'ERA. L.T. | MEY'ERA. | Syngenesia Polygamia Superflua. |
| sessilis, s.s. (ye.) sessile-flowered. | W. Indies. | 7. 8. S. ȝ. |
| ECLI'PTA. W. | ECLI'PTA. | Syngenesia Polygamia Superflua. L. |
| 1 erécta, w. (wh.) upright. | S. America. | 1690. 7. 9. H. ○. |
| 2 punctata, w. (wh.) dotted. | W. Indies. | 1822. — S. ○. |
| 3 prostrata, w. (wh.) trailing. | E. Indies. | 1732. — S. ○. |
| 4 procumbens, M. (wh.) procumbent. | Carolina. | 1819. — H. ○. |
| 5 lineáris, Ot. (wh.) linear-leaved. | | 1825. — S. ○. |
| CHRYSANTHE'LLUM. CHRYSANTHE'LLUM. | Syngenesia Polygamia Superflua. P.S. | |
| procumbens, p.s. (ye.) procumbent. | W. Indies. | 1768. 6. 7. S. ○. |
| <i>Verbesina mutica</i> , w. | | Swartzobs. 314.t.8.f.1. |
| SIEGESBE'CKIA. W. | SIEGESBE'CKIA. | Syngenesia Polygamia Superflua. L. |
| 1 orientalis, w. (ye.) oriental. | E. Indies. | 1730. 8. 10. H. ○. |
| 2 triangulàris, C.I. (ye.) triangular-leav'd. | Mexico. | — H. ○. |
| 3 ibérica, w. (ye.) Iberian. | Iberia. | 1823. — H. ○. |
| 4 droseroides, B.F.G. (ye.) Sundew-like. | Mexico. | 1825. — F. ȝ. |
| 5 flosculosa, w. (ye.) small-flowered. | Peru. | 1784. 6. 7. H. ○. |
| VERBESINA. W. | VERBESINA. | Syngenesia Polygamia Superflua. L. |
| 1 alata, w. (or.) wing-stalked. | S. America. | 1699. 5. 10. G. ȝ. |
| 2 chinensis, w. (or.) Chinese. | China. | 1814. — G. ȝ. |
| 3 virginica, w. (wh.) white-flowered. | N. America. | 1812. 7. 9. H. ȝ. |
| 4 virgata, w. (ye.) slender. | Mexico. | 1825. — H. ȝ. |
| 5 gigantea, w. (wh.) tree. | W. Indies. | 1758. — S. h. |
| 6 Boswellia, L. (ye.) eatable. | E. Indies. | 1818. — S. ○. |
| 7 salicifolia, K.S. (ye.) Willow-leaved. | Mexico. | 1824. 10. 11. G. ȝ. |
| 8 tridentata, s.s. (ye.) three-toothed. | W. Indies. | 1759. 6. 8. S. h. |
| 9 atriplicifolia, s.s. (st.) Atriplex-leav'd. | Trinidad. | 1822. 9. 11. S. h. |
| 10 buphthalmoides, s.s. (ye.) procumbent. | Monte Video. | 1827. 6. 8. F. ᲃ. |
| 11 serrata, w. (ye.) saw-leaved. | Mexico. | 1803. 7. 10. G. ȝ. |
| 12 Siegesbeckia, w. (ye.) American. | Virginia. | 1731. 10. 11. H. ȝ. |
| 13 dichotoma, w. (ye.) forked. | E. Indies. | 1789. 6. 7. S. ○. |
| 14 sativa, B.M. (ye.) oil-seed. | | 1806. 8. 9. S. ○. |
| 15 biflora, w. (ye.) two-flowered. | | 1818. 7. 9. S. ○. |
| 16 calendulacea, w. (ye.) Marygold-flower'd. | Ceylon. | 1739. — S. ○. |
| 17 fruticosa, w. (ye.) shrubby. | W. Indies. | 1759. 6. 8. S. h. |
| 18 pinnatifida, w. (st.) pinnatifid. | Mexico. | 1824. 8. 10. F. ȝ. |
| SYNEDRE'LLA. P.S. | SYNEDRE'LLA. | Syngenesia Polygamia Superflua. |
| nodiflora, p.s. (ye.) sessile-flowered. | W. Indies. | 1726. 6. 7. S. ○. |
| ENC'ELIA. J. | ENC'ELIA. | Syngenesia Polygamia Superflua. P.S. |
| 1 canescens, p.s. (ye.) canescent. | Mexico. | 1786. 7. 9. G. ᲃ. |
| <i>Pallásia halimifolia</i> , w. <i>Coreopsis limensis</i> . Jacq. ic. 3. t. 594. | | |
| 2 halimifolia, p.s. (ye.) great-flowered. | Mexico. | 1825. — G. ᲃ. |
| <i>Pallásia grandiflora</i> , w. | | |
| GALINS'OGEA. K.S. | GALIN'S'OGEA. | Syngenesia Polygamia Superflua. W. |
| 1 trilobata, w. (ye.) three-lobed. | Peru. | 1797. 8. 11. H. ○. |
| 2 balbisioïdes, K.S. (ye.) Mexican. | Mexico. | 1825. — H. ○. |
| PTILOSTE'PHIUM. K.S. PTILOSTE'PHIUM. | Syngenesia Polygamia Superflua. | |
| 1 coronopifolium, K.S. (ye.) Buckshorn-leav'd. | Mexico. | 1823. 8. 10. H. ○. |
| 2 trifidum, K.S. (ye.) trifid-leaved. | | — H. ○. |
| WIBO'RGIA. K.S. | WIBO'RGIA. | Syngenesia Polygamia Superflua. |
| parviflora, K.S. (wh.) small-flowered. | S. America. | 1796. 6. 8. H. ○. |
| <i>Galinsoga parviflora</i> , w. <i>Wiborgia Acmella</i> . Roth. | | |
| PASC'ALIA. W. | PASC'ALIA. | Syngenesia Polygamia Superflua. W. |
| glaucia, w. (ye.) glaucous-leav'd. | Chili. | 1799. 6. 8. H. ȝ. |
| Andr. deposit. t. 549. | | |

HELIO'PSIS. P.S.	HELIO'PSIS. Syngenesia Polygamia Superflua. P.S.		
1 canescens. k.s. (<i>ye.</i>) hoary-leaved.	S. America. 1820. 8. 10. H. ȝ.		Bot. reg. t. 592.
2 lævis. p.s. (<i>ye.</i>) smooth.	Mexico. 1714. 7. 10. H. ȝ.		L'Herit. stirp. t. 45.
3 helianthoides. (<i>ye.</i>) roughish.	N. America. — — H. ȝ.		
<i>Buphthalmum helianthoides.</i> L.			
4 scabra. s.s. (<i>ye.</i>) rough-leaved.	— — — — H. ȝ.		
DIOM'EDEA. Cas.	DIOM'EDEA. Syngenesia Polygamia Superflua.		
1 glabrata. k.s. (<i>ye.</i>) smooth.	S. America. 1699. 5. 9. G. ḥ.		Dill. elth. t. 38. f. 43.
<i>Buphthalmum arboréns.</i> w.			
2 bidentata. s.s. (<i>ye.</i>) two-toothed.	Carolina. 1696. 6. 8. G. ḥ.		Catesb. carol. 1. t. 93.
<i>Buphthalmum fruticosum.</i> Ph.—Dill. elth. t. 28. f. 4.			
3 argentea. k.s. (<i>ye.</i>) silvery-leaved.	Cuba. 1823. — G. ḥ.		
BUPHTHA'LUM.	OX-EYE. Syngenesia Polygamia Superflua. L.		
1 stenophyllum. s.s. (<i>ye.</i>) linear-leaved.	Canaries. 1816. 5. 7. G. ḥ.		
2 elegans. s.s. (<i>ye.</i>) elegant.	— — — — G. ḥ.		
3 laevigatum. s.s. (<i>ye.</i>) smooth-leaved.	Teneriffe. 1824. — G. ḥ.		
4 flosculosum. s.s. (<i>ye.</i>) rayless.	Mesopotamia. 1827. — G. ḥ.		
5 sericeum. w. (<i>ye.</i>) silky.	Canaries. 1779. 5. 7. G. ḥ.		Vent. h. cels. t. 25.
6 spinosum. w. (<i>ye.</i>) prickly.	Spain. 1570. 6. 9. H. ⊖.		Bot. mag. t. 1836.
7 aquanticum. w. (<i>ye.</i>) sweet-scented.	S. Europe. 1731. 7. 8. H. ⊖.		Barrel. icon. 551.
8 maritimum. w. (<i>ye.</i>) sea.	Sicily. 1640. 7. 9. H. ȝ.		Breyn. cent. t. 77.
9 salicifolium. w. (<i>ye.</i>) Willow-leaved.	Austria. 1759. 6. 10. H. ȝ.		Bocc. mus. t. 129.
10 grandiflorum. w. (<i>ye.</i>) great-flowered.	— — — — H. ȝ.		Jacq. aust. 4. t. 370.
11 speciosissimum. s.s. (<i>ye.</i>) handsome.	Italy. 1828. — — H. ȝ.		Moris. s. 6. t. 7. f. 52.
12 cordifolium. w. (<i>ye.</i>) heart-leaved.	Hungary. 1739. 6. 8. H. ȝ.		Arduin. spec. 1. t. 12.
13 procumbens. Df. (<i>ye.</i>) procumbent.	— — — — H. ȝ.		W. et K. hung. 2. t. 113.
ZEX'MENIA. D.D.	ZEXM'ENIA. Syngenesia Polygamia Necessaria.		
tagetiflora. d.d. (<i>ye.</i>) Tagetes-flowered.	Mexico. 1828. 8. 10. F. ḥ.		
WED'ELIA. K.S.	WED'ELIA. Syngenesia Polygamia Necessaria. W.		
1 acapulcensis. k.s. (<i>ye.</i>) Acapulco.	Mexico. 1823. 6. 8. S. ḥ.		
2 hispida. k.s. (<i>ye.</i>) hispid.	New Spain. 1819. — — S. ḥ.		Bot. reg. t. 543.
3 aurea. d.d. (<i>ye.</i>) golden-flower'd.	Mexico. 1828. 8. 10. F. ḥ.		
4 radiosa. b.r. (<i>ye.</i>) rayed.	S. America. 1822. — — S. ḥ.		Bot. reg. t. 610.
GYMNOL'OMA. K.S.	GYMNOL'OMA. Syngenesia Polygamia Necessaria.		
1 triplinerve. k.s. (<i>ye.</i>) triple-nerved.	S. America. 1825. 6. 9. S. ḥ.		
2 maculatum. b.r. (<i>ye.</i>) spotted-stalked.	— — — — S. ḥ.		Bot. reg. t. 662.
ACTINOM'ERIS. N.	ACTINOM'ERIS. Syngenesia Polygamia Frustranea.		
1 squarrosa. n. (<i>ye.</i>) squarrose.	N. America. 1640. 9. 11. H. ȝ.		Jacq. vind. 2. t. 110.
<i>Coreopsis alternifolia.</i> Jacq. <i>Verbesina Coreopsis.</i> Ph.			
2 præcera. n. (<i>ye.</i>) tall.	N. America. 1765. 9. 10. H. ȝ.		
<i>Coreopsis præcera.</i> H.K.			
3 alba. n. (<i>wh.</i>) white-flowered.	— — — — H. ȝ.		
4 alata. n. (<i>ye.</i>) wing-stalked.	Louisiana. 1825. — — H. ȝ.		
<i>Coreopsis alata.</i> W.	Mexico. 1803. 9. 11. H. ȝ.		Cavan. ic. 3. t. 260.
5 ovata. n. (<i>ye.</i>) oval-leaved.	— — — — H. ȝ.		Cavan. ic. 3. t. 280.
HELIANTHUS. W.	SUNFLOWER. Syngenesia Polygamia Frustranea. L.		
1 annuus. w. (<i>ye.</i>) annual.	S. America. 1596. 6. 10. H. ⊖.		
2 indicus. w. (<i>ye.</i>) dwarf annual.	Egypt. 1785. — — H. ⊖.		Tabern. ic. 764.
3 lenticularis. b.r. (<i>ye.</i>) freckled.	N. America. 1827. 8. 9. H. ⊖.		Bot. reg. t. 1265.
4 petiolaris. n. (<i>ye.</i>) long-petioled.	Arkansa. 1826. 7. 11. H. ⊖.		Swt. br. fl. gar. ser. 2.
5 asper. Roth.			
6 dentatus. w. (<i>ye.</i>) tooth-leaved.	Mexico. 1798. 9. 11. F. ȝ.		Cavan. ic. 3. t. 220.
7 multiflorus. w. (<i>ye.</i>) many-flowered.	N. America. 1597. 8. 10. H. ȝ.		Bot. mag. t. 227.
β plenus.	(<i>ye.</i>) double-flowered.	— — — — H. ȝ.	
8 pubescens. b.r. (<i>ye.</i>) pubescent.	Jerusalem Artichoke. Brazil. 1617. 9. 10. H. ȝ.		Jacq. vind. 2. t. 161.
9 atrorubens. w. (<i>ye.</i>) dark eyed.	N. America. 1795. 7. 10. H. ȝ.		Bot. reg. t. 524.
10 diffusus. b.m. (<i>ye.</i>) spreading.	— — — — H. ȝ.		Dill. elth. t. 94. f. 110.
11 laetiflorus. p.s. (<i>ye.</i>) pale-flowered.	— — — — H. ȝ.		Bot. mag. t. 2020.
12 divaricatus. w. (<i>ye.</i>) divaricate.	— — — — H. ȝ.		
13 trachelifolius. w. (<i>ye.</i>) Trachelium-leav'd.	— — — — H. ȝ.		
14 longifolius. Ph. (<i>ye.</i>) long-leaved.	Georgia. 1812. — — H. ȝ.		
15 pauciflorus. n. (<i>ye.</i>) few-flowered.	Louisiana. 1823. — — H. ȝ.		
16 giganteus. w. (<i>ye.</i>) gigantic.	N. America. 1714. 9. 10. H. ȝ.		
17 altissimus. w. (<i>ye.</i>) tallest.	— — — — H. ȝ.		Jacq. vind. 2. t. 160.
18 strumosus. w. (<i>ye.</i>) Carrot-rooted.	— — — — H. ȝ.		Bocc. sic. t. 27. f. 4.
19 prostratus. w. (<i>ye.</i>) rough.	— — — — H. ȝ.		
20 decapetalus. w. (<i>ye.</i>) ten-rayed.	— — — — H. ȝ.		Rob. ic. 235.

COMPOSITÆ.

21	móllis. w.	(ye.) soft.	N. America.	1805.	7. 10. H. 4.	
22	macrophyllus. w.	(ye.) large-leaved.	_____	1800.	8. 10. H. 4.	Willd. hort. ber. t. 70.
23	angustifolius. w.	(ye.) narrow-leaved.	_____	1789.	9. 10. H. 4.	Bot. mag. t. 2051.
24	excelsus. w.	(ye.) lofty. giganteus. Cav. non w.	Mexico.	1824.	— H. 4.	Cav. ic. 3. t. 219.
25	linearis. w.	(ye.) linear-leaved	_____	1818.	8. 10. H. 4.	Bot. reg. t. 523.
26	parviflorus. k.s.	(ye.) small-flowered.	_____	1824.	— H. 4.	H. et B. 4. t. 378.
27	cornifolius. k.s.	(ye.) Dogwood-leaved.	_____	—	— H. 4.	
28	trilobatus. L.en.	(ye.) three-lobed.	_____	1823.	— H. 4.	
VIGITERA. K.S.						
		VIGI'ERA. Syngenesia Polygamia Frustranea.				
	helianthoides. k.s.	(ye.) Sunflower-like. Cuba.	1825.	6. 9. S. 4.		H. et B. 4. t. 379.
GAILLARDIA. W.						
		GAILLA'R'DIA. Syngenesia Polygamia Frustranea. W.				
1	bicolor. w.	(ye. re.) two-coloured.	Carolina.	1787.	7. 10. H. 4.	Bot. mag. t. 1602.
2	aristata. Ph.	(ye.) long-awned.	N. America.	1812.	— H. 4.	Bot. reg. t. 1186.
RUDBECKIA. D.D.						
		RUDBE'C'KIA. Syngenesia Polygamia Frustranea. L.				
1	pinnata. Ph.	(ye.) fragrant.	N. America.	1803.	8. 9. H. 4.	Swt. br. fl. gar. t. 146.
2	digitata. Ph.	(ye.) narrow-jagged-leav'd.	1759.	— H. 4.	Moris. s. 6. t. 6. f. 54.	
3	laciñata. Ph.	(ye.) broad jagged-leav'd.	1640.	7. 9. H. 4.	— s. 6. t. 6. f. 53.	
4	columnaris. Ph.	(ye.) high-crowned.	1811.	8. 9. H. 4.	Bot. mag. t. 1601.	
5	lævigata. Ph.	(ye.) smooth.	Georgia.	1812.	7. 8. H. 4.	Jacq. icon. 3. t. 592.
6	amplexifolia. Ph.	(ye.) stem-clasping.	Louisiana.	1793.	7. 9. H. ⊙.	
CENTROCARPHA. D.D.						
		CENTROCA'R'PH'A. Syngenesia Polygamia Frustranea.				
1	subtomentosa. Ph.	(ye.) downy-lobed.	N. America.	1802.	8. 9. H. 4.	
2	triloba. d.d.	(ye.) three-lobed.	1699.	7. 10. H. 4.	Bot. reg. t. 525.	
		Rudbeckia triloba. B.R.				
3	hirta. d.d.	(ye.) great hairy.	1714.	6. 11. H. 4.	Swt. br. fl. gar. t. 82.	
		Rudbeckia hirta. B.F.G.				
4	fulgida. d.d.	(ye.) small hairy.	1760.	7. 8. H. 4.	Bot. mag. t. 1996.	
5	acutifolia. B.F.G.	(ye.) pointed-leaved.	1822.	7. 11. H. 4.	Swt. br. fl. gar. s. 2.	
6	spathulata. w.	(ye.) spatulate-leav'd.	1825.	8. 11. H. 4.		
7	gracilis. n.	(ye.) slender hairy.	—	— H. 4.		
ECHINACEA. D.D.						
		ECHIN'ACEA. Syngenesia Polygamia Frustranea.				
1	purpurea. d.d.	(pu.) reflexed-rayed.	N. America.	1699.	7. 10. H. 4.	Bot. mag. t. 2.
		Rudbeckia purpurea. B.M.				
2	serotina. d.d.	(li.) hispid-stalked.	1816.	8. 10. H. 4.	Swt. br. fl. gar. t. 4.	
		Rudbeckia serotina. B.F.G.				
3	natifolia.	(ro.) Rape-leaved.	New Spain.	1824.	7. 10. F. 4.	
		Rudbeckia natifolia. K.S.				
4	heterophylla. d.d.	(li.) various-leaved.	Mexico.	1828.	9. 11. H. 4.	Swt. br. fl. gar. s. 2. t. 32.
CO'SMOS. C.I.						
		Co'smos. Syngenesia Polygamia Frustranea. W.				
1	sulphureus.	(st.) Southernwood-leav'd.	Mexico.	1799.	7. 8. H. ⊙.	Jacq. ie. 3. t. 595.
2	tenellus. k.s.	(ro.) slender.	Mexico.	1825.	10. 11. G. ⊙.	Bot. mag. t. 1535.
3	bipinnatus. c.i.	(pu.) purple-flower'd.	—	1799.	— G. ⊙.	Jacq. schoen. 3. t. 374.
4	crithmifolius. k.s.	(ro.) Samphire-leav'd.	—	1825.	— G. ⊙.	
5	parviflorus. w.	(wh.) white-flowered.	—	1800.	— H. ⊙.	
GEORGINA. W.						
		GEORG'INA. Syngenesia Polygamia Superfluo—Frustranea.				
1	Cervantessii. LG.	(sc.) Cervantes'.	Mexico.	1820.	8. 10. H. 4.	Swt. br. fl. gar. s. 2. t. 22.
	β Aurðra. s.F.G.	(sc.) double-flowered.	—	1828.	— H. 4.	Swt. flor. guid. t. 117.
2	crocata. LG.	(sc.) saffron-coloured.	—	1818.	— H. 4.	Swt. br. fl. gar. t. 282.
3	coccinea. w.	(sc.) scarlet-flowered.	—	1802.	— H. 4.	Cavan. icon. t. 266.
4	ròsea. c.i.	(ro.) bipinnate-leaved.	—	—	7. 10. H. 4.	t. 265.
5	astrantiæflòra.	(pu.) Astrantia-flower'd.	—	1812.	— H. 4.	
6	variabilis. w.	(va.) variable.	—	1789.	6. 11. H. 4.	Cavan. icon. 1. t. 80.
	Dahlia pinnata. c.i.	supérflua. H.K.				
a	purpurea.	(pu.) purple-flowered.	—	—	— H. 4.	Ann. mus. 3. t. 37. f. 1.
β	lilacina.	(li.) lilac-coloured.	—	—	— H. 4.	Sal. par. lond. t. 16.
γ	pallida.	(bh.) blush-flowered.	—	—	— H. 4.	
δ	alba.	(wh.) white-flowered.	—	—	— H. 4.	
ε	flava.	(ye.) yellow-flowered.	—	—	— H. 4.	
ζ	cuprea.	(co.) copper-coloured.	—	—	— H. 4.	
η	rùbra.	(re.) red-flowered.	—	—	— H. 4.	Bot. mag. t. 1885. A.
ς	coccinea.	(sc.) scarlet-flowered.	—	—	— H. 4.	
ι	punicea.	(cr.) crimson-flowered.	—	—	— H. 4.	Bot. reg. t. 55.
κ	atrosanguinea.	(d.cr.) dark crimson.	—	—	— H. 4.	
λ	atropurpurea.	(d.pu.) dark purple.	—	—	— H. 4.	
μ	nigra.	(da.) dark-coloured.	—	—	— H. 4.	
ν	variegata.	(v.) variegated-flower'd.	—	—	— H. 4.	
ξ	nana.	(li.) double dwarf lilac.	—	—	— H. 4.	
ο	supérba.	(pu.) double purple.	—	—	— H. 4.	Andr. deposit. t. 483.
			—	— H. 4.	Bot. mag. t. 1885. B.

π <i>Junoniæna</i> . (d.pu.) <i>Juno</i> .	Mexico.	6. 11. H. ¼.	Swt. flor. guid. t. 102.
ρ <i>imperiðsa</i> . (d.pu.) <i>Dennis's imperial</i> .	1827.	—	H. ¼.	v. 1. t. 71.
σ <i>stellariðs</i> . (sc.) <i>morning star</i> .	1826.	—	H. ¼.	v. 1. t. 65.
τ <i>sphaerocphala</i> . (d.cr.) <i>globe-flowered</i> .	1827.	—	H. ¼.	v. 2. t. 115.
ν <i>Belladonna</i> . (pi.) <i>painted Lady</i> .	—	—	H. ¼.	v. 2. t. 110.

The varieties of the present species are almost endless.

CALLIO'PSIS. S.S. *CALLIO'PSIS*. Syngenesia Polygamia Frustranea.

1 <i>tinctoria</i> . N. (ye.pu.) two-coloured.	Arkansa.	1823.	6. 10. H. ⊙.	Swt. br. fl. gar. t. 72.
bicolor. s.s. <i>Coreópsis tinctoria</i> . N.	—	—	—	—

2 <i>palmata</i> . s.s. (ye.) palmate-leaved.	Louisiana.	—	8. 10. H. ⊙.	Swt. br. fl. gar. t. 72.
---	------------	---	--------------	--------------------------

3 <i>rosea</i> . s.s. (ro.) rose-coloured.	N. America.	—	—	H. ¼.
--	-------------	---	---	-------

Coreópsis rosea. N.

COREO'PSIS. W. *COREO'PSIS*. Syngenesia Polygamia Frustranea. L.

1 <i>grandiflora</i> . n. (ye.) large-flowered.	N. America.	1826.	8. 9. H. ¼.	Swt. br. fl. gar. t. 175.
---	-------------	-------	-------------	---------------------------

2 <i>verticillata</i> . w. (ye.) whorl-leaved.	—	1759.	7. 10. H. ¼.	Bot. mag. t. 156.
--	---	-------	--------------	-------------------

3 <i>tenuifolia</i> . w. (ye.) slender-leaved.	—	1780.	7. 8. H. ¼.	Pluk. man. t. 344. f. 4.
--	---	-------	-------------	--------------------------

4 <i>trichosperma</i> . M. (ye.) pinnate leaved.	Carolina.	1822.	8. 10. G. ♂.	Plum. ic. 53. f. 1.
--	-----------	-------	--------------	---------------------

5 <i>chrysanthra</i> . w. (ye.) Angelica-leav'd. W. Indies.	—	1752.	7. 9. S. ♂.	Bot. reg. t. 1228.
---	---	-------	-------------	--------------------

6 <i>aurea</i> . w. (ye.) Hemp-leaved.	N. America.	1785.	8. 9. H. ¼.	Moris. s. 6. t. 3. f. 44.
--	-------------	-------	-------------	---------------------------

7 <i>tripteris</i> . Ph. (ye.) three-leaved.	—	1737.	8. 10. H. ¼.	—
--	---	-------	--------------	---

8 <i>discolor</i> . L.en. (ye.) two-coloured.	—	1818.	—	H. ¼.
---	---	-------	---	-------

9 <i>senifolia</i> . Ph. (ye.) sessile-leaved.	—	1812.	—	H. ¼.
--	---	-------	---	-------

10 <i>amplexicaulis</i> . K.S. (ye.) stem-clasping.	—	1806.	7. 8. H. ¼.	—
---	---	-------	-------------	---

Simsia amplexicaulis. P.S.

11 <i>álba</i> . w. (wh.) climbing.	Jamaica.	1699.	6. 7. S. ¼. ⊖.	Herm. parad. 124.
-------------------------------------	----------	-------	----------------	-------------------

12 <i>incisa</i> . B.R. (ye.) jagged-leaved.	W. Indies.	9. 12. S. ¼. ⊖.	Bot. reg. t. 7.
--	------------	------	-----------------	-----------------

13 <i>réptans</i> . w. (ye.) trailing.	—	1792.	7. 9. S. ⊖.	Smith spic. t. 22.
--	---	-------	-------------	--------------------

14 <i>auriculata</i> . w. (ye.) ear-leaved.	N. America.	1699.	8. 10. H. ¼.	Schk. hand. 3. t. 260.
---	-------------	-------	--------------	------------------------

15 <i>latifolia</i> . w. (ye.) broad-leaved.	—	1786.	8. 9. H. ¼.	—
--	---	-------	-------------	---

16 <i>arguta</i> . Ph. (ye.) sharp-notched.	Carolina.	—	H. ¼.
---	-----------	------	---	-------

17 <i>lanceolata</i> . w. (ye.) spear-leaved.	—	1724.	7. 9. H. ¼.	Swt. br. fl. gar. t. 10.
---	---	-------	-------------	--------------------------

18 <i>crassifolia</i> . w. (ye.) thick-leaved.	—	1786.	8. 10. H. ¼.	—
--	---	-------	--------------	---

19 <i>angustifolia</i> . w. (ye.) narrow-leaved.	N. America.	1778.	6. 8. H. ¼.	—
--	-------------	-------	-------------	---

B'IDENS. K.S. *B'IDENS*. Syngenesia Polygamia Frustranea.

1 <i>nodiflora</i> . w. (ye.) sessile-flowered.	E. Indies.	1732.	7. 8. S. ⊖.	Dill. elth. t. 44. f. 52.
---	------------	-------	-------------	---------------------------

2 <i>tripartita</i> . w. (ye.) trifid.	Britain.	7. 9. H. ⊖.	Eng. bot. t. 1113.
--	----------	------	-------------	--------------------

3 <i>cernua</i> . w. (ye.) nodding.	—	—	— t. 1114.
-------------------------------------	---	------	---	------------

4 <i>chrysanthemoides</i> . w. (ye.) large yellow.	N. America.	1827.	8. 10. H. ⊖.	Orteg. dec. 8. t. 12.
--	-------------	-------	--------------	-----------------------

5 <i>arguta</i> . K.S. (ye.) sharp-toothed.	Mexico.	1825.	—	—
---	---------	-------	---	---

6 <i>triplinervia</i> . K.S. (ye.) triply-nerved.	—	—	—	—
---	---	---	---	---

7 <i>heterophylla</i> . w. (ye.) various-leaved.	—	1803.	8. 9. F. ¼.	Moris. his. s. 6. t. 5. f. 21.
--	---	-------	-------------	--------------------------------

8 <i>luxurians</i> . W.en. (ye.) luxuriant.	—	1823.	—	—
---	---	-------	---	---

9 <i>frondosa</i> . w. (ye.) smooth-stalked.	N. America.	1710.	7. 8. H. ⊖.	—
--	-------------	-------	-------------	---

10 <i>foliosa</i> . W.en. (ye.) leafy.	—	1823.	—	H. ⊖.
--	---	-------	---	-------

11 <i>connata</i> . w. (ye.) connate.	N. America.	—	—	H. ⊖.
---------------------------------------	-------------	---	---	-------

12 <i>leucantha</i> . w. (wh.) white-flowered.	S. America.	—	—	H. ⊖.
--	-------------	---	---	-------

13 <i>striata</i> . B.F.G. (std.) stripe-flowered.	Mexico.	1827.	8. 11. H. ⊖.	—
--	---------	-------	--------------	---

14 <i>chinensis</i> . w. (wh.) Chinese.	China.	1801.	6. 7. H. ⊖.	—
---	--------	-------	-------------	---

15 <i>diversifolia</i> . w. (ye.) different-leav'd.	S. America.	1811.	9. 10. H. ⊖.	Bot. mag. t. 1689.
---	-------------	-------	--------------	--------------------

16 <i>Cósmea lutea</i> . B.M. <i>Coreópsis diversifolia</i> . Jacq.	—	—	—	Dill. elth. t. 43. f. 51.
---	---	---	---	---------------------------

17 <i>ciliata</i> . (ye.) fringed.	Mexico.	1827.	8. 10. H. ⊖.	Cavan. ic. 3. t. 229.
------------------------------------	---------	-------	--------------	-----------------------

18 <i>chrysanthra</i> . Ort. (ye.) golden-flowered.	—	—	—	— t. 13.
---	---	---	---	----------

19 <i>coronata</i> . F. (ye.) crowned.	—	1829.	—	Herm. parad. t. 123.
--	---	-------	---	----------------------

20 <i>parviflora</i> . s.s. (ye.) small-flowered.	Siberia.	1823.	—	Arduin. spec. 2. t. 18.
---	----------	-------	---	-------------------------

21 <i>bullata</i> . w. (ye.) rough-leaved.	Italy.	1759.	—	—
--	--------	-------	---	---

22 <i>reflexa</i> . L.en. (ye.) reflexed.	Mexico.	1824.	—	—
---	---------	-------	---	---

23 <i>crithmifolia</i> . K.S. (ye.) Samphire-leav'd.	Quito.	—	9. 10. F. ¼.	—
--	--------	---	--------------	---

24 <i>angustifolia</i> . K.S. (or.) narrow-leaved.	New Spain.	—	—	—
--	------------	---	---	---

25 <i>procera</i> . B.R. (ye.) tall.	S. America.	1818.	—	Bot. reg. t. 684.
--------------------------------------	-------------	-------	---	-------------------

26 <i>fernæfolia</i> . (ye.) Fennel-leaved.	Mexico.	1799.	10. 11. F. ¼.	Jacq. scheenb. 3. t. 373.
---	---------	-------	---------------	---------------------------

2

3 camphorina. w. (wh.)	Camphor-scent'd.	C. B. S.	1794.	4. 7. G. h.	Seb. mus. 1. t. 90. f. 2.
4 dentata. w.	(wh.) tooth-leaved.	—	1820.	— G. h.	
SCLEROCA'RPUS. W. SCLEROCA'RPUS.	Syngenesia Polygamia Frustranea.	W.			
africânum. w. (ye.) African.	Guinea.	1812.	7. 8. G. ♂.	Jacq. ic. 1. t. 176.	
CULL'UMIA. H.K.	CULL'UMIA. Syngenesia Polygamia Frustranea. H.K.				
1 ciliaris. H.K. (ye.) ciliated.	C. B. S.	1774.	5. 6. G. h.	Burm. afr. t. 54. f. 1.	
2 setosa. H.K. (ye.) recurved smooth-leav'd.—	—	1780.	6. 8. G. h.	Comm. hort. 2. t. 28.	
3 squarrôsa. H.K. (ye.) recurved-awl-leaved.—	—	1786.	— G. h.	Th. act. hafn. 3. t. 5.	
BERCK'H'EYA. H.K. BERCK'H'EYA. Syngenesia Polygamia Frustranea. H.K.					
1 cynaroïdes. w. (ye.) Artichoke-cup'd.	C. B. S.	1789.	6. G. ♀.		
2 obovata. w. (ye.) smooth shrubby.	—	1794.	6. 8. G. h.	Hout. n. h. 6. t. 34. f. 2.	
3 incâna. w. (ye.) hoary.	—	1739.	7. 8. G. h.	Jacq. ic. 3. t. 591.	
4 cuneata. w. (ye.) wedge-leaved.	—	1812.	6. 8. G. h.	Th. act. hafn. 3. t. 10.	
5 palmata. w. (ye.) palmated.	—	1800.	— G. h.	3. t. 13.	
6 grandiflora. w. (ye.) large-flowered.	—	1812.	— G. h.	Bot. mag. t. 1844.	
7 uniflora. w. (ye.) single-flowered.	—	1815.	— G. h.	Th. act. hafn. 3. t. 7.	
8 cérnua. H.K. (ye.) drooping-flower'd.	—	1774.	5. 7. G. ♂.	Meerb. ic. 1. t. 40.	
DIDE'LTA. H.K. DIDE'LTA. Syngenesia Polygamia Frustranea. H.K.					
1 carnosum. H.K. (ye.) alternate-leav'd.	C. B. S.	1774.	6. 7. G. h.	L'Her. stirp. t. 28.	
2 spinosum. H.K. (ye.) opposite-leaved.	—	—	— G. h.	Wendl. obs. t. 4. f. 32.	
GORT'ERIA. H.K. GORT'ERIA. Syngenesia Polygamia Frustranea. H.K.					
personata. H.K. (ye.) procumbent.	C. B. S.	1774.	7. 8. G. ♂.	Jacq. coll. 4. t. 21. f. 1.	
GAZ'ANIA. H.K. GAZ'ANIA. Syngenesia Polygamia Frustranea. H.K.					
1 uniflora. B.M. (ye.) golden.	C. B. S.	1816.	7. 8. G. h.	Bot. mag. t. 2270.	
2 rigens. H.K. (ye.) great-flowered.	—	1755.	5. 9. G. h.	— t. 90.	
3 heterophylla.w.(ye.)various-leaved.	—	1812.	— G. ♀.	Willd. hort. ber. t. 97.	
4 Pavonia. H.K. (ye.) Peacock.	—	1804.	6. 7. G. ♀.	Bot. reg. t. 35.	
5 subulata. H.K. (ye.) awl-leaved.	—	1792.	7. 8. G. ♀.		
CRYPTOSTE'MMA. H.K. CRYPTOSTE'MMA. Syngenesia Polygamia Frustranea.					
1 calendulaceum. H.K. (ye.) Marygold-flow'r'd.C.B.S.	1752.	6. 8. H. ♂.	Bot. mag. t. 2252.		
2 hypochondriacum. H.K. (ye.) divided-rayed.	—	1731.	7. 8. H. ♂.		
3 runcinatum. H.K. (ye.) Dandelion-leaved.	—	1794.	— H. ♂.		
ARCTOTH'ECA. W. ARCTOTH'ECA. Syngenesia Polygamia Frustranea. H.K.					
1 rēpens. w. (ye.) creeping.	C. B. S.	1793.	7. 8. G. ♀.	Jacq. schoen. 3. t. 306.	
2 hirta. L.en. (ye.) hairy.	—	1818.	7. 9. G. ♀.		
SPHENO'GYNE. H.K. SPHENO'GYNE. Syngenesia Polygamia Frustranea. H.K.					
1 anthemoides. H.K. (ye.) white-crowned.	C. B. S.	1774.	7. 9. H. ♂.	Bot. mag. t. 544.	
2 crithmifolia.H.K.(ye.) Samphire-leav'd.	—	1768.	4. 8. G. h.	Burm. afr. t. 65. f. 1.	
3 scariosa. H.K. (ye.) scaly-cupped.	—	1774.	— G. h.		
4 abrotanifolia.H.K.(ye.)Southernwood-leav'd.	—	1789.	5. 8. G. h.		
5 dentata. H.K. (ye.) small-leaved.	—	1787.	6. 7. G. h.	Burm. afr. t. 64.	
6 pilifera. B.R. (ye.) hair toothed.	—	1820.	— G. h.	Bot. reg. t. 604.	
7 serrata. s.s. (ye.) saw-leaved.	—	1826.	— G. h.		
8 foeniculacea. (ye.) Fennel-leaved.	—	1825.	5. 9. H. ♂.	Jacq. schoen. 2. t. 156.	
Arctotis foeniculacea. Jacq.					
9 leucanthemoïdes. (ye.) Ox-eye Daisy-leav'd.	—	—	— H. ♂.		2. t. 164.
Arctotis leucanthemoïdes. Jacq.					
10 odorata. H.K. (ye.) smooth-seeded.	—	1774.	4. 6. G. h.		
TITH'ONIA. W. TITH'ONIA. Syngenesia Polygamia Frustranea. W.					
1 tagetiflora. w. (or.) Tagetes-flower'd.W. Indies.	—	1821.	8. 9. S. ♀.	Bot. reg. t. 591.	
2 tubæformis. d.p. (ye.) tube-flowered. Mexico.	—	1799.	9. 11. H. ♂.	Jacq. schoen. 3. t. 375.	
Helianthus tubæformis. Jacq.					
ARCT'OTIS. H.K. ARCT'OTIS. Syngenesia Polygamia Necessaria. L.					
1 acaulis. w. (ye.) dwarf.	C. B. S.	1759.	4. 7. G. ♀.	Bot. reg. t. 122.	
2 tricolor. w. (pu.wh.) three-coloured.	—	1794.	5. 7. G. ♀.	— t. 131.	
3 undulata. w. (or.) wave-leaved.	—	1795.	4. 6. G. ♀.	Jacq. schoen. 2. t. 160.	
4 grandiflora.w.(or.cr.)great-flowered.	—	1774.	3. 5. G. h.	— t. 378.	
5 Massoniâna. s.s.(st.re.)Masson's.	—	—	— G. ♂.		
grandiflora. H.K. non Jacq.					
6 speciosa. w. (pu.ye.) showy.	—	1812.	6. 8. G. ♀.	Bot. mag. t. 2182.	
7 glaucophylla.w.(or.pi.)Sea-green-leav'd.	—	1794.	5. 8. G. ♀.	Jacq. schoen. 2. t. 170.	
8 plantaginea.w.(ye.pu.)Plantain-leaved.	—	1768.	6. 8. G. ♀.		
9 argéntea. w. (v.) silver-leaved.	—	1774.	8. G. ♂.		
10 rosea. w. (ro.) Rose.	—	1793.	7. 9. G. ♀.	Jacq. schoen. 2. t. 162.	
11 decumbens.w.(wh.pi.)decumbent.	—	1790.	— G. ♀.	— t. 381.	
12 angustifolia. w. (or.) narrow-leaved.	—	1739.	— G. ♀.		
13 flaccida. w. (wh.re.) bending-stalked.	—	1794.	5. 7. G. ♂.	Jacq. schoen. 2. t. 163.	
14 virgata. w. (ye.pu.) slender-twigg'd.	—	1816.	7. 9. H. ♂.	— t. 307.	
15 flammæa. J.F. (or.) flame-coloured.	—	1822.	— G. ♀.	Jacq. frag. t. 47. f. 1.	

16 amœna. J.F. (wh.pu.) handsome.	C. B. S.	1822.	7. 9. G. ȝ.	Jacq. frag. t. 118.
17 paniculâta.w. (wh.cr.) paniced.	_____	_____	G. h.	Jacq. schœn. 3. t. 380.
18 décurrens.w. (wh.pi.) decurrent.	_____	_____	6. 7. G. h.	_____ 2. t. 165.
19 melanocicla.w. (wh.re.) various-colour'd.	_____	1812.	G. h.	_____
20 réptans. w. (st.br.) creeping.	_____	1795.	7. 9. G. ȝ.	Jacq. schœn. 3. t. 382.
21 auriculâta.w. (ye.pu.) ear-leaved.	_____	_____	6. 8. G. h.	_____ 2. t. 169.
22 fastuosa. w. (or.cr.) Orange-flower'd.	_____	_____	5. 7. G. ♂.	_____ 2. t. 166.
23 spinulosa. w. (or.bk.) thorny-leaved.	_____	_____	5. 8. G. ♂.	_____ 2. t. 167.
24 maculâta. w. (st.pu.) spotted.	_____	1812.	G. h.	Bot. reg. t. 130.
25 áspera. B.R. (wh.ro.) broad rough-leav'd.	_____	1710.	7. 9. G. h.	_____ t. 34.
26 elátior. w. (ye.pu.) tall.	_____	1816.	G. h.	Jacq. schœn. 2. t. 172.
27 arboréscens.w. (wh.pi.) arborescent.	_____	_____	G. h.	_____ 2. t. 171.
28 aurélula. B.R. (or.) narrow rough-leav'd.	_____	1710.	G. h.	Bot. reg. t. 32.
29 revoluta. w. (ye.bk.) revolute-leaved.	_____	1824.	8. 10. H. ♂.	Jacq. schœn. 2. t. 173.
30 cùprea. w. (co.pu.) copper-coloured.	_____	_____	7. 8. G. h.	_____ 2. t. 176.
31 squarrosa. w. (or pu.) squarrose.	_____	1825.	G. h.	_____ 2. t. 177.
32 Cineraria. w. (ye.or.) black-eyed.	_____	_____	G. ȝ.	_____ 2. t. 174.
33 bicolor. W.en. (wh.re.) two-coloured.	_____	1812.	G. ȝ.	_____
34 glabrata. w. (ye.pu.) smooth-leaved.	_____	1816.	7. 9. G. h.	Jacq. schœn. 2. t. 175.
35 purpurea. W.en.(pu.) purple.	_____	1827.	G. h.	_____

CALENDULA. W. MARYGOLD. Syngenesia Polygamia Necessaria. L.

1 arvensis. w. (ye.) field.	Europe.	1597.	6. 9. H. ♂.	Ger. herb. 603. f. 10.
2 sícula. W.en. (ye.) Sicilian.	Sicily.	1816.	H. ♂.	_____
3stellata. w. (ye.) starry.	Barbary.	1795.	H. ♂.	Schk. hand 3. t. 265.
4 officinalis. w. (ye.) common.	S. Europe.	1573.	H. ♂.	_____
β plena. (ye.) double-flowered.	_____	_____	H. ♂.	_____
5 ægyptiaca. P.S. (ye.) Egyptian.	Egypt.	1820.	H. ♂.	_____
6 suffruticosa. w. (ye.) suffrutescent.	S. Europe.	1823.	G. h.	_____
7 denticulata. s.s. (ye.) toothed.	Barbary.	1821.	6. 8. F. h.	_____
8 sancta. w. (st.) pale-flowered.	Levant.	1731.	5. 9. H. ♂.	_____
9 incana. w. (ye.) hoary.	Barbary.	1796.	6. 8. H. ♂.	Desf. atl. 2. t. 245.
tomentosa. Df. non L.	_____	_____	_____	_____

10 pluvialis. w. (wh.pu.) small Cape.

C. B. S.	1699.	H. ♂.	Mill. ic. t. 75. f. 2.
11 hybrida. w. (wh.pu.) great Cape.	1752.	6. 7. H. ♂.	Swt. br. fl. gar. t. 39.
12 nudicauis. w. (wh.) naked-stalked.	1731.	6. 8. H. ♂.	Comm. hort. 2. t. 33.
13 graminifolia.w.(wh.pu.) Grass-leaved.	_____	5. 9. G. ȝ.	Bot. reg. t. 289.
14 Trágus. w. (pu.wh.) bending-stalked.	1774.	5. 6. G. h.	Bot. mag. t. 408.
β fláccida. v. (or.) flaccid.	_____	6. 8. G. h.	Bot. reg. t. 28.
15 viscósa. H.K. (or.) viscous.	1790.	6. 9. G. h.	Andr. deposit. t. 412.

Arctotis glutinosa. B.M. 1343.

16 dentata. A.R. (wh.) tooth-leaved.	1790.	5. 6. G. h.	Andr. deposit. c. ic.
17 oppositifolia. w. (ye.) glaucous-leaved.	1774.	8. G. h.	_____
18 fruticosa. w. (wh.) shrubby.	1752.	6. 7. G. h.	Mill. ic. 2. t. 283.
19 chrysanthemifolia. v. (ye.) large-flower'd.	1790.	3. 8. G. h.	Bot. reg. t. 40.
20 arboréscens. w. (ye.) rough-leaved.	1774.	12. G. h.	Jacq. ic. 3. t. 596.
21 muricata. w. (ye.) muricate.	1816.	G. h.	_____

POLY'MNIA. W. POLY'MNIA. Syngenesia Polygamia Necessaria. L.

1 canadensis. w. (ye.) Canadian.	N. America.	1768.	7. 8. H. ȝ.	Lin. am. ac. 3. t. 1. f. 5.
2 Uvedalia. w. (ye.) broad-leaved.	_____	1699.	8. 10. H. ȝ.	Moris. s. 6. t. 7. f. 55.
3 maculata. c.I. (ye.) spotted-stalked.	Mexico.	1823.	9. 11. H. ȝ.	Cav. ic. 3. t. 227.
4 abyssinica. w. (ye.) upright.	Africa.	1775.	4. 5. S. ♂.	_____

MELAMP'ODIUM. W. MELAMP'ODIUM. Syngenesia Polygamia Necessaria. W.

1 americanum. w. (ye.) American.	Vera Cruz.	1733.	8. 10. S. ♂.	Reliq. Houst. 9. t. 21.
2 humile. w. (ye.) dwarf.	Jamaica.	1782.	6. 10. S. ♂.	_____
3 hispidum. K.S. (ye.) hispid.	New Spain.	1825.	9. 11. H. ♂.	H. et B. 4. t. 399.
4 perfoliatum. K.S. (ye.) perfoliate.	_____	1796.	8. 10. H. ♂.	Cavan. ic. 1. t. 15.
<i>Alcina perfoliata</i> . Cav. <i>Wedelia perfoliata</i> . w.	_____	_____	_____	_____

5 longifolium. W.en.(ye.)long-leaved.

6 ovatifolium. R.I. (ye.) oval-leaved.

Wedelia ovatifolium. w. *Alcina ovalifolia*. LG.

7 sericeum. s.s. (ye.) silky-leaved.

New Spain. 1825. — H. ♂.

MILL'ERIA. L. MILL'ERIA. Syngenesia Polygamia Necessaria. L.

1 quinqueflora. w. (ye.) five flowered.	Vera Cruz.	1731.	7. 10. H. ♂.	Cavan. ic. 1. t. 82.
2 biflora. w. (ye.) two-flowered.	S. America.	1730.	H. ♂.	Mart. dec. 47. f. 1.

FLAV'ERIA. J. FLAV'ERIA. Syngenesia Polygamia Necessaria. P.S.

1 repanda. LG. (ye.) repand.	S. America.	1799.	7. 9. H. ♂.	Cavan. ic. 1. t. 4.
2 Contrayérba. K.S. (ye.) Peruvian.	Peru.	1794.	8. 10. S. ♂.	_____
3 angustifolia. K.S. (ye.) narrow-leaved.	Mexico.	1825.	H. ȝ.	3. t. 223.

COMPOSITÆ.

4 lineáris. LG.	(ye.) linear-leaved.	S. America.	1825.	8. 10. S. ○.
5 marítima. K.S.	(ye.) sea-side.	Cuba.	1822.	— S. ○.
BALTIM'ORA. W.	BALTIM'ORA.	Syngenesia Polygamia Necessaria. L.		
1 recta. w.	(ye.) upright.	Vera Cruz.	1699.	6. 7. H. ○.
2 álba. p.s.	(wh.) white.	1824.	— H. ○.
ERIO'COMA. K.S.	ERIO'COMA.	Syngenesia Polygamia Frustranea.		
1 floribunda. K.S.	(wh.) many-flowered.	Mexico.	1828.	9. 11. F. ™.
2 frágans. D.D.	(wh.) sweet-scented.	F. ™.
SI'LPHIUM. W.	SI'LPHIUM.	Syngenesia Polygamia Necessaria. L.		
1 laciniátum. w.	(ye.) jagged-leaved.	N. America.	1781.	7. 9. H. ○.
2 compótum. w.	(ye.) scolop-leaved.	1789.	— H. ¼.
3 therebinthináceum.	(ye.) broad-leaved.	1765.	8. 9. H. ¼.
4 perfoliátum.	(ye.) perfoliate.	1766.	7. 10. H. ¼.
5 conjúntum. W.en.	(ye.) conjoined.	H. ¼.
6 connátum. w.	(ye.) round-stalked.	1765.	— H. ¼.
7 Asteriscus. w.	(ye.) hairy-stalked.	1732.	7. 9. H. ¼.
8 scábrum. n.	(ye.) rough-leaved.	1823.	— H. ¼.
9 pùmulum. w.	(ye.) dwarf woolly.	H. ¼.
10 trifoliátum. w.	(ye.) three-leaved.	1755.	7. 10. H. ¼.
11 ternátum. w.	(ye.) various-leaved.	1806.	— H. ¼.
12 atropurpureum. w.	(ye.) purple-stalked.	1812.	— H. ¼.
OSTEOSPE'R'MUM.	OSTEOSPE'R'MUM.	Syngenesia Polygamia Necessaria.	L.	
1 spinosum. H.K.	(ye.) rough-leaved.	C. B. S.	1700.	2. 10. G. ™.
2 spinéscens. H.K.	(ye.) smooth-leaved.	1793.	3. 6. G. ™.
3 pisiferum. w.	(ye.) smooth.	1757.	3. 5. G. ™.
4 moniliférum. w.	(ye.) Poplar-leaved.	1714.	7. 8. G. ™.
5 hirsútum. w.	(ye.) hairy.	1822.	— G. ™.
6 ciliátum. w.	(ye.) fringed.	1818.	— G. ™.
7 imbricátum. w.	(ye.) imbricated.	1824.	— G. ™.
8 calenduláceum. w.	(ye.) Marygold-like.	1823.	— G. ™.
9 ilicifólium. w.	(ye.) Holly-leaved.	1816.	— G. ™.
10 rígidum. w.	(ye.) rigid.	1774.	4. 7. G. ™.
11 cœréleum. w.	(bl.) blue-flowered.	6. 9. G. ™.
12 perfoliátum. w.	(ye.) perfoliate.	1816.	5. 8. G. ™.
13 níveum. w.	(ye.) white-leaved.	1820.	— G. ™.
14 corymbósium. w.	(ye.) corymbed.	G. ™.
15 incánum. w.	(ye.) hoary-leaved.	1816.	7. 8. G. ™.
16 polygaloides. w.	(ye.) Milkwort-leaved.	1759.	— G. ™.
				Burm. afr. 172. t. 62.
				Jacq. ic. 1. t. 179.
				Pluk. mant. t. 382.

Subordo VII. AMBROSIACEÆ. Link enum. 2. p. 366.

PARTH'ENIUM. W.	PARTH'ENIUM.	Syngenesia Polygamia Necessaria.	L.	
1 integrifólium. w.	(wh.) entire-leaved.	Virginia.	1661.	6. 10. H. ¼.
2 incánum. K.S.	(wh.) hoary.	Mexico.	1824.	— H. ○.
3 Hysteróphorus. w.	(wh.) cut-leaved.	W. Indies.	1728.	7. 10. H. ○.
Argyroch'etu bipinnatifida.	Cavan. icon. 4. t. 378.			
'IVA. W.	'IVA.	Syngenesia Polygamia Necessaria. L.		
1 ciliáta. w.	(ye.) ciliated.	N. America.	1820.	7. 8. H. ○.
2 ánnua. w.	(ye.) annual.	S. America.	1768.	— H. ○.
3 xanthiifólia. n.	(ye.) Xanthium-leaved.	Missouri.	1827.	— H. ○.
4 axillaris. Ph.	(ye.) axillary-flowered.	5. 6. H. ¼.
5 frutíscens. w.	(ye.) shrubby.	N. America.	1711.	8. H. ™.
AMBR'OSIA. W.	AMBR'OSIA.	Monocëa Pentandria. L.		
1 integrifólia. w.	(gr.) entire-leaved.	N. America.	1816.	7. 9. H. ○.
2 trípida. w.	(gr.) trifid-leaved.	1699.	— H. ○.
3 elátior. w.	(gr.) tall.	1696.	7. 8. H. ○.
4 artemisiifólia. w.	(gr.) Mugwort-leaved.	1759.	— H. ○.
5 paniculáta. w.	(gr.) paniced.	1811.	7. 9. H. ○.
6 marítima. w.	(gr.) sea.	Italy.	1570.	7. 8. H. ○.
7 cumanénis. K.S.	(gr.) Cumaná.	S. America.	1823.	9. 11. G. ¼.
8 peruviana. w.	(gr.) Peruvian.	Peru.	1821.	— G. ¼.
XANTH'IUM. W.	XANTH'IUM.	Monocëa Pentandria. L.		
1 Strumarium. w.	(gr.) small burdock.	England.	7. 9. H. ○.
2 orientále. w.	(gr.) oriental.	China.	1685.	— H. ○.
3 canadénse. s.s.	(gr.) Canadian.	Canada.	H. ○.
4 echinátum. w.	(gr.) hedgehog.	1818.	— H. ○.
				Eng. bot. v. 36. t. 2544.
				Linn. dec. 33. t. 17.

5 cathárticum. k.s. (gr.)	divided-leaved.	S. America.	1825.	9. 10. H. ⊙.	
6 spinòsum. w.	(gr.) spiny.	S. Europe.	1713.	7. 9. H. ⊙.	Herm. parad. 246.
FRANS'ERIA. W.	FRANS'ERIA.	Monocia Pentandria. W.			
1 artemisioides. w. (gr.)	Mugwort-leaved. Peru.		1759.	7. 9. G. h.	Willd. hort. ber. t. 2.
2 ambrosioídes. w. (gr.)	Ambrosia-leaved. Mexico.		1796.	— G. h.	Cavan. ic. 2. t. 200.

Subordo VIII. ANTHEMIDEÆ. Kth. synops. 2. p. 513.

CEPHALO'PHORA. W. CEPHALO'PHORA.	Syngenesia Polygamia Æqualis. W.				
glauca. w. (wh.) glaucous.	Chili.	1798.	7. 8. H. ⊙.	Cavan. ic. 6. t. 599.	
Græmia aromatica. Hook. ex. flor. 189.					
ETH'ULIA. W.	ETH'ULIA. Syngenesia Polygamia Æqualis. W.				
1 conyoídes. w. (li.) paniced.	India.	1776.	7. 8. S. ⊙.	Linn. fil. dec. t. 1.	
2 divaricata. w. (pu.) spreading.	—	1815.	— S. ⊙.	Lam. ill. t. 699.	
Epáltes divaricata. Cas.					
3 ramosa. H.B. (pu.) branching.	—	1818.	— S. ⊙.		
4 brasiliensis. L.en. (pu.) Brazilian.	Brazil.	1822.	— S. ȳ.		
SPARGANO'PHORUS. G. SPARGANO'PHORUS.	Syngenesia Polygamia Æqualis.				
1 Vaillantii. p.s. (pu.) Vaillant's.	E. Indies.	1824.	6. 8. S. ȳ.	Gært. fr. 2. t. 165. f. 4.	
Ethulia Sparganophorus. L.					
2 Struchioides. p.s. (pu.) axillary-flower'd. W. Indies.	—	1826.	— S. ȳ.	Brown. jam. t. 34. f. 2.	
3 verticillatūs. m. (pu.) whorl-leaved.	N. America.	1827.	8. 10. H.w. ȳ.		
TARCHONA'NTHUS. W. AFRICAN FLEABANE.	Syngenesia Polygamia Æqualis. L.				
1 camphoratus. w. (ye.) strong-scented.	C. B. S.	1690.	6. 10. G. h.	Lam. ill. t. 671.	
2 dentatus. p.s. (ye.) scentless.	—	1816.	— G. h.		
3 ellípticus. p.s. (ye.) elliptic-leaved.	—	1823.	— G. h.		
PODA'NTHUS. LG.	PODA'NTHUS. Syngenesia Polygamia Æqualis.				
1 míliqui. H.C. (ye.) Chilian.	Chile.	1826.	7. 9. F. h.		
2 gratus. D.D. (ye.) grateful.	—	1825.	— F. h.	Hor. phys. ber. 75. t. 16.	
Euxenia grata. s.s.					
OTA'NTHUS. L.en.	OTA'NTHUS. Syngenesia Polygamia Æqualis.				
marítimus. L.en. (ye.) sea-side.	England.	7. 9. H. h.	Eng. bot. v. 2. t. 141.	
Santolina maritima. E.B.					
SANTOLINA. W.	LAVENDER-COTTON. Syngenesia Polygamia Æqualis. L.				
1 Chámæ-Cyparissus. w. (ye.) common.	S. Europe.	1573.	7. H. h.	Lam. ill. t. 671. f. 3.	
2 ericoïdes. p.s. (ye.) Heath-like.	—	1827.	7. 9. H. h.		
3 squarrosa. w. (ye.) hoary.	—	1570.	7. 8. H. h.	Moris. s. 6. t. 3. f. 17.	
4 viridis. w. (ye.) dark-green.	—	1727.	7. H. h.		
5 rosmarinifolia. w. (ye.) Rosemary-leaved.	—	1683.	7. 9. H. h.	Sm. exot. bot. 2. t. 62.	
6 pectinata. s.s. (ye.) pectinate.	Spain.	1825.	— H. h.		
7 viscosa. s.s. (ye.) viscous.	—	—	— H. h.		
8 erithmifolia. W.en. (ye.) Samphire-leaved. S. Europe.	—	1823.	— H. ȳ.		
9 pinnata. Viv. (wh.) wing-leaved.	—	1791.	— F. h.		
10 anthemoides. w. (ye.) Chamomile-leaved.	—	1727.	7. 8. H. ȳ.	Flor. græc. t. 854.	
11 alpina. L. (ye.) alpine.	—	1798.	7. 9. H. ȳ.	Barr. rar. t. 522.	
ATHAN'ASIA. W.	ATHAN'ASIA. Syngenesia Polygamia Æqualis. L.				
1 crenata. w. (ye.) crenated.	C. B. S.	1816.	6. 8. G. h.		
2 punctata. w. (ye.) punctated.	—	1822.	5. 7. G. h.	Petiv. gaz. t. 81. f. 6.	
3 capitata. w. (ye.) hairy.	—	1774.	1. 3. G. h.	Moris. s. 6. t. 3. f. 48.	
4 pubescens. w. (ye.) villous-leaved.	—	1768.	6. 8. G. h.	Comm. hort. 2. t. 47.	
5 canescens. w. (ye.) canescent.	—	1822.	— G. h.	Cavan. ic. 1. t. 3.	
lanuginosa. Cav.					
6 cuneifolia. p.s. (ye.) wedge-leaved.	—	1816.	7. 9. G. h.	Lam. ill. t. 670. f. 3.	
7 ánnua. w. (ye.) annual.	Barbary.	1686.	7. 8. H. ⊙.	Bot. mag. t. 2276.	
8 dentata. w. (ye.) tooth-leaved.	C. B. S.	1759.	— G. h.	Comm. rar. t. 41.	
9 trifurcata. w. (ye.) trifid-leaved.	—	1710.	— G. h.	Comm. hort. 2. t. 49.	
10 virgata. w. (ye.) twiggy.	—	1815.	— G. h.	Jacq. schœn. 2. t. 148.	
11 tomentosa. w. (ye.) Lavender-leaved.	—	1774.	5. 6. G. h.	Lam. ill. t. 670. f. 1.	
cineraria. L.					
12 filiformis. w. (ye.) fine-leaved.	—	1787.	8. G. h.		
13 crithmifolia. w. (ye.) Samphire-leav'd.	—	1723.	7. 8. G. h.	Comm. hort. 2. t. 50.	
14 parviflora. w. (ye.) small-flowered.	—	1731.	4. G. h.	Jacq. schœn. 2. t. 149.	
15 tricuspid. p.s. (ye.) three-pointed.	—	1816.	— G. h.		
16 pectinata. w. (ye.) pectinated.	—	1774.	5. 6. G. h.		
17 pinnata. w. (ye.) pinnated.	—	1824.	6. 8. G. h.	Lam. ill. t. 670. f. 4.	
BALSAM'ITA. W.	COSTMARY. Syngenesia Polygamia Æqualis. W.				
1 virgata. w. (ye.) twiggy.	Italy.	1791.	6. 7. H. ȳ.	Jacq. obs. 4. t. 81.	

COMPOSITÆ.

2 ageratifòlia. w. (<i>ye.</i>) Ageratum-leav'd.	Candia.	1605.	6. 10.	G. $\frac{1}{2}$.	Alp. exot. t. 326.
3 vulgàris. w. (<i>ye.</i>) common.	Italy.	1568.	8. 9.	H. $\frac{1}{2}$.	Schk. hand. 3. t. 240.
4 ànnua. DC. (<i>ye.</i>) annual.	S. Europe.	1629.	7. 8.	H. \odot .	Mill. ic. 2. t. 227. f. 1.
Tanacètum ànnuum. w.					
PE'NTZIA. Th.	PE'NTZIA.	Syngenesia Polygamia Æqualis. W.			
flabellifòrmis. w. (<i>ye.</i>) fan-leaved.	C. B. S.	1774.	5. 8.	G. $\frac{1}{2}$.	Bot. mag. t. 212.
Tanacètum flabellifòrmis. B.M.					
TANACETUM. W.	TANSY.	Syngenesia Polygamia Superflua. L.			
1 vestitum. w. (<i>ye.</i>) imbricated.	C. B. S.	1816.	8. 9.	G. $\frac{1}{2}$.	
2 linifòlium. w. (<i>ye.</i>) Flax-leaved.	—	1774.	8.	G. $\frac{1}{2}$.	
3 suffruticòsum. w. (<i>ye.</i>) shrubby.	—	1751.	5. 9.	G. $\frac{1}{2}$.	Comm. hort. 2. t. 100.
4 canariénse. DC. (<i>ye.</i>) Canary.	Canaries.	1816.	—	G. $\frac{1}{2}$.	
5 incànum. s.s. (<i>ye.</i>) hoary.	Levant.	1827.	—	H. $\frac{1}{2}$.	Gmel. sib. 2. t. 65. f. 2.
6 sibiricum. w. (<i>ye.</i>) Siberian.	Siberia.	1823.	—	H. $\frac{1}{2}$.	
7 argénteum. w. (<i>ye.</i>) silvery.	Levant.	1812.	—	H. $\frac{1}{2}$.	
8 grandifòrum. w. (<i>ye.</i>) great-flowered.	C. B. S.	1825.	—	G. $\frac{1}{2}$.	
9 myriophyllum. w. (<i>ye.</i>) many-leaved.	Levant.	1818.	—	H. $\frac{1}{2}$.	
Achillea bipinnata. L.					
10 paucifòrum. s.s. (<i>ye.</i>) few-flowered.	N. America.	1827.	—	H. $\frac{1}{2}$.	
11 vulgàre. w. (<i>ye.</i>) common.	Britain.	—	—	H. $\frac{1}{2}$.	Eng. bot. v. 18. t. 1229.
β crispum. (<i>ye.</i>) curled.	—	—	—	H. $\frac{1}{2}$.	
ARTEMISIA. W.	WORMWOOD.	Syngenesia Polygamia Superflua. L.			
1 jùdàica. w. (<i>ye.</i>) Judean.	Levant.	1688.	8.	G. $\frac{1}{2}$.	Pluk. alm. t. 73. f. 2.
2 valentina. w. (<i>ye.gr.</i>) Spanish.	Spain.	1739.	7. 8.	F. $\frac{1}{2}$.	Barr. ic. t. 485.
3 arragonénis. w. (<i>ye.gr.</i>) snowy-white.	—	1816.	—	F. $\frac{1}{2}$.	Asso arrag. t. 8. f. 1.
4 salsoloides. s.s. (<i>ye.gr.</i>) saltwort-like.	Siberia.	1827.	—	F. $\frac{1}{2}$.	
5 subcanescens. w. (<i>ye.gr.</i>) hoary-leav'd.	S. Europe.	—	—	F. $\frac{1}{2}$.	
6 Abrotanum.w. (<i>ye.gr.</i>) Southernwood.	—	1548.	8. 10.	H. $\frac{1}{2}$.	Blackw. t. 555.
7 viridifòlia. s.s. (<i>ye.gr.</i>) green-leaved.	Siberia.	1828.	—	H. $\frac{1}{2}$.	Lobel. ic. 768.
8 pròcera. w. (<i>ye.gr.</i>) tall. <i>paniculata</i> . Lam.	S. Europe.	1816.	—	F. $\frac{1}{2}$.	Dill. elth. t. 33. f. 37.
9 tenuifòlia. w. (<i>ye.gr.</i>) slender-leaved.	—	1732.	9. 12.	G. $\frac{1}{2}$.	Flor. græc. t. 856.
10 àfra. w. (<i>wh.gr.</i>) Cape.	C. B. S.	1816.	—	G. $\frac{1}{2}$.	
11 arboréscens. w. (<i>ye.gr.</i>) tree.	Levant.	1640.	6. 8.	H. $\frac{1}{2}$.	
12 argéntea. w. (<i>ye.</i>) silvery.	Madeira.	1777.	6. 7.	G. $\frac{1}{2}$.	
13 lednicénsis. s.s. (<i>ye.gr.</i>) linear-lobed.	Hungary.	1829.	—	H. $\frac{1}{2}$.	
14 paucifòlia. s.s. (<i>ye.</i>) few-flowered.	Siberia.	1824.	—	H. $\frac{1}{2}$.	
15 glaciàlis. w. (<i>ye.gr.</i>) silky.	Switzerland.	1739.	7. 8.	H. $\frac{1}{2}$.	Jacq. aust. 5. t. ap. 35.
16 rupéstris. w. (<i>br.</i>) stone.	N. Europe.	—	—	H. $\frac{1}{2}$.	All. ped. t. 9. f. 1.
17 mutellina. w. (<i>ye.gr.</i>) Alpine.	Alps of Europ.	1815.	—	H. $\frac{1}{2}$.	Vill. daup. 3. t. 35.
18 spléndens. s.s. (<i>ye.</i>) glossy.	Caucasus.	1827.	—	H. $\frac{1}{2}$.	
19 furcata. s.s. (<i>br.</i>) forked.	Siberia.	1829.	—	H. $\frac{1}{2}$.	
20 lanata. s.s. (<i>wh.ye.</i>) woolly.	S. Europe.	1827.	—	H. $\frac{1}{2}$.	
21 alpina. M.B. (<i>ye.gr.</i>) alpine.	Caucasus.	—	—	H. $\frac{1}{2}$.	
22 jeniseánsis. s.s. (<i>ye.gr.</i>) leafy-panicled.	Siberia.	1824.	—	H. $\frac{1}{2}$.	Jacq. aust. 5. t. ap. 34.
23 pedemontana. L.en. (<i>br.</i>) Swiss.	Switzerland.	1823.	—	H. $\frac{1}{2}$.	Flor. dan. t. 801.
24 caucásica. w. (<i>ye.gr.</i>) Caucasian.	Caucasus.	1804.	6. 7.	H. $\frac{1}{2}$.	
25 spicata. w. (<i>br.</i>) spiked.	Switzerland.	1790.	—	H. $\frac{1}{2}$.	
26 norvègica. s.s. (<i>br.</i>) Norwegian. rupestrìs. Flor. dan. <i>nei aitorum</i> .	Norway.	1823.	—	H. $\frac{1}{2}$.	
27 grønländica. s.s. (<i>br.</i>) Greenland.	Greenland.	1827.	—	H. $\frac{1}{2}$.	
28 ludoviciàna. N. (<i>br.</i>) St. Louis.	Missouri.	—	—	H. $\frac{1}{2}$.	
29 desertorum. s.s. (<i>br.</i>) desert. <i>leucanthemifòlia</i> . Hort.	Tartary.	1817.	7. 9.	H. $\frac{1}{2}$.	Pall. it. 3. t. Hh. f. 2.
30 pectinata. w. (<i>br.</i>) comb-leaved.	Dauria.	1806.	—	H. \odot .	All. ped. 1. t. 10. f. 3.
31 tanacetifòlia. w. (<i>br.</i>) Tansy-leaved.	Siberia.	1768.	7. 8.	H. $\frac{1}{2}$.	
32 laciniata. w. (<i>gr.ye.</i>) jagged.	Dauria.	1823.	—	H. $\frac{1}{2}$.	Vill. delph. 3. t. 35.
33 insipida. w. (<i>br.</i>) insipid.	S. Europe.	—	—	H. $\frac{1}{2}$.	
34 mollissima. D.P. (<i>ye.</i>) soft-leaved.	Nepaul.	—	8. 11.	H. $\frac{1}{2}$.	
35 potentillæfòlia. s.s. (<i>gr.st.</i>) Cinquefoil-leav'd.	Siberia.	1827.	7. 9.	H. $\frac{1}{2}$.	
36 armeniaca. s.s. (<i>gr.ye.</i>) Armenian.	Armenia.	1828.	—	H. $\frac{1}{2}$.	
37 canescens. s.s. (<i>ye.ye.</i>) canescent.	—	—	—	H. $\frac{1}{2}$.	
38 Santonica. w. (<i>wh.gr.</i>) Tartarian.	Siberia.	1596.	9. 11.	H. $\frac{1}{2}$.	
39 scoparia. w. (<i>wh.gr.</i>) besom.	Hungary.	1796.	7. 9.	H. \odot .	
40 campétris. w. (<i>br.</i>) field.	England.	—	8.	H. $\frac{1}{2}$.	
41 herbacea. w. (<i>ye.gr.</i>) herbaceous.	Siberia.	1817.	—	H. $\frac{1}{2}$.	
42 Marschalliana. s.s. (<i>gr.ye.</i>) M. Bieberstein's. <i>inodora</i> . M.B. <i>pauciflora</i> . w.	—	1800.	8. 9.	H. $\frac{1}{2}$.	
43 virgata. s.s. (<i>br.</i>) twiggy.	N. America.	1827.	7. 9.	H. $\frac{1}{2}$.	

44	palústris. w.	(ye.) marsh.	Siberia.	1801.	7. 8. H. ⊖.	Gmel. sib. 2. t. 55.
45	neglécta. W.en.	(gr.ye.) neglected.	—	1815.	— H. ȳ.	Lob. ic. t. 769. f. 1.
46	camphorata. w.	(ye.gr.) Camphor-scent'd.	S. Europe.	1820.	— H. ȳ.	
47	álbida. W.en.	(st.) white-leaved.	Siberia.	—	— H. ȳ.	
48	húmilis. W.en.	(ye.gr.) dwarfish.	Carniola.	—	8. 10. H. ȳ.	
49	crithmifolia. w.	(br.) Samphire-leav'd.	Portugal.	1739.	— H. ȳ.	
50	saxatilis. w.	(wh.) rock.	Hungary.	1816.	6. 8. H. ȳ.	
51	glaúca. w.	(gr.) glaucous.	Siberia.	1806.	— H. ȳ.	
52	nútans. w.	(br.gr.) nodding.	Tartary.	1819.	7. 9. H. ȳ.	W. et K. hung. 1. t. 75.
53	monogyna. w.	(ye.) one-styled.	Hungary.	1816.	— H. ȳ.	
54	palmata. w.	(gr.ye.) palmated.	S. Europe.	1739.	6. 7. G. ȳ.	
55	nivea. W.en.	(gr.ye.) snowy.	Siberia.	1815.	— H. ȳ.	
56	marítima. w.	(br.) drooping-flower'd.	Britain.	—	8. 9. H. ȳ.	Eng. bot. v. 24. t. 1706.
57	gólica. w.	(br.) upright-flower'd.	—	—	— H. ȳ.	— t. 1001.
58	salina. w.	(wh.gr.) sea-side.	Germany.	1817.	— H. ȳ.	Spreng. halens. t. 12.
59	frágans. w.	(st.) Lavender-leav'd.	Armenia.	1739.	6. 7. H. ȳ.	
60	austriaca. w.	(br.) Austrian.	Austria.	1597.	8. 10. H. ȳ.	Jacq. aust. 1. t. 100.
61	frígora. s.s.	(gr.ye.) frigid.	Siberia.	—	— H. ȳ.	
62	vallesiaca. w.	(st.) downy.	Italy.	1739.	7. 8. H. ȳ.	
63	taúrica. w.	(wh.gr.) Taurian.	Tauria.	1824.	— H. ȳ.	
64	orientális. w.	(br.) oriental.	Armenia.	—	8. 10. H. ȳ.	
65	serícea. w.	(wh.) silky-leaved.	—	1796.	6. 7. H. ȳ.	Gmel. sib. t. 64. f. 1.
66	repens. w.	(br.) creeping.	—	1805.	— H. ȳ.	
67	ramòsa. s.s.	(st.) branching.	Canaries.	1816.	6. 8. G. ȳ.	
68	bal-amita. W.en.	(ye.) Balsam-scented.	—	—	— H. ȳ.	
69	póntica. w.	(ye.) Roman.	Austria.	1570.	9. H. ȳ.	Jacq. aust. 1. t. 99.
70	chamaemelifolia. w.(br.)	Chamomile-leav'd.	S. Europe.	1739.	7. 8. F. ȳ.	Vill. dauph. 3. t. 35.
71	ánnua. w.	(wh.gr.) annual.	Siberia.	1741.	— H. ⊖.	Am. ruth. t. 196. f. 23.
72	bienniás. Ph.	(ye.gr.) biennial.	Missouri.	1804.	— H. ȳ.	Bot. mag. t. 2472.
73	Absínthium. w.	(ye.) common.	Britain.	—	7. 10. H. ȳ.	Eng. bot. v. 18. t. 1230.
74	Sieversiana. w.(br.gr.)	Sievers's.	Siberia.	1800.	7. 8. H. ȳ.	
75	vulgáris. w.	(pu.) Mugwort.	Britain.	—	8. 9. H. ȳ.	
76	procumbens. dc.	(ye.gr.) procumbent.	Siberia.	1823.	— H. ȳ.	Eng. bot. v. 14. t. 978.
77	incisa. W.en.	(ye.pu.) cut-leaved.	—	—	— H. ȳ.	
78	fasciculàta. M.B.	(ye.pu.) fascicled.	Iberia.	1820.	— H. ȳ.	
79	élégans. F.	(ye.) elegant.	China.	—	7. 9. H. ȳ.	
80	camtschática. F.	(gr.st.) Kamtschatka.	Kamtschatka.	1827.	8. 10. H. ȳ.	
81	sacrórum. Led.	(gr.st.) sacred.	Siberia.	1828.	7. 9. H. ȳ.	
82	Mertensiàna. Wh.(gr.st.)	Mertens's.	Hungary.	1826.	— H. ȳ.	Rheed. mal. 10. t. 45.
83	índica. w.	(ye.gr.) Indian.	E. Indies.	1796.	9. 10. F. ȳ.	Gmel. sib. 2. t. 48. f. 1.2.
84	integrifolia.w.	(ye.gr.) entire-leaved.	Siberia.	1759.	7. 8. H. ȳ.	
85	japónica. w.	(wh.) Japanese.	Japan.	1804.	10. 11. H. ȳ.	
86	dracunculoides. Ph.	(gr.) Tarragon-like.	N.America.	1811.	8. 10. H. ȳ.	
87	cérnua. N. nútans.	Fraser.	—	—	—	
88	serràta. N.	(gr.) saw-leaved.	Missouri.	1827.	— H. ȳ.	
89	gnaphalioïdes. N.(br.)	Cudweeed-like.	—	—	— H. ȳ.	
90	cæruléscens. w.	(ye.) blueish.	England.	—	8. 10. H. ȳ.	Eng. bot. v. 34. t. 2426.
91	inodòra. w.	(ye.) inodorous.	Siberia.	1548.	7. 8. H. ȳ.	Gmel. sib. 2. t. 59&60. f. 1.
92	Dracunculus. w.	(wh.gr.) Tarragon.	S. Europe.	—	— H. ȳ.	Gmel. sib. 2. t. 61. f. 1.2.
ERIOCEPHALUS. W. WOOLHEAD.			Syngenesia Polygamia Necessaria. L.			
1	africánus. w.	(ye.) cluster-leaved.	C. B. S.	1732.	1. 4. G. ȳ.	Bot. mag. t. 833.
	frutéscent. H.K.					
2	decussátus. B.T.	(ye.) cross-leaved.	—	1816.	3. 6. G. ȳ.	
3	purpureus. B.T.	(pu.) purple.	—	—	— G. ȳ.	
4	racemosus. w.	(ye.) silver-leaved.	—	1739.	3. 4. G. ȳ.	
5	spinéscent. B.T.	(ye.) spiny.	—	1816.	— G. ȳ.	
HI'PPIA. W.	HI'PPIA.		Syngenesia Polygamia Necessaria. L.			
1	frutéscent. w.	(ye.) shrubby.	C. B. S.	1710.	2. 8. G. ȳ.	Bot. mag. t. 1855.
2	integrifolia. w.	(ye.) annual.	E. Indies.	1777.	7. 8. S. ⊖.	Hout. n. hist. t. 67. f. 2.
SOL'IVA. L.T.	SOL'IVA.		Syngenesia Polygamia Necessaria.			
1	anthemifolia.	(gr.) Chamomile-leav'd.	S. America.	1812.	4. 12. H. ⊖.	J. ann. mus. t. 61. f. 1.
	Gymnóstyles anthemifolia. P.S.					
2	pterospérma. J.	(gr.) wing-seeded.	Brazil.	1827.	5. 9. H. ⊖.	
3	stolonifera. L.T.	(gr.) stoloniferous.	Portugal?	1818.	4. 12. H. ⊖.	
	Hippia stolonifera. P.S.	Gymnóstyles lusitánica. s.s.				
4	minùta.	(ye.) fringe-leaved.	S. America.	1825.	— H. ⊖.	
	Hippia minùta. L.	Gymnóstyles minùta. s.s.				
5	pedicellata. P.S.	(gr.) pedicelled.	S. America.	—	— H. ⊖.	

LIDBE'CKIA. W.	LIDBE'CKIA.	Syngenesia Polygamia Superflua. W.			
1 lobata. w. (ye.) lobe-leaved.	C. B. S.	1816. 5. 6. G. ™.			Lam. ill. t. 701. f. 3.
<i>Lancisia lobata.</i> p.s. <i>Cotula quinqueloba.</i> L.					Berg. cap. 306. t. 5. f. 9.
2 pectinata. w. (ye.) silver-leaved.	C. B. S.	1774. 5. 6. G. ™.			Lam. ill. t. 701. f. 1.
C'ENIA. J.	C'ENIA.	Syngenesia Polygamia Superflua.			
turbinata. p.s. (wh. re.) turbinated.	C. B. S.	1713. 7. 8. H. ⊙.			
<i>Lidbeckia turbinata.</i> L. <i>Cotula turbinata.</i> L.					Dill. elth. t. 23. f. 25.
C'OTULA. W.	C'OTULA.	Syngenesia Polygamia Superflua. W.			
1 anthemoïdes. w. (ye.) Anthemis-like.	St. Helena.	1696. 7. 8. S. ⊙.			Houtt. syst. t. 69. f. 4.
2 aurea. w. (ye.) golden.	S. Europe.	1823. 7. 9. H. ⊙.			Lam. ill. t. 700. f. 1.
3 nudicaulis. w. (ye.) naked-stalked.	C. B. S.	1816. — G. ¼.			
4 coronopifolia. w. (ye.) Buck's Horn.	—	1683. — G. ⊙.			Pluk. man. t. 430. f. 7.
5 umbellata. w. (ye.) umbel-flowered.	—	1822. — H. ⊙.			
6 viscosa. w. (wh.) clammy.	Vera Cruz.	1739. 8. S. ¼.			
7 tanacetifolia. w. (ye.) Tansy-leaved.	C. B. S.	1780. 6. 8. H. ⊙.			
8 sericea. w. (ye.) silky.	—	1816. — G. ¼.			
9 sphæranthus. L.en. (ye.) round-headed.	Congo.	1820. — S. ¼.			
GRA'NGEA. J.	GRA'NGEA.	Syngenesia Polygamia Superflua.			
1 cuneifolia. Lk. (br. ye.) wedge-leaved.	China.	1816. 7. 9. H. ⊙.			Lam. ill. t. 699. f. 2.
2 minima. w. (ye.) least.	—	1765. — H. ⊙.			Burm. ind. t. 58. f. 3.
3 decumbens. Df. (ye.) decumbent.	N. Holland.	1815. — H. ⊙.			Lam. ill. t. 699. f. 3.
4 maderaspatana. s. (ye.) Madras.	E. Indies.	1780. 7. 8. S. ⊙.			Lam. ill. t. 699. f. 1.
<i>Cotula maderaspatana.</i> W.					
5 bicolor. Df. (ye.) two-coloured.	—	1804. — S. ⊙.			
<i>Cotula bicolor.</i> w. <i>latifolia.</i> P.S.					
6 cinerea. L.en. (ye.) grey.	Egypt.	1825. — H. ⊙.			
ANAC'YCLUS. W.	ANAC'YCLUS.	Syngenesia Polygamia Superflua. W.			
1 créticus. w. (wh.) trailing.	Candia.	1759. 6. 8. H. ⊙.			Ann. mus. 11. t. 22.
2 orientalis. w. (wh.) oriental.	Levant.	1731. — H. ⊙.			Boer. lugd. 1. t. 110.
3 aureus. w. (ye.) golden-flower'd.	—	1570. — H. ⊙.			Lam. ill. t. 700. f. 2.
4 radiatus. L.en. (ye.) rayed.	—	1823. — H. ⊙.			
5 alexandrinus. w. (wh.) Egyptian.	Egypt.	1828. — H. ⊙.			
6 clavatus. P.S. (wh.) clavate.	Barbary.	1823. — H. ⊙.			
7 Pyréthrum. L.en. (w.) Pelliitory of Spain. S. Europe.	1570.	6. 7. H. ¼.			
<i>Anthemis Pyréthrum.</i> w.					Bot. mag. t. 462.
8 valentinus. w. (ye.) fine-leaved.	Spain.	1656. — H. ⊙.			Schk. han. 3. t. 254. b.
9 purpurascens. P.S. (ye. cr.) two-coloured.	S. Europe.	1829. — H. ⊙.			
10 officinarum. Hayn. (wh.) officinal.	—	1828. — H. ⊙.			
11 inflatus. Leh. (wh.) inflated.	Siberia.	1829. — H. ⊙.			
12 tomentosus. Leh. (wh.) woolly.	—	— — H. ⊙.			
13 divaricatus. Cav. (wh.) spreading.	—	1824. — H. ⊙.			
ZALUZ'ANIA. P.S.	ZALUZ'ANIA.	Syngenesia Polygamia Superflua.			
triloba. p.s. (ye.) three-lobed.	Mexico.	1798. 7. 9. F. ™.			
<i>A'nthemis triloba.</i> w.					
FERDINA'NDIA. LG. FERDINA'NDIA.	Syngenesia Polygamia Superflua.				
1 augusta. LG. (ye.) august.	Mexico.	1816. 8. 10. F. ™.			
2 integrifolia. D.D. (ye.) entire-leaved.	—	1828. — F. ™.			
ANTHEMIS. W.	CHAMOMILE.	Syngenesia Polygamia Superflua. L.			
1 rigescens. W.en. (wh.) rigid.	Caucasus.	1805. 7. 9. H. ¼.			W. hort. ber. 1. t. 62.
<i>nigricans.</i> Hort.					
2 Cota. w. (wh.) Venetian.	Italy.	1714. 7. 8. H. ⊙.			
3 altissima. w. (wh.) tall.	S. Europe.	1731. 8. H. ⊙.			Pluk. alm. t. 17. f. 5.
4 punctata. w. (wh.) dotted.	Levant.	1825. 7. 9. G. ™.			Flor. græc. t. 881.
5 maritima. w. (wh.) sea.	England. 7. 8. H. ⊙.			Desf. atl. 2. t. 139.
6 tomentosa. w. (wh.) downy.	Levant.	1795. 7. 10. H. ¼.			Eng. bot. v. 33. t. 2370.
7 ruthénica. M.B. (wh.) Russian.	Tauria.	1823. — H. ⊙.			
8 fruticulosa. M.B. (wh.) frutescent.	—	— H. ™.			
9 pubescens. w. (wh.) pubescent.	S. Europe.	1803. 7. 8. H. ¼.			
10 biaristata. s.s. (wh.) two-awned.	S. France.	1822. — H. ¼.			
11 revoluta. s.s. (wh.) revolute.	Canaries.	1816. 8. 10. G. ™.			
12 secundiramea. s.s. (wh.) secund-branch'd.	Sicily.	1824. 7. 9. H. ⊙.			
13 mixta. w. (wh.) simple-leaved.	France.	1731. 7. 8. H. ⊙.			
14 Kitaibeli. (wh.) Kitaibel's.	Hungary.	1807. — H. ¼.			
15 coronopifolia. w. (wh.) Buckshorn-leav'd.	Spain.	— — H. ¼.			
16 australis. w. (wh.) procumbent.	S. France.	1818. — H. ¼.			
17 ibérica. M.B. (wh.) Iberian.	Iberia.	1829. — H. ¼.			
18 Barrelieri. s.s. (wh.) Barrelier's	Naples.	— — 8. 10. H. ⊙.			
19 saxatilis. DC. (wh.) rock.	S. France.	1807. 7. 8. H. ¼.			
20 alpina. w. (wh.) Alpine.	Europe.	— — H. ¼.			Jacq. aust. 6. ap. t. 30.

21	carpática. w.	(wh.) Carpathian.	Carpathian-m.	1818.	7. 8. H. ¼.		
22	corymbosa. w.	(wh.) corymbose.	Austria.	1821.	— H. ¼.		
23	retusa. s.s.	(wh.) retuse.	1824.	— H. ¼.		
24	ramosa. s.s.	(wh.) branching.	—	9. 11. H. ¼.		
25	mucronulata.s.s.(wh.)	mucronulate.	Italy.	1829.	8. 10. H. ⊖.		
26	Chamomilla. W.	(wh.) various-leaved.	S. Europe.	—	— H. ¼.		
27	chia. w.	(wh.) cut-leaved.	Chio.	1731.	6. 10. H. ⊖.		
28	nobilis. w.	(wh.) common.	Britain.	7. 9. H. ¼.	Eng. bot. v. 14. t. 908.	
29	aryensis. w.	(wh.) corn.	—	6. 8. H. ½.	— v. 9. t. 602.	
30	austriaca. w.	(wh.) Austrian.	Austria.	1759.	5. 8. H. ⊖.	Jacq. aust. 5. t. 444.	
31	Cótula. w.	(wh.) stinking.	Britain.	6. 9. H. ⊖.	Eng. bot. v. 25. t. 1772.	
32	fállax. W.en.	(wh.) false Chamomile.	—	—	H. ⊖.		
33	fuscata. w.	(wh.) brown-scaled.	Portugal.	1805.	7. 8. H. ⊖.		
34	peregrina. w.	(wh.) close-branched.	Piedmont.	1817.	— H. ⊖.		
35	candidissima.s.s.(wh.)	silvery-leaved.	Caucasus.	1812.	— H. ¼.		
36	montana. w.	(wh.) mountain.	Italy.	1759.	7. 10. H. ¼.	Ger. prov. t. 8.	
37	incrassata. s.s.(wh.)	thick-stalked.	S. France.	1823.	— H. ⊖.		
38	pedunculata.s.s.(wh.)	large-stalked.	Portugal.	—	8. 10. H. ⊖.		
39	petraea. L.en.	(wh.) long-stalked.	Naples.	—	— H. ⊖.		
40	apiifolia. B.R.	(wh.) Parsley-leaved.	—	1764.	8. 9. H. ¼.	Bot. reg. t. 527.	
41	minuta. Leh.	(wh.) minute.	—	1829.	— H. ⊖.		
42	suffruticosa. Leh.	(wh.) suffrutescent.	—	—	6. 9. H. ½.		
43	globosa. w.	(ye.) globe.	Naples.	—	8. 9. G. ¼.	Jacq. schoen. 3. t. 371.	
44	valentina. w.	(ye.) purple-stalked.	S. Europe.	1596.	7. 8. H. ⊖.	Breyn. cent. t. 75.	
45	tinctoria. w.	(ye.) Ox Eye.	Britain.	6. 11. H. ¼.	Eng. bot. v. 21. t. 1472.	
46	discoidea. w.	(ye.) saw-leaved.	Italy.	1800.	7. 8. H. ⊖.		
47	Marschalliana.w.(ye.)	M. Bieberstein's.	Caucasus.	1828.	— H. ¼.		
48	Rudolphiana.M.B.(ye.)	Rudolph's.	—	—	— H. ¼.		
CLADA'NTHUS. Cas.	CLADA'NTHUS.	—	Syngenesia	Polygamia	Superflua.		
arábicus. Cas.	(ye.) Arabian.	Barbary.	1759.	7. 8. H. ⊖.		Smith spic. 9. t. 10..	
A'nthemis arábica. w.							
MATRIC'ARIA. W.	MATRIC'ARIA.	—	Syngenesia	Polygamia	Superflua. L.		
1	suavolens. w.	(wh.) sweet.	Europe.	1781.	6. 8. H. ⊖.		
2	Chamomilla. w.	(wh.) Wild Chamomile.	Britain.	5. 7. H. ⊖.	Eng. bot. v. 18. t. 1232.	
3	capensis. w.	(wh.) Cape.	C. B. S.	1699.	7. 9. G. ½.	Seb. thes. 1. t. 16. f. 2.	
4	pusilla. W.en.	(wh.) small.	—	—	— H. ⊖.		
ACHILL'EA. W.	MILFOIL.	—	Syngenesia	Polygamia	Superflua. L.		
1	lingulata. w.	(wh.) tongue-leaved.	Hungary.	1815.	7. 8. H. ¼.		
2	Hérba-rota. w.	(wh.) Herbarota.	France.	1640.	6. 7. H. ¼.	W. et K. hung. 1. t. 2.	
3	grandiflora. M.B.(wh.)	great-flowered.	Caucasus.	1815.	7. 8. H. ¼.	All. ped. 1. t. 9. f. 3.	
4	Ptámica. w.	(wh.) Sneezewort.	Britain.	—	7. 11. H. ¼.	Eng. bot. v. 11. t. 757.	
	β flòre pleno.	(wh.) double-flowered.	—	—	— H. ¼.		
5	cristata. w.	(wh.) slender-branch'd.	Italy.	1784.	7. 8. H. ¼.		
6	Ageratum. w.	(ye.) Sweet Maudlin.	S. Europe.	1570.	8. 10. H. ¼.		
7	decolorans.W.en.(st.)	pale yellow.	England.	1798.	6. 8. H. ¼.	Eng. bot. v. 36. t. 2531.	
	serrata. E.B. nec aliorum.						
8	speciosa. W.en.	(wh.) spear-leaved.	—	1804.	7. 9. H. ¼.		
9	biserrata. M.B.(wh.)	two-toothed.	Iberia.	1825.	— H. ¼.		
10	salicifolia. F.	(wh.) Willow-leaved.	Siberia.	1827.	— H. ¼.		
11	dentifera. Jac.	(wh.) tooth-leaved.	—	1824.	— H. ¼.		
12	alpina. w.	(wh.) Alpine.	Siberia.	1731.	7. 11. H. ¼.	Bocc. mus. 144. t. 101.	
13	serrata. w.	(wh.) saw-leaved.	Switzerland.	1686.	8. 9. H. ¼.		
14	coronopifolia.w.(st.)	buckshorn-leav'd.	Levant.	1816.	— H. ¼.		
15	Clavénna. w.	(wh.) silver-leaved.	Austria.	1656.	6. 7. H. ¼.	Bot. mag. t. 1287.	
16	Thomasiána.DC.(wh.)	Thomas's.	S. France.	1828.	— H. ¼.		
17	impatiens. w.	(wh.) impatient.	Siberia.	1759.	6. 9. H. ¼.	Gmel. sib. 2. t. 83. f. 1.	
18	pectinata. w.	(st.) comb-leaved.	Hungary.	1801.	8. 9. H. ¼.	W. et K. hung. 1. t. 34.	
	ochroleúca. w.K. non w.						
19	Gérberi. w.	(ye.) various-leaved.	Siberia.	1816.	— H. ¼.	Gmel. sib. 2. t. 83. f. 2.	
20	pilosa. w.	(ye.) hairy.	—	1826.	— H. ¼.		
21	squarrrosa. w.	(wh.) rough-headed.	—	1775.	7. 8. H. ¼.		
22	falcata. w.	(wh.) sickle-leaved.	Levant.	1739.	6. 9. H. ¼.	Lam. ill. t. 683. f. 3.	
23	santolinoides.LG.(w.)	Santolina-like.	Spain.	1824.	— H. ¼.		
24	tenuifolia. w.	(ye.) slender-leaved.	Levant.	1733.	6. 8. H. ½.	Flor. græc. t. 891.	
25	Santolina. w.	(ye.) Lavender-cotton-leav'd.—	—	1759.	— H. ¼.		
26	álbida. s.s.	(st.) white woolly.	—	1819.	6. 10. H. ¼.		
27	anthemoídes.w.	(ye.) Chamomile-like.	—	—	— H. ¼.		
28	decumbens. s.s.(ye.)	decumbent.	Kamtschatka.	1800.	6. 8. H. ¼.		
29	atrata. w.	(wh.) black-cupped.	Austria.	1596.	7. 9. H. ¼.	Jacq. aust. 1. t. 77.	

COMPOSITÆ.

30 moschàta. w. (wh.) musk.		Italy.	1775.	6. 7. H. ¼.	Jacq. aust. 5. t. ap. 83.
31 nàna. w. (wh.) dwarf.		Candia.	1759.	6. 8. H. ¼.	All. ped. 1. t. 9. f. 3.
32 crética. w. (wh.) Cretan.		Levant.	1739.	7. 8. H. ¼.	Bocc. mus. t. 34.
33 aëgyptiaca. w. (ye.) Egyptian.		Italy.	1640.	7. 9. G. ¼.	Flor. græc. t. 892.
34 macrophylla. w. (wh.) large-leaved.		Greece.	1710.	7. 8. H. ¼.	Triumf. obs. t. 23.
35 holosericea. F.G. (wh.) silky-leaved.		Naples.	1823.	7. 9. H. ¼.	
36 punctata. T.N. (st.) dotted.		Tauria.	1828.	— H. ¼.	
37 leptophylla. M.B. (ye.) narrow-leaved.		—	1823.	— H. ¼.	
38 taúrica. M.B. (st.) Taurian.		—	1829.	— H. ¼.	Buxb. cent. 2. t. 19.
39 àurea. w. (go.) golden-flower'd.	Levant.	1739.	6. 9. H. ¼.		
40 Eupatòrium. w. (go.) Caspian.	CaspianShores.	1803.	7. 8. H. ¼.		
41 filipendulina. s.s. (go.) short-rayed.	—	1812.	— H. ¼.		
42 compácta. w. (ye.) compact.	—	—	— H. ¼.		
43 pubescens. w. (st.) downy.	Levant.	1739.	6. 9. H. ¼.		
44 crithmifòlia. w. (wh.) Samphire-leav'd.	Hungary.	1804.	7. 8. H. ¼.		
45 tanacetifòlia. w. (ro.) Tansy-leaved.	Switzerland.	1658.	— H. ¼.		
46 distans. w. (wh.) branching.	Italy.	1804.	— H. ¼.		
47 lanàta. W.en. (wh.) woolly.	—	—	— H. ¼.		
48 magna. w. (wh.) great.	S. Europe.	1683.	6. 11. H. ¼.		
49 Millefolium. w. (wh.) Yarrow.	Britain.	—	6. 10. H. ¼.		
β rùbra. (re.) red-flowered.	—	—	— H. ¼.		
50 asplenifòlia. p.s. (ro.) Rose-coloured.	N. America.	1803.	6. 8. H. ¼.		
51 micrantha. w. (st.) small-flowered.	Levant.	1805.	6. 10. H. ¼.		
52 tomentosa. w. (ye.) tomentose.	S. Europe.	1648.	5. 10. H. ¼.		
53 ochroleuca. w. (st.) cream-coloured.	—	1804.	7. 9. H. ¼.		
54 montana. Len. (wh.) mountain.	Switzerland.	1798.	— H. ¼.		
55 bannática. w.k. (wh.) Hungarian.	Hungary.	1823.	— H. ¼.		
56 microphylla. w. (wh.) small-leaved.	Spain.	1800.	— H. ¼.		
57 ligística. w. (wh.) Ligurian.	Italy.	1791.	6. 8. H. ¼.		
58 nobilis. w. (wh.) showy.	Germany.	1640.	— H. ¼.		
59 myriophylla. s.s. (wh.) many-leaved.	—	1798.	7. 9. H. ¼.		
60 odorata. w. (wh.) sweet-scented.	Spain.	1729.	6. 8. H. ¼.		
61 setacea. w. (wh.) bristly.	Hungary.	1805.	5. 6. H. ¼.		
62 abrotanifòlia. w. (ye.) Southernwood-leav'd.	Levant.	1739.	6. 8. H. ¼.		
63 chamomelifòlia. DC. (wh.) Chamomile-ld.	Pyrenees.	1820.	— H. ¼.		
64 mongólica. F. (wh.) Mongolic.	Tartary.	1823.	— H. ¼.		
65 sibírica. Df. (wh.) Siberian.	Siberia.	1800.	— H. ¼.		
66 rosea. w.k. (ro.) rosy.	Hungary.	1803.	— H. ¼.		
67 dubia. F. (wh.) doubtful.	Siberia.	1800.	— H. ¼.		
68 heterophylla. s. (wh.) different-leaved.	—	1804.	7. 9. H. ¼.		
69 intermédia. s. (wh.) intermediate.	—	1824.	— H. ¼.		
70 paradóxa. L.C. (wh.) uncertain.	—	1822.	— H. ¼.		
CHRYSANTHEMUM. W. CHRYSANTHEMUM.	Syngenesia	Polygamia	Superflua. L.		
1 atránum. w. (wh.) fleshy-leaved.	Austria.	1731.	7. 8. H. ¼.		
2 Leucánthemum. w. Ox-eye Daisy.	Britain.	—	6. 7. H. ¼.	Eng. bot. v. 9. t. 601.	
3 montànum. w. (wh.) mountain.	France.	1759.	— H. ¼.	Jacq. obs. 4. t. 91.	
4 heterophylum. w. (wh.) various-leaved.	Switzerland.	1806.	7. 8. H. ¼.		
5 sylvestre. W.en. (wh.) wood.	Portugal.	1804.	6. 7. H. ¼.		
6 hybrídum. Gus. (st.) hybrid.	Sicily.	1827.	— H. ¼.		
7 paludósium. w. (wh.) marsh.	Barbary.	1817.	8. 9. H. ¼.		
8 rotundifòlium. w. (wh.) round-leaved.	Hungary.	1819.	7. 8. H. ¼.		
9 mexicánum. K.S. (wh.) Mexican.	Mexico.	1827.	8. 11. H. ¼.		
10 graminifòlium. w. (wh.) Grass-leaved.	Montpelier.	1739.	5. 7. H. ¼.		
11 integrifòlium. s.s. (wh.) entire-leaved.	N. America.	1827.	6. 8. H. ¼.		
12 perpusillum. s.s. (wh.) small.	Corsica.	—	— H. ¼.		
13 anomálum. P.S. (wh.) anomalous.	Spain.	1824.	7. 9. H. ½.		
14 daucifòlium. P.S. (wh.) carrot-leaved.	—	—	— H. ¼.		
15 monspeliense. w. (wh. pu.) Montpelier.	—	1739.	6. 9. H. ¼.	Jacq. obs. 4. t. 93.	
16 Achilléæ. w. (wh.) Milfoil-leaved.	Italy.	1775.	6. 8. H. ¼.	Mich. gen. 34. t. 29.	
17 argéntum. w. (wh.) silver-leaved.	Levant.	1731.	7. 8. H. ¼.	Willd. hort. ber. t. 33.	
18 árticum. w. (bh.) northern.	Kamtschatka.	1801.	6. 8. H. ¼.	Bot. mag. t. 508.	
19 carinátum. w. (wh. pu.) three-coloured.	Barbary.	1796.	7. 10. H. ○.		
20 pùmilum. W.en. (wh.) small.	C. B. S.	1806.	— H. ○.		
21 coronárium. w. (wh. ye.) garden.	Sicily.	1629.	— H. ○.		
α album. (wh.) white-flowered.	—	—	— H. ○.		
β àureum. (ye.) golden.	—	—	— H. ○.		
22 ségetum. w. (ye.) corn.	Britain.	—	6. 8. H. ○.	Eng. bot. v. 8. t. 540.	
23 Mycónis. w. (ye.) tongue-leaved.	Italy.	1775.	7. 8. H. ○.	Jacq. obs. 4. t. 94.	
24 itálicum. w. (ye.) Italian.	—	1796.	6. 7. H. ¼.		

25	tripartitum. B.F.G. (ye.) three-parted.	China.	1820.	10. 12. F. 4.	Swt. br. fl. gar. t. 193.
26	índicum. P.S. (ye.) Indian.	_____	1819.	— G. ḥ.	Hort. trans. v. 4. pl.12.
27	sinense. L.T. (va.) variable chinese.	_____	1790.	— H. ḥ.	
	<i>A'nthemis artemisia folia. w. Chrysanthemum índicum.</i> B.M. non L.				
β	purpureum. (pu.) old purple.	_____		— H. ḥ.	
γ	variabile. (wh.) changeable white.	_____	1802?	— H. ḥ.	Bot. mag. t. 327.
δ	tubulosum álbum. (wh.) quilled white.	_____	1808.	— H. ḥ.	— t. 2042.
ε	supérbum. (wh.) superb white.	_____	1817.	— H. ḥ.	Bot. reg. t. 4.
ζ	tessellatum. (wh.) tasselled white.	_____	1816.	— H. ḥ.	— t. 455.
η	tubulosum luteum. (ye.) quilled yellow.	_____	1802.	— H. ḥ.	
ς	sulphureum. (st.) straw-coloured.	_____	—	— H. ḥ.	
ι	aureum. (go.) golden yellow.	_____	—	— H. ḥ.	Bot. reg. t. 4. f. sup.
κ	díscolor. (ti.) large lilac.	_____	1808.	— H. ḥ.	
λ	lilacinum. (pi.) pink or lilac.	_____	1798.	— H. ḥ.	
μ	cípreum. (co.) buff or copper-colour'd.	_____		— H. ḥ.	
ν	fúlvum. (br.) Spanish brown.	_____	1806.	— H. ḥ.	
ξ	flámeum. (ye.) quilled flame yellow.	_____	1819.	— H. ḥ.	Hort. trans. v. 4. pl.14.
ο	tubulosum róseum. (hi.) quilled pink.	_____	—	— H. ḥ.	Bot. reg. t. 616.
π	atropurpureum. (cr.) early crimson.	_____	1820.	— H. ḥ.	Hort. trans. v.5. pl.3.inf.
ρ	aurantiacum. (or.) large quilled orange.	_____	—	— H. ḥ.	— v.5. pl.3.f.sup.
σ	expansum. (li.) expanded light purple.	_____	—	— H. ḥ.	Bot. mag. t. 2556.
τ	purpurascens. (li.) quilled light purple.	_____	—	— H. ḥ.	
υ	involutum. (li.) curled lilac.	_____	—	— H. ḥ.	Swt. br. fl. gar. t. 7.
φ	fasciculatum. (ye.) superb clustered yellow.	—	—	— H. ḥ.	— t. 14.
χ	tubulosum cárneum. semidouble quilled pink.	—	—	— H. ḥ.	Hort. tr. v.5. pl.17*. inf.
ψ	álbum semidúplex. semidouble quilled white.	—	—	— H. ḥ.	Hort. tr. v.5. pl.17**.inf.
ω	tubulosum aurántium. semidouble quilled orange.	—	—	— H. ḥ.	Hort. t.v.5.pl.17**.sup.
aa	serotinum. (p. pu.) late pale purple.	—	—	— H. ḥ.	Hort. t.v.5.pl.17**.f.sup.
ββ	salmoneum. (sa.) quilled salmon-colour'd.	—	—	— H. ḥ.	
γγ	párvulum. (ye.) small yellow.	—	—	— H. ḥ.	
δδ	papyráceum. (wh.) paper white.	—	1821.	— H. ḥ.	
εε	pállidum. (li.) late pale pink.	—	1822.	— H. ḥ.	
ζζ	chrysocómum. (ye.) tasselled yellow.	—	1824.	— H. 4.	
ηη	Waratáh. (ye.) yellow Waratah.	—	—	— H. 4.	
ςς	Sabiñi. (ye.) golden Lotus.	—	—	— H. 4.	
ιι	chryseides. (ye.) double Indian yellow.	—	—	— H. 4.	
κκ	Pákkii. (ye.) Park's small yellow.	—	—	— H. 4.	
λλ	pállens. (p.or.) semidouble quilled pale orange.	—	—	— H. 4.	
μμ	stramineum. (st.) pale buff.	—	—	— H. 4.	
νν	mutábile. (st.) changeable pale buff.	—	—	— H. 4.	
ξξ	bicolor. (v.) two-coloured incurved.	—	—	— H. 4.	
οο	versicolor. (re.) two-coloured red.	—	—	— H. 4.	
ππ	stellatum. (pu.) starry purple.	—	—	— H. 4.	
ρρ	ornátum. (ti.) tasselled lilac.	—	—	— H. 4.	
σσ	fulvescens. (pu.) brown purple.	—	—	— H. 4.	
ττ	verecundum. (bh.) early blush.	—	—	— H. 4.	
vv	blándum. (bh.) blush.	—	—	— H. 4.	
φφ	leucánthum. (wh.) double Indian white.	—	—	— H. 4.	

CENTROSPE'RMUM. S.S. CENTROSPE'RMUM. Syngenesia Polygamia Superflua.

chrysánthemum.s.s.(ye.)yellow-flower'd. Spain.

Heteránthemis hírtta. Schot.

PYR'ETHRUM. W. FEVERFEW. Syngenesia Polygamia Superflua. W.

1	féniculaceum. W.en. (wh.) Fennel-leav'd. Teneriffe.	1815.	1. 12. G. ḥ.	Bot. reg. t. 272.	
2	erithmifolium. W.en. (wh.) Samphire-leav'd.	—	— G. ḥ.		
3	anethifolium. W.en. (wh.) Dill-leaved.	—	— G. ḥ.		
4	frutéscentia. W.en. (wh.) shrubby.	Canaries.	1699.	— G. ḥ.	
5	coronopifolium. W.en. (wh.) Buckshorn-leav'd.	—	— G. ḥ.	
6	grandiflorum. W.en. (wh.) great-flowered.	—	1815.	— G. ḥ.	
7	speciosum. W.en. (wh.) large-flowered.	—	—	— G. ḥ.	
8	Broussonetiæ. s.s. (wh.) Broussonet's.	—	1829.	7. 10. G. ḥ.	
9	pinnatifidum. L.en. (wh.) cut-leaved.	Madeira.	1778.	5. 8. G. ḥ.	
	<i>Chrysánthemum pinnatifidum. s.s.</i>				
10	ptarmicæfólium. w. (wh.) Sneezewort-leav'd. Caucasus.	1803.	7. 8. H. 4.		
11	serotinum. w. (wh.) creeping-rooted. N. America.	1731.	9. 10. H. 4.	Jacq. obs. 4. t. 90.	
12	uliginosum. w. (wh.) marsh.	Hungary.	1816.	7. 9. H. 4.	Bot. mag. t. 2706.
13	alpinum. w. (wh.) Alpine.	Switzerland.	1759.	7. 8. H. 4.	
14	saxatíle. b.c.	(wh.) rock.	1816.	— H. 4.	Lodd. bot. cab. t. 126.
15	Barrelieri. Lk.	(wh.) Barrelier's.	Alps Europe.	
16	Halléri. W.en.	(wh.) Haller's.	Switzerland.	1820.	— H. 4.

17 ceratophylloides. w. (wh.)	Stags-horn-ld. Piedmont.	1808.	6. 7. H. 4.	All. ped. t. 37. f. 1.	
<i>Chrysanthemum ceratophylloides</i> . All.					
18 simplicifolium. w. (wh.)	simple-leaved. S. America.	1825.	8. 10. H. ⊖.		
19 maximum. DC.	(wh.) broad-leaved. Pyrenees.	1816.	5. 8. H. 4.		
<i>latifolium</i> . w.					
20 Balsamita. w. (wh.)	various-leaved. Levant.	1779.	— H. 4.	Jacq. obs. 4. t. 89.	
21 oppositifolium. s.s. (wh.)	opposite-leaved. Portugal.	1828.	7. 9. H. 4.		
22 tomentosum. s.s. (wh.)	woolly. Corsica.	1829.	— H. 4.		
23 palustre. w. (wh.)	marshy. Armenia.	1824.	7. 10. H. ⊖.		
24 macrophyllum. w. (wh.)	large-leaved. Hungary.	1803.	— H. 4.	W. et K. hung. 1. t. 94.	
25 corymbosum. w. (wh.)	mountain. Germany.	1596.	6. 8. H. 4.	Jacq. aust. 4. t. 379.	
26 pulverulentum. W. en. (wh.)	powdered. Iberia.	1806.	— H. 4.		
27 cinerarifolium. s.s. (wh.)	Ragwort-leav'd. Dalmatia.	1824.	— H. 4.		
28 Parthenium. w. (wh.)	common. Britain.	6. 9. H. 4.	Eng. bot. v. 18. t. 1231.	
β flôre pleno. (wh.)	double-flowered.	— H. 4.		
29 parthenifolium. w. (wh.)	Parthenium-leav'd. Caucasus.	1804.	6. 7. H. 4.	Vent. cels. t. 43.	
30 caucasicum. w. (wh.)	Caucasian.	—	7. 8. H. 4.		
31 tenuifolium. T.N. (wh.)	slender-leaved. Naples.	1806.	— H. 4.		
32 sericeum. M.B.	(wh.) silky. Iberia.	1824.	— H. 4.		
33 inodorum. w. (wh.)	scentless. Britain.	8. 9. H. ⊖.	Eng. bot. v. 10. t. 676.	
34 maritimum. w. (wh.)	sea.	6. 10. H. 4.	— v. 14. t. 979.	
35 parviflorum. w. (wh.)	small-flowered. Caucasus.	—	6. 7. H. ⊖.		
36 præcox. M.B.	(wh.) early-flowered.	1827.	5. 7. H. 4.		
37 incanum. Led.	(wh.) hoary. Siberia.	—	7. 8. H. 4.		
38 glabrum. LG.	(wh.) smooth. Spain.	1825.	— H. 4.		
39 leptophyllum. s.s. (wh.)	narrow-leaved. Caucasus.	1829.	— H. 4.		
40 diversifolium. H.E.F. (wh.)	different-leaved. N. Holland.	1824.	5. 7. G. $\frac{1}{2}$.	Hook. ex. flor. t. 215.	
41 carneum. M.B.	(pi.) flesh-coloured. Caucasus.	1804.	8. 9. H. 4.	Bot. mag. t. 1080.	
<i>Chrysanthemum coccineum</i> . B.M.					
42 roseum. M.B.	(ro.) rose-coloured.	—	1818.	— H. 4.	
<i>Chrysanthemum coccineum</i> . w.					
43 achillæfolium. M.B. (ye.)	Milfoil-like.	Cumana.	1824.	6. 7. H. ⊖.	
44 millefoliatum. w. (ye.)	Milfoil-leaved.	Siberia.	1731.	5. 9. H. 4.	Mill. ic. I. t. 9.
45 bipinnatum. w. (ye.)	wing-leaved.	—	1796.	6. 7. H. 4.	Gmel. sib. 2. t. 85. f. 1.
46 indicum. H.K.	(ye.) Indian.	E. Indies.	1810.	6. 9. H. ⊖.	Bot. mag. t. 1521.
ERIOPHYLLUM. LG.	ERIOPHYLLUM.	Syngenesia	Polygamia	Superflua.	
1 lanatum. (ye.)	woolly.	N. America.	1827.	6. 8. H. 4.	Bot. reg. t. 1167.
<i>cæspitosum</i> . B.R.	<i>Actinella lanata</i> . Ph.	<i>Trichophyllum lanatum</i> . n.	<i>Helennium lanatum</i> .	s.s.	
2 oppositifolium. N. (ye.)	opposite-leaved.	N. America.	1826.	— H. 4.	
3 trolliifolium. LG. (ye.)	Trollius-leaved.	New Spain.	1828.	8. 10. F. 4.	

ORDO CXXI.

LOBELIACEÆ. Juss. ann. mus. 18. p. 1.

LOB'ELIA. R.S.	LOB'ELIA.	Pentandria	Monogynia.	R.S.
1 linearis. R.S.	(bl.) linear-leaved.	C. B. S.	1791.	5. 8. G. $\frac{1}{2}$.
2 setacea. R.S.	(bl.) bristle-leaved.	—	1816.	— G. 4.
3 commutata. R.S.	(bl.) notch-leaved.	—	—	Act. Gorenk. 1811. f. 11.
4 simplex. R.S.	(bl.) simple-stalked.	—	1794.	— G. $\frac{1}{2}$.
5 unidentata. R.S. (d.pu.)	single-toothed.	—	—	Bot. mag. t. 1484.
6 pinifolia. R.S.	(bl.) Pine-leaved.	—	1752.	— G. 4.
7 Dortmanna. R.S.	(bl.) water.	Britain.	Andr. deposit. t. 240.
8 paludosæ. R.S.	(bl.) marsh.	N. America.	1823.	— H.w. 4.
9 Tùpa. R.S.	(sc.) Mullein-leav'd.	Peru.	1824.	Eng. bot. v. 2. t. 140.
10 salicifolia. H.S.L.	(or.) Willow-leaved.	Chili.	1794.	Swt. br. fl. gar. t. 284.
<i>Tùpa</i> . H.K. non L. <i>gigantea</i> . B.M. non Cav. <i>arguta</i> .	B.R. t. 973.			Bot. mag. t. 1325.
11 Kálmiæ. R.S.	(bl.) Kalm's.	N. America.	1820.	7. 9. H. $\frac{1}{2}$.
12 Nuttallæ. R.S.	(bl.) Nuttall's.	—	—	Bot. mag. t. 2238.
gracilis. N. nec aliorum.				
13 pauciflora. K.S.	(bl.) few-flowered.	Mexico.	1824.	F. 4.
14 chinensis. R.S.	(p.bl.) Chinese.	China.	1822.	H. ⊖.
15 bellidifolia. R.S.	(bl.) Daisy-leaved.	C. B. S.	1790.	5. 8. G. 4.
16 rhizophyta. R.S. (wh.)	rooting branched.	—	1814.	G. 4.
				Bot. mag. t. 2519.

17 decumbens. B.M. (<i>bl.</i>) decumbent.	C. B. S.	1820.	5. 8. G. 4.	Bot. mag t. 2277.
18 triquètra. R.S. (<i>bl.</i>) triangular.	_____	1774.	— G. 4.	
19 cinérea. R.S. (<i>wh.</i>) gray.	_____	1824.	6. 10. H. ⊖.	
20 fenestrális. R.S. (<i>bl.</i>) windowed.	Mexico.	1826.	9. 11. H. ⊖.	Cavan. icon. 6.t. 512.f.1.
21 gigántea. R.S. (<i>or.</i>) gigantic.	S. America.	1828.	8. 11. S. ™.	_____ 6. t. 513.
22 longiflòra. R.S. (<i>wh.</i>) long-flowered.	Jamaica.	1752.	5. 8. S. 4.	Bot. mag. t. 2563.
23 tomentósa. R.S. (<i>bl.</i>) tomentose.	C. B. S.	1816.	— G. 4.	
24 secúnda. R.S. (<i>wh.</i>) side-flowering.	_____	1790.	— G. 4.	
25 trialáta. D.P. (<i>bl.</i>) three-winged.	Nepaul.	1820.	— G. 4.	Hook. ex. flor. t. 44.
micránta. H.E.F. non K.S.				
26 alàta. B.P. (<i>bl.</i>) winged-stalked.	N. S. W.	1804.	— G. 4.	
27 acuminàta. R.S. (<i>st.</i>) taper-pointed.	Jamaica.	1822.	6. 9. S. ™.	Lab. nov. hol. 1. t. 72.
28 nicotianæfólia. R.S. (<i>st.</i>) Tobacco-leaved.	E. Indies.	_____	— S. 4.	Sloan. his. 1. t. 95. f. 2.
29 circiifólia. R.S. (<i>st.</i>) Cirsium-leaved.	W. Indies.	1825.	8. 10. S. 4.	Plum. sp. v. 4. t. 116.
30 persicifólia. R.S. (<i>pu.</i>) Peach-leaved.	_____	1818.	6. 9. S. 4.	
31 Cavanillesiàna. R.S. (<i>re.</i>) Cavanilles'.	New Spain.	1825.	— S. 4.	Cavan. ic. 6. t. 518.
persicifólia. Cav. non Lam.				
32 Claytoniàna. R.S. (<i>p.bl.</i>) Clayton's.	N. America.	1799.	6. 8. H. 4.	Willd. hort. ber. t. 30.
goodenioïdes. w. pallida. Roth.				
33 assúrgens. R.S. (<i>pu.</i>) purple.	W. Indies.	1787.	6. 10. S. ™.	Andr. reposit. t. 553.
34 pyramidális. D.P. (<i>or.</i>) pyramidal.	Nepaul.	1819.	7. 10. G. 4.	Bot. mag. t. 2387.
35 fulgén. R.S. (<i>cr.</i>) fulgent.	Mexico.	1809.	5. 9. F. 4.	Bot. reg. t. 165.
36 spléndens. R.S. (<i>sc.</i>) splendid.	_____	1814.	— F. 4.	_____ t. 60.
37 cardinalís. R.S. (<i>sc.</i>) Cardinal flower.	Virginia.	1629.	— F. 4.	Bot. mag. t. 320.
38 syphilítica. R.S. (<i>bl.</i>) blue cardinal.	_____	1665.	8. 10. H. 4.	Bot. reg. t. 537.
39 débilis. R.S. (<i>bl.</i>) feeble.	C. B. S.	1774.	4. 8. H. ⊖.	
40 grácilis. B.P. (<i>bl.</i>) slender.	N. S. W.	1801.	7. 10. G. ⊖.	Bot. mag. t. 741.
41 purpurásca. B.P. (<i>pu.</i>) purplish.	_____	1809.	6. 8. G. 4.	
42 surinamensis. R.S. (<i>bl.</i>) Surinam.	W. Indies.	1786.	1. 7. S. ™.	Bot. mag. t. 225.
β rúbra. (<i>re.</i>) red-flowered.	Brazil.	1820.	12. 4. S. ™.	Lodd. bot. cab. t. 749.
43 Cliffortiàna. s.s. (<i>pu.</i>) purple-flowered.	S. America.	1733.	7. 8. H. ⊖.	Lin. h. clif. 426. t. 26.
44 Michauxii. n. (<i>pu.</i>) Michaux's.	N. America.	— H. ⊖.	
Cliffortiàna. M. non L.				
45 infláta. R.S. (<i>bl.</i>) bladder-podded.	_____	1759.	— H. ⊖.	Swt. br. fl. gar. t. 99.
46 ürens. R.S. (<i>bl.</i>) acrid.	England.	6. 7. H. 4.	Eng. bot. v. 14. t. 953.
47 pubérula. R.S. (<i>bl.</i>) downy.	N. America.	1819.	— H. 4.	
48 am'ena. R.S. (<i>bl.</i>) bright blue.	_____	1812.	6. 8. H. 4.	Ann. mus. 18. t. 1. n. 1.
49 minúta. R.S. (<i>bh.</i>) small.	C. B. S.	1772.	6. 9. G. 4.	Bot. mag. t. 2590.
50 mímina. B.M. (<i>wh.</i>) least.	_____	6. 10. G. 4.	_____ t. 2077.
51 Lauréntia. R.S. (<i>bl.</i>) Italian.	Italy.	1778.	7. 8. H. ⊖.	Mich. gen. 18. t. 14.
52 tenélla. R.S. (<i>vi.</i>) slender-leaved.	Sicily.	1820.	6. 8. H. ⊖.	
53 campanuloïdes. R.S. (<i>bl.</i>) campanulate.	Japan.	1819.	7. 9. G. 4.	Bot. reg. t. 733.
54 Erínus. R.S. (<i>bl.</i>) ascending.	C. B. S.	1752.	6. 9. H. 4.	Bot. mag. t. 901.
55 Bre'ni. R.S. (<i>bl.</i>) Breynius's.	_____	1823.	— G. 4.	Breyn. cent. t. 89.
56 erinóides. R.S. (<i>bl.</i>) trailing.	_____	1759.	6. 8. G. 3.	Herm. lugd. t. 109.
57 bicolor. R.S. (<i>bl. wh.</i>) spotted.	_____	1795.	— H. ⊖.	Bot. mag. t. 514.
58 Schránkii. (<i>bl.</i>) Schrank's.	_____	1824.	— G. 4.	Schránk. h. mon. t. 31.
secúnda. Sck. non Thunb. decumbens. L. enum. non B.M.				
59 corymbosa. B.M. (<i>wh. pu.</i>) corymbose.	C. B. S.	1825.	— G. 4.	Bot. mag. t. 2693.
60 ilicifólia. B.M. (<i>wh. pu.</i>) Holly-leaved.	_____	1815.	5. 9. G. 4.	_____ t. 1896.
61 ánceps. R.S. (<i>bl.</i>) flat-edged.	_____	1824.	7. 10. H. ⊖.	
62 pubéscentes. R.S. (<i>wh.</i>) downy-leaved.	_____	1780.	6. 8. G. 4.	Jacq. schœn. 2. t. 178.
63 hirsuta. R.S. (<i>bl.</i>) blue-flower'd hairy.	_____	1759.	5. 9. G. 4.	Burn. afr. t. 40. f. 2.
64 zeylánica. R.S. (<i>pu.</i>) Ceylon.	E. Indies.	1820.	— S. 4.	
65 lutea. R.S. (<i>ye.</i>) yellow.	C. B. S.	1774.	6. 7. G. 4.	Bot. mag. t. 1319.
66 variifólia. B.M. (<i>ye.</i>) various-leaved.	_____	1812.	— G. 4.	_____ t. 1692.
67 coronopifólia. R.S. (<i>bl.</i>) Buckshorn-leaved.	_____	1752.	7. 8. G. 4.	_____ t. 644.
68 cœrulea. B.M. (<i>bl.</i>) blue-flowered.	_____	1823.	5. 8. G. ™.	_____ t. 2701.
69 Thunbergii. (<i>bl.</i>) upright-raceme-flower'd.	_____	1812.	7. 10. G. 4.	
coronopifólia. Th. non B.M.				
70 crenáta. R.S. (<i>bl.</i>) notched-leaved.	_____	1794.	4. 5. G. 4.	
71 Símsii. (<i>bl.</i>) long-peduncled.	_____	1819.	8. 11. G. 4.	Bot. mag. t. 2251.
pedunculáta. B.M. non B.P.				
72 verbasifólia. Sm. (<i>pu.</i>) Mullein-leaved.	Nepaul.	1822.	5. 7. G. 4.	
73 racemosa. B.M. (<i>gr.</i>) racemed.	W. Indies.	1818.	7. 9. S. ™.	Bot. mag. t. 2137.
74 críspa. Gr. (<i>bl.</i>) curled-leaved.	Mexico.	1825.	5. 6. F. 4.	
75 heteromälla. Jac. (<i>bl.</i>) woolly-leaved.	1829.	— G. 4.	
76 móllis. Gr. (<i>pu.</i>) soft.	Dominica.	1828.	5. 8. S. 4.	
77 rugulosa. Gr. (<i>wh.</i>) rugged.	New Zealand.	—	— G. 4.	

LOBELIACEÆ.

78 cóncolor. B.P.	(bl.) self-coloured.	N. S. W.	1828.	5. 6. G. ¼.	
79 inundàta. B.P.	(bl.) inundated.	_____	1822.	5. 8. G. w. ¼.	
80 pedunculàta. B.P.	(bl.) long-stalked.	_____	—	— G. ¼.	
81 dentàta. B.P.	(bl.) toothed.	_____	1819.	— G. ¼.	Cavan. ic. 6. t. 522.
82 gibbosa. B.P.	(bl.) gibbous.	_____	1824.	— G. ¼.	Lab. nov. hol. 1. t. 71.
83 senecioïdes. B.M.	(bl.) Groundsel-like.	_____	—	7. 10. G. ™.	Bot. mag. t. 2702.
<i>Isotoma axillaris.</i> Bot. reg. t. 964.					

MONO'PSIS. Sal.	MONO'PSIS.	Pentandria Monogynia.			
1 conspicuá. H.T.	(bl.) conspicuous.	C. B. S.	1812.	7. 8. H. ⊙.	Bot. mag. t. 1499.

Lobelia spiculum. B.M.

2 inconspicuá. H.T. (bl.) inconspicuous. — H. ⊙.

CLINTONIA. B.R.	CLINTONIA.	Pentandria Monogynia.			
élegans. B.R.	(bl.) elegant.	Columbia.	1827.	6. 9. H. ⊙.	Bot. reg. t. 1241.

CYPHIA. R.S.	CYPHIA.	Pentandria Monogynia. W.			
1 volùbilis. R.S.	(pi.) climbing.	C. B. S.	1795.	6. 8. S. ¼. —	
2 digitàta. R.S.	(li.) finger-leaved.	_____	1820.	7. 10. G. ¼.	
3 incisa. R.S.	(bh.) cut-leaved.	_____	—	— G. ¼.	
4 bulbòsa. R.S.	(pi.) bulb-bearing.	_____	1791.	8. 9. S. ¼.	Burm. afr. t. 38. f. 2.
5 Phyteùma. R.S.	(pi.) Phyteuma-leav'd.	_____	1820.	2. 5. S. ¼.	Bot. reg. t. 625.
6 Cardámines. R.S.	(bh.) Cardamine-leaved.	_____	—	— S. ¼.	

ORDO CXXII.

STYLI'DIUM. Brown prodr. 1. p. 565.

STYLI'DIUM. B.P.	STYLI'DIUM.	Gynandria Diandria.			
1 graminifòlium. B.P. (ro.) grass-leaved.	_____	N. S. W.	1803.	4. 8. G. ¼.	Bot. reg. t. 90.
2 linéare. B.P.	(li.) linear-leaved.	_____	1820.	— G. ¼.	Sm. exot. bot. 2. t. 67.
<i>Venentatia minor.</i> Ex. b.					
3 scàndens. B.P.	(wh.) climbing.	N. Holland.	1803.	7. 8. G. ¼. —	
4 fruticòsum. B.P.	(wh.) frutescent.	_____	—	5. 10. G. ™.	Sal. par. lond. t. 77.
<i>glandulosum.</i> P.L.					
5 laricifòlium. B.R. (ro.) Larch-leaved.	_____	N. S. W.	1818.	— G. ™.	Bot. reg. t. 550.
<i>tenuifòlium.</i> B.M. 2249.					
6 adnátum. B.P.	(bh.) adnate.	N. Holland.	1824.	— G. ™.	Bot. reg. t. 914.

ORDO CXXIII.

GOODENOVIÆ. Brown prodr. 1. p. 573.

GOOD'ENIA. B.P.	GOOD'ENIA.	Pentandria Monogynia. W.			
1 bellidifòlia. B.P.	(ye.) Daisy-leaved.	N. S. W.	1823.	6. 9. G. ¼.	
2 stellígera. B.P.	(ye.) starry-haired.	_____	1825.	— G. ¼.	
3 paniculàta. B.P.	(ye.) paniced.	_____	—	— G. ¼.	
4 grátilis. B.P.	(ye.) slender.	N. Holland.	1824.	5. 8. G. ¼.	
5 decurréns. B.P.	(ye.) decurrent.	N. S. W.	1825.	— G. ¼.	
6 ovàta. B.P.	(ye.) oval-leaved.	_____	1793.	6. 10. G. ™.	
7 grandifòra. B.P.	(ye.) large-flowered.	_____	1802.	6. 8. G. ™.	
8 hederacea. B.P.	(ye.) Ivy-leaved.	_____	1823.	— G. ¼.	
<i>E'UTHALES. B.P.</i>					
<i>E'UTHALES.</i> Pentandria Monogynia. R.S.					
trinérvis. B.P. (ye. pu.) three-nerved.	_____	N. Holland.	1803.	5. 9. G. ¼.	
<i>Goodénia tenélla.</i> A.R. 466. <i>Velleia trinérvis.</i> Lab. n. hol. t. 77.					

VELLEIA. B.P.	VELLEIA.	Pentandria Monogynia. R.S.			
1 paradóxa. B.P.	(ye.) paradoxical.	N. S. W.	1823.	6. 8. G. ¼.	
2 lyràta. B.P.	(ye.) lyrate-leaved.	_____	1819.	4. 6. G. ¼.	Bot. reg. t. 971.
3 spatulàta. B.P.	(ye.) spatulate.	_____	1823.	— G. ¼.	— t. 551.
<i>LECHENA'ULTIA. B.P.</i>					
<i>LECHENA'ULTIA.</i> Pentandria Monogynia. R.S.					
1 formòsa. B.P.	(sc.) handsome.	N. Holland.	1823.	6. 9. G. ™.	
2 oblàta. S.F.A.	(ye.) oblate-petaled.	_____	—	— G. ™.	Swt. flor. aust. t. 26.
<i>SCÆVOLA. B.P.</i>					
<i>SCÆVOLA.</i> Pentandria Monogynia. R.S.					
1 Koenigii. R.S.	(wh.) Konig's.	E. Indies.	1818.	8. 10. S. ™.	Bot. mag. t. 2732.

2 <i>Plumiéri. r.s.</i> (<i>wh.</i>) <i>Plumier's.</i>	<i>W. Indies.</i>	1724.	8. 10.	S. <i>h.</i>	Jacq. amer. t. 179. f. 88.
<i>Lobélia Plumiéri.</i> Jacq.					
3 <i>ivæfòlia. Lht.</i> (<i>wh.</i>) <i>Iva-leaved.</i>	<i>Trinidad.</i>	1820.	—	S. <i>h.</i>	
4 <i>Taccàda. F.I.</i> (<i>wh.</i>) <i>Indian.</i>	<i>E. Indies.</i>	1818.	—	S. <i>h.</i>	Gärt. fruct. 1. t. 25.
5 <i>crassifòlia. B.P.</i> (<i>bl.</i>) <i>thick-leaved.</i>	<i>N. Holland.</i>	1805.	8. 10.	G. <i>h.</i>	Lab. nov. hol. 1. t. 79.
6 <i>cuneifòrmis. B.P.</i> (<i>bl.</i>) <i>wedge-leaved.</i>	<i>V. Diem. Isl.</i>	1823.	—	G. <i>u.</i>	— 1. t. 80.
7 <i>microcárpa. B.P.</i> (<i>bl.</i>) <i>small-fruited.</i>	<i>N. S. W.</i>	1790.	5. 9.	G. <i>u.</i>	Cavan. ic. 6. t. 509.
<i>Goodénia lævigatæ. B.M.</i> 287.					
8 <i>suavèolens. B.P.</i> (<i>bl.</i>) <i>sweet-scented.</i>	—	1798.	—	G. <i>u.</i>	Andr. reposit. t. 22.
<i>Goodénia calendulacea. A.R.</i>					
9 <i>hispida. B.P.</i> (<i>li.</i>) <i>branching.</i>	<i>N. S. W.</i>	1820.	5. 8.	G. <i>u.</i>	Cavan. ic. 6. t. 510.
<i>Goodénia ramosissima. Sm.</i> new holl.	15. t. 5.				
DAMPI'ERA. B.P.	DAMP'I'ERA.	Pentandria Monogynia. R.S.			
1 <i>undulatæ. B.P.</i> (<i>bl.</i>) <i>wave-leaved.</i>	<i>N. S. W.</i>	1824.	5. 8.	G. <i>h.</i>	
2 <i>ovalifòlia. B.P.</i> (<i>pu.</i>) <i>oval leaved.</i>	—	1823.	5. 9.	G. <i>h.</i>	
3 <i>purpùrea. B.P.</i> (<i>pu.</i>) <i>purple-flowered.</i>	—	1825.	—	G. <i>h.</i>	
4 <i>stricta. B.P.</i> (<i>bl.</i>) <i>upright.</i>	<i>N. S. W.</i>	1814.	6. 8.	G. <i>u.</i>	

ORDO CXXIV.

CAMPANULACEÆ. *Juss. gen. 163.*

CANARINA. W.	CANARINA.	Hexandria Monogynia. W.			
<i>Campánula. w. (or.)</i> <i>Canary.</i>	<i>Canaries.</i>	1696.	1. 3.	G. <i>u.</i>	Bot. mag. t. 444.
MICHA'UXIA. W.	MICHA'UXIA.	Octandria Monogynia. W.			
1 <i>campanuloïdes. w. (li.)</i> <i>rough-leaved.</i>	<i>Levant.</i>	1787.	6. 8.	G. <i>♂.</i>	Bot. mag. t. 219.
2 <i>decándra. F.</i> <i>decandrous.</i>	<i>Persia.</i>	1829.	H. <i>♂?</i>	
LIGHTFO'OTIA. R.S.	LIGHTFO'OTIA.	Pentandria Monogynia. R.S.			
1 <i>oxycocoïdes. R.S. (pa.)</i> <i>lance-leaved.</i>	C. B. S.	1787.	7.	G. <i>h.</i>	
2 <i>tenélia. B.C.</i> (<i>pa.</i>) <i>slender.</i>	—	1824.	6. 8.	H. <i>h.</i>	Sm. exot. bot. 2. t. 69. Lodd bot. cab. t. 1038.
3 <i>lanceolatæ. L.en. (w.bl.)</i> <i>spear-leaved.</i>	—	1826.	—	G. <i>u.</i>	
4 <i>muscósa. L.en. (wh.bl.)</i> <i>moss-like.</i>	—	—	—	G. <i>u.</i>	
5 <i>subulatæ. R.S.</i> (<i>bl.</i>) <i>awl-leaved.</i>	—	—	—	G. <i>u.</i>	
ADENO'PHORA. F.	ADENO'PHORA.	Pentandria Monogynia.			
1 <i>verticillatæ. F.</i> (<i>bl.</i>) <i>whorled.</i>	<i>Dahuria.</i>	1783.	6. 8.	H. <i>u.</i>	Pall. it. 3. t. G. f. 1.
2 <i>marsupiiflòra. F.</i> (<i>bl.</i>) <i>bellied.</i>	—	1815.	7. 8.	H. <i>u.</i>	Bot. reg. t. 149.
<i>Campánula coronatæ. B.R.</i>					
3 <i>Gmelini. F.</i> (<i>bl.</i>) <i>Gmelin's.</i>	—	1823.	5. 8.	H. <i>u.</i>	Gmel. sib. 3. t. 33.
4 <i>coronopifòlia. F.</i> (<i>bl.</i>) <i>buckshorn-leav'd.</i>	—	1822.	—	H. <i>u.</i>	Swt. br. fl. gar. t. 104.
5 <i>denticulatæ. F.</i> (<i>bl.</i>) <i>toothed.</i>	—	—	5. 7.	H. <i>u.</i>	— t. 116.
<i>Campánula tricuspidatæ. R.S.</i>					
6 <i>latifòlia. F.</i> (<i>bl.</i>) <i>broad-leaved.</i>	<i>Siberia.</i>	1827.	—	H. <i>u.</i>	
<i>Campánula pereskiaefòlia. R.S.</i>					
7 <i>Lamáckii. F.</i> (<i>bl.</i>) <i>Lamarck's.</i>	—	1826.	—	H. <i>u.</i>	Gmel. sib. 3. n. 18. t. 26.
8 <i>stylòsa. F.</i> (<i>bl.</i>) <i>long-styled.</i>	—	1820.	—	H. <i>u.</i>	— 3. n. 20. t. 27.
9 <i>reticulatæ. F.</i> (<i>bl.</i>) <i>netted.</i>	—	1829.	—	H. <i>u.</i>	
10 <i>communis. F.</i> (<i>p.bl.</i>) <i>common.</i>	—	1784.	5. 9.	H. <i>u.</i>	Jacq. schoenb. 3. t. 335.
<i>Campánula liliifòlia. B.R.</i>					
β <i>suavèolens. R.S. (p.bl.)</i> <i>sweet-scented.</i>	<i>Hungary.</i>	1818.	—	H. <i>u.</i>	Bot. reg. t. 236.
γ <i>hy'bridæ. R.S. (p.bl.)</i> <i>upright-flower'd.</i>	—	—	—	H. <i>u.</i>	
11 <i>intermédia. (p.bl.)</i> <i>intermediate.</i>	<i>Siberia.</i>	1820.	—	H. <i>u.</i>	
<i>Campánula intermédia. R.S.</i>					
WAHLENBERGIA. Sc. WAHLENBE'RGLIA.	Pentandria Monogynia.				
1 <i>grandifòra. Sc.</i> (<i>bl.</i>) <i>great-flowered.</i>	<i>Siberia.</i>	1782.	6. 8.	H. <i>u.</i>	Bot. mag. t. 252.
<i>Campánula grandifòra. B.M.</i>					
2 <i>péndula. Sc.</i> (<i>bl.</i>) <i>pendulous.</i>	<i>Canaries.</i>	1777.	7. 8.	H. <i>○.</i>	
<i>Campánula lobelioides. R.S.</i>					
3 <i>elongatæ. Sc.</i> (<i>p.bl.</i>) <i>elongated.</i>	1816.	—	H. <i>○.</i>	
CAMPÀNULA. R.S.	BELL-FLOWER.	Pentandria Monogynia. L.			
1 <i>cenisia. R.S.</i> (<i>bl.</i>) <i>ciliated.</i>	<i>Switzerland.</i>	1775.	6. 7.	H. <i>u.</i>	All. ped. 1. t. 6. f. 2.
2 <i>uniflòra. R.S.</i> (<i>bl.</i>) <i>single-flowered.</i>	<i>Lapland.</i>	1827.	—	H. <i>u.</i>	Linn. fl. lap. t. 9. f. 5.6.
3 <i>Kitaibeliana. R.S. (bl.)</i> <i>Kitaibel's.</i>	<i>Hungary.</i>	1828.	—	H. <i>u.</i>	
4 <i>Bellárdii. R.S.</i> (<i>bl.</i>) <i>Bellardi's.</i>	<i>Italy.</i>	1813.	—	H. <i>u.</i>	All. ped. t. 85. f. 5.
5 <i>Stevènii. M.B.</i> (<i>bl.</i>) <i>Steven's.</i>	<i>Siberia.</i>	1828.	—	H. <i>u.</i>	
6 <i>púlla. R.S.</i> (<i>d.pu.</i>) <i>dark-flowered.</i>	<i>Austria.</i>	1779.	—	H. <i>u.</i>	Bot. mag. t. 2492.

7	Zóysii. R.S.	(bl.) blunt-leaved.	Carniola.	1813.	6. 8.	H. ¼.	Jacq. ic. 2. t. 334.
8	diffusa. R.S.	(bl.) diffuse.	Italy.	1818.	—	H. ¼.	Cyr. pl. rar. n. t. 11.f. 2.
	β frágilis. L.en.	(bl.) brittle.	—	—	—	H. ¼.	Bot. mag. t. 117.
9	carpáthica. R.S.	(bl.) Carpathian.	Carp. Alps.	1774.	—	H. ¼.	Eng. bot. v. 13. t. 866.
10	dasycárpia. R.S.	(bl.) fleshy-podded.	Hungary.	1827.	—	H. ¼.	
11	rotundifolia. R.S.	(bl.) round-leaved.	Britain.	—	—	H. ¼.	Lodd. bot. cab. t. 561.
	α cærulea.	(bl.) blue-flowered.	—	—	—	H. ¼.	Scheuch. it. 4. f. 4.
	β albiflora.	(wh.) white-flowered.	—	—	—	H. ¼.	Bot. mag. t. 512.
12	excisa. R.S.	(bl.) cut-flowered.	Switzerland.	1816.	—	H. ¼.	Scop. carn. ed. 2. t. 4.
13	pusilla. R.S.	(bl. wh.) small.	Europe Alps.	—	—	H. ¼.	Bot. mag. t. 691.
14	pùmila. R.S.	(bl. wh.) dwarf.	Switzerland.	—	—	H. ¼.	Sm. exot. bot. t. 45.
	α cærulea.	(bl.) blue-flowered.	—	—	—	H. ¼.	
	β álba.	(wh.) white-flowered.	—	—	—	H. ¼.	
15	cæspitosa. R.S.	(bl.) tufted.	Carniola.	1818.	—	H. ¼.	Lab. nov. hol. 1. t. 70.
16	pubescens. R.S.	(bl.) pubescent.	Bohemia.	1813.	—	H. ¼.	Lodd. bot. cab. t. 1406.
17	gracilis. B.M.	(bl.) slender.	N. S. W.	1794.	4. 8.	G. ½.	W. et K. hung. t. 136.
18	erecta.	(bl.) upright.	—	—	—	G. ½.	All. ped. t. 47. f. 2.
	stricta. Sm. non L.						Vill. delph. 1. t. 10.
19	littoralis. Lab.	(bl.) sea-side.	V. Diem. Isl.	1820.	—	G. ½.	Lodd. bot. cab. t. 485.
20	capillaris. B.C.	(bl.) capillary.	N. Holland.	1824.	—	G. ½.	Eng. bot. v. 4. t. 283.
21	quadrifida. B.P.	(bl.) quadrifid.	N. S. W.	1822.	—	G. ½.	Flor. dan. t. 1087.
22	saxicola. B.P.	(bl.) small rock.	V. Diem. Isl.	1825.	5. 10.	G. ½.	Flor. græc. t. 205.
23	Waldsteiniàna. R.S. (vi.)	Waldstein's.	Hungary.	—	6. 8.	H. ¼.	Bot. mag. t. 397.
24	lanceolata. R.S.	(bl.) spear-leaved.	Pyrenees.	1827.	—	H. ¼.	Andr. deposit. t. 396.
25	divaricata. R.S.	(bl.) divaricate.	N. America.	—	—	H. ¼.	Jacq. schœn. 3. t. 336.
26	virgata. R.S.	(bl.) twiggy.	—	1823.	—	H. ¼.	Dodart. mem. 4. t. 113.
27	linifolia. R.S.	(bl.) Flax-leaved.	Sweden.	1816.	—	H. ¼.	Bot. reg. t. 57.
28	Scheuchzèri. s.s.	(bl.) Scheuchzer's.	Alps Europe.	1813.	—	H. ¼.	Duham. ed. nov. t. 41.
29	pátula. R.S.	(bl.) spreading.	Britain.	—	7. 8.	H. ½.	Jacq. schœn. 4. t. 472.
30	neglecta. R.S.	(bl.) neglected.	Europe.	1818.	—	H. ¼.	Barrel. ic. 567.
	crenata. L.en. ucránica. Hort.						
31	Rapunculus. R.S. (bl.)	Rampion.	Britain.	—	—	H. ¼.	Eng. bot. v. 5. t. 302.
32	persicifolia. R.S. (va.)	Peach-leaved.	Europe.	1596.	7. 9.	H. ¼.	Flor. dan. t. 85.
	α cærulea.	(bl.) blue-flowered.	Scotland.	—	—	H. ¼.	Bot. mag. t. 2553.
	β álba.	(wh.) white-flowered.	Europe.	—	—	H. ¼.	
	γ máxima. B.M.	(bl.) large-flowered.	—	—	—	H. ¼.	
	δ álbo-plena.	(wh.) double white.	—	—	—	H. ¼.	
	ϵ cæruleo-plena.	(bl.) double blue.	—	—	—	H. ¼.	
33	pyramidalis. R.S. (bl.)	pyramidal.	Carniola.	1596.	—	H. ¼.	
	α cærulea.	(bl.) common blue.	—	—	—	H. ¼.	
	β albiflora.	(wh.) white flowered.	—	1816.	—	H. ¼.	
34	versicolor. R.S. (bl.)	various-color'd.	Greece.	1788.	—	H. ¼.	
	planiflora. W.en. non Lam.	Willdenowiàna. R.S.	—	—	—		
35	obliqua. R.S.	(bl.) oblique-flower'd.	N. Europe.	1813.	6. 7.	H. ½.	
36	americana. R.S.	(bl.) American.	N. America.	1763.	7.	H. ½.	
37	nítida. H.K.	(bl. wh.) glossy.	—	1731.	7. 8.	H. ¼.	
	α cærulea.	(bl.) blue-flowered.	—	—	—	H. ¼.	
	β álba.	(wh.) white-flowered.	—	—	—	H. ¼.	
38	aurea. R.S.	(ye.) golden-flower'd.	Madeira.	1777.	7. 9.	G. ½.	
	α latifolia. R.S.	(ye.) broad-leaved.	—	—	—	G. ½.	
	β angustifolia. R.S. (ye.)	narrow-leaved.	—	—	—	G. ½.	
39	rhomboidea. R.S. (bl.)	Germander-l'd.	Switzerland.	1775.	7.	H. ¼.	
40	asteroides. R.S.	(bl.) star-flowered.	—	1824.	—	H. ¼.	
41	Alpini. R.S.	(bl.) Alpinus'.	Italy.	1823.	7. 9.	H. ¼.	
42	Ottoniàna. R.S.	(bl.) Otto's.	C. B. S.	1822.	5. 7.	G. ½.	
43	undulata. R.S.	(bl.) waved.	—	—	—	G. ¼.	
44	latifolia. R.S.	(bl.) giant.	Britain.	—	7.	H. ¼.	
	α cærulea.	(bl.) blue-flowered.	—	—	—	H. ¼.	
	β álba.	(wh.) white-flowered.	—	—	—	H. ¼.	
45	macrántha. L.en.	(bl.) great-flowered.	Russia.	1822.	6. 7.	H. ¼.	
46	eriocárpa. R.S.	(bl.) woolly-capsul'd.	Caucasus.	1824.	—	H. ¼.	
47	urticifolia. R.S.	(bl.) Nettle-leaved.	Germany.	1800.	8.	H. ¼.	
	β flòre pleno.	(wh.) double white.	—	—	—	H. ¼.	
48	Trachélium. R.S.	(bl.) throatwort.	Britain.	—	6. 8.	H. ¼.	
	α cærulea.	(bl.) blue-flowered.	—	—	—	H. ¼.	
	β álba.	(wh.) white-flowered.	—	—	—	H. ¼.	
	γ múltiplex.	(bl. wh.) double-flowered.	—	—	—	H. ¼.	
49	rapunculoides. R.S. (bl.)	creeping.	England.	—	6. 7.	H. ¼.	Eng. bot. v. 20. t. 1369.
	infundibulifòrmis.	Bot. mag. t. 2632.					

50 ucrâniæ. R.S. (bl.) Ukraine.	Ukraine.	1819.	6. 7. H. ¼.	
51 trachelioïdes. R.S. (bl.) throatwort-like.	Caucasus.	1818.	— H. ¼.	
52 bononiensis. R.S. (bl.) paniced.	Italy.	1773.	8. 9. H. ¼.	Moris. s. 5. t. 4. f. 38.
53 simplex. DC. (bl.) simple-stalked.	—	1823.	7. 8. H. ¼.	
54 lychnitis. H.H. (bl.) Lychnis-like.	—	—	— H. ¼.	
55 obliquifolia. R.S. (bl.) oblique-leaved.	Italy.	1820.	— H. ¼.	Ten. fl. neap. t. 17.
56 trichocalycina. R.S. (bl.) long-pointed-calyx'd.	—	1824.	— H. ¼.	— t. 16.
57 ruthénica. R.S. (bl.) Russian.	Russia.	1815.	6. 8. H. ¼.	Bot. mag. t. 2653.
58 graminifolia. R.S. (vi.) grass-leaved.	Hungary.	1823.	— H. ¼.	W. et K. hung. 2. t. 154.
59 tenuifolia. R.S. (vi.) slender-leaved.	—	1820.	5. 9. H. ¼.	— t. 155.
60 glomerata. R.S. (bl.) clustered.	Britain.	—	— H. ¼.	Eng. bot. v. 2. t. 90.
β albiflora. (wh.) white-flowered.	—	—	— H. ¼.	
61 elliptica. R.S. (bl.) elliptic-leaved.	Hungary.	1819.	— H. ¼.	Bocc. mus. 70. t. 58.
62 cephalantha. F. (p.bl.) pale-flowered.	Siberia.	1824.	— H. ¼.	
63 farinosa. Bes. (bl.) mealy.	Poland.	—	— H. ¼.	
64 aggregata. R.S. (bl.) crowded-flower'd.	Bavaria.	—	— H. ¼.	
65 speciosa. R.S. (bl.) specious.	Siberia.	—	— H. ¼.	Bot. mag. t. 2649.
glomerata β dahurica.	Bot. reg. 620.			
66 nicænsis. R.S. (bl.) Nice.	Italy.	1820.	— H. ¼.	All. ped. t. 39. f. 1.
67 foliosa. R.S. (bl.) leafy.	—	—	6. 9. H. ¼.	Ten. fl. neap. t. 18.
68 petræa. R.S. (bl.) woolly-leaved.	Mt. Baldo.	1818.	— H. ¼.	
69 Rainéri. R.S. (bl.) Rainer's.	Italy.	1825.	— H. ¼.	
70 Cervicaria. R.S. (bl.) wave-leaved.	Germany.	1768.	7. H. ♂.	Flor. dan. t. 787.
71 cervicarioïdes. R.S. (bl.) intermediate.	Italy.	1819.	— H. ¼.	
72 multiflora. R.S. (vi.) many-flowered.	Hungary.	1814.	6. 7. H. ♂.	W. et K. hung. t. 263.
macrostachya. W.en.				
73 collina. R.S. (bl.) Sage-leaved.	Caucasus.	1803.	7. 8. H. ¼.	Bot. mag. t. 927.
74 azurea. R.S. (bl.) azure.	Switzerland.	1778.	6. 7. H. ¼.	— t. 551.
75 Lörei. R.S. (bl.) Pollini's.	Mt. Baldo.	1819.	6. 9. H. ⓠ.	— t. 2581.
76 lactiflora. R.S. (st.) milk-coloured.	Caucasus.	1814.	7. 9. H. ¼.	Bot. reg. t. 241.
77 thyrsoidæa. R.S. (st.) long-spiked.	Switzerland.	1785.	6. 8. H. ♂.	Bot. mag. t. 1290.
78 peregrina. R.S. (bl.) rough-leaved.	C. B. S.	1794.	— G. ♂.	— t. 1257.
79 tomentosa. R.S. (wh.) tomentose.	Levant.	1820.	6. 9. H. ¼.	
80 aparinoïdes. R.S. (p.bl.) Aparine-like.	N. America.	1825.	7. 9. H. ⓠ.	
81 cernua. R.S. (wh.) nodding-flower'd.	C. B. S.	1804.	— G. ♂.	
82 capensis. R.S. (vi.) Cape.	—	1803.	— G. ⓠ.	Bot. mag. t. 782.
Roëlla decurrens. And. rep. t. 238.				
83 paniculata. R.S. (wh.) paniced.	C. B. S.	1822.	6. 8. G. ♂.	
84 spathulata. F.G. (bl.) spathulate.	Greece.	1824.	— G. ¼.	Flor. græc. t. 203.
85 velutina. R.S. (p.bl.) velvety.	Gibraltar.	—	— F. ¼.	Desf. atl. 1. t. 51.
86 microphylla. R.S. (p.bl.) small-leaved.	Spain.	1820.	— H. ¼.	
87 Allioni. R.S. (p.bl.) Allion's.	S. Europe.	1818.	— H. ¼.	All. ped. n. 418. t. 6. f. 3.
88 ligularis. R.S. (bl.) strap-leaved.	Alps.	1825.	— H. ¼.	Bot. mag. t. 1258.
89 barbata. R.S. (p.bl.) bearded.	Italy.	1752.	— H. ¼.	— t. 1723.
90 punctata. R.S. (wh.) dotted-flower'd.	Siberia.	1813.	5. 6. H. ¼.	Knor. delic. 1. t. 9.
91 Médiump. R.S. (bl.) Canterbury-bell.	Germany.	1597.	6. 9. H. ♂.	Garid. aix. p. 76. t. 18.
β albidæa. (wh.) white-flowered.	—	—	— H. ♂.	Lapeyr. pyren. t. 6.
92 longifolia. R.S. (bl.) long-leaved.	Pyrenees.	1824.	— H. ♂.	Flor. græc. t. 210.
93 betonicifolia. R.S. (bl.) Betony-leaved.	Olympus.	1823.	— H. ♂.	Swt. br. fl. gar. t. 280.
94 dichotoma. R.S. (bl.) forked.	Levant.	1827.	7. 10. H. ⓠ.	All. ped. 1. t. 46. f. 2.
95 spicata. R.S. (bl.) spiked.	Switzerland.	1786.	7. H. ♂.	Bot. mag. t. 957.
96 alpina. R.S. (pa.bl.) Alpine.	—	1779.	— H. ¼.	— t. 404.
97 mollis. R.S. (bl.) soft.	Syria.	1788.	5. 8. G. ¼.	Barrel. ic. t. 813.
98 saxatilis. R.S. (bl.) rock.	Candia.	1768.	— G. ¼.	Sal. par. lond. t. 26.
99 alliariæfolia. R.S. (wh.) Alliaria-leaved.	Caucasus.	1803.	8. 9. H. ¼.	
macrophylla. B.M. t. 912.				
100 sarmatica. B.R. (pa.bl.) gum-bearing.	Siberia.	—	6. 8. H. ¼.	Bot. reg. t. 237.
gummifera. R.S. betonicaefolia. M.B.				
101 lamüfolia. R.S. (st.) Lamium-leaved.	Caucasus.	1818.	— H. ¼.	Buxb. cent. 5. t. 18.
102 pendula. R.S. (st.) pendulous.	—	1824.	— H. ¼.	
103 sibirica. R.S. (bl.) Siberian.	Siberia.	1783.	7. 9. H. ¼.	Bot. mag. t. 659.
104 divergens. R.S. (bl.) divergent.	Hungary.	1814.	6. 7. H. ♂.	W. et K. hung. t. 258.
spathulata. W.K.				
105 violæfolia. R.S. (wh.) Violet-leaved.	Siberia.	1827.	6. 8. H. ¼.	
106 A'dami. R.S. (bl.) Adam's.	Caucasus.	1829.	— H. ¼.	
107 Pallasiæna. R.S. (bl.) Pallas's.	Siberia.	1827.	— H. ¼.	
108 lingulata. R.S. (vi.) tongue-leaved.	Hungary.	1804.	7. 8. H. ¼.	W. et K. hung. 1. t. 64.
109 caucasicæa. R.S. (bl.) Caucasian.	Caucasus.	—	— H. ¼.	
110 laciniata. R.S. (bl.) jagged-leaved.	Greece.	1788.	5. 8. G. ¼.	Andr. deposit. t. 385.
		Y 4		

CAMPANULACEÆ.

111	lyrata. R.S.	(bl.) lyre-leaved.	Levant.	6. 8.	H. ¼.	Moris. 2. v. 5. t. 3. f. 51.
112	rupéstris. R.S.	(bl.) various-leaved.	1827.	—	H. ½.		Flor. græc. t. 213.
113	strícta. R.S.	(bl.) straight.	Egypt.	—	—	H. ¼.	Tour. an. mu. p. 137. t. 13.
114	cichorácea. R.S.	(bl.) Succory-leaved.	Greece.	1824.	—	F. ½.	Flor. græc. t. 209.
115	capitata. R.S.	(bl.) headed.	—	1768.	6. 7.	H. ¼.	Bot. mag. t. 811.
116	lanuginósa. R.S.	(bl.) woolly.	—	1824.	—	H. ½.	
117	Erinus. R.S. (pa. bl.)	forked.	S. Europe.	1768.	7. 8.	H. ¼.	Moris. s. 5. t. 3. f. 25.
118	drabifolia. R.S. (bl.)	Draba-leaved.	Athens.	1823.	—	H. ¼.	Flor. græc. t. 215.
119	erinoïdes. R.S. (bl.)	Erinus-like.	Africa.	—	—	H. ¼.	Herm. lugd. t. 111?
120	Elátines. R.S. (pa. pu.)	trailing-stalked.	S. Europe.	—	7. 9.	H. ¼.	All. ped. n. 422. t. 7. f. 1.
121	hederácea. R.S. (pa. bl.)	Ivy-leaved.	England.	5. 6.	H. ¼.	Eng. bot. v. 2. t. 73.
122	arvática. R.S. (bl.)	Spanish.	Spain.	1825.	—	H. ¼.	Pluk. phyt. t. 23. f. 1.
123	hispidula. R.S. (bl. wh.)	hispid-calyxed.	C. B. S.	1816.	7. 9.	H. ¼.	Comm. hort. 2. t. 37.
PRISMATOCAR'PUS. L'H. VENUS-LOOKING-GLASS. Pentandria Monogynia.							
1	fruticosus. L'H.	(bl.) shrubby.	C. B. S.	1787.	8.	G. ½.	
2	Hookeri.	(wh.) Hooker's.	—	1823.	—	G. ½.	Bot. mag. t. 2733.
<i>Campánula Prismatocárpus. B.M. nec L'Her.</i>							
3	nítidus. L'H.	(bl.) shining.	—	—	6. 8.	H. ¼.	L'Her. sert. ang. 2. t. 3.
<i>Campánula Prismatocárpus. w.</i>							
4	hispidula. R.S. (bl. wh.)	bristly.	—	1822.	—	G. ¼.	Comm. hort. 2. t. 37.
5	interruptus. L'H. (bl.)	interrupted.	—	1820.	—	G. ¼.	
6	Spéculum. L'H. (bl.)	common.	S. Europe.	1596.	5. 8.	H. ¼.	Bot. mag. t. 102.
	β álbis.	(wh.) white-flowered.	—	—	—	H. ¼.	
7	hirsútus. T.N.	(bl.) hairy.	Italy.	1822.	—	H. ¼.	Ten. fl. neap. t. 19.
8	falcátus. T.N.	(bl.) falcate.	—	—	—	H. ¼.	t. 20.
9	hybridus. L'H.	(bl.) corn.	England.	—	H. ¼.	Eng. bot. v. 6. t. 375.
10	pentagonius. L'H. (bl.)	five-angled.	Turkey.	1686.	—	H. ¼.	Bot. reg. t. 56.
11	perfoliatus. (bl.)	perfoliate.	N. America.	1680.	—	H. ¼.	Moris. s. 5. t. 2. f. 23.
<i>Campánula perfoliata. R.S. amplexicaulis. M.</i>							
ROE'LLA. R.S. ROE'LLA. Pentandria Monogynia. L.							
1	ciliata. R.S. (bl. wh.)	ciliate.	C. B. S.	1774.	6. 9.	G. ½.	Bot. mag. t. 378.
2	pedunculata. R.S. (bl.)	peduncled.	—	1827.	—	G. ½.	
3	filiformis. R.S. (bl.)	filiform.	—	1816.	—	G. ½.	Lam. ill. t. 123. f. 2.
4	decurrens. R.S. (bl.)	decurrent.	—	1787.	7. 9.	H. ¼.	L'Her. sert. ang. 4. t. 6.
5	squarròsa. R.S. (bl.)	trailing.	—	—	7. 8.	G. ¼.	
6	muscòsa. R.S. (bl.)	moss-like.	—	1802.	8. 10.	H. ¼.	
7	spicata. R.S. (wh.)	spiked.	—	1824.	—	G. ½.	
PHYTÉ'UMA. R.S. PHYTÉ'UMA. Pentandria Monogynia. L.							
1	pauciflòrum. R.S. (bl.)	few-flowered.	Germany.	1823.	6. 8.	H. ¼.	Bot. mag. t. 1797.
2	globulariæfólium. R.S. (bl.)	Globularia-leav'd.	S. Europe.	—	—	H. ¼.	t. 2271.
3	Scheuchzéri. R.S. (bl.)	Scheuchzer's.	Switzerland.	1813.	5. 6.	H. ¼.	All. ped. n. 427. t. 7. f. 3.
4	scorzonérifólium. R.S. (bl.)	Scorzonera-leav'd.	Europe.	1817.	—	H. ¼.	Jacq. ic. 2. t. 333.
5	Micheli. R.S. (bl.)	Michel's.	Pyrenees.	1822.	6. 7.	H. ¼.	Jacq. aust. ap. t. 50.
6	hemiphæricum. R.S. (bl.)	linear-leaved.	Switzerland.	1752.	7.	H. ¼.	Eng. bot. v. 2. t. 142.
7	comosum. R.S. (da. bl.)	tufted.	Austria.	—	6. 7.	H. ¼.	Vill. delph. 2. t. 11. f. 3. A.
8	orbiculare. R.S. (bl.)	round-headed.	England.	6. 8.	H. ¼.	
9	Charmeli. R.S. (bl.)	long-bracted.	Pyrénées.	1819.	—	H. ¼.	
10	Siebèrii. s.s.	(bl.) Sieber's.	Levant.	1827.	7. 8.	H. ¼.	
11	inaequátum. R.S. (bl.)	unequal-leaved.	Hungary.	1826.	6. 7.	H. ¼.	
12	cordátum. B.M. (bl.)	heart-leaved.	S. Europe.	1804.	7. 8.	H. ¼.	Bot. mag. t. 1466.
13	ellipticum. R.S. (bl.)	elliptic-leaved.	Switzerland.	1820.	—	H. ¼.	Vill. delph. 2. t. 11. f. 2.
14	nigrum. R.S. (da. pu.)	black-flowered.	Bohemia.	—	—	H. ¼.	
15	betonicifólium. R.S. (pa.)	Betony-leaved.	Switzerland.	1816.	6. 8.	H. ¼.	Bot. mag. t. 2066.
16	spicatum. R.S. (st.)	spiked.	Europe.	1597.	—	H. ¼.	Flor. dan. t. 362.
17	Halléri. R.S. (pa. vi.)	Haller's.	Switzerland.	1819.	—	H. ¼.	
	β ovatum.	(pa.) oval-spiked.	—	1814.	—	H. ¼.	
18	stylósum. Bes.	(bl.) long-styled.	Poland.	1827.	—	H. ¼.	
19	strictum. B.M.	(bl.) upright.	S. Europe.	1819.	—	H. ¼.	
20	virgátum. R.S.	(bl.) slender-branched.	Levant.	1823.	7. 9.	H. ¼.	Bot. mag. t. 2145.
21	campanuloïdes. R.S. (bl.)	Campanula-like.	Caucasus.	1804.	6. 8.	H. ¼.	Lab. syr. 2. p. 11. t. 6.
22	canescens. R.S.	(li.) canescens.	Hungary.	—	—	H. ¼.	Bot. mag. t. 1015.
23	pinnátum. R.S.	(bl.) winged-leaved.	Candia.	1640.	—	G. ¼.	W. et K. hung. 1. t. 14.
TRACH'ELIUM. R.S. THROATWORT. Pentandria Monogynia. L.							
1	céruleum. R.S.	(bl.) blue.	Italy.	1640.	7. 9.	H. ½.	Venten. cels. t. 52.
2	diffusum. R.S.	(bl.) shrubby.	C. B. S.	1787.	—	G. ½.	
JAS'IONE. R.S. SHEEP'S SCABIOUS. Pentandria Monogynia. W.							
1	montána. R.S.	(bl.) mountain.	Britain.	6. 7.	H. ¼.	Eng. bot. v. 13. t. 82.
2	perennis. R.S.	(bl.) perennial.	France.	1787.	—	H. ¼.	Bot. reg. t. 505.
3	foliòsa. s.s.	(bl.) leafy.	S. Spain.	1826.	—	H. ¼.	

ORDO CXXV.

GESNEREÆ. Rich. et Juss. ann. mus. 5. p. 428.

GLOXI'NIA. S.S.	GLOXI'NIA.	Didynamia Angiospermia. L.				
1 <i>hirsuta</i> . B.R.	(<i>bl.</i>) hairy.	Brazil.	1825.	6. 8.	S. $\frac{1}{4}$.	Bot. reg. t. 1004.
2 <i>speciosa</i> . B.R.	(<i>bl.</i>) many-flowered.	1815.	6. 11.	S. $\frac{1}{4}$.	— t. 213.	
β <i>alba</i> .	(<i>wh.</i>) white-flowered.	1822.	—	S. $\frac{1}{4}$.		
3 <i>caulescens</i> . B.R.	(<i>vi.</i>) caulescent.	Pernambuca.	1820.	8. 10.	S. $\frac{1}{4}$.	Bot. reg. t. 1127.
4 <i>maculata</i> , s.s.	(<i>pu.</i>) spotted-stalked.	S. America.	1739.	7. 10.	S. $\frac{1}{4}$.	Bot. mag. t. 1191.
SINNI'NGIA. Nees.	SINNI'NGIA.	Didynamia Angiospermia.				
1 <i>guttata</i> . B.R.	(<i>gr.ye.</i>) spotted-throated.	Brazil.	1825.	6. 8.	S. $\frac{1}{2}$.	Bot. reg. t. 1134.
2 <i>Helléri</i> . B.R.	(<i>gr.ye.</i>) Heller's.	1824.	—	S. $\frac{1}{2}$.	— t. 997.	
3 <i>velutina</i> . B.R.	(<i>gr.ye.</i>) velvety.	—	—	S. $\frac{1}{2}$.	Lodd. bot. cab. t. 1398.	
4 <i>villosa</i> . B.R.	(<i>gr.ye.</i>) shaggy.	1826.	—	S. $\frac{1}{2}$.	Bot. reg. t. 1134.	
CODONOPHORA. B.R. CODONOPHORA.		Didynamia Angiospermia.				
1 <i>grandiflora</i> . B.R.	(<i>gr.</i>) large-flowered.	Brazil.	1816.	5. 6.	S. $\frac{1}{2}$.	Bot. reg. t. 428.
<i>Gesneria prasinata</i> . Bot. reg.						
2 <i>lanceolata</i> . B.R.	(<i>gr.</i>) spear-leaved.	—	1826.	—	S. $\frac{1}{2}$.	
GESNERIA. L.	GESNERIA.	Didynamia Angiospermia. L.				
1 <i>tomentosa</i> , s.s.	(<i>gr.pu.</i>) woolly.	S. America.	1752.	11. 6.	S. $\frac{1}{2}$.	Bot. mag. t. 1023.
2 <i>scabra</i> , s.s.	(<i>pu.</i>) rough.	Jamaica.	1822.	6. 8.	S. $\frac{1}{2}$.	
3 <i>corymbosa</i> , s.s.	(<i>ye.</i>) corymbose.	—	—	S. $\frac{1}{2}$.		
4 <i>Douglasii</i> . B.R.	(<i>sp.</i>) Douglas's.	Brazil.	1825.	5. 8.	S. $\frac{1}{2}$.	Bot. reg. t. 1110.
<i>verticillata</i> . Bot. mag. t. 2776. non Cavan.						
5 <i>aggregata</i> , s.s.	(<i>sc.</i>) clustered.	—	1816.	6. 10.	S. $\frac{1}{4}$.	Bot. reg. t. 329.
6 <i>pendulina</i> . B.R.	(<i>sc.</i>) drooping-flowered.	—	1825.	6. 9.	S. $\frac{1}{4}$.	— t. 1032.
7 <i>rutila</i> . B.R.	(<i>sc.</i>) brilliant.	—	1826.	6. 10.	S. $\frac{1}{4}$.	— t. 1158.
β <i>atrosanguinea</i> . B.R. (<i>cr.</i>) dark-crimson.	—	—	—	S. $\frac{1}{4}$.	— t. 1279.	
8 <i>latifolia</i> . M.N.	(<i>sc.</i>) long-spiked.	—	1825.	7. 10.	S. $\frac{1}{4}$.	— t. 1202.
<i>macrostachya</i> . B.R.						
9 <i>tubiflora</i> . C.I.	(<i>sc.</i>) tube-flowered.	S. America.	1815.	2. 6.	S. $\frac{1}{4}$.	Cavan. ic. 6. t. 584.
10 <i>bulbosa</i> , s.s.	(<i>sc.</i>) bulbous-rooted.	Brazil.	1816.	5. 8.	S. $\frac{1}{4}$.	Bot. reg. t. 343.
11 <i>acaulis</i> , s.s.	(<i>sc.</i>) stemless.	Jamaica.	1793.	—	S. $\frac{1}{4}$.	Sloan. jam. 1. t. 102. f. 1.
BESLERIA. L.	BESLERIA.	Didynamia Angiospermia. L.				
1 <i>cristata</i> , s.s.	(<i>ye.</i>) crested.	W. Indies.	1739.	6. 8.	S. $\frac{1}{2}$.	Jacq. amer. t. 119.
2 <i>coccinea</i> , s.s.	(<i>ye.</i>) scarlet-calyxed.	Guiana.	1824.	5. 8.	S. $\frac{1}{2}$.	Aub. gui. 2. t. 255.
3 <i>dichrus</i> , s.s.	(<i>ye.re.</i>) two-coloured.	Brazil.	1823.	—	S. $\frac{1}{2}$.	
4 <i>pulchella</i> , s.s.	(<i>or.</i>) large-flowered.	W. Indies.	1806.	—	S. $\frac{1}{2}$.	Bot. mag. t. 1146.
5 <i>melittifolia</i> , s.s.	(<i>pu.</i>) Balm-leaved.	Guiana.	1739.	6. 7.	S. $\frac{1}{2}$.	Sm. exot. bot. 1. t. 54.
6 <i>serrulata</i> , s.s.	(<i>st.</i>) saw-leaved.	W. Indies.	1806.	—	S. $\frac{1}{2}$.	Jacq. schœn. 3. t. 290.
7 <i>lutea</i> , s.s.	(<i>ye.</i>) yellow-flowered.	Guiana.	1739.	7. 8.	S. $\frac{1}{2}$.	Plum. ic. 49. f. 1.
8 <i>grandifolia</i> , s.s.	(<i>ye.</i>) large-leaved.	Brazil.	1823.	—	S. $\frac{1}{2}$.	
9 <i>hirtella</i> , s.s.	(<i>ye.</i>) hairy-leaved.	—	—	—	S. $\frac{1}{2}$.	
10 <i>incarnata</i> , s.s.	(<i>fl.</i>) flesh-coloured.	Guiana.	—	—	S. $\frac{1}{2}$.	Aub. gui. 2. t. 256.

ORDO CXXVI.

VACCINIEÆ. DC. theor. ed. 1. p. 216.

VACCI'NIUM. W.	WHORTLE-BERRY.	Oct-Decandria Monogynia.				
1 <i>Myrtillus</i> . w.	(<i>bh.</i>) Blea-berry.	Britain.	4. 6.	H. $\frac{1}{2}$.	Eng. bot. 7. t. 456.
2 <i>stamineum</i> . Ph.	(<i>wh.</i>) long-stamened.	N. America.	1772.	5. 6.	H. $\frac{1}{2}$.	Pluk. alm. t. 339. f. 3.
3 <i>álbum</i> . Ph.	(<i>wh.</i>) white-flowered.	—	—	—	H. $\frac{1}{2}$.	Andr. deposit. t. 263.
<i>stamineum</i> . A.R. non Ph.						
4 <i>pallidum</i> . w.	(<i>wh.</i>) pale.	—	—	—	H. $\frac{1}{2}$.	
5 <i>uliginosum</i> . w.	(<i>ro.</i>) great Bilberry.	Britain.	4. 5.	H. $\frac{1}{2}$.	Eng. bot. v. 9. t. 581.
6 <i>diffusum</i> . w.	(<i>bh.</i>) tree.	N. America.	1765.	5. 11.	H. $\frac{1}{2}$.	Bot. mag. t. 1607.
<i>arboreum</i> . M.						
7 <i>domosum</i> . B.M.	(<i>bh.</i>) bushy.	—	1774.	5. 6.	H. $\frac{1}{2}$.	Bot. mag. t. 1106.
<i>hirtellum</i> . H.K.						
β <i>hämile</i> . W.D.	(<i>wh.</i>) dwarf.	—	—	—	H. $\frac{1}{2}$.	Wats. dend. brit. t. 32.

VACCINIEÆ.

8 frondosum. w. (wh.) blunt-leaved.	N. America.	1761.	5. 6. H. h.	Andr. deposit. t. 140.
9 venustum. h.k. (wh.) red-twiggled.	1770.	— H. h.	Wang. am. t. 30. f. 69.	
10 resinosum. w. (st.) clammy.	1772.	— H. h.	Bot. mag. t. 1288.	
β rubescens. (re.) red-flowered.	—	— H. h.	Andr. deposit. t. 125.	
11 parviflorum. A.R. (re.) small-flowered.	1804.	— H. h.	Wats. dend. brit. t. 123.	
12 corymbosum. w. (wh.) corymbose.	1806.	6. 7. H. h.	Bot. reg. t. 400.	
13 amœnum. w. (wh.) broad-leaved.	1765.	5. 6. H. h.	Andr. deposit. t. 181.	
14 canadense. s.s. (wh.) Canadian.	1827.	5. 7. H. h.	Wats. dend. brit. t. 34.	
15 virgatum. w. (pa.re.) twiggy.	1767.	4. 5. H. h.		
16 Watsoni. (bh.) Watson's.	1812.	— H. h.		
virgatum β angustifolium. w.d.				
17 mariannum. w.d. (wh.) Maryland.	—	5. 6. H. h.		
18 grandiflorum. w.d. (wh.) great-flowered.	—	7. 8. H. h.		
19 elongatum. w.d. (bh.) elongated.	—	— H. h.		
20 minutiflorum. w.d. (bh.) minute-flowered.	—	— H. h.		
21 glabrum. w.d. (re.) smooth.	—	— H. h.		
22 mucronatum. L. (wh.) mucronate.	—	... 5. 6. H. h.		
23 fuscatum. w. (re.) cluster-flowered.	1770.	— H. h.	Bot. reg. t. 302.	
formosum. A.R. 97.				
β angustifolium. Ph. (re.) narrow-leaved.	—	— H. h.		
24 galézans. M. (st.) Gale-leaved.	1806.	— H. h.		
25 ligustrinum. w. (pu.re.) Privet-leaved.	—	— H. h.		
26 tenellum. w. (ro.) Pensylvanian.	1772.	— H. h.	Wats. dend. brit. t. 35.	
pensylvanicum. M.				
27 angustifolium. w. (wh.) narrow-leaved.	1776.	4. 5. H. h.		
myrtilloides. M.				
28 Arctostáphylos. w. (re.) oriental.	Levant.	1777.	6. 8. H. h.	Bot. mag. t. 974.
29 maderense. s.s. (re.) Madeira.	Madeira.	1810.	7. 2. H. h.	Andr. deposit. t. 30.
Arctostáphylos. A.R.				
30 meridionale. w. (re.) Jamaica.	Jamaica.	1778.	3. 6. G. h.	
31 salicinum. s.s. (bh.) Willow-like.	N. America.	1827.	6. 8. H. h.	
32 crassiſolum. s.s. (pi.) thick-leaved.	Carolina.	1787.	6. 7. F. h.	Bot. mag. t. 1152.
33 myrtifolium. M. (pi.) Myrtle-leaved.	—	1812.	5. 7. F. h.	
34 nítidum. s.s. (pi.) glossy-leaved.	—	1794.	5. 6. F. h.	Andr. rep. t. 480.
35 decumbens. B.M. (pi.) decumbent.	—	—	— F. h.	Bot. mag. t. 1550.
36 myrsinoides. M. (pu.) Myrsine-leaved.	—	... —	— F. h.	
37 buxifolium. s.s. (bh.) Box-leaved.	N. America.	1794.	— H. h.	Bot. mag. t. 928.
38 Vitis Idæa. w. (bh.) Cow-berry.	Britain.	—	4. 6. H. h.	Eng. bot. v. 9. t. 598.
α minor. (bh.) smaller.	—	—	— H. h.	Lodd. bot. cab. t. 1023.
β major. (ro.) larger.	N. America.	—	— H. h.	— t. 616.
39 ovatum. Ph. (bh.) oval-leaved.	—	1827.	5. 7. H. h.	
40 obtusum. Ph. (bh.) blunt-creeping.	—	—	— H. h.	
41 halleriæfolium. l.c. (wh.) Halleria-leaved.	—	—	5. 6. H. h.	
OXYCO'CCUS. P.S. CRANBERRY.	Octandria Monogynia.			
1 palustris. p.s. (pi.) common.	Britain.	—	5. 6. H. h.	Eng. bot. v. 5. t. 319.
Vaccinium Oxyccos. E.B. Schölleria Oxyccos. Roth.				
2 macrocarpus. Ph. (pi.) large-fruited.	N. America.	1760.	5. 8. H. h.	Wats. dend. brit. t. 122.
3 erectus. Ph. (ro.) upright.	—	1806.	6. 8. H. h.	— t. 31.
Vaccinium erythrocarpum. M.				
4 hispidulus. p.s. (pi.) Thyme-leaved.	—	1776.	5. 7. H. h.	Mich. amer. 1. t. 23.
Vaccinium hispidulum. w. Arbùtus thymifolia. Ait. Gaultheria serpyllifolia. Ph.				

ORDO CXXVII.

ERICÆ. Don Prodr. p. 148.

Sect. I. ERICEÆ VERÆ.

ARCTOSTA'PHYLOS. K.S. ARCTOSTA'PHYLOS. Decandria Monogynia.

1 Uva-úrsi. s.s. (bh.) Bear-berry.	Britain.	4. 6. H. h.	Eng. bot. v. 10. t. 714.
A'rbutus Uva-úrsi. E.B. Mairanía Uva-úrsi. Desv.				
2 alpina. s.s. (pi.) Alpine.	Scotland.	4. 5. H. h.	Eng. bot. v. 29. t. 2030.

A'R BUTUS. K.S.	STRAWBERRY-TREE.	Decandria Monogynia. L.						
1 Unèdo. s.s. (<i>wh.re.</i>) common.	Ireland.	9. 12.	H. $\frac{1}{2}$.			Eng. bot. v. 34. t. 2377.	
β rùbra. (<i>re.</i>) red-flowered.	—	H. $\frac{1}{2}$.			Lodd. bot. cab. t. 123.	
γ flòre plèno. (<i>wh.</i>) double-flowered.	—	H. $\frac{1}{2}$.				
δ schizopétala. (<i>wh.</i>) jagged-petaled.	—	H. $\frac{1}{2}$.				
ϵ integrifòlia. (<i>wh.</i>) entire-leaved.	—	H. $\frac{1}{2}$.				
ζ críspa. (<i>wh.</i>) curled-leaved.	—	H. $\frac{1}{2}$.				
η salicifòlia. (<i>wh.</i>) Willow-leaved.	—	H. $\frac{1}{2}$.				
2 canarién sis. s.s. (<i>ro.</i>) long-leaved.	Canaries.	1796.	5. 6.	G. $\frac{1}{2}$.			Bot. mag. t. 1577.	
3 hy'brida. s.s. (<i>wh.</i>) hybrid.	Hybrid.	—	H. $\frac{1}{2}$.			Bot. reg. t. 619.	
<i>serratifòlia.</i> B.C. 580. <i>andrachnoïdes.</i> L.en.								
4 Milléri.	Miller's.	1825.	H. $\frac{1}{2}$.				
5 Andrâchne. s.s. (<i>wh.</i>) oriental.	Levant.	1724.	3. 4.	F. $\frac{1}{2}$.			Bot. reg. t. 113.	
6 pròcerâ. D.	(<i>wh.</i>) tall.	N. America.	1827.	H. $\frac{1}{2}$.			
7 Menziësi. Ph.	(<i>wh.</i>) Mr. Menzies's.	—	—	H. $\frac{1}{2}$.			
8 tomentòsa. Ph.	(<i>wh.</i>) woolly.	—	—	H. $\frac{1}{2}$.			
9 laurifòlia. s.s.	(<i>wh.</i>) Laurel-leaved.	Mexico.	1825.	F. $\frac{1}{2}$.			
10 phillyreæfòlia. s.s. (<i>wh.</i>) Phillyrea-leaved.	Peru.	1812.	G. $\frac{1}{2}$.				
11 ? pùmila. s.s.	(<i>wh.</i>) dwarf.	Magellan.	1824.	G. $\frac{1}{2}$.			
GAULTHERIA. S.S. GAULTHERIA.	Decandria Monogynia. L.							
1 procùmbens. s.s. (<i>bh.</i>) trailing.	N. America.	1762.	7. 9.	H. $\frac{1}{2}$.			Andr. deposit. t. 116.	
2 frágantissima. s.s. (<i>wh.</i>) fragrant.	Nepaul.	1824.	H. $\frac{1}{2}$.			Wal. as. res. 13. p. 297. ic.	
3 Shallon. Ph.	(<i>bh.</i>) Shallon.	N. America.	1827.	5. 6.	H. $\frac{1}{2}$.		Lodd. bot. cab. t. 1732.	
ENKIA'NTHUS. B.M. ENKIA'NTHUS.	Decandria Monogynia.							
1 quinqueflòrus. B.R. (<i>pi.</i>) Canton.	China.	1812.	10. 2.	G. $\frac{1}{2}$.			Bot. reg. t. 884.	
2 reticulatòs. B.R. (<i>bh.</i>) netted-leaved.	1820.	—	G. $\frac{1}{2}$.			——— t. 885.	
ANDRO'MEDA. N. ANDRO'MEDA.	Decandria Monogynia. L.							
1 hypnoides. w.	(<i>wh.</i>) Moss-like.	Lapland.	1798.	6. 7.	H. $\frac{1}{2}$.		Bot. mag. t. 2936.	
2 polifòlia. w.	(<i>ro.</i>) wild rosemary.	Britain.	—	H. $\frac{1}{2}$.		Eng. bot. v. 10. t. 713.	
3 glaucophylla. Len. (<i>wh.</i>) glaucous-leaved.	N. America.	5. 9.	H. $\frac{1}{2}$.				
4 rosmarinifòlia. Ph. (<i>bh.</i>) Rosemary-leaved.	—	—	H. $\frac{1}{2}$.			Pall. ross. 2. t. 70. f. B.	
5 subulatâ. P.S.	(<i>ro.</i>) awl-leaved.	Europe.	—	H. $\frac{1}{2}$.			
6 angustifòlia. Ph. (<i>wh.</i>) narrow-leaved.	N. America.	2. 4.	H. $\frac{1}{2}$.				
<i>crispâ.</i> Link enum.								
7 calyculatâ. w.	(<i>wh.</i>) various-leaved.	Russia.	1748.	2. 4.	H. $\frac{1}{2}$.		Pall. ross. 2. t. 72. f. 1.	
α ventricosa. H.K. (<i>wh.</i>) globe-flowered.	—	—	H. $\frac{1}{2}$.			Bot. mag. t. 1286.	
β nàna. B.C.	(<i>wh.</i>) dwarf.	—	H. $\frac{1}{2}$.			Lodd. bot. cab. t. 862.	
γ latifòlia. H.K. (<i>wh.</i>) broad-leaved.	Newfoundland.	—	H. $\frac{1}{2}$.			——— t. 530.	
8 japónica. w.	(<i>wh.</i>) Japan.	Japan.	1806.	5. 6.	H. $\frac{1}{2}$.		Thunb. jap. 181. t. 22.	
9 ovalifòlia. D.P.	(<i>bh.</i>) oval-leaved.	Nepaul.	1823.	—	G. $\frac{1}{2}$.		Wall. as. res. 13. p. 39. ic.	
10 buxifòlia. P.S.	(<i>er.</i>) Box-leaved.	Mauritius.	1822.	4. 6.	G. $\frac{1}{2}$.		Bot. mag. t. 2660.	
11 fasciculatâ. w.	(<i>wh.</i>) bundle-flower'd.	Jamaica.	—	G. $\frac{1}{2}$.			
12 jamaicén sis. w.	(<i>wh.</i>) Jamaica.	1793.	—	G. $\frac{1}{2}$.			
13 mariâna. w.	(<i>bh.</i>) Maryland.	N. America.	1736.	6. 7.	H. $\frac{1}{2}$.		Pluk. mant. t. 448. f. 6.	
α ovâlis.	(<i>bh.</i>) oval-leaved.	—	—	H. $\frac{1}{2}$.		Bot. mag. t. 1579.	
β oblóngâ.	(<i>bh.</i>) oblong-leaved.	—	—	H. $\frac{1}{2}$.			
14 speciòsa. Ph.	(<i>wh.</i>) large-flowered.	Carolina.	1800.	6. 9.	H. $\frac{1}{2}$.		Bot. mag. t. 970.	
α nítida. Ph.	(<i>wh.</i>) smooth-leaved.	—	—	H. $\frac{1}{2}$.			
<i>cassinaefòlia.</i> B.M.								
β glauca. W.D.	(<i>wh.</i>) glaucous-leaved.	—	—	H. $\frac{1}{2}$.		Wats. dend. brit. t. 26.	
Andrómeda dealbátâ.	Bot. reg. t. 1010.							
γ pulvérulenta. Ph. (<i>wh.</i>) mealy-leaved.	—	—	H. $\frac{1}{2}$.			Bot. mag. t. 667.	
15 coriacea. w.	(<i>pi.</i>) thick-leaved.	N. America.	1765.	6. 8.	H. $\frac{1}{2}$.		——— t. 1095.	
α nítida. M. lúcida. Lam. mariâna. Jacq. ic. 3. t. 79. nec aliorum.								
16 acuminatâ. w.	(<i>wh.</i>) acute-leaved.	N. America.	—	8.	H. $\frac{1}{2}$.		Sm. exot. bot. 2. t. 89.	
17 Catesb'ei. w.	(<i>wh.</i>) spiny-leaved.	1793.	6. 7.	H. $\frac{1}{2}$.		Bot. mag. t. 1955.	
<i>spinulosa.</i> Ph.								
18 axillaris. w.	(<i>wh.</i>) axil-flowering.	1765.	5. 8.	H. $\frac{1}{2}$.		Duham. ar.ed. nov. t. 39.	
β longifòlia.	(<i>wh.</i>) long-leaved.	—	—	H. $\frac{1}{2}$.		Bot. mag. t. 2357.	
19 floribunda. Ph.	(<i>wh.</i>) many-flowered.	Georgia.	1812.	5. 6.	H. $\frac{1}{2}$.		——— t. 1566.	
20 racemosa. w.	(<i>wh.</i>) branching.	N. America.	1736.	7.	H. $\frac{1}{2}$.			
21 spicata. w.d.	(<i>wh.</i>) spiked.	—	6. 7.	H. $\frac{1}{2}$.		Wats. dend. brit. t. 36.	
22 arbórea. w.	(<i>wh.</i>) Sorrel-tree.	1752.	7. 9.	H. $\frac{1}{2}$.		Bot. mag. t. 905.	
LYONIA. N.	LYONIA.	Decandria Monogynia. N.						
1 ferruginea. N.	(<i>wh.</i>) rusty-leaved.	Georgia.	1784.	6. 7.	G. $\frac{1}{2}$.		Venten. malm. t. 80.	
<i>Andrómeda ferruginea.</i> w.								
2 rigida. N.	(<i>wh.</i>) rigid.	1774.	4. 5.	G. $\frac{1}{2}$.		Lodd. bot. cab. t. 430.	
3 salicifòlia. W.D.	(<i>wh.</i>) Willow-leaved.	N. America.	6. 7.	H. $\frac{1}{2}$.		Wats. dend. brit. t. 38.	
4 paniculatâ. N.	(<i>wh.</i>) panicle.	1748.	5. 6.	H. $\frac{1}{2}$.		——— t. 37.	

5 capreæfòlia. w.d. (wh.)	tendril-leaved.	N. America.	1812.	7.	H. h.	Wats. dend. brit. t. 127
6 multiflòra. w.d. (wh.)	many-flowered.	—	—	—	H. h.	— t. 128.
7 frondòsa. n.	(wh.) bristly-flowered.	—	1806.	5. 6.	H. h.	
MYLOCARYUM. W.en. BUCKWHEAT-TREE.	Decandria Monogynia. W.					
ligustrinum. Ph. (wh.)	Privet-leaved.	Georgia.	1786.	5. 6.	G. h.	
Cliftonia ligustrina. s.s.	—	—	—	—	G. h.	
CLE'THRA. W.	CLE'THRA.	Decandria Monogynia. W.				
1 alnifòlia. s.s.	(wh.) Alder-leaved.	N. America.	1731.	8. 10.	H. h.	Lam. ill. t. 369.
2 tomentòsa. s.s.	(wh.) woolly-leaved.	—	—	—	H. h.	Wats. dend. brit. t. 39.
3 scábra. s.s.	(wh.) rough-leaved.	Georgia.	1806.	—	H. h.	
4 acuminatà. s.s.	(wh.) acute-leaved.	Carolina.	—	—	H. h.	
5 paniculatà. s.s.	(wh.) paniced.	N. America.	1770.	—	H. h.	
6 arbórea. s.s.	(wh.) tree.	Madeira.	1784.	—	G. h.	Bot. mag. t. 1057.
β variegatà.	(wh.) variegated.	—	—	—	G. h.	
γ minor.	(wh.) dwarf.	—	—	—	G. h.	
7 ferruginea. L.en. (wh.)	ferruginous.	S. America.	1818.	—	G. h.	
CYRILLA. L.	CYRILLA.	Pentandria Monogynia. L.				
racemiflòra. L.	(wh.) Carolina.	Carolina.	1765.	6. 8.	F. h.	Bot. mag. t. 2456.
caroliniana. P.S.	Itea Cyrilla. w.	—	—	—	—	
BROSSÆA. L.	BROSSÆA.	Pentandria Monogynia. L.				
coccinea. L.	(sc.) scarlet.	S. America.	—	—	S. h.	Lam. ill. t. 111. f. 3.
BLÆRIA. W.	BLÆRIA.	Tetrandria Monogynia. W.				
1 ericoides. s.s.	(pu.) heath-leaved.	C. B. S.	1774.	8. 10.	G. h.	Wendl. coll. 1. t. 25.
2 muscosa. s.s.	(wh.) Moss-leaved.	—	—	6. 8.	G. h.	
3 fasciculatà. s.s.	(wh.) close-headed.	—	1812.	—	G. h.	
4 paucifolia. s.s.	(fl.) few-leaved.	—	—	—	G. h.	Wendl. coll. 2. t. 43.
5 articulatà. s.s.	(bh.) jointed.	—	1795.	5. 6.	G. h.	— 2. t. 44.
6 scábra. s.s.	(wh.) rough.	—	1806.	—	G. h.	— 1. t. 31.
7 depréssâ. R.S.	(pu.) depressed.	—	1816.	—	G. h.	
8 purpùrea. s.s.	(pu.) purple-flowered.	—	1791.	—	G. h.	
9 dumosa. s.s.	(ro.) bushy.	—	1806.	6. 8.	G. h.	Wendl. coll. 2. t. 38.
10 glabellâ. s.s.	(wh.) smooth.	—	1816.	—	G. h.	Seb. thes. 1. t. 20. f. 2.
11 ciliâris. s.s.	(wh.) fringed.	—	1795.	—	G. h.	Wendl. coll. 2. t. 49.
SYMPI'EAZA. R.S.	SYMPI'EAZA.	Tetrandria Monogynia. R.S.				
capitellatâ. R.S.	(pu.) headed.	C. B. S.	1812.	6. 8.	G. h.	Wendl. coll. 2. t. 37.
Blæria bracteata. Wend.	—	—	—	—	—	
CALL'UNA. Sal.	CALL'UNA.	Octandria Monogynia.				
vulgàris. W.en. (pi.) common.	Britain.	—	—	6. 8.	H. h.	Eng. bot. v. 15. t. 1013.
Erica vulgàris. E.B.	—	—	—	—	—	
α carnæa.	(pi.) flesh-coloured.	—	—	—	H. h.	
β álba.	(wh.) white-flowered.	—	—	—	H. h.	
γ multiplex.	(pi.) double-flowered.	—	—	—	H. h.	
ER'ICA. W.	HEATH.	Octandria Monogynia. L.				
1 absinthoides. H.K. (pu.)	Wormwood-like.	C. B. S.	1792.	3. 6.	G. h.	
2 actæa. L.en.	(wh.) cluster-leaved.	—	1810.	—	G. h.	Andr. heath. v. 3.
3 acuminatâ. A.H.	(pi.) pointed-leaved.	—	1798.	5. 6.	G. h.	
β pallida.	(pa.) pale-flowered.	—	—	—	G. h.	
4 acutâ. A.H.	(re.) pointed-cupped.	—	1799.	5. 8.	G. h.	Andr. heath. v. 2.
5 acutangulâris. L.C. (re.)	acute-angled.	—	1823.	—	G. h.	
6 adenophora. s.s. (wh.)	gland-haired.	—	1810.	—	G. h.	
7 aggregatâ. s.s.	(li.) clustered.	—	—	—	G. h.	Wendl. er. f. 13. n. 5.
8 Aitoniana. H.K. (wh. re.)	Aiton's.	—	1790.	6. 9.	G. h.	Bot. mag. t. 429.
9 álbens. H.K.	(wh.) pallid.	—	1789.	3. 8.	G. h.	— t. 440.
10 álbida. L.C.	(wh.) white-flowered.	—	1823.	5. 8.	G. h.	
11 alopecuroïdes. s.s. (li.)	fox-tail.	—	1812.	—	G. h.	Lodd. bot. cab. t. 874.
12 ambigua. s.s.	(pi.) ambiguous.	—	1798.	—	G. h.	Andr. heath. v. 1.
cylindrica. A.H. non Thunb.	hy'bida. Hort.	—	—	—	—	
13 am'œna. H.K.	(pu.) feathery.	C. B. S.	1795.	3. 7.	G. h.	Wend. eric. 17. p. 73. ic.
plumosa. A.H.	—	—	—	—	—	
14 ampullacea. H.K. (bh.)	flask.	—	1790.	6. 8.	G. h.	Bot. mag. t. 303.
15 andromedaflòra. H.K. (pi.)	Andromeda-fld.	—	1803.	3. 6.	G. h.	— t. 1250.
16 anthina. s.s.	(pu.) full-flowering.	—	1811.	—	G. h.	
flòrida. Hort. non W.	—	—	—	—	—	
17 aperta. s.s.	(re.) open-flowered.	—	—	—	G. h.	
18 Aphænes. s.s.	(wh.) Parsley-piert.	—	1820.	—	G. h.	
19 arborea. s.s.	(wh.) tree.	S. Europe.	1658.	2. 6.	G. h.	
β stylosa. P.S.	(wh.) long-styled.	—	—	—	G. h.	
20 arbúscula. B.C.	(re.) little-tree.	C. B. S.	1818.	4. 8.	G. h.	Lodd. bot. cab. t. 843.
21 arctata. L.C.	(wh.) arched.	—	—	—	G. h.	

22	Archeriana. H.K. (re.)	Lady Archer's.	C. B. S.	1796.	8. 11. G. h.	Andr. heath. v. 2.
23	árdens. H.K.	(sc.) ardent.	_____	1800.	4. 6. G. h.	Bot. reg. t. 115.
24	argentiflóra. s.s. (wh.)	silvery-flowered.	_____	1816.	— G. h.	Andr. heath. v. 4.
25	aristata. H.K.	(cr.) awned.	_____	1801.	3. 8. G. h.	Bot. mag. t. 1249.
26	assúrgens. L.en. (wh.)	assurgent.	_____	1810.	5. 8. G. h.	
27	aúrea. H.K.	(ye.) golden.	_____	1799.	7. 9. G. h.	Andr. heath. v. 2.
28	auriculáris. L.T.	(re.) eared.	_____	1810.	3. 7. G. h.	
29	austrális. H.K.	(re.) Spanish.	Spain.	1769.	4. 7. H. h.	Andr. heath. v. 3.
30	axilláris. w.	(pu.) axillary.	C. B. S.	1798.	5. 7. G. h.	
31	azaleæfólia. L.T.	(li.) Azalea-leaved.	_____	—	5. 8. G. h.	
32	báccans. H.K.	(li.) Arbutus-flower'd.	_____	1774.	4. 7. G. h.	Bot. mag. t. 358.
33	Bandoniana. L.C. (bh.)	Lord Bandon's.	_____	1816.	5. 8. G. h.	Andr. heath. v. 4.
34	Bánksii. H.K.	(st.) Banks's.	_____	1787.	2. 7. G. h.	— v. 1.
35	barbáta. H.K.	(wh.) bearded.	_____	1799.	5. 8. G. h.	— v. 2.
36	Beaumontia. A.H. (wh.)	Mrs. Beaumont's.	_____	1820.	6. 8. G. h.	— v. 4.
37	bélla. s.s.	(pu.) pretty.	_____	—	— G. h.	
38	Bergiána. H.K. (re.)	Bergius's.	_____	1787.	4. 8. G. h.	Lodd. bot. cab. t. 939.
	quadriflóra. A.H.	v. 2.				
39	bícolor. w. (re.gr.)	two-coloured.	_____	1790.	8. 3. G. h.	
	calathiflóra. L.T.					
40	biflóra. B.C.	(wh.) two-flowered.	_____	1820.	5. 8. G. h.	Lodd. bot. cab. t. 683.
41	bíflora. H.K.	(pi.) charming.	_____	1800.	4. 9. G. h.	Andr. heath. v. 3.
42	Blandfordiana. H.K. (ye.)	Blandford's.	_____	1803.	3. 6. G. h.	Bot. mag. t. 1793.
43	Bonplandiána. B.M. (wh.)	Bonpland's.	_____	1816.	— G. h.	— t. 2126.
44	borboniæfólia. L.T. (wh.)	Borbonia-leaved.	_____	—	— G. h.	
45	Bowieána. B.C. (wh.)	Bowie's.	_____	—	3. 9. G. h.	Lodd. bot. cab. t. 842.
	Bauèria. Andr. heath. v. 4.					
46	brachiális. L.T.	(re.) armed.	_____	1792.	— G. h.	
47	bracteáta. w.	(re.) red-bracted.	_____	1800.	5. 7. G. h.	
48	bracteoláris. P.S. (wh.)	bracteate.	_____	—	3. 7. G. h.	
49	brevifólia. L.T.	(wh.) short-leaved.	_____	—	8. 1. G. h.	
50	Broadleyána. A.H. (pi.)	Broadley's.	_____	1810.	5. 7. G. h.	Andr. heath. v. 3.
51	bruniána. w. (wh.)	Brunia-like.	_____	1790.	4. 7. G. h.	
52	bucciniflóra. B.M. (li.wh.)	trumpet-flowered.	_____	1816.	5. 9. G. h.	Bot. mag. t. 2465.
53	cæsia. L.T.	(wh.) grey.	_____	1795.	2. 5. G. h.	
54	cáffra. A.H.	(wh.) sweet-scented.	_____	1774.	8. 10. G. h.	Lodd. bot. cab. t. 196.
55	caledònica. s.s. (ro.)	Caledonian.	_____	1816.	5. 9. G. h.	
56	callósa. s.s. (li.)	callous-leaved.	_____	1799.	6. 9. G. h.	Andr. heath. v. 3.
	canaliculáta β minor. A.H.					
57	calycina. w. (pi.)	calycine.	_____	—	— G. h.	
58	calyculáta. P.S. (br.)	rusty-flowered.	_____	1800.	5. 7. G. h.	
59	campanuláta. H.K. (ye.)	bell-flowered.	_____	1791.	4. 8. G. h.	Andr. heath. v. 1.
60	campylophylla. s.s. (li.)	hook-leaved.	_____	1802.	3. 5. G. h.	
	incírvula. A.H.					
61	canaliculáta. H.K. (li.)	channelled.	_____	1799.	8. 2. G. h.	Andr. heath. v. 3.
62	canéscens. H.K.	(li.) canescent.	_____	1790.	5. 8. G. h.	— v. 2.
	eriocéphala. A.H. non Lam.					
63	cápax. L.T.	(re.wh.) capacious.	_____	1806.	— G. h.	
64	capitáta. H.K.	(st.) downy-headed.	_____	1774.	3. 7. G. h.	Andr. heath. v. 1.
65	carduiflóra. L.T.	(li.) Carduus-leaved.	_____	1806.	5. 7. G. h.	
66	carinátä. B.C.	(pi.) keeled.	_____	—	6. 9. G. h.	
67	cárnea. L.	(pi.) early dwarf.	Germany.	1763.	1. 8. G. h.	Lodd. bot. cab. t. 1071.
68	carniúla. B.C.	(fl.) flesh-coloured.	C. B. S.	1816.	5. 8. G. h.	Jacq. aust. t. 32.
69	catervæfólia. L.T.	(li.) huddled-leaved.	_____	1790.	4. 7. G. h.	Lodd. bot. cab. t. 926.
70	Celsiána. L.C.	(re.) Cels's.	_____	—	— G. h.	
71	cephalòtes. w.	(pu.) purple-headed.	_____	1812.	3. 7. G. h.	
72	cerinthoides. H.K. (sc.)	honeywort-flower'd.	_____	1774.	5. 9. G. h.	Bot. mag. t. 220.
	glabriúscula. (sc.)	smoothish.	_____	—	— G. h.	
	hispida. (sc.)	bristly.	_____	—	— G. h.	
73	cérnua. H.K.	(bh.) drooping-flowered.	_____	1791.	8. 12. G. h.	Mont.ac. ups.2. t.9. f.3.
74	cerviciflóra. L.T.	(ye.) Stag's-horn-flower'd.	_____	—	3. 7. G. h.	
	inapértä. Hort.					
75	chlamydiflóra. L.T.	(re.) cloaked-flower'd.	_____	1801.	5. 9. G. h.	
76	ciliáris. w.	(pi.) ciliated.	Portugal.	1759.	7. 9. H. h.	Bot. mag. t. 484.
77	ciliúlfólia. L.T.	(wh.) fringe-flower'd.	C. B. S.	—	5. 8. G. h.	
78	cineráscente. w.	(pu.) greyish.	_____	—	— G. h.	
79	cinérea. w.	(pu.) fine-leaved.	Britain.	—	6. 9. H. h.	Eng. bot. v. 15. t. 1015.
	atropurpurea. (d.pu.)	dark-purple.	_____	—	— H. h.	Lodd. bot. cab. t. 1409.
	rubra. (re.)	red-flowered.	_____	—	— H. h.	

γ cárnea.	(fl.) flesh-coloured.	Britain.	6. 9. H. h.	
δ álba.	(wh.) white-flowered.	C. B. S.	— H. h.	
80 cistifólia. L.en.	(wh.) Cistus-leaved.	C. B. S.	1799.	5. 8. G. h.	Andr. heath. v. 2.
barbáta β mīnor. A.H.					
81 clavæflóra. H.K.	(gr.) club-flowered.	—	1779.	8. 10. G. h.	Andr. heath. v. 2.
sessiliflóra. A.H. non L.					
82 claváta. s.s.	(gr.) clubbed.	—	1812.	— G. h.	Andr. heath. ic.
83 Clintóniæ. Lee.	(ro.) Lady Clinton's.	—	1816.	5. 8. G. h.	
84 coarctáta. s.s.	(li.) close.	—	—	— G. h.	
85 coccinea. H.K.	(sc.) scarlet-flowered.	—	1783.	1. 12. G. h.	
86 cólorans. B.R.	(wh.) changing-coloured.	—	1812.	1. 6. G. h.	
87 comósâ. H.K.	(re.wh.) tufted.	—	1787.	4. 8. G. h.	
a rúbra.	(re.) red-flowered.	—	—	— G. h.	
β álba.	(wh.) white-flowered.	—	—	— G. h.	
88 complanáta. s.s.	(re.) flattened.	—	—	— G. h.	
89 Comptoniâna. s.s.(re.)	Lady Northampton's.	—	1820.	— G. h.	
90 concáva. B.C.	(pu.) concave-flowered.	—	1808.	2. 5. G. h.	
91 concínna. H.K.	(bh.) blush.	—	1773.	7. 10. G. h.	
92 cóncolor. L.C.	(wh.) self-coloured.	—	—	3. 6. G. h.	
93 confécta. H.K.	(wh.) crowded-flowered.	—	1800.	10. 2. G. h.	
94 confertiflóra. s.s.	(re.) crowded-leaved.	—	—	5. 8. G. h.	
95 congéstâ. s.s.	(wh.) close-headed.	—	1812.	— G. h.	
96 cónica. L.C.	(re.) conical.	—	1820.	— G. h.	
97 conspícuâ. H.K.	(ye.) conspicuous.	—	1774.	— G. h.	
98 constântia. Lee.	(re.) constant-flowered.	—	1810.	— G. h.	
99 cordâta. H.K.	(wh.) heart-leaved.	—	1799.	4. 7. G. h.	
100 coriflóra. H.K.	(bh.) Coris-leaved.	—	1774.	8. 12. G. h.	
articulâris. B.M.	<i>calycinâ. A.H.</i>				
101 cory'dalis. L.T.	(re.) winged anthered.	—	1812.	2. 5. G. h.	
102 costâta. H.K.	(re.) ribbed-flowered.	—	1795.	2. 6. G. h.	
β supérba. A.H. (bh.) superb.			1820.	4. 5. G. h.	
103 Coventryâna. B.C. (ro.)	Lord Coventry's.	—	1808.	5. 6. G. h.	
104 crassiâ. A.H.	(li.) thick-leaved.	—	1826.	— G. h.	
105 crinitâ. B.C.	(wh.) long-haired.	—	1825.	— G. h.	
106 cristâflóra. L.T.	(li.) crested.	—	1803.	— G. h.	
107 crossotâ. s.s.	(wh.) fringed-calyxed.	—	1820.	— G. h.	
108 crucifórmis. s.s.	(ye.) cross-like.	—	1818.	— G. h.	
109 cruenta. H.K.	(cr.) bloody-flowered.	—	1774.	9. 5. G. h.	
110 cùbica. H.K.	(pu.) cube-flowered.	—	1790.	4. 7. G. h.	
β mīnor. B.C.	(pu.) lesser cube-flowered.	—	—	— G. h.	
111 cumuliflóra. L.T.	(wh.) heap-flowered.	—	1801.	5. 9. G. h.	
112 curviflóra. H.K.	(ye.) curve-flowered.	—	1774.	7. 10. G. h.	
113 curvirôstris. L.T.	(li.) curve-styled.	—	1790.	— G. h.	
declinâta. Hort.					
114 Cushiniâna. Lee. (re.)	Cushin's.	—	1816.	— G. h.	
115 cuspidigera. L.T.	(li.) cuspidate.	—	—	2. 5. G. h.	
116 cylindrica. Th.	(wh.) white cylindrical.	—	1810.	3. 5. G. h.	
117 cyrillæflóra. L.T.	(li.) Cyrilla-flowered.	—	1800.	2. 8. G. h.	
118 daphnaeflóra. B.C. (bh.)	Daphne-flowered.	—	1791.	4. 5. G. h.	
119 daphnoïdes. B.C. (ro.)	Daphne-like.	—	—	— G. h.	
120 decipiens. s.s.	(fl.) slender woolly.	—	1822.	— G. h.	
121 décôlorans. W.	(ro.wh.) discoloured.	—	1812.	— G. h.	
122 décora. H.K.	(li.) graceful.	—	1790.	11. 1. G. h.	Andr. heath. v. 3.
123 defléxa. L.C.	(wh.) deflexed.	—	1812.	5. 8. G. h.	
124 demissa. L.C.	(re.) low.	—	1810.	— G. h.	
125 dénâsa. s.s.	(bh.) dense.	—	—	— G. h.	
126 densiflóra. W.	(re.) dense-leaved.	—	1811.	3. 8. G. h.	Andr. heath. ic.
127 denticulâta. W.	(wh.) tooth-calyxed.	—	—	— G. h.	
fastigiâta. A.H. nec aliorum.					Lodd. bot. cab. t. 1090.
128 depressa. H.K.	(wh.) depressed.	—	1789.	6. 8. G. h.	Andr. heath. v. 2.
μ pêstris. A.H.					
129 dianthiflóra. L.T.	(re.) pink-leaved.	—	1796.	3. 8. G. h.	
130 diaphâna. s.s.	(fl.) transparent.	—	1800.	5. 8. G. h.	Andr. heath. v. 4.
transpârens. B.C. 177. non Th.					
131 dichrus. s.s.	(re.gr.) Andrew's 2-color'd.	—	1790.	8. 5. G. h.	Andr. heath. v. 4.
hicolor. A.H. non W.					
132 Dickensoniâna. L.C. (va.)	Mr. Dickenson's.	—	1809.	— G. h.	
α álba.	(wh.) white-flowered.	—	—	— G. h.	
β rúbra.	(re.) red-flowered.	—	—	— G. h.	

133	diosmæfòlia. L.T. (<i>li.</i>)	Diosma-leaved.	C. B. S.	1792.	3. 7.	G. h.	
134	diotæflòra. L.T. (<i>wh.</i>)	Diotis-flowered.	—	1795.	3. 8.	G. h.	
135	díscolor. H.K.	(<i>bh.</i>) different-coloured.	—	1788.	10. 3.	G. h.	Andr. heath. v. 1.
136	dístans. s.s.	(<i>vi.</i>) distant.	—	1816.	—	G. h.	
137	Donniána. Lee.	(<i>re.</i>) Donn's.	—	1812.	5. 9.	G. h.	
138	droseróides. A.H.	(<i>pu.</i>) Sundew-like. <i>droseróides</i> β <i>minor</i> . A.H.	—	—	7. 10.	G. h.	Andr. heath. v. 4.
139	dumosâ. A.H.	(<i>re.</i>) bushy.	—	—	—	G. h.	Andr. heath. v. 4.
140	echiifòra. H.K.	(<i>pi.</i>) Echium-flowered.	—	1798.	2. 7.	G. h.	— v. 3.
	β <i>purpurea</i> . A.H.	(<i>pu.</i>) purple-flowered.	—	1812.	—	G. h.	— v. 4.
141	elátâ. H.K.	(<i>ye.</i>) tall.	—	1790.	7. 9.	G. h.	— v. 2.
142	élegans. H.K.	(<i>pi.</i>) elegant.	—	1799.	11. 5.	G. h.	Bot. mag. t. 960.
143	elongátâ. B.C.	(<i>wh.</i>) elongated.	—	1820.	3. 6.	G. h.	Lodd. bot. cab. t. 738.
144	emarginatâ. A.H.	(<i>wh.</i>) hairy cupped.	—	1800.	6. 10.	G. h.	Andr. heath. ic.
145	embothriifòlia. L.T. (<i>re.</i>)	Embothrium-leav'd.	—	—	2. 5.	G. h.	Bot. mag. t. 447.
146	empetrifòlia. H.K.	(<i>re.</i>) crowberry-leaved.	—	1774.	4. 7.	G. h.	Andr. heath. v. 2.
147	empetroides. H.K.	(<i>pu.</i>) close-flowered.	—	1758.	5. 8.	G. h.	Lodd. bot. cab. t. 1186.
148	epistómia. L.C.	(<i>wh.</i>) spigget.	—	1800.	—	G. h.	
149	equisetifòlia. L.T. (<i>re.</i>)	Equisetum-leaved. <i>articuláris</i> . Hort.	—	—	—	G. h.	
150	eriocéphala. B.C.	(<i>wh.</i>) woolly-headed.	—	1816.	6. 8.	G. h.	Lodd. bot. cab. t. 1270.
151	erósâ. B.C.	(<i>bh.</i>) gnawed-petal'd.	—	—	—	G. h.	— t. 133.
152	erubéscens. H.K.	(<i>bh.</i>) reddish-flowered.	—	1800.	3. 8.	G. h.	Andr. heath. v. 3.
153	Ewerâna. H.K.	(<i>cr.</i>) Ewer's. <i>a glâbra.</i> (cr.) smooth-leaved. <i>β longiflòra.</i> (cr.) long-flowered. <i>γ pilosa.</i> (cr.) hairy-leaved.	—	1793.	7. 11.	G. h.	— v. 2.
154	exígua. L.T.	(<i>pu.</i>) small downy.	—	1790.	3. 9.	G. h.	
155	exímia. L.C.	(<i>ro.</i>) choice.	—	1811.	—	G. h.	Lodd. bot. cab. t. 1105.
156	exsérta. L.C.	(<i>wh.</i>) exserted.	—	1820.	—	G. h.	
157	expánsa. L.C.	(<i>re.</i>) expanded.	—	1811.	—	G. h.	
158	exprómta. s.s.	(<i>re.</i>) woolly-branched.	—	—	—	G. h.	
159	exsúdans. B.C.	(<i>pi.</i>) sweating.	—	1810.	8. 10.	G. h.	Lodd. bot. cab. t. 287.
160	exsúrgens. H.K.	(<i>or. re.</i>) quiver-formed.	—	1792.	1. 12.	G. h.	Andr. heath. v. 1.
161	fábrilis. L.T.	(<i>re.</i>) tiled-leaved.	—	1791.	4. 8.	G. h.	
162	fállax. L.T.	(<i>wh.</i>) viscous-flower'd.	—	—	11. 3.	G. h.	
163	fasciculáris. H.K.	(<i>ro. gr.</i>) cluster-flower'd. <i>coronata</i> . A.H. <i>radiflòra</i> . L.T.	—	1787.	2. 7.	G. h.	Andr. heath. v. 1.
164	fastigiâta. W.	(<i>bh.</i>) crowded. <i>Walkeri</i> . A.H. v. 1. <i>nec</i> v. 4.	—	1792.	2. 7.	G. h.	Andr. heath. v. 1.
165	faústa. L.T.	(<i>re.</i>) bristly-leaved.	—	1795.	4. 6.	G. h.	
166	ferruginea. H.K.	(<i>cr. bh.</i>) rusty.	—	1798.	5. 7.	G. h.	Andr. heath. v. 3.
167	fésta. L.T.	(<i>bh.</i>) angular-stalked.	—	—	3. 5.	G. h.	
168	fibula. L.en.	(<i>li.</i>) button.	—	1812.	5. 6.	G. h.	Bot. reg. t. 6.
169	filamentòsa. H.K.	(<i>re.</i>) long-peduncled.	—	1800.	1. 12.	G. h.	
170	filifòrmis. L.T.	(<i>re.</i>) filiform.	—	—	2. 5.	G. h.	
171	fimbriâta. A.H.	(<i>re.</i>) fringed.	—	1800.	3. 7.	G. h.	Andr. heath. ic.
172	finitíma. L.C.	(<i>pu.</i>) related.	—	—	—	G. h.	
173	fistulæflòra. L.T. (<i>wh.</i>) white slender-flower'd. <i>tenuiflòra</i> β <i>álba</i> . A.H. <i>Cliffordiana</i> . B.C. 34.	—	—	1800.	7. 11.	G. h.	Andr. heath. v. 3.
174	fláccida. L.en.	(<i>wh.</i>) flaccid.	—	1810.	—	G. h.	
175	flagellàris. L.en.	(<i>st.</i>) whipcord.	—	—	—	G. h.	
176	flagellifòrmis. A.H.	(<i>re.</i>) whipcord-like.	—	1812.	—	G. h.	Andr. heath. v. 4.
177	flámmea. H.K.	(<i>ye.</i>) flame-flowered. <i>bibax</i> . L.T.	—	1798.	10. 5.	G. h.	— v. 2.
178	flávâ. H.K.	(<i>ye.</i>) 3-leaved yellow.	—	1795.	9. 4.	G. h.	Andr. heath. v. 2.
	β <i>imbricâta</i> . B.M.	(<i>ye.</i>) imbricated.	—	—	—	G. h.	Bot. mag. t. 1815.
179	flexicáulis. H.K.	(<i>re.</i>) crook-stalked. <i>glandulosa</i> . A.H. <i>nec</i> <i>aliorum</i> .	—	1800.	5. 1.	G. h.	Andr. heath. v. 2.
180	flexuôsa. H.K.	(<i>wh.</i>) zigzag.	—	1792.	4. 7.	G. h.	Andr. heath. v. 1.
181	floribùnda. B.C.	(<i>i.</i>) many-flowered.	—	1800.	11. 3.	G. h.	Lodd. bot. cab. t. 176.
182	flòrida. H.K.	(<i>pu.</i>) florid.	—	1803.	5. 8.	G. h.	Thumb. eric. n. 64. t. 6.
183	foliacea. A.H.	(<i>ye.</i>) foliaceous.	—	1822.	5. 7.	G. h.	Andr. heath. v. 4.
184	foliculâris. H.K.	(<i>ye.</i>) yellow pencilled. <i>Petiveriana</i> . A.H.	—	1794.	2. 7.	G. h.	— v. 1.
185	formòsa. W.	(<i>va.</i>) handsome.	—	1795.	3. 8.	G. h.	Thunb. eric. n. 82. t. 3.
	α <i>álba</i> .	(<i>wh.</i>) white-flowered. <i>reflexa alba</i> . Hort.	—	—	—	G. h.	Andr. heath. v. 4.
	β <i>rúbra</i> .	(<i>re.</i>) red-flowered. <i>reflexa rubra</i> . Hort.	—	—	—	G. h.	— v. 4.
186	frágrans. H.K.	(<i>pu.</i>) fragrant.	C. B. S.	1803.	3. 7.	G. h.	Bot. mag. t. 2181.

187	fùgax. L.T.	(pu.) fugacious.	C. B. S.	1800.	3. 5. G. h.	
188	fúrfurósa. H.K.	(re.) columnar-threaded. <i>monadelpha. A.H. non B.M.</i>	—	1789.	8. 12. G. h.	Andr. heath. v. 1.
189	gélida. H.K.	(gr.) green verticillate. <i>gilva. w. alveiflòra. L.T.</i>	—	1790.	4. 7. G. h.	Andr. heath. v. 2.
190	gemífera. B.M. (or.) jewel.	—	1802.	7. 10. G. h.	Bot. mag. t. 2266.	
191	genistæfòlia. L.T. (re.) Genista-leaved.	—	—	10. 5. G. h.		
192	glábra. L.en.	(wh.) glabrous.	—	1802.	5. 8. G. h.	
193	glandulòsa. Th.	(cr.) glandular-haired.	—	1801.	— G. h.	
194	gláuca. H.K.	(pu.) glaucous-leaved.	—	1792.	— G. h.	Bot. mag. t. 580.
195	globòsa. w.	(bh.) globular-flowered.	—	1789.	7. 9. G. h.	Andr. heath. v. 4.
196	glomeràta. s.s.	(li.) glomerate.	—	1812.	— G. h.	— v. 4.
197	glomiflòra. L.T. (wh.) round-flowered.	—	—	—	4. 9. G. h.	
198	glutinòsa. H.K. (pu.) glutinous. <i>drosoroídes. A.H.</i>	—	1787.	7. 10. G. h.	Icon. hort. kew. t. 17.	
199	gnaphalòdes. w. (wh.) cudweed-like.	—	1812.	2. 4. G. h.	Pluk. m.68. t. 346. f.11.	
200	grácilis. H.K.	(re.) gracile.	—	1794.	2. 6. G. h.	Wend. eric. 8. p. 9. ic.
201	grandiflòra. H.K. (ye.) great-flowered.	—	—	1775.	5. 9. G. h.	Bot. mag. t. 189.
202	grandinòsa. B.C. (wh.) hail-flowered.	—	—	1820.	— G. h.	Lodd. bot. cab. t. 627.
203	halicácaba. H.K. (wh.) bladder-flowered.	—	—	1780.	5. 8. G. h.	Andr. heath. v. 2.
204	helianthemifòlia. L.T. plain-leaved.	—	—	1796.	6. 9. G. h.	
205	herbácea. L.	(fl.) trailing.	S. Europe.	1763.	1. 8. H. h.	Bot. mag. t. 471.
206	Hibbertiàna. H.K. (cr.gr.) Hibbert's.	C. B. S.	—	1800.	6. 9. G. h.	Andr. heath. v. 3.
207	hirsùta. B.C. (wh. pu.) hairy.	—	—	1812.	5. 8. G. h.	Lodd. bot. cab. t. 754.
208	hírta. w.	(re.) hairy-leaved.	—	1795.	4. 7. G. h.	Thunb. eric. n. 56. t. 2.
209	hirtiflòra. B.M.	(li.) hairy-flowered.	—	1790.	— G. h.	Bot. mag. t. 481.
210	híspida. s.s.	(re.) hispid.	—	1791.	7. 9. G. h.	Andr. heath. ic.
211	hispídula. w.	(pu.) bristly-stemmed.	—	—	6. 8. G. h.	
212	holosericea. L.T. (wh.) velvetty.	—	—	—	2. 7. G. h.	Andr. heath. v. 2.
213	horizontalis. H.K. (wh.) horizontal-leaved.	—	—	1800.	7. 9. G. h.	Lodd. bot. cab. t. 389.
214	Humeána. B.C. (pi.) Sir A. Hume's.	—	—	1808.	5. 7. G. h.	
215	humifúsia. L.T.	(pi.) low.	—	—	6. 7. G. h.	
216	hyacinthóides. H.K. (pi.) Hyacinth-flower'd.	—	—	1798.	6. 8. G. h.	Andr. heath. v. 3.
217	hyssopifòlia. L.T. (re.) Hyssop-leaved.	—	—	1800.	5. 9. G. h.	
218	ignéscens. H.K. (or.sc.) fiery.	—	—	1792.	3. 7. G. h.	Andr. heath. v. 2.
219	imbecilla.	(re.) feeble.	—	1793.	6. 7. G. h.	
	grácilis. L.T. nec aliorum.	—	—	—	—	
220	imbricáta. w.	(pu.) imbricated.	—	1786.	5. 8. G. h.	Lodd. bot. cab. t. 1243.
221	imperiális. s.s.	(ro.) imperial.	—	1816.	— G. h.	Andr. heath. v. 4.
222	incána. s.s.	(wh. re.) hoary.	—	—	— G. h.	Wend. er. f. 18. n. 4. ic.
223	incarnàta. w.	(fl.) flesh-coloured.	—	1791.	— G. h.	Andr. heath. v. 1.
224	incúrva. w.	(wh.) incurved.	—	—	— G. h.	
225	infatáa. w.	(ro.) inflated.	—	1800.	5. 9. G. h.	Thunb. dis. n. 67. t. 2. f.2.
226	infundibulifòrmis. (ro.wh.) funnel-flower'd.	—	—	1812.	— G. h.	Lodd. bot. cab. t. 589.
227	intertéxta. B.C. (wh.) interwoven.	—	—	1810.	— G. h.	— t. 1034.
228	intervallàris. L.T. (bh.) distant-leaved.	—	—	—	5. 11. G. h.	
229	Irbyána. A.H.	(bh.) Irby's.	—	1800.	6. 10. G. h.	Andr. heath. v. 3.
230	jasminifòlia. H.K. (wh.) Jasmine-flower'd.	—	—	1794.	— G. h.	v. 1.
231	jubátia. s.s.	(re.) crest-flowered.	—	1816.	— G. h.	Lodd. bot. cab. t. 799.
232	Juliána. s.s.	(re.) Julian's.	—	1812.	5. 10. G. h.	
233	labiális. L.T.	(pi.) lipped.	—	—	— G. h.	Andr. heath. v. 3.
234	lachneáfòlia. H.K. (pu.) Lachnea-leaved.	—	—	1793.	5. 7. G. h.	Lodd. bot. cab. t. 901.
235	lactifòlia. B.C. (st.) milk-coloured. tricolor. s.s.	—	—	1816.	— G. h.	
236	l'avis. A.H.	(wh.) smooth.	—	1790.	3. 6. G. h.	Andr. heath. v. 3.
237	Lambertiàna. H.K. (wh.) Lambert's.	—	—	1800.	5. 8. G. h.	— v. 2.
238	lanátia. w.	(sn.) woolly.	—	1775.	2. 5. G. h.	Wend. eric. f. 5. n. 2.
239	lanceolàta. P.S.	(wh.) spear-leaved.	—	1791.	6. 12. G. h.	— fasc. 8. p. 13. ic.
240	lanuginòsa. H.K.	(br.) large brown-flower'd.	—	1803.	9. 1. G. h.	Andr. heath. v. 3.
241	laricina. s.s.	(pi.) Larch-leaved.	—	1824.	6. 8. G. h.	
242	lasíophylla. s.s.	(li.) downy-leaved.	—	1816.	5. 8. G. h.	
243	laterális. H.K.	(fl.) side-flowered.	—	1791.	3. 7. G. h.	Andr. heath. v. 1.
244	latifòlia. H.K.	(re.) broad-leaved.	—	1800.	5. 8. G. h.	— v. 2.
245	lavandulafòlia. L.T. (re.) Lavender-leaved.	—	—	1795.	— G. h.	Bot. mag. t. 1720.
246	Lawsoni. B.M.	(re.) red slender-flower'd.	—	1802.	4. 7. G. h.	Andr. heath. v. 3.
247	láxa. H.K.	(li.) loose-flowered.	—	1800.	9. 2. G. h.	— v. 1.
248	Leeána. H.K.	(ye.) Lee's.	—	1788.	8. 1. G. h.	Andr. heath. v. 2.
249	leptocárpha. s.s. (re.) slender chaffed.	—	—	1824.	4. 7. G. h.	
250	leucántha. L.en.	(wh.) white-blossomed. leucanthèra. A.H. non L.	—	1803.	1. 5. G. h.	

	C. B. S.	1803.	2. 6. G. h.	
251 leucanthèra. w. (wh.) white-tipped.	—	1790.	1. 5. G. h.	Andr. heath. v. 2.
252 Linnæâna. H. K. (pu. wh.) Linnæus's. <i>Linnæoides</i> . A. H.	—	1822.	— G. h.	Andr. heath. v. 4.
β supérba. A. H. (pu. wh.) superb.	—	4. 8. G. h.	Lodd. bot. cab. t. 983.
2 longiflora. B. C. (ye.) long-flowered.	—	1787.	2. 7. G. h.	Wend. eric. t. 3.
254 longifolia. w. (re.) long-leaved. pinea. Wend.	—	1818.	3. 8. G. h.	Lodd. bot. cab. t. 103.
255 longipedunculàta. (li.) long-stalked.	—	1800.	4. 1. G. h.	Andr. heath. v. 2.
256 lúcida. H. K. (pi.) lucid.	—	1774.	2. 5. G. h.	— v. 1.
257 lútea. H. K. (ye.) yellow.	—	1790.	— G. h.	Andr. heath. v. 4.
258 lyrígera. L. T.	(re.) lyre-bearing.	1816.	4. 8. G. h.	— v. 1.
259 magnífica. s. s. (ro.) magnificent.	—	1762.	7. 10. G. h.	Lodd. bot. cab. t. 125.
260 mammòsa. w. (pu.) nipple. a purpurea. (pu.) purple-flowered.	—	—	— G. h.	— t. 951.
β pálida. (li.) pale-flowered.	—	—	— G. h.	Andr. heath. v. 1.
261 margaritacea. H. K. (wh.) pearl-flowered.	—	1775.	5. 9. G. h.	— v. 1.
262 mariòlia. H. K. (wh.) Marum-leaved.	—	1773.	5. 7. G. h.	Bot. mag. t. 336.
263 Massoni. H. K. (or. gr.) Masson's.	—	1787.	7. 10. G. h.	— t. 471.
264 mediterrânea. w. (re.) Mediterranean. Portugal.	—	1648.	3. 5. H. h.	Lodd. bot. cab. t. 867.
265 melanthèra. w. (li.) dark-anthered. C. B. S.	—	1803.	6. 8. G. h.	— t. 333.
266 melastoma. A. H. (ye.) black-anthered.	—	1795.	4. 7. G. h.	Andr. heath. v. 1.
267 mellifera. s. s. (pu.) honey-bearing.	—	1816.	4. 8. G. h.	Bot. mag. t. 612.
268 metulæflora. H. K. (re.) nine-pin.	—	1798.	6. 8. G. h.	— v. 4.
269 microphylla. L. C. (re.) small-leaved.	—	—	— G. h.	Andr. heath. v. 4.
270 micròstoma. s. s. (re.) aggregate-flower'd.	—	1816.	— G. h.	— v. 1.
271 minutæflora. A. R. (pu.) minute-flowered.	—	1822.	5. 7. G. h.	Andr. heath. v. 4.
272 mirabilis. A. H. (pu.) wonderful.	—	—	3. 8. G. h.	— v. 4.
273 modésta. L. T. (bh.) modest.	—	—	— G. h.	Andr. heath. v. 4.
274 molleàris. P. S. (pu.) soft-leaved.	—	1803.	4. 10. G. h.	Bot. mag. t. 1370.
275 móllis. s. s. (pu.) soft.	—	1816.	— G. h.	Andr. heath. v. 4.
276 mollissima. L. C. (wh.) softest.	—	—	— G. h.	Bot. mag. t. 1915.
277 monadélphia. B. M. (wh.) monadelphous. <i>Banksia</i> β purpurea. A. H.	—	1789.	5. 7. G. h.	Andr. heath. v. 4.
278 Monsoniæ. H. K. (wh.) Lady Monson's.	—	1787.	4. 9. G. h.	— v. 1.
279 montàna. L. C. (li.) mountain.	—	1816.	— G. h.	Lodd. bot. cab. t. 1212.
280 moschata. A. H. (li.) musky.	—	1805.	5. 7. G. h.	Andr. heath. v. 4.
281 mucòsa. H. K. (pu.) mucous.	—	1787.	2. 4. G. h.	— v. 2.
282 mucosoïdes. L. C. (li.) mucous-like.	—	1800.	3. 4. G. h.	Andr. heath. v. 4.
283 mucronata. s. s. (re.) mucronate.	—	1812.	4. 8. G. h.	— v. 2.
284 multiflora. H. K. (bh.) many-flowered. France.	—	1731.	6. 11. G. h.	Lodd. bot. cab. t. 114.
285 múnida. L. T. (bh.) neat-flowered. C. B. S.	—	—	5. 7. G. h.	Andr. heath. v. 1.
286 mündula. B. C. (pi.) neat.	—	1816.	3. 8. G. h.	— v. 3.
287 Muscari. H. K. (st.) musk.	—	1790.	— G. h.	— v. 2.
288 mutabilis. H. K. (pu.) mutable.	—	1798.	2. 10. G. h.	Andr. heath. v. 2.
289 nàna. L. T. (ye.) dwarf. depréssia. A. H.	—	1792.	5. 8. G. h.	Lodd. bot. cab. t. 764.
290 nidiflora. L. T. (bh.) nest-flowered.	—	—	10. 5. G. h.	Andr. heath. v. 1.
291 nidularia. B. C. (pi.) nest-like.	—	1816.	5. 8. G. h.	— v. 3.
292 nigricans. s. s. (wh.) black-stamened.	—	—	— G. h.	— v. 4.
293 nigrita. H. K. (wh.) black-tipped.	—	1790.	3. 7. G. h.	Andr. heath. v. 2.
294 nítida. H. K. (wh.) shining.	—	1800.	7. 10. G. h.	— v. 3.
295 nivalis. A. H. (wh.) snowy-flowered.	—	1820.	5. 6. G. h.	— v. 4.
296 nivea. s. s. (wh.) snowy.	—	1816.	5. 9. G. h.	Andr. heath. v. 1.
297 Nivéni. H. K. (re.) Niven's.	—	1799.	2. 7. G. h.	— v. 2.
298 nolæflora. L. T. (re.) bell-shaped.	—	—	2. 5. G. h.	Andr. heath. v. 2.
299 notabilis. s. s. (re.) notable.	—	1816.	3. 8. G. h.	— v. 3.
300 nudiflora. w. (ye.) small-bracted.	—	1783.	7. 8. G. h.	— v. 4.
301 obbâta. H. K. (re. wh.) bottle.	—	1796.	4. 7. G. h.	Andr. heath. v. 2.
302 obcordata. s. s. (wh.) obcordate.	—	1812.	— G. h.	Andr. heath. v. 1.
303 obliqua. H. K. (pu.) oblique-leaved.	—	1789.	8. 10. G. h.	Lodd. bot. cab. t. 1027.
304 obtusa. B. C. (re.) obtuse.	—	1816.	5. 8. G. h.	Andr. heath. v. 3.
305 octogôna. L. C. (re.) eight-angled.	—	—	— G. h.	— v. 4.
306 odorata. H. K. (wh.) perfumed.	—	1804.	4. 7. G. h.	Andr. heath. v. 1.
307 ollùa. s. s. (ro.) close smooth-leav'd.	—	1812.	— G. h.	Lodd. bot. cab. t. 1343.
308 onosmæflora. L. T. (ye.) Onosma-flowered. glutinosa. A. H. viscòsa. Wend.	—	1789.	9. 5. G. h.	— t. 1060.
309 oppositifolia. A. H. (wh.) opposite-leaved. a álba. (wh.) white-flowered.	—	1804.	3. 5. G. h.	Z
β rùbra. B. C. (re.) red-flowered.	—	—	— G. h.	

310 ostrina, B.C. (sc.) scarlet-flowered.	C. B. S.	1824. 3. 6. G. h.	Lodd. bot. cab. t. 1218.
311 ovata, B.C. (re.) ovate.	_____	1791. 2. 5. G. h.	_____ t. 417.
312 oxyccifolia, L.T. (bh.) Cranberry-leaved.	_____	_____ G. h.	Andr. heath. v. 4.
313 pâllens, s.s. (st.) pale.	_____	1812. 3. 8. G. h.	Lodd. bot. cab. t. 1355.
314 pâllida, B.C. (pu.) pallid.	_____	_____ G. h.	Andr. heath. v. 2.
315 paliflora, L.T. (pa.) pale-flowered.	_____	_____ 2. 8. G. h.	Lodd. bot. cab. t. 1194.
316 palustris, H.K. (bh.) marsh.	_____	1799. 5. 10. G. h.	Andr. heath. v. 2.
317 paniculata, w. (pu.) paniced.	_____	1774. 2. 8. G. h.	Lodd. bot. cab. t. 197.
318 pannosa, L.T. (wh.) cloth-flowered.	_____	_____ 2. 6. G. h.	Petiv. gaz. t. 3. f. 7.
319 pârilis, L.T. (re.) matched.	_____	1789. 5. 8. G. h.	Andr. heath. v. 3.
320 Parmentieri, B.C. (wh.) Parmentier's.	_____	1816. _____ G. h.	Andr. heath. v. 3.
β rosea, L.C. (ro.) rose-coloured.	_____	_____ G. h.	_____ v. 4.
321 parviflora, L.T. (pu.) small-flld. downy.	_____	1790. 3. 9. G. h.	Lodd. bot. cab. t. 902.
322 Passerina, w. (wh.) sparrow-wort.	_____	1800. 11. 5. G. h.	Andr. heath. v. 2.
323 pátnes, A.H. (pu.) spreading.	_____	_____ 3. 6. G. h.	Wend. eric. 4. p. 5. ic.
324 Patersoni, H.K. (ye.) Paterson's.	_____	1791. 3. 8. G. h.	Andr. heath. v. 2.
325 pavettaeflora, L.T. (wh.) Pavetta-flowered.	_____	_____ 5. 8. G. h.	_____ v. 4.
326 pectinifolia, L.T. (bh.) pectinated.	_____	1800. 11. 6. G. h.	Bot. mag. t. 342.
327 pedunculata, s.s. (re.) peduncled.	_____	1810. 5. 9. G. h.	Wend. eric. 1. p. 7. ic.
328 pellucida, A.H. (wh.) pellucid.	_____	1800. 10. 6. G. h.	Andr. heath. v. 3.
β rûbra, A.H. (re.) red-flowered.	_____	1806. _____ G. h.	_____ v. 4.
329 peltata, A.H. (gr. pu.) peltate-stigma'd.	_____	1804. 4. 8. G. h.	Andr. heath. v. 3.
330 pendula, w. (pu.) pendulous.	_____	1791. 7. 8. G. h.	Lam. ill. t. 288. f. 3.
331 penicillata, A.H. (re.) pencilled.	_____	1774. 4. 7. G. h.	Lodd. bot. cab. t. 265.
332 penicilliflora, L.T. (wh.) white-pencilled.	_____	1792. 5. 8. G. h.	Bot. mag. t. 443.
333 periplocaefolia, L.T. (li.) Periploca-leaved.	_____	_____ 8. 12. G. h.	Lodd. bot. cab. t. 606.
334 persoluta, H.K. (re.) garland.	_____	1774. 2. 5. G. h.	Lodd. bot. cab. t. 1259.
335 perspicua, H.K. (wh.) clear-flowered.	_____	1790. 3. 6. G. h.	Andr. heath. v. 1.
β Linnæa, A.H. v. 2.	_____	1774. 3. 7. G. h.	Andr. heath. v. 1.
336 petiolata, H.K. (wh.) rosemary-leaved.	_____	_____ G. h.	Lam. ill. t. 288. f. 3.
337 Petiveriana, H.K. (ye.) Petiver's.	_____	_____ 5. 8. G. h.	Lodd. bot. cab. t. 265.
338 Peziza, B.C. (wh.) button-flowered.	_____	1800. 4. 7. G. h.	Bot. mag. t. 443.
339 phylcoïdes, w. (pu.) Phyllica-like.	_____	1788. 3. 7. G. h.	Lodd. bot. cab. t. 606.
340 physodes, H.K. (wh.) sticky.	_____	1800. _____ G. h.	Andr. heath. v. 4.
341 picta, L.C. (wh. ye.) painted.	_____	1789. 4. 5. G. h.	Andr. heath. v. 4.
342 pilosa, s.s. (gr.) pilose.	_____	1790. 8. 12. G. h.	Bot. mag. t. 443.
343 pilulifera, w. (pu.) ball-bearing.	_____	_____ G. h.	Lodd. bot. cab. t. 606.
344 pînea, s.s. (wh.) Pine-like.	_____	1790. 6. 12. G. h.	Andr. heath. v. 1.
β purpurea, L.C. (pu.) purple.	_____	1820. _____ G. h.	_____ v. 4.
345 pinifolia, A.H. (re.) Pine-leaved.	_____	1790. _____ G. h.	_____ v. 3.
pitophylla, s.s.	_____	1774. 4. 7. G. h.	_____ v. 1.
a coccinea, A.H. (sc.) scarlet.	_____	1804. 7. 10. G. h.	Andr. heath. v. 4.
β discolor, A.H. (wh. pu.) two-coloured.	_____	1805. 2. 5. G. h.	Lodd. bot. cab. t. 1413.
346 planifolia, A.H. (re.) flat-leaved.	_____	1796. 5. 7. G. h.	Andr. heath. v. 3.
347 Plukenetiâna, H.K. (sc.) Plukenet's.	_____	_____ G. h.	_____ v. 4.
β pallida, L.C. (pa.) pale-flowered.	_____	1802. 4. 7. G. h.	Bot. mag. t. 1548.
γ albens, A.H. (wh.) white-flowered.	_____	1800. _____ G. h.	Andr. heath. v. 2.
348 polytricha, (sc. gr.) many-haired.	_____	1791. _____ G. h.	Bot. mag. t. 2140.
hirta, Lod. bot. cab. nec aliorum pilosa, Hort.	_____	1816. _____ G. h.	Wend. eric. f. 3. p. 5. ic.
349 præcox, L.C. (li.) early-dwarf.	_____	1800. 7. 8. G. h.	Thunb. dis. n. 24. t. 4. f. 2.
350 prægnans, A.H. (bh.) swelled.	_____	1816. 5. 8. G. h.	Andr. heath. v. 4.
β coccinea, (sc.) scarlet-flowered.	_____	1790. 2. 12. G. h.	Andr. heath. v. 4.
351 præstans, s.s. (bh.) excelling.	_____	1792. 1. 5. G. h.	Andr. heath. v. 4.
352 primuloides, A.H. (re. wh.) Cowslip-flower'd.	_____	1812. 3. 8. G. h.	Andr. heath. v. 4.
353 princeps, H.K. (pi.) fine red.	_____	1820. 7. 10. G. h.	Andr. heath. v. 4.
354 procera, s.s. (ye.) lofty.	_____	1820. 7. 10. G. h.	Andr. heath. v. 4.
355 procumbens, s.s. (re.) procumbent.	_____	1820. 7. 10. G. h.	Andr. heath. v. 4.
356 propendens, H.K. (pu.) pendent.	_____	1820. 7. 10. G. h.	Andr. heath. v. 4.
357 protrudens, L.en. (wh.) protruding.	_____	1820. 7. 10. G. h.	Andr. heath. v. 4.
358 pubescens, H.K. (li.) pale downy.	_____	1820. 7. 10. G. h.	Andr. heath. v. 4.
359 pubígera, L.T. (li.) pubescent-flowered.	_____	1820. 7. 10. G. h.	Andr. heath. v. 4.
360 pubidûnda, L.T. (bh.) nodding.	_____	1820. 7. 10. G. h.	Andr. heath. v. 4.
nûtans, Wendland.	_____	1820. 7. 10. G. h.	Andr. heath. v. 4.
361 pulchella, w. (re.) neat.	_____	1820. 7. 10. G. h.	Andr. heath. v. 4.
362 pulcherrima, (ro. wh.) beautiful.	_____	1820. 7. 10. G. h.	Andr. heath. v. 4.
jasminiflora β minor, A.H. v. 4. Irbyana, Hort. non Andr.	_____	1820. 7. 10. G. h.	Andr. heath. v. 4.
363 pulverulenta, L.C. (wh.) powdered.	_____	1820. 7. 10. G. h.	Andr. heath. v. 4.
364 pulviniformis, L.T. (bh.) cushion.	_____	1820. 7. 10. G. h.	Andr. heath. v. 4.
365 pùmila, s.s. (li.) short.	_____	1820. 7. 10. G. h.	Andr. heath. v. 4.

366 pùra. B.C. (wh.) clear.	C. B. S.	1812.	5. 8. G. h.	Lodd. bot. cab. t. 72.
367 purpùrea. H.K. (pu.) purple-flowered.	1789.	1. 12. G. h.	Andr. heath. v. 2.	
368 pusilla. L.T. (bh.) low.	6. 7. G. h.		
369 pyramidális. H.K. (bh.) pyramidal.	1787.	2. 5. G. h.	Bot. mag. t. 366.	
370 pyramidifórmis. s.s. (bh.) pyramid-formed.	1818.	— G. h.		
371 pyrolæflóra. L.T. (wh.) Pyrola-flowered.	1790.	5. 7. G. h.	Andr. heath. v. 4.	
andromedæflóra. β áiba. A.H. v. 4.				
372 quadraeflóra. L.T. (re.) four-flowered.	—	3. 8. G. h.		
373 quadranguláris. A.H. (fl.wh.) square-tubed.	1812.	5. 8. G. h.	Andr. heath. v. 4.	
374 radiáta. H.K. (re.) rayed.	1798.	8. 11. G. h.	— v. 1.	
β discolor. A.H. (re.wh.) two-coloured.	1820.	— G. h.	— v. 4.	
375 racemòsa. w. (fl.) racemed.	1795.	4. 5. G. h.		
376 racemífera. H.K. (re.) compact-flowered.	1803.	4. 6. G. h.	Andr. heath. v. 2.	
377 ramentácea. H.K. (pu.) slender-branched.	1786.	7. 12. G. h.	— v. 1.	
378 ramulòsa. Viv. (ii.) Italian.	1826.	5. 8. H. h.		
379 recurvàta. B.C. (wh.) recurved.	C. B. S.	1812.	— G. h.	Lodd. bot. cab. t. 1093.
380 refléxa. s.s. (pu.wh.) reflexed.	—	— G. h.		
381 réfulgens. A.H. (ro.gr.) fulgent.	1816.	5. 9. G. h.	Andr. heath. v. 4.	
382 régérminans. w. (li.) clustered-flower'd.	1791.	5. 8. G. h.		
383 resinòsa. H.K. (or.) varnished.	1803.	— G. h.	Bot. mag. t. 1139.	
vèrnix. A.H. v. 3.				
384 retorta. H.K. (ro.) recurved-leaved.	1787.	— G. h.	Bot. mag. t. 362.	
385 retrofleáxa. H.K. (re.) jointed.	—	7. 9. G. h.	Andr. heath. v. 1.	
pulchellá. A.H. non Th. articulàris. Th. non L.				
386 rígida. B.C. (re.) rigid.	1820.	5. 8. G. h.	Lodd. bot. cab. t. 1286.	
387 rigidifòlia. s.s. (re.) rigid-leaved.	1816.	6. 8. G. h.		
388 Rollisóni. (bh.) Rollison's.	1823.	5. 8. G. h.	Andr. heath. v. 4.	
exstírgens β hy'brida. A.H. v. 4.				
389 ròsea. H.K. (ro.) rose-coloured.	1798.	6. 10. G. h.	Andr. heath. v. 2.	
390 rubélia. B.M. (re.) reddish.	1812.	5. 8. G. h.	Bot. mag. t. 2165.	
391 rùbens. H.K. (re.) red-flowered.	1798.	6. 9. G. h.		
392 rùbida. B.C. (re.) ruby.	1820.	— G. h.	Lodd. bot. cab. t. 1166.	
393 rubrosépala. (re.wh.) red-calyxed.	1825.	— G. h.	Andr. heath. v. 4.	
ruber-cályx. A.H. v. 4.				
394 rugòsa. s.s. (ye.pi.) wrinkled.	1812.	— G. h.	Andr. heath. v. 4.	
395 Russeliána. B.C. (ro.) Duke of Bedford's.	1824.	— G. h.	Lodd. bot. cab. t. 1013.	
396 Sainsburyàna. s.s. (ro.) Sainsbury's.	1804.	7. 9. G. h.	Andr. heath. v. 4.	
397 sá lax. L.T. (re.) broad-stigma'd.	1796.	5. 9. G. h.		
398 Salisburýana. s.s. (ro.) Salisbury's.	1815.	— G. h.	Andr. heath. v. 4.	
399 sanguínea. B.C. (cr.) bloody.	3. 8. G. h.	Lodd. bot. cab. t. 36.	
400 sanguinolénta. B.M. (pu.) dark-flowered.	1810.	— G. h.	Bot. mag. t. 2263.	
pygmæa. Andr. heath. v. 4.				
401 Savileána. (re.) Savile's.	1800.	6. 7. G. h.	Lodd. bot. cab. t. 96.	
402 scabriúscula. B.C. (wh.) roughish.	—	— G. h.	— t. 517.	
403 scaríosa. B.C. (li.) scarose.	—	— G. h.	— t. 477.	
404 Schóllii. B.C. (pu.) Scholl's.	1790.	5. 9. G. h.	— t. 538.	
405 scopària. w. (gr.) small green-fld. S. Europe.	1770.	4. 5. F. h.	Linn. eric. n. 14. f. fl.	
406 Sebána. H.K. (re.) Seba's.	C. B. S.	1774.	3. 6. G. h.	Andr. heath. v. 1.
a fusca. Lod. (br.) brown-flowered.	—	— G. h.		
β lutea. Lod. (ye.) yellow-flowered.	—	— G. h.	Lodd. bot. cab. t. 266.	
γ minor. Lod. (or.) smaller.	—	— G. h.		
407 selaginifòlia. L.T. (re.) Selago-leaved.	1801.	2. 6. G. h.		
408 serpylifòlia. B.C. (li.) Serpyllum-leaved.	1812.	5. 8. G. h.	Lodd. bot. cab. t. 744.	
409 serratifòlia. H.K. (ye.) saw-leaved.	1790.	8. 12. G. h.	Andr. heath. v. 1.	
410 serrulàta. L.C. (ye.) sawed.	1814.	6. 8. G. h.		
411 setacea. A.H. (bh.) bristly-leaved.	1796.	2. 8. G. h.	Andr. heath. v. 1.	
412 sexfària. H.K. (wh.) six-angled.	1774.	5. 8. G. h.	— v. 2.	
413 Shannoniana. A.H. (bh.) Earl Shannon's.	1800.	6. 7. G. h.	Lodd. bot. cab. t. 168.	
414 sicæfòlia. L.T. (re.) dagger-leaved.	1791.	2. 7. G. h.		
415 simpliciflóra. H.K. (y.re.) single-flowered.	1774.	3. 7. G. h.	Wend. eric. 17. p. 69. ic.	
416 Smithiana. L.C. (re.) Smith's.	1810.	— G. h.		
417 socciiflóra. L.T. (gr.) green-pencilled.	1799.	4. 5. G. h.	Andr. heath. v. 1.	
Sebána viridis. A.H.				
418 Solándri. H.K. (re.) Solander's.	1800.	3. 9. G. h.	Andr. heath. v. 2.	
419 sórdida. H.K. (br.) sordid.	1790.	8. 5. G. h.	— v. 1.	
laniflóra. Wend.				
420 Sparrmánni. H.K. (ye.) Sparrman's	1794.	9. 3. G. h.	Andr. heath. v. 3.	
áspera. A.H. hystriciflóra. L.T.	1800.	3. 8. G. h.	Lodd. bot. cab. t. 1467.	
421 spársla. L.C. (li.) scattered.	Z 2			

422 speciosa. H.K. (<i>or.re.</i>) specious.	C. B. S.	1800.	6. 9. G. h.	Andr. heath. v. 2.
423 spicata. H.K. (<i>gr.</i>) spiked.	_____	1789.	1. 12. G. h.	_____ v. 1.
424 spiralis. L.C. (<i>wh.</i>) spiral.	_____	1816.	5. 8. G. h.	Wend. eric. 8. p. 5. ic.
425 splendens. w. (<i>sc.</i>) splendid.	_____	1792.	4. 9. G. h.	Lodd. bot. cab. t. 566.
426 spumosa. H.K. (<i>st.</i>) spounous.	_____	1786.	5. 8. G. h.	Andr. heath. v. 1.
427 spuria. H.K. (<i>pi.</i>) spurious.	_____	1796.	4. 8. G. h.	Andr. heath. v. 3.
428 squamæflora. L.T. (<i>pi.</i>) scaly-flowered.	_____	5. 11. G. h.	Andr. heath. v. 3.
429 squamosa. H.K. (<i>li.</i>) scaly-cupped.	_____	1794.	4. 6. G. h.	Andr. heath. v. 3.
430 squarrösa. L.T. (<i>pu.</i>) squarrose.	_____	3. 8. G. h.	Andr. heath. v. 3.
431 stagnalis. L.T. (<i>bh.</i>) stagnant.	_____	1790.	3. 6. G. h.	Andr. heath. v. 3.
432 staminea. A.H. (<i>st.</i>) reflex-stamened.	_____	1799.	6. 9. G. h.	Lodd. bot. cab. t. 893.
433 stellata. B.C. (<i>wh.</i>) starry. <i>solandroides</i> . Andr. heath. v. 4.	_____	1806.	— G. h.	Andr. heath. v. 4.
434 stellaria. s.s. (<i>bh.</i>) star-bearing.	_____	—	4. 8. G. h.	_____ v. 4.
435 stenantha. (<i>bh.</i>) slender-blush-flower'd.	_____	1820.	5. 8. G. h.	Andr. heath. v. 2.
<i>tenuiflora γ carnnea</i> . A.H. v. 4.				
436 stricta. H.K. (<i>li.</i>) straight-branch'd. S. Europe.	1765.	8. 11. G. h.	Lodd. bot. cab. t. 24.	
437 strigosa. H.K. (<i>li.</i>) dwarf downy.	C. B. S.	1775.	3. 8. G. h.	Bot. mag. t. 1984.
438 struthiolæflora. L.C. (<i>wh.</i>) Struthiola-flower'd.	1812.	5. 8. G. h.	Andr. heath. c. ic.	
439 stylaris. s.s. (<i>re.</i>) prominent-styled.	—	—	G. h.	Andr. heath. v. 1.
440 stylosa. L.T. (<i>li.</i>) long-styled.	1789.	1. 8. G. h.	_____ v. 2.	
441 suavæolens. L.C. (<i>li.</i>) sweet.	1800.	5. 11. G. h.	_____ v. 4.	
442 subulata. s.s. (<i>re.</i>) awl-leaved.	1818.	6. 8. G. h.	Bot. mag. t. 480.	
443 sulphurea. B.M. (<i>st.</i>) sulphur-coloured.	1812.	3. 5. G. h.	Andr. heath. v. 1.	
444 supèra. (<i>re.</i>) superb. <i>formosa</i> . A.H. <i>neè aliorum.</i>	1795.	6. 9. G. h.	Andr. heath. v. 2.	
<i>α coccinea</i> . (<i>sc.</i>) scarlet.	—	—	G. h.	Andr. heath. v. 3.
<i>β rösea</i> . (<i>ro.</i>) rose-coloured.	—	—	G. h.	Andr. heath. v. 4.
<i>γ bicolor</i> . (<i>ye.gr.</i>) two-coloured.	—	—	G. h.	Seb. mus. 1. t. 73. f. 6.
445 Swainsoniana. s.s. (<i>pi.</i>) Swainson's.	1810.	6. 7. G. h.	Wend. eric. 6. p. 9. ic.	
446 tardiflora. L.T. (<i>li.</i>) pubescent.	1790.	3. 8. G. h.	Andr. heath. v. 3.	
<i>pubescens</i> . B.M. <i>non L.</i>				Eng. bot. v. 15. t. 1014.
447 taxifolia. H.K. (<i>bh.</i>) yew-leaved.	1788.	7. 11. G. h.	Andr. heath. v. 1.	
448 tegulæfolia. L.T. (<i>wh.</i>) tiled.	1. 5. G. h.	Andr. heath. v. 4.	
449 Templeæ. A.H. (<i>ro.</i>) Lady Temple's.	1820.	5. 8. G. h.	_____ v. 2.	
450 tenella. H.K. (<i>re.</i>) delicate.	1791.	8. 5. G. h.	_____ v. 3.	
451 tenuiflora. A.H. (<i>st.</i>) slender yellow-flower'd.	1800.	4. 6. G. h.	Seb. mus. 1. t. 73. f. 6.	
452 tenuifolia. H.K. (<i>re.</i>) slender-leaved.	1794.	4. 8. G. h.	Andr. heath. v. 3.	
453 tenuis. L.T. (<i>wh.</i>) slender.	1790.	8. 2. G. h.	Eng. bot. v. 15. t. 1014.	
454 tenuissima. s.s. (<i>re.</i>) very-slender.	1803.	2. 8. G. h.	Andr. heath. v. 4.	
455 tetragona. H.K. (<i>ye.</i>) square-flowered.	1789.	7. 9. G. h.	Lodd. bot. cab. t. 1294.	
456 Tétralix. H.K. (<i>li.</i>) cross-leaved.	Britain.	H. h.	Bot. mag. t. 1214.
<i>α rübra</i> . (<i>re.</i>) red-flowered.	1812.	5. 9. G. h.	Andr. heath. v. 2.	
<i>β carnæa</i> . (<i>li.</i>) flesh-coloured.	—	—	H. h.	Andr. heath. v. 3.
<i>γ alba</i> . (<i>wh.</i>) white-flowered.	—	—	H. h.	Bot. mag. t. 1626.
457 teucriifolia. s.s. (<i>pu.</i>) Teucrium-leaved.	C. B. S.	1810.	— G. h.	Lodd. bot. cab. t. 962.
458 thalictriflora. s.s. (<i>wh.</i>) meadow-rue-flower'd.	—	1794.	5. 8. G. h.	Andr. heath. v. 3.
459 Thunbergii. H.K. (<i>or.</i>) Thunberg's.	—	1789.	— G. h.	Andr. heath. v. 3.
460 thymifolia. H.K. (<i>re.</i>) Thyme-leaved.	1800.	— G. h.	— v. 3.	
461 tiaræflora. H.K. (<i>re.</i>) turban-flower'd.	1812.	6. 8. G. h.	Bot. mag. t. 1626.	
462 togata. B.M. (<i>re.</i>) large-cupped.	2. 8. G. h.		
463 tomentosa. L.T. (<i>wh.</i>) woolly-flower'd.	1816.	3. 6. G. h.		
464 tortuosa. L.C. (<i>wh.</i>) twisted.	1822.	— G. h.		
465 totta. s.s. (<i>wh.</i>) bristly-branched.	5. 8. G. h.		
466 tragulifera. L.T. (<i>li.</i>) broad-anthered.	1797.	— G. h.		
467 translucens. Lee. (<i>bh.</i>) translucent.	1800.	— G. h.		
468 transparens. Th. (<i>bh.</i>) transparent.	1809.	— G. h.		
469 triceps. B.C. (<i>wh.</i>) three-headed.	1803.	6. 9. G. h.	Andr. heath. v. 3.	
470 tricolor. H.S.L. (<i>or.r.gr.</i>) three-coloured. <i>aristata minor</i> . A.H. <i>non tricolor</i> . s.s.	—	1774.	3. 6. G. h.	Wend. eric. 12. p. 13.
471 triflora. H.K. (<i>wh.</i>) three-flowered.	—	1812.	— G. h.	Lodd. bot. cab. t. 257.
472 triphylla. L.en. (<i>re.ye.</i>) three-leaved.	—	—	5. 10. G. h.	— t. 668.
473 triumphans. B.C. (<i>wh.</i>) triumphant.	—	—	— G. h.	
474 trüssula. B.C. (<i>wh.pi.</i>) dapper.	—	1790.	2. 8. G. h.	Andr. heath. v. 1.
475 tuberculæris. L.T. (<i>pu.</i>) tubercled.	—	1775.	4. 7. G. h.	Lodd. bot. cab. t. 1157.
476 tubiflora. H.K. (<i>pi.</i>) tube-flowered.	—	1822.	— G. h.	Bot. reg. t. 65.
477 tubiæcula. B.C. (<i>re.</i>) small-tubed.	—	1812.	5. 9. G. h.	
478 tumida. B.R. (<i>sc.</i>) tumid.	—	1793.	10. 2. G. h.	
479 turbiniæflora. L.T. (<i>wh.</i>) top-flowered.	—	—	—	

	C. B. S.	1800.	3. 6. G. h.	Andr. heath. v. 4.
480 <i>túrgida</i> . L.en. (<i>pi.</i>) turgid. <i>taxisfolia</i> β <i>májor</i> . A.H. v. 4.	—	1796.	6. 9. G. h.	
481 <i>turrígera</i> . L.T. (<i>li.</i>) Cypress. <i>cypressina</i> . Hort.	—	1782.	5. 7. G. h.	Andr. heath. v. 2.
482 <i>umbellátæ</i> . H.K. (<i>re.</i>) umbelled. Portugal.	—	1824.	5. 8. G. h.	— v. 4.
483 <i>undulátæ</i> . A.H. (<i>cr.</i>) crumple-flower'd. C. B. S.	—	1812.	5. 7. G. h.	— v. 4.
484 <i>única</i> , s.s. (<i>bh.</i>) one-flowered. <i>pedunculátæ</i> . A.H. <i>neq; aliorum</i> .	—	1778.	— G. h.	Icon. hort. kew. t. 16.
485 <i>urceoláris</i> . H.K. (<i>wh.</i>) pitcher-flowered.	—	1812.	4. 8. G. h.	
486 <i>ursina</i> . Lee. (<i>wh.</i>) shaggy.	—	7. 8. H. h.	Eng. bot. v. 1. t. 3.
487 <i>vágans</i> . H.K. (<i>bh.</i>) Cornish. Cornwall. β <i>álba</i> . (<i>wh.</i>) white-flowering.	—	— H. h.	
488 <i>vària</i> . B.C. (<i>va.</i>) variable. C. B. S.	—	1820.	5. 8. G. h.	Lodd. bot. cab. t. 1325.
489 <i>velitáris</i> . L.T. (<i>bh.</i>) javelin-flowered.	—	1790.	1. 6. G. h.	
490 <i>velleriflóra</i> . H.K. (<i>bh.</i>) woolly-flowered. <i>bruniádes</i> . A.H.	—	1774.	2. 6. G. h.	Andr. heath. v. 1.
491 <i>ventricosa</i> . H.K. (<i>bh.</i>) porcelain.	—	1787.	4. 9. G. h.	Bot. mag. t. 350.
492 <i>ventrosa</i> . (<i>wh.</i>) tun-bellied. <i>obbáta</i> β <i>umbellátæ</i> . A.H. v. 4.	—	1822.	5. 8. G. h.	Andr. heath. v. 4.
493 <i>verecúnda</i> . L.T. (<i>bh.</i>) blushing. <i>cérnua</i> . A.H. <i>non</i> Thunb.	—	1791.	4. 9. G. h.	Andr. heath. v. 1.
494 <i>verníciflúa</i> . L.T. (<i>wh.</i>) viscous-leaved.	—	1804.	3. 9. G. h.	
495 <i>versicolor</i> . H.K. (<i>sc.or.</i>) various-colored.	—	1790.	11. 5. G. h.	Andr. heath. v. 1.
496 <i>verticillátæ</i> . A.H. (<i>sc.</i>) verticillate.	—	1774.	7. 10. G. h.	— v. 1.
497 <i>vesiculáris</i> . L.T. (<i>bh.</i>) bladdered.	—	1796.	2. 8. G. h.	
498 <i>vestíflua</i> . L.T. (<i>re.</i>) clothed.	—	—	3. 6. G. h.	
499 <i>vestita</i> . H.K. (<i>va.</i>) tremulous. <i>a álba</i> . H.K. (<i>wh.</i>) white. β <i>incarnáta</i> . H.K. (<i>fl.</i>) flesh-coloured.	—	1789.	1. 12. G. h.	Andr. heath. v. 1.
γ <i>purpurea</i> . H.K. (<i>pu.</i>) purple-flowered. δ <i>rósea</i> . H.K. (<i>ro.</i>) rose-coloured. ϵ <i>fulgida</i> . H.K. (<i>sc.</i>) bright red. ζ <i>coccinea</i> . H.K. (<i>sc.</i>) scarlet. η <i>lutea</i> . H.K. (<i>ye.</i>) yellow.	—	—	— G. h.	— v. 2.
500 <i>villòsa</i> . H.K. (<i>wh.</i>) villous.	—	1800.	2. 6. G. h.	— v. 1.
501 <i>virgáta</i> , s.s. (<i>wh.</i>) twiggy.	—	1824.	— G. h.	— v. 2.
502 <i>viridescens</i> . B.C. (<i>gr.</i>) greenish.	—	1. 6. G. h.	— v. 2.
503 <i>viridiflóra</i> . A.H. (<i>gr.</i>) green-flowered.	—	1820.	5. 8. G. h.	Lodd. bot. cab. t. 233.
504 <i>viridipurpùrea</i> . W. (<i>gr.</i>) green & purple. Portugal.	—	5. 8. F. h.	Andr. heath. v. 4.
505 <i>viridis</i> . H.K. (<i>gr.</i>) green-flowered. C. B. S.	—	1800.	— G. h.	Andr. heath. v. 2.
506 <i>viscaria</i> . H.K. (<i>li.</i>) clammy-flower'd.	—	1774.	3. 7. G. h.	Icon. hort. kew. t. 1.
507 <i>Walkeriana</i> . (<i>re.</i>) Walker's. <i>Walkèria</i> β <i>rúbræ</i> . A.H. v. 4.	—	1806.	6. 8. G. h.	Andr. heath. v. 4.
508 <i>xeranthemifólia</i> . L.T. Xeranthemum-ld.	—	1812.	5. 8. G. h.	
509 <i>Zeyheri</i> . s.s. (<i>li.</i>) Zeyher's.	—	1824.	— G. h.	
MENZTESIA Sm. MENZTESIA. Octandria Monogynia.				
1 <i>ferruginea</i> , s.s. (<i>co.</i>) ferruginous. N. America.	1811.	5. 6. H. h.	Sm. ic. ined. 1. t. 56.	
2 <i>globularis</i> , s.s. (<i>co.</i>) globular-flower'd.	1806.	— H. h.	Sal. par. lond. t. 44.	
<i>ferruginea</i> β . B.M. 1571.				
3 <i>polifólia</i> , s.s. (<i>pu.</i>) Irish. <i>Erica Daboëcia</i> . E.B.	Ireland.	6. 9. H. h.	Eng. bot. v. 1. t. 35.
β <i>nana</i> . (<i>pu.</i>) dwarf.	— H. h.	
4 <i>cærulea</i> , s.s. (<i>pu.</i>) yew-leaved.	Scotland.	6. 7. H. h.	Eng. bot. v. 35. t. 2169.
PYROLA S.S. PYROLA. WINTER-GREEN. Decandria Monogynia. L.				
1 <i>uniflóra</i> , s.s. (<i>wh.</i>) single-flower'd. Britain.	6. 7. H. $\frac{1}{2}$.	Eng. bot. t. 146.	
2 <i>secunda</i> , s.s. (<i>wh.</i>) serrated.	— H. $\frac{1}{2}$.	— t. 517.	
3 <i>mínor</i> , s.s. (<i>bh.</i>) lesser.	— H. $\frac{1}{2}$.	— t. 158.	
β <i>rósea</i> . E.B. (<i>ro.</i>) rose-coloured.	England.	7. 8. H. $\frac{1}{2}$.	— t. 2543.
4 <i>média</i> , s.s. (<i>bh.</i>) intermediate.	6. 7. H. $\frac{1}{2}$.	— t. 945.
5 <i>dentata</i> . D.D. (<i>st.</i>) toothed.	N. America.	1827.	— H. $\frac{1}{2}$.	
6 <i>elliptica</i> . D.D. (<i>wh.</i>) elliptic-leaved.	—	1826.	— H. $\frac{1}{2}$.	
7 <i>occidentalis</i> . D.D. (<i>st.</i>) western.	—	1827.	— H. $\frac{1}{2}$.	
8 <i>rotundifólia</i> , s.s. (<i>st.</i>) round-leaved.	Britain.	— H. $\frac{1}{2}$.	Eng. bot. t. 213.
9 <i>americana</i> . (<i>wh.</i>) American. <i>rotundifolia</i> . Ph. non Eng. bot.	N. America.	— H. $\frac{1}{2}$.	
10 <i>asarifolia</i> , s.s. (<i>st.</i>) Asarum-leaved.	—	1822.	— H. $\frac{1}{2}$.	
11 <i>chlorántha</i> , s.s. (<i>st.</i>) cream-colour'd.	—	—	— H. $\frac{1}{2}$.	
CHIMA'PHILA Ph. CHIMA'PHILA. Decandria Monogynia.				
1 <i>umbellátæ</i> , s.s. (<i>wh.re.</i>) umbelled.	N. America.	1752.	6. 8. H. $\frac{1}{2}$.	Bot. mag. t. 778.
<i>corymbosa</i> . Ph. <i>Pyrola umbellátæ</i> . L.			Z 3	

- 2 maculata. s.s. (*wh.*) spotted-leaved. N. America. 1752. 6. 8. H. ¼. Bot. mag. t. 897.
DIAPENIA. W. **DIAPENIA.** Pentandria Monogynia. L.
 lappónica. w. (*wh.*) obtuse-leaved. Lapland. 1801. 2. 3. H. ¼. Bot. mag. t. 1108.
PYXIDANTH'ERA. M. **PYXIDANTH'ERA.** Pentandria Monogynia.
 barbulata. m. (*wh.*) barbed. Carolina. 1806. 7. G. ¼. Mich. amer. t. 17.
Diapénzia cuneifolia. Sal. par. lond. t. 104.

Sect. II. **MONOTROPEÆ.** *Don prodr. p. 151.*

- MONOTROPA. N.** **YELLOW BIRD'S-NEST.** Decandria Monogynia. L.
 1 uniflora. n. (*st.*) one-flowered. N. America. 1822. 6. 8. H. ¼. Hook. ex. flor. t. 85.
 2 Hypopitys. E.B. (*st.*) common. Britain. 6. 7. H. ¼. Eng. bot. v. 1. t. 69.
Hypopithys europaea. N.

Sect. III. **RHODORACEÆ.** *Don prodr. p. 152.*

- KALMIA. W.** **KALMIA.** Decandria Monogynia. L.
 1 latifolia. w. (*pi. wh.*) broad-leaved. N. America. 1734. 5. 7. H. ½. Bot. mag. t. 175.
 2 angustifolia. w. (*re.*) narrow-leaved. 1736. — H. ½. — t. 331.
 β variegata. (*li.*) variegated. — — H. ½.
 γ minima. (*re.*) least. — — H. ½.
 δ nana. (*re.*) dwarf. — — H. ½.
 ϵ pumila. (*re.*) small. — — H. ½.
 ζ rosea. (*ro.*) rose-coloured. — — H. ½.
 η rubra. B.C. (*re.*) red-flowered. — — H. ½.
 ϑ ovata. Ph. (*re.*) oval-leaved. New Jersey. — — H. ½.
 3 glauca. w. (*re.*) glaucous. N. America. 1767. 4. 5. H. ½. Lodd. bot. cab. t. 502.
 4 rosmarinifolia. Ph. (*re.*) Rosemary-leaved. — 1812. — H. ½. Bot. mag. t. 177.
 5 hirsuta. w. (*ro.*) hairy. — 1786. 8. 9. H. ½. Bot. mag. t. 138.
EPIGAEA. W. **EPIGAEA.** Decandria Monogynia. L.
 repens. w. (*wh.*) creeping. N. America. 1736. 7. 8. H. ½. Bot. reg. t. 201.
RHODODENDRON. W. **RHODODENDRON.** Pent-Decandria Monogynia. L.
 1 maximum. w. (*bh.*) large. N. America. 1736. 6. 8. H. ½. Bot. mag. t. 951.
 2 album. Ph. (*wh.*) white-flowered. — 1811. — H. ½.
 3 purpureum. Ph. (*pu.*) purple-flowered. — — H. ½.
 4 ponticum. w. (*va.*) common. Gibraltar. 1763. 5. 6. H. ½. Bot. mag. t. 650.
 α purpureum. (*pu.*) purple. — — H. ½.
 β roseum. (*ro.*) rose-coloured. — — H. ½.
 γ album. (*wh.*) white. — — H. ½.
 δ variegatum. (*li.*) variegated. — — H. ½.
 ϵ marginatum. (*li.*) silver-edged. — — H. ½.
 ζ multiplex. (*pu.*) double-blossom'd. — — H. ½.
 5 punctatum. w. (*li.*) dotted-leaved. N. America. 1786. 6. 8. H. ½. Andr. deposit. t. 36.
 α minus. W.D. (*li.*) smallest. — — H. ½. Wats. dend. brit. t. 162.
 β major. B.R. (*li.*) larger. — 1812. — H. ½. Bot. reg. t. 37.
 6 obtusum. W.D. (*li.*) blunt-leaved. Hybrid. — H. ½. Wats. dend. brit. t. 162.
 7 azaleoides. s.s. (*li.*) Azalea-like. — — H. ½. Andr. deposit. t. 379.
 ponticum β subdeciduum. A.R. — — — — Andr. deposit. t. 36.
 8 hybridum. B.R. (*li.*) Herbert's hybrid. — H. ½. Bot. reg. t. 195.
 9 Goweniænum. B.F.G. (*pu.*) Mr. Gowen's. — 1825. — H. ½. Swt. br. fl. gar. t. 263.
 10 Smithii. Smith's. Arboreo-pont. 1826. H. ½.
 11 Carnarvoni Lord Carnarvon's. Arb-catawba. H. ½.
 12 catawbiense. M. (*ro.*) Catawba. N. America. 1809. 6. 8. H. ½.
 13 arboreum. Sm. (*sc.*) tree. India. 1817. 4. 6. F. ½. Bot. mag. t. 1671.
 β roseum. (*ro.*) rose-coloured. — — F. ½. Swt. br. fl. gar. t. 250.
 γ niveum. (*wh.*) snowy white. — — F. ½. Bot. reg. t. 1240.
 14 campanulatum. D.P. (*ro.*) bell-flowered. Nepaul. 1824. F. ½.
 15 anthopogon. D.P. (*ro.*) bearded-flower'd. — — G. ½.
 16 cinnamomeum. Wal. (*ro.*) cinnamon-color'd. — 1826. G. ½.
 17 setosum. D.P. (*pu.*) bristly. — 1820. G. ½.
 18 dauricum. w. (*pu.*) Daurian. Siberia. 1780. 2. 4. H. ½. Bot. mag. t. 636.
 β altaicum. f. (*li.*) Altay. — 1824. — H. ½. Bot. reg. t. 194.
 γ atrovirens. B.R. (*d. pu.*) dark green. — — H. ½.

19	<i>myrtifolium</i> . B.C. (<i>pu.</i>) Myrtle-leaved.	Hybrid.	6. 8. H. $\frac{1}{2}$.	Lodd. bot. cab. t. 908.
20	<i>frugiferum</i> . w. (<i>ro.</i>) rusty-leaved.	Switzerland.	1752.	5. 7. H. $\frac{1}{2}$.	Jacq. aust. 3. t. 255.
21	<i>hirsutum</i> . w. (<i>ro.</i>) hairy-leaved.		1656.	— H. $\frac{1}{2}$.	Bot. mag. t. 1853.
22	<i>chrysanthum</i> . w. (<i>ye.</i>) yellow. <i>officinale</i> . P.L.	Siberia.	1796.	6. 7. H. $\frac{1}{2}$.	Sal. par. lond. t. 80.
23	<i>caucasicum</i> . w. (<i>ro. wh.</i>) Caucasian.	Caucasus.	1803.	5. 7. H. $\frac{1}{2}$.	Bot. mag. t. 1145.
24	<i>Catesbeia</i> . L.C. (<i>ro.</i>) Catesby's.	N. America.	— H. $\frac{1}{2}$.	Pall. ros. 1. t. 33.
25	<i>camtschaticum</i> . w. (<i>pi.</i>) Kamtschatka.	Kamtschatka.	1802. H. $\frac{1}{2}$.	Bot. mag. t. 488.
26	<i>chamaecistus</i> . w. (<i>pi.</i>) Thyme-leaved.	Siberia.	1786.	5. 6. H. $\frac{1}{2}$.	Flor. dan. t. 966.
27	<i>lapponicum</i> . s.s. (<i>ro.</i>) Lapland.	Lapland.	1827. H. $\frac{1}{2}$.	
	<i>Azalea lapponica</i> . L.				
28	<i>Rhodora</i> . (<i>pi.</i>) Canadian. <i>Rhodora canadensis</i> . w.	Canada.	1767.	4. 5. H. $\frac{1}{2}$.	Bot. mag. t. 474.
29	<i>indicum</i> . (<i>va.</i>) Indian.	China.	1808.	3. 5. G. $\frac{1}{2}$.	
	<i>Azalea indica</i> . B.M.				
	α <i>punicum</i> . (<i>pi.</i>) red-flowered.	—	—	— G. $\frac{1}{2}$.	Bot. mag. t. 1480.
	β <i>phænicum</i> . (<i>pu.</i>) purple-flowered.	—	1824.	— G. $\frac{1}{2}$.	— t. 2667.
	γ <i>Smithii</i> . (<i>li.</i>) Smith's.	Hybrid.	1826.	4. 7. G. $\frac{1}{2}$.	
	δ <i>purpureum</i> . (<i>pu.</i>) double-purple.	China.	1819.	3. 5. G. $\frac{1}{2}$.	Bot. mag. t. 2509.
	ϵ <i>luteum</i> . (<i>ye.</i>) double yellow.	—	1826.	— G. $\frac{1}{2}$.	
	ζ <i>álbum</i> . (<i>wh.</i>) white-flowered.	—	1819.	3. 6. G. $\frac{1}{2}$.	Bot. reg. t. 811.
	<i>Azalea ledifolia</i> . Bot. mag. t. 2901.				
30	<i>sinense</i> . B.F.G. (<i>ye.</i>) Chinese.	—	1824.	— H. $\frac{1}{2}$.	
	α <i>flámmatum</i> . B.F.G. (<i>or.</i>) flame-coloured.	—	—	— H. $\frac{1}{2}$.	Lodd. bot. cab. t. 885.
	β <i>flavescens</i> . B.F.G. (<i>ye.</i>) pale yellow.	—	1826.	— H. $\frac{1}{2}$.	Swt. br. fl. gar. t. 290.
31	<i>luteum</i> . (<i>ye.</i>) yellow.	Turkey.	1793.	5. 6. H. $\frac{1}{2}$.	Bot. mag. t. 433.
	<i>Azalea póntica</i> . B.M.				
	β <i>cùpreum</i> . (<i>co.</i>) copper-coloured.	—	—	— H. $\frac{1}{2}$.	
	γ <i>glauicum</i> . (<i>pa.</i>) glaucous.	—	—	— H. $\frac{1}{2}$.	
	δ <i>pállidum</i> . (<i>pa.</i>) pale.	—	—	— H. $\frac{1}{2}$.	
	ϵ <i>albiflórum</i> . (<i>wh.</i>) white-flowered.	—	—	— H. $\frac{1}{2}$.	
32	<i>tricolor</i> . (<i>ye. ro. wh.</i>) three-coloured.	Hybrid.	1816.	— H. $\frac{1}{2}$.	Bot. mag. t. 2383.
33	<i>Mortéria</i> . B.F.G. (<i>fl. ye.</i>) Morter's.	—	1828.	— H. $\frac{1}{2}$.	Swt. br. fl. gar. s. 2. t. 10.
	α <i>cárneum</i> . (<i>fl.</i>) flesh-coloured.	—	—	— H. $\frac{1}{2}$.	
	β <i>préstans</i> . (<i>co. bh.</i>) handsome.	—	—	— H. $\frac{1}{2}$.	
34	<i>calendulàceum</i> . (<i>ye.</i>) yellow American. N. America.	1806.	—	— H. $\frac{1}{2}$.	
	<i>Azalea calendulacea</i> . M.				
	α <i>cròicum</i> . (<i>sn.</i>) saffron-coloured.	—	—	— H. $\frac{1}{2}$.	Bot. mag. t. 1721.
	β <i>chrysoléctum</i> . (<i>go.</i>) golden.	—	—	— H. $\frac{1}{2}$.	
	γ <i>grandiflòrum</i> . (<i>or.</i>) large-flowered.	—	—	— H. $\frac{1}{2}$.	
	δ <i>cùpreum</i> . (<i>co.</i>) copper-coloured.	—	—	— H. $\frac{1}{2}$.	
	ϵ <i>spléndens</i> . (<i>co. ye.</i>) shining.	—	—	— H. $\frac{1}{2}$.	
	ζ <i>flámmatum</i> . (<i>co. ye.</i>) flame-coloured.	—	1812.	— H. $\frac{1}{2}$.	Bot. reg. t. 145.
	η <i>triumphans</i> . (<i>co. ye.</i>) triumphant.	—	—	— H. $\frac{1}{2}$.	
	\varnothing <i>ignescens</i> . (<i>co. ye.</i>) fiery-flowered.	—	—	— H. $\frac{1}{2}$.	
35	<i>canescens</i> . M. (<i>ro.</i>) canescens.	—	1812.	— H. $\frac{1}{2}$.	
36	<i>speciosum</i> . (<i>va.</i>) large-calyxed.	—	—	— H. $\frac{1}{2}$.	
	α <i>májor</i> . (<i>sc.</i>) large scarlet.	—	—	— H. $\frac{1}{2}$.	
	β <i>crispum</i> . (<i>sc.</i>) curled.	—	—	— H. $\frac{1}{2}$.	
	γ <i>aurantium</i> . (<i>or.</i>) orange.	—	—	— H. $\frac{1}{2}$.	
	δ <i>obliquum</i> . (<i>re.</i>) oblique-leaved.	—	—	— H. $\frac{1}{2}$.	
	ϵ <i>prunifolium</i> . (<i>re.</i>) Plum-leaved.	—	—	— H. $\frac{1}{2}$.	
	ζ <i>cucullatum</i> . (<i>re.</i>) hollow-leaved.	—	—	— H. $\frac{1}{2}$.	
	η <i>revolutum</i> . (<i>re.</i>) revolute.	—	—	— H. $\frac{1}{2}$.	
	\varnothing <i>ciliatum</i> . (<i>re.</i>) fringed.	—	—	— H. $\frac{1}{2}$.	
	ι <i>acutifolium</i> . (<i>re.</i>) acute-leaved.	—	—	— H. $\frac{1}{2}$.	
	κ <i>undulatum</i> . (<i>re.</i>) wave-leaved.	—	—	— H. $\frac{1}{2}$.	
	λ <i>tortulifolium</i> . (<i>re.</i>) twisted-leaved.	—	—	— H. $\frac{1}{2}$.	
37	<i>nudiflòrum</i> . (<i>va.</i>) small calyxed.	—	1734.	— H. $\frac{1}{2}$.	
	<i>Azalea nudiflora</i> . W.				
	α <i>álbum</i> . (<i>wh.</i>) early white.	—	—	— H. $\frac{1}{2}$.	
	β <i>álbo-plènum</i> . (<i>wh.</i>) double white.	—	—	— H. $\frac{1}{2}$.	
	γ <i>blándum</i> . (<i>bh.</i>) blush-flowered.	—	—	— H. $\frac{1}{2}$.	
	δ <i>cárneum</i> . (<i>fl.</i>) pale red.	—	1734.	— H. $\frac{1}{2}$.	Bot. reg. t. 120.
	ϵ <i>carolinianum</i> . (<i>fl.</i>) Carolina.	—	—	— H. $\frac{1}{2}$.	Bot. mag. t. 180.
	ζ <i>coécineum</i> . (<i>sc.</i>) scarlet.	—	—	— H. $\frac{1}{2}$.	
	η <i>corymbosum</i> . (<i>wh.</i>) corymb-flowered.	—	—	— H. $\frac{1}{2}$.	
	\varnothing <i>crispum</i> . (<i>bh.</i>) curled.	—	—	— H. $\frac{1}{2}$.	

ERICEÆ.

ι <i>cumulatum.</i>	(wh.) bundled.	N.America.	5. 6. H. $\frac{1}{2}$.
κ <i>dicolor.</i>	(re.wh.) two-coloured.	H. $\frac{1}{2}$.
λ <i>fastigiatum.</i>	(bh.) crowded.	H. $\frac{1}{2}$.
μ <i>floridum.</i>	(bh.) abundant-flower'd.	H. $\frac{1}{2}$.
ν <i>globosum.</i>	(bh.) globose.	H. $\frac{1}{2}$.
ξ <i>glomeratum.</i>	(bh.) clustered.	H. $\frac{1}{2}$.
\circ <i>incanum.</i>	(wh.) hoary-leaved.	H. $\frac{1}{2}$.
π <i>incarnatum.</i>	(fl.) flesh-coloured.	H. $\frac{1}{2}$.
ρ <i>mirabile.</i>	(re.) wonderful.	H. $\frac{1}{2}$.
σ <i>montanum.</i>	(fl.) mountain.	H. $\frac{1}{2}$.
τ <i>pallidum.</i>	(li.wh.) pale-flowered.	H. $\frac{1}{2}$.
υ <i>paludosum.</i>	(li.wh.) marsh.	H. $\frac{1}{2}$.
ϕ <i>papilionaceum.</i>	(fl.) Butterfly.	H. $\frac{1}{2}$.
χ <i>partitum.</i>	(fl.wh.) five-parted.	H. $\frac{1}{2}$.
ψ <i>parviflorum.</i>	(bh.) small-flowered.	H. $\frac{1}{2}$.
ω <i>proliferum.</i>	(bh.) proliferous.	H. $\frac{1}{2}$.
$\alpha\alpha$ <i>pumilum.</i>	(bh.) dwarf.	H. $\frac{1}{2}$.
$\beta\beta$ <i>purpurascens.</i>	(fl.) purplish.	H. $\frac{1}{2}$.
$\gamma\gamma$ <i>purpureum.</i>	(pu.) purple.	H. $\frac{1}{2}$.
$\delta\delta$ <i>purpureo-plenum.</i>	(pu.) double purple.	H. $\frac{1}{2}$.
$\epsilon\epsilon$ <i>rösseum.</i>	(ro.) rosy.	H. $\frac{1}{2}$.
$\zeta\zeta$ <i>ruberrimum.</i>	(re.) dark-red.	H. $\frac{1}{2}$.
$\eta\eta$ <i>rubescens.</i>	(pi.) reddish.	H. $\frac{1}{2}$.
$\theta\theta$ <i>rubicundum.</i>	(re.) rubicund.	H. $\frac{1}{2}$.
$\iota\iota$ <i>ruberum. B.C.</i>	(re.) red-flowered.	H. $\frac{1}{2}$.
$\kappa\kappa$ <i>ruber-plenum.</i>	(re.) double-red.	H. $\frac{1}{2}$.
$\lambda\lambda$ <i>rufum.</i>	(re.) rufous.	H. $\frac{1}{2}$.
$\mu\mu$ <i>rutilus.</i>	(re.) deep red.	H. $\frac{1}{2}$.
$\nu\nu$ <i>semiduplex.</i>	(wh.) semidouble.	H. $\frac{1}{2}$.
$\xi\xi$ <i>stamineum.</i>	(wh.) long-stamened.	H. $\frac{1}{2}$.
oo <i>stellatum.</i>	(wh.) starry.	H. $\frac{1}{2}$.
$\pi\pi$ <i>tricolor.</i>	(wh.re.) three-coloured.	H. $\frac{1}{2}$.
$\rho\rho$ <i>variabile.</i>	(re.wh.) variable.	H. $\frac{1}{2}$.
$\sigma\sigma$ <i>variegatum.</i>	(std.) variegated.	H. $\frac{1}{2}$.
$\tau\tau$ <i>versicolor.</i>	(re.wh.) various-coloured.	H. $\frac{1}{2}$.
$\upsilon\upsilon$ <i>violaceum.</i>	(vi.) violet.	H. $\frac{1}{2}$.
38 <i>bicolor.</i>	(wh.re.) two-coloured.	1734.	H. $\frac{1}{2}$. Lodd. bot. cab. t. 51.
<i>Azalea bicolor.</i> Ph.				Trew. ehret. t. 48.
39 <i>viscidum.</i>	(wh.) viscid.	7. 8. H. $\frac{1}{2}$.	Meerb. ic. 2. t. 9.
<i>Azalea viscosa.</i> Ph.				
α <i>odoratum.</i>	(wh.) common white.	H. $\frac{1}{2}$.
β <i>crispum.</i>	(wh.) curled.	H. $\frac{1}{2}$.
γ <i>dealbatum.</i>	(wh.) powdered.	H. $\frac{1}{2}$.
δ <i>vittatum.</i>	(re.wh.) striped-flowered.	H. $\frac{1}{2}$.
ϵ <i>penicillatum.</i>	(re.wh.) pencilled.	H. $\frac{1}{2}$.
ζ <i>variegatum.</i>	(re.wh.) variegated.	H. $\frac{1}{2}$.
η <i>rubescens.</i>	(re.) reddish-flowered.	H. $\frac{1}{2}$.
θ <i>pubescens.</i>	(re.wh.) pubescent.	H. $\frac{1}{2}$.
ι <i>fissum.</i>	(re.wh.) narrow-petaled.	H. $\frac{1}{2}$.
40 <i>arborescens.</i>	(re.) arborescent.	1818. 5. 7. H. $\frac{1}{2}$.	Lodd. bot. cab. t. 441.
<i>Azalea arborescens.</i> Ph.				
41 <i>nítidum.</i>	(re.wh.) glossy-leaved.	1812. 6. 8. H. $\frac{1}{2}$.	Bot. reg. t. 414.
<i>Azalea nítida.</i> B.R.				
42 <i>glaucum.</i>	(wh.) glaucous dwarf.	1784. 6. H. $\frac{1}{2}$.	Wats. dend. brit. t. 5.
<i>Azalea glauca.</i> Ph.				
43 <i>hispidum.</i>	(wh.) tall glaucous.	6. 8. H. $\frac{1}{2}$.
<i>Azalea hispida.</i> Ph.				Wats. dend. brit. t. 6.
AZ'ALEA. D.P.	AZ'ALEA.	Pentandria Monogynia.	R.S.	
<i>procumbens. E.B. (ro.)</i>	trailing.	Britain. 4. 5. H. $\frac{1}{2}$.	Eng. bot. v. 13. t. 865.
<i>Loiseleuria procumbens.</i> R.S.	<i>Chamaelidon procumbens.</i> L.en.			
AMMYRS'INE. Ph.	AMMYRS'INE.	Decandria Monogynia.	Ph.	
1 <i>buxifolia.</i> Ph.	(wh.) upright.	N. America.	1736. 4. 5. H. $\frac{1}{2}$.	Bot. reg. t. 531.
	<i>Lédiophylloides buxifolium.</i> w.	<i>Leiophylloides thymifolium.</i> s.s.		
2 <i>Lyoni.</i>	(wh.) spreading.	Carolina.	1812. — H. $\frac{1}{2}$.	
L'EDUM. Ph.	L'EDUM.	Decandria Monogynia.	L.	
1 <i>palustre. w.</i>	(wh.) marsh.	Europe.	1762. 4. 5. H. $\frac{1}{2}$.	Flor. dan. t. 1031.
2 <i>decumbens. l.c.</i>	(wh.) decumbent.	Hudson's Bay.	— H. $\frac{1}{2}$.	
3 <i>canadense. l.c.</i>	(wh.) dwarf.	Canada.	1824. — H. $\frac{1}{2}$.	Lodd. bot. cab. t. 1049.
4 <i>latifolium. w.</i>	(wh.) broad-leaved.	N. America.	1763. — H. $\frac{1}{2}$.	Jacq. ic. 3. t. 464.

BEFARIA. L.	BEFARIA. Dodecandria Monogynia. L.					
racemosa. s.s. (bh.) sweet-scented.	Florida.	1810.	6. 7. G. h.	Vent. cels. t. 51.		
paniculata. M.	<i>Bejaria racemosa.</i> Ph.					
'ITEA. L.	'ITEA. Pentandria Monogynia. L.					
virginica. L.	(wh.) Virginian.	N. America.	1744.	6. 8. H. h.	Wats. dend. brit. t. 12.	

ORDO CXXVIII.

GALACINEÆ. *D. Don in Edinb. phil. journ. Dec. 1828. p. 53.*

G'ALAX. W.	G'ALAX. Pentandria Monogynia. W.					
aphylla. w.	(bh.) heart-leaved.	N. America.	1786.	6. 7. H. 4.	Bot. mag. t. 754.	
cordifolia. h.k.	<i>Blandfordia cordata.</i> Andr. rep. t. 343.					

ORDO CXXIX.

EPACRIDEÆ. *Brown prodr. I. p. 535.*

Sect I. EPACRIDEÆ VERÆ.

DRA'COPHY'LLUM. B.P.	DRACOPHY'LLUM. Pentandria Monogynia.					
1 secundum. b.p.	(wh.) side-flowering.	N. S. W.	1824.	4. 6. G. h.		
2 longifolium.r.s.	(wh.) long-leaved.	N. Zealand.		— G. h.		
SPHENO'TOMA. S.F.A.	SPHENO'TOMA.	Pentandria Monogynia.				
gracilis. s.f.a.	(wh.) slender.	N. Holland.	1825.	5. 10. G. h.	Fors.ch.gen.t.10.f.i-m.	
SPRENG'ELIA. Sm.	SPRENG'ELIA.	Pentandria Monogynia.				
incarnata. b.p.	(pi.) flesh-coloured.	N. S. W.	1793.	4. 6. G. h.	Swt. flor. aust. t. 44.	
RONCELETIA. B.P.	RONCELETIA.	Pentandria Monogynia.				
sprengelioides.b.p.(w.)marsh.		N. S. W.	1826.	4. 6. G. h.	Bot. mag. t. 1719.	
ANDERSONIA. B.P.	ANDERS'ONIA.	Pentandria Monogynia.				
sprengelioides.b.p.(w.)Sprengelia-like.		N. Holland.	1803.	3. 7. G. h.	Bot. mag. t. 1645.	
COS'MELIA. B.P.	COSM'ELIA.	Pentandria Monogynia.				
rubra. b.p.	(re.) red-flowered.	N. Holland.	1826.	4. 6. G. h.		
LYSIN'EMA. B.P.	LYSIN'EMA.	Pentandria Monogynia.				
1 pentapetalum.b.p.(w.)five-petaled.		N. Holland.	1826.	4. 6. G. h.		
2 conspicuum. b.p.(wh.)conspicuous.			1824.	— G. h.		
3 pungens. b.p.	(wh.) pungent.	N. S. W.	1804.	3. 6. G. h.	Cav. ic. 4. t. 346.	
Epácris attenuata. Lodd. bot. cab. t. 38.						
4 roseum. b.c.	(ro.) rose-coloured.		1804.	— G. h.	Lodd. bot. cab. t. 863.	
Epácris rosea. b.c. pungens. Bot. mag. t. 1199.						
EPACRIS. B.P.	EPACRIS.	Pentandria Monogynia.				
1 purpurascens.b.p.(li.)rigid-leaved.		N. S. W.	1803.	2. 6. G. h.	Bot. mag. t. 844.	
pungens. B.M.						
α lilacina.	(li.) pale-flowered.			— G. h.	Lodd. bot. cab. t. 237.	
β rubra.	(re.) red-flowered.			— G. h.	— t. 876.	
2 pulchella. B.P.	(wh.) sweet-scented.		1804.	3. 6. G. h.	— t. 170.	
3 microphylla.B.P.(wh.)small-leaved.			1817.	— G. h.		
4 grandiflora. B.P.	(cr.) crimson.		1803.	1. 6. G. h.	Bot. mag. t. 982.	
5 ruscifolia. B.P.	(wh.) Ruscus-leaved.	V. Diem. Isl.	1824.	3. 5. G. h.		
6 impressa. B.P.	(ro.) impressed.			2. 5. G. h.	Swt. flor. aust. t. 4.	
7 sparsa. B.P.	(wh.) scattered-flower'd.	N. S. W.	1825.	3. 5. G. h.		
8 obtusifolia. B.P.	(wh.) blunt-leaved.		1804.	4. 6. G. h.	Sm. exot. bot. I. t. 40.	
9 heteronema. B.P.(wh.)hairy-branched.		V. Diem. Isl.	1824.	— G. h.	Labill. n. hol. I. t. 56.	
10 paludosa. B.P.	(wh.) marsh.	N. S. W.		— G. h.	Lodd. bot. cab. t. 1226.	
11 apiculata. F.T.	(wh.) pointed.		1823.	5. 8. G. h.		
12 onosmæflora.F.T.	(re.) Onosma-flowered.			— G. h.		
13 exserta. B.P.	(wh.) exserted.	V. Diem. Isl.	1812.	4. 6. G. h.		
14 mucronulata.B.P.(wh.)short-pointed.			1824.	— G. h.		

Sect. II. STYPHELIÆ.

TROCHOCÀ'RPA. B.P.	TROCHOCÀ'RPA.	Pentandria Monogynia.				
laurina. B.P.	(wh.) Laurel-leaved.	N. S. W.	1820.	G. h.	Linn. trans. 8. t. 9.
<i>Styphèlia cornifòlia</i> . L.T.						
ACRO'TRICHE. B.P.	ACRO'TRICHE.	Pentandria Monogynia.				
1 cordàta. B.P.	(wh.) heart-leaved.	N. Holland.	1825.	4. 6.	G. h.	Labill. n. hol. 1. t. 63.
2 ovalifòlia. B.P.	(wh.) oval-leaved.		1824.	—	G. h.	
3 serrulàta. B.P.	(wh.) saw-leaved.	V. Diem. Isl.	1816.	—	G. h.	Labill. n. hol. 1. t. 62.
<i>Styphèlia serrulàta</i> . Labill.						
4 divaricàta. B.P.	(wh.) divaricate.	N. S. W.	1822.	—	G. h.	
MONO'TOCA. B.P.	MONO'TOCA.	Pentandria Monogynia.				
1 scopària. B.P.	(wh.) broom.	N. S. W.	1823.	4. 7.	G. h.	Smith new hol. 43.
<i>Styphèlia scopària</i> . Smith.						
2 lineàta. B.P.	(wh.) lineated.	V. Diem. Isl.	1804.	5. 8.	G. h.	Lab. nov. hol. 1. t. 61.
<i>Styphèlia glauca</i> . Labill.						
3 álbens. B.P.	(wh.) white-leaved.	N. S. W.	1823.	—	G. h.	
4 elliptica. B.P.	(wh.) elliptic-leaved.		1802.	—	G. h.	Smith new hol. 49.
LEUCOP'OGON. B.P.	LEUCOP'OGON.	Pentandria Monogynia.				
1 lanceolàtus. B.P.	(wh.) small-flowered.	N. S. W.	1790.	5. 8.	G. h.	Swt. flor. aust. t. 47.
<i>Styphèlia parviflora</i> . Andr. deposit. t. 287. <i>gnidium</i> . Vent. malm. t. 23.						
2 Richéi. B.P.	(wh.) oblong-leaved.	N. S. W.	1824.	—	G. h.	Lab. nov. hol. 1. t. 60.
3 müticus. B.P.	(wh.) awnless.		1826.	—	G. h.	
4 ericoides. B.P.	(wh.) heath-leaved.		1815.	4. 7.	G. h.	Smith new hol. 48.
<i>Epácris spuria</i> . Cavan. ic. 4. t. 347. f. 1.						
5 virgátus. B.P.	(wh.) twiggy.			—	G. h.	Lab. nov. hol. 1. t. 64.
6 collinus. B.P.	(wh.) hill.	V. Diem. Isl.	1822.	5. 6.	G. h.	— 1. t. 65.
7 amplexicaùlis. B.P.	(w.) stem-clasping.	N. S. W.	1815.	4. 8.	G. h.	Linn. trans. 8. t. 8.
8 microphyllus. B.P.	(w.) small-leaved.		1818.	—	G. h.	Cavan. ic. 4. t. 349. f. 2.
<i>Perajea microphylla</i> . Cav.						
9 striátus. B.P.	(wh.) striated.	N. Holland.	1824.	—	G. h.	
10 attenuátus. F.T.	(wh.) attenuated.	N. S. W.	—	—	G. h.	
11 bifòrûs. B.P.	(wh.) two-flowered.		1826.	—	G. h.	
12 setíger. B.P.	(wh.) bristle-pointed.		1824.	—	G. h.	
13 apprèssus. B.P.	(wh.) pressed-leaved.			4. 6.	G. h.	
14 juniperinus. B.P.	(wh.) Juniper-leaved.		1804.	—	G. h.	
15 interrúptus. B.P.	(wh.) interrupted.	N. Holland.	1825.	3. 6.	G. h.	
16 polystachyus. B.P.	(wh.) many-spiked.			6. 7.	G. h.	
LISSA'NTHE. B.P.	LISSA'NTHE.	Pentandria Monogynia.				
1 sápida. B.P.	(wh.) recurved-flowered.	N. S. W.	1823.	11. 1.	G. h.	Lodd. bot. eab. t. 447.
2 subulàta. B.P.	(wh.) awl-leaved.			4. 7.	G. h.	— t. 1451.
3 strigósa. B.P.	(wh.) strigose.			—	G. h.	— t. 1436.
4 daphnoïdes. B.P.	(wh.) Daphne-leaved.	V. Diem. Isl.	1818.	6. 7.	G. h.	
5 ciliàta. B.P.	(wh.) fringed.		1823.	—	G. h.	
CYATH'ODES. B.P.	CYATH'ODES.	Pentandria Monogynia.				
1 glauca. B.P.	(wh.) glaucous.	V. Diem. Isl.	1818.	4. 6.	G. h.	Bot. reg. t. 1275.
2 oxycedrus. B.P.	(wh.) linear-leaved.		1822.	5. 8.	G. h.	Sm. new hol. 48.
3 abietina. B.P.	(wh.) Fir-like.		1824.	8. 12.	G. h.	Lodd. bot. cab. t. 466.
MELI'CHRUS. B.P.	MELI'CHRUS.	Pentandria Monogynia.				
1 rotátus. B.P.	(re.) rotate-flowered.	N. S. W.	1822.	5. 8.	G. h.	Lab. nov. hol. 1. t. 81.
<i>Venentaria procumbens</i> . Cav.						— 1. t. 69.
2 mèdius. F.T.	(re.) intermediate.		1823.	—	G. h.	— 1. t. 68.
STENANTH'ERA. B.P.	STENANTH'ERA.	Pentandria Monogynia.				
pinifòlia. B.P.	(sc.) Pine-leaved.	N. S. W.	1811.	5. 7.	G. h.	
ASTROL'OMA. B.P.	ASTROL'OMA.	Pentandria Monogynia.				
1 humifùsum. B.P.	(sc.) Juniper-leaved.	N. S. W.	1807.	5. 10.	G. h.	Bot. reg. t. 218.
2 denticulàtum. B.P.	(re.) toothed.		1825.	—	G. h.	Bot. mag. t. 1439.
STYPH'ELIA. B.P.	STYPH'ELIA.	Pentandria Monogynia.				
1 tubiflòra. B.P.	(cr.) crimson.	N. S. W.	1802.	5. 8.	G. h.	Sm. new hol. 45. t. 14.
2 triflòra. B.P.	(cr.gr.) three-flowered.		1796.	—	G. h.	Bot. mag. t. 1297.
3 viridiflòra. B.P.	(gr.) green-flowered.		1791.	4. 6.	G. h.	Swt. flor. aust. t. 50.
4 latifòlia. B.P.	(gr.) broad-leaved.		1823.	—	G. h.	
5 adscéndens. B.P.	(gr.) ascending.	V. Diem. Isl.	1822.	—	G. h.	
6 læta. B.P.	(gr.) fruitful.	N. S. W.	—	—	G. h.	
7 longifòlia. B.P.	(gr.) long-leaved.		1807.	—	G. h.	Bot. reg. t. 24.

ORDO CXXX.

PENÆACEÆ.

PENÆA. R.S.	PENÆA. Tetrandria Monogynia. R.S.	C. B. S.	1816.	5. 7.	G. h.	
1 myrtoides. w.	(ye.) Myrtle-like.	1822.	—	G. h.		
2 fruticulosa. R.S.	(re.) blunt-leaved.	1787.	—	G. h.	Vent. malm. t. 87.	
3 mucronata. R.S.	(re.) sharp-pointed.	1823.	4. 6.	G. h.	Bot. reg. t. 106.	
4 squamosa. R.S.	(re.) scaly.	1824.	—	G. h.	Bot. mag. t. 2809.	
5 imbricata. B.M.	(re.) imbricate.	1816.	5. 7.	G. h.	Th.m.n.c.ber.1.t.3.f.1.	
6 acuta. R.S.	(re.) acute-leaved.	—	—	G. h.		
7 Sarcocolla. R.S.	(ye.) rhomb-leaved.	—	—	G. h.		
8 marginata. R.S.	(wh.) thick-edged.	—	—	G. h.		

SUBCLASSIS III. COROLLIFLORÆ. DC.

ORDO CXXXI.

SYMPLOCINEÆ. *Don Prodr. p. 144.*

SYMPLOCOS. L'H.	SYMPLOCOS. Polyadelphia Polyandria. L.				
1 coccinea. K.S.	(sc.) scarlet.	Mexico.	1825.	S. h.
2 tinctoria. w.	(ye.) Laurel-leaved.	Carolina.	1780.	4. 6.	F. h.
	<i>Höpea tinctoria. Ph.</i>				Catesb. car. 1. t. 54.
3 sinica. B.R.	(wh.) Chinese.	China.	1822.	4. 6.	F. h.
4 crataegoides. D.P.	(wh.) Hawthorn-like.	Nepaul.	1824.	G. h.
SCHO'EPFIA. D.P.	SCHO'EPFIA. Pentandria Monogynia.				
frâgrans. D.P.	(ye.) fragrant.	Nepaul.	1827.	G. h.

ORDO CXXXII.

STYRACINEÆ. *Kth. synops. 2. p. 315.*

STYRAX. W.	STORAX. Decandria Monogynia. L.				
1 officinale. w.	(wh.) officinal.	Italy.	1597.	7.	H. h.
2 grandifolium. w.	(wh.) great-leaved.	N. America.	1765.	—	H. h.
3 pulverulentum. M.	(wh.) powdered.		1794.	6. 7.	H. h.
4 glabrum. N.	(wh.) smooth.		1765.	7. 8.	H. h.
	<i>lävigatum. Bot. mag. t. 921.</i>				Andr. deposit. t. 631.
HAL'ESIA. W.	SNOWDROP-TREE. Dodecandria Monogynia. L.				Wats. dend. brit. t. 129.
1 tétraptera. w.	(wh.) four-winged.	Carolina.	1756.	4. 5.	H. h.
2 parviflora. B.R.	(wh.) small-flowered.		1812.	5.	H. h.
3 diptera. w.	(wh.) two-winged.	N. America.	1758.	4. 5.	H. h.

— t. 41.
— t. 40.

Bot. mag. t. 910.
Bot. reg. t. 952.
Lodd. bot. cab. t. 1172.

ORDO CXXXIII.

MYRSINEÆ. *Brown prodr. 532.*

MÆSA. J.	MÆSA. Pentandria Monogynia.				
1 indica. D.P.	(wh.) Indian.	E. Indies.	1812.	2. 6.	S. h.
	<i>Bæobætris indica. F.I.</i>				Bot. mag. t. 2052.
2 tomentosa. D.P.	(wh.) tomentose.	Nepaul.	1818.	—	S. h.
3 argentea. F.I.	(wh.) silvery-leaved.	E. Indies.	—	—	S. h.
4 macrophylla. F.I.	(wh.) large-leaved.		1823.	—	S. h.

JACQUI'NIA. W.	JACQUI'NIA.	Pentandria Monogynia. W.
1 arbórea. w. (wh.) blunt-leaved.	W. Indies.	1820. S. h.
2 armilláris. w. (wh.) armed-leaved.	——	1768. 4. 6. S. h.
3 aurantiaca. H.K. (or.) orange-color'd.	Sandwich Isl.	1796. 4. 9. S. h.
4 ruscifolia. w. (or.) Ruscus-leaved.	S. America.	1729. — S. h.
5 macrocárpa. c.i. (or.) large capsuled.	——	1820. — S. h.
ARDI'SIA. W.	ARDI'SIA.	Pentandria Monogynia. W.
1 acuminata. w. (pi.) acuminate.	Guiana.	1803. 7. 8. S. h.
<i>Myrsine Icacorea. R.S.</i>		
2 tiniifolia. R.S. (pi.)	Tinus-leaved. Jamaica.	1824. — S. h.
3 solanacea. w. (ro.)	Nightshade-like. E. Indies.	1798. 6. 9. S. h.
4 paniculata. F.I. (ro.)	panicked.	1818. 3. 8. S. h.
5 colorata. F.I. (ro.)	red-flowered.	1816. — S. h.
6 umbellata. F.I. (pi.)	umbel-flowered.	1809. 6. 9. S. h.
<i>littoralis.</i> Andr. repos. t. 630.		
7 lanceolata. F.I. (pi.)	leathery-leaved.	1818. — S. h.
8 élégans. A.R. (pi.)	elegant.	1809. 7. 9. S. h.
<i>crenata. F.I.</i>		
9 punctata. B.R. (pi.)	spotted-flowered. China.	1823. 6. 9. S. h.
10 lentiginosa. B.R. (wh.)	freckled.	1809. 3. 12. G. h.
<i>crenulata. B.C. 2. non Vent. crenata.</i> Bot. mag. t. 1950.		
11 glandulosa. F.I. (wh.)	glandular. E. Indies.	1827. — S. h.
12 macrocarpa. F.I. (pi.)	large-fruited. Nepaul.	1824. 5. 9. S. h.
13 complanata. F.I. (pi.)	pubescent. Pinang.	—— — S. h.
14 thyrsiflora. D.P. (pi.)	thyrsse-flower'd. Nepaul.	—— 4. 8. S. h.
15 pyramidalis. Cav. (pi.)	pyramidal. Santa Cruz.	1818. 4. 9. S. h.
16 excelsa. w. (gr.)	Laurel-leaved. Madeira.	1784. 7. 8. G. h.
<i>Myrsine Heberdenii. R.S.</i>		
17 coriacea. R.S. (ro.)	leathery-leaved. W. Indies.	1822. 4. 8. S. h.
18 serrulata. w. (pi.)	saw-leaved.	1820. 7. 8. S. h.
19 crenulata. w. (re.)	crenulated.	1809. 6. 9. S. h.
20 lateriflora. w. (pi.)	side-flowering.	1793. — S. h.
21 canaliculata. w. (pi.)	channeled.	NewGranada 1824. 4. 8. S. h.
EMBELIA. L.	EMBELIA.	Pentandria Monogynia. L.
1 robusta. F.I. (gr.wh.)	robust. E. Indies.	1823. S. h.
2 floribunda. F.I. (ye.gr.)	sweet-scented. Nepaul.	1826. S. h.
MYRS'INE. B.P.	MYRS'INE.	Polygamia Dicecia. L.
1 africana. s.s. (gr.pu.)	African. C. B. S.	1691. 3. 5. G. h.
2 retusa. s.s. (gr.pu.)	retuse-leaved. Azores.	1778. 6. G. h.
3 rotundifolia. R.S. (w.pu.)	round-leaved.	C. B. S. 1816. — G. h.
4 semiserrata. F.I. (bh.)	semiserrate. Nepaul.	1822. G. h.
5 capitellata. F.I. (gr.)	axillary headed.	—— — G. h.
6 bifaria. F.I. (bh.)	bifarious.	—— — G. h.
7 subspinosa. D.P. (wh.)	spiny-leaved.	1823. G. h.
8 coriacea. B.P. (wh.)	leathery-leaved. Jamaica.	1822. S. h.
<i>Samára coriacea. Sw.</i>		
9 Samára. s.s. (wh.)	oval-leaved.	C. B. S. 1770. 11. 2. G. h.
<i>Samára pentandra. w.</i>		
10 melanophleos. s.s. (wh.)	Laurel-leaved.	—— 1783. 3. 6. G. h.
<i>Sideróxylon melanóphleum. w.</i>		
11 mitis. s.s. (wh.)	pointed-leaved.	—— 6. 9. G. h.
<i>Sideróxylon mítē. B.M. Scleróxylum mítē. W.en. Mangúlla Milleriána. P.S.</i>		
12 venulosa. s.s. (wh.)	veined-leaved.	1816. — G. h.
13 canariense. s.s. (wh.)	Canary.	—— — G. h.
14 variabilis. B.P. (wh.)	variable.	N. S. W. 1820. — G. h.
ÆGIC'ERAS. G.	ÆGIC'ERAS.	Pentandria Monogynia.
frágrans. B.P. (wh.)	fragrant.	N. Holland. 1824. G. h.
		Kon.an.bot.1.p.129.ic.

ORDO CXXXIV.

SAPOTEÆ. Brown prodr. 1. p. 528.

INOCA'RPUS. W.	OTAHEITE-CHESNUT.	Decandria Monogynia. W.
édulis. w. (wh.)	eatable.	South-sea Isl. 1793. S. h.
BUM'ELIA. S.S.	BUM'ELIA.	Pentandria Monogynia. W.
1 lycioídes. s.s. (wh.)	Boxthorn-leav'd.N.America.	1758. 8. H. h.
		Duham. arb. 2. t. 68.

2	<i>oblongifolia</i> . N.	(<i>wh.</i>) oblong-leaved.	Mississippi.	1818.	7. 8. H. $\frac{1}{2}$.		
3	<i>tēnax</i> . R.S.	(<i>wh.</i>) silvery-leaved.	Carolina.	1765.	— — H. $\frac{1}{2}$.		<i>Wats. dend. brit. t. 10.</i>
	<i>chrysophylloides</i> . W.D.	<i>Sideroxylon tēnax</i> . L.					
4	<i>lanuginōsa</i> . R.S.	(<i>wh.</i>) woolly-leaved.	— —	— —	H. $\frac{1}{2}$.		
5	<i>reclināta</i> . s.s.	(<i>wh.</i>) reclined.	— —	1806.	— — H. $\frac{1}{2}$.		<i>Venten. choix. t. 22.</i>
6	<i>strigōsa</i> . s.s.	(<i>wh.</i>) strigose.	— —	— —	G. $\frac{1}{2}$.		
		<i>Sideroxylon strigosum</i> . W.en.					
7	<i>nīgra</i> . s.s.	(<i>wh.</i>) black.	W. Indies.	1806.	6. 8. S. $\frac{1}{2}$.		
8	<i>salicifolia</i> . s.s.	(<i>wh.</i>) Willow-leaved.	S. America.	1758.	— — S. $\frac{1}{2}$.		<i>Sloan. hist. 2. t. 206. f. 2.</i>
9	<i>fetidissima</i> . s.s. (<i>wh.</i>) fetid.		W. Indies.	— —	5. 7. S. $\frac{1}{2}$.		
10	<i>rotundifolia</i> . s.s. (<i>wh.</i>) round-leaved.		— —	1816.	— — S. $\frac{1}{2}$.		
SERSAL'SIA . B.P.	SERSAL'SIA .				Pentandria Monogynia.		
<i>sericea</i> . B.P.	(<i>wh.</i>) silky-leaved.		N. Holland.	1772.	5. 6. G. $\frac{1}{2}$.		
	<i>Sideroxylon sericeum</i> . w.						
SIDEROXYLON . W. IRON-WOOD.					Pentandria Monogynia. L.		
1	<i>inérme</i> . w.	(<i>wh.</i>) smooth.	C. B. S.	1692.	7. 8. G. $\frac{1}{2}$.		<i>Lam. ill. 2. t. 120. f. 1.</i>
2	<i>tomentosum</i> . w. (<i>wh.</i>) tomentose.		E. Indies.	1818.	— — S. $\frac{1}{2}$.		<i>Roxb. corom. I. t. 28.</i>
ARG'ANIA . R.S.	ARG'ANIA .				Pentandria Monogynia.		
	<i>Sideroxylum</i> . R.S. (<i>ye.</i>) spiny.		Morocco.	1711.	7. 8. H $\frac{1}{2}$.		
	<i>Sideroxylum spinosum</i> . L. <i>Elæodendron A'rgan</i> . w.						<i>Comm. hort. I. t. 83.</i>
CHRYSOPHYLLUM . W. STAR-APPLE.					Pentandria Monogynia. L.		
1	<i>Cainito</i> . w.	(<i>wh.</i>) broad-leaved.	W. Indies.	1737.	5. 6. S. $\frac{1}{2}$.		
β	<i>jamaicēnse</i> .	(<i>wh.</i>) purple-fruited.	— —	— —	S. $\frac{1}{2}$.		<i>Jacq. amer. t. 37. f. 1.</i>
γ	<i>cæruleum</i> .	(<i>wh.</i>) blue-fruited.	— —	— —	S. $\frac{1}{2}$.		<i>Jacq. amer. pict. t. 52.</i>
δ	<i>microphyllum</i> .	(<i>wh.</i>) small-leaved.	— —	— —	S. $\frac{1}{2}$.		<i>Jacq. amer. t. 37.</i>
2	<i>monopyrenum</i> . w. (<i>wh.</i>) acute-leaved.		— —	1816.	— — S. $\frac{1}{2}$.		<i>Jacq. amer. pict. t. 53.</i>
3	<i>argéntea</i> . w. (<i>wh.</i>) silvery.		Martinico.	1758.	— — S. $\frac{1}{2}$.		<i>Burm. amer. t. 69.</i>
4	<i>macrophyllum</i> . G.D. (<i>wh.</i>) large-leaved.		Sierra Leone.	1822.	— — S. $\frac{1}{2}$.		<i>Jacq. amer. pict. t. 51.</i>
5	<i>obovatum</i> . G.D. (<i>wh.</i>) obovate.		— —	— —	— — S. $\frac{1}{2}$.		
6	<i>glabrum</i> . w. (<i>wh.</i>) smooth-leaved.		Martinico.	1818.	4. 6. S. $\frac{1}{2}$.		<i>Jacq. amer. pict. t. 55.</i>
NYCTERISITION . R.P. <i>Nycterisition</i> .					Pentandria Monogynia.		
	<i>ferrugineum</i> . R.P. (<i>wh.</i>) ferruginous.		S. America.	1823.	— — S. $\frac{1}{2}$.		<i>Flor. peruv. 2. t. 187.</i>
A'CHRAS . J.	SAPOTA .				Hexandria Monogynia. S.S.		
1	<i>Sapota</i> . w.	(<i>wh.</i>) common.	S. America.	1731.	4. 6. S. $\frac{1}{2}$.		
β	<i>Zapotilla</i> . Jacq. (<i>wh.</i>) <i>Naseberry-tree</i> .		— —	— —	S. $\frac{1}{2}$.		<i>Jacq. amer. 57. t. 41.</i>
2	<i>australis</i> . B.P.	(<i>wh.</i>) New Holland.	N. S. W.	1823.	— — G. $\frac{1}{2}$.		<i>—— 57. t. 41. b.</i>
LUC'UMA . J.	LUC'UMA .				Pentandria Monogynia.		
1	<i>mammosum</i> . K.S.	(<i>wh.</i>) Mamme.	S. America.	1739.	5. 8. S. $\frac{1}{2}$.		<i>Jacq. amer. t. 182. f. 19.</i>
2	<i>Bonplāndii</i> . K.S.	(<i>wh.</i>) Bonpland's.	Havannah.	1822.	— — S. $\frac{1}{2}$.		
3	<i>obovatum</i> . K.S.	(<i>wh.</i>) obovate-leaved.	Peru.	— —	— — S. $\frac{1}{2}$.		<i>Flor. peruv. 3. t. 239.</i>
4	<i>salicifolium</i> . K.S. (<i>wh.</i>)	Willow-leaved.	Mexico.	1823.	— — S. $\frac{1}{2}$.		
MI'MUSOPS . W.	MI'MUSOPS .				Octandria Monogynia. L.		
1	<i>Eléngi</i> . s.s.	(<i>wh.</i>) pointed-leaved.	E. Indies.	1796.	5. 7. S. $\frac{1}{2}$.		<i>Roxb. cor. I. t. 14.</i>
2	<i>dissécta</i> . s.s.	(<i>wh.</i>) oval-leaved.	South-sea Isl.	1804.	— — S. $\frac{1}{2}$.		
3	<i>hexándra</i> . s.s.	(<i>wh.</i>) hexandrous.	E. Indies.	1816.	— — S. $\frac{1}{2}$.		<i>Roxb. cor. I. t. 15.</i>
4	<i>Kaūki</i> . s.s.	(<i>wh.</i>) obtuse-leaved.	— —	1796.	5. 8. S. $\frac{1}{2}$.		<i>Rumph. amb. 3. t. 8.</i>
IMBRIC'ARIA . J.	IMBRIC'ARIA .				Octandria Monogynia.		
	<i>borbonica</i> . g.	(<i>wh.</i>) Bourbon.	Bourbon.	1820.	— — S. $\frac{1}{2}$.		
BA'SSIA . L.	BA'SSIA .				Dodecadria Monogynia. L.		
1	<i>longifolia</i> . w.	(<i>wh.</i>) long-leaved.	E. Indies.	1811.	— — S. $\frac{1}{2}$.		<i>Lam. ill. t. 398.</i>
2	<i>latifolia</i> . w.	(<i>wh.</i>) broad-leaved.	— —	1799.	— — S. $\frac{1}{2}$.		<i>Roxb. cor. I. t. 19.</i>
3	<i>butyracea</i> . D.P.	(<i>wh.</i>) butter-tree.	Nepaul.	1823.	— — S. $\frac{1}{2}$.		

ORDO CXXXV.

EBENACEÆ. *Brown prodr. I. p. 524.*

MABA . J.	MABA .	<i>Diœcia Hexandria</i> . W.					
	<i>buxifolia</i> . P.S.	(<i>wh.</i>) Box-leaved.	E. Indies.	1810.	— — S. $\frac{1}{2}$.		<i>Roxb. cor. I. t. 45.</i>
	<i>Ferreola buxifolia</i> . Roxb.						
CARGIL'LLIA . B.P.	CARGIL'LLIA .	<i>Polygamia Diœcia</i> .					
	<i>australis</i> . B.P.	(gr.) southern.	N. S. W.	1816.	— — G. $\frac{1}{2}$.		
DIO'SPYROS . S.S.	DATE PLUM .	<i>Polygamia Dicecia</i> . W.					
1	<i>virginiana</i> . s.s. (<i>p.ye.</i>)	Persimon.	N. America.	1629.	6. 7. H. $\frac{1}{2}$.		<i>Wats. dend. brit. t. 146.</i>
2	<i>pubescens</i> . s.s. (<i>p.ye.</i>)	pubescent.	— —	1812.	4. 5. H. $\frac{1}{2}$.		

EBENACEÆ.

3 reticulàta. s.s. (wh.) netted.	Mauritius.	1823.	4. 5. S. h.
4 Ebenáster. s.s. (wh.) Ebenus-like.	E. Indies.	1792.	6. 8. S. h.
5 melanóxylon. s.s. (wh.) black-wooded.	_____	1817.	____ S. h.
6 E'benum. s.s. (wh.) smooth.	_____	1792.	____ S. h.
7 montâna. s.s. (wh.) mountain.	_____	1819.	____ S. h.
8 obovata. s.s. (wh.) four-seeded.	W. Indies.	1796.	5. 7. S. h.
9 sylvática. s.s. (wh.) wood.	E. Indies.	1812.	____ S. h.
10 Embryópteris. s.s. (gr.) glutiniferous.	_____	1796.	5. 8. S. h.
Embryópteris glutinifera. B.C. 1. t. 70.			
11 lobata. s.s. (wh.) lobed-fruited.	China.	1822.	____ G. h.
12 vaccinioides. L.C. (wh.) Vaccinium-like.	_____	____	____ G. h.
13 édulis. L.C. (wh.) eatable.	E. Indies.	_____	____ S. h.
14 Lótus. s.s. (pa.ye.) European.	Italy.	1596.	6. 7. H. h.
15 Kákì. s.s. (wh.) Japan.	Japan.	1789.	____ G. h.
16 hirsùta. s.s. (wh.) hairy.	Ceylon.	1816.	____ S. h.
17 chloróxylon. s.s. (wh.) cluster-flowered.	E. Indies.	1822.	____ S. h.
18 cordifolia. s.s. (wh.) heart-leaved.	_____	1794.	5. 7. S. h.
ROY'ENA. W.	ROY'ENA. Decandria Dignynia. L.		
1 lúcida. w. (wh.) shining-leaved.	C. B. S.	1690.	5. 6. G. h.
2 villòsa. w. (wh.) heart-leaved.	_____	1774.	6. 7. G. h.
3 pállens. w. (wh.) pale.	_____	1789.	____ G. h.
4 myrtifolia. W.en. (wh.) Myrtle-leaved.	_____	1812.	____ G. h.
5 glábra. w. (wh.) smooth.	_____	1731.	7. 9. G. h.
6 pubescens. W.en. (wh.) pubescent.	_____	1752.	____ G. h.
7 hirsùta. W.en. (wh.) hairy-leaved.	_____	1789.	____ G. h.
8 angustifolia. w. (wh.) willow-leaved.	_____	1789.	6. 7. G. h.
9 latifolia. W.en. (wh.) broad-leaved.	_____	1816.	____ G. h.
10 ambigua. v. (wh.) obovate-leaved.	_____	1815.	6. 7. G. h.
11 polyandra. w. (wh.) oval-leaved.	_____	1774.	____ G. h.
VI'SNEA. W.	VI'SNEA. Dodecandria Trigynia. W.		
Mocanèra. w. (wh.) Canary.	Canaries.	1815.	8. 12. G. h.

ORDO CXXXVI.

OLEINÆ. Brown prodr. 1. p. 522.

FRA'XINUS. W.

	ASH-TREE. Polygamia Diœcia. L.			
1 excélsior. w.	(gr.) common.	Britain.	4. 5. H. 5.
β argéntea.	(gr.) silver-striped.	_____	____ H. 5.
γ péndula.	(gr.) weeping.	_____	____ H. 5.
δ horizontális.	(gr.) horizontal.	_____	____ H. 5.
ε jáspea.	(gr.) coloured branch'd.	_____	____ H. 5.
ζ eròsa.	(gr.) erose-leaved.	_____	____ H. 5.
2 àurea. W.en.	(gr.) yellow barked.	_____	____ H. 5.
3 atrovirens. P.S.	(gr.) greencurled-leav'd.	_____	____ H. 5.
4 nàna. W.en.	(gr.) dwarf.	_____	____ H. h.
5 verrucosa. L.en.	(gr.) warted.	_____	4. 5. H. 5.
6 heterophyl'a. s.s.(gr.)	various-leaved.	England.	____ H. 5.
simplicifolia. E.B.				
7 polemonifolia. R.S. (gr.)	Polemonium-ld.	N. America.	____ H. 5.
8 parvifolia. s.s.	(gr.) small-leaved.	Levant.	1816.	____ H. 5.
9 lentiscifolia. s.s.	(gr.) Lentiscus-leaved.	Aleppo.	1710.	5. 6. H. 5.
10 argénte'a. s.s.	(gr.) silver-leaved.	Corsica.	1825.	____ H. 5.
11 sambucifolia. s.s.	(gr.) Elder-leaved.	N. America.	1800.	____ H. 5.
12 oxycárp'a. s.s.	(gr.) sharp-leaved.	Tauria.	1815.	5. 6. H. 5.
oxyphylla. M.B.				
13 pállida. s.s.	(gr.) pale.	N. America.	1818. H. 5.
14 rufa. s.s.	(gr.) rufous.	_____	1822. H. 5.
15 fúsc'a. s.s.	(gr.) brown branched.	_____	____ H. 5.
16 nigra. s.s.	(gr.) black-twigg'd.	_____	1818.	4. 6. H. 5.
17 elliptica. s.s.	(gr.) elliptic-leaved.	_____	1824. H. 5.
18 ovata. s.s.	(gr.) oval-leaved.	_____	____ H. 5.
19 Richárd'i. s.s.	(gr.) Richard's.	_____	____ H. 5.
20 álba. s.s.	(gr.) grey branched.	_____	1825. H. 5.
21 cinérea. s.s.	(gr.) cinereous.	_____	1818. H. 5.

Rumph. amb. 1. t. 6.
Roxb. cor. 1. t. 46.
Rotb. act. hafn. 2. t. 5.
Roxb. cor. 1. t. 48.
Jacq. schen. 3. t. 312.
Roxb. cor. 1. t. 47.
Bot. reg. t. 499.

Hook. ex. flor. t. 139.

Mill. ic. t. 116.
Kæmp. amoen. t. 806.
Roxb. cor. 1. t. 49.
____ 1. t. 50.

Lam. ill. t. 370. f. 1.

Comm. hort. 1. t. 65.
Bot. reg. t. 500.
Lam. ill. t. 370. f. 2.

Vent. malm. t. 17.

Eng. bot. v. 24. t. 1692.

Eng. bot. v. 35. t. 2476.

Pluk. alm. t. 182. f. 4.

22	víridis. s.s.	(gr.) green-branched.	N. America.	1816.	H.5.	
23	longifolia. s.s.	(gr.) long-leaved.	1825.	H.5.		
24	tamariscifolia. R.S.	(gr.) Tamarisk-leaved.	Levant.	—	H.5.	
25	mexicana.	(gr.) Mexican.	Mexico.	1825.	H.5.	
26	rubicunda. s.s.	(gr.) red-veined.	N. America.	—	—	H.5.	
27	pulverulenta. s.s.	(gr.) powdered.	—	—	—	H.5.	
28	mixta. s.s.	(gr.) mixed.	—	—	—	H.5.	
29	expansa. s.s.	(gr.) expanded	—	—	—	H.5.	
30	platycarpa. s.s.	(gr.) flat-seeded.	—	1816.	4. 6.	H.5.	Catesb. car. t. 80.
31	quadranglelata. s.s.	(gr.) four-sided.	—	1811.	—	H.5.	Mich. f. arb. ic.
32	epiptera. s.s.	(gr.) dotted-stalked.	—	—	—	H.5.	— t. 33.
	discolor. M.						
33	lácea. Bosc.	(gr.) lance-leaved.	—	1823.	—	H.5.	
34	pannosa. s.s.	(gr.) cloth-leaved.	—	1818.	—	H.5.	
35	pubescens. s.s.	(gr.) pubescent.	—	1783.	5.	H.5.	Du Roi. ed. 2. t. 1.
36	cúrvidens. R.S.	(gr.) curved-toothed.	Carolina.	1811.	4. 5.	H.5.	
37	caroliniana. s.s.	(gr.) Carolina.	N. America.	—	4. 6.	H.5.	
38	juglandifolia. s.s.	(gr.) Walnut-leaved.	—	1783.	—	H.5.	
39	acuminata. s.s.	(gr.) taper-pointed.	—	1723.	—	H.5.	
40	americana. s.s.	(gr.) white.	—	—	—	H.5.	
O'RNU'S. P.S.	FLOWERING ASH.	Diandria Monogynia. P.S.					
1	americana. Ph.	(wh.) American.	N. America.	1812.	5. 6.	H.5.	
2	euro'pæa. P.S.	(wh.) European.	Italy.	1710.	—	H.5.	Flor. græc. 1. t. 4.
	Fráxinus O'rnu's. w.						
3	rotundifolia. P.S.	(wh.) manna.	—	1697.	4. 5.	H.5.	
4	striata.	(wh.) streak-barked.	—	1818.	—	H.5.	
	Fráxinus striata. s.s.						
5	floribunda.	(wh.) many-flowered.	Nepaul.	—	G.5.	
	Fráxinus floribunda. F.I.						
CHIONA'NTHUS. S.S.	FRINGE-TREE.	Diandria Monogynia. L.					
1	virginica. w.	(wh.) smooth-leaved.	N. America.	1736.	5. 7.	H.5.	Catesb. car. 1. t. 68.
2	maritima.	(wh.) pubescent-leaved.	—	—	—	H.5.	Wats. dend. brit. t. 1.
	virginica β maritima.	Ph. virginica β angustifolia. W.D.					
3	axillaris. B.P.	(wh.) axil-flowering.	N. Holland.	1810.	G. h.	
LINOCTERA. S.S.	LINOC'ERA.	Diandria Monogynia. S.S.					
1	ligustrina. s.s.	(wh.) Privet-like.	W. Indies.	1820.	S. h.	
2	cotinifolia. s.s.	(wh.) Cotinus-leaved.	Ceylon.	1818.	S. h.	
3	compacta. s.s.	(wh.) Caribbean.	W. Indies.	1793.	S. h.	Jacq. coll. 2. t. 6. f. 1.
FONTANESIA. S.S.	FONTAN'ESIA.	Diandria Monogynia. W.					
	phillyraeoides. s.s. (wh.)	Phillyrea-leaved.	Syria.	1787.	6. 8.	H. h.	
NOTE'L'EA. B.P.	NOTE'L'EA.	Diandria Monogynia. S.S.					
1	longifolia. B.P.	(wh.) long-leaved.	N. S. W.	1790.	3. 6.	G. h.	Venten. choix. t. 25.
	'Olea apetala. A.R.	316.					
2	ovata. B.P.	(wh.) oval-leaved.	—	1818.	—	G. h.	
3	ligustrina. B.P.	(wh.) Privet-leaved.	V. Diem. Isl.	1807.	7. 8.	G. h.	
4	rígida. R.S.	(wh.) rigid-leaved.	—	1820.	—	G. h.	
OLEA. B.P.	OLIVE.	Diandria Monogynia. L.					
1	euro'pæa. s.s.	(wh.) European.	S. Europe.	1570.	6. 8.	G. h.	
	a Oleaster. R.S.	(wh.) spiny-branched.	Portugal.	—	—	G. h.	Flor. græc. 1. t. 3.
	β longifolia. R.S.	(wh.) long-leaved.	S. Europe.	—	—	G. h.	Lodd. bot. cab. t. 456.
	γ ferruginea. R.S.	(wh.) ferruginous.	—	—	—	G. h.	
	δ latifolia. R.S.	(wh.) broad-leaved.	—	—	—	G. h.	
	ε obliqua. R.S.	(wh.) oblique-leaved.	—	—	—	G. h.	
	ζ buxifolia. R.S.	(wh.) Box-leaved.	—	—	—	G. h.	
2	verrucosa. s.s.	(wh.) warted.	C. B. S.	1814.	—	G. h.	
3	americana. s.s.	(wh.) American.	N. America.	1758.	6.	G. h.	Catesb. car. 1. t. 61.
4	frágans. s.s.	(wh.) fragrant.	China.	1771.	6. 8.	G. h.	Bot. mag. t. 1552.
5	lácea. s.s.	(wh.) lance-leaved.	Mauritius.	1818.	—	S. h.	
6	dioica. F.I.	(wh.) dioecious.	E. Indies.	—	—	S. h.	
7	capensis. s.s.	(wh.) Cape.	C. B. S.	1730.	6. 9.	G. h.	Bot. reg. t. 613.
8	undulata. s.s.	(wh.) wave-leaved.	—	—	—	G. h.	Lodd. bot. cab. t. 379.
9	emarginata. R.S.	(wh.) emarginate.	Mauritius.	1816.	S. h.	
10	excelsa. s.s.	(wh.) Laurel-leaved.	Madeira.	1784.	5. 6.	G. h.	Jacq. schœn. 3. t. 251.
11	cernua. s.s.	(wh.) drooping-flower'd.	Mauritius.	1816.	S. h.	
12	paniculata. B.P.	(wh.) panicled.	N. S. W.	1823.	4. 6.	G. h.	
13	Roxburghii.	(wh.) Roxburgh's.	E. Indies.	1828.	S. h.	
	paniculata. F.I. non B.P.						
14	robusta.	(wh.) large Phillyrea.	—	1818.	S. h.	
	Phillyrea robusta. F.I.						

OLEINÆ.

15 chinensis.	(wh.) Chinese.	China.	1820.	4. 6. H. \ddot{h} .	
	<i>Phillyrea paniculata</i> . F.I.				
16 angustifolia.	(wh.) narrow-l'd. <i>Phillyrea</i> . S. Europe.	1597.	5. 6. H. \ddot{h} .	Lam. ill. t. 8. f. 3.	
	<i>Phillyrea angustifolia</i> . s.s.				
α lanceolata.	(wh.) spear-leaved.	—	—	H. \ddot{h} .	
β rosmarinifolia.	(wh.) Rosemary-leaved.	—	—	H. \ddot{h} .	
γ brachiata.	(wh.) cross-branched.	—	—	H. \ddot{h} .	
17 media.	R.S.	(wh.) twiggy <i>Phillyrea</i> .	—	H. \ddot{h} .	
	β buxifolia.	R.S. (wh.) Box-leaved.	—	H. \ddot{h} .	
18 ligustrifolia.	R.S. (wh.)	Privet-leaved.	—	H. \ddot{h} .	
19 pendula.	R.S. (wh.)	pendulous.	—	H. \ddot{h} .	
20 oleæfolia.	R.S. (wh.)	Olive-leaved <i>Phillyrea</i> .	—	H. \ddot{h} .	
21 lœvis.	R.S. (wh.)	smooth-leaved.	—	H. \ddot{h} .	Pluk. alm. t. 310. f. 5.
22 latifolia.	R.S. (wh.)	broad-leaved	—	H. \ddot{h} .	Duham. arb. t. 125.
23 obliqua.	R.S. (wh.)	oblique-leaved.	—	H. \ddot{h} .	Flor. græc. t. 2.
24 spinosa.	R.S. (wh.)	Holly-leaved.	—	H. \ddot{h} .	
	<i>Phillyrea ilicifolia</i> . W.en.				
LIGU'STRUM. W.	PRIVET.	Diandria Monogynia. L.			
1 vulgare.	R.S. (wh.) common.	Britain.	6. 7. H. \dot{h} .	Eng. bot. v. 11. t. 764.
β variegatum.	(wh.) striped-leaved.	—	H. \dot{h} .	
γ angustifolium.	(wh.) narrow-leaved.	—	H. \dot{h} .	
δ leucocarpum.	(wh.) white-berried.	—	H. \dot{h} .	
ϵ italicum.	Mill. (wh.) evergreen.	Italy.	H. \dot{h} .	
2 lùcum.	R.S. (wh.) Wax-tree.	China.	1794.	6. 9. F. \ddot{h} .	Wats. dend. brit. t. 137.
SYRI'NGA. R.S.	LILAC.	Diandria Monogynia. L.			
1 vulgaris. w.	(va.) common.	Persia.	1597.	5. H. \dot{h} .	Schk. hand. 1. t. 2.
α lilacina.	(li.) pale-flowered.	—	—	H. \dot{h} .	Bot. mag. t. 183.
β purpurea.	(pu.) purple-flowered.	—	—	H. \dot{h} .	
γ violacea.	(vi.) violet.	—	—	H. \dot{h} .	
δ alba.	(wh.) white-flowered.	—	—	H. \dot{h} .	
ϵ variegata.	(li.) variegated-leaved.	—	—	H. \dot{h} .	
2 chinensis. w.	(pu.) Siberian.	China.	1795.	H. \dot{h} .	
3 persica. w.	(li.) Persian.	Persia.	1640.	H. \dot{h} .	Bot. mag. t. 486.
α lilacina.	(li.) pale-flowered.	—	—	H. \dot{h} .	
β alba.	(wh.) white-flowered.	—	—	H. \dot{h} .	
γ laciniata.	(wh.) cut-leaved.	—	—	H. \dot{h} .	Schmidt arb. 2. t. 79.

ORDO CXXXVII.

JASMINEÆ. Brown prodr. 1. p. 520.

NYCTA'NTHES. W.	NYCTA'NTHES.	Diandria Monogynia. L.			
árbor tristis. w.	(wh.ye.) square-stalked.	E. Indies.	1781.	6. 9. S. \dot{h} .	Bot. reg. t. 399.
JASMINUM. W.	JASMINE.	Diandria Monogynia. L.			
1 Sámbac. s.s.	(wh.) single Arabian.	E. Indies.	1665.	1. 12. S. \dot{h} .	Bot. reg. t. 1.
β mūltiplex.	(wh.) semidouble.	—	1700.	S. \dot{h} .	Andr. reposit. t. 497.
γ trifoliatum.	(wh.) Tuscan.	—	1730.	S. \dot{h} .	Bot. mag. t. 1785.
2 undulatum. s.s.	(wh.) wave-leaved.	—	1812.	2. 6. S. \dot{h} .	Bot. reg. t. 436.
3 hirsutum. s.s.	(wh.) hairy Indian.	—	1759.	5. 12. S. \dot{h} .	Sm. ex. bot. 2. t. 118.
	<i>multiflorum</i> . And. rep. t. 436.				
4 angustifolium. F.I.	(wh.) narrow-leaved.	—	1816.	1. 12. S. \dot{h} .	Rheed. mal. 6. t. 53.
5 laurifolium. F.I.	(wh.) Laurel-leaved.	—	—	S. \dot{h} .	Bot. reg. t. 521.
6 trinérve. s.s.	(wh.) three-nerved.	Java.	1804.	4. 6. S. \dot{h} .	— t. 918.
7 glaucum. s.s.	(wh.) glaucous.	C. B. S.	1774.	6. 8. S. \dot{h} .	Sal. stirp. rar. t. 8.
8 bracteatum. s.s.	(wh.) bracteate.	Sumatra.	1820.	S. \dot{h} .	
9 latifolium. s.s.	(wh.) broad-leaved.	E. Indies.	1817.	S. \dot{h} .	
10 arboréscens. s.s.	(wh.) arborescent.	—	1824.	S. \dot{h} .	
11 scándens. s.s.	(wh.) scandent.	—	—	S. \dot{h} .	
12 simplicifolium. s.s.	(wh.) simple-leaved.	Society Isl.	3. 6. S. \dot{h} .	Bot. mag. t. 980.
13 elongatum. s.s.	(wh.) long-branched.	E. Indies.	1523.	S. \dot{h} .	Berg. act. an. 1772. t. 11.
14 grácie. s.s.	(wh.) slender.	Norfolk Isl.	1791.	1. 12. G. \dot{h} .	Andr. reposit. t. 127.
15 acuminatum. s.s.	(wh.) acuminate.	N. Holland.	1820.	6. 9. G. \dot{h} .	Bot. reg. t. 1296?
16 volùbile. R.S.	(wh.) twisting.	C. B. S.	1816.	4. 7. G. \dot{h} .	Jacq. schœn. 3. t. 321.
17 auriculatum. s.s.	(wh.) auriculate.	E. Indies.	1790.	5. 9. G. \dot{h} .	Bot. reg. t. 264.
18 paniculatum. s.s.	(wh.) variable-leaved.	China.	1812.	2. 12. S. \dot{h} .	— t. 690.

19 fléxile. s.s. (wh.) acute 3-leaved.	E. Indies.	1820.	4. 8. S. h.~.
20 tortuosum. s.s. (wh.) twisting.	C. B. S.	—	— S. h.~. Jacq. schœn. 4. t. 490.
fléxile. Jacq. non w.			
21 campanulatum. s.s. (wh.) bell-shaped calyx'd.	—	—	S. h.~.
22 capénse. s.s. (wh.) Cape.	C. B. S.	1816.	G. h.
23 azoricum. s.s. (wh.) white Azorian.	Madeira.	1724.	4. 11. G. h. Bot. reg. t. 89.
24 dispérnum. s.s. (wh.) two-seeded.	Nepaul.	1825. G. h.~.
25 fruticans. s.s. (ye.) common yellow.	S. Europe.	1570.	4. 10. H. h. Bot. mag. t. 461.
26 humili. s.s. (ye.) Italian.	—	1656.	6. 9. H. h. Bot. reg. t. 350.
27 odoratissimum. s.s. (ye.) yellow Azorian.	Madeira.	—	5. 11. G. h. Bot. mag. t. 285.
28 revolutum. s.s. (ye.) yellow Nepaul.	Nepaul.	1812.	3. 10. H. h. Bot. reg. t. 178.
chrysanthemum. F.I.			
29 pubigerum. D.P. (ye.) pubescent.	—	1828.	H. h.~.
30 officinale. s.s. (wh.) common white.	E. Indies.	1548.	6. 10. H. h.~. Bot. mag. t. 31.
β leucophyllum. (wh.) white-edged.	—	H. h.~.
γ chrysophyllum. (wh.) yellow-edged.	—	H. h.~.
δ multiplex. (wh.) double-blossomed.	—	1823.	H. h.~.
31 grandiflorum. s.s. (wh.) Catalonian.	—	1629.	G. h.~. Bot. reg. t. 91.

ORDO CXXXVIII.

APOCINEÆ. Brown prodr. 1. p. 465.

ALY'XIA. B.P.	ALY'XIA.	Pentandria Monogynia.	
Richardsoni.	(wh.) Richardson's.	N. Holland.	1823. 4. 6. S. h.
RAUWO'LFI A. S.S.	RAUWO'LFI A.	Pentandria Monogynia. W.	
1 nítida. s.s.	(wh.) shining.	S. America.	1752. 6. 9. S. h.
2 ternifolia. s.s.	(wh.) three-leaved.	—	1822. — S. h.
3 canescens. s.s.	(fl.) hoary.	Jamaica.	1739. — S. h.
4 tomentosa. s.s.	(wh.) woolly.	S. America.	1823. — S. h.
5 spinosa. r.s.	(st.) spiny.	Peru.	1827. — S. h.
VALL'ESIA. R.P.	VALL'ESIA.	Pentandria Monogynia.	
cymbæfolia. r.s. (wh.) boat-leaved.	S. America.	1816. 6. 8. S. h.	Cavan. icon. 6. t. 526.
Rauwolfa glabra. w.			
OPHIO'XYLON. W.	OPHIO'XYLON.	Polygamia Monoecia. W.	
serpentinum. w. (wh.) serpent-wood.	E. Indies.	1690. 5. 6. S. h.	Bot. mag. t. 784.
STRY'CHNOS. S.S.	STRY'CHNOS.	Pentandria Monogynia. W.	
1 Nux vómica. s.s. (wh.) Poison-nut.	E. Indies.	1778. S. h.	Roxb. cor. 1. t. 4.
2 colubrina. s.s. (gr.ye.) muricate-fruited.	—	1818. 5. 8. S. h.	Rheed. mal. 8. t. 24.
3 potatorum. s.s. (wh.) clearing-nut.	—	1794. 4. 6. S. h.	Roxb. cor. 1. t. 5.
4 madagascariensis. s.s. (wh.) Madagascar.	Madagascar.	1818. S. h.	
5 spinosa. s.s. (gr.ye.) spiny.	—	1824. S. h.	
6 axillaris. s.s. (wh.) axillary-flowered.	E. Indies.	1818. 5. 7. S. h.	Linn. trans. v. 12. t. 15.
THEOPHRA'STA. S.S. THEOPHRA'STA.		Pentandria Monogynia.	
1 longifolia. r.s.	(or.) long-leaved.	Caracas.	1827. S. h.
2 macrophylla. r.s. large-leaved.	Brazil.	— S. h.
3 americana. r.s. American.	S. America.	— S. h.
4 Jussieai. s.s.	(wh.) Jussieu's.	—	1818. 1. 4. S. h.
CLAVI'JA. R.P.	CLAVI'JA.	Pentandria Monogynia.	
macrophylla. R.P. (wh.) long-leaved.	S. America.	1816. S. h.	Flor. peruv. gen. t. 30.
FAGR'EA. W.	FAGR'EA.	Pentandria Monogynia.	
1 zeylanica. w.	(wh.) Ceylon.	Ceylon.	1824. 4. 7. S. h.
2 frágans. F.I.	(wh.) fragrant.	China.	1826. 5. 7. S. h.
CARISSA. L.	CARISSA.	Pentandria Monogynia. L.	
1 Carándas. s.s.	(wh.) Jasmine-flower'd.	E. Indies.	1790. 7. 8. S. h.
2 spinárum. s.s.	(wh.) Box-leaved.	—	1809. 8. 12. S. h.
3 ovata. B.P.	(wh.) oval-leaved.	N. Holland.	1817. 6. 9. G. h.
4 Xylopícoron. s.s.	(wh.) nerved-leaved.	Mauritius.	1820. — S. h.
ARDU'INA. L.	ARDU'INA.	Pentandria Monogynia. L.	
bispinosa. s.s.	(wh.) two-spined.	C. B. S.	1760. 3. 8. G. h.
Carissa Arduina. P.S.			Mill. ic. 2. t. 300.
GELS'EMIUM. J.	GELS'EMIUM.	Pentandria Monogynia.	
semperfíren. p.s. (ye.) evergreen.	N. America.	1640. 6. 7. G. h.~.	Catesb. car. 1. t. 53.
nítidum. r.s. Bigñonia semperfíren. w.			

APOCINEÆ.

CE'RBERA. W.	CE'RBERA.	Pentandria Monogynia. L.				
1 Ahouai. s.s. (ye.) oval-leaved.	Brazil.	1739.	6. 7.	S. h.	Bot. mag. t. 737.	
2 ovâta. s.s. (ye.) ovate-leaved.	N. Spain.	—	S. h.	Cavan. ic. 3. t. 270.	
3 fruticosa. s.s. (re.) purple-flowered.	Pegu.	1817.	3. 9.	S. h.	Bot. reg. t. 391.	
4 Mângas. L.T. (wh.) blunt-leaved.	E. Indies.	1759.	4. 8.	S. h.	Burm. zeyl. t. 70. f. 1.	
5 Odâllam. L.T. (wh.re.) spear-leaved.	—	—	6. 9.	S. h.	Bot. mag. t. 1845.	
Mângas. B.M. non L.T.						
6 lactaria. L.T. (wh.) Milk-tree.	—	—	S. h.	Rumph. amb. 2. t. 81.	
Mângas. G. 2. t. 123 et 124. f. 1. non L.T.						
7 Tângin. B.M. (bh.re.) poison-nut.	Madagascar.	1826.	S. h.	Bot. mag. t. 2968.	
8 lauriifolia. B.C. (wh.) Laurel-leaved.	E. Indies.	1819.	6. 9.	S. h.	Lodd. bot. cab. t. 989.	
9 maculata. s.s. (wh.re.) wave-leaved.	Bourbon.	1782.	6. 7.	S. h.	Andr. deposit. t. 130.	
10 dichotoma. B.C. (wh.) forked.	E. Indies.	1827.	—	S. h.	Lodd. bot. cab. t. 1516.	
11 thevetioides. s.s. (ye.) revolute-margin'd.	N. Spain.	1820.	S. h.		
12 Thévetia. s.s. (ye.) linear-leaved.	S. America.	1735.	S. h.	Jacq. amer. 48. t. 34.	
13 cuneifolia. K.S. (ye.) wedge-leaved.	Mexico.	1826.	4. 6.	S. h.		
OCHR'OSIA. J.	OCHR'OSIA.	Pentandria Monogynia. R.S.				
borbônica. R.S. (st.) Bourbon.	Bourbon.	1823.	S. h.		
Cérbera borbônica. s.s. <i>Ophiòxylon Ochròsia</i> . P.S.						
DISSOLENA. R.S.	DISSOLENA.	Pentandria Monogynia. R.S.				
verticillata. R.S. (wh.) whorl-leaved.	China.	1812.	S. h.		
ALLAMA'NDA. S.S.	ALLAMA'NDA.	Pentandria Monogynia. L.				
1 cathartica. s.s. (ye.) Willow-leaved.	Guiana.	1785.	6. 7.	S. h.	Bot. mag. t. 338.	
2? verticillata. s.s. whorl-leaved.	E. Indies.	1812.	S. h.		
<i>Nérium tinctòrium</i> . Hort.						
VI'NCA. R.S.	PERIWINKLE.	Pentandria Monogynia. L.				
1 herbacea. R.S. (bl.) herbaceous.	Hungary.	1816.	6. 7.	H. ȝ.	Bot. reg. t. 301.	
2 minor. R.S. (bl.) lesser.	Britain.	3. 9.	H. h.	Eng. bot. v. 13. t. 917.	
α cærulea.	(bl.) blue-flowered.	—	—	H. h.		
β ábla.	(wh.) white-flowered.	—	—	H. h.		
γ atropurpurea.	(pu.) dark-purple.	—	—	H. h.		
δ máltiplex.	(pu.) double-flowered.	—	—	H. h.		
ε variegata.	(wh.) variegated.	—	—	H. h.		
3 major. R.S. (bl.) greater.	England.	—	H. h.	Eng. bot. v. 8. t. 514.	
β máltiplex.	(bl.) double-blue.	—	—	H. h.		
4 pusilla. R.S. (st.) small-flowered.	E. Indies.	1778.	8.	S. ⓧ.	Rheed. mal. 9. t. 33.	
parviflora. W.						
5 rosea. R.S. (ro.) Madagascar.	—	1756.	3. 10.	S. h.	Bot. mag. t. 248.	
β ábla.	(wh.) white-flowered.	—	—	S. h.		
γ ocellata.	(wh.re.) red-eyed.	—	—	S. h.		
WRI'GHTIA. S.S.	WRI'GHTIA.	Pentandria Monogynia. S.S.				
1 antídysentérica. s.s. (wh.) oval-leaved.	E. Indies.	1778.	S. h.	Burm. zeyl. t. 77.	
2 zeylanica. s.s. (wh.) spear-leaved.	—	S. h.	— t. 12. f. 2.	
3 dubia. s.s. (sc.) doubtful.	—	5. 8.	S. h.	Bot. mag. t. 1646.	
<i>Camerària dàbia</i> . B.M.						
4 coccinea. B.M. (sc.) scarlet.	—	1820.	6. 8.	S. h.	Bot. mag. t. 2696.	
<i>Nérium coccineum</i> . Bot. cab. t. 894.						
5 tinctoria. s.s. (wh.) dyer's.	—	1812.	—	S. h.	Bot. reg. t. 933.	
STROPH'A'NTHUS. DC.	STROPH'A'NTHUS.	Pentandria Monogynia. P.S.				
1 dichotomus. DC. (ro.) forked.	E. Indies.	1816.	2. 10.	S. h.	Burm. ind. 68. t. 26.	
2 divulgans. Gr. (ye.) spreading.	China.	—	—	S. h.	Bot. reg. t. 469.	
dichotomus. B.R. non DC.						
3 sarmentosus. DC. (re.) sarmentose.	Sierra Leon.	1822.	S. h.	Desf. an. mus. 1. t. 27.	
N'ERIUM. R.S.	OLEANDER.	Pentandria Monogynia. L.				
1 Oleánder. R.S. (ro.) common.	S. Europe.	1596.	6. 10.	G. h.	Flor. græc. t. 248.	
β ábla.	(wh.) white-flowered.	—	—	G. h.	Lodd. bot. cab. t. 700.	
γ spléndens.	(ro.) double-hybrid.	Hybrid.	1815.	—	Bot. reg. t. 74.	
2 odorum. R.S. (ro.) sweet-scented.	E. Indies.	1683.	6. 8.	S. h.	Rheed. mal. 9. t. 2.	
β carneum.	(fl.) flesh-coloured.	—	1816.	—	Bot. mag. t. 2032.	
γ flore pleno.	(ro.) double-flowered.	—	—	S. h.	Rheed. mal. 9. t. 1.	
3 flavescens. R.S. (st.) yellowish.	—	1816.	—	S. h.		
TABER'EMONT'ANA. TABERN'EMONT'ANA.	Pentandria Monogynia. L.					
1 citrifolia. R.S. (st.) Citron-leaved.	Jamaica.	1734.	5. 8.	S. h.	Plum. ic. t. 248. f. 2.	
2 ábla. R.S. (wh.) white-flowered.	W. Indies.	1780.	—	S. h.	Jacq. amer. t. 175. f. 13.	
<i>citrifolia</i> . Jacq. non L.						
3 lauriifolia. R.S. (st.) Laurel-leaved.	—	1768.	—	S. h.	Bot. reg. t. 716.	
4 cymosa. R.S. (wh.) cyme-flowered.	S. America.	1820.	—	S. h.	Jacq. amer. t. 181. f. 14.	
5 amygdalifolia. R.S. (wh.) Peach-leaved.	—	1780.	4. 8.	S. h.	Bot. reg. t. 338.	
6 discolor. R.S. (st.) two-coloured.	Jamaica.	1822.	—	S. h.		

7 persicariæfolia. R.S. (st.)	Persicaria-leav'd. Mauritus.	1826.	4. 8.	S. h.	Jacq. icon. rar. t. 320.	
8 arcuata. R.S. (st.)	arch'd follicled. Peru.	1820.	—	S. h.	Flor. peruv. 2. t. 143.	
9 crista. L.T. (wh.)	wave-leaved. E. Indies.	1818.	5. 10.	S. h.	Rheed. mal. 1. t. 46.	
10 coronaria. L.T. (wh.)	broad-leaved. ———	1710.	6. 10.	S. h.	Lodd. bot. cab. t. 406.	
β flôre pleno.	(wh.) double-flowered. ———	—	—	S. h.	Bot. mag. t. 1865.	
Nérüm coronárium. B.M.						
11 densiflora. B.R. (wh.)	close-flowered. Ceylon.	1824.	6. 7.	S. h.	Bot. reg. t. 1273.	
12 undulata. R.S. (st.)	waved. Trinidad.	—	—	S. h.		
13 odorata. s.s. (ye.)	yellow-flowered. Cayenne.	1793.	10. 11.	S. h.	Aub. guian. 1. t. 102.	
Cameraria lutea. w.	Cameraria Tamaquarina. Aublet.					
CAMERARIA. S.S.	BASTARD MANGANEEL. Pentandria Monogynia. L.					
1 latifolia. s.s. (wh.)	broad-leaved. Havannah.	1733.	8.	S. h.	Andr. deposit. t. 261.	
2 angustifolia. s.s. (wh.)	narrow-leaved. S. America.	1752.	9.	S. h.	Plum. ic. t. 72. f. 2.	
AMS'ONIA. R.S.	AMS'ONIA. Pentandria Monogynia. P.S.					
1 latifolia. R.S. (bl.)	broad-leaved. N. America.	1759.	5. 6.	H. y.	Bot. reg. t. 151.	
2 salicifolia. R.S. (pa.bl.)	Willow-leaved. ———	1812.	—	H. y.	Bot. mag. t. 1873.	
3 angustifolia. R.S. (bl.)	hairy-stalked. ———	1774.	—	H. y.	Venten. choix. t. 29.	
PLUM'ERIA. S.S.	PLUM'ERIA. Pentandria Monogynia. L.					
1 rübra. R.S. (re.)	red-flowered. Jamaica.	1690.	7. 8.	S. h.	Bot. reg. t. 780.	
2 purpurea. R.S. (pu.)	purple. Peru.	1820.	—	S. h.	Flor. peruv. 2. t. 137.	
3 acuminata. H.K. (wh.)	acute-leaved. E. Indies.	1790.	6. 9.	S. h.	Bot. reg. t. 114.	
acutifolia. R.S.						
4 obtusa. R.S. (wh.)	blunt-leaved. W. Indies.	1733.	7. 8.	S. h.	Catesb. car. 1. t. 93.	
5 alba. R.S. (wh.)	white. Jamaica.	—	—	S. h.	Jacq. amer. t. 174. f. 2.	
6 incarnata. R.P. (fl.)	flesh-coloured. Peru.	—	—	S. h.	Flor. peruv. t. 138.	
7 carinata. R.P. (wh.re.)	keel-leaved. ———	1824.	—	S. h.		
8 bicolor. R.P. (ye.wh.)	two-coloured. S. America.	1815.	6. 9.	S. h.	Bot. reg. t. 480.	
9 tricolor. R.P. (wh.re.)	three-coloured. ———	—	—	S. h.	————— t. 510.	
10 lutea. R.P. (ye.)	yellow-flowered. ———	—	—	S. h.	Flor. peruv. 2. t. 142.	
11 aurantia. H.S.S. (or.)	orange-flowered. ———	1816.	—	S. h.		
12 pudica. R.S. (st.)	wax-flowered. ———	—	7. 8.	S. h.		
13 longifolia. R.S.	long-leaved. Madagascar.	1816.	—	S. h.		
14 tuberculata. B.C. (wh.)	tubercl. S. America.	1820.	6. 8.	S. h.	Lodd. bot. cab. t. 681.	
15 tenuifolia. L.C.	slender-leaved. ———	—	—	S. h.		
16 Northiana. L.C.	North's. ———	—	—	S. h.		
17 mexicana. B.C. (wh.)	Mexican. Mexico.	—	6. 9.	S. h.	Lodd. bot. cab. t. 1024.	
18 Cowani. D.D. (wh.)	Cowan's. ———	1824.	—	S. h.		
19 Blandfordiana. L.C.	Blandford's. S. America.	—	—	S. h.		
20 nivea. L.C. (wh.)	snowy white. ———	—	6. 8.	S. h.		
21 conspicua. L.C.	conspicuous. ———	—	—	S. h.		
22 macrophylla. L.C.	large-leaved. ———	—	—	S. h.		
23 leucantha. L.C. (wh.)	white-blossomed. ———	—	—	S. h.		
24 parvifolia. H.S.S. (wh.)	small-leaved. W. Indies.	1813.	7. 8.	S. h.		
PREST'ONIA. S.S.	PREST'ONIA. Pentandria Monogynia. S.S.					
1 tomentosa. s.s. (ye.)	tomentose. Brazil.	1820.	5. 8.	S. h. —		
2 glabrata. s.s. (ye.)	smooth. S. America.	1823.	—	S. h. —		
CRYPTOL'EPIS. B.	CRYPTOL'EPIS. Pentandria Monogynia.					
élegans. L.C.	elegant. E. Indies.	1823.	—	S. h. —		
ALST'ONIA. S.S.	ALST'ONIA. Pentandria Monogynia. S.S.					
1 scholariæ. s.s. (wh.)	whorl-leaved. E. Indies.	1816.	—	S. h.	Rheed. mal. 1. t. 45.	
2 venenata. s.s. (wh.)	poisonous. ———	1823.	—	S. h.		
THENA'R'DIA. K.S.	THENA'R'DIA. Pentandria Monogynia.					
floribunda. K.S. (gr.)	abundant-flowered. Mexico.	1823.	—	S. h. —	H. et B. 3. t. 240.	
ECH'ITES. R.S.	ECH'ITES. Pentandria Monogynia. L.					
1 longiflora. s.s. (wh.)	long-flowered. Brazil.	1816.	5. 8.	S. h. —		
2 bispinosa. s.s. (fl.)	two-spined. C. B. S.	1795.	7. 11.	S. h.		
3 succulenta. s.s. (re.wh.)	succulent. ———	1820.	4. 6.	S. h.	Jacq. fragm. t. 117.	
tuberosa. H.S.S.						
4 repens. R.S. (re.)	creeping.	St. Domingo.	1824.	6. 8.	S. h.	Jacq. am. pict. t. 35.
5 biflora. s.s. (wh.)	twin-flowered. W. Indies.	1793.	7. 8.	S. h. —	Jacq. amer. 30. t. 21.	
6 umbellata. s.s. (wh.)	umbelled. Jamaica.	1733.	7.	S. h. —	————— 30. t. 22.	
7 torosa. s.s. (ye.)	climbing. ———	1778.	6. 8.	S. h. —	————— 33. t. 27.	
8 antidysenterica. s.s. (pi.)	medicinal.	E. Indies.	1821.	—	S. h.	
9 pubescens. L.T. (wh.)	pubescent.	—	1822.	—	S. h.	Rheed. mal. 1. t. 49. ?
10 suberecta. s.s. (ye.)	Savanna-flower. W. Indies.	1759.	6. 8.	S. h. —	Bot. mag. t. 1064.	
11 caryophyllata. s.s. (st.)	Clove-scented. E. Indies.	1812.	9. 11.	S. h. —	————— t. 1919.	
12 difformis. s.s. (ye.)	deformed. Carolina.	1806.	6. 7.	S. h. —		
13 grandiflora. R.S. (pi.)	large-flowered.	E. Indies.	1823.	—	S. h.	
macrantha. s.s.						

14 reticulata. s.s. (wh.) netted-leaved.	E. Indies.	1818.	6. 7.	G. h.~.
15 Héynii. s.s. (wh.) Heyne's.	_____	____	____	S. h.~.
16 frutescens. H.B. (wh.) frutescent.	_____	1816.	____	S. h.~.
17 paniculata. H.B. (wh.) panicled.	_____	____	____	S. h.~.
BEAUMONTIA. B.R. BEAUMONTIA. Pentandria Monogynia.				
1 grandiflora. B.R. (wh.) oval-leaved.	E. Indies.	1812.	5. 7.	S. h.~. Bot. reg. t. 911.
2 longifolia. L.C. (wh.) long-leaved.	_____	1818.	____	S. h.~.
HÆMADI'CYON. H.T. HÆMADI'CYON. Pentandria Monogynia.				
venosum. H.T. (ye.) red-veined.	W. Indies.	1820.	4. 8.	S. h.~. Bot. mag. t. 2473.
<i>Echites nūtans.</i> B.M.				
VALLARIS. S.S. VALLARIS. Pentandria Monogynia. S.S.				
Pergulānus. s.s. (wh.) smooth-leaved.	E. Indies.	1818.	____	S. h.~. Rumph. am. 5. t. 29.
<i>Pergularia glabra.</i> w. <i>Emericia Pergularia.</i> R.S.				
ICHNOCA'RPUS. S.S. ICHNOCA'RPUS. Pentandria Monogynia. S.S.				
frutescens. s.s. (pu.) shrubby.	E. Indies.	1759.	7. 8.	S. h. Burm. zeyl. t. 12. f. 1.
PARSONSIA. R.S. PARSONSIA. Pentandria Monogynia. R.S.				
1 corymbosa. r.s. (re.) corymb-flowered. W. Indies.	1820.	6. 8.	S. h.~.	Jacq. amer. pict. t. 37.
<i>Echites corymbosa.</i> Jacq.				
2 floribunda. r.s. (wh.) many-flowered. Jamaica.	_____	____	____	S. h.
<i>Echites floribunda.</i> w.				
WILLUGHBE'IA. S.S. WILLUGHBE'IA. Pentandria Monogynia. S.S.				
ēdulis. s.s. (bh.) eatable.	E. Indies.	1818.	6. 8.	S. h.~. Roxb. corom. 3. t. 280.
LYO'NSIA. B.P. LYO'NSIA. Pentandria Monogynia. S.S.				
straminea. b.p. (st.) climbing.	N. S. W.	1820.	6. 8.	S. h.~.
MELOD'INUS. R.S. MELOD'INUS. Pentandria Mono-Digynia.				
1 scāndens. r.s. (wh.) climbing.	N. Caledonia.	1775.	____	S. h.~. Lam. ill. t. 179.
2 parvifolius. L.C. (wh.) small-leaved.	E. Indies.	_____	____	S. h.
3 undulatus. L.C. (wh.) wave-leaved.	_____	____	____	S. h.
4 monogynus. B.R. (wh.) one-styled.	_____	1816.	4. 8.	S. h.~. Bot. reg. t. 834.
CARPOD'INUS. D.D. PISHAMIN. Pentandria Digynia.				
1 dūcīs. D.D. (wh.) sweet.	Sierra Leone.	1823.	____	S. h.~.
2 ácidus. D.D. (wh.) sour.	_____	____	____	S. h.~.
APO'CYNUM. S.S. DOG'S-BANE. Pentandria Monogynia. R.S.				
1 androsæmifolium. L.C. (bh.) Tutsan-leaved. N. America.	1688.	7. 9.	H. Ȑ.	Bot. mag. t. 280.
2 cannabīnum. s.s. (gr.) Hemp-like.	_____	1699.	____	H. Ȑ. Moris. s. 15. t. 3. f. 14.
3 hypericifolium. s.s. (wh.) Hypericum-leav'd.	_____	1758.	6. 7.	H. Ȑ. Jacq. vind. 3. t. 66.
4 venētum. s.s. (re.) Venetian.	Adriatic Isl.	1690.	____	H. Ȑ. Lob. ic. 372. f. 1. 2.
β álbum.	(wh.) white-flowered.	_____	____	H. Ȑ.

ORDO .CXXXIX.

ASCLEPIADEÆ. *Brown prodr. 1. p. 458.*

PERIPL'OCÀ. R.S. PERIPL'OCÀ. Pentandria Digynia. L.				
1 gr'æca. R.S. (pu.) Virginian-silk.	Syria.	1597.	7. 8.	S. h.~. Bot. reg. t. 803.
2 mauritiāna. R.S. (pu.) long-leaved.	Mauritius.	1823.	____	S. h.~. Rheed. mal. 9. t. 11.
3 levigata. R.S. (ye.) smooth.	Canaries.	1779.	6. 8.	G. h. Cavan. ic. 3. t. 217. <i>punicafolia.</i> Cav. icon.
HEMIDE'SMUS. R.S. HEMIDE'SMUS. Pentandria Digynia. R.S.				
indicus. R.S. (gr.) Indian.	Ceylon.	1796.	____	S. h. Burm. zeyl. t. 83. f. 1.
CRYPTOST'EGIA. B.R. CRYPTOST'EGIA. Pentandria Digynia. S.S.				
grandiflora. B.R. (ro.) large-flowered.	E. Indies.	1812.	6. 8.	S. h.~. Bot. reg. t. 435.
SECAM'ONE. R.S. SECAM'ONE. Pentandria Digynia. R.S.				
1 emética. R.S. (wh.) narrow-leaved.	India.	1816.	____	S. h. Willd. phyt. 1. t. 5. f. 2.
2 ægyptiaca. H.K. (wh.) Egyptian.	Egypt.	1752.	7.	G. h. Alp. ægyp. t. 134. <i>Alpini.</i> r.s. <i>Periploca Secamone.</i> w.
DUVAL'IA. H.S. DUVAL'IA. Pentandria Digynia. H.S.				
1 reclinata. H.S. (pu.ye.) reclined.	C. B. S.	1795.	8. 10.	D.S. h. Bot. mag. t. 1397.
2 elegans. H.S. (d.pu.) elegant.	_____	____	____	D.S. h. t. 1184. <i>Stapēlia Elegans.</i> B.M.
3 hirtella. s.s. (d.pu.) hairy.	_____	____	____	D.S. h. Jacq. stap. c. ic. <i>Stapēlia hirtella.</i> Jacq.
4 cæpitosa. H.S. (d.pu.) tufted.	_____	1790.	8. 11.	D.S. h. Mass. stap. 20. t. 29.

5 radiàta. H.S. (d.pu.) radiated.	C. B. S.	1795.	9. 11.	D.S. $\ddot{\text{h}}$.	Bot. mag. t. 619.
6 replicàta. s.s. (br.pu.) replicate. <i>Stapèlia replicàta.</i> Jacq.	_____	1806.	8. 10.	D.S. $\ddot{\text{h}}$.	Jacq. stap. c. ic.
7 mastòdes. s.s. (d.pu.) teat.	_____	1821.	—	D.S. $\ddot{\text{h}}$.	Jacq. stap. c. ic.
8 Jacquiniana. R.S.(d.pu.)Jacquin's. <i>Stapèlia radiàta.</i> Jacq. non B.M.	_____	1802.	7. 9.	D.S. $\ddot{\text{h}}$.	— t. 37.
9 tuberculàta. H.S.s.(d.p.)tubercled.	_____	1774.	7. 11.	D.S. $\ddot{\text{h}}$.	
10 laevigàta. H.S.s. (d.p.) smooth.	_____	1808.	—	D.S. $\ddot{\text{h}}$.	
11 glomeràta. H.S.s. (d.p.) clustered.	_____	—	—	D.S. $\ddot{\text{h}}$.	
12 compàcta. H.S.s. (d.p.) compact.	_____	1800.	8. 11.	D.S. $\ddot{\text{h}}$.	
PECTIN'ARIA. H.S.S. PECTIN'ARIA.	Pentandria Digynia. H.S.				
1 mammillàris. (br.re.) prickly. <i>Stapèlia mammillaris.</i> W.	C. B. S.	1774.	6. 8.	D.S. $\ddot{\text{h}}$.	Burm. afr. p. 27. t. 11.
2 articulàta. H.S.s. (d.r.)jointed. <i>Stapèlia articulata.</i> Mas.	_____	—	8. 9.	D.S. $\ddot{\text{h}}$.	Mass. stap. 20. t. 30.
O'RBEA. H.S. O'RBEA. Pentandria Digynia. H.S.					
1 maculòsa. H.S. (y.r.) spotted.	C. B. S.	1804.	6. 9.	D.S. $\ddot{\text{h}}$.	Bot. mag. t. 1833.
2 mixta. H.S. (vi.ye.) mixed. <i>Stapèlia mixta.</i> Mas.	_____	1795.	—	D.S. $\ddot{\text{h}}$.	Mass. stap. 23. t. 38.
3 quinquenérvia. H.S.(s.p.)five-nerved.	_____	1810.	10. 12.	D.S. $\ddot{\text{h}}$.	
4 trisulca. J.S. (ye.pu.) trisulcate.	_____	1806.	—	D.S. $\ddot{\text{h}}$.	Jacq. stap. c. io.
5 bisulca. H.S. (s.p.) bisulcate.	_____	—	—	D.S. $\ddot{\text{h}}$.	
6 variegàta. H.S. (s.p.) variegated. <i>Stapèlia variegata.</i> L. non B.M.	_____	1727.	6. 10.	D.S. $\ddot{\text{h}}$.	Moris. s. 15. t. 3. f. 4.
7 Curtisi. H.S. (s.p.) Curtis's. <i>Stapèlia variegata.</i> B.M. non L.	_____	1787.	—	D.S. $\ddot{\text{h}}$.	Bot. mag. t. 26.
8 lépida. H.S.s. (st.cr.) pretty.	_____	1805.	7. 8.	D.S. $\ddot{\text{h}}$.	Jacq. stap. c. ic.
9 orbicularis. H.S.s. (br.) large orbred.	_____	1799.	7. 11.	D.S. $\ddot{\text{h}}$.	Andr. reposit. t. 448.
10 bufònica. H.S. (y.bk.) toad-like. <i>Stapèlia Bufonis.</i> B.M.	_____	1806.	8. 10.	D.S. $\ddot{\text{h}}$.	Bot. mag. t. 1676.
11 clypéata. H.S.s. (y.p.) shield-flowered.	_____	1799.	6. 11.	D.S. $\ddot{\text{h}}$.	Jacq. stap. c. ic.
12 planiflòra. H.S.s.(ye.bk.)flat-flowered.	_____	1805.	7. 11.	D.S. $\ddot{\text{h}}$.	Lodd. bot. cab. t. 191.
13 inodòra. H.S.s. (s.p.) scentless.	_____	1788.	6. 8.	D.S. $\ddot{\text{h}}$.	
14 rugòsa. s.s. (gr.br.) rugged.	_____	1805.	5. 8.	D.S. $\ddot{\text{h}}$.	Jacq. stap. c. ic.
15 Wendlandiana. R.S.(s.br.)Wendland's. <i>Stapèlia rugosa.</i> Wend. non Jacq.	_____	—	6. 9.	D.S. $\ddot{\text{h}}$.	Wendl. coll. 2. t. 52.
16 marginàta. R.S. (s.vi.) margined.	_____	1805.	—	D.S. $\ddot{\text{h}}$.	
17 conspurcàta. R.S.(s.vi.) dirtied.	_____	1795.	6. 10.	D.S. $\ddot{\text{h}}$.	Jacq. stap. c. ic.
18 normális. R.S. (s.p.) sulphur-coloured.	_____	1821.	—	D.S. $\ddot{\text{h}}$.	Bot. reg. t. 755.
19 marmoràta. R.S. (s.da.) marbled.	_____	—	—	D.S. $\ddot{\text{h}}$.	
20 retusa. H.S. (s.da.) retuse.	_____	1808.	9. 10.	D.S. $\ddot{\text{h}}$.	
21 anguinea. H.S. (y.da.) snake.	_____	1806.	8. 9.	D.S. $\ddot{\text{h}}$.	Lodd. bot. cab. t. 828.
22 picta. H.S. (st.d.p.) painted.	_____	1799.	6. 9.	D.S. $\ddot{\text{h}}$.	Bot. mag. t. 1169.
23 Woodfordiana.H.S.(s.p.)Woodford's.	_____	1805.	—	D.S. $\ddot{\text{h}}$.	
STAP'ELIA. H.S. CARRION-FLOWER. Pentandria Digynia. L.					
1 grandiflòra. R.S. (d.p.) great-flowered.	C. B. S.	1795.	9. 12.	D.S. $\ddot{\text{h}}$.	Mass. stap. 13. t. 11.
2 spectabilis. H.S. (d.p.) showy. <i>grandiflora.</i> B.M. non Mas.	_____	1802.	—	D.S. $\ddot{\text{h}}$.	Bot. mag. t. 585.
3 ambigua. R.S. (re.pu.) ambiguous. β fulva. (br.) brown-flowered. <i>ambigua varietas.</i> Jacq.	_____	1795.	6. 11.	D.S. $\ddot{\text{h}}$.	Mass. stap. 13. t. 12.
4 soròria. R.S. (re.pu.) sister.	_____	—	—	D.S. $\ddot{\text{h}}$.	Jacq. stap. c. ic.
5 elongàta. (br.) long-branched. <i>soròria varietas.</i> Jacq.	_____	1797.	6. 8.	D.S. $\ddot{\text{h}}$.	Mass. stap. 23. t. 39.
6 lunàta. (pu.) lunate-spotted. <i>soròria varietas alia.</i> Jacq.	_____	—	—	D.S. $\ddot{\text{h}}$.	Jacq. stap. c. ic.
7 pátula. R.S. (d.p.) spreading. <i>soròria.</i> Jacq. non Mas.	_____	—	D.S. $\ddot{\text{h}}$.	Jacq. stap. c. ic.
8 defléxa. B.M. (ye.re.) reflexed. <i>refléxa.</i> H.S. <i>Gonostémone defléxum.</i> s.s.	_____	1806.	—	D.S. $\ddot{\text{h}}$.	Bot. mag. t. 1890.
9 lúcida. R.S. (pu.) glossy.	_____	1821.	—	D.S. $\ddot{\text{h}}$.	
10 breviróstris.W.en.(br.)short-beaked.	_____	—	D.S. $\ddot{\text{h}}$.	
11 Massóni. H.S. (d.p.) Masson's.	_____	1808.	—	D.S. $\ddot{\text{h}}$.	
12 Astèrias. R.S. (d.p.ye.) Starfish.	_____	1795.	5. 11.	D.S. $\ddot{\text{h}}$.	Bot. mag. t. 536.
13stellàris. H.S.s. (br.p.) starry.	_____	1804.	8. 10.	D.S. $\ddot{\text{h}}$.	Jacq. stap. c. ic.
14 hirsùta. R.S. (s.d.vi.) hairy. β atra. B.R. (da.pu.) dark-flowered.	_____	1710.	6. 8.	D.S. $\ddot{\text{h}}$.	Jacq. misc. 1. t. 3.
15 lanífera. H.S.s. (d.br.) woolly-flowered.	_____	1814.	6. 9.	D.S. $\ddot{\text{h}}$.	Bot. reg. t. 756.

ASCLEPIADEÆ.

16	flavicomàta. H.S. (br.)	yellow-haired.	C. B. S.	1814.	6. 9.	D.S. $\ddot{\eta}$.	
17	hamàta. R.S.	(pu.re.) hooked.	_____	1805.	—	D.S. $\ddot{\eta}$.	Lodd. bot. cab. t. 242.
18	comàta. R.S.	(d.p.ye.) very-hairy.	_____	1821.	—	D.S. $\ddot{\eta}$.	
19	multiflòra. DC.	(pu.re.) many-flowered.	_____	—	—	D.S. $\ddot{\eta}$.	
20	rúfa. R.S.	(bl.re.) rusty brown.	_____	1795.	6. 11.	D.S. $\ddot{\eta}$.	Lodd. bot. cab. t. 239.
21	pulvinàta. R.S.	(r.p.) cushioned.	_____	1821.	—	D.S. $\ddot{\eta}$.	Bot. mag. t. 1240.
22	fissirostris. R.S. (gr.pu.)	clove-beaked.	_____	1821.	—	D.S. $\ddot{\eta}$.	Jacq. stap. c. ic.
23	concínna. R.S.	(br.) spruce.	_____	1795.	6. 8.	D.S. $\ddot{\eta}$.	Mass. stap. 15. t. 18.
24	glandulifera. R.S. (y.br.)	hairy-glanded.	_____	—	—	D.S. $\ddot{\eta}$.	
25	hispidula. R.S.	(br.) hispid.	_____	1821.	—	D.S. $\ddot{\eta}$.	
26	glanduliflòra. R.S. (st.)	gland-flowered.	_____	1795.	8. 11.	D.S. $\ddot{\eta}$.	Mass. stap. 16. t. 19.
27	acuminàta. R.S. (d.p.)	acuminated.	_____	—	7. 9.	D.S. $\ddot{\eta}$.	— 15. t. 17.
28?	cordàta. H.S. heart-shaped.	_____	1805.	6. 8.	D.S. $\ddot{\eta}$.	
29?	canéscent. H.S. canescent.	_____	1795.	—	D.S. $\ddot{\eta}$.	
30?	ophiúncula. H.S. serpent-like.	_____	1805.	—	D.S. $\ddot{\eta}$.	
TRIDE'NTEA. H.S.		TRIDE'NTEA.	Pentandria	Digynia.	H.S.		
1	gemmiflòra. H.S. (bk.)	gem-flowered.	C. B. S.	1795.	10. 11.	D.S. $\ddot{\eta}$.	Mass. stap. 14. t. 15.
2	stygia. H.S.	(bk.) Stygian.	_____	1800.	8. 10.	D.S. $\ddot{\eta}$.	Bot. mag. t. 1839.
	<i>Stapèlia gemmiflòra. B.M.</i>						
3	moschàta. H.S.S. (d.p.)	musk-scented.	_____	1799.	9. 11.	D.S. $\ddot{\eta}$.	Lodd. bot. cab. t. 1051.
	<i>Stapèlia hircòsa. Jacq. stap. c. ic.</i>						
4	véluta. H.S.	(d.p.) purple smooth.	_____	1793.	5. 11.	D.S. $\ddot{\eta}$.	Mass. stap. 15. t. 16.
5	Símsii. H.S.	(d.p.) Sims's.	_____	1800.	—	D.S. $\ddot{\eta}$.	Bot. mag. t. 1234.
	<i>Stapèlia véluta. B.M. non Mass.</i>						
6	paniculàta. R.S.	(re.) paniced.	_____	1805.	6. 9.	D.S. $\ddot{\eta}$.	
7	juvéncula. S.S.	(d.br.) short-flowered.	_____	—	D.S. $\ddot{\eta}$.	Jacq. stap. c. ic.
8	depréssà. R.S.	(re.p.) depressed.	_____	—	D.S. $\ddot{\eta}$.	— c. ic.
TROMOTRÍCHE. H.S. TROMO'TRÍCHE.		Pentandria	Digynia.	H.S.			
1	fuscàta. H.S.S. (br.)	brown-flowered.	C. B. S.	1814.	6. 10.	D.S. $\ddot{\eta}$.	Jacq. stap. c. ic.
	<i>Stapèlia fuscàta. Jacq.</i>						
2	glauca. H.S.S.	(cr.) large-glaucous.	_____	1799.	6. 11.	D.S. $\ddot{\eta}$.	Jacq. stap. c. ic.
3	revolùta. H.S.	(cr.) revolute-flowered.	_____	1790.	6. 9.	D.S. $\ddot{\eta}$.	Bot. mag. t. 724.
	<i>Stapèlia revolùta. B.M.</i>						
4	pruinòsa. H.S.	(bh.) frosted.	_____	1795.	6. 8.	D.S. $\ddot{\eta}$.	Mass. stap. 24. t. 41.
5	obliqua. R.S.	(ye.vi.) oblique-flowered.	_____	1805.	6. 9.	D.S. $\ddot{\eta}$.	
6	mutàbilis.	(gr.br.) changeable.	_____	1823.	—	D.S. $\ddot{\eta}$.	Jacq. stap. c. ic.
	β variabilis. (st.re.) variable.		_____	—	—	D.S. $\ddot{\eta}$.	— c. ic.
	<i>Stapèlia mutàbilis varietas. Jacq.</i>						
7	ciliàta.	(ye.) fringed.	_____	1795.	10. 12.	D.S. $\ddot{\eta}$.	Mass. stap. 9. t. 1.
	<i>Stapèlia ciliàta. Mass.</i>						
PODA'NTHES. H.S.		PODA'NTHES.	Pentandria	Digynia.	H.S.		
1	púlchra. H.S.	(st.) beautiful sulphur.	C. B. S.	1800.	8. 10.	D.S. $\ddot{\eta}$.	
	β major. (st.d.p.) larger.		_____	—	—	D.S. $\ddot{\eta}$.	Bot. mag. t. 786.
	<i>Stapèlia verrucosa. B.M. nec aliorum.</i>						
2	irroràta. H.S.	(st.cr.) dewy.	_____	1795.	7. 9.	D.S. $\ddot{\eta}$.	Mass. stap. 12. t. 9.
3	verrucòsa. H.S. (st.br.)	warted.	_____	—	8. 10.	D.S. $\ddot{\eta}$.	— 11. t. 8.
4	roriflùa. R.S.	(st.cr.) dew-bearing.	_____	1802.	—	D.S. $\ddot{\eta}$.	Jacq. stap. c. ic.
	<i>Stapèlia roriflùa. Jacq.</i>						
5	pulchella. H.S. (st.cr.)	beautiful.	_____	1795.	5. 11.	D.S. $\ddot{\eta}$.	Mass. stap. 22. t. 36.
6	incarnàta. S.S.	(fl.) flesh-coloured.	_____	1793.	4. 8.	D.S. $\ddot{\eta}$.	— 22. t. 34.
OB'ESIA. H.S.		O'BESIA.	Pentandria	Digynia.	H.S.		
1	serrulàta. s.s. (st.br.)	sawed.	C. B. S.	1805.	6. 8.	D.S. $\ddot{\eta}$.	Jacq. stap. c. ic.
	<i>Stapèlia serrulata. Jacq.</i>						
2	geminàta. H.S. (ye.br.)	twin-flowered.	_____	1795.	5. 11.	D.S. $\ddot{\eta}$.	Bot. mag. t. 1326.
	<i>Stapèlia geminata. B.M.</i>						
3	decoràta. H.S.	(ye.pu.) neat.	_____	—	—	D.S. $\ddot{\eta}$.	Mass. stap. 19. t. 26.
4?	árida.	(ye.br.) dry.	_____	1795.	8.	D.S. $\ddot{\eta}$.	— 21. t. 33.
	<i>Stapèlia árida. Mass.</i>						
GONOST'EMON. H.S. GONOST'EMON.		Pentandria	Digynia.	H.S.			
1	divaricatum. H.S. (fl.)	divaricated.	C. B. S.	1793.	6. 11.	D.S. $\ddot{\eta}$.	Bot. mag. t. 1007.
	<i>Stapèlia divaricata. B.M.</i>						
2	stríctum. H.S.S. (fl.)	upright.	_____	1819.	10. 12.	D.S. $\ddot{\eta}$.	Bot. mag. t. 2037.
	<i>Stapèlia strícta. B.M.</i>						
3	pálidum. R.S.	(pu.) pale-flowered.	_____	—	—	D.S. $\ddot{\eta}$.	Wendl. coll. 2. t. 51.
	<i>Stapèlia pálida. Wendl.</i>						
CARUNCUL'ARIA. H.S. CARUNCUL'ARIA.		Pentandria	Digynia.	H.S.			
1	Símsii.	(re.bk.) Sims's.	C. B. S.	1790.	6. 11.	D.S. $\ddot{\eta}$.	Sims bot. mag. t. 793.
	<i>Stapèlia pedunculata. B.M. pedunculata Simsii. Jacq. stap. c. ic.</i>						

2 Massòni.	(br.) Masson's.	C. B. S.	1790.	6. 11.	D.S. $\frac{1}{2}$.	Mass. stap. p. 17. t. 21.
	<i>Stapelia pedunculata</i> . Mass. <i>pedunculata</i> <i>Massoni</i> .	Jacq. stap. c. ic.				
3 Jacquiní.	(co.) Jacquin's.	—	—	—	D.S. $\frac{1}{2}$.	Jacq. stap. c. ic.
	<i>Stapelia pedunculata</i> . Jacq.					
4 penduliflora.	(br.) pendulous-flower'd.	—	—	—	D.S. $\frac{1}{2}$.	Jacq. stap. c. ic.
	<i>Stapelia pedunculata alia</i> . Jacq.					
5 aperta.	(p.pu.) open-flowered.	—	1795.	7. 8.	D.S. $\frac{1}{2}$.	Mass. stap. 23. t. 37.
	<i>Stapelia aperta</i> . Mass.					
PIARA'NTHUS. B.	PIARA'NTHUS.	Pentandria Digynia. H.S.				
1 punctatus. R.S.	(fl.) purple dotted.	C. B. S.	1795.	7. 11.	D.S. $\frac{1}{2}$.	Mass. stap. 18. t. 24.
2 púllus. R.S.	(d.p.) many-flowered.	—	1774.	8. 9.	D.S. $\frac{1}{2}$.	Bot. mag. t. 1648.
3 ramosus.	(bk.) branched.	—	1795.	6. 8.	D.S. $\frac{1}{2}$.	Mass. stap. 21. t. 32.
	<i>Stapelia ramosa</i> . Mass.					
4 parviflorus. s.s.	(pu.) small-flowered.	—	—	—	D.S. $\frac{1}{2}$.	Mass. stap. 22. t. 35.
5 piliferus.	(bk.) hairy-tubercl.	—	1799.	—	D.S. $\frac{1}{2}$.	— 17. t. 23.
HO'ODIA.	HO'ODIA.	Pentandria Digynia.				
Gordoni.	(br.) Gordon's.	C. B. S.	1796.	—	D.S. $\frac{1}{2}$.	Mass. stap. 24. t. 40.
	<i>Stapelia Gordoni</i> . Mass.					
HUE'RRIA. B.	HUE'RRIA.	Pentandria Digynia. R.S.				
1 reticulata. h.s. (d.pu.) netted.	—	C. B. S.	1793.	7. 8.	D.S. $\frac{1}{2}$.	Bot. mag. t. 1662.
	<i>Stapelia reticulata</i> . B.M.					
β deformed. J.S.(wh.pu.) deformed.	—	—	—	—	D.S. $\frac{1}{2}$.	Jacq. stap. c. ic.
2 campanulata. h.s.(st.pu.) bell-shaped.	—	—	1795.	7. 10.	D.S. $\frac{1}{2}$.	Bot. mag. t. 1227.
3 clavigera. h.s.s.(st.cr.) club-haired.	—	—	—	—	D.S. $\frac{1}{2}$.	— t. 1661.
	<i>Stapelia campanulata</i> . B.M. nec aliorum.					
4 venusta. h.s. (st.re.) showy.	—	—	—	6. 7.	D.S. $\frac{1}{2}$.	Mass. stap. 10. t. 3.
5 lentiginosa. h.s.(st.pu.) freckled.	—	—	—	7. 11.	D.S. $\frac{1}{2}$.	Bot. mag. t. 506.
6 guttata. h.s. (st.cr.) red-spotted.	—	—	—	8. 11.	D.S. $\frac{1}{2}$.	Mass. stap. 10. t. 4.
7 humiliis. h.s. (st.pu.) humble.	—	—	—	—	D.S. $\frac{1}{2}$.	— 10. t. 5.
8 tubatæ. h.s. (st.re.) tube-flowered.	—	—	1805.	—	D.S. $\frac{1}{2}$.	Jacq. stap. c. ic.
9 duodecimfida. J.S.(st.re.) eleven-cleft.	—	—	—	—	D.S. $\frac{1}{2}$.	— c. ic.
10 crispa. h.s. (st.pu.) curled.	—	—	—	—	D.S. $\frac{1}{2}$.	—
11 barbata. h.s. (st.pu.) bearded.	—	—	1795.	—	D.S. $\frac{1}{2}$.	Mass. stap. 11. t. 7.
12 ocellata. r.s. (st.pu.) eyed.	—	—	1816.	—	D.S. $\frac{1}{2}$.	—
BRACHYSTE'LMA. B.R. BRACHYSTE'LMA.	Pentandria Digynia. S.S.					
tuberösüm. B.R. (pu.) tuberous-rooted.	C. B. S.	1820.	4. 10.	S. $\frac{1}{2}$.	Bot. reg. t. 722.	
CARALL'UMA. H.S. CARALL'UMA.	Pentandria Digynia. S.S.					
1 adscéndens. h.s.s. (pu.) ascending.	E. Indies.	1804.	7. 11.	D.S. $\frac{1}{2}$.	Roxb. cor. 1. t. 30.	
2 umbellata. h.s. (fl.) umbelled.	—	—	—	D.S. $\frac{1}{2}$.	— 3. t. 241.	
3 crenulata. Wa. (ye.pu.) crenulate.	—	1824.	6. 9.	D.S. $\frac{1}{2}$.	Wall pl. as. rar. t. 7.	
4 fimbriata. Wa. (pu.) fringed.	—	—	—	D.S. $\frac{1}{2}$.	— t. 8.	
HO'YA. B.P.	HO'YA.	Pentandria Digynia. R.S.				
1 crassiñolia. h.s. (wh.) large-leaved.	China.	1817.	10. 1.	D.S. $\frac{1}{2}$.	—	
2 carnosa. H.K. (bh.) fleshy-leaved.	—	1802.	7. 10.	D.G. $\frac{1}{2}$.	Bot. mag. t. 788.	
3 pálida. B.R. (bh.) pale.	—	1820.	—	D.S. $\frac{1}{2}$.	Bot. reg. t. 951.	
acuta. H.R.						
4 Póttii. H.T. (bh.) Potts's.	—	1822.	—	D.S. $\frac{1}{2}$.	—	
5 trinervis. H.T. (bh.) three-nerved.	—	1824.	—	D.S. $\frac{1}{2}$.	—	
6 lanceolata. D.P. (bh.) spear-leaved.	Nepaul.	1825.	—	D.G. $\frac{1}{2}$.	—	
7 viridiñflora. B.P. (gr.) green-flowered.	E. Indies.	—	—	S. $\frac{1}{2}$.	Rheed. mal. 9. t. 15.	
FISCH'ERIA. DC.	FISCH'ERIA.	Pentandria Digynia.				
scándens. DC. (gr.ye.) climbing.	S. America.	1826.	—	S. $\frac{1}{2}$.	DC.hort.mon.ined.t.67.	
TYLO'PHORA. B.P.	TYLO'PHORA.	Pentandria Digynia. R.S.				
1 grandiflora. B.P. (br.gr.) great-flowered.	N. S. W.	1822.	6. 8.	G. $\frac{1}{2}$.	—	
2 barbata. B.P. (gr.br.) bearded.	—	—	—	G. $\frac{1}{2}$.	—	
3 exilis. L.T. (pu.) slender.	E. Indies.	1823.	—	S. $\frac{1}{2}$.	Linn. trans. 12. t. 16.	
CEROP'EGIA. R.S.	CEROP'EGIA.	Pentandria Digynia. R.S.				
1 aphylla. h.s. (st.) leafless.	Canaries.	1808.	—	S. $\frac{1}{2}$.	—	
2 stapediæformis. H.P. (br.) Stapedia-like.	C. B. S.	1826.	7. 8.	D.S. $\frac{1}{2}$.	—	
3 torulosæ. H.R. (st.) straddling.	—	1821.	—	D.S. $\frac{1}{2}$.	—	
4 dichotoma. H.R. (st.) forked.	E. Indies.	1806.	8.	D.S. $\frac{1}{2}$.	—	
5 júncea. R.S. (st.pu.) rushy.	—	1820.	4. 6.	D.S. $\frac{1}{2}$.	Roxb. cor. 1. t. 10.	
6 africana. B.R. (br.) African.	C. B. S.	1821.	4. 11.	D.S. $\frac{1}{2}$.	Bot. reg. t. 626.	
7 bulbosa. R.S. (pu.) bulbous-rooted.	E. Indies.	—	—	D.S. $\frac{1}{2}$.	Roxb. cor. 1. t. 7.	
8 tuberosa. r.s. (pu.) tuberous-rooted.	—	—	—	D.S. $\frac{1}{2}$.	— 1. t. 9.	
PERGUL'ARIA. R.S.	PERGUL'ARIA.	Pentandria Digynia. R.S.				
1 odoratissima. r.s. (st.) large.	E. Indies.	1784.	6. 7.	S. $\frac{1}{2}$.	Bot. reg. t. 412.	
2 minor. r.s. (st.) smaller.	—	1790.	5. 8.	S. $\frac{1}{2}$.	Bot. mag. t. 755.	
3 sanguinolenta. B.M. (st.) bloody-juiced.	Sierra Leon.	1822.	8. 9.	S. $\frac{1}{2}$.	— t. 2532.	

MARSD'ENIA. B.P.	MARSD'ENIA.	Pentandria Digynia. R.S.						
1 suavèolens. B.P. (<i>wh.</i>) sweet-scented.	N. S. W.	1816.	3. 8.	G. h.~.	Bot. reg. t. 489.			
2 erécta. R.S. (<i>wh.</i>) upright.	Syria.	1597.	7. 8.	G. h.~.	Jacq. vind. 1. t. 337.			
DISCHI'DIA. B.P.	DISCHI'DIA.	Pentandria Digynia. S.S.						
bengalénis. L.T. (<i>st.</i>) Bengal.	E. Indies.	1818.	6. 10.	S. h.~.	Bot. mag. t. 2916.			
GYMN'EMA. B.P.	GYMN'EMA.	Pentandria Digynia. R.S.						
1 sylvestre. R.S. (<i>wh.</i>) netted-leaved.	Ceylon.	1816.	6. 8.	S. h.~.	Willd. phyt. 1. t. 5. f. 3.			
2 tingens. s.s. (<i>ye.</i>) pointed-leaved.	E. Indies.	1823.	—	S. h.~.	Roxb. corom. 3. t. 239.			
<i>Asclepias tingens.</i> R.C.								
3 tenacissimum. s.s. (<i>ye.</i>) tough.		1806.	—	S. h.~.	Roxb. corom. 3. t. 240.			
SARCO'LOBUS. R.S.	SARCO'LOBUS.	Pentandria Digynia. R.S.						
1 globosus. R.S. (<i>wh.</i>) globular-fruit'd.	E. Indies.	1823.	—	S. h.~.	Wall. as. res. 12. t. 4.			
2 carinatus. R.S. (<i>gr.ye.</i>) keeled-fruited.		—	—	S. h.~.	—	—	—	12. t. 5.
GONO'LOBUS. R.S.	GONO'LOBUS.	Pentandria Digynia. R.S.						
1 marítimus. R.S. (<i>gr.p.</i>) sea-side.	Caracas.	1823.	6. 10.	S. h.~.	Bot. reg. t. 931.			
2 suberosus. R.S. (<i>pu.</i>) cork-barked.	America.	1732.	7. 9.	S. h.~.	Dill. elt. t. 229. f. 296.			
3 crisipiflórus. R.S. (<i>wh.gr.</i>) curled-flower'd.	S. America.	1741.	7. 8.	S. h.~.	Plum. ic. t. 216. f. 1.			
4 guianénis. s.s. (<i>gr.</i>) green-flowered.	Guiana.	1826.	7. 9.	S. h.~.				
<i>viridiflórus.</i> R.S. non Nutt.	<i>Cynáncum viridiflórum.</i>	Meyr.						
5 grandiflórus. R.S. (<i>ye.</i>) large-flowered.	S. America.	1825.	8. 10.	S. h.~.	Bot. reg. t. 1053.			
6 rostrátus. R.S. (<i>gr.</i>) beaked-petal'd.	Trinidad.	1823.	—	S. h.~.				
7 niger. R.S. (<i>bk.</i>) black-flowered.	Mexico.	1825.	9. 11.	G. y.~.	Bot. mag. t. 2799.			
<i>Cynáncum nigrum.</i> Cav. ic. 2. t. 159.								
8 uniflórus. K.S. (<i>pu.</i>) solitary-flowered.		—	—	S. h.~.	H. et B. 3. t. 238.			
9 carolinénis. R.S. (<i>d.p.</i>) blunt-flowered.	N. America.	1820.	7. 9.	H. y.~.	Jacq. ic. 2. t. 342.			
10 hirsutus. R.S. (<i>d.pu.</i>) hairy.		1806.	6. 9.	H. y.~.	Swt. br. fl. gar. t. 1.			
11 macrophyllus. N. (<i>p.pu.</i>) large-leaved.		1822.	—	H. y.~.				
12 discolor. R.S. (<i>d.pu.</i>) Virginian.		1809.	7. 9.	H. y.~.	Bot. mag. t. 1273.			
<i>Cynáncum discolor.</i> B.M.								
13 obliquus. R.S. (<i>pu.</i>) oblique-flower'd.	Carolina.	1818.	—	H. y.~.	Jacq. ic. 2. t. 341.			
14 Baldwyniánus. (<i>st.</i>) Baldwyn's.	Savannah.	1822.	7. 9.	H. y.~.	Lodd. bot. cab. t. 365?			
<i>hirsutus.</i> L. bot. cab. ? nec aliorum. <i>caroliniánus.</i> N. non Jacq.								
15 hirtus. R.S. (<i>pu.</i>) hairy-leaved.	America.	—	—	S. h.~.	Moris. his. 3. s. 15. t. 3. f. 61			
16 undulátus. R.S. (<i>br.pu.</i>) wave-leaved.	S. America.	1814.	—	S. h.~.	Jacq. amer. 85. t. 58.			
17 viridiflórus. N. (<i>gr.</i>) Nuttall's.	Mississippi.	1822.	—	H. y.~.				
<i>Nuttallianus.</i> s.s.								
18 l'ævis. R.S. (<i>d.pu.</i>) smooth.		1806.	6. 7.	H. y.~.				
19 diadématus. B.R. (<i>st.</i>) red-crowned.	Mexico.	1812.	9. 10.	S. h.~.	Bot. reg. t. 252.			
20 barbátus. R.S. (<i>ye.</i>) bearded.		1826.	—	G. h.~.	H. et B. N. gen. 3. t. 239.			
21 mucronátus. (<i>br.re.</i>) sharp-pointed.	Trinidad.	1804.	7. 8.	S. h.~.	Andr. deposit. t. 515.			
<i>Cynáncum mucronatum.</i> A.R. non K.S.								
ENSL'ENIA. N.	ENSL'ENIA.	Pentandria Digynia.						
álbida. N.	(<i>st.</i>) straw-coloured.	Virginia.	1828.	7. 9.	H. y.~.			
ASCL'EPIAS. R.S.	SWALLOW-WORT.	Pentandria Digynia. R.S.						
1 syriaca. Lam. (<i>li.</i>) Syrian.	Astracan.	1629.	7. 8.	H. y.~.	Blackw. t. 521.			
<i>Beid el Ossar.</i> Veslingius. Linn. trans. v. 14. p. 239.								
2 obovata. E.C. (<i>pu.</i>) obovate-leaved.	Carolina.	1826.	—	H. y.~.				
3 tomentosa. E.C. (<i>pu.</i>) woolly.		—	—	H. y.~.				
4 exaltata. L.en. (<i>pu.</i>) tall.	N. America.	—	—	H. y.~.				
5 phytolaccoides. R.S. (<i>pu.</i>) Phytolacca-like.		1812.	—	H. y.~.				
6 parviflóra. R.S. (<i>wh.</i>) small flowered.		1774.	7. 10.	H. y.~.	Jacq. f. ecl. t. 28.			
7 salicifólia. B.C. (<i>wh.</i>) Willow-leaved.	S. America.	1816.	6. 9.	G. y.~.	Lodd. bot. cab. t. 272.			
8 angustifólia. R.S. (<i>fl.</i>) narrow-leaved.	Mexico.	1824.	—	F. y.~.				
9 curassávica. R.S. (<i>or.</i>) Curassavian.	S. America.	1692.	6. 10.	S. y.~.	Bot. reg. t. 81.			
10 nivea. R.S. (<i>wh.</i>) Almond-leaved.	N. America.	1730.	7. 9.	H. y.~.	Bot. mag. t. 1181.			
11 virgata. R.S. (<i>li.</i>) slender.	Mexico.	1825.	—	F. y.~.				
12 incarnata. R.S. (<i>pu.</i>) flesh-coloured.	N. America.	1710.	7. 8.	H. y.~.	Bot. reg. t. 250.			
13 púlchra. R.S. (<i>pu.</i>) hairy.		—	—	H. y.~.	Swt. br. fl. gar. s. 2. t. 18.			
14 am'œna. R.S. (<i>pu.</i>) oval-leaved.		1732.	—	H. y.~.	Dill. elt. t. 27. f. 30.			
15 purpurascens. R.S. (<i>pu.</i>) purple.		—	—	H. y.~.	— t. 28. f. 31.			
16 citrifólia. R.S. (<i>wh.</i>) Citron-leaved.	Florida.	1812.	—	F. y.~.	Jacq. icon. t. 345.			
17 variegata. R.S. (<i>wh.re.</i>) variegated.		1597.	—	H. y.~.	Bot. mag. t. 1182.			
18 obtusifólia. R.S. (<i>pu.</i>) blunt-leaved.		1823.	—	H. y.~.				
19 amplexicaulis. R.S. (<i>re.</i>) stem-clasping.		1812.	—	H. y.~.				
20 laurifolia. M. (<i>pu.</i>) Laurel-leaved.		1823.	—	H. y.~.	Pluk. amat. t. 358. f. 2.			
<i>periplocæfólia.</i> N.								
21 acuminata. R.S. (<i>wh.pu.</i>) taper-pointed.		1812.	—	H. y.~.				
22 paupérula. R.S. (<i>re.</i>) remote-leaved.		—	—	6. 7. H. y.~.				
23 pedicellata. R.S. (<i>gr.ye.</i>) dwarf.		—	—	H. y.~.				

24 adrifolia. R.S. (bh.) four-leaved.	N. America.	1823.	8. 9. H. 4.	Jacq. obs. 2. t. 33.
25 aciculata. R.S. (bh.) whorl-leaved.	1759.	8. 10. H. 4.	Swt. br. fl. gar. t. 144.	
26 mexicana. R.S. (wh.) Mexican.	Mexico.	1823.	7. 9. G. ½.	Cavan. ic. 1. t. 58.
27 rosea. K.S. (ro.) rose-coloured.	1824.	— G. 4.		
28 linifolia. R.S. (wh.) Flax-leaved.	—	— G. ½.		
29 Linaria. R.S. (wh.) toadflax-leaved.	1802.	— G. 4.	Cavan. ic. 1. t. 57.	
30 rufa. R.S. (re.) red-flowered.	N. America.	—	H. 4.	
31 tuberosa. R.S. (or.) tuberous-rooted.	1690.	7. 9. H. 4.	Bot. reg. t. 76.	
32 decumbens. w. (or.) decumbent.	1731.	— H. 4.	Swt. br. fl. gar. s. 2.t.24.	
ACER'ATES. E.C. ACER'ATES. Pentandria Dignynia.				
longifolia. E.C. (p.pu.) long-leaved.	N. America.	1816.	7. 8. H. 4.	
Asclepias longifolia. R.S. <i>Gomphocarpus longifolius</i> . s.s.				
ANANTHERIX. N. ANANTHERIX. Pentandria Dignynia. N.				
viridis. n. (gr.) green-flowered.	N. America.	1812.	8. 10. H. 4.	
STYLA'NDRA. N. STYLA'NDRA. Pentandria Dignynia. N.				
pumila. n. (st.) dwarf.	Carolina.	1812.	7. 8. G. 4.	
<i>Podostigma pubescens</i> , s.s. <i>Asclepias pedicellata</i> , Ph.				
GOMPHOCA'Rpus. R.S. GOMPHOCA'Rpus. Pentandria Dignynia. R.S.				
1 arborescens. R.S. (wh.) broad-leaved.	C. B. S.	1714.	12. G. ½.	Jacq. schœnb. 1. t. 50.
2 crispus. R.S. (pu.) curled-leaved.	—	1774.	7. G. ½.	Comm. rar. t. 17.
3 fruticosus. R.S. (wh.) Willow-leaved.	—	1714.	6. 9. G. ½.	Bot. mag. t. 1628.
<i>Asclepias fruticosa</i> . w.				
OXYSTE'LMA. R.S. OXYSTE'LMA. Pentandria Dignynia. R.S.				
esculentum. R.S. (wh.ro.) esculent.	E. Indies.	1816. S. 4. —	Roxb. corom. 1. t. 11.
XYSMAL'OBium. R.S. XYSMAL'OBium. Pentandria Dignynia. R.S.				
1 undulatum. R.S. (gr.) waved-leaved.	C. B. S.	1783.	7. 8. G. ½.	Comm. rar. t. 16.
2 grandiflorum. R.S. (pu.) large-flowered.	—	1823.	— G. ½.	
CALOTROPIS. R.S. CALOTROPIS. Pentandria Dignynia. R.S.				
1 præcera. R.S. (wh.) bell-flowered.	Persia.	1714.	7. 9. S. ½.	Andr. reposit. t. 271.
<i>Asclepias gigantea</i> . A.R.				
2 gigantea. R.S. (p.pu.) curled-flower'd.	E. Indies.	1690.	— S. ½.	Bot. reg. t. 58.
OXYPET'ALUM. R.S. OXYPET'ALUM. Pentandria Dignynia.				
1 appendiculatum. M.N. appended.	Brazil.	1824.	6. 8. S. ½. —	Mart. n. pl. bras. t. 30.
• 2 Banksii. R.S. Banks's.	—	1826.	— S. ½. —	t. 29.
CYNA'NCHUM. R.S. CYNA'NCHUM. Pentandria Dignynia. L.				
1 acutum. R.S. (wh.) acute-leaved.	Europe.	1596.	7. H. 4. —	Flor. græc. t. 250.
2 monspeliacum. R.S. (bh.) Montpellier.	S. Europe.	—	8. 9. H. 4. —	t. 251.
3 pauciflorum. R.S. (gr.w.) few-flowered.	E. Indies.	1822.	5. 8. S. ½. —	Will.phy.l.n.23.t.5.f.3.
<i>Periplœca tunicata</i> . w.				
4 pilosum. R.S. (wh.) hairy.	C. B. S.	1726.	6. 9. G. ½. —	Bot. reg. t. 111.
<i>Periplœca africana</i> . A.R. 557.				
5 crassifolium. R.S. (gr.w.) thick-leaved.	—	1816. G. ½. —	
6 capense. R.S. (wh.) Cape.	—	1820. G. ½. —	
7 roseum. R.S. (ro.) rosy.	Dahuria.	1816.	7. 8. H. 4.	Gmel. sib. 4. t. 42.
<i>Asclepias davurica</i> . w.				
8 nigrum. R.S. (bk.) black.	S. Europe.	1596.	5. 8. H. 4. —	Bot. mag. t. 2390.
9 medium. R.S. (br.) middle.	—	—	— H. 4. —	
10 Vincetoxicum. R.S. (w.) officinal.	Europe.	1596.	— H. 4.	Flor. dan. t. 849.
<i>Asclepias Vincetoxicum</i> . w.				
11 fuscatum. s.s. (st.) yellow-flowered.	—	—	H. 4.	
12 sibiricum. R.S. (st.) Siberian.	Siberia.	1775.	7. 8. H. 4.	Murr. gott. 2. t. 7.
13 fœtidum. R.S. (st.) fetid.	Mexico.	1823.	— G. ½. —	Cavan. ic. 2. t. 158.
14 viridiflorum. B.M. (gr.re.) green-flowered.	E. Indies.	1814.	10. 12. S. ½. —	Bot. mag. t. 1929.
15 undatum. A.R. (gr.) wave-leaved.	—	1803.	7. 8. S. ½. —	Andr. reposit. t. 410.
HARRISONIA. B.M. HARRISONIA. Pentandria Dignynia.				
loniceroides. B.M. (cr.) Honeysuckle.	Brazil.	1824.	9. 10. S. ½.	Bot. mag. t. 2699.
DÆMIA. R.S. DÆMIA. Pentandria Dignynia. R.S.				
1 extensa. R.S. (st.) smooth-leaved.	E. Indies.	1778.	7. 8. S. ½. —	Jacq. ic. 1. t. 54.
2 bicolor. (gr.pu.) two-coloured.	—	1806.	— S. ½. —	Andr. reposit. t. 562.
<i>Cynanchum bicolor</i> . A.R.				
DIPLOL'EPIS. R.S. DIPLOL'EPIS. Pentandria Dignynia.				
1 vomitoria. H.T. (st.) dull-leaved.	China.	1820.	6. 9. S. ½. —	
2 apiculata. H.T. (st.) glossy-leaved.	—	—	S. ½. —	
SARCOSTE'MMA. R.S. SARCOSTE'MMA. Pentandria Dignynia. R.S.				
1 viminalis. R.S. (wh.) twisting.	E. Indies.	1731.	7. S. ½. —	
2 Swartzianum. R.S. (wh.) Swartz's.	W. Indies.	—	— S. ½. —	Sloan. hist. 1.t.131.f.1.
EUST'EGIA. R.S. EUST'EGIA. Pentandria Dignynia. R.S.				
hastata. R.S. (gr.wh.) halberd-leaved.	C. B. S.	1816.	6. 8. G. ½.	Spr. n. ent. t. 1. f. 5-10.
<i>Apocynum hastatum</i> . w.				

METASTE'LMA. R.S.	METASTE'LMA.	Pentandria Digynia.				
parviflórum. r.s.(wh.)	small-flowered.	W. Indies.	1826.	5. 6.	S. h. .	Plum. ic. 215. f. 1.
MICROL'OMA. R.S.	MICROL'OMA.	Pentandria Digynia. R.S.				
1 sagittátum. r.s. (sc.)	arrow-leaved.	C. B. S.	1775.	7. 8.	G. h. .	Jacq. schoen. 1. t. 38.
2 lineáre. r.s. (cr.)	linear-leaved.	1816.			G. h. .	Pluk. man. 17. t. 335. f. 5.
ASTEPHA'NUS. R.S.	ASTEPHA'NUS.	Pentandria Digynia. R.S.				
1 triflórus. r.s. (wh.)	three-flowered.	C. B. S.	1816.	7. 9.	G. 2. .	
2 lineáris. r.s. (wh.)	panicked.	1822.			G. 2. .	
3 lanceolátus. r.s. (wh.)	spear-leaved.	1822.			G. 2. .	

ORDO CXL.

GENTIANEÆ. *Brown prodr. 1. p. 449.*

GENTIA'NA. R.S.	GENTIAN.	Tetr-Pentandria Digynia.				
1 lútea. R.S.	(ye.) yellow.	Alps Europe.	1596.	6. 7.	H. 2. .	Mill. ic. t. 139.
2 hy'brida. R.S.	(ye.re.) hybrid.	Switzerland.	1825.	—	H. 2. .	
3 purpùrea. R.S.	(pu.) purple.	—	1768.	—	H. 2. .	Andr. reposit. t. 117.
4 pannònica. R.S. (pa.pu.)	round-petaled.	—	—	—	H. 2. .	Jacq. aust. 2. t. 136.
β albiflòra.	(wh.) white-flowered.	1826.	—	—	H. 2. .	
5 punctàta. R.S. (st.pu.)	dotted-flowered.	1775.	—	—	H. 2. .	Jacq. aust. 5. t. ap. 28.
6 campanulàta. R.S. (ye.)	bell-flowered.	1824.	—	—	H. 2. .	— ap. t. 29.
7 septémfida. R.S. (bl.)	crested.	Levant.	1804.	—	H. 2. .	Bot. mag. t. 1229.
β guttátâ.	(bl.wh.) spotted.	—	—	—	H. 2. .	— t. 1410.
8 asclepiadèa. R.S. (bl.)	swallowwort-ld.	Austria.	1629.	7. 8.	H. 2. .	— t. 1078.
9 cruciàta. R.S. (bl.)	Crosswort.	—	1596.	6. 7.	H. 2. .	Jacq. aust. 4. t. 372.
10 macrophylla. R.S. (bl.)	long-leaved.	Siberia.	1796.	7. 8.	H. 2. .	Bot. mag. t. 1414.
11 adscéndens. s.s.	(bl.) ascending.	—	1799.	6. 7.	H. 2. .	— t. 705.
β decúmbens. R.S. (bl.)	decumbent.	—	—	—	H. 2. .	— t. 723.
12 frígida. R.S. (wh.)	frigid.	Carpathian m.	1816.	—	H. 2. .	Froel. gent. n. 13. ic.
13 algída. R.S. (st.bl.)	narrow-leaved.	Siberia.	1808.	—	H. 2. .	Pall. ros. 2. t. 95.
14 gélida. R.S. (st.)	cream-coloured.	Caucasus.	1816.	—	H. 2. .	
15 Pneumonánthe. R.S. (bl.)	Calathian violet.	England.	8. 9.	H. 2. .	Eng. bot. v. 1. t. 20.
β guttátâ. B.M. (bh.wh.)	spotted-flowered.	—	—	H. 2. .	Bot. mag. t. 1101.
16 pseudo-pneumonanthe. R.S. (bl.)	Canadian.	N.America.	1828.	—	H. 2. .	
	Pneumonánthe. Ph.					
17 lineáris. R.S.	(bl.) linear-leaved.	—	1816.	—	H. 2. .	
18 triflòra. R.S.	(p.bl.) three-flowered.	Siberia.	1807.	6. 7.	H. 2. .	Pall. ross. 1. t. 93. f. 1.
19 Saponària. B.M.	(bl.) barrel-flower'd.	N. America.	1776.	8. 9.	H. 2. .	Bot. mag. t. 1039.
20 Catesb'æi. N.	(bl.) Catesby's.	—	—	6. 8.	H. 2. .	Andr. reposit. t. 418.
21 intermédia. B.M. (pu.)	intermediate.	—	1820.	8. 9.	H. 2. .	Bot. mag. t. 2303.
22 incarnâta. B.M.	(fl.) flesh-coloured.	—	1812.	9. 11.	H. 2. .	— t. 1856.
23 ochroleúca. R.S.	(st.) pale-flowered.	—	1803.	8. 9.	H. 2. .	— t. 1551.
24 quinqueflòra. R.S. (pu.)	five-flowered.	—	1823.	7. 9.	H. ♂.	
	amareloides. M.					
25 àurea. R.S.	(wh.ye.) large involucred.	Norway.	—	—	H. ♂.	Flor. dan. t. 344.
quinquefòlia. F.D.	(involuta.)	Rottb. ac. havn. 10. p. 434. t. 1. f. 2. A.B.				
26 umbellàta. M.B. (pa.bl.)	umbel-flowered.	Caucasus.	1823.	8. 10.	H. ⓧ.	
27 acáulis. R.S.	(bl.) dwarf.	Wales.	3. 6.	H. 2. .	Eng. bot. v. 23. t. 1594.
28 alpina. R.S.	(bl.) Alpine.	Switzerland.	1818.	—	H. 2. .	Lodd. bot. cab. t. 476.
29 altaïca. R.S.	(bl.) Altay.	Siberia.	1826.	—	H. 2. .	Pall. ross. 2. t. 97. f. 1.
30 pyrenâica. R.S.	(bl.) Pyrenean.	Pyrenees.	1825.	—	H. 2. .	W. et K. hung. t. 207.
31 pùmila. R.S.	(bl.) small.	Switzerland.	—	—	H. 2. .	Jacq. obs. 2. t. 49.
32 vérna. R.S.	(bl.) spring.	England.	—	H. 2. .	Eng. bot. v. 7. t. 493.
33 aestìva. R.S.	(bl.) summer.	Germany.	1826.	—	H. 2. .	Rœm. arch. 1. t. 3. f. 8.
34 bavârica. R.S.	(bl.) Bavarian.	—	1775.	7.	H. 2. .	Vill. delph. 2. t. 10.
35 imbricâta. R.S.	(bl.) imbricated	Switzerland.	1825.	—	H. 2. .	Barrel. ic. 101. f. 2.
36 angulòsa. R.S.	(bl.) angular-calyx'd.	Caucasus.	1826.	4. 6.	H. 2. .	Marsch. cent. rar. 1. t. 47.
37 utriculòsa. R.S.	(bl.) bladdered.	—	—	7. 9.	H. ♂.	W. et K. hung. t. 206.
38 nivalis. R.S.	(bl.) small alpine.	Scotland.	8. 9.	H. ⓧ.	Eng. bot. v. 13. t. 896.
39 aquática. R.S.	(bl.) water-side.	Dahuria.	1827.	7. 9.	H. ♂.	

40	hùmìlis. R.S.	(pa.bll.) dwarf.	Caucasus.	1827.	7. 9.	H. ♂.	Pall. ross. 2. t. 97. f. 2.	
41	uniflòra. R.S.	(vi.) one-flowered.	M. Carpathian.	1828.	—	H. ♂.	Rœm. ar. 1.p. 23.t. 2.f. 4.	
42	germánica. R.S.	(vi.) German.	Germany.	1825.	8. 9.	H. ♂.	Mayrh. fl. mon. t. 278.	
43	Amarélla. R.S.	(vi.) autumnal.	Britain.	8. 10.	H. ♂.	Eng. bot. v. 4. t. 236.	
44	uliginòsa. R.S.	(bl.) marsh.	Germany.	1827.	—	H. ♂.	Rœm. ar. 1.p. 22.t. 2.f. 3.	
45	obtusifòlia. R.S. (bl.st.)	blunt-leaved.	Switzerland.	1828.	—	H. ♂.	Bot. mag. t. 1038.	
46	praténisia. R.S.	(vi.) field.	Siberia.	1824.	—	H. ♂.	Eng. bot. v. 4. t. 237.	
47	caucásica. R.S.	(bl.) Caucasean.	Caucasus.	1804.	7. 8.	H. ♂.	Sch. rom. arc. 1.t. 2.f. 5.	
48	campéstris. R.S.	(vi.) field.	Britain.	8. 10.	H. ♂.	Reich. ic. t. 271.	
49	angustifòlia. R.S.	(bl.) slender-leaved.	N. America.	1820.	8. 9.	H. ♀.	Pall. ross. 2. t. 89. f. 3.	
50	glaciáris. R.S.	(bl.) northern.	N. Europe.	1826.	—	H. ♂.	Jacq. aust. t. 113.	
51	conférta. F.	(bl.) crowded.	Altay.	1827.	—	H. ♂.	Swt. br. fl. gar. t. 139.	
52	Gebléri. F.	(bl.) Gebler's.	—	—	—	H. ♂.	Bot. mag. t. 639.	
53	rotátæ. R.S.	(bl.) rotate.	Siberia.	—	—	H. ♂.	ciliáta. B.M. nec aliorum.	
54	ciliáta. R.S.	(bl.) fringe-flowered.	S. Europe.	1759.	—	H. ♀.	Reich. ic. t. 271.	
55	fimbriáta. R.S.	(bl.) fimbriate.	Caucasus.	1824.	7. 9.	H. ♀.	Pall. ross. 2. t. 89. f. 3.	
56	criníta. R.S.	(bl.) jagged-flower'd.	N. America.	1804.	7. 10.	H. ♂.	Jacq. aust. t. 113.	
	fimbriáta. Andr. rep. t. 509. non w.						Swt. br. fl. gar. t. 139.	
57	barbáta. R.S.	(bl.) bearded.	Siberia.	8. 9.	H. ♂.	Bot. mag. t. 639.	
	ciliáta. B.M. nec aliorum.							
SWERTIA. R.S.								
	FELWORT. Tetr-Pentandria Dignynia.							
1	perénnis. R.S. (br.pu.)	marsh.	England.	7. 8.	H. ♀.	Eng. bot. v. 21. t. 1441.	
2	Michauxiána. R.S. (st.)	American.	N. America.	1824.	—	H. ♂.	Pall. ross. 2. t. 91.	
3	dichótoma.R.S.(gr.re.)	forked.	Siberia.	1827.	7. 9.	H. ♂.		
CHL'ORA. S.S.								
	YELLOW-WORT. Octandria Monogynia. L.							
1	imperfoliáta. s.s. (ye.)	sessile-leaved.	Italy.	1826.	6. 7.	H. ⓧ.	Eng. bot. v. 1. t. 60.	
2	perfoliáta. s.s. (ye.)	perfoliate.	Britain.	—	H. ⓧ.		
FRAS'ERA. M.								
	FRAS'ERA. Tetrandria Monogynia. R.S.							
	Waltéri. M.	(ye.) Walter's.	N. America.	1795.	7. 8.	H. ♂.		
	carolinén sis. Walt.							
ERYTHR'EA. R.S.								
	ERYTHR'EA. Pentandria Monogynia. P.S.							
1	Centárium. R.S. (ro.)	common.	Britain.	7. 8.	H. ⓧ.	Eng. bot. v. 6. t. 417.	
2	latifòlia. s.s.	(ro.) broad-leaved.	Lancashire.	—	H. ⓧ.		
3	pulchéllea. R.S.	(ro.) dwarf branching.	England.	8. 9.	H. ⓧ.	Eng. bot. v. 7. t. 458.	
4	littorális. R.S.	(ro.) simple dwarf.	Britain.	6. 8.	H. ⓧ.	— v. 33. t. 2305.	
5	conférta. R.S.	(ro.) close-flowered.	Spain.	1818.	5. 11.	H. ♀.		
6	spicáta. R.S.	(ro.) spiked.	S. Europe.	—	H. ♂.	Flor. græc. t. 238.	
	β albiflòra. (wh.) white-flowered.	—	—	—	—	H. ♂.	Matth. comm. p. 488.f.2.	
7	austrális. B.P.	(ro.) New Holland.	New Holland.	1823.	5. 8.	G. ♂.	W. et K. pl. hung. t. 238.	
8	uliginòsa. R.S.	(ro.) marsh.	Hungary.	1825.	6. 8.	H. ♂.	Swt. br. fl. gar. t. 137.	
9	aggregátæ. B.F.G.	(ro.) cluster-branch'd.	Nepaul.	1824.	3. 11.	H. ♀.	Flor. græc. t. 237.	
10	marítima. F.G.	(ye.) sea-side.	S. Europe.	1777.	7. 8.	H. ⓧ.		
11	Massóni.	(ye.) Masson's.	Azores.	—	—	F. ♀.		
	marítima. H.K. nec aliorum, vid. obs. in Flor. Græc.							
12	lútea. R.S.	(ye.) yellow-flowered.	S. Europe.	—	H. ⓧ.	Barrel. ic. 468.	
SABB'ATIA. R.S.								
	SABB'ATIA. Pentandria Monogynia. R.S.							
1	grácilis. R.S.	(ro.) slender.	N. America.	7. 9.	H. ♂.	Sal. par. lond. t. 32.	
2	stelláris. R.S.	(ro.ye.) starry.	—	1827.	6. 8.	H. ♂.	Bartr. ic. ined. t. 13.	
3	angularis. R.S.	(pu.) angular-stalked.	—	1823.	7. 8.	H. ♂.	Bot. mag. t. 1600.	
4	calycòsa. R.S.	(ro.) dichotomous.	—	1812.	6. 8.	H. ♂.		
5	chloróides. R.S.	(ro.) Chlorla-like.	—	1817.	8. 9.	H. ♂.		
6	paniculáta. R.S.	(wh.) paniced.	—	—	—	H. ♀.		
VOHI'RIA. R.S.								
	VOHI'RIA. Pentandria Monogynia.							
1	rósea. R.S.	(ro.) rose-coloured.	Trinidad.	1822.	6. 8.	S. ♀.	Aubl. guian. 1.t. 83. f. 1.	
	Lita rósea. W.							
2	cörulea. R.S.	(bl.) blue-flowered.	—	1824.	—	S. ♀.	Aubl. guian. 1.t. 83. f. 2.	
3	uniflòra. R.S.	(ye.) one-flowered.	W. Indies.	—	—	S. ♀.	Jacq. amer. 87.t. 60.f.3.	
CHIR'ONIA. R.S.								
	CHIR'ONIA. Pentandria Monogynia. L.							
1	jasminoïdes. R.S.	(ro.) Jasmine-like.	C. B. S.	1812.	4. 7.	G. ♂.	Bot. reg. t. 197.	
2	lychnoïdes. R.S.	(pu.) Lychnis-flowered.	—	1816.	—	G. ♂.	Linn. trans. 7. t. 12. f. 3.	
3	nudicáulis. R.S.	(pu.) naked-stalked.	—	—	—	G. ♂.	Bot. mag. t. 511.	
4	linoïdes. R.S.	(pi.) flax-leaved.	—	1787.	7. 9.	G. ♂.	Linn. trans. 7. t. 12. f. 2.	
5	tetragóna. R.S.	(pu.) four-sided.	—	1822.	—	G. ♂.	Bot. mag. t. 233.	
6	baccífera. R.S.	(pi.) berry-bearing.	—	1759.	6. 7.	G. ♂.	— t. 818.	
7	angustifòlia. B.M.	(ro.) narrow-leaved.	—	1800.	6. 8.	G. ♂.	— t. 37.	
8	frutéscent. R.S.	(ro.) frutescent.	—	1756.	6. 9.	G. ♂.		
	β albiflòra. (wh.) white-flowered.	—	—	—	—	G. ♂.		
9	decussáta. R.S.	(ro.) cross-leaved.	—	1789.	—	G. ♂.	Bot. mag. t. 707.	

HIPPION. S.S.	HIPPION.	Pentandria Monogynia. S.S.			
viscōsum. s.s. (ye.) clammy.	Canaries.	1781.	6. 7. G. ♂.		
<i>Exācum viscōsum</i> . Sm. <i>Gentiana viscōsa</i> . Bot. mag. t. 2135.					Sm. ic. fas. 3. t. 18.
SEB'EA. B.P.	SEB'EA.	Tetr-Pentandria Monogynia. S.S.			
1 ovāta. B.P. (st.) oval-leaved.	N. S. W.	1820.	8. 10. G. ⓧ.		Lab. nov. hol. 1. t. 52.
2 albēns. B.P. (wh.) white-flowered.	C. B. S.	—	— G. ⓧ.	Burm. afr. t. 74. f. 4.	— t. 74. f. 5.
3 cordāta. B.P. (ye.) heart-leaved.	—	1815.	7. 8. H. ⓧ.		
CICE'NDIA. Adans.	CICE'NDIA.	Tetrandria Monogynia.			Eng. bot. v. 4. t. 235.
filiforme. (ye.) slender-stalk'd. Britain.	—	—	— H. ⓧ.		
<i>Exācum filiforme</i> . E.B. <i>Microcāle filiforme</i> . Lk.					
EX'ACUM. B.P.	EX'ACUM.	Tetrandria Monogynia. L.			
tetragōnum. D.P. (ye.) square-stalked.	Nepaul.	1820.	8. 10. H. ⓧ.		
COUTOUB'EA. P.S.	COUTOUB'EA.	Tetrandria Monogynia.			Cavan. icon. 4. t. 328.
1 ternifolia. p.s. (wh.) three-leaved.	Trinidad.	1820.	6. 10. S. ⓧ.	Aubl. guian. 1. t. 28.	
2 ramōsa. Aub. (pu.) purple-flowered.	—	1823.	— S. ⓧ.		
<i>purpūrea</i> . Lam. ill. t. 79. <i>Exācum ramōsum</i> . R.S.					
MITRASA'CME. B.P.	MITRASA'CME.	Tetrandria Monogynia. R.S.			
1 polymorpha. B.P. (wh.) variable.	N. S. W.	1822.	6. 8. G. ⓧ.		Bot. mag. t. 370.
2 canescens. B.P. (wh.) canescent.	—	1820.	— G. ⓧ.		— t. 2822.
3 serpylliifolia. B.P. (wh.) creeping.	—	1826.	— G. ⓧ.		
HOUSTONIA. R.S.	HOUSTONIA.	Tetrandria Monogynia. L.			
1 cœrulea. R.S. (pa.bl.) blue-flowered.	N. America.	1785.	5. 8. H. ⓧ.		
2 serpylliifolia. R.S. (wh.) Thyme-leaved.	—	1827.	— H. ⓧ.		
3 temēlla. R.S. (pu.) slender.	Carolina.	1812.	— H. ⓧ.		
4 albibiflora. N. (wh.) white-flowered.	N. America.	1828.	— H. ⓧ.		
5 purpūrea. R.S. (li.) purple-flowered.	—	1800.	— H. ⓧ.		
SPIG'ELIA. R.S.	WORM-GRASS.	Pentandria Monogynia. L.			
1 Anthēmia. R.S. (bl.) annual.	W. Indies.	1759.	7. S. ⓧ.	Bot. mag. t. 2359.	
2 maryāndica. R.S. (sc.) perennial.	N. America.	1694.	7. 8. H. ⓧ.	— t. 80.	
EUST'OMA. P.L.	EUST'OMA.	Pentandria Monogynia.			Sal. par. lond. t. 241.
sileneifolium. P.L. (pu.) Silene-leaved.	Providence Isl.	1804.	7. S. ⓧ.		
<i>Lisianthus glaucifolius</i> . Jacq. ic. 1. t. 33.					
LISIA'NTHUS. R.S.	LISIA'NTHUS.	Pentandria Monogynia. L.			
1 longifolius. R.S. (ye.) long-leaved.	Jamaica.	1793.	7. 9. S. ⓧ.	Bot. reg. t. 880.	
2 grandiflorus. R.S. (st.) large-flowered.	Trinidad.	1820.	— S. ⓧ.	Aubl. guian. 1. t. 79.	
3 exsertus. R.S. (st.) oval-leaved.	W. Indies.	1793.	— S. ⓧ.		
4 latifolius. R.S. (ye.) broad-leaved.	—	1822.	— S. ⓧ.		
5 umbellatus. R.S. (ye.) umbel-flowered.	—	—	— S. ⓧ.		
6 cordifolius. R.S. (ye.) heart-leaved.	—	1816.	6. 8. S. ⓧ.	Brown. jam. 2. t. 9. f. 2.	
7 acutangulus. R.S. (ye.) acute-angled.	Peru.	1820.	— S. ⓧ.	Flor. per. 2. t. 122. f.a.	
8 angustifolius. R.S. (gr.) narrow-leaved.	S. America.	1824.	7. 10. S. ⓧ.		
LOG'ANIA. B.P.	LOG'ANIA.	Pentandria Monogynia. R.S.			
1 latifolia. B.P. (wh.) broad-leaved.	N. Holland.	1816.	4. 6. G. ⓧ.	Lab. nov. hol. 1. t. 61.	
<i>Exācum vagināle</i> . Lab.					
2 floribunda. B.P. (wh.) many-flowered.	N. S. W.	1797.	4. 5. G. ⓧ.	Andr. deposit. t. 520.	
<i>Euōsma albiflora</i> . A.R.					
3 revoluta. B.P. (wh.) revolute-leaved.	—	1822.	4. 6. G. ⓧ.		
VILLA'RSIA. R.S.	VILLA'RSIA.	Pentandria Monogynia. R.S.			
1 nymphoides. R.S. (ye.) common.	England.	—	6. 7. H.w. 4.	Eng. bot. v. 4. t. 217.	
<i>Menyánthes nymphoides</i> . E.B.					
2 indica. R.S. (ye.) Indian.	E. Indies.	1790.	6. 8. S.w. 4.	Bot. mag. t. 658.	
3 sarmentosa. R.S. (ye.) sarmentose.	N. S. W.	1806.	— G.w. 4.	— t. 1323.	
<i>Menyánthes sarmentosa</i> . B.M.					
4 lacunosa. V. (wh.) smooth-flowered.	N. America.	1812.	6. 7. G.w. 4.	Venten. choix. 9.	
5 reniformis. B.P. (ye.) kidney-leaved.	N. S. W.	1820.	— G.w. 4.		
6 parnassiifolia. B.P. (ye.) tall.	—	1805.	6. 10. G.w. 4.	Bot. mag. t. 1029.	
<i>Menyánthes exaltata</i> . B.M. <i>Swertia parnassiifolia</i> . Lab. n. hol. t. 97.					
7 ovata. R.S. (ye.) oval-leaved.	C. B. S.	1786.	5. 7. G.w. 4.	Bot. mag. t. 1909.	
MENYA'NTHES. R.S.	BUCK-BEAN.	Pentandria Monogynia. L.			
1 trifoliata. E.B. (li.) common.	Britain.	—	6. 7. H.w. 4.	Eng. bot. v. 7. t. 495.	
2 americana. (li.) American.	N. America.	—	4. 6. H.w. 4.		
<i>trifoliata</i> β mīnor. P.S.					

ORDO CXLI.

BIGNONIACEÆ. *Brown prodr. 1. p. 470.*

CHILO'PSIS. D.D.	CHILO'PSIS.	Didynamia Angiospermia.				
lineáris.	(<i>ro.</i>) linear-leaved.	Mexico.	1825.	S. h.	Cavan. ic. 3. t. 269.
<i>Bignonia?</i> lineáris. Cav.						
CATA'LPA. J.	CATA'LPA.	Diandria Monogynia. S.S.				
1 syringifolia. s.s. (<i>wh.</i>) common.	N. America.	1726.	6. 8.	H. h.	Bot. mag. t. 1094.	
<i>Bignonia Catálpæ.</i> L.						
2 longissima. s.s. (<i>wh.pu.</i>) wave-leaved.	W. Indies.	1777.	S. h.	Plum. ic. t. 57.	
3 microphylla. s.s. (<i>bh.</i>) small-leaved.	Hispaniola.	1820.	S. h.	— t. 55.	
BIGNONIA. S.S.	TRUMPET-FLOWER.	Didynamia Angiospermia. L.				
1 U'nguis. s.s. (<i>ye.</i>) Barbadoes.	W. Indies.	1759.	S. h.	Plum. amer. t. 94.	
2 stamínea. s.s. (<i>ye.pu.</i>) long-stamened.	Hispaniola.	1820.	6. 9.	S. h.	Plum. ic. 56. f. 2.	
3 capreolata. s.s. (<i>ye.pu.</i>) four-leaved.	N. America.	1710.	6. 8.	H. h.	Bot. mag. t. 864.	
4 pubescens. s.s. (<i>ye.</i>) downy.	New Spain.	1759.	6. 7.	S. h.	—	
5 æquinoctialis. s.s. (<i>ye.</i>) equinoctial.	W. Indies.	1768.	6. 8.	S. h.	Plum. ic. t. 55. f. 1.	
6 Chamberla'nyii. B.M. (<i>ye.</i>) Chamberlayne's. S. America.	1818.	—	—	S. h.	Bot. mag. t. 2148.	
7 chrysoleuca. s.s. (<i>ye.wh.</i>) yellow & white.	—	1824.	—	S. h.	—	
8 alliacea. s.s. garlick-scented.	W. Indies.	1790.	S. h.	—	
9 spectabilis. s.s. (<i>pu.</i>) showy.	S. America.	1820.	S. h.	—	
10 picta. k.s. (<i>wh.</i>) flat-stemmed.	—	—	—	S. h.	—	
11 obliqua. k.s. (<i>re.</i>) oblique-leaved.	Caracas.	1828.	—	S. h.	—	
12 laurifolia. s.s. (<i>li.</i>) Laurel-leaved.	Guiana.	1804.	S. h.	—	
13 lactiflora. s.s. (<i>wh.</i>) milk-coloured.	Trinidad.	1822.	4. 6.	S. h.	Vahl. symb. 3. t. 66.	
14 villosa. s.s. (<i>pu.</i>) villous.	S. America.	1820.	S. h.	—	
15 ophthalmica. A. (<i>ye.</i>) Anderson's.	St. Vincent.	1812.	S. h.	—	
16 rigescens. s.s. (<i>fl.ye.</i>) rigid-leaved.	S. America.	1816.	6. 8.	S. h.	Jacq. schoenb. 2. t. 210.	
17 grandifolia. B.R. (<i>ye.</i>) large-leaved.	—	—	4. 7.	S. h.	Bot. reg. t. 418.	
18 latifolia. s.s. (<i>ye.</i>) broad-leaved.	Cayenne.	1824.	S. h.	—	
19 elongata. s.s. (<i>ye.</i>) long-racemed.	S. America.	1820.	S. h.	Vahl. ecl. 2. t. 16.	
20 mollissima. k.s. (<i>pu.</i>) soft-leaved.	Caracas.	1812.	S. h.	—	
21 diversifolia. k.s. different-leav'd.	Mexico.	1824.	S. h.	—	
22 floribunda. k.s. abundant-flower'd.	—	—	—	S. h.	—	
23 cändicans. s.s. (<i>pu.</i>) white-leaved.	Trinidad.	1822.	9. 12.	S. h.	—	
24 quadrangularis. D.L. square-stalked.	—	—	—	S. h.	—	
25 colorata. s.s. (<i>ye.</i>) coloured calyxed.	—	1824.	S. h.	—	
26 crucigera. s.s. (<i>or.</i>) cross-bearing.	S. America.	1759.	6. 8.	S. h.	Plum. ic. t. 58.	
27 Cherére. B.R. (<i>or.</i>) various-leaved.	Guiana.	1820.	6. 10.	S. h.	Bot. reg. t. 1301.	
28 heterophylla. w.						
29 venusta. s.s. (<i>or.</i>) comely.	S. America.	1816.	9. 12.	S. h.	Bot. reg. t. 249.	
29 incarnata. s.s. (<i>ro.</i>) flesh-coloured.	Guiana.	1818.	6. 9.	S. h.	Aub. gui. 3. t. 261.	
30 mollis. s.s. (<i>ye.</i>) soft.	—	—	—	S. h.	Vahl. ecl. 2. t. 10.	
30 tomentosa. Rich. non Th.						
31 littoralis. k.s. (<i>re.</i>) sea-side.	Mexico.	1824.	S. h.	—	
32 viridiflora. B.C. (<i>gr.</i>) green-flowered.	S. America.	1820.	6. 9.	S. h.	Lodd. bot. cab. t. 1026.	
33 triphylla. s.s. (<i>wh.</i>) three-leaved.	Mexico.	1733.	—	S. h.	—	
34 pallida. B.R. (<i>li.</i>) pale.	S. America.	1823.	7. 8.	S. h.	Bot. reg. t. 965.	
35 fluviatilis. s.s. (<i>wh.</i>) marsh.	—	1818.	—	S. h.	Aub. gui. 2. t. 267.	
36 Leucóxylon. s.s. (<i>pi.</i>) white-wooded.	W. Indies.	1759.	6. 7.	S. h.	Andr. reposit. t. 43.	
37 chrysántha. s.s. (<i>ye.</i>) golden-flower'd.	Caracas.	1822.	S. h.	Jacq. schoenb. 2. t. 211.	
38 lepidota. k.s. (<i>ro.</i>) silver-spotted.	Cuba.	1828.	S. h.	—	
39 esculifolia. k.s. (<i>or.</i>) <i>Æsculus</i> -leaved.	Mexico.	1824.	S. h.	—	
40 serratifolia. s.s. (<i>ye.</i>) saw-leaved.	Trinidad.	1822.	S. h.	—	
41 bijuga. s.s. Madagascar.	Madagascar.	—	S. h.	—	
42 Clématis. k.s. (<i>wh.ye.</i>) Clematis-leaved.	Caracas.	—	S. h.	—	
43 chelonoides. s.s. (<i>ye.</i>) tree.	E. Indies.	1808.	S. h.	Rheed. mal. 6. t. 26.	
44 variabilis. s.s. (<i>gr.ye.</i>) variable-leaved.	Caracas.	1820.	6. 8.	S. h.	Jacq. schoenb. 2. t. 212.	
45 indica. w. (<i>wh.std.</i>) Indian.	E. Indies.	1775.	S. h.	Rheed. mal. 1. t. 43.	
46 jasminifolia. k.s. Jasmine-leaved.	S. America.	1824.	S. h.	—	
47 crenata. L.C. crenated.	1825.	S. h.	—	
48 decipiens. L.C. doubtful.	—	S. h.	—	
49 lúcida. L.C. glossy.	S. America.	1824.	S. h.	—	
50 álba. w. (<i>wh.</i>) white-flowered.	Guiana.	1823.	S. h.	Aub. gui. 2. t. 266.	

51 <i>echinàta</i> . w. (bh.)	bristly-fruited.	Guiana.	1804.	S. h. ~.	Aub. gui. 2. t. 264.
52 <i>pùrpùrea</i> . L.C. (pu.)	purple.	S. America.	1822.	S. h.	
53 <i>tomentòsa</i> . Th. tomentose.	Japan.	1823.	S. h. ~.	Kæmpf. amœn. t. 860.
54 <i>stipulàta</i> . H.B. stipuled.	E. Indies.	1825.	S. h.	
55 <i>undulàta</i> . H.B. wave-leaved.	——	1827.	S. h.	
56 <i>suavéolens</i> . H.B. sweet-scented.	——	1825.	S. h.	
57 <i>císpà</i> . H.B. curled.	——	1822.	S. h.	
58 <i>xylócarpa</i> . H.B. woody-podded.	——	1825.	S. h.	
59 <i>aménica</i> handsome.	——	1823.	S. h.	
60 <i>spicátâ</i> . D.L. spiked-flowered.	Trinidad.	1822.	S. h. ~.	
61 <i>multipùgâ</i> . R. many-leaved.	E. Indies.	1826.	S. h.	
MILLINGTONIA. L.	MILLINGTONIA.	Didynamia Angiospermia. L.			
horténisia. L. (wh.)	cork-barked.	E. Indies.	1818.	S. h.	Roxb. corom. 3. t. 214.
<i>Bignònìa suberòsa</i> . Roxb.					
TEC'OMA. J.	TEC'OMA.	Didynamia Angiospermia. S.S.			
1 <i>spléndida</i> . s.s. (or.)	splendid.	Brazil.	1826.	S. h.	
2 <i>pentaphylla</i> . s.s. (wh.)	five-leaved.	Jamaica.	1733.	S. h.	Marc. bras. t. 118.
3 <i>digitatâ</i> . K.S. digitate-leaved.	S. America.	1818.	S. h.	
4 <i>astrális</i> . B.P. (bh.)	broad-leaved.	N. S. W.	1793.	G. h. ~.	Bot. mag. t. 865.
<i>Bignònìa Pandòræ</i> . B.M.					
5 <i>meonántha</i> . (bh.)	narrow-leaved.	——	1815.	G. h. ~.	
<i>Bignònìa meonántha</i> . L.en.					
6 <i>mollis</i> . K.S.	(ye.) soft-leaved.	Mexico.	1824.	S. h.	
7 <i>radicans</i> . s.s. (or.)	Ash-leaved.	N. America.	1640.	H. h. ~.	
<i>a majeor</i> . H.K.	(or.) great.	——	——	H. h. ~.	Bot. mag. t. 485.
<i>θ minor</i> . H.K.	(or.) smaller.	——	——	H. h. ~.	Catesb. car. 1. t. 65.
8 <i>grandiflòra</i> . (or.)	large-flowered.	China.	1800.	G. h. ~.	Bot. mag. t. 1398.
<i>Bignònìa grandiflòra</i> . B.M.	<i>Incarvilléa grandiflòra</i> . s.s.				
9 <i>stáns</i> . s.s. (ye.)	upright-ash-ld.	S. America.	1730.	S. h.	Plum. ic. t. 54.
10 <i>incisa</i> . D.L. (ye.)	cut-leaved.	Trinidad.	1824.	S. h.	
11 <i>capénsis</i> . B.R. (or.)	Cape.	C. B. S.	——	G. h. ~.	Bot. reg. t. 1117.
JACARA'NDA. S.S.	JACARA'NDA.	Didynamia Angiospermia. S.S.			
1 <i>tomentosa</i> . D.D. (pu.)	tomentose.	Brazil.	1824.	S. h.	
2 <i>mimosifòlia</i> . B.R. (bl.)	Mimosa-leaved.	——	1818.	S. h.	Bot. reg. t. 631.
<i>ovalifòlia</i> . Bot. mag. t. 2327.					
3 <i>bahaménsis</i> . B.M. (bl.)	Bahama.	Bahama Isl.	1724.	S. h.	Catesb. car. 1. t. 42.
<i>Bignònìa cerùleà</i> . w.					
4 <i>Copaia</i> . D.D. (bl.)	Box-leaved.	Guiana.	1793.	S. h.	Aub. gui. 2. t. 265.
<i>Bignònìa pròcera</i> . w.					
5 <i>rhombifòlia</i> . s.s. (pu.)	(pu.) rhomb-leaved.	Surinam.	1823.	S. h.	
<i>filicifòlia</i> . D.D.					
SPATH'ODEA. S.S.	SPATH'ODEA.	Didynamia Angiospermia. S.S.			
1 <i>uncata</i> . s.s. (ye.)	hooked.	Guiana.	1804.	G. h. ~.	Bot. mag. t. 1511.
<i>Bignònìa uncata</i> . B.M.					
2 <i>corymbòsa</i> . s.s. (ye.)	corymb-flower'd.	Trinidad.	1823.	S. h.	
3 <i>Rheèdii</i> . s.s. (wh.)	Rheede's.	E. Indies.	1794.	S. h.	Roxb. cor. 2. t. 144.
<i>Bignònìa spathacea</i> . R.C.					
4 <i>Roxburghii</i> . s.s. (ro.)	Roxburgh's.	——	1820.	S. h.	Roxb. cor. v. 2. t. 145.
<i>Bignònìa quadrilocularis</i> . Roxb.					
5 <i>fraxinifòlia</i> . K.S. (ye.)	Ash-leaved.	Caracas.	1821.	S. h. ~.	
<i>Bignònìa fraxinifòlia</i> . s.s.					
AMPHIL'OPHIUM. K.S.	AMPHIL'OPHIUM.	Didynamia Angiospermia. S.S.			
paniculàtum. K.S. (pu.)	paniced.	S. America.	1788.	S. h. ~.	Jacq. amer. t. 116.
<i>Bignònìa paniculàta</i> . Jacq.					
CALAMPÉLIS. D.D.	CALAMPÉLIS.	Didynamia Angiospermia.			
scábra. B.F.G. (or.)	rough-podded.	Chile.	1823.	H. h. ~.	Swt. br. fl. gar. s. 2. t. 30.
<i>Eccremocárpus scáber</i> . Bot. reg. t. 939.					
ECCREMOCA'RPUIS.	ECCREMOCA'RPUIS.	Didynamia Angiospermia. S.S.			
longiflorus. K.S. (ye.)	long-flowered.	Peru.	1824.	G. h. ~.	H. et B. pl. æq. 1. t. 65.
FI'ELDIA. Cun.	FI'ELDIA.	Didynamia Angiospermia.			
austrális. Cun. (gr.st.)	New Holland.	N. Holland.	1824.	G. h.	Hook. ex. flor. t. 232.

ORDO CXLII.

COBŒACEÆ. *D. Don in Edinb. phil. journ.*

COB'ŒA. Cav. **COB'ŒA.** Pentandria Monogynia.
 scándens. Cav. (*bl.*) climbing. Mexico. 1792. 5. 10. G. h. — Bot. mag. t. 851.

ORDO CXLI.

PEDALINÆ. *Brown prodr.* 519.

PED'ALIUM. S.S. **PED'ALIUM.** Didynamia Angiospermia. S.S.
 Mùrex. s.s. (*ye.*) prickly-fruited. E. Indies. 1778. 8. 9. S. ⊙. Lam. ill. t. 538.

ORDO CXLIV.

DIDYMOCARPEÆ. *D. Don.*

STREPTOCA'RPU'S. B.R. **STREPTOCA'RPU'S.** Diandria Monogynia.
 Réxi. B.R. (*bl.*) Cape. C. B. S. 1824. 2. 10. S. ¼. Bot. reg. t. 1173.
Didymocárpus Réxi. Hook. ex. flor. t. 227.

ORDO CXLV.

SESAMEÆ. *Brown.*

TOUR'ETIA. S.S. TOUR'ETIA. Didynamia Angiospermia. S.S.				
lappacea. s.s. (<i>sc.</i>) scarlet-flowered. Peru. 1788. 6. 8. S. ⊙. — Sal. stirp. rar. 5. t. 3.				
<i>Dombéya lappacea.</i> Sal.				
MARTY'NIA. J. MARTY'NIA. Didynamia Angiospermia. L.				
1 diándra. s.s. (<i>re.</i>) two-stamed. New Spain. 1731. 7. 8. S. ⊙. Andr. deposit. t. 575.				
2 proboscídea. s.s. (<i>li.</i>) horn-capsuled. America. 1738. 6. 8. G. ⊙. Bot. mag. t. 1056.				
<i>ánnuas.</i> L.				
3 lútea. B.R. (<i>ye.</i>) yellow. S. America. 1824. 7. 9. H. ⊙. Bot. reg. t. 934.				
4 longiflóra. s.s. (<i>pu.</i>) long-flowered. C. B. S. 1781. 7. 8. G. ⊙. Meerb. ic. 1. t. 7.				
CRANIO'LARIA. J. CRANIO'LARIA. Didynamia Angiospermia. L.				
<i>ánnuas.</i> k.s. (<i>wh.sp.</i>) annual. S. America. 1733. 7. 8. S. ⊙. Jacq. amer. t. 110.				
<i>Marty'nia Craniolária.</i> W.				
SESAMUM. S.S. OILY-GRAIN. Didynamia Angiospermia. L.				
1 orientále. s.s. (<i>bh.</i>) oriental. E. Indies. 1731. 7. 8. S. ⊙. Rheed. mal. 9. t. 54.				
2 indicum. s.s. (<i>bh.</i>) Indian. — — — — S. ⊙. Bot. mag. t. 1788.				
3 lútemum. s.s. (<i>ye.</i>) yellow. — — — — S. ⊙.				

ORDO CXLVI.

POLEMONIACEÆ. *D. Don in Edinb. phil. journ.*

POLEM'ONIUM. S.S. GREEK VALERIAN. Pentandria Monogynia. L.				
1 cerúleum. s.s. (<i>bl.</i>) common. Britain. 6. H. ¼. Eng. bot. t. 14.				
β álbum. (<i>wh.</i>) white-flowered. — — — — H. ¼.				
γ maculátum. s.s. (<i>sp.</i>) spot-flowered. — — — — H. ¼.				

2 grâcile. W.en. (<i>wh.</i>) slender.	Dahuria.	1818.	6. 7. H. Ȑ.	
3 mexicânum. s.s. (<i>bl.</i>) Mexican.	Mexico.	1817.	4. 6. H. Ȑ.	Bot. reg. t. 460.
4 réptans. s.s. (<i>bl.</i>) creeping.	N. America.	1758.	4. 5. H. Ȑ.	Bot. mag. t. 1887.
<i>cæruleum</i> β <i>piliferum</i> . Bot. reg. t. 1303.				
5 villôsum. G.R. (<i>bl.</i>) dwarf hairy.	Siberia.	1827.	6. 9. H. Ȑ.	Swt. br. fl. gar. t. 266.
6 Richardsoni. B.M. (<i>bl.</i>) Richardson's.	N. America.	—	— H. Ȑ.	Bot. mag. t. 2800.
7 hûmîle. B.M. (<i>bl.</i>) dwarf.	—	—	— H. Ȑ.	Bot. reg. t. 1304.
8 sibíicum. D.D. (<i>wh.</i>) bipinnate.	Siberia.	1800.	6. H. Ȑ.	Swt. br. fl. gar. t. 182.
PHLO'X. R.S. LÝCHNIDEA.	Pentandria Monogynia. L.			
1 paniculâta. R.S. (<i>pu.</i>) paniced.	N. America.	1732.	8. 9. H. Ȑ.	Dill. elth. t. 166. f. 203.
<i>β álba</i> . (<i>wh.</i>) white-flowered.	—	1813.	— H. Ȑ.	
2 undulâta. R.S. (<i>pu.</i>) wave-leaved.	—	1759.	7. 8. H. Ȑ.	
3 acuminâta. R.S. (<i>pu.</i>) cross-leaved.	—	1812.	6. 8. H. Ȑ.	Bot. mag. t. 1880.
<i>decussâta</i> . Lyon. <i>latifolia</i> . Hort. non M.				
4 virginica. L.C. (<i>pu.</i>) Virginian.	—	—	— H. Ȑ.	
5 intermédia. L.C. (<i>pu.</i>) intermediate.	—	—	7. 9. H. Ȑ.	
6 penduliflora. B.F.G. (<i>pu.</i>) drooping-flower'd.	—	1824.	— H. Ȑ.	Swt. br. fl. gar. s. 2.
7 corymbosa. B.F.G. (<i>li.</i>) corymb-flower'd.	—	—	— H. Ȑ.	
8 acutifolia. B.F.G. (<i>pu.</i>) sharp-leaved.	—	1825.	— H. Ȑ.	Swt. br. fl. gar. t. 224.
9 odorâta. B.F.G. (<i>re.</i>) sweet-scented.	—	1824.	— H. Ȑ.	— t. 248.
10 scâbra. B.F.G. (<i>li.</i>) rough-leaved.	—	1812.	— H. Ȑ.	
<i>Sickmanni</i> . Lehman.				
11 maculatâ. R.S. (<i>pu.</i>) spotted-stalked.	—	1740.	7. 8. H. Ȑ.	Jacq. vind. 2. t. 147.
12 pyramidális. R.S. (<i>li.</i>) pyramidal.	—	1800.	6. 8. H. Ȑ.	Swt. br. fl. gar. t. 233.
13 latifolia. R.S. (<i>pu.</i>) broad-leaved.	—	1812.	— H. Ȑ.	
14 carolinâ. R.S. (<i>re.pu.</i>) rough-stemmed.	—	1728.	7. 9. H. Ȑ.	Bot. mag. t. 1344.
15 cordâta. E.C. (<i>li.</i>) heart-leaved.	Carolina.	1826.	6. 9. H. Ȑ.	Swt. br. fl. gar. s. 2.t.13.
16 excélsa. H.E. (<i>pu.</i>) tall upright.	N. America.	1824.	— H. Ȑ.	
17 Wheeleriana. B.F.G. (<i>pi.</i>) Wheeler's.	Hybrid.	—	— H. Ȑ.	
18 Shephérddii. H.E. (<i>pu.</i>) Shepherd's.	—	—	— H. Ȑ.	
19 glaberrima. R.S. (<i>pu.</i>) smooth-stalked.	N. America.	1725.	6. 8. H. Ȑ.	Swt. br. fl. gar. s. 2 t. 36.
20 triflora. R.S. (<i>bh.</i>) flesh-coloured.	—	1816.	7. 9. H. Ȑ.	Swt. brit. fl. gar. t. 29.
<i>cárnea</i> . Bot. mag. t. 2155.				
21 suavélens. R.S. (<i>wh.</i>) white-flowered.	—	1766.	7. 8. H. Ȑ.	
22 longiflora. B.F.G. (<i>wh.</i>) long-flowered.	—	1827.	7. 10. H. Ȑ.	Swt. br. fl. gar. s. 2. t. 31.
23 reflexâ. B.F.G. (<i>d.pu.</i>) reflexed-leaved.	—	1824.	6. 9. H. Ȑ.	Swt. br. fl. gar. t. 232.
24 suffruticosa. B.R. (<i>d.pu.</i>) glossy-leaved.	—	1790.	7. 9. H. Ȑ.	Bot. reg. t. 68.
25 Listoniâna. H.E. (<i>pu.</i>) Liston's.	—	1816.	5. 7. H. Ȑ.	
26 ovâta. w. (<i>pu.</i>) ovate-leaved.	—	1759.	— H. Ȑ.	Bot. mag. t. 528.
27 stolonifera. B.M. (<i>bl.</i>) creeping.	—	1800.	6. 9. H. Ȑ.	— t. 563.
<i>prostrâta</i> . H.K. <i>réptans</i> . M. <i>stolonifera</i> has the right of priority.				
28 pilosa. B.M. (<i>p.bl.</i>) hairy-leaved.	N. America.	1759.	5. 6. H. Ȑ.	Bot. mag. t. 1307.
<i>aristâta</i> . M.				
29 amœna. B.M. (<i>ro.</i>) Fraser's hairy.	—	1809.	6. 7. H. Ȑ.	Bot. mag. t. 1308.
<i>pilosâ</i> . β . R.S.				
30 canadén sis. B.F.G. (<i>pa.bl.</i>) Canadian.	Canada.	1826.	5. 7. H. Ȑ.	Swt. br. fl. gar. t. 221.
31 divaricâta. R.S. (<i>pa.bl.</i>) early-flowering.	N. America.	1746.	4. 6. H. Ȑ.	Bot. mag. t. 163.
32 procumbens. B.F.G. (<i>bl.</i>) procumbent.	—	1829.	5. 6. H. Ȑ.	Swt. br. fl. gar. s. 2. t. 7.
33 subulâta. R.S. (<i>li.pu.</i>) awl-leaved.	—	1786.	4. 6. H. $\ddot{\circ}$.	Bot. mag. t. 411.
34 nivâlis. B.C. (<i>wh.</i>) snowy white.	—	1820.	— H. $\ddot{\circ}$.	Swt. br. fl. gar. t. 185.
35 setâcea. R.S. (<i>pi.</i>) bristle-leaved.	—	1786.	4. 5. H. $\ddot{\circ}$.	Bot. mag. t. 415.
COLL'OMIA. N. COLL'OMIA. Pentandria Monogynia.				
1 grandiflora. D. (<i>bf.</i>) large-flowered.	N. America.	1827.	6. 7. H. \odot .	Bot. reg. t. 1174.
2 lineâris. D. (<i>fl.</i>) linear-leaved.	—	—	— H. \odot .	— t. 1166.
3 grâcilis. D. (<i>ro.</i>) slender.	—	—	5. 7. H. \odot .	Bot. mag. t. 2924.
<i>Gilia gracilis</i> . B.M.				
4 heterophylla. B.M. (<i>ro.</i>) various-leaved.	—	—	5. 8. H. \odot .	Bot. mag. t. 2895.
GI'LIA. R.P. GI'LIA. Pentandria Monogynia. S.S.				
1 inconspicua. B.M. (<i>bl.</i>) small blue.	America.	1793.	9. 11. H. \odot .	
<i>Ipomopsis inconspecta</i> . Sim. exot. bot. 1. t. 14.				
2 capitâta. D. (<i>bl.</i>) cluster-flowered.	—	1826.	5. 9. H. \odot .	
IPOMO'PSIS. M. IPO'MO'PSIS. Pentandria Monogynia.				
1 elegans. Ex.B. (<i>sc.</i>) elegant.	Carolina.	1726.	8. 9. G. δ .	
<i>Cantua coronopifolia</i> . Andr. rep. t.				
2 pulchella. D. (<i>sc.</i>) pretty.	N. America.	1827.	— G. δ .	
<i>elegans</i> . Bot. reg. non M.				
HOITZIA. J. HOITZIA. Pentandria Monogynia. S.S.				
1 coccinea. K.S. (<i>sc.</i>) scarlet.	Mexico.	1826.	5. 7. G. δ .	Cavan. ic. 4. t. 365.
2 cœrula. K.S. (<i>bl.</i>) blue-flowered.	—	1827.	— G. δ .	— 4. t. 366.

CALD'ASIA. S.S.

CALD'ASIA. Pentandria Monogynia.

heterophylla. s.s. (*bl.*) various-leaved. New Spain. 1813. 5. 12. S. (c). Bot. reg. t. 92.*Bonplandia geminiflora*. Cav. ic. 6. t. 532.

ORDO CXLVII.

HYDROLEACEÆ. *Kth. synops.* 2. p. 234.

HYDR'OLEA. S.S.	HYDR'OLEA.	Pentandria Digynia. L.			
1 spinosa. s.s.	(<i>bl.</i>) thorny.	S. America. 1791.	7. 8.	S. h.	Bot. reg. t. 566.
2 quadrivalvis. s.s.	(<i>bl.</i>) four-valved.	Carolina. 1824.	—	G. 4.	
WIGA'NDIA. K.S.	WIGA'NDIA.	Pentandria Digynia.			
1 urens. K.S.	{ <i>vi.</i> stinging.	Mexico. 1827.	4. 6.	G. 4.	Flor. peruv. 3. t. 243.
2 caracasana. K.S.	{ <i>vi.</i> Caracas.	Caracas. 1825.	—	S. h.	
N'AMA. S.S.	N'AMA.	Pentandria Digynia. L.			
1 jamaicensis. s.s. (<i>wh. bl.</i>) Jamaica.	Jamaica.	1812.	6. 9.	S. (c).	Lam. ill. t. 184.
2 undulata. K.S.	{ <i>vi.</i> wave-leaved.	Mexico. 1826.	—	G. 4.	

ORDO CXLVIII.

CONVOLVULACEÆ. *Brown prodr.* 1. p. 481.

? RETZIA. R.S.	RE'TZIA.	Pentandria Monogynia. R.S.			
spicata. R.S.	{ <i>br.</i> spik'd.	C. B. S.	5. 6.	G. h.	Lam. ill. t. 103.
CONVO'LVLUS. B.P.	BINDWEED.	Pentandria Monogynia. L.			
1 arvensis. R.S. (<i>pi. wh.</i>) small.	Britain.	6. 9.	H. 4.	Eng. bot. v. 5. t. 312.
2 chinensis. B.R. (<i>pi.</i>) Chinese.	China.	1816.	—	H. 4.	Bot. reg. t. 322.
3 Malcolmi. F.I. (<i>wh.</i>) Malcolm's.	Persia.	1824.	7. 10.	H. 4.	
4 emarginatus. s.s. (<i>re.</i>) emarginate.	—	6. 9.	H. 4.	
5 bicuspidiatus. s.s. (<i>bh.</i>) two-pointed.	Davuria.	—	—	H. 4.	
6 Scammonia. R.S. (<i>st. bh.</i>) Scammony.	Levant.	1596.	7. 8.	H. 4.	Flor. græc. t. 192.
7 erubescens. R.S. (<i>bh.</i>) maiden blush.	N. S. W.	1803.	7. 9.	G. 4.	Bot. mag. t. 1067.
8 quinqueflorus. R.S. (<i>bh.</i>) five-flowered.	Mauritius.	1826.	—	S. h.	
9 pannifolius. R.S. (<i>bl.</i>) cloth-leaved.	Canaries.	1805.	6. 9.	G. h.	Bot. reg. t. 222.
10 canariensis. R.S. (<i>pi.</i>) Canary.	1690.	—	G. h.	Bot. mag. t. 1228.
11 Massoni. R.S. (<i>wh.</i>) Masson's.	1788.	—	G. h.	Bot. reg. t. 133.
Dryändri. s.s. <i>suffruticosus</i> . B.R.					
12 canticans. R.S. hoary.	E. Indies.	1818.	S. h.	
13 bonariensis. R.S. (<i>std.</i>) stripe-flowered.	Buenos Ayres.	1827.	6. 8.	G. 4.	Cavan. icon. 5. t. 48. f. 2.
14 tilaceus. R.S. (<i>wh.</i>) Lime-tree-leaved.	Brazil.	1820.	—	S. 4.	
15 guianensis. R.S. (<i>bh.</i>) Guiana.	Guiana.	—	—	S. 4.	Aub. gui. 1. t. 52.
16 ciliatus. R.S. (<i>bh.</i>) fringed-leaved.	Cayenne.	—	—	S. h.	
17 pentanthus. R.S. (<i>bl.</i>) five-flowered.	W. Indies.	1808.	4. 10.	S. h.	Bot. reg. t. 439.
18 farmosus. R.S. (<i>bh.</i>) mealy-stalked.	Madeira.	1777.	5. 6.	G. 4.	Sal. par. lond. t. 45.
19 Hermánniae. s.s. (<i>wh.</i>) Peruvian.	Peru.	1799.	8. 9.	G. 4.	L'Herit. stirp. t. 33.
20 teuniissimus. R.S. (<i>li.</i>) slender-leaved.	Levant.	—	6. 10.	G. 4.	Flor. græc. t. 195.
21 althaeoides. R.S. (<i>li.</i>) Althaea-leaved.	—	1597.	6. 9.	F. 4.	t. 194.
22 bryoniaefolius. R.S. (<i>pu.</i>) Bryony-leaved.	China.	1802.	7. 8.	G. 4.	Bot. mag. t. 943.
23 italicus. R.S. (<i>pi.</i>) Italian.	Italy.	—	—	F. 4.	Ten. fl. neap. 1. t. 15.
hirsutus. T.N. non M.B.					
24 hirsutus. M.B. (<i>pu. ye.</i>) hairy.	Tauria.	1824.	—	H. 4.	
25 alceifolius. R.S. (<i>st. pu.</i>) Hollyhock-leav'd.	C. B. S.	1823.	—	G. 4.	
26 macrocarpus. R.S. (<i>pu.</i>) long-fruited.	S. America.	1752.	—	S. (c).	Burm. amer. t. 91. f. 1.
27 quinquefolius. R.S. (<i>w. pu.</i>) five-leaved.	Vera Cruz.	—	—	S. (c).	Pluk. alm. t. 167. f. 6.
28 glaber. R.S. (<i>wh.</i>) smooth.	Cayenne.	1806.	—	S. h.	Aub. gui. 1. t. 53.
29 tenellus. R.S. (<i>wh.</i>) slender.	N. America.	1812.	6. 8.	H. (c).	Pluk. alm. t. 166. f. 4.
30 aquaticus. N. (<i>wh.</i>) water-side.	—	—	—	H. (c).	
31 siccus. R.S. (<i>bl.</i>) small-flowered.	S. Europe.	1640.	—	H. (c).	Bot. reg. t. 445.
32 pseudosiccus. R.S. (<i>wh.</i>) long-pedunc'l'd.	Canaries.	1815.	7. 8.	H. (c).	t. 498.
elongatus. W.enum.					
33 Imperati. R.S. (<i>st.</i>) trailing.	Naples.	1824.	6. 9.	H. 4.	Cyr. pl. rar. fasc. 1. t. 5.

34 sagittifolius. F.G. (bh.)	Sibthorp's.	Samos.	1823.	7. 8. H. 4.	Flor. græc. t. 193.
35 hirtus. R.S.	(bl.) hairy-stalked.	E. Indies.	1806.	6. 8. S. ⊙.	Cavan. ic. 5. t. 482.
36 platycáros. R.S. (bh.)	flat-podded.	Mexico.	1827.	8. 10. F. 4.	
37 multífidus. R.S. (bh.)	multifid.	C. B. S.	1822.	6. 8. G. 4.	Gmel. iter. 3. t. 7.
38 pérsicus. M.B. (wh.)	Persian.	Persia.	1829.	— H. 4.	Bot. mag. t. 27.
39 tricolor. R.S. (bl.ye.)	three-coloured.	S. Europe.	1629.	7. 9. H. ⊙.	
β albiflórus.	(wh.) white-flowered.	—	—	— H. ⊙.	
40 meonanthus. R.S. (bl.ye.)	pale blue.	Portugal.	1827.	— H. ⊙.	Flor. græc. t. 197.
41 pentapetaloides. R.S. (bl.)	Majorca.	Majorca.	1789.	6. 7. H. ⊙.	— t. 199.
42 lineátus. R.S. (bh.)	dwarf.	S. Europe.	1714.	— H. 4.	Barrel. ic. t. 311.
43 Gerárdi. R.S. (bh.)	Gerard's.	—	—	— H. 4.	Cavan. ic. 3. t. 277. f. 1.
44 undulátus. R.S. (wh.re.)	waved.	—	1816.	6. 10. H. ⊙.	
humilis. Jacq. col. 4. t. 22.	f. 3. evolvuloïdes. F.G. t. 198.	ciliátus. P.S.			Flor. græc. t. 202.
45 lanátus. R.S. (wh.)	woolly-leaved.	Levant.	1829.	5. 6. G. ™.	
46 intermédia. R.S. (bh.)	intermediate.	S. Europe.	1825.	6. 7. H. 4.	Bot. mag. t. 459.
47 Cneórum. R.S. (bh.)	silvery-leaved.	Levant.	1640.	5. 9. G. ™.	Bocc. mus. 138. t. 96.
48 saxatilis. R.S. (bh.)	rock.	S. Europe.	1796.	6. 8. F. 4.	Jacq. ic. 1. t. 34.
49 floridus. R.S. (bh.)	many-flowered.	Teneriffe.	1779.	8. 9. G. ™.	Jacq. aust. 3. t. 296.
50 Cantábrica. R.S. (fl.)	flax-leaved.	S. Europe.	1680.	5. 9. H. 4.	
51 salvifolius. L.en. (bh.)	Sage-leaved.	Palestine.	1825.	— H. 4.	Bot. mag. t. 289.
52 lineáris. R.S. (wh.re.)	narrow-leaved.	S. Europe.	1770.	— G. ™.	Desf. atl. 1. t. 48.
53 suffricticósus. R.S. (bh.)	suffrutescent.	Levant.	1827.	6. 8. F. ™.	Flor. græc. t. 201.
54 Dorycnium. R.S. (ro.)	silky-leaved.	—	1806.	6. 7. H. ™.	Venten. choix. t. 24.
55 holoseríceus. R.S. (st.)	silky.	Tauria.	1824.	— H. 4.	
56 scopárius. R.S. (wh.)	broom.	Canaries.	1733.	8. 9. G. ™.	
57 terréstiris. F. (wh.)	small.	Altay.	1828.	7. 9. H. 4.	
CALYSTEGIA. B.P. BEARBIND.					
1 sèpium. R.S. (wh.)	great hedge.	Pentandria Monogynia. R.S.	6. 9. H. 4. ⊙.	Eng. bot. v. 5. t. 313.
2 sylvétris. R.S. (wh.)	wood.	Britain.	— H. 4. ⊙.	W. et K. hung. t. 261.
<i>Convolvulus sylvestris</i> . W.en.					
3 infláta. Df. (re.)	inflated.	N. America.	— H. 4. ⊙.	Bot. mag. t. 732.
sèpium. Ph. <i>Convolvulus sèpium</i> β incarnàta. B.M.					
4 dahurica. B.M. (pi.)	Dahurian.	Dahuria.	1820.	7. 8. H. 4. ⊙.	Bot. mag. t. 2609.
5 Catesbiána. R.S. (ro.)	Catesby's.	Carolina.	1816.	7. 9. H. 4. ⊙.	
6 marginàta. B.P. (wh.)	margined.	N. S. W.	1824.	7. 8. F. 4. ⊙.	
7 stáns. M. (wh.)	woolly.	N. America.	1818.	6. 7. H. 4.	
<i>tomentòsa</i> . Ph. <i>Convolvulus stáns</i> . M.					
8 spithamæa. R.S. (wh.)	small upright.	—	1696.	7. 8. H. 4.	Hook. ex. flor. t. 97.
9 Soldanélla. R.S. (pi.)	sea-side.	Britain.	6. 7. H. 4.	Eng. bot. v. 5. t. 34.
<i>Convolvulus Soldanella</i> . E.B.					
10 reniformis. B.P. (pi.)	kidney-leaved.	N. S. W.	1822.	6. 8. F. 4.	Swt.br.fl.gar.v.2.t.181.
11 hederaéca. F.I. (ro.)	Ivy-leaved.	Nepaul.	1826.	— F. 4. ⊙.	
12 Keriana. (bh.)	Mr. Ker's.	Mauritius.	1818.	— S. ⊙.	Bot. reg. t. 318.
<i>Convolvulus involucratus</i> . B.R. non Beau. <i>Convolvulus bicolor</i> . B.M. 2205. nec aliorum.					
IPOMŒA. B.P. IPOM'œA. Pentandria Monogynia. R.S.					
1 Quamoclit. R.S. (sc.)	wing-leaved.	E. Indies.	1629.	7. 9. S. ⊙. ⊙.	Bot. mag. t. 244.
β albiflóra.	(wh.) white-flowered.	—	—	— S. ⊙. ⊙.	
2 dissécta. R.S. (wh.)	cut-leaved.	Tropic's.	1813.	6. 9. S. 4. ⊙.	Willd. phyt. 1. t. 2. f. 3.
3 tuberòsa. R.S. (st.pu.)	tuberous-rooted.	W. Indies.	1731.	8. 10. S. 4. ⊙.	
4 digitáta. R.S. (sc.)	fingered.	S. America.	— S. 4. ⊙.	Plum. ic. 92. f.
5 sinuàta. R.S. (wh.)	sinuated.	N. America.	1813.	6. 9. G. 4. ⊙.	Jacq. vind. t. 1. 9.
<i>dissécta</i> . Ph. non. w. <i>Convolvulus disséctus</i> . M.					
6 tuberculáta. R.S. (li.)	tubercled.	Bourbon.	1818.	— S. ™. ⊙.	Jacq. schœn. 2. t. 199.
<i>stipulacea</i> . Jacq.					
7 caírica.	(pu.) jagged-leaved.	Egypt.	1680.	— S. ™. ⊙.	Bot. mag. t. 699.
<i>Convolvulus caíricus</i> . B.M. <i>Ipomœa palmata</i> . R.S.					
8 pendula. B.P. (pu.)	pendulous-flower'd.	N. S. W.	1808.	5. 10. G. ™. ⊙.	Bot. reg. t. 632.
9 dasyspérmäa. J.E. (ye.pu.)	warted.	E. Indies.	1815.	8. 9. G. ⊙. ⊙.	— t. 86.
<i>tuberulata</i> . B.R. non R.S. <i>Convolvulus pedatus</i> . F.I.					
10 cóptica. R.S. (wh.)	palmate-leaved.	E. Indies.	1824.	8. 10. S. ⊙. ⊙.	
11 pés-tigris. R.S. (bh.)	palmated.	—	1732.	— S. ⊙. ⊙.	Dill. elth. t. 318. f. 411.
12 platénensis. B.R. (pu.)	Plata.	S. America.	1817.	6. 9. S. 4. ⊙.	Bot. reg. t. 33.
13 setosa. B.R. (pu.)	bristly-stalked.	Brazil.	—	8. 10. S. ™. ⊙.	— t. 335.
14 paniculáta. R.S. (pu.)	panicked.	E. Indies.	1799.	6. 9. S. ™. ⊙.	— t. 62.
15 insignis. A.R. (pu.)	large purple.	—	1814.	— S. ™. ⊙.	— t. 75.
<i>gossypifolia</i> . R.S.					
16 senegalénsis. R.S. (pu.)	Senegal.	Senegal.	1822.	6. 10. S. ™. ⊙.	
17 spléndens. G.D. (pu.)	splendid.	Sierra Leon.	—	6. 9. S. ™. ⊙.	
18 parviflora. R.S. (pu.)	small-flowered.	Santa Cruz.	—	7. 10. S. ⊙. ⊙.	Sloan, hist. 1. t. 97. f. 1.

- 19 heterophylla. R.S. (*bl.*) various-leaved. S. America. 1817. 7. 10. S. 4. — Jacq. frag. t. 42. f. 4.
 20 macrorhizos. R.S. (*sc.*) large-rooted. W. Indies. — S. 4. — Plum. cat. mss.v.2.t.58.
 21 umbellata. R.S. (*sc.*) umbelled. S. America. 1739. 6. 7. S. 4. — Burm. amer. t. 92. f. 2.
 22 carolina. R.S. (*pu.*) Carolina. Carolina. 1732. 7. 8. G. 4. — Dill. elth. t. 84. f. 98.
 23 pulchella. R.S. (*vi.*) neat. E. Indies. 1824. 5. 8. S. 4. —
 24 venosa. R.S. (*pu.*) veined-leaved. Mauritius. 1820. 7. 8. S. 4. —
 25 pentaphylla. R.S. (*wh.*) five-leaved. W. Indies. 1739. 8. 9. S. 4. — Jacq. ic. 2. t. 319.
 26 Cavanillesii. R.S. (*bh.*) Cavanille's. S. America. 1815. — S. 4. — Cavan. ic. 3. t. 256.
pentaphylla. Cav. non Jacq.
 27 armata. R.S. (*re.vi.*) armed-calyxed. Mexico. 1824. 7. 10. S. 4. — Cavan. ic. 5. t. 478. f. 2.
muricata. Cav. non Jacq.
 28 ternata. R.S. (*wh.*) ternate-leaved. S. America. — 6. 8. S. 4. — Jacq. schoenb. 1. t. 37.
 29 coccinea. R.S. (*sc.*) bright-scarlet. W. Indies. 1713. 6. 9. S. 4. — Andr. deposit. t. 99.
 30 lutorea. R.S. (*or.*) common scarlet. Guatimala. 1759. — H. 4. — Bot. mag. t. 221.
coccinea. B.M. nec *aliorum*.
 31 angularis. R.S. (*cr.*) angular-leaved. E. Indies. 1806. — S. 4. —
phaenicea. F.I.
 32 hastigera. K.S. (*fl.*) halbert-bearing. Mexico. 1824. 6. 8. S. 4. —
 33 longiflora. K.S. (*wh.*) long-flowered. Cuba. 1803. — S. 4. —
 34 sagittata. R.S. (*ro.*) sagittate. S. Europe. 1826. 7. 9. H. 4. — Cavan. icon. 2. t. 107.
 35 bona nox. R.S. (*wh.*) large white. America. 1773. 6. 9. G. 4. — Bot. mag. t. 752.
 36 lacunosa. R.S. (*wh.pu.*) starry. N. America. 1640. 7. 8. H. 4. — Dill. elth. t. 87. f. 102.
 37 Batatas. R.S. (*wh.pu.*) sweet potatoe. India. 1597. — S. 4. — Rheed. mal. 7. t. 50.
Convolvulus Batatas. W.
 38 Turpethum. R.S. (*wh.*) square-stalked. Ceylon. 1752. 3. 6. S. 4. — Bot. reg. t. 279.
 39 chrysoides. B.R. (*ye.*) golden. China. 1817. 6. 10. S. 4. — — t. 270.
 40 ochracea. B.R. (*ye.*) pale yellow. Benin. 1826. 8. 10. S. 4. — — t. 1060.
 41 Jálapa. (*ro.*) true Jalap. S. America. 1733. 8. 11. S. 4. — Bot. mag. t. 1572.
Jálapa. ß rðsea. B.R. 621. *Convolvulus Jálapa*. B.M.
 42 Michauxii. (*wh.bh.*) Michaux's. Georgia. 1815. — G. 4. — Bot. reg. t. 342.
Jálapa. a. B.R. *macrorhiza*. M. non R.S.
 43 sanguinea. R.S. (*sc.*) blood-flowered. W. Indies. 1812. 2. 11. S. 4. — Bot. reg. t. 9.
 44 repanda. R.S. (*sc.*) scalloped. — 1793. 5. 10. S. 4. — Sal. par. lond. t. 81.
 45 varia. R.S. (*bl.*) variable-leaved. S. America. 1816. — S. 4. —
Convolvulus pubescens. S.S.
 46 scabra. R.S. (*wh.*) rough. Egypt. 1804. — H. 4. —
 47 scabrida. R.S. (*wh.*) roughish. S. America. 1824. — H. 4. —
 48 hederacea. B.R. (*bl.*) deep-lobed. N. America. 1729. 8. 10. H. 4. — Bot. reg. t. 85.
barbata. Roth.
 49 cuspidata. R.S. (*pu.*) sharp-pointed. Peru. 1732. 6. 7. H. 4. — Flor. per. 2. t. 119. f.a.
 50 Dilleniæ. R.S. (*bl.*) Dilleniæ's. Guinea. — S. 4. — Dill. elth. t. 81. f. 93.
 51 Nil. P.S. (*bl.*) blue. S. America. 1597. 7. 9. S. 4. — Bot. mag. t. 188.
 52 cœrulea. B.R. (*bl.*) East Indian. E. Indies. 1815. — S. 4. — Bot. reg. t. 276.
 53 cœrulescens. F.I. (*bl.*) light blue. — 1820. 6. 8. S. 4. —
 54 pudibunda. B.R. (*ro.*) modest. W. Indies. 1825. 7. 10. S. 4. — Bot. reg. t. 999.
 55 triloba. R.S. (*vi.*) three-lobed. — 1752. 6. 7. S. 4. —
 56 hederifolia. R.S. (*vi.*) Ivy-leaved. S. America. 1773. 7. H. 4. — Plum. ic. t. 93. f. 2.
 57 hepaticifolia. R.S. (*bl.*) Hepatica-leav'd. E. Indies. 1759. 8. 9. S. 4. — Burm. ind. t. 20. f. 2.
 58 pi actata. R.S. (*vi.*) spotted-calyx'd. S. America. — 7. 11. S. 4. — Dill. elth. t. 83. f. 96.
 59 mutabilis. B.R. (*bl.*) changeable. — 1812. 5. 9. S. 4. — Bot. reg. t. 39.
 60 ae inata. R.S. (*pu.*) taper-pointed. — ... S. 4. —
 61 c mutata. R.S. (*li.*) hairy-capsuled. N. America. carolinæ. Ph. *Convolvulus carolinæ*. M. non L. — H. 4. — Dill. elth. t. 84. f. 98.
 62 gemella. R.S. (*wh.*) twin-flowered. E. Indies. 1827. 7. 9. S. 4. — Burm. ind. t. 21. f. 1.
 63 gangética. (*pi.*) Ganges. — 1812. 5. 9. S. 4. —
Convolvulus gangeticus. F.I.
 64 semidigyna. F.I. (*wh.*) cleft-styled. — 1826. — S. 4. —
 65 muricata. R.S. (*pu.*) rough. — 1777. 7. 8. S. 4. — Jacq. schoen. 3. t. 323.
bõng nox β *purpurascens*. B.R. 290. *Convolvulus muricatus*. W. — 1827. 7. 9. S. 4. —
 66 pseudomuricata. L.en. (*pu.*) rough-stem'd. — 1799. 8. 9. S. 4. — Jacq. amer. 26. t. 18.
 67 carnea. R.S. (*fl.*) flesh-coloured. S. America. 1822. 8. 10. S. 4. — Beauv. fl. d'ow. 2. t. 89.
 68 involucrata. R.S. (*pu.*) involucred. Sierra Leon. 1824. 7. 10. H. 4. — Swt. br. fl. gar. t. 86.
 69 barbigena. B.F.G. (*bl.*) bearded-calyx'd. America. 1629. 6. 9. H. 4. — Bot. mag. t. 113.
 70 purpurea. R.S. (*pu.*) purple. — — H. 4. —
β incarnata. (*fl.*) flesh-coloured. — — H. 4. —
γ leucantha. (*wh.*) white-flowered. — — H. 4. —
δ varia. B.M. (*std.*) striped-flowered. — — H. 4. — Bot. mag. t. 1682.
ε elatior. B.M. (*wh.bl.*) spotted-flowered. — — H. 4. — — t. 1005.
discolor. Jacq.

CONVOLVULACEÆ.

71 violacea. R.S. (vi.) purple-flower'd. S. America.	1732. 8. 9. S. ⊙. —. Plum. ic. t. 93. f. 1.
72 corymbosa. R.S. (wh.) corymbose. India.	1823. — S. h. — t. 89. f. 2.
73 campanulata. R.S. (pu.w.) bell-flowered. E. Indies.	1800. — S. ⊙. Rheed. mal. 2. t. 56.
74 polyanthes. R.S. (ye.gr.) umbel-flowered. W. Indies. <i>Convolvulus umbellatus. w.</i>	1739. 6. 7. S. ♀. —. Plum. amer. 88. t. 102.
75 multiflora. R.S. (pu.) many-flowered. Jamaica. — S. ♀. —. Pluk. phyt. t. 167. f. 1.
76 malabatica. R.S. (pu.st.) Malabar. E. Indies.	1823. S. h. —. Rheed. mal. 2. t. 51.
77 solanifolia. R.S. (ro.) Nightshade-ld. America.	1759. 7. 8. S. ♀. —. Plum. ic. 94. f. 1.
78 leucantha. R.S. (wh.) white-flowered. S. America.	1823. 8. 10. H. ⊙. —. Jacq. ic. 2. t. 318.
79 obscura. R.S. (wh.pu.) hairy. E. Indies.	1732. 6. 8. S. ⊙. —. Bot. reg. t. 239.
80 fastigiata. (pu.) crowded.	1816. — S. ♀. —.
<i>Convolvulus fastigiatus. F.I.</i>	
81 maxima. (pu.re) great Ceylon. Ceylon.	1799. 7. 8. S. h. —. Rheed. mal. 2. t. 53.
82 hirsutula. J.E. (wh.) hairy-stemmed. S. America.	1828. — S. ⊙. —. Jacq. f. eclog. t. 44.
83 blanda. (wh.) fair-flowered.	1820. S. h. —.
<i>Convolvulus blandus. F.I.</i>	
84 laurifolia. (ro.) Laurel-leaved.	1822. 5. 8. S. h. —.
<i>Convolvulus laurifolius. F.I.</i>	
85 sphærocéphala. (ro.) round-headed.	1816. 8. 10. S. ⊙. —.
<i>Convolvulus sphærocéphalus. F.I.</i>	
86 sibirica. R.S. (wh.) Siberian. Siberia.	1779. 7. 8. H. ⊙. —. Pall. it. 3. p. 723. t. K.
87 Hardwickii. s.s. (wh.) fringe-calyxed. E. Indies.	1816. 5. 8. S. ♀. —.
<i>Convolvulus calycinus. F.I. non Kth.</i>	
88 tamnifolia. R.S. (bl.) Tamus-leaved. Carolina.	1732. 7. H. ⊙. —. Dill. elth. t. 318. f. 410.
89 capitata. R.S. (bl.) headed. E. Indies.	1823. 7. 10. S. ♀. —.
90 triquètra. R.S. (pu.) triquetrous. Santa Cruz. — S. h. —.
91 pilosa. (pi.) various-ld. hairy. E. Indies.	1815. — S. ⊙. —.
<i>Convolvulus pilosus. F.I.</i>	
92 denticulata. B.P. (ye.) denticulate.	1778. 7. 8. S. ⊙. —. Bot. reg. t. 317.
<i>Convolvulus Médium. w. non L.</i>	
93 bicolor. (ye.pu.) two-coloured.	1815. — S. ♀. —.
<i>Convolvulus bicolor. F.I.</i>	
94 atropurpurea. F.I. (d.pu.) dark purple.	Nepaul. 1824. 7. 9. H. ⊙. —.
95 dentata. F.I. (ye.) toothed-lobed.	E. Indies. 1820. — S. ♀. —.
96 digitifolia. (pu.) finger-leaved.	1815. 6. 8. S. h. —.
<i>Convolvulus digitatus. F.I.</i>	
97 polytricha. (wh.) many-haired.	1826. 6. 10. S. ⊙. —.
<i>Convolvulus hirsutus. F.I.</i>	
98 heptaphylla. F.I. (pi.) seven-leaved.	1827. S. h. —.
99 bogotensis. K.S. (pu.) Bogota.	S. America. 1820. 5. 8. S. h. —.
100 sidæfolia. K.S. (wh.) Sida-leaved.	1812. — S. h. —.
101 albivænia. (wh.) white-veined.	Algoa Bay. 1825. 9. 10. S. h. —. Bot. reg. t. 1116.
<i>Convolvulus albivénus. B.R.</i>	
102 scrobiculata. B.R. (wh.pu.) furrowed.	Trinidad. 1822. 3. 6. S. h. —. Bot. reg. t. 1076.
103 grandiflora. R.S. (wh.) great-flowered.	E. Indies. 1802. 9. S. h. —. Andr. reposit. t. 188.
104 noctiluca. B.R. (wh.) night-flowered.	1811. 9. 10. S. h. —. Bot. reg. t. 889.
105 latiflora. R.S. (wh.) broad-flowered.	W. Indies. — S. h. —. Jacq. vind. 3. t. 69.
106 filiformis. R.S. (pu.) filiform.	1823. — S. h. —. Jacq. amer. pict. t. 26.
107 salicifolia. F.I. (wh.) Willow-leaved.	E. Indies. 1816. 5. 7. S. ♀. —.
108 Roxburghii. (ro.) Roxburgh's.	1817. — S. h. —.
<i>multiflora. F.I. nec aliorum.</i>	
109 sepiaria. F.I. (ro.) hedge. — S. h. —. Rheed. mal. 11. t. 53.
110 pandurata. B.R. (w.re.) Virginian.	N. America. 1732. 6. 9. G. ♀. —. Bot. reg. t. 588.
<i>Convolvulus pandurus. B.M. 1939.</i>	
111 canticans. (wh.) hoary. 6. 8. G. ♀. —. Bot. mag. t. 1603.
<i>Convolvulus canticans. B.M.</i>	
112 glaucifolia. R.S. (fl.) glaucous-leaved. Mexico.	1732. 5. 7. G. ♀. —. Dill. elth. t. 87. f. 101.
113 reptans. R.S. (li.) creeping. E. Indies.	1806. — S. ♀. —. Rumph.am. 5.t.155.f.1.
114 angustifolia. R.S. (ye.) narrow-leaved.	1800. 7. 8. S. ⊙. —. Jacq. ic. rar. t. 317.
115 tridentata. R.S. (pu.st.) trifid.	1778. — S. ⊙. —. Rheed. mal. 2. t. 65.
116 stans. R.S. (vi.) Mexican.	Mexico. 1824. — S. ♀. —. Cavan. icon. 3. t. 250.
117 bignonioides. B.M. (d.pu.) trumpet-flower'd. Cayenne.	1823. — S. ♀. —. Bot. mag. t. 2645.
118 sagittifolia. B.R. (ro.) arrow-leaved.	Carolina. 1819. 6. 9. G. ♀. —. Bot. reg. t. 437.
<i>Convolvulus sagittifolius. R.S.</i>	
119 vitifolia. (ye.) Vine-leaved.	E. Indies. 1820. — S. h. —. Burm. ind. t. 18. f. 1.
<i>Convolvulus vitifolius. R.S.</i>	
120 stipulacea. (wh.) stipuled.	1805. 8. 10. S. ♀. —.
<i>Convolvulus stipulaceus. F.I.</i>	
121 pentagona. F.I. (wh.) five-angled.	1824. 5. 8. S. ♀. —.
122 cæspitosa. F.I. (st.) tufted.	1827. — S. ♀. —.

123	<i>Beladámboe.</i> R.S. (<i>wh.</i>) whipcord. <i>Convólvulus flagellifórmis.</i> F.I.	E. Indies.	1809.	7. 9. S. ¼.	Rheed. mal. 11. t. 58.
124	<i>reníformis.</i> F.I. (<i>ye.</i>) kidney-leaved.	—	1825.	— S. ¼.	
125	<i>bilobáta.</i> F.I. (<i>pu.</i>) two-lobed.	—	1809.	6. 7. S. ¼.	Rumph. amb. 5.t.159.f.1
126	<i>brasiliénsis.</i> (<i>pu.</i>) three-flowered. Brazil. <i>Convólvulus brasiliénsis.</i> w.	—	1726.	— S. ¼.	Plum. amer. 89. t. 104.
127	<i>pés-cápria.</i> (<i>pu.</i>) single-flowered. E. Indies. <i>marítima.</i> B.R. <i>Convólvulus pés-cápria.</i> F.I.	—	1770.	— S. ¼.	Bot. reg. t. 319.
128	<i>arboréscens.</i> (<i>wh. pu.</i>) tree.	Mexico.	1818.	... S. h.	
ARGYREIA. S.S.	<i>ARGYREIA.</i> Pentandria Monogynia. S.S.				
1	<i>spléndens.</i> (<i>ro.</i>) glossy-leaved. E. Indies.	—	1814.	11. 1. S. h. —	Bot. mag. t. 2628.
	<i>Ipomœa spléndens.</i> B.M. <i>Lettsonia spléndens.</i> F.I.				
2	<i>speciosa.</i> (<i>pu.</i>) broad-leaved. E. Indies.	—	1778.	7. 9. S. h. —	Bot. mag. t. 2446.
	<i>Ipomœa speciosa.</i> B.M. <i>Convólvulus speciosa.</i> w. <i>Lettsonia nervosa.</i> F.I.				
3	<i>argéntea.</i> (<i>pi.</i>) silvery-leaved. E. Indies.	—	1820.	... S. h. —	
	<i>Lettsonia argéntea.</i> F.I.				
4	<i>setosa.</i> F.I. (<i>pi.</i>) bristly-stemmed.	—	—	... S. h. —	
5	<i>cuneáta.</i> B.R. (<i>d. pu.</i>) wedge-leaved.	—	1817.	6. 9. S. h. —	Bot. reg. t. 661.
	<i>Lettsonia cuneáta.</i> F.I. <i>Ipomœa atrosanguinea.</i> B.M. 2170.				
6	<i>cymosa.</i> (<i>hi.</i>) cyme-flowered. E. Indies.	—	1823.	... S. h. —	
	<i>Lettsonia cymosa.</i> F.I.				
7	<i>pomácea.</i> F.I. (<i>pi.</i>) Apple-fruited.	—	1828.	... S. h. —	
8	<i>bóna-nóx.</i> F.I. (<i>wh.</i>) night-flowering.	—	1799.	6. 9. S. h. —	
9	<i>uniflora.</i> F.I. (<i>wh.</i>) kidney-leaved.	—	1817.	... S. h. —	
10	<i>ornátæ.</i> F.I. (<i>wh.</i>) large white-flower'd.	—	1824.	... S. h.	
11	<i>virgáta.</i> slender-twigg'd.	—	1825.	... S. h. —	
DIN'ETUS. B.F.G.	DIN'ETUS. Pentandria Monogynia.				
1	<i>racemosus.</i> B.F.G. (<i>wh.</i>) cluster-flower'd. Nepaul.	—	1823.	7. 11. H. ⊙. —	Swt. br. fl. gar. t. 127.
	<i>Poránia racemosa.</i> F.I.				
2	<i>paniculátus.</i> (<i>wh.</i>) paniced.	E. Indies.	—	... S. h. —	Roxb. corom. 3. t. 235.
	<i>Poránia paniculata.</i> F.I.				
POR'ANA. W.	POR'ANA. Pentandria Digynia.				
	<i>volubilis.</i> w. (<i>wh.</i>) climbing.	E. Indies.	1823.	... S. h. —	Burm. ind. t. 21. f. 1.
EVO'LVLUS. S.S.	EVO'LVLUS. Pentandria Digynia. R.S.				
1	<i>nummulárius.</i> R.S. (<i>bl.</i>) round-leaved.	Jamaica.	1816.	10. S. ⊙.	Jacq. amer. t. 260. f.23.
2	<i>gangéticus.</i> R.S. (<i>bl.</i>) Ganges.	E. Indies.	1823.	— S. ⊙.	
3	<i>emarginátus.</i> R.S. (<i>bl.</i>) kidney-leaved.	—	1816.	— S. ⊙.	Burm. ind. t. 30. f. 1.
4	<i>alsinoides.</i> R.S. (<i>bl.</i>) chickweed-leaved.	—	1733.	6. 7. S. ¼.	Rheed. mal. 11. t. 64.
5	<i>hirsútus.</i> R.S. (<i>pa. bl.</i>) hairy.	Cumana.	1825.	6. 8. S. ¼.	Lam. ill. t. 216. f. 2.
6	<i>lanceolátus.</i> Poi. (<i>pa. bl.</i>) spear-leaved.	S. America.	1818.	— S. ¼.	
7	<i>linifólius.</i> R.S. (<i>bl.</i>) flax-leaved.	Jamaica.	1782.	8. 9. S. ⊙.	Brown. jam. t. 10. f. 2.
8	<i>sericeus.</i> R.S. (<i>wh.</i>) silky.	—	1827.	8. 10. S. ⊙.	
9	<i>incánuus.</i> R.S. (<i>bl.</i>) hoary.	Peru.	1824.	— S. ¼.	
10	Nuttallianus. R.S. (<i>ye.</i>) Nuttall's. argénteus. N.	Missouri.	1821.	5. 6. H. ¼.	
11	<i>latifólius.</i> B.R. (<i>wh.</i>) broad-leaved.	Brazil.	1819.	6. 7. S. ¼.	Bot. reg. t. 401.
CRE'SSA. R.S.	CRE'SSA. Pentandria Digynia. L.				
	<i>crética.</i> R.S. (<i>wh.</i>) Cretan.	Levant.	1822.	7. 9. H. ⊙.	Flor. græc. t. 256.
DICHO'NDRA. R.S.	DICHO'NDRA. Pentandria Digynia. L.				
1	<i>répens.</i> R.S. (<i>wh.</i>) creeping.	N. S. W.	1803.	6. 8. G. ¼.	Sm. ic. ined. 1. t. 8.
2	<i>rotundifolia.</i> L.en. (<i>wh.</i>) round-leaved.	Persia.	1816.	— H. ¼.	
3	<i>caroliniana.</i> R.S. (<i>gr.</i>) Carolina.	Carolina.	1812.	— G. ¼.	
4	<i>sericea.</i> s.s. (<i>wh.</i>) silky.	W. Indies.	1793.	— S. ¼.	Swartz. ic. t. 10.
5	<i>argéntea.</i> R.S. (<i>wh.</i>) silvery.	New Spain.	1818.	— S. ¼.	Willd. hort. ber. t. 81.
FAL'KIA. R.S.	FA'LKIA. Pentandria Digynia. L.				
	<i>répens.</i> R.S. (<i>wh.</i>) creeping.	C. B. S.	1774.	5. 8. G. h.	Andr. deposit. t. 257.
CUSC'UTA. R.S.	DODDER. Pentandria Digynia. R.S.				
1	<i>europæa.</i> R.S. (<i>wh.</i>) great.	England.	8. 9. H. ⊙. —	Eng. bot. v. 6. t. 378.
2	<i>Epithýmum.</i> R.S. (<i>wh.</i>) lesser.	Britain.	7. 10. H. ¼. —	v. 1. t. 55.
3	<i>epilinum.</i> s.s. (<i>wh.</i>) Flax.	Germany.	1821.	— H. ⊙. —	
4	<i>monogyna.</i> R.S. (<i>wh.</i>) one-styled.	Levant.	—	— H. ⊙. —	Flor. græc. t. 257.
5	<i>lupuliformis.</i> R.S. (<i>wh.</i>) rough dotted.	Silesia.	1824.	— H. ⊙. —	Krock. siles. t. 36.
6	<i>macrocárpæ.</i> Led. (<i>wh.</i>) large-podded.	Siberia.	1827.	— H. ⊙. —	
7	<i>chinénensis.</i> R.S. (<i>wh.</i>) Chinese.	China.	1803.	— G. ⊙. —	
8	<i>austrális.</i> B.P. (<i>wh.</i>) New Holland.	N. Holland.	1818.	— G. ¼. —	
9	<i>chilénensis.</i> B.R. (<i>wh.</i>) Chili.	Chili.	1821.	1. 12. G. ¼. —	Bot. reg. t. 603.
10	<i>odoráta.</i> R.S. (<i>wh.</i>) sweet-scented.	Lima.	1820.	— H. ¼. —	
11	<i>verrucósa.</i> B.F.G. (<i>wh.</i>) warted-calyxed.	Nepaul.	1822.	9. 11. H. ¼. —	Swt. br. fl. gar. t. 6.
12	Hookéri. (<i>wh.</i>) Hooker's. refléxa, var. <i>verrucósa.</i> H.E.F. excl. synon.	E. Indies.	1823.	— H. ⊙.	Hook. ex. flor. t. 150.
			2 B 3		

ORDO CXLIX.

BORAGINEÆ. Brown prodr. I. p. 492.

TIARI'DIUM. S.S.	TIARI'DIUM.	Pentandria Monogynia. S.S.				
1 indicum. s.s. (li.) Indian.	W. Indies.	1713.	6. 8.	S. ♂.	Bot. mag. t. 1837.	
<i>Heliotropium indicum.</i> B.M.						
2 velutinum. R.S. (wh.) villous-stalked.	E. Indies.	—	S. ♂.	Rheed. mal. 10. t. 48.	
HELIOTR'OPIUM. S.S. TURNSOLE.	Pentandria Monogynia. L.					
1 peruvianum. s.s. (wh.) Peruvian.	Peru.	1757.	5. 9.	G. ♀.		
β hybridum. (wh.) More's hybrid.	Hybrid.	1815.	—	G. ♀.		
2 corymbosum. s.s. (bl.) large-flowered.	—	1808.	—	G. ♀.	Bot. mag. t. 1609.	
<i>grandiflorum.</i> Len.						
3 maroccanum. s.s. (wh.) Morocco.	Morocco.	1821.	—	G. ♀.		
4 parviflorum. s.s. (wh.) small-flowered.	India.	1732.	7. 9.	S. ♂.	Dill. elth. t. 146. f. 175.	
5 europæum. s.s. (wh.) European.	S. Europe.	1562.	6. 10.	H. ♂.	Jacq. aust. 3. t. 207.	
6 commutatum. R.S. (wh.) smaller.	—	—	H. ♂.		
7 oblongifolium. L.en. (w.) oblong-leaved.	—	1827.	—	H. ♂.		
8 suavolens. s.s. (wh.) sweet-scented.	Caucasus.	1823.	—	H. ♂.		
9 coromandelianum. w. (w.) Coromandel.	E. Indies.	1812.	7. 10.	S. ♂.		
10 curassavicum. s.s. (wh.) glaucous.	S. America.	1731.	6. 7.	S. ♂.	Bot. mag. t. 2669.	
11 chenopodioides. W.en. (w.) smooth-leaved.	—	1820.	—	S. ♂.		
12 anisophyllum. B.G. (w.) unequal-leaved.	Africa.	1822.	—	S. ♂.	Beau. fl. d. ow. 2. t. 94.	
13 zeylanicum. R.S. (wh.) Ceylon.	Ceylon.	1815.	—	S. ♂.	Burm. ind. t. 16. f. 2.	
14 linifolium. R.S. (wh.) flax-leaved.	C. B. S.	—	—	G. ♀.		
<i>Myosotis fruticosa.</i> w.						
15 demissum. R.S. (wh.) dwarf.	S. America.	1752.	5. 6.	S. ♀.	Plum. ic. 227. f. 2.	
<i>Tournefortia humilis.</i> w.						
16 supinum. s.s. (wh.) trailing.	S. Europe.	1640.	6. 7.	H. ♂.	Flor. græc. t. 157.	
LITHOSPER'MUM. S.S. GROMWELL.	Pentandria Monogynia. L.					
1 fruticosum. s.s. (re.) shrubby.	S. Europe.	1683.	5. 6.	H. ♀.		
2 prostratum. s.s. (pu.) prostrate.	—	1825.	—	H. ♀.	Flor. græc. t. 161.	
3 sericeum. s.s. (st.) silky.	N. America.	1824.	—	H. ♀.	Lois. fl. gall. add. I. t. 4.	
4 divaricatum. s.s. (st.) spreading.	Africa.	1829.	7. 9.	H. ♀.		
5 hispidulum. s.s. (bl.) bristly.	Greece.	1828.	—	F. ♀.		
6 arvensis. s.s. (st.) corn.	Britain.	—	F. ♂.		
7 officinale. s.s. (ye.) officinal.	—	5. 6.	H. ♂.	Eng. bot. v. 2. t. 123.	
8 latifolium. s.s. (st.) broad-leaved.	America.	1816.	—	H. ♀.	—	v. 2. t. 134.
9 distichum. s.s. (st.) two-spiked.	Mexico.	1806.	—	G. ♀.		
10 orientale. s.s. (ye.) yellow-flower'd.	Levant.	1713.	—	H. ♀.	Jacq. fragm. t. 48. f. 3.	
11 lineatum. s.s. (st.) lined.	—	1816.	—	H. ♀.	Bot. mag. t. 515.	
12 apulum. s.s. (ye.) small.	S. Europe.	1768.	6. 7.	H. ♂.	Flor. græc. t. 158.	
13 crassifolium. s.s. (st.) thick-leaved.	Levant.	1825.	—	H. ♀.		
14 decumbens. s.s. (st.) decumbent.	Persia.	—	7. 9.	H. ♂.	Vent. cels. t. 37.	
15 villsum. s.s. (pu.) villous.	S. Europe.	1820.	6. 7.	H. ♀.		
16 tintorium. R.S. (st.) dyer's.	—	1596.	6. 10.	H. ♀.	Eng. bot. v. 2. t. 117.	
17 purpureo-cœruleum. (bl.) creeping.	England.	5. 6.	H. ♀.	Viv. frag. ital. t. 5.	
18 graminifolium. s.s. (bl.) grass-leaved.	S. Europe.	1826.	—	H. ♀.	Jacq. ic. 2. t. 313.	
19 tenuiflorum. s.s. (wh.) slender-flower'd.	Egypt.	1796.	—	H. ♂.	Linn. dec. 1. t. 7.	
20 disperrimum. s.s. (li.) two-seeded.	Spain.	1799.	6. 7.	H. ♂.	Swt. br. fl. gar. t. 121.	
21 davuricum. s.s. (bl.) Daurian.	Dauria.	1812.	5. 6.	H. ♀.		
<i>Pulmonaria davurica.</i> B.M. 1743.						
22 denticulatum. s.s. (pu.) toothed.	N. America.	1801.	—	H. ♀.		
23 sibiricum. s.s. (pu.) Siberian.	Siberia.	—	6. 7.	H. ♀.	Gmel. sib. 4. n. 15. t. 39.	
24 maritimum. s.s. (bl.) sea-side.	Britain.	—	H. ♀.	Eng. bot. v. 6. t. 368.	
<i>Pulmonaria maritima.</i> E.B.						
25 parviflorum. (bl.) small-flowered.	N. America.	1824.	—	H. ♀.		
26 púlchrum. s.s. (bl.) Virginian.	—	1699.	3. 5.	H. ♀.	Bot. mag. t. 160.	
<i>Pulmonaria virginica.</i> B.M.						
27 paniculatum. s.s. (pu.) paniced.	Hudson's Bay.	1778.	5. 6.	H. ♀.	Bot. reg. t. 146.	
<i>Pulmonaria paniculata.</i> B.R.						
28 lanceolatum. (bl.) margined.	Louisiana.	1813.	—	H. ♀.		
<i>Pulmonaria lanceolata.</i> Ph. marginata. N.						
29 simplicissimum. s.s. (bl.) simple-stalked.	Siberia.	1829.	—	H. ♀.		
BA'TSCHIA. M.	BA'TSCHIA.	Pentandria Monogynia.				
1 Gmelini. M. (ye.) Carolina.	Carolina.	1812.	5. 7.	H. ♀.		
<i>caroliniana.</i> R.S. <i>Lithospérnum hirtum.</i> s.s.						

2 canescens. M.	(ye.) canescent.	N. America.	5. 7. H. 4.	Mich. amer. 1. t. 14.
3 longiflora. N.	(ye.) long-flowered.	Missouri.	1812.	— H. 4.	Moris. his. s. 11. t. 29. f. 5.
4 decumbens. N.	(ye.) decumbent.	N. America.	1826.	— H. 4.	Eng. bot. v. 23. t. 1628.
5 conspicua. B.	(ye.) conspicuous.		1827.	— H. 4.	Swt. br. fl. gar. ic.
PULMONARIA. S.S.	LUNGWORT.	Pentandria Monogynia. L.			Eng. bot. v. 2. t. 118.
1 azurea. s.s.	(bl.) azure.	Hungary.	1823.	5. H. 4.	Moris. s. 11. t. 29.
2 angustifolia. s.s.	(pu.) narrow-leaved.	Britain.	4. 5. H. 4.	DC. ic. ined. t. 64.
3 tuberosa. s.s.	(pu.) tuberous-rooted.	Hungary.	1824.	5. H. 4.	
4 mollis. s.s.	(pu.) soft-leaved.		1816.	— H. 4.	
5 officinalis. R.S.	(pu.) common.	England.	— H. 4.	
6 saccharata. L.en.(wh.)	white-flowered.	Germany.	1816.	— H. 4.	
7 grandiflora. R.S.	(pu.) large-flowered.		...	— H. 4.	
8 oblongata. Schr.	(pu.) oblong-leaved.	Germany.	1820.	— H. 4.	
ONOSM'ODIUM. M.	ONOSM'ODIUM.	Pentandria Monogynia. R.S.			
1 hispidum. R.S.	(st.) hispid.	N. America.	1759.	6. H. 4.	Moris. s. 11. t. 28. f. 3.
2 molle. R.S.	(wh.) soft.		1812.	6. 8. H. 4.	Mich. amer. t. 15.
ONO'SMA. R.S.	ONO'SMA.	Pentandria Monogynia. L.			Flor. græc. t. 172.
1 echoides. s.s.	(ye.) hairy.	S. Europe.	1683.	3. 6. H. 4.	Colum. ecphr. p. 183. f. 2.
2 giganteum. s.s.	(ye.) gigantic.	Tauria.	1818.	— H. 4.	W. et K. hung. 3. t. 279.
3 montanum. R.S.	(ye.) mountain.	Levant.	1827.	— H. 4.	2. t. 173.
4 arenarium. R.S.	(ye.) sand.	Hungary.	1804.	3. 7. H. 4.	Bot. mag. t. 889.
5 stellulatum. W.K.	(ye.) starry-haired.		1816.	— H. 4.	
6 taureum. B.M.	(go.) golden-flower'd.	Tauria.	1801.	4. 6. F. 4.	
7 tinctorium. R.S.	(st.) dyer's.	Volga.	1828.	— H. 4.	Flor. græc. t. 173.
8 rupreste. R.S.	(ye.) rock.	Caucasus.	1823.	— H. 4.	
9 erectum. R.S.	(ye.) upright.	Crete.	1828.	— F. 4.	
10 orientale. s.s.	(st.) oriental.	Levant.	1752.	5. 6. G. 4.	Pall. it. 2. app. t. 50.
11 divaricatum. s.s.	(st.) spreading.	Caucasus.	1818.	4. 6. H. 4.	Bot. mag. t. 2248.
12 simplicissimum. R.S. (st.)	linear-leaved.	Siberia.	1768.	— H. 4.	
13 trimérivum. s.s.	(st.) three-nerved.	Mexico.	1825.	— G. 4.	
14 sericeum. R.S.	(ye.) silky-leaved.	Levant.	1752.	6. 7. F. 4.	
15 synanthemum. F.	(ye.) connected.	Altay.	1827.	— H. 4.	
MO'LTKIA. R.S.	Mo'LTKIA.	Pentandria Monogynia.			
cerulea. R.S.	(bl.) blue-flowered.	Persia.	1829.	4. 6. H. 4.	
SY'MPHYTUM. R.S.	COMFREY.	Pentandria Monogynia. L.			Eng. bot. v. 12. t. 817.
1 officinale. R.S.	(wh.) common.	Britain.	5. 7. H. 4.	
β purpurascens.	(pu.) purple.		— H. 4.	
2 bohemicum. P.S.	(re.) red-flowered.	Bohemia.	— H. 4.	Eng. bot. v. 23. t. 1502.
3 caucasicum. R.S.	(bl.) Caucasian.	Caucasus.	1816.	— H. 4.	Reich. icon. t. 220.
4 tuberosum. R.S.	(wh.) tuberous-root'd.	Scotland.	5. 10. H. 4.	Bot. mag. t. 1912.
5 bulbosum. s.s. bulbous-rooted.	Levant.	1829.	5. 7. H. 4.	— t. 1787.
6 orientale. R.S.	(wh.) eastern.	Turkey.	1752.	— H. 4.	
7 bullatum. s.s.	(wh.) blistered.	Tauria.	1806.	— H. 4.	
tauricum. B.M.					
8 cordatum. W.K.	(st.) heart-leaved.	Hungary.	1813.	— H. 4.	W. et K. hung. 6. t. 7.
9 asperimum. R.S.	(bl.) roughest.	Caucasus.	1799.	5. 9. H. 4.	Bot. mag. t. 929.
10 echinatum. R.S.	(pu.) bristly.		— H. 4.	
11 peregrinum. s.s.	(wh.) oblique-leaved.	Poland.	1816.	5. 7. H. 4.	
CERINTHE. R.S.	HONEYWORT.	Pentandria Monogynia. L.			
1 major. R.S.	(ye. pu.) great.	S. Europe.	1596.	7. 8. H. ⊖.	Bot. mag. t. 333.
2 áspéra. R.S.	(pu. ye.) rough.		1633.	— H. ⊖.	Flor. græc. t. 170.
3 retorta. F.G.	(pu. ye.) bent topped.	Levant.	1828.	— H. ⊖.	— t. 171.
4 minor. R.S.	(ye.) small.	Austria.	1570.	6. 10. H. ⊖.	Jacq. aust. 2. t. 124.
5 alpina. R.S.	(pu. ye.) alpine.	M. Carpathian.	1827.	— H. ⊖.	
6 maculata. R.S.	(ye. pu.) spotted.	Tauria.	1804.	— H. ♂.	
'ECHIUM. R.S.	VIPER'S-BUGLOSS.	Pentandria Monogynia. L.			
1 fruticosum. R.S. (wh. bh.)	frutescent.	C. B. S.	1759.	5. 6. G. ♂.	Bot. reg. t. 86.
2 hispidum. R.S.	(wh.) hispid-leaved.		1816.	— G. ♂.	
3 verrucosum. R.S.	(wh.) warted.		1820.	5. 7. G. ♂.	
4 laevigatum. R.S.	(wh.) smooth-stalked.		1774.	6. 7. G. ♂.	
5 glaucophyllum. R.S. (bl.)	glaucous-leaved.		1792.	6. 8. G. ♂.	Jacq. ic. t. 312.
6 glábrum. R.S.	(wh.) smooth.		1791.	5. 6. G. ♂.	
7 paniculatum. R.S. (wh.)	panicled.		1815.	— G. ♂.	
8 grandiflorum. R.S. (ro.)	great-flowered.		1787.	6. 7. G. ♂.	Bot. reg. t. 124.
formosum. P.S.					
9 ferocissimum. A.R.	(bl.) prickly-stalked.		1794.	— G. ♂.	Andr. reposit. t. 39.
10 petræum. s.s.	(bl.) rock.	Dalmatia.	1828.	— H. ♂.	
11 calycinum. R.S.	(bl. ye.) large-calyxed.	S. Europe.	1829.	6. 8. H. ⊖.	Viv. frag. ital. 1. t. 4.
			2 B 4		

BORAGINEÆ.

12 spicatum. R.S. (wh.) spiked dwarf.	C. B. S.	1799.	3. 5.	G. h.
13 caudatum. R.S. (re.) tailed.	—	1815.	6. 9.	G. h.
14 capitatum. R.S. (re.) headed.	—	1822.	—	G. h.
15 spherocephalon. s.s.(w.) round-headed.	—	1824.	—	G. h.
16 argenteum. R.S. (bl.) silvery.	—	1789.	6. 7.	G. h.
17 lasiophyllum. L.en.(w.)woolly-leaved.	—	1819.	—	G. h.
18 Swartzii. R.S. (bl.) Swartz's.	C. B. S.	1816.	—	G. h.
19 incanum. R.S. (bl.) hoary.	—	—	4. 6.	G. h.
20 strigosum. R.S. (vi.) strigose.	—	1819.	6. 7.	G. h.
21 giganteum. R.S. (wh.) gigantic.	Canaries.	1779.	7. 11.	G. h.
22 aculeatum. R.S. (st.) prickly.	—	1815.	5. 8.	G. h.
23 molle. R.S. (wh.) soft-leaved.	—	—	—	G. h.
24 nervosum. H.K. (bl.) sinewy-leaved.	Madeira.	1777.	6. 8.	G. h.
25 bifrons. DC. (bh.) two-fronted.	Canaries.	1815.	—	G. h.
26 ambiguum. DC. (bh.) ambiguous.	—	—	5. 7.	G. h.
27 virescens. DC. (pa.bl.) green-leaved.	—	—	—	G. h.
28 candicans. R.S. (bl.) hoary tree.	Madeira.	1777.	5. 6.	G. h.
29 cynoglossoides. R.S. (bl.)hound's-tongue-ld.	Canaries.	1816.	—	G. h.
30 densiflorum. DC. (li.) dense-flowered.	—	—	—	G. h.
31 fastuosum. H.K. (bl.) fastuous.	—	1779.	4. 8.	G. h.
32 strictum. R.S. (bl.) upright.	—	—	5. 12.	G. δ.
33 lineatum. R.S. (bh.) lined.	—	1815.	5. 6.	G. h.
34 macrophyllum. R.S.(bl.)large-leaved.	—	1812.	—	G. h.
35 foliosum. R.S. (wh.) leafy.	—	—	—	G. h.
36 simplex. DC. (wh.) simple-stalked.	Canaries.	—	5. 12.	G. δ.
37 brachyanthum. R.S.(w.)short-flowered.	Madeira.	1820.	5. 7.	G. h.
38 scabrum. R.S. (bl.) rough.	C. B. S.	—	—	G. h.
39 plantagineum. R.S.(pu.)Plantain-leav'd. Italy.	—	1776.	7. 10.	H. ⊙.
40 lusitanicum. R.S. (bl.) Portugal.	S. Europe.	1731.	7. 8.	H. 4.
41 plantaginoides. R.S.(bl.)Plantain-like.	—	1825.	—	H. ⊙.
42 prostratum. R.S. (re.) prostrate.	Egypt.	1824.	—	H. 4.
43 macranthum. R.S. (vi.) large-flowered.	Barbary.	—	7. 10.	H. ⊙.
grandiflorum. Desf. non Andr.	—	—	—	Flor. græc. t. 179.
44 australis. R.S. (re.pu.) oval-leaved.	S. Europe.	1824.	—	H. ⊙.
45 Símsii. (re.bl.) Sims's.	—	1816.	—	H. ⊙.
creticum. B.M. non Flor. græc.	—	—	—	Swt. br. fl. gar. t. 101.
46 maritimum. R.S. (bl.) sea-side.	Italy.	1815.	—	H. ⊙.
47 italicum. R.S. (wh.) white.	Jersey.	—	7. 8.	H. δ.
48 pyrenaicum. R.S. (bl.) Pyrenean.	Pyrenees.	1815.	—	H. δ.
49 asperimum. M.B. (bl.) roughest.	Caucasus.	1826.	—	H. δ.
50 thyrsoidéum. R.S. (bl.) bunch-flowered.	—	1823.	—	H. δ.
51 rúbrum. R.S. (re.) red-flowered.	Hungary.	1791.	—	H. 4.
52 vulgare. R.S. (bl.) common.	Britain.	—	—	H. δ.
β ábum. (wh.) white-flowered.	—	—	—	H. δ.
53 tuberculatum. R.S.(bl.) tubercled.	S. Europe.	1826.	—	H. δ.
54 violaceum. R.S. (vi.bl.) violet-flowered.	Austria.	1658.	—	H. ⊙.
55 parviflorum. H.K. (bl.) small-flowered.	Barbary.	1798.	—	H. ⊙.
micranthum. R.S.	—	—	—	Bot. mag. t. 1822.
56 tenuis. s.s. (bl.) slender.	—	1824.	—	H. ⊙.
57 Sibthórii. R.S. (ro.) Sibthorp's.	Naples.	—	—	Eng. bot. v. 29. t. 2081.
hispidum. F.G. nec aliorum. Elegans. Lehm.	—	—	—	Bocc. mus. 2. t. 78.
58 diffusum. R.S. (re.) spreading.	Crete.	1827.	7. 9.	H. ⊙.
59 creticum. R.S. (pu.) Cretan.	Levant.	1683.	—	H. ⊙.
60 orientale. R.S. (pu.) oriental.	—	1780.	7. 8.	H. ⊙.
61 dahuricum. F. (bl.) Dahurian.	Dahuria.	1827.	—	H. δ.
62 angustifolium. s.s. (bl.) narrow-leaved.	Spain.	1826.	—	H. ⊙.
63 Lagascánum. R.S. (li.) Lagasca's.	—	—	—	H. 4.
64 salmanticum. R.S.(pu.) long-stamened.	—	1824.	—	H. ⊙.
N'ONEA. DC.	N'ONEA. Pentandria Monogynia. R.S.	—	—	Flor. græc. t. 181.
1 violacea. DC. (vi.) violet-coloured.	S. Europe.	1686.	5. 7.	H. ⊙.
Lycópsis vesicaria. w.	—	—	—	Moris. s. 11. t. 26. f. 11.
2 nigricans. DC. (d.pu.) black-flowered.	—	—	—	Zanon. hist. 56. t. 38.
3 púlla. DC. (d.pu.) dark-flowered.	Germany.	1648.	6. 7.	H. 4.
4 ciliata. R.S. (ye.) fringed.	Levant.	1804.	—	H. ⊙.
5 rosea. Len. (ro.) rose-coloured.	Caucasus.	1824.	—	H. ⊙.
Anchusa ròsea. M.B. Lycópsis ròsea. s.s.	—	—	—	Jacq. aust. 2. t. 188.
6 setosa. R.S. (ye.) bristly.	Iberia.	1826.	—	H. ⊙.
7 versicolor. (bl.wh.) various-colored.	—	1820.	—	H. ⊙.
Anchusa versicolor. R.S.	—	—	—	Andr. deposit. t. 154.
			Vent. malm. t. 71.	

8 lùtea. n.c.	(ye.) pale yellow.	S. Europe.	1805.	6. 7. H. ⊖.	Nocc. hort. tran. t. 3.
Anchùsa lùtea. R.S.	<i>Lycópsis lùtea</i> . s.s.				
9 pícta.	(bl.std.) painted.	Tauria.	1818.	— H. ⊖.	
Anchùsa pícta. M.B.	<i>Lycópsis pícta</i> . s.s.				
LYCOP'SIS. L.	WILD BUGLOSS.	Pentandria Monogynia. L.			
arvén's. L.	(bl.) common.	Britain.	5. 8. H. ⊖.	Eng. bot. v. 14. t. 938.
ASPERUGO. L.	GERMAN MADWORT.	Pentandria Monogynia. L.			
procumbens. L.	(bl.) procumbent.	Britain.	4. 5. H. ⊖.	Eng. bot. v. 10. t. 661.
ANCH'USA. R.S.	BUGLOSS.	Pentandria Monogynia. L.			
1 Barrelieri. R.S.	(bl.) Barrelier's.	Italy.	1815.	6. 7. H. Ȑ.	Barrel. ic. 333.
2 paniculàta. R.S.	(bl.) paniced.	Levant.	1777.	6. 10. H. Ȑ.	Flor. græc. t. 163.
3 itálica. B.R.	(bl.) Italian.	S. Europe.	1597.	— H. Ȑ.	Bot. reg. t. 483.
4 Millèri. R.S.	(ro.) Miller's.	— H. ⊖.	
5 myosotidiflòra. s.s.	(bl.) Myosotis-flower'd.	Caucasus.	1828.	— H. Ȑ.	MB. cent. pl. rar. t. 23.
6 stylòsa. R.S.	(bl.) long-styled.	Tauria.	1824.	— H. ⊖.	
7 pròcera. s.s.	(bl.) tall.	Poland.	1822.	— H. Ȑ.	
8 lycosoides. L.en.	(vi.) Lycopsis-like.	—	— H. Ȑ.	
9 variegatà. s.s.	(bl.cr.) variegated.	Levant.	1683.	6. 7. H. ⊖.	Flor. græc. t. 178.
Lycópsis variegatà. F.G.					
10 ovàta. s.s.	(bl.) oval-leaved.	1796.	— H. ⊖.	Buxb. cent. 5. t. 30.
Lycópsis orientális. R.S.					
11 tinctoria. R.S.	(bl.) dyer's.	1596.	6. 10. H. Ȑ.	Flor. græc. t. 166.
12 parviflòra. R.S.	(bl.) small-flowered.	Levant.	1827.	— H. ⊖.	
13 aggregatà. s.s.	(bl.) clustered.	—	— H. ⊖.	Flor. græc. t. 107.
parviflòra. Flor. græc. non w.					
14 hispida. R.S.	(vi.) hispid.	Egypt.	1823.	— H. Ȑ.	
15 tenélia. R.S.	(wh.) slender.	China.	1816.	— H. ⊖.	Jacq. f. ecl. t. 29.
zeylánica. Jacq.					
16 caespitosa. F.G.	(bl.) tufted.	Levant.	1828.	— H. Ȑ.	Flor. græc. t. 169.
17 officinalis. R.S.	(vi.) common.	Britain.	— H. Ȑ.	Eng. bot. v. 10. t. 662.
18 incarnatà. Schr.	(fl.) flesh-coloured.	Europe.	— H. Ȑ.	
19 undulatà. R.S.	(pu.) wave-leaved.	S. Europe.	1739.	6. 8. H. Ȑ.	Bot. mag. t. 2119.
20 ochroleuca. R.S.	(st.) pale-flowered.	Caucasus.	1810.	7. 8. H. Ȑ.	— t. 1608.
21 angustifòlia. R.S.	(bl.) narrow-leaved.	Europe.	1640.	5. 6. H. Ȑ.	Flor. græc. t. 164.
22 capénsis. R.S.	(vi.) Cape.	C. B. S.	1800.	7. 8. G. ♂.	Andr. reposit. t. 336.
23 hybrida. R.S.	(pu.) villous.	Naples.	1818.	7. 10. H. ⊖.	Ten. fl. neap. t. 11.
24 Gmelini. s.s.	(bl.) Gmelin's.	Siberia.	1829.	— H. Ȑ.	
25 alpéstris. s.s.	(bl.) alpine.	Caucasus.	— H. Ȑ.	
26 Agárdhii. R.S.	(pu.) tubercle-leav'd.	Siberia.	1823.	7. 9. H. Ȑ.	
27 verrucosà. R.S.	(wh.) warted.	Egypt.	7. 10. H. ⊖.	Jacq. vind. 3. t. 21.
Asperugo ægyptiaca. Jacq.					
28 rupéstris. B.P.	(bl.) rock.	Siberia.	1802.	6. 8. H. Ȑ.	Pall. it. 3. ap. t. E. f. 3.
Myosotis rupéstris. W.					
MYOSOTIS. R.S.	SCORPION-GRASS.	Pentandria Monogynia. L.			
1 palústris. E.B.	(bl.) marsh.	Britain.	6. 9. H.w. Ȑ.	Eng. bot. v. 28. t. 1973.
scorpioides. W.					
2 caespitosa. s.s.	(bl.) tufted.	5. 7. H.w. Ȑ.	
3 intermédia. s.s.	(bl.) intermediate.	4. 6. H. ♂.	Fl. dan. 583. f. maj.
4 sylvática. R.S.	(blye.) wood.	6. 8. H. Ȑ.	Raii syn. t. 9. f. 2.
5 commutatà. R.S.	(bl.) changed.	Europe.	— H. ♂.	
6 alpéstris. R.S.	(bl.) rock.	Scotland.	— H. Ȑ.	Eng. bot. v. 36. t. 2559.
rupicola. E.B.					
7 lithospermifòlia. R.S. (bl.) Gromwell-leav'd.	Caucasus.	1823.	— H. Ȑ.		
8 suavolens. R.S.	(bl.) sweet-scented.	Europe.	—	— H. Ȑ.	
9 sericea. s.s.	(bl.) silky-leaved.	Siberia.	1802.	— H. Ȑ.	
10 villòsa. s.s.	(bl.) villous.	—	1828.	— H. Ȑ.	
11 sparsiflòra. R.S.	(wh.) loose-flowered.	Europe.	1818.	— H. ⊖.	
12 stricta. R.S.	(bl.ye.) upright.	—	—	— H. ⊖.	
13 arvén's. E.B.	(bl.) corn.	Britain.	6. 9. H. ⊖.	Eng. bot. v. 36. t. 2558.
14 versicolor. E.B. (bl.ye.) various-colour'd.	—	4. 6. H. ⊖.	— t. 480. f. 1.
15 nàna. R.S.	(bl.) dwarf.	—	6. 8. H. Ȑ.	Hacq. pl. alp. car. t. 2.f. 6.
16 peduncularis. s.s.	(bl.) peduncled.	Astracan.	1822.	— H. ⊖.	
17 clavatà. s.s.	(bl.) clavate.	Siberia.	1829.	— H. ⊖.	
18 australis. B.P.	(bl.) southern.	N. S. W.	1824.	— G. ⊖.	
19 ungulatà. F.	(bl.) clawed.	Siberia.	1822.	— H. ⊖.	
ECHINOSPERMUM. S.S. ECHINOSPERMUM.	Pentandria Monogynia. S.S.				
1 zeylanicum. s.s.	(bl.) Ceylon.	Ceylon.	1826.	6. 9. S. ⊖.	
2 virginianum. s.s.	(pa.) Virginian.	Virginia.	1699.	6. 7. H. ♂.	Moris. s. 11. t. 30. f. 9.
Myosotis virginiana. w. Rochelia virginiana. R.S.					

BORAGINEÆ.

3 deflêxum. s.s. (bl.) deflexed.	Hungary.	1823.	6. 7. H. ♂.	Wahl. a. holm. 1810. t. 4.
4 Láppula. s.s. (bl.) common.	Europe.	1656.	4. 8. H. ♂.	Flor. dan. t. 692.
5 squarròsum. L.en.(bl.) squarrose. pátulum. s.s.	Siberia.	1802.	— H. ♂.	
6 Redówskii. L.A. (bl.) Redowski's.	Russia.	1821.	6. 7. H. ♂.	
7 barbátum. s.s. (bl.) bearded.	Tauria.	1819.	— H. ♂.	
8 marginátum. s.s. (bl.) margined.	Astracan.	—	— H. ♂.	
9 heteracanthum. Led.(bl.) various-spined.	Siberia.	1826.	— H. ♂.	
10 grandifòrum. Led.(bl.) large-flowered.	—	1827.	— H. ♂.	
OMPHAL'ODES. L.A. OMPHAL'ODES. Pentandria	Monogynia. L.en.			
1 nítida. L.A. (bl.) shining.	Portugal.	1812.	4. 6. H. ♀.	Bot. mag. t. 2529.
Cynoglòssum nítidum. B.M. Picòtia nítida. R.S.				
2 verna. L.en. (bl.) Comfrey-leav'd. S. Europe.		1633.	3. 5. H. ♀.	Bot. mag. t. 7.
Cynoglòssum Omphalòdes. B.M. Picotia verna. R.S.				
3 brassicæfòlia. L. (wh.) stem clasping.	Spain.	1824.	6. 8. H. ♂.	Lehm. m. ber. 8. 2. t. 6.
amplexicaulis. Lehm. Picotia brassicæfòlia. R.S.				
4 littoralis. L.A. (wh.) sea-side.	France.	1827.	— H. ♂.	
5 linifòlia. L.A. (wh.) Venus Navelwort. Portugal.		1648.	— H. ♂.	Barrel. ic. 1234.
6 lithospermifòlia. L.A. (bl.) loose-racemed.	Levant.	1828.	4. 6. H. ♀.	
7 scorpioides. L.A. (bl.) variable-leaved.	Bohemia.	1823.	6. 8. H. ♂.	Lehm. m. ber. 8. 2. t. 7.
8 sempervirens. D.P. (bl.) evergreen.	Britain.	5. 6. H. ♀.	Eng. bot. v. 1. t. 45.
Anchusa sempervirens. E.B.				
RIND'ERA. R.S. RIND'ERA. Pentandria Monogynia. R.S.				
1 laevigata. R.S. (wh.) smooth.	Siberia.	1824.	6. 7. H. ♀.	Pall. ross. 2. t. 88.
tetráspis. Pall. Cynoglòssum Rindèra. Pall. Cynoglòssum laevigatum. w.				
2 magellánensis. R.S. (vi.) woolly.	Italy.	1823.	5. 7. H. ♀.	
Cynoglòssum magellánense. Tenore.				
3 Colúmnæ. R.S. (pu.) Columna's.		1824.	— H. ♂.	Column. ephr. 1.t. 178.
4 cristata. R.S. (pu.) crested.	Armenia.	1829.	— H. ♂.	
MA'TTIA. R.S. MA'TTIA. Pentandria Monogynia. R.S.				
1 umbellatæ. R.S. (re.ye.) umbel-flowered.	Hungary.	1818.	6. 7. H. ♀.	W. et K. hung. 2.t.148.
2 lanata. R.S. (ro.) woolly.	Levant.	1800.	— G. ♀.	Tourn. an. mus. 10.t.37.
Cynoglòssum lanatum. w.				
CYNOGL'OSUM. R.S. HOUND'S-TONGUE. Pentandria Monogynia. L.				
1 officinale. R.S. (ro.) common.	Britain.	6. 7. H. ♂.	Eng. bot. v. 13. t. 921.
2 Ha'nkii. R.S. (pu.) rough-leaved.	Hungary.	1815.	— H. ♂.	
3 sylvaticum. R.S. (pu.) green-leaved.	Britain.	— H. ♂.	Eng. bot. v. 23. t. 1642.
4 Dioscóridis. R.S. (bl.) Dioscoride's.	Dauphiny.	1818.	— H. ♂.	
5 bicolor. R.S. (wh.re.) two-coloured.	Germany.	—	— H. ♂.	
6 píctum. R.S. (re.bl.) Madeira.	Madeira.	1658.	8. G. ♂.	
7 lanceolatum. L.A. (wh.bl.) spear-leav'd.	Africa.	1806.	7. 8. H. ♂.	
8 virginicum. s.s. (bl.wh.) stem-clasping.	N. America.	1812.	6. 7. H. ♀.	
amplexicaule. M.				
9 clandestinum. R.S. (vi.) hidden-flower'd.	Barbary.	1823.	— G. ♂.	
10 canescens. R.S. (bl.) canescent.	E. Indies.	1815.	7. 10. G. ♂.	Desf. atl. 1. t. 42.
racemosum. F.I.				Jacq. scheenb. 4. t. 489.
11 apenninum. R.S. (vi.) Apennine.	Italy.	1731.	4. 7. H. ♂.	Sabb. hort. rom. 2. t. 36.
12 tomentosum. L.A. (vi.) woolly-leaved.	—	1823.	5. 8. H. ♀.	
13 magellánense. T.S. (ro.) woolly.	—	—	5. 7. H. ♀.	
14 holosericeum. s.s. (vi.) silky.	Caucasus.	1827.	5. 9. H. ♂.	
15 diffusum. L.A. (wh.) spreading.	E. Indies.	1820.	7. 10. H. ♂.	
16 australe. B.P. (vi.) New Holland.	N. S. W.	—	5. 7. G. ♀.	
17 cheirifolium. s.s. (pu.) silvery-leaved.	Levant.	1596.	6. 7. H. ♂.	
Anchusa lanata. R.S.				
TRICHO'DE'SMA. B.P. TRICHO'DE'SMA. Pentandria Monogynia. R.S.				
1 indicum. R.S. (bl.) Indian.	E. Indies.	1759.	6. 10. H. ♂.	Pluk. alm. 30. t. 76. f. 3.
Borago indica. L.				
2 africánum. R.S. (bl.) African.	C. B. S.	—	7. 8. H. ♂.	Isn. act. par. 1718. t. 11.
3 zeylánicum. R.S. (bl.) Ceylon.	E. Indies.	1799.	— S. ♂.	Jacq. ic. 2. t. 314.
BOR'AGO. R.S. BORAGE. Pentandria Monogynia. L.				
1 officinalis. R.S. (bl.) common.	Britain.	6. 9. H. ♂.	Eng. bot. v. 1. t. 36.
β albiflora.	(wh.) white-flowered.	— H. ♂.	
2 orientalis. R.S. (bl.) oriental.	Turkey.	1752.	3. 5. H. ♀.	Bot. reg. t. 288.
3 crética. R.S. (bh.) Cretan.	Greece.	1823.	5. 6. H. ♀.	Flor. græc. t. 176.
4 laxiflora. R.S. (bl.) bell-flowered.	Corsica.	1813.	5. 8. H. ♂.	Bot. mag. t. 1798.
5 crassifolia. R.S. (fl.) thick-leaved.	Persia.	1824.	— H. ♀.	Venten. cels. t. 100.
TOURNEFO'R'TIA. R.S. TOURNEFO'R'TIA. Pentandria Monogynia. L.				
1 cymosa. R.S. (fl.) broad-leaved.	Jamaica.	1777.	7. S. ♂.	Jacq. ic. 1. t. 31.
2 foetidissima. R.S. (fl.) Tobacco-leav'd.	—	1739.	9. S. ♂.	Plum. ic. 226. t. 230.

3 bicolor. R.S. (<i>wh.gr.</i>) two-coloured.	Jamaica.	1812.	S. h.	
4 suffruticosa. R.S. (<i>wh.</i>) hoary-leaved.	—	1759.	6. 9.	S. h.	Sloan. hist. 2. t. 162. f.4.
5 mutabilis. R.S. (<i>wh.</i>) changeable.	Java.	1820.	—	S. h.	Venten. choix. t. 3.
6 maculata. R.S. (<i>st.</i>) spotted-fruited.	Carthagena.	1828.	—	S. h.	
7 gnaphalodes. R.S. (<i>wh.</i>) Gnaphalium-like.	W. Indies.	1820.	—	S. h.	Jacq. amer. t. 173. f. 11.
<i>Heliotropium gnaphalodes.</i> w.					
8 velutina. K.S. (<i>wh.</i>) velvetty.	Mexico.	1826.	—	G. h.	H. et B. am. 3. t. 201.
9 umbellata. K.S. (<i>wh.</i>) umbel-flower'd.	—	—	—	G. h.	— 3. t. 202.
10 fruticosa. R.S. (<i>wh.</i>) shrubby.	Canaries.	1779.	6. 10.	G. h.	Bot. reg. t. 464.
<i>Messerschmidia fruticosa.</i> w.					
11 angustifolia. R.S. (<i>pa. bl.</i>) narrow-leav'd.	—	—	—	G. h.	
12 argentea. R.S. (<i>wh.</i>) silvery-leaved.	E. Indies.	1822.	—	S. h.	Rumph. amb. 4. t. 55.
13 Arguzia. R.S. (<i>wh.</i>) herbaceous.	Siberia.	1780.	—	H. 4.	Gmel. it. 2. t. 27.
MESSERSCHMIDIA. R.S. MESSERSCHMIDIA. Pentandria Monogynia. R.S.					
1 caracasana. K.S. (<i>wh.</i>) Caracas.	Caracas.	1828.	5. 9.	S. h.	
2 hirsutissima. R.S. (<i>wh.</i>) very hairy.	S. America.	1818.	5. 8.	S. h.	Sloan. hist. 2. t. 212. f.1.
<i>Tournefortia hirsutissima.</i> w.					
3 volubilis. R.S. (<i>gr.</i>) climbing.	Jamaica.	1752.	7. 8.	S. h.	Sloan. hist. 1. t. 143. f.2.
<i>Tournefortia volubilis.</i> w.					
4 scandens. R.S. (<i>gr.</i>) scandent.	S. America.	1816.	—	S. h.	Flor. peruv. 2. t. 148.
5 laurifolia. R.S. (<i>st.</i>) Laurel-leaved.	W. Indies.	1823.	—	S. h.	Venten. choix. t. 2.
BEURR'ERIA. G. BEURR'ERIA. Pentandria Monogynia. R.S.					
1 succulenta. s.s. (<i>wh.</i>) oval-leaved.	W. Indies.	1758.	5. 7.	S. h.	Jacq. obs. 2. t. 26.
2 exsucca. s.s. (<i>wh.</i>) dry-fruited.	—	1804.	—	S. h.	Jacq. amer. t. 173. f. 17.
EHR'ETIA. S.S. EHR'ETIA. Pentandria Monogynia. L.					
1 serrata. F.I. (<i>wh.</i>) saw-leaved.	Nepaul.	1812.	6. 9.	G. h.	
2 tinfolia. s.s. (<i>wh.</i>) Tinus-leaved.	Jamaica.	1734.	6. 7.	S. h.	Trew Ehret. t. 24.
3 acuminata. B.P. (<i>wh.</i>) acuminate.	N. S. W.	1820.	—	G. h.	
4 laxa. s.s. (<i>wh.gr.</i>) loose-flowered.	Mauritius.	1821.	—	S. h.	Jacq. schœn. 1. t. 41.
5 internodis. s.s. (<i>wh.</i>) axillary-flower'd.	—	—	—	S. h.	L'Herit. stirp. 1. t. 24.
6 lœvis. s.s. (<i>wh.</i>) smooth.	E. Indies.	1816.	—	S. h.	Roxb. corom. t. 56.
7 buxifolia. s.s. (<i>wh.</i>) Box-leaved.	—	—	—	S. h.	— 1. t. 57.
8 microphylla. s.s. (<i>wh.</i>) small-leaved.	Nepaul.	1818.	6. 8.	S. h.	Pluk. phyt. t. 31. f. 1.
9 áspera. s.s. (<i>wh.</i>) rough-leaved.	E. Indies.	1795.	—	S. h.	Roxb. corom. 1. t. 55.
10 divaricata. DC. (<i>wh.</i>) divaricate.	Havannah.	1820.	—	S. h.	
CO'R'DIA. B.P. CO'R'DIA. Pentandria Monogynia. L.					
1 latifolia. F.I. (<i>wh.</i>) broad-leaved.	E. Indies.	1824.	6. 8.	S. h.	Rumph. amb. 2. t. 75.
2 Myxa. R.S. (<i>wh.</i>) smooth-leaved.	—	1640.	—	S. h.	Rheed. mal. 4. t. 37.
3 monóica. R.S. (<i>wh.</i>) Birch-leaved.	—	1799.	3. 4.	S. h.	Roxb. corom. 1. t. 58.
4 Collococca. R.S. (<i>gr.</i>) long-leaved.	Jamaica.	1759.	—	S. h.	Sloan. hist. 2. t. 203. f.2.
5 micrantha. R.S. (<i>gr.</i>) small-flowered.	—	1822.	—	S. h.	
6 serrata. R.S. (<i>wh.</i>) saw-leaved.	E. Indies.	1824.	—	S. h.	
7 tetraphylla. R.S. (<i>wh.</i>) four-leaved.	Trinidad.	1823.	4. 7.	S. h.	Aubl. guian. 1. t. 88.
8 gerasanthoides. K.S. (<i>w.</i>) taper-leaved.	Cuba.	1824.	—	S. h.	
9 Geraschanthus. R.S. (<i>w.</i>) Spanish Elm.	W. Indies.	1789.	5.	S. h.	Jacq. amer. t. 175. f.16.
10 grandid. F.I. (<i>wh.</i>) great.	E. Indies.	1818.	—	S. h.	
11 nodosa. R.S. (<i>wh.</i>) hairy.	Guiana.	1803.	6. 7.	S. h.	Aub. gui. 1. t. 86.
<i>hirsuta.</i> w. <i>Collococcus.</i> Aub. non L.					
12 flavescens. R.S. (<i>wh.</i>) yellow-fruited.	—	1818.	—	S. h.	Aub. gui. 1. t. 89.
13 spinescens. R.S. (<i>wh.</i>) spiny.	E. Indies.	—	—	S. h.	
14 macrophylla. R.S. (<i>wh.</i>) broad-leaved.	W. Indies.	1752.	—	S. h.	Sloan. hist. 2. t. 221. f.1.
15 campanulata. F.I. (<i>ye.</i>) bell-flowered.	E. Indies.	1824.	—	S. h.	Rumph. amb. 2. t. 75.
16 Sebastena. R.S. (<i>or.</i>) rough-leaved.	W. Indies.	1728.	7. 8.	S. h.	Bot. mag. t. 794.
17 lœvis. R.S. (<i>bh.</i>) smooth-leaved.	Caracas.	1816.	4. 8.	S. h.	Jacq. schœn. 1. t. 40.
18 reticulata. R.S. (<i>wh.</i>) netted-leaved.	E. Indies.	1820.	—	S. h.	
<i>angustifolia.</i> F.I.					
19 polygamia. F.I. (<i>wh.</i>) polygamous.	—	1812.	6. 8.	S. h.	
20 dentata. R.S. (<i>wh.</i>) toothed.	Caracas.	1823.	—	S. h.	
21 elliptica. R.S. (<i>wh.</i>) elliptic-leaved.	W. Indies.	1804.	6. 8.	S. h.	
22 dichotoma. R.S. (<i>wh.</i>) forked.	N. Holland.	1824.	—	S. h.	
23 scaberrima. K.S. (<i>wh.</i>) rough.	Peru.	—	—	S. h.	
24 angustifolia. R.S. (<i>wh.</i>) narrow-leaved.	Santa Cruz.	1808.	6. 8.	S. h.	
25 curassavica. R.S. (<i>wh.</i>) long-spiked.	S. America.	1759.	—	S. h.	Centur. amer. t. 56.
<i>Varronia curassavica.</i> w.					
26 martinicensis. R.S. (<i>wh.</i>) Martinico.	Martinico.	1795.	8. 9.	S. h.	Jacq. amer. 41. t. 32.
27 bullata. R.S. (<i>wh.</i>) blister-leaved.	W. Indies.	1823.	—	S. h.	Jacq. amer. pict. t. 43.
28 monosperma. R.S. (<i>wh.</i>) single-seeded.	Caracas.	1820.	5. 7.	S. h.	Jacq. schœn. 1. t. 39.
29 lineata. R.S. (<i>wh.</i>) round-spiked.	W. Indies.	1793.	6. 9.	S. h.	Brown. jam. t. 13. f. 2.
<i>Varronia lineata.</i> w.					

30 globosa. R.S. (wh.) globe-headed. Cuba. 1820.	7. 8. S. h.	Sloan, his. 2. t. 194. f. 2.
31 mirabiloides. R.S. (wh.) jointed. Hispaniola. 1798.	9. S. h.	Jacq. amer. 41. t. 33.
32 álba. R.S. (wh.) white-flowered. Carthagena. 1818. S. h.	Comm. hort. 1. t. 80.
33 bifurcata. R.S. (wh.) forked. Peru. — S. h.	Flor. peruv. 2.t.146.f.a.
Varronia dichotoma. R.P. parviflora. P.S.		
PATAG'ONULA. R.S. PATAG'ONULA. Pentandria Monogynia. R.S.		
americana. R.S. (wh.) spear-leaved. S. America. 1732.	6. 8. G. h.	Dill. elth. t. 226. f. 293.
Cordia Patagonula. W.		

ORDO CL.

HYDROPHYLLEÆ. Brown.

HYDROPHYLLUM. L. WATER-LEAF. Pentandria Monogynia. L.		
1 virginicum. L. (wh.bl.) Virginian. N. America. 1739.	5. 6. H. 4.	Bot. reg. t. 331.
2 canadense. L. (wh.) Canadian. Canada. 1759.	— H. 4.	— t. 242.
PHAC'ELIA. B.P. PHAC'ELIA. Pentandria Monogynia. R.S.		
1 bipinnatifida. M. (bl.) bipinnatifid. N. America. 1827.	5. 6. H. 4.	Mich. amer. 1. t. 16.
2 circinnata. J.E. (wh.) woolly. Magellan. 1816.	6. 10. H. 4.	Jacq. f. ecl. t. 91.
Hydrophyllum magellanicum. Lam. Aldæa circinnata. W.en.		
3 Aldæa. B.P. (wh.) various-leaved. Peru. 1820.	— G. 4.	Flor. peruv. 2. t. 14.
Aldæa pinnata. R.P.		
NEMO'PHILA. N. NEMO'PHILA. Pentandria Monogynia. S.S.		
1 phacelioides. N. (bl.) Phacelia-like. N. America. 1822.	6. 9. H. ⊖.	Swt. br. fl. gar. t. 32.
2 paniculata. s.s.(pa.bl.) paniced. — 1812.	5. 6. H. 4.	
Hydrophyllum appendiculatum. M.		
ELL'ISIA. R.S. ELL'ISIA. Pentandria Monogynia. W.		
1 Nyctelæa. R.S. (wh.) cut-leaved. Virginia. 1755.	4. 8. H. ⊖.	Trew pl. select. t. 99.
2 ambigua. N. (wh.) ambiguous. N. America. 1825.	4. 6. H. ⊖.	
EUTOCA. B. E'UTOCA. Pentandria Monogynia.		
1 multiflora. B.R. (pu.) many-flowered. N. America. 1826.	5. 6. H. ⊖.	Bot. reg. t. 1180.
2 Franklinii. s.s. (pu.) Franklin's. Mississippi. 1827.	— H. ⊖.	

ORDO CLI.

SOLANEÆ. Brown prodr. 1. p. 443.

Sect. I. PERICARPIUM CAPSULARE.

ANTHOCE'RCIS. B.P. ANTHOCE'RCIS. Didynamia Angiospermia. S.S.		
1 littorea. B.P. (ye.) yellow. N. Holland. 1803.	5. 8. G. h.	Swt. flor. aust. t. 17.
2 albicans. F.T. (wh.) white-flowered. — 1823.	4. 6. G. h.	— t. 16.
3 viscosa. B.P. (wh.) viscous. — — — G. h.		Bot. mag. t. 2961.
4 ilicifolia. B.M. (wh.) Holly-leaved. — 1829. G. h.	
CE'LSIA. S.S. CE'LSIA. Didynamia Angiospermia.		
1 orientalis. s.s. (ye.) oriental. Levant. 1713.	7. 8. H. ⊖.	Lam. ill. t. 532.
2 Arcturus. w. (ye.) scollop-leaved. Candia. 1780.	7. 9. F. ♂.	Bot. mag. t. 1962.
3 coronandelina. w. (ye.) Coronandel. E. Indies. 1783.	7. 8. S. ⊖.	Bot. mag. t. 964.
4 crética. s.s. (ye.pu.) great-flowered. Crete. 1752.	7. 9. F. ♂.	
5 viscosa. s.s. (ye.) clammy. 1816.	— F. ⊖.	
6 betonicæfolia. s.s.(ye.) Betony-leaved. Barbary. 1824.	— F. ♂.	
7 sublanata. B.R. (ye.) woolly. 1818.	— H. 4.	
8 lanceolata. s.s. (ye.) spear-leaved. Levant. 1816.	— F. ♂.	
VERBA'SCUM. L. MULLEIN. Pentandria Monogynia. L.		
1 Thapsus. R.S. (st.) Shepherd's club. Britain.	7. 8. H. ♂.	Eng. bot. v. 8. t. 549.
2 elongatum. W.en. (wh.) long-spiked. Europe. 1813.	— H. ♂.	Tabern. Kraeut. 956. ic.
3 thapsoides. R.S. (ye.) branching. Portugal.	— H. ♂.	Schrad. verb. t. 5. f. 2.
4 thapsiforme. R.S. (ye.) simple-stalked. Europe.	— H. ♂.	
5 indicum. Wal. (ye.) Indian. Nepaul. 1827.	— F. ♂.	
6 crassifolium. R.S. (ye.) thick-leaved. S. Europe. 1816.	— H. ♂.	
7 cuspidatum. R.S. (ye.) sharp-pointed. Austria. —	— H. ♂.	H. et Lk. fl. port. t. 26.
		Schrad. verb. t. 1. f. 1.

8 gossypinum. M.B. (ye.) cottony.	Iberia.	1824.	6. 8. H. ♂.	Ten. fl. neap. t. 22.
9 níveum. R.S. (ye.) snowy.	Naples.	1825.	— H. ♂.	
10 candidissimum. DC. (ye.) very white-l'd.	S. Europe.	1825.	— H. ♂.	
11 densiflòrum. R.S. (ye.) dense-flowered.	Italy.	—	— H. ♂.	
12 macránthum. R.S. (ye.) large-flowered.	Portugal.	1823.	— H. ♂.	H. et Lk. fl. port. t. 27.
13 austrâle. R.S. (ye.) southern.	S. Europe.	1816.	— H. ♂.	Schrad. verb. t. 2.
14 phlomoides. R.S. (ye.) woolly.	Turkey.	1739.	— H. ♂.	Flor. græc. t. 224.
15 rugulósum. w. (ye.) rugged-leaved.	S. Europe.	1820.	— H. ♂.	
16 conocárpon. s.s. (ye.) conical-capsuled.	Sardinia.	1829.	— H. ♂.	
17 condensátum. R.S. (ye.) close-flowered.	Austria.	1818.	— H. ♂.	Schrad. verb. t. 3.
18 nemorósium. R.S. (ye.) wood.	—	—	— H. ♂.	— t. 1. f. 2.
19 montánum. R.S. (st.) mountain.	Switzerland.	1816.	— H. ♂.	
20 collinum. R.S. (ye.) hill.	Germany.	1820.	— H. ♂.	Schrad. verb. t. 5. f. 1.
21 versiflòrum. R.S. (br.re.) scatter'd-flower'd.	—	1815.	— H. ♂.	
22 ramigerum. R.S. (ye.) small-branched.	—	1824.	— H. ♂.	Schrad. verb. t. 4.
23 Bastárdi. R.S. (ye.) Bastard's.	S. France.	—	— H. ♂.	
24 auriculátum. F.G. (ye.) eared-leaved.	Levant.	1826.	— E. ♂.	Flor. græc. t. 225.
25 sinuátum. R.S. (ye.) scollop-leaved.	S. Europe.	1570.	7. 8. H. ♂.	— t. 227.
26 undulátum. M.B. (ye.) waved-leaved.	Caucasus.	1816.	— H. ♂.	
27 plicátum. F.G. (ye.) plaited-leaved.	Greece.	—	— F. 4.	Flor. græc. t. 226.
28 bipinnatífidum. B.M. (ye.) bipinnatifid.	Tauria.	1813.	— H. ♂.	Bot. mag. t. 1777.
29 pinnatífidum. R.S. (ye.) pinnatifid.	Archipelago.	1788.	5. 6. G. 4.	Flor. græc. t. 228.
30 Chaíxi. R.S. (ye.) French.	France.	1816.	7. 8. H. ♂.	Vill. delph. 2. t. 13.
31 Boerhaavii. R.S. (ye.) annual.	S. Europe.	1731.	— H. ♂.	Mill. ic. 2. t. 273.
32 compáctum. R.S. (ye.) compact-flower'd.	Tauria.	1824.	— H. ♂.	
33 gnaphalòdes. M.B. (ye.) Gnaphalium-like.	—	—	— H. ♂.	
34 ovalifólium. B.M. (ye.) oval-leaved.	Caucasus.	1804.	7. 9. H. 4.	Bot. mag. t. 1037.
35 formosum. B.R. (ye. pu.) beautiful.	Russia.	1814.	7. 8. H. ♂.	Bot. reg. t. 558.
36 Lychnitis. R.S. (wh.) white.	Britain.	6. 8. H. ♂.	Eng. bot. v. 1. t. 58.
37 austriacum. R.S. (st.) Austrian.	Austria.	1819.	— H. ♂.	
38 pulveréntum. R.S. (ye.) powdered.	England.	— H. ♂.	Eng. bot. v. 7. t. 487.
39 majále. DC. (ye.) purple-stalked.	Montpelier.	1822.	5. 7. H. ♂.	
40 Stevénii. F. (ye.) Steven's.	Siberia.	1820.	6. 8. H. ♂.	
41 fasciculátum. E. (ye.) clustered.	Mount Sinai.	1826.	5. 8. F. 4.	
42 æthiòpicum. E. (ye.) Ethiopian.	—	1825.	— F. 4.	
43 pyramidátum. R.S. (ye.) pyramidal.	Caucasus.	1804.	7. 8. H. 4.	Swt. br. fl. gar. t. 31.
44 ovátum. R.S. (ye.) oval-leaved.	Spain.	—	— H. ♂.	Schrad. h. gott. t. 15.
45 rotundifólium. R.S. (ye.) round-leaved.	Naples.	1826.	— H. ♂.	Ten. fl. neap. t. 23.
46 floccosum. R.S. (ye.) wool-bearing.	Hungary.	1805.	6. 7. H. ♂.	W. et K. hung. 1. t. 79.
47 leptostächyum. R.S. (ye.) slender-spiked.	S. France.	1824.	7. 9. H. ♂.	
48 angustifólium. R.S. (ye.) narrow-leaved.	Naples.	1822.	— H. ♂.	
49 longifólium. R.S. (ye.) long-leaved.	—	—	— H. ♂.	Ten. fl. neap. t. 21.
50 nigrum. R.S. (ye.) black-rooted.	England.	6. 8. H. 4.	Eng. bot. v. 1. t. 59.
51 orientâle. R.S. (ye.) oriental.	Caucasus.	1820.	— H. 4.	
52 Alopecúrus. R.S. (ye.) fox-tail.	France.	1818.	— H. 4.	
53 virgátum. R.S. (ye.) slender.	Britain.	8. H. ♂.	Eng. bot. v. 8. t. 550.
54 speciosum. R.S. (ye.) splendid.	Austria.	1820.	6. 8. H. 4.	Schrad. h. gott. t. 16.
55 grandiflòrum. R.S. (ye.) great-flowered.	—	1816.	— H. 4.	— t. 15.
56 monspessulânum. s.s. (ye.) Montpelier.	S. France.	1824.	— H. ♂.	Schrad. ver.p.2.t.2.f.2.
57 lanátum. s.s. (ye.) woolly-leaved.	S. Europe.	1828.	— H. ♂.	— pars. 2. t. 2. f. 1.
58 hyosérifólium. s.s. (ye.) Hyoseris-leav'd.	Levant.	1829.	— F. 4.	— pars. 2. t. 3. f. 1.
59 chrysýnum. s.s. (ye.) golden.	Palestine.	1827.	— F. 4.	
60 Schottiânum. s.s. (ye.) Schott's.	Austria.	1828.	— H. ♂.	Schrad. ver.p.2.t.3.f.2.
61 Blattâria. R.S. (ye.) yellow moth.	Britain.	7. 9. H. ♂.	Eng. bot. v. 6. t. 393.
62 glábrum. Mil. (wh.) white moth.	—	—	— H. ♂.	
63 blattarioides. R.S. (ye.) moth-like.	S. Europe.	1816.	— H. ♂.	
64 bannáticum. s.s. (ye.) Hungarian.	Hungary.	1820.	— H. ♂.	
65 repándum. R.S. (ye.) waved.	—	1813.	7. 8. H. ♂.	
66 hæmorrhoidâle. R.S. (ye.) Madeira.	Madeira.	1777.	6. 8. G. ♂.	Bot. mag. t. 1226.
67 cùpreum. R.S. (co.) copper-colour'd. Hybrid.	—	1798.	5. 9. H. 4.	— t. 885.
68 phœnicium. R.S. (d. pu.) purple-flower'd. S. Europe.	—	1596.	5. 8. H. 4.	Schrad. h. gott. t. 14.
69 puníceum. R.S. (pi.) pink-flower'd.	—	1815.	— H. ♂.	W. et K. hung. 2. t. 197.
70 rubiginósum. R.S. (re. br.) brown-flower'd. Hungary.	—	—	— H. ♂.	Trew Ehret. t. 16. f. 1.
71 ferrugineum. R.S. (br.) rusty.	S. Europe.	1633.	— H. 4.	Andr. deposit. t. 162.
72 triste. R.S. (da. br.) dark-flowered.	Levant.	1788.	— H. 4.	
73 ferrugineum. A.R. nec alicorū.	—	—	— H. ♂.	
73 spectabilis. M.B. (ye.) showy.	Tauria.	1822.	6. 8. H. ♂.	
74 betonicæfólium. R.S. (ye.) Betony-leav'd.	Levant.	1828.	— F. ♂.	Desf. ann. mus. 11. t. 4.
75 urticæfólium. R.S. (ye. re.) spotted-flower'd.	—	1827.	— H. ♂.	

SOLANEÆ.

76 lyratūm. R.S. (ye.) lyrate-leaved.	Spain.	1816.	6. 8. H. ♂.	
77 Osbeckii. R.S. (gr.ye.) Osbeck's.	1752.	7. 8. H. ♂.	Buxb. cent. 5. t. 32.	
78 ceratophyllum.s.s.(ye.)buckshorn-leav'd.	Levant.	1829.	— H. ♂.	Schrab. verb.p.2.t.1.f.2.
79 spinosum. R.S. (ye.) spiny.	Candia.	1824.	5. 8. G. h.	Flor. græc. t. 229.
RAMONDA. P.S. RAMO'nda.	Pentandria Monogynia. P.S.			
pyrenaica. P.S. (pu.) Borage-leaved.	Pyrenees.	1731.	5. H. ♀.	Bot. mag. t. 236.
Vérbascum Myconi. B.M.				
HYOSC'YAMUS. R.S. HENBANE.	Pentandria Monogynia. L.			
1 niger. R.S. (ye.bk.) common.	Britain.	...	6. 7. H. ♂.	Eng. bot. v. 9. t. 591.
2 agréstis. R.S. (gr.pu.) field.	Hungary.	1816.	6. 9. H. ⓪.	Swt. br. fl. gar. t. 27.
3 pálidus. R.S. (st.) pale-flowered.	—	1815.	— H. ⓪.	
4 reticulátus. R.S. (re.) Egyptian.	Egypt.	1640.	7. 9. H. ⓪.	Camer. hort. 77. t. 22.
5 ábus. R.S. (st.pu.) white.	Greece.	1570.	7. 8. H. ⓪.	Flor. græc. t. 230.
6 Clusi. F.G. (wh.) Clusius's.	S. Europe.	—	— H. ⓪.	Clus. hist. 2. p. 84. f. 1.
ábus β minor. R.S.				
7 àureus. R.S. (ye.pu.) golden.	Levant.	1640.	3. 10. G. h.	Bot. mag. t. 87.
8 Seneciônis. R.S. (ye.pu.) yellow-flowered.	Egypt.	1812.	— G. ♀.	
9 canariénis. B.R. (ye.pu.) various-leaved.	Canaries.	1816.	1. 12. G. ♂.	Bot. reg. t. 180.
10 müticus. R.S. (wh.pu.) three-coloured.	Egypt.	1818.	5. 8. G. ♂.	
11 Datôra. F.E. (ye.pu.) Egyptian.	—	1829.	— F. ♀.	
12 pusillus. R.S. (ye.bk.) dwarf.	Persia.	1691.	7. H. ⓪.	Pluk. alm. t. 37. f. 5.
13 auriculátus. T.N. (ye.pu.) eared.	Naples.	1828.	— H. ⓪.	
14 micránthus. Led. (st.pu.) small-flowered.	Siberia.	1829.	— H. ♂.	
15 májor. Mil. (st.pu.) large.	Levant.	...	— F. ♂.	
16 minor. Mil. (ye.) lesser.	—	...	— F. ♂.	
17 physaloides. R.S. (d.pu.) purple-flowered.	Siberia.	1777.	3. 4. H. ♀.	Swt. br. fl. gar. t. 13.
18 orientális. R.S. (li.) oriental.	Iberia.	1815.	— H. ♀.	— t. 12.
SCOP'OLIA. Jacq.	Scop'olia.	Pentandria Monogynia. R.S.		Jacq. obs. 1. t. 20.
carniolica. Jacq.	Nightshade-leav'd. Carniola.	1780.	4. 5. H. ♀.	
atropoides. R.S. <i>Hyosc'ymus Scop'olia</i> .	Bot. mag. t. 1126.			
NICOTTANA. R.S. TOBACCO.	Pentandria Monogynia. L.			
1 chinénis. R.S. (ro.) Chinese.	China.	1815.	6. 8. G. h.	Mill. icon. t. 185. f. 1.
2 fruticosa. R.S. (ro.) frutescent.	C. B. S.	1699.	7. 9. G. h.	
3 álba. Mil. (wh.) white-flowered.	Tobago.	—	8. 10. H. ⓪.	
4 latíssima. DC. (ro.) broad-leaved.	America.	...	7. 9. H. ⓪.	
macrophylla. R.S.				
5 Tabácum. R.S. (ro.) Virginian.	—	1570.	— H. ⓪.	Blackw. t. 146.
α attenuata. (re.) slender.	—	...	— H. ⓪.	
β palléscens. (bh.) pale-flowered.	—	...	— H. ⓪.	
γ álipes. (ro.) winged.	—	...	— H. ⓪.	
δ serotina. (ro.) late-flowered.	—	...	— H. ⓪.	
ε grátilipes. (ro.) slender-stalked.	—	...	— H. ⓪.	
ζ Vérda. (ro.) Verda.	—	...	— H. ⓪.	
η lingua. (ro.) tongue-leaved.	—	...	— H. ⓪.	
6 nepalénis. Ot. (ro.) Nepaul.	Nepaul.	1829.	— H. ⓪.	
7 brasiliénis. Ot. (ro.) Brazilian.	Brazil.	1825.	— H. ⓪.	
8 decúrrens. Ag. (ro.) decurrent.	S. America.	1820.	— H. ⓪.	
9 petiolata. Ag. (ro.) petiolate.	—	—	— H. ⓪.	
10 sanguínea. Ot. (sc.) scarlet-flowered.	Brazil.	1829.	— H. ⓪.	
11 angustifólia. R.S. (gr.) narrow-leaved.	Chili.	1822.	— H. ⓪.	Flor. peruv. 2.t.130.f.a.
12 viscósa. R.S. (gr.) viscous.	Buenos Ayres.	1824.	— H. ⓪.	
13 pusilla. R.S. (gr.) Primrose-leaved.	Vera Cruz.	1733.	8. S. ♂.	Mill. ic. t. 185. f. 2.
14 glutinosa. R.S. (or.) clammy.	Peru.	1759.	7. 9. H. ⓪.	Swt. br. fl. gar. t. 107.
15 rústica. R.S. (gr.) common.	America.	1570.	— H. ⓪.	Blackw. t. 437.
α asiática. R.S. (gr.) Asiatic.	—	—	— H. ⓪.	
β brasiliá. R.S. (gr.) Brazil.	Brazil.	—	— H. ⓪.	
γ pùmila. R.S. (gr.) dwarf.	S. America.	—	— H. ⓪.	
16 húmilis. L.en. (gr.) lesser.	America.	—	— H. ⓪.	
17 paniculata. R.S. (gr.) paniced.	Peru.	1752.	— H. ⓪.	Flor. peruv. 2.t.129.f.b.
18 cerinthoídes. R.S. (gr.) honeywort.	America.	1816.	— H. ⓪.	Lehm. nicot. t. 2.
19 Langsdórfii. s.s. (gr.ye.) Langsdorff's.	S. America.	1824.	— H. ⓪.	Bot. mag. t. 2555.
20 glauca. B.M. (st.) glaucous.	Buenos Ayres.	1827.	— H. ⓪.	— t. 2837.
21 repánda. R.S. (bh.) stem-clasping.	Havannah.	1822.	8. 10. H. ⓪.	— t. 2484.
lyrata. K.S.				
22 plumbaginifólia. R.S. curled-leaved.	America.	1816.	5. 10. G. ♀.	Vivian. elench. t. 5.
crispá. Jacq. fragm. t.84.				
23 pulmonarioídes. K.S. (wh.) Pulmonaria-like.	S. America.	1828.	— H. ⓪.	
24 bonariénis. R.S. (wh.) Buenos Ayres.	Buenos Ayres.	1821.	— H. ⓪.	
25 suaveolens. R.S. (wh.) sweet-scented.	N. S. W.	1800.	5. 9. G. ♀.	Lehm. nicot. n. 8. t. 1.
undulata. Vent. malm. t. 10. non R.P.				Bot. mag. t. 673.

26	vincæflòra. s.s.	(wh.) Vinca-flowered. S. America.	1825.	8. 10. H.○.		
27	dilatàta. L.en.	(wh.) dilated-leaved.	—	—	H.○.	
28	noctiflòra. B.M.	(wh.) night-flowering. Andes.	1826.	7. 10. F.‡.		Swt. br. fl. gar. t. 262.
29	acuminàta. B.M.	(wh.) taper-pointed. BuenosAyres.	—	—	F.‡.	Bot. mag. t. 2919.
30	nàna. B.R.	(wh.) dwarf.	S. America.	1822.	8. 10. H.○.	Bot. reg. t. 833.
31	quadriùalvis. R.S.	(wh.) four-valved.	N. America.	1811.	7. 8. H.○.	Bot. mag. t. 1778.
32	multiválvis. B.R.	(wh.) many-valved.	—	1826.	9. 10. H.○.	Bot. reg. t. 1057.
PET'UNIA. J.	PET'UNIA.	Pentandria Monogynia. R.S.				
nyctaginiflòra. J.(wh.)	large-flowered.	S. America.	1824.	4. 10. G.‡.		Swt. br. fl. gar. t. 119.
SALPIGLO'SSIS. R.P.	SALPIGLO'SSIS.	Didynamia Angiospermia.				
1	stramínea. H.E.F.(st.)	straw-coloured. Chile.	1826.	6. 9. F.‡.		Swt. br. fl. gar. t. 231.
2	pícta. B.F.G.	(wh.bl.) painted-flowered.	—	—	F.‡.	— v. 3. t. 258.
3	atropurpurea. B.M.(d.pu.)	dark purple.	—	—	F.‡.	— v. 3. t. 271.
4	sinuàta. R.P.	(cr.) crimson.	—	1829.	F.‡.	Flor.per.v.5.t.515.ined.
DAT'URA. R.S.	THORN-APPLE.	Pentandria Monogynia. L.				
1	férox. R.S.	(wh.) Chinese.	China.	1731.	7. 10. H.○.	Zanon. hist. t. 162.
2	Stramoniùm. R.S.(wh.)	common.	England.	—	H.○.	Eng. bot. v. 18. t. 1288.
	β canéscens. F.I.	(st.) canescens.	Nepaul.	1824.	H.○.	
3	Tátula. R.S.	(bl.) light blue.	S. America.	1629.	H.○.	Swt. br. fl. gar. t. 83.
4	muricàta. S.S.	(wh.) muricated.	—	1819.	H.○.	
5	hybrida. T.P.	(wh.) hybrid.	Hybrid.	1826.	6. 9. F.‡.	
6	Métel. R.S.	(wh.) downy.	Asia.	1596.	6. 10. H.○.	Bot. mag. t. 1440.
7	fruticòsa. L.en.	(wh.) frutescent.	S. America.	1825.	S.‡.	
8	quercifòlia. K.S.	(li.) Oak-leaved.	Mexico.	1824.	H.○.	Rumph. am.5.t.243.f.2.
9	fastuòsa. R.S. (pu.wh.)	purple.	Egypt.	1629.	H.○.	
	β florè pleno. (pu.wh.)	double-flowered.	—	—	G.○.	
10	dùbia. R.S.	(wh.) doubtful.	—	—	H.○.	
11	l'ævis. R.S.	(wh.) smooth-fruited.	Africa.	1780.	H.○.	Jacq. vind. 3. t. 82.
12	ceratocauña. R.S.(wh.pu.)	horn-stalked.	Cuba.	1805.	8. 10. H.○.	Jacq. schoen. 3. t.339.
13	guayaquilénis. K.S.(wh.)	Guayaquil.	S. America.	1824.	H.○.	
BRUGMA'NSIA. P.S.	BRUGMA'NSIA.	Pentandria Monogynia. R.S.				
1	cándida. P.S.	(wh.) downy-stalked.	Peru.	1818.	8. 9. S.‡.	Flor. peruv. 2. t. 128.
	Datura arbòrea. L.					
2	suavèolens.	(wh.) smooth-stalked.	S. America.	1733.	— S.‡.	
	Datura arbòrea. Hort. suavèolens. W.enum.					
VE'STIA. W.	VESTIA.	Pentandria Monogynia. W.				
lycioïdes. W.en.(gr.ye.)	Boxthorn-like. Chili.	1815.	1. 12. G.‡.		Bot. reg. t. 299.	
? CODON. W.	CODON.	Decandria Monogynia. L.				
Royèni. w.	(re.wh.) prickly.	C. B. S.	1801.	9. 10. G.♂.	And. repos. t. 325.	

Sect. II. PERICARPIUM BACCATUM.

ANIS'ODUS. S.S.	ANIS'ODUS.	Pentandria Monogynia. S.S.				
lùridus. s.s.	(br.) lurid.	Nepaul.	1823.	8. 10. H.‡.	Swt. br. fl. gar. t. 125.	
Whitlèya stramonifolia B.F.G.						
NECTO'UXIA. K.S.	NECTO'UXIA.	Pentandria Monogynia.				
formòsa. K.S.	(ye.) pretty.	Mexico.	1826.	7. 10. H.○.	H.etB.n.g.am.3.t.193.	
ATROPA. R.S.	A'TROPA.	Pentandria Monogynia. L.				
Belladónna. r.s. (br.pu.)	deadly nightshade.	Britain.	—	6. 7. H.‡.	Eng. bot. v. 9. t. 592.	
SA'RACHA. R.S.	SA'RACHA.	Pentandria Monogynia.				
1 procumbens. R.P. (st.)	procumbent.	Peru.	1822.	7. 9. H.‡.	Cavan. ic. 1. t. 72.	
2 umbellata. DC.	(st.) umbel-flowered.	Peru.	—	H.‡.	Swt. br. fl. gar. t. 85.	
MANDRAG'ORA. J.	MANDRAKE.	Pentandria Monogynia. R.S.				
1 pr'æcox. B.F.G. (st.bh.)	blister-leaved.	Switzerland.	1818.	3. 4. H.‡.	Swt. br. fl. gar. t. 198.	
2 vernális. s.s.	(bh.) large-leaved.	S. Europe.	1562.	5. 6. H.‡.		
3 autumnális. s.s.	(li.) autumn-flowered.	Levant.	—	8. 9. H.‡.	Flor. græc. t. 232.	
NOL'ANA. R.S.	NOL'ANA.	Pentandria Monogynia. L.				
1 prostrata. R.S.	(bl.) trailing.	Peru.	1761.	7. 9. H.○.	Bot. mag. t. 731.	
2 tenella. H.C.	(bl.) slender.	Chile.	1824.	F.‡.	Bot. reg. t. 865.	
3 paradoxa. B.R.	(bl.) violet-coloured.	Chili.	1822.	H.○.		
NICA'NDRA. R.S.	NICA'NDRA.	Pentandria Monogynia. R.S.				
1 physaloïdes. R.S.	(bl.) American.	America.	1759.	7. 9. H.○.	Bot. mag. t. 2458.	
2 indica. R.S.	(bl.) Indian.	Nepaul.	1823.	H.○.	Rumph.amb.6.t.26.f.1.	
PHY'SALIS. R.S.	WINTER CHERRY.	Pentandria Monogynia. L.				
arboréscens. w.	(ye.) arborescent.	Campeachy.	1733.	6. 8. S.‡.	Mill. ic. 2. t. 206. f. 2.	

SOLANEÆ.

2 frutéscens. DC. (<i>ye.</i>) frutescent.	Spain.	1737.	1. 3. G. h.	Cavan. ic. 2. t. 102.
<i>A'tropa frutéscens</i> . R.S.	Canaries.	1779.	5. 8. G. h.	
3 aristáta. w. (<i>ye.</i>) bearded.	Levant.	1596.	7. 8. G. h.	Flor. græc. t. 233.
<i>A'tropa aristáta</i> . R.S.	Mexico.	1826.	— G. h.	Jacq. f. ecl. t. 23.
4 somnifera. R.S. (<i>st.</i>) clustered.	E. Indies.	1759.	— G. h.	Pluk. alm. t. 111. f. 5.
β <i>aristáta</i> . (<i>st.</i>) awned.	S. America.	1699.	6. 9. S. 4.	
5 flexuosa. R.S. (<i>gr.wh.</i>) flexuose.	C. B. S.	1816.	— G. h.	Bot. mag. t. 2625.
6 curassávica. R.S. (<i>st.</i>) Curassavian.	America.	1732.	7. H. 4.	
7 tomentosa. R.S. (<i>ye.</i>) woolly.	N. America.	1780.	— H. 4.	Jacq. vind. 2. t. 136.
8 viscósa. R.S. (<i>st.</i>) clammy.	—	1726.	7. 9. H. 4.	
9 lanceoláta. R.S. (<i>ye.</i>) spear-leaved.	—	—	— H. 4.	
10 pensylvánica. R.S. (<i>ye.</i>) Pennsylvanian.	—	—	— H. 4.	
11 Jacquiní. L.en. (<i>ye.</i>) Jacquin's.	—	—	— H. 4.	
<i>viscósa</i> . Jacq. <i>non L.</i>	—	—	— H. 4.	
12 Alkekéngi. R.S. (<i>wh.</i>) common.	S. Europe.	1548.	— H. 4.	Flor. græc. t. 234.
13 tuberósa. R.S. (<i>ye.pu.</i>) tuberous-rooted.	—	1816.	— S. 4.	
14 chenopodífolia. R.S. (<i>ye.</i>) goose-foot-leav'd.	S. America.	1798.	7. 8. S. h.	
15 peruviana. R.S. (<i>ye.pu.</i>) Peruvian.	Peru.	1773.	— S. 4.	
16 édulis. B.M. (<i>ye.pu.</i>) eatable.	S. America.	—	— S. 4.	Bot. mag. t. 1068.
17 pubescens. R.S. (<i>ye.pu.</i>) pubescent.	America.	1640.	— H. 4.	Feuil. per. 3. t. 1.
18 micránta. L.en. (<i>ye.</i>) small-flowered.	—	1822.	— H. 4.	
19 nodosa. R.S. (<i>ye.</i>) knotted.	—	—	— H. 4.	
20 barbadensis. R.S. (<i>st.</i>) Barbadoes.	W. Indies.	1798.	— H. 4.	
21 dúbia. L.en. (<i>ye.</i>) dubious.	Brazil.	1819.	— H. 4.	Jacq. ic. 1. t. 39.
22 foetidissima. R.S. (<i>ye.br.</i>) stinking.	New Spain.	—	— H. 4.	
23 f'etens. R.S. (<i>ye.br.</i>) strong-scented.	Peru.	1827.	— H. 4.	
24 anguláta. R.S. (<i>wh.</i>) angular-branch'd.	India.	1732.	6. 9. H. 4.	Dill. elth. t. 12. f. 12.
25 Rothiana. R.S. (<i>wh.</i>) Roth's.	E. Indies.	1827.	— S. 4.	
26 philadéphica. R.S. (<i>ye.</i>) Philadelphian.	N. America.	1824.	— H. 4.	
27 atriplicifolia. R.S. (<i>ye.</i>) Atriplex-leaved.	—	1798.	7. 8. H. 4.	Jacq. frag. t. 85. f. a.
<i>chenopodífolia</i> . w. <i>non Lamarck.</i>	—	—	— H. 4.	
28 mímina. R.S. (<i>ye.</i>) least.	E. Indies.	1759.	— H. 4.	Rheed. mal. 10. t. 70.
29 pruinosa. R.S. (<i>ye.</i>) hairy annual.	America.	1726.	— H. 4.	Dill. elth. 10. t. 9. f. 9.
30 Lagascii. R.S. (<i>ye.</i>) Lagasca's.	New Spain.	1824.	— H. 4.	
31 páltula. R.S. (<i>wh.</i>) spreading.	Vera Cruz.	1731.	7. 9. S. 4.	
32 villósa. R.S. (<i>st.</i>) villous.	—	—	— S. 4.	
33 cordáta. R.S. (<i>ye.gr.</i>) heart-leaved.	—	—	— S. 4.	
34 máxima. R.S. (<i>st.</i>) large annual.	—	—	— S. 4.	
35 abyssinica. Ot. (<i>ye.</i>) Abyssinian.	Abyssinia.	1828.	— S. 4.	
36 prostrata. R.S. (<i>bl.</i>) trailing.	Peru.	1782.	— H. 4.	
CA'PSICUM. R.S.	CA'PSICUM.	Pentandria Monogynia. L.		Andr. deposit. t. 75.
1 ánnuum. R.S. (<i>wh.</i>) annual.	India.	1548.	6. 7. G. 4.	Rheed. mal. 2. t. 35.
2 lóngum. R.S. (<i>wh.</i>) long-fruited.	—	—	— G. 4.	Dodon. pemph. 716.f.3.
3 cordiforme. R.S. (<i>wh.</i>) heart-shaped.	India.	1731.	— G. 4.	
4 tetragónum. R.S. (<i>wh.</i>) four-sided.	—	—	— G. 4.	
5 angulósum. R.S. (<i>wh.</i>) angular-fruited.	—	—	— S. 4.	
6 sphæricum. R.S. (<i>wh.</i>) round-fruited.	—	1807.	4. 7. S. 4.	
7 ovátum. R.S. (<i>wh.</i>) oval-fruited.	—	—	— S. h.	
8 pén'dulum. R.S. (<i>wh.</i>) pendulous.	—	1804.	— S. h.	
9 lúteum. R.S. (<i>wh.</i>) yellow-fruited.	E. Indies.	1820.	— S. h.	
10 gróssum. R.S. (<i>wh.</i>) various-fruited.	India.	1759.	5. 8. S. h.	Besl. eyst. 2. t. 2. f. I.
α <i>globosum</i> . R.S. (<i>wh.</i>) globular-fruited.	—	—	— S. h.	
β <i>lúteum</i> . R.S. (<i>wh.</i>) yellow oval-fruited.	—	—	— S. h.	
γ <i>bifidum</i> . (<i>wh.</i>) cleft-fruited.	—	—	— S. h.	
11 conoides. R.S. (<i>st.</i>) conical-fruited.	India.	1728.	— S. h.	
12 pyramidale. R.S. (<i>wh.</i>) pyramidal.	Egypt.	1732.	— S. h.	
13 cerasifórme. R.S. (<i>st.</i>) Cherry-fruited.	W. Indies.	1759.	— S. h.	
14 Millérii. R.S. (<i>st.</i>) Miller's.	New Spain.	—	— S. h.	
<i>cerasifórme</i> . Mill. <i>non w.</i>	—	—	— S. h.	
15 purpureum. F.I. (<i>pu.</i>) purple.	E. Indies.	1824.	— S. h.	
16 frutéscens. R.S. (<i>wh.</i>) shrubby.	India.	1656.	6. 9. S. h.	Rheed. mal. 2. t. 56.
β <i>torulósum</i> . (<i>wh.</i>) twisted.	—	—	— S. h.	
17 bicolor. R.S. (<i>pu.</i>) purple-flower'd.	W. Indies.	1804.	— S. h.	Bot. mag. t. 1835.
<i>nigrum</i> . W.en. <i>violaceum</i> . DC.	—	—	— S. h.	
18 cæruléscens. R.S. (<i>pu.</i>) bluish.	—	1827.	— S. h.	
19 baccátum. R.S. (<i>wh.</i>) Bird pepper.	W. Indies.	1731.	— S. h.	Sloan. j. 1. t. 146. f. 2.
20 minali. F.I. (<i>wh.</i>) small-berried.	E. Indies.	1728.	— S. h.	
21 sinénsis. R.S. (<i>wh.</i>) oval Chinese.	China.	1807.	— S. h.	Jacq. vind. 3. t. 67.
22 havanénsis. K.S. (<i>wh.</i>) Havannah.	Havannah.	1826.	— S. h.	

23 microcarpum. R.S. (wh.) small-fruited.	S. America.	1824.	6. 9. S. h.	
24 micranthum. R.S. (wh.) small-flowered.	Brazil.	—	— S. h.	
LYCOPE'RSICUM. R.S. LOVE-APPLE.	Pentandria Monogynia. R.S.			
1 pimpinellifolium. R.S. (ye.) Pimpinella-lid.	S. America.	...	6. 8. S. y.	
2 regulare. R.S. (ye.) regular-leaved.	—	...	6. 9. G. (○)	
3 peruvianum. R.S. (ye.) Peruvian.	Peru.	...	6. 8. G. y.	Bot. mag. t. 2814.
<i>Solanum peruvianum</i> . Jacq. icon. t. 327.				
4 dentatum. R.S. (ye.) tooth-leaved.	S. America.	...	— H. (○)	Dunal solan. 2. ined. t. 82.
5 Humboldtii. R.S. (ye.) Humboldt's.	—	1815.	— H. (○)	Willd. hort. ber. 1. t. 27.
6 pyrifforme. R.S. (ye.) Pear-fruited.	—	...	— H. (○)	Dunal solan. t. 26.
7 cerasiforme. R.S. (ye.) Cherry-fruited.	Peru.	1800.	7. 9. H. (○)	Jacq. vind. 1. t. 11.
<i>Solanum pseudolycopersicum</i> . Jacq.				
α ruber. (ye.) red-fruited.	—	—	— H. (○)	
β luteum. (ye.) yellow-fruited.	—	—	— H. (○)	
8 esculentum. R.S. (ye.) Tomato.	S. America.	1596.	— H. (○)	Dunal solan. t. 3. f. 3.
α erythrocarpum. (ye.) red-fruited.	—	—	— H. (○)	
β chrysocarpum. (ye.) yellow-fruited.	—	—	— H. (○)	
γ leucocarpum. (ye.) white-fruited.	—	—	— H. (○)	
9 procumbens. R.S. (ye.) procumbent.	—	...	— H. (○)	
SOL'ANUM. R.S. NIGHTSHADE.	Pentandria Monogynia. L.			
1 tuberosum. R.S. (bl. wh.) Potatoe.	Peru.	1597.	6. 8. H. y.	Plenck. off. t. 121.
2 appendiculatum. R.S. (wh.) appendaged.	Mexico.	1823.	— S. h.	Dunal solan. ed. 2. t. 84.
3 Seaforthianum. R.S. (pi.) Seaforth's.	Barbadoes.	1804.	7. 9. S. h. (○)	Andr. deposit. t. 504.
4 betaceum. R.S. (bh.) Beet-leaved.	New Spain.	1803.	6. 7. S. h.	t. 511.
5 muricatum. R.S. (wh. vi.) warted.	Peru.	1785.	7. 8. S. h.	Feuil. peruv. 2. t. 26.
6 diversifolium. R.S. (wh.) different-leaved.	Caracas.	1824.	— S. h.	Dunal solan. ed. 2. t. 88.
7 Bulbocastanum. R.S. (wh.) tuberous-rooted.	Mexico.	1826.	6. 9. H. y.	— ed. 2. t. 3.
8 oliganthum. R.S. (wh.) climbing.	Orinoco.	1824.	7. 8. S. h. (○)	— ed. 2. t. 90.
9 reclinatum. R.S. (bl.) reclining.	Peru.	...	— S. h.	Lam. ill. t. 145. f. 4.
10 laciniatum. B.P. (ll.) cut-leaved.	N. Holland.	1772.	— G. h.	Bot. mag. t. 349.
α fruticosum. (bl.) frutescent.	N. S. W.	—	— G. h.	
β herbaceum. (bl.) herbaceous.	V. Diem. Isl.	—	— H. (○)	
11 quercifolium. R.S. (vi.) Oak-leaved.	Peru.	1787.	6. 7. H. y.	Feuil. per. 2. t. 15 ?
12 radicans. R.S. (li.) rooting.	—	1771.	7. 8. G. y.	Linn. dec. 1. t. 10.
13 corymbosum. R.S. (vi.) corymb-flowered.	—	1786.	— S. y.	Jacq. ic. 1. t. 40.
14 Dulcamara. R.S. (bl.) bitter-sweet.	Britain.	...	6. 8. H. h. (○)	Eng. bot. v. 8. t. 565.
β album. (wh.) white-flowered.	—	...	— H. h. (○)	
γ variegatum. (pa.) variegated.	—	...	— H. h. (○)	
15 littoralis. L.en. (bl.) sea-side.	Europe.	...	— H. h.	
16 Tegore. R.S. (gr.) hairy-stemmed.	Guiana.	1820.	— S. h.	Aubl. guian. 1. t. 84.
17 quitense. s.s. (wh. pu.) Lima.	Lima.	1825.	— G. h.	Bot. mag. t. 2739.
18 macrocarpon. s.s. (bl.) large-fruited.	Peru.	1759.	5. 9. S. h.	Mill. ic. 2. t. 294.
19 calycinum. R.S. (bl.) large-calyxed.	Mexico.	1825.	— S. h.	Dunal sol. ed. 2. t. 29.
20 racemiflorum. R.S. (bl.) racemed.	S. America.	1818.	7. 9. H. (○)	Jacq. schoen. 3. t. 333.
scabrum. Jacq. non Vahl.				
21 aethiopicum. R.S. (wh.) Ethiopian.	Ethiopia.	1597.	— H. (○)	Jacq. vind. 1. t. 12.
22 Zuccagnianum. R.S. (wh.) Zuccagni's.	—	1818.	— H. (○)	Dunal solan. t. 11.
23 pseudo-capsicum. (wh.) Winter-cherry.	Madeira.	1596.	6. 9. G. h.	Sabb. h. rom. t. 59.
24 nodiflorum. R.S. (wh.) knot-flowered.	Guiana.	1815.	— G. (○)	Jacq. ic. 2. t. 326.
25 Dilleniæ. R.S. (wh.) Dillenius's.	Hungary.	...	— H. (○)	Dill. elth. t. 275. f. 355.
26 guineense. R.S. (gr. wh.) large-berried.	Guinea.	...	— H. (○)	— t. 274. f. 354.
27 fistulosum. R.S. (wh.) fistulous-stalked.	Mauritius.	1826.	— H. (○)	
28 nigrum. R.S. (wh.) black-berried.	Britain.	...	— H. (○)	Eng. bot. v. 8. t. 566.
29 oleraceum. R.S. (wh.) cultivated.	W. Indies.	...	— S. (○)	Piso lib. 4. c. 50. f. 3.
30 pterocaule. R.S. (wh. vi.) wing-stalked.	America.	...	— H. (○)	Dill. elth. t. 275. f. 256.
31 judaicum. R.S. (wh.) rough-stalked.	Europe.	...	— H. (○)	
32 suffruticosum. R.S. (wh.) suffrutescent.	Barbary.	1804.	5. 9. G. h.	
33 quadrangularis. R.S. (wh.) quadrangular.	C. B. S.	1812.	— H. (○)	
34 triangulare. R.S. (wh.) three-sided.	E. Indies.	...	6. 9. S. (○)	
35 rubrum. R.S. (wh.) red-berried.	S. America.	...	— H. (○)	
36 incertum. R.S. (wh.) uncertain.	E. Indies.	1810.	— H. (○)	
37 miniatum. R.S. (wh.) pubescent-stalk'd.	Europe.	...	— H. (○)	
38 humile. R.S. (wh.) small.	S. Europe.	...	— H. (○)	
39 flavum. R.S. (st.) yellow-berried.	Hungary.	1816.	— H. (○)	
ochroleucum. Dun.				
40 villosum. R.S. (wh.) villous annual.	S. Europe.	...	— H. (○)	Dill. elth. t. 274. f. 353.
41 hirsutum. R.S. (wh.) hairy annual.	Egypt.	...	— H. (○)	
42 decemdentatum. F.I. (wh.) ten-toothed.	China.	1826.	— H. (○)	
43 Rumphii. R.S. (wh.) Rumphius's.	E. Indies.	...	— H. (○)	Rumph. amb. 6. t. 26. f. 1.

44	Besséri. R.S. (wh.) Besser's.	America.	1824.	6. 9. H. ⊖.	
45	cestrifolium. s.s. (wh.) Cestrum-leaved.	1826.	4. 7. S. h.	
46	longiflorum. R.S. (bl.) long-flowered.	Guiana.	1823.	6. 8. S. h.	Dunal solan. t. 9.
	<i>longifolium</i> . Dun.				
47	triquêtrum. R.S. (wh.) triquetrous.	New Spain.	1821.	— S. h.	Cavan. ic. 3. t. 259.
48	bombéne. R.S. (wh.) Bomba.	S. America.	1818.	— S. h.	Jacq. f. eclog. t. 24.
49	pubigerum. R.S. (wh.) downy.	Mexico.	1823.	— S. h.	Dunal sol. t. 6.
	<i>Cervantésii</i> . Lag.				
50	pyrifolium. R.S. (wh.) Pear-leaved.	W. Indies.	1824.	6. 8. S. h.	Dun. sol. ed. 2. t. 34.
51	macranthérum. R.S. (vi.) large-anthered.	Mexico.	1826.	— G. h. ⊖.	H. et B.gen.am.3.t.195.
52	laurifolium. R.S. (wh.) Laurel-leaved.	S. America.	1820.	— S. h.	Dunal sol. t. 8.
53	spirál. F.I. (wh.) spiral-racemed.	E. Indies.	—	4. 6. S. h.	
54	Hookerianum. s.s. (bl.) Hooker's.	Mexico.	1825.	6. 9. G. h.	Bot. mag. t. 2708.
	<i>coriaceum</i> . Bot. mag. nec aliorum.				
55	dealbátum. Lind. (wh.) white-leaved.	Chile.	1826.	10. 11. G. h.	Bot. mag. t. 2697.
	<i>saponáceum</i> . B.M. non Dunal.				
56	salicifolium. D.L. (wh.) Willow-leaved.	Trinidad.	1822.	6. 9. S. h.	
57	verbascifolium. R.S. (w.) Mullein-leaved.	W. Indies.	1749.	6. 8. S. h.	Jacq. vind. 1. t. 13.
58	auriculatum. R.S. (vi.) ear-leaved.	Madagascar.	1773.	— S. h.	Scop. del. 3. t. 8.
59	pubescens. R.S. (pu.) pubescent.	E. Indies.	1816.	6. 8. S. h.	Willd. phyt. 1. t. 3.
60	argénteum. R.S. (wh.) silvery.	Brazil.	—	— S. h.	Dunal sol. ed. 2. t. 39.
61	nudum. K.S. (wh.) naked.	Mexico.	1824.	3. 6. S. h.	
62	diphyllum. R.S. (wh.) two-leaved.	W. Indies.	1699.	6. 7. S. h.	Jacq. ic. 2. t. 322.
63	arbóreum. R.S. (wh.) tree.	Cumana.	1819.	— S. h.	Dunal sol. ed. 2. t. 198.
64	havanéne. R.S. (bl.) Havannah.	W. Indies.	1793.	7. 8. S. h.	Jacq. amer. pict. t. 48.
65	aggregatum. R.S. (li.) crowded.	Guinea.	—	— S. h.	Jacq. ic. 2. t. 323.
66	Plukentii. R.S. (wh.) Plukenet's.	S. America.	—	6. 8. S. h.	Pluk. phyt. t. 227. f. 2.
67	eriocályx. R.S. (wh.) woolly-calyxed.	Carthagena.	1731.	5. 8. S. h.	
68	umbellátum. R.S. (wh.) umbelled.	S. America.	—	— S. h.	
69	americanum. R.S. (pu.) American.	Virginia.	1727.	7. 9. H. ⊖.	
70	axilliflórum. R.S. (bl.) axil-flowered.	Vera Cruz.	1728.	6. 9. S. h. ⊖.	
	scándens. Miller.				
71	memphiticum. R.S. (wh.) tooth-stalked.	1823.	— S. 4.	
72	pérsicum. R.S. (bl.) Persian.	Persia.	1829.	6. 9. H. h. ⊖.	
73	capsicátrum. Ot. (wh.) Capsicum-like.	1827.	— S. h.	
74	lycioídes. R.S. (li.) spiny.	Peru.	1791.	5. 6. S. h.	Jacq. ic. 1. t. 46.
75	monánthum. R.S. (fl.) one-flowered.	New Spain.	1816.	— S. 4.	
76	fugax. R.S. (wh.ye.) fugacious.	Caracas.	—	— S. h.	Jacq. ic. 2. t. 324.
77	stellátum. R.S. (bl.) starry.	S. America.	—	— S. h.	— 2. t. 325.
78	virgátum. R.S. (vi.) twiggy.	W. Indies.	1819.	— S. h.	Dunal sol. ed. 2. t. 40.
79	lérntum. R.S. (vi.) climbing woolly.	New Spain.	1823.	6. 8. S. h. ⊖.	Cavan. ic. 4. t. 308.
80	scándens. R.S. (pu.) climbing.	W. Indies.	1822.	— S. h. ⊖.	
81	atrosanguíneum. Sck. (c.) dark crimson.	1827.	— S. h.	
82	elæagnifolium. R.S. (bl.) Elæagnos-leaved.	Chile.	1823.	6. 8. S. h.	Cavan. ic. 3. t. 243.
83	obtusifolium. K.S. (bl.) blunt-leaved.	Mexico.	1826.	— G. h.	Dunal sol. ed. 2. t. 119.
84	stelligerum. R.S. (li.) starry-haired.	N. Holland.	1820.	— S. h.	Sm. exot. bot. 2. t. 88.
85	racemósium. R.S. (wh.) wave-leaved.	W. Indies.	1781.	7. 8. S. h.	Jacq. amer. pict. t. 50.
86	igneum. R.S. (wh.) red-spined.	S. America.	1714.	3. 11. S. h.	Jacq. vind. 1. t. 14.
87	subarmatum. R.S. (wh.) partly armed.	1823.	— S. h.	
88	bahaméne. R.S. (wh.vi.) Bahama.	Baháma.	1732.	6. 7. S. h.	Dill. elth. t. 271. f. 350.
89	hirtum. R.S. (wh.) hairy-calyxed.	Trinidad.	1823.	— S. h.	Vahl ic. rar. d. 3. t. 21.
90	tomentosum. R.S. (bl.) tomentose.	S. America.	1662.	— S. h.	Bocc. sic. 8. t. 5.
91	hy'bridum. R.S. (pa.bl.) hybrid.	Guinea.	1815.	— S. h.	Jacq. vind. 2. t. 113.
92	coccineum. R.S. (bh.) scarlet-fruited.	1820.	— S. h.	Jacq. ic. 1. t. 43.
93	jamaicénse. R.S. (li.) Jamaica.	Jamaica.	—	— S. h.	
94	cnefífolium. R.S. (wh.) wedge-leaved.	W. Indies.	—	— S. h.	Dunal sol. t. 22.
95	heterophyllum. R.S. (bh.) variable-leaved.	Guiana.	1823.	7. 9. S. h.	— ed. 2. t. 62.
96	volutíle. R.S. (bl.) twining.	Hispaniola.	—	— S. h. ⊖.	Plum. MSS. t. 4. f. 32.
97	lanceafolium. R.S. (wh.) lance-leaved.	S. America.	—	— S. h. ⊖.	Jacq. ic. 2. t. 329.
98	coriáceum. R.S. (bl.) leathery-leav'd.	Trinidad.	1824.	6. 9. S. h. ⊖.	Dunn. sol. p. 197. t. 14.
99	bonariénse. R.S. (wh.) Buenos Ayres.	BuenosAyres.	1727.	— H. h.	Dill. elth. t. 272. f. 351.
100	fastigiátum. R.S. (pa.bl.) fastigiate.	S. America.	1820.	— S. h.	Jacq. f. eclog. t. 6.
101	oporinum. R.S. (bl.) large-leaved.	—	— S. h.	Dunal sol. t. 16.
	<i>macrophyllum</i> . Dun.				
102	subinérme. R.S. (bl.) spear-leaved.	W. Indies.	1752.	7. 8. S. h.	Jacq. amer. t. 40. f. 3.
103	lanceolátum. R.S. (bl.) lanceolate.	New Spain.	—	6. 7. S. h.	Cavan. ic. 3. t. 245.
104	astroítés. Jacq. (bl.) Aster-like.	S. America.	1823.	— S. h.	Jacq. f. eclog. t. 65.
105	glutinósium. R.S. (bl.) glutinous.	1818.	6. 8. S. h.	Dunal sol. ed. 2. t. 54.
106	Brownii. R.S. (vi.) Brown's.	N. S. W.	1820.	— S. h.	
	<i>violaceum</i> . B.P. non Jacq.				

107	giganteum. R.S. (<i>wh. vi.</i>) tall.	C. B. S.	1792.	6. 8. G. h.	Bot. mag. t. 1921.
108	tórvum. R.S. (<i>wh.</i>) curved-prickled.	W. Indies.	1816.	— S. h.	Jacq. schœn. 3. t. 334.
109	ferrugíneum. R.S. (<i>wh.</i>) ferruginous.	1824.	— S. h.	.
110	élatum. L.en. (<i>bl.</i>) tall.	1824.	6. 9. S. h.	.
111	ochroneúrum. L.en. (<i>bl.</i>) rusty-branched.	Brazil.	—	6. 8. S. h.	.
112	môle. K.S. (<i>wh.</i>) soft-leaved.	Caracas.	1826.	— G. h.	.
113	Lichtensteíni. R.S. .. Lichtenstein's.	C. B. S.	1824.	— G. h.	.
114	esculentum. R.S. (<i>li.</i>) eatable.	Tropics.	1597.	6. 9. G. ⊙.	Dunal sol. t. 3. f. E.
115	ovígerum. R.S. (<i>pa. bl.</i>) Egg-plant.	Arabia.	—	— G. ⊙.	Blackw. t. 549.
	à violaceum. (<i>pa. bl.</i>) violet-fruited.	—	—	— G. ⊙.	
	β álbum. (<i>pa. bl.</i>) white-fruited.	—	—	— G. ⊙.	
	γ luteum. (<i>pa. bl.</i>) yellow-fruited.	—	—	— G. ⊙.	
	δ rüber. (<i>pa. bl.</i>) red-fruited.	—	—	— G. ⊙.	
116	lóngum. F.I. (<i>vi.</i>) long-fruited.	E. Indies.	1816.	— S. h.	
117	insánum. W. (<i>pu.</i>) insane.	—	1815.	8. 10. S. h.	Pluk. alm. t. 226. f. 3.
118	melanoxylon. s.s. (<i>vi.</i>) black-wooded.	—	1825.	7. 10. S. ⊙.	
119	undátum. R.S. (<i>pa. bl.</i>) waved.	E. Indies.	1818.	6. 9. S. h.	Rheed. mal. 2. t. 37.
120	fuscátem. R.S. (<i>vi.</i>) brown-branched.	America.	—	7. 8. G. ⊙.	Jacq. ic. 1. t. 42.
121	sodómeum. R.S. (<i>pu.</i>) black-spined.	Africa.	1688.	6. 7. G. h.	Flor. græc. t. 235.
122	incánum. R.S. (<i>re.</i>) hoary.	Mauritius.	1823.	— S. h.	Scop. del. 1. t. 1.
123	índicum. R.S. (<i>bl.</i>) Indian.	India.	1732.	7. S. h.	Dill. elth. t. 270. f. 349.
124	coágulans. R.S. (<i>bl.</i>) scollop-leaved.	Arabia Felix.	1802.	— S. h.	Jacq. schœn. 4. t. 469.
125	sánctum. R.S. (<i>bl. pu.</i>) oblique-leaved.	Egypt.	—	— S. h.	
126	marginátum. R.S. (<i>li.</i>) white margined.	Africa.	1775.	6. 9. G. h.	Bot. mag. t. 1928.
127	campeñéne. L.en. (<i>bl.</i>) Congo.	—	1815.	— G. h.	
128	campechiéne. R.S. (<i>li.</i>) purple-spined.	America.	1732.	7. G. h.	Dill. elth. t. 268. f. 347.
129	plataniólium. B.M. (<i>vi.</i>) Plane-leaved.	S. America.	1822.	7. 9. S. h.	Bot. mag. t. 2618.
130	myriacánthum. R.S. (<i>w.</i>) many-spined.	—	—	— S. h.	
131	aculeatíssimum. R.S. (<i>w.</i>) very prickly.	—	1800.	— S. h.	Jacq. ic. 1. t. 41.
132	spinosisíssimum. D.L. (<i>w.</i>) very spiny.	Trinidad.	1820.	— S. h.	
133	lividum. s.s. (<i>wh.</i>) livid.	Madagascar.	1821.	— S. h.	
134	lasiocárpus. R.S. (<i>wh.</i>) four-celled.	E. Indies.	1824.	6. 8. S. ⊗.	Rheel. mal. 2. t. 35.
	hirsútum. F.I. non Dunal. <i>quadriloculare</i> . s.s.	—	—	—	
135	mammósium. R.S. (<i>pa. bl.</i>) nipple.	W. Indies.	1699.	7. 8. H. ⊙.	Merian. surin. t. 27.
136	ciliátum. R.S. (<i>pa. bl.</i>) ciliated.	—	1816.	— H. ⊙.	Dunal sol. t. 18.
137	stramonifólium. R.S. (<i>bl.</i>) broad-leaved.	E. Indies.	1778.	6. 9. S. h.	Jacq. ic. 1. t. 44.
138	flavéscentis. R.S. (<i>wh.</i>) yellowish.	Trinidad.	1822.	— S. h.	
139	férox. R.S. (<i>li. pu.</i>) Malabar.	Malabar.	1795.	8. 9. S. ⊗.	Dunal sol. ed. 2. t. 68.
140	crassipétalum. F.I. (<i>li.</i>) thick-petaled.	Nepaul.	1824.	4. 7. G. h.	
141	lysimachioídes. F.I. (<i>w.</i>) Loosestrife-like.	E. Indies.	1823.	6. 9. S. ⊗.	
142	campanulátum. B.P. (<i>pu.</i>) bell-flowered.	N. S. W.	1818.	8. 9. G. ⊙.	
143	armátum. B.P. (<i>pu.</i>) armed.	—	1819.	— S. ⊗.	Dunal sol. ed. 2. t. 69.
144	pungétium. B.P. (<i>vi.</i>) pungent.	—	1823.	— G. ⊙.	— ed. 2. t. 70.
145	cíneum. B.P. (<i>wh.</i>) grey.	—	—	— G. ⊗.	
146	Milléri. R.S. (<i>wh.</i>) Miller's.	C. B. S.	1762.	7. 8. G. h.	Jacq. ic. 2. t. 330.
147	trilobátum. R.S. (<i>wh.</i>) three-lobed.	W. Indies.	1759.	8. S. h.	Pluk. alm. t. 316. f. 5.
148	acetosafólium. R.S. (<i>w.</i>) Sorrel-leaved.	E. Indies.	—	— S. h.	Burm. ind. t. 22. f. 2.
149	rigéscens. R.S. (<i>vi.</i>) rigid.	C. B. S.	1815.	6. 8. G. h.	Jacq. schœn. I. t. 42.
150	carolinéne. R.S. (<i>w. bl.</i>) Carolina.	Carolina.	1732.	7. 9. G. ⊗.	Jacq. ic. 2. t. 331.
151	violáceum. R.S. (<i>bl.</i>) violet.	E. Indies.	1823.	6. 9. S. h.	Jacq. fragm. t. 133. f. 1.
152	pyracántum. B.M. (<i>pu.</i>) orange-thorn'd.	Madagascar.	1789.	8. 9. S. h.	Sm. exot. bot. t. 64.
	β inérmis. (<i>pu.</i>) smooth-calyxed.	—	—	— S. h.	Bot. mag. t. 2547.
153	virgiñánum. R.S. (<i>vi.</i>) Virginian.	Virginia.	1662.	5. 8. H. ⊙.	Dill. elth. t. 267. f. 346.
154	Jacquiní. R.S. (<i>bl. pu.</i>) Jacquin's.	E. Indies.	1804.	9. 11. S. ⊙.	Jacq. ic. 2. t. 332.
155	xanthocárpus. R.S. (<i>bl.</i>) yellow-fruited.	Ethiopia.	1828.	6. 10. H. ⊙.	Wend. s. han. 1. t. 2.
156	Balbísi. R.S. (<i>wh.</i>) Balbis's.	S. America.	1816.	4. 9. S. h.	Bot. mag. t. 2568.
	decurréns. B.R. 140. brancafólium. Jacq. f. ecl. t. 7. viscósum. DC.	—	—	—	
157	téctum. R.S. (<i>ye.</i>) villous-calyxed.	Mexico.	1826.	7. 10. G. h.	Cavan. icon. 4. t. 309.
158	capénse. R.S. (<i>wh.</i>) Cape prickly.	C. B. S.	1816.	6. 10. G. h.	
159	Houstoní. R.S. (<i>wh.</i>) Houstoun's.	Vera Cruz.	1731.	— S. h.	
160	Angírium. R.S. (<i>ye.</i>) procumbent.	—	—	— S. h.	
NYCT'ERIUM. V.		Pentandria Monogynia.			
1	cordifól. v. (<i>vi.</i>) Canary.	Canaries.	1779.	3. 4. G. h.	Venten. malm. t. 85.
	Solánum vespertilio. R.S.	—	—	—	
2	Amazónium. B.M. (<i>bl.</i>) Lambert's.	Brazil.	1814.	5. 9. S. h.	Bot. mag. t. 1801.
	Solánum Amazónium. Bot. reg. t. 71.	—	—	—	
3	cornútum. L.en. (<i>ye.</i>) horned.	S. America.	1825.	— S. h.	Juss. ann. mus. 3. t. 9.
4	lobátum. (<i>ye.</i>) Missouri.	Missouri.	1813.	7. 9. H. ⊙.	Pursh fl. amer. 1. t. 7.
	Solánum heterándrum. Ph. Androcéra lobata. N.	—	—	—	

SOLANEÆ.

5 rostratum. L.en. (<i>ye.</i>) beaked.	Mexico.	1823.	7. 9. H. ⊙.	Dunal sol. t. 24.
6 Fontanesianum. (<i>ye.</i>) Desfontaine's.	Brazil.	1813.	— H. ⊙.	Bot. reg. t. 177.
7 heterodoxum. L.en. (<i>bl.</i>) very prickly.	Mexico.	1824.	— H. ⊙.	Dunal sol. t. 25.
WITHERINGIA. R.S. WITHERINGIA.	Tetr-Pentandria Monogynia.			
1 solanacea. R.S. (<i>st.</i>) yellow-flowered.	S. America.	1742.	5. 9. S. ȳ.	L'Herit. sert. angl. t. 1.
2 stramonifolia. K.s. (<i>st.</i>) Stramonium-ld.	Mexico.	1823.	— S. h.	Dill. elth. t. 273. f. 352.
3 crassifolia. R.S. (<i>li.</i>) thick-leaved.	C. B. S.	1729.	— G. h.	Bot. mag. t. 2768.
4 montana. R.S. (<i>wh. bl.</i>) mountain.	Peru.	1822.	— G. ȳ.	Cavan. ic. 4. t. 359. f. 1.
5 phyllantha. R.S. (<i>bl.</i>) annual.	—	—	7. 9. H. ⊙.	Flor. peruv. 2.t.170.f.b.
6 pinnatifida. R.S. (<i>li.</i>) pinnatifid.	—	—	6. 8. G. ȳ.	
CE'STRUM. R.S.	CE'STRUM. Pentandria Monogynia.	L.		
1 laurifolium. R.S. (<i>ye.</i>) Laurel-leaved.	W. Indies.	1691.	5. 8. S. h.	Smith spicil. 2. t. 2.
2 citrifolium. R.S. (<i>st.</i>) Citron-leaved.	—	1816.	— S. h.	Venten. choix. t. 18.
3 macrophyllum. R.S. (<i>st.</i>) large-leaved.	W. Indies.	1812.	— S. h.	Dill. elth. t. 153. f. 185.
4 venenatum. R.S. (<i>st.</i>) poisonous.	C. B. S.	1787.	2. 4. G. h.	Bot. mag. t. 1770.
5 nocturnum. R.S. (<i>gr. ye.</i>) night-smelling.	E. Indies.	1732.	11. S. h.	Jacq. schœn. 4. t. 452.
6 hirtum. R.S. (<i>gr. wh.</i>) hairy.	Jamaica.	1822.	5. 8. S. h.	L'Herit. s. nov. 1. t. 35.
7 Párqui. R.S. (<i>st.</i>) Willow-leaved.	S. America.	1787.	6. 7. G. h.	Jacq. schœn. 3. t. 332.
8 thyrsoides. K.s. (<i>st.</i>) bunch-flowered.	Mexico.	1826.	5. 7. S. h.	Flor. peruv. 2. t. 155.
9 suberosum. R.S. (<i>gr.</i>) cork-barked.	—	1815.	6. 7. S. h.	
10 auriculatum. w. (<i>wh.</i>) eared-leaved.	Peru.	1774.	— S. h.	Bot. mag. t. 2929.
11 exstipulatum. R.S. (<i>gr.</i>) exstipulate.	S. America.	1824.	— S. h.	Jacq. schoen. 3. t. 324.
12 tinctorium. R.S. (<i>wh.</i>) dying.	Caracas.	1823.	5. 6. S. h.	————— 3. t. 325.
13 undulatum. R.S. (<i>st.</i>) waved-leaved.	Peru.	—	— S. h.	————— 3. t. 326.
14 multiflorum. R.S. (<i>gr.</i>) many-flowered.	S. America.	1825.	— S. h.	————— 3. t. 327.
15 paniculatum. K.s. (<i>gr. ye.</i>) paniced.	Caracas.	1819.	— S. h.	Sloan. hist. 2. t. 204. f. 2.
16 latifolium. R.S. (<i>st.</i>) broad-leaved.	Trinidad.	1822.	5. 6. S. h.	Jacq. schoen. 3. t. 329.
17 alaternoïdes. R.S. (<i>gr. ye.</i>) Alaternus-like.	S. America.	1820.	4. 6. S. h.	Bot. mag. t. 2974.
18 hirsutum. R.S. (<i>st.</i>) hairy-stalked.	—	1822.	5. 6. S. h.	Swt. br. fl. gar. ser. 2.
19 caulinorum. R.S. (<i>wh.</i>) stem-flowering.	Martinica.	1818.	— S. h.	Bot. mag. t. 1729.
20 salicifolium. R.S. (<i>st.</i>) Sallow-leaved.	Caracas.	—	3. 6. S. h.	Jacq. schœn. 3. t. 331.
21 pendulinum. R.S. (<i>wh.</i>) pendulous.	—	1820.	— S. h.	Dill. elth. t. 154. f. 186.
22 vespertinum. R.S. (<i>wh.</i>) cluster-flowered.	W. Indies.	1759.	5. 7. S. h.	Lodd. bot. cab. t. 618.
23 pallidum. R.S. (<i>gr. ye.</i>) pale-flowered.	Jamaica.	1732.	— S. h.	Bot. reg. t. 354.
24 fetidissimum. R.S. (<i>ye.</i>) stinking.	—	—	5. 8. S. h.	Trew Ehret. t. 24. f. 1.
25 bracteatum. B.M. (<i>st.</i>) large-bracted.	S. America.	1828.	12. 2. S. h.	
26 acuminatum. (<i>ye. gr.</i>) tapering-leaved.	Mexico.	1824.	7. 10. F. h.	
27 fastigiatum. R.S. (<i>wh.</i>) Honeysuckle.	W. Indies.	—	10. 12. S. h.	
28 odontospermum. R.S. (<i>w.</i>) smooth.	S. America.	1793.	7. 9. S. h.	
29 diurnum. R.S. (<i>wh.</i>) day smelling.	W. Indies.	1732.	10. 11. S. h.	
30 roseum. R.S. (<i>ro.</i>) rose-coloured.	Mexico.	1829.	— G. h.	
31 tomentosum. R.S. (<i>st.</i>) downy.	S. America.	1790.	6. 7. S. h.	
32 angustifolium. B.C. (<i>st.</i>) narrow-leaved.	—	1820.	— S. h.	
LYCIUM. R.S.	Box-THORN.	Pentandria Monogynia.	L.	
1 áfrum. R.S. (<i>vi.</i>) African.	C. B. S.	1712.	6. 7. H. h.	
2 tenué. R.S. (<i>pu.</i>) slender.	—	—	— H. h.	
3 rigidum. R.S. (<i>vi.</i>) rigid.	—	1795.	— H. h.	
<i>carnosum</i> . P.S. <i>italicum</i> Miller.				
4 microphyllum. R.S. (<i>vi.</i>) small-leaved.	—	—	— H. h.	Pall. ros. 1. t. 49.
5 ruthenicum. R.S. (<i>ro.</i>) Russian.	Siberia.	1804.	— H. h.	
6 tetrandrūm. R.S. (<i>pu.</i>) tetrandrous.	C. B. S.	—	— H. h.	
7 barbārum. R.S. (<i>re.</i>) Willow-leaved.	Barbary.	1696.	5. 8. H. h.	Wats. dend. brit. t. 9.
8 Shawii. R.S. (<i>wh.</i>) white-flowered.	—	—	— H. h.	Shaw afric. f. 349.
9 Trewiānum. R.S. (<i>pu.</i>) Trew's.	China.	—	— H. h.	Trew Ehret. t. 68.
<i>chinense</i> . P.S. <i>non</i> Mill.				
10 chinense. R.S. (<i>re.</i>) Chinese.	—	—	— H. h.	Wats. dend. brit. t. 8.
<i>ovatum</i> . P.S. <i>barbārum</i> β <i>chinense</i> . w.				
11 lanceolatum. P.S. (<i>vi.</i>) spear-leaved.	S. Europe.	—	— H. h.	Duham. arb. nov. t. 32.
12 turbinatum. P.S. (<i>re.</i>) top-shaped.	China.	1709.	— H. h.	————— t. 31.
13 cinereum. R.S. (<i>pu.</i>) gray.	C. B. S.	1815.	6. 8. H. h.	Flor. græc. t. 236.
14 europæum. R.S. (<i>li.</i>) European.	S. Europe.	1730.	5. 8. H. h.	L'Herit. stirp. t. 83.
15 hóridum. R.S. (<i>wh.</i>) succulent-leav'd.	C. B. S.	1791.	7. 8. H. h.	Flor. peruv. 2. t. 182.f.a.
16 boerhaaviaefolium. (<i>bh.</i>) glaucous-leaved.	Peru.	1780.	4. 5. G. h.	
17 arborescens. s.s. (<i>wh.</i>) arborescent.	—	1819.	6. 7. G. h.	
<i>A'tropo arboréscens</i> . R.S.				
18 caroliniānum. R.S. (<i>pu.</i>) Carolina.	Carolina.	1806.	7. 9. H. h.	Jacq. schœn. 1. t. 45.
19 capsulare. R.S. (<i>pu.</i>) capsular.	Mexico.	1730.	6. 8. G. h.	
SOLA'NDRA. R.S.	SOLA'NDRA.	Pentandria Monogynia.	R.S.	
1 grandiflora. R.S. (<i>li.</i>) large-flowered.	Jamaica.	1781.	3. 8. S. h.	

2 nitida. R.S.	(st.) glossy-leaved.	Jamaica.	1781.	3. 8. S. h.	Bot. mag. t. 1874.
<i>grandiflora.</i> B.M. non Jacq.					
3 viridiflora. s.s.	(gr.) green-flowered.	Brazil.	1816.	5. 6. S. h.	Bot. mag. t. 1948.
4 oppositifolia.	opposite-leaved.	W. Indies.	1824. S. h.	
BRUNEL'SIA. W.	BRUNEL'SIA.	<i>Didynamia Angiospermia.</i> L.			
1 undulata. w. (wh.st.)	large-flowered.	Jamaica.	1780.	6. 7. S. h.	Bot. reg. t. 228.
2 americana. w.(wh.st.)	American.	W. Indies.	1735.	— S. h.	Bot. mag. t. 393.
<i>a latifolia.</i> (wh.st.) broad-leaved.				S. h.	
<i>β angustifolia.</i> (wh.st.) narrow-leaved.				S. h.	
3 violacea. B.C. (wh.st.)	purple-stalked.		1820.	— S. h.	Lod. bot. cab. t. 792.
4 uniflora. D.D. (bl.wh.)	single-flowered.	Brazil.	1827.	1. 12. S. h.	Bot. mag. t. 2829.
<i>Franciscea Hopéana.</i> B.M.					
CRESCENTIA. W.	CALABASH-TREE.	<i>Didynamia Angiospermia.</i> L.			
1 Cujête. w.	(li.ye.) oval-fruited.	Jamaica.	1690. S. h.	Jacq. amer. t. 111.
2 acuminata. K.S.	(li.) acuminate.	Cuba.	1822.	— S. h.	
3 cucurbitina. w. (li.ye.)	round-fruited.	W. Indies.	1733. S. h.	Plum. ic. t. 109.
4 alata. K.S.	(li.) winged-leaved.	Mexico.	1825. S. h.	
TRIPINNARIA. S.S.	TRIPINNARIA.	<i>Didynamia Angiospermia.</i>			
africana. s.s.	(re.) African.	Mauritius.	1823. S. h.	Jacq. coll. 3. t. 18.
<i>Crescentia pinnata.</i> Jacq. <i>Tan'acium pinnatum.</i> w.					

ORDO CLII.

SCROPHULARINÆ. Brown prodr. 1. p. 433.

Sect. I. STAMINA (4) ANTERIFERA.

BU'DDLEA. S.S.	BU'DDLEA.	<i>Tetrandria Monogynia.</i> L.			
1 madagascariensis. (or.)	Madagascar.	Madagascar.	1824.	9. 10. S. h.	Bot. mag. t. 2824.
2 saligna. s.s.	(wh.) Willow-like.	C. B. S.	1816.	8. 9. G. h.	Jacq. schœn. 1. t. 29.
3 paniculata. s.s.	(wh.) paniced.	Nepaul.	1823.	— G. h.	
4 americana. R.S.	(ye.) American.	W. Indies.	1824.	7. 10. S. h.	Sloan. hist. 2. t. 173. f.1.
5 occidentalis. R.S. (wh.)	occidental.	Peru.	1730. G. h.	Flor. peruv. 1. t. 83.
6 salvifolia. s.s. (wh.cr.)	Sage-leaved.	C. B. S.	1760.	8. 9. G. h.	Jacq. schœn. 1. t. 28.
7 Neemda. s.s.	(wh.) Nepaul.	Nepaul.	1822.	— G. h.	
8 brasiliensis. (or.)	Brazilian.	Brazil.	—	9. 11. S. h.	Bot. mag. t. 2713.
9 heterophylla. B.R. (ye.)	different-leaved.	S. America.	1826.	1. 5. S. h.	Bot. reg. t. 1259.
10 globosa. s.s.	(ye.) round-headed.	Chile.	1774.	5. 7. H. h.	Bot. mag. t. 174.
11 connata. s.s.	(or.) connate-leaved.	Peru.	1826.	5. 6. G. h.	— t. 2853.
12 salicifolia. s.s.	(ye.) Willow-leaved.	S. America.	1823. S. h.	
13 diversifolia. R.S. (wh.)	various-leaved.	Java.	— S. h.	
SCOPARIA. L.	SCOPARIA.	<i>Tetrandria Monogynia.</i> L.			
1 dulcis. L.	(wh.) sweet.	S. America.	1730.	6. 9. S. h.	Herm. parad. t. 241.
2 flava. Ot.	(ye.) yellow.		1829.	— S. h.	
CAPR'ARIA. S.S.	GOAT-WEED.	<i>Didynamia Angiospermia.</i> L.			
1 undulata. s.s.	(ro.) wave-leaved.	C. B. S.	1774.	3. 7. G. h.	Bot. mag. t. 1556.
2 cuneata. s.s.	(ro.) wedge-leaved.	S. America.	1759.	6. 8. S. h.	
3 lanceolata. s.s.	(ye.) spear-leaved.	C. B. S.	1774.	5. 7. G. h.	Pl. sel. h. ber. t. 4.
4 biflora. s.s.	(wh.) two-flowered.	S. America.	1752.	7. 8. S. h.	Jacq. amer. t. 115.
5 humiliis. w.	(ro.) dwarf.	E. Indies.	1781.	— S. ⊖.	
TE'EDIA. S.S.	TE'EDIA.	<i>Didynamia Angiospermia.</i> S.S.			
1 lúcida. s.s.	(pu.) shining.	C. B. S.	1774.	4. 7. G. h.	Bot. reg. t. 210.
<i>Capraria lúcida.</i> w. <i>Borkhausenia lúcida.</i> Roth.					
2 pubescens. s.s.	(pu.) pubescent.	C. B. S.	1816.	5. 10. G. h.	Bot. reg. t. 214.
HALL'ERIA. S.S.	HALL'ERIA.	<i>Didynamia Angiospermia.</i> L.			
1 lúcida. s.s.	(sc.) shining-leaved.	C. B. S.	1752.	6. 8. G. h.	Bot. mag. t. 1744.
2 elliptica. s.s.	(sc.) elliptic-leaved.		1816.	— G. h.	Burm. afr. t. 84. f. 1.
STEMODIA. S.S.	STEMODIA.	<i>Didynamia Angiospermia.</i> W.			
1 suffruticosa. s.s.	(bl.) suffrutescent.	S. America.	1823.	5. 8. S. h.	Roxb. Corom. t. 163.
2 viscósa. s.s.	(vi.) viscous.	E. Indies.	1818.	7. 10. S. ⊖.	
3 verticillaris. s.s.	(pu.) whorled.	Brazil.	1822.	— H. ⊖.	
4 parviflora. H.K.	(wh.) small-flowered.	S. America.	1759.	7. 8. S. ⊖.	
5 marítima. s.s.	(bl.) sea-side.	Cuba.	1822.	— S. ⊖.	Jacq. amer. t. 174. f.66.

SCROPHULARINÆ.

RUSS'ELIA. S.S.	RUSS'ELIA. Didynamia Angiospermia. W.
1 floribunda. s.s. (sc.) abundant-flowering.	Mexico. 1824. 6. 8. S. 4.
2 multiflora. s.s. (sc.) many-flowered.	1812. — S. 4.
TREVIR'ANIA. S.S.	TREVIR'ANIA. Didynamia Angiospermia. S.S.
coccinea. s.s. (sc.) scarlet.	Jamaica. 1778. 8. 10. S. 4.
Cyrilla pulchella. B.M. Achimenes coccinea. P.S.	
COLU'MNEA. S.S.	COLU'MNEA. Didynamia Angiospermia. L.
1 scàndens. s.s. (sc.) climbing.	W. Indies. 1759. 8. 9. S. ½.
2 hirsuta. s.s. (li.) hairy.	1780. 8. 11. S. ½.
3 trifoliata. L.en. (bl.) three-leaved.	S. America. 1825. — S. ½.
4 rutilans. s.s. (ye.re.) red-leaved.	Jamaica. 1823. — S. ½.
5 hispida. s.s. (sc.) bristly.	1824. — S. ½.
MAURA'NDIA. S.S.	MAURA'NDIA. Didynamia Angiospermia. W.
1 semperflorans. s.s. (ro.) red-flowered.	Mexico. 1796. 1. 12. G. ½. — Bot. mag. t. 460.
2 antirrhiniflora. s.s. (bl.) blue-flowered.	1814. — G. ½. — t. 1643.
3 Barclayana. B.R. (bl.) Mr. Barclay's.	1826. 4. 11. G. ½. — Bot. reg. t. 1108.
LOPHOSPERMUM. D.D. LOPHOSPERMUM. Didynamia Angiospermia.	
scàndens. L.T. (pu.) climbing.	Mexico. 1830. 6. 8. G. ½. —
DODA'R'TIA. S.S.	DODA'R'TIA. Didynamia Angiospermia. L.
orientalis. s.s. (d.pu.) oriental.	Levant. 1752. 7. 8. H. 4.
CYMBARIA. S.S.	CYMBARIA. Didynamia Angiospermia. L.
davurica. s.s. (ye.) Daurian.	Davuria. 1796. 6. 7. H. 4.
COLLI'NSIA. N.	COLLI'NSIA. Didynamia Angiospermia.
1 verna. n. (bl.pu.) large-flowered.	N. America. 1827. 4. 11. H. ⊙.
2 parviflora. d. (bl.) small-flowered.	— 6. 9. H. ⊙.
NE'MESIA. S.S.	NE'MESIA. Didynamia Angiospermia. S.S.
1 chamædryfolia. s.s. (w.bl.) Chamædry's-ld. C. B. S.	1787. 4. 9. G. 4.
Antirrhinum mucrocárpum. w.	
2 bicorne. s.s. (wh.bl.) horned.	1774. 7. 8. G. ⊙.
3 foetens. s.s. (wh.bl.) foetid.	1798. 4. 9. G. 4.
4 linearis. s.s. (sn.ro.) linear-leaved.	1822. — G. ⊙.
ANARRH'INUM. S.S.	ANARRH'INUM. Didynamia Angiospermia. S.S.
1 pubescens. E. (bl.) pubescent.	Mount Sinai. 1826. 5. 8. H. 4.
2 bellidifolium. s.s. (bl.) Daisy-leaved.	France. 1629. 6. 8. H. 3.
ANTIRRH'INUM. S.S.	SNAP-DRAGON. Didynamia Angiospermia. L.
1 mājus. s.s. (va.) great.	England. 6. 8. H. 4.
a albiflorum. (wh.) white-flowered. — H. 4.
β ròseum. (ro.) rose-coloured. — H. 4.
γ coccineum. (sc.) scarlet. — H. 4.
δ bicolor. (re.wh.) two-coloured. — H. 4.
ε multiplex. (sc.) double-flowered. — H. 4.
2 meonanthum. L.en. (re.) smaller-flower'd. S. Europe.	1820. — H. 4.
3 angustifolium. Urv. (re.) narrow-leaved.	Italy. 1818. — H. 4.
4 siculum. s.s. (st.) Sicilian.	Sicily. 1804. 7. 8. H. 4.
5 Oróntium. s.s. (bh.) Calf's-snout.	Britain. 7. 9. H. ⊙.
6 calycinum. L.en. (wh.) long-calyxed.	S. Europe. — H. ⊙.
7 Asarina. s.s. (wh.) heart-leaved.	Italy. 1699. 7. F. 4.
8 molle. s.s. (wh.) soft-leaved.	Spain. 1752. 7. 10. F. 4.
LIN'ARIA. S.S.	TOAD-FLAX. Didynamia Angiospermia. S.S.
1 fruticosa. s.s. (ye.) shrubby.	Barbary. 1822. 5. 6. G. ½.
2 frutescens. s.s. (ye.) frutescent.	C. B. S. 1816. 5. 7. G. ½.
Antirrhinum frutescens. Th.	
3 dalmatica. s.s. (ye.) Dalmatian.	Levant. 1731. 6. 7. G. ½.
4 scoparia. s.s. (ye.) broom-like.	Teneriffe. 1815. — G. ½.
5 Cymbalaria. s.s. (bh.) Ivy-leaved.	England. 5. 11. H. 4.
6 pallida. t.s. (li.) pale-flowered.	Italy. 1824. — H. 4.
pubescens. T.P. non P.S.	
7 pilosa. s.s. (li.) hairy-leaved.	1800. 6. 9. H. 4.
8 hepaticæfolia. s.s. (pu.) Hepatica-leav'd.	Corsica. 1828. — F. 4.
9 aequitribola. s.s. (d.pu.) equal 3-lobed.	Sardinia. — — F. 4.
10 Elatine. s.s. (bl.) sharp-pointed.	England. 7. 11. H. ⊙.
11 cirrhosa. s.s. (bl.wh.) cirrhose.	Egypt. 1777. 7. 8. H. ⊙.
12 spuria. s.s. (ye.vi.) round-leaved.	England. 7. 9. H. ⊙.
Antirrhinum spuriu'm. E.B.	
13 lanigera. s.s. (st.) woolly.	S. Europe. 1820. — H. ⊙.
14 dealbata. s.s. (ye.) villous-leaved.	Portugal. — 7. 9. H. ⊙.
15 latifolia. s.s. (ye.) broad-leaved.	Barbary. 1800. 6. 7. H. ⊙.
16 triphylla. s.s. (bl.ye.) three-leaved.	Sicily. 1596. 6. 9. H. ⊙.

Bot. reg. t. 805.
Brown jam. t. 30. f. 3.

Bot. mag. t. 1528.

Bot. mag. t. 374.

Desf. atl. 2. t. 134.

Bot. mag. t. 324.

Burm. afr. t. 75. f. 3.
Venten. malm. t. 41.

Bot. mag. t. 2056.

Eng. bot. v. 2. t. 129.

Lk. fl. port. 1. t. 56.

Eng. bot. v. 17. t. 1155.

Bot. mag. t. 902.

Desf. atl. 2. t. 135.

Buxb. cent. 1. t. 24.

Eng. bot. v. 7. t. 502.

Jacq. obs. 2. t. 48.

Bot. mag. t. 2941.

Eng. bot. v. 10. t. 692.

Jacq. vind. 1. t. 82.

Eng. bot. v. 10. t. 691.

17. <i>reflexa</i> . s.s. (wh.) reflex-flowered. S. Europe.	1824. 6. 9. H. ⊙.	Flor. græc. t. 593.
18. <i>Organifolia</i> . s.s. (bl.) Marjoram-leav'd. —	1785. — F. 4.	Barrel. ic. 598.
19. <i>cretacea</i> . s.s. (bl.ye.) crowded-flower'd. Siberia.	1826. — H. ⊙.	
20. <i>virgata</i> . s.s. (bl.) slender-stalked. Barbary.	1824. — H. ⊙.	Desf. atl. 2. t. 135.
21. <i>thymifolia</i> . s.s. (ye.) Thyme-leaved. S. Europe.	— — H. ⊙.	
22. <i>villosa</i> . s.s. (wh.) villous. Spain.	1786. 7. 8. F. 4.	Barrel. ic. 597.
23. <i>rubrifolia</i> . s.s. (bl.) red-leaved. S. France.	1824. — F. 4.	
24. <i>saxatilis</i> . P.S. (ye.) rock. S. Europe.	1827. 6. 8. H. 4.	
25. <i>pubescens</i> . P.S. (ye.) pubescent. —	1824. 6. 9. H. ⊙.	
26. <i>hirta</i> . s.s. (ye.) shaggy-leaved. Spain.	1759. — H. ⊙.	Jacq. ic. 1. t. 117.
27. <i>triornithophora</i> . (pu.) three bird's. Portugal.	1710. — F. 4.	Bot. mag. t. 523.
28. <i>parviflora</i> . s.s. (ye.) small flowered. Sardinia.	1829. — H. ⊙.	Desf. atl. 2. t. 137.
29. <i>albifrons</i> . F.G. (wh.) white fronted. Greece.	1828. — H. ⊙.	Flor. græc. t. 588.
30. <i>bipartita</i> . s.s. (bl.) cloven flowered. Barbary.	1801. — H. ⊙.	Swt. br. fl. gar. t. 30.
<i>speciosa</i> . Jacq. fil. eclog. t. 95.		
31. <i>minor</i> . s.s. (pu.ye.) little erect. England. 6. 11. H. ⊙.	Eng. bot. v. 28. t. 2014.
<i>Antirrhinum minus</i> . E.B.		
32. <i>littoralis</i> . s.s. (bl.) shore. Austria.	1824. — H. ⊙.	
33. <i>chalepensis</i> . s.s. (wh.) white-flowered. Levant.	1680. 6. 7. H. ⊙.	Flor. græc. t. 592.
34. <i>tristis</i> . s.s. (bk.br.) melancholy. Spain.	1727. 7. 8. G. 4.	Bot. mag. t. 74.
35. <i>versicolor</i> . s.s. (ye.bl.) various-colour'd. France.	1777. 7. 9. H. ⊙.	Jacq. ic. 1. t. 116.
36. <i>linarioides</i> . s.s. (ye.) toadflax-like. S. Europe.	1816. 5. 8. H. 4.	
37. <i>micrantha</i> . s.s. (ye.) small-flowered. Spain.	1825. 5. 7. H. ⊙.	Flor. græc. t. 587.
38. <i>viscosa</i> . s.s. (br.ye.) clammy. —	1786. 7. H. ⊙.	Bot. mag. t. 368.
39. <i>purpurea</i> . s.s. (pu.) purple. S. Europe.	1648. 7. 9. H. 4.	Flor. græc. t. 589.
40. <i>arenaria</i> . s.s. (ye.) sand. Hungary.	1825. — H. ⊙.	
41. <i>alpina</i> . s.s. (bl.) Alpine. Austria.	1570. 7. 11. H. 4.	Bot. mag. t. 205.
42. <i>lusitanica</i> . s.s. (ye.) Portugal. S. Europe.	1820. 6. 7. H. 4.	
43. <i>pyrenaica</i> . s.s. (ye.) Pyrenean. Pyrenees.	1824. 6. 8. H. ⊙.	
44. <i>Hælava</i> . F.E. (ye.) hairy calyxed. Egypt.	1803. 7. 9. H. ⊙.	
45. <i>pelisseriana</i> . s.s. (vi.) violet-coloured. S. Europe.	1640. 6. 9. H. ⊙.	Flor. græc. t. 591.
46. <i>elegans</i> . P.S. (bl.) elegant. Spain.	1812. — H. ⊙.	
47. <i>répens</i> . H.K. (std.) creeping rooted. England. 7. 10. H. 4.	Eng. bot. v. 18. t. 1253.
48. <i>amethystina</i> . s.s. (bl.) bright blue. S. Europe.	1816. — H. ⊙.	Cavan. ic. 1. t. 33. f. 1.
<i>Antirrhinum bipunctatum</i> . Cav. nec <i>aliorum</i> .		
49. <i>supina</i> . s.s. (ye.) trailing. Spain.	1728. — H. ⊙.	Flor. græc. t. 595.
50. <i>bipunctata</i> . s.s. (ye.) two-spotted. —	1749. 6. 8. H. ⊙.	
51. <i>spartea</i> . s.s. (ye.) branching. —	1772. 6. 10. H. ⊙.	Bot. mag. t. 200.
52. <i>simplex</i> . s.s. (bl.) upright. S. Europe.	1816. 7. 8. H. ⊙.	Jacq. ic. 3. t. 499.
53. <i>arvensis</i> . s.s. (ye.) corn.	... — H. ⊙.	Flor. græc. t. 590.
54. <i>diffusa</i> . s.s. (ye.bl.) spreading. Portugal.	1824. — H. ⊙.	
55. <i>multicaulis</i> . s.s. (wh.bl.) many-stalked. Levant.	1728. 5. 7. H. ⊙.	Bocc. sic. t. 19. f. 1.
56. <i>macroura</i> . L.en. (ye.) large-spurred. Caucasus.	1823. 6. 9. H. 4.	
57. <i>incarnata</i> . s.s. (li.) flesh-coloured. S. Europe.	1826. — H. ⊙.	
58. <i>reticulata</i> . s.s. (ye.pu.) netted-flower'd. Algiers.	1788. 5. 7. F. 4.	Smith ic. pict. t. 2.
59. <i>glauca</i> . s.s. (ye.) glaucous-leaved. S. Europe.	1800. 6. 8. H. ⊙.	Buxb. cent. 4. t. 37.
60. <i>ægyptiaca</i> . s.s. (ye.) Egyptian. Egypt.	1771. 7. H. ⊙.	
61. <i>sileneifolia</i> . Lk. (ye.) Silene-leaved. Portugal.	1820. 7. 9. H. ⊙.	
62. <i>genistifolia</i> . s.s. (ye.) Broom-leaved. Austria.	1704. 7. 8. H. 4.	Flor. græc. t. 596.
63. <i>linifolia</i> . s.s. (ye.) flax-leaved. Italy.	1824. — H. 4.	
64. <i>vulgaria</i> . s.s. (ye.) common. Britain. 6. 9. H. 4.	Eng. bot. v. 10. t. 658.
<i>Antirrhinum Linaria</i> . E.B.		
β <i>Peloria</i> . (ye.) regular-flowered. — — H. 4.	Eng. bot. v. 4. t. 260.
65. <i>Loeselii</i> . s.s. (ye.bl.) Loesel's. Russia.	1824. — H. ⊙.	
66. <i>odora</i> . M.B. (ye.bl.) sweet-scented. Caucasus.	1827. — H. ⊙.	
67. <i>stricta</i> . s.s. (ye.) upright. Levant.	1829. — H. ⊙.	Flor. græc. t. 594.
68. <i>junccea</i> . s.s. (ye.) rush-stalked. Spain.	1780. 7. 8. H. ⊙.	
69. <i>canadensis</i> . s.s. (vi.) Canadian. N. America.	1812. 6. 8. H. ⊙.	Venten. cels. t. 49.
70. <i>acutifolia</i> . F. (ye.bl.) sharp-leaved. Levant.	1826. — H. ⊙.	
71. <i>fasciculata</i> . Ot. (ye.) clustered. 1829. — H. ⊙.		
SCROPHULARIA. L. FIGWORT. Didynamia Angiospermia. L.		
1. <i>marilandica</i> . s.s. (g.br.) Maryland. N. America.	1759. 5. 7. H. 4.	
2. <i>peregrina</i> . s.s. (pu.) Nettle-leaved. Italy.	1640. 6. 8. H. ♂.	Flor. græc. t. 597.
3. <i>nodosa</i> . s.s. (d.pu.) knotty-rooted. Britain. 5. 7. H. 4.	Eng. bot. v. 22. t. 1544.
4. <i>aquatica</i> . s.s. (d.pu.) water.	— v. 12. t. 854.
5. <i>glabrata</i> . s.s. (pu.ye.) spear-leaved. Canaries.	1779. 4. 5. G. ♂.	Jacq. schœn. 2. t. 209.
6. <i>biserrata</i> . s.s. (pu.ye.) double-serrate.	1816. 5. 7. H. 4.	
7. <i>arguta</i> . s.s. (re.) slender upright. Canaries.	1778. — H. ⊙.	
8. <i>vernalis</i> . s.s. (ye.) yellow. Britain. 3. 5. H. ♂.	Eng. bot. v. 8. t. 567.

SCROPHULARINÆ.

9 Smithii. L.en. (<i>br.ye.</i>) Smith's.	Canaries.	1815.	4. 8. G. 4.	
10 glandulösa. s.s. (<i>br.ye.</i>) glandular.	Hungary.	1806.	6. 9. H. 4.	W. et K. hung. t. 214.
11 rugösa. w.k. (<i>br.ye.</i>) rugged.	—	1816.	— H. 4.	Scop. carn. 2. t. 32.
12 Scopoliæ. P.S. (<i>br.ye.</i>) Scopoli's.	Germany.	—	— H. 4.	Eng. bot. v. 31. t. 22.
13 Scorodonia. s.s. (<i>pu.</i>) balm-leaved.	Britain.	7. 8. H. 4.	Mur. com. got. 4. t. 2.
14 grandidentata. T.S. (<i>pu.</i>) great-toothed.	Italy.	1829.	6. 8. H. 4.	Barrel. ic. 274.
15 altáica. s.s. (<i>st.</i>) white-flowered.	Siberia.	1786.	5. 6. H. 4.	
16 betonicaefolia. s.s. (<i>pu.</i>) Betony-leaved.	Spain.	1752.	6. 8. H. 4.	Lobel. ic. 533.
17 hirsuta. L.en. (<i>pu.ye.</i>) hairy.	S. Europe.	1822.	— H. 4.	
18 appendiculata. s.s. (<i>pu.</i>) heart-leaved.	Morocco.	1805.	7. H. 4.	Jacq. schœn. 3. t. 286.
19 Balbisii. s.s. (<i>pu.ye.</i>) Balbis's.	Italy.	1823.	6. 8. H. 4.	W. et K. hung. 2. t. 170.
20 auriculata. s.s. (<i>br.ye.</i>) ear-leaved.	Spain.	1772.	7. 8. H. 4.	
21 micrantha. s.s. (<i>pu.</i>) small-flowered.	Levant.	1828.	— H. 4.	
22 heterophylla. s.s. (<i>pu.ye.</i>) various-leaved.	—	1827.	— G. ½.	
23 trifoliata. s.s. (<i>ye.br.</i>) three-leaved.	Africa.	1731.	5. 9. H. 5.	
24 laciniata. w.k. (<i>br.ye.</i>) jag-leaved.	Hungary.	1806.	7. H. 4.	Herm. lugd. t. 547.
25 verbenaefolia. Poir. (<i>br.ye.</i>) Verbain-leav'd.	—	1816.	7. 8. H. 4.	Schk. hand. 2. t. 173.
26 cæsia. s.s. (<i>ye.br.</i>) grey.	Levant.	1828.	— F. 5.	
27 adscendens. W.en. (<i>br.ye.</i>) ascending.	—	1816.	— H. 4.	
28 variegata. s.s. (<i>pu.ye.</i>) variegated.	Caucasus.	—	— H. 4.	
29 ebullifolia. s.s. (<i>ye.pu.</i>) Elder-leaved.	—	1824.	— H. 4.	
30 grandiflora. DC. (<i>br.ye.</i>) large-flowered.	—	1816.	— H. 4.	
31 frutescens. s.s. (<i>br.ye.</i>) shrubby.	Portugal.	1768.	6. 8. F. ½.	
32 rupestris. s.s. (<i>br.ye.</i>) rock.	Tauria.	1826.	6. 9. H. 4.	
33 orientalis. s.s. (<i>br.ye.</i>) oriental.	Levant.	1710.	7. 8. H. 4.	
34 melissæfolia. s.s. (<i>br.ye.</i>) balm-like.	—	1823.	— H. 4.	
35 lanceolata. s.s. (<i>st.</i>) spear-leaved.	Pensylvania.	1826.	— H. 4.	
36 cretacea. s.s. (<i>ye.</i>) narrow-leaved.	Siberia.	1828.	— H. 4.	
37 lyraea. s.s. (<i>br.pu.</i>) lyrate-leaved.	Portugal.	1816.	— H. 4.	
38 mellifera. s.s. (<i>br.pu.</i>) honey-bearing.	Barbary.	1786.	— H. 4.	
39 sambucifolia. s.s. (<i>br.g.</i>) Elder-like.	Spain.	1640.	7. 9. H. 4.	
40 hispida. s.s. (<i>br.ye.</i>) hispid.	Levant.	1820.	— H. 4.	
41 tanacetifolia. s.s. (<i>br.ye.</i>) Tansey-leaved.	Tauria.	1804.	— H. 4.	
42 canina. s.s. (<i>or.re.</i>) wing-leaved.	S. Europe.	1683.	6. 8. F. 5.	
43 chrysanthemifolia. (<i>br.pu.</i>) Chrysanthemum-l'd.	Tauria.	1816.	— H. 4.	
44 lúcida. s.s. (<i>br.ye.</i>) shining-leaved.	Levant.	1596.	— H. 4.	
45 filicifolia. s.s. (<i>d.pu.</i>) Fern-leaved.	Crete.	— H. 4.	
46 multifida. s.s. (<i>br.ye.</i>) multifid-leaved.	—	1816.	— H. 4.	
47 glauca. F.G. (<i>pu.</i>) glaucous.	Greece.	1827.	— H. 4.	
48 pubescens. F. (<i>pu.</i>) pubescent.	—	—	— H. 4.	
49 pallida. F. (<i>pa.</i>) pale-flowered.	Iberia.	—	— H. 4.	
50 obtusifolia. F. (<i>pu.</i>) blunt-leaved.	Hungary.	1821.	— H. 4.	
DIGITALIS. S.S. FOXGLOVE. Didynamia Angiospermia. L.				
1 Scéptrum. s.s. (<i>or.</i>) Madeira shrubby.	Madeira.	1777.	7. 8. G. ½.	Sm. exot. bot. 2. t. 73.
2 canariensis. s.s. (<i>ye.</i>) Canary.	Canaries.	1698.	6. 7. G. ½.	Bot. reg. t. 48.
3 obscura. s.s. (<i>or.</i>) Willow-leaved.	Spain.	1778.	7. 8. H. 4.	Lind. digit. t. 26.
4 lutea. s.s. (<i>st.</i>) small yellow.	France.	1629.	— H. 4.	— t. 23.
5 hybrida. Sv. (<i>pi.wh.</i>) hybrid.	Hybrid.	— H. 4.	Salv. nou. bull. 3.337.t.6.
<i>lutea</i> γ Lindl. digit. t. 25.				
6 fucata. P.S. (<i>pu.</i>) painted.	—	— H. 4.	
<i>lutea</i> δ Lindl. digit. t. 24.				
7média. s.s. (<i>ye.</i>) middle yellow.	Germany.	1816.	— H. 4.	
8 micrantha. s.s. (<i>st.</i>) small-blossom'd.	Switzerland.	1824.	— H. 4.	Elm. mon. 46. t. 2.
9 tubiflora. L.D. (<i>st.</i>) tube-flowered.	S. Europe.	1816.	— H. 4.	Lind. digit. t. 22.
10 lutescens. L.D. (<i>st.</i>) pale yellow.	—	—	— H. 4.	— t. 21.
11 purpurascens. s.s. (<i>pi.</i>) blush-flowered.	Italy.	1776.	— H. 4.	— t. 20.
<i>erubescens</i> . H.K.				
12 rígida. L.D. (<i>pu.ye.</i>) rigid.	S. Europe.	1824.	— H. 4.	Lind. digit. t. 19.
13 viridiflora. L.D. (<i>gr.</i>) green-flowered.	Levant.	1827.	— H. 4.	— t. 18.
14 parviflora. s.s. (<i>pu.</i>) small-flowered.	—	1798.	6. 8. H. 4.	— t. 17.
15 laciniata. B.R. (<i>ye.</i>) cut-leaved.	Malaga.	1826.	— H. 4.	Bot. reg. t. 1201.
16 orientalis. s.s. (<i>li.wh.</i>) oriental.	Levant.	1818.	7. 8. H. 4.	— t. 554.
17 lanata. s.s. (<i>wh.br.</i>) woolly.	Hungary.	1789.	6. 7. H. 4.	Swt. br. fl. gar. t. 291.
18 leucophæa. s.s. (<i>wh.br.</i>) broad-lipped.	Greece.	1788.	6. 10. H. 4.	Lind. digit. t. 14.
19 aurea. s.s. (<i>or.</i>) golden.	—	1815.	7. 8. H. 4.	— t. 13.
20 ferruginea. s.s. (<i>br.</i>) iron-coloured.	Italy.	1597.	— H. 4.	t. 12.
21 sibirica. L.D. (<i>ye.re.</i>) Siberian.	Siberia.	1820.	— H. 4.	t. 11.
22 lavigata. s.s. (<i>ye.</i>) shining-leaved.	Hungary.	1810.	— H. 4.	t. 10.
23 fulva. s.s. (<i>co.</i>) tawny.	S. Europe.	1820.	6. 8. H. 4.	t. 9.

24	ambigua. w. <i>grandiflora</i> . s.s.	(ye.) greater yellow. Switzerland.	1596.	7. 8. H. 4.	Bot. reg. t. 64.
25	<i>ochroleuca</i> . s.s.	(st.) sulphur-color'd. Germany.	1816.	— H. 4.	Lind. digit. t. 8.
26	<i>occidentis</i> . s.s.	(br.) netted-flower'd. Hungary.	1824.	— H. 4.	Lind. digit. t. 5. 6.
27	<i>minor</i> . s.s.	(pu.) dwarf. Spain.	1789.	6. 7. H. 4.	Lind. digit. t. 5. 6.
28	<i>erostachya</i> . Bes.	(pu.) woolly spiked.	1826.	— H. 4.	Bot. mag. t. 2194.
29	<i>tomentosa</i> . s.s.	(pu.) tomentose. Portugal.	1820.	6. 8. H. 4.	Sm. exot. bot. I. t. 43.
30	<i>Thapsia</i> . s.s.	(pu.) Mullein. Spain.	1752.	5. 8. H. 4.	Eng. bot. v. 19. t. 1297.
31	<i>purpurea</i> . s.s.	(pu.) common. Britain.	6. 9. H. 5.	
	<i>β alba</i> .	(wh.) white-flowered.	— H. 5.	
CHEL'ONE. S.S.					
	CHEL'ONE. Didynamia Angiospermia. L.				
1	glabra. w.	(wh.) white-flowered. N. America.	1730.	8. 10. H. 4.	Trew pl. select. t. 88.
2	obliqua. w.	(pu.) red-flowered.	1752.	— H. 4.	Bot. reg. t. 175.
3	Lyoni. s.s.	(re.) Lyon's.	1812.	7. 9. H. 4.	Swt. br. fl. gar. t. 293.
	<i>major</i> . Bot. mag. t. 1864.				
4	nemorosa. D.	(pu.) wood. N. America.	1827.	7. 9. H. 4.	Bot. reg. t. 1211.
5	Scoulérii. D.	(bl.) Scouler's. Columbia.	—	5. 7. H. 4.	— t. 1277.
6	erianthèra. N.	(bl.) blue-flowered. Louisiana.	1811.	7. 9. F. 4.	Bot. mag. t. 1672.
	<i>Pentstemon glabra</i> . B.M.				
7	barbata. s.s.	(sc.) scarlet. Mexico.	1794.	6. 9. H. 4.	Bot. reg. t. 116.
8	rösea. B.F.G.	(ro.) rose-coloured.	1825.	— H. 5.	Swt. br. fl. gar. t. 230.
	<i>Pentstemon angustifolium</i> . B.R. t. 1122. non Pursh.				
9	campanulata. s.s. (ro.)	bell-flowered. Mexico.	1794.	3. 10. F. 5.	Bot. mag. t. 1878.
	<i>Pentstemon campanulata</i> . Jacq. schœn. 3. t. 362.				
10	atropurpurea. B.F.G. (d. pu.)	dark purple. Mexico.	1824.	— H. 5.	Swt. br. fl. gar. t. 235.
11	pulchella. B.R.	(pu.) pretty.	1825.	5. 10. H. 5.	Bot. reg. t. 1138.
12	lævigata. s.s.	(pi.) smooth. N. America.	1776.	7. 9. H. 4.	Bot. mag. t. 1425.
13	hirsuta. s.s.	(bl. pi.) narrow-l'd hairy.	1758.	— H. 4.	Moris. s. 11. t. 21. f. 3.
14	pubescens.	(bl. pu.) broad-l'd hairy.	—	— H. 4.	Bot. mag. t. 1424.
	<i>Pentstemon pubescens</i> . B.M.				
15	gracilis. N.	(li.) slender.	1825.	— H. 4.	Bot. mag. t. 2945.
16	glaucia. D.	(li.) glaucous.	1827.	7. 9. H. 4.	Bot. reg. t. 1286.
17	digitalis. B.F.G.	(wh.) foxglove-flower'd. Arkansa.	1824.	6. 8. H. 4.	Swt. br. fl. gar. t. 120.
	<i>Pentstemon digitalis</i> . Bot. mag. t. 2587.				
18	cristata. s.s.	(vi.) crested. Louisiana.	1811.	7. 9. F. 4.	
19	cœrulea. s.s.	(bl.) slender blue.	—	— F. 4.	
	<i>Pentstemon angustifolium</i> . Ph. non K.S.				
20	grandiflora. s.s. (pi. wh.)	great-flowered.	—	— F. 4.	
21	álbida. s.s.	(wh.) dwarf.	—	— F. 4.	
22	diffusa. D.	(pu.) spreading. Columbia.	1827.	6. 10. H. 4.	Bot. reg. t. 1132.
23	ovata. D.	(bl. pu.) oval-leaved. N. America.	—	6. 8. H. 4.	Bot. mag. t. 2903.
24	venusta. D.	(pu.) handsome.	—	7. 9. H. 4.	Bot. reg. t. 1309.
25	deusta. D.	(pu.) blasted.	—	— H. 4.	
26	decussata. D.	(pu.) cross-leaved.	—	— H. 4.	
27	glandulosa. D.	(pu.) glandular.	—	— H. 4.	Bot. reg. t. 1262.
28	acuminata. D.	(pu.) pointed-leaved.	—	6. 8. H. 4.	— t. 1285.
29	Richardsoni. D.	(pu.) Richardson's.	—	7. 10. H. 4.	— t. 1121.
30	attenuata. D.	(st.) taper-pointed.	—	7. 9. H. 4.	— t. 1295.
31	conferta. D.	(st.) crowded-flower'd. Columbia.	—	— H. 4.	— t. 1260.
32	pruinosa. D.	(bl.) blue-leaved. N. America.	—	6. 8. H. 4.	— t. 1280.
33	procera. D.	(vi.) tall.	—	— H. 4.	Bot. mag. t. 2954.
34	speciosa. D.	(bl.) beautiful.	—	6. 9. H. 4.	Bot. reg. t. 1270.
35	trifylla. D.	(li.) three-leaved.	—	7. 9. H. 4.	— t. 1245.
36	serrulata. D.	(pu.) saw-leaved.	—	— H. 4.	
37	suffruticosa. D.	(pu.) suffrutescent.	—	— H. 5.	
HERPE'STIS. G.					
	HERPE'STIS. Didynamia Angiospermia. S.S.				
1	stricta. s.s.	(bl.) upright. Brazil.	1823.	6. 8. S. 4.	
2	Monnieria. K.S.	(wh.) thyme-leaved. S. America.	1772.	7. 9. S. 4.	Ehret. pict. t. 14. f. 2.
3	portulacacea. B.M. (w.)	Purslane-leaved.	1822.	— S. 4.	Bot. mag. t. 2557.
4	cuneifolia. Ph.	(bl.) wedge-leaved. N. America.	1812.	8. 9. H. 4.	
5	amplexicaulis. s.s. (bl.)	stem-clasping. Carolina.	1822.	6. 8. H. 4.	
6	micrantha. Ph.	(wh.) small-flowered. N. America.	1827.	7. 8. H. 4.	
7	semiserrata. Ot.	(wh.) semiserrate. Mexico.	1824.	— G. 4.	
8	chrysanthia. Ot.	(ye.) yellow-flowered.	1829.	— G. 4.	
MAZUS. B.P.					
	MAZUS. Didynamia Angiospermia. S.S.				
	rugosus. B.P.	(wh.) China. China.	1780.	5. 10. H. 4.	Swt. br. fl. gar. t. 36.
	<i>Lindernia japonica</i> . Th.	<i>Hornemannia bicolor</i> . W.en.			
TOR'ENIA. B.P.					
	TOR'ENIA. Didynamia Angiospermia. S.S.				
1	ovata.	(wh.) oval-leaved. Manila.	1823.	5. 10. H. 4.	Pl. sel. hort. ber. t. 9.
	<i>Hornemannia ovata</i> . L.en.	<i>Titmannia ovata</i> . s.s.			

SCROPHULARIACE.

2 viscósa.	(wh.) viscous.	E. Indies.	1823.	5. 10. S. ⊖.
Hornemánnia viscósa.	W.en. Tittmánnia viscósa. s.s.			
LINDE'RNIA. B.P.	LINDE'RNIA.	Didynamia Angiospermia. L.		
Pyxidária. w.	(pu.) European.	S. Europe.	1789.	6. 8. H. ⊖.
MI'MULUS. L.	MONKEY-FLOWER.	Didynamia Angiospermia. L.		
1 glutinósum. s.s.	(ye.) orange-colored.	California.	1794.	1. 12. G. h.
2 ríngens. s.s.	(li.) gaping.	N. America.	1759.	7. 8. H. Ȑ.
3 alátus. s.s.	(li.) oval-leaved.		1783.	— H. Ȑ.
4 guttátus. DC.	(ye.) spotted-flowered.	America.	1812.	6. 9. F. Ȑ.
lúteus. B.M. non L.				
5 lúteus. B.R.	(ye.) yellow-flower'd.	Chile.	1825.	— H. Ȑ.
β rivularis. B.R.	(ye. pu.) dark-spotted.	Columbia.	—	— H. Ȑ.
6 floribúnus. B.R.	(ye.) many-flowered.		1826.	8. 10. H. ⊖.
7 moschátus. B.R.	(ye.) musk-scented.		—	7. 10. H. Ȑ.
8 parviflórus. B.R.	(st.) small-flowered.	Mexico.	1824.	6. 9. H. ⊖.
BROWA'LLIA. L.	BROWA'LLIA.	Didynamia Angiospermia. L.		
1 eláta. w.	(bl. wh.) upright.	Peru.	1768.	6. 9. G. ⊖.
2 demíssia. w.	(bl. wh.) spreading.	S. America.	1735.	— G. ⊖.
ANGEL'ONIA. K.S.	ANGEL'ONIA.	Didynamia Angiospermia. L.		
salicariæfólia. k.s. (bl.)	Salicaria-leav'd.	S. America.	1818.	7. 10. S. Ȑ.
ALONS'OA. R.P.	ALONS'OA.	Didynamia Angiospermia. H.K.		
1 incisifolia. R.P.	(sc.) Nettle-leaved.	Chile.	1795.	5. 10. G. Ȑ.
Célsia urticáefólia. B.M.	Hémimeris urticáefólia. w.			
2 acutifolia. R.P.	(sc.) acute-leaved.	Peru.	1790.	— G. Ȑ.
3 intermédia. B.C.	(or.) intermediate.		—	— G. Ȑ.
4 lineáris. H.K.	(sc.) linear-leaved.		1790.	— G. Ȑ.
5 caulinálata. R.P.	(sc.) wing-stemmed.	Chile.	1823.	5. 10. G. Ȑ.
DIA'SCIA. L.en.	DIA'SCIA.	Didynamia Angiospermia. S.S.		
Bergiána. L.en. (re.) Bergius's.		C. B. S.	1816.	7. 10. H. ⊖.
Hémimeris diffusa. w.				
VANDE'LLIA. S.S.	VANDE'LLIA.	Didynamia Angiospermia. L.		
diffusa. s.s.	(wh.) spreading.	S. America.	1824.	7. 9. S. ⊖.
LIMOSE'LLIA. L.	MUDWORT.	Didynamia Angiospermia. L.		
aquática. L.	(fl.) water.	Britain.	—	7. 9. H. ⊖.
DISA'NDRA. L.	DISA'NDRA.	Heptandria Monogynia. L.		
prostráta. L.	(ye.) trailing.	Madeira.	1771.	5. 8. G. Ȑ.
SIBTHO'RPIA. L.	SIBTHO'RPIA.	Didynamia Angiospermia. L.		
européa. L.	(wh.) Cornishmoney-wort.	England.	—	7. 8. H. Ȑ.
ERINUS. S.S.	ERINUS.	Didynamia Angiospermia. L.		
1 Lychnidéa. s.s. (wh. pu.)	Phlox-flower'd.	C. B. S.	1821.	3. 10. G. Ȑ.
2 frágrans. H.K. (wh. pu.)	fragrant.		1776.	5. 6. G. Ȑ.
3 alpinus. P.S.	(pu.) smooth-leaved.	Pyrenees.	1739.	3. 4. H. Ȑ.
4 hispánicus. P.S.	(pu.) hairy-leaved.	Spain.	—	— H. Ȑ.
MAN'ULEA. S.S.	MAN'ULEA.	Didynamia Angiospermia. L.		
1 viscósa. s.s.	(re.) clammy.	C. B. S.	1774.	6. 11. G. Ȑ.
Buchnèra viscósa. B.M.				
2 tormentosa. s.s.	(or.) tomentose.		1774.	5. 12. G. Ȑ.
3 angustifólia. L.en.	(or.) narrow-leaved.		1822.	— G. Ȑ.
4 rhynchántha. s.s. (or.)	wedge-leaved.		1816.	6. 9. G. Ȑ.
5 oppositifólia. v.	(wh.) opposite-leaved.		—	5. 12. G. Ȑ.
6 villosa. P.S.	(wh.) villous.		1783.	6. 7. G. ⊖.
Buchnèra capénensis. w.				
7 cerúlea. s.s.	(bl.) blue-flowered.		1822.	— G. Ȑ.
8 rúbra. s.s.	(re.) red.		1790.	4. 9. G. Ȑ.
9 Cheiránthus. s.s. (ye.)	Wallflower-like.		1795.	6. 8. G. Ȑ.
10 argéntea. s.s.	(wh.) silvery.		1801.	7. 11. G. ⊖.
11 pendulculáta. P.S. (wh.)	solitary-flower'd.		1790.	6. 11. G. Ȑ.
12 f'œtida. s.s.	(wh.) stinking.		1794.	6. 9. G. ⊖.
Buchnèra f'œtida. A.R.				
13 violacea. s.s.	(vi.) violet-flowered.		1816.	— G. Ȑ.
14 cordata. s.s.	(wh.) trailing heart-ld.		—	— G. Ȑ.
BUCHN'ERA. S.S.	BUCHN'ERA.	Didynamia Angiospermia. L.		
americana. s.s.	(bl.) American.	N. America.	1733.	6. 8. H. Ȑ.
EUPHR'ASIA. W.	EYE-BRIGHT.	Didynamia Angiospermia. L.		
1 linifolia. s.s.	(li.) flax-leaved.	S. France.	1823.	7. 9. H. ⊖.
2 lútea. s.s.	(ye.) yellow.	S. Europe.	1816.	— H. ⊖.
3 alpina. s.s.	(pu.) Alpine.		1823.	— H. ⊖.
4 latifolia. s.s.	(re.) broad-leaved.		1826.	— H. ⊖.
5 mímina. P.S.	(wh. pu.) least.	Switzerland.	1828.	— H. ⊖.

Lam. ill. t. 522.

Bot. mag. t. 354.

—— t. 283.

Bot. mag. t. 1501.

Feuill. peruv. 2. t. 34.

Bot. reg. t. 1030.

—— t. 1125.

—— t. 1118.

—— t. 874.

Bot. mag. t. 34.

—— t. 1136.

Bot. reg. t. 415.

Bot. mag. t. 417.

Lodd. bot. cab. t. 1456.

Bot. mag. t. 210.

Pluk. phyt. 320. f. 5.

Marc. bras. 1. c. 15. f. 1.

Eng. bot. v. 5. t. 357.

Bot. mag. t. 218.

Eng. bot. v. 10. t. 619.

Bot. reg. t. 748.

Burm. afr. t. 49. f. 4.

Bot. mag. t. 310.

Bot. mag. t. 217.

Bot. mag. t. 322.

Pl. sel. hort. ber. t. 20.

Vent. malm. t. 15.

Burm. afr. t. 50. f. 2.

Comm. hort. 2. t. 42.

Andr. reposit. t. 84.

—— t. 80.

Jacq. aust. t. 398.

Magn. monsp. t. 94.

6 tricuspidatæ. s.s. (<i>w.pu.</i>) three-pointed.	Italy.	1828.	6. 9. H. ⊖.	Pluk. alm. t. 177. f. 1.
7 officinalis. s.s. (<i>wh.pu.</i>) common.	Britain.	7. 9. H. ⊖.	Eng. bot. v. 20. t. 1416.
BA'RTSIA. S.S.	BA'RTSIA.	Didynamia Angiospermia. L.		
1 viscosa. s.s. (<i>ye.</i>) yellow viscid.	Britain.	7. 8. H. ⊖.	Eng. bot. v. 15. t. 1045.
2 alpina. s.s. (<i>vi.</i>) Alpine.	— H. ⊖.	— v. 6. t. 361.
3 Odontites. s.s. (<i>re.</i>) red-flowered.	7. 9. H. ⊖.	— v. 20. t. 1415.
4 Trixago. s.s. (<i>ye.</i>) simple-stalked.	S. Europe.	1820.	— H. ⊖.	Flor. græc. t. 585.
Trixago <i>rhinanthina</i> . L. enum.				
5 latifolia. F.G. (<i>re.wh.</i>) broad-leaved.	Greece.	1827.	6. 9. H. ⊖.	Flor. græc. v. 5. t. 586.
CASTILLEJA. S.S.	CASTILL'EJA.	Didynamia Angiosperma. S.S.		
1 integrifolia. s.s. (<i>wh.</i>) entire-leaved.	S. America.	1825. G. ḥ.	Sm. ic. ined. t. 39.
2 toluccensis. s.s. (<i>wh.</i>) procumbent.	Mexico.	— G. ḥ.	Bot. reg. t. 925.
3 septentrionalis. B.R. (<i>st.</i>) northern.	N. America.	1824.	7. 8. H. ⊖.	Gmel. sib. 3. t. 42.
4 pálida. s.s. (<i>st.</i>) pale-flowered.	Siberia.	1782.	6. 9. H. ȳ.	
Bártzia pálida. w.				
5 coccinea. s.s. (<i>or.</i>) vermilion.	N. America.	1787.	7. 9. H. ⊖.	Bot. reg. t. 1136.
Bártzia coccinea. w. <i>Euchroma coccinea</i> . N.				
6 sessiliflora. Ph. (<i>bh.</i>) sessile-flowered.	Louisiana.	1811.	7. 8. H. ȳ.	
grandiflora s.s. <i>Euchroma grandiflorum</i> . n.				
RHINA'NTHUS. W. YELLOW RATTLE. Didynamia Angiospermia. L.				
1 Crista-gálli. w. (<i>ye.</i>) Cock's-comb.	Britain.	6. 8. H. ⊖.	Eng. bot. v. 10. t. 657.
2 mágior. E.F. (<i>ye.</i>) larger.	England.	— H. ⊖.	
3 Alecterolóphus. L.en. villous.	Germany.	1816.	— H. ⊖.	Bull. herb. t. 125.
villous. P.S.				
MELAMP'YRUM. S.S. COW-WHEAT. Didynamia Angiospermia. L.				
1 cristatum. E.B. (<i>ye.pu.</i>) crested.	England.	7. 8. H. ⊖.	Eng. bot. v. 1. t. 41.
2 arvénse. E.B. (<i>ye.pu.</i>) corn-field.	—	6. 7. H. ⊖.	— v. 1. t. 53.
3 nemorosum. s.s. (<i>ye.</i>) coloured-bracted.	Europe.	1823.	— H. ⊖.	Flor. dan. t. 305.
4 pratense. E.B. (<i>ye.</i>) common.	Britain.	7. 8. H. ⊖.	Eng. bot. v. 2. t. 113.
5 sylvaticum. E.B. (<i>co.</i>) wood.	—	— H. ⊖.	— v. 12. t. 804.
PEDICUL'ARIS. S.S. LOUSEWORT. Didynamia Angiospermia. L.				
1 incarnata. s.s. (<i>fl.</i>) flesh-coloured.	Austria.	1796.	6. 7. H. ȳ.	Jacq. aust. 2. t. 140.
2 rostrata. s.s. (<i>pu.</i>) beaked.	S. Europe.	1823.	— H. ȳ.	— t. 205.
3 tuberosa. s.s. (<i>ye.</i>) tuberous.	Switzerland.	1799.	7. 8. H. ȳ.	Hall. helv. n. 323. t. 10.
4 ascéndens. Schl. (<i>ye.</i>) ascending.	—	1825.	— H. ȳ.	
5 gyroflexa. s.s. (<i>pu.</i>) bending.	—	—	— H. ȳ.	Vill. delph. 2. t. 9.
6 atrorubens. DC. (<i>d.re.</i>) dark-red.	—	—	— H. ȳ.	
7 asplenifolia. P.S. (<i>pu.</i>) Fern-leaved.	—	1828.	— H. ȳ.	Willd. bot. zeit. c. ic.
8 uncinata. s.s. (<i>fl.ye.</i>) hooked-flowered.	Siberia.	1815.	— H. ȳ.	Gmel. sib. 3. t. 45.
9 compácta. s.s. (<i>ye.</i>) close-headed.	—	—	— H. ȳ.	
10 resupinata. s.s. (<i>pu.</i>) resupinate.	—	1816.	5. 7. H. ȳ.	Gmel. sib. 3. t. 44.
11 palústris. E.B. (<i>pu.</i>) marsh.	Britain.	6. 7. H. ȳ.	Eng. bot. v. 6. t. 399.
12 euphrasioïdes. s.s. (<i>ye.re.</i>) eyebright-leaved.	Siberia.	1816.	— H. ȳ.	Gmel. sib. 3. t. 43.
13 striata. s.s. (<i>ye.cr.</i>) stripe-flowered.	Dahuria.	1826.	— H. ȳ.	Pall.it.3.a.n.98.t.R.f.2.
14 canadensis. s.s. (<i>ye.</i>) Canadian.	N. America.	1800.	7. 8. H. ȳ.	Swt. br. fl. gar. t. 37.
15 sylvatica. E.B. (<i>ro.</i>) common.	Britain.	5. 7. H. ȳ.	Eng. bot. v. 6. t. 400.
16 myriophylla. s.s. (<i>ye.</i>) Milfoil-leaved.	Dahuria.	1816.	— H. ȳ.	Pall. it. 3. ap. t. 8. f. 1.
17 rübens. s.s. (<i>pu.</i>) red-flowered.	—	1827.	— H. ⊖.	
18 sudética. s.s. (<i>pu.</i>) Swedish.	N. Europe.	—	— H. ȳ.	Hall. helv. t. 8. f. 3.
19 flámmea. s.s. (<i>ye.sc.</i>) upright.	—	1775.	7. H. ȳ.	
20 Oèderi. s.s. (<i>ye.</i>) Oeder's.	—	1827.	— H. ȳ.	
21 versicolor. s.s. (<i>ye.br.</i>) various-coloured.	Switzerland.	1825.	7. 8. H. ȳ.	All. ped. 1. t. 4. f. 1.
22 achillefolia. s.s. (<i>ye.</i>) Milfoil-leaved.	Siberia.	1827.	— H. ȳ.	
23 comosa. s.s. (<i>ye.</i>) spiked.	Italy.	1775.	— H. ȳ.	Pall.it.3.a.n.100.t.8.f.2.
24 pálida. s.s. (<i>st.</i>) pale yellow.	N. America.	1826.	— H. ȳ.	Flor. dan. t. 26.
25 spicata. s.s. (<i>pu.</i>) spike-flowered.	Dahuria.	1827.	— H. ȳ.	Pall.it.3.a.n.97.t.R.f.1.
26 Scéptrum. s.s. (<i>ye.cr.</i>) sceptred.	Sweden.	1793.	— H. ȳ.	Jacq. aust. 3. t. 258.
27 flava. s.s. (<i>ye.</i>) yellow.	Siberia.	1828.	— H. ȳ.	— 2. t. 139.
28 recutita. s.s. (<i>pu.</i>) jagged-leaved.	Austria.	1787.	— H. ȳ.	— 3. t. 206.
29 foliosa. s.s. (<i>ye.</i>) leafy.	—	1786.	— H. ȳ.	
30 verticillata. s.s. (<i>pu.</i>) whorled.	—	1790.	5. 6. H. ȳ.	
31 proboscidea. s.s. proboscis-flower'd.	Siberia.	1827. H. ȳ.	
32 élégans. Led.	(<i>pu.</i>) elegant.	—	6. 8. H. ȳ.	
33 speciosa. Led.	(<i>pu.</i>) beautiful.	—	— H. ȳ.	
34 elata. s.s.	(<i>pu.</i>) lofty.	—	— H. ⊖.	
35 exaltata. Led.	(<i>pu.</i>) tall.	—	— H. ȳ.	
36 rósea. s.s.	(<i>ro.</i>) rose-coloured.	S. Europe.	1825.	7. 8. H. ȳ.
GERARDIA. S.S.	GERA'DIA.	Didynamia Angiospermia.		Jacq. ic. 1. t. II15.
1 delphinifolia. s.s. (<i>ro.</i>) Larkspur-leav'd.	E. Indies.	1800.	6. 7. S. ȳ.	Roxb. corom. 1. t. 90.

2 purpurea.s.s.	(pu.) purple.	N. America.	1772.	7. 8. H. ♂.	
3 tenuifolia. s.s.	(pu.) slender-leaved.	—	1812.	—	H. ♂.
4 maritima. s.s.	(li.) sea-side.	—	1823.	—	H. ♂.
5 flava. s.s.	(ye.) yellow.	—	1796.	—	H. ♀.
6 queratifolia. s.s.	(ye.) oak-leaved.	—	1812.	—	H. ♀.
7 Pedicularia. s.s.	(ye.) variable.	—	1826.	6. 8. H. ♂.	
SEYMERIA. S.S.	SEYMERIA.	Didynamia Angiospermia. S.S.			
1 tenuifolia. s.s.	(ye.) slender-leaved.	Carolina.	7. 9. H. ⓧ.	
Afzelia cassioidea. Gmel.	Gerardia Afzelia. M.				
2 pectinata. s.s.	(ye.) pectinated.	Carolina.	—	H. ⓧ.
<hr/>					
Sect. II. STAMINA (2) ANTERIFERA.					
CALCEOLARIA. R.S. SLIPPERWORT.	Diandria Monogynia. L.				
1 Fothergillii. R.S. (pu.) Fothergill's.	Falkland. Isl.	1777.	5. 8. G. ♀.		
2 plantaginea. R.S. (ye.) Plantain-leaved.	Chile.	1826.	6. 9. F. ♀.		
3 corymbosa. R.S. (ye.) corymb-flower'd.	—	1823.	4. 8. G. ♀.		
4 arachnoidea. B.M. (pu.) cobweb.	—	1827.	6. 10. F. ♀.		
5 purpurea. B.M. (pu.) purple-flowered.	—	1826.	—	F. ♀.	
6 petiolaris. R.S. (ye.) much branched.	—	1827.	6. 9. G. ♀.		
floribunda. B.R. connata.	Bot. mag. t. 2876.				
7 ascendens. B.R. (ye.) ascending.	—	1826.	—	G. ♂.	
8 polifolia. B.M. (ye.) white-leaved.	—	—	—	G. ♀.	
9 rugosa. R.P. (ye.) rugged-leaved.	—	1823.	6. 8. G. ♂.		
crenata. Bot. reg. t. 790.					
10 integrifolia. R.P. (ye.) entire-leaved.	—	—	—	G. ♂.	
rugosa. Bot. mag. t. 2523.					
11 angustifolia. (ye.) narrow-leaved.	—	1826.	—	G. ♂.	
12 thysiflora. B.M. (ye.) tufted-flowered.	—	1827.	—	G. ♂.	
13 pinnata. R.S. (st.) wing-leaved.	Peru.	1773.	7. 9. H. ⓧ.		
14 scabiosæfolia. R.S. (st.) Scabious-leaved.	—	1823.	—	G. ♀.	
SCHIZA'NTHUS. R.P. SCHIZA'NTHUS.	Diandria Monogynia.				
1 pinnatus. H.E.F. (li.pu.) winged-leaved.	Chile.	1823.	7. 10. H. ⓧ.		
2 porrigena. H.E.F. (wh.pu.) spreading.	—	—	—	H. ⓧ.	
SCHWE'NKIA. R.S. SCHWE'NKIA.	Diandria Monogynia. W.				
1 americana. R.S. (vi.) American.	Cayenne.	1781.	8. 9. S. ♂.		
2 brawallioides. K.S. (gr.pu.) Browallia-like.	Caracas.	1824.	—	S. ♀.	
3 Hilariâna. DC. (gr.pu.) St. Hilaire's.	Brazil.	1826.	—	S. ⓧ.	
BONNA'YA. S.S.	BONNA'YA.	Diandria Monogynia. S.S.			
1 brachiata. s.s. (wh.) brachiate.	E. Indies.	1825.	6. 9. H. ⓧ.		
2 brachycarpa. Ot. (vi.) short-capsuled.	—	1829.	—	S. ⓧ.	
3 veronicaefolia. s.s. (vi.) Speedwell-leaved.	—	1798.	—	S. ♂.	
Gratiola veronicaefolia. w.					
4 reptans. s.s. (li.) creeping.	—	1820.	—	S. ⓧ.	
GRATIOLA. B.P.	HEDGE-HYSSOP.	Diandria Monogynia. L.			
1 latifolia. B.P. (wh.) broad-leaved.	N. S. W.	1822.	6. 8. G. ♀.		
2 officinalis. R.S. (wh.) officinal.	Europe.	1568.	5. 8. H. ♀.		
3 virginica. R.S. (st.) Virginian.	Virginia.	1759.	8. —	H. ♀.	
4 aurea. Ph. (ye.) golden pert.	N. Amerika.	1826.	7. 9. H. ♀.		
5 quadridentata. Ph. (st.) four-toothed.	—	—	6. 8. H. ♀.		
6 acuminata. Ph. (st.) sharp-pointed.	—	1828.	7. 8. H. ♀.		
7 pilosa. Ph. (wh.) hairy.	—	1827.	—	H. ♀.	
WULFENIA. R.S.	WULFENIA.	Diandria Monogynia. R.S.			
carinthiaca. R.S. (bl.) blue-flowered.	Carinthia.	1817.	6. 9. H. ♀.		
PÆDEROTA. R.S.	PÆDEROTA.	Diandria Monogynia. L.			
1 Agèria. R.S. (st.) sulphur-coloured.	Italy.	1823.	6. 7. H. ♀.		
2 Bonarota. R.S. (vi.) violet-coloured.	Germany.	1826.	—	H. ♀.	
LEPTA'NDRA. N.	LEPTA'NDRA.	Diandria Monogynia. N.			
1 virginica. (wh.) Virginian.	Virginia.	1714.	7. 9. H. ♀.		
Veronica virginica. w.					
2 incarnata. G.D. (li.) lilac-flowered.	—	—	—	H. ♀.	
3 sibirica. N. (wh.) Siberian.	Siberia.	1779.	7. 8. H. ♀.		
VERONICA. R.S.	SPEEDWELL.	Diandria Monogynia. L.			
1 foliosa. R.S. (bl.) leafy.	Hungary.	1805.	7. 9. H. ♀.		
2 crenulata. R.S. (bl.) notch-flowered.	S. Europe.	1804.	—	H. ♀.	
3 maritima. R.S. (bl.) sea-side.	—	1570.	—	H. ♀.	
4 Hostii. R.S. (bl.) Host's.	Italy.	—	H. ♀.	

Bot. mag. t. 2048.
Bart. fl. amer. 3. t. 82.
Pluk. amal. t. 389. f. 3.
Pursh amer. 1. t. 19.
Lam. ill. t. 529. f. 2.

Bot. mag. t. 348.
— t. 2805.
Bot. reg. t. 723.
Bot. mag. t. 2874.
— t. 2775.
Bot. reg. t. 1214.

Bot. reg. t. 1215.
Bot. mag. t. 2897.
Hook. ex. flor. t. 99.

Bot. reg. t. 744.

Bot. reg. t. 1083.
Bot. mag. t. 2915.
— t. 41.
— t. 2405.

Swt. br. fl. gar. t. 63.
— t. 76.

Schw. h. m. haag. t. 1.
H. et B. n. g. am. 2. t. 181.
DC. pl. rar. genev. t. 10.

Pl. sel. h. ber. t. 11.

Roxb. corom. 2. t. 154.

Flor. dan. t. 363.

Lodd. bot. cab. t. 1399.
Lam. ill. 1. t. 16. f. 2.

Swt. br. fl. gar. t. 66.

Jacq. vind. 2. t. 121.
Jacq. aust. 5. t. 39.

Pluk. alm. t. 70. f. 2.

Amm. ruth. 20. t. 4.

W. et K. hung. 2. t. 102.
Hoff. phyt. p. 95. t. E.f. 3.
Schrad. ver. t. 1. f. 1.
Tabern. n. kreut. 2. f. 4.

5 spūria. R.S.	(bl.) spurious.	Siberia.	1731.	7. 9. H. 4.	Gmel. it. 1. t. 39.
6 paniculāta. R.S.	(bl.) paniced.	Tartary.	1797.	6. 7. H. 4.	
7 Stephaniana. R.S.	(bl.) Stephan's.	Persia.	1821.	— H. 4.	
8 complicatā. R.S.	(bl.) bundled.	S. Europe.	1812.	9. 10. H. 4.	Hoff. phyt. p. 98. t. E.f. 4.
9 azūrea. R.S.	(bl.) azure.	1818.	6. 9. H. 4.	
10 glábria. R.S.	(bl.) smooth.	S. Europe.	1804.	7. 9. H. 4.	Schrad. ver. p. 25. t. 1. f. 4.
β albiflōra. (wh.) white-flowered.	—	— H. 4.	
11 brachyphylla. R.S. (bl.) short-leaved.	1818.	— H. 4.	
γ brevijōdia. L.en.	—	— H. 4.	
12 elatiōr. R.S.	(bl.) tall.	S. Europe.	1808.	— H. 4.	
13 exaltatā.	(bl.) lofty.	Siberia.	1816.	— H. 4.	
14 falcatā. R.S.	(bl.) falcate-leaved.	1820.	6. 7. H. 4.	
15 acūta. R.S.	(bl.) acute-leaved.	—	7. 9. H. 4.	
16 argūta. R.S.	(bl.) sharp-notched.	S. Europe.	1812.	— H. 4.	Schrad. ver. n. 7. t. 2. f. 2.
17 nitens. R.S.	(bl.) glossy-leaved.	1816.	— H. 4.	
18 longibracteāta. R.S. (bl.) long-bracted.	—	— H. 4.	
β latifolia.	(bl.) broad-leaved.	—	— H. 4.	
19 mēdia. R.S.	(bl.) long-spiked.	Germany.	1804.	— H. 4.	Schrad. ver. n. 8. t. 1. f. 2.
20 persicifolia. R.S.	(bl.) Peach-leaved.	1819.	— H. 4.	
21 australis. R.S.	(bl.) pubescent.	S. Europe.	1812.	— H. 4.	Schrad. ver. n. 9. t. 2. f. 3.
22 longifolia. R.S.	(bl.) long-leaved.	1731.	— H. 4.	— p. 26. t. 2. f. 1.
23 grōssa. R.S.	(bl.) stout.	S. Russia.	1827.	6. 9. H. 4.	
24 ambigūa. R.S.	(bl.) ambiguous.	England.	— H. 4.	
25 neglēcta. R.S.	(bl.) canescent.	Siberia.	1797.	7. 9. H. 4.	
26 angustifolia. s.s.	(bl.) narrow-leaved.	1823.	— H. 4.	
27 incāna. R.S.	(bl.) hoary.	Russia.	1759.	— H. 4.	Hoff. c. gott. 15. t. 6.
28 rigēns. R.S.	(bl.) rigid.	1824.	— H. 4.	
29 élégans. R.S.	(ro.) elefant.	Italy.	— H. 4.	
30 spicatā. R.S.	(bl.) spikēd.	England.	— H. 4.	Eng. bot. v. 1. t. 2.
31 Clūsii. R.S.	(bl.) Clusius's.	N. Europe.	— H. 4.	Flor. dan. t. 52.
32 menthaefolia. R.S.	(bl.) Mint-leaved.	Austria.	1818.	— H. 4.	
33 Barrelieri. R.S.	(bl.) Barrelier's.	S. Europe.	— H. 4.	Barrel. ic. t. 682.
34 orchidēa. R.S.	(bl.) Orchis-like.	Europe.	— H. 4.	
35 crassifolia. R.S.	(vi.) thick-leaved.	Hungary.	1818.	— H. 4.	
36 hybrida. R.S.	(bl.) Welsh.	England.	— H. 4.	Eng. bot. v. 10. t. 673.
37 confusa. R.S.	(bl.) confused.	1819.	6. 7. H. 4.	
38 longiflōra. R.S.	(li.) long-flowered.	1823.	7. 9. H. 4.	
39 ruthénica. R.S.	(bl.) Russian.	Russia.	— H. 4.	
40 grāndis. R.S.	(wh.) large white.	Siberia.	1826.	— H. 4.	
41 Pònæ. R.S.	(bl.) dwarf.	Pyrenees.	1823.	— H. 4.	Gouan. ill. t. 1. f. 1.
42 polystachya. R.S.	(bl.) many-spiked.	1819.	— H. 4.	
43 præalta. L.C.	(bl.) tall upright.	1817.	6. 8. H. 4.	
44 incarnatā. G.D.	(pi.) flesh-coloured.	— H. 4.	
45 cárnea. Ræu.	(pi.) pale red.	— H. 4.	
46 corymbosā. G.D.	(bl.) corymbose.	1817.	6. 7. H. 4.	
47 melancholica. Df.	(bl.) melancholy.	1820.	— H. 4.	
48 leucántha. F.	(wh.) white-flowered.	Siberia.	1824.	— H. 4.	
49 ascēndens. R.S.	(bl.) ascending.	1828.	— H. 4.	
50 villōsa. R.S.	(bl.) villous.	S. Europe.	1800.	— H. 4.	Schrad. ver. p. 13. t. 1. f. 3.
51 pinnatā. R.S.	(bl.) wing-leaved.	Siberia.	1776.	6. 8. H. 4.	Hoff. c. gott. 15. t. 10.
52 incīsa. R.S.	(bl.) cut-leaved.	1779.	— H. 4.	
53 laciniatā. R.S.	(bl.) jagged-leaved.	1780.	— H. 4.	Jung. ic. rar. C. 1. f. 2.
54 gentianoiōdes. F.G.	(vi.) Gentian-like.	Levant.	1748.	5. 6. H. 4.	Flor. græc. 1. t. 5.
55 gentianifolia. (pa.bl.) Gentian-leaved.	—	— H. 4.	Bot. mag. t. 1002.
<i>Veronica gentianoides. B.M. non Flor. græc.</i>					
56 pallida. R.S.	(pa.) pale.	1816.	— H. 4.	
57 bellidioïdes. R.S.	(bl.) Daisy-leaved.	Switzerland.	1775.	6. 7. H. 4.	Hall. hist. t. 15. f. 1.
58 fruticulōsa. R.S.	(pu.) flesh-coloured.	Scotland.	6. 8. H. 4.	Eng. bot. v. 15. t. 1028.
59 saxatilis. R.S.	(bl.) blue rock.	6. 7. H. 4.	— v. 15. t. 1027.
60 crética. R.S.	(bl.) Cretan.	Crete.	1819.	5. 6. H. 4.	
61 Baumgartēnii. R.S. (pa.bl.) Baumgarten's.	Transylvania.	1826.	— H. 4.		
62 nummularia. R.S.	(bl.) moneywort.	Pyrenees.	1820.	6. 8. H. 4.	Gouan. ill. 1. t. 1. f. 2.
63 alpina. R.S.	(bl.) Alpine.	Scotland.	5. 6. H. 4.	Eng. bot. v. 7. t. 484.
α acutifolia.	(bl.) acute entire-leav'd.	Europe.	— H. 4.	Allion. ped. t. 22. f. 5.
β pùmila. All.	(bl.) saw-leaved.	—	— H. 4.	Eng. bot. v. 7. t. 484.
γ obtusifolia.	(bl.) blunt entire-leav'd.	Scotland.	— H. 4.	Krocker siles. t. 3.
δ integrifolia. w.	(bl.) entire-leaved.	Europe.	— H. 4.	
ε heterophylla.	(bl.) various-leaved.	—	— H. 4.	
ζ rotundifolia.	(bl.) round-leaved.	—	— H. 4.	Braun. salisb. n. 15. t. 2.

SCROPHULARINÆ.

64 Wormskiöldii. R.S. (bl.)	Wormskiold's.	Greenland.	1819.	5. 6. H. ♀.	Bot. mag. t. 2975.
65 grandifolia. R.S. (bl.)	large-leaved.	Italy.	1828.	— H. ♀.	W. et K. hung. 3. t. 245.
66 depauperata. R.S. (bl.)	few-flowered.	Hungary.	1819.	— H. ♀.	Eng. bot. v. 15. t. 1075.
67 serpyllifolia. R.S. (bl. fl.)	smooth. α orata. (bl. fl.) common.	Britain.	4. 7. H. ♀.	
β tenella. Schm. (bl.)	roundish-leaved.	Europe.	— H. ♀.	
γ neglecta. Schm. (pa.)	neglected.	—	— H. ♀.	
δ quaternata. (pa.li.)	four-leaved.	—	— H. ♀.	
ε nummulariaefolia. (pa.)	moneywort-leaved.	—	— H. ♀.	
ζ humifusa. L.T. (pa.)	trailing.	—	— H. ♀.	
68 hirsuta. R.S. (pa.)	bristly. setigera. s.s.	Scotland.	— H. ♀.	
69 tenella. R.S. (pu.)	slender.	Pyrenees.	1820.	5. 6. H. ♀.	Allion. ped. t. 22. f. 1.
70 microphylla. R.S. (pu.)	small-leaved.	Hungary.	1824.	— H. ♀.	
71 parviflora. R.S. (bl.)	small-flowered.	N. Zealand.	1822.	6. 8. H. ♀.	
72 undulata. R.S. (wh.)	wave-leaved.	E. Indies.	1827.	— G. ♀.	
73 decussata. R.S. (wh.)	cross-leaved.	Falkland Isl.	1776.	— F. h.	
74 aphylla. R.S. (bl.)	naked-stalked.	Italy.	1775.	5. H. ♀.	Bot. mag. t. 242.
75 Beccabunga. R.S. (bl.)	brook-lime.	Britain.	5. 6. H.w. ♀.	Seguier. ver. t. 3. f. 2.
76 Anagallis. R.S. (pa.bl.)	long-leaved water.	—	6. 8. H.w. ♀.	Eng. bot. v. 10. t. 655.
77 scutellata. R.S. (li.)	marsh.	—	— H.w. ♀.	— v. 11. t. 781.
78 Parmularia. R.S. (li.)	tooth-leav'd marsh.	Austria.	1816.	— H.w. ♀.	Trattin. fl. aust. b. 20.
79 caroliniana. R.S. (bl.)	Carolina.	Carolina.	1821.	6. 7. H.w. ♀.	Flor. græc. v. 1. t. 6.
80 thymifolia. F.G. (bl.)	thyme-leaved.	Crete.	1827.	5. 7. F. ♀.	Buxb. C. I. t. 39. f. 1.
81 pectinata. R.S. (bl.)	pectinated.	Levant.	1820.	6. 8. H. ♀.	Sal. stirp. rar. 7. t. 4.
82 orientalis. R.S. (bl.)	various-leaved.	—	1748.	— H. ♀.	
83 trichocarpa. R.S. (bl.)	hairy-capsuled. pilocarpa. Len.	—	1820.	— H. ♀.	
84 Jacquinii. R.S. (bl.)	Jacquin's.	Austria.	1748.	— H. ♀.	Jacq. aust. 4. t. 329.
austriaca. Jacq. non L.					
85 austriaca. R.S. (bl.)	Austrian.	—	—	— H. ♀.	Moris. h. 2. s. 3. t. 23. f. 12.
polymorpha. W.en.					
86 tenuifolia. R.S. (bl.)	slender-leaved.	Iberia.	1821.	6. 7. H. ♀.	
87 pinnatifida. w. (bl.)	pinnatifid.	—	1817.	5. 6. H. ♀.	Bot. mag. t. 1679.
88 multifida. R.S. (bl.)	multifid-leaved.	Siberia.	1748.	6. 8. H. ♀.	Lodd. bot. cab. t. 911.
89 caucasica. R.S. (wh. bl.)	Caucasean.	Caucasus.	1815.	— H. ♀.	Allion. ped. 1. t. 46. f. 3.
90 taurea. w. (bl.)	Taurian.	Tauria.	—	— H. ♀.	Eng. bot. v. 11. t. 765.
91 Allioni. R.S. (bl.)	shining-leaved.	S. Europe.	1748.	— H. ♀.	
92 officinalis. R.S. (bl.)	officinal.	Britain.	4. 7. H. ♀.	
β rigidula. Gr. (bl.)	rigid-leaved.	Scotland.	— H. ♀.	
γ albiloba. (wh.)	white-flowered.	Britain.	— H. ♀.	
93 Mulleriæ. R.S. (bl.)	Muller's.	Styria.	1825.	5. 7. H. ♀.	
94 Tournefortii. R.S. (bl.)	Tournefort's.	S. France.	1821.	5. 6. H. ♀.	
95 Chaixi. R.S. (bl.)	Chaix's.	S. Europe.	1825.	5. 7. H. ♀.	Bauh. hist. 3. p. 287. ic.
96 prostrata. R.S. (bl.)	trailing.	Germany.	1774.	5. 6. H. ♀.	
97 pilosa. R.S. (bl.)	hairy.	Bohemia.	1819.	— H. ♀.	
98 acutiflora. R.S. (li.)	acute-flowered.	France.	1821.	— H. ♀.	Pohl tent. fl. b. p. 15. f. 1.
99 pliata. R.S. (bl.)	plaited-leaved.	Bohemia.	1823.	— H. ♀.	Hoff. f. port. t. 57.
100 micrantha. R.S. (wh. pu.)	soft-leaved. mollis. Zea.	Portugal.	—	— H. ♀.	
101 latifolia. R.S. (bl.)	broad-leaved.	Austria.	1748.	6. 7. H. ♀.	
102 Teucrium. R.S. (bl.)	saw-leaved.	Europe.	1596.	6. 8. H. ♀.	Swt. br. fl. gar. t. 23.
103 crinita. R.S. (bl.)	very hairy.	Hungary.	1826.	— H. ♀.	Lodd. bot. cab. t. 425.
104 lutetiæna. R.S. (pa. bl.)	Portugal.	Portugal.	—	— H. ♀.	
105 dentata. R.S. (bl.)	tooth-leaved.	S. Europe.	1816.	— H. ♀.	
106 Schmidtii. R.S. (bl.)	Schmidt's.	Bohemia.	1797.	6. 7. H. ♀.	
paniculata. w. non Pall.					
107 peduncularis. R.S. (bl.)	long-peduncled.	Caucasus.	1825.	— H. ♀.	
108 petraea. R.S. (pa. bl.)	Caucasean rock.	—	1821.	— H. ♀.	
109 umbræa. M.B. (bl.)	shady wood.	Tauria.	1827.	— H. ♀.	
110 Chamædrys. R.S. (bl.)	Germanander.	Britain.	5. 8. H. ♀.	M. B. cent. ros. 1. t. 7.
111 stolonifera. R.S. (bl.)	creeping.	N. America.	1827.	— H. ♀.	Eng. bot. v. 9. t. 623.
112 divaricata. R.S. (bl.)	divaricate.	Sweden.	1828.	— H. ♀.	
113 Vestiæna. R.S. (bl.)	Vest's.	—	1820.	— H. ♀.	
114 maxima. R.S. (pa. bl.)	large.	Iberia.	1824.	— H. ♀.	
115 urticæfolia. R.S. (pa. bl.)	Nettle-leaved.	Austria.	1776.	6. 7. H. ♀.	Buxb. cent. 1. t. 34.
116 montana. R.S. (pa.)	mountain.	Britain.	7. 8. H. ♀.	Jacq. aust. 1. t. 59.
117 plebeia. B.P. (bl.)	plebeian.	N. S. W.	1822.	6. 8. G. ♀.	Eng. bot. v. 11. t. 766.
arguta. R.S.					

118 <i>Brownii.</i> B.P. (<i>bl.</i>) sharp-toothed.	N. S. W.	1822.	6. 8. G. ȝ.	
119 <i>gracilis.</i> B.P. (<i>bl.</i>) slender.	—	—	— G. ȝ.	
120 <i>perfoliata.</i> B.P. (<i>bl.</i>) perfoliate.	—	1815.	7. 8. G. ȝ.	Bot. mag. t. 1936.
121 <i>labiata.</i> B.P. (<i>pa.bl.</i>) labiated.	N. Holland.	1802.	4. 7. F. ȝ.	— t. 1660.
<i>Derwentina.</i> Andr. reposit. t. 531.				
122 <i>mollissima.</i> R.S. (<i>bl.</i>) softest.	—	1819.	6. 8. H. ȝ.	
123 <i>reniformis.</i> Ph. (<i>pa.bl.</i>) kidney-leaved.	Missouri.	1827.	5. 8. H. ȝ.	
124 <i>Waldsteiniana.</i> R.S. (<i>bl.</i>) Waldstein's.	—	1819.	6. 8. H. ȝ.	
125 <i>verna.</i> R.S. (<i>pa.bl.</i>) vernal.	Britain.	—	4. 5. H. ○.	Eng. bot. v. 1. t. 25.
126 <i>digitata.</i> R.S. (<i>bl.</i>) fingered.	Spain.	1805.	6. 7. H. ○.	Reich. ic. t. 36.
127 <i>triphylos.</i> R.S. (<i>bl.</i>) variable-leaved.	Britain.	—	4. 5. H. ○.	Eng. bot. v. 1. t. 26.
128 <i>hederifolia.</i> R.S. (<i>pa.bl.</i>) Ivy-leaved.	—	—	3. 10. H. ○.	— v. 11. t. 784.
129 <i>Cymbalaria.</i> R.S. (<i>wh.</i>) fleshy-leaved.	Levant.	1823.	4. 8. H. ○.	Flor. græc. t. 9.
130 <i>peregrina.</i> R.S. (<i>pa.</i>) knotgrass-leaved.	N. Europe.	1680.	5. 6. H. ○.	Flor. dan. t. 407.
131 <i>depræssa.</i> R.S. (<i>bl.</i>) depressed.	Hungary.	1826.	4. 8. H. ○.	
132 <i>glauca.</i> F.G. (<i>bl.</i>) glaucous.	Greece.	—	— H. ○.	Flor. græc. v. 1. t. 7.
133 <i>amœna.</i> R.S. (<i>bl.</i>) fine blue.	Iberia.	1823.	5. 6. H. ○.	MB. cent. ros. t. 1. f. 18.
134 <i>biloba.</i> R.S. (<i>wh.</i>) two-lobed.	Levant.	1819.	4. 6. H. ○.	Buxb. cent. 1. t. 36.
135 <i>præcox.</i> R.S. (<i>bl.</i>) early-flowered.	Europe.	—	3. 5. H. ○.	Allion. ped. t. 1. f. 1.
136 <i>acinifolia.</i> R.S. (<i>pa.bl.</i>) Basil-leaved.	S. Europe.	1788.	4. 5. H. ○.	Poit. et Turp. par. t. 23.
137 <i>arvensis.</i> R.S. (<i>bl.</i>) wall.	Britain.	—	4. 7. H. ○.	Eng. bot. v. 11. t. 734.
138 <i>agræstis.</i> R.I. (<i>va.</i>) field.	—	—	1. 12. H. ○.	Reich. ic. C. 3. t. 277.
<i>puclchella.</i> DC. non Bernhard. <i>versicolor.</i> Fries.				
139 <i>opaca.</i> R.I. (<i>pa.bl.</i>) dull-leaved.	Britain.	—	— H. ○.	Reic. ic. C. 3. p. 67. t. 278.
140 <i>polita.</i> R.I. (<i>bl.</i>) polished.	—	—	— H. ○.	— C. 3. p. 45. t. 246.
141 <i>didyma.</i> T.N. (<i>bl.</i>) toothed-calyxed.	Naples.	1827.	— H. ○.	
142 <i>pulchella.</i> R.S. (<i>wh.ro.</i>) neat.	—	1823.	3. 6. H. ○.	
143 <i>Buxbaumii.</i> s.s. (<i>bl.</i>) Buxbaum's.	S. Europe.	1820.	— H. ○.	Ten. fl. nap. t. 1.
144 <i>filiformis.</i> s.s. (<i>bl.</i>) long-stalked.	Levant.	1780.	5. H. ○.	Buxb. cent. 1. t. 40. f. 1.
145 <i>persica.</i> s.s. (<i>bl.</i>) Persian.	Persia.	1816.	1. 12. H. ○.	
DIPLOPHYLLUM. S.S. DIPLOPHYLLUM. Diandria Monogynia. S.S.				
veronicaeformis. s.s. (<i>bl.</i>) Speedwell-like.	Caucasus.	1813.	4. 6. H. ○.	Linn. trans. v. 11. t. 31.
<i>Veronica crista galli.</i> L.T.				
HEMIMERIS. L. HEMIMERIS. Diandria Monogynia.				
montana. L. (<i>re.</i>) mountain.	C. B. S.	1816.	6. 8. G. ȝ.	Lam. ill. t. 532.

ORDO CLIII.

LABIATÆ. *Brown prodr.* 1. p. 499.Tribus I. MENTHOIDEÆ. *Bentham in Bot. reg. fol.* 1282.

LYCOPUS. R.S.	WATER-HOREHOUND.	Diandria Monogynia. L.		
1 <i>australis.</i> B.P. (<i>wh.</i>) New Holland.	N. S. W.	1823.	5. 7. G. ȝ.	
2 <i>europeus.</i> R.S. (<i>wh.</i>) Europæan.	Britain.	—	7. 8. H. ȝ.	Eng. bot. v. 16. t. 1105.
3 <i>intermedius.</i> (<i>wh.</i>) intermediate.	Europe.	1816.	— H. ȝ.	
4 <i>exaltatus.</i> R.S. (<i>wh.</i>) tall.	Italy.	1739.	— H. ȝ.	Flor. græc. 1. t. 12.
5 <i>virginicus.</i> R.S. (<i>wh.</i>) Virginian.	N. America.	1760.	8. 9. H. ȝ.	
6 <i>sinuatus.</i> E.C. (<i>wh.</i>) sinuate-leaved.	—	1812.	— H. ȝ.	
ISA'NTHUS. M.	ISA'NTHUS.	Didynamia Gymnospermia. S.S.		
cæruleus. M. (<i>bl.</i>) blue.	N. America.	1732.	6. 8. H. ♂.	Dill. elt. t. 285. f. 369.
<i>Trichostema brachiata.</i> L.				
AUDIBERTIA. B.R. AUDIBERTIA. Didynamia Gymnospermia.				
pusilla. B.R. (<i>pu.</i>) small.	Corsica.	1829.	5. 8. H. ȝ.	
<i>Thymus parviflorus.</i> Requien.				
ME'NTHA. L.	MINT.	Didynamia Gymnospermia. L.		
1 <i>viridis.</i> w. (<i>pu.</i>) spear.	Britain.	—	8. H. ȝ.	Eng. bot. v. 34. t. 2424.
2 <i>levigata.</i> W.en. (<i>pu.</i>) smooth-leaved.	Germany.	—	6. 8. H. ȝ.	
3 <i>balsamea.</i> W.en. (<i>pu.</i>) Balsam-scent'd.	Italy.	1804.	7. 8. H. ȝ.	
4 <i>crispata.</i> W.en. (<i>pu.</i>) crumpled.	Europe.	1807.	— H. ȝ.	
5 <i>piperita.</i> E.B. (<i>pu.</i>) pepper.	England.	—	8. 9. H. ȝ.	Eng. bot. v. 10. t. 687.
6 <i>niliaca.</i> S.S. (<i>pu.</i>) Egyptian.	Egypt.	1796.	7. 8. H. ȝ.	Jacq. vind. 3. t. 87.
7 <i>tenuis.</i> M. (<i>wh.</i>) slender.	N. America.	1826.	— H. ȝ.	

LABIATÆ.

8 sylvestris. E.B. (<i>pu.</i>) wild.	Britain.	7. . H. 2.	Eng. bot. v. 10. t. 686.
9 nemorosa. W.en. (<i>pu.</i>) wood.	—	— H. 2.	Flor. dan. v. 84.
10 hirta. W.en. (<i>pu.</i>) hairy.	Europe.	— H. 2.	Rumph. amb. 6. t. 16.
11 pubescens. W.en. (<i>pu.</i>) pubescent.	—	— H. 2.	
12 Auricularia. w. (<i>hi.</i>) Indian.	E. Indies. 1796.	— S. 2.	
13 crista. w. (<i>pu.</i>) curled.	Siberia. 1640.	— H. 2.	
14 pyramidalis. T.N. (<i>pu.</i>) pyramidal.	Naples. 1824.	— H. 2.	
15 undulata. W.en. (<i>pu.</i>) wave-leaved.	Europe. 1816.	— H. 2.	
16 incana. W.en. (<i>pu.</i>) hoary.	—	— H. 2.	
17 lavandulacea. W.en. (<i>pu.</i>) Lavender-leaved.	Spain. 1825.	— H. 2.	
18 capensis. w. (<i>pu.</i>) Cape.	C. B. S. 1816.	— G. 2.	
19 rotundifolia. E.B. (<i>pu.</i>) round-leaved. β variegata. (<i>pu.</i>) variegated.	Britain.	8. 9. H. 2.	Eng. bot. v. 7. t. 446.
20 macrostachya. T.N. (<i>pu.</i>) long-spiked.	Naples. 1820.	— H. 2.	
21 gratissima. w. (<i>pu.</i>) oblong-leaved.	Britain.	7. 8. H. 2.	
22 citrata. s.s. (<i>pu.</i>) Bergamot. odorata. E.B.	England.	— H. 2.	Eng. bot. v. 15. t. 1025.
23 hirsuta. E.B. (<i>pu.</i>) hairy water.	Britain.	7. 9. H.w. 2.	Eng. bot. v. 7. t. 447.
24 acutifolia. E.B. (<i>pu.</i>) acute-leaved.	England.	9. H.w. 2.	— v. 34. t. 2415.
25 sativa. E.B. (<i>re.</i>) tall red.	—	8. 9. H.w. 2.	— v. 7. t. 448.
26 rubra. E.B. (<i>re.</i>) common red.	Britain.	9. H.w. 2.	— v. 20. t. 1413.
27 gentilis. E.B. (<i>re.</i>) bushy red.	—	7. 8. H.w. 2.	— v. 30. t. 118.
28 gracilis. H.K. (<i>re.</i>) narrow-leaved.	—	— H.w. 2.	— v. 7. t. 449.
29 palustris. S.M. (<i>pu.</i>) marsh.	—	9. H.w. 2.	
30 villosa. S.M. (<i>pu.</i>) horse.	—	7. 8. H.w. 2.	Sole's Mints. t. 17.
31 pratensis. S.M. (<i>pu.</i>) meadow.	—	— H.w. 2.	
32 rivalis. S.M. (<i>pu.</i>) river.	—	— H.w. 2.	
33 paludosa. S.M. (<i>pu.</i>) brook.	—	9. H.w. 2.	
34 arvensis. E.B. (<i>li.</i>) corn.	—	7. 9. H. 2.	Eng. bot. v. 30. t. 2119.
35 præcox. S.M. (<i>li.</i>) early-flowering.	—	6. H. 2.	Sole's Mints. ic.
36 agrætis. E.B. (<i>li.</i>) field.	—	7. 8. H. 2.	Eng. bot. v. 30. t. 2120.
37 austriaca. s.s. (<i>pu.</i>) Austrian.	Austria. 1818.	— H. 2.	
38 dentata. w. (<i>pu.</i>) toothed.	Germany. 1816.	— H. 2.	
39 canadensis. w. (<i>pu.</i>) Canadian.	N. America. 1801.	— H. 2.	
40 borealis. M. (<i>li.</i>) northern.	—	1824.	
41 elegans. s.s. (<i>pu.</i>) elegant.	S. Europe. 1828.	— H. 2.	
42 badensis. s.s. (<i>pu.</i>) Baden.	Italy. —	— H. 2.	
43 rugosa. Roth. (<i>pu.</i>) rugged-leaved.	Germany. 1824.	— H. 2.	
44 glabrata. s.s. (<i>pu.</i>) smooth.	Egypt. 1802.	— H. 2.	
45 Pulegium. s.s. (<i>li.</i>) Pennyroyal.	Britain.	8. 9. H. 2.	Eng. bot. v. 15. t. 1026.
46 cervina. s.s. (<i>wh.</i>) Hyssop-leaved.	S. France. 1648.	6. 8. H. 2.	Moris. s. p. t. 7. f. 7.
COLEBRO'OKIA. S.S. COLEBRO'OKIA. Didynamia Gymnospermia.			
1 oppositifolia. D.P. (<i>wh.</i>) opposite-leav'd.	Nepaul. 1816.	6. 9. S. 2.	Sm. exot. bot. t. 115.
2 ternifolia. s.s. (<i>wh.</i>) three-leaved.	E. Indies. —	— S. 2.	Arduin. spec. 2. t. 13.
PERILLA. W. PERILLA. Didynamia Gymnospermia. L.			
1 ocyoides. w. (<i>wh.</i>) Basil-leaved.	India. 1770.	7. 8. H. 2.	
2 fruticosa. D.P. (<i>wh.</i>) frutescent.	Nepaul. 1818.	— S. 2.	Bot. mag. t. 2560.
ELSHO'LTZIA. S.S. ELSHO'LTZIA. Didynamia Gymnospermia. W.			
1 cristata. s.s. (<i>li.</i>) crested.	Siberia. 1789.	5. 7. H. 2.	DC. pl. rar. gen. t. 8.
2 ocyoides. s.s. (<i>li.</i>) Basil-like.	E. Indies. 1823.	— S. 2.	
APHANOCHILUS. B.R. APHANOCHILUS. Didynamia Gymnospermia.			
1 blandus. (<i>wh.</i>) agreeable.	Nepaul. 1820.	10. 12. G. 2.	
Mentha blanda. DC.			
2 incisa. B.R. (<i>wh.</i>) cut-leaved.	—	1818. 6. 10. G. 2.	
Mentha blanda. H.C. non DC.			
DYSOPHYLLA. B.R. DYSOPHYLLA. Didynamia Gymnospermia.			
1 quadrifolia. B.R. (<i>li.</i>) four-leaved.	Nepaul. 1818.	6. 8. G. 2.	
2 stellata. B.R. (<i>li.</i>) starry-leaved.	—	1816. — G. 2.	
3 verticillata. B.R. (<i>li.</i>) whorl-leaved.	—	1828. — G. 2.	
4 pumila. B.R. (<i>li.</i>) dwarf.	—	— G. 2.	
Mentha verticillata. B.M. pumila. Graham.			
POGOSTEMON. S.S. POGOSTEMON. Didynamia Gymnospermia. S.S.			
plectranthoides. s.s. (<i>w.</i>) Plectanthrus-like.	1818.	6. 8. S. 2.	Desf. m. mus. 2. t. 6.

Tribus II. SATUREINEÆ. *Benth. in Bot. reg.*

BYSTROP'OGON. S.S. BYSTROP'OGON. Didynamia Gymnospermia. S.S.

1	origanifolius. s.s.	(<i>li.</i>) Marjoram-leav'd. Teneriffe.	1815.	7. 8. G. h.
2	plumosus. s.s.	(<i>li.</i>) woolly-flower'd. Canaries.	1779.	6. 7. G. h.
3	canariensis. s.s.	(<i>li.</i>) Canary.	1714.	6. 8. G. h.
4	punctatus. s.s.	(<i>li.</i>) cluster-flower'd. Madeira.	1775.	7. 9. G. h.

PYCNA'NTHEMUM. M. PYCNA'NTHEMUM. Didynamia Angiospermia. S.S.

1	linifolium. s.s.	(<i>w.</i>) Flax-leaved. N. America.	1739.	7. 8. H. 4.
2	lanceolatum. s.s.	(<i>w.</i>) spear-leaved.	1812.	— H. 4.
3	aristatum. s.s.	(<i>w.</i>) awned.	1752.	8. H. 4.
4	verticillatum. s.s.	(<i>w.</i>) whorl-leaved.	1816.	7. 8. H. 4.
5	Monardella. s.s.	(<i>li.</i>) Monarda-like.	—	— H. 4.
6	incanum. s.s.	(<i>bh.</i>) hoary.	1732.	7. 10. H. 4.
				Dill. elt. t. 74. f. 85.

SATUR'EJA. S.S. SAVORY. Didynamia Gymnospermia. L.

1	approximata. s.s.	(<i>li.</i>) bundle-leaved. Sicily.	1825.	5. 7. F. h.
2	spinosa. F.G.	(<i>w.</i>) spiny. Crete.	1827.	— G. h.
3	viminea. s.s.	(<i>li.</i>) Pennyroyal tree. Jamaica.	1783.	— S. h.
4	montana. s.s.	(<i>li.</i>) winter. S. Europe.	1562.	6. 7. H. h.
5	europæstris. s.s.	(<i>w.</i>) rock. Carnioia.	1798.	— H. 4.
6	obovata. s.s.	(<i>li.</i>) obovate-leaved. Spain.	1824.	— H. h.
7	tenuifolia. s.s.	(<i>li.</i>) slender-leaved. Italy.	1820.	— H. h.
8	Julianna. s.s.	(<i>li.</i>) linear-leaved.	1596.	5. 9. F. h.
9	græca. s.s.	(<i>li.</i>) Grecian.	1759.	6. 7. F. 4.
10	hirsuta. s.s.	(<i>li.</i>) hairy.	1824.	— F. 4.
11	Teneriffæ. s.s.	(<i>pu.</i>) Teneriffe.	1815.	— F. 4.
12	hortensis. s.s.	(<i>w.</i>) summer. Italy.	1652.	6. 8. H. ⊖.
13	capitata. w.	(<i>li.</i>) ciliated.	1596.	6. 10. H. h.
14	nervosa. s.s.	(<i>li.</i>) nerved.	1824.	— H. h.
15	Thymbra. s.s.	(<i>li.</i>) whorl-flowered. Candia.	1640.	5. 7. G. h.
16	congesta. s.s.	(<i>li.</i>) close-flowered.	1825.	— G. h.

THYMUS. S.S. THYME. Didynamia Gymnospermia. L.

1	villus. s.s.	(<i>li.</i>) hairy.	Portugal.	1759.	6. 7. F. h.
2	cephalotus. s.s.	(<i>li.</i>) great headed.	—	—	7. 8. H. h.
3	capitatus. L.en.	(<i>li.</i>) headed.	1823.	—	H. h.
4	odoratissimus. s.s.	(<i>li.</i>) very sweet.	Tauria.	—	H. h.
5	acicularis. s.s.	(<i>li.</i>) needle-shaped.	Hungary.	1806.	6. 8. H. h.
6	hirsutus. s.s.	(<i>li.</i>) hairy.	—	1816.	— H. h.
7	Zygis. s.s.	(<i>w.</i>) Spanish.	Spain.	1771.	8. H. h.
8	Marschallianus. s.s.	(<i>li.</i>) Marschall's.	Russia.	1815.	6. 8. H. h.
9	glabrescens. L.en.	(<i>li.</i>) smooth.	S. Europe.	1823.	— H. 4.
10	angustifolius. s.s.	(<i>li.</i>) narrow-leaved.	Europe.	— H. h.
11	Serpillum. s.s.	(<i>li.</i>) wild or mother of. Britain.	— H. h.
12	exserens. s.s.	(<i>li.</i>) exserted.	— H. h.
13	citriodorus. L.en.	(<i>li.</i>) Lemon.	Europe.	— H. h.
14	collinus. s.s.	(<i>li.</i>) hill.	Tauria.	1823.	— H. h.
15	montanus. W.K.	(<i>li.</i>) mountain.	Hungary.	1800.	6. 7. H. h.
16	nummularius. M.B.	(<i>li.</i>) moneyswort.	Caucasus.	1818.	— H. h.
17	pannonicus. W.en.	(<i>li.</i>) Hungarian.	Hungary.	—	— H. h.
18	lanuginosus. s.s.	(<i>li.</i>) woolly.	Europe.	6. 8. H. h.
19	tomentosus. s.s.	(<i>li.</i>) tomentose.	Spain.	1816.	— H. h.
20	albicans. s.s.	(<i>li.</i>) white-leaved.	Portugal.	—	— H. h.
21	adscendens. L.en.	(<i>li.</i>) ascending.	S. Europe.	— H. h.
22	lucidus. s.s.	(<i>li.</i>) shining-leaved.	1816.	— H. h.
23	vulgäris. s.s.	(<i>li.</i>) common.	S. Europe.	1548.	5. 8. H. h.
	β latifolius.	(<i>li.</i>) broad-leaved.	—	—	— H. h.
24	Masticina. s.s.	(<i>li.</i>) Mastick.	Spain.	1596.	7. 9. F. h.
25	elongatus. L.en.	(<i>li.</i>) elongated.	1816.	— F. h.
26	graveolens. F.G.	(<i>li.</i>) strong-scented.	Levant.	1823.	— F. h.
27	Tragoricanum. s.s.	(<i>li.</i>) goat's.	Candia.	1640.	5. 6. F. h.
28	filiformis. s.s.	(<i>li.</i>) filiform.	Minorca.	1770.	6. 7. G. h.
29	fruticosus. s.s.	(<i>li.</i>) frutescent.	Sicily.	1823.	— H. h.
30	Teneriffæ. s.s.	(<i>li.</i>) Teneriffe.	Teneriffe.	1828.	— F. h.
31	incanum. s.s.	(<i>w.</i>) hoary.	Levant.	—	— F. 4.
32	croaticus. P.S.	(<i>li.</i>) oval-leaved.	Hungary.	1802.	7. 8. H. h.
	Piperella. W.K. non L.	—	—	—	Flor. græc. t. 577.
33	ericæfolius. W.en.	(<i>li.</i>) Heath-leaved.	Spain.	1806.	6. 8. H. h.

LABIATÆ.

ORIGANUM. S.S.

MARJORAM.

Didynamia Gymnospermia. L.

1 vulgare. s.s.	(ro.) common.
2 humile. s.s.	(wh.) dwarf.
3 virens. s.s.	(ro.) green.
4 heracleoticum. s.s.	(li.) winter, sweet.
5 craticum. s.s.	(li.) Cretan.
6 megastachyum. L.en.	(li.) large-spiked.
7 macrostachyon. s.s.	(li.) long-spiked.
8 hirtum. L.en.	(wh.) hairy.
9 oblongatum. L.en.	(wh.) oblong-spiked.

10 smyrnæum. L. (wh.) Smyrna.

11 Onites. s.s. (wh.) pot.

12 Mari. w. (li.) Lavender-scented. Crete.

13 Majoranoïdes. w. (wh.) shrubby sweet.

14 Majorana. s.s. (wh.) knotted.

15 indicum. s.s. (wh.) Indian knotted. E. Indies.

16 salvifolium. Roth. (li.) Sage-leaved.

17 pallidum. P.S. (pa.) pale.

18 stoloniferum. F. (li.) creeping.

19 Tournefortii. s.s. (ro.) Tournefort's.

20 sipylycum. s.s. (ro.) Mount Sipylus.

21 Dictamnus. s.s. (li.) Dittany of Crete. Candia.

22 ægyptiacum. s.s. (li.) Egyptian.

LOPHANTHUS. B.R. LOPHANTHUS. Didynamia Gymnospermia.

1 chinensis. B.R. (ye.) Mint-leaved.

Hyssopus Lophanthus. s.s.

2 nepetoides. B.R. (ye.) square-stalked. N. America.

3 scrophulariifolius. B.R. (re.) Figwort-leaved.

4 anisatus. B.R. (bl.) Anise-like.

5 urticifolius. B.R. (pu.) Nettle-leaved.

6 multifidus. B.R. (wh.) multifid.

Nepeta multifida. w.

HYSSOPUS. B.R. HYSSOP. Didynamia Gymnospermia.

1 officinalis. s.s. (bl.) common.

2 orientalis. w. (bl.) narrow-leaved.

angustifolius. M.B.

3 discolor. F. (bl.) two-coloured.

4 septemfidus. E. seven-cleft.

5 septemcrenatus. E. seven-notched.

WESTRÍNGIA. S.S. WESTRÍNGIA. Didynamia Gymnospermia. S.S.

1 rosmariniformis. B.P. (w.) Rosemary like.

2 Dampieri. B.P. (wh.) Dampier's.

3 angustifolia. B.P. (wh.) narrow-leaved.

4 longifolia. B.P. (wh.) long-leaved.

N. S. W. 1791. 5. 8. G. h.

N. Holland. 1803. 5. 7. G. h.

V. Diem. Isl. 1823. G. h.

N. S. W. 1826. G. h.

Tribus III. AJUGOIDEÆ. Benth. in Bot. reg.

LEUCOSCEPTRUM. S.S. LEUCOSCEPTRUM. Didynamia Gymnospermia.

cânum. s.s. (wh.) white-flowered. Nepaul. 1825. 7. 9. F. 4.

TEUCRIUM. S.S. GERMANDER. Didynamia Gymnospermia. L.

1 fruticans. w. (bl.) narrow-l'd. shrubby. Spain. 1640. 6. 9. H. h.

2 latifolium. L. (li.) broad-l'd. shrubby.

3 heterophyllum. s.s. (li.) various-leaved.

4 Laxmanni. s.s. (ye. pu.) Laxmann's.

5 alpestre. F.G. (wh.) alpine.

6 brevifolium. s.s. (li.) short-leaved.

7 asiaticum. s.s. (bh.) Asiatic.

8 cubense. s.s. (li.) Cuba.

9 Nissolianum. s.s. (pu.) Nissole's.

10 levigatum. s.s. (li.) smooth.

11 campanulatum. s.s. (bh.) bell-flowered.

12 régium. s.s. (pu.) royal.

13 multiflorum. s.s. (pu.) many-flowered.

14 lúcidum. s.s. (re.) shining.

15 Chamædrys. s.s. (pu.) wall.

16 Scordium. s.s. (pu.) water.

Buenos Ayres. 1829. G. 4.

Levant. 1728. 7. 8. H. 4.

Spain. 1699. 5. 10. F. h.

.... 1731. 7. 9. H. 4.

S. Europe. 1730. 6. 9. H. 4.

England. 5. 8. H. 4.

.... 1752. 6. 7. F. 4.

Bocca. mus. t. 117.

Flor. græc. t. 532.

Eng. bot. v. 10. t. 680.

Pluk. alm. t. 65. f. 1.

Andr. deposit. t. 214.

Jacq. vind. 3. t. 41.

Jacq. obs. 2. t. 30.

Moris. s. t. 22. f. 19.

Flor. græc. t. 527.

Bot. mag. t. 245.

Flor. græc. t. 528.

Jacq. vind. 3. t. 528.

Jacq. obs. 2. t. 30.

Flor. græc. t. 532.

Eng. bot. v. 12. t. 828.

Bot. mag. t. 2605.

Eng. bot. v. 16. t. 1143.

Hoff. et Lk. fl. p. 1. t. 9.

Lobel. ic. 492.

Schk. hand. 2. t. 164.

Flor. græc. t. 571.

_____. t. 572.

Moris. s. 11. t. 3. f. 1.

Jacq. vind. 2. t. 182.

Jacq. vind. 1. t. 68.

Herm. parad. t. 106.

Bot. reg. t. 1282.

Gmel. sib. 3. t. 55.

Jacq. aust. 3. t. 254.

Bot. mag. t. 2299.

17 scordioïdes. s.s. (<i>pu.</i>) likened.	S. Europe.	1816.	7. 8. H. 4.		
18 bracteatum. s.s. (<i>pu.</i>) bracted.	Barbary.	1828.	— G. 4.	Desf. atl. 2. t. 120.	
19 spinosum. s.s. (<i>wh.</i>) thorny.	Spain.	1640.	5. 6. H. ½.	Flor. græc. t. 539.	
20 Bôtrys. s.s. (<i>pu.</i>) cut-leaved.	S. Europe.	1633.	7. 9. H. ⊖.	Ger. emac. 525. f. 2.	
21 trifidum. s.s. (<i>re.</i>) trifid-leaved.	C. B. S.	1791.	6. 8. G. ½.	Barrel. rar. t. 1080.	
22 angustissimum. s.s. (<i>re.</i>) narrow-leaved.	Spain.	1818.	— H. ½.	— t. 1092.	
23 pùmulum. s.s. (<i>re.</i>) small.	—	1816.	7. 8. F. ½.	Cavan. ic. 2. t. 118.	
24 Libanitis. s.s. (<i>ye. bh.</i>) Spanish.	Spain.	1824.	6. 8. H. 4.	Jacq. aust. 5. t. 417.	
25 thymifòlium. s.s. (<i>pu.</i>) Thyme-leaved.	—	—	6. 10. F. ½.	Flor. græc. t. 534.	
26 supinum. s.s. (<i>wh.</i>) procumbent.	Austria.	1752.	— H. ½.	— t. 535.	
27 montanum. s.s. (<i>st.</i>) dwarf mountain.	S. Europe.	1710.	7. 10. F. ½.	— t. 537.	
28 Pòlium. s.s. (<i>wh.</i>) Poley.	—	1562.	7. 9. F. ½.	Col. ecphr. 1. t. 67.	
29 cuneifòlium. F.G. (<i>wh.</i>) wedge-leaved.	Levant.	1828.	6. 8. F. ½.	Flor. græc. t. 536.	
30 Pseudohyssòpus. s.s. (<i>wh.</i>) Hyssop-leaved.	Italy.	1804.	6. 7. H. ½.	Barrel. rar. t. 1096.	
31 capitatum. s.s. (<i>ro.</i>) round-headed.	Spain.	1731.	7. 8. F. ½.	— t. 1048.	
32 lusitanicum. s.s. (<i>wh.</i>) Portugal.	Portugal.	1816.	— F. ½.	— t. 1083.	
33 pycnophyl'lum. s.s. (<i>wh.</i>) clustered.	Spain.	—	— H. 4.	Cavan. ic. 2. t. 117.	
34 valentinum. s.s. (<i>wh.</i>) Valencia.	—	1826.	— F. ½.	Barrel. rar. t. 1073.	
35 gnaphalodes. s.s. (<i>re.</i>) woolly-calyx'd.	—	—	7. 9. F. ½.	Bot. mag. t. 2013.	
36 aùreum. s.s. (<i>ye.</i>) golden Poley.	S. Europe.	1731.	6. 7. F. ½.	Bot. reg. t. 1255.	
37 flavescens. P.S. (<i>ye.</i>) yellow Poley.	—	—	7. 9. F. ½.	Schkuhr. handb. 160.	
38 hyrcanicum. s.s. (<i>pu.</i>) Betony-leaved.	Persia.	1763.	8. 10. H. 4.	Flor. græc. t. 531.	
39 inflatulum. s.s. (<i>pu.</i>) thick-spiked.	Jamaica.	1778.	— S. 2.	Jacq. schœn. 3. t. 358.	
40 orchioïdes. B.R. (<i>bh.</i>) Orchis-flowered.	Chile.	1826.	— F. ½.	Eng. bot. v. 22. t. 1543.	
heterophyl'l'a. Cavan. icon. 6. t. 577. non L'Herit.	—	—	—	Bot. mag. t. 1114.	
41 virginicum. s.s. (<i>pu.</i>) Virginian.	N. America.	1768.	5. 6. H. 4.	Flor. græc. t. 533.	
42 canadénse. s.s. (<i>pu.</i>) Canadian.	—	—	8. 9. H. 4.	Jacq. vind. 1. t. 94.	
43 Arduini. s.s. (<i>wh.</i>) Arduini's.	Levant.	1827.	— H. 4.	Desf. atl. 2. t. 117.	
44 abutiloides. s.s. (<i>ye.</i>) Mulberry-leaved.	Madeira.	1777.	4. 5. G. ½.	Park. theat. 17. f. 2.	
45 Scorodònia. s.s. (<i>st.</i>) wood sage.	Britain.	—	7. H. 4.	Flor. græc. t. 529.	
46 betonicum. s.s. (<i>pu.</i>) hoary.	Madeira.	1775.	5. 8. G. ½.	Flor. græc. t. 529.	
47 flàvum. s.s. (<i>st.</i>) yellow-flowered.	S. Europe.	1640.	7. 9. H. ½.	Bot. mag. t. 1279.	
48 massiliénse. s.s. (<i>li.</i>) sweet scented.	France.	1731.	6. 7. H. ½.	Bot. mag. t. 2448.	
49 resupinátum. s.s. (<i>st.</i>) resupinate.	Barbary.	1801.	7. 8. H. ⊖.	Lam. ill. t. 515.	
50 Mârum. s.s. (<i>pu.</i>) Cat-thyme.	Spain.	1640.	7. 9. F. ½.	TRICHOSTEMA. R.S. AMETHY'STEA. Diandria Monogynia. L.	
51 subspindosum. s.s. (<i>pu.</i>) Minorca.	Minorca.	1816.	— G. ½.	TRICHOSTEMA. S.S. TRICHOSTEMA. Didynamia Gymnospermia. L.	
52 quadratulum. s.s. (<i>ro.</i>) many-branched.	Levant.	1828.	— F. ½.	Didynamia Gymnospermia. L.	
53 créticum. s.s. (<i>ro.</i>) Cretan.	Crete.	1825.	— G. ½.	Didynamia Gymnospermia. L.	
54 pyrenaicum. s.s. (<i>pu.</i>) Pyrenean.	Pyrenees.	1731.	6. 8. H. 4.	Didynamia Gymnospermia. L.	
55 orientale. s.s. (<i>bl.</i>) great-flowered.	Levant.	1752.	7. 8. H. 4.	Didynamia Gymnospermia. L.	
AMETHY'STEA. R.S. AMETHY'STEA. Diandria Monogynia. L.	—	—	—	Didynamia Gymnospermia. L.	
coerulea. R.S. (<i>bl.</i>) blue-flowered.	Siberia.	1759.	6. 7. H. ⊖.	Didynamia Gymnospermia. L.	
créticula. R.S. (<i>bl.</i>) blue-flowered.	Siberia.	1759.	6. 7. H. ⊖.	Didynamia Gymnospermia. L.	
pyrenaicum. s.s. (<i>pu.</i>) Pyrenean.	Pyrenees.	1731.	6. 8. H. 4.	Didynamia Gymnospermia. L.	
orientale. s.s. (<i>bl.</i>) great-flowered.	Levant.	1752.	7. 8. H. 4.	Didynamia Gymnospermia. L.	
TRICHOSTEMA. R.S. TRICHOSTEMA. Didynamia Gymnospermia. L.	—	—	—	Didynamia Gymnospermia. L.	
1 dichótoma. w.	(<i>bl.</i>) Marjoram-leav'd.	N. America.	1759.	6. 7. H. ⊖.	Didynamia Gymnospermia. L.
2 lineáris. s.s.	(<i>bl.</i>) linear-leaved.	—	— H. ⊖.	Didynamia Gymnospermia. L.	
AJUGA. S.S. BUGLE. Didynamia Gymnospermia. L.	—	—	—	Didynamia Gymnospermia. L.	
1 Iva. s.s. (<i>ye.</i>) musky.	S. Europe.	1759.	7. 8. H. ⊖.	Flor. græc. t. 525.	
2 réptans. s.s. (<i>bl.</i>) common.	Britain.	—	5. 6. H. 4.	Eng. bot. v. 7. t. 480.	
α cærulea.	(<i>bl.</i>) blue-flowered.	—	— H. 4.		
β rubra.	(<i>pu.</i>) red-flowered.	—	— H. 4.		
γ álba.	(<i>wh.</i>) white-flowered.	—	— H. 4.		
3 álpina. s.s. (<i>bl.</i>) Alpine.	England.	—	5. 7. H. 4.	Eng. bot. v. 7. t. 477.	
4 genevénis. s.s. (<i>bl.</i>) Geneva.	Switzerland.	1656.	5. 6. H. 4.	Bull. herb. t. 361.	
5 pyramidális. s.s. (<i>li.</i>) pyramidal.	Britain.	—	— H. 4.	Eng. bot. v. 18. t. 1270.	
6 oblongata. s.s. (<i>bl.</i>) oblong-leaved.	Caucasus.	1829.	— H. 4.	Swt. br. fl. gar. ser. 2.	
7 orientalis. s.s. (<i>bl.</i>) oriental.	Levant.	1732.	— H. 4.	Eng. bot. v. 2. t. 77.	
8 Chamaépitys. s.s. (<i>ye.</i>) ground pine.	England.	—	4. 7. H. ⊖.	Flor. græc. t. 524.	
9 Chia. F.G.	(<i>ye.</i>) Grecian.	Levant.	—	— H. ⊖.	— t. 526.
10 salicifòlia. F.G.	(<i>ye.</i>) Willow-leaved.	—	— H. ⊖.	Burm. zeyl. t. 71. f. 1.	
ANISOMELES. B.P. ANISOMELES. Didynamia Gymnospermia. S.S.	—	—	—	Bot. mag. t. 2071.	
1 ovata. s.s. (<i>vi. wh.</i>) broad-leaved.	E. Indies.	1783.	7. 8. S. ⊖.		
2 malabárica. s.s. (<i>vi.</i>) Malabar.	—	1817.	— S. ½.		

Tribus IV. MONARDEÆ. Benth. in Bot. reg.

MONARDA. R.S. MONARDA. Diandria Monogynia. L.	N. America.	1656.	6. 8. H. 4.	Mill. ic. t. 122. f. 2.
1 fistulosa. R.S. (<i>pu.</i>) fistulous.	—	—	6. 9. H. 4.	Swt. br. fl. gar. t. 98.
2 media. R.S. (<i>da. pu.</i>) purple-bracted.	—	—	—	

LABIATÆ.

3 mollis. R.S. (<i>pa.pu.</i>) soft.	N. America.	6. 7. H. Ȑ.
4 oblongata. R.S. (<i>pa.pu.</i>) long-leaved.	1761.	7. 9. H. Ȑ.
5 altissima. R.S. (<i>li.</i>) tallest.	1818.	— H. Ȑ.
6 affinis. L.en. (<i>li.</i>) affined.	—	— H. Ȑ.
7 menthaefolia. B.M. (<i>pu.</i>) Mint-leaved.	1827.	— H. Ȑ.
8 rugosa. R.S. (<i>wh.</i>) white-flowered.	1761.	— H. Ȑ.
9 purpurea. R.S. (<i>pu.</i>) purple-flowered. <i>fistulosa</i> . var. <i>crimson</i> . B.M.	1789.	6. 8. H. Ȑ.
10 Kalmiana. R.S. (<i>sc.</i>) pubescent-flower'd.	1813.	— H. Ȑ.
11 didyma. R.S. (<i>sc.</i>) Oswego Tea.	1752.	— H. Ȑ.
12 Russelliæna. N. (<i>wh.re.</i>) Russell's.	1823.	— H. Ȑ.
13 clinopodium. Ph. (<i>st.</i>) wild basil-leav'd.	1771.	7. H. Ȑ.
14 gracilis. R.S. (<i>st.</i>) slender.	1824.	7. 8. H. Ȑ.
15 punctata. R.S. (<i>ye.re.</i>) spotted.	1714.	6. 10. F. Ȑ.
16 citriodora. R.S. (<i>li.</i>) Citron-scented. New Spain.	1823.	— F. Ȑ.
BLEPHYLIA. B.R. BLEPHYLIA. Diandria Monogynia.		
1 ciliata. B.R. (<i>bl.</i>) blue-flowered.	N. America. 1798.	7. 8. H. Ȑ.
<i>Monarda ciliata</i> . R.S.		
2 hirsuta. B.R. (<i>li.</i>) hairy.	—	7. 9. H. Ȑ.
ZIZIPHORA. S.S. ZIZIPHORA. Diandria Monogynia. L.		
1 clinopodioides. R.S. (<i>li.</i>) headed.	Siberia. 1799.	7. 8. H. ⊖.
<i>Cùnìa capitata</i> . L.		
2 acinoides. W.en. (<i>ro.</i>) Thyme-leaved.	—	1786. — H. ⊖.
3 media. L.en. (<i>li.</i>) middle.	—	1816. — H. Ȑ.
4 capitata. R.S. (<i>ro.</i>) oval-leaved.	Syria.	1752. — H. ⊖.
5 hispanica. R.S. (<i>li.</i>) Spanish.	Spain.	1759. 6. H. ⊖.
6 tenuior. R.S. (<i>li.</i>) spear-leaved.	Levant.	1752. 6. 7. H. ⊖.
7 taurica. R.S. (<i>li.</i>) Taurian.	Tauria.	1816. 7. 9. H. ⊖.
8 serpyllacea. R.S. (<i>li.</i>) sweet-scented.	Caucasus.	1803. 7. 8. H. Ȑ.
9 dasyantha. R.S. (<i>li.</i>) hairy-flowered.	Siberia.	— 6. 8. H. Ȑ.
<i>Pouschkini</i> . B.M.		
C'UNILA. N. C'UNILA. Diandria Monogynia. L.		
<i>marianna</i> . N. (<i>pu.</i>) Mint-leaved.	N. America. 1759.	7. 9. H. Ȑ.
HEDE'OMA. N. HEDE'OMA. Diandria Monogynia. P.S.		
1 pulegioides. N. (<i>wh.vi.</i>) Pennyroyal-lid.	N. America. 1777.	6. 8. H. ⊖.
2 thymoides. P.S. (<i>bh.</i>) Thyme-leaved.	France.	1699. — H. ⊖.
ROSMAR'INUS. R.S. ROSEMARY. Diandria Monogynia. L.		
1 officinalis. R.S. (<i>pa.bl.</i>) common.	S. Europe. 1548.	1. 4. H. Ȑ.
α <i>angustifolius</i> . (<i>pa.bl.</i>) narrow-leaved.	—	— H. Ȑ.
β <i>latifolius</i> . (<i>pa.bl.</i>) broad-leaved.	—	— H. Ȑ.
γ <i>argenteus</i> . (<i>pa.bl.</i>) silver-striped.	—	— H. Ȑ.
δ <i>aureus</i> . (<i>pa.bl.</i>) gold-striped.	—	— H. Ȑ.
2 chilensis. R.S. (<i>wh.</i>) Chili.	Chili.	1795. 7. G. Ȑ.
SYNA'NDRA. N. SYNA'NDRA. Didynamia Gymnospermia.		
<i>grandiflora</i> . N. (<i>st.</i>) great-flowered.	N. America. 1827.	6. 7. H. Ȑ.

Tribus V. NEPETÆ. *Benth. in Bot. reg.*

LEON'OTIS. S.S.	LION'S-EAR. Didynamia Gymnospermia. S.S.		
1 Leonurus. s.s. (<i>or.</i>) narrow-leaved.	C. B. S. 1712.	10. 12. G. Ȑ.	Bot. mag. t. 478.
<i>Phlomis Leonurus</i> . w.			
2 ovata. s.s. (<i>or.</i>) dwarf shrubby.	—	1713. 6. 7. G. Ȑ.	Mill. ic. 2. t. 162. f. 1.
3 intermedia. s.s. (<i>or.</i>) intermediate.	S. Africa. 1823.	— G. Ȑ.	Bot. reg. t. 850.
4 nepetifolia. s.s. (<i>or.</i>) Catmint-leaved.	E. Indies. 1778.	9. 10. S. ⊖.	— t. 281.
LEUCAS. B.P.	LEUCAS. Didynamia Gymnospermia. S.S.		
1 chinensis. s.s. (<i>wh.</i>) Chinese.	China. 1822.	6. 8. S. Ȑ.	Jacq. ic. 1. t. 111.
2 zeylanica. s.s. (<i>wh.</i>) Ceylon.	E. Indies. 1777.	6. 10. S. ⊖.	
3 linifolia. s.s. (<i>wh.</i>) Flax-leaved.	—	1816. — S. ⊖.	
4 Pluknetii. s.s. (<i>wh.</i>) Pluknet's.	—	1824. — S. ⊖.	
5 cephalotes. s.s. (<i>wh.pu.</i>) large-headed.	—	1811. — S. ⊖.	
6 nuntans. s.s. (<i>wh.</i>) nodding.	—	1826. — S. ⊖.	
7 hirta. s.s. (<i>wh.</i>) hairy.	—	1824. — S. ⊖.	
8 áspera. s.s. (<i>wh.</i>) rough.	Persia.	1823. — S. ⊖.	
9 indica. s.s. (<i>lh.</i>) Indian	E. Indies.	1789. 7. 8. S. ⊖.	
10 urticafolia. s.s. (<i>wh.</i>) Nettle-leaved.	—	1810. — S. ⊖.	

11	martiniensis.	(wh.) West-Indian. <i>Phlomis martinicensis</i> . w.	W. Indies.	1781.	7. 8. S.○.	Jacq. ic. 1. t. 110.
12	biflora. s.s.	(wh.) two-flowered.	E. Indies.	1820.	— S.○.	Burm. zeyl. t. 63. f. 1.
PHL'OMIS. S.S.		JERUSALEM-SAGE.	Didynamia Gymnospermia. L.			
1	fruticosa. s.s.	(ye.) shrubby.	Spain.	1596.	6. 7. H. ½.	Flor. græc. t. 563.
2	ferruginea. s.s.	(ye.) ferruginous.	Naples.	— H. ½.	
3	lanata. s.s.	(ye.) small shrubby.	S. Europe.	1596.	— H. ½.	
4	crética. s.s.	(ye.) Cretan.	Crete.	1823.	— G. ½.	
5	scariosa. s.s.	(ye.) scariose-bracted.	Italy.	—	— F. ½.	
6	virens. s.s.	(ye.) green shrubby.	Levant.	— F. ½.	
7	microphylla. Sie.	(ye.) small-leaved.	—	1827.	— F. ½.	
8	purpurea. s.s.	(pu.) purple.	S. Europe.	1661.	6. 8. F. ½.	Smith spicil. 6. t. 3.
9	italica. s.s.	(pu.) Italian.	Italy.	—	— F. ½.	
10	Nissolia. s.s.	(ye.) Nissole's.	Levant.	1757.	6. 7. F. ¼.	Mill. ic. 2. t. 204.
11	armeniaea. s.s.	(ye.) Armenian.	Armenia.	1829.	— F. ½.	
12	Lychnites. s.s.	(ye.) lamp-wick.	S. Europe.	1658.	6. 8. G. ½.	Bot. mag. t. 999.
13	fluccosa. B.R.	(ye.) woolly.	Egypt.	1828.	8. 11. F. ½.	Bot. reg. t. 1300.
14	salviefolia. p.s.(ye.re.)	Sage-leaved.	—	1822.	6. 9. F. ½.	Jacq. scheenb. 3. t. 359.
15	agraria. Led.	(pu.) field.	Siberia.	1827.	— H. ¼.	
16	Sâmia. s.s.	(pu.) Samian.	N. Africa.	1714.	6. 7. H. ¼.	Flor. græc. t. 564.
17	Russelliana. Lag.	(ye.) Russell's. <i>lunaria</i> β <i>Russelliana</i> . B.M. 2542.	Levant.	1821.	— H. ¼.	Russ. alep. 2. t. 16.
18	Hérba-vénti. s.s.	(pu.) rough-leaved.	S. Europe.	1596.	7. 9. H. ¼.	
19	pungens. s.s.	(pu.) pungent.	Levant.	1818.	6. 8. H. ¼.	Swt. br. fl. gar. t. 33.
20	crinita. s.s.	(or.) white woolly.	Spain.	1829.	— H. ¼.	Cavan. icon. 3. t. 247.
21	alpina. s.s.	(pu.) Alpine.	Siberia.	1802.	6. 9. H. ¼.	Pall. act. pet. 2. t. 13.
22	tuberosa. s.s.	(pu.) tuberous-root'd.	—	1759.	6. 10. H. ¼.	Bot. mag. t. 1555.
23	laciniatâ. s.s.	(ye.) jagged-leaved.	Levant.	1731.	7. H. ¼.	Swt. br. fl. gar. t. 24.
BALL'OTA. S.S.		STINKING HOEHOUND.	Didynamia Gymnospermia. L.			
1	vulgâris. s.s.	(pu.) common.	Europe.	—	7. 9. H. ¼.	
2	fœtida. L.en.	(pu.) strong-scented.	Britain.	—	— H. ¼.	Eng. bot. v. 1. t. 46.
<i>a nigra.</i>	(pu.) purple-flowered.	—	—	— H. ¼.		
<i>β alba.</i>	(wh.) white-flowered.	—	—	— H. ¼.		
BERING'ERIA. B.R.	BERING'ERIA.	Didynamia Gymnospermia.				
1	cinerea. B.R.	(li.) cinereous.	S. Europe.	1824.	6. 9. H. ¼.	
		<i>Marrubium cinereum</i> . s.s.				
2	acetabulosum. B.R.(li.)	saucer-leaved.	Candia.	1676.	6. 8. F. ¼.	Barrel. icon. 129.
3	pseudo-dictamnus. B.R.(li.)	shrubby white.	—	1596.	7. 8. F. ½.	Flor. græc. t. 562.
		<i>Marrubium pseudo-dictamnus</i> . F.G.				
4	africanum. B.R.	(li.) African.	C. B. S.	1710.	7. 9. G. ¼.	Comm. hort. 2. t. 90.
5	crispum. B.R.	(li.) curled-leaved.	S. Europe.	1714.	7. 8. H. ¼.	Herm. parad. t. 200.
6	hispanicum. B.R.	(li.) Spanish.	Spain.	—	— H. ¼.	— t. 201.
7	hirsutum. B.R.	(li.) hairy.	S. Europe.	6. 7. H. ¼.	
8	orientale. B.R.	(li.) oriental.	Levant.	1828.	— H. ¼.	
MOLUCCE'LLA. S.S.	MOLUCCA-BALM.	Didynamia Gymnospermia. L.				
1	tuberosa. s.s.	(wh.) tuberous-root'd.	Tartary.	1796.	7. H. ¼.	Pall. it. 3. app. t. T.
2	lævis. s.s.	(wh.re.) smooth.	Syria.	1570.	7. 8. H.○.	Bot. mag. t. 1852.
CHASM'ONIA. B.R.	CHASM'ONIA.	Didynamia Gymnospermia.				
incisa. B.R.	(wh.re.) spiny.	Levant.	1596.	7. 8. H.○.	Bot. reg. t. 1244.	
		<i>Moluccella spinosa</i> . Flor. græc. t. 567.				
LEON'URUS. S.S.	MOTHERWORT.	Didynamia Gymnospermia. L.				
1	Cardiaca. s.s.	(li.wh.) common.	Britain.	7. 8. H. ¼.	Eng. bot. v. 4. t. 286.
2	villous. s.s.	(li.wh.) villous.	Tauria.	1824.	— H. ¼.	Murr. c. gott. 8. t. 4.
3	crispus. s.s.	(li.) curl-leaved.	Siberia.	1658.	— H. ¼.	
4	supinus. s.s.	(wh.) procumbent.	—	1816.	6. 8. H. ¼.	
5	glaucescens. Led.	(li.) glaucous-leaved.	—	1827.	7. 9. H. ¼.	
6	tataricus. s.s.	(li.) Tartarian.	Tartary.	1756.	8. 10. H. ♂.	Mill. ic. 1. t. 80.
7	sibiricus. s.s.	(ro.) Siberian.	Siberia.	1759.	6. 8. H. ♂.	Swt. br. fl. gar. t. 204.
8	condensatus. s.s.	(li.) condensed.	—	1824.	— H. ¼.	Swt. br. fl. gar. t. 197.
9	heterophyllos. B.F.G.(ro.)	various-leav'd.	S. America.	—	— H.○.	
10	multifidus. Desf.	(li.) multifid.	—	—	— H.○.	
11	neglectus. Sch.	(li.) neglected.	—	—	— H. ¼.	
12	altaicus. s.s.	(li.) Altaic.	Altay.	1826.	— H. ¼.	
13	lanatus. s.s.	(st.) woolly.	Siberia.	1752.	— H. ¼.	Gmel. sib. 3. t. 54.
		<i>Ballota lanata</i> . w.				
GALEO'BDOLON. S.S.	DEAD-NETTLE.	Didynamia Angiospermia. S.S.				
		luteum. E.B.	(ye.) yellow.	Britain. 5. 6. H. ¼.	Eng. bot. v. 11. t. 787.
GALEO'PSIS. S.S.	HEMP-NETTLE.	Didynamia Gymnospermia. L.				
1	Ládanum. E.B.	(ro.) red.	Britain.	7. 9. H.○.	Eng. bot. v. 13. t. 884.

LABIATÆ.

2 angustifolia. p.s. (ro.) narrow-leaved.	Europe.	1820.	7. 9. H. ⊙.
3 canescens. s.s. (li.) canescent.	Germany.	1827.	— H. ⊙.
4 pubescens. s.s. (wh.re.) pubescent.	—	1820.	— H. ⊙.
5 ochroleuca. s.s. (ye.) downy.	Britain.	7. 8. H. ⊙.
villosa. E.B.			
6 Tetràhit. E.B. (wh.pu.) common.	—	— H. ⊙.
7 versicolor. E.B. (ye.vi.) various-colour'd.	—	— H. ⊙.
cannabina. w.			
L'AMÍCICA. S.S.	ARCHANGEL.	Didynamia Gymnospermia. L.	
1 longiflòrum. t.n. (pu.) long-flowered.	S. Europe.	4. 9. H. ¼.
maculatum. Eng. bot. non Flor. Græc.			
2 rugosum. s.s. (pu.) rough.	Italy.	1766.	7. 8. H. ¼.
3 maculatum. E.B. (pu.) spotted.	Britain.	4. 5. H. ¼.
4 album. E.B. (wh.) white.	—	—	4. 9. H. ¼.
5 striatum. F.G. (wh.std.) striped-flower'd. Levant.	1829.	—	H. ¼.
6 bifidum. s.s. (wh.) bifid-helmetted. Naples.	1825.	—	H. ¼.
7 moschatum. s.s. (wh.) musk-scented. Levant.	1789.	7. 9. H. ¼.	
8 mollé. s.s. (wh.) Pellitory-leaved.	Naples.	1824.	4. 8. H. ¼.
9 flexuosum. T.N. (wh.) flexuous.	Italy.	1729.	7. 8. H. ¼.
10 garganicum. s.s. (pu.) woolly.	—	1711.	3. 10. H. ¼.
11 lœviatulum. s.s. (pu.) smooth.	—	—	2. 8. H. ⊙.
12 purpureum. s.s. (pu.) purple.	Britain.	—	— H. ⊙.
β albidum. (wh.) white-flowered.	—	—	— H. ⊙.
13 incisum. s.s. (li.) cut-leaved.	England.	—	5. 7. H. ⊙.
14 amplexicaulé. s.s. (pu.) Henbit.	Britain.	—	— H. ⊙.
15 multifidum. s.s. (pu.) multifid-leaved. Levant.	1752.	4. 5.	H. ⊙.
ORVALA. DC.	ORV'ALA.	Didynamia Gymnospermia.	
1 lamioïdes. DC. (pu.bh.) balm-leaved.	Italy.	1596.	4. 7. H. ¼.
<i>Lâmiu' Orvâla. B.M.</i>			
2 garginica. DC. (pu.) Garganian.	—	—	H. ¼.
PHYSOSTEGIA. B.R.	PHYSOST'EGIA.	Didynamia Angiospermia.	
1 virginiana. B.R. (cr.) Virginian.	N. America.	1683.	7. 9. H. ¼.
2 speciosa. B.F.G. (ro.) beautiful.	—	1820.	— H. ¼.
<i>Dracocéphalum speciosum. B.F.G.</i>			
3 denticulata. B.R. (ro.) toothed.	—	1787.	8. 9. H. ¼.
<i>Dracocéphalum denticulatum. B.M.</i>			
4 variegata. B.R. (re.wh.) variegated.	—	1812.	— H. ¼.
SPHA'CELE. B.R.	SPHA'CELE.	Didynamia Gymnospermia.	
Lindlèi. B.R. (li.) Lindley's.	Valparaiso.	1825.	7. 9. G. ḥ.
<i>Stachys Sálviae. B.R.</i>			
BETO'NICA. S.S.	BETONY.	Didynamia Gymnospermia. L.	
1 officinalis. s.s. (pu.) wood.	Britain.	7. 8. H. ¼.
2 strícta. s.s. (pu.) Danish.	Denmark.	1592.	6. 7. H. ¼.
3 incana. s.s. (pi.) hoary.	Italy.	1759.	— H. ¼.
4 orientalis. s.s. (pu.) oriental.	Levant.	1737.	— H. ¼.
5 alopecuroides. s.s. (st.) Foxtail.	S. Europe.	1759.	7. H. ¼.
6 hirsuta. s.s. (pu.) hairy.	Italy.	1710.	6. 7. H. ¼.
7 grandiflora. s.s. (pu.) great-flowered.	Siberia.	1800.	— H. ¼.
ST'ACHYS. S.S.	HEDGE-NETTLE.	Didynamia Gymnospermia. L.	
1 spinosa. s.s. (wh.) thorny.	Candia.	1840.	7. F. ḥ.
2 glutinosa. s.s. (wh.) clammy.	—	1729.	6. 7. H. ¼.
3 fruticosus. s.s. (pu.) frutescent.	Caucasus.	1825.	— H. ḥ.
4 æthiopica. s.s. (li.) Ethiopian.	C. B. S.	1770.	4. 7. G. ¼.
5 cósrica. s.s. (li.) Corsican.	Corsica.	1825.	— H. ¼.
6 angustifolia. s.s. (li.) narrow-leaved.	Tauria.	1823.	6. 7. H. ¼.
7 rugosa. s.s. (st.) rough.	C. B. S.	1774.	7. 8. G. ḥ.
8 betonicaefolia. s.s. (ye.) Betony-leaved.	Crete.	1812.	6. 7. H. ¼.
9 stenophylla. s.s. (ye.) linear-leaved.	S. Spain.	1819.	— F. ḥ.
10 orientalis. s.s. (st.) oriental.	Levant.	1768.	— H. ¼.
11 marítima. s.s. (ye.) sea-side.	S. Europe.	1714.	7. H. ¼.
12 ánnua. s.s. (wh.pu.) annual.	—	1713.	6. 8. H. ⊙.
13 hirta. s.s. (ye.pu.) procumbent.	Spain.	1725.	— H. ¼.
14 arvensis. s.s. (pu.) corn.	Britain.	7. 8. H. ⊙.
15 arenaria. s.s. (pu.) sand.	Barbary.	1824.	6. 8. H. ¼.
16 diffusa. Schweig. (re.) diffuse.	S. Europe.	1804.	— H. ¼.
arenaria. B.M. non Desf. rúbra. Donn.			
17 ibérica. B.M. (pu.) Iberian.	Iberia.	1823.	— H. ¼.
18 scordifolia. W.en. (pu.) wedge-leaved.	1816.	7. 8. H. ¼.
19 áspera. M. (pu.) harsh-leaved.	N. America.	—	H. ¼.

Eng. bot. v. 33. t. 2353.

Eng. bot. v. 3. t. 207.

——— v. 10. t. 667.

Eng. bot. v. 36. t. 2550.

Flor. græc. t. 555.

——— t. 556.

Eng. bot. v. 11. t. 768.

Flor. græc. t. 557.

Cyr. rar. fasc. 1. t. 7.

Ten. flor. nap. ic.

Sm. exot. bot. 1. t. 48.

Pluk. alm. t. 198. f. 1.

Eng. bot. v. 11. t. 769.

Eng. bot. v. 27. t. 1933.

——— v. 11. t. 770.

Comm. rar. t. 26.

Bot. mag. t. 172.

Bot. mag. t. 467.

Swt. br. fl. gar. t. 93.

Bot. mag. t. 214.

Vent. cels. t. 44.

Bot. reg. t. 1226.

Eng. bot. v. 16. t. 1142.

Bot. mag. t. 2125.

Lam. ill. t. 507. f. 2.

Jacq. aust. t. 1. t. 78.

Mur. com. got. 2. t. 3.

Bot. mag. t. 700.

Flor. græc. t. 559.

Moris. s. 11. t. 4. f. 17.

Jacq. obs. 4. t. 77.

Swt. br. fl. gar. 2. t. 180.

Jacq. ic. 3. t. 493.

Flor. græc. t. 560.

Jacq. vind. 1. t. 70.

Jacq. aust. 4. t. 360.

All. ped. 1. t. 2. f. 3.

Eng. bot. v. 17. t. 1154.

Desf. atl. 2. t. 126.

Bot. mag. t. 1915.

LABIATÆ.

20 híspida. s.s.	(<i>pu.</i>) hispid.	N. America.	1827.	7. 8. H. ȝ.	
21 itálica. Mil.	(<i>pu.</i>) Italian.	Italy.	1747.	— H. ♂.	
22 Balbisii. L.en.	(<i>ye.</i>) Balbis's.		1823.	— H. ȝ.	
23 nepetafòlia. s.s.	(<i>pu.</i>) Catmint-leaved.	Levant.	1805.	6. 8. H. ȝ.	
24 arábica. s.s.	(<i>re.</i>) Arabian.	Arabia.	1820.	— H. ⊙.	Eng. bot. v. 30. t. 2089.
25 ambígua. s.s.	(<i>pu.</i>) ambiguous.	Britain.	6. 7. H. ȝ.	— v. 6. t. 416.
26 sylvática. s.s.	(<i>pu.</i>) common.		7. 8. H. ȝ.	L'Herit. s. nov. t. 26.
27 circinàta. s.s.	(<i>pu.</i>) blunt-leaved.	Barbary.	1777.	5. 7. H. ȝ.	
canarién sis. Jacq. ic. 1. t. 108.					
28 mollissima. W.en. (<i>li.</i>) soft-leaved.	Corfu.	1806.	7. 8. H. ȝ.	W. hort. ber. t. 60.	
29 pubescens. T.s.	(<i>st.</i>) pubescent.	Italy.	1826.	— H. ȝ.	
30 decumbens. P.s.	(<i>st.</i>) decumbent.	S. Europe.	1816.	5. 7. H. ȝ.	
31 coccinea. s.s.	(<i>sc.</i>) scarlet.	S. America.	1798.	6. 8. F. ȝ.	Bot. mag. t. 666.
32 récta. s.s.	(<i>ye.</i>) upright.	S. Europe.	1683.	— H. ȝ.	Jacq. aust. 4. t. 359.
33 herálea. w.	(<i>pu.</i>) Phlomis-leaved.	Italy.	1816.	6. 7. H. ȝ.	All. ped. t. 84. f. 1.
phlomoides. W.en.					
34 oblíqua. P.s.	(<i>st.</i>) oblique-leaved.	Hungary.	—	— H. ȝ.	W. et K. hung. t. 133.
35 palústris. s.s.	(<i>pu.</i>) Clown's all-heal.	Britain.	8. H. ȝ.	Eng. bot. v. 24. t. 1675.
36 germánica. s.s.	(<i>pu.</i>) downy.	England.	7. 8. H. ȝ.	— v. 12. t. 829.
37 intermédia. w.	(<i>pu.</i>) oblong-leaved.	Carolina.	1762.	6. 7. H. ȝ.	Swt. br. fl. gar. t. 100.
38 sibírica. L.en.	(<i>pu.</i>) Siberian.	Siberia.	1822.	6. 8. H. ȝ.	
39 polystachya. s.s.	(<i>pu.</i>) many-spiked.	Naples.	1824.	— H. ȝ.	
40 salviæfòlia. s.s.	(<i>pu.</i>) Sage-leaved.		—	— H. ȝ.	
41 crética. s.s.	(<i>pu.</i>) Cretan.	Candia.	1640.	— F. ȝ.	Flor. græc. t. 558.
42 dasyántha. Raf.	(<i>pu.</i>) woolly-flower'd.	Sicily.	1829.	— H. ȝ.	
43 exaltàta. Ot.	(<i>pu.</i>) tall.	—	— H. ȝ.	
44 alpina. s.s.	(<i>pu.</i>) Alpine.	Germany.	1597.	— H. ȝ.	Lap. pyr. 1. t. 8.
45 lanáta. s.s.	(<i>pu.</i>) woolly.	Siberia.	1782.	6. 9. H. ȝ.	Jacq. ic. 1. t. 107.
46 latifòlia. s.s.	(<i>pu.</i>) broad-leaved.	1775.	6. 7. H. ȝ.	
47 biénnis. Roth.	(<i>pu.</i>) biennial.	Germany.	1801.	6. 8. H. ♂.	
48 lusitánica.	(<i>pu.</i>) Portugal.	Portugal.	1800.	— H. ȝ.	
<i>Eriostemon lusitanicum.</i> Lk.					
ZIET'ENIA. P.S.	ZIET'ENIA.	Didynamia Gymnospermia. P.S.			
orientális. P.S.	(<i>li.</i>) Lavender-leaved.	Levant.	1816.	5. 7. H. ȝ.	
lavandulifòlia. L.en.	<i>Stachys lavandulifòlia.</i> w.				
CHAIT'URUS. B.R.	CHAIT'URUS.	Didynamia Gymnospermia.			
marrubiástrum. Ehr.	(<i>li.</i>) small-flowered.	Austria.	1710.	6. 8. H. ⊙.	Jacq. aust. 5. t. 405.
<i>Leonurus marrubiastrum.</i> w.					
CRANIO'TOME. B.R.	CRANIO'TOME.	Didynamia Gymnospermia.			
versicolor. B.R. (<i>re.wh.</i>) various-coloured.	Nepaul.	1823.	7. 8. G. ȝ.		
<i>Ajuga furcata.</i> L.enum.	<i>Anisomèles nepalensis.</i> s.s.				
NE'PETA. L.	CATMINT.	Didynamia Gymnospermia. L.			
1 serpyllifòlia. s.s.	(<i>li.</i>) Thyme-leaved.	Tauria.	1826.	6. 8. H. ȝ.	
2 parvifòra. s.s.	(<i>li.</i>) small-flowered.	—	1828.	— H. ȝ.	
3 sibírica. s.s.	(<i>bl.</i>) Siberian.	Siberia.	1804.	7. H. ȝ.	
4 grandifòra. s.s.	(<i>bl.</i>) large-flowered.	Caucasus.	1806.	7. 8. H. ȝ.	
5 melissæfòlia. s.s.	(<i>bl.</i>) Balm-leaved.	Candia.	1752.	— H. ȝ.	
6 coloràta. W.en.	(<i>vi.</i>) coloured.	Caucasus.	1806.	— H. ȝ.	
7 longifòra. H.K.	(<i>bl.</i>) long-flowered.	Persia.	1802.	6. 8. H. ȝ.	Venten. cels. t. 66.
8 Mussini. H.K.	(<i>bl.</i>) scollop-leav'd.	Siberia.	1804.	5. 8. H. ȝ.	Bot. mag. t. 923.
longifòra. B.M. non v.					
9 incána. s.s.	(<i>bl.</i>) hoary.	Levant.	1723.	8. H. ȝ.	
10 gravéolens. s.s.	(<i>bl.</i>) strong-smelling.	S. Europe.	1804.	7. 8. H. ȝ.	All. ped. t. 2. f. 1.
11 amethystina. Poir.	(<i>bl.</i>) amethystine.	—	1820.	— H. ȝ.	
12 Nepetélla. s.s.	(<i>li.</i>) small.	—	1758.	— H. ȝ.	
13 teucriifòlia. s.s.	(<i>bl.</i>) Teucrium-leav'd.	Armenia.	1816.	— H. ȝ.	
14 lamiifòlia. s.s.	(<i>bl.</i>) Lamium-leaved.	—	1806.	— H. ȝ.	
15 citriodòra. Lag.	(<i>wh.</i>) Citron-scented.	Spain.	1826.	— H. ȝ.	
16 lophántha. F.	(<i>bl.</i>) crested.	Altay.	1821.	— H. ȝ.	
17 angustifòlia. s.s.	(<i>wh.</i>) narrow-leaved.	Spain.	1798.	6. 7. H. ȝ.	Eng. bot. v. 2. t. 137.
18 Catária. s.s.	(<i>wh.</i>) common.	Britain.	7. 9. H. ȝ.	
19 ucránica. s.s.	(<i>vi.</i>) Ukrainian.	Ukraine.	1789.	7. 8. H. ȝ.	
20 nùda. L.	(<i>li.</i>) naked.	S. Europe.	1710.	— H. ȝ.	Flor. græc. t. 547.
21 pannònica. L.	(<i>li.</i>) close-flowered.	Hungary.	1683.	8. 10. H. ȝ.	Jacq. aust. 2. t. 129.
22 violacea. s.s.	(<i>vi.</i>) violet-coloured.	Spain.	1723.	7. 9. H. ȝ.	Bocc. mus. t. 36.
23 coerulea. s.s.	(<i>bl.</i>) blue-flowered.	—	1777.	5. 6. H. ȝ.	
24 latifòlia. s.s.	(<i>pu.</i>) broad-leaved.	Pyrenees.	1818.	7. 8. H. ȝ.	
25 macrùra. s.s.	(<i>li.</i>) long-tailed.	Siberia.	—	— H. ȝ.	
26 críspa. s.s.	(<i>bl.</i>) curl-leaved.	Levant.	1800.	— H. ȝ.	

27 diffusa. s.s.	(bl.) spreading.	Siberia.	1820.	7. 8. H. ȝ.	
28 imbricata. s.s.	(pu.) imbricated.	Spain.	1825.	— H. ȝ.	
29 teucroides. s.s.	(br.) Teucrium-like.	Levant.	1819.	— H. ȝ.	
30 italica. w.	(wh.sp.) Italian.	Italy.	1640.	6. 8. H. ȝ.	Flor. græc. t. 548.
31 marrubioides. W.en.	(pu.) horehound-leav'd.	Barbary.	1823.	— H. ȝ.	Desf. atl. 2. t. 123.
32 multibracteata. s.s.	(vi.) many-bracted.	S. Europe.	1774.	5. 6. H. ȝ.	Jacq. obs. 3. t. 75.
33 lanata. s.s.	(vi.) woolly.	Spain.	1683.	6. 8. H. ȝ.	Barrel. ic. t. 602.
34 tuberosa. s.s.	(vi.) tuberous-root'd.	Morocco.	1801.	7. 8. H. ȝ.	Desf. atl. 2. t. 124.
35 reticulata. s.s.	(li.) netted.	Crete.	1823.	— G. ȝ.	Alp. exot. t. 283.
36 Scordotis. s.s.	(wh.) sessile-spiked.	1804.	— H. ȝ.	Cavan. ic. 1. t. 49.
37 suavolens. Rœm.	(wh.) sweet-scented.	1779.	6. 7. H. ȝ.	
38 botryoides. w.	(wh.) annual.	
	bipinnata. Cav.				
GLE'CHOMA. L.	GROUND-IVY.	Didynamia Gymnospermia. L.			
1 hederaea. E.B.	(bl.) common.	Britain.	3. 5. H. ȝ.	Eng. bot. v. 12. t. 853.
2 hirsuta. P.S.	(pu.) hairy.	Hungary.	— H. ȝ.	W. et K. hung. t. 119.
HEMIC'ENIA. B.P.	HEMIC'ENIA.	Didynamia Gymnospermia.			
	purpurea. B.P.	N. S. W.	1824.	4. 8. G. ȝ.	Jacq. vind. 3. t. 30.
SIDE'RITIS. S.S.	IRONWORT.	Didynamia Gymnospermia. L.			Comm. hort. 2. t. 99.
1 canariensis. s.s.	(ye.) Canary.	Canaries.	1697.	5. 8. G. ȝ.	Flor. græc. t. 551.
2 cändigans. s.s.	(ye.) Mulinleaved.	Madeira.	1714.	4. 7. G. ȝ.	Murr. com. got. 1. t. 4.
3 crética. s.s.	(wh.) Cretan.	Crete.	1823.	— G. ȝ.	Flor. græc. t. 552.
4 montana. s.s.	(ye.) mountain.	Austria.	1752.	7. 8. H. ȝ.	—— t. 550.
5 elegans. s.s.	(wh.bl.) dark-flowered.	1787.	7. H. ȝ.	
6 romana. s.s.	(wh.) Roman.	Italy.	1759.	6. 8. H. ȝ.	
7 syriaca. s.s.	(ye.) Syrian.	Levant.	1597.	6. 9. G. ȝ.	
8 taúrica. W.en.	(ye.) Taurian.	Tauria.	1818.	— H. ȝ.	
9 brütia. T.S.	(ye.) woolly.	Naples.	1827.	— F. ȝ.	
10 perfoliata. s.s.	(ye.) perfoliate.	Levant.	1731.	8. 11. H. ȝ.	Vent. cels. 1. t. 98.
11 pulvillans. s.s.	(wh.ye.) brownish.	1828.	— F. ȝ.	Cavan. ic. 2. t. 304.
12 leucantha. s.s.	(wh.) white-flowered.	Spain.	— F. ȝ.	—— 2. t. 186.
13 incana. s.s.	(ye.) Lavender-leav'd.	1752.	7. 8. H. ȝ.	
14 angustifolia. s.s.	(ye.) narrow-leaved.	1825.	— H. ȝ.	Cavan. ic. 3. t. 209.
15 spinosa. s.s.	(ye.) spiny.	6. 9. H. ȝ.	
16 ilicifolia. s.s.	(ye.) Holly-leaved.	Levant.	— H. ȝ.	
17 serrata. s.s.	(ye.) saw-leaved.	Spain.	1818.	— H. ȝ.	Schk. hand. 2. t. 158.
18 hyssopifolia. s.s.	(ye.) Hyssop-leaved.	Pyrenees.	1597.	6. 11. H. ȝ.	
19 f'ostida. L.en.	(ye.) strong-scented.	— H. ȝ.	
20 pyrenaica. s.s.	(ye.) Pyrenean.	Pyrenees.	1816.	— H. ȝ.	
21 chamædryfolia. s.s.	(ye.) Germanander-ld.	Spain.	— H. ȝ.	Cavan. ic. 4. t. 301.
22 scordioides. s.s.	(ye.) scollop-leaved.	France.	1597.	8. 11. H. ȝ.	Barrel. ic. t. 343.
23 hirsuta. s.s.	(ye.) hairy.	S. Europe.	1731.	6. 7. H. ȝ.	Cavan. ic. 4. t. 302.
24 crispata. s.s.	(ye.) curled-leaved.	Gibraltar.	1816.	6. 9. F. ȝ.	
MARRUBIUM. S.S.	HOREHOUND.	Didynamia Gymnospermia. L.			
1 Aly'ssum. s.s.	(pu.) plaited-leaved.	Spain.	1597.	7. 8. H. ȝ.	Ger. herb. 379. f. 1.
2 astracanicum. s.s.	(bl.) Astracan.	Levant.	1816.	— H. ȝ.	Jacq. ic. 1. t. 109.
3 peregrinum. w.	(wh.) Sicilian.	Sicily.	1640.	7. 9. H. ȝ.	Jacq. aust. 2. t. 160.
4 candidissimum. s.s.	(wh.) woolly white.	Levant.	1732.	— H. ȝ.	Dill. elt. 274. f. 214.
5 creticum. w.	(wh.) Cretan.	1596.	— H. ȝ.	
6 affine. L.en.	(pu.) related.	Siberia.	1816.	— H. ȝ.	Bocc. mus. 2. t. 96.
7 supinum. s.s.	(wh.) procumbent.	S. Europe.	1714.	8. 10. H. ȝ.	Flor. græc. t. 561.
8 catariaefolium. s.s.	(wh.) Catmint-leav'd.	Levant.	1823.	7. 9. H. ȝ.	Eng. bot. t. 6. f. 410.
9 velutinum. F.C.	(ye.) velvety.	1827.	— F. ȝ.	
10 vulgare. s.s.	(wh.) common.	Britain.	6. 9. H. ȝ.	
LAVA'NDULA. S.S.	LAVENDER.	Didynamia Gymnospermia. L.			
1 Spica. s.s.	(bl.) common.	S. Europe.	1568.	7. 9. H. ȝ.	Schk. hand. 2. t. 157.
β alba.	(wh.) white-flowered.	— H. ȝ.	
2 latifolia. s.s.	(bl.) broad-leaved.	— H. ȝ.	
3 St'echas. s.s.	(bl.) French.	S. Europe.	1562.	5. 7. G. ȝ.	Flor. græc. t. 549.
4 viridis. s.s.	(bl.) Madeira.	Madeira.	1777.	— G. ȝ.	Hoff. et Lk. lus. 1. t. 4.
5 dentata. s.s.	(bl.) tooth-leaved.	Spain.	1597.	6. 9. G. ȝ.	Bot. mag. t. 401.
6 pinnata. s.s.	(bl.) pinnated.	Madeira.	1777.	4. 8. G. ȝ.	—— t. 400.
7 heterophylla. s.s.	(bl.) various-leaved.	1816.	— G. ȝ.	
8 formosa. L.en.	(bl.) handsome.	Canaries.	— G. ȝ.	
9 multifida. s.s.	(bl.) cut-leaved.	1597.	7. 9. G. ȝ.	Lobel. ic. 432.
10 abortivoides. s.s.	(bl.) Southernwood-ld.	1699.	6. 9. G. ȝ.	Comm. rar. t. 27.
DRACOC'EPHALUM. DRAGON'S-HEAD.	Didynamia Gymnospermia. L.				
1 Ruyschiæna. s.s.	(bl.) Hyssop-leaved.	N. Europe.	1699.	6. 9. H. ȝ.	Flor. dan. t. 121.
2 austriacum. s.s.	(bl.) Austrian.	Austria.	1597.	6. 7. H. ȝ.	Jacq. ic. 1. t. 112.

3	origanoïdes. s.s.	(<i>bl.</i>) Marjoram-like.	Siberia.	1827.	6. 7. H. ȝ.		
4	botryoïdes. s.s.	(<i>bl.</i>) woolly-leaved.	Caucasus.	1823.	— H. ȝ.		
5	palmatum. s.s.	(<i>bl.</i>) palmated.	Siberia.	1815.	6. 8. H. ȝ.		
6	canariense. s.s.	(<i>li.</i>) Balm of Gilead.	Canaries.	1697.	7. 9. G. ḥ.		
7	pinnatum. s.s.	(<i>bl.</i>) wing-leaved.	Siberia.	1827.	— H. ḥ.		
8	peregrinum. s.s.	(<i>bl.</i>) prickly-leaved.	—	1759.	7. 8. H. ȝ.		
9	argunense. s.s.	(<i>bl.</i>) Fischer's.	—	1822.	7. 9. H. ȝ.		
10	integerrifolium. Led.	(<i>bl.</i>) entire-leaved.	—	1827.	— H. ȝ.		
11	grandiflorum. s.s.	(<i>bl.</i>) great-flowered.	—	1759.	— H. ⊙.	Comm. hort. 2. t. 41.	
12	altaïense. s.s.	(<i>bl.</i>) Betony-leaved.	Georgia.	1787.	— H. ȝ.		
13	Moldávica. s.s.	(<i>bl.</i>) Moldavian.	Moldavia.	1596.	7. 8. H. ⊙.		
14	canescens. s.s.	(<i>bl.</i>) hoary-leaved.	Levant.	1711.	— H. ⊙.		
15	ibéricum. s.s.	(<i>bl.</i>) Iberian.	Iberia.	1827.	— H. ⊙.		
16	peltatum. s.s.	(<i>bl.</i>) Willow-leaved.	Levant.	1711.	— H. ⊙.		
17	thymiflorum. s.s.	(<i>bl.</i>) small-flowered.	Siberia.	1752.	6. 9. H. ⊙.		
18	nútans. s.s.	(<i>bl.</i>) nodding.	—	1731.	7. 8. H. ȝ.		
19	chamædryoides. s.s.	(<i>bl.</i>) Chamædrys-like.	—	1828.	— H. ȝ.		
20	sibiricum. s.s.	(<i>bl.</i>) Siberian.	Siberia.	1760.	6. 8. H. ȝ.		
HORM'INUM. P.S.							
	pyrenäicum. s.s.	(<i>bl.</i>) Pyrenean.	Pyrenees.	1820.	6. 7. H. ȝ.		
MELISSA. S.S.							
	BALM.	Didynamia Gymnospermia. L.					
1	officinalis. w.	(<i>bh.</i>) common.	S. Europe.	1573.	6. 10. H. ȝ.		
2	hirsuta. H.H.	(<i>bh.</i>) hairy.	—	...	— H. ȝ.		
3	cordifolia. P.S.	(<i>bh.</i>) heart-leaved.	—	...	— H. ȝ.		
4	altissima. s.s.	(<i>bh.</i>) tall.	—	1823.	— H. ȝ.		
LEPECHI'NIA. S.S.							
	LEPECHI'NIA.	Didynamia Gymnospermia. W.					
	spicata. s.s.	(<i>st.</i>) spiked.	Mexico.	1800.	7. 8. H. ȝ.		
THY'MBRA. S.S.							
	THY'MBRA.	Didynamia Gymnospermia. L.					
1	spicata. L.	(<i>re.</i>) spiked-flower'd.	Levant.	1699.	6. 7. G. ḥ.		
2	verticillata. L.	(<i>re.</i>) whorl-flowered.	Spain.	1702.	— G. ḥ.		
A'CYNOS. P.S.							
	A'CYNOS.	Didynamia Gymnospermia. P.S.					
1	vulgaris. P.S.	(<i>vi.</i>) Basil-leaved.	Britain.	6. 8. H. ⊙.		
Thymus A'cinos. E.B.							
2	graveolens. L.en.	(<i>vi.</i>) strong-scented.	Tauria.	1823.	— H. ḥ.		
3	villosum. P.S.	(<i>vi.</i>) villous.	Germany.	1817.	— H. ⊙.		
4	exiguus. F.G.	(<i>pu.</i>) small.	Levant.	1828.	— H. ⊙.		
5	alpinus. P.S.	(<i>vi.</i>) Alpine.	Austria.	1731.	6. 9. H. ḡ.		
6	purpurascens. P.S. (<i>pu.</i>)	purple.	Spain.	1820.	— H. ḡ.		
7	patavinus. P.S.	(<i>pu.</i>) Marjoram-leav'd.	S. Europe.	1776.	6. 8. H. ḡ.		
8	rotundifolius. P.S. (<i>pu.</i>)	round-leaved.	Spain.	1820.	— H. ḥ.		
GARD'OQUIA. S.S.							
	GARD'OQUIA.	Didynamia Gymnospermia.					
1	discolor. K.S.	(<i>pu.</i>) two-coloured.	Caracas.	1827.	4. 7. S. ḥ.		
2	origanoïdes. s.s.	(<i>pu.</i>) Marjoram-like.	Trinidad.	1824.	— S. ḥ.		
CALAMI'NTHA. Ph.							
	CALAMINT.	Didynamia Gymnospermia.					
1	marifolia. P.S. (<i>wh.sp.</i>)	Marum-leaved.	Spain.	1788.	6. 7. H. ȝ.		
	Népeta marifolia.	Cav. Thymus marifolius. W.en.					
2	crética. P.S.	(<i>pu.</i>) Cretan.	S. Europe.	1596.	— F. ḥ.		
3	fruticosa. P.S.	(<i>pu.</i>) shrubby.	Spain.	1752.	7. 9. F. ḥ.		
4	álba. L.en.	(<i>wh.</i>) white.	Hungary.	1818.	— H. ȝ.		
5	Népeta. P.S.	(<i>ii.</i>) lesser.	England.	7. 10. H. ȝ.		
	Mélissa Népeta.	L. Thymus Népeta. E.B.					
6	vulgaris.	(<i>ii.</i>) common.	England.	7. 8. H. ȝ.		
	Mélissa Calaminta.	L. Thymus Calaminta. E.B.					
7	grandiflora. P.S.	(<i>ro.</i>) great-flowered.	Italy.	1596.	6. 9. H. ȝ.		
	Melissa grandiflora.	B.M.					
8	caroliniana. N.	(<i>ii.</i>) Carolina.	Carolina.	1804.	6. 7. F. ḥ.		
	Thymus grandiflorus.	B.M.					
CLINOP'ODIUM. S.S. WILD-BASIL.							
	Didynamia Gymnospermia. L.						
1	vulgare. s.s.	(<i>ro.</i>) common.	Britain.	6. 8. H. ȝ.		
2	atropurpureum. H.H.	(<i>pu.</i>) dark purple.	S. Europe.	1820.	— H. ȝ.		
3	ægyptiacum. s.s.	(<i>pu.</i>) Egyptian.	Egypt.	1759.	— H. ȝ.		
MELITTIS. S.S.							
	BASTARD-BALM.	Didynamia Gymnospermia. L.					
1	Melissophyllum. s.s. (<i>pu.</i>)	common.	England.	6. 8. H. ȝ.		
	β alpina.	(<i>pu.wh.</i>) Alpine.	Switzerland.	— H. ȝ.		
2	grandiflora. s.s. (<i>wh.vi.</i>)	great-flowered.	England.	5. H. ȝ.		
MACB'RIDEA. E.C.							
	MACB'RIDEA.	Didynamia Gymnospermia.					
	pulchra. N.	(<i>re.std.</i>) pretty.	Carolina.	1804.	7. 9. G. ḥ.		
	Thymbra caroliniana.	Walter.					

Swt. br. fl. gar. s. 2.

Lam. ill. t. 513. f. 1.

Swt. br. fl. gar. t. 38.

Lam. ill. t. 513. f. 2.

Gmel. sib. 3. t. 50.

Bot. reg. t. 841.

Bot. mag. t. 2185.

Swt. br. fl. gar. t. 252.

Flor. græc. t. 579.

Bot. reg. t. 1292.

Flor. græc. t. 546.

Jacq. obs. 4. t. 87.

Eng. bot. v. 6. t. 411.

Cavan. ic. 6. t. 576.

Barrel. ic. 1166.

Eng. bot. v. 20. t. 1414.

Eng. bot. v. 24. t. 1676.

Bot. mag. t. 208.

Bot. mag. t. 997.

Eng. bot. v. 20. t. 1401.

Eng. bot. v. 9. t. 577.

Eng. bot. v. 9. t. 636.

PRUNE'LLA. S.S.	SELF-HEAL.	Didynamia Gymnospermia. L.		
1 vulgàris. s.s. (bl.) common.	(bl.) common.	Britain.	7. 8. H. ¼.	
β álba.	(wh.) white-flowered.	— H. ¼.	
2 austrális.	(bl.) New Holland.	N. S. W. 1826.	— F. ¼.	
vulgàris. B.P. nec aliorum.				
3 pensylvánica. s.s. (bl.) Pensylvanian.	N. America. 1801.	7. 9. H. ⓠ.	Eng. bot. v. 14. t. 961.	
4 grandiflòra. s.s. (bl.) great-flowered.	Austria. 1596.	— H. ¼.		
5 intermédia. s.s. (bl.) intermediate.	S. Europe. 1790.	— H. ¼.		
grandiflòra. B.M.				
6 incisa. L.en.	(bl.) cut-leaved.	— H. ¼.	
7 laciniàta. L.	(ye.) yellow-flower'd.	Austria. 1713.	— H. ⓠ.	
8 hastàta. s.s.	(bl.) halberd-leaved.	Portugal.	— H. ¼.	
9 longifòlia. P.S.	(bl.) long-leaved.	France.	— H. ¼.	
10 hyssopifòlia. s.s. (pu.) hyssop-leaved.	S. Europe. 1731.	— H. ¼.	W. hort. ber. I. t. 9.	
CLEONIA. S.S.	CLEONIA.	Didynamia Gymnospermia. L.	Bot. mag. t. 2014.	
lusitánica. s.s. (pa.bl.) sweet-scented.	Portugal. 1710.	6. 7. H. ⓠ.	— t. 337.	
SCUTELL'ARIA. S.S.	SKULLCAP.	Didynamia Gymnospermia. L.	Lam. ill. t. 516. f. 2.	
1 galericulàta. s.s. (bl.) common.	Britain.	6. 9. H. ¼.	Moris. s. 11. t. 5. f. 7.	
2 hastifòlia. s.s. (pu.) hastate-leaved.	Germany. 1798.	6. 7. H. ¼.	Mill. ic. 1. t. 70.	
3 minor. s.s. (ro.) lesser.	Britain.	7. 8. H. ¼.	Eng. bot. v. 8. t. 523.	
4 párvula. s.s. (bl.) slender.	N. America. 1823.	— H. ¼.	Eng. bot. v. 8. t. 524.	
5 havanénis. s.s. (wh.bl.) Havannah.	Havannah. 1793.	5. 6. S. ¼.	Hook. ex. flor. t. 106.	
6 húmilib. B.P.	(pu.) dwarf.	N. S. W. 1822.	6. 8. G. ¼.	Jacq. obs. 2. t. 29.
7 grandiflòra. s.s. (re.ye.) large-flowered.	Siberia. 1804.	7. 8. H. ¼.	Bot. mag. t. 635.	
8 lateriflòra. s.s. (pu.) Virginian.	N. America. 1752.	6. 9. H. ¼.		
9 nervòsa. s.s. (bl.) nerved-leaved.	1827.	— H. ¼.		
10 orientalís. s.s. (ye.) yellow-flower'd.	Levant. 1729.	7. 9. H. ¼.	Flor. græc. t. 580.	
11 altàica. F.	(bl.wh.) Altay.	Siberia. 1816.	— H. ¼.	Swt. br. fl. gar. t. 45.
12 víscida. s.s. (bl.wh.) viscid.	Iberia. 1820.	— H. ¼.		
13 variegàta. s.s. (st.bl.) variegated.	— H. ¼.		
14 alpina. s.s. (bl.) Alpine.	Hungary. 1752.	6. 10. H. ¼.	Swt. br. fl. gar. t. 90.	
15 peregrina. s.s. (d.pu.) Florentine.	Italy. 1783.	— H. ¼.	Flor. græc. t. 582.	
16 altíssima. s.s. (d.pu.) tall.	Levant. 1731.	7. 8. H. ¼.		
17 pálida. s.s. (pa.) pale-flowered.	Tauria. 1824.	— H. ¼.		
18 Colúmnæ. s.s. (d.pu.) heart-leaved.	Italy. 1806.	6. 8. H. ¼.	Swt. br. fl. gar. t. 52.	
19 rubicúnda. s.s. (re.) rubicund.	1815.	— H. ¼.	
20 nigréscens. s.s. (d.pu.) dark.	Crete. 1826.	— F. ¼.		
21 àlbida. s.s. (wh.) hairy white.	Levant. 1771.	6. 7. H. ¼.	Flor. græc. t. 581.	
22 lupulina. s.s. (st.) Tartarian.	Tartary. 1739.	6. 9. H. ¼.	Schmidel ic. t. 73.	
23 integrifòlia. s.s. (bl.) entire-leaved.	N. America. 1731.	— H. ¼.	Pluk. alm. t. 441. f. 6.	
24 hyssopifòlia. W.	(bl.) Hyssop-leaved.	— H. ¼.	
25 purpuráscens. s.s. (pu.) purplish.	W. Indies. 1823.	— S. ¼.		
26 caroliniana. s.s. (pu.) Carolina.	Carolina. 1811.	— H. ¼.	Lam. ill. t. 515. f. 3.	
27 macrántha. F.	(pu.) large-flowered.	Dahuria. 1827.	— H. ¼.	
28 pilòsa. s.s. (d.pu.) pilose.	N. America. 1800.	— H. ¼.	Andr. reposit. t. 494.	
29 serrata. s.s. (d.pu.) saw-leaved.	— H. ¼.		
30 scordifòlia. F. (pu.wh.) Scordium-leaved.	Siberia. 1823.	— H. ¼.		
31 vérna. Bess. (wh.pu.) vernal.	5. 6. H. ¼.		
SA'LVIA. R.S.	SAGE.	Diandria Monogynia. L.		
1 pomifera. R.S.	(li.) Apple-bearing.	Candia. 1699.	7. 8. H. ¼.	Flor. græc. t. 15.
2 calycina. R.S.	(li.) broad-calyxed.	Greece. 1823.	— G. ½.	— t. 16.
3 canariénis. R.S.	(li.) Canary.	Canaries. 1697.	6. 9. G. ½.	Trew. pl. rar. 2. t. 19.
4 àurea. R.S.	(br.) golden.	C. B. S. 1731.	4. 11. G. ½.	Bot. mag. t. 182.
5 dentàta. R.S.	(bl.) tooth-leaved.	1774. 12. 1. G. ½.	
6 Réglia. R.S.	(sc.) Regla.	Mexico. 1828.	9. 11. G. ½.	Cavan. icon. 5. t. 455.
7 spléndens. s.s.	(sc.) splendid.	Brazil. 1823.	8. 3. S. ½.	Bot. reg. t. 687.
8 bracteàta. s.s.	(li.) long-bracted.	Syria. 1788.	6. 8. G. ½.	Bot. mag. t. 2320.
9 pilàntha. L.en.	(bl.) hairy-flowered.	1823. 6. 9. G. ½.	
10 incarnàta. R.S.	(fl.) flesh-coloured.	Levant. 1729.	— G. ¼.	Pluk. phyt. t. 194. f. 6.
11 interrúpta. R.S. (wh.bl.) interrupted.	Barbary.	1825.	5. 6. H. ½.	Swt. br. fl. gar. t. 169.
12 pinnàta. R.S.	(pu.) winged-leaved.	Levant. 1731.	7. H. ½.	Boer. lugd. I. t. 167.
13 ríngens. F.G.	(li.) ringent.	1827.	Flor. græc. t. 18.
14 rosaefòlia. R.S.	(pu.) Rose-leaved.	— G. ½.	Smith ic. ined. 1. t. 5.
15 scabiosæfòlia. R.S. (wh.li.) Scabious-leav'd.	Peru.	1826.	— G. ¼.	Jacq. f. ecl. t. 3.
16 Habliziàna. R.S. (w.re.) Taurian.	Tauria.	1795.	8. H. ½.	Bot. mag. t. 1429.
17 vulnerariefòlia. R.S. (w.bl.) Armenian.	Armenia.	1829.	6. 9. F. ½.	Jacq. f. ecl. t. 13.
18 lanceolàta. R.S.	(bl.) lance-leaved.	S. America.	1818.	Jacq. schœn. 3. t. 252.
19 hirsùta. R.S.	(bl.) hirsute.	Mexico.	1801.	Cavan. ic. 1. t. 24.
20 leucántha. R.S. (wh.) white-flowered.	1825.	7. 10. G. ¼.	

21	angustifolia. R.S. (<i>bl.</i>) narrow-leaved. <i>réptans</i> . Jacq. schœn. t. 319.	Mexico.	1806.	6. 7. G. 4.	Cavan. ic. 4. t. 317.
22	trichostemmoides. R.S. (<i>bl.</i>) Missouri.	Missouri.	1827.	6. 9. H. ⊖.	
23	azurea. R.S. (<i>bl.</i>) azure-flowered.	Carolina.	1806.	8. 9. F. 4.	Bot. mag. t. 1728.
24	bengalensis. F.I. (<i>wh.</i>) Bengal.	E. Indies.	1825.	6. 9. S. h.	
25	occidentalis. R.S. (<i>wh.</i>) West Indian.	W. Indies.	1820.	— S. 4.	
26	pseudococcinea. (sc.) pale scarlet.	S. America.	1797.	6. 8. S. h.	Jacq. ic. 2. t. 209.
27	amœna. B.M. (<i>bl.</i>) fine blue. <i>Boosiâna</i> . Jacq. f. ecl. I. t. 47. <i>amœna</i> , B.R. 446.	W. Indies.	1793.	9. 12. S. h.	Bot. mag. t. 1294.
28	rôsea. R.S. (<i>ro.</i>) rose-coloured.	E. Indies.	1816.	6. 9. S. h.	
29	tubifera. R.S. (<i>sc.</i>) long-tubed.	Mexico.	1824.	— S. 4.	Cavan. ic. 1. t. 25.
30	papilionacea. R.S. (<i>bl.</i>) butterfly-like.	New Spain.	1825.	7. 10. S. h.	— 4. t. 319.
31	mexicana. R.S. (<i>bl.</i>) Mexican.	Mexico.	1724.	5. 7. G. h.	— 1. t. 26.
32	chamaedryoides. R.S. (<i>re.</i>) Germanander-like.	—	1795.	6. 9. G. h.	Bot. mag. t. 808.
33	cæsia. R.S. (<i>pa.bl.</i>) grey.	S. America.	1813.	— G. h.	
34	involucrata. R.S. (<i>pu.</i>) large-bracted.	Mexico.	1825.	8. 11. G. h.	Bot. reg. t. 1205.
35	purpurea. R.S. (<i>pu.</i>) purple.	—	—	5. 7. S. 4.	Cavan. ic. 2. t. 166.
36	hispanica. R.S. (<i>bl.</i>) Spanish.	Spain.	1739.	6. 8. H. ⊖.	Bot. reg. t. 359.
37	præcox. s.s. (<i>pu.</i>) early-flowered.	N. Africa.	1826.	3. 6. F. 4.	
38	scordioïdes. R.S. (<i>bl.</i>) glandular.	—	1824.	5. 8. S. 4.	
39	papillosa. R.S. (<i>wh.</i>) papillose.	—	1827.	6. 8. G. h.	
40	serotina. R.S. (<i>bl.</i>) late-flowering.	Chio.	1803.	8. 10. G. h.	Jacq. ic. 1. t. 3.
41	dominica. R.S. (<i>wh.</i>) Dominican.	W. Indies.	1759.	7. S. 4.	Swartz. obs. p. 18. t. 1. f. 1.
42	tenella. R.S. (<i>bl.</i>) slender.	Jamaica.	1821.	6. 9. S. ⊖.	Swartz. icon. t. 2.
43	tiliaeïflora. R.S. (<i>bl.</i>) Lime-leaved.	S. America.	1793.	6. 8. S. 4.	Jacq. schœn. 3. t. 254.
44	polystachya. R.S. (<i>bl.</i>) many-spiked.	New Spain.	1822.	9. 12. G. h.	Cavan. ic. 1. t. 27.
45	lamiaïflora. R.S. (<i>bl.</i>) Lamium-leaved.	S. America.	1826.	— G. h.	Jacq. schœn. 3. t. 318.
46	micrantha. R.S. (<i>wh.</i>) small-flowered.	—	1818.	6. 8. S. 4.	
47	leoniroides. R.S. (<i>sc.</i>) shining-leaved. <i>formosa</i> . B.M. 376.	Peru.	1783.	4. 10. G. h.	Lam. ill. t. 20. f. 3.
48	fulgens. R.S. (<i>sc.</i>) glossy.	Mexico.	1827.	1. 12. G. h.	
49	coccinea. R.S. (<i>sc.</i>) scarlet-flowered.	S. America.	1774.	4. 10. G. h.	Swt. br. fl. gar. s. 2.
50	pulchella. R.S. (<i>sc.</i>) pretty.	—	—	8. 12. G. h.	Mirb. an. m. 15. t. 15. f. 8.
51	amethystina. R.S. (<i>bl.</i>) amethystine.	—	1824.	6. 9. S. h.	Coll. hort. rip. t. 16.
52	amarissima. R.S. (<i>bl.</i>) bitter.	Mexico.	1803.	7. 8. G. 4.	Smith ic. ined. 2. t. 27.
53	glutinosa. R.S. (<i>st.</i>) glutinous.	Europe.	1596.	6. 9. H. 4.	Bot. reg. t. 347.
54	nubicola. B.F.G. (<i>st.</i>) Nepaul.	Nepaul.	1824.	8. 10. H. 4.	Sabb. hort. rom. t. 3.
55	ægyptiaca. R.S. (<i>wh.</i>) Egyptian.	Egypt.	1770.	6. 7. H. ⊖.	Swt. br. fl. gar. t. 140.
56	crética. R.S. (<i>vi.</i>) Cretan.	Crete.	1760.	6. 8. F. h.	Jacq. vind. 2. t. 108.
57	paniculata. R.S. (<i>pu.</i>) paniced.	C. B. S.	1758.	— G. h.	Rivin. monop. t. 128.
58	africana. R.S. (<i>bl.</i>) African.	—	1731.	4. 6. G. h.	Mill. ic. t. 225. f. 1.
59	colorata. R.S. (<i>bl. cr.</i>) coloured-calyxed.	—	1758.	7. 8. G. h.	Comm. hort. 2. t. 91.
60	officinalis. R.S. (<i>bl.</i>) garden.	S. Europe.	1597.	6. 7. H. h.	Mill. ic. t. 225. f. 2.
61	chromatica. Hg. (<i>bl.</i>) variegated.	—	—	— H. h.	Lam. ill. t. 20. f. 1.
62	tenuior. Df. (<i>bl.</i>) narrow-leaved.	Spain.	—	— H. h.	
63	hispanorum. Lg. (<i>bl.</i>) Spanish.	—	—	— H. h.	
64	lavandulæfolia. R.S. (<i>bl.</i>) Lavender-leaved.	—	—	— H. h.	
65	rosmarinifolia. Lg. (<i>bl.</i>) Rosemary-leaved.	—	—	— H. h.	
66	Spielmanni. R.S. (<i>bl.</i>) Spielman's.	Caucasus.	1813.	— H. 4.	Jacq. f. ecl. 2. t. 15.
67	phlomoides. R.S. (<i>pu.</i>) Mullein-leaved.	Spain.	1805.	5. 6. H. ♂.	Røm. script. his. t. 1. f. 1.
68	urticifolia. R.S. (<i>bl.</i>) Nettle-leaved.	N. America.	1799.	6. 7. H. 4.	Moris. his. 3. t. 13. f. 31.
69	élégans. R.S. (<i>cr.</i>) elegant.	Mexico.	1820.	7. 10. G. h.	H. et B. pl. am. 2. t. 144.
	<i>incarnata</i> . K.S. non w.				
70	lævigata. K.S. (<i>pu.</i>) smooth.	—	1826.	— G. h.	H. et B. pl. am. 2. t. 147.
71	bullata. W. en. (<i>re.</i>) blistered.	Spain.	1804.	7. 8. H. 4.	Jacq. f. ecl. t. 38.
	<i>lusitanica</i> . J.E. <i>elongata</i> . Spr.				
72	rugosa. R.S. (<i>wh.</i>) wrinkle-leaved.	C. B. S.	1775.	— H. h.	
73	verticillata. R.S. (<i>li.</i>) whorled.	Germany.	1658.	6. 11. H. 4.	Mirb. an. m. 15. t. 15. f. 10.
	β <i>parviflora</i> . (<i>li.</i>) small-flowered.		—	— H. 4.	
74	indica. R.S. (<i>bl. wh.</i>) Indian.	India.	1731.	5. 7. H. 4.	Bot. mag. t. 395.
75	Tenori. R.S. (<i>bl.</i>) Tenore's.	Italy.	1820.	5. 8. H. 4.	Swt. br. fl. gar. t. 26.
76	candidissima. R.S. (<i>p. bl.</i>) hoary-leaved. <i>crassifolia</i> . F.G. non Cav.	Levant.	1804.	6. 7. G. 4.	Flor. græc. t. 26.
77	verbascifolia. R.S. (<i>wh. ye.</i>) Verbascum-ld.	Iberia.	1825.	— H. 4.	
78	odorata. R.S. (<i>wh.</i>) sweet-scented.	Bagdad.	1804.	7. F. h.	Jacq. f. ecl. t. 16.
79	compræssa. R.S. (<i>wh.</i>) compressed.	Levant.	1816.	7. 8. H. 4.	Venten. cels. t. 59.
80	mollis. R.S. (<i>ro.</i>) soft.	Siberia.	1596.	6. 10. H. 4.	Jacq. f. ecl. t. 37.
81	grandiflora. R.S. (<i>bl.</i>) great-flowered.	S. Europe.	1616.	6. 9. H. h.	— t. 36.
82	crassifolia. R.S. (<i>bl.</i>) thick-leaved.	Barbary.	1804.	— F. h.	

83 pratensis. R.S. (bl.) meadow.	England.	5. 11. H. ȝ.	Eng. bot. v. 3. t. 153.	
84 variegata. R.S. (bl. wh.) variegated.	Hungary.	1814.	6. 8. H. ȝ.	Barrel. ic. t. 185.	
85 haematodes. R.S. (bl.) bloody-veined.	Italy.	1699.	7. 8. H. ȝ.		
86 nepetifolia. R.S. (bl.) Catmint-leaved.	1823.	— H. ♂.		
87 aspidophylla. R.S. (bl.) slender-leaved.	Peru.	1824.	7. 10. H. ♂.		
88 lineatifolia. Lg. (bl.) line-leaved.	Mexico.	1823.	6. 9. G. ḥ.		
89 melissodora. Ig. (bl.) balm-scented.	New Spain.	1826.	— G. ḥ.		
90 agglutinans. Lg. (sc.) clammy-stalked.	1827.	— G. ḥ.		
91 viscosa. R.S. (wh. cr.) clammy.	Italy.	1773.	5. 6. H. ȝ.	Jacq. ic. 1. t. 5.	
92 disermis. R.S. (wh.) long-spiked.	Syria.	—	7. H. ȝ.	Barrel. ic. 187.	
93 nutans. R.S. (bl.) nodding.	Russia.	1780.	6. 8. H. ȝ.	Bot. mag. t. 2436.	
94 betonicæfolia. R.S. (bl.) Betony-leaved.	1804.	6. 7. H. ȝ.		
95 pendula. R.S. (bl.) drooping.	—	6. 8. H. ȝ.		
β grandiflora. (d.bl.) large-flowered.	—	— H. ȝ.		
96 amplexicaulis. R.S. (li.) stem-clasping.	Levant.	1813.	7. 9. H. ȝ.		
97 parviflora. R.S. (st.) small-flowered.	—	1816.	— F. ȝ.		
98 austriaca. R.S. (st.) Austrian.	Austria.	1776.	6. 7. H. ȝ.	Jacq. aust. 2. t. 112.	
99 syriaca. R.S. (wh.) Syrian.	Levant.	1759.	7. G. ḥ.	Arduin. spec. 1. t. 1.	
100 nubia. R.S. (bl.) Nubian.	Africa.	1784.	6. 7. G. ȝ.	Murr. c. gott. 1778. t. 3.	
101 virgata. R.S. (wh. ro. bl.) long-branched.	Armenia.	1758.	7. 11. H. ȝ.	Jacq. schœn. 1. t. 37.	
102 campêstris. R.S. (bl.) field.	Tauria.	1813.	6. 7. H. ȝ.		
103 sylvestris. R.S. (bl.) wood.	Germany.	1759.	6. 10. H. ȝ.	Jacq. aust. 3. t. 212.	
104 nemorosa. R.S. (va.) spear-leaved.	1728.	— H. ȝ.		
105 pâtula. R.S. (bh.) spreading.	Portugal.	1805.	5. 7. H. ȝ.		
106 tingitana. R.S. (st.) Tangier.	Barbary.	1796.	7. G. ḥ.	Rivin. monop. 1. t. 52.	
107 Sclarea. R.S. (bh.) common Clary.	Italy.	1562.	7. 9. H. ♂.	Flor. græc. 1. t. 25.	
108 spinosa. R.S. (wh.) thorny-calyxed.	Egypt.	1789.	6. G. ♂.	Jacq. ic. 1. t. 7.	
109 Æthiopis. R.S. (wh.) woolly.	Austria.	1570.	5. 6. H. ♂.	Jacq. aust. 3. t. 211.	
110 argentea. R.S. (wh.) silvery.	Crete.	1759.	5. 7. H. ♂.	Flor. græc. 1. t. 27.	
111 applanata. R.S. (wh.) short-spined.	1818.	6. 8. H. ♂.		
112 Horminum. R.S. (pu.) annual Clary. α violacea.	S. Europe.	1596.	— H. ♂.	Flor. græc. 1. t. 20.	
β rûbra.	(bl.) purple-topped. (re.) red-topped.	—	— H. ♂.		
113 viridis. R.S. (li.) green topped.	Italy.	1759.	— H. ♂.		
114 pyramidalis. R.S. (li.) pyramidal.	—	1825.	— H. ȝ.	Flor. græc. 1. t. 19.	
115 rûbra. s.s.	(re.) red-flowered.	Egypt.	1827.	6. 8. H. ȝ.	
116 Simsiana. B.R. (bl. wh.) Sims's.	Russia.	1820.	6. 7. H. ȝ.	Bot. reg. t. 1003.	
117 verbenacea. R.S. (bl.) wild clary.	Britain.	6. 10. H. ȝ.	Eng. bot. v. 3. t. 154.	
118 oblongata. J.E. (bl.) oblong-leaved.	Tauria.	1820.	— H. ȝ.	Jacq. f. ecl. 1. t. 14.	
119 Sibthorpii. F.G. (li.) Sibthorp's.	Levant.	1829.	6. 9. F. ȝ.	Flor. græc. t. 22.	
120 triloba. R.S.	S. Europe.	1596.	6. 7. H. ḥ.	— 1. t. 17.	
121 Clusi. R.S. (wh.) Clusius's.	Levant.	1827.	6. 9. F. ḥ.	Jacq. schœn. 2. t. 195.	
122 gigantea. R.S. (wh.) giant.	1816.	6. 9. H. ♂.		
123 scorodoniamæfolia. R.S. villous.	1824.	— F. ḥ.		
124 lyrata. R.S. (pu.) lyre-leaved.	N. America.	1728.	— H. ȝ.	Moris. s. 11. t. 13. f. 27.	
125 abyssinica. R.S. (pu.) Abyssinian.	Abyssinia.	1775.	6. 7. G. ȝ.	Jacq. ic. 1. t. 6.	
126 nilotica. R.S. (bl.) Nilotic.	Egypt.	1780.	6. 8. H. ♂.	Jacq. vind. 3. t. 92.	
127 Forskohlii. R.S. (bl.) Forskohl's.	Levant.	1800.	— H. ȝ.	Bot. mag. t. 988.	
128 napifolia. R.S. (li.) Rape-leaved.	Italy.	1776.	6. 7. H. ȝ.	Jacq. vind. 2. t. 262.	
129 aurita. R.S. (li.) ear-leaved.	C. B. S.	1795.	5. 6. G. ḥ.		
130 bicolor. R.S. (bl. ye.) two-coloured.	Barbary.	1793.	6. 7. H. ȝ.	Bot. mag. t. 1774.	
131 Barrelieri. R.S. (bl.) Barrelier's.	S. Europe.	1821.	5. 8. H. ȝ.	Barrel. ic. 186.	
132 lacinata. R.S. (bl. wh.) jagged-leaved.	1818.	— H. ȝ.	Jacq. schœn. 1. t. 8.	
133 runcinata. R.S. (bl.) runcinate.	C. B. S.	1795.	— G. ḥ.		
134 scabra. R.S. (pu.) rough-leaved.	—	1774.	4. 9. G. ḥ.	Barrel. ic. 220.	
135 erosa. R.S. (st.) erose-leaved.	1823.	6. 9. H. ♂.	Flor. græc. 1. t. 23.	
136 polymorpha. R.S. (bl.) variable.	Portugal.	1821.	— H. ȝ.	— 1. t. 24.	
137 multifida. F.G. (pa. bl.) multifid.	S. Europe.	1823.	5. 6. H. ȝ.	Pluk. alm. t. 194. f. 5.	
138 clandestina. R.S. (li.) cut-leaved.	Italy.	1739.	4. 7. H. ♂.	Arduin. spec. 2. t. 2.	
139 ceratophylla. R.S. (li.) stag's-horn-leav'd.	Persia.	1699.	7. 8. H. ♂.		
140 ceratophylloides. (li.) branching.	Egypt.	1771.	6. 8. H. ♂.		
141 prismatica. R.S. (bl.) slender-spiked.	Mexico.	1822.	— H. ♂.		
COLLINS'ONIA. R.S. COLLINS'ONIA.	Diandria Monogynia. L.				
1 canadensis. R.S. (ye.) Nettle-leaved.	N. America.	1735.	8. 10. H. ȝ.	Linn. hort. clif. t. 5.	
2 scabriuscula. R.S. (re. ye.) rough-stalked.	Florida.	1776.	7. 9. H. ȝ.	Bot. mag. t. 1213.	
3 ovalis. R.S. (ye.) oval-leaved.	Carolina.	1812.	8. H. ȝ.		
4 tuberosa. M. (ye.) tuberous.	—	1806.	— H. ȝ.		
5 anisata. R.S. (ye.) Anise-scented.	—	—	10. H. ȝ.		

Tribus VI. PRASIEÆ. *Benth. in Bot. reg.*

PRASIUM. S.S.	PRASIUM. Didynamia Gymnospermia. L.				
1 majus. s.s. (wh.) great.	S. Europe. 1699.	6. 8. G. ḥ.	Flor. græc. t. 584.		
2 minus. s.s. (wh.) smaller.	1752.	— F. ḥ.			
PHRYMA. S.S.	PHRYMA. Didynamia Gymnospermia. L.				
leptoſtachya. s.s. (wh.) slender-spiked.	N. America. 1802.	8. 9. H. ȳ.	Lam. ill. t. 516.		

Tribus VII. OCYMOIDEÆ. *Benth. in Bot. reg.*

MOSCHO'SMA. B.R.	MOSCHO'SMA. Didynamia Gymnospermia.				
1 ocymoïdes. B.R. (wh.) Basil-like.	1823.	6. 8. S. ⊙.			
<i>Lumnitzera ocymoïdes</i> . s.s. <i>O'cymum polycladum</i> . Link.					
2 polystachya. B.R. (wh.) many-spiked.	E. Indies. 1783.	7. 8. S. ⊙.	Murr. c. gott. 3. t. 3.		
O'CIMUM. W.	BASIL. Didynamia Gymnospermia. L.				
1 thyrsiflōrum. s.s. (wh.) thyrsse-flowered.	E. Indies. 1806.	7. 8. S. ♂.	Jacq. vind. 3. t. 72.		
2 gratissimum. s.s. (wh.) shrubby.	1752.	— S. ḥ.	Jacq. ic. 3. t. 495.		
3 Basiliicum. w. (wh.) common sweet.	India. 1548.	— H. ⊙.	Blackw. t. 104.		
4 pilōsum. W.en. (wh.) hairy.	1816.	7. 9. H. ⊙.			
5 micrānthum. s.s. (wh.) small-flowered.	—	— S. ⊙.			
6 viride. s.s. (wh.) green.	E. Indies. —	— S. ḥ.			
7 suave. s.s. (wh.) sweet-scented.	—	— S. ḥ.			
8 cānum. B.M. (wh.) hoary.	China. 1822.	— H. ⊙.	Bot. mag. t. 2452.		
9 sāctum. s.s. (wh.) purple-stalked.	E. Indies. 1758.	— S. ⊙.	Rheed. mal. 10. t. 92.		
10 febrifugum. s.s. (wh.) Feverwort.	Sierra Leon. 1821.	— S. ⊙.	Bot. reg. t. 753.		
11 ábum. s.s. (wh.) large white-fld.	E. Indies. 1816.	— S. ⊙.			
12 americanum. s.s. (wh.) American.	S. America. 1789.	7. 8. S. ⊙.	Jacq. vind. 3. t. 86.		
13 caryophyllātum. H.B. (w.) Clove-scented.	E. Indies. 1817.	— S. ȳ.			
14 tenuiflōrum. w. (li.) slender-spiked.	—	— S. ♂.	Rumph. amb. 5.t.92.f.2.		
15 menthoïdes. s.s. (wh.) Mint-leaved.	—	— S. ⊙.			
16 inodōrum. Lam. (wh.) scentless.	—	— S. ⊙.			
17 cristātum. H.B. (wh.) crested.	—	— S. ⊙.			
18 grandiflōrum. s.s. (wh.) great-flowered.	Abyssinia. 1802.	9. 10. G. ḥ.	L'Herit. s. nov. t. 43.		
19 mīnimum. s.s. (wh.) bush.	Ceylon. 1573.	— H. ⊙.	Schk. handb. 2. t. 166.		
20 carnōsum. s.s. (wh.) thick-leaved.	Monte Video. 1828.	6. 8. H. ⊙.			
ORTHOSIPHON. B.R. ORTHOSIPHON.	Didynamia Gymnospermia.				
1 adscēndens. B.R. (wh.) ascending.	E. Indies. 1828.	6. 8. S. ȳ.			
<i>O'cymum adscēndens</i> . w.					
2 ásperum. B.R. (wh.) rough.	—	1827. — S. ȳ.			
<i>O'cymum ásperum</i> . Roth. <i>Plectránthus ásper</i> . s.s.					
COLEUS. B.R.	COLEUS. Didynamia Gymnospermia.				
1 Forskhlæi. B.R. (vi. wh.) Forskhol's.	Abyssinia. 1806.	10. 11. S. ḥ.	Bot. mag. t. 2036.		
<i>Plectránthus Forskhlai</i> . B.M. <i>barbátus</i> . Andr. reposit. t. 594.					
2 comosus. B.R. (bl.) tufted.	Nepaul.	1820. 6. 9. G. ḥ.	Bot. mag. t. 2318.		
3 scutellarioides. B.R. (bl.w.) Scullcap-like.	E. Indies.	1764. 7. 8. S. ⊙.	— t. 1416.		
4 amboficus. Lour. (bl.) Amboyna.	—	1826. — S. ḥ.	Rumph. amb. 8. t. 72.		
PLECTRA'NTHUS. S.S. PLECTRA'NTHUS.	Didynamia Gymnospermia. S.S.				
1 fruticosus. s.s. (bl.) shrubby.	C. B. S.	1774. 6. 9. G. ḥ.	L'Herit. stirp. 85. t. 41.		
2 galeatus. s.s. (bl.) helmetted.	Nepaul.	1820. 7. 8. S. ⊙.			
3 graveolens. B.P. (bl.) strong-scented.	N. S. W.	— — G. ḥ.			
4 australis. B.P. (bl.) New Holland.	—	1823. — G. ⊙.			
5 incānus. s.s. (bl.) hoary.	Nepaul.	— — G. ȳ.			
6 monachòrum. s.s. (bh.) monk's basil.	E. Indies.	1796. — S. ⊙.			
<i>O'cymum monachòrum</i> . w.					
7 móllis. s.s. (li.) heart-leaved.	—	1781. 9. 10. S. ⊙.			
8 strobiliferus. H.B. (bl.) thick-spiked.	—	1804. 6. 9. S. ⊙.	Rheed. mal. 10. t. 90.		
9 secundus. H.B. (bl.) side-flowering.	—	1816. — S. ⊙.			
10 punctatus. s.s. (wh.) dotted.	Africa.	1775. 1. 5. G. ♂.	L'Herit. stirp. 1. t. 41.		
11 rotundifolius. s.s. (bl.) round-leaved.	Mauritius.	1823. 6. 8. S. ⊙.			
12 parviflōrus. s.s. (bl.) small-flowered.	S. America.	1805. 6. 9. G. ḥ.			
13 cordifolius. D.P. (bl.) heart-leaved.	Nepaul.	1823. — G. ḥ.			
14 rubicundus. D.P. (pu.) rubicund.	—	1826. — G. ḥ.			
15 virgatus. D.P. (bl.) slender.	—	— G. ḥ.			
16 coloratus. D.P. (bl.) coloured bracted.	—	1823. — G. ḥ.			

17 Coëtsa. D.P. (bl.) canescent.	Nepaul.	1823.	6. 9. G. ¼.	
18 ternifoliūs. D.P. (bl.) three-leaved.	———	1825.	——— G. ½.	
19 ternatūs. B.M. (bl.) Omime-root.	Madagascar.	1820.	11. S. ¼.	Bot. mag. t. 2460.
GENIO'SPORUM. B.R. GENIO'SPORUM.	Didynamia Gymnospermia.			
prostratum. B.R. (pa.bl.) prostrate.	E. Indies.	1812.	6. 9. S. Ⓛ.	
O'cymum prostratum. w.				
ACROCE'PHALUS. B.R. ACROCE'PHALUS.	Didynamia Gymnospermia.			
1 capitātus. B.R. (wh.) headed.	China.	1806.	7. 9. H. Ⓛ.	
Lumnitzera capitata. s.s. O'cymum capitellatum. w.				
2 villōsus. B.R. (wh.) villous.	E. Indies.	1817.	7. 8. S. ¼.	
ANISOCH'ILUS. B.R. ANISOCH'ILUS.	Didynamia Gymnospermia.			
carnosās. B.R. (li.) thick-leaved.	E. Indies.	1788.	6. 7. S. Ⓛ.	
Lavändula carnosa. w. Plectránthus dùbius. s.s.				
PYCNOST'ACHYS. PYCNOST'ACHYS.	Didynamia Gymnospermia.			
cœrûlea. H.E.F. (bl.) blue-flowered.	Madagascar.	1825.	8. 10. F. Ⓛ.	
ÆOLLA'NTHUS. M.P. ÆOLLA'NTHUS.	Didynamia Gymnospermia.			
suavèolens. M.P.B. (wh.) sweet-scented.	Brazil.	1829.	7. 10. S. Ⓛ.	
HYPTIS. S.S. HYPTIS.	Didynamia Gymnospermia. S.S.			
1 ebracteata. s.s. (bl.) small-headed.	W. Indies.	1778.	10. 1. S. Ⓛ.	
Balòta suavèolens. Jacq. schœnb. 3. t. 42.				
2 radiata. s.s. (wh.) Carolina.	Carolina.	1690.	6. 7. G. ¼.	
3 capitata. s.s. (wh.) Jamaica.	W. Indies.	1714.	——— S. Ⓛ.	
4 polyanthes. s.s. (wh.) many-flowered.	S. America.	1819.	——— S. Ⓛ.	
5 brèvipes. s.s. (li.) short-peduncled.	W. Indies.	1822.	——— S. Ⓛ.	
6 recurvata. s.s. (wh.) recurved.	Cayenne.	———	——— S. ¼.	
7 álbida. s.s. (bl.) white-leaved.	Mexico.	1825.	——— S. ½.	
8 pérsica. P.S. (st.) Persian.	Persia.	1800.	7. G. ½.	
Brotéra pérsica. L.T.				
9 spicata. s.s. (bl.) spiked.	S. America.	1819.	7. 9. G. Ⓛ.	
10 verticillata. s.s. (wh.) whorled.	———	1824.	——— S. Ⓛ.	
11 Plumieri. K.S. (bl.) Plumier's.	———	1819.	——— S. Ⓛ.	
12 stachyodes. s.s. (wh.) whorl-spiked.	Brazil.	1820.	——— S. Ⓛ.	
13 pectinata. s.s. (ye.) pectinated.	W. Indies.	1776.	12. 1. S. ¼.	
PROSTANTH'ERA. B.P. PROSTANTH'ERA.	Didynamia Gymnospermia. S.S.			
1 lasiánthos. B.P. (wh.) villous-flowered.	N. S. W.	1808.	6. 7. G. ½.	
2 cœrûlea. B.P. (bl.) blue-flowered.	———	1824.	4. 7. G. ½.	
3 prunelloïdes. B.P. (pu.) Prunella-like.	———	1826.	——— G. ½.	
4 incisa. B.P. (wh.) cut-leaved.	———	1825.	——— G. ½.	
5 lineáris. B.P. (wh.) linear-leaved.	———	1823.	——— G. ½.	
6 dentículata. B.P. (wh.) tooth-leaved.	———	———	——— G. ½.	
7 marifòlia. B.P. (wh.) Marum-leaved.	———	1827.	4. 6. G. ½.	
8 saxícola. B.P. (wh.) rock.	———	1823.	——— G. ½.	
9 violacea. B.P. (bl.) violet-coloured.	———	———	——— G. ½.	

Bot. reg. t. 1072.

ORDO CLIV.

SELAGINEÆ. Choisy mem. fam. Selag. 1823. p. 19.

POLYC'ENIA. C.S. POLYC'ENIA.	Didynamia Gymnospermia.			
hebenstretioides. c.s. (w.) spike-flowered.	C. B. S.	1816.	8. 10. G. Ⓛ.	Chois. sel. t. 2. f. 1.
HEBENSTREITIA. C.S. HEBENSTREITIA.	Didynamia Angiospermia.			
1 dentata. w. (wh.) dwarf tooth-ld.	C. B. S.	1739.	5. 9. G. Ⓛ.	Bot. mag. t. 483.
2 integrifolia. w. (ye.) golden.	———	1792.	4. 6. G. ½.	Andr. reposit. t. 252.
aurea. Andr.				
3 fruticosa. B.M. (wh.) frutescent.	———	1816.	5. 11. G. Ⓛ.	Bot. mag. t. 1970.
4 albiflora. L.en. (wh.) white-flowered.	———	1815.	——— G. ½.	
5 chamædryfòlia. L.en. Chamædrys-leav'd.	———	1816.	——— G. ½.	
6 tenuifolia. H.H. (wh.) slender-leaved.	———	1826.	——— G. Ⓛ.	
7 scabra. w. (wh.) rough-leaved.	———	———	5. 9. G. ½.	
8 cordata. w. (wh.re.) heart-leaved.	———	1774.	7. 8. G. ½.	
DISCHI'SMA. C.S. DISCHI'SMA.	Didynamia Angiospermia.			
1 ciliatūm. c.s. (wh.) ciliated.	C. B. S.	1815.	5. 7. G. Ⓛ.	
2 erinoïdes. (wh.) Erinus-leaved.	———	1816.	5. 11. G. Ⓛ.	
Hebenstreitia erinoïdes. w.				

3 hispidum. (wh.) long-spiked. <i>Hebenstreitia hispida</i> , Lam.	C. B. S.	1816.	5. 11. G. ⊙.	Burm. afr. 109. t. 41. f. 1.
4 spicatum. c.s. (wh.) spiked. 5 cattatum. c.s. (ye.) headed.	1815. — 1822.	5. 7. G. ⊙. — G. ⊙.		
AGATHE'LPIS. C.S. <i>AGATHE'LPIS.</i> Diandria Monogynia.				
1 angustifolia. c.s. (wh.) narrow-leaved. <i>Eranthemum angustifolium</i> . L. <i>Selago angustifolia</i> . w.	C. B. S.	1823.	5. 8. G. ™.	Burm. afr. t. 47. f. 3.
2 parvifolium. c.s. (wh.) small-leaved. MICRODON. C.S. <i>MICRODON.</i> Didynamia Gymnospermia.	C. B. S.	1816.	— G. ™.	Lam. ill. t. 17. f. 2.
1 lúcidum. c.s. (wh.) glossy. 2 ovatum. c.s. (wh.) oval-headed. <i>Selago ovata</i> , B.M.	C. B. S.	1812. — 1774.	6. 7. G. ™. — G. ™.	Venten. malm. t. 26. Bot. mag. t. 186.
SELAGO. C.S. <i>SELAGO.</i> Didynamia Gymnospermia. L.				
1 micrantha. c.s. (ye.) small-flowered. 2 divaricata. c.s. (wh.) spreading. 3 fruticosa. c.s. (wh.) frutescent. 4 minutissima. c.s. (ye.) small. 5 corymbosa. c.s. (wh.) fine-leaved. 6 cinerea. c.s. (vi.) cinereous. 7 hispida. c.s. (ye.) hispid. 8 adpressa. c.s. (wh.) close-flowered. 9 canescens. s.s. (wh.) canescent. 10 spinea. L.en. (wh.) reflex-leaved. 11 diffusa. s.s. (wh.) diffuse. 12 spicata. L.en. (wh.) spiked. 13 teretifolia. L.en. (wh.) cylinder-leaved. 14 rapunculoides. s.s. (vi.) Rampion-like. 15 fulvomaculata. L.en. (vi.) brown-spotted. 16 spuria. s.s. (vi.) linear-leaved. 17 ramulosa. L.en. (wh.) branching. 18 fasciculata. s.s. (bl.) cluster-flowered. 19 heterophylla. s.s. (wh.) various-leaved. 20 polystachya. s.s. (wh.) many-spiked. 21 bracteata. s.s. (wh.) imbricate-bracted. 22 polygaloides. s.s. (vi.) milkwort-like. 23 rotundifolia. s.s. (wh.) round-leaved. 24 pterophylla. Ot. winged-leaved. 25 ciliata. c.s. (wh.) fringed.	C. B. S.	1824. — 1816. — 1822. — 1816. — 1699. — 1816. — 1823. — 1816. — 1812. — 1816. — 1807. — 1816. — 1819. — 1790. — 1816. — 1779. — 1815. — 1774. — 1829. — 1823. — 1816. — 1807. — 1816. — 1828. — 1824.	5. 7. G. ™. — G. ™. 6. 8. G. ™. — G. ™. 6. 9. G. ™. — G. ™. 7. 11. G. ™. — G. ™. 7. 8. G. ™. — G. ™. 6. 9. G. ™. — G. ™. 6. 9. G. ™. — G. ™. 7. 11. G. ™. — G. ™. 7. 10. G. ™. — G. ™. 6. 9. G. ™. — G. ™. 6. 8. G. ™. — G. ™. 6. 9. G. ™. — G. ™. 7. 11. G. ™. — G. ™. 7. 9. G. ™. — G. ™. — G. ™.	Chois. sel. t. 3. Comm. hort. 2. t. 40. Chois. sel. t. 4. Chois. sel. t. 5.

ORDO CLV.

VERBENACEÆ. *Brown prodr. 1. p. 510.***CLERODE'NDRUM. B.P.CLERODE'NDRUM.** Didynamia Angiospermia. L.

1 pubescens. B.R. (wh.) downy-leaved.	W. Indies.	1824.	7. 9. S. ℡.	Bot. reg. t. 1035.
2 ligustrinum. s.s. (wh.) Privet-leaved.	Mauritius.	1789.	8. 11. S. ℡.	Jacq. col. sup. t. 5. f. 1.
3 inerme. s.s. (wh.) smooth.	E. Indies.	1692.	— S. ℡.	— t. 4. f. 1.
4 angustifolium. s.s. (wh.) narrow-leaved.	Jamaica.	1823.	— S. ℡.	Lam. ill. t. 544. f. 2.
5 coramendlinum. s.s. (wh.) Coramandel.	E. Indies.	1824.	— S. ℡.	
6 emirnense. B.M. (wh.) small-flowered.	Madagascar.	1823.	6. 12. S. ℡. —	Bot. mag. t. 2925.
<i>floribundum.</i> Bot. reg. t. 1035. non B.P.				
7 volubile. s.s. (wh.) climbing.	Guinea.	1822.	8. 11. S. ℡. —	
8 buxifolium. s.s. (wh.) Box-leaved.	1818.	7. 8. S. ℡.	
<i>Volkameria buxifolia</i> . W.en.				
9 heterophyllum. s.s. (wh.) various-leaved.	Mauritius.	1805.	8. 9. S. ℡.	Andr. deposit. t. 554.
<i>Volkameria angustifolia</i> . A.R.				
10 macrophyllum. B.M. (st. bl.) large-leaved.	1822.	— S. ℡.	Bot. mag. t. 2536.
11 serratum. s.s. (wh.) saw-leaved.	E. Indies.	—	— S. ℡.	Rheed. mal. 4. t. 29.
12 lividum. B.R. (wh.) livid-bracted.	China.	1824.	9. 11. S. ℡.	Bot. reg. t. 945.
13 cénum. Wal. (wh.) drooping.	E. Indies.	1822. S. ℡.	
13 Siphonanthus. s.s. (wh.) whorled-leaved.	1796. S. ℡.	Burm. ind. t. 43. f. 1.
<i>Siphonanthus indica</i> et <i>Oviéda mitis</i> . L.				
15 verticillatums. D.P. (wh.) four-leaved.	Nepaul.	1818. S. ℡.	
16 hastatum. B.R. (wh.) halbert-leaved.	E. Indies.	1824.	7. 11. S. ℡.	Bot. reg. t. 1307.

17	odoratum. D.P.	(re.) odoriferous.	Nepaul.	1820.	7. 8.	S. h.
18	fœtidum. D.P.	(wh.) strong-scented.	—	—	—	S. h.
19	cordatum. D.P.	(wh.) heart-leaved.	—	1826.	—	S. h.
20	nūtans. D.P.	(wh.) nodding.	—	—	—	S. h.
21	scândens. s.s.	(wh.) scandent.	Guinea.	1822.	7. 9.	S. h.
22	trichotomum. s.s.(wh.)	three-forked.	Japan.	1800.	—	G. h.
23	fortunatum. s.s. (wh.)	spear-leaved.	E. Indies.	1784.	7. 8.	S. h.
24	attenuatum. B.P.(wh.)	attenuated.	N. S. W.	1824.	5. 8.	G. h.
25	squamatum. s.s. (sc.)	scarlet,	China.	1790.	6. 9.	S. h.
26	tomentosum. s.s. (wh.)	downy.	N. S. W.	1794.	3. 5.	G. h.
27	infortunatum. s.s.(wh.)	long-flowered.	E. Indies.	—	—	S. h.
28	phlomoides. s.s. (wh.)	Phlomis like.	—	1820.	5. 7.	S. h.
29	frâgrans. s.s. (wh.)	fragrant.	China.	1790.	12. 8.	S. h.
	β multiplex.	(wh.) double-flowered.	—	—	—	S. h.
30	viscosum. s.s. (wh.)	clammy.	E. Indies.	1796.	5. 8.	S. h.
31	dentatum. H.B. (wh.)	tooth-leaved.	—	1826.	—	S. h.
32	nerifolium. H.B.(wh.)	Oleander-leaved.	—	1824.	—	S. h.
33	paniculatum. s.s. (sc.)	panicked.	Java.	1809.	7. 10.	S. h.
34	urticifolium. H.B.(wh.)	Nettle-leaved.	E. Indies.	1822.	—	S. h.
35	violaceum. L.en. (vi.)	violet-coloured.	—	—	—	S. h.
36	japonicum. (wh.)	Japan.	Japan.	1823.	—	S. h.
	Volkamèria japonica, Th.					
37	costatum. B.P.	(wh.) ribbed-leaved.	N. S. W.	1823.	5. 8.	G. h.
	VOLKAM'ERIA. S.S.	VOIKAM'ERIA.	Didynamia	Angiospermia. L.		
	aculeata. s.s. (wh.)	prickly.	W. Indies.	1739.	8. 10.	S. h.
	ÆGI'PHILA. R.S.	ÆGI'PHILA.	Tetrandria	Monogynia. L.		
1	martinicensis. r.s.(st.)	Martinico.	W. Indies.	1780.	11. 1.	S. h.
2	arborens. R.S.	(st.) arborescent.	Trinidad.	1820.	—	S. h.
3	lævis. R.S.	(ye.) smooth.	Guiana.	1824.	6. 7.	S. h.
4	elata. R.S.	(ye.) tall.	W. Indies.	1823.	7. 11.	S. h.
5	fœtida. R.S.	(ye.) fetid.	—	1800.	6. 7.	S. h.
6	trifida. R.S.	(wh.) trifid-flowered.	Jamaica.	1826.	—	S. h.
7	diffusa. R.S.	(st.) diffuse.	W. Indies.	1804.	7. 8.	S. h.
8	obovata. R.S.	(st.) oval-leaved.	—	—	—	S. h.
	CALLICA'RPA. R.S.	CALLICA'RPA.	Tetrandria	Monogynia. L.		
1	americana. R.S.	(re.) American.	Carolina.	1724.	6. 7.	G. h.
2	câna. R.S.	(li.) hoary.	E. Indies.	1799.	—	S. h.
3	lanata. R.S.	(pu.) woolly	—	1788.	—	S. h.
4	incâna. F.I.	(ro.) white-leaved.	—	1816.	6. 10.	S. h.
5	macrophylla. R.S.	(ro.) large-leaved.	—	1808.	—	S. h.
6	lanceolaria. F.I.	(pu.) lance-leaved.	—	1816.	—	S. h.
7	ferruginea. R.S.	(li.) rusty.	Jamaica.	1794.	6. 7.	S. h.
8	reticulata. R.S.	(wh.) netted-leaved.	—	1822.	—	S. h.
9	longifolia. R.S.	(wh.) long-leaved.	China.	—	—	S. h.
10	rubella. B.R.	(re.) red-flowered.	—	—	—	S. h.
11	purpurea. F.I.	(pu.) purple-flowered.	—	—	7. 9.	S. h.
12	arborea. F.I.	(p.pu.) tree.	Nepaul.	—	—	S. h.
	VITEX. S.S.	CHASTE-TREE.	Didynamia	Angiospermia. L.		
1	ovata. s.s.	(bl.) oval-leaved.	China.	1796.	7. 8.	S. h.
2	triflora. s.s.	(vi.) three-flowered.	Cayenne.	1823.	—	S. h.
3	latifolia. s.s.	(bl.) broad-leaved.	E. Indies.	1820.	—	S. h.
4	altissima. s.s.	(vi.) tall.	Ceylon.	1802.	—	S. h.
5	alata. s.s.	(vi.) winged.	E. Indies.	1818.	6. 8.	S. h.
6	gigantea. K.S.	(vi.) giant.	Guayaquil.	1826.	—	S. h.
7	bicolor. s.s.	(bl.wh.) two-coloured.	E. Indies.	1810.	6. 8.	S. h.
8	trifolia. s.s.	(vi.) three-leaved.	—	1759.	—	S. h.
9	Negundo. s.s.	(bl.) Ash-leaved.	—	—	—	S. h.
10	umbrosa. s.s.	(vi.) bushy.	W. Indies.	1824.	—	S. h.
11	Agnus-castus. w.	(wh.) common.	Sicily.	1570.	9.	H. h.
	β latifolia.	(wh.) broad-leaved.	—	—	—	H. h.
12	incisa. s.s.	(bl.) cut-leaved.	China.	1758.	7. 9.	G. h.
	Negundo. B.M.					
13	arborea. H.B.	(bl.) tree.	E. Indies.	1820.	—	S. h.
14	bignonioïdes. K.S.	(bl.) trumpet-flowered.	Caracas.	1826.	—	S. h.
15	saligna. H.B.	(bl.) Willow-leaved.	E. Indies.	1823.	—	S. h.
16	Domiana.	(vi.) Don's.	Sierra Leon.	1824.	—	S. h.
	umbrosa. H.T. non Sw.					
17	heterophylla. H.B.	(bl.) various-leaved.	E. Indies.	1820.	—	S. h.
18	capitata. P.S.	(bl.) headed-flowered.	Trinidad.	1822.	6. 8.	S. h.

Banks ic. Kämp. t. 22.
Osb. it. t. 11.Jacq. ic. 3. t. 500.
Bot. mag. t. 1518.Burm. ind. t. 45. f. 1.
Venten. malm. t. 70.
Bot. mag. t. 1834.
— t. 1805.

Bot. reg. t. 406.

Brown jam. t. 20. f. 2.

Lodd. bot. cab. t. 132.
Aub. gui. 1. t. 24.
— 1. t. 25.
Bot. reg. t. 946.Andr. deposit. t. 578. f. 1.
— t. 578. f. 2.

Catesb. car. 2. t. 47.

Gært. fruct. 1. t. 94.

Hook. ex. flor. t. 133.
Bot. reg. t. 883.

Rheed. mal. 5. t. 2.

Bot. mag. t. 2187.
Rheed. mal. 2. t. 11.

Bot. mag. t. 364.

Vahl. ecl. 2. t. 18.

WALLROTHIA. S.S. **WALLROTHIA.** Didynamia Angiospermia. S.S.
Leucóxylon. s.s. (*pu.*) white-wooded. Ceylon. 1793. S. h.
Vitex Leucóxylon. L.

CHLOA'NTHES. B.P. **CHLOA'NTHES.** Didynamia Angiospermia. S.S.
1 stœchadis. B.P. (*st.*) white-leaved. N. S. W. 1822. 6. 8. G. h.
2 glandulosa. B.P. (*st.*) glandular-leaved. — 1824. — G. h.
PRE'MNA. S.S. **PRE'MNA.** Didynamia Angiospermia. S.S.

1 integrifolia. s.s. (*wh.*) entire-leaved. E. Indies. 1822. S. h.
2 tomentosa. s.s. (*wh.*) woolly. — 1825. S. h.
3 reticulata. s.s. (*wh.*) netted-leaved. Jamaica. 1826. S. h.
4 spinosa. H.B. (*wh.*) spiny. — 1822. S. h.
5 esculenta. H.B. (*wh.*) esculent. — 1824. S. h.
6 latifolia. H.B. (*wh.*) broad-leaved. E. Indies. 1827. S. h.
7 serratifolia. s.s. (*wh.*) saw-leaved. — — — S. h.

HOLMSKIOLDIA. HOLMSKIOLDIA. Didynamia Angiospermia. S.S.
1 sanguinea. s.s. (*sc.*) scarlet. E. Indies. 1796. 10. 4. S. h. Rumph. amb. 3. t. 133.

Hastignia coccinea. Sm. ex. bot. t. 80. — 1824. — S. h. Bot. reg. t. 692.

2 scandens. (*pi.*) climbing. — 1824. — S. h. *Hastignia scandens.* H.B.

PETR'EA. S.S. **PETR'EA.** Didynamia Angiospermia. L.

1 volubilis. s.s. (*pu.*) climbing. Vera Cruz. 1733. 7. 8. S. h. Bot. mag. t. 628.
2 rugosa. K.S. (*pu.*) rough-leaved. Caracas. 1824. S. h.
3 erecta. L.C. (*pu.*) upright. S. America. — S. h.

HO'STA. Jacq. HO'STA. Diandria Monogynia.

1 cerulea. Jacq. (*bl.*) blue-flowered. S. America. 1733. 6. 9. S. h. Bot. reg. t. 1204.
Cornutia punctata. B.M. 2611.

2 latifolia. K.S. (*vi.*) broad-leaved. Mexico. 1824. — S. h.

CORNUTIA. P.S. **CORNUTIA.** Diandria Monogynia. S.S.

pyramidalis. P.S. (*bl.*) pyramidal. W. Indies. 1733. S. h.

GMEL'INA. S.S. **GMEL'INA.** Didynamia Angiospermia. L.

1 arborea. s.s. (*br.ye.*) tree. E. Indies. 1812. S. h.

2 villosa. s.s. (*ye.*) villous. — 1818. S. h.

3 asiatica. s.s. (*ye.*) oval-leaved. — 1792. S. h.

4 parviflora. P.S. (*ye.*) small-flowered. — 1817. S. h.

parvifolia. s.s. — — — S. h.

5 sinuata. L.en. (*ye.*) sinuated. — 1824. S. h.

CITHAREXYLUM. W. FIDDLE-WOOD. Didynamia Angiospermia. L.

1 caudatum. s.s. (*wh.*) upright. Jamaica. 1763. S. h.

erectum. Jacq. — — — S. h.

2 villosum. s.s. (*wh.*) hairy-leaved. W. Indies. 1784. S. h.

3 subseratum. s.s. (*wh.*) subserrate. Hispaniola. 1819. S. h.

4 quadrangulare. s.s. (*wh.*) square-stalked. Jamaica. 1759. 6. 8. S. h.

5 cinereum. s.s. (*wh.*) ash-coloured. W. Indies. 1739. — S. h.

6 molle. s.s. (*wh.*) soft-leaved. — 1822. — S. h.

7 pentandrum. s.s. (*pu.*) pentandrous. Portorico. 1815. — S. h.

DURA'NTA. S.S. **DURA'NTA.** Didynamia Angiospermia. L.

1 Plumieri. B.R. (*bl.*) smooth. S. America. 1733. 6. 10. S. h.

2 Ellisia. B.M. (*bl.*) prickly. W. Indies. 1739. 8. S. h.

3 inermis. Mill. (*bl.*) unarmed. S. America. — — S. h.

4 xalapensis. K.S. (*bl.*) Mexican. Mexico. 1822. S. h.

5 Mutisia. K.S. (*bl.*) Mutis's. S. America. — 6. 8. S. h.

AMAS'ONIA. S.S. **AMAS'ONIA.** Didynamia Angiospermia. S.S.

punicea. s.s. (*ye.*) puniceous. Trinidad. 1823. S. h.

LANT'ANA. S.S. **LANT'ANA.** Didynamia Angiospermia. L.

1 aculeata. s.s. (*or.*) changeable-color'd. W. Indies. 1692. 4. 11. S. h.

2 purpurea. s.s. (*pu.*) purple-flowered. S. America. 1822. — S. h.

3 scabrida. s.s. (*or.*) rough. W. Indies. 1774. 7. 10. S. h.

4 mista. w. (*st.li.*) Nettle-leaved. — 1732. 8. 10. S. h.

5 melissæfolia. s.s. (*ye.*) Balm-leaved. — — — 7. 9. S. h.

6 nivea. s.s. (*wh.*) snowy white. E. Indies. 1810. — — S. h.

7 stricta. s.s. (*ro.*) narrow-leaved. Jamaica. 1733. — — S. h.

8 Rádula. w. (*pu.*) Rasp-leaved. W. Indies. 1803. — — S. h.

9 salviaefolia. s.s. (*re.*) Sage-leaved. C. B. S. 1823. — — G. h.

10 récta. s.s. (*bh.*) upright. Jamaica. 1758. 6. 8. S. h.

11 involucrata. s.s. (*ti.*) round-leaved. W. Indies. 1690. 5. 7. S. h.

12 odorata. s.s. (*wh.*) sweet-scented. — 1758. 5. 11. S. h.

13 Cámrara. s.s. (*ye.*) various-coloured. — 1691. 4. 9. S. h.

14 trifolia. s.s. (*ro.*) three-leaved. — 1733. 6. 9. S. h.

15 ánnua. s.s. (*ro.*) annual. S. America. — 7. 8. S. h.

Rumph. amb. 3. t. 133.

Bot. reg. t. 692.

Bot. reg. t. 1204.

Lam. ill. t. 541.

Roxb. corom. 3. t. 246.

Lam. ill. t. 542.

Roxb. corom. 1. t. 32.

Jacq. ic. 3. t. 501.

Jacq. ic. 1. t. 118.

Jacq. vind. 1. t. 22.

Jacq. amer. t. 118.

Vent. cels. t. 47.

Bot. reg. t. 244.

Bot. mag. t. 1759.

Bot. mag. t. 96.

Lodd. bot. cab. t. 68.

Dill. elt. t. 57. f. 66.

Bot. mag. t. 1946.

Sloan. jam. 2. t. 195. f. 4.

Jacq. schœnbl. 3. t. 285.

— 3. t. 360.

Pluk. alm. t. 114. f. 5.

Plum. ic. 71. f. 2.

Dill. elt. t. 56. f. 65.

Bot. mag. t. 1449.

— t. 1022.

VERBENACEÆ.

16	fucata. s.s.	(ro.) painted.	Brazil.	1822.	7. 8.	S. h.	Bot. reg. t. 798.
17	lavandulacea.s.s.(wh.)	Lavender-like.	—	—	S. h.	Jacq. schœn. t. 473.
18	crocæa. L.en.	(or.) saffron-coloured.	Jamaica.	—	—	S. h.	
19	violacea. Desf.	(vi.) violet-coloured:	S. America.	1824.	—	S. h.	
20	brasiliensis. L.en.(wh.)	Brazilian.	Brazil.	—	7. 8.	S. h.	
21	alba. L.en.	(wh.) white-flowered.	S. America.	—	—	S. h.	
22	hirta. Gr.	(wh.) hairy white.	—	1825.	8. 11.	S. h.	
23	horrifica. k.s.	(ye.) horrid.	Mexico.	—	7. 8.	S. h.	
24	Sellœ. L.O.	(ro.) Sello's.	Monte Video.	1829.	12. 4.	S. h.	Bot. mag. t. 2981.
SPIELMANIA. S.S.		SPIELMANIA.	Didynamia Angiospermia. S.S.				Bot. mag. t. 1899.
	africana. s.s.	(wh.) African.	C. B. S.	1710.	2. 11.	G. h.	Roxb. corom. 1. t. 6.
TECTONA. W.		TEAK-WOOD.	Pentandria Monogynia. L.				Roxb. corom. 2. t. 146.
	grandis. w.	(wh.) great.	E. Indies.	1777.	—	S. h.	
STREPTIUM. S.S.		STREPTIUM.	Didynamia Angiospermia. S.S.				
	asperum. s.s.	(wh.) rough.	E. Indies.	1799.	7. 8.	S. y.	
<i>Tortula aspera</i> w. <i>Priva leptostachya</i> . p.s.							
PRIVA. S.S.		PRIVA.	Didynamia Angiospermia. S.S.				
1	lappulacea. p.s.	(pu.) bristly-fruited.	S. America.	1822.	6. 8.	S. y.	Jacq. obs. 1. t. 24.
	echinata. k.s.						
2	mexicana. p.s.	(li.) Mexican.	Mexico.	1726.	8. 9.	S. y.	Dill. elth. t. 302. f. 389.
	hispida. k.s.	<i>Verbena mexicana</i> . L.					
TAMONEA. S.S.		TAMONEA.	Diandria Monogynia.				
1	verbenacea. s.s.	(bl.) thorny-fruited.	W. Indies.	1733.	8.	S. ⊙.	Banks. vel. Hous. t. 2.
	curassavica. p.s.	<i>Ghinia spinosa</i> . w.					
2	spicata. k.s.	(bl.) spiky.	Trinidad.	1824.	8. 10.	S. ⊙.	Aubl. guian. t. 268.
STACHYTRPHETA. BASTARD VERVAIN.		Diandria Monogynia. S.S.					
1	angustifolia. r.s.	(vi.) narrow-leaved.	S. America.	—	6. 9.	S. ⊙.	Jacq. obs. 4. t. 86.
2	elatior. r.s.	(vi.) tall.	Brazil.	1821.	8. 11.	S. ⊙.	
3	indica. r.s.	(vi.) Indian.	Ceylon.	1732.	8. 9.	S. ⊙.	
4	urticifolia. B.M.	(bl.) Nettle-leaved.	S. America.	—	5. 9.	S. ♂.	Bot. mag. t. 1848.
5	jamaicensis. r.s.	(bl.) Jamaica.	W. Indies.	1714.	6. 9.	S. ♂.	— t. 1860.
6	hirta. k.s.	(vi.) hairy-leaved.	New Granada.	1821.	5. 10.	S. h.	
7	cayennensis. s.s.	(bl.) Cayenne.	Cayenne.	1819.	—	S. h.	
8	umbrosa. k.s.	(bl.) shade.	Cumana.	1829.	—	S. h.	
9	prismatica. r.s.	(bl.) Germanander-leav'd.	W. Indies.	1699.	5. 6.	S. ♂.	Jacq. ic. 2. t. 208.
10	orubica. r.s.	(vi.) Orubian.	S. America.	—	6. 8.	G. ♂.	Ehr. pict. t. 5. f. 1.
11	mutabilis. r.s.	(ro.) changeable-flower'd.	—	1801.	3. 9.	S. h.	Bot. mag. t. 976.
	<i>Verbena mutabilis</i> . w. <i>Cymburus mutabilis</i> .	Par. lond. t. 49.					
12	crassifolia. r.s.	(vi.) thick-leaved.	Brazil.	1826.	6. 9.	S. h.	Brown jam. t. 3. f. 1.
13	Zuccagni. r.s.(ro.vi.)	Zuccagni's.	—	1824.	—	S. h.	Flor. græc. t. 553.
14	hirsutissima. L.en.(bl.)	hairy.	Brazil.	1821.	—	S. y.	
ZAPANIA. J.		ZAPANIA.	Didynamia Angiospermia. P.S.				
1	stoechadifolia. p.s.(wh.)	oval-spiked.	W. Indies.	1732.	8. 9.	S. y.	
2	nodiflora. B.P.	(pi.) knot-flowered.	America.	1664.	7. 8.	G. y.	Bot. mag. t. 367.
	<i>Verbena nodiflora</i> . L. <i>Lippia nodiflora</i> . s.s.						
ALOYSIA. P.S.		ALOYSIA.	Didynamia Angiospermia. P.S.				
	citrodora. p.s.*	(li.) Lemon-scented.	Chile.	1784.	5. 9.	G. h.	
	<i>Veronica triphylla</i> . B.M. <i>Lippia citrodora</i> . k.s.						
VERBENA. S.S.		VERVAIN.	Didynamia Angiospermia. S.S.				
1	angustifolia. s.s.	(bl.) narrow-leaved.	N. America.	1802.	6. 8.	H. y.	
	rugosa. w.						
2	paniculata. s.s.	(pu.) panicled.	—	1800.	7. 8.	H. y.	Bot. reg. t. 1102.
3	bonariensis. s.s.	(bl.) cluster-flower'd.	Buenos Ayres.	1732.	7. 10.	H. ♂.	Dill. elt. t. 300. f. 387.
4	caroliniana. s.s.	(pi.) Carolina.	N. America.	—	6. 9.	H. y.	— t. 301. f. 388.
5	urticafolia. s.s.	(wh.) Nettle-leaved.	—	1683.	7. 9.	H. y.	Rob. ic. 26.
6	stricta. s.s.	(bl.) upright.	—	1802.	7. 8.	H. y.	Bot. mag. t. 1976.
7	lasiostachys. s.s.	(bl.) woolly-spiked.	California.	1823.	—	H. y.	
8	bracteosa. s.s.	(bl.) long-bracted.	N. America.	1812.	7.	H. y.	Bot. mag. t. 2910.
9	Aubletia. s.s.	(ro.) rose.	—	1774.	6. 8.	F. ♂.	— t. 308.
10	Lamberti. B.M.	(ro.) Lambert's.	Peru.	1816.	—	F. y.	— t. 2200.
11	chamædryfolia. p.s.(sc.)	scarlet-flower'd.	Buenos Ayres.	1827.	4. 10.	F. y.	Swt. br. fl. gar. ser. 2. t. 9.
	<i>Melindres</i> . Bót. reg. t. 1184. <i>Erinus peruvianus</i> . L.						
12	barbata. Gr.	(li.) bearded.	Mexico.	—	9. 12.	G. h.	Herm. parad. t. 242.
13	hastata. s.s.	(pu.) halberd-leaved.	Canada.	1710.	—	H. y.	
14	scabra. s.s.	(vi.) rough.	Mexico.	1822.	—	H. y.	
15	alata. B.F.G.	(pu.) winged-stalked.	Monte Video.	1827.	7. 10.	F. y.	Swt. br. fl. gar. ser. 2. t. 41.
16	officinalis. E.B.	(li.) common.	Britain.	—	6. 9.	H. y.	Eng. bot. v. 11. t. 769.
17	sororia. D.P.	(li.) sister.	Nepaul.	1823.	8. 10.	H. y.	Swt. br. fl. gar. t. 202.
18	spuria. s.s.	(li.) jagged-leaved.	N. America.	1731.	7. 8.	H. ♂.	

19 erinoïdes. s.s. <i>multifida</i> . R.P.	(ro.) Erinus-like.	Peru.	1820.	7. 8. H. ⊙.	Flor. peruv. I. t. 33. f.c.
20 pulchella. B.F.G.	(li.) pretty.	Buenos Ayres.	1827.	4. 10. F. 4.	Swt. br. fl. gar. s. 2. t. 295.
21 canescens. s.s.	(vi.) canescent.	Mexico.	1820.	7. 8. H. 4.	H. et B. 2. t. 136.
22 supina. s.s.	(li.) trailing.	Spain.	1640.	6. 7. H. ⊙.	Flor. græc. t. 554.
β <i>hirsuta</i> . E.	(li.) hairy.	Egypt.	1829.	— H. ⊙.	
23 prostrata. H.K.	prostrate.	N. America.	1794.	— H. 4.	
LI'PIA. J.	LI'PIA.	Didynamia Angiospermia. S.S.			
purpurea. s.s.	(pu.) purple.	Mexico.	1823.	6. 8. G. h.	Jacq. f. eclog. t. 85.
XENOP'OMA. S.S.	XENOP'OMA.	Didynamia Angiospermia.			
obovatum. s.s.	(wh.) obovate-leaved.	China.	1816.	5. 8. G. h.	Coll. hort. rip. t. 25.

ORDO CLVI.

MYOPORINÆ. Brown prodr. p. 514.

MYOP'ORUM. B.P.	MYOP'ORUM.	Didynamia Angiospermia. S.S.			
1 ellípticum. B.P.	(wh.) smooth-leaved.	N. S. W.	1789.	1. 3. G. h.	Andr. repôsit. t. 283.
Pogònia glabra. A.R.					
2 acuminatum. B.P.	(wh.) acuminate.	—	1812.	— G. h.	
3 montanum. B.P.	(wh.) mountain.	—	1822.	2. 6. G. h.	
4 parvifolium. B.P.	(wh.) small-leaved.	N. Holland.	1803.	3. 9. G. h.	
5 débile. B.P.	(bl.) procumbent.	N. S. W.	1793.	5. 8. G. h.	
6 diffusum. B.P.	(bl.) diffuse.	N. Holland.	—	8. 2. G. h.	
7 tuberculatum. B.P.	(wh.) tubercled.	—	1803.	5. 8. G. h.	
8 viscósum. B.P.	(wh.) viscid.	—	—	— G. h.	
9 crassifolium. s.s.	(wh.) thick-leaved.	N. Zealand.	1822.	— G. h.	
10 oppositifolium. s.s.	(wh.) opposite-leaved.	N. Holland.	1803.	1. 12. G. h.	
STENOCH'ILUS. B.P.	STENOCH'ILUS.	Didynamia Angiospermia. S.S.			
1 glaber. B.P.	(sc.) smooth-leaved.	N. Holland.	1803.	1. 12. G. h.	
2 maculatus. B.R.	(cr. sp.) spotted-flowered.	—	1820.	4. 8. G. h.	Bot. mag. t. 1691.
3 viscósum. B.M.	(ye.) clammy.	—	1825.	— G. h.	t. 1830.
BO'NTIA. S.S.	BO'NTIA.	Didynamia Angiospermia. L.			
daphnoïdes. s.s.	(st.) Barbadoes.	W. Indies.	1690.	6. S. h.	
AVICE'NNIA. B.P.	AVICE'NNIA.	Didynamia Angiospermia. L.			Dill. elth. t. 49. f. 57.
tomentosa. B.P.	(wh.) downy-leaved.	India.	1793.	— S. h.	Palisot. d'Owar. t. 47.

ORDO CLVII.

ACANTHACEÆ. Brown prodr. I. p. 472.

ACA'NTHUS. S.S.	BEARS-BREECH.	Didynamia Angiospermia. L.			
1 mollis. w.	(wh.) soft-leaved.	Italy.	1548.	7. 9. H. 4.	Lam. ill. t. 550.
2 niger. L.en.	(wh.) shining-leaved.	Portugal.	1759.	— H. 4.	
3 spinosus. B.M.	(wh.) prickly-leaved.	S. Europe.	1629.	— H. 4.	Bot. mag. t. 1808.
4 spinosissimus. P.S.	(wh.) white-spined.	—	—	— H. 4.	
5 carduifolius. s.s.	(wh.) Cape.	C. B. S.	1816.	— G. 4.	Rheed. mal. 2. t. 48.
6 ilicifolius. s.s.	(wh.) Holly-leaved.	E. Indies.	1759.	— S. h.	
1 capensis. s.s.	(wh.) simple-spined.	C. B. S.	1816.	5. 9. G. h.	
2 furcata. s.s.	(wh.) fork-spined.	—	—	6. 8. G. h.	
3 procumbens. s.s.	(wh.) procumbent.	—	1825.	— G. h.	
BLEPH'ARIS. J.	BLEPH'ARIS.	Didynamia Angiospermia. P.S.			
1 capensis. s.s.	(wh.) simple-spined.	C. B. S.	1816.	5. 9. G. h.	
2 furcata. s.s.	(wh.) fork-spined.	—	—	6. 8. G. h.	
3 procumbens. s.s.	(wh.) procumbent.	—	1825.	— G. h.	
THUNBER'GIA. S.S.	THUNBER'GIA.	Didynamia Angiospermia. S.S.			
1 grandiflora. s.s.	(bl.) large-flowered.	E. Indies.	1822.	3. 8. S. h. —	Bot. reg. t. 495.
2 frágans. s.s.	(wh.) white Indian.	—	1796.	5. 9. S. h. —	Bot. mag. t. 1881.
3 cordata. C.R.	(wh.) heart-leaved.	Brazil.	1823.	3. 8. S. h. —	Coll. h. rip. t. 21.
4 capensis. s.s.	(wh.) Cape.	C. B. S.	1816.	— G. h. —	
5 alata. B.M.	(or.) winged.	Zanzebar.	1825.	1. 12. S. h. —	Bot. mag. t. 2591.
6 angulata. H.E.F.	(pu.) angular-stalked.	Mauritius.	1824.	5. 8. S. h. —	Hook. ex. flor. t. 166.
7 coccinea. D.P.	(sc.) scarlet-flowered.	Nepaul.	—	6. 2. S. h. —	t. 195.

BAR'L'ERIA. S.S.

	BAR'L'ERIA.	Didynamia Angiospermia. L.	
1 buxitòlia. s.s.	(bl.) Box-leaved.	E. Indies.	1763. 6. 7. S. h.
2 Prionitis. s.s.	(ye.) thorny.	—	1759. 7. 8. S. h.
3 spicata. H.B.	(pu.) spiked-flowered.	—	1822. 4. 8. S. h.
4 longifolia. s.s.	(vi.) long-leaved.	—	1781. 7. 9. S. d.
5 cristata. s.s.	(bl.) crested.	—	1796. 4. 9. S. h.
6 alba. B.C.	(wh.) white-flowered.	—	1818. — S. h.
7 ciliata. H.B.	(pu.) fringed.	—	1822. 4. 10. S. h.
8 tomentosa. s.s.	(bl.) woolly.	—	1825. — S. h.
9 flava. s.s.	(ye.) yellow-flowered.	Arabia.	1816. — S. h.
mitis. B.R. 191.	Justicia flava. w. Eranthemum flavum. W.en.		

10 longiflora. s.s.	(vi.) long-flowered.	E. Indies.	1816. 6. 9. S. h.
11 purpurea. B.C.	(pu.) purple.	—	1817. — S. h.
12 dichotoma. H.B.	(pu.) forked.	—	1822. — S. h.
13 cœrulea. H.B.	(bl.) blue-flowered.	—	— S. h.

HYGRO'PHILA. B.P.	HYGRO'PHILA.	Didynamia Angiospermia. S.S.	
rín:ens. s.s.	(li.) gaping-flowered.	E. Indies.	1807. 7. 8. S. h.

GEISSOM'ERIA. B.R.	GEISSOM'ERIA.	Didynamia Angiospermia.	
1 longiflora. B.R.	(sc.) long-flowered.	Brazil.	1825. 8. 10. S. h.
2 fúlgida.	(sc.) bright-flowered.	W. Indies.	1804. 7. 8. S. h.

Ruélia fúlgida. A.R.

RUE'LLIA. S.S.	RUE'LLIA.	Didynamia Angiospermia. L.	
1 ocymoides. s.s.	(hl.) Basil-like.	Mexico.	1815. 7. 8. S. h.
2 ciliata. s.s.	(li.) fringed.	E. Indies.	1806. — S. h.
3 australis. s.s.	(re.) New Holland.	N. S. W.	1820. 6. 9. G. h.
4 foetida. s.s.	(vi.) strong-scented.	Mexico.	1824. — G. h.
5 biflora. s.s.	(bl.) two-flowered.	Carolina.	1765. 7. G. 4.
6 violacea. s.s.	(bl.) violet-flowered.	Guiana.	1822. 6. 9. S. 4.
7 salicifolia. s.s.	(vi.) Willow-leaved.	E. Indies.	— — S. h.
8 ovata. s.s.	(bl.) oval-leaved.	Mexico.	1800. 7. 8. S. 4.
9 pátula. s.s.	(vi.) spreading.	E. Indies.	1774. — S. h.
10 Sabiniâna. B.R.	(vi.) Sabine's.	—	1827. 4. 8. S. h.
11 brasila. Mart.	(vi.) Brazil.	Brazil.	1828. 8. 9. S. h.
12 anisophylla. H.E.F. (bl.)	unequal-leaved.	E. Indies.	1823. 9. 4. S. h.
persicifolia. Bot. reg. t. 955.			

13 formosa. s.s.	(sc.) splendid.	Brazil.	1808. 6. 9. S. h.
14 macrophylla. s.s.	(re.) large-leaved.	S. America.	1820. — S. h.
15 tuberosa. s.s.	(vi.) tuberous-root'd.	Jamaica.	1752. 7. 8. S. 4.
16 strépens. s.s.	(pa.bl.) whorl-flowered.	N. America.	1726. — G. 4.
17 láctea. s.s.	(li.) white.	Mexico.	1796. 6. 8. G. 4.
18 clandestina. s.s.	(bl.) three-flowered.	Barbadoes.	1728. 7. 8. S. 4.
19 paniculata. s.s.	(li.) paniced.	W. Indies.	1768. 8. 10. S. 4.
20 obovata. H.B.	(pu.) obovate-leaved.	E. Indies.	1820. 7. 11. S. ⊙.
21 undulata. s.s.	(vi.) wave-leaved.	—	1817. — S. 4.
22 hirta. s.s.	(vi.) hairy.	—	— S. 4.
23 pubescens. P.S.	(bl.) pubescent.	C. B. S.	1816. 5. 7. G. h.
24 tetragona. s.s.	(bl.) four-sided.	Brazil.	1818. 6. 9. S. 4.
25 longifolia. s.s.	(pu.) long-leaved.	E. Indies.	1821. — S. h.
26 cérnua. H.B.	(vi.) cernuous.	—	— S. h.
27 semitetrandra. Jac.	... semitetrandrous.	—	1829. ... S. h.

BL'ECHUM. B.P.	BL'ECHUM.	Didynamia Angiospermia. S.S.	
1 Brównei. s.s.	(li.) dense-spiked.	W. Indies.	1780. 6. S. ⊙.
2 laxiflorum. s.s.	(wh.) loose-flowered.	—	1822. 6. 8. S. 4.
3 angustifolium. s.s.	(bl.) narrow-leaved.	—	1826. 7. 10. S. 4.

APHELA'NDRA. B.P.	APHELA'NDRA.	Didynamia Angiospermia. S.S.	
cristata. s.s.	(sc.) dense-spiked.	W. Indies.	1733. 6. 9. S. h.

Ruélia cristata. A.R. 506.	Justicia pulcherrima. L.		
----------------------------	--------------------------	--	--

CROSSA'NDRA. P.S.	CROSSA'NDRA.	Didynamia Angiospermia.	
undulatifolia. P.L.	(sc.) wave-leaved.	E. Indies.	1800. 6. 1. S. h.

Ruélia infundibuliformis. A.R. 542.	Harrachia speciosa. Jacq. ecl. t. 21.		
-------------------------------------	---------------------------------------	--	--

PHAYLO'PSIS. W.	PHAYLO'PSIS.	Didynamia Angiospermia. W.	
1 longifolia. B.M.	(wh.) long-leaved.	Sierra Leon.	1822. 3. 5. S. h.

2 imbricata.	(wh.) imbricated.	Mauritius.	— — S. h.
--------------	-------------------	------------	-----------

Aethilema imbricata. s.s.			
---------------------------	--	--	--

3 parviflora. w.	(wh.) small-flowered.	E. Indies.	1825. 4. 8. S. 4.
------------------	-----------------------	------------	-------------------

LEPIDAG'ATHIS. W.	LEPIDAG'ATHIS.	Didynamia Angiospermia. W.	
-------------------	----------------	----------------------------	--

cristata. w.	(pu.) crested.	E. Indies.	1817. 5. 8. S. 4.
--------------	----------------	------------	-------------------

ELYTRARIA. R.S.	ELYTRARIA.	Diandria Monogynia. S.S.	
-----------------	------------	--------------------------	--

1 crenata. r.s.	(wh.) notched-leav'd.	E. Indies.	1819. 11. 2. S. 4.
-----------------	-----------------------	------------	--------------------

Justicia acaulis. F.I.			
------------------------	--	--	--

Rheed. mal. 2. t. 47.	
— 9. t. 41.	

Pluk alm. t. 133. f. 4.	
Bot. mag. t. 1615.	

Lodd. bot. cab. t. 360.	
-------------------------	--

Jacq. f. ecl. t. 46.	
----------------------	--

Vahl symb. 1. t. 16.	
----------------------	--

Lodd. bot. cab. t. 344.	
-------------------------	--

Bot. reg. t. 1045.	
--------------------	--

Andr. deposit. t. 527.	
------------------------	--

Cavan. ic. 5. t. 456.	
-----------------------	--

Cavan. ic. 6. t. 586. f. 1.	
-----------------------------	--

Aub. gui. 2. t. 271.	
----------------------	--

Cavan. ic. 3. t. 254.	
-----------------------	--

Jacq. ic. 1. t. 119.	
----------------------	--

Bot. reg. t. 1238.	
--------------------	--

Hook. ex. flor. 191.	
----------------------	--

Bot. mag. t. 1400.	
--------------------	--

Vahl symb. 2. t. 39.	
----------------------	--

Sloan. jam. 1. t. 95. f. 1.	
-----------------------------	--

Schlk. hand. 2. t. 177.	
-------------------------	--

Cavan. ic. 3. t. 255.	
-----------------------	--

Dill. elth. 2. t. 248. f. 320.	
--------------------------------	--

Bot. reg. t. 585.	
-------------------	--

Vahl symb. 3. t. 67.	
----------------------	--

Sloan. jam. 1. t. 109. f. 1.	
------------------------------	--

Bot. reg. t. 69.	
------------------	--

Bot. mag. t. 2433.	
--------------------	--

Roxb. corom. 2. t. 127.	
-------------------------	--

2 <i>lyrata</i> . R.S.	(<i>wh.</i>) lyrate-leaved.	E. Indies.	1825.	11. 2. S. 4.		
3 <i>virgata</i> . R.S.	(<i>wh.</i>) twiggy.	Carolina.	1813.	7. 8. H. 4.	Mich. amer. 1. t. 1.	
4 <i>ramosa</i> . K.S.	(<i>bl.</i>) branching.	Mexico.	1825.	— G. 4.		
5 <i>caulescens</i> . R.S. (<i>wh.</i>) caulescent.		Manilla.	1828.	— G. 4.		
JUSTICIA. R.S.	JUSTICIA.	<i>Diandria Monogynia</i> , L.				
1 <i>bicalyculata</i> . R.S. (<i>li.</i>) Malabar.	E. Indies.	1785.	8. 10. S. 4.	Cavan. ic. 1. t. 71.		
2 <i>hispidia</i> . R.S. (<i>wh.</i>) hispid.	Sierra Leon.	1824.	4. 8. S. 4.			
3 <i>Ecbolium</i> . R.S. (<i>pa.bl.</i>) long-spiked.	E. Indies.	1759.	3. 8. S. 4.	Bot. mag. t. 1847.		
4 <i>ventricosa</i> . B.M. (<i>sp.</i>) hop-flowered.	China.	1825.	4. 6. S. 4.	— t. 2766.		
5 <i>calycotricha</i> . s.s. (<i>ye.</i>) yellow-flower'd. Brazil. <i>flavícoma</i> . Bot. reg. t. 1027.		—	1. 3. S. 4.	— t. 2816.		
6 <i>thyrsiflora</i> . F.I. (<i>or.</i>) thyrs-flower'd. E. Indies.		1812.	5. 9. S. 4.			
7 <i>vittellina</i> . F.I. (<i>pi.ye.</i>) reflex-leaved.		1818.	— S. 4.			
8 <i>álba</i> . F.I. (<i>wh.</i>) white-flowered.		1816.	4. 8. S. 4.			
9 <i>coccinea</i> . R.S. (<i>sc.</i>) scarlet.	S. America.	1770.	12. 4. S. 4.	Bot. mag. t. 432.		
10 <i>polysperma</i> . F.I. (<i>bl.</i>) many-seeded.	E. Indies.	1821.	8. 10. S. 4.	Rheed. mal. 9. t. 46.		
11 <i>echoïdes</i> . R.S. (<i>wh.pu.</i>) <i>Echium</i> -like.		1812.	6. 9. S. 4.	Sal. par. lond. t. 50.		
12 <i>quadripida</i> . R.S. (<i>sc.</i>) twiggy. <i>virgularis</i> . P.L.	Mexico.	1795.	3. 9. S. 4.			
13 <i>pumila</i> . R.S. (<i>sc.</i>) dwarf.	S. America.	1820.	— S. 4.			
14 <i>nigricans</i> . R.S. (<i>wh.re.</i>) black-striped.	China.	1819.	6. 9. G. 4.	Andr. deposit. t. 570.		
15 <i>nítida</i> . R.S. (<i>wh.sp.</i>) glossy.	W. Indies.	1790.	3. 9. S. 4.	Jacq. ic. t. 205.		
16 <i>bracteolata</i> . R.S. (<i>pu.</i>) bracteolate.	Caracas.	1824.	6. 8. S. 4.	Rheed. mal. 6. t. 60.		
17 <i>picta</i> . R.S. (<i>cr.</i>) painted. <i>a álba</i> . (<i>cr.</i>) white-marked.	E. Indies.	1780.	7. 8. S. 4.	Bot. reg. t. 1227.		
18 <i>paniculata</i> . R.S. (<i>ro.</i>) paniced.		—	— S. 4.	Bot. mag. t. 1870.		
19 <i>salicina</i> . R.S. (<i>re.</i>) Willow-like.	Peru.	1819.	— S. 4.	Jacq. vinct. 2. t. 34.		
20 <i>secunda</i> . R.S. (<i>pu.</i>) side-flowering.	W. Indies.	1793.	5. 9. S. 4.	Bot. mag. t. 2060.		
21 <i>geniculata</i> . B.M. (<i>sc.</i>) jointed-stalked.		1822.	7. 10. S. 4.	— t. 2487.		
22 <i>nodosa</i> . B.M. (<i>cr.pu.</i>) swoln-jointed.	Brazil.	1826.	8. 10. S. 4.	— t. 2914.		
23 <i>oblongata</i> . L.O. (<i>pi.</i>) oblong-leaved.	Java.	—	9. 10. S. 4.			
24 <i>ciliáris</i> . R.S. (<i>wh.</i>) ciliated.	Cumana.	1780.	7. 8. S. 4.	Jacq. vind. 2. t. 104.		
25 <i>lúcida</i> . R.S. (<i>sc.</i>) shining-leaved.	W. Indies.	1795.	— S. 4.	Bot. mag. t. 1014.		
26 <i>salviæflora</i> . R.S. (<i>sc.</i>) Sage-flowered.	Mexico.	1823.	— S. 4.			
27 <i>Gandarússa</i> . R.S. (<i>vi.</i>) Willow-leaved.	E. Indies.	1800.	6. 7. S. 4.	Bot. reg. t. 635.		
28 <i>carthaginensis</i> . R.S. (<i>sc.</i>) Carthagorean.	Carthagena.	1792.	— S. 4.	— t. 797.		
29 <i>pedunculosa</i> . R.S. (<i>pu.</i>) long-peduncled. <i>americana</i> . w.	N. America.	1759.	7. 8. H. 4.	Bot. mag. t. 2367.		
30 <i>quadrangularis</i> . B.M. (<i>pu.</i>) square-stalked.	Mauritius.	—	1. 3. S. 4.	Bot. mag. t. 2845.		
31 <i>procumbens</i> . R.S. (<i>ro.</i>) procumbent.	E. Indies.	1798.	7. 8. S. 4.	Pluk. alm. t. 56. f. 3.		
32 <i>quinquangularis</i> . F.I. (<i>ro.</i>) five-angled.		1820.	— S. 4.			
33 <i>comáta</i> . R.S. (<i>li.</i>) Balsam-herb.	Jamaica.	1795.	— S. 4.	Sloan. jam. 1. t. 103. f. 2.		
34 <i>eustachiána</i> . R.S. (<i>pu.</i>) Eustachian.	St. Eustachia.	1799.	8. 9. S. 4.	Jacq. amer. pict. t. 5.		
35 <i>Keriána</i> . (<i>sc.or.</i>) Ker's.	China.	1815.	6. 9. S. 4.	Bot. reg. t. 309.		
<i>eustachiána</i> . B.R. Bot. mag. t. 2076. non Jacq.						
36 <i>nasuta</i> . R.S. (<i>wh.</i>) white-flowering.	E. Indies.	1790.	2. 10. S. 4.	Bot. mag. t. 325.		
37 <i>lanceolária</i> . F.I. (<i>ro.</i>) lance-leaved.		1818.	8. 12. S. 4.			
38 <i>speciosa</i> . F.I. (<i>pu.</i>) beautiful.		1823.	1. 12. S. 4.	Bot. mag. t. 2722.		
39 <i>Roxburghiana</i> . R.S. (<i>pi.</i>) Roxburgh's. <i>tinctória</i> . F.I. non Lour. <i>báplica</i> . s.s.		1815.	6. 9. S. 4.	Rumph. amb. 6.t.22.f.1.		
40 <i>pectoralis</i> . R.S. (<i>pu.</i>) Garden Balsam.	W. Indies.	1787.	5. 6. S. 4.	Bot. reg. t. 796.		
41 <i>periplocæfolia</i> . R.S. (<i>pu.</i>) Periploca-leav'd.	S. America.	1799.	6. 8. S. 4.	Jacq. coll. s. t. 7. f. 2.		
42 <i>furcata</i> . R.S. (<i>vi.</i>) forked. <i>peruviana</i> . B.M. Cavan. ic. 1. t. 28.	Peru.	1795.	4. 8. S. 4.	Bot. mag. t. 430.		
43 <i>lithospermifolia</i> . (<i>vi.</i>) Gromwell-leav'd.		1796.	— S. 4.	Jacq. schoen. 1. t. 4.		
44 <i>caracassana</i> . R.S. (<i>vi.</i>) Caracas.	Caracas.	1818.	— S. 4.	Jacq. ic. 2. t. 206.		
45 <i>leucantha</i> . R.S. (<i>wh.</i>) dirty white.	Brazil.	1824.	— S. 4.			
46 <i>Adhátoda</i> . R.S. (<i>wh.pu.</i>) Malabar-nut.	Ceylon.	1699.	5. 7. G. 4.	Bot. mag. t. 861.		
47 <i>decussata</i> . F.I. (<i>li.</i>) cross-leaved.	E. Indies.	1824.	2. 4. S. 4.			
48 <i>Betónica</i> . R.S. (<i>wh.</i>) Betony-leaved.		1739.	— S. 4.	Rheed. mal. 2. t. 21.		
49 <i>ramosissima</i> . F.I. (<i>wh.</i>) branching.		1812.	8. 12. S. 4.			
50 <i>plumbaginifolia</i> . R.S. (<i>re.</i>) Plumbago-leav'd.		1819.	6. 9. S. 4.	Jacq. f. ecl. t. 12.		
51 <i>formosa</i> . R.S. (<i>pu.</i>) handsome.		—	— S. 4.			
52 <i>maculata</i> . B.C. (<i>pu.</i>) spotted-leaved.	W. Indies.	1823.	— S. 4.	Lodd. bot. cab. t. 626.		
53 <i>divaricata</i> . R.S. (<i>wh.</i>) divaricate.	C. B. S.	1822.	3. 6. G. 4.			
54 <i>pátula</i> . R.S. (<i>wh.</i>) spreading-branch'd.		1824.	— G. 4.			
55 <i>hyssopifolia</i> . R.S. (<i>st.</i>) Hyssop-leaved.	Canaries.	1690.	3. 8. G. 4.	Mill. ic. 9. t. 13.		
56 <i>orchioïdes</i> . R.S. (<i>bh.</i>) Broom-leaved.	C. B. S.	1774.	8. G. 4.	Vent. malm. t. 51.		

ACANTHACEÆ.

57	tranquebariænsis. R.S. (bh.)	Tranquebar. E. Indies.	1818.	4. 8. S. ¼.	
58	glâbra. F.I. (ro.ye.)	smooth.	1824.	6. 9. S. ¼.	
59	vestita. R.S. (vi.)	clothed.	1827.	S. ¼.	
DICLIPTERA. J.	DICLIPTERA.	Diandria Monogynia. R.S.			
1	spinôsa. B.C. (ye.)	spiny.	Madagascar.	4. 6. S. ½.	Lodd. bot. cab. t. 1244.
2	chinensis. R.S. (ro.)	Chinese.	China.	8. 12. S. ½.	Burm. ind. t. 4. f. 1. ?
3	bivalvis. R.S. (ro.)	two-valved.	E. Indies.	1820. S. ½.	
4	multiflôra. R.S. (vi.)	many-flowered.	Quito.	1821. 6. 9. S. ½.	Flor. peruv. 1. t. 14.f.b.
5	retusa. R.S. (pu.)	retuse.	Santa Cruz.	1816. 7. 9. S. ¼.	Lodd. bot. cab. t. 724.
6	congesta. K.S. (pu.)	crowded-flowered.	Mexico.	1825. S. ¼.	
7	resupinata. R.S. (vi.)	resupinate.	N. Spain.	1805. 5. 8. S. ¼.	Cavan. ic. 3. t. 203.
	<i>Justicia sexangularis.</i> Cav. non L.				
8	pectinata. R.S. (bl.)	pectinated.	E. Indies.	1801. 7. 9. S. ¼.	Roxb. corom. 2. t. 153.
	<i>Justicia parviflora.</i> w.				
9	rêpens. R.S. (li.)	creeping.	1822.	— S. ¼.	Roxb. corom. 2. t. 152.
10	brasiliensis. R.S. (pu.)	Brazilian.	Brazil.	1824. — S. ¼.	
11	scorpioides. R.S. (pu.)	Scorpion-grass-ld.	Vera Cruz. — S. ½.	Reliqu. Houst. t. 1.
12	sexangularis. R.S. (re.)	chickweed-ld.	S. America.	1733. 7. S. ⊙.	Pluk. alm. t. 270. f. 6.
13	assurgens. R.S. (sc.)	assurgent.	W. Indies.	1827. — S. ¼.	
NELSONIA. B.P.	NELSONIA.	Diandria Monogynia.			
lamiafolia. F.I. (pu.)	Lamium-leaved.	E. Indies.	1824.	6. 8. S. ¼.	
HYPOESTES. R.S.	HYPÖESTES.	Diandria Monogynia. R.S.			
1	purpurea. R.S. (pu.)	purple-flower'd.	China.	1822. 6. 8. S. ¼.	
	<i>Justicia purpurea.</i> L.				
2	verticillaris. R.S. (bh.)	whorl-flowered.	C. B. S.	1823. 4. 6. G. ¼.	
3	sérpens. R.S. (li.)	creeping-stalked.	Mauritius.	1823. 6. 8. S. ¼.	
4	involucrata. R.S. (wh.)	involucred.	E. Indies.	1811. 7. 8. S. ½.	
ERA'NTHEMUM. R.S.	ERA'NTHEMUM.	Diandria Monogynia. R.S.			
1	variabile. B.P. (pu.)	variable.	N. S. W.	1820. 6. 9. G. ½.	
2	montánum. F.I. (bl.)	mountain.	E. Indies.	1825. 7. 2. S. ½.	
3	pulchellum. F.I. (bl.)	nervose.	1796.	1. 10. S. ½.	Andr. deposit. t. 88.
	<i>Justicia nervosa.</i> B.M. 1358.				
4	racemosum. F.I. (bh.)	racemed.	1826.	7. 2. S. ½.	
5	barlerioïdes. F.I. (pa.bl.)	Barleria-like.	1824.	— S. ½.	
6	stríctum. F.I. (bl.)	upright.	1822.	— S. ½.	Bot. reg. t. 867.
7	crenulatum. B.R. (li.)	crenulate.	1824.	10. 2. S. ½.	t. 879.
8	bicolor. s.s. (wh.pu.)	two-coloured.	Philippines.	1802. 5. 8. S. ½.	Bot. mag. t. 1423.
	<i>Justicia bicolor.</i> B.M.				
9	spinosum. R.S. (wh.)	thorny.	W. Indies.	1733. 7. 8. S. ½.	Jacq. amer. 2. t. 2. f. 1.
10	acanthophorum. R.S. (li.)	spiny.	China.	1822. — S. ½.	
11	ambiguum. R.S.	bunch-leaved.	1823. S. ½.	
	<i>parviflorum.</i> L. enum. nec aliorum.				

ORDO CLVIII.

OROBANCHEÆ. Ventenat.

LATHRÆA. S.S.	TOOTHWORT.	Didynamia Angiospermia. L.		
squamaria. s.s. (pu.)	scaly.	Britain. 4. H. ¼.	Eng. bot. v. 1. t. 50.	
OROBANCHE. S.S.	BROOM-RAPE.	Didynamia Angiospermia. L.		
1	màjor. E.B. (br.pu.)	greater.	Britain. 6. 7. H. ¼.	Eng. bot. v. 6. t. 421.
2	elàtior. E.B. (br.)	taller. 7. 8. H. ¼.	— v. 8. t. 568.
3	mìnòr. E.B. (pa.pu.)	smaller. — H. ¼.	— v. 6. t. 422.
4	rùbra. E.B. (pu.re.)	red.	Ireland. 8. H. ¼.	— v. 25. t. 1786.
5	cœrûlea. E.B. (bl.)	blue.	Britain. 7. H. ¼.	— v. 6. t. 423.
6	ramosa. E.B. (pu.bl.)	branching. 8. 9. H. ¼.	— v. 3. t. 184.

ORDO CLIX.

LENTIBULARIÆ. Brown prodr. 1. p. 429.

PINGUICULA. L.	BUTTERWORT.	Diandria Monogynia. L.		
1	vulgâris. s.s. (vi.)	common.	Britain. 5. H.w. ¼.	Eng. bot. v. 1. t. 70.
2	alpina. s.s. (wh.st.)	Alpine.	Europe. 1794. 4. H.w. ¼.	Flor. dan. t. 453.

3 grandiflora, s.s.	(<i>bl.</i>) great-flowered.	Ireland.	5. 6. H.w. 4.	Eng. bot. v. 31. t. 2184.
4 lusitánica, s.s.	(<i>fl.</i>) pale.	Britain.	6. 7. H.w. 4.	——— v. 3. t. 145.
5 lútea, B.R.	(<i>ye.</i>) yellow.	Carolina.	1816.	——— H.w. 4.	Bot. reg. t. 126.
6 edéntula, H.E.F.	(<i>ye.</i>) toothless.	N. America.	1821.	4. 5. H.w. 4.	Hook. ex. flor. t. 16.
UTRICUL'ARIA. L.	HOODED MILFOIL.	Diandria Monogynia. L.			
1 vulgàris, E.B.	(<i>ye.</i>) common.	Britain.	7. H. 4.	Eng. bot. v. 4. t. 253.
2 intermédia, E.B.	(<i>ye.</i>) intermediate.	Ireland.	——— H. 4.	——— v. 35. t. 2489.
3 mìnora, E.B.	(<i>st.</i>) lesser.	Britain.	——— H. 4.	——— v. 4. t. 254.

ORDO CLX.

PRIMULACEÆ. Brown prodr. p. 427.

CY'CLAMEN. L.	CY'CLAMEN.	Pentandria Monogynia. L.			
1 còum, R.S.	(<i>re.</i>) green, round-ld.	S. Europe.	1596.	1. 4. H. 4.	Bot. mag. t. 4.
2 vérnum, B.F.G.	(<i>re.</i>) spring.	———	1814.	2. 5. H. 4.	Swt. br. fl. gar. t. 9.
3 europ'æum, R.S.	(<i>li.</i>) round-leaved.	Europe.	1596.	7. 8. H. 4.	——— t. 176.
4 pérsicum, R.S.	(<i>wh.</i>) Persian.	Cyprus.	1731.	2. 4. G. 4.	Bot. mag. t. 44.
	<i>a inodorum.</i> (<i>wh.pu.</i>) scentless.	———	———	G. 4.	
	β <i>odoratum.</i> (<i>wh.pu.</i>) sweet-scented.	———	———	G. 4.	
	γ <i>albiflorum.</i> (<i>wh.</i>) pure white.	———	———	G. 4.	
	δ <i>laciniatum.</i> B.R. (<i>wh.pu.</i>) jagged.	———	———	G. 4.	
5 hederifolium, E.B.	(<i>li.</i>) Ivy-leaved.	Britain.	6. 8. H. 4.	Bot. reg. t. 1095.
	α <i>purpurascens.</i> (<i>li.</i>) purplish-flower'd.	———	H. 4.	Eng. bot. v. 8. t. 548.
	β <i>ábidum.</i> (<i>wh.</i>) white-flowered.	———	H. 4.	
6 neapolitanum, T.N. (<i>re.</i>) Neapolitan.		Naples.	1826.	——— H. 4.	
7 répandum, F.G.	(<i>re.</i>) angular-leaved.	Greece.	3. 4. F. 4.	Clus. hist. 1. p. 265. ic.
8 latifolium, F.G.	(<i>li.</i>) broad-leaved.	———	1823.	4. 5. F. 4.	Swt. br. fl. gar. t. 117.
DODEC'ATHEON. L.	AMERICAN COWSLIP.	Pentandria Monogynia. L.			Flor. græc. t. 185.
Méadia, R.S.	(<i>li.</i>) Mead's.	Virginia.	1744.	4. 6. H. 4.	
β <i>albiflora.</i> (<i>wh.</i>) white-flowered.	———	1820.	——— H. 4.	Bot. mag. t. 12.	
SOLDANE'LLA. R.S.	SOLDANE'LLA.	Pentandria Monogynia. L.			
1 montâna, R.S.	(<i>bl.</i>) mountain.	Bohemia.	1816.	4. H. 4.	Swt. br. fl. gar. t. 11.
2 alpina, R.S.	(<i>bl.</i>) Alpine.	Switzerland.	1656.	——— H. 4.	Bot. mag. t. 49.
3 púsilla, B.F.G.	(<i>vi.</i>) lesser.	S. Europe.	1824.	——— H. 4.	Swt. br. fl. gar. s. 2. t. 48.
4 mímina, R.S.	(<i>bl.</i>) least.	Carp. mount.	1820.	——— H. 4.	——— s. 2. t. 53.
CORT'USA. L.	BEARS-EAR SANICLE.	Pentandria Monogynia. L.			
Mathioli, R.S.	(<i>li.</i>) short calyxed.	Austria.	1596.	4. 6. H. 4.	Bot. mag. t. 987.
PRI'MULA. L.	PRIMROSE.	Pentandria Monogynia. L.			Bot. reg. t. 539.
1 prænitens, B.R.	(<i>pi.</i>) glossy.	China.	1820.	1. 12. F. 4.	
	<i>sinensis.</i> Lind. col. non Lour.				
	<i>a rosea.</i> (<i>ro.</i>) rose-coloured.	———	———	F. 4.	Bot. mag. t. 2564.
	β <i>albiflora.</i> B.F.G. (<i>wh.</i>) white-flowered.	———	———	F. 4.	Swt. br. fl. gar. t. 196.
	γ <i>dentiflora.</i> (<i>ro.</i>) jagged-flowered.	———	———	F. 4.	Lindl. coll. bot. t. 7.
2 cortusoïdes, R.S.	(<i>re.</i>) Cortusa-leaved.	Siberia.	1794.	5. 7. H. 4.	Bot. mag. t. 399.
3 dentiflora, A.R.	(<i>re.</i>) tooth-flowered.	———	H. 4.	Andr. reposit. t. 451.
4 suavèolens, R.S.	(<i>ye.</i>) sweet-scented.	Naples.	1826.	——— H. 4.	Ten. fl. neap. 1. t. 13.
	<i>Colúnnea.</i> T.N.				
5 inflata, R.S.	(<i>ye.</i>) bellied-calyxed.	Hungary.	1824.	——— H. 4.	Swt. br. fl. gar. s. 2.
6 brevistyla, R.S.	(<i>st.</i>) short-styled.	France.	——— H. 4.	
	β <i>versicolor.</i> (<i>vu.</i>) various-coloured.	———	H. 4.	
7 vèris, R.S.	(<i>re.</i>) Cowslip.	Britain.	5. 6. H. 4.	Eng. bot. v. 1. t. 5.
	β <i>rùbra.</i> (<i>re.</i>) red cowslip.	———	H. 4.	
8 acaulis, R.S.	(<i>st.</i>) common.	Britain.	3. 5. H. 4.	Eng. bot. v. 1. t. 4.
	<i>vulgàris.</i> E.B.				
	β <i>álbo-plena.</i> (<i>wh.</i>) double white.	H. 4.	
	γ <i>sulphureo-plena.</i> (<i>st.</i>) double-sulphur.	H. 4.	
	δ <i>cùpreo-plena.</i> (<i>co.</i>) double-copper.	H. 4.	
	ϵ <i>salmoneo-plena.</i> (<i>pa.</i>) double-salmon.	H. 4.	
	ζ <i>lilacino-plena.</i> (<i>li.</i>) double-lilac.	H. 4.	
	η <i>rùbro-plena.</i> (<i>re.</i>) double Scotch.	H. 4.	
	ϑ <i>purpureo-plena.</i> (<i>d.pu.</i>) double dark purple.	H. 4.	Bot. mag. t. 229.

PRIMULACEÆ.

9	elàtor. R.S.	(st.) Oxlip.	Britain.	3. 5.	H. ȝ.	Eng. bot. v. 8. t. 518.			
	β polyántha.	(va.) Polyanthus.	—	H. ȝ.				
	γ Fletchéri.	(ve.ye.) Fletcher's defiance.	—	H. ȝ.	Swt. flor. guid. t. 49.			
	δ Burnardiàna.	(ve.ye.) Burnard's Formosa	—	H. ȝ.	— t. 97.			
	ε Shadiàna.	(pu.ye.) Shad's telegraph.	—	H. ȝ.	— t. 125.			
	ζ calycántha.	(va.) coloured-calyxed.	—	H. ȝ.				
10	Perreiniána. R.S. (pu.)	Perrein's.	Spain.	—	H. ȝ.	Flugg. an. mus. 12.t.37.			
	Flugeána.	Lehm. mon. prim. p. 36. t. 2.								
11	Pallásii. R.S.	(st.) Pallas's.	Altay.	1820.	—	H. ȝ.	Bot. reg. t. 896.			
12	Aurícula. R.S.	(va.) Auricula.	Switzerland.	1596.	4. 5.	H. ȝ.	Jacq. aust. 5. t. 415.			
	β lútea.	(ye.) yellow-flowered.	—	H. ȝ.				
	γ calycántha.	(ye.) yellow-calyxed.	—	H. ȝ.	Trattin. tabul. t. 431.			
	ð integérriana.	(va.) entire-leaved.	—	H. ȝ.	— t. 432.			
	ε horténsis.	(va.) garden.	—	H. ȝ.	Swt. flor. guid. t. 84.			
	ζ venústa.	(pu.) Howe's Venus.	—	H. ȝ.	— t. 69.			
	η Redmâni.	(bl.) Redman's metropolitan.	—	H. ȝ.	— t. 63.			
	θ melanocycla. (bk.wh.)	Wild's black and clear.	—	H. ȝ.	— t. 3.			
	ι albifólia.	(pu.st.) Taylor's glory.	—	H. ȝ.	— t. 22.			
	κ Hedgeána.	(pu.st.) Hedge's Britannia.	—	H. ȝ.	— t. 131.			
	λ dilécta.	(pu.wh.) Wood's delight.	—	H. ȝ.	μ Warrisíana. (ve.gr.)	Warris's union.	— t. 134.
	ν Moreána.	(pu.gr.) More's Navarino.	—	H. ȝ.	— t. 58.			
	ο Quiragána.	(pu.gr.) Burnard's Gen. Quiraga.	—	H. ȝ.	— t. 129.			
	π Hilliana.	(pu.gr.) Page's Lord Hill.	—	H. ȝ.	— t. 118.			
	ρ violácea.	(vi.gr.) Moore's Violet.	—	H. ȝ.	— t. 105.			
	σ Oldenbúrgħii.	(ve.gr.) Page's Dutchess of Oldenburgh	—	H. ȝ.	— t. 2.			
	τ Alexándri.	(ve.gr.) Stretch's Alexander.	—	H. ȝ.	— t. 10.			
	υ undulátā.	(ve.gr.) Page's champion.	—	H. ȝ.	— t. 15.			
	φ Grimesii.	(ve.gr.) Grimes's privateer.	—	H. ȝ.	— t. 29.			
	χ Lawrieána.	(ve.gr.) Lawrie's Hertfordshire hero.	—	H. ȝ.	— t. 94.			
	ψ indépendans. (bk.gr.)	Booth's freedom.	—	H. ȝ.	— t. 91.			
	w obscura.	(pu.gr.) Cockup's Eclipse.	—	H. ȝ.	— t. 45.			
	aa Pollítiana.	(ve.gr.) Pollit's highland boy.	—	H. ȝ.	— t. 52.			
	ββ Smíthii.	(ve.gr.) Smith's Waterloo.	—	H. ȝ.	— t. 76.			
	γγ gloriósa.	(ve.gr.) Lawrie's glory of Cheshunt.	—	H. ȝ.	— t. 79.			
13	Palinúri. R.S.	(ye.) unequal-bracted.	Naples.	1816.	—	H. ȝ.	Swt. br. fl. gar. t. 8.			
14	Balbisii. R.S.	(ye.) Balbis's.	Italy.	1826.	—	H. ȝ.	Moret. fl. vicent. p. 7. ic.			
15	marginátā. B.M.	(ro.) silver-edged.	Switzerland.	1777.	3. 4.	H. ȝ.	Bot. mag. t. 191.			
	crenátā. R.S.									
16	longiflóra. R.S.	(li.) long-flowered.	1818.	4. 6.	H. ȝ.	Jacq. aust. 5. t. 46.			
17	longiflóra. B.M.	(li.) long-leaved.	Levant.	1790.	4. 5.	H. ȝ.	Bot. mag. t. 392.			
	auriculátā. R.S.									
18	fariñósa. R.S.	(re.) Bird's-eye.	Britain.	4. 7.	H. ȝ.	Eng. bot. v. 1. t. 6.			
19	scótica. s.s.	(pu.) Scotch bird's-eye.	Scotland.	7.	H. ȝ.	Hook. fl. lond. t. 133.			
20	strícta. R.S.	(ii.) Horneman's.	N. Europe.	1822.	6. 7.	H. ȝ.	Flor. dan. t. 1385.			
	Hornemanniana. s.s.									
21	davúrica. R.S.	(ro.) Siberian bird's-eye.	Davuria.	1806.	5. 6.	H. ȝ.	Bot. mag. t. 1219.			
	internmédia. B.M.									
22	pusílla. s.s.	(li.) American bird's-eye.	N. America.	1819.	—	H. ȝ.	Hook. ex. flor. t. 68.			
23	altaica. R.S.	(li.) pale-flowered.	Altay.	1826.	3. 5.	H. ȝ.	Lehm. prim. t. 5.			
24	finmárchica. R.S.	(vi.) Norwegian.	Norway.	1798.	—	H. ȝ.	Flor. dan. t. 188.			
25	nivális. R.S.	(pu.) snowy mountain.	Davuria.	1790.	4. 5.	H. ȝ.	Pall. it. 3. app. t. 9. f. 2.			
26	viscósia. R.S.	(pu.) clammy.	Piedmont.	1792.	4.	H. ȝ.	Trattin. tabul. t. 434.			
27	glutinósa. R.S.	(vi.) glutinous.	S. Europe.	1820.	3. 6.	H. ȝ.	Jacq. aust. t. 26.			
28	carniòlica. R.S.	(vi.) elliptic-leaved.	Carniola.	1826.	—	H. ȝ.	— 5. t. 4.			
29	glaucescens. s.s.	(pu.) glaucescent.	Switzerland.	—	—	H. ȝ.	Swt. br. fl. gar. t. 254.			
30	integrifólia. R.S.	(pu.) entire-leaved.	Pyrenees.	—	6. 7.	H. ȝ.	Bot. mag. t. 942.			
31	Símsii.	(st.) Sims's.	Switzerland.	4. 6.	H. ȝ.	— t. 1161.			
	villòsa β flòre álbo. B.M.									
32	nívea. f.	(wh.) snowy white.	Siberia.	—	H. ȝ.				
	nivális. Hort. nec aliorum.									
33	helvética. B.C.	(pu.) Swiss.	Switzerland.	1790.	—	H. ȝ.	Lodd. bot. cab. t. 348.			
34	pubescens. R.S.	(pu.) pubescent.	Pyrenees.	1768.	—	H. ȝ.	Trattin. tabul. t. 427.			
	villòsa. B.M. 14. non Jacq.									
35	villòsa. R.S.	(pu.) villous.	S. Europe.	—	—	H. ȝ.	Jacq. aust. 5. t. 27.			
36	ciliáta. R.S.	(pu.) ciliated.	Switzerland.	—	H. ȝ.	Swt. br. fl. gar. s. 2.			
	decora. Bot. mag. t. 1922.									
37	mínima. R.S.	(pu.) smallest.	—	1819.	4. 5.	H. ȝ.	Bot. reg. t. 581.			

38 involucrata. L.O. (<i>ye.</i>) involucred. <i>verticillata</i> . B.M. non Vahl. nec Forsk.	Egypt.	1826.	3. 4. F. 4.	Bot. mag. t. 2842.
ANDROS'ACE. L.	ANDROS'ACE.	Pentandria Monogynia. L.		
1 máxima. R.S. (<i>wh.</i>) oval-leaved.	Austria.	1597.	3. 6. H. ⊙.	Jacq. aust. 4. t. 331.
2 elongata. R.S. (<i>wh.</i>) cluster-flowered.	—	1776.	4. 5. H. ⊙.	— 4. t. 330.
3 nana. R.S. (<i>wh.</i>) dwarf.	S. Europe.	1825.	— H. ⊙.	
4 septentrionalis. R.S. (<i>wh.</i>) tooth-leaved.	Russia.	1755.	— H. ⊙.	Bot. mag. t. 2021.
5 alismoides. H.H. (<i>wh.</i>) Alisma-like.	Siberia.	1827.	— H. ⊙.	
6 coronopifolia. A.R. (<i>wh.</i>) Buckshorn-ld. <i>lactiflora</i> . R.S.	—	1806.	6. 9. H. ♂.	Andr. deposit. t. 647.
7 filiformis. R.S. (<i>wh.</i>) filiform.	—	1823.	— H. ⊙.	Gmel. sib. 4. t. 44. f. 4.
8 villosa. R.S. (<i>bh.</i>) villous.	Pyrenees.	1790.	6. 7. H. 4.	Bot. mag. t. 743.
9 chamaejasme. R.S. (<i>bh.</i>) grass-leaved.	Austria.	1768.	6. 8. H. 4.	Lodd. bot. cab. t. 232.
10 brevifolia. R.S. (<i>wh.</i>) short-leaved.	Siberia.	1827.	— H. ♂.	Vill. delph. 2. t. 15.
11 macrocarpa. Led. (<i>wh.</i>) large-capsuled.	—	—	— H. ⊙.	
12 lactea. R.S. (<i>wh.</i>) white-flowered.	—	1752.	— H. 4.	Bot. mag. t. 868.
13 carnea. R.S. (<i>ro.</i>) awl-leaved.	Switzerland.	1768.	7. 8. H. 4.	Lodd. bot. cab. t. 40.
14 obtusifolia. R.S. (<i>bh.</i>) blunt-leaved.	—	1820.	— H. 4.	All. ped. 1. t. 46. f. 1.
15 albana. R.S. (<i>bh.</i>) capitata.	Caucasus.	1827.	— H. 4.	Linn. trans. 11. t. 33.
16 acaulis. Ot. (<i>wh.</i>) stalkless.	Siberia.	1825.	— H. ♂.	
AR'ETIA. W.	AR'ETIA.	Pentandria Monogynia. L.		
1 helvetica. w. (<i>st.</i>) imbricated.	Switzerland.	1775.	5. 6. H. 4.	Schk. hand. 1. t. 32.
2 argentea. (<i>st.</i>) silvery.	—	1826.	— H. 4.	Lap. pyr. ic. ined. t. 67.
<i>Androsace argentea</i> . R.S.				
3 alpina. w. (<i>pi.</i>) linear-leaved.	—	1775.	— H. 4.	Jacq. aust. 5. app. t. 18.
4 pubescens. L.C. (<i>pi.</i>) pubescent.	—	1820.	— H. 4.	Lodd. bot. cab. t. 1273.
5 Vitaliana. w. (<i>ye.</i>) grass-leaved.	Pyrenees.	1787.	— H. 4.	— t. 166.
<i>Primula sedifolia</i> . P.L. 107. <i>Androsace Vitaliana</i> . R.S.				
TRIENTALIS. W.	WINTER-GREEN.	Heptandria Monogynia. L.		
1 europaea. w. (<i>wh.</i>) oval-leaved.	Britain.	—	5. 6. H. 4.	Eng. bot. v. 1. t. 15.
2 americana. Ph. (<i>wh.</i>) spear-leaved.	N. America.	1816.	7. 8. H. 4.	Bart. flor. amer. 2. t. 48.
C'ORIS. L.	C'ORIS.	Pentandria Monogynia. L.		
monspeliensis. L. (<i>vi.</i>) Montpelier.	S. Europe.	1640.	6. 7. G. ♂.	
HOTT'ONIA. L.	WATER-VIOLET.	Pentandria Monogynia. L.		
palustris. E.B. (<i>pi.</i>) marsh.	England.	—	7. 8. H.w. 4.	Eng. bot. v. 6. t. 364.
LYSIMACHIA. L.	LOOSE-STRIFE.	Pentandria Monogynia. L.		
1 vulgaris. R.S. (<i>ye.</i>) common.	Britain.	—	7. 9. H. 4.	Eng. bot. v. 11. t. 761.
2 Ephemerum. R.S. (<i>wh.</i>) Willow-leaved.	Spain.	1730.	— H. 4.	Bot. mag. t. 2346.
3 atropurpurea. R.S. (<i>d.pu.</i>) dark purple.	Levant.	—	— H. ⊙.	Flor. græc. t. 187.
4 dūbia. R.S. (<i>li.</i>) pale purple.	—	1759.	— H. ♂.	— t. 188.
5 strīcta. R.S. (<i>ye.</i>) upright.	N. America.	1781.	7. 8. H. 4.	Bot. mag. t. 104.
bulbifera. B.M. <i>racemosa</i> . M.				
6 capitata. R.S. (<i>ye.</i>) headed.	—	1813.	5. 7. H. 4.	
7 thyrsiflora. R.S. (<i>ye.</i>) tufted.	England.	—	— H. 4.	Eng. bot. v. 3. t. 176.
8 quadrifolia. R.S. (<i>ye.</i>) four-leaved.	N. America.	1798.	7. 8. H. 4.	Lam. ill. 1. t. 101. f. 2.
<i>hirsuta</i> . M.				
9 longifolia. R.S. (<i>ye.</i>) long-leaved.	—	—	— H. 4.	Bot. mag. t. 660.
<i>quadriflora</i> . B.M.				
10 lanceolata. R.S. (<i>ye.</i>) spear-leaved.	—	1826.	— H. 4.	
11 verticillata. R.S. (<i>ye.</i>) whorl-leaved.	Hungary.	1818.	6. 8. H. 4.	Swt. br. fl. gar. t. 21.
12 ciliata. R.S. (<i>ye.</i>) ciliated.	N. America.	1732.	— H. 4.	Wachend. hort. t. 12.
13 punctata. R.S. (<i>ye.</i>) punctated.	S. Europe.	1658.	7. 8. H. 4.	Jacq. aust. 4. t. 366.
14 angustifolia. R.S. (<i>ye.</i>) narrow-leaved.	N. America.	1803.	7. 9. H. 4.	
15 hybrida. R.S. (<i>ye.</i>) hybrid.	—	1806.	— H. 2.	
16 nemorum. R.S. (<i>ye.</i>) wood.	Britain.	—	5. 7. H. 4.	Eng. bot. v. 8. t. 527.
17 anagalloides. F.G. (<i>ye.</i>) Pimpernell-fl'd.	Crete.	1789.	— F. 4.	Flor. græc. t. 198.
18 maculata. B.P. (<i>ye.</i>) spotted.	N. S. W.	1822.	— G. 4.	
19 Nummularia. R.S. (<i>ye.</i>) Moneywort.	Britain.	—	6. 7. H. 4.	Eng. bot. v. 8. t. 528.
LUBI'NIA. S.S.	LUBI'NIA.	Pentandria Monogynia. S.S.		
atropurpurea. s.s. (<i>pu.</i>) dark-purple.	C. B. S.	1823.	7. 9. H. 4.	Swt. br. fl. gar. s.2. t. 34.
ASTEROLINON. LK.	ASTEROLINON.	Pentandria Monogynia. L.en.		
stellatum. L.en. (<i>bh.</i>) starry.	Italy.	1658.	6. H. ⊙.	Flor. græc. t. 189.
<i>Lysimachia Linum-stellatum</i> . F.G.				
ANAGA'LLIS. L.	PIMPERNELL.	Pentandria Monogynia. L.		
1 arvensis. E.B. (<i>sc.</i>) red-flowered.	Britain.	—	6. 9. H. ⊙.	Eng. bot. v. 8. t. 529.
<i>phœnicea</i> . R.S.				
2 carnea. Schl. (<i>fl.</i>) flesh-coloured.	Switzerland.	—	— H. ⊙.	Swt. br. fl. gar. t. 132.
3 indica. B.F.G. (<i>bl.</i>) Indian.	E. Indies.	1824.	— H. ⊙.	Eng. bot. v. 26. t. 1823.
4 cœrulea. R.S. (<i>bl.</i>) blue-flowered.	Britain.	—	— H. ⊙.	

PRIMULACEÆ.

5 collina. R.S.	(sc.) large-flowered. Morocco.	1803.	5. 7. G. ♂.	Bot. mag. t. 831.
fruticōsa. B.M.	<i>grandiflōra</i> . A.R. 367.			
6 latifòlia. R.S.	(bl.) broad-leaved. Spain.	1759.	— G. ♂.	Bot. mag. t. 2380.
7 Monéllia. R.S.	(bl.) blue Italian. Italy.	1648.	5. 9. G. ♀.	t. 319.
8 linifòlia. R.S.	(bl.) Flax-leaved. Portugal.	1796.	6. 7. G. ♀.	Moris. s. 5. t. 26. f. 3.
9 tenellà. R.S.	(ro.) bog. Britain.	7. 8. H. ♀.	Eng. bot. v. 8. t. 530.
CENTU'NCULUS. L.	BASTARD PIMPERNELL.	Tetrandria Monogynia. L.		Eng. bot. v. 8. t. 531.
mínimus. E.B.	(fl.) least. Britain.	6. 7. H. ♂.	
? SAM'OLUS. B.P.	BROOK-WEED.	Pentandria Monogynia. L.		
1 Valerándi. E.B. (wh.) common.	Britain.	6. 8. H. ♀.	Eng. bot. v. 10. t. 703.
2 littorális. B.P. (wh.) sea-side.	N. S. W.	1806.	7. 9. G. ♀.	Lab. n. hcll. I. t. 54.
3 campanuloides. s.s. (w.)	Campanula like. C. B. S.	1816.	— G. ♀.	
	Campánula poròsa. w.			

ORDO CLXI.

GLOBULARIÆ. *Lam. et DC. fl. fr. ed. 3. v. 3. p. 427.*

GLOBULARIA. L.	GLOBULARIA.	Tetrandria Monogynia. L.		
1 longifòlia. w.	(bl.) Willow-leaved. Madeira.	1775.	7. 8. G. ♂.	Bot. reg. t. 685.
salicina. R.S.				
2 Alypum. R.S. (pa.bl.) three-toothed.	S. Europe.	1640.	8. 9. G. ♂.	Garid. aix. t. 42.
β integrifòlia. (pa.bl.) entire-leaved.	—	—	— G. ♂.	Røem. et Ust. b.m. 7.t.5.
3 linifòlia. R.S.	(bl.) Flax-leaved. Spain.	1818.	5. 6. H. ♀.	
4 vulgáris. R.S.	(bl.) common. Europe.	1640.	— H. ♀.	Swt. br. fl. gar. t. 20.
5 spinosa. R.S.	(bl.) prickly-leaved. Spain.	—	— F. ♀.	
6 cordifòlia. R.S.	(bl.) wedge-leaved. Germany.	1633.	6. 7. H. ♀.	Swt. br. fl. gar. t. 34.
7 nána. R.S.	(li.) dwarf. S. France.	1818.	— H. ♀.	DC. ic. pl. gal. rar. 1.t.3.
8 nudicaulis. R.S.	(bl.) naked-stalked. Germany.	1629.	— H. ♀.	Jacq. aust. 3. t. 230.
9 bellidifòlia. R.S.	(bl.) Daisy-leaved. Italy.	1824.	— H. ♀.	
10 incanescens. R.S. (pu.) hoary.	—	1828.	— H. ♀.	Viv. frag. it. 1. t. 3.

***** SUBCLASSIS IV. MONOCHLAMYDEÆ.



ORDO CLXII.

PLUMBAGINEÆ. *Brown prodr. 1. p. 425.*

STATICE. T.	THRIFT.	Pentandria Pentagynia. L.		
1 vulgáris.	(pa. li.) common.	Europe.	6. 8. H. ♀.
Armèria vulgáris. R.S.				Schk. handb. t. 87.
2 arenaria. P.S.	(li.) sand.	France.	5. 8. H. ♀.
3 alpina.	(vi.) flattish-stalked.	Carinthia.	— H. ♀.
4 littorális.	(vi.) ciliate-leaved.	Portugal.	— H. ♀.
Armèria littorális. R.S.				
5 marítima. Mill.	(ro.) sea-side.	Britain.	5. 7. H. ♀.
Armèria. E.B. Armèria marítima. R.S.				Eng. bot. v. 4. t. 226.
6 hùmilis. P.S.	(li.) dwarf.	Portugal.	— H. ♀.
7 juniperifòlia. w.	(li.) Juniper-leaved.	Spain.	1818.	— H. ♀.
8 pinifòlia. Brot.	(li.) Pine-leaved.	Portugal.	— H. ♀.
9 púngens. Brot.	(li.) pungent-leav'd.	—	— H. ♀.
10 alliacea. w.	(wh.) Garlic-leaved.	Spain.	1798.	5. 6. H. ♀.
11 scorzonerifòlia.	(li.) spear-leaved.	S. Europe.	1816.	— H. ♀.
Armèria scorzonerifòlia. s.s.				Flor. græc. t. 29.

12 lusitánica. P.S.	(sc.) large-leaved.	S. Europe.	1814.	5. 6. F. 4.		
13 latifòlia.	(li.) broad-leaved.	Portugal.	1740.	5. 7. H. 4.	Jacq. vind. 1. t. 42.	
cephalòtes. H.K.	<i>Armèria latifòlia</i> . R.S.					
14 plantagínea.	(li.) Plantain-leav'd.	S. Europe.	1816.	5. 6. H. 4.		
15 denticulàta. Brot.	(fl.) tooth-leaved.	Italy.	—	— H. 4.		
16 dianthóides. B.F.G.	(li.) Pink-like.	S. Europe.	1824.	5. 7. H. 4.	Swt. br. fl. gar. ser. 2.	
	<i>Armèria dianthóides</i> . s.s.					
17 hírtia.	(li.) hairy.	Gibraltar.	1818.	— H. 4.		
	<i>Armèria hírtia</i> . R.S.					
18 fasciculàta. P.S.	(li.) bundle-leaved.	Portugal.	4. 8. H. 4.	Venten. cels. t. 38.	
TAXA'NTHEMA. B.P.	SEA LAVENDER.	Pentandria Pentagynia.				
1 graminifòlia. R.S.	(sc.) Grass-leaved.	Siberia.	1780.	6. 7. H. 4.		
2 Limónium. R.S.	(bl.) common.	England.	5. 8. H. 4.	Eng. bot. v. 2. t. 102.	
	<i>Státice Limónium</i> . E.B.					
3 elàta. s.s.	(bl.) tall.	Siberia.	1824.	— H. 4.		
4 Gmelini. R.S.	(bl.) Gmelin's.	—	1796.	6. 8. H. 4.	Gmel. sib. 2. t. 90.	
5 caroliniàna. R.S.	(bl.) Carolina.	N. America.	1824.	— H. 4.		
6 scopària. R.S.	(bl.) Broom.	Siberia.	1796.	— H. 4.		
7 latifòlia. R.S.	(bl.) broad-leaved.	—	1791.	5. 7. H. 4.	Hoff. c. gott. 1796. t. 1.	
8 purpuráta. R.S.	(pu.) purple.	C. B. S.	1800.	6. 7. G. 4.	Flor. græc. t. 295.	
9 bellidifòlia. R.S.	(bl.) Daisy-leaved.	Greece.	1823.	— G. 4.	— t. 296.	
10 globulariæfòlia. R.S.	(li.) Globularia-leav'd.	S. Europe.	—	— F. 4.	Lobel. ic. p. 295. f. 2.	
11 oleæfòlia. R.S.	(vi.) Olive-leaved.	Italy.	1688.	5. 8. H. 4.	Willd. h. ber. t. 63.	
12 Willdenowiâna. R.S.	(vi.) Willdenow's.	S. France.	1804.	6. 8. H. 4.		
	<i>Státice spathulâta</i> . w. non Desf.					
13 auriculæfòlia. R.S.	(bl.) Auricula-leav'd.	Barbary.	1781.	7. 8. F. 4.		
14 incàna. B.F.G.	(re.) hoary.	Levant.	1823.	— H. 4.	Swt. br. fl. gar. t. 272.	
α sangüínea.	(cr.) crimson-flowered.	—	—	— H. 4.	— t. 272.f.1.	
β rosea.	(ro.) rose-coloured.	—	—	— H. 4.	— t. 272.f.2.	
γ álba.	(wh.) white-flowered.	—	—	— H. 4.		
15 spatulâta. R.S.	(bl.) spatula-leaved.	Barbary.	1804.	6. 8. F. 4.	Bot. mag. t. 1617.	
16 cordâta. R.S.	(bl.) retuse-leaved.	S. Europe.	1752.	5. 7. F. 4.	Barrel. ic. 805.	
17 emarginâta. R.S.	(bl.) emarginate.	Gibraltar.	—	— F. 4.		
18 cásplia R.S.	(pa. bl.) Caspian.	Siberia.	—	— F. 4.	Gmel. sib. 2. t. 89. f. 2.	
19 scábra. R.S.	(bl.) rough-branch'd.	C. B. S.	1788.	5. 7. G. 4.		
20 tetragôna. R.S.	(bl.) square-stalked.	—	1816.	— G. 4.		
21 reticulâta. R.S.	(bl.) matted.	England.	—	7. 8. H. 4.	Eng. bot. v. 5. t. 328.	
	<i>Státice reticulâta</i> . E.B.					
22 dichótoma. R.S.	(bl.) forked.	Spain.	1818.	— H. 4.	Cavan. ic. 1. t. 50.	
23 virgâta. R.S.	(bl.) twiggly.	—	—	— H. 4.		
24 vimínea. R.S.	(bl.) slender.	—	1823.	— H. 4.		
25 echoídeas. F.G.	(li.) rough-leaved.	S. Europe.	1752.	— F. 4.	Flor. græc. t. 299.	
26 rórida. F.G.	(pa. bl.) frosted.	Levant.	—	— F. 4.	— t. 298.	
	<i>echoídeas</i> . Prodr. fl. græc. non L.					
27 palmâris. F.G.	(ro.) rosy-coloured.	—	1826.	— F. 4.	Flor. græc. t. 297.	
28 speciòsa. B.F.G.	(re.) Plantain-leaved.	Russia.	1776.	— H. 4.	Swt. br. fl. gar. t. 105.	
	<i>Státice speciòsa</i> . Bot. mag. 656.					
29 conspícuâ. R.S.	(re.) conspicuous.	—	1804.	— H. 4.	Bot. mag. t. 1629.	
	<i>Státice conspícuâ</i> . B.M.					
30 tatárica. B.F.G.	(re.) Tartarian.	—	1731.	6. 7. H. 4.	Swt. br. fl. gar. t. 37.	
31 aceròsa. M.B.	(re.) needle-leaved.	M. Ararat.	1829.	— F. 4.	Buxb. cent. 2. p. 18. t. 10.	
32 flexuòsa. R.S.	(ro.) zigzag.	Siberia.	1791.	7. 8. H. 4.	Gmel. sib. 2. t. 89. f. 1?	
33 minûta. R.S.	(li.) small.	S. Europe.	1658.	6. 7. H. 4.	Bocc. sic. t. 13. f. 3.	
34 pectinâta. R.S.	(bl.) triangular-stalk'd.	Canaries.	1780.	9. 10. F. 4.		
35 cineràa. R.S.	(wh.) ash-coloured.	C. B. S.	1822.	— G. 4.		
36 macrophylla. s.s.	(bl. w.) large-leaved.	Canaries.	1816.	7. 9. F. 4.		
37 suffruticòsa. R.S.	(li.) narrow-leaved.	Siberia.	1779.	5. 9. H. 4.	Gmel. sib. 2. t. 88. f. 2.3.	
38 monopétala. R.S.	(pu.) Sicilian shrubby.	Sicily.	1731.	7. 8. F. 4.	Bocc. sic. t. 16. 17.	
39 cylindrifòlia. R.S.	(vi.) round-leaved.	N. Africa.	1738.	— G. 4.		
40 ferulacea. R.S.	(ye.) Fennel-like.	Siberia.	1796.	5. — H. 4.	Moris. s. 15. t. 1. f. 23.	
41 sinuâta. R.S.	(bl. ye.) scollop-leaved.	Levant.	1629.	5. 9. F. 4.	Bot. mag. t. 71.	
42 Thouïni. R.S.	(bl. ye.) Thouin's.	Africa.	1738.	6. 10. F. 4.	Mart. cent. t. 84.	
43 alâta. R.S.	(ye.) winged.	—	1806.	— F. 4.		
44 spicâta. R.S.	(wh.) spiked.	Caucasus.	1828.	8. 10. H. 4.	Gmel. sib. 2. t. 91. f. 2.	
45 ægyptiaca. R.S.	(wh.) Egyptian.	Egypt.	1816.	6. 10. F. 4.	Bot. mag. t. 2363.	
46 mucronâta. R.S.	(li.) curled.	Barbary.	1784.	— F. 4.	L'Herit. stirp. t. 13.	
	<i>PLUMB'AGO</i> . L.	LEADWORT.	Pentandria Monogynia. L.			
1 europaea. R.S.	(vi.) European.	S. Europe.	1596.	9. 10. H. 4.	Bot. mag. t. 2139.	
2 lapathifòlia. R.S.	(wh.) Dock-leaved.	Iberia.	1824.	— H. 4.		

3 capénsis. R.S.	(<i>bl.</i>) blue-flowered.	C. B. S.	1818.	9. 3. S. h.	Bot. reg. t. 417.
4 tristis. H.K.	(<i>br.</i>) dark-flowered.	—	1792.	5. 6. G. h.	Rheed. mal. 10. t. 8.
5 zeylánica. R.S.	(<i>wh.</i>) Ceylon.	E. Indies.	1731.	4. 9. S. h.	
6 occidentális.	(<i>wh.</i>) West Indian.	W. Indies.	1817.	— S. h.	
zeylánica β. R.S.	<i>scándens</i> . Hort.				
7 mexicána. K.S.	(<i>vi.</i>) Mexican.	Mexico.	1829.	4. 8. S. 4.	
8 rósea. R.S.	(<i>ro.</i>) rose-coloured.	E. Indies.	1777.	3. 7. S. h.	Bot. mag. t. 230.
9 scándens. R.S.	(<i>wh.</i>) climbing.	W. Indies.	1699.	7. 8. S. h.	Jacq. amer. pict. t. 13.
10 rhomboídea. B.M.	(<i>bl.</i>) rhomboid-leaved.	—	1828.	8. 10. S. ⊖.	Bot. mag. t. 2917.
11 micránta. F.	(<i>pu.</i>) small-flowered.	Persia.	1829.	6. 8. H. ⊖.	

ORDO CLXIII.

PLANTAGINEÆ. *Juss. gen. 89.*

GLA'UX. L.

marítima. R.S.

PLANTAGO. L.

màjor. R.S.

 β rósea.

2 asiática. R.S.

3 Cornuti. R.S.

Gouani. s.s.

4 Brútia. R.S.

5 gentianóides. S.S.

6 exaltata. R.S.

7 cordáta. R.S.

8 máxima. R.S.

9 crássia. s.s.

crispus. Jacq.

10média. R.S.

11 camtschática. s.s.

12 lanceolata. R.S.

13 sibérica. R.S.

14 eriophóra. R.S.

15 montaña. R.S.

16 capénsis. R.S.

17 altíssima. R.S.

18 Schöttii. R.S.

19 Lagópus. R.S.

20 eriostachya. R.S.

21 sphærocéphala. R.S.

w. round-headed.

alpina. Jacq. non L.

22 lusitánica. R.S.

23 pseudo-lusitánica. R.S.

24 virgínica. R.S.

25 depréssia. R.S.

26 sparsiflóra. R.S.

27 saxátilis. R.S.

28 hungárica. R.S.

29 sericea. R.S.

argentea. DC.

30 álbicans. R.S.

31 mexicána. s.s.

32 patagònica. R.S.

33 victoriális. R.S.

34 holoserícea. R.S.

35 microcéphala. R.S.

wh. small-headed.

36 Bellárdi. R.S.

(br.) Bellárdi's.

37 crética. R.S.

(st.) Cretan.

38 gramínea. R.S.

wh. grass-leaved.

39 marítima. R.S.

gr. sea-side.

40 salsa. R.S.

wh. saltwort.

41 crassifolia. R.S.

br. thick-leaved.

42 hirsuta. R.S.

wh. hairy.

BLACK SALTWORT. Pentandria Monogynia. L.

(li.) sea-side. Britain. 5. 6. H. 4.

PLANTAIN. Tetrandria Monogynia. L.

(gr.) greater. Britain. 5. 7. H. 4.

(gr.) rose-bracted. — — H. 4.

(li.) broad-leaved. Siberia. 1787. 7. H. 4.

(wh.) Cornuti's. S. Europe. 1801. 7. 8. H. 4.

Calabria. 1824. — H. 4.

Greece. 1826. 5. 8. H. 4.

(wh.) tallest. 1824. 7. 8. H. 4.

(wh.) heart-leaved. N. America. 1818. — H. 4.

(li.) hollow-leaved. Siberia. 1763. — H. 4.

(wh.) succulent. S. Europe. 1793. 6. 7. H. 4.

Britain. 5. 7. H. 4.

Kamtschatka. 1819. — H. 4.

Britain. — H. 4.

Siberia. 1823. — H. 4.

(wh.) woolly-leaved. Portugal. 1825. — H. 4.

(wh.) mountain. S. Europe. 1823. — H. 4.

(wh.) Cape. C. B. S. 1788. 5. 8. G. 4.

(wh.) tall. Italy. 1774. 6. 7. H. 4.

(br.) Schott's. Dalmatia. 1829. — H. 4.

(br.) round-headed. Spain. 1683. — H. 4.

(wh.) woolly-spiked. Naples. 1825. 7. 8. H. ⊖.

(wh.) round-headed. Austria. 1774. 6. 7. H. 4.

Portugal. 1781. 7. 8. H. 4.

1819. — H. 4.

N. America. 1688. 6. 9. H. ⊖.

(br.) depressed. — 1818. — H. 4.

(br.) scattered-flowered. — — H. ⊖.

Iberia. 1820. 6. 8. H. 4.

Hungary. 1824. — H. 4.

1820. — H. 4.

S. Europe. 1776. 6. 9. H. 4.

Mexico. 1820. 6. 7. H. 4.

Patagonia. 1793. 6. 9. H. ⊖.

S. Europe. 1824. — H. 4.

Switzerland. — — H. 4.

.... H. ⊖.

S. Europe. 1797. — H. ⊖.

Bellárdi's. 1711. 6. 7. H. ⊖.

Candia. 1711. 6. 7. H. ⊖.

France. 1804. — H. 4.

Britain. 7. H. 4.

Tauria. 1819. — H. 4.

Egypt. 1824. — H. 4.

C. B. S. 1801. 6. 7. G. 3.

Eng. bot. v. 1. t. 13.

Eng. bot. v. 22. t. 1558.

Gmel. sib. 4. t. 37?

Corn. canad. p. 163. ic.

Jacq. f. ecl. t. 72.

Jacq. ic. 1. t. 26.

Jacq. col. sup. 34. t. 16.

Eng. bot. v. 22. t. 1559.

Eng. bot. v. 8. t. 507.

Jacq. obs. 4. t. 83.

Flor. græc. t. 144.

Jacq. vind. 2. t. 125.

Barrel. ic. t. 745.

Moris. s. 8. t. 15. f. 8.

W. et K. hung. 3. t. 203.

t. 151.

Flor. græc. t. 145.

Jacq. ic. 2. t. 306.

Flor. græc. t. 146.

t. 147.

Dod. pent. 108. ic.

Eng. bot. v. 3. t. 175.

Jacq. schœn. 3. t. 258.

43 <i>tùmida</i> . L.en.	(<i>wh.</i>) tumid-capsuled. Chile.	1819.	6. 7. H. ⊖.	
44 <i>alpina</i> . R.S.	(<i>li.</i>) Alpine. S. Europe.	1774.	— H. ȳ.	
45 <i>tenuiflòra</i> . R.S.	(<i>br.</i>) slender-flowered. Hungary.	1819.	— H. ⊖.	W. et K. hung. t. t. 39.
46 <i>bidentata</i> . R.S.	(<i>wh.</i>) white margined. Switzerland.	1824.	— H. ȳ.	
47 <i>serpentina</i> . R.S.	(<i>wh.</i>) serpentine. S. France.	1826.	— H. ȳ.	
48 <i>recurvata</i> . R.S.	(<i>wh.</i>) recurved-leaved. S. Europe.	1799.	— H. ♂.	Murr. c. gott. 1780. t. 6.
49 <i>Wulfeni</i> . R.S.	(<i>wh.</i>) Wulfen's. Carinthia.	1802.	— H. ȳ.	Lobel. ic. 429.
50 <i>subulàta</i> . R.S.	(<i>wh.</i>) awl-leaved. S. Europe.	1596.	7. H. ȳ.	Jacq. coll. 1. t. 10.
51 <i>Holosteum</i> . R.S.	(<i>wh.</i>) Scopoli's. Carniola.	1825.	7. 8. H. ȳ.	
52 <i>elongatà</i> . R.S.	(<i>wh.</i>) long-spiked. Louisiana.	1821.	— H. ȳ.	
53 <i>humifusa</i> . Ber.	(<i>wh.</i>) trailing.	1825.	— H. ȳ.	
54 <i>canescens</i> . Schr.	(<i>wh.</i>) canescent.	1826.	7. 9. H. ⊖.	
55 <i>micrantha</i> . Led.	(<i>gr.</i>) small-flowered. Siberia.	1829.	6. 8. H. ȳ.	
56 <i>Corónopus</i> . R.S.	(<i>gr.</i>) star of the earth. Britain.	4. 9. H. ⊖.	Eng. bot. v. 13. t. 892.
57 <i>Jacquinii</i> . R.S.	(<i>wh.</i>) Jacquin's. S. Europe.	1801.	7. 8. H. ȳ.	Jacq. ic. 1. t. 27.
<i>Cornuti</i> . Jacq.				
58 <i>Serraria</i> . R.S.	(<i>wh.</i>) saw-leaved. Barbary.	1640.	6. 7. H. ȳ.	Moris. s. 8. t. 16. f. 19.
59 <i>macrorhiza</i> . R.S.	(<i>br.</i>) large-rooted. Morocco.	1798.	7. 8. H. ȳ.	Bocc. sic. t. 15. f. 2.
60 <i>Lœflingii</i> . R.S.	(<i>br.</i>) narrow-leaved. Spain.	— H. ⊖.	Jacq. vind. 2. t. 126.
61 <i>amplexicaulis</i> . R.S.	(<i>br.</i>) stem-clasping.	1797.	6. 7. H. ⊖.	Cavan. ic. 2. t. 125.
62 <i>villosa</i> . R.S.	(<i>gr.</i>) villous. Germany.	1804.	— H. ⊖.	
63 <i>vaginatà</i> . R.S.	(<i>li.</i>) sheathed. Canaries.	1816.	— G. ḥ.	Venten. cels. t. 29.
64 <i>Cynops</i> . R.S.	(<i>gr.</i>) shrubby. S. Europe.	1596.	5. 8. H. ḥ.	Jacq. fragm. t. 182.
65 <i>áfra</i> . R.S.	(<i>wh.</i>) Barbary.	1640.	6. F. ḥ.	Moris. s. 8. t. 17. f. 4.
66 <i>divaricatà</i> . R.S.	(<i>wh.</i>) spreading.	1824.	— H. ȳ.	
67 <i>squarròsa</i> . R.S.	(<i>wh.</i>) leafy-spiked. Egypt.	1787.	8. 9. H. ⊖.	Jacq. ic. 1. t. 28.
68 <i>Psylium</i> . R.S.	(<i>bh.</i>) Fleawort. S. Europe.	1562.	7. 8. H. ⊖.	Flor. græc. t. 149.
69 <i>arenaria</i> . R.S.	(<i>wh.</i>) sand. Hungary.	1804.	5. 8. H. ⊖.	W. et K. hung. 1. t. 51.
70 <i>strícta</i> . R.S.	(<i>bh.</i>) upright. Morocco.	—	7. 8. H. ⊖.	Moris. s. 3. 8. t. 17. f. 2.
71 <i>sícula</i> . R.S.	(<i>wh.</i>) Sicilian. Sicily.	1820.	7. 10. H. ⊖.	
72 <i>parviflora</i> . R.S.	(<i>br.</i>) small-flowered. Barbary.	1826.	— H. ȳ.	
73 <i>Ispaghul</i> . R.S.	(<i>wh.</i>) East Indian. E. Indies.	1819.	7. 8. S. ⊖.	
74 <i>índica</i> . R.S.	(<i>wh.</i>) Indian. India.	1780.	— H. ⊖.	
75 <i>pùmila</i> . R.S.	(<i>wh.</i>) dwarf. S. Europe.	1790.	— H. ⊖.	Murr. c. gott. 1778. t. 5.
76 <i>nítida</i> . R.S.	(<i>bl.</i>) glossy.	1822.	7. 10. H. ⊖.	Bot. mag. t. 2616.
77 <i>brasiliensis</i> . B.M.	(<i>wh.</i>) Brazilian. Brazil.	—	— S. ȳ.	Eng. bot. v. 7. t. 468.
LITTORE'LLA. L.	SHORE-WEED. <i>Monocècia Tetrandria</i> . L.			
lacústris. E.B.	(<i>gr.</i>) Plantain-leaved. Britain.	6. 9. H. ȳ.	

ORDO CLXIV.

NYCTAGINEÆ. *Juss. gen. 90.*OXYBA'PHUS. R.S. UMBRELLA-WORT. *Triandria Monogynia* R.S.

1 <i>viscosus</i> . R.S.	(<i>pu.</i>) viscid. Pern.	1793.	5. 9. H. ȳ.	
2 <i>glabrifolius</i> . R.S.	(<i>pu.</i>) smooth-leaved. New Spain.	1811.	7. 8. H. ȳ.	
3 <i>ovatus</i> . R.S.	(<i>re.</i>) oval-leaved. Peru.	1820.	— H. ȳ.	
4 <i>Cervantésii</i> . Lag.	(<i>pu.</i>) Cervantes'. Mexico.	1822.	7. 10. H. ȳ.	
5 <i>expansus</i> . R.S.	(<i>pu.</i>) expanded. Lima.	—	— H. ȳ.	
6 <i>aggregátus</i> . R.S.	(<i>re.</i>) aggregate. New Spain.	1811.	— H. ȳ.	
7 <i>nyctaginea</i> . N.	(<i>re.</i>) heart-leaved. Missouri.	1823.	7. 9. H. ȳ.	
<i>Alliònia nyctaginea</i> . M. <i>Calymènia nyctaginea</i> . N.				
8 <i>álbida</i> . N.	(<i>li.</i>) white-leaved. Carolina.	1824.	— H. ⊖.	
9 <i>pilosa</i> . N.	(<i>re.</i>) pilose. Missouri.	1812.	— H. ⊖.	
<i>Calymènia pilosa</i> . N. <i>Alliònia ovata</i> . Ph.				
10 <i>hirsuta</i> . N.	(<i>re.</i>) hairy. Louisiana.	—	— H. ȳ.	
11 <i>angustifolia</i> .	(<i>li.</i>) narrow-leaved.	—	— H. ȳ.	
<i>Alliònia linearis</i> . Ph. <i>Calymènia angustifolia</i> . N.				
12 <i>décumbens</i> . N.	(<i>li.</i>) decumbent. Missouri.	1818.	— H. ȳ.	
MIRA'BILIS. R.S.	MARVEL OF PERU. <i>Pentandria Monogynia</i> . L.			
1 <i>dichótoma</i> . R.S.	(<i>re. wh.</i>) forked. Mexico.	1640.	7. 8. G. ȳ.	
<i>a ràbra</i> .	(<i>re.</i>) red-flowered.	—	— F. ȳ.	Plenck ic. off. t. 139.
<i>β álba</i> .	(<i>wh.</i>) white-flowered.	—	— F. ȳ.	

Bot. mag. t. 434.
Cavan. ic. 4. t. 379.
Flor. peruv 1. t. 75. f. b.
Swt. br. fl. gar. t. 84.
Flor. peruv. 1. t. 75. f. a.
Cavan. ic. 5. t. 437.

2 Jalápa. R.S.	(va.) common.	India.	1596.	6. 9. H. Ȑ.	Bot. mag. t. 371.
α álba.	(wh.) white-flowered.	—	—	H. Ȑ.	
β fláva.	(ye.) yellow-flowered.	—	—	H. Ȑ.	
γ rúbra.	(re.) red-flowered.	—	—	H. Ȑ.	
δ rúbro-fláva.	(re.ye.) red and yellow.	—	—	H. Ȑ.	
ε rúbro-álba.	(re.wh.) red and white.	—	—	H. Ȑ.	
3 hybrida. R.S.	(re.) hybrid.	Hybrid.	1818.	6. 9. H. Ȑ.	Sm. exot. bot. 1. t. 23.
4 longiflòra. R.S.	(wh.) long-flowered.	Mexico.	1759.	—	H. Ȑ.
5 suavéolens. K.S.	(wh.) sweet-scented.	—	1824.	—	H. Ȑ.
ABR'ONIA. R.S.	ABR'ONIA. Pentandria Monogynia. R.S.				
1 umbellátæ. R.S.	(bh.) flesh-coloured.	California.	1823.	6. 8. H. Ȑ.	
2 melliféra. B.M.	(wh.) honey-scented.	—	1827.	5. 9. H. Ȑ.	Hook. ex. flor. t. 194. Bot. mag. t. 2879.
ALL'ONIA. K.S.	ALL'ONIA. Tetrandria Monogynia. L.				
1 violácea. K.S.	(vi.) violet-coloured.	Cumana.	1820.	7. 9. H. ⊙.	L'Herit. stirp. t. 31.
2 incarnátæ. K.S.	(bh.) flesh-coloured.	—	—	H. ⊙.	
BOERHA'AVIA. L.	HOGWEED. Mono-Di-Tetrandria Monogynia.				
1 erécta. R.S.	(wh.) upright.	W. Indies.	1733.	7. 9. S. Ȑ.	Jacq. vind. 1. t. 5. 6.
2 procúmbens. F.I. (w.re.) procumbent.	E. Indies.	—	—	S. Ȑ.	Rheed. mal. 7. t. 56.
3 diffusa. R.S.	(sc.) spreading.	W. Indies.	1690.	8. 9. S. Ȑ.	
4 adscéndens. R.S.	(re.) ascending.	Guinea.	1824.	6. 9. S. Ȑ.	
5 hirsúta. R.S.	(sc.) scarlet-trailing.	Jamaica.	1733.	5. 8. S. Ȑ.	Jacq. vind. 1. t. 7.
6 plumbaginea. R.S. (ro.)	Plumbago-leaved.	Spain.	1822.	—	Cavan. ic. 2. t. 112.
7 discolor. K.S.	(bh.) two-coloured.	Guayaquil.	1825.	—	
8 viscosa. K.S.	(sc.) viscous.	—	—	S. Ȑ.	
9 scándens. R.S.	(gr.) climbing.	Jamaica.	1691.	4. 9. S. Ȑ.	Jacq. vind. 1. t. 4.
10 excélsa. R.S.	(pu.) lofty.	S. America.	1820.	—	
PIS'ONIA. L.	PIS'ONIA. Heptandria Monogynia. L.				
1 grándis. B.P.	(wh.) superb.	N. Holland.	1805.	—	G. Ȑ.
2 hirtélla. K.S.	(wh.) hairy.	Mexico.	1825.	—	G. Ȑ.
3 frágans. s.s.	(gr.) fragrant.	—	—	S. Ȑ.	
4 macrophylla. L.en. (w.) long-leaved.	—	—	1820.	—	S. Ȑ.
5 obováta. L.en.	(gr.) obovate-leaved.	S. America.	—	—	S. Ȑ.
6 nígricans. s.s.	(wh.) black.	W. Indies.	1806.	5. 7. S. Ȑ.	
7 obtusáta. s.s.	(gr.) blunt-leaved.	—	1822.	—	S. Ȑ.
8 mexicána. L.en. (ro.)	Mexican.	Mexico.	1825.	6. 8. H. Ȑ.	
	Boerhaavia arboréscens. P.s.				
9 nítida. W.en.	(li.) glossy.	Madagascar.	1824.	—	S. Ȑ.
10 aculeáta. s.s.	(ye.) prickly.	Jamaica.	1739.	3. 4. S. Ȑ.	Lam. ill. t. 861.
BOLD'OA. S.S.	BOLD'OA. Triandria Monogynia. S.S.				
lanceolátæ. s.s.	(re.) lance-leaved.	Mexico.	1824.	5. 7. G. Ȑ.	H. et B. pl. æq. 1. t. 44.
Salpiánthus arenárius. K.S.					

ORDO CLXV.

AMARANTHACEÆ. Brown prodr. p. 413.

AMARA'NTHUS. L.	AMARANTH. Monœcia Tri-Pentandria. L.				
1 tenuifolius. w.	(gr.) fine-leaved.	E. Indies.	1801.	7. 9. H. ⊙.	
2 angustifolius. w.	(gr.) narrow-leaved.	Levant.	1723.	—	H. ⊙.
3 pálidus. s.s.	(pa.) pale.	Tauria.	1823.	—	H. ⊙.
4 persicarioïdes. s.s. (re.)	Persicaria-like.	Nepaul.	1818.	9. 11. H. ⊙.	
5 Acroglochin. s.s. (gr.)	spiny-bracted.	—	1820.	—	H. ⊙.
6 álbis. w.	(wh.) white.	N. America.	1778.	7. 9. H. ⊙.	
7 græcizans. w.	(gr.) Pellitory-leaved.	—	1759.	—	H. ⊙.
8 polygamus. w.	(gr.) hermaphrodite.	E. Indies.	1780.	—	H. ⊙.
9 Blitum. E.B.	(gr.) wild.	England.	—	6. 8. H. ⊙.	
10 víridis. w.	(gr.) green.	Brazil.	1768.	8. 9. H. ⊙.	
11 campéstris. w.	(gr.) field.	E. Indies.	1818.	—	H. ⊙.
12 polygonoides. w.	(gr.) spotted-leaved.	Jamaica.	1778.	7. 8. H. ⊙.	
13 melanchólico. w.	(pu.) melancholy.	E. Indies.	1731.	6. 9. H. ⊙.	
14 tricolor. w.	(st.) three-coloured.	—	1548.	—	G. ⊙.
15 bicolor. w.	(st.) two-coloured.	—	1802.	7. 9. H. ⊙.	
16 lineáetus. B.P.	(gr.) line-leaved.	N. Holland.	1828.	—	H. ⊙.
17 gangéticus. w.	(gr.) oval-spiked.	E. Indies.	1778.	—	H. ⊙.
18 mangostanus. w.	(gr.) rhomb-leaved.	—	1801.	—	H. ⊙.

W. amar. 9. t. 1. f. 2.
— 8. t. 4. f. 7.
Rumph. amb. 5. t. 82.
Eng. bot. v. 31. t. 2212.
W. amar. 18. t. 8. f. 16.

W. amar. 11. t. 6. f. 12.
— 15. t. 9. f. 18.
Knor. thes. 2. t. A. 3. 5.

W. amar. 16. t. 6. f. 11.
— 13. t. 12.

19	polystachys. w. (gr.) many-spiked.	E. Indies.	1816.	7. 9. H. ⓧ.	
20	tristis. w. (pu.) round-headed.	China.	1759.	6. 8. H. ⓧ.	W. amar. 21. t. 5. f. 10.
21	inamoenus. w. (gr.) inelegant.	—	1816.	— H. ⓧ.	
22	incómpetus. W.en. (gr.) slovenly.	—	—	— H. ⓧ.	
23	lividus. w. (br.) livid.	N. America.	1759.	7. 9. H. ⓧ.	W. amar. 20. t. 1. f. 1.
24	oleraceus. w. (li.) eatable.	E. Indies.	1764.	7. 8. H. ⓧ.	— 17. t. 5. f. 9.
25	lanceæfolius. H.B. (gr.) lance-leaved.	—	1816.	8. 10. H. ⓧ.	
26	atropurpureus. H.B. (pu.) dark-purple.	—	1821.	— H. ⓧ.	
27	gracilis. s.s. (gr.) oval-leaved.	Guinea.	1806.	7. 8. H. ⓧ.	Jacq. ic. 2. t. 344.
	<i>Chenopodium caudatum.</i> Jacq.				
28	prostratus. w. (gr.) trailing.	France.	1739.	7. 9. H. ⓧ.	
29	scandens. w. (gr.) climbing.	S. America.	1796.	7. 8. S. ⓧ.	
30	deflexus. w. (gr.) bending.	—	1805.	— H. ⓧ.	
31	fasciatus. H.B. (gr.) fasciate.	E. Indies.	1820.	8. 10. H. ⓧ.	W. amar. 10. t. 10. f. 20.
32	bullatus. s.s. (gr.) blistered.	—	—	7. 9. H. ⓧ.	
33	Arardhanus. (gr.) Indian.	E. Indies.	1825.	— G. ⓧ.	
34	Berchtholdi. H.(pi.gr.) Berchthold's.	S. Europe.	1826.	— H. ⓧ.	
35	bahiensis. Lk. (gr.) Bahia.	Bahia.	1823.	— H. ⓧ.	
36	purpurascens. Ot. (pu.) purplish.	—	1828.	— H. ⓧ.	
37	littoralis. Ot. (gr.) sea-side.	—	1823.	— H. ⓧ.	
38	rigidus. s. (gr.) rigid.	—	—	— H. ⓧ.	
39	tortuosus. H.H. (gr.) twisted.	—	—	— H. ⓧ.	
40	laxifolius. Ot. (gr.) loose-leaved.	—	—	— H. ⓧ.	
41	glomeratus. Ot. (gr.) glomerate.	—	1829.	— H. ⓧ.	
42	spicatus. P.S. (gr.) spiked.	France.	—	— H. ⓧ.	
43	pumilus. s.s. (gr.) dwarf.	N. America.	—	6. 9. H. ⓧ.	
44	curvifolius. s.s. (gr.) curved-leaved.	Caucasus.	1827.	— H. ⓧ.	
45	hydrus. w. (re.) clustered.	—	1656.	— H. ⓧ.	W. amar. 26. t. 9. f. 17.
46	hécticus. w. (pi.) oval-spiked.	—	1796.	8. 9. H. ⓧ.	— 25. t. 7. f. 13.
47	lætus. w. (re.) blunt-leaved.	Italy.	1799.	7. 9. H. ⓧ.	— 28. t. 8. f. 15.
48	strictus. w. (gr.) upright.	—	1793.	— H. ⓧ.	— 27. t. 3. f. 5.
49	paniculatus. w. (re.) paniced.	N. America.	1798.	— H. ⓧ.	— 32. t. 2. f. 4.
50	celosioides. K.S. (wh.) Spinach-like.	S. America.	1824.	9. 11. H. ⓧ.	
51	caracasanus. K.S. (wh.) Caracas.	Caracas.	1821.	— H. ⓧ.	
52	retroflexus. w. (gr.) hairy.	N. America.	1759.	7. 9. H. ⓧ.	W. amar. 33. t. 11. f. 21.
53	flavus. w. (st.) pale yellow.	India.	—	— H. ⓧ.	— 35. t. 3. f. 6.
54	chlorostachys. w. (ye.) nodding.	—	1796.	— H. ⓧ.	— 34. t. 10. f. 19.
55	parisenis. s.s. (gr.) Paris.	France.	1819.	— H. ⓧ.	
56	hypochondriacus. w. (re.) Prince's feather.	Virginia.	1684.	— H. ⓧ.	
57	sanguineus. w. (re.) spreading.	Bahama Isl.	1775.	— H. ⓧ.	W. amar. 31. t. 2. f. 3.
58	cruentus. w. (cr.) various-leaved.	China.	1728.	6. 8. H. ⓧ.	
59	speciosus. D.P. (sc.) showy.	Nepaul.	1819.	9. 11. H. ⓧ.	Bot. mag. t. 2227.
60	caudatus. w. (re.) Love-lies bleeding.	E. Indies.	1596.	8. 9. H. ⓧ.	
β maximus.	(re.) tree.	—	—	— H. ⓧ.	
61	spinosus. w. (gr.) prickly.	W. Indies.	1683.	7. 9. H. ⓧ.	W. amar. 38. t. 4. f. 8.
62	frumentaceus. H.B. (gr.) cultivated.	E. Indies.	1819.	— H. ⓧ.	
LECANOCA'RPUS. N.E. LECANOCA'RPUS.	Monocotyledonous.	Monocot. Pentandria.			
nepalensis. N.E. (gr.) Nepaul.	Nepaul.	Nepaul.	1816.	8. 10. H. ⓧ.	Nees ab Ess. ic. sel. t. 2.
Amaranthus caulinodus. Lk.					

CEL'OSIA. R.S.	COCKS-COMB.	Pentandria Monogynia. L.			
1	argentea. R.S. (bh.) silvery-spiked.	China.	1714.	6. 9. S. ⓧ.	Mart. cent. t. 7.
β linearis.	(bh.) narrow-leaved.	E. Indies.	—	— S. ⓧ.	Rheed. mal. 10. t. 39.
2	pyramidalis. R.S. (wh.) pyramidal.	—	—	— S. ⓧ.	Burm. ind. 65. t. 25. f. 1.
3	margaritacea. R.S. (wh. re.) pearl-flowered.	India.	—	— S. ⓧ.	Rheed. mal. 10. t. 38.
4	cristata. R.S. (re.) common.	Asia.	1570.	— S. ⓧ.	Rumph. amb. 5. t. 84.
a elata.	(re.) tall red.	—	—	— S. ⓧ.	Knor. del. 1. t. H. 5. 6.
β compacta.	(re.) dwarfed.	—	—	— S. ⓧ.	
γ flavescens.	(st.) yellow.	—	—	— S. ⓧ.	
5	comosa. R.S. (re.) tufted.	E. Indies.	1802.	— S. ⓧ.	
6	coccinea. R.S. (sc.) scarlet.	China.	1597.	— S. ⓧ.	Barrel. rar. t. 1195.
7	castrénis. R.S. (re.) branched.	E. Indies.	1739.	— S. ⓧ.	Andr. reposit. t. 635.
8	cérnua. A.R. (sc.) drooping.	—	1809.	7. 8. S. ⓧ.	Pluk. alm. 11. t. 334.f.4.
9	Monsoniæ. R.S. (bh.) downy.	—	1778.	7. 9. S. ⓧ.	Jacq. vind. 1. t. 98.
	<i>Achyranthes Monsoniæ.</i> s.s. <i>Illécebrum Monsoniæ.</i> L.				
10	nodiflora. R.S. (wh.) knotted.	E. Indies.	1780.	— S. ⓧ.	Jacq. ic. 2. t. 339.
LESTIBUD'ESIA. R.S. LESTIBUD'ESIA.	Pentandria Monogynia. R.S.				
1	virgata. R.S. (gr.) wave-leaved.	W. Indies.	1815.	8. 10. S. h.	
Celosia virgata. Jacq.					

- 2 paniculâta. r.s. (st.) panicle. Jamaica. 1733. 6. 9. S. ♂.
 3 trigyna. r.s. (wh.) oval-leaved. Senegal. 1777. 8. 10. S. ♂.
DEERÍNGIA. R.S. **DEERÍNGIA.** Pentandria Trigynia.
 1 índica. s.s. (wh.) Berry-bearing. E. Indies. 1804. — S. ♂.
 2 celosíoides. B.P. (wh.) Cockscomb-like. N. Holland. 1822. — S. ♀.
CHAMISS'OA. K.S. **CHAMISS'OA.** Pentandria Monogynia. R.S.
 altissima. k.s. (st.) tall. S. America. 1819. 5. 6. S. ♀. H. et B. n. gen. 2. t. 125.
CLADOSTACHYS. D.P. **CLADOSTACHYS.** Pentandria Monogynia.
 1 frutescens. D.P. (gr.) frutescent. E. Indies. 1777. 8. 11. S. ♀.
Achyranthes muricata. L. *Chamissoa muricata.* s.s.
 2 alternifolia. (pu.) alternate-leav'd. E. Indies. 1789. 7. 8. S. ⊙.
Achyranthes alternifolia. w. *Desmodium alternifolia.* DC.
ACHYRA'NTHES. R.S. **ACHYRA'NTHES.** Pentandria Monogynia. L.
 1 argéntea. r.s. (ro.) upright. Sicily. 1713. 5. 10. H. ♀.
 2 áspera. r.s. (gr.) rough. India. 1751. — S. ♀.
 3 obtusifolia. Lam. (gr.) blunt-leaved. — — S. ♀.
 4 crispa. r.s. (gr.) curl-leaved. — 1816. — S. ♀.
 5 virgata. r.s. (gr.) slender. Portorico. 1823. — S. ♀.
 6 fruticosa. r.s. (wh.) shrubby. E. Indies. 1816. — S. ♀.
 7 nivea. r.s. (wh.) white. Canaries. 1780. 5. 7. G. ♀.
 8 pubescens. r.s. (wh.) pubescent. — 1824. — S. ♀.
 9 brachiata. r.s. (wh.) armed. E. Indies. 1829. 8. 10. S. ⊙.
 10 globulifera. f. (wh.) globe-headed. — 1828. — S. ⊙.
DESMOCH'ÆTA. R.S. **DESMOCH'ÆTA.** Pentandria Monogynia. R.S.
 1 atropurpurea. DC. (pu.) dark-purple. E. Indies. 1759. 8. 10. S. ♀.
Achyranthes lappacea. L. *Pupalia lappacea.* J.
 2 flavescens. DC. (st.) yellowish. E. Indies. — — S. ♀.
 3 pátula. r.s. (wh.) spreading. — 1824. — S. ♀.
 4 prostrata. DC. (pu.) prostrate. — 1793. 7. 8. S. ♀.
 5 sanguinolenta. Lk. (gr.) bloody-leaved. — 1824. — S. ♀.
ALTERNANTH'ERA. **ALTERNANTH'ERA.** Pentandria Monogynia. R.S.
 1 sessilis. r.s. (wh.) sessile-flowered. E. Indies. 1778. 7. 10. S. ♂.
 2 ficoides. r.s. (wh.) creeping-stalk'd. S. America. 1819. — S. ♀.
Gomphrena ficoides. Jacq.
 3 spinosa. r.s. (st.) spiny. — — — S. ♀.
axillaris. DC. *Achyranthes spinosa.* Horn.
 4 denticalata. B.P. (wh.) tooth-leaved. N. S. W. 1822. — G. ♀.
 5 tenella. DC. (wh.) slender. E. Indies. 1828. 7. 9. H. ⊙.
 6 Achyrantha. R.S. (br.) creeping. Buenos Ayres. 1732. 6. 8. S. ♀.
BUCHO'LZIA. M.P. **BUCHO'LZIA.** Pentandria Monogynia.
 1 polygonoides. M.P. (w.) Persicaria-leaved. S. America. 1731. — G. ♀.
Illecebrum polygonoides. w. *Alternanthera polygonoides.* R.S.
 2 frutescens. M.P. (wh.) frutescent. Peru. 1822. — S. ♀.
ÆR'UA. J. **ÆR'UA.** Pentandria Monogynia. R.S.
 1 lanata. r.s. (wh.) woolly. E. Indies. 1691. 4. 8. S. ♂.
Illecebrum lanatum. L.
 2 javanica. r.s. (wh.) spear-leaved. — 1786. — S. ♀.
PHILO'XERUS. B.P. **PHILO'XERUS.** Pentandria Monogynia. R.S.
 1 vermicularis. R.S. (wh.) creeping. S. America. — — S. ♀.
 2 aggregatus. R.S. (wh.) clustered. Cumana. 1819. — S. ♀.
MOGIPH'ANES. M.P. **MOGIPH'ANES.** Pentandria Monogynia.
 1 brasiliensis. M.P. (wh.) upright. Brazil. 1790. 7. 10. S. ♀.
Gomphrena brasiliensis. Jacq. ic. 2. t. 346. *Philoxerus brasiliensis.* B.P.
 2 straminea. M.P. (st.) straw-coloured. Brazil. 1790. 7. 10. S. ♀.
GOMPHR'ENA. R.S. **GLOBE AMARANTH.** Pentandria Monogynia. L.
 1 globosa. r.s. (re.) annual. India. 1714. 5. 10. S. ⊙.
β alba. (wh.) white-flowered. — — — S. ⊙.
 2 decumbens. R.S. (pu. wh.) decumbent. Mexico. — — S. ⊙.
 3 perennialis. R.S. (st.) perennial. S. America. 1732. 7. 10. S. ♀.
 4 arboriscens. R.S. (st.) tree. — 1802. — S. ♀.
 5 villosa. s.s. (st.) villous. Monte Video. 1826. 6. 8. S. ♀.
 6 nigricans. M.P. (bk.) dark-flowered. Brazil. 1827. — S. ♀.
 7 calliosa. s.s. calious. S. America. 1829. — S. ♀.
 8 carnea. Jac. (fl.) flesh-coloured. — — — S. ♀.
OPLOTH'ECA. N. **OPLOTH'ECA.** Pentandria Monogynia.
 1 lactea. M.P. (wh.) milk-coloured. S. America. 1818. 7. 10. S. ♀.
 2 interrupta. N. (wh.) trailing. W. Indies. 1733. — S. ♂. DC. h. mons. ined. t. 93.
Gomphrena interrupta. R.S. Jacq. icon. 1. t. 51.
 3 floridana. N. (wh.) Florida. Florida. 1823. 8. 10. F. ♀. Bot. mag. t. 2003.

Sloan. hist. 1. t. 91. f. 2.
 Jacq. vind. 3. t. 15.

Rumph. amb. 5. t. 83.

Flor. græc. t. 244.
 Mill. ic. t. 11. f. 2.

Pluk. alm. 2. t. 334. f. 5.

Rheed. mal. 10. t. 59.

DC. h. mons. ined. t. 79.

Rumph. amb. 6. t. 11.
 — — t. 27. f. 2.

Rheed. mal. 10. t. 11.
 Jacq. amer. pict. t. 90.

Mart. pl. bras. 2. t. 152.

Herm. parad. t. 17.

L'Herit. stirp. 1. t. 37.

Mill. ic. 1. t. 11. f. 1.

Burm. ind. t. 65. f. 2.

Herm. parad. t. 15.

Mart. pl. bras. 2. t. 133.

Mart. pl. bras. 2. t. 135.

Bot. mag. t. 2815.

Jacq. schœn. t. 482.

Bot. mag. t. 2614.

Mart. pl. bras. 2. t. 113.

DC. h. mons. ined. t. 93.

Jacq. icon. 1. t. 51.

Bot. mag. t. 2003.

BRAND'ESIA. M.P. **BRAND'ESIA.** Pentandria Monogynia.

1 pubérola. M.P. (*wh.*) pubescent. Brazil. 1825. 6. 9. S. $\frac{1}{2}$.

Achyranth s. c. ituliflora. Colla hort. rip. t. 18.

2 villosa. M.P. (*wh.*) villous. Brazil. 1826. — S. \odot .

3 pórrigens. M.P. (*re.*) crimson-flowered. Peru. 1802. 4. 8. G. $\frac{1}{2}$.

Achyranthes pórrigens. B.M.

PFAFFIA. M.P. **PFAFFIA.** Pentandria Monogynia.

gnaphaloïdes. M.P. (*w.*) Cudweed-like. Brazil. 1822. 6. 8. S. $\frac{1}{2}$.

Celosia gnaphaloïdes. L.

IRES'INE. S.S. **IRES'INE.** Diœcia Pentandria. L.

1 celosioïdes. s.s. (*wh.*) Florida. America. 1733. 7. 8. F. $\frac{1}{2}$.

2 diffusa. s.s. (*wh.*) spreading. S. America. 1812. 7. 9. S. $\frac{1}{2}$.

3 elongata. s.s. (*wh.*) long-stalked. — 1822. — S. $\frac{1}{2}$.

R'OSEA. M.P. **R'OSEA.** Polygamia Monœcia.

elatior. M.P. (*st.*) tall. S. America. 1820. 8. 10. S. \odot .

Iresine elatior. W.

Mart. bras. 2. t. 128.

Bot. mag. t. 830.

IRES'INE. S.S. **IRES'INE.** Diœcia Pentandria. L.

1 celosioïdes. s.s. (*wh.*) Florida. America. 1733. 7. 8. F. $\frac{1}{2}$.

2 diffusa. s.s. (*wh.*) spreading. S. America. 1812. 7. 9. S. $\frac{1}{2}$.

3 elongata. s.s. (*wh.*) long-stalked. — 1822. — S. $\frac{1}{2}$.

R'OSEA. M.P. **R'OSEA.** Polygamia Monœcia.

elatior. M.P. (*st.*) tall. S. America. 1820. 8. 10. S. \odot .

Sloan. hist. 1. t. 90.

ORDO CLXVI.

PHYTOLACEÆ. Brown.

PHYTOLA'CCA. W. **PHYTOLA'CCA.** Decandria Decagynia. L.

1 octandra. w. (*wh.*) white-flowered. Mexico. 1732. 7. 11. G. $\frac{1}{2}$.

2 abyssinica. w. (*wh.*) African. Africa. 1775. 5. 6. S. $\frac{1}{2}$.

3 heptandra. s.s. (*wh.*) 7-anthered. 1819. 7. 10. G. $\frac{1}{2}$.

4 bogotensis. k.s. (*wh.*) variable anther'd. S. America. 1823. — G. $\frac{1}{2}$.

5 decandra. w. (*bh.*) Virginian poke. Virginia. 1615. 8. 9. H. $\frac{1}{2}$.

6 dodecandra. w. (*bh.*) recurved-leaved. 5. 6. S. $\frac{1}{2}$.

7 mexicana. (*wh.*) Mexican. Mexico. 1824. 7. 10. G. $\frac{1}{2}$.

icosandra. B.M. non w.

8 icosandra. w. (*wh.*) red-stemmed. E. Indies. 1758. 7. 11. S. $\frac{1}{2}$.

9 dioica. w. (*wh.*) tree. S. America. 1768. S. $\frac{1}{2}$.

RIVINA. R.S. **RIVINA.** Tetrandria Monogynia. L.

1 humiliis. r.s. (*wh.*) downy. W. Indies. 1699. 1. 10. S. $\frac{1}{2}$.

2 lœvis. r.s. (*bh.*) smooth. — 1733. 2. 9. S. $\frac{1}{2}$.

3 latifolia. r.s. (*pu.*) broad-leaved. Madagascar. 1826. 7. 9. S. \odot .

4 brasiliensis. r.s. (*gr.*) wave-leaved. Brazil. 1790. 6. 7. S. $\frac{1}{2}$.

5 purpurascens. r.s. (*pu.*) purple. W. Indies. 1815. 5. 8. S. $\frac{1}{2}$.

6 lanceolata. L.en. (*wh.*) spear-leaved. Brazil. — — — S. $\frac{1}{2}$.

7 octandra. r.s. (*wh.*) climbing. W. Indies. 1752. 5. 6. S. $\frac{1}{2}$. \odot . Plum. ic. 241.

MICR'OTEA. K.S. **MICR'OTEA.** Pentandria Dignynia. R.S.

débiles. k.s. (*wh.*) slender. S. America. 1824. 7. 9. S. \odot .

ANCISTROCA'RPU'S. K.S. **ANCISTROCA'RPU'S.** Octandria Tetragynia.

mayapurensis. k.s. (*wh.*) branching. S. America. 1822. 5. 6. H. \odot .

PETIV'ERIA. W. **PETIV'ERIA.** Heptandria Monogynia. L.

1 alliacea. w. (*wh.*) Garlic-scented. Jamaica. 1759. 6. 7. S. $\frac{1}{2}$.

2 octandra. w. (*wh.*) dwarf. W. Indies. 1737. — S. $\frac{1}{2}$.

GIS'EKIA. W. **GIS'EKIA.** Pentandria Pentagynia. L.

pharnacioïdes. w. (*wh.*) trailing. E. Indies. 1783. 6. S. \odot .

Dill. elt. t. 239. f. 308.

Hoff. c. get. t. 2.

Bot. mag. t. 931.

Bot. mag. t. 2633.

Mill. ic. t. 207.

L'Herit. st. nov. t. 70.

Bot. mag. t. 1781.

Lam. ill. t. 81. f. 2.

Schrad. gen. nov. t. 5.

Plum. ic. 241.

H. et B. pl. am. 2.t.122.

Lodd. bot. cab. t. 148.

Plum. ic. 213. t. 219.

Roxb. corom. 2. t. 183.

ORDO CLXVII.

CHENOPODEÆ. Brown prodr. p. 405.

BASELLA. R.S.

MALABAR-NIGHTSHADE. Pentandria Trigynia. L.

1 rubra. r.s. (*re.*) red-flowered. E. Indies. 1731. 7. 9. S. δ . \sim . Rheed. mal. 7. t. 24.

2 alba. r.s. (*wh.*) white-flowered. — 1688. 7. 11. S. δ . \sim . Pluk. alm. t. 63. f. 1.

3 lúcida. r.s. (*wh.*) shining. — 1802. — S. \odot . \sim .

4 cordifolia. r.s. (*re.*) heart-leaved. — — S. \odot . \sim .

5 tuberosa. r.s. (*st.*) tuberous-rooted. S. America. 1824. — S. $\frac{1}{2}$. \sim .

2 F

CHENOPODEÆ.

6 marginata. r.s. (wh.) red-margined.	S. America. 1824.	7. 11. S. 4. —.
7 japonica. r.s. (bh.) dwarf.	China. 1814.	— G. (○). Burm. Ind. t. 39. f. 4.
HABLI'TZIA. M.B. HABLI'TZIA.	Pentandria Monogynia.	
tamnoïdes. m.b. (wh.) Tamus-like.	Caucasus. 1828.	7. 10. H. 4. —.
ANRED'ERA. J. ANRED'ERA.	Pentandria Trigynia.	
spicata. k.s. (bh.) spiky.	Cuba. 1749.	7. 9. S. 4. —. Sloan. hist. 1. t. 90. f. 1.
Basella vesicaria. Lam. ill. t. 215. f. 1. Polygonum scandens. w.		
ANABA'SIS. L. ANABA'SIS.	Pentandria Digynia. L.	
1 aphylla. r.s. (gr.) leafless.	Levant. 1823.	5. 7. F. (h.).
2 florida. m.b. (wh.) flowering.	Persia. 1829.	7. 11. H. (○).
3 cretacea. r.s. (gr.) long-rooted.	Siberia. 1828.	6. 8. H. 4.
4 spinosissima. r.s. (gr.) spiny-branched.	Iberia. 1829.	— H. (h.).
5 tamariscifolia. r.s. (gr.) Tamarisk-leaved.	Spain. 1752.	6. 7. F. (h.).
6 foliosa. s.s. (gr.) leafy.	Volga. 1828.	7. 10. H. (○).
Salsola clavifolia. M.B. bacifera. Pall.		
SALSOLA. R.S. SALTWORT. Pentandria Digynia. L.		
1 Kali. r.s. (fl.) prickly.	Britain.	7. 8. H. (○).
2 Tragus. r.s. (ro.) smooth.	S. Europe. 1816.	— H. (○).
3 tamariscina. r.s. (br.) Tamarisk-like.	Tauria. 1828.	7. 9. H. (○).
4 crassa. r.s. (re.wh.) thick-leaved.	Caucasus. 1823.	8. 10. H. (○).
5 rosacea. r.s. (pu.) rose-coloured.	Asia. 1759.	7. 8. H. (○).
6 oppositifolia. r.s. (ro.) opposite-leaved.	Sicily. 1829. H. (h.).
7 brachiata. r.s. (pu.) armed.	Tauria. 1828.	7. 10. H. (○).
8 glauca. r.s. (ro.) glaucous.	Armenia. 1829. H. (h.).
9 vermiculata. r.s. (gr.) suffruticose.	Siberia. 1759.	7. 8. H. (h.).
10 microphylla. r.s. (re.) small-leaved.	Spain. —	— H. (○).
11 spissa. r.s. (ro.) dwarf.	Caucasus. 1829.	7. 10. H. (○).
12 verrucosa. r.s. (gr.) warted.	—	1822. 7. 8. H. (h.).
13 ericoides. r.s. (br.) Heath-like.	—	1829. H. (h.).
14 laniflora. r.s. (re.) woolly-flowered.	Siberia. 1797.	6. 8. H. (○).
15 sativa. r.s. (pi.) cultivated.	Spain. 1783.	7. 8. H. 4.
16 obtusifolia. Led. (br.) blunt-leaved.	Siberia. 1827. H. (○).
17 marginata. F. (gr.) thick margined.	Armenia. 1829. H. (h.)?
18 Sorda. r.s. (wh.) long fleshy-leav'd.	S. Europe. 1683.	7. 8. H. (○).
K'OCHIA. R.S. K'OCHIA. Pentandria Digynia. R.S.		
1 arenaria. r.s. (wh.) sand.	Hungary. 1819.	7. 9. H. (○).
2 dasyantha. r.s. (wh.) woolly-flowered.	Caucasus. 1829.	— H. (○).
3 prostrata. r.s. (gr.) trailing.	S. Europe. 1780.	6. 8. F. (h.).
Salsola prostrata. Jacq.		
4 trigyna. L.en. (gr.) slender-leaved.	Spain. 1804.	7. 8. H. (○).
Salsola altissima. Cav. Chenopodium trigynum. R.S.		
5 hyssopifolia. r.s. (gr.re.) Hyssop-leaved.	Siberia. 1801.	6. 8. H. (○).
6 sedoides. r.s. (gr.) Stonecrop-like.	Tauria. 1823.	— H. 4.
7 eriophora. r.s. (gr.) woolly.	Spain. 1816.	— H. (○).
8 scoparia. r.s. (gr.) Summer Cypress.	Greece. 1629.	6. 9. H. 4.
Chenopodium Scoparia. w.		
CHEN'OLEA. L. CHEN'OLEA. Pentandria Monogynia. L.		
diffusa. w. (wh.) silky.	C. B. S. 1758.	8. 9. G. (h.).
Salsola sericea. H.K. Kochia sericea. r.s. Chenopodium sericeum. s.s.		
ENCHYL'ENA. B.P. ENCHYL'ENA. Pentandria Di-Trigynia.		
tomentosa. b.p. (wh.) woolly.	N. Holland. 1824.	6. 8. G. (h.).
RHAG'ODIA. B.P. RHAG'ODIA. Polygamia Monococcia.		
1 Billardieri. b.p. (gr.) narrow-leaved.	N. S. W. 1822.	6. 7. G. (h.).
2 hastata. b.p. (gr.) halberd-leaved.	— 1803.	— G. (h.).
3 nutans. b.p. (gr.) nodding-branch'd.	V.Diem. Isl. 1824.	7. 9. G. 4.
ACROGLO'CHIN. Schr. ACROGLO'CHIN. Pentandria Digynia.		
chenopodioides. f. (gr.) Goose-foot-like.	Siberia. 1823.	5. 9. H. (○).
CHENOP'DIUM. R.S. GOOSE-FOOT. Pentandria Digynia. L.		
1 Bonus-Henricus. (gr.) English Mercury.	Britain.	5. 8. H. 4.
2 urbiculum. e.b. (gr.) upright.	—	8. H. (○).
3 Atriplicis. r.s. (cr.) purple.	China. 1780.	8. 9. H. (○).
4 punctulatum. r.s. (re.) white-spotted.	— 1820.	— H. (○).
5 giganteum. d.p. (gr.) gigantic.	Nepaul.	9. 11. H. (○).
6 rubrum. r.s. (fl.) red.	Britain. 8. 9. H. (○).
7 chrysomelanospurum. r.s. black-seeded.	Galicia. 1825.	— H. (○).
8 carthagensem. r.s. (gr.re.) Carthage.	Carthage. 1824.	6. 8. H. (○).
9 rhombifolium. r.s. (gr.) angular-leaved.	N. America. 1807.	7. 9. H. (○).
10 guineense. w. (gr.) Guinea.	Guinea. 1790.	8. 9. H. (○).
11 murale. e.b. (gr.) nettle-leaved.	Britain. — H. (○).
Jacq. ic. 2. t. 345.		
Eng. bot. v. 24. t. 1722.		
Eng. bot. v. 15. t. 1033.		
v. 10. t. 717.		
Jacq. vind. 3. t. 80.		
Scop. del. ins. 1. t. 11.		
Eng. bot. v. 24. t. 1721.		
Loud. enc. pl. f. 14270.		
Lab. nov. hol. 1. t. 96.		
Loud. enc. pl. f. 14270.		

12	Quinòa. R.S.	(gr.) green Quinoa.	Chile.	1822.	7. 9. H.○.	Feuil. peruv. t. 10.
	β rùbrum.	(re.) red Quinoa.		—	— H.○.	Feuil. per. ed. germ. t. 10.
13	serotinum. R.S.	(gr.) late-flowering.	S. Europe.	8. 9. H.○.	
14	ficifolium. E.B.	(gr.) Fig-leaved.	England.	— H.○.	Eng. bot. v. 24. t. 1724.
15	álbum. E.B.	(gr.) white.	Britain.	6. 9. H.○.	— v. 24. t. 1723.
	β subrotundum.	(gr.) round-leaved.	—	— H.○.	Petiv. h. brit. t. 8. f. 4.
	γ víride. w.	(gr.) green.	—	— H.○.	
	ð integrifolium.	(gr.) entire-leaved.	—	— H.○.	
	ε crassifolium.	(gr.) thick-leaved.	—	— H.○.	
16	opulifolium. R.S.	(gr.) Guelder-rose-ld.	S. Europe.	— H.○.	Bast. j. bot. 1814. t. 3.
17	hy'bridum. E.B.	(gr.) Maple-leaved.	Britain.	8. 9. H.○.	Eng. bot. v. 27. t. 1919.
18	blítoides. R.S.	(gr.) Blitum-like.	France.	1827.	— H.○.	Flor. græc. t. 253.
19	Bòtrys. R.S.	(gr.) cut-leaved.	S. Europe.	1548.	6. 9. H.○.	
20	f'etidum. W.en.	(gr.) strong-scented.	1823.	6. 8. H.○.	Dill. elt. t. 66. f. 77.
21	múltifidum. R.S.	(gr.) multifid.	BuenosAyres.	1732.	6. 10. G. ¼.	Eng. bot. v. 32. t. 2247.
22	botryoïdes. E.B.	(re.) many-many-clustered.	Britain.	—	6. 8. H.○.	Moris. s. 5. t. 35. f. 8.
23	ambrosioïdes. R.S.	(gr.) Mexican.	Mexico.	1640.	6. 10. H.○.	
24	suffruticósum. R.S.	(gr.) suffrutescent.	—	1819.	— G. ½.	Dill. elt. t. 66. f. 76.
25	anthelmínticum. w.(gr.)	American.	America.	1732.	7. 8. G. ½.	
26	gravéolens. R.S.	(gr.) strong-scented.	Mexico.	1819.	8. 11. H.○.	
27	incisum. R.S.	(gr.) jagged.	—	1826.	— H.○.	
28	glaúcum. R.S.	(gr.) Oak-leaved.	England.	7. 8. H.○.	Eng. bot. v. 21. t. 1454.
29	crassifolium. R.S.	(cr.) fleshy-leaved.	—	1820.	— H.○.	
30	humifúsrum. R.S.	(pu.) trailing.	—	1824.	— H.○.	
31	sépium. R.S.	(re.) hedge.	Bohemia.	1822.	— H.○.	
32	radiatum. R.S.	(gr.) rayed.	N. America.	1803.	6. 8. H.○.	W. hort. ber. 1. t. 28.
	Kochia dentata. w.	Salsola platyphylla. M.				
33	ólidum. E.B.	(gr.) stinking.	Britain.	7. 8. H.○.	Eng. bot. v. 15. t. 1034.
	f'etidum. R.S.	Vulvária. w.				
34	polyspermum. R.S.(gr.)	Allseed.	—	— H.○.	Eng. bot. v. 21. t. 1480.
35	acutifolium. E.B.	(gr.) acute-leaved.	—	— H.○.	— v. 21. t. 1481.
36	marginátum. R.S.	(gr.) bony-edged.	—	1824.	— H.○.	
37	cáudatum. R.S.	(gr.) long-tailed.	Guinea.	1806.	— G. ½.	Jacq. ic. 2. t. 341.
38	laterale. R.S.	(gr.) oblong-leaved.	—	1781.	8. 9. S.○.	
39	strictum. R.S.	(gr.) upright.	E. Indies.	—	9. 11. H.○.	Gmel. sib. 3. t. 15. f. 1.
40	aristatum. R.S.	(gr.) bearded.	Siberia.	1771.	6. 9. H.○.	
41	lanceolatum. R.S.	(gr.) spear-leaved.	Pensylvania.	1809.	7. H.○.	
42	fruticósum. R.S.	(gr.) shrubby.	England.	8. 9. H. ½.	Eng. bot. v. 9. t. 635.
	Salsola fruticosa. E.B.					
43	parvifolium. R.S.	(gr.) small-leaved.	Caucasus.	1829.	— H. ½.	Pall. ill. 3. t. 44.
44	altissimum. R.S.	(wh.) Grass-leaved.	Russia.	1775.	7. 8. H.○.	Schrad. d. hal. t. 1. f. 3
	Salsola altissima. w.	Suaeda altissima.	Pall. ill. t. 42.			
45	trig'ynum. R.S.	(gr.) trigynous.	Spain.	—	— H.○.	Cavan. icon. 3. t. 289.
46	hirsutum. R.S.	(gr.) hairy.	Denmark.	1791.	— H.○.	Flor. dan. t. 187.
	Salsola hirsuta. F.D.					
47	sálsum. R.S.	(gr.) Saltwort.	Astracan.	1782.	8. 9. H.○.	Jacq. vind. 3. t. 83.
48	spicatum. R.S.	(gr.) spiked.	Spain.	1816.	8. 10. H.○.	Cavan. ic. 3. t. 290.
49	setigerum. R.S.	(gr.) setiferous.	—	—	— H.○.	DC. h. mons. ined. t. 87.
50	maritimum. R.S.	(gr.) sea blite.	Britain.	8. H.○.	Eng. bot. v. 9. t. 633.
A	TRIPLEX. R.S.	ORACHE.	Polygamia Monœcia. L.			
1	Hálimus. R.S.	(wh.) tall shrubby.	Spain.	1640.	7. 8. H. ½.	Park. theat. 724. f. 2.
2	portulacoïdes. R.S.	(gr.) dwarf shrubby.	Britain.	— H. ½.	Eng. bot. v. 4. t. 261.
3	verruciferum. R.S.	(gr.) warted-fruited.	Caucasus.	1827.	— H. ½.	
4	álbicans. R.S.	(bh.) white.	C. B. S.	1774.	6. 7. G. ½.	
5	glaúcum. w.	(gr.) glaucous.	S. Europe.	1732.	7. 8. F. ½.	Dill. elt. t. 40. f. 46.
6	linifolium. K.S.	(gr.) Flax-leaved.	S. America.	1819.	— G. ½.	
7	polygamum. R.S.	(gr.) polygamous.	New Spain.	1826.	— F. ½.	
8	coriaceum. R.S.	(gr.) leathery.	Egypt.	1828.	6. 9. F. ½.	
9	ròseum. R.S.	(gr.) rose-like.	S. Europe.	1739.	6. 7. H.○.	Schk. hand. 3. t. 350.
10	foliòsum. R.S.	(gr.) leafy.	Portugal.	1824.	— H.○.	
11	sibiricum. R.S.	(wh.) Siberian.	Siberia.	1783.	7. 8. H.○.	Schk. hand. 3. t. 350.
12	tatáricum. R.S.	(gr.) Tartarian.	Tartary.	1778.	— H.○.	— 3. p. 539. t. 349.
13	bengálense. R.S.	(gr.) Bengal.	E. Indies.	—	— H.○.	
14	hortense. R.S.	(gr.) garden.	Tartary.	1548.	— H.○.	Blackw. t. 99 et 552.
	β intermédia.	(re.) intermediate.	—	—	— H.○.	
	γ rùbrum.	(re.) red garden.	—	—	— H.○.	
15	nítens. R.S.	(gr.) glossy.	Hungary.	1746.	— H.○.	Schkahr hand. t. 348.
16	acuminátum. R.S.	(gr.) acuminate.	—	1820.	— H.○.	W. et K. hung. t. 103.
17	vénatum. R.S.	(gr.) Venetian.	Venice.	—	— H.○.	

CHENOPODEÆ.

18 incànum. R.S.	(gr.) hoary-leaved.	Spain.	1818.	7. 8. H. ⊙.
19 ábum. R.S.	(gr.) white-leaved.	Italy.	1826.	— H. ⊙.
20 incisum. M.B.	(gr.) cut-leaved.	Tauria.	1829.	— H. ⊙.
21 laciñatum. R.S.	(gr.) frosted sea.	Britain.	...	— H. ⊙.
22 Besseriànum. R.S.	(gr.) Besser's.	Hungary.	1825.	— H. ⊙.
23 hastatum. R.S.	(gr.) halberd-leaved.	Europe.	...	— H. ⊙.
24 prostratum. DC.	(gr.) prostrate.		...	— H. ⊙.
25 obtusum. R.S.	(gr.) blunt-leaved.	Siberia.	1825.	— H. ⊙.
26 microspérnum. R.S.	(w.) small-seeded.	Hungary.	1800.	— H. ⊙.
27 oblongifolium. R.S.	(gr.) oblong-leaved.		—	— H. ⊙.
28 pátulum. R.S.	(gr.) spreading.	Britain.	...	6. 9. H. ⊙.
29 angustifolium. R.S.	(gr.) narrow-leaved.		—	6. 8. H. ⊙.
30 eréctum. R.S.	(gr.) upright.	England.	...	8. H. ⊙.
31 littorale. E.B.	(gr.) grass-leaved.	Britain.	...	8. 9. H. ⊙.
32 pedunculatum. R.S.	(gr.) pedunculated.	England.	...	7. 9. H. ⊙.
33 oppositifolium. R.S.	(gr.) opposite-leaved.	France.	1827.	— H. ⊙.
34 verticillatum. R.S.	(gr.) whorl-leaved.	Spain.	1822.	— H. ⊙.
35 sulcatum. R.S.	(gr.) sulcate.		—	— H. ⊙.

B'ETA. R.S. BEET. Pentandria Digynia. L.

1 vulgáris. R.S.	(gr.) common.	S. Europe.	1548.	— H. ♂.
α viridis.	(gr.) green.		—	— H. ♂.
β rúbra.	(gr.) red.		—	— H. ♂.
γ lútea.	(gr.) yellow.		—	— H. ♂.
δ macrorhíza.	(gr.) mangold-wortzel.		—	— H. ♂.
2 Cicla. R.S.	(gr.) white.	Portugal.	1570.	— H. ♂.
3 críspa. R.S.	(gr.) curled-leaved.		—	— H. ♂.
4 trigyna. R.S.	(wh.) Hungarian.	Hungary.	1796.	7. 8. H. ♀.
5 marítima. E.B.	(gr.) sea-side.	Britain.	...	6. 8. H. ♀.
6 macrorhíza. R.S.	(gr.) large-rooted.	Caucasus.	1827.	— H. ♀.
7 foliòsa. E.	(gr.) leafy.	Egypt.	1826.	7. 9. H. ⊙.
8 pátula. R.S.	(gr.) spreading.	Madeira.	1778.	8. 9. G. ♂.

SPINACIA. W. SPINAGE. Dioccia Pentandria. L.

oleracea. w.	(gr.) common.	...	1568.	3. 10. H. ⊙.
α spinosa. w.	(gr.) prickly.	—	—	— H. ⊙.
β inermis.	(gr.) round.	—	—	— H. ⊙.

ACNIDA. W. VIRGINIAN HEMP. Dioccia Pentandria. L.

1 cannabiná. w.	(gr.) common.	N. America.	1640.	6. 7. H. ⊙.
2 ruscocárpa. w.	(gr.) rugged-capsuled.	Virginia.	1824.	— H. ⊙.

B'OSEA. R.S. GOLDEN-ROD. Pentandria Digynia. L.

yervamóra. R.S.	(br.) tree.	Canaries.	1728.	6. 9. G. ♂.
-----------------	-------------	-----------	-------	-------------

CORPR'MUM. L. TICK-SEED. Monandria Digynia. L.

1 hyssopifolium. R.S.	(wh.) Hyssop-leaved.	Europe.	1739.	7. H. ⊙.
2 nítidum. R.S.	(wh.) glossy.	Hungary.	1827.	8. 9. H. ⊙.
3 Redo'wskii. R.S.	(wh.) Redowski's.	Caucasus.	1828.	— H. ⊙.
4 spicatum. R.S.	(wh.) spiked.		—	— H. ⊙.
5 canescens. R.S.	(wh.) canescent.	Hungary.	1822.	7. 9. H. ⊙.
6 intermédia. R.S.	(wh.) intermediate.	Prussia.	—	— H. ⊙.
7 tenué. L.en.	(wh.) slender.		1823.	— H. ⊙.
8 squarrósum. R.S.	(wh.) rough-spiked.	Russia.	1759.	8. 9. H. ⊙.
9 Marschállii. R.S.	(wh.) Marschall's.	Caucasus.	1829.	— H. ⊙.
10 latifolium. Lk.	(wh.) broad-leaved.		1826.	— H. ⊙.
11 sabulósüm. Led.	(wh.) gravel.	Siberia.	1827.	— H. ⊙.

BLITUM. R.S. STRAWBERRY-BLITE. Monandria Digynia. L.

1 capitatum. R.S.	(re.) berry-headed.	Austria.	1633.	5. 8. H. ⊙.
2 virgátum. R.S.	(re.) slender.	S. Europe.	1680.	5. 9. H. ⊙.
3 petiolare. L.en.	(re.) petiolate.		1818.	— H. ⊙.
4 chemopodioïdes. R.S.	(gr.) Goosefoot-like.		1826.	— H. ⊙.
5 marítimum. N.	(gr.) sea-side.	N. America.	1823.	— H. ⊙.

CERATOCAR'PUS. L. CERATOCAR'PUS. Monœcia Monandria. L.

arenarius. L.	(gr.) sand.	Tartary.	1757.	6. 7. H. ⊙.
---------------	-------------	----------	-------	-------------

SALICO'RNIÁ. L. GLASSWORT. Monandria Monogynia. L.

1 herbácea. s.s.	(gr.) marsh.	Britain.	...	8. 9. H. ⊙.
2 procumbens. E.B.	(gr.) procumbent.	England.	...	— H. ⊙.
3 rádicans. E.B.	(gr.) rooting.	Britain.	...	9. H. ♀.
4 pygmæa. R.S.	(gr.) small.	Caucasus.	1827.	— H. ⊙.
5 perennans. R.S.	(gr.) perennial.	Siberia.	1823.	8. 9. H. ♀.
6 fruticosa. E.B.	(gr.) shrubby.	Britain.	—	— H. ♂.
7 arábica. R.S.	(gr.) Arabian.	Arabia.	1758.	— G. ♂.
8 foliòsa. R.S.	(gr.) leafy.	Siberia.	1827.	— H. ♂.

Eng. bot. v. 3. t. 165.

Flor. dan. t. 1286.

W. et K. hung. t. 250.

t. 221.

Eng. bot. v. 13. t. 936.

v. 25. t. 1774.

v. 31. 2223.

v. 10. t. 708.

v. 4. t. 232.

Cavan. H. R. M. t. 21.

Schk. hand. 1. t. 56.

Plenck off. t. 169.

Kern. abbild. t. 235.

Plenck off. t. 170.

W. et K. hung. 1. t. 35.

Eng. bot. v. 4. t. 285.

Schk. hand. 3. t. 324.

Mich. fl. amer. 2. t. 50.

Jacq. f. eclog. t. 25.

Flor. græc. 1. t. 1.

Pall. ros. 2. t. 99.

Poif. et T. fl. p. 1. t. 2.

Bot. mag. t. 276.

Buxb.in. act. pet. 1. t. 9.

Eng. bot. v. 6. t. 415.

v. 35. t. 2475.

v. 24. t. 1691.

Pall. it. 1. ap. T. D.F.I.

Eng. bot. v. 35. t. 2467.

Moris. s. 2. t. 33. f. 7.

HALOCN'EMUM. M.B. HALOCN'EMUM.	Monandria Digynia.				
caspicum. R.S. (gr.) Caspian.	Caucasus.	1827.	7. 8. H. h.	Pall. it. 1. ap. t. A. f. 2.	
POLYCNC'EMUM. R.S. POLYCN'EMUM.	Triandria Monogynia. L.				
1 arvénse. R.S. (wh.) trailing.	S. Europe.	1640.	7. H. ⊖.	Jacq. aust. 4. t. 365.	
2 recúrvum. R.S. (wh.) recurved-leaved.	Switzerland.	1824.	7. 8. H. ⊖.	Pall. ill. t. 50.	
3 sálsum. R.S. (gr.) saltwort.	Tartary.	1828.	— H. ⊖.	— t. 51.	
4 sibíricum. R.S. (wh.) Siberia.	Siberia.	1829.	— H. ⊖.	Pall. ill. ap. T. E. f. 2.	
5 oppositifólium. R.S. (wh.) opposite-leaved.	Tartary.	1820.	— H. ⊖.	Pall. ill. t. 55.	
6 crassifólium. R.S. (wh.) thick-leaved.	Astrakan.	1828.	— H. ⊖.	— t. 56.	
7 sclerospérnum. R.S. (w.) hard-seeded.		1821.	— H. ⊖.		
8 pubéscent. Led. (wh.) pubescent.	Siberia.	1827.	— H. ⊖.		
CAMPHOROSMA. L. CAMPHOROSMA.	Tetrandria Monogynia. L.				
1 monspeliacum. w.(wh.) hairy.	S. Europe.	1640.	8. 9. G. h.	Schk. hand. 1. t. 26.	
peréne. Pall. ill. pl. imp. cogn. t. 57.					
2 ovatum. R.S. (wh.) Hungarian.	Hungary.	1823.	— H. ⊖.	W. et K. hung. 1. t. 63.	
GAL'ENIA. W.	GAL'ENIA. Octandria Digynia. L.				
africana. w. (pa.) African.	C. B. S.	1752.	6. 8. G. h.	Lam. ill. t. 314.	
AX'YRIS. W.	AX'YRIS. Monococcia Triandria. L.				
1 amaranthoides. w. (gr.) simple-spiked.	Siberia.	1758.	6. 7. H. ⊖.	Gmel. sib. t. 2. f. 2.	
2 hy'bida. w. (gr.) bastard.		1780.	6. 8. H. ⊖.	— t. 4. f. 1.	
3 prostrata. w. (gr.) trailing.		1798.	7. 8. H. ⊖.	— t. 4. f. 2.	
DI'OTIS. W.	DI'OTIS. Monococcia Tetrandria. L.				
ceratoídes. w. (wh.) shrubby.	Siberia.	1780.	8. H. h.	Jacq. ic. 1. t. 189.	
<i>Ax'yris ceratoídes.</i> Jacq. <i>Ceratospérnum papposum.</i> P.s.					

ORDO CLXVIII.

BEGONIACEÆ. *Ventenat. et Don prodr.* p. 223.

BEG'ONIA. L.	BEG'ONIA. Monococcia Polyandria. L.				
1 nítida. s.s.	(bh.) shining-leaved.	Jamaica.	1777.	5. 12. S. h.	Sal. par. lond. t. 72.
2 dichótoma. s.s.	(wh.) forked.	Caracas.	1800.	7. 8. S. h.	Jacq. ic. 3. t. 619.
3 renifórmis. s.s.	(wh.) kidney-leaved.	Brazil.	1823.	— S. h.	Linn. trans. t. 14. f. 1.2.
4 spatulátâ. s.s.	(wh.) spatula-leaved.	W. Indies.	1813.	8. 10. S. h.	Lodd. bot. cab. t. 17.
5 odorâta. s.s.	(wh.) sweet-scented.	S. America.	1823.	6. 8. S. h.	
6 sanguínea. s.s.	(pi.) bloody.	Brazil.	1828.	6. 10. S. ȳ.	
7 Sellöii. L.O.	(wh.) Sello's.			— S. h.	
8 undulátâ. B.M.	(wh.) wave-leaved.		1825.	8. 12. S. h.	Bot. mag. t. 2723.
9 semperflorens. B.C. (wh.) ever-blown.		Mexico.	1829.	1. 12. S. h.	Lodd. bot. cab. t. 1439.
10 Hookéri. (pi.) Hooker's.			1827.	— S. h.	Bot. mag. t. 2920.
semperflorens. B.M. non. B.C.					
11 dísticha. s.s.	(wh.) distichous.	S. America.	1820.	6. 8. S. h.	
12 macrophylla. s.s. (bh.) large-leaved.		Jamaica.	1793.	5. 9. S. h.	
13 hirtellâ. s.s.	(wh.) fringed.	W. Indies.	1818.	— S. h.	
14 Evansiâna. A.R.	(pi.) Evans's.	China.	1804.	— S. ȳ.	Bot. mag. t. 1473.
discolor. H.K. published 1813.	<i>Evansiâna. A.R. published 1811.</i>	B.M. 1812.			
15 picta. Sm.	(pi.) painted.	Nepaul.	1823.	5. 9. S. ȳ.	Sm. exot. bot. t. 101.
16 palmâta. D.P.	(pi.) palmated.		1825.	— S. ȳ.	
17 vitifólia. L.O. Vine-leaved.	Brazil.	1829. S. h.	
18 platanifólia. L.O. Plane-leaved.		 S. h.	
19 diversifólia. B.M.	(pi.) various-leaved.	Mexico.	1827.	9. 12. S. h. —	Bot. mag. t. 2966.
20 villòsa. B.R.	(wh.) villous.	Brazil.		8. 10. S. h.	Bot. reg. t. 1252.
21 pauciflòra. B.R.	(bh.) few-flowered.	W. Indies.	1816.	5. 9. S. h.	— t. 471.
22 pátula. H.S.S.	(bh.) spreading.	Brazil.	1811.	— S. h.	
23 bulbifera. L.O.	(pi.) bulb-bearing.		1829.	9. 11. S. ȳ.	
24 monóptera. L.O.	(pi.) one-winged.			— S. ȳ.	
25 tuberosâ. s.s.	(pi.) tuberous.	Amboyna.	1810.	7. 9. S. ȳ.	Rumph. am. 5.t. 169.f.2.
26 acutifólia. s.s.	(wh.) acute-leaved.	W. Indies.	1822.	— S. h.	Sloan. jam. t. 127. f. 1.2.
27 acuminâta. s.s.	(wh.) pointed-leaved.	Jamaica.	1790.	5. 12. S. h.	Bot. reg. t. 364.
28 lúcida. H.R.	(wh.) polished.	Trinidad.	1820.	— S. ⊖.	
29 argyrostigmâ. B.R. (pi.) silver-spotted.		Brazil.	1819.	7. 9. S. h.	Bot. reg. t. 666.
maculâta. s.s.					
30 dipétala. B.M.	(pi.) two-petaled.	E. Indies.	1826.	4. 8. S. h.	Bot. mag. t. 2849.
31 papillôsa. B.M.	(pi.) papillose.	S. America.	1823.	— S. h.	— t. 2846.

32 insignis. B.M.	(pi.) handsome.	Brazil.	1828.	1. 12.	S. h.	Bot. mag. t. 2900.
33 incarnata. L.O.	(pi.) flesh-coloured.	—	1829.	—	S. h.	
34 Martiana. L.O.	(pi.) Martius's.	—	—	—	S. h.	
35 suaveolens. B.C.	(wh.) fragrant.	W. Indies.	1816.	5. 9.	S. h.	Lodd. bot. cab. t. 69.
36 humilis. H.E.F.	(wh.) dwarf.	—	1788.	10.	S. d.	Hook. ex. flor. t. 17.
37 hirsuta. s.s.	(wh.) shaggy-leaved.	—	1789.	5. 6.	S. d.	Aub. gui. 2. t. 348.
38 ulmifolia. s.s.	(bh.) Elm-leaved.	S. America.	1820.	6. 1.	S. h.	Hook. ex. flor. t. 57.
39 scandens. s.s.	(gr.) climbing.	Jamaica.	1824.	7. 9.	S. h. —	Aub. gui. 2. t. 349.

ORDO CLXIX.

POLYGONEÆ. Brown prodr. p. 419.

COCCOL'OBIA. L.	SEASIDE-GRAPE. Octandria Trigynia. L.	W. Indies.	1690.	...	S. h.	Jacq. amer. t. 73.
1 uvifera. s.s.	(wh.) round-leaved.	—	—	...	S. h.	Pluk. phyt. 222. f. 8.
2 pubescens. s.s.	(wh.) downy.	—	—	...	S. h.	Plum. ic. t. 146. f. 1.
3 excoriata. s.s.	(wh.) oval-leaved.	—	1733.	...	S. h.	
4 nivea. s.s.	(st.) snowy.	Jamaica.	1826.	...	S. h.	
5 punctata. s.s.	(wh.) spear-leaved.	W. Indies.	1733.	...	S. h.	Jacq. amer. 114. t. 77.
6 laurifolia. W.en.	(wh.) Laurel-leaved.	S. America.	1816.	8. 9.	S. h.	
7 diversifolia. s.s.	(wh.) different-leaved.	Hispaniola.	—	—	S. h.	Hook. ex. flor. t. 102.
8 obtusifolia. s.s.	(wh.) blunt-leaved.	S. America.	1823.	...	S. h.	Jacq. amer. t. 74.
9 microstachya. s.s.	(wh.) small-spiked.	W. Indies.	—	—	S. h.	
10 barbadensis. s.s.	(wh.) Barbadoes.	Barbadoes.	1790.	...	S. h.	Jacq. obs. 1. t. 8.
11 latifolia. s.s.	(wh.) broad-leaved.	S. America.	1812.	...	S. h.	Lam. ill. t. 316. f. 4.
12 acuminata. K.S.	(wh.) sharp-pointed.	—	1824.	...	S. h.	
13 longifolia. L.en.	(wh.) long-leaved.	—	1818.	...	S. h.	
14 tenuifolia. s.s.	(wh.) small-leaved.	Jamaica.	1824.	...	S. h.	Brown. jam. 210. f. 3.
BRUNNICHIA. S.S.	BRUNNICHIA. Decandria Trigynia. W.	Carolina.	1787.	...	G. h. —	Gært. s. 1. t. 48. f. 2.
cirrhosa. s.s.	(gr.) Carolina.	Carolina.	—	—	—	
POLY'GONUM. L.	PERSICARIA. Tetr-Octandria Di-Trigynia.	Britain.	...	5. 9. H. 4.	Eng. bot. v. 8. t. 509.	
1 Bistorta. w.	(pi.) Snake-weed.	N. America.	1828.	6. 7. H. 4.	Eng. bot. v. 10. t. 669.	
2 bistortoides. Ph.	(bh.) Snakeweed-like.	Britain.	...	5. 9. H. 4.		
3 viviparum. E.B.	(gr.) alpine bistort.	Nepaul.	1823.	—	H. 4.	
4 petiolatum. D.P.	(pi.) petiolate.	—	1822.	—	H. 4.	
5 macrophyllum. D.P.	(pu.) large-leaved.	—	1823.	—	H. 4.	
6 affine. D.P.	(ro.) related.	N. America.	1818.	6. 9. H. ⊖.		
7 mite. P.S.	(li.) tasteless.	—	1824.	—	H. ⊖.	
8 hydropperoïdes. Ph.	(w.) acrid.	Sicily.	1826.	—	H. 4.	
9 tenuiforum. s.s.	(li.) slender-flowered.	N. America.	1749.	7. 9. H. 4. —	Pluk. alm. t. 177. f. 7.	
10 scandens. w.	(pi.) climbing.	England.	—	7. 8. H. ⊖.	Eng. bot. v. 15. t. 1044.	
11 Fagopyrum. w.	(pi.) Buck-wheat.	Nepaul.	1796.	—	H. ⊖.	Bot. reg. t. 1065.
12 emarginatum. w.	(pi.) Indian buck-wheat.	Siberia.	1759.	—	H. ⊖.	Gmel. sib. 3. t. 13. f. 1.
13 tataticum. w.	(gr.) Tartarian.	Britain.	—	—	—	
14 Convolvulus. w.	(bh.) common climbing.	—	5. 9. H. ⊖. —	Eng. bot. v. 14. t. 941.		
15 dumetorum. w.	(li.) bush.	S. Europe.	1803.	—	H. ⊖.	Flor. dan. t. 759.
16 adpressum. B.P.	(pi.) polygamous.	N. S. W.	1822.	—	G. h. —	Lab. n. hol. 1. t. 127.
17 sagittatum. w.	(wh.) arrow-leaved.	N. America.	1759.	7. 8. H. ⊖.	Bart. fl. amer. t. 101.	
18 chinense. w.	(pi.) Chinese.	China.	1795.	—	H. ⊖.	Burm. ind. t. 30. f. 3.
19 pâtens. D.P.	(pi.) spreading.	Nepaul.	1823.	—	H. 4.	
20 élégans. s.s.	(wh.) elegant.	Naples.	1824.	—	H. h.	Bocc. mus. 2. t. 27.
21 alpinum. w.	(wh.) Alpine.	Italy.	1816.	—	H. 4.	Gmel. sib. 3. t. 10.
22 undulatum. w.	(wh.) wave-leaved.	Siberia.	1789.	6. 7. H. 4.	—	3. t. 11. f. 1.
23 divaricatum. w.	(wh.) divaricated.	—	1759.	7. 8. H. 4.		
24 salignum. w.	(wh.) Willow-like.	—	1816.	5. 8. H. 4.		
25 aciculatum. w.	(wh.) narrow-leaved.	—	—	6. 7. H. 4.		
26 cymosum. s.s.	(wh.) cyme-flowered.	Naples.	1827.	7. 9. H. 4.		
27 aviculare. E.B.	(ro.) Knot-grass.	Britain.	—	4. 10. H. ⊖.	Eng. bot. v. 18. t. 1252.	
28 Dryandri. s.s.	(bh.) Dryander's.	E. Indies.	1796.	4. 8. S. 4.		
élégans. H.K. nec alliorum.						
29 arenarium. w.k.	(pu.) sand.	Hungary.	1807.	5. 8. H. ⊖.	W. et K. hung. t. 67.	
30 ramosum. L.en.	(wh.) branching.	Siberia.	1822.	6. 8. H. 4.		
31 littorale. L.en.	(wh.) sea-side.	S. Europe.	1816.	5. 8. H. h.		
32 Bellardii. P.S.	(wh.) Bellardi's.	—	1822.	—	H. 4.	All. ped. t. 90. f. 2.
33 pátulum. M.E.	(wh.) spreading-branch'd.	Tauria.	—	—	H. 4.	

34 marítimum. w. (wh.)	sea-shore.	S. Europe.	1816.	6. 8. H. ḥ.	Flor. græc. t. 363.
35 acetōsum. M.B.	(li.) Sorrel-leaved.	Astracan.	1823.	— H. ȳ.	
36 eréctum. w.	(wh.) upright.	N. America.	1792.	7. 8. H. ȳ.	Bart.fl. amer. 3. t. 95. f. 2.
37 linifólium. B.A.	(bh.) Flax-leaved.	—	1824.	6. 8. H. ○.	
38 herniaríoïdes. s.s.(wh.)	Egyptian.	Egypt.	1822.	— F. ḥ.	
39 pensylvánicum.w.(ro.)	Pensylvanian.	N. America.	1819.	7. 8. H. ○.	
40 orientálē. w.	(re.) common.	E. Indies.	1707.	7. 10. H. ○.	Bot. mag. t. 213.
β álbūm.	(wh.) white-flowered.	—	—	— H. ○.	Rheed. mal. 12. t. 77.
41 barbátum. s.s.	(re.) bearded.	China.	1822.	6. 9. F. ȳ.	
42 tinctorium. w.	(re.) dyer's.	—	1776.	7. 8. G. ḡ.	
43 decípiens. B.P.	(ro.) decipient.	N. S. W.	1822.	— G. ȳ.	
44 plebéium. B.P.	(bh.) linear-leaved.	—	—	— G. ȳ.	
45 Persicária.	(li.) spotted.	Britain.	—	— H. ○.	Eng. bot. v. 11. t. 756.
46 lapathifólium. s.s.(gr.)	Dock-leaved.	—	—	— H. ○.	— v. 20. t. 1382.
47 incánum. W.en.	(li.) hoary.	Germany.	1804.	— H. ○.	Schrink. bav. 1. t. 3.
48 persicarioídes.k.s.(w.)	Mexican.	Mexico.	1826.	— H. ȳ.	
49 Hydropíper. E.B.(gr.)	Water pepper.	Britain.	—	7. 9. H. ○.	Eng. bot. v. 14. t. 989.
50 glábrum. w.	(gr.) smooth.	E. Indies.	1822.	— S. ȳ.	
51 mínuſ. E.B.	(li.) slender.	England.	—	— H. ○.	Eng. bot. v. 15. t. 1043.
52 rēpens.	(wh.) rooting-branched.	Fulham.	—	8. 10. H. ○.	
53 tortuósom. D.P.	(wh.) twisting.	Nepaul.	1824.	7. 10. F. ȳ.	Flor. græc. t. 364.
54 equisetifórmē.s.s.(wh.)	Equisetum-like.	Britain.	—	— H. ○.	
55 oxyspérmum.Led.(w.)	sharp-seeded.	Siberia.	1827.	— H. ȳ.	
56 arifólium. w.	(bh.) Arum-leaved.	N. America.	1816.	5. 10. H. ○.	Bart. fl. amer. t. 100.
57 ocreánum. w.	(bh.) spear-leaved.	Siberia.	1780.	6. 9. H. ȳ.	Gmel. sib. 3. t. 8.
58 virginíānum. w. (pa.)	Virginian.	N. America.	1640.	8. 9. H. ȳ.	Park. theat. 857. f. 6.
59 amphíbium. w.	(ro.) amphibious.	Britain.	—	6. 8. H.w. ȳ.	Eng. bot. v. 7. t. 436.
60 gráclile. B.P.	(pi.) slender-stalked.	N. S. W.	1822.	— G. ȳ.	
61 acetoséfólium. P.S.(re.)	Brazilian.	Brazil.	—	— S. h. ○.	Venten. cels. t. 88.

TRAGOP'YRUM. M.B. TRAGOP'YRUM.

Octandria

Trigynia.

1 lanceolátum. M.B.(pi.)	spear-leaved.	Siberia.	1770.	7. 8. H. ḥ.	Bot. reg. t. 255.
<i>Poly'gonum frutéscens. B.R.</i>					
2 buxifólium. M.B. (pi.)	Box-leaved.	Caucasus.	1800.	— H. ḥ.	Bot. mag. t. 1065.
<i>Poly'gonum críspulum. B.M.</i>					
ATRAPHA'XIS. L.	ATRAPHA'XIS.	Hexandria	Digynia.	L.	
1 spinósa. s.s.	(gr.) prickly.	Levant.	1732.	8. G. ȳ.	L'Herit. stirp. 1. t. 14.
2 unduláta. s.s.	(gr.) waved-leaved.	C. B. S.	—	6. 7. G. ḥ.	Dill. elt. t. 32. f. 36.
OXYRIA. Sm.	MOUNTAIN SORREL.	Hexandria	Digynia.	Sm.	
reníformis. Sm.	(gr.) kidney-leaved.	Britain.	—	6. 7. H. ȳ.	Eng. bot. v. 13. t. 910.
<i>Rumex dig'ynus. E.B.</i>					

R'UMEX. L. DOCK. Hexandria Trigynia. L.

1 Hydrolapáthum.s.s.(gr.)	great water.	Britain.	—	7. 8. H.w. ȳ.	Eng. bot. v. 30. t. 2104.
<i>aquáticus. E.B.</i>					
2 Británnica. s.s.	(gr.) Virginian.	N. America.	—	6. 7. H. ȳ.	
3 máximus. s.s.	(gr.) great.	Europe.	—	— H. ȳ.	Eng. bot. v. 11. t. 724.
4 acútus. E.B.	(gr.) sharp.	Britain.	—	— H. ȳ.	
5 Nemolapáthum.w.(gr.)	flat-valved.	Europe.	1816.	— H. ȳ.	
6 nemorósus. W.en. (gr.)	wood.	—	—	— H. ȳ.	
7 sanguíneus. E.B.	(gr.) bloody-veined.	England.	—	6. 7. H. ȳ.	Eng. bot. v. 22. t. 1533.
8 condylódes. M.B. (gr.)	spreading.	Caucasus.	1823.	— H. ȳ.	
9 críspus. E.B.	(gr.) curled.	Britain.	—	— H. ȳ.	Eng. bot. v. 28. t. 1998.
10 Patiéntia. w.	(gr.) Patience.	Italy.	1573.	— H. ȳ.	
11 confértus. W.en. (gr.)	close-leaved.	—	1816.	— H. ȳ.	
12 brasilién's. L.en. (gr.)	Brazilian.	Brazil.	1823.	— G. ȳ.	
13 verticillátus. s.s.	(gr.) whorled.	N. America.	1818.	— H. ȳ.	
14 spathulátus. P.S.	(gr.) spatulate-leav'd.	C. B. S.	1824.	8. 2. G. ȳ.	
15 purpúreus. P.S.	(gr.) purple-veined.	Europe.	—	6. 7. H. ȳ.	
16 linearis. s.s.	(gr.) linear-leaved.	C. B. S.	1824.	8. 12. G. ȳ.	
17 salicifólius. s.s.	(gr.) Willow-leaved.	California.	1821.	7. 10. F. ȳ.	
18 lavigátus. s.s.	(gr.) smooth.	—	1828.	— H. ȳ.	
19 sylvestris. s.s.	(gr.) wood.	Germany.	1826.	6. 8. H. ȳ.	
20 obtusifólius. E.B.	(gr.) broad-leaved.	Britain.	—	— H. ȳ.	Eng. bot. v. 28. t. 1999.
21 nepalén's. s.s.	(gr.) Nepalese.	Nepaul.	1823.	— H. ȳ.	
22 palústris. E.B.	(ye.) yellow marsh.	England.	—	7. 8. H. ȳ.	Eng. bot. v. 27. t. 1932.
23 marítimus. E.B.	(ye.) sea-side.	Britain.	—	— H. ȳ.	v. 11. t. 725.
24 divaricátus. w.	(gr.) divaricate.	Italy.	1793.	— H. ȳ.	Till. pis. t. 37. f. 2.
25 strictus. L.en.	(gr.) upright.	—	1824.	— H. ȳ.	
26 persicarioídes.w.	(gr.) Persicaria-like.	N. America.	1773.	6. 7. H. ȳ.	
27 dentátus. w.	(gr.) dentated.	Egypt.	1782.	7. 8. H. ○.	Dill. elt. t. 158. f. 191.

28 ægyptiacus. w. (gr.)	Egyptian.	Egypt.	1734.	6. 7. H. ⊙.	Till. pis. t. 37. f. 1.
29 ucranicus. s.s. (gr.)	Ukraine.	Ukraine.	1820.	— H. ⊙.	
30 crispátulus. m.	(gr.) curled-leaved.	N. America.	—	— H. ⊙.	
31 cristatus. DC.	(gr.) crested.	—	— H. ¼.	
32 púlcher. E.B.	(gr.) fiddle.	Britain.	6. 8. H. ¼.	Eng. bot. v. 22. t. 1576.
33 aquáticus. s.s. (gr.)	water.	Europe.	1818.	— H. w. ¼.	Jacq. ic. 1. t. 67.
34 scutátus. w. (gr.)	French sorrel.	France.	1596.	6. 7. H. ¼.	Moris. s. 5. t. 28. f. 7.
35 hastifoliás. M.B. (gr.)	hastate-leaved.	Caucasus.	1816.	— H. ¼.	Zanon, hist. 9. t. 6.
36 vesicárius. w. (gr.)	bladder.	Africa.	1656.	7. 8. H. ⊙.	
37 tingitánus. w. (gr.)	Tanger.	Barbary.	1680.	6. 8. H. ¼.	
38 longifoliás. K.S. (gr.)	long-leaved.	S. America.	1827.	7. 10. F. ¼.	
39 sarcorhizus. L.en.(gr.)	fleshy-rooted.	C. B. S.	1824.	8. 12. G. ¼.	
40 Burchellii. s.s. (gr.)	Burchell's.	—	1816.	8. 10. G. ™.	
41 Lunária. w. (gr.)	tree.	Canaries.	1698.	6. 7. G. ℡.	Pluk. alm. t. 252. f. 3.
42 orientális. Bern. (gr.)	oriental.	Levant.	1821.	7. 9. H. ¼.	
43 conspersus. Ot. (gr.)	scattered.	1824.	— H. ¼.	
44 domésticus. s.s. (gr.)	various-leaved.	Sweden.	1823.	— H. ¼.	
45 róseus. w. (gr.)	rose.	Egypt.	1737.	7. 8. H. ⊙.	Cavan. ic. 1. t. 41. f. 1.
46 lácerus. s.s. (gr.)	torn-leaved.	—	1825.	— H. ⊙.	
47 bucephalóphorus. (gr.)	Basil-leaved.	Italy.	1683.	6. H. ⊙.	
48 fimbriátus. s.s. (gr.)	fringe-valved.	C. B. S.	1816.	7. 9. G. ¼.	
49 Brównii. s.s. (gr.)	Brown's fringed.	N. S. W.	1824.	— G. ¼.	
50 Acetosa. E.B. (gr.)	common Sorrel.	Britain.	6. 8. H. ¼.	Eng. bot. v. 2. t. 127.
51 Acetosella. E.B. (st.)	sheep's sorrel.	—	5. 7. H. ¼.	— v. 24. t. 1674.
52 abyssinicus. Jacq.(gr.)	Abyssinian.	Abyssinia.	1823.	— G. ™.	Jacq. vind. 3. t. 98.
53 tuberósus. w. (gr.)	tuberous-rooted.	S. Europe.	1752.	6. 8. H. ¼.	
54 arifoliás. s.s. (gr.)	Arum-leaved.	Africa.	1775.	12. 4. G. ™.	Zorn. ic. 261.
55 triangulàris. DC. (gr.)	triangular-leaved.	S. Europe.	1824.	— F. ¼.	
56 amplexicaulis.s.s.(gr.)	stem-clasping.	Pyrenees.	1816.	6. 8. H. ¼.	
57 luxuriáns. s.s. (gr.)	luxuriant.	S. Europe.	— H. ¼.	
58 alpinus. w. (gr.)	Alpine.	France.	1597.	6. 7. H. ¼.	
59 frutéscens. s.s. (gr.)	frutescent.	Tristan d'Ac.	1816.	6. 9. G. ™.	
60 gigánteus. H.K. (gr.)	tall.	Sandwich Isl.	1796.	6. 8. G. ¼.	
EMEX. S.S.	'EMEX. Hexandria Trigynia. S.S.	—	—	—	
spinosus. s.s. (gr.)	spiny-seeded.	Candia.	1656.	6. 7. H. ⊙.	
Rùmex spinòsus. w.	—	—	—	—	
PODO'PTERUS. K.S. Podo'PTERUS.	Hexandria Trigynia. S.S.	—	—	—	
mexicanus. k.s. (wh.)	Mexican.	Mexico.	1825.	... G. ™.	H. et B. pl. am. 2.t.107.
TRIPLARIS. K.S.	TRIPLARIS.	Dicecia Enneandria. L.	—	—	
americana.k.s.(ro.pu.)	American.	S. America.	1823.	5. 8. S. ™.	Jacq. amer. 13.t.173.f.5.
RH'EUM. L.	RHUBARB.	Enneandria Trigynia. L.	—	—	
1 Rhapóticum.w.(wh.)	common.	Asia.	1573.	5. 6. H. ¼.	Sabb. hort. 1. t. 34.
2 undulátum. w. (wh.)	wave-leaved.	China.	1734.	— H. ¼.	L. amoen. acad. 3. t. 4.
3 críspum. And. (wh.)	curl-leaved.	—	1820.	— H. ¼.	
4 compáctum. w. (wh.)	thick-leaved.	Tartary.	1758.	— H. ¼.	Mill. ic. 2. t. 218.
5 tatáricum. w. (wh.)	Tartarian.	—	1793.	— H. ¼.	
6 hy'bridum. w. (wh.)	bastard.	Asia.	1778.	— H. ¼.	Murr. c. gott. t. 1.
7 austriacum. And. (wh.)	Austrian.	Austria.	1819.	— H. ¼.	
8 sibíicum. And. (wh.)	Siberian.	Siberia.	—	— H. ¼.	
9 fenestrátum. F. (wh.)	windowed.	—	1760.	— H. ¼.	
10 rotundifoliúm. And. (w.)	round-leaved.	—	—	— H. ¼.	
11 nútans. F. (wh.)	nodding.	Asia.	1824.	— H. ¼.	Ann. mus. 1. t. 49.
12 Ríbes. w. (wh.)	warted-leaved.	Levant.	1724.	— H. ¼.	Linn. fasc. 7. t. 4.
13 palmátum. w. (wh.)	palmated.	China.	1763.	4. 5. H. ¼.	Swt. br. fl. gar. t. 269.
14 austrále. D.P. (d.pu.)	Nepaul.	Nepaul.	1826.	— H. ¼.	
ERIO'GONUM. M.	ERIO'GONUM.	Enneandria Trigynia. N.	—	—	
1 tomentósüm. M. (wh.)	woolly.	Carolina.	1811.	5. 6. H. ¼.	Mich. amer. v. 1. t. 24.
2 flávum. N. (ye.)	ilkly.	Missouri.	—	7. H. ¼.	
sericeum. Ph.	—	—	—	—	
3 pauciflórum. Ph. (wh.)	few-flowered.	Louisiana.	1812.	— H. ¼.	
4 longifoliúm. N. (wh.)	long-leaved.	Arkansa.	1824.	— H. ¼.	
CALLIGONUM. W.	CALLIGONUM.	Dodecandra Tetragynia. L.	—	—	Pall. ros. 2. t. 77. 78.
1 Pallásia. w. (wh.gr.)	Caspian.	Caspian shores.	1780. 8.	H. ™.	
2 Pandéri. w. (gr.wh.)	Fischer's.	—	1824.	7. 9. H. ™.	
3 comosum. P.S. (gr.wh.)	tufted.	Persia.	1829. H. ™.	
KŒNIGIA. W.	KŒNIGIA.	Triandria Trigynia. L.	—	—	
islándica. w. (gr.)	Iceland.	Iceland.	1773.	4. H. ⊙.	Lam. ill. gen. t. 51.

ORDO CLXX.

LAURINÆ. Brown prodr. p. 401.

LA'URUS. W.	LA'URUS. Enneandria Monogynia. L.	S. Europe.	1561.	4. 5. H. h.	Flor. græc. t. 365.
1 nòbilis. w.	(gr.ye.) Sweet Bay.	—	1561.	—	—
β undulata.	(gr.ye.) wave-leaved.	—	—	H. h.	
γ salicifolia.	(gr.ye.) Willow-leaved.	—	—	H. h.	
δ variegata.	(gr.ye.) variegated.	—	—	H. h.	
2 coriacea. s.s.	(gr.ye.) leathery-leaved.	Jamaica.	1822.	—	S. h.
3 salicifolia. s.s.	(gr.) Willow-leaved.	W. Indies.	—	—	S. h.
4 Catesb'xi. P.S.	(wh.) Catesby's.	Florida.	1811.	4. 6. F. h.	Catesb. car. 1. t. 2b.
5 carolinensis. M.	(gr.ye.) Carolina.	N. America.	1806.	—	F. h.
a glabra.	(gr.ye.) smooth-leaved.	—	—	F. h.	
β pubescens.	(gr.ye.) downy.	—	—	F. h.	
γ obtusa.	(gr.ye.) blunt-leaved.	—	—	F. h.	
6 Borbònia. w.	(gr.ye.) Bourbon.	S. America.	1739.	—	F. h.
7 thrysiflora. s.s.	(gr.ye.) bunch-flowered.	Madagascar.	1824.	—	S. h.
8 indica. w.	(gr.ye.) royal bay.	Madeira.	1665.	10. 3. G. h.	Pluk. alm. t. 304. f. 1.
9 f'etens. w.	(gr.ye.) Madeira or Til.	—	1760.	—	G. h.
10 canariensis. W.en.(st.)	Canary.	Canaries.	1815.	—	G. h.
11 aggregata. B.M.	(wh.) clustered.	China.	1812.	—	S. h.
12 Chloroxylon.w.	(gr.ye.) Cogwood-tree.	Jamaica.	1778.	—	S. h.
13 pendula. w.	(gr.ye.) pendulous.	—	1822.	—	S. h.
14 glauca. w.	(gr.ye.) glaucous.	Japan.	1806.	—	G. h.
15 aestivalis. w.	(gr.ye.) veined-leaved.	N. America.	1765.	4. 5. G. h.	Comm. hort. 1. t. 97.
16 Benzoin. w.	(gr.ye.) Benjamin-tree.	—	1683.	—	H. h.
17 Dióspyrus.P.S.	(gr.ye.) twiggy.	—	1810.	—	H. h.
18 geniculata. M.	(gr.ye.) flexuose.	—	1759.	—	H. h.
19 Sassafras. w.	(gr.ye.) Sassafras-tree.	—	1633.	5. 6. H. h.	Brown. jam. t. 7. f. 1.
20 álbida. n.	(gr.ye.) white Sassafras.	—	—	—	Catesb. car. 1. t. 55.
PERSEA. G.	ALLIGATOR-PEAR. Enneandria Monogynia. S.S.	W. Indies.	1739.	4. 6. S. h.	Bot. reg. t. 1258.
gratissima. G.	(st.) common.	—	—	—	
Laurus Pérsæa. w.	—	—	—	—	
CINNAM'OMUM. B.P. CINNAMON.	Enneandria Monogynia.	—	—	—	
1 Cássia. s.s.	(gr.) pubescent.	E. Indies.	1768.	5. 9. S. h.	Andr. deposit. t. 596.
Laurus Cinnamomum. A.R. Pérsæa Cássia. s.s.	—	—	—	—	
2 vérum.	(gr.) true.	Ceylon.	1763.	6. 9. S. h.	Bot. mag. t. 1636.
Laurus Cássia. B.M. Pérsæa Cinnamomum. s.s.	—	—	—	—	
3 dulcis. s.s.	(gr.wh.) sweet.	E. Indies.	1824.	—	S. h.
4 nitidum. H.E.F.	(gr.) glossy-leaved.	Sumatra.	—	6. 9. S. h.	Hook. ex. flor. t. 176.
5 Bejolghôta.	(gr.) square-branch'd.	E. Indies.	1805.	—	S. h.
Laurus Bejolghôta. L.T. Laùrus malabáthrica. H.B?	—	—	—	—	
6 Culitlaban.	(gr.) three-flowered.	Moluccas.	—	—	Rumph. amb. 2. t. 14.
Laurus Culitlaban. H.B.	—	—	—	—	
7 gráclie. Hort.	(gr.) slender.	E. Indies.	1805.	—	S. h.
8 Camphora.	(wh.) Camphire-tree.	Japan.	1727.	3. 6. G. h.	Bot. mag. t. 2658.
Laurus Camphora. B.M. Pérsæa Camföra. s.s.	—	—	—	—	
OC'OTEA. K.S.	OC'OTEA. Dodecadria Monogynia.	—	—	—	
psychotrioides. K.S.(gr.)	spear-leaved.	Mexico.	1828.	—	G. h.
CRYPTOC'ARYA. B.P. CRYPTOC'ARYA.	Dodecadria Monogynia.	—	—	—	
1 glaucescens. B.P.	(gr.) glaucous-leaved.	N. S. W.	1820.	—	G. h.
2 obovata. B.P.	(gr.) obovate-leaved.	—	1823.	—	G. h.
TETRANTH'ERA. B.P. TETRANTH'ERA.	Diœcia Enneandria.	—	—	—	
1 laurifolia.B.R.	(gr.wh.) Laurel-leaved.	Japan.	1823.	1. 5. G. h.	Bot. reg. t. 893.
2 apetala. B.P.	(gr.wh.) petalless.	E. Indies.	—	—	Roxb. corom. 2. t. 147.
3 dealbata. B.P.	(gr.) white-leaved.	N. S. W.	1824.	—	G. h.
4 involucrata. Rox.	(gr.) involucred.	E. Indies.	1820.	1. 5. G. h.	
CASS'YTHA. L.	CASS'YTHA.	Dodecadria Monogynia.	—	—	
1 filiformis. L.	(wh.) slender.	C. B. S.	—	5. 8. G. ȳ. —.	Rumph. amb. 5. t. 184. f. 4
2 glabælla. B.P.	(wh.) smooth.	N. S. W.	1822.	—	G. ȳ. —.
3 paniculata. B.P.	(wh.) panicled.	—	1824.	—	G. ȳ. —.

LAURINIS AFFINE.

GYROCA'RPUS. B.P.	GYROCA'RPUS.	Polygamia Monœcia.				
1 americanus. w.	(ye.) American.	W. Indies.	1816.	S. h.	Jacq. amer. t. 178. f. 80.
2 asiaticus. w.	(ye.) Asiatic.	E. Indies.	1812.	S. h.	Roxb. corom. 1. t. 1.
<i>Jacquini.</i> R.C.						
HERNA'NDIA. L.	JACK IN A BOX.	Monœcia Triandria. L.				
1 sonòra. w.	(wh.) peltate-leaved.	W. Indies.	1693.	S. h.	Rumph. amb. 2. t. 85.
2 guianensis. w.	(wh.) Guiana.	Guiana.	1823.	S. h.	Aub. gni. 2. t. 309.
3 ovígera. w.	(wh.) oval-leaved.	E. Indies.	1820.	S. h.	Rumph. amb. 3. t. 123.

ORDO CLXXI.

MYRISTICÆÆ. *Brown prodr. p. 399.*

MYRI'STICA. W.	NUTMEG.	Dicæcia Monadelphia. L.				
1 moschata. w.	(ye.) true.	E. Indies.	1795.	S. h.	Hook. ex. flor. t. 155-156
<i>officinalis.</i> H.E.F.	<i>aromática.</i> Lam.					
2 sebifera. s.s.	(ye.) woolly-fruited.	S. America.	1829.	S. h.	
3 fátua. s.s.	(ye.) tasteless.	Surinam.	1812.	S. h.	Pluk. alm. t. 250. f. 6.

ORDO CLXXII.

PROTEACEÆ. *Brown Linn. trans. 10. p. 46.*

A'ULAX. P.L.	A'ULAX.	Dicæcia Tetrandria. H.K.				
1 pinifolia. L.T.	(ye.) Pine-leaved,	C. B. S.	1780.	7. 9.	G. h.	Andr. deposit. t. 76.
<i>Prôtea pinifolia.</i> A.R.						
2 umbellata. L.T.	(ye.) umbelled.		1774.	6. 8.	G. h.	Andr. deposit. t. 248.
LEUCADE'NDRON. L.T.	LEUCADE'NDRON.	Dicæcia Tetrandria. H.K.				
1 argénteum. L.T.	(st.) Silver-tree.	C. B. S.	1693.	6. 8.	G. h.	Bot. reg. t. 979.
<i>Prôtea argénteæ.</i> w.						
2 plumosum. L.T.	(ye.) feathered.		1774.	—	G. h.	Thunb. diss. 40. t. 4.
<i>Prôtea parviflora.</i> Th.						
3 reticulatum. L.T.	(ye.) retuse-leaved.		1825.	—	G. h.	
4 spathulatum. L.T.	(ye.) spathulate.		1812.	6. 8.	G. h.	
5 angustatum. L.T.	(ye.) narrow-leaved.		1824.	—	G. h.	
6 imbricatum. L.T.	(ye.) imbricated.		1790.	—	G. h.	
7 buxifolium. L.T.	(ye.) Box-leaved.		1812.	—	G. h.	
8 Levisanus. L.T.	(ye.) short-leaved.		1774.	4. 6.	G. h.	
9 linifolium. L.T.	(ye.) Flax-leaved.		1809.	—	G. h.	
10stellatum. L.T.	(ye.) starry.		—	5. 6.	G. h.	
<i>Prôtea stellariæ.</i> B.M. squarrösa. K.P.						
11 fusciflorum. L.T.	(br.) brown-flowered.		1812.	—	G. h.	Jacq. schœnb. I. t. 27.
12 tortum. L.T.	(ye.) twisted-leaved.		1790.	3. 5.	G. h.	Bot. reg. t. 826.
13 cinereum. L.T.	(ye.) gray-leaved.		1774.	7. 8.	G. h.	
14 corymbosum. L.T.	(ye.) corymbed.		1790.	4. 7.	G. h.	Bot. reg. t. 402.
15 decorum. L.T.	(ye.) decorous.		—	5. 8.	G. h.	
16 squarrosum. L.T.	(ye.) squarrose.		1812.	—	G. h.	Boerh. lug. 2.p. 197.c.ic.
<i>Prôtea strobilina.</i> L.						
17 cóncolor. L.T.	(ye.) one-coloured.		1774.	3. 6.	G. h.	Andr. deposit. t. 307.
<i>Prôtea globosa.</i> B.M. 878. <i>strobilina.</i> Schrad. s. han. I. t. 1.						
18 grandiflorum. L.T.	(ye.) great-flowered.	C. B. S.	1789.	4. 6.	G. h.	
19 ovale. L.T.	(ye.) oval-leaved.		1818.	—	G. h.	
20 venosum. L.T.	(ye.) veined-leaved.		1815.	—	G. h.	
21 decurrent. L.T.	(ye.) decurrent.		1812.	6. 8.	G. h.	
22 glábrum. L.T.	(ye.) smooth.		—	3. 6.	G. h.	

Wend. h. herr. t. 14?

t. 1.

Jacq. schœnb. I. t. 26.

Bot. mag. t. 881.

Jacq. schœnb. I. t. 27.

Bot. reg. t. 826.

Sal. par. lond. t. 105.

23 stríctum. L.T.	(ye.) upright.	C. B. S.	1795.	4. 6. G. h.	Sal. par. lond. t. 75.
	<i>Euryspérnum salicifolium</i> . P.L. <i>Prôtea confíera</i> . Andr. rep. t. 541.				
24 virgátum. L.T.	(ye.) slender.	C. B. S.	G. h.	
25 adscéndens. L.T.	(ye.) ascending.		1774.	6. 8. G. h.	Pluk. mant. t. 229. f. 6.
26 concínum. L.T.	(ye.) neat.		1800.	— G. h.	
27 salígnum. L.T.	(ye.) Willow-leaved.		1774.	4. 6. G. h.	Boer. lugd. 2.p. 204. ic.
28 uliginósus. L.T.	(ye.) swamp.		1795.	— G. h.	Breyn. cent. 21. t. 9.
29 floridum. L.T.	(ye.) florid.		—	— G. h.	Pluk. phyt. t. 229. f. 4.
	<i>Prôtea salígnæ</i> mas. et fem. Andr. rep. t. 572?				
30 platyspérnum. L.T. (ye.) flat-seeded.			1818.	— G. h.	
31 comosum. L.T.	(ye.) tufted.		1812.	5. 8. G. h.	
32 æmulum. L.T.	(ye.) related.		1789.	6. 9. G. h.	
	<i>Prôtea incúrvæ</i> . Andr. rep. t. 429.				
33 abietinum. L.T.	(ye.) Pine-leaved.		—	7. 9. G. h.	Andr. deposit. t. 461.
	<i>Prôtea teretifolia</i> . Andr. rep. t. 461.				
34 scábrum. L.T.	(ye.) rough.		1812.	4. 6. G. h.	
35 sericeum. L.T.	(ye.) silky-leaved.		1818.	— G. h.	
36 Globulária. L.T.	(ye.) globe-headed.		1810.	— G. h.	
37 pubésens. L.T.	(ye.) pubescent.		1822.	— G. h.	Lam. ill. 1. t. 53.
38 ericifolium. L.T.	(ye.) Heath-leaved.		—	— G. h.	
39 infléxum. L.en.	(ye.) inflexed.		1812.	— G. h.	
40 caudátum. L.en.	(ye.) tailed.		—	— G. h.	
41 marginátum. L.en.	(ye.) villous-margined.		1818.	— G. h.	
42 crassifolium. L.T.	(ye.) thick-leaved.		1816.	— G. h.	
43 cartilagínænum. L.T.	(ye.) bony-margined.		—	— G. h.	
PETRO' PHILA. L.T.	PETRO' PHILA.	Tetrandria Monogynia. R.S.			
1 teretifolia. L.T.	(wh.) round-leaved.	N. Holland.	1824. G. h.	
2 filifolia. L.T.	(wh.) thread-leaved.		— G. h.	
3 aciculáris. L.T.	(wh.) needle-leaved.		— G. h.	
4 pulchella. L.T.	(wh.) Fennel-leaved.	N. S. W.	1790.	7. 8. G. h.	
5 pedunculátæ. L.T. (st.) peduncled.			1824.	4. 8. G. h.	Bot. mag. t. 796.
6 diversifolia. L.T. (wh.) various-leaved.		N. Holland.	1803.	4. 6. G. h.	
7 trifida. L.T.	(wh.) trifid-leaved.		1824. G. h.	
ISOP' OGON. L.T.	Isor' oeon.	Tetrandria Monogynia. R.S.			
1 teretifolius. L.T. (wh.) round-leaved.		N. Holland	1824.	4. 8. G. h.	
2 anethiifolius. L.T. (wh.) Dill-leaved.		N. S. W.	1796.	3. 6. G. h.	Cavan. ic. 6. t. 549.
3 formosus. L.T.	(li.) handsome.	N. Holland.	1805.	— G. h.	Bot. reg. t. 1288.
4 anomóniifolius. L.T. (st.) Anemone-leav'd.		N. S. W.	1791.	7. 9. G. h.	Bot. mag. t. 697.
5 divaricátæ. A.R. (ye.) spreading.			—	— G. h.	Andr. rep. t. 465.
6 trilobus. L.T.	(wh.) three-lobed.	N. Holland.	1803.	5. 6. G. h.	
7 longifolius. L.T.	(ye.) long-leaved.		1820.	— G. h.	Bot. reg. t. 900.
8 propinquus. M.C. (ye.) related.			1824. G. h.	
9 attenuátus. L.T. (wh.) attenuated.			—	5. 6. G. h.	
10 polycéphalus. L.T. (wh.) many-headed.			— G. h.	
11 axillaris. L.T. axillary.		— G. h.	
PR' OTEA. L.T.	PR' OTEA.	Tetrandria Monogynia. L.			
1 cynaroïdes. L.T.	(re.) Artichoke-fl'd.	C. B. S.	1774.	11. 3. G. h.	Bot. mag. t. 770.
2 latifolia. L.T.	(sc.) ray-flowered.		1806.	7. 9. G. h.	— t. 1717.
	<i>a coccinea.</i>		—	— G. h.	Andr. deposit. t. 646.
	<i>β viridiflora.</i>		—	— G. h.	— t. 646.
3 compácta. L.T.	(pu.) compact.		1810.	— G. h.	
4 longiflora. L.T.	(st.) long-flowered.		1795.	1. 4. G. h.	Sm. exot. bot. 2. t. 81.
	<i>ochroleuca</i> . Sm. <i>lacticolor</i> . Par. lond. t. 27.				
5 coccinea. L.T.	(sc.) scarlet-flowered.		1810.	5. 6. G. h.	
6 speciosa. L.T.	(wh. br.) splendid.		1786.	3. 6. G. h.	Knight's prot. c. ic.
7 obtusa.	(pa. pi.) obtuse.		—	— G. h.	Andr. deposit. t. 110.
	<i>speciosa</i> . B.M. 1183. <i>Erodéndrum obtusum</i> . K.P.				
8 macrophylla. L.T.	(wh.) large-leaved.		1810.	— G. h.	Bot. mag. t. 1713.
9 formosa. L.T.	(sc.) handsome.		1789.	5. 6. G. h.	
	<i>coronata</i> . A.R. 469. <i>Erodéndrum formosum</i> . Par. lond. t. 76.				
10 melaleuca. L.T.	(bk.) black-fringed.		1786.	3. 7. G. h.	Andr. deposit. t. 103.
11 Lepidocárpon. L.T.	(bk.) crested.		1806.	— G. h.	Wein. phyt. 4. t. 895.
12 nerifolia. L.T.	(pu.) Oleander-leaved.		—	2. 4. G. h.	Bot. reg. t. 208.
13 pulchella. L.T.	(pu.) wave-leaved.		1795.	3. 8. G. h.	— t. 20.
	<i>a citiata.</i>		—	— G. h.	Andr. deposit. t. 270.
	<i>β glabra.</i>		—	— G. h.	— t. 277.
	<i>γ speciosa.</i>		—	— G. h.	— t. 442.
14 pátens. L.T.	(st. bk.) spreading.		1789.	3. 6. G. h.	— t. 543.
15 magnifica. A.R. (pu. bk.) magnificent.			—	— G. h.	— t. 438.

16 incómpcta. L.T. (<i>wh.</i>) villous-branched. C. B. S.	1807.	5. 7. G. h.	
17 longifolia. L.T. (<i>st. bk.</i>) long-leaved.	1798.	3. 8. G. h.	Bot. reg. t. 47.
18 umbonális. (<i>st. pu.</i>) embossed.		— G. h.	Andr. deposit. t. 144.
longifolia. var. <i>cone turbinato</i> . A.R. <i>Erodéndrum umbonálē</i> . K.P.			
19 ligulaefolia. (<i>st. pu.</i>) strap-leaved.	1798.	3. 8. G. h.	Andr. deposit. t. 133.
longifolia ferruginoso-purpurea. A.R. <i>Erodéndrum ligulaefolium</i> . K.P.			
20 mellifera. L.T. (<i>ro. wh.</i>) honey-bearing.	1774.	12. 5. G. h.	Bot. mag. t. 346.
β álba. (<i>wh.</i>) white-flowered.		— G. h.	Andr. deposit. ic.
21 grandiflora. L.T. (<i>ro. w.</i>) great-flowered.	1787.	5. 6. G. h.	Bot. mag. t. 2447.
β marginata. (<i>ro. wh.</i>) red margined.		— G. h.	Bot. reg. t. 569.
22 Scólymus. L.T. (<i>ro. wh.</i>) small-flowered.	1780.	— G. h.	Bot. mag. t. 698.
23 mucronifolia. L.T. (<i>wh. ro.</i>) dagger-leaved.	1803.	7. 12. G. h.	Andr. rep. t. 500.
24 nána. L.T. (<i>cr.</i>) dwarf.	1787.	4. 7. G. h.	Sm. exot. bot. 1. t. 44.
rosacea. Sm. <i>acufolia</i> . Par. lond. t. 2.			
25 péndula. L.T. (<i>wh. gr.</i>) pendulous.	1806.	3. 6. G. h.	
26 ténax. L.T. (<i>st.</i>) tough.	1801.	2. 5. G. h.	Sal. par. lond. t. 70.
27 canaliculata. L.T. (<i>pi.</i>) channel-leaved.	1800.	12. 2. G. h.	Andr. deposit. t. 437.
28 acuminata. B.M. (<i>cr.</i>) sharp-pointed.	1809.	3. 6. G. h.	Bot. mag. t. 1694.
29 acaulis. L.T. (<i>st. br.</i>) short-stalked.	1802.	5. 9. G. h.	Wein. phyt. 4. t. 898.
30 glaucophylla. P.L. (<i>st. br.</i>) glaucous-leaved.		— 7. 9. G. h.	Sal. par. lond. t. 11.
31 elongata. L.T. (<i>st. br.</i>) elongated.	1810.	— G. h.	
32 angustata. L.T. (<i>st.</i>) narrow-leaved.	1809.	— G. h.	
33 revoluta. L.T. (<i>wh.</i>) revolute-leaved.	1818.	— G. h.	
34 tenuifolia. L.T. (<i>wh. br.</i>) slender-leaved.	1816.	— G. h.	
35 l'avis. L.T. (<i>ro. gr.</i>) smooth-leaved.	1806.	5. 7. G. h.	Bot. mag. t. 2439.
longifolia. Par. lond. t. 37.			
36 scábia. L.T. (<i>wh. br.</i>) rough-leaved.	1809.	— G. h.	
37 répens. L.T. (<i>re. wh.</i>) creeping.	1800.	6. 8. G. h.	Wein. phyt. 4. t. 897.a.
38 lórea. L.T. (<i>wh.</i>) thong-leaved.	1812.	— G. h.	
39 turbiniflora. L.T. (<i>ro.</i>) turfy.	1803.	4. 5. G. h.	Sal. par. lond. t. 103.
40 Scopoléndrium. L.T. (<i>wh. br.</i>) Hart's-tongue.	1802.	— G. h.	
41 cordáta. L.T. (<i>cr.</i>) heart-leaved.	1790.	3. 5. G. h.	Andr. deposit. t. 289.
42 amplexicaulis. L.T. (<i>re.</i>) stem-clasping.	1802.	1. 3. G. h.	Sal. par. lond. t. 67.
43 humilius. L.T. (<i>pu.</i>) low-flowering.		6. 8. G. h.	Andr. deposit. t. 532.
44 acerosa. L.T. (<i>cr.</i>) Pine-leaved.	1803.	3. 5. G. h.	Bot. reg. t. 351.
LEUCOSPER'MUM. L.T. LEUCOSPER'MUM. Tetrandria Monogynia. R.S.			
1 lineáre. L.T. (<i>ye.</i>) linear-leaved.	C. B. S.	1774.	8. 9. G. h.
2 attenuatum. L.T. (<i>ye.</i>) attenuated.		1810.	— G. h.
3 Tótum. L.T. (<i>ye.</i>) smooth-bracted.		1774.	6. 8. G. h.
4 médium. L.T. (<i>ye.</i>) oval-leaved.		1794.	— G. h.
<i>Prótea formosa</i> . A.R.?			Andr. deposit. t. 17?
5 elliptic. L.T. (<i>ye.</i>) elliptic.		1803.	5. 8. G. h.
6 Conocárpum. L.T. (<i>go.</i>) many-toothed.		1774.	— G. h.
7 grandiflorum. L.T. (<i>go.</i>) great-flowered.		1800.	5. 7. G. h.
<i>Leucadéndron grandiflorum</i> . P.L.			
8 pùberum. L.T. (<i>ye.</i>) downy-leaved.		1774.	5. 8. G. h.
9 buxifolium. L.T. (<i>ye.</i>) Box-leaved.		1822.	— G. h.
10 pátulum. L.T. (<i>ye.</i>) spreading.		1816.	— G. h.
11 spathulatum. L.T. (<i>ye.</i>) spathulate.		1810.	— G. h.
12 tomentosum. K.P. (<i>ye.</i>) cottony.		1789.	8. 9. G. h.
13 párlle. K.P. (<i>ye.</i>) matched.		1790.	— G. h.
14 cándicans. (<i>or.</i>) Rose-scented.		—	— G. h.
<i>Prótea cándicans</i> . A.R.			
15 Hypophyllum. L.T. (<i>ye.</i>) trifid-leaved.		1787.	— G. h.
16 crinitum. L.T. (<i>ye. li.</i>) villous-bracted.		1824.	— G. h.
17 oleafolium. L.T. (<i>ye.</i>) Olive-leaved.		1822.	4. 6. G. h.
18 diffusum. L.T. (<i>ye.</i>) diffuse.		1823.	— G. h.
MIMETES. L.T. MIMETES. Tetrandria Monogynia. R.S.			
1 hírta. L.T. (<i>ye. cr.</i>) hairy.	C. B. S.	1774.	6. 8. G. h.
2 capitulata. L.T. (<i>re.</i>) small-headed.		1322.	— G. h.
3 pauciflora. L.T. (<i>re.</i>) few-flowered.		1816.	— G. h.
4 palústris. K.P. (<i>re.</i>) marsh.		1802.	— G. h.
5 cucullata. L.T. (<i>re.</i>) three-toothed.		1789. G. h.
<i>Prótea cucullata</i> . W.			
6 Hartogii. L.T. (<i>re.</i>) woolly-leaved.		1810. G. h.
7 divaricata. L.T. (<i>wh.</i>) divaricate.		1795.	6. 9. G. h.
8 vacciniifolia. (<i>re. wh.</i>) Vaccinium-leaved.		1800.	— G. h.
<i>Diastélla vacciniifolia</i> . K.P.			
9 purpurea. L.T. (<i>pu.</i>) Heath-leaved.		1789.	11. 12. G. h.

SERR'URIA. L.T.	SERR'URIA.	Tetrandria Monogynia. R.S.
1 acrocárpa. L.T.	(<i>li.</i>) sharp-fruited.	C. B. S. 1822. 4. 6. G. h.
2 elevàta. L.T.	(<i>li.</i>) sandy-hill.	1821. — G. h.
3 Aitòni. L.T.	(<i>li.</i>) Aiton's.	1823. — G. h.
4 diffusa. L.T.	(<i>li.</i>) diffuse.	1810. 6. 8. G. h.
5 abrotanifòlia. K.P.	(<i>li.</i>) Southernwood-leaved.	1803. — G. h.
Pròtea abrotanifòlia hírta. A.R.		Andr. deposit. t. 522.
6 millefòlia. K.P.	(<i>li.</i>) thousand-leaved.	— — G. h.
7 artemisiæfòlia. K.P.	(<i>li.</i>) Wormwood-leaved.	1789. — G. h.
8 pinnàta. L.T.	(<i>li.</i>) slender.	1803. — G. h.
9 arenària. L.T.	(<i>li.</i>) sand.	— — G. h.
10 cyanoïdes. L.T.	(<i>li.</i>) trifid-leaved.	— — G. h.
11 scariòsa. L.T.	(<i>li.</i>) scaly-bracted.	1816. 5. 7. G. h.
12 pedunculàta. L.T.	(<i>li.</i>) woolly-headed.	1789. 5. 8. G. h.
Pròtea glomeràta. A.R.		Pluk. mant. t. 345. f. 6.
13 scopària. L.T.	(<i>li.</i>) Broom.	1809. — G. h.
14 Nivèni. L.T.	(<i>li.</i>) Niven's.	1800. — G. h.
Pròtea decúmbens. A.R. non Th.		Andr. deposit. t. 349.
15 villòsa. L.T.	(<i>li.</i>) villous.	1810. 4. 7. G. h.
16 fèniculàcea. L.T.	(<i>li.</i>) Fennel-leaved.	1816. — G. h.
17 ciliàta. L.T.	(<i>li.</i>) ciliated.	1803. — G. h.
18 congestà. L.T.	(<i>li.</i>) crowded.	1816. — G. h.
19 squarròsa. L.T.	(<i>li.</i>) squarrose.	1810. — G. h.
20 phyllicoïdes. L.T.	(<i>li.</i>) Phylica-flowered.	1788. — G. h.
Pròtea abrotanifòlia. A.R.		Andr. deposit. t. 507. f. 4.
21 àmulà. L.T.	(<i>li.</i>) grey-branched.	1803. — G. h.
22 pàrilis. K.P.	(<i>li.</i>) matched.	— — G. h.
23 odoràta.	(<i>li.</i>) sweet-scented.	— — G. h.
Pròtea abrotanifòlia odoràta. A.R.		Andr. deposit. t. 507.
24 emarginàta.	(<i>li.</i>) emarginate.	1800. — G. h.
arenària. K.P. non Br. abrotanifòlia minor. A.R.		Andr. deposit. t. 536.
25 decúmbens. L.T.	(<i>li.</i>) decumbent.	1816. — G. h.
26 flagellàris. L.T.	(<i>li.</i>) procumbent.	— — G. h.
27 glomeràta. L.T.	(<i>li.</i>) many-headed.	1789. — G. h.
28 decipiens. L.T.	(<i>li.</i>) decipient.	1806. — G. h.
29 Roxburghii. L.T.	(<i>li.</i>) Roxburgh's.	— — G. h.
30 Burmánni. L.T.	(<i>li.</i>) Burmann's.	1786. — G. h.
Pròtea Serrària. L.		Burm. afr. t. 99. f. 1.
31 triteràta. L.T.	(<i>wh.</i>) silvery-flowered.	1802. — G. h.
Pròtea argentiflòra. A.R.		Andr. deposit. t. 447.
32 elongàta. L.T.	(<i>li.</i>) long-peduncled.	1800. — G. h.
33 crithmifòlia. L.T.	(<i>li.</i>) Samphire-leaved.	1816. — G. h.
NIV'ENIA. L.T.	NIV'ENIA.	Tetrandria Monogynia. R.S.
1 Scéptrunn. L.T.	(<i>pu.</i>) sceptre-like.	C. B. S. 1790. 5. 6. G. h.
2 spathulàta. L.T.	(<i>pu.</i>) maiden hair-leaved.	— 7. 8. G. h.
3 parvifòlia. L.T.	(<i>pu.</i>) small-leaved.	1810. — G. h.
4 spicàta. L.T.	(<i>pu.</i>) spiked.	1786. 6. 8. G. h.
5 crithmifòlia. L.T.	(<i>pu.</i>) Samphire-leaved.	1797. — G. h.
Pròtea Lagòpus. A.R. non Thunb.		Andr. deposit. t. 243.
6 mèdia. L.T.	(<i>bh.</i>) middle.	1803. — G. h.
Pròtea spicata. A.R. ?		Andr. deposit. t. 234?
7 Lagòpus. L.T.	(<i>pu.</i>) Hare's-foot.	1810. — G. h.
SOROC'E'PHALUS. L.T. SOROC'E'PHALUS.		Tetrandria Monogynia. R.S.
1 imbérbis. L.T.	(<i>pu.</i>) smooth.	C. B. S. 1806. 6. 8. G. h.
2 spatallòides. L.T.	(<i>pu.</i>) club-bearing.	1803. — G. h.
3 tenuifòlia. L.T.	(<i>pu.</i>) slender-leaved.	1802. — G. h.
4 lanàtus. L.T.	(<i>pu.</i>) woolly.	1790. 6. 9. G. h.
5 imbricàtus. L.T.	(<i>ro.</i>) imbricated.	1794. 4. 7. G. h.
6 diversifòlius. L.T.	(<i>pu.</i>) various-leaved.	1807. — G. h.
SPATA'LLA. L.T.	SPATA'LLA.	Tetrandria Monogynia. R.S.
1 pedunculàta. L.T.	(<i>pu.</i>) long-peduncled.	C. B. S. 1822. 4. 8. G. h.
2 nívea. L.T.	(<i>li.</i>) snowy-bearded.	1806. 4. 8. G. h.
3 ramulòsa. L.T.	(<i>pu.</i>) branching.	1812. — G. h.
4 bracteàta. L.T.	(<i>pu.</i>) bracteate.	— — 5. 9. G. h.
5 prolifera. L.T.	(<i>pu.</i>) proliferous.	1800. 6. 8. G. h.
6 pyramidalis. L.T.	(<i>pu.</i>) pyramidal.	1821. — G. h.
7 incúrvà. L.T.	(<i>pu.</i>) incurved-leaved.	1789. 5. 6. G. h.
8 caudàta. L.T.	(<i>pu.</i>) tailed.	1812. — G. h.
9 Thunbergii. L.T.	(<i>pu.</i>) Thunberg's.	1806. — G. h.
		Thunb. dis. n. 27. t. 4.
		Thunb. dis. n. 22. t. 3.
		sec. ic. t. 2.

PROTEACEÆ.

ADENA'NTHOS. L.T.	ADENA'NTHOS.	Tetrandria Monogynia.				
1 obovâta. L.T.	(li.) obovate-leaved.	N. Holland.	1824.	G. h.	Labil. n. hol. 1. t. 37.
2 cuneâta. L.T.	(li.) wedge-leaved.	—	—	G. h.	— v. 1. t. 36.
3 sericea. L.T.	(li.) silky-leaved.	—	—	G. h.	— v. 1. t. 38.
CONOSPERMUM. L.T.	CONOSPERMUM.	Triandria Monogynia.				
1 ellipticum. L.T.	(wh.) elliptic-leaved.	N. S. W.	1822.	5. 7.	G. h.	
2 taxifolium. L.T.	(st.) Yew-leaved.	—	1823.	4. 6.	G. h.	Bot. mag. t. 2724.
3 ericifolium. L.T.	(bh.) Heath-leaved.	—	1820.	—	G. h.	— t. 2850.
4 longifolium. L.T.	(wh.) long-leaved.	—	1824.	G. h.	Sm. exot. bot. 2. t. 82.
5 acinacifolium. Gr.	(wh.) symeter-leaved.	—	1824.	4. 6.	G. h.	
6 tenuifolium. L.T.	(wh.) slender-leaved.	—	—	—	G. h.	
AGAST'ACHYS. L.T.	AGAST'ACHYS.	Tetrandria Monogynia. R.S.				
odorâta. L.T.	(ye.) sweet-scented.	V. Diem. Isl.	1824.	4. 7.	G. h.	
PERSO'ONIA. L.T.	PERSO'ONIA.	Tetrandria Monogynia. R.S.				
1 teretifolia. L.T.	(ye.) round-leaved.	N. Holland.	1824.	5. 6.	G. h.	Linn. trans. 10. t. 16. f. 1.
2 pinifolia. L.T.	(ye.) Pine-leaved.	N. S. W.	1818.	5. 7.	G. h.	Lab. n. holl. 1. t. 45.
3 juniperina. L.T.	(ye.) Juniper-leaved.	V. Diem. Isl.	1823.	—	G. h.	Lodd. bot. cab. t. 327.
4 hirsuta. L.T.	(ye.) hairy.	N. S. W.	1800.	—	G. h.	
5 mollis. L.T.	(ye.) soft.	—	1818.	—	G. h.	Bot. mag. t. 760.
6 linearis. L.T.	(ye.) linear-leaved.	—	1794.	7. 8.	G. h.	
7 pallida. Gr.	(st.) pale.	N. S. W.	1824.	9. 10.	G. h.	
8 lucida. L.T.	(ye.) shining.	—	—	6. 8.	G. h.	Lodd. bot. cab. 922.
9 flexifolia. L.T.	(ye.) bending-leaved.	—	1822.	—	G. h.	
10 chamaepitys. F.T.	(ye.) ground-pine.	N. S. W.	1824.	—	G. h.	
11 scabra. L.T.	(st.) rough.	N. Holland.	—	—	G. h.	
12 nutans. L.T.	(ye.) nodding.	N. S. W.	1821.	—	G. h.	
13 pruinosa. M.C.	(ye.) frosted.	N. Holland.	1824.	—	G. h.	
14 lanceolata. L.T.	(st.) spear-leaved.	—	1791.	6. 7.	G. h.	Andr. deposit. t. 74.
β latifolia. B.P.	(st.) broad-leaved.	—	—	—	G. h.	— t. 280?
15 salicina. L.T.	(st.) Willow-leaved.	—	1795.	6. 7.	G. h.	Cavan. ic. 4. t. 389?
16 ferruginea. L.T.	(ye.) ferruginous.	—	1818.	—	G. h.	Sm. exot. bot. 2. t. 83.
BRAEJUM. L.T.	AFRICAN ALMOND.	Polygamia Monococcia. L.				
stellatifolium. L.T.	common.	C. B. S.	1731.	3. 4.	G. h.	Hou. n. h. par. 2. t. 6. f. 37.
GREVILLEA. L.T.	GREVILLEA.	Tetrandria Monogynia. R.S.				
1 punicea. L.T.	(sc.) scarlet-flowered.	N. S. W.	1822.	4. 9.	G. h.	Bot. reg. t. 1319.
2 dubia. L.T.	(ro.) doubtful.	—	—	—	G. h.	Bot. mag. t. 862.
3 sericea. L.T.	(ro.) silky.	—	1790.	—	G. h.	Andr. deposit. t. 272.
4 linearis. L.T.	(li.) linear-leaved.	—	—	—	G. h.	Bot. mag. t. 2661.
α incarnata. B.M.	(pi.) flesh-coloured.	—	—	—	G. h.	Lodd. bot. cab. t. 858.
β alba. B.C.	(bh.) white-flowered.	—	—	—	G. h.	
5 stricta. L.T.	(li.) upright.	—	1822.	—	G. h.	Swt. flor. aust. t. 30.
6 riparia. L.T.	(li.) river-side.	—	1791.	4. 5.	G. h.	Lodd. bot. cab. t. 1003.
7 rosmarinifolia. F.T.	(li.) Rosemary-leaved.	—	1824.	—	G. h.	Swt. flor. aust. t. 7.
8 parviflora. L.T.	(st.) small-flowered.	—	1824.	—	G. h.	Swt. flor. aust. t. 55.
9 juniperina. L.T. (gr. st.)	Juniper-like.	—	1821.	—	G. h.	Lodd. bot. cab. t. 857.
10 concinna. L.T. (st. ro.)	pretty.	N. Holland.	1824.	3. 8.	G. h.	Swt. flor. aust. t. 38.
11 arenaria. L.T.	(li.) sand.	N. S. W.	1803.	4. 9.	G. h.	Hook. ex. flor. t. 216.
12 montana. L.T.	(gr.) mountain.	—	—	—	G. h.	
13 acuminata. L.T.	(gr.) acute-leaved.	—	1805.	—	G. h.	
14 cinerea. L.T.	(gr.) grey.	—	1821.	—	G. h.	
15 mucronulata. L.T.	(gr.) mucronulate.	—	1809.	—	G. h.	
16 Baueri. L.T.	(bh.) Bauer's.	—	1823.	4. 8.	G. h.	
pubescens. Hooker, et Lodd. bot. cab. t. 1229,						
17 sulphurea. F.T.	(st.) sulphur-coloured.	—	1823.	—	G. h.	
18 sphacelata. L.T.	(br.) sphacelate.	—	1825.	—	G. h.	
19 phylloides. L.T.	(br.) Phyllica-like.	—	—	—	G. h.	
20 collina. K.P.	(br.) hill.	—	1812.	—	G. h.	
21 buxifolia. L.T.	(pu.) Box-leaved.	—	1790.	2. 9.	G. h.	Bot. reg. t. 443.
22 podalyriæfolia. c.c. (pu.)	Podalyria-leaved.	N. Holland.	1821.	5. 8.	I. h.	
23 podocarpifolia.	Podocarpus-leav'd.	—	1823.	I. h.	
24 berberifolia. c.c.	Berry-leaved.	—	—	—	I. h.	
25 acanthifolia. F.T. (gr.)	Acanthus-leaved.	—	—	—	I. h.	Bot. mag. t. 2807.
26 Gaudichaudii. B.	(ye.) Gaudichaud's.	—	1823.	I. h.	
27 trifurcata. c.c.	three-forked.	—	—	—	I. h.	
28 asplenifolia. L.T. (st.)	Spleenwort-leaved.	N. S. W.	1806.	I. h.	
29 Flindersii. M.C.	Flinders's.	N. Holland.	1823.	I. h.	
H'AKEA. L.T.	H'AKEA.	Tetrandria Monogynia. R.S.				
1 pugioniformis. L.T. (wh.)	dagger-fruited.	N. S. W.	1796.	5. 6.	G. h.	Lodd. bot. cab. t. 353.
2 pârilis. K.P.	(wh.) matched.	V. Diem. Isl.	—	—	G. h.	

3	epiglottis. L.T.	(wh.) curved-fruited.	V. Diem. Isl.	1822.	5. 6. G. h.	Lab. nov. hol. 1. t. 40.
4	nodosa. L.T.	(wh.) knot-podded.	N. Holland.	1824.	— G. h.	
5	flexilis. L.T.	(wh.) bending.	—	—	— G. h.	
6	obliqua. L.T.	(wh.) oblique-flowered.	N. Holland.	1803.	6. 9. G. h.	
7	subulata. ψ.	(wh.) awl-leaved.	—	1824.	— G. h.	
8	petrophiloides. ψ. (wh.)	Petrophila-like.	—	1825.	— G. h.	
9	lissopérrma. L.T.	(wh.) smooth-seeded.	V. Diem. Isl.	1824.	— G. h.	
10	gibbosa. L.T.	(st.) gibbos-fruited.	N. S. W.	1790.	5. 6. G. h.	Cavan. ic. 6. t. 534.
11	aciculáris. L.T.	(wh.) needle-leaved.	—	—	— G. h.	Venten. malm. t. 111.
	<i>Cónchium aciculare.</i> V.					
12	suavéolens. L.T.	(wh.) sweet-scented.	N. Holland.	1803.	11. 1. G. h.	
13	Lamberti. N.L.F.	(wh.) Lambert's.	—	1823.	— G. h.	
14	microcárpa. L.T.	(wh.) small-fruited.	V. Diem. Isl.	1818.	4. 8. G. h.	Bot. reg. t. 475.
15	trifurcata. L.T.	(wh.) trifurcate.	N. Holland.	1824.	— G. h.	
16	echinata. ψ.	(wh.) bristly-podded.	—	1825.	— G. h.	
17	angustifolia. ψ.	(wh.) narrow-leaved.	—	1824.	— G. h.	
18	lineáris. L.T.	(wh.) linear-leaved.	—	—	— G. h.	Swt. flor. aust. t. 43.
19	florida. L.T.	(wh.) many-flowered.	—	1803.	5. 6. G. h.	Bot. mag. t. 2579.
20	ilicifolia. L.T.	(wh.) Holly-leaved.	—	—	7. 9. G. h.	
21	nitida. L.T.	(wh.) glossy.	—	—	6. 8. G. h.	Bot. mag. t. 2246.
22	amplexicaulis. L.T.	(wh.) smooth-stalked.	—	—	— G. h.	
23	prostrata. L.T.	(wh.) pubescent.	—	—	— G. h.	
24	cératophylla. L.T.	(wh.) horn-leaved.	—	—	— G. h.	
25	acanthophylla. L.en.	(w.) Acanthus-leaved.	—	—	— G. h.	
26	undulata. L.T.	(wh.) wave-leaved.	—	—	6. 8. G. h.	
27	repanda. ψ.	(wh.) repand.	—	1824.	— G. h.	
28	propinqua. F.T.	(wh.) related.	—	1823.	— G. h.	
29	oleifolia. L.T.	(wh.) Olive-leaved.	—	1794.	6. 7. G. h.	
30	saligna. L.T.	(wh.) Willow-leaved.	—	1791.	3. 7. G. h.	Swt. flor. aust. t. 27.
31	marginata. L.T.	(wh.) marginate.	—	1824.	4. 7. G. h.	
32	ruscifolia. L.T.	(st.) Ruscus-leaved.	—	—	— G. h.	
33	cineræa. L.T.	(wh.) hoary-leaved.	—	1803.	6. 7. G. h.	Lab. nov. holl. 1. t. 39.
34	canescens. L.en.	(wh.) canescent.	N. S. W.	1800.	6. 8. G. h.	
35	dactyloides. L.T.	(wh.) nerved-leaved.	—	1790.	— G. h.	Cavan. ic. 6. t. 535.
36	elliptica. L.T.	(wh.) oval-leaved.	N. Holland.	1794.	— G. h.	
37	ferruginea. S.F.A.	(wh.) rusty-stemmed.	—	1824.	4. 8. G. h.	Swt. flor. aust. t. 45.
38	cucullata. ψ.	(wh.) hollow-leaved.	—	—	— G. h.	
39	clavata. L.T.	(wh.) clavate.	—	1824.	— G. h.	Lab. nov. holl. 1. t. 41.
	EMBO'THRIUM.	EMBO'THRIUM.	Tetrandria Monogynia.			
	stroblinum. Lab.	(st.) cone-flowered.	N. Holland.	1824.	3. 6. G. h.	
	LAMBE'R'TIA. L.T.	LAMBE'R'TIA.	Tetrandria Monogynia. R.S.			
1	uniflora. L.T.	(re.) single-flowered.	N. Holland.	1824.	— G. h.	Bauer ic. ined.
2	inermis. L.T.	(or.) unarmed.	—	—	— G. h.	— ic. ined.
3	formosa. L.T.	(ro.) handsome.	N. S. W.	1788.	6. 8. G. h.	Bot. reg. t. 528.
4	echinata. L.T.	(ro.) lobe-leaved.	N. Holland.	1824.	— G. h.	Bauer ic. ined.
	XYLO'MELUM. L.T.	WOODEN-PEAR-TREE.	Tetrandria Monogynia. R.S.			
	pyriforr. L.T.	(wh.) prickly-leaved.	N. S. W.	1789.	6. 8. G. h.	Cavan. ic. 6. t. 536.
	RHOP'ALA. L.T.	RHOP'ALA.	Tetrandria Monogynia. R.S.			
1	montana. L.T.	(st.) mountain.	Guiana.	1823.	4. 6. S. h.	Aub. gui. 1. t. 32.
2	nitida. L.T.	(st.) glossy.	—	1826.	— S. h.	Rudge pl. gui. 1. t. 39.
3	dentata. L.T.	(st.) tooth-leaved.	—	1802.	5. 8. S. h.	
4	sessilifolia. L.T.	(st.) sessile-leaved.	—	1803.	— S. h.	Rudg. gui. 1. t. 31.
	TELOP'EA. L.T.	WARATAH.	Tetrandria Monogynia. R.S.			
	speciosissima. L.T.	(cr.) splendid.	N. S. W.	1789.	5. 7. G. h.	Bot. mag. t. 1128.
	<i>Embóthrium speciosissimum.</i> B.M.					
	LOM'ATIA. L.T.	LOM'ATIA.	Tetrandria Monogynia. R.S.			
1	silaefolia. L.T.	(wh.) cut-leaved.	N. S. W.	1792.	6. 8. G. h.	Bot. mag. t. 1272.
2	tinctoria. L.T.	(st.) various-leaved.	V. Diem. Isl.	1822.	— G. h.	Lab. n. holl. 1. t. 42-43.
3	polymorpha. L.T.	(st.) variable.	—	1823.	— G. h.	
4	ilicifolia. L.T.	(st.) Holly-leaved.	N. Holland.	1824.	— G. h.	Bot. reg. t. 442.
5	longifolia. L.T.	(wh.) long-leaved.	N. S. W.	1816.	5. 8. G. h.	Flor. peruv. 1. t. 94. a.
6	dentata. L.T.	(st.) tooth-leaved.	Chile.	1824.	— G. h.	
	STENOCA'RPUS. L.T.	STENOCA'RPUS.	Tetrandria Monogynia. R.S.			
	salignus. L.T.	(wh.) sweet-scented.	N. S. W.	1815.	6. 9. G. h.	Bot. reg. t. 441.
	BOTRYC'ERAS. W.	BOTRYC'ERAS.	Tetrandria Monogynia. R.S.			
	laurinum. w.	(gr.) Laurel-like.	N. Holland.	1820.	6. 9. G. h.	W.m.b. 1811. t. 10. f. 10.
	BA'NKSIA. L.T.	BA'NKSIA.	Tetrandria Monogynia. R.S.			
1	pulchella. L.T.	(st.) small-flowered.	N. Holland.	1803.	2. 8. G. h.	
2	sphaerocarpa. L.T.	(st.) round-fruited.	—	1803.	— G. h.	

PROTEACEÆ.

3 nūtans. L.T.	(st.) nodding-flowered.	N. Holland.	1803.	4. 6. G. h.
4 ericifolia. L.T.	(st.) Heath-leaved.	N. S. W.	1788.	1. 12. G. h.
5 spinulōsa. L.T.	(st.) spiny-leaved.	——	——	12. 5. G. h.
6 collina. L.T.	(st.) hill.	——	1800. G. h.
7 occidentalis. L.T.	(st.) west-coast.	N. Holland.	1803.	4. 8. G. h.
8 littorālis. L.T.	(ye.) sea-side.	——	——	4. 7. G. h.
9 margināta. L.T.	(gr.) various-leaved.	N. S. W.	1804.	5. 8. G. h.
10 australis. L.T.	(gr.) southern.	V. Diem. Isl.	1816.	—— G. h.
marcēscens. Hooker bot. mag. 2803. nec aliorum.				
11 insulāris. L.T.	(er.) Island.	——	1823. G. h.
12 Cunninghamii. s.s. (st.)	Cunningham's.	N. Holland.	1822.	—— G. h.
13 integrifolia. L.T.	(st.) entire-leaved.	N. S. W.	1788.	5. 8. G. h.
14 compár. L.T.	(st.) related.	N. Holland.	1823. G. h.
15 verticillāta. L.T.	(st.) whorl-leaved.	——	1794.	7. 11. G. h.
16 coccinea. L.T.	(sc.) scarlet-flowered.	——	1803. G. h.
17 paludōsa. L.T.	(st.) marsh.	N. S. W.	1805.	1. 4. G. h.
18 intermédia. ψ.	(st.) intermediate.	N. Holland.	1824. G. h.
19 oblongifolia. L.T.	(st.) oblong-leaved.	N. S. W.	1788.	5. 8. G. h.
20 latifolia. L.T.	(st.) broad-leaved.	——	1802.	—— G. h.
21 marcēscens. L.T.	(st.) short-leaved.	N. Holland.	1794.	1. 12. G. h.
præmōrsa. Andr. rep. t. 258. <i>asplenifolia</i> . K.P.				
22 attenuāta. L.T.	(st.) attenuated, smooth-fld.	——	1794.	1. 10. G. h.
23 serrāta. L.T.	(st.) saw-leaved.	N. S. W.	1788.	7. 9. G. h.
undulāta. Bot. reg. t. 1316.				
24 'æmula. L.T.	(st.) deeply-sawed.	——	——	1. 6. G. h.
25 dentāta. L.T.	(st.) tooth-leaved.	N. Holland.	1822. S. h.
26 querifolia. L.T.	(st.) Oak-leaved.	——	1805. G. h.
27 dryandroïdes. s.F.A. (br.)	Dryandra-like.	——	1824.	3. 6. G. h.
28 speciōsa. L.T.	(wh.) long-leaved.	——	1805.	5. 8. G. h.
29 grāndis. L.T.	(st.) great-flowered.	——	1794. G. h.
30 prostrāta. ψ.	(st.) prostrate.	——	1824. G. h.
31 rēpens. L.T.	(st.) creeping.	——	1803. G. h.
32 ilicifolia. L.T.	(st.) Holly-leaved.	——	1824. G. h.
DRYAN'DRA. L.T. DRYAN'DRA. Tetrandria Monogynia. R.S.				
1 floribunda. L.T.	(ye.) many-flowered.	N. Holland.	1803.	1. 12. G. h.
2 cuneāta. L.T.	(st.) wedge-leaved.	——	——	11. 2. G. h.
<i>a brevifolia.</i>	(st.) short-leaved.	——	——	—— G. h.
<i>β longifolia.</i>	(st.) long-leaved.	——	——	—— G. h.
3 armata. L.T.	(ye.) acute-leaved.	——	——	1. 12. G. h.
4 falcāta. L.T.	(ye.) falcate-leaved.	——	1824. G. h.
5 formōsa. L.T.	(or.) Apricot-scented.	——	1803.	1. 12. G. h.
6 mucronulāta. L.T.	(ye.) pointed-leaved.	——	1824. G. h.
7 plumōsa. L.T.	(ye.) feathered.	——	1803.	1. 8. C. I.
8 obtusa. L.T.	(ye.) blunt-leaved.	——	—— G. h.
9 nivea. L.T.	(ye.) white-leaved.	——	——	7. 9. G. h.
10 longifolia. L.T.	(ye.) long-leaved.	——	1805.	1. 12. G. h.
11 Baxtēri. ψ.	(ye.) Baxter's.	——	1824. G. h.
12 tenuifolia. L.T.	(ye.) slender-leaved.	——	1803.	3. 5. G. h.
13 nervosa. s.F.A.	(ye.) nerved-leaved.	——	1824.	3. 12. G. h.
14 pteridifolia. L.T.	(ye.) Pteris-leaved.	——	——	— G. h.

ORDO CLXXIII.

THYMELÆÆ. Brown prodr. p. 358.

DI'RCA. W.	LEATHER-WOOD. Octandria Monogynia. L.			
palustris. w.	(ye.) marsh.	Virginia.	1750.	3. 4. H. h.
LAGETTA. J.	LACE-BARK. Octandria Monogynia. S.S.			Bot. reg. t. 292.
lintearia. j.	(wh.) Jamaica.	Jamaica.	1793.	12. 1. S. h.
DA'PHNE. W.	DAPHNE. Octandria Monogynia. L.			Lam. ill. t. 289.
1 Mezēreum. w. (re.wh.)	Mezereon.	England.	Eng. bot. v. 20. t. 1381.
a álbum.	(wh.) white-flowered.	——	— H. h.
β rubrum.	(re.) red-flowered.	——	— H. h.
γ autumnale.	(re.) autumnal-flowered.	——	8. 12. H. h.

THYMELEÆ.

2 Laurèola. w. (gr.)	spurge-laurel.	Britain.	1. 3. H. $\frac{1}{2}$.	Eng. bot. v. 2. t. 119.
3 póntica. w. (gr.ye.)	Pontic.	Pontus.	1759.	4. 5. H. $\frac{1}{2}$.	Bot. mag. t. 1282.
4 tinifòlia. w. (st.)	Bonace-bark.	Jamaica.	1733.	5. 7. S. $\frac{1}{2}$.	Lodd. bot. cab. t. 150.
5 Gnídiu. w. (wh.)	Flax-leaved.	Spain.	1597.	6. 8. H. $\frac{1}{2}$.	Bot. mag. t. 1587.
6 odòra. w. (bh.)	sweet-scented.	China.	1771.	12. 3. G. $\frac{1}{2}$.	Swt. br. fl. gar. t. 200.
7 hy'brida. B.F.G. (ro.)	Dauphin.	Hybrid.	1826.	1. 6. F. $\frac{1}{2}$.	Lodd. bot. cab. t. 66.
8 alpina. w. (wh.)	Alpine.	Italy.	1759.	5. 7. H. $\frac{1}{2}$.	Bot. mag. t. 1875.
9 altaïca. w. (wh.)	Altaic.	Siberia.	1796.	4. 5. H. $\frac{1}{2}$.	——— t. 313.
10 Cneòrum. w. (ro.)	trailing.	Austria.	1752.	4. 9. H. $\frac{1}{2}$.	Bot. mag. t. 428.
11 strìata. s.s. (pi.)	streaked.	Hungary.	— H. $\frac{1}{2}$.	
12 collìna. s.s. (re.)	hairy.	Italy.	1752.	1. 6. H. $\frac{1}{2}$.	Bot. mag. t. 1917.
13 sericea. s.s. (li.)	silky-leaved.	Levant.	1818.	— H. $\frac{1}{2}$.	Lodd. bot. cab. t. 719.
14 oleoides. s.s.	Olive-leaved.	Crete.	1815.	1. 12. H. $\frac{1}{2}$.	Pluk. alm. t. 229. f. 2.
15 napolitana. B.C. (re.)	Neapolitan.	Italy.	1823.	1. 6. H. $\frac{1}{2}$.	——— t. 318. f. 6.
<i>collina</i> β <i>neapolitana</i> . B.R. 822.					
16 Thymel'æa. w. (st.)	smooth-leaved.	Spain.	1815.	2. 4. H. $\frac{1}{2}$.	
17 Tártón-ráira. w. (st.)	silvery-leaved.	France.	1640.	5. 7. H. $\frac{1}{2}$.	
GN'DIA. L. <i>GNI'DIA. Octandria Monogynia. L.</i>					
1 pinifòlia. w. (wh.)	Pine-leaved.	C. B. S.	1768.	5. 6. G. $\frac{1}{2}$.	Bot. reg. t. 19.
2 radiàta. B.C. (wh.)	radiate.	———	1818.	— G. $\frac{1}{2}$.	Lodd. bot. cab. t. 29.
3 imberbis. H.K. (st.)	smooth-scaled.	———	1792.	8. 4. G. $\frac{1}{2}$.	Bot. mag. t. 1463.
4 simplex. H.K. (ye.)	Flax-leaved.	———	1786.	5. 6. G. $\frac{1}{2}$.	——— t. 812.
5 juniperifòlia. w. (wh.)	Juniper-leaved.	———	1822.	— G. $\frac{1}{2}$.	
6 capitâta. w. (wh.)	purple-twiggled.	———	1788.	6. 7. G. $\frac{1}{2}$.	
7 imbricâta. w. (st.)	imbricated.	———	1822.	4. 8. G. $\frac{1}{2}$.	Bot. reg. t. 757.
<i>denudata</i> . B.R.					
8 oppositifòlia. H.K. (ye.)	opposite-leaved.	———	1783.	5. 7. G. $\frac{1}{2}$.	Bot. reg. t. 2.
9 tomentosa. s.s. (st.)	tomentose.	———	1822.	— G. $\frac{1}{2}$.	
10 sericea. w. (ye.)	silky.	———	1786.	— G. $\frac{1}{2}$.	Andr. deposit. t. 225.
<i>oppositifolia</i> . A.R. nec aliorum.					
LACHN'EA. W. <i>LACHN'EA. Octandria Monogynia. L.</i>					
1 buxifòlia. Lam.	(wh.) green box-leaved.	C. B. S.	1800.	5. 7. G. $\frac{1}{2}$.	Bot. mag. t. 1657.
2 glauca. H.K.	(wh.) glaucous.	———	———	— G. $\frac{1}{2}$.	——— t. 1658.
3 purpurea. w. (pu.)	purple-flowered.	———	———	— G. $\frac{1}{2}$.	——— t. 1594.
4 eriocéphala. w. (wh.)	woolly-headed.	———	1793.	6. 7. G. $\frac{1}{2}$.	——— t. 1295.
5 conglomerata. w. (wh.)	clustered.	———	1773.	— G. $\frac{1}{2}$.	
PASSER'INA. W. <i>PASSER'INA. SPARROW-WORT. Octandria Monogynia. L.</i>					
1 unifòlia. w. (wh.)	one-flowered.	C. B. S.	1759.	4. 7. G. $\frac{1}{2}$.	Wend. obs. t. 2. f. 18.
2 láxa. w. (wh.)	loose.	———	1804.	6. 7. G. $\frac{1}{2}$.	Lodd. bot. cab. t. 755.
3 tenuifòlia. w. (st.)	slender-flowered.	———	1812.	— G. $\frac{1}{2}$.	
4 filiformis. w. (st.)	filiform.	———	1752.	6. 8. G. $\frac{1}{2}$.	Wend. obs. t. 2. f. 15.
5 capitâta. w. (wh.)	headed.	———	1789.	6. 10. G. $\frac{1}{2}$.	——— t. 2. f. 17.
6 hirsuta. w. (st.)	hairy.	S. Europe.	1759.	— G. $\frac{1}{2}$.	Bot. mag. t. 1949.
7 striâta. Lam.	(wh.) striate.	C. B. S.	1822.	— G. $\frac{1}{2}$.	
8 ericoïdes. w. (st.)	Heath-like.	———	———	— G. $\frac{1}{2}$.	
9 ciliâta. w. (wh.)	fringed.	———	1819.	— G. $\frac{1}{2}$.	Seb. mus. 2. t. 12. f. 9.
10 grandifòra. w. (wh.)	great-flowered.	———	1789.	5. 6. G. $\frac{1}{2}$.	Bot. mag. t. 292.
11 spicâta	(wh.) spiked.	———	1787.	5. 7. G. $\frac{1}{2}$.	Lodd. bot. cab. t. 311.
12 anthylloides. w. (st.)	Anthyllus-like.	———	1816.	— G. $\frac{1}{2}$.	
STELL'ERA. W. <i>STELL'ERA. Octandria Monogynia. L.</i>					
1 Passerina. w. (ye.)	Flax-leaved.	S. Europe.	1759.	7. 8. H. \odot .	Jacq. ic. 1. t. 68.
<i>Passerina annua</i> . s.s.					
2 altaïca. P.S.	(wh.) Altay.	Altay.	1824.	— H. \odot .	
3 dichótoma. F.	(wh.) forked.	Dahuria.	———	— H. \odot .	
D'AIS. L. <i>D'AIS. Decandria Monogynia. L.</i>					
1 cotinifòlia. w. (pi.)	Cotinus-leaved.	C. B. S.	1776.	6. 7. G. $\frac{1}{2}$.	Bot. mag. t. 147.
STRUTH'iola. L. <i>STRUTH'iola. Tetrandria Monogynia. L.</i>					
1 virgâta. H.K.	(re.) twiggy.	C. B. S.	1779.	4. 8. G. $\frac{1}{2}$.	Andr. deposit. t. 139.
2 ciliâta. A.R.	(st.) ciliated.	———	———	— G. $\frac{1}{2}$.	——— t. 149.
3 lateriflòra. R.S.	(wh.) side-flowering.	———	1816.	— G. $\frac{1}{2}$.	
4 pubéscent. R.S.	(re.) downy.	———	1790.	— G. $\frac{1}{2}$.	Bot. mag. t. 1212.
5 longiflòra. R.S.	(wh.) long-flowered.	———	1824.	— G. $\frac{1}{2}$.	Lam. ill. t. 78.
6 glâbra. R.S.	(wh.) upright smooth.	———	1798.	4. 9. G. $\frac{1}{2}$.	Wendl. obs. t. 2. f. 10.
<i>erecta</i> . W. <i>stricta</i> . Donn.					
7 juniperina. R.S.	(wh.) drooping.	———	1758.	— G. $\frac{1}{2}$.	Bot. mag. t. 222.
<i>erecta</i> . B.M. nec aliorum.					
8 angustifòlia. R.S.	(st.) narrow-leaved.	———	1816.	— G. $\frac{1}{2}$.	
9 incâna. B.C.	(ye.) hoary.	———	1802.	7. 8. G. $\frac{1}{2}$.	Lodd. bot. cab. t. 11.
10 imbricâta. R.S.	(ye.) tiled-leaved.	———	1794.	4. 8. G. $\frac{1}{2}$.	Andr. deposit. t. 113.

THYMELÆÆ.

11 ovata. R.S.	(wh.) oval-leaved.	C. B. S.	1792.	2. 6. G. h.	Andr. deposit. t. 119.
12 tomentosa. R.S.	(wh.) downy-leaved.	_____	1799.	8. 9. G. h.	_____ t. 334.
PIMEL'EA. B.P.	PIMEL'EA. Diandria Monogynia. R.S.				
1 linifolia. B.P.	(wh.) flax-leaved.	N. S. W.	1793.	2. 8. G. h.	
2 paludosa. B.P.	(wh.) marsh.	_____	1826.	— G. h.	Bot. mag. t. 891.
3 glauca. B.P.	(wh.) glaucous-leaved.	_____	1822.	— G. h.	Linn. trans. 10. t. 13. f. 2.
4 ligustrina. B.P.	(wh.) Privet-like.	_____	1823.	— G. h.	Lab. nov. holl. I. t. 3.
5 decussata. B.P.	(re.) cross-leaved.	_____	—	4. 8. G. h.	Swt. flor. aust. t. 8.
ferruginea. Lab.					
6 rosea. B.P.	(ro.) rose-coloured.	N. Holland.	1800.	3. 9. G. h.	Bot. mag. t. 1458.
7 linoïdes. F.T.	(wh.) flax-like.	N. S. W.	1824.	— G. h.	Linn. trans. 10. t. 14. f. 1.
8 filamentosa. L.T.	(wh.) filamentose.	N. Holland.	1826.	— G. h.	Lodd. bot. cab. t. 179.
9 pauciflora. B.P.	(ye.) few-flowered.	V. Diem. Isl.	1812.	— G. h.	Bot. reg. t. 1268.
10 humilis. B.P.	(wh.) dwarf.	_____	1825.	6. 9. G. h.	
11 clavata. B.P.	(ye.) clubbed.	N. Holland.	1826.	3. 6. G. h.	
12 incana. B.P.	(st.) hoary.	V. Diem. Isl.	—	4. 8. G. h.	
13 drupacea. B.P.	(wh.) drupe-bearing.	_____	1820.	— G. h.	Swt. flor. aust. t. 52.
14 spicata. B.P.	(wh.) spiked.	N. S. W.	1824.	— G. h.	Linn. trans. 10. t. 14. f. 2.
? TR'OPHIS. W.	RAMOON-TREE. Diœcia Tetrandria. W.				Brown. jam. t. 37. f. 1.
1 americana. w.	(gr.) American.	W. Indies.	1789.	4. 5. S. h.	
2 áspera. w.	(gr.) rough-leaved.	E. Indies.	1802.	— S. h.	
3 spinosa. s.s.	(gr.) spiny.	_____	1826.	... S. h.	

ORDO CLXXIV.

OSYRIDEÆ. Brown.

OS'YRIS. W.	POET's CASSIA. Diœcia Triandria. L.				
ábla. w.	(wh.) white.	S. Europe.	1739.	... G. h.	Lam. ill. t. 802.
EXOCA'RPOS. B.P.	EXOCA'RPOS. Pentandria Monogynia.				
1 cupressiformis. B.P.	(wh.) Cypress-like.	N. Holland.	1820.	... G. h.	Labill. voy. 1. t. 14.
2 humifusa. B.P.	(gr.) low-procumbent.	V. Diem. Isl.	1822.	... G. h.	
3 stricta. B.P.	(gr.) upright.	N. S. W.	—	... G. h.	

ORDO CLXXV.

SANTALACEÆ. Brown prodr. p. 350.

SA'NTALUM. B.P.	SANDAL-WOOD. Tetrandria Monogynia. W.				
1 ábum. F.I.	(ye. pu.) white.	E. Indies.	1804.	... S. h.	Rumph. amb. 2. t. 11.
2 myrtifolium. F.I.	(li.) Myrtle-leaved.	_____	1819.	— G. h.	Roxb. corom. 1. t. 2.
3 obtusifolium. B.P.	blunt-leaved.	N. S. W.	1823.	— G. h.	
FUS'ANUS. B.P.	FUS'ANUS. Tetrandria Monogynia.				
compræssus. H.K.	(gr.) flat-stalked.	C. B. S.	1776.	... G. h.	Berg. cap. 38. t. 1. f. 1.
LEPTOM'ERIA. B.P.	LEPTOM'ERIA. Pentandria Monogynia. R.S.				
1 ácida. B.P.	(wh.) acid.	N. S. W.	1823.	— G. h.	
2 Billardieri. B.P.	(wh.) Labillardiere's.	_____	—	— G. h.	Lab. nov. hol. 1. t. 93.
	<i>Thesium drupaceum</i> . Lab.				
TH'ESIUM. R.S.	THE'SIUM. Pentandria Monogynia. L.				
1 linophyllum. B.S.	(gr.) common.	England.	...	6. 7. H. 4.	Eng. bot. v. 4. t. 247.
2 intermedium. R.S.	(wh.) intermediate.	Germany.	1818.	— H. 4.	Schkuhr hand. t. 51.
3 montanum. R.S.	(wh.) mountain.	Europe.	...	— H. 4.	Hayn. sc. j. t. 119.
4 alpinum. R.S.	(wh.) Alpine.	_____	1814.	— H. 4.	Jacq. aust. 5. t. 416.
5 ramosum. R.S.	(wh.) branching.	_____	1824.	— H. 4.	Hayn. sc. j. p. 30. t. 7.
6 bracteatum. R.S.	(gr.) blunt-leaved.	Germany.	—	— H. 4.	— l. c. t. 7.
7 ? amplexicaulé. R.S.	(gr.) stem-clasping.	C. B. S.	1787.	— G. h.	Pluk. mant. t. 342. f. 1.
COMA'NDRA. N.	COMA'NDRA. Pentandria Monogynia. N.				
umbellata. n.	(wh.) umbellied.	N. America.	1782.	6. H. 4.	
	<i>Thesium umbellatum</i> . w.				

HAMILTONIA. W.	OIL-NUT.	Polygamia Dioccia. W.				
oleífera. w.	(gr.ye.)	Olive-bearing.	N. America.	1800. G. h.	Pursh. amer. I. t. 13.
<i>Pyrułaria pùbera.</i> M.						
NYSSA. R.S.	TUPELO.	Polygamia Dioccia. L.				
1 villòsa. R.S.	(gr.)	hairy-leaved.	N. America.	1806.	5. H. h.	Mich. arb. 2. t. 21.
2 aquática. R.S.	(gr.)	water.	—	1739.	4. 5. H. h.	— 2. t. 22.
biflòra. w.						
3 canticans. R.S.	(gr.)	Ogechee-lime.	—	1812. H. h.	Mich. arb. 2. t. 20.
4 capitata. H.K.	(gr.)	headed.	Carolina.	1806. H. h.	
5 tomentosa. R.S.	(gr.)	tomentose.	—	1812. H. h.	Mich. arb. 2. t. 19.
6 denticulata. R.S.	(gr.)	toothed.	N. America.	1795.	5. 6. H. h.	Catesb. car. I. t. 60.

ORDO CLXXVI.

ELÆAGNEÆ. *Don prodr. p. 67.*

HIPPO'PHAE. L.	SEA-BUCKTHORN.	Dioccia Tetrandria. L.				
1 rhamnoïdes. W.	(st.) common.	England.	4. 5. H. h.	Eng. bot. t. 425.	
2 salicifolia. D.P.	(st.) Willow-leaved.	Nepaul.	1819. H. h.		
SHEPHE'R'DIA. N.	SHEPHE'R'DIA.	Dioccia Octandria. N.				
1 argéntea. N.	(st.) silvery.	Missouri.	1818. H. h.		
<i>Hippóphæ argéntea.</i> Ph.						
2 canadensis. N.	(st.) Canadian.	—	1759.	4. 5. H. h.		
<i>Hippóphæ canadensis.</i> W.						
ELÆA'GNUS. L.	OLEASTER.	Tetrandria Monogynia. L.				
1 angustifolia. R.S. (wh.)	narrow-leaved.	S. Europe.	1633.	7. H. h.	Bot. reg. t. 1156.	
2 songáricus. F.	(st.) Fischer's.	Siberia.	1821.	— H. h.		
3 orientalis. R.S.	(wh.) oriental.	Levant.	1748.	7. 8. G. h.	Pall. ros. I. t. 5.	
4 spinosa. R.S.	(wh.) spiny.	Egypt.	1826.	— F. h.		
5 argéntea. Ph.	(wh.) silvery.	N. America.	1813.	— H. h.	Wats. dend. brit. t. 16.	
6 arbórea. F.I.	(wh.) tree.	Nepaul.	1819. H. h.		
7 conferta. F.I.	(st.) close-flowered.	—	— H. h.		
8 latifolia. R.S.	(wh.) broad-leaved.	E. Indies.	1712.	7. 8. S. h.	Burm. zeyl. t. 39. f. 2.	
9 triflora. F.I.	(wh.) three-flowered.	Nepaul.	1825. H. h.	— t. 39. f. 1.	
10 umbellata. D.P.	(wh.) umbelled.	—	1829. H. h.	Thunb. jap. t. 14.	

ORDO CLXXVII.

ASARINÆ. *Kunth synops. I. p. 442.*

A'SARUM. L.	ASARABACCA.	Dodecadria Monogynia. L.				
1 europæum. s.s.	(pu.) common.	England.	5. H. ¼.	Eng. bot. v. 16. t. 1083.	
2 canadense. s.s.	(pu.) Canadian.	Canada.	1713.	4. 7. H. ¼.	Swt. br. fl. gar. t. 95.	
3 virginicum. w.	(pu.) Virginian.	Virginia.	1759.	4. 5. H. ¼.	— t. 18.	
4 arifolium. s.s.	(pu.) Arum-leaved.	Carolina.	1818.	5. 6. H. ¼.	Hook. ex flor. t. 40.	
ARISTOL'OCHIA. W.	BIRTHWORT.	Gynandria Hexandria. L.				
1 bilobata. w.	(pu.) two-lobed.	W. Indies.	1822.	6. 8. S. h.	Plum. amer. t. 106.	
2 trilobata. w.	(pu.) three-lobed.	S. America.	1775.	6. 7. S. h.		
3 rígens. w.	(br.) gaping.	—	1820.	— S. h.	Vahl. symb. 2. t. 47.	
4 labiata. B.R.	(br.) lipped.	—	—	5. 9. S. h.	Jacq. ic. pict. t. 224.	
5 pentandra. w.	(pu.) pentandrous.	Cuba.	1828. S. h.		
6 panduriiformis. w.	(pu.) fiddle-leaved.	Caracas.	1823. S. h.		
7 máxima. w.	(pu.) great.	New Spain.	1759.	7. S. h.	Jacq. amer. t. 146.	
8 Siphon. w.	(br.) broad-leaved.	N. America.	1763.	6. 7. H. h.	Bot. mag. t. 534.	
9 tomentosa. B.M.	(ye.) downy.	—	1799.	— H. h.	— t. 1369.	
10 odoratissima. w.	(pu.) sweet-scented.	Jamaica.	1737.	7. S. h.	Sloan. jam. I. t. 104. f. 1.	
11 barbata. w.	(pu.) bearded.	Caracas.	1796.	6. 8. S. h.	Jacq. ic. 3. t. 608.	
12 bilabiata. w.	(pu.) two-lipped.	Cuba.	1829. S. h.		
13 Índica. w.	(pu.) Indian.	E. Indies.	1780.	6. 7. S. h.	Rheed. mal. 8. t. 25.	
14 acuminata. w.	(pu.) taper-pointed.	Mauritius.	1820. S. h.		

15 fœtida. k.s.	(pu.) fetid.	Mexico.	1822.	6. 8. S. h. —. H. et B. 2. t. 114.
16 hastatæ. k.s.	(pu.) halberd-leaved.	Havannah.	— S. h. —. —. 2. t. 116.
17 b'etica. w.	(pu.) Spanish.	Spain.	1596.	5. 6. H. h. —. Moris. s. 12. t. 17. f. 6.
18 glauca. w.	(ye.) glaucous.	Barbary.	1785.	6. 8. G. h. —. Bot. mag. t. 1115.
19 sempervirens. w. (pu.) evergreen.		Candia.	1727.	5. 6. G. h. —. —. t. 1116.
20 longa. w.	(pu.) long-rooted.	S. Europe.	1548.	6. 10. H. 4. Mill. ic. t. 51. f. 2.
21 microphylla. W.en.	(pu.) small-leaved.	1826.	—. G. 4.
22 Serpentaria. w.	(pu.) Snake-root.	N. America.	1632.	6. 7. H. 4. Jacq. schoen. 3. t. 385.
23 sagittata. Muhl.	(pu.) arrow-leaved.	—	1819.	—. H. 4.
24 bracteata. w.	(pu.) bracteate.	E. Indies.	1793.	7. S. 4.
25 Pistolochia. w.	(pu.) small.	S. Europe.	1597.	6. 7. H. 4.
26 rotunda. w.	(pu.) round-rooted.	—	1596.	3. 10. G. 4.
27 pallida. w.	(pa.) pale-flowered.	Italy.	1640.	5. 8. H. 4.
28 hirta. w.	(pu.) hairy.	Chio.	1759.	5. 6. G. 4.
29 Clematitís. w. (ye.) common.		England.	5. 8. H. 4.
30 arboréscens.w. (pu.ye.) tree.		America.	1737.	6. 7. G. h.

ORDO CLXXVIII.

CYTINEÆ. Brogniart.

NEPE'NTHES. L.	PITCHER-PLANT.	Dicæcia Monadelphia. L.
1 distillatòria. w. (gr.)	cylindrical.	Ceylon. 1789. 4. 5. S.w. 4. Burm. zeyl. 42. t. 17.
2 Phyllamphora.w.(gr.)	ventricose.	China. 1822. 6. 8. S.w. 4. Bot. mag. t. 2629.

ORDO CLXXIX.

STACKHOUSEÆ. Brown.

STACKHO'USIA. Lab. STACKHO'USIA. Pentandria Trigynia.

1 linariifolia. F.T.	(st.) Linaria-leaved.	N. S. W. 1823. 3. 6. G. h.
2 spathulata. s.s.	(wh.) spathulate-leaved.	— 1825. 4. 6. G. h.

ORDO CLXXX.

EUPHORBIACEÆ. Juss. gen. 384.

MERCURI'ALIS. W. MERCURY. Dicæcia Enneandria. L.

1 perennis. w.	(gr.) perennial.	Britain.	4. 5. H. 4.	Eng. bot. v. 26. t. 1872.
2 ambigua. w.	(gr.) doubtful.	Spain. 1806.	7. 8. H. (○)	Linn. f. dec. 1. t. 8.
3 annua. w.	(gr.) annual.	Britain.	7. 9. H. (○)	Eng. bot. v. 8. t. 559.
4 elliptica. w.	(gr.) oval-leaved.	Portugal. 1802.	5. 7. G. h.	Venten. cels. t. 12.
5 tomentosa. w.	(gr.) woolly.	Spain. 1640.	7. 9. H. h.	

PEDILA'NTHUS. Neck. SLIPPER-SPURGE. Dodecadria Trigynia. Len.

1 tithymaloides. B.R. (sc.)	Myrtle-leaved.	S. America. 1699.	6. 7. D.S. h.	Bot. reg. t. 837.
myrtifolius. L.en.	Crepidaria myrtifolia. H.S.S.			
2 carinatus.	(sc.) keel-leaved.	W. Indies. 1809.	— D.S. h.	Bot. mag. t. 2514.
Euphorbia carinata. B.M.	Crepidaria carinata. H.S.S.			
3 subcarinatus. H.R. (sc.)	small-keeled.	W. Indies. 1819.	— D.S. h.	
4 canaliculatus. (sc.)	channelled-leaved.	—	— D.S. h.	Lodd. bot. cab. t. 727.
Euphorbia canaliculata. B.C.				
5 padifolius. H.S.	(sc.) Padus-leaved.	— 1699.	— D.S. h.	Dill. elt. t. 288. f. 372.
6 cordellatus. (sc.)	small-heart-leaved.	— 1795.	— D.S. h.	
Crepidaria cordellata. H.S.				

EUPHOR'BIA. W.	SPURGE. Dodecandra Trigynia. L.			
1 uncinatæ. DC. (gr.ye.) twin-spined.	C. B. S.	1795.	6. 8. D.G. h.	DC. plant. gras. t. 151.
<i>Scolopéndria</i> . H.s. procumbens. Meerb. ic. rar. t. 55. ex H.P.				
2 squarrosa. H.P. (gr.ye.) squarrose.	C. B. S.	1825.	— D.G. h.	
3 trigona. h.s. (gr.ye.) upright 3-sided. E. Indies.		1768.	7. 8. D.S. h.	
4 antiquorum. H.s.(gr.ye.) spreading 3-sided.		1688.	— D.S. h.	Comm. hort. 1. t. 12.
5 láctea. H.s. (gr.ye.) marbled.		1804.	— D.S. h.	
6 grandidéns. H.P. (gr.ye.) large-toothed.	C. B. S.	1820.	— D.G. h.	
7 canariénis. w. (br.re.) Canary.	Canaries.	1697.	3. 4. D.G. h.	DC. plant. gras. t. 140.
8 cæruléscens. H.P. (st.) blue-stemmed.	C. B. S.	1825.	— D.G. h.	
9 tetragóna. H.P. (gr.ye.) four-sided.			— D.G. h.	
10 heptágona. w. (br.re.) seven-angled.		1731.	7. 11. D.G. h.	Bradl. suc. 2. t. 13.
a nigra. H.s. (br.re.) black-spined.			— D.G. h.	
β rúbra. H.s. (br.re.) red-spined.			— D.G. h.	
11 pentágona. H.P. (gr.ye.) five-sided.		1825.	— D.G. h.	
12 enneágona. H.s.(gr.ye.) nine-angled.		1790.	7. 10. D.G. h.	
13 mammillaris. w. (gr.ye.) warty-angled.		1759.	7. 8. D.G. h.	Comm. præl. t. 9.
14 cereifórmis. w. (gr.ye.) naked.		1731.	6. 7. D.G. h.	Burm. afr. t. 9. f. 3.
15 officinárum. w. (ye.) officinal.	Africa.	1597.	— D.G. h.	DC. plant. gras. t. 77.
16 odontophylla. W.en. (gr.) white-spined.	C. B. S.	1824.	— D.G. h.	
17 stellaspina. H.P. (gr.ye.) starry-spined.			— D.G. h.	
18 polygona. H.s. (gr.ye.) many-angled.		1790.	7. 9. D.G. h.	Lodd. bot. cab. t. 1334.
19 nereifólia. w. (re.) Oleander-leaved.	India.	1690.	6. 7. D.S. h.	DC. plant. gras. t. 46.
20 Hy'strix. w. (st.) Porcupine.	C. B. S.	1695.	— D.G. h.	Jacq. schœn. 2. t. 207.
loricata. P.s. <i>Treisia</i> <i>Hy'strix</i> . H.s.				
21 varians. H.s. (gr.ye.) variable-stemm'd.	E. Indies.	1800.	6. 9. D.S. h.	
22 grandifolia. H.s.(gr.ye.) great-leaved.	Sierra Leon.	1798.	— D.S. h.	
23 erosa. W.en. (gr.ye.) erose.		1818.	— D.S. h.	
24 tribuloides. s.s. (wh.) Caltrops-like.	Canaries.	1816.	6. 9. D.G. h.	
25 Millii. Desm. (sc.) splendid.	Madagascar.	1828.	— D.S. h.	Bot. mag. t. 2902.
splendens. B.M. ann. 1829. <i>Millii</i> . Desm. ann. 1826.				
26 cucumerina. w. (gr.) Cucumber-like.	C. B. S.		5. 9. D.G. h.	Vail. it. t. 5.
27 meloformis. w. (ye.) Melon-like.		1774.	— D.G. h.	Andr. reposit. t. 617.
28 clava. w. (gr.) club.			3. 12. D.G. h.	Jacq. ic. 1. t. 85.
<i>Treisia</i> <i>tuberculata</i> . H.s.s.				
29 Haworthii. (gr.) Haworth's.			— D.G. h.	
<i>Treisia</i> <i>clava</i> . H.s. excl. syn. w.				
30 bupleurifolia. w. (gr.) cone-shaped.		1791.	6. 7. D.G. h.	Jacq. schœn. 1. t. 106.
31 Cáput medusæw. (gr.) great Medusa's head.	Africa.	1731.	8. D.G. h.	Lodd. bot. cab. t. 1315.
<i>Medusa</i> <i>màjor</i> . H.s.				
β humiliis. (gr.) dwarfer.		1768.	— D.G. h.	Burm. afr. t. 8.
32 tessellata. H.s. (gr.) chequer'd-M'-head.		1788.	— D.G. h.	
33 fructis-pini. H.s.(gr.) small M'-head.	C. B. S.	1731.	— D.G. h.	DC. plant. gras. t. 150.
β geminata. (gr.) twin-branched.			— D.G. h.	Burm. afr. 18. t. 9. f. 1.
34 procumbens. H.s.(gr.) least M'-head.		1768.	— D.G. h.	
35 pátula. Mill. (br.gr.) spreading.			— D.G. h.	t. 10. f. 1.
<i>Dactylanthes</i> <i>pátula</i> . H.s.				
36 anacantha. w. (br.gr.) scaly.		1727.	9. 10. D.G. h.	Bot. mag. t. 2520.
37 globosa. B.M. (br.gr.) roundish-jointed.		1822.	6. 11. D.G. h.	— t. 2624.
<i>Dactylanthes</i> <i>globosa</i> . H.p.				
38 tuberculata. w. (wh.) tuberculate.		1805.	6. 8. D.G. h.	Jacq. schœn. 2. t. 208.
39 hamata. H.s. (br.gr.) hooked.		1795.	— D.G. h.	Burm. afr. t. 6. f. 3.
40 Commelinæ. DC. (br.gr.) Commelin's.	Africa.		— D.S. h.	
41 Ornithopus. W.en.(w.) Bird's-foot.		1816.	— G. h.	Jacq. fragm. t. 120.
42 clandestina. s.s. (gr.) clandestine.		1824.	— D.G. h.	
43 lophogóna. w. (gr.) curl-winged.	Madagascar.		— D.S. h.	DC. plant. gras. ic.
44 péndula. L.en. (gr.) pendulous.		1808.	— D.S. h.	
<i>Tithymalus</i> <i>péndulus</i> . H.				
45 Tirucálli. w. (wh.) Indian tree.	India.	1690.	— D.S. h.	Rheed. mal. 2. t. 34.
46 mauritánica. w. (ye.) Barbary.	Africa.	1732.	6. 8. D.G. h.	Dill. elt. t. 289. f. 373.
47 Lamárckii. H.S.L.(gr.) Lamarck's.		1808.	— D.G. h.	
mauritánica. Lam. nec aliorum. <i>virgata</i> . P.s. non W.K.				
48 aphylla. W.en. (gr.) leafless.	Teneriffe.	1815.	— D.G. h.	
49 atropurpurea. W.en.(pu.)dark purple.			— D.G. h.	
50 piscatória. w. (gr.ye.) smooth spear-l'd.	Canaries.	1777.	— D.G. h.	
51 balsamifera.w.(gr.ye.) balsam.		1779.	— D.G. h.	
52 laurifolia. P.s. (gr.ye.) Laurel-leaved.	Peru.	1820.	— D.G. h.	
53 bracteata. s.s. (gr.) bracteated.		1809.	— D.S. h.	Jacq. schœn. b. t. 276.
54 dendroides. w. (ye.) tree-like.	Italy.	1768.	— H. h.	Barrel. ic. 910.

EUPHORBIACEÆ.

55 l'æta. w.	(st.) Mezereon-leaved. S. Europe.	1758.	6. 8. D.G. h.	Jacq. ic. 1. t. 87.
56 punicea. w.	(cr.) scarlet-flowered. Jamaica.	1778.	1. 9. D.S. h.	Bot. reg. t. 190.
57 heterophylla. w. (sc.)	various-leaved. W. Indies.	1690.	4. 9. D.S. h.	Plum. ic. t. 251. f. 3.
58 cyathophora. w. (ye.pi.)	coloured. S. America.	1806.	7. 8. D.S. h.	Jacq. ic. 3. t. 480.
59 repánda. (gr.ye.)	waved. E. Indies.	1808.	8. D.S. h.	
Tithymalus repándus. H.S.				
60 biglandulösa. H.S. (gr.)	twin-glanded. Isl. Bourbon.	—	9. D.S. h.	
61 geniculata. P.S. (st.)	jointed. Cuba.	1823.	7. 10. S. (●)	Jacq. schoen. 3. t. 277.
prunifolia. H.K.	Tithymalus prunifolius. H.S.			
62 nudiflora. s.s.	(st.) naked-flowered. Jamaica.	1800.	8. D.S. h.	Jacq. ic. 3. t. 470.
63 verticillata. s.s.	(gr.) whorled. W. Indies.	1826.	— D.S. h.	
64 cotinifolia. w.	(wh.) Cotinus-leaved. S. America.	1690.	7. 8. D.S. h.	Hook. ex. flor. t. 59.
65 petiolaris. H.K.	(ye.) long-stalked. W. Indies.	1800.	5. 6. D.S. h.	Bot. mag. t. 883.
66 mellifera. w. (pu.gr.)	honey-bearing. Madeira.	1784.	4. 5. G. h.	— t. 1305.
67 glabra. s.s. (gr.ye.)	smooth-leaved. W. Indies.	1824.	— D.S. h.	
68 toxicaria. L.en. (gr.)	poisonous.	1825.	— D.S. h.	
69 picta. w. (gr.)	painted. S. America.	1789.	5. 7. S. 4.	Jacq. ic. 3. t. 477.
70 silenifolia. (gr.ye.)	Silene-leaved. C. B. S.	1821.	— G. 4.	
Tithymalus silenifolius. H.R.				
71 crista. H.R. (gr.ye.)	curl-leaved. —	—	G. 4.	
72 tuberosa. w. (gr.ye.)	tuberous. Ethiopia.	1800.	10. 12. G. h.	Burm. afr. 9. t. 4.
73 affinis. DC.	(gr.ye.) affined. —	1823.	6. 8. G. h.	Jacq. ic. 1. t. 86.
74 linaria. w. (ye.)	toad-flax-leaved.	1794.	— S. h.	
75 linifolia. w. (ye.)	Flax-leaved. W. Indies.	—	— S. h.	
76 ocyoidea. w. (ye.)	Basil-leaved. S. America.	1733.	— S. (●)	
77 dentata. M.	(ye.) dented. N. America.	1806.	6. 7. H. (●)	
78 hypericifolia. w. (wh.)	Hypericum-leaved. America.	1727.	6. 9. H. (●)	
79 Humboldtia. W.en.(gr.)	Humboldt's. S. America.	1809.	7. 10. S. (♂)	Hook. ex. flor. t. 36.
80 prostrata. w. (pu.)	trailing-red. W. Indies.	1758.	— S. (●)	
81 rosea. w. (ro.)	rosy. E. Indies.	1808.	8. S. (●)	
82 maculata. w. (ye.)	spotted. S. America.	1660.	7. H. (●)	
83 humifusa. W.en. (ye.)	trailing.	1824.	6. 8. H. (●)	
84 pilulifera. w. (gr.)	globular. E. Indies.	1800.	— S. 4.	
85 hyssopifolia. w. (gr.)	Hyssop-leaved. W. Indies.	1787.	8. 9. S. (●)	Jacq. vind. 2. t. 186.
86 scordifolia. s.s. (wh.)	Scordium-leaved. S. America.	1826.	6. 8. S. h.	Jacq. icon. t. 476.
87 tannensis. s.s. (gr.)	Tanna. N. Hebrides.	—	— G. h.	
88 thymifolia. w. (wh.)	Thyme-leaved. India.	1699.	7. 8. S. (●)	
89 parviflora. w. (wh.)	small-flowered. E. Indies.	1818.	— S. (●)	
90 variegata. B.M. (wh.)	variegated. Louisiana.	1811.	6. 8. H. (●)	
91 canescens. w. (wh.re.)	canescent. Spain.	1818.	7. 8. H. (●)	
92 Chamæsyc. w. (wh.)	scollop-leaved. S. Europe.	1752.	— H. (●)	
93 Péplis. w. (ye.)	purple. Devonshire.	— H. (●)	
94 polygonifolia. w. (st.)	knotgrass-leaved. N. America.	1704.	6. 7. H. (●)	
95 Ipêcacuanha. w. (gr.)	Ipêcacuanha.	—	1812.	— H. 4.
96 Péplus. w. (gr.)	petty. Britain.	7. 8. H. (●)	
97 peploides. DC.	(gr.) least. S. Europe.	1800.	7. 9. H. (●)	
Europa minima. H.S.				
98 falcata. w. (ye.)	sickle-leaved. —	1699.	6. 8. H. (●)	
99 acuminata. M.B. (gr.)	acuminate. Iberia.	1823.	— H. (●)	
100 exigua. w. (gr.)	dwarf. Britain.	7. H. (●)	
101 diffusa. w. (gr.ye.)	diffuse. Austria.	1798.	7. 8. H. (●)	
102 undulata. M.B. (wh.)	undulate. Caucasus.	1820.	— H. 4.	
103 micrantha. w. (gr.)	small-flowered. Persia.	1803.	7. 9. H. (●)	
104 Lagascana. s.s. (gr.)	Lagasca's. Spain.	1827.	7. 10. H. (●)	
105 Láthyrus. w. (gr.)	Caper. England.	5. 10. H. (♂)	
106 terracina. w. (ye.)	retuse-leaved. S. Europe.	1818.	7. 10. H. (●)	
107 valentina. P.S. (ye.)	Spanish. Spain.	1804.	7. 8. H. (●)	
108 'Arios. w. (ye.)	pear-rooted. Candia.	1596.	6. 7. H. 4.	
109 genistoides. w. (ye.)	Genista-like. C. B. S.	1808.	7. 8. G. h.	
110 condensata. s.s. (ye.)	tufted. Iberia.	1828.	6. 8. H. 4.	
111 spinosa. w. (ye.)	prickly. Levant.	1710.	6. 7. F. h.	
112 fruticosa. L.en. (ye.)	shrubby. Sicily.	1825.	— F. h.	
113 epithymoides. w. (ye.)	broad-leaved. Austria.	1805.	5. 6. H. 4.	
114 villosa. w.k.	(sn.) villous. Hungary.	1818.	6. 9. H. 4.	
115 temnifolia. s.s. (ye.)	slender-leaved. S. Europe.	1827.	— H. (●)	
116 echinocarpa. s.s. (ye.)	prickly-podded. Crete.	1828.	— H. (♂)	
117 condylocarpa. M.B. (ye.)	warted-fruited. Caucasus.	1824.	— H. 4.	
118 obtusata. Ph. (ye.)	blunt-bracted. N. America.	1826.	— H. 4.	
119 merculiâna. Ph. (ye.)	Mercury-like. —	1824.	— H. 4.	

Wats. dend. brit. t. 45.
Biv. pl. sic. cent. 1. 35.
Andr. deposit. t. 616.
W. et K. hung. t. 93.

EUPHORBIACEÆ.

120 marginata. Ph. (wh.) white-margined. N. America.	1816.	6. 9. H. ⊖.	
121 dulcis. w. (pu.) sweet. S. Europe.	1759.	5. 6. H. Ȑ.	Jacq. aust. 3. t. 213.
122 ambigua. w.k. (ye.) ambiguous. Hungary.	1824.	— H. Ȑ.	W. et K. hung. t. 135.
123 carniolica. w. (ye.) Carniolian. Carniola.	1795.	8. H. Ȑ.	Scop. carn. t. 21.
124 angulata. w. (ye.) angular-stalked. Austria.	1819.	7. 8. H. Ȑ.	Jacq. ic. t. 481.
125 nummulariaefolia. (ye.) moneywort-leaved.	1800.	7. G. h.	
126 erythrina. L.en. (gr.) purple-margined. C. B. S.	1816.	6. 8. G. Ȑ.	Ten. fl. neap. t. 42.
127 neapolitana. T.N. (wh.) Neapolitan. Naples.	1825.	7. 8. H. Ȑ.	Bocc. sic. t. 5.
128 Pithyusa. w. (gr.) Juniper-leaved. S. Europe.	1741.	6. 7. F. h.	Eng. bot. v. 7. t. 441.
129 portlandica. w. (ye.) Portland. Britain.	5. 9. H. Ȑ.	Jacq. aust. t. 345.
130 saxatilis. w. (ye.) rock. Austria.	1816.	6. 8. H. Ȑ.	Eng. bot. v. 3. t. 195.
131 glareosa. M.B. (ye.) stony. Caucasus.	1819.	— H. Ȑ.	
132 Paralias. w. (ye.) sea. England.	7. 9. H. Ȑ.	
β suffruticosa. (ye.) shrubby sea.	— F. h.	Jacq. schœn. 1. t. 107.
133 junccea. w. (ye.) rushy. Madeira.	1779.	7. 8. G. Ȑ.	
134 congesta. W.en. (ye.) fiddle-leaved. Spain.	1817.	— H. Ȑ.	
135 aleppica. w. (ye.) Aleppo. Crete.	1739.	— F. Ȑ.	Alp. exot. t. 64.
136 taurinensis. W.en. (ye.) Italian. Italy.	1818.	— H. Ȑ.	All. ped. 2. t. 83. f. 2.
137 segetalis. w. (ye.) corn. S. Europe.	1699.	— H. ⊖.	Jacq. aust. 5. t. 450.
138 gracilis. s.s. (ye.) slender. Ukraine.	1826.	— H. Ȑ.	
139 cæspitosa. T.N. (ye.) tufted. Naples.	—	— H. Ȑ.	
140 pinea. s.s. (ye.) Pine-leaved. —	—	— H. Ȑ.	
linifolia. T.N. non w.			
141 biumbellata. DC. (ye.) double-umbel'd. Barbary.	1780.	8. H. ⊖.	Poir. it. ed. germ. t. 1.
142 angustifolia. (ye.) narrow-leaved.	—	7. 8. H. Ȑ.	
'Esula angustifolia. H.S.			
143 multicorymbosa. (ye.) many-flowered.	1805.	— H. Ȑ.	
144 provincialis. w. (ye.) linear-leaved. S. Europe.	1800.	8. 11. H. ⊖.	
145 juncoides. H.S. (ye.) Rush-like.	—	7. 8. H. ⊖.	
146 helioscopia. w. (ye.) Wartwort. Britain.	— H. ⊖.	Eng. bot. v. 13. t. 883.
147stellulata. Sn. (ye.) starry. Europe.	1823.	— H. ⊖.	
148 bialata. L.en. (ye.) two-winged.	1819.	— H. ⊖.	
149 serrata. w. (ye.) narrow notch-ld. S. Europe.	1710.	7. G. Ȑ.	Jacq. ic. 3. t. 483.
150 uralensis. F. (ye.) Fischer's. Tauria.	1824.	— H. Ȑ.	
151 cratica. w. (ye.) Cretan hoary. Levant.	1768.	— H. Ȑ.	
152 rigens. (ye.) stiff.	1795.	7. 8. H. h.	
Galarhæus rigidus. H.S.			
153 verrucosa. w. (ye.) warted. France.	1800.	6. H. Ȑ.	Moris. s. 10. t. 3. f. 3.
154 corollata. w. (wh.) great-flowered. N. America.	1803.	7. 9. H. Ȑ.	Lodd. bot. cab. t. 390.
155 spathulæfolia. H.S. (gr.) spatula-leaved.	1800.	8. G. h.	
156 corallioides. w. (ye.) Coral-stalked. S. Europe.	1752.	6. 9. H. Ȑ.	
157 mollis. Gm. (ye.) soft.	1823.	— H. Ȑ.	
158 androsæmifolia. H.S. (ye.) Tutsan-leav'd. Hungary.	1804.	7. 9. H. Ȑ.	
159 pilosa. w. (ye.) hairy. Siberia.	1758.	5. 8. H. Ȑ.	
160 orientalis. w. (st.) oriental. Levant.	1739.	6. 7. H. Ȑ.	
161 fragifera. L.en. (ye.) Strawberry-like. Italy.	1825.	— H. Ȑ.	
162 virgata. w.k. (ye.) twiggy. Hungary.	1807.	7. H. Ȑ.	W. et K. hung. t. 162.
163 platyphylla. H.S. (ye.) annual warted. England.	7. 8. H. ⊖.	Jacq. aust. t. 376.
164 stricta. E.B. (ye.) upright warted.	— H. ⊖.	Eng. bot. v. 5. t. 333.
165 literata. w. (st.) blotch-leaved.	1790.	8. H. ⊖.	Jacq. ic. 3. t. 482.
166 crispata. L.en. (ye.) crisp-leaved.	1819.	7. 9. H. Ȑ.	
167 pubescens. s.s. (ye.) pubescent. Barbaria.	1827.	— H. Ȑ.	Jacq. f. ecl. t. 66.
168 lutescens. Led. (ye.) yellowish. Siberia.	—	— H. Ȑ.	
169 Ehrenbergii. (ye.) Ehrenberg's. Egypt.	1826.	6. 9. H. ⊖.	
heterophylla. Ehr. non w.			
170 trigonocarpa. F. (ye.) three-sided podded. Russia.	1821.	— H. ⊖.	
171 retusa. c.l. (gr.) notch-leaved. S. Europe.	—	— H. ⊖.	Cavan. ic. t. 34. f. 3.
172 'Esula. w. (gr.) leafy-branched. Britain.	5. 7. H. Ȑ.	Eng. bot. v. 20. t. 1399.
173 Gerardiæna. w. (ye.) Gerard's. Germany.	1801.	7. H. Ȑ.	Jacq. aust. t. 436.
174 Cyparissias. w. (ye.) Cypress. England.	5. 9. H. Ȑ.	Eng. bot. v. 12. t. 840.
175 nicænus. w. (ye.) sharp-leaved. Spain.	1809.	— F. Ȑ.	Jacq. ic. 3. t. 485.
176 myrsinoides. w. (ye.) glaucous. S. Europe.	1570.	4. 6. H. Ȑ.	
177 rigida. M.B. (ye.) rigid. Tauria.	1819.	6. 8. H. Ȑ.	Bot. reg. t. 274.
178 imbricata. w. (ye.) imbricated. Portugal.	1804.	8. 9. F. h.	
179 glaucescens. W.en. (ye.) glaucescent.	1819.	— H. Ȑ.	
180 agraria. M.B. (ye.) field. Tauria.	—	6. 8. H. ⊖.	
181 palustris. w. (ye.) marsh. Sweden.	1570.	5. 8. H. Ȑ.	Flor. dan. t. 866.
182 pallida. w. (ye.) pale. Hungary.	1818.	6. 8. H. Ȑ.	
183 lucida. w.k. (ye.) glossy. —	1820.	— H. Ȑ.	W. et K. hung. t. 55.

184 pròcera. M.B. (ye.) tall.	Terek.	1824.	6. 8. H. ¼.	
185 ceratocárpa. T.N. (ye.) horn-capsuled.	Naples.	—	— H. ¼.	Ten. fl. neap. t. 43.
186 emarginàta. w. (ye.) freckled.	Italy.	1758.	7. 8. H. ¼.	
187 híberna. E.B. (ye.) Irish.	Ireland.	—	5. 6. H. ¼.	Eng. bot. v. 19. t. 1337.
188 salicifòlia. w. (ye.) Willow-leaved.	Hungary.	1804.	— H. ¼.	Lodd. bot. cab. t. 973.
189 sylvàtica. w. (pi.) wood.	S. Europe.	1768.	7. 9. F. ½.	Jacq. aust. t. 375.
190 amygdaloïdes. w. (st.) Almond-leaved.	England.	—	3. 6. H. ¼.	Eng. bot. v. 4. t. 256.
191 Characias. w. (ye. pu.) upright red.	—	—	— H. ½.	— v. 7. t. 442.
192 véneta. L.en. (ye.) venomous.	Europe.	1824.	— F. ½.	
193 Bonplândii. (wh.) Bonpland's.	N. Spain.	1823.	4. 6. S. ¼.	
marginàta. k.s. non Nutt.				
CI'CCA. L.	Ci'CCA. Monœcia Tetrandria. L.			
dísticha. w. (gr.) long-leaved.	E. Indies.	1796.	4. 6. S. ½.	Jacq. schœn. 2. t. 194.
BU'XUS. W.	Box-tree. Monœcia Tetrandria. L.			
1 chinénsis. L.en. (ye.) China.	China.	1812.	4. 8. H. ½.	
2 baleàrica. w. (ye.) Minorca.	Minorca.	1780.	7. H. ½.	
3 semperfivens. w. (ye.) common.	England.	—	4. H. ½.	Eng. bot. v. 19. t. 1341.
α arborèscens. Mill. (ye.) tree.	—	—	— H. ½.	
† argéntea. (ye.) silvery-leaved.	—	—	— H. ½.	
‡ àurea. (ye.) golden-leaved.	—	—	— H. ½.	
marginàta. (ye.) margined.	—	—	— H. ½.	
β angustifòlia. Mill. (ye.) narrow-leaved.	—	—	— H. ½.	
† variezàta. (ye.) variezated.	—	—	— H. ½.	
γ myrtifòlia. Mill. (ye.) myrtle-leaved.	—	—	— H. ½.	
δ suffruticòsa. (ye.) dwarf.	—	—	— H. ½.	
SARCÓCO'CCA. B.R. SARCOCO'CCA.	Monœcia Tetrandria.			
coriacea. (wh.) leathery.	Nepaul.	1820.	4. 6. G. ½.	Bot. reg. t. 358.
prunifòrmis. B.R.	Pachysandra coriacea.	Hook. ex. flor. t. 148.		
PACHYSA'NDRA. M.	PACHYSA'NDRA.	Monœcia Tetrandria. W.		
1 procumbens. m. (bh.) procumbent.	N. America.	1800.	3. 4. H. ¼.	Bot. reg. t. 33.
ACID'OTON. W.	ACID'OTON.	Monœcia Polyandria.		
ürens. w. (gr.) stinging.	Jamaica.	1793.	4. 6. S. ½.	Sloan. jam. 1. t. 83. f. 1.
PLUKEN'ETIA. W.	PLUKEN'ETIA.	Monœcia Monadelphia. L.		
1 volùbilis. w. (gr.) twining.	W. Indies.	1739.	7. 8. S. ½.	Plum. ic. t. 226.
2 corniculàta. s.s. (gr.) horned.	E. Indies.	1827.	— S. ½.	Rumph. amb. 1.t.79.f.2.
DALECHA'MPIA. W.	DALECHA'MPIA.	Monœcia Monadelphia.		
1 scàndens. w. (gr.) climbing.	W. Indies.	1739.	6. 7. S. ½.	Jacq. amer. 252. t. 160.
2 brasiliéns. w. (gr.) Brazilian.	Brazil.	1820.	— S. ½.	
3 fimbriata. k.s. (gr.) fringed.	Mexico.	1829.	— S. ½.	
ACA'LYPHA. W.	ACA'LYPHA.	Monœcia Monadelphia. W.		
1 diversifòlia. w. (gr.) various-leaved.	Caracas.	1824.	6. 8. S. ½.	
2 hermandifòlia. s.s. (gr.) Hernandia-leaved.	Jamaica.	1825.	— S. ½.	Schk. hand. 3. t. 311.
3 virgínica. w. (gr.) Virginian.	N. America.	1759.	7. 8. H. ¼.	Lam. ill. t. 780. f. 2.
4 caroliniana. w. (gr.) Carolina.	—	1811.	— H. ¼.	Vahl symb. 1. t. 20.
5 ciliàta. w. (gr.) ciliated.	E. Indies.	1799.	— S. ¼.	Sloan. hist. 1. t. 82. f. 3.
6 réptans. w. (gr.) creeping.	Jamaica.	1822.	— S. ½.	Rheed. mal. 10. t. 81.
7 índica. w. (gr.) Indian.	E. Indies.	1759.	7. 9. S. ¼.	Jacq. schoen. 2. t. 243.
8 hispida. w. (gr.) hispid.	—	1820.	— S. ¼.	
9 cuspidàta. w. (gr.) cuspidate.	Caracas.	—	— S. ½.	
10 rúbra. W.en. (re.) red.	S. America.	1823.	6. 9. S. ¼.	
11 brachystàcha. W.en. (gr.) short-spiked.	China.	1816.	7. 8. H. ¼.	
12 pacifìcola. W.en. (gr.) few-flowered.	—	—	— H. ¼.	
13 alopecuroidæa. w. (gr.) Fox-tail.	Venezuela.	1804.	7. 9. H. ¼.	Jacq. ic. 3. t. 620.
14 anemoiòdes. k.s. (gr.) Anemia-like.	Mexico.	1829.	— S. ½.	
15 purpurascens. k.s. (pu.) purple-bracted.	—	1827.	— S. ½.	
16 elliptica. s.s. (gr.) elliptic-leaved.	Jamaica.	1825.	— S. ½.	
17 lævigàta. w. (gr.) smooth.	—	1819.	— S. ½.	
18 virgàta. w. (gr.) twiggy.	—	1822.	— S. ½.	
19 macrostàcha. w. (gr.) long-spiked.	Caracas.	1824.	— S. ½.	Brown. jam. t. 36. f. 2.
20 polystàcha. w. (gr.) many-spiked.	—	—	— S. ½.	Jacq. schoen. 2. t. 245.
21 scaròsà. w. (gr.) rough.	Jamaica.	1822.	— S. ½.	— 2. t. 246.
22 integrifòlia. s.s. (gr.) entire-leaved.	Madagascar.	1823.	— S. ½.	
OMALA'NTHUS. J.	OMALA'NTHUS.	Monœcia Monadelphia.		
populifòlius. B.M.(wh.) Poplar-leaved.	N. Holland.	1824.	6. 8. G. ½.	Bot. mag. t. 2780.
CAPER'ONIA. Hilar.	CAPER'ONIA.	Monœcia Monadelphia.		
1 castaneafòlia. s.s.(gr.) Chestnut-leaved.	S. America.	1828.	7. 8. S. ¼.	
2 palústris. s.s. (gr.) marsh.	Vera Cruz.	1731.	— S. ¼.	Mart. dec. 4. t. 88.
Crôton palûstris. w.				

CROTON. W.	CR'OTON. Monœcia Monadelphia. L.	E. Indies.	1804.	6. 9.	S. h.	Rheed mal. 6. t. 61.
1 variegatus. w.	(gr.) variegated.	—	—	—	S. h.	Rumph. amb. 4.t.26.f.2.
α latifolius. H.B.	(gr.) broad-leaved.	—	—	—	S. h.	— 4. t. 25.
β crispus. H.B.	(gr.) curl-leaved.	—	—	—	S. h.	Lodd. bot. cab. t. 870.
γ medius. H.B.	(gr.) narrow-leaved.	—	1820.	—	S. h.	Jacq. amer. t. 162. f. 4.
2 pictus. B.C.	(gr.) painted-leaved.	—	—	—	S. h.	Catesb. car. 2. t. 46.
3 linearis. s.s.	(gr.) Willow-leaved.	W. Indies.	1733.	7.	S. h.	Sloan. jam. 2. t. 174. f. 2.
4 Cascarilla. s.s.	(gr.) Cascarilla.	S. America.	1778.	7. 8.	S. h.	L'Herit. stirp. 17. t. 9.
5 maritimus. s.s. (gr.wh.)	sea-side.	Carolina.	1786.	—	G. h.	
6 gracilis. k.s.	(gr.) slender.	Mexico.	1824.	7. 8.	S. h.	
7 glabellus. s.s. (gr.wh.)	Laurel-leaved.	Jamaica.	1778.	5. 8.	S. h.	
8 hirtus. s.s. (gr.wh.)	hairy.	Trinidad.	1821.	—	S. ⊙.	
9 argenteus. s.s. (gr.)	silvery-leaved.	S. America.	1733.	7. 8.	S. ⊙.	Rheed. mal. 2. t. 33.
10 Tiglum. w. (wh.)	purging.	E. Indies.	1796.	8. 9.	S. h.	Jacq. ic. 1. t. 194.
11 glandulosus. s.s. (gr.)	glandular.	W. Indies.	1818.	—	S. ⊙.	Aubl. guian. 2. t. 339.
12 subluteus. s.s. (gr.ye.)	yellowish.	Trinidad.	1822.	—	S. h.	
13 populifolius.s.s.(gr.wh.)	Poplar-leaved.	Jamaica.	1824.	—	S. h.	Plum. mss. 4. t. 123.
14 tiliæfolius. P.S. (gr.wh.)	Lime-tree-leaved.	St. Vincent.	1825.	—	S. h.	Sloan. hist. 2. t. 174. f. 2.
15 Elutèria. s.s. (gr.wh.)	sea-side Balsam.	Jamaica.	1748.	—	S. h.	
16 lúcidus. s.s. (gr.wh.)	shining.	—	1822.	7. 9.	S. h.	
17 micanus. s.s. (gr.ye.)	glittering.	—	1815.	—	S. h.	Pluk. alm. t. 220. f. 5.
18 pungens. s.s. (gr.ye.)	pungent.	Caracas.	1791.	—	S. h.	Jacq. ic. 3. t. 622.
19 penicillatus. s.s. (gr.)	pencilled.	Cuba.	1799.	7. 8.	S. h.	Venten. choix. t. 12.
20 flavens. s.s. (ye.gr.)	pale yellow.	W. Indies.	1822.	—	S. h.	
21 macrophyllus.s.s.(wh.)	large-leaved.	Jamaica.	—	—	S. h.	Rumph. amb. 3. t. 126.
22 aromaticus. s.s. (wh.)	aromatic.	Ceylon.	1793.	—	S. h.	
23 humilius. s.s. (wh.gr.)	humble.	Jamaica.	1799.	—	S. h.	
24 farinosis. s.s. (wh.)	mealy.	Madagascar.	1826.	—	S. h.	
25 Astroites. w. (gr.wh.)	woolly-leaved.	W. Indies.	1782.	7. 8.	S. h.	
26 lobatus. s.s. (wh.)	various-leaved.	Vera Cruz.	1730.	—	S. ⊙.	Mart. dec. 5. t. 46.
27 gossypifolius. s.s. (wh.)	Cotton-leaved.	Trinidad.	1823.	—	S. h.	Vahl. symb. t. 49.
28 tomentosus. L.en.	tomentose.	—	—	—	S. h.	
CROZO'PHORA. J.	CROZO'PHORA. Monœcia Pentandria.					
1 plicata. J.	(wh.gr.) plaited.	E. Indies.	1824.	6. 8.	G. ⊙.	Burm. ind. t. 62. f. 1.
2 tinctoria. J.	(wh.gr.) officinal.	S. Europe.	1570.	7. 8.	H. ⊙.	Niss. act. par. 1712. t. 17.
Croton tinctorium. w.						
3 verbascifolia.J.(wh.gr.)	Mullein-leaved.	Greece.	1826.	—	H. ⊙.	
ROTTL'ERA. W.	ROTTL'ERA. Diœcia Icosandria. W.					
1 brasiliensis. s.s. (gr.)	Brazilian.	Brazil.	1820.	—	S. h.	
2 tinctoria. s.s. (gr.)	dyer's.	E. Indies.	1810.	—	S. h.	Roxb. corom. 2. t. 168.
GEL'ONIUM. W.	GEL'ONIUM. Diœcia Icosandria. W.					
1 bifarium. w. (gr.)	oval-leaved.	E. Indies.	1793.	6. 8.	S. h.	
2 lanceolatum. H.B. (gr.)	spear-leaved.	—	1825.	—	S. h.	
TRE'WIA. W.	TRE'WIA. Diœcia Polyandria. W.					
nudiflora. w. (gr.)	naked-flowered.	E. Indies.	1796.	—	S. h.	Rheed. mal. 1. t. 42.
HY'ENA'NCHE. H.K.	HY'ENA-POISON. Diœcia Dodecandraia.					
globosa. H.K. (gr.wh.)	Cape.	C. B. S.	1783.	4. 9.	G. h.	Lamb. cinch. 52. t. 10.
E'UCLEA. W.	E'UCLEA. Diœcia Dodecandraia. W.					
1 racemosa. w. (wh.)	round-leaved.	C. B. S.	1722.	11. 12.	G. h.	Jacq. fragm. 3. t. 1. f. 5.
2 undulata. w. (wh.)	wave-leaved.	—	1794.	6. 10.	G. h.	
JA'TROPHA. W.	PHYSIC-NUT. Monœcia Monadelphia. L.					
1 gossypifolia. w. (sc.)	Cotton-leaved.	W. Indies.	1690.	5. 8.	S. h.	Bot. reg. t. 746.
2 panduræfolia. w. (sc.)	fiddle-leaved.	Cuba.	1800.	—	S. h.	Bot. mag. t. 604.
3 integriflora. w. (sc.)	spicy.	—	1809.	—	S. h.	— t. 1464.
4 Cúrcas. w. (gr.)	angular-leaved.	S. America.	1731.	—	S. h.	Jacq. vind. 3. t. 63.
5 multifida. w. (sc.)	multifid.	—	1696.	6. 8.	S. h.	Sal. par. lond. t. 91.
6 coccinea. L.en. (sc.)	scarlet.	—	1822.	—	S. h.	
CNIDO'SCOLUS. P.B.	CNIDO'SCOLUS. Monœcia Monadelphia.					
1 urens. P.B.	(wh.) stinging.	Brazil.	1690.	5. 7.	S. h.	Jacq. vind. 1. t. 21.
Játröpha urens. Jacq.						
2 stimulosus. Ph.	(wh.) stimulous.	N. America.	1812.	—	F. 4.	Pluk. alm. t. 220. f. 3.
Játröpha stimulosa. Ph.						
3 napæifolia. w. (wh.)	Napæa-leaved.	Antilles.	1823.	7. 8.	S. h.	
4 frágrans. k.s.	(wh.) fragrant.	Cuba.	1822.	6. 8.	S. h.	Reliq. houst. 6. t. 15.
5 herbacea. w. (wh.)	annual.	Vera Cruz.	1759.	7. 8.	S. ⊙.	Pohl. flor. bras. 1. t. 50.
6 Maregraviæ. P.B. (wh.)	Marcgrave's.	Brazil.	1823.	—	S. h.	— 1. t. 52.
7 vitifolius. P.B.	(wh.) Vine-leaved.	—	—	—	S. h.	
MA'NIHOT. P.B.	MA'NIHOT. Monœcia Decandria.					
1 fœtida. P.B.	(bl.br.) strong-scented.	Mexico.	1824.	7. 8.	S. h.	

EUPHORBIACEÆ.

2 Jánipha. P.B. (<i>br.gr.</i>) Læffling's.	S. America.	1824.	7. 8.	S. h.	Jacq. amer. t. 162. f. 1.
<i>Játróphæ carthaginénsis.</i> Jacq. Jánipha. w. <i>Jánipha Læfflingii.</i> K.S.					
3 grátilis. P.B. (<i>br.gr.</i>) slender.	Brazil.	1822.	7. 8.	S. h.	Pohl. pl. bras. 1. t. 16.
4 anómala. P.B. (<i>br.</i>) anomalous.		—	—	S. h.	— 1. t. 21.
5 pruinosa. P.B. (<i>bl.br.</i>) frosted.		1824.	—	S. h.	— 1. t. 22.
6 digitiformis. P.B. (<i>bl.br.</i>) finger-formed.		1826.	—	S. h.	— 1. t. 27.
7 digitata. finger-leaved. N. S. W.	1810.	G. h.		
<i>Cróton digitatum.</i> F.					
8 diversifolia. different-leaved. N. Holland.	1822.	G. h.		
9 tenuifolia. P.B. (<i>bl.br.</i>) slender-leaved. Brazil.	—	6. 8.	S. h.	Pohl. pl. bras. 1. t. 29.	
10 aesculifolia. K.S. (<i>bl.br.</i>) Æsculus-leaved. S. America.	1825.	5. 8.	S. h.	H. et B. 2. t. 109.	
11 cannabina. (<i>bl.br.</i>) Cassava.	1739.	7. 8.	S. h.	Sloan. jam. 1. t. 85.	
<i>Játróphæ Mánihot.</i> w. <i>Jánipha Mánihot.</i> K.S.					
12 caricaefolia. P.B. (<i>bl.br.</i>) Papaw-leaved. Brazil.	1822.	—	S. h.	Pohl. pl. bras. 1. t. 32.	
13 Dalechampiæformis. P.B. (<i>br.</i>) Dalechampia-like.	1818.	—	S. h.	— 1. t. 36.	
14 sinuata. P.B. (<i>br.</i>) sinuate-leaved.	1824.	—	S. h.	— 1. t. 41.	
ALEURITES. W. ALEURITES. <i>Monœcia Monadelphia.</i> L.					
1 triloba. w. (<i>wh.</i>) three-lobed. Society Isles. 1793.					
2 Ambinux. P.S. (<i>wh.</i>) Molucca. Ceylon. 1803.					
<i>Cróton moluccanum.</i> w.					
RI'CINUS. W. PALMA-CHRISTI. <i>Monœcia Monadelphia.</i>					
1 communis. w. (<i>re.st.</i>) Castor-oil plant. E. Indies.	1548.	7. 8.	H. ⓧ.	Lam. ill. t. 792.	
2 Krappa. L.O. (<i>wh.fl.</i>) Krappa.	1827.	8. 11.	H. ⓧ.		
3 leucocarpus. L.O. (<i>wh.bh.</i>) white-capsuled.		—	F. ⓧ.		
4 macrophyllus. L.O. (<i>w.r.</i>) large-leaved.		—	F. ⓧ.		
5 paniculatus. L.O. (<i>re.wh.</i>) paniced.	1824.	8. 10.	F. ⓧ.		
6 rutilans. L.O. (<i>re.wh.</i>) red-stalked.	1827.	—	F. ⓧ.		
7 undulatus. L.O. (<i>re.wh.</i>) wave-leaved.		—	F. ⓧ.		
8 viridis. w. (<i>bh.wh.</i>) green. E. Indies.	1802.	8.	H. ⓧ.		
9 africanus. w. (<i>re.wh.</i>) African. Africa.	7. 8.	G. h.		
10 lividus. w. (<i>re.wh.</i>) livid-leaved. C. B. S.	1795.	—	G. h.		
11 inermis. w. (<i>re.st.</i>) smooth-capsuled. India.	1755.	—	S. ⓧ.		
12 armatus. A.R. (<i>re.wh.</i>) rough-capsuled. Malta.	18. 7.	—	S. ⓧ.		
MA'PPA. J. MA'PPA. <i>Monœcia Monadelphia.</i>					
1 moluccana. s.s. (<i>wh.</i>) Molucca. Moluccas. 1828.	...	S. h.			
2 Tanaria. s.s. (<i>wh.</i>) scollop-leaved. E. Indies. 1809.	7. 9.	S. h.			
<i>Ricinus Tanarius.</i> w.					
OMPHALEA. W. OMPHALEA. <i>Monœcia Monadelphia.</i> L.					
triandra. w. (<i>wh.</i>) long-leaved. Jamaica. 1763.	6. 7.	S. h.			
HIPPO'MANE. W. MANCHINEEL-TREE. <i>Monœcia Monadelphia.</i> L.					
1 Mancinella. w. (<i>gr.</i>) common. W. Indies. 1690.	S. ⓧ.			
2 spinosa. s.s. (<i>gr.</i>) spiny.	1820.	S. ⓧ.	ot. cab. t. 519.	
<i>Sapium ilicijolium.</i> w.					
S'APIUM. W. S'APIUM. <i>Monœcia Monadelphia.</i> L.					
1 aequiparium. s.s. single-spiked. W. Indies. 1692.	S. h.	J. Amer. t. 158.		
2 Hippomane. s.s. (<i>gr.</i>) two-glanded.	1824.	S. h.		
3 indicum. w. (<i>gr.</i>) Indian. E. Indies. 1817.	S. h.	tesb. car. 2. t. 95.		
SIPHONIA. W. ELASTIC-GUM-TREE. <i>Monœcia Monadelphia.</i> W.					
Cahúchu. w. (<i>gr.</i>) common. S. America. 1823.	S. h.	1. ic. 171. f. 1.		
<i>elástica.</i> P.S.					
AGYN'EJA. W. AGYN'EJA. <i>Monœcia Monadelphia.</i> W.					
1 impubes. w. (<i>gr.</i>) smooth. China. 1820.	7. 9.	H. ⓧ.			
2 pùbera. w. (<i>gr.</i>) woolly.	1823.	S. h.		
STILLI'NGIA. W. STILLI'NGIA. <i>Monœcia Monadelphia.</i> W.					
1 sylvatica. w. (<i>ye.</i>) wood. Carolina. 1787.	7. 8.	G. 4.			
2 ligustrina. w. (<i>st.</i>) Privet-leaved. N. America. 1812.	—	G. h.			
3 sebifera. w. (<i>st.</i>) Tallow-tree. China. 1703.	9.	S. h.			
H'URA. W. SANDBOX-TREE. <i>Monœcia Monadelphia.</i> W.					
1 strépens. W.en. (<i>gr.</i>) unequal-toothed. S. America.	7. 9.	S. h.		
2 crépitans. W.en. (<i>gr.</i>) equal-toothed.	1733.	—	S. h.		
BRAD'L'EJA. W. BRAD'L'EJA. <i>Monœcia Monadelphia.</i> W.					
1 sinica. w. (<i>wh.</i>) Chinese. China. 1816.	7. 9.	S. h.			
2 nítida. H.B. (<i>wh.</i>) glossy. E. Indies. 1820.	—	S. h.			
3 multilocularis. s.s. (<i>wh.</i>) multilocular.		—	S. h.		
PHYLЛА'NTHUS. W. PHYLЛА'NTHUS. <i>Monœcia Monadelphia.</i> W.					
1 obovatus. w. (<i>wh.</i>) annual. N. America. 1803.	7. 8.	H. ⓧ.			
2 maderaspátensis. w. (<i>gr.</i>) Madras. E. Indies. 1783.	—	S. h.			
3 grátilis. H.B. (<i>gr.</i>) slender.	1820.	—	S. 4.		
4 longifolius. w. (<i>gr.</i>) long-leaved. Isl. Bourbon. 1822.	—	S. h.			
				Lam. ill. t. 156. f. 1.	

5 lanceolatus. w. (gr.)	spear-leaved.	Isl. Bourbon.	1822.	6. 8. S. h.	
6 nútans. w. (bh. st.)	nodding.	Jamaica.	—	— S. h.	Jacq. schœn. 2. t. 193.
7 grandifolius. w. (gr.)	great-leaved.	Portorico.	1771.	— S. h.	
8 virósus. w. (gr.)	venomous.	E. Indies.	1802.	— S. h.	
9 turbinatus. B.M. (gr.)	shining-leaved.	China.	7. 9. S. h.	Bot. mag. t. 1862.
10 Conámi. w. (gr.)	Brazilian.	Brazil.	1791.	— S. h.	Aub. gui. 2. t. 354.
11 obcordatus. H.B. (gr.)	obcordate.	E. Indies.	1818.	— S. 4.	
12 retusus. H.B. (gr.)	r. tuse-leaved.	—	1822.	— S. 4.	Lodd. bot. cab. t. 577.
13 reticulatus. B.C. (bh.)	netted.	—	1818.	8. 9. S. h.	
14 lúcens. w. (gr.)	thick-veined.	China.	1822.	7. 9. S. h.	
15 cantoniensis. L.en.(gr.)C	ton.	—	1819.	7. 8. H.○.	
16 cuneatus. W.en. (gr.)	v. dge-leaved.	—	—	— H.○.	Rheed. mal. 10. t. 15.
17 Niruri. w. (gr.) I.	tian.	E. Indies.	1692.	6. 9. S. ○.	Rumph. amb. 6.t.17.f.2.
18 Urinaria. w. (gr.)	s. ging.	—	1819.	— S. ○.	
19 microphyllus. K.S. (gr.)	s. ill-leaved.	S. America.	1824.	— S. h.	
20 polyphyllus. w. (gr.)	n. iy-leaved.	E. Indies.	1805.	7. 9. S. h.	Lodd. bot. cab. t. 721.
21 mimosoides. w. (gr.)	M. osa-like.	W. Indies.	1820.	— S. h.	
22 rhamnoïdes. w. (gr.)	Rt. us-like.	E. Indies.	—	— S. h.	Lodd. bot. cab. t. 731.
23 stríctus. H.B. (gr.)	up.	—	1823.	— S. h.	
24 fraxiniifolius. B.C. (gr.)	As' ed.	—	1818.	— S. h.	
25 juglandifolius. W.en.(gr.)V.	nutt-leaved.	New Spain.	—	6. 9. S. h.	
26 scándens. Roxb. (gr.)	climbing.	E. Indies.	1803.	— S. h.	Lodd. bot. cab. t. 548.
E'MBLICA. L.T.	E'MBLICA.	Monoclea Syngenesia. L.T.			
1 officinalis. L.T.	(st.) officinal.	E. Indies.	1768.	6. 9. S. h.	
2 racemosa. s.s.	(st.) racemed.	—	1793.	7. 8. S. h.	

Phyllanthus racemosus. w.

KIRGAN'ELIA. J. KIRGAN'ELIA. Monoclea Monadelphia.
élegans. J. (gr.) elegant. Mauritius. 1820. 6. 9. S. h.

Phyllanthus Kirganelia. w.

XYLOPHYLLA. W. XYLOPHYLLA. Monoclea Monadelphia. W.

1 longifolia. w. (re)	long-leaved.	E. Indies.	1816.	1. 12. S. h.
-----------------------	--------------	------------	-------	--------------

Phyllanthus cerámica. P.S.

2 latifolia. w. (wh.)	Seaside Laurel.	S. America.	1783.	8. 10. S. h.
-----------------------	-----------------	-------------	-------	--------------

3 speciosa. (wh.)	handsome.	Jamaica.	1818.	— S. h.
-------------------	-----------	----------	-------	---------

4 arbúscula. w. Phyllanthus speciosus. Jacq.				
--	--	--	--	--

5 angustifolia. w. (wh.)	narrow-leaved.	—	1789.	7. 8. S. h.
--------------------------	----------------	---	-------	-------------

6 linearis. P.S. (re)	linear-leaved.	—	1822.	— S. h.
-----------------------	----------------	---	-------	---------

7 elongata. B.C. (re)	elongated.	—	—	— S. h.
-----------------------	------------	---	---	---------

8 montana. w. (or.)	mountain.	—	1820.	11. 4. S. h.
---------------------	-----------	---	-------	--------------

9 falcatá. w. (re)	te-leaved.	Bahama Isl.	1699.	7. 9. S. h.
--------------------	------------	-------------	-------	-------------

10 ramiflora. w. (u.)	leaved.	Siberia.	1785.	7. 8. H. h.
-----------------------	---------	----------	-------	-------------

10 obovata. W.en. (re)	ovate-leaved.	—	—	— S. h.
------------------------	---------------	---	---	---------

ANDRA'CHNE. W. telephoides. w.	ASTARD ORPINE. Monoclea Gynandria. L.			
--------------------------------	---------------------------------------	--	--	--

BRID'ELIA. W. (wh.)	climbing.	E. Indies.	1804.	7. 9. S. h.
---------------------	-----------	------------	-------	-------------

2 montana. s.s. (st.)	mountain.	—	1825.	— S. h.
-----------------------	-----------	---	-------	---------

3 fruticosa. P.S. (st.)	frutescent.	—	1827.	— S. h.
-------------------------	-------------	---	-------	---------

CLU'YTIA. W. (st.)	hill.	E. Indies.	1807. S. h.
--------------------	-------	------------	-------	------------

2 pátula. w. (st.)	spreading.	—	1812. S. h.
--------------------	------------	---	-------	------------

3 pulchélla. w. (st.)	broad-leaved.	C. B. S.	1739.	1. 6. G. h.
-----------------------	---------------	----------	-------	-------------

4 heterophylla. w. (st.)	various-leaved.	—	1816.	2. 4. G. h.
--------------------------	-----------------	---	-------	-------------

5 tomentosa. w. (st.)	tomentose.	—	1812.	4. 6. G. h.
-----------------------	------------	---	-------	-------------

6 pubescens. w. (st.)	pubescent.	—	1817.	— G. h.
-----------------------	------------	---	-------	---------

7 polifolia. w. (wh.)	Poley-leaved.	—	1790.	— G. h.
-----------------------	---------------	---	-------	---------

8 ericoides. w. (wh.)	Heath-leaved.	—	—	— G. h.
-----------------------	---------------	---	---	---------

9 daphnoides. w. (wh.)	Daphne-like.	—	1731.	5. 6. G. h.
------------------------	--------------	---	-------	-------------

10 polygonoides. (wh.)	Polygonum-like.	—	1790.	12. 3. G. h.
------------------------	-----------------	---	-------	--------------

11 alaternoides. w. (wh.)	narrow-leaved.	—	1692.	— G. h.
---------------------------	----------------	---	-------	---------

EXCCE'ARIA. W. EXCÆ'ARIA. Dioclea Monadelphia. L.				
---	--	--	--	--

1 lúcida. s.s. (wh.)	shining-leaved.	Jamaica.	1824.	4. 6. S. h.
----------------------	-----------------	----------	-------	-------------

2 glandulosa. s.s. (wh.)	glandular.	—	—	— S. h.
--------------------------	------------	---	---	---------

3 Agallócha. s.s. (gr.ye.)	Indian.	E. Indies.	1823. S. h.
----------------------------	---------	------------	-------	------------

4 serrata. H.K. (wh.)	saw-leaved.	Chile.	1796.	11. 2. G. h.
-----------------------	-------------	--------	-------	--------------

AD'ELIA. W. AD'ELIA. Dioclea Monadelphia. L.				
--	--	--	--	--

1 Bernárdia. w. (gr.wh.)	villous-leaved.	Jamaica.	1768.	7. 8. S. h.
--------------------------	-----------------	----------	-------	-------------

2 Ricinélla. w. (gr.wh.)	smooth-leaved.	—	—	6. 8. S. h.
--------------------------	----------------	---	---	-------------

3 Acidótón. w. (gr.wh.)	Box-leaved.	—	—	6. 7. S. h.
-------------------------	-------------	---	---	-------------

Jacq. schœn. 2. t. 193.

Bot. mag. t. 1862.

Aub. gui. 2. t. 354.

Rheed. mal. 10. t. 15.

Rumph. amb. 6.t.17.f.2.

LOURE'IRA. W.	LOURE'IRA. Diœcia Monadelphia. W.					
1 cuneifolia. w. (bh.) wedge-leaved.	Mexico. 1824.	6. 8. G. h.	Cavan. ic. 5. t. 429.			
<i>Mozinna spathulata.</i> Ort. dec. 8. t. 13.						
2 glandulosa. w. (bh.) glandulous.	1799.	— G. h.	Cavan. ic. 5. t. 430.			
B'ORYA. W.	B'ORYA. Diœcia Diandria. W.					
1 porulosa. w. (gr. wh.) Florida.	Florida. 1806.	7. 8. H. h.				
2 ligustrina. w. (gr. wh.) Privet-leaved.	N. America. 1812.	— H. h.				
3 prinoïdes. W.en. (gr. wh.) Prinos-like.	1816.	— H. h.				
4 nítida. W.en. (gr. wh.) glossy-leaved.	—	— H. h.				
5 retusa. W.en. (gr. wh.) retuse-leaved.	1812.	— H. h.				
6 acuminata. w. (gr. wh.) pointed.	—	— H. h.				
<i>Adelia acuminata.</i> M.						
SECURI'NEGA. J.	SECURI'NEGA. Monœcia Pentandria.					
nítida. P.S. (wh.)	Otaheite Myrtle. Otaheite. 1793.	3. 8. S. h.	Lindl. collect. t. 9.			
TR'AGIA. W.	TR'AGIA. Monœcia Triandria. W.					
1 volubilis. w. (gr.) twining.	W. Indies. 1739.	6. 7. S. h.	Trew. pl. rar. 2. t. 15.			
2 involucrata. w. (gr.) involured.	E. Indies. 1759.	— S. ⊙.	Jacq. ic. 1. t. 190.			
3 mercurialis. w. (gr.) Mercury-like.	—	— S. ⊙.	Rheed. mal. 10. t. 82.			
4 urticæfolia. s.s. (gr.) nettle-leaved.	N. America. 1812.	— H. ⊙.				
5 hirsuta. D.L. (gr.) hairy.	Trinidad. 1820.	— S. ⊙.				
6 urens. w. (gr.) stinging.	Virginia. 1699.	8. H. ⊙.				
7 cannabina. w. (gr.) Hemp-leaved.	—	1699. 6. 7. S. ♀.				
MICROST'ACHYS. J.	MICROST'ACHYS. Monœcia Triandria.					
1 corniculata. (gr.) two-horned.	Trinidad. 1820.	6. 7. S. ⊙.				
<i>Tragia corniculata.</i> Vahl. <i>Cnemidostachys Vahlii.</i> s.s.						
2 Chamælæa. j. (gr.) lance-leaved.	E. Indies. 1793.	— S. ⊙.	Rheed. mal. 2. t. 34.			

ORDO CLXXXI.

EMPETREÆ. *D. Don on the affinities of the Empetreae.* p. 3.

EMP'ETRUM. D.D.	EMP'ETRUM. Diœcia Triandria. L.				
nigrum. w. (gr.) black-berried.	Britain.	4. 5. H. h.	Eng. bot. v. 8. t. 526.		
COR'EMA. D.D.	COR'EMA. Diœcia Triandria.				
ábum. D.D. (gr.) white-berried.	Portugal. 1774.	4. 6. H. h.			
<i>Empetrum ábum.</i> w.					
CERAT'IOLA. M.	CERAT'IOLA. Monœcia Diandria. N.				
ericoides. M. (st.) Heath-like.	N. America. 1810.	4. 6. H. h.	Pursh. fl. amer. 1. t. 13.		

ORDO CLXXXII.

ANTIDESMEÆ.

ANTIDE'SMA. W.	ANTIDE'SMA. Diœcia Pentandria. L.				
1 alexiteria. w. (gr.) Laurel-leaved.	E. Indies. 1793.	5. 6. S. h.	Rheed. mal. 5. t. 11.		
2 pubescens. w. (gr.) pubescent.	1818.	— S. h.	Roxb. corom. 2. t. 167.		
3 paniculata. w. (gr.) paniced.	1800.	— S. h.			
STIL'AGO. W.	STIL'AGO. Diœcia Triandria. W.				
1 Bünius. w. (gr.) Laurel-leaved.	E. Indies. 1757.	8. 10. S. h.	Rheed. mal. 4. t. 56.		
2 diandra. w. (gr.) diandrous.	1800.	— S. h.	Roxb. corom. 2. t. 166.		

ORDO CLXXXIII.

URTICEÆ. *Juss. gen. 400.*

FICUS. L.	FIG-TREE. Polygamia Diœcia. W.				
1 religiosa. R.S. (gr.) Poplar-leaved.	E. Indies. 1731. S. h.	Rheed. mal. 1. t. 27.		
2 superstitionis. L.en. (gr.) superstitious.	— S. h.			

3	<i>nymphaeifolia</i> , R.S. (gr.)	Water-lily-leaved.	E. Indies.	1759.	S. h.	
4	<i>cordata</i> , R.S. (gr.)	heart-leaved.	C. B. S.	1802.	4. 6.	S. h.	Thunb. fic. p. 8. c. ic.
5	<i>oblongata</i> , L.en. (gr.)	oblong-leaved.	———	1816.	S. h.	
6	<i>crassinervia</i> , R.S. (gr.)	thick-nerved.	S. America.	———	S. h.	
7	<i>ruberinervia</i> , L.en. (gr.)	red-nerved.	Brazil.	1824.	S. h.	
8	<i>rubicinosa</i> , R.S. (gr.)	ferrugineous.	N. S. W.	1789.	3. 6.	G. h.	Venten. malm. t. 114.
	<i>australis</i> , W.						
9	<i>costata</i> , R.S. (gr.)	rib-leaved.	E. Indies.	1763.	S. h.	
10	<i>laevigata</i> , R.S. (gr.)	smooth.	W. Indies.	1822.	S. h.	
11	<i>lentiginosa</i> , R.S. (gr.)	veined-leaved.	———	1826.	S. h.	
12	<i>citrifolia</i> , R.S. (gr.)	Citron-leaved.	———	1824.	S. h.	
13	<i>coronata</i> , R.S. (gr. re.)	red-crowned.	N. S. W.	1807.	3. 6.	G. h.	Colla hort. rip. t. 8.
14	<i>lucida</i> , R.S. (gr.)	shining-leaved.	E. Indies.	1772.	4. 6.	S. h.	
15	<i>scandens</i> , R.S. (gr.)	climbing.	S. America.	1816.	—	S. h.	
16	<i>stipulata</i> , R.S. (gr.)	trailing.	China.	1771.	—	G. h.	
17	<i>pedunculata</i> , R.S. (gr.)	Willow-leaved.	S. America.	1776.	5. 7.	S. h.	Pluk. alm. t. 178. f. 4.
18	<i>venosa</i> , R.S. (gr.)	vein-leaved.	E. Indies.	1763.	6. 8.	S. h.	Rheed. mal. 3. t. 64.
19	<i>leucotoma</i> , R.S. (gr.)	white-spotted.	———	—	S. h.	W. hort. ber. 1. t. 36.
	<i>venosa</i> , W. non H.K.						
20	<i>tinctoria</i> , R.S. (gr.)	Otaheite.	Society Isl.	1783.	5. 6.	S. h.	
21	<i>septica</i> , R.S. (gr.)	rough.	E. Indies.	1826.	S. h.	Rumph. amb. 3. t. 96.
22	<i>benjamina</i> , R.S. (gr.)	oval-leaved.	———	1757.	S. h.	Rheed. mal. 1. t. 26.
23	<i>padifolia</i> , K.S. (gr.)	Padus-leaved.	Mexico.	1826.	S. h.	
24	<i>complicata</i> , K.S. (gr.)	complicated.	———	—	S. h.	
25	<i>glabrata</i> , K.S. (gr.)	smooth.	Teneriffe.	1816.	G. h.	
26	<i>benghalensis</i> , R.S. (gr.)	Bengal.	E. Indies.	1690.	4. 5.	S. h.	Rheed. mal. 1. t. 28.
27	<i>microcarpa</i> , R.S. (gr.)	small-fruited.	Guinea.	1823.	—	S. h.	
28	<i>americana</i> , R.S. (gr.)	American.	Jamaica.	1822.	S. h.	Plum. spec. t. 132. f. 2.
29	<i>racemosa</i> , R.S. (gr.)	clustered.	E. Indies.	1759.	4. 6.	S. h.	Rheed. mal. 1. t. 25.
30	<i>pertusa</i> , R.S. (gr.)	Laurel-leaved.	S. America.	1780.	—	S. h.	
31	<i>terebrita</i> , R.S. (gr.)	obovate-leaved.	Mauritius.	1822.	S. h.	Bory it. t. 17.
32	<i>pumila</i> , R.S. (gr.)	dwarf.	China.	1759.	4. 6.	G. h.	Kämpf. amen. t. 804.
33	<i>comosa</i> , R.S. (gr.)	tufted.	E. Indies.	1825.	S. h.	Roxb. corom. 2. t. 125.
34	<i>glomerata</i> , R.S. (gr.)	crowded.	———	1818.	S. h.	— 2. t. 123.
35	<i>Hookeri</i> , (gr.)	Hooker's.	W. Indies.	1816.	1. 4.	S. h.	Hook. ex. flor. t. 111.
	<i>nitida</i> , H.E.F. nec aliorum.						
36	<i>nitida</i> , R.S. (gr.)	glossy-leaved.	E. Indies.	1786.	3. 6.	S. h.	Rheed. mal. 3. t. 55.
37	<i>ciliolosa</i> , R.S. (gr.)	fringe stipuled.	———	1816.	—	S. h.	
38	<i>rubra</i> , R.S. (re.)	vermilion-fruited.	E. Indies.	1826.	S. h.	Rumph. amb. 3. t. 85?
39	<i>cotinifolia</i> , K.S. (gr.)	Cotinus-leaved.	Mexico.	1824.	3. 6.	S. h.	
40	<i>obtusifolia</i> , K.S. (gr.)	blunt-leaved.	———	—	S. h.	
41	<i>coriacea</i> , R.S. (gr.)	leathery-leaved.	E. Indies.	1772.	S. h.	
42	<i>populnea</i> , R.S. (gr.)	Poplar-leaved.	———	—	S. h.	
43	<i>macrophylla</i> , R.S. (gr.)	large-leaved.	N. Holland.	1816.	4. 6.	G. h.	
44	<i>elastica</i> , H.B. (gr.)	Indian rubber.	E. Indies.	1815.	G. h.	
45	<i>Brassii</i> , H.T. (gr.)	Brass's.	Sierra Leon.	1822.	S. h.	
46	<i>lasiophylla</i> , L.en. (gr.)	downy-leaved.	———	1816.	S. h.	
47	<i>glaucocephylla</i> , R.S. (gr.)	glaucous-leaved.	S. America.	1824.	S. h.	
48	<i>insipida</i> , R.S. (gr.)	insipid.	Caracas.	1829.	S. h.	
49	<i>retusa</i> , R.S. (gr.)	retuse-leaved.	E. Indies.	1793.	4. 6.	S. h.	
50	<i>laurifolia</i> , R.S. (gr.)	Laurel-leaved.	W. Indies.	1759.	—	S. h.	Sloan. jam. 2. t. 223.
	<i>martiniensis</i> , W.						
51	<i>calyculata</i> , R.S. (gr.)	calyculate.	Vera Cruz.	1760.	4. 6.	S. h.	
52	<i>brasiliensis</i> , L.en. (gr.)	Brazilian.	Brazil.	1820.	S. h.	
53	<i>cuneata</i> , R.S. (gr.)	wedge-leaved.	———	1826.	S. h.	
54	<i>indica</i> , R.S. (gr.)	Banyan-tree.	E. Indies.	1759.	S. h.	Rheed. mal. 3. t. 63.
55	<i>salicifolia</i> , R.S. (gr.)	Willow-leaved.	Arabia.	1818.	S. h.	
56	<i>aurantiaca</i> , Lod. (or)	orange-coloured.	—	S. h.	
57	<i>pendula</i> , L.en. (gr.)	pendulous.	1824.	5. 7.	S. h.	
58	<i>myrtifolia</i> , L.en. (gr.)	Myrtle-leaved.	———	—	S. h.	
59	<i>pubescens</i> , R.S. (gr.)	pubescent.	E. Indies.	———	S. h.	
60	<i>nervosa</i> , R.S. (gr.)	strong-nerved.	———	1826.	S. h.	
61	<i>elliptica</i> , K.S. (gr.)	elliptic-leaved.	S. America.	1824.	S. h.	
62	<i>acuminata</i> , L.C. (gr.)	acute-leaved.	—	S. h.	
63	<i>longifolia</i> , L.C. (gr.)	long-leaved.	—	S. h.	
64	<i>caudata</i> , L.C. (gr.)	tailed.	—	S. h.	
65	<i>dumosa</i> , L.C. (gr.)	bushy.	—	S. h.	
66	<i>viscidifolia</i> , L.C. (gr.)	Viscum-leaved	—	S. h.	
67	<i>lanceolata</i> , H.B. (gr.)	spear-leaved.	E. Indies.	1820.	S. h.	

68 obtusata. L.en. (gr.) obtuse-toothed.	1820.	S. h.	
69 muntia. L.en. (gr.) rough-leaved. N. Holland. 1818.	G. h.		Willden. sic. t. 2.
70 ulmifolia. R.S. (gr.) Elm-leaved. Philippine. 1825.	S. h.		Roxb. corom. 2. t. 124.
71 capensis. R.S. (gr.) Cape. C. B. S. 1816.	G. h.		
72 exasperata. R.S. (gr.) very rough. scabra. w. non Forst. Guinea. 1822.	S. h.		
73 hispida. R.S. (gr.) opposite-leaved. E. Indies. 1802.	S. h.		
oppositifolia. Rox. scabra. Jacq. schoen. 3. t. 315.				
74 Lichtensteinii. L.en. (gr.) Lichtenstein's. C. B. S. 1824.	G. h.		
75 aquatica. w. (gr.) water. E. Indies. 1758.	4. 6.	S. h.		Rheed. mal. 3. t. 62.
76 heterophylla. R.S. (gr.) various-leaved. ——— 1816.	S. h.		
77 repens. R.S. (gr.) creeping-stemmed. ——— 1805.	S. h.		
78 virgata. H.B. (gr.) twiggy. ——— 1817.	G. h.		Trew-Ehret. t. 73. 74.
79 Carica. R.S. (gr.) common. S. Europe. 1548.	6. 7.	H. h.		
BROSIMUM. W. BREAD-NUT. Polygamia Diecia. W.				Sw. fl. ind. oc. 1. f. 1. f. 1.
1 Alicastrum. w. (gr.) Jamaica. Jamaica. 1776.	S. h.		
2 spirum. w. (gr.) Milkwood. ——— 1789.	S. h.		
3 Galactodendron. D.D. (gr.) Cow-tree. Caracas. 1829.	S. h.		
Galactodendron utilis. k.s. Palo de Vacca. Humboldt.				
ARTOCA'RPUS. W. BREAD-FRUIT. Monocelia Monandria. W.				
1 incisa. w. (gr.) true. S. Sea Isles. 1793.	S. h.		Rumph. amb. 1. t. 33.
a nufera. (gr.) nut-bearing. ——— ———	S. h.		
β apyrēna. (gr.) seedless. ——— ———	S. h.		
2 integrifolia. w. (gr.) Jaca-tree. E. Indies. 1778.	6.	S. h.		Rumph. amb. 1. t. 32.
β heterophylla. Lam. various-leaved. ——— ———	S. h.		Rheed. mal. 3. t. 26-28.
MACLURA. N. MACLURA. Diocia Tetrandria. N.				Rumph. amb. 1. t. 30.
1 aurantiaca. n. (gr.) Osage Apple. Missouri. 1824.	H. h.		
2 tinctoria. D.D. (gr.) Fustick-wood. W. Indies. 1739.	S. h.		Lamb. pin.v.2. ap. t. 12.
Morus tinctoria. w. Broussonetia tinctoria. k.s.				Plum. ic. t. 204.
BROUSSON'ETIA. W. PAPER MULBERRY. Diocia Tetrandria. W.				
papyrifera. w. (gr. br.) common. Japan. 1751.	4. 7.	H. h.		
M'ORUS. W. MULBERRY. Monocelia Tetrandria. L.				
1 alba. w. (gr.) white. China. 1596.	6.	H. h.		Andr. reposit. t. 488.
2 tatrica. w. (gr.) Tartarian. Tartary. 1784.	—	H. h.		
3 italicia. w. (gr.) Italian. Italy. ———	—	H. h.		
4 constantinopolitana. (gr.) Constantinople. Constantinople. ———	—	H. h.		
5 nigra. w. (gr.) common. Italy. 1548.	—	H. 5.		
6 rubra. w. (gr.) red. N. America. 1629.	6. 7.	H. 5.		
7 scabra. W.en. (gr.) rough. ——— 1812.	—	H. h.		
8 pensylvanica. L.C. (gr.) Pennsylvanian. ——— ———	—	H. h.		
9 pumila. L.C. (gr.) dwarf. ——— ———	—	H. h.		
10 indica. w. (gr.) Indian. E. Indies. 1820.	S. h.		Rheed. mal. 1. t. 49.
11 sinensis. L.C. (gr.) Chinese. China. ———	G. h.		Jacq. ic. 3. t. 617.
12 mauritianna. w. (gr.) Mauritius. Mauritius. ——— ———	—	S. h.		
CONOCE'PHALUS. B.J. CONOCE'PHALUS. Monocelia Tetrandria.				Bot. reg. t. 1203.
naucleiflorus. B.R. (ye.) Nauclea-flower'd. E. Indies. 1816.	10. 4.	S. h.		
BÖHM'ERIA. W. BÖHM'ERIA. Monocelia Tetrandria. W.				
1 caudata. w. (gr.)tailed. W. Indies. 1822.	4. 8.	S. h.		
2 cylindrica. w. (gr.) cylindrical. Virginia. 1759.	6. 8.	H. 4.		
3 macrophylla. D.P. (gr.) large-leaved. Nepaul. 1820.	—	G. h.		
4 rotundifolia. D.P. (gr.) round-leaved. ——— 1819.	—	G. 4.		
5 platiphylla. D.P. (gr.) rough-leaved. ——— 1818.	6. 10.	G. h.		
6 spicata. s.s. (gr.) spiked. Japan. 1824.	—	H. 4.		
7 ramiflora. w. (gr.) branch-flowering. W. Indies. 1822.	—	S. h.		
Catürus ramiflorus. L. Pröcris ramiflorus. Lam.				
8 elongata. L.en. (gr.) elongated. Brazil. ——— 7. 9. H. 4.				Hout. syst. 10. t. 72. f. 2.
9 lateriflora. w. (gr.) side-flowering. N. America. 1816.	6. 8.	H. 4.		Lam. ill. t. 763.
10 rubescens. w. (gr.) tree. Canaries. 1779.	2. 5.	G. h.		
Urtica arborea. L'Her. stirp. t. 20.				
11 salicifolia. D.P. (gr.) Willow-leaved. Nepaul. 1826.	7. 10.	G. h.		Jacq. frag. n. 30. t. 5. f. 1.
12 frondosa. D.P. (gr.) frondose. ——— 1818.	—	G. h.		
13 ternifolia. D.P. (gr.) three-leaved. ——— 1827.	—	G. h.		
14 frutescens. w. (gr.) frutescent. ——— 1819.	—	G. h.		
URTICA. W. NETTLE. Monocelia Tetrandria. L.				
1 pilulifera. w. (gr.) Roman. England. 6. 8. H. 4.				Eng. bot. v. 3. t. 148.
2 balearica. w. (gr.) Balearic. Balearic Isl. 1731.	6. 7.	H. 4.		Blackw. t. 321. f. 1.
3 Dodartia. w. (gr.) Dodart's. S. Europe. 1683.	7. 8.	H. 4.		
4 pumila. w. (gr.) dwarf. N. America. 1817.	—	H. 4.		
5 crassifolia. w. (gr.) thick-leaved. S. America. 1822.	—	S. h.		

URTICEÆ.

6 grandifolia. w.	(gr.) great-leaved.	Jamaica.	1793.	7. 8. S. h.	Sloan. jam. 1. t. 83. f. 2.
7 reticulatæ. w.	(gr.) net-leaved.	W. Indies.	1809.	10. 11. S. ȳ.	Bot. mag. t. 2481.
8 involucrata. B.M.	(gr.) imbossed.			8. 10. S. h.	
9 diffusa. w.	(gr.) spreading.				
10 rufa. w.	(gr.) rusty.	Jamaica.	1793.	6. 9. S. h.	
11 scabrélla. L.en.	(gr.) rough.	E. Indies.	1815.	10. 12. G. h.	
12 scripta. D.P.	(gr.) black-lined.	Nepaul.	1823.	6. 8. H. ȳ.	
13 rupestræ. s.s.	(gr.) rock.	Sicily.	1828.	— H. h.	
14 convéxa. H.H.	(gr.) convex-leaved.		1824.	6. 9. H. ⊖.	
15 ürens. w.	(gr.) small.	Britain.	6. 10. H. ⊖.	Eng. bot. v. 18. t. 1236.
16 dioica. w.	(gr.) common.		7. 9. H. ȳ.	— v. 25. t. 1750.
17 chamaedryoides. Ph.	(gr.) Germanander-like.	Georgia.	1813.	— H. ⊖.	
18 membranacea. w.	(gr.) membranaceous.	S. Europe.	1818.	— H. ȳ.	
19 angustifolia. F.	(gr.) narrow-leaved.	Russia.	1824.	— H. ȳ.	
20 árdens. L.en.	(gr.) burning.	Nepaul.	1819.	8. 11. H. ⊖.	
21 cannabina. w.	(gr.) Hemp-leaved.	Siberia.	1749.	7. 9. H. ȳ.	Amm. ruth. 249. t. 25.
22 rugosa. w.	(gr.) rough-leaved.	Jamaica.	1793.	5. 7. S. ȳ.	
23 grátilis. w.	(gr.) slender-stalked.	Hudson's Bay.	1782.	6. 8. H. ȳ.	
24 Parietaria. w.	(gr.) Pellitory-leaved.	Jamaica.	1793.	7. 9. S. h.	Sloan. j. 1. t. 93. f. 1.
25 ciliata. w.	(gr.) ciliated.		1815.	— S. h.	
26 rádicans. s.s.	(gr.) rooting.		1809.	— H. h.	
27 nummulariifolia. w.	(gr.) moneywort-leaved.		1822.	— S. h.	Sloan. hist. 1. t. 131. f. 4.
28 depræssa. w.	(gr.) depressed.		—	— S. ȳ.	
29 microphylla. w.	(gr.) small-leaved.	W. Indies.	1793.	4. 5. S. ȳ.	Sloan. j. 1. t. 93. f. 2.
30 trianthemoïdes. P.S.	(gr.) branching.		1824.	— S. ȳ.	
31 serrulata. s.s.	(gr.) saw-leaved.	Jamaica.	—	— S. ȳ.	
32 lúcida. s.s.	(gr.) glossy-leaved.		1808.	— S. h.	
33 recurvæ. H.H.	(gr.) recurved.		1826.	6. 9. H. ⊖.	Jacq. schœn. 3. t. 388.
34 macrostachya. L.O.	(gr.) long-spiked.	S. America.	1824.	— S. h.	
35 pulchella. L.en.	(gr.) neat.	E. Indies.	1816.	— S. h.	Pluk. alm. t. 239. f. 2.
36 æstuans. w.	(gr.) Surinam.	Surinam.	1803.	6. 7. S. ȳ.	Jacq. vind. 2. t. 166.
37 capitata. w.	(gr.) headed.	N. America.	1820.	— H. ȳ.	Jacq. schœn. 3. t. 386.
38 canadensis. w.	(gr.) Canadian.	Canada.	1656.	8. 10. H. ȳ.	
39 nivea. w.	(gr.) white-leaved.	China.	1739.	8. 9. H. ȳ.	
40 caracasana. w.	(gr.) Caracas.	Caracas.	1820.	— S. h.	
41 caravellana. L.en.	(gr.) pointed-leaved.	S. America.	—	8. 10. H. ⊖.	Andr. reposit. t. 454.
42 baccifera. w.	(gr.) berry-bearing.		1793.	7. 8. S. h.	
43 elongata. w.	(gr.) elongated.	Philippines.	1823.	— S. h.	
44 capitellata. s.s.	(gr.) small-headed.	Java.	1816.	4. 8. S. h.	
45 ferocissima.	(gr.) ferocious.	Nepaul.	1805.	8. 10. H. ⊖.	
horrída. L.en. non k.s.					
46 pentandra. H.B.	(gr.) pentandrous.	E. Indies.	—	— S. ȳ.	
47 arborescens. L.en.	(gr.) arborescent.	Manila.	1822.	— S. h.	Jacq. vind. 1. t. 48.
48 diversifolia. L.en.	(gr.) different-leaved.	E. Indies.	1820.	— S. h.	
49 heterophylla. D.P.	(gr.) various-leaved.	Nepaul.	1823.	6. 8. H. ȳ.	
FORSKÖHLEA. L.	FORSKÖHLEA.	Monœcia	Monandria. J.		
1 fruticosa. w.	(gr.) frutescent.	Teneriffe.	1816.	6. 8. G. h.	
2 tenacissima. w.	(gr.) clammy.	Egypt.	1767.	7. 8. H. ⊖.	
3 cándida. w.	(gr.) rough.	C. B. S.	1774.	6. 7. G. ȳ.	
4 víridis. E.	(gr.) green.	Arabia.	1828.	— G. ⊖.	
5 angustifolia. w.	(gr.) narrow-leaved.	Teneriffe.	1779.	7. 8. H. ⊖.	Murr. c. gott. p. 24. t. 2.
PARIETARIA. W.	PELLITORY.	Polygamia	Monœcia	L.	
1 índica. w.	(gr.) Indian.	E. Indies.	1790.	4. 5. S. ȳ.	
2 officinalis. w.	(gr.) common.	Britain.	6. 9. H. ȳ.	Eng. bot. v. 13. t. 879.
3 judaica. w.	(gr.) Basil-leaved.	Germany.	1728.	— H. ȳ.	Schkuh. hand. 3. t. 34.
4 pensylvánica. w.	(gr.) Pensylvanian.	N. America.	1816.	— H. ⊖.	
5 urticafolia. w.	(gr.) Nettle-leaved.	Isl. Bourbon.	1700.	— S. ⊖.	
6 lusitánica. w.	(gr.) chickweed-leaved.	Spain.	1710.	7. 8. H. ⊖.	Bocc. sic. t. 24. f. 13.
7 crética. w.	(gr.) Cretan.	Crete.	1818.	— H. ⊖.	
8 prostrata. L.en.	(gr.) prostrate.		1823.	— H. ȳ.	
9 polygonoides. w.	(gr.) Polygonum-leav'd.	Armenia.	1728.	— H. ⊖.	Mart. cent. pl. rar. t. 8.
PI'LEA. L.col.	PI'LEA.	Monœcia	Tetrandria.	L.	
muscosa. L.col.	(gr.) mossy.	W. Indies.	1793.	4. 5. S. ȳ.	Lindl. col. t. 4.
CA'NNABIS. W.	HEMP.	Diœcia	Tetrandria.	L.	
sativa. w.	(gr.) common.	India.	6. 7. H. ⊖.	Schk. hand. 3. t. 325.
H'UMULUS. W.	HOP.	Diœcia	Tetrandria.	L.	
Lùpulus. w.	(gr.) common.	Britain.	6. 8. H. ȳ.	Eng. bot. v. 6. t. 427.
THELY'GONUM. W.	DOG'S-CABBAGE.	Monœcia	Polyandria.	L.	
Cynocrámbe. w.	(gr.) common.	S. Europe.	1710.	7. H. ⊖.	Schk. hand. 3. t. 299.

GN'ETUM. W.	GN'ETUM.	Monœcia Monadelphia. L.				
Gnèmon. w.	(gr.) Indian.	India.	1815.	S. b.	Rumph. amb. 1. t. 71.
CECR' OPIA. W.	SNAKE-WOOD.	Diœcia Diandria. L.				Lam. ill. t. 800.
1 peltata. w.	(gr.) peiltated.	Jamaica.	1778.	S. b.	
2 palmata. w.	(gr.) palmated.	Brazil.	1823.	S. b.	
3 cóncolor. w.	(gr.) self-coloured.	—	—	—	S. b.	
GUNN'ERA. W.	GUNN'ERA.	Gynandria Diandria. L.				
perpensa. w.	(gr.) marsh-marygold-ld. C. B. S.	1688.	4. 6. G. 4.			Bot. mag. t. 2376.
DORSTENIA. W.	DORST'ENIA.	Monœcia Diandria.				Swart. obs. t. 7. f. 2.
1 cordifolia. w.	(gr.) heart-leaved.	W. Indies.	1822.	5. 8. S. u.		
2 brasiliensis. w.	(gr.) Brazilian.	S. America.	1792.	3. 4. S. u.		
3 Houstoni. w.	(gr.) Houstoun's.	—	1747.	6. 7. S. u.		
4 arifolia. w.	(gr.) Arum-leaved.	Brazil.	1823.	—	S. u.	
5 Contrayérra. w.	(gr.) Contrajerva-root.	S. America.	1748.	5. 8. S. u.		
6 Drakena. w.	(gr.) round-rooted.	Vera Cruz.	1729.	—	S. u.	
7 tubicina. B.M.	(gr.) tubed.	Trinidad.	1824.	7. 9. S. u.		
8 ceratosanthes. B.C. (gr. pu.) horned.		S. America.	—	4. 6. S. u.		
MERTE'NSIA. K.S.	MERTE'NSIA.	Polygamia Monœcia. R.S.				H. et B. n. gen. 2.t.103.
1 laevigata. K.S.	(ye.) Mexican.	Mexico.	1825.	S. b.	
2 iguánea. R.S.	(ye.) Jamaica.	Jamaica.	1791.	4. 6. S. b.		
<i>Rhamnus iguáneus.</i> L.	<i>Zizyphus iguáneus.</i> P.S.	<i>Celtis aculeata.</i> w.				
3 rhamnoïdes. R.S. (ye.)	Rhamnus-like.	—	1791.	4. 6. S. b.		
CELTIS. K.S.	NETTLE-TREE.	Polygamia Monœcia. L.				Cavan. ic. 3. t. 294.
1 micrantha. w.	(gr.) smooth.	Jamaica.	1739.	8. 9. S. b.		
2 orientalis. w.	(gr.) oriental.	E. Indies.	1820.	—	S. b.	Plum. ic. t. 206. f. 1.
3 pumila. Ph.	(gr.) dwarf.	N. America.	1812.	5.	H. b.	Rheed. mal. 4. t. 1.
4 crassifolia. w.	(gr.) heart-leaved. <i>cordifolia.</i> Duh.	—	—	5. 6. H. 5.		Duham. arb. 2. t. 9.
5 occidentalis. w.	(gr.) American.	—	1656.	4. 5. H. 5.		Wats. dend. brit. t. 147.
β cordata. R.S.	(gr.) heart-leaved.	—	—	—	H. 5.	
γ scabriuscula. R.S. (gr.) rough-leaved.	—	—	—	—	H. 5.	
6 Willdenowiæ. R.S. (gr.) Willdenow's.		China.	—	—	H. 5.	Tourn. it. 2. t. 41.
7 sinensis. R.S.	(gr.) Chinese.	—	—	—	H. 5.	Wats. dend. brit. t. 105.
8 laevigata. R.S.	(gr.) smooth.	Louisiana.	1812.	4. 5. H. 5.		
9 Tournefortii. w.	(gr.) Tournefort's.	Levant.	1738.	—	H. 5.	
10 australis. w.	(gr.) European.	S. Europe.	1796.	5.	H. 5.	
PLAN'ERA. M.	PLAN'ERA.	Tetrandria Diginia. S.S.				
1 Richardi. M.	(br. gr.) Richard's.	N. America.	1766.	4. 5. H. 5.		
<i>Ulmus nemorælis.</i> H.K.						
2 carpinifolia. w.D. (gr.)	Hornbeam-leaved.	Siberia.	—	—	H. 5.	Wats. dend. brit. t. 106.
3 parvifolia.	(gr.) small-leaved.	Caucasus.	—	—	H. 5.	Jacq. scheenb. t. 262.
<i>Ulmus parvifolia.</i> Jacq.						
4 Gmelini. M.	(br. gr.) Gmelin's. <i>aquatica.</i> N.	N. America.	1816.	—	H. 5.	
ULMUS. R.S.	ELM-TREE.	Pentandria Diginia. L.				
1 campéstris. E.B.	(pu.) common English.	Britain.	—	4. 5. H. 5.		Eng. bot. v. 27. t. 1886.
2 suberosa. E.B.	(pu.) cork-barked.	—	—	3. 4. H. 5.		— v. 31. t. 2161.
3 glabra. E.B.	(pu.) smooth.	—	—	—	H. 5.	— v. 32. t. 2248.
4 mäjor. E.B.	(br.) declining-branch'd.	—	—	—	H. 5.	— v. 36. t. 2542.
5 montana. E.B.	(gr.) Wych.	—	—	4. 5. H. 5.		— v. 27. t. 1887.
6 effusa. w.	(gr.) spreading-flowered.	—	—	—	H. 5.	Schk. hand. t. 57.
7 álba. w.	(br.) white.	Hungary.	1816.	—	H. 5.	
8 crispa. w.	(br.) curl-leaved.	N. America.	—	—	H. 5.	
9 americana. w.	(gr.) large-leaved.	—	1752.	—	H. 5.	
10 pendula. W.en.	(gr.) pendulous.	—	—	—	H. 5.	
11 fulva. M.	(br.) slippery.	—	—	—	H. 5.	
12 scabra. Mill.	(br.) rough.	—	—	—	H. 5.	
13 pumila. w.	(br.) dwarf.	Siberia.	1771.	—	H. b.	
14 microphylla. P.S.	(br.) jagged-leaved.	—	—	—	H. b.	Pall. ros. 1. t. 48.
15 alata. M.	(br.) winged.	N. America.	1816.	—	H. 5.	Pall. r. 1. t. 48. f.A.B.C.
16 chinensis. P.S.	Chinese.	China.	—	—	G. b.	Mich. arb. 3. t. 5.

ORDO CLXXXIV.

JUGLANDEÆ. DC. theor. ed. 1. p. 215.

JU'GLANS. N.	WALNUT-TREE.	Monœcia	Polyandria.	L.		
1 regia. w.	(gr.) common.	Persia.	1562.	4. 5. H.5.	Lam. ill. t. 781.	
β <i>máxima</i> . P.S.	(gr.) double-fruited.	_____	_____	— H.5.		
γ <i>serotina</i> . P.S.	(gr.) late-fruited.	_____	_____	— H.5.		
2 nigra. w.	(gr.) black.	N. America.	1629.	— H.5.	Wats. dend. brit. t. 158.	
3 cinerea. w.	(gr.) shell bark.	_____	1656.	— H.5.	_____ v. 1. t. 192.	
<i>cathartica</i> . Mich. arb. 1. t. 2.						
4 fraxinifolia. s.s.	(gr.) Ash-leaved.	_____	_____	— H.5.		
5 pterocarpa. s.s.	(gr.) winged-fruited.	Caucasus.	1828. H.5.		
<i>Rhú obscurum</i> . M.B.						
C'A ^T N.	HICKORY NUT.	Monœcia	Tetr-Hexandria.			
1 <i>virmis</i> . N.	(gr.) Pecan nut.	N. America.	4. 5. H.5.	Mich. arb. 1. t. 3.	
2 angustifolia. H.K.	(gr.) narrow-leaved.	_____	1766.	— H.5.	Mich. arb. 1. t. 8.	
3 sulcata. N.	(gr.) shell-bark.	_____	1804.	— H.5.	_____ 1. t. 7.	
4 alba. N.	(gr.) shag-bark.	_____	1730.	— H.5.		
<i>Juglans compræssa</i> . w.						
5 tomentosa. N.	(gr.) mocker nut.	_____	1629.	— H.5.	Mich. arb. 1. t. 6.	
<i>Juglans alba</i> . w. non M.						
β <i>máxima</i> . N.	(gr.) large-fruited.	_____	1824.	— H.5.		
6 pubescens. L.en.	(gr.) pubescent.	_____	— H.5.		
7 microcarpa. N.	(gr.) small-fruited.	_____	1824.	— H.5.		
8 amara. N.	(gr.) bitter nut.	_____	1800.	5. H.5.		
9 porcina. N.	(gr.) Pig nut.	_____	1812.	4. 5. H.5.	Mich. arb. 1. t. 9.f. 3-4.	
<i>Juglans obcordata</i> . w.						
10 glabra. w.	(gr.) smooth-leaved.	_____	1799. H.5.	Mich. arb. 1. t. 9. f. 1-2.	
<i>Juglans porcina</i> β <i>ficifórmis</i> . M.						
11 aquatica. N.	(gr.) water.	Carolina.	1826. H.5.	Mich. arb. 1. t. 5.	
12 integrifolia. s.s.	(gr.) entire-leaved.	Louisiana.	— H.5.		
<i>Hicorius integrifolius</i> . Rafin.						

ORDO CLXXXV.

CUPULIFERÆ. Kth. synops. 1. p. 353.

QUE'RCUS. W.	OAK.	Monœcia	Polyandria.	L.		
1 Phéllos. w.	(y.gr.) Willow-leaved.	N. America.	1723.	5. 6. H.5.	Mich. arb. 1. t. 12.	
β <i>humilis</i> . Ph.	(y.gr.) dwarf.	_____	1812.	— H. $\frac{1}{2}$.	Catesb. car. 1. t. 22.	
2 maritima. w.	(y.gr.) sea.	_____	1811.	— H. $\frac{1}{2}$.	Mich. querc. t. 13. f. 1.	
3 sericea. w.	(y.gr.) running.	_____	1724.	— H. $\frac{1}{2}$.	Mich. arb. 2. t. 15.	
4 virens. w.	(y.gr.) live.	_____	1739.	5. H.5.	_____ 2. t. 11.	
5 cinerea. w.	(y.gr.) ash-coloured.	_____	1789.	5. 6. H.5.	_____ 2. t. 14.	
6 imbricaria. w.	(y.gr.) tile-cupped.	_____	1786.	— H.5.	_____ 2. t. 13.	
7 laurifolia. w.	(y.gr.) Laurel-leaved.	_____	—	5. H.5.	Mich. querc. t. 17.	
β <i>obtusa</i> . Ph.	(y.gr.) blunt-leaved.	_____	—	— H.5.	_____ t. 18.	
8 lutea. w.	(y.gr.) yellow-leaved.	Mexico.	1825. H.5.	Banks ic. Kämp. t. 17.	
9 glauca. w.	(y.gr.) glaucous.	Japan.	1822. H.5.		
10 grammintia. w.	(y.gr.) Holly-leaved.	S. France.	1730.	6. H. $\frac{1}{2}$.		
11 Ballota. w.	(y.gr.) Barbary.	Barbary.	1818.	— H. $\frac{1}{2}$.		
12 rotundifolia. w.	(y.gr.) round-leaved.	Spain.	—	— H. $\frac{1}{2}$.		
13 <i>Ilex</i> . w.	(y.gr.) evergreen.	S. France.	1581.	5. 6. H. $\frac{1}{2}$.		
α <i>integrifolia</i> . H.K.	(y.gr.) entire-leaved.	_____	—	— H. $\frac{1}{2}$.		
β <i>serrata</i> . H.K.	(y.gr.) saw-leaved.	_____	—	— H. $\frac{1}{2}$.		
γ <i>oblonga</i> . H.K.	(y.gr.) long-leaved.	_____	—	— H. $\frac{1}{2}$.		
δ <i>fagifolia</i> . L.C.	(y.gr.) Beech-leaved.	_____	—	— H. $\frac{1}{2}$.		
ϵ <i>crispa</i> . L.C.	(y.gr.) curl-leaved.	_____	—	— H. $\frac{1}{2}$.		
14 variifolia.	(y.gr.) different-leaved.	S. Europe.	— H. $\frac{1}{2}$.		
<i>heterophylla</i> . Lam. non Ph.						

15 crenāta. Lam. (<i>y.gr.</i>) crenated.	S. Europe.	5. 6.	H. $\frac{1}{2}$.	
16 lusitânica. Lam. (<i>y.gr.</i>) Portugal.	Portugal.	1824.	—	H. $\frac{1}{2}$.	Cavan, icon. 2. t. 129.
17 calycina. Poir. (<i>y.gr.</i>) calycine.	S. Europe.	—	H. $\frac{1}{2}$.	
18 expânsa. Poir. (<i>y.gr.</i>) spreading.	—	—	—	H. $\frac{1}{2}$.	
19 Lezermiâna. Poir. (<i>y.gr.</i>) glossy-leaved.	—	—	—	H. $\frac{1}{2}$.	
20 castellâna. Bosc. (<i>y.gr.</i>) subdeciduous.	—	—	—	H. $\frac{1}{2}$.	
21 Süber. w. (<i>y.gr.</i>) Cork-tree.	S. France.	1699.	6.	H. $\frac{1}{2}$.	Wats. dend. brit. t. 89.
22 coccífera. w. (<i>y.gr.</i>) Kernes.	—	1683.	5.	H. $\frac{1}{2}$.	— t. 91.
23 prasina. Bosc. (<i>y.gr.</i>) glaucous-leaved.	Spain.	1824.	—	H. $\frac{1}{2}$.	
24 húmilis. P.S. (<i>y.gr.</i>) low.	S. Europe.	—	H. $\frac{1}{2}$.	
25 infectória. w. (<i>y.gr.</i>) Levant.	Levant.	1812.	H. $\frac{1}{2}$.	Lobel. ic. 2. p. 157.
26 Turnéni. W.en. (<i>y.gr.</i>) Turner's.	—	—	—	H. 5.	
27 Prinus. w. (<i>y.gr.</i>) Chestnut-leaved.	N. America.	1730.	5. 6.	H. 5.	Mich. arb. 2. t. 7.
28 bicolor. w. (<i>y.gr.</i>) two-coloured.	—	1811.	5.	H. 5.	Mich. arb. 2. t. 8.
29 montâna. w. (<i>y.gr.</i>) rock Chestnut.	—	1800.	—	H. 5.	— 2. t. 6.
30 Micháuxii. n. (<i>y.gr.</i>) swamp white. <i>Prinos discolor.</i> M.	—	1812.	H. 5.	Mich. arb. 2. t. 9.
31 Castânea. w. (<i>y.gr.</i>) yellow Chestnut.	—	1816.	5.	H. 5.	— 2. t. 10.
32 prinoídes. w. (<i>y.gr.</i>) dwarf Chestnut. <i>Chinquapin.</i> Ph.	—	—	—	H. $\frac{1}{2}$.	
33 agrifolia. w. (<i>y.gr.</i>) spiny-leaved.	—	1827.	H. 5.	Pluk. phyt. t. 196. f. 3.
34 heterophylla. Ph. (<i>y.gr.</i>) various-leaved.	—	—	—	H. 5.	Mich. arb. 2. t. 16.
35 aquática. Ph. (<i>y.gr.</i>) water.	—	1723.	—	H. 5.	— 2. t. 17.
36 hemisp'hærica. w. (<i>y.gr.</i>) hemispherical.	—	1816.	—	H. 5.	Mich. querc. t. 20. f. 2.
37 nâna. w. (<i>y.gr.</i>) dwarf.	—	1738.	—	H. $\frac{1}{2}$.	Abb. ins. 2. t. 59.
38 trilòba. w. (<i>y.gr.</i>) three-lobed.	—	1800.	—	H. 5.	Mich. querc. t. 26.
39 nigra. w. (<i>y.gr.</i>) black.	—	1739.	—	H. $\frac{1}{2}$.	Mich. arb. 2. t. 18.
40 tinctoria. w. (<i>y.gr.</i>) dyer's.	—	1800.	—	H. 5.	Mich. querc. t. 24.
41 discolor. w. (<i>y.gr.</i>) two-coloured.	—	1763.	—	H. 5.	— t. 25.
42 lanuginosa. D.P. (<i>y.gr.</i>) Nepaul.	Nepaul.	1818.	G. 5.	
43 Phullâta. D.P. (<i>y.gr.</i>) slender-pointed.	—	1824.	G. 5.	
44 coccinea. w. (<i>y.gr.</i>) scarlet.	N. America.	1691.	5.	H. 5.	Mich. arb. 2. t. 23.
45 ambigua. M. (<i>y.gr.</i>) gray.	—	1800.	—	H. 5.	— 2. t. 24.
46 rûbra. w. (<i>y.gr.</i>) champion.	—	1739.	—	H. 5.	— 2. t. 26.
47 Catesbæi. w. (<i>y.gr.</i>) barren scrub.	—	1820.	—	H. $\frac{1}{2}$.	— 2. t. 20.
48 falcatâ. M. (<i>y.gr.</i>) downy-leaved. <i>elongâta.</i> w.	—	1763.	—	H. 5.	— 2. t. 21.
49 palústris. w. (<i>y.gr.</i>) marsh.	—	1800.	—	H. 5.	Mich. arb. 2. t. 25.
50 Banistéri. M. (<i>y.gr.</i>) Banister's. <i>ilicifolia.</i> w.	—	—	—	H. $\frac{1}{2}$.	— 2. t. 19.
51 stellata. w. (<i>y.gr.</i>) starred. <i>obtusiloba.</i> M.	—	—	—	H. 5.	Mich. arb. 2. t. 4.
52 macrocárpa. w. (<i>y.gr.</i>) overcup white.	—	1800.	—	H. 5.	Mich. arb. 2. t. 3.
53 pseudo-süber. w. (<i>y.gr.</i>) cork-like.	Spain.	1818.	—	H. $\frac{1}{2}$.	Sant. itin. 156. t. 4.
54 Ægilops. w. (<i>y.gr.</i>) Velenida.	S. Europe.	1731.	H. 5.	Mill. dict. n. 7. t. 215.
55 álba. w. (<i>y.gr.</i>) white.	N. America.	1724.	5.	H. 5.	Mich. arb. 2. t. 1.
56 repânda. L.en. (<i>y.gr.</i>) repand.	—	—	—	H. 5.	Mich. querc. t. 5. f. 2.
57 E'sculus. w. (<i>y.gr.</i>) Italian.	S. Europe.	1739.	—	H. 5.	Eng. bot. v. 26. t. 1845.
58 Rôbor. w. (<i>y.gr.</i>) sessile-fruited. <i>sessiliiflora.</i> E.B.	Britain.	4. 5.	H. 5.	
59 pedunculâta. w. (<i>y.gr.</i>) common. <i>Robur. E.B. non w.</i>	—	—	H. 5.	Eng. bot. v. 19. t. 1342.
60 pubescens. w. (<i>y.gr.</i>) Durmast.	England.	—	H. 5.	
61 fastigiâta. Lam. (<i>y.gr.</i>) Cypress.	Pyrenees.	1822.	H. 5.	Second. d. chen. t. 5.
62 conglomerâta. P.S. (<i>y.gr.</i>) clustered.	Europe.	4. 5.	H. 5.	
63 olivâformis. M. (<i>y.gr.</i>) mossy-cup.	N. America.	1811.	5.	H. 5.	Mich. arb. 2. t. 2.
64 lyrâta. w. (<i>y.gr.</i>) lyre-leaved.	—	1786.	—	H. 5.	— 2. t. 5.
65 fagínea. P.S. (<i>y.gr.</i>) Beech-like.	S. Europe.	1824.	H. 5.	
66 haliphleos. P.S. (<i>y.gr.</i>) hispid-cupped.	France.	4. 5.	H. 5.	
67 Taùzin. P.S. (<i>y.gr.</i>) soft jagged-leav'd.	S. Europe.	—	H. 5.	
β laciniâta. Desv. (<i>y.gr.</i>) cut-leaved.	—	—	—	H. 5.	
γ digitâta. Desv. (<i>y.gr.</i>) fingered.	—	—	—	H. 5.	
68 Cérrius. w. (<i>y.gr.</i>) Turkey. <i>a frondosa.</i> H.K. (<i>y.gr.</i>) common.	Turkey.	1735.	—	H. 5.	Wats. dend. brit. t. 92.
β bullâta. H.K. (<i>y.gr.</i>) rough-leaved.	—	—	—	H. 5.	
γ sinuâta. H.K. (<i>y.gr.</i>) narrow-leaved.	—	—	—	H. 5.	
? δ dentâta. w.D. (<i>y.gr.</i>) Fulham.	—	—	—	H. 5.	
69 Lucombeâna. (<i>y.gr.</i>) Lucombe's. <i>exoniensis.</i> L.C.	Levant.	6.	H. 5.	Wats. dend. brit. t. 93.
		5. 6.	H. 5.	

70 austriaca. W. en. (<i>y.gr.</i>) Austrian.	Austria.	4. 5. H.5.	Clus. hist. 1. p. 20. c. ic.
71 mexicana. k.s. (<i>y.gr.</i>) Mexican.	Mexico. 1824. G. b.	H. et B. pl. æq. 2. t. 82.
72 spicata. k.s. (<i>y.gr.</i>) spiked. G. b.	2. t. 89.
73 Bonplandiæna. (<i>y.gr.</i>) Bonpland's. ambigua. k.s. non M. G. b. 2. t. 93.
FAGUS. W. BEECH. Monœcia Polyandria. L.	N. America.	5. H.5.	Mich. arb. 2. t. 8.
1 americana. (<i>y.gr.</i>) white. <i>sylvatica</i> β <i>americana</i> . N.	
2 sylvatica. w. (<i>y.gr.</i>) common. β <i>purpurea</i> . H.K. (<i>pu.</i>) dark purple.	Britain.	4. 5. H.5.	Eng. bot. v. 26. t. 1846.
3 ferruginea. w. (<i>y.gr.</i>) ferruginous.	Germany.	— H.5.	
4 comptoniaefolia. Df. (<i>y.gr.</i>) Fern-leaved. <i>sylvatica</i> γ <i>asplenifolia</i> . H.S.L.	N. America. 1796.	5. 6. H.5.	Mich. arb. 2. t. 9.
CASTANEA. G. CHESTNUT. Monœcia Polyandria. W.	N. America.	5. 6. H.5.	Mich. arb. 2. t. 6.
1 americana. P.S. (<i>y.gr.</i>) American. <i>væsca</i> β <i>americana</i> . N.	
2 vésca. w. (<i>y.gr.</i>) common. β <i>variegata</i> . (<i>y.gr.</i>) striped-leaved. γ <i>asplenifolia</i> . (<i>y.gr.</i>) fern-leaved.	England.	— H.5. — H.5. — H. b.	Eng. bot. v. 13. t. 886.
3 pumila. w. (<i>y.gr.</i>) dwarf.	N. America. 1699.	7. H. b.	Mich. arb. 2. t. 7.
CO'RYLUS. W. NUT-TREE. Monœcia Polyandria. L.	
1 Avellana. w. (<i>y.re.</i>) common Hazel. Britain.	2. 4. H.5.	Eng. bot. v. 11. t. 723.	
β <i>alba</i> . H.K. (<i>y.re.</i>) white Filbert.	— H.5.	
γ <i>rubra</i> . H.K. (<i>y.re.</i>) red Filbert.	— H.5.	
δ <i>ovata</i> . Lam. (<i>y.re.</i>) oval-fruited.	— H.5.	
ε <i>barcelonensis</i> . L.C. (<i>y.re.</i>) Barcelona.	Spain.	— H.5.	
ζ <i>grandis</i> . H.K. (<i>y.re.</i>) Cob-nut.	— H.5.	
η <i>glomerata</i> . H.K. (<i>y.re.</i>) clustered.	— H.5.	
2 tubulosa. w. (<i>y.re.</i>) Lambert's.	S. Europe. 1759.	3. 4. H.5.	Lam. ill. t. 780.
3 heterophylla. F. (<i>y.re.</i>) various-leaved.	Danube. 1829. H. b.	Wang. amer. t. 29. f. 63.
4 americana. w. (<i>st.re.</i>) Cuckold.	N. America. 1798.	3. 4. H.5.	
5 humili. w. (<i>st.re.</i>) dwarf cuckold.	— H. b.	Willd. arb. t. 1. f. 2.
6 rostrata. w. (<i>st.re.</i>) common cuckold.	— H.5.	
7 Colurna. w. (<i>st.re.</i>) Constantinople. Constantinople. 1665.	— H.5.	Wats. dend. brit. t. 99.

ORDO CLXXXVI.

BETULINEÆ. Kth. synops. 1. p. 363.

O'STRYA. W.	HOP-HORNBEAM. Monœcia Polyandria. W.		
1 vulgaris. w. (<i>gr.</i>) common.	Italy. 1724.	5. H.5.	Wats. dend. brit. t. 143.
2 virginica. w. (<i>gr.</i>) American.	N. America. 1692.	5. 6. H.5.	Abb. ins. 2. t. 75.
CARPINUS. W.	HORNBEAM. Monœcia Polyandria. L.		
1 Bétulus. w. (<i>gr.</i>) common.	Britain.	3. 5. H.5.	Eng. bot. v. 29. t. 2032.
β <i>variegata</i> . (<i>gr.</i>) striped-leaved.	— H.5.	
γ <i>incisa</i> . H.K. (<i>gr.</i>) cut-leaved.	— H.5.	
δ <i>quercifolia</i> . Desf. (<i>gr.</i>) Oak-leaved.	— H.5.	
2 americana. w. (<i>gr.</i>) American.	N. America. 1812.	— H.5.	Wats. dend. brit. t. 157,
3 orientalis. w. (<i>gr.</i>) eastern.	Levant. 1739.	5. 6. H.5.	— t. 98.
BETULA. W.	BIRCH. Monœcia Polyandria. L.		
1 álba. w. (<i>gr.</i>) common.	Britain.	4. 6. H.5.	Eng. bot. v. 31. t. 2198.
β <i>verrucosa</i> . Ehr. (<i>gr.</i>) warted.	— H.5.	
γ <i>pendula</i> . Roth. (<i>gr.</i>) weeping.	— H.5.	
δ <i>dalecarlica</i> . L. (<i>gr.</i>) palmate-leaved.	— H.5.	
ε <i>macrocarpa</i> . (<i>gr.</i>) large-fruited.	— H.5.	
2 pontica. Df. (<i>gr.</i>) eastern.	Asia.	— H.5.	
3 pubescens. P.S. (<i>gr.</i>) pubescent.	Europe.	4. 6. H.5.	Wats. dend. brit. t. 94.
4 populifolia. w. (<i>gr.</i>) Poplar-leaved.	N. America. 1750.	7. H.5.	Wats. dend. brit. t. 151,
5 excelsa. w. (<i>gr.</i>) tall.	1767.	— t. 95.
6 davurica. w. (<i>gr.</i>) Daurian.	Dauria. 1786.	7. H.5.	Pall. ros. 1. t. 39.
7 nigra. w. (<i>gr.</i>) black.	N. America. 1736.	7. 8. H.5.	Wats. dend. brit. t. 153.
8 lutea. m. (<i>gr.</i>) yellow.	1816.	
9 lanulosa. m. (<i>gr.</i>) woolly.	1812.	Willd. arb. t. 2. f. 1.
<i>rubra</i> . Mich. arb. 2. t. 3.			

BETULINEÆ.

10 papyrâcea. w. (gr.) paper.	N. America.	1750.	5. 6.	H. 5.	Wats. dend. brit. t. 151.
11 lénâta. w. (gr.) soft.	—	1759.	7.	H. 5.	— t. 144.
12 carpínifòlia. W.en.(gr.) Hornbeam-leaved.	—	1825.	—	H. 5.	
13 carpâthica. Kit. (gr.) Carpathian.	Carpathian mt.	1825.	—	H. 5.	
14 ovâta. w. (gr.) oval-leaved.	Europe.	—	6. 7.	H. h.	Wats. dend. brit. t. 96.
viridis. Vill. <i>Alnus viridis</i> . n.c.					
15 fruticôsa. w. (gr.) shrubby.	Siberia.	1816.	—	H. h.	Wats. dend. brit. t. 154.
16 glandulôsa. M. (gr.) glandular.	N. America.	—	5.	H. h.	
17 pùmila. w. (gr.) hairy dwarf.	—	1762.	4. 5.	H. h.	Wats. dend. brit. t. 97.
18 nâna. w. (gr.) smooth dwarf.	Scotland.	—	5.	H. h.	Eng. bot. v. 33. t. 2326.
β macrophylla. Sch.(gr.) large-leaved.	Switzerland.	1823.	—	H. h.	
19 tristis. Len. (gr.) dark.	Kamtschatka.	1824.	—	H. h.	
A'LNU.S. W.	ALDER. Monœcia Tetrandria. W.				
1 glutinôsa. w. (gr.) common.	Britain.	—	3. 4.	H. 5.	Eng. bot. v. 21. t. 1508.
β emarginâta. (gr.) emarginate.	—	—	—	H. 5.	
γ incisa. w. (gr.) cut-leaved.	—	—	—	H. 5.	
δ laciniâta. w. (gr.) jagged-leaved.	—	—	—	H. 5.	
ε querçifòlia. (gr.) Oak-leaved.	—	—	—	H. 5.	
2 incâna. w. (gr.) hoary-leaved.	Europe.	1780.	6.	H. 5.	
β angulâta. H.K. (gr.) angular-leaved.	—	—	—	H. 5.	
γ pinnâta. (gr.) pinnated.	—	—	—	H. 5.	
3 oblongâta. w. (gr.) oblong-leaved.	S. Europe.	1730.	7.	H. 5.	
β elliptica. (gr.) elliptic-leaved.	—	—	—	H. 5.	
4 undulâta. w. (gr.) undulate.	N. America.	1782.	5. 6.	H. h.	
crispâ. Ph. <i>Betula crispâ</i> . H.K.					
5 glauca. s.s. (gr.) glaucous.	—	1812.	3. 4.	H. 5.	
6 serrulâta. w. (gr.) notch-leaved.	—	1769.	3.	H. h.	Abb. ins. 2. t. 92.
7 rugosa. s.s. (gr.) rugged-leaved.	—	—	—	H. h.	
8 sibírica. F.	Siberia.	1820.	—	H. h.	
9 cordifolia. T.P. (gr.) heart-leaved.	Naples.	—	3. 4.	H. h.	Ten. flor. nap. t. 99.
10 macrophylla. Df. (gr.) large-leaved.	—	—	—	H. h.	
11 rûbra. Df. (gr.) red.	—	—	—	H. h.	
12 pùmila. L.C. (gr.) dwarf.	—	—	—	H. h.	
13 oxyacanthifòlia.L.C.(gr.)Hawthorn-leaved.	—	1812.	—	H. h.	
14 subrotunda. Df. (gr.) roundish-leaved.	—	1822.	—	H. h.	

ORDO CLXXXVII.

SALICEÆ. Kth. synops. I. p. 364.

SA'LIX. W.	WILLOW. Diœcia 1-2-3-5-andra-Monadelpha.				
1 præcox. s.s. (g.ye.) early.	Europe.	—	2. 3.	H. 5.	Salic. woburn. t. 26.
2 pomerânea. W.en.(g.ye.)Pomeranian.	Pomerania.	1816.	3. 5.	H. 5.	Eng. bot. v. 27. t. 1897.
3 cinèrea. E.B. (g.ye.) gray.	Britain.	—	5.	H. 5.	Salic. woburn. t. 48.
4 angustâta. Ph. (g.ye.) narrow-leaved.	Pensylvania.	1811.	3. 4.	H. 5.	Hoffm. salic. I. t. 17. f. 2.
5 Wulfeniâna.s.s. (br.st.)Wulfen's.	Scotland.	—	4. 5.	H. 5.	Salic. woburn. t. 35.
6 silesiaca. s.s. (g.ye.) Silesian.	Silesia.	1820.	—	H. 5.	Eng. bot. v. 23. t. 1617.
7 Ammanniâna.s.s.(g.ye.)Amman's.	Carinthia.	—	—	H. 5.	Flor. dan. t. 1238.
8 acutifòlia. s.s. (g.ye.) acute-leaved.	Caucasus.	1824.	2. 3.	H. 5.	
9 hastâta. w. (g.ye.) halberd-leaved.	Lapland.	1780.	5.	H. 5.	
10 malifòlia. E.B. (g.ye.) Apple-leaved.	England.	—	4.	H. 5.	
11 serrulâta. w. (g.ye.) saw-leaved.	Lapland.	1820.	5.	H. 5.	
hastâta. F.D.					
12 Pontederâna.w.(g.ye.)Pontedera's.	Germany.	1818.	—	H. 5.	Salic. woburn. t. 43.
13 phyllicifòlia.E.B.(g.ye.)Tea-leaved.	Scotland.	—	—	H. h.	Eng. bot. v. 28. t. 1958.
14 Andersoniâna.E.B.(g.ye.)Anderson's.	—	—	—	H. h.	— v. 33. t. 2343.
15 tenuifòlia. E.B. (g.ye.) slender-leaved.	Britain.	—	5. 6.	H. h.	— v. 31. t. 2186.
16 coruscans. w. (g.ye.) glittering.	Switzerland.	1824.	—	H. 5.	Jacq. aust. t. 408.
Arbúscula. Jacq. non E.B.					
17 frágilis. E.B. (g.ye.) crack.	Britain.	—	4. 5.	H. 5.	Eng. bot. v. 26. t. 1807.
18 Russelliâna.E.B.(g.ye.)Bedford.	England.	—	—	H. 5.	— v. 26. t. 1808.
19 Trevirâni. L.en. (g.ye.) Treviranus's.	Germany.	1825.	—	H. 5.	
20 hippophæfòlia.L.en.(g.ye.)Sea-buckthorn-ld.—	—	1823.	—	H. 5.	
21 viréscens. s.w. (g.ye.) greenish.	Switzerland.	—	—	H. 5.	Salic. woburn. t. 7.

22 triándra. E.B. (g.ye.) long-leaved.	Britain.	5. 8.	H.5.	Eng. bot. v. 20. t. 1435.	
23 amygdalina. E.B. (g.ye.) Almond-leaved.	4. 5.	H.5.	v. 27. t. 1936.	
24 Hoppeána. w. (g.ye.) Hoppe's.	Switzerland.	1817.	—	H.5.		
25 ambíguā. Ph. (gr.ye.) ambiguous.	N. America.	1813.	3. 4.	H.5.		
26 decípiens. E.B. (gr.ye.) varnished.	England.	5.	H.5.	Eng. bot. v. 27. t. 1937.	
27 vitellina. E.B. (gr.ye.) yellow twigged.	3. 5.	H.5.	v. 20. t. 1389.	
28 álba. E.B. (gr.ye.) common white.	Britain.	4. 5.	H.5.	v. 34. t. 2430.
29 cœrulaea. E.B. (gr.ye.) blue-leaved.	England.	5. 8.	H.5.	v. 34. t. 2431.
30 pedicelláris.s.s.(gr.ye.) pedicellated.	N. America.	1811.	4. 5.	H.5.		
pensylvánica. Salic. Woburn. t. 95. ?						
31 babylónica. w. (g.st.) weeping.	Levant.	1692.	5. 6.	H.5.	Salic. woburn. t. 22.	
32 annuláris. s.w. (g.st.) annular-leaved.	1823.	—	H.5.	t. 21.
33 myricóides. s.s. (g.ye.) Myrica-like.	N. America.	1811.	4.	H.5.	Ann. bot. 2. t. 5. f. 2.	
34 lúcida. s.s. (g.ye.) shining.	—	5.	H.5.	2. t. 5. f. 7.
35 rígida. s.s. (g.ye.) rigid.	—	4. 5.	H.5.	2. t. 5. f. 4.
36 cordátā. s.s. (gr.ye.) heart-leaved.	—	—	H. h.	2. t. 5. f. 8.
37 nigra. s.s. (g.ye.) black.	—	5.	H.5.	2. t. 5. f. 5.
38 riparia. w. (g.ye.) rivulet.	Europe.	1820.	—	H.5.		
39 incána. DC. (g.ye.) hoary.	—	4.	H.5.	Salic. woburn. t. 90.	
40 incanéscens. Sch. (g.ye.) incandescent.	Switzerland.	1823.	3. 4.	H.5.	t. 120.
41 lanáta. E.F. (gr.ye.) woolly.	Scotland.	—	—	H. h.	t. 71.
42 pentándra. E.B. (gr.ye.) Bay-leaved.	Britain.	5. 6.	H. h.	Eng. bot. v. 26. t. 1805.	
43 Meyeriana. s.s. (g.st.) Meyer's.	Germany.	1823.	4. 5.	H. h.	Salic. woburn. t. 33.	
44 Forbesii. (g.st.) Forbes's.	Switzerland.	—	—	H.5.	t. 32.
lúcida. s.w. nec aliorum.						
45 tetraspérra. w. (g.ye.) four-seeded.	E. Indies.	1796.	—	S. h.	Roxb. corom. 1. t. 97.	
46 gariepina. B.T. (g.ye.) Cape.	C. B. S.	1816.	—	G. h.		
47 Humboldtiana. w. (bl.) Humboldt's.	Peru.	1821.	G. h.	H. et B. 2. t. 99, 100.	
48 Bonplandiána. k.s. (br.) Bonpland's.	Mexico.	—	G. h.	2. t. 101, 102.
49 Houstoniana. Ph. (gr.ye.) Houston's.	America.	1812.	4. 5.	H. h.	Salic. woburn. t. 11.	
50 Lyoni. s.w. (g.ye.) Lyon's.	Switzerland.	1816.	—	H. h.	t. 10.
51 finmárchica. s.s. (g.ye.) Finnmark.	Sweden.	1821.	—	H. h.	Flor. dan. t. 1051.	
52 Villarsiána. w. (g.ye.) Villars's.	S. France.	1818.	5. 6.	H. h.	Salic. woburn. t. 17.	
53 myrtilloïdes. s.s. (br.ye.) Myrtle-leaved.	Sweden.	1772.	4. 8.	H. h.	t. 66.
54 retusa. w. (g.ye.) blunt-leaved.	Italy.	1763.	—	H. h.	f. 139.
55 serpyllifòlia. w. (g.ye.) Thyme-leaved.	Switzerland.	1818.	—	H. h.	t. 65.
56 Kitaibeliána. w. (g.ye.) Kitaibel's.	Carpathian m.	1816.	—	H. h.	t. 64.
57 'Uva-úrsi. s.s. (g.ye.) Bearberry-like.	Labrador.	1811.	4. 5.	H. h.	f. 151.
58 berberifòlia. w. (g.ye.) Barberry-leaved.	Davuria.	1824.	—	H. h.	Pall. ros. 2. t. 82.	
59 herbácea. E.B. (g.ye.) least.	Britain.	6.	H. h.	Eng. bot. v. 27. t. 1907.	
60 monándra. Ard. (g.ye.) monandrous.	Europe.	3. 4.	H. h.	Salic. woburn. t. 4.	
61 purpùrea. E.B. (g.ye.) bitter purple.	England.	3.	H. h.	Eng. bot. v. 20. t. 1388.	
62 Hélix. E.B. (g.ye.) Rose.	Britain.	3. 4.	H. h.	v. 19. t. 1343.
63 Lambertiána. E.B. (g.ye.) Boyton.	England.	—	H. h.	v. 19. t. 1359.
64 rúbra. E.B. (g.ye.) red-stemmed.	—	4. 3.	H. h.	v. 16. t. 1145.
65 Forbiána. E.B. (g.ye.) basket osier.	—	4.	H. h.	v. 19. t. 1344.
66 viminális. E.B. (g.ye.) common osier.	Britain.	4. 5.	H. h.	v. 27. t. 1898.
67 mollíssima. s.s. (g.ye.) softest.	Germany.	3. 4.	H. h.	Salic. woburn. t. 134.	
68 Smithiána. s.s. (g.ye.) Smith's.	England.	—	H. h.		
mollissima. Eng. bot. v. 21. t. 1509. nec aliorum.						
69 acumináta. E.B. (g.ye.) acuminate.	Britain.	4.	H.5.	Eng. bot. v. 20. t. 1434.	
70 holosericea. w. (g.ye.) holosericeous.	Germany.	1822.	—	H.5.		
71 aurita. w. (g.ye.) eared.	Britain.	4. 5.	H.5.	Hoff. sal. 1. t. 22. f. 1.	
72 uliginósa. W.en. (g.ye.) marsh.	—	4. 6.	H.5.	Eng. bot. v. 21. t. 1487.	
aurita. E.B. non w.						
73 paludósa. L.en. (g.ye.) fen.	Germany.	—	H.5.		
74 aquática. E.B. (g.ye.) water.	Britain.	4.	H.5.	Eng. bot. v. 20. t. 1437.	
75 spatuláta. W.en. (g.ye.) spatulate.	Germany.	—	H.5.		
76 cáprea. E.B. (g.ye.) great round-leaved.	Britain.	4. 5.	H.5.	Eng. bot. v. 21. t. 1488.	
77 sphacélata. E.B. (g.ye.) withered-pointed.	Scotland.	—	H. h.	v. 33. t. 2333.
78 laurína. w. (g.ye.) Laurel-leaved.	England.	—	H. h.	v. 26. t. 1806.
bicolor. E.B. non Ehr.						
79 bicolor. w. (g.ye.) two-coloured.	Germany.	1818.	—	H. h.		
80 Forsteriana. E.B. (g.ye.) Forster's.	Scotland.	—	H.5.	Eng. bot. v. 33. t. 1403.	
81 nígricans. E.B. (g.ye.) dark broad-leaved.	England.	4.	H.5.	v. 17. t. 1213.
82 violacea. s.s. (g.ye.) violet.	Russia.	3. 4.	H.5.	Andr. deposit. t. 581.	
83 conformis. s.w. (g.ye.) uniform-leaved.	—	2. 3.	H.5.	Salic. woburn. t. 24.	
84 Weigliána. s.s. (g.ye.) long-styled.	Silesia.	1820.	—	H.5.	t. 51.
cotinifòlia. E.B. (g.ye.) Quince-leaved.	Britain.	4.	H.5.	Eng. bot. v. 20. t. 1403.	

SALICEÆ.

86 conifera. s.s. (g.ye.) cone-bearing.	N. America.	1800.	5.	H. 5.	Wang. amer. t. 31. f. 72.
87 prinoïdes. s.s. (g.ye.) Prinos-like.	1811.	3. 4.	H. 5.	Salic. woburn. t. 40.	
88 fuscata. s.s. (g.ye.) brown-stemmed.	—	—	H. 5.		
89 recurvata. s.s. (g.ye.) recurve-flowered.	—	4.	H. h.		
90 tristis. s.s. (g.ye.) linear-leaved.	1765.	3. 4.	H. h.	Salic. woburn. f. 150.	
91 Muhlenbergiana.(re.ye.)Muhlenberg's.	1811.	4.	H. h.	Ann. bot. 2. t. 5. f. 9.	
92 grisea. W.en. (br.ye.) brown.	1816.	—	H. h.	Eng. bot. v. 20. t. 1436.	
93 lanceolata. E.B. (g.ye.) sharp-leaved.	England.	4.	5. H. h.	— v. 17. t. 1214.	
94 stipularis. E.B. (g.ye.) auricled.	Britain.	3.	H. h.	Salic. woburn. t. 13.	
95 undulata. s.s. (g.ye.) wave-leaved.	Germany.	1819.	4. 5.	— t. 12.	
96 virgata. s.w. (g.ye.) twiggy.	—	5. 6.	H. h.	— t. 16.	
97 Hoffmanniana. (g.ye.) Hoffman's.	Switzerland.	—	5.	H. h.	— t. 19.
98 montana. w. (g.ye.) mountain.	—	—	5. 6. H. 5.	— t. 30.	
99 monspeliensis.s.w.(g.ye.)Montpelier.	Montpelier.	—	4. 5.	H. h.	— t. 39.
100 pataens. s.w. (g.ye.) spreading.	—	—	5. 8.	H. h.	— t. 41.
101 Willdenowiana.s.w.(g.ye.)Willdenow's.	—	—	4. 8.	H. h.	— t. 42.
102 crispa. s.w. (g.st.) curled.	—	—	3.	H. h.	— t. 44.
103 nitens. s.w. (g.ye.) glossy.	Scotland.	—	4. 5.	H. h.	— t. 45.
104 Borreriiana.s.w.(g.st.)Borrer's.	—	—	4.	H. h.	— t. 47.
105 Davalliana.s.w.(br.st.)Davall's.	—	—	—	H. h.	Eng. bot. v. 16. t. 1147.
106 petiolaris.E.B.(br.st.)dark long-leaved.	N. America.	—	4.	H. h.	Eng. bot. v. 20. t. 1402.
107 pataula. s.s. (g.ye.) spreading.	Italy.	1818.	—	H. h.	Salic. woburn. t. 91.
108 oleifolia. E.B. (g.ye.) Olive-leaved.	Britain.	—	3.	H. 5.	Ann. bot. 2. t. 5. f. 1.
109 candida. s.s. (r.st.) hoary.	N. America.	1811.	2. 3.	H. h.	Eng. bot. v. 26. t. 1809.
110 discolor. s.s. (g.ye.) brown-branched.	—	—	4.	H. h.	— v. 3. t. 183.
111 arenaria. E.B. (g.ye.) sand.	Scotland.	—	5.	H. h.	Salic. woburn. t. 79.
112 repens. E.B. (br.ye.) creeping.	Britain.	—	5.	H. h.	Eng. bot. v. 19. t. 1365.
113 incubacea.w. (bh.ye.) trailing.	—	1775.	—	H. h.	— v. 19. t. 1366.
114 rosmarinifolia.E.B.(g.ye.)Rosemary-leaved.	—	—	4. 5.	H. h.	— v. 28. t. 1960.
115 arbúscula.E.B.(g.ye.)little tree.	Scotland.	—	4.	H. h.	— v. 28. t. 1961.
116 fusca. E.B. (g.st.) brown.	Britain.	—	5.	H. h.	— v. 28. t. 1959.
117 parvifolia.E.B.(g.ye.)small-leaved.	England.	—	4. 5.	H. h.	
118 adscendens.E.B.(g.ye.)ascending.	—	—	5.	H. h.	
119 prostrata.E.B. (g.ye.) prostrate.	Britain.	—	5.	H. h.	
120 canescens.W.en.(g.ye.)canescent.	Germany.	1815.	4. 5.	H. h.	
121 leucophylla.W.en.(g.ye.)white-leaved.	—	—	—	H. h.	
122 elaeagnoides.L.en.(g.ye.)Elaeagnus-like.	Switzerland.	1823.	4. 8.	H. h.	Salic. woburn. t. 69.
123 argentea. E.B. (g.ye.) silky sand.	Britain.	—	5.	H. h.	Eng. bot. v. 19. t. 1364.
124 procumbens.s.w.(g.ye.)procumbent.	Scotland.	—	—	H. h.	Salic. woburn. t. 61.
125 ramifusa.s.w.(g.ye.)brown branched.	Britain.	—	4. 7.	H. h.	— t. 53.
126 floribunda.s.w.(g.ye.)many flowered.	—	—	—	H. h.	— t. 54.
127 versifolia. s.s. (g.ye.) various-leaved.	N. Europe.	1824.	5.	H. h.	Eng. bot. v. 19. t. 1361.
128 prunifolia.E.B.(g.ye.)Plum-leaved.	Scotland.	—	4. 5.	H. h.	— v. 20. t. 1390.
129 Dicksoniana.E.B.(g.ye.)Dickson's.	—	—	5.	H. h.	
130 humiliis. W.en. (g.ye.) humble.	—	1824.	—	H. h.	
131 formosa.w. (g.ye.) handsome.	Switzerland.	1820.	—	H. h.	Eng. bot. v. 19. t. 1360.
132 myrsinifolia.E.B.(g.ye.)Whortle-leaved.	Scotland.	—	4. 6.	H. h.	— v. 19. t. 1362.
133 venulosa. E.B. (g.ye.) veiny-leaved.	—	—	4. 5.	H. h.	— v. 33. t. 2341.
134 vaccinifolia.E.B.(g.ye.)Bilberry-leaved.	—	—	—	H. h.	— v. 19. t. 1363.
135 carinata. E.B. (g.ye.) folded-leaved.	—	—	—	H. h.	
136 arbutifolia. w. (g.ye.) Arbutus-leaved.	Switzerland.	1818.	—	H. h.	Eng. bot. v. 33. t. 2342.
137 rupestris. E.B. (g.ye.) rock.	Scotland.	—	4.	H. h.	Salic. woburn. t. 73.
138 lappónum. w. (g.ye.) Lapland.	Lapland.	—	—	H. h.	
139 helvética. Vil. (g.ye.) Swiss.	Switzerland.	1824.	4. 8.	H. h.	
140 leucophylla.w.(g.ye.)white-leaved.	—	—	—	H. h.	
141 lívida. s.s. (g.ye.) livid.	Lapland.	—	4.	H. h.	Salic. woburn. t. 63.
142 polaris. s.s. (g.ye.) polar.	—	—	4. 7.	H. h.	Ann. bot. 2. t. 5. f. 6.
143 longifolia. s.s. (g.ye.) long-leaved.	N. America.	1819.	4. 5.	H. h.	Eng. bot. v. 26. t. 1810.
144 glauca. E.B. (g.ye.) glaucous.	Scotland.	—	5.	H. h.	Salic. woburn. t. 74.
145 sericea. w. (wh.ye.) silky.	Alps Europe.	1816.	—	H. h.	Eng. bot. v. 36. t. 2586.
146 Stuartiana.E.B.(w.ye.)Stuart's.	Scotland.	—	7. 8.	H. h.	— v. 27. t. 1908.
147 reticulata. E.B. (re.) wrinkled.	Britain.	—	6. 7.	H. h.	— v. 20. t. 1404.
148 hírtia. E.B. (g.ye.) hairy-branched.	England.	—	4. 5.	H. 5.	— v. 16. t. 1146.
149 Croweaña.E.B.(g.ye.)Crowe's.	—	—	—	H. h.	Salic. woburn. f. 143.
150 fagifolia. w. (g.ye.) Beech-leaved.	Hungary.	1819.	—	H. h.	Salic. woburn. f. 148.
151 cordifolia. Ph. (g.ye.) heart-shaped.	N. America.	1811.	—	H. h.	— t. 49.
152 planifolia. Ph. (g.ye.) flat-leaved.	Labrador.	—	—	H. h.	
153 falcata. Ph. (g.ye.) sickle-leaved.	N. America.	—	—	H. h.	
154 tetrapla. L.en. (g.ye.) elegant.	Scotland.	1824.	4.	H. h.	

155 obtusa. L.en. (g.ye.) blunt-leaved.	Switzerland.	1820.	4.	H. h.	
156 eriántha. L.en. (g.ye.) woolly-flowered.	—	—	—	H. h.	Vill.delph.3. t. 50. f.11.
157 cæsia. Vill. (g.ye.) cæsious.	N. Europe.	—	4. 5.	H. h.	
158 ulmifolia.W.en.(g.ye.)Elm-leaved.	Switzerland.	—	4.	H. h.	
159 foliosa. w. (g.ye.) leafy.	Lapland.	1818.	—	H. h.	
160 obtusifolia. w. (g.ye.) obtuse-leaved.	—	—	—	H. h.	
161 atrovirens.s.w.(g.ye.)dark-green.	Switzerland.	1824.	—	H. h.	
162 australis. s.w. (g.ye.) southern.	—	—	5.	H. h.	Salic. woburn. t. 108.
163 carpiniifolia.s.w.(g.ye.)Hornbeam-leaved.	Germany.	—	3. 4.	H. h.	t. 103.
164 cydoniifolia.Sch.(g.ye.)Cydonia-leaved.	Switzerland.	—	—	H. h.	
165 subalpina.s.w.(g.ye.)subalpine.	—	—	4. 5.	H. h.	Salic. woburn. t. 93.
166 proteæfolia.s.w.(g.ye.)Protea-leaved.	—	1820.	—	H. h.	t. 75.
167 strípida. s.w. (g.ye.) creaking.	—	—	3. 4.	H. h.	t. 100.
168 pyrenæica. s.s. (g.ye.) Pyrenean.	Pyrenees.	1823.	4.	H. h.	
169 Jacquiniana.w.(g.ye.)Jacquin's. fusca. Jacquin. nec aliorum.	Austria.	1818.	—	H. h.	Jacq. aust. t. 409.
170 Schraderianum.w.(g.ye.)Schrader's.	Germany.	1820.	—	H. h.	
171 finnárhica.w.(g.ye.)Swedish.	Sweden.	1825.	4. 5.	H. h.	
172 serrulata. w. (g.ye.) serrulate.	Lapland.	1822.	—	H. h.	
173 versicolor.s.w.(g.ye.) various-coloured.	Switzerland.	—	5.	H. h.	Salic. woburn. t. 77.
174 Doniana.E.F. (br.ye.) Mr. Don's.	Scotland.	4. 5.	H. h.	t. 88.
175 linearis.s.w. (br.st.) linear-leaved.	Switzerland.	1820.	—	H. h.	t. 89.
176 villosa. s.w. (g.ye.) villous.	—	—	H. h.	t. 92.
177 reflexa. s.w. (g.ye.) reflexed.	—	3.	H. h.	t. 94.
178 petræa. s.w. (g.ye.) rock-side.	Britain.	4.	H. h.	t. 97.
179 grisonensis.s.w.(g.ye.)Grisons.	Grisons.	1820.	3. 4.	H. h.	t. 99.
180 sórdida. s.w. (g.st.) sordid.	Switzerland.	—	4.	H. h.	t. 101.
181 glabrata. Sch. (g.ye.) smooth.	—	1823.	4. 5.	H. h.	
182 decumbens.s.w.(g.ye.)decumbent.	—	—	5.	H. h.	Salic. woburn. t. 88.
183 heterophylla.Deb.(g.ye.)different-leaved.	—	—	—	H. h.	
184 murina. Sch. (g.ye.) wall.	Switzerland.	1824.	3. 4.	H. h.	
185 nervosa. Sch. (g.ye.) nerved.	—	—	—	H. h.	
186 pannosa. s.w. (g.ye.) cloth-leaved.	—	—	4. 5.	H. h.	Salic. woburn. t. 123.
187 pallescens.Sch.(g.st.) pallid.	—	—	—	H. h.	
188 pallida. s.w. (g.ye.) pale.	—	—	5.	H. h.	Salic. woburn. t. 96.
189 alaternoïdes.s.w.(ye.) Alaternus-like.	—	1823.	4. 5.	H. h.	t. 76.
190 albescens.Sch. (g.st.) whitish.	—	—	—	H. h.	
191 rivularis.s.w. (g.ye.) rivulet.	—	1824.	5.	H. h.	Salic. woburn. t. 102.
192 rotundata.s.w. (g.ye.) rounded-leaved.	—	—	4. 5.	H. h.	t. 104.
193 dura. s.w. (g.ye.) hardy.	—	—	—	H.5.	t. 105.
194 firma. s.w. (g.ye.) firm.	—	—	—	3. 4. H. h.	t. 106.
195 Ansoniana.s.w.(g.ye.)Anson's.	—	—	—	H. h.	t. 107.
196 coriacea. s.w. (g.ye.) leathery.	—	1822.	3.	H. h.	t. 112.
197 crassifolia.s.w.(g.ye.) thick-leaved.	—	—	4. 5.	H. h.	t. 115.
198 vaudensis.s.w.(g.ye.) Vaudois.	Switzerland.	1824.	3. 4.	H. h.	t. 117.
199 latifolia. s.w. (g.ye.) broad-leaved.	—	—	3.	H. h.	t. 118.
200 ferruginea.s.w.(br.st.)ferruginous.	Britain.	4.	H. h.	t. 128.
201 geminata. s.w. (g.ye.) twin-flowered.	—	—	3.	H. h.	t. 129.
202 macrostipulacea.s.w.(g.y.)large-stipul'd.	Switzerland.	1822.	4. 5.	H.5.	t. 130.
203 Micheliana.s.w.(g.ye.)Michel's.	—	—	4.	H.5.	t. 135.
204 atropurpurea.s.w.(g.ye.)dark-purple.	—	1824.	—	H. h.	
205 damascena.s.w.(g.ye.) Damask.	—	—	3. 4.	H. h.	
206 mutabilis. s.w. (g.ye.) changeable.	—	—	—	H. h.	
207 cerasifolia.Sch.(g.ye.)Cherry-leaved.	—	—	4.	H. h.	
208 cinnamomea.Sch.(g.ye.)Cinnamon-scented.	—	—	4. 5.	H. h.	
209 clethraefolia.Sch.(g.ye.)Clethra-leaved.	—	—	—	H. h.	
210 grisophylla.s.w.(g.ye.)gray-leaved.	—	—	4.	H. h.	Salic. woburn. t. 119.
211 lacustris. s.w. (g.ye.) lake.	—	—	3.	H. h.	t. 116.
212 mespiliifolia.Sch.(g.ye.)Mespilus-leaved.	—	—	4. 5.	H. h.	
213 obtusiferrata.Sch.(g.ye.)bluntly-sawed.	—	—	—	H. h.	
214 pyrifolia. Sch. (g.ye.) Pear-leaved.	—	—	—	H. h.	
215 Schleicheriana.s.w.(g.ye.)Schleicher's.	—	—	—	H. h.	Salic. woburn. t. 98.
216 velutina.W.en.(g.ye.)velvety.	—	1826.	3. 4.	H. h.	
217 Starkeana. w. (g.ye.) Starke's.	Silesia.	1820.	4. 5.	H. h.	
PO'PULUS. L.	POPLAR.	Dicacia Octandria. L.			
1 álba. E.B. (fl.) Abele-tree.	Britain.	3. 4.	H.5.	Eng. bot. v. 23. t. 1618.
2 canescens. E.B. (fl.) gray.	England.	—	H.5.	v. 23. t. 1619.
3 nivea. w. (fl.) snowy.	Europe.	—	H.5.	
4 trépida. w. (fl.) trembling.	N. America.	1812.	—	H.5.	Mich. arb. 3. t. 8. f. 1.

5 trémula. w.	(fl.) Aspen.	Britain.	3. 4.	H.5.	Eng. bot. v. 27. t. 1909.
6 lavigata. w.	(fl.) smooth.	N. America.	1769.	—	H.5.	Mich. arb. 3. t. 11.
7 tremuloïdes. M.	(fl.) trembling-like.	—	1811.	—	H.5.	Duham. arb. 2. t. 53.
8 gr'eca. w.	(fl.) Athenian.	Archipelago.	1779.	—	H.5.	— 184. t. 54.
9 nigra. w.	(fl.) black.	Britain.	—	H.5.	Eng. bot. v. 27. t. 1910.
10 acladæsa. Lind.	(re.) black Italian.	N. America.	5.	H.5.	Mich. arb. 3. t. 10. f. 1.
11 hudsonica. m.	(fl.) American black.	—	—	H.5.	
<i>betulifolia. Ph.</i>						
12 dilatata. w.	(sc.) Lombardy.	Italy.	1758.	—	H.5.	
13 viminea. Df.	(fl.) slender-twigg'd.	N. America.	1824.	4. 5.	H.5.	
14 monilifera. w.	(fl.) Canadian.	Canada.	1772.	3. 5.	H.5.	Wats. dend. brit. t. 102.
15 angulata. w.	(fl.) Carolina.	Carolina.	1738.	3.	H.5.	Mich. arb. 3. t. 12.
16 grandidentata. M.	(fl.) large-toothed.	N. America.	1812.	4.	H.5.	
<i>β pendula. N.</i>						
17 balsamifera. w.	(gr.) Tacamahac.	—	1692.	—	H.5.	Mich. arb. 3. t. 13. f. 1.
18 canticans. w.	(gr.) heart-leaved.	—	1772.	3.	H.5.	Catesb. car. 1. t. 34.
19 heterophylla. w. (gr.)	various-leaved.	—	1765.	4. 5.	H.5.	Mich. arb. 3. t. 9.
20 laurifolia. Led.	(fl.) Laurel-leaved.	Altay.	1827.	H.5.	

ORDO CLXXXVIII.

PLATANEÆ.

PLATANUS. W.	PLANE-TREE.	Monœcia Polyandria. L.				
1 orientalis. w.	(gr.) oriental.	Levant.	1548.	4. 5.	H.5.	Wats. dend. brit. t. 101.
2 cuneata. w.	(gr.) wave-leaved.	—	1739.	—	H.5.	
3 acerifolia. w.	(gr.) Maple-leaved.	—	1724.	—	H.5.	Wats. dend. brit. t. 100.
4 occidentalis. w.	(gr.) American.	N. America.	1640.	—	H.5.	
5 mexicana.	(gr.) Mexican.	Mexico.	1826.	F.5.	
LIQUIDA'MBAR. W.	SWEET-GUM-TREE.	Monœcia Polyandria. L.				
1 Styraciflua. w. (gr. wh.)	Maple-leaved.	N. America.	1683.	3. 4.	H.5.	Catesb. car. 2. t. 65.
2 imberbe. w. (gr. wh.)	oriental.	Levant.	1759.	H.5.	

ORDO CLXXXIX.

MYRICEÆ. Kth. synops. 1. p. 361.

COMPTONIA. W.	COMPTONIA.	Monœcia Triandria. W.				
asplenifolia. w. (br.)	Fern-leaved.	N. America.	1714.	3. 5.	H. h.	Wats. dend. brit. t. 166.
MYRICA. W.	CANDLEBERRY-MYRTLE.	Dioecia Tetrandria. L.				
1 Gale. w.	(br.) Sweet Gale.	Britain.	5.	H. h.	Eng. bot. v. 8. t. 562.
2 cerifera. w.	(br.) common.	N. America.	1699.	5. 6.	H. h.	Catesb. car. 1. t. 69.
3 carolinensis. Ph.	(br.) broad-leaved.	—	1730.	5.	H. h.	1. t. 13.
4 pensylvanica. Ph.	(br.) Pennsylvanian.	—	—	—	H. h.	Duham. arb. ed. n. 2. t. 55.
5 Faya. w.	(br.) Azorian.	Azores.	1777.	6. 7.	F. h.	— 2. t. 56.
6 mexicana. W. en.	(br.) Mexican.	Mexico.	1821.	6. 8.	G. h.	
7 esculenta. D.P.	(br.) eatable.	Nepaul.	—	—	G. h.	Jacq. ic. 3. t. 625.
8 segregata. w.	(br.) netted-leaved.	S. America.	—	—	G. h.	Pluk. alm. t. 48. f. 8.
9 æthiopica. w.	(br.) African.	C. B. S.	1795.	6. 7.	G. h.	Burm. afr. t. 98. f. 1.
10 serrata. w.	(br.) saw-leaved.	—	1793.	8.	G. h.	Jacq. frag. 2. t. 1. f. 4.
11 lacinata. W. en.	(br.) smooth Oak-ld.	—	1752.	6. 7.	G. h.	Pluk. alm. t. 319. f. 7.
12 quercifolia. W. en.	(br.) hairy Oak-leav'd.	—	—	—	G. h.	
13 cordifolia. w.	(br.) heart-leaved.	—	1759.	5. 7.	G. h.	
14 integrifolia. H. B.	(br.) entire-leaved.	E. Indies.	1824.	—	S. h.	
NAGEIA. W.	NAGEIA.	Dioecia Tetrandria.				
1 japonica. w. (gr.)	Japan.	Japan.	1812.	2. 4.	G. h.	
<i>Buxus dioica. Hort. Myrica Nagi. Thunb.</i>						
2 Putranjiva. H. B.	(gr.) grey-barked.	E. Indies.	1822.	—	S. h.	
CASUARINA. W.	SHRUBBY HORSETAIL.	Monœcia Monandria. W.				
1 equisetifolia.w. (br. re.)	Horse-tail-leav'd. S. Sea Island.	1793.	S. h.	Rumph. amb. 3. t. 57.	
2 nodiflora. w.	(br. re.) knotted-flower'd.	—	1818.	S. h.	

3 glauca. s.s. (<i>br.re.</i>) glaucous.	N. S. W.	1818.	11. 4.	G. h.	
4 paludosa. s.s. (<i>br.re.</i>) marsh.	_____	1822.	—	G. h.	
5 stricta. w. (<i>br.re.</i>) upright.	_____	1775.	11. 2.	G. h.	Andr. deposit. t. 346.
6 distyla. w. (<i>br.re.</i>) two-styled.	_____	1812.	—	G. h.	Vent. dec. pl. nov. t. 62.
7 quadrivalvis. P.S. (<i>br.re.</i>) four-valved.	N. Holland.	—	—	G. h.	Lab. nov. hol. 2. t. 218.
8 torulosa. w. (<i>br.re.</i>) cork-barked.	N. S. W.	1772.	4. 6.	G. h.	
9 muricata. L.en. (<i>br.re.</i>) muricate.	E. Indies.	1816.	S. h.	
10 tenuissima. s.s. (<i>br.re.</i>) slender.	N. S. W.	1824.	4. 6.	G. h.	
11 nana. s.s. (<i>br.re.</i>) dwarf.	_____	—	—	G. h.	

ORDO CXC.

HAMAMELIDEÆ. Brown.

HAMAM'ELIS. W.	WITCH-HAZEL.	Tetrandria Digynia. L.			
1 virginica. w. (<i>ye.</i>) Virginian.	N. America.	1736.	11. 5.	H. h.	Lodd. bot. cab. t. 598.
2 macrophylla. Ph. (<i>ye.</i>) large-leaved.	_____	1812.	—	H. h.	
FOTHERGILLA. W.	FOTHERGILLA.	Polyandria Digynia. L.			
1 alnifolia. w. (<i>wh.</i>) obtuse-leaved.	N. America.	1765.	4. 6.	H. h.	Bot. mag. t. 1341.
2 major. B.M.	(<i>wh.</i>) large-leaved.	_____	—	5. 6.	H. h. t. 1342.
3 Gardéni. Jacq.	(<i>wh.</i>) acute-leaved.	_____	—	—	H. h. Jacq. ic. 1. t. 100.
4 serotina. B.M.	(<i>wh.</i>) green-leaved.	_____	—	—	H. h.

ORDO CXCI.

CONIFERÆ. Juss. gen. 411.

SUBORDO I. TAXINÆ. Richard.

SALISBURIA. R.	MAIDENHAIR-TREE.	Monœcia Polyandria. H.K.				
adiantifolia. L.T. (g.st.) Japan.	Japan.	1754.	4. 5.	H.5.	Wats. dend. brit. t. 168.	
Gingko biloba. L.						
PHYLLOCLADUS. R.	PHYLLOCLADUS.	Monœcia Monadelphia.				
rhomboïdalis. R. (<i>ye.</i>) Fern-leaved.	V. Diem. Isl.	1825.	G. h.	Rich. conif. pl. 3.	
Podocarpus asplenifolia. Labill. nov. holl. 2. t. 221.						
PODOCA'RPU'S. L'H.	PODOCA'RPU'S.	Monœcia Monadelphia.				
1 Chilinus. R.	(<i>ye.</i>) sharp-leaved.	C. B. S.	1774.	7.	G. h.	
2 elongatus. R.	(<i>ye.</i>) Chinese.	China.	—	5. 7.	H. h.	
Taxus chinensis. H.B.						
3 coriaceus. R.	(<i>ye.</i>) leathery-leaved.	_____	1824.	—	G. h.	
4 verticillatus.	(<i>ye.</i>) whorl-leaved.	Japan.	1812.	G. h.	
5 macrophyllus. L.en. (<i>ye.</i>) large-leaved.	China.	1804.	7. 8.	G. h.	Kämpf. ic. t. 24.	
6 nereifolius. L.P.	(<i>ye.</i>) Wax-Dammar.	P.Wales Isl.	1809.	S. h.	Rumph. amb. 3. t. 26.
TA'XUS. W.	YEW-TREE.	Dicecia Monadelphia. L.				
1 baccata. w.	(<i>ye.</i>) common.	Britain.	2. 4.	H.5.	
2 hibernica.	(<i>ye.</i>) Irish.	Ireland.	—	H. h.	
3 canadensis. w.	(<i>ye.</i>) Canadian.	Canada.	1818.	—	H.5.	
4 nucifera. w.	(<i>ye.</i>) Acorn-bearing.	China.	1764.	G. h.	
DACRY'DIUM. L.P.	DACRY'DIUM.	Monœcia Monadelphia.				
cupressinum. L.P. (<i>ye.</i>) N. Zealand spruce.	N. Zealand.	1825.	G. h.	Kämpf. amœn. t. 815.	
Thalamia cupressina. s.s.					Lamb. pin. v. 1. t. 41.	

SUBORDO II. CUPRESSINÆ. Richard.

CALLITRIS. R.	CALLITRIS.	Monœcia Monadelphia.			
1 quadrivalvis. R.	(<i>ye.</i>) jointed.	Barbary.	1815.	2. 5.	F. h.
Thuya articulata. Vahl. symb. 2. t. 48.					Rich. conif. pl. 8. f. 1.
2 arenosa.	(<i>ye.</i>) sand.	N. Holland.	1824.	F. h.
3 pyramidalis.	(<i>ye.</i>) pyramidal.	_____	—	F. h.

CONIFERÆ.

4 australis.	(ye.) southern.	N. S. W.	4. 5. F. $\frac{1}{2}$.	
Cupréssus australis. P.S.					
5 oblonga. R.	(ye.) oblong-scaled.	—	1822.	— G. $\frac{1}{2}$.	Rich. conif. pl. 18. f. 2.
6 rhomboidea. R.	(ye.) rhomboid-scal'd.	N. Holland.	—	— G. $\frac{1}{2}$.	— pl. 18. f. 1.
TH'UJA. W.	ARBOR-VITÆ.	Monœcia Monadelphia. L.			
1 occidentalis. w.	(ye.) American.	N. America. 1596.	5.	H. $\frac{1}{2}$.	Wats. dend. brit. t. 150.
2 pyramidalis. T.P.	(ye.) pyramidal.	—	1824.	— H. $\frac{1}{2}$.	Wats. dend. brit. t. 149.
3 orientalis. w.	(ye.) Chinese.	China. 1752.	2. 3.	H. $\frac{1}{2}$.	Lamb. pin. v. 2. p. 19. t.
4 plicata. L.P.	(ye.) Nee's.	Nootka sound. 1796.	—	H. $\frac{1}{2}$.	Lamb. pin. app. t. 1.
5 cupressoides. w.	(ye.) African.	C. B. S. 1799.	—	G. $\frac{1}{2}$.	Wats. dend. brit. t. 156.
6 dolabrata. w.	(ye.) ovate-leaved.	China. 1822.	—	H. $\frac{1}{2}$.	
7 sphaeroidea. s.s. (ye.) White Cedar.		N. America. 1736.	4. 5.	H. $\frac{1}{2}$.	
Cupréssus thyoides. w.					
8 pendula. L.P.	(ye.) weeping.	Tartary. 1810.	2. 5.	H. $\frac{1}{2}$.	Lamb. pin. v. 2. t. 51.
CUPRE'SSUS. W.	CYPRESS.	Monœcia Monadelphia. L.			
1 sempervirens. w.	(ye.) common.	Candia. 1548.	5.	H. $\frac{1}{2}$.	Pall. ros. 2. t. 53.
α stricta.	(ye.) upright.	—	—	H. $\frac{1}{2}$.	
β horizontalis.	(ye.) horizontal.	—	—	H. $\frac{1}{2}$.	
2 torulosa. D.P.	(ye.) twisting.	Nepaul.	1826.	— H. $\frac{1}{2}$.	
3 lusitánica. w.	(ye.) Cedar of Goa.	Goa. 1683.	4. 5.	F. $\frac{1}{2}$.	Lamb. pin. 95. t. 49.
4 pendula. Th.	(ye.) pendulous.	Japan. 1818.	—	F. $\frac{1}{2}$.	— t. 50.
5 bacicórmis. W.en. (ye.) berry-like.		—	—	4. 5. H. $\frac{1}{2}$.	
6 juniperoides. w.	(ye.) African.	C. B. S. 1756.	—	G. $\frac{1}{2}$.	
TAX'ODIUM. Rich.	TAX'ODIUM.	Monœcia Monadelphia.			
disticum. Rich.	(ye.) deciduous.	America. 1640.	5.	H.5.	Mich. arb. 3. t. 1.
Cupréssus disticha. w.	Schubertia. Mirb.				
α pátens. H.K.	(ye.) spreading.	—	—	H.5.	
β nūtans. H.K.	(ye.) long-leaved.	—	—	H.5.	
JUNI'PERUS. W.	JUNIPER.	Diœcia Monadelphia. L.			
1 thurifera. w.	(ye.) Spanish.	S. Europe. 1752.	5. 6.	H. $\frac{1}{2}$.	Pluk. alm. t. 197. f. 4.
2 barbadensis. w.	(ye.) Barbadoes.	Barbadoes. 1811.	—	G. $\frac{1}{2}$.	Herm. lugd. t. 347.
3 bermudiána. w.	(ye.) Bermudas Cedar.	Bermudas. 1683.	—	F. $\frac{1}{2}$.	
4 chinénsis. w.	(ye.) Chinese.	China. 1804.	—	H. $\frac{1}{2}$.	
5 excélsa. w.	(ye.) tall.	Siberia. 1806.	—	H. $\frac{1}{2}$.	
6 Hermánni. s.s.	(ye.) Hermann's.	—	—	H. $\frac{1}{2}$.	
7 Sabina. w.	(ye.) Savin.	S. Europe. 1548.	—	H. $\frac{1}{2}$.	
β variegata.	(ye.) variegated.	—	—	H. $\frac{1}{2}$.	
8 tamariscifolia.	(ye.) Tamarisk-leav'd.	—	—	H. $\frac{1}{2}$.	
9 prostrata. P.S.	(ye.) prostrate.	N. America. ...	—	H. $\frac{1}{2}$.	
10 daúrica. A.R.	(ye.) Daurian.	Dauria. 1791.	6. 8.	H. $\frac{1}{2}$.	Andr. deposit. t. 534.
11 virginiana. w.	(ye.) red Cedar.	N. America. 1664.	5. 6.	H. $\frac{1}{2}$.	Mich. arb. 3. t. 5.
12 glauca. W.en.	(ye.) glaucous.	China. 1814.	—	H. $\frac{1}{2}$.	
13 squamata. D.P.	(ye.) scaly-branched.	Nepaul. 1824.	—	H. $\frac{1}{2}$.	
14 recúrva. D.P.	(ye.) recurve-branch'd.	—	1822.	5. 6. H. $\frac{1}{2}$.	
15 commúnis. w.	(ye.) common.	Britain. ...	—	H. $\frac{1}{2}$.	Eng. bot. v. 16. t. 1100.
16 suécica. Mill.	(ye.) Swedish.	N. Europe. ...	—	H. $\frac{1}{2}$.	
17 nána. w.	(ye.) dwarf.	Siberia. 1815.	—	H. $\frac{1}{2}$.	Pall. ros. 2. t. 54. f.A.B.
sibírica. Burgsd. <i>communis</i> γ . H.K.					
18 hemisphærica. s.s. (ye.) hemispherical.	Sicily.	1828.	—	H. $\frac{1}{2}$.	
19 oblonga. M.B.	(ye.) oblong.	Armenia. 1829.	—	H. $\frac{1}{2}$.	
20 Oxycedrus. w.	(ye.) brown-berried.	Spain. 1739.	—	H. $\frac{1}{2}$.	Duham. arb. 2. t. 128.
21 phœnica. w.	(ye.) Phœnician.	S. Europe. 1683.	—	H. $\frac{1}{2}$.	Pall. ros. 2. t. 57.
22 ly'cia. w.	(ye.) Lycian.	—	1693.	— H. $\frac{1}{2}$.	— 2. t. 56.
23 capensis. Lam.	(ye.) Cape.	C. B. S. 1816.	—	G. $\frac{1}{2}$.	
24 drupacea. Lab.	(ye.) drupaceous.	—	—	H. $\frac{1}{2}$.	Lab. pl. syr. 2. t. 8.
EPH'EDRA.	EPH'EDRA.	Diœcia Monadelphia. L.			
1 distachya. w.	(st.) great.	France. 1570.	6. 7.	H. $\frac{1}{2}$.	Schk. hand. 3. t. 339.
2 monostachya. w.	(st.) small.	Siberia. 1722.	9. 11.	H. $\frac{1}{2}$.	Wats. dend. brit. t. 142.
3 altissima. w.	(st.) tall.	Barbary. 1823.	—	H. $\frac{1}{2}$.	Desf. atl. 2. t. 253.

SUBORDO III. ABIETINÆ. Richard.

DA'MMARA. L.P.	DAMMAR PINE.	Monœcia Monadelphia. L.en.			
1 australis. L.P.	(ye.) Cowrie-tree.	N. Zealand. 1823.	G. $\frac{1}{2}$.	Lamb. pin. 2. p. 14. t. 6.
2 orientalis. L.P.	(ye.) Amboyna pitch-tree.	Amboyna. 1804.	S. $\frac{1}{2}$.	— v.1.t.38.-v.2.p.16.
loranthifolia. L.en.	<i>Pinus Dammara</i> . w.	<i>A'gathis loranthifolia</i> . Sal.			
ARAUCARIA. J.	ARAUC'RIA.	Diœcia Monadelphia. W.			
1 imbricata. w.	(ye.) Chile Pine.	Chile. 1796.	H.5.	Lamb. pin. 2. p. 9. t.4.B.

2 brasiliæna. L.P.	(ye.) Brazil Pine.	Brazil.	1816.	G.5.	Lamb. pin. 2. p.12. t. 5.
3 excelsa. H.K.	(ye.) Norfolk Island.	Norfolk Isl.	1793.	G.5.	——— 1. t. 39. 40.
Dombéya excelsa. L.P.						
4 Cunninghamii.	(ye.) Cunningham's.	N. Holland.	1827.	G.5.	
B'ELIS. L.T.	B'ELIS.	Monœcia Monadelphia.				
lanceolata.	(ye.) spear-leaved.	China.	1804.	12. 1.	G. $\frac{1}{2}$.	Lamb. pin. 1. t. 34.
jaculifolia. L.T.	<i>Pinus lanceolata.</i> L.P.	<i>Cunninghamia lanceolata.</i> Bot. mag. t. 2743.				
P'INUS. W.	PINE OF FIR.	Monœcia Monadelphia. W.				
P'INUS.	PINE-TREE.					
1 sylvestr. w.	(ye.) Scotch.	Scotland.	5.	H.5.	Lamb. pin. 1. t. 1.
2 Laricio. P.S.	(ye.) Corsican.	Corsica.	1814.	4. 5.	H.5.	——— v. 2. p. 28. t. 9.
3 uncinata. DC.	(ye.) hooked.	Pyrenees.	1820.	—	H.5.	
4 Pumilio. w.	(ye.) upright-coned.	Carniola.	1779.	—	H.5.	Lamb. pin. p. 5. t. 2.
5 Mughus. w.	(ye.) nodding-coned.	Austria.	—	H.5.	Jacq. ic. 1. t. 193.
6 pungens. m.	(ye.) prickly-coned.	N. America.	1804.	H.5.	Mich. arb. 1. t. 5.
7 Banksiana. Ph.	(ye.) Hudson's Bay.	Hudson's bay.	1785.	5. 6.	H.5.	Lamb. pin. p. 7. t. 3.
8 Pallasiæna. L.P.	(ye.) Pallas's.	Crimea.	1804.	—	H.5.	——— v. 2. p. 1. t. 1.
9 Pinaster. w.	(ye.) cluster.	S. Europe.	1596.	4. 5.	H.5.	——— v. 1. p. 9. t. 4. 5.
10 Pineæ. w.	(ye.) stone.	———	1548.	5.	H.5.	——— v. 1. p. 11. t. 6. 7. 8.
11 maritima. w.	(ye.) maritime.	———	1759.	5. 6.	H.5.	——— v. 2. p. 30. t. 10.
12 halepensis. w.	(ye.) Aleppo.	Levant.	1683.	5.	H.5.	——— v. 1. p. 15. t. 11.
13 înops. Ph.	(ye.) Jersey.	N. America.	1739.	—	H.5.	——— v. 1. p. 18. t. 13.
14 resinosa. Ph.	(ye.) pitch.	———	1756.	—	H.5.	——— v. 1. p. 20. t. 14.
15 variabilis. Ph.	(ye.) 2 and 3-leaved.	———	1739.	5. 6.	H.5.	——— v. 1. p. 22. t. 15.
16 canariensis. DC.	(ye.) Canary.	Canaries.	1815.	3. 4.	G.5.	DC. pl. rar. gen. t. 1-2.
17 Tæda. Ph.	(ye.) frankincense.	N. America.	1713.	5. 6.	H. $\frac{1}{2}$.	Lam.p.v.1.p.23.t.16.17.
18 lutea. Walt.	(ye.) yellow.	———	H.5.	
19 serotina. Ph.	(ye.) Fox-tail.	———	1713.	5. 6.	H.5.	Mich. arb. 1. t. 7.
20 rigida. Ph.	(ye.) three-leaved.	———	1759.	—	H.5.	Lamb. pin. p. 25. t. 18. 19.
21 ponderosa. D.	(ye.) ponderous.	———	1827.	H.5.	
22 palustris. Ph.	(ye.) swamp.	———	1730.	H.5.	Lamb. p. 27. t. 20.
23 longifolia. w.	(ye.) long-leaved.	E. Indies.	1801.	S.5.	——— p. 29. t. 21.
24 Gerardi. Wal.	(ye.) Gerard's.	Nepaul.	1824.	H.5.	
25 occidentalis. w.	(ye.) West Indian.	W. Indies.	1820.	S. $\frac{1}{2}$.	Plum. sp. 17. t. 161.
26 aduncâa. Bosc.	(ye.) crooked.	1822.	H.5.	
27 româa. h. Bel.	(ye.) Roman.	Italy.	H.5.	
28 sibîrica. Dut.	(ye.) Siberian.	Siberia.	—	H.5.	
STR'OBUS.	WEYMOUTH PINE.					
29 Cembra. w.	(ye.) Siberian stone.	Siberia.	1746.	5.	H.5.	Lamb. pin. p. 34. t. 23. 24.
30 pygmæa. F.	(ye.) pygmy.	———	1816.	H. $\frac{1}{2}$.	
31 Strôbus. w.	(ye.) Weymouth.	N. America.	1705.	4.	H.5.	Lamb. pin. p. 31. t. 22.
32 Lambertiæna. L.P.	(ye.) Lambert's.	California.	1827.	H.5.	——— pin. c. icon.
33 excelsa. D.P.	(ye.) Bhotan.	Nepaul.	1822.	H.5.	——— v. 2. p. 5. t. 3.
C'EDRUS.	CEDAR.					
34 Cèdrus. w.	(ye.) Cedar of Lebanon.	Levant.	1683.	5.	H.5.	Lamb. pin. v.1.p.59.t.37
35 Deodâra. L.P.	(ye.) Indian Cedar.	Nepaul.	1822.	H.5.	——— v. 2. p. 8. t. 4.
L'ARIX.	LARCH.					
36 pendula. w.	(ye.re.) black Larch.	N. America.	1739.	5.	H.5.	Lamb. pin. v.1.p.56.t.36.
37 microcarpa. w.	(ye.re.) red Larch.	———	1760.	—	H.5.	——— p. 58. t. 37.
38 Lärix. w.	(ye.re.) common Larch.	Germany.	1629.	3. 4.	H.5.	——— p. 53. t. 35.
39 dahurica. F.	(ye.re.) Dahurian.	Dahuria.	1827.	H.5.	
40 intermédia. F.	(ye.re.) intermediate.	Altay.	1828.	H.5.	
PE'UCE.	SILVER FIR.					
41 canadensis. w.	(ye.) hemlock Spruce.	N. America.	1736.	5.	H.5.	Lamb. pin. p. 50. t. 32.
42 dumosa. D.P.	(ye.) bushy.	Nepaul.	1820.	H.5.	
43 taxifolia. Ph.	(ye.) Yew-leaved.	Columbia.	1822.	H.5.	Lamb. pin. v.1.p.51.t.33
44 spectabilis. D.P.	(ye.) purple-coned.	Nepaul.	1825.	H.5.	——— v. 2. p. 3. t. 2.
45 Picea. w.	(ye.) Silver-Fir.	Germany.	1603.	5.	H.5.	——— 1. p. 36. t. 40.
46 Pichta. F.	(ye.) Fischer's.	Altay.	1824.	H. $\frac{1}{2}$.	
47 Douglasiæna. D.	(ye.) Douglas's.	N. America.	1827.	H.5.	
48 Balsamea. w.	(ye.) Balm of Gilead.	———	1696.	5.	H.5.	Lamb. pin. v.1. p.48.t.31
49 Fraseri. Ph.	(ye.) double Balsam.	Penssylvania.	1811.	H.5.	
ABIES.	SPRUCE FIR.					
50 nigra. Ph.	(ye.) black Spruce.	N. America.	1700.	5.	H.5.	Lamb. pin. v.1.p.41.t.27
51 rubra. Ph.	(ye.) red Spruce.	———	1755.	—	H.5.	——— v. 1. p. 43. t. 28.
52 Clanbrassiliæna. L.	en. (ye.) dwarf.	———	1810.	H. $\frac{1}{2}$.	
53 álba. Ph.	(ye.) white Spruce.	———	1700.	5. 6.	H.5.	Lamb. pin. v.1.p.39.t.26
54 'Abies. w.	(ye.) Norway Spruce.	N. Europe.	1548.	4.	H.5.	——— v. 1. p. 37. t. 25.
55 orientalis. w.	(ye.) oriental.	Levant.	1824.	H.5.	——— v. 1. p. 45. t. 29.

CLASSIS II.

MONOCOTYLEDONEÆ seu ENDOGENÆ. DC.

* SUBCLASSIS 1. PHANEROGANEÆ. DC.

ORDO CXII.

CYCADEÆ. *Brown prodr.* 346.

ZAMIA. W.	ZAMIA. Diœcia Polyandria. W.	C. B. S.	1775.	6. 8.	G. ḥ.	Till. pis. 129. t. 45.
1 púngens. w.	(br.) needle.	—	—	—	—	Jacq. frag. 1. t. 25. 26.
2 cycadifolia. w.	(br.) Cycas-leaved.	—	—	—	—	
3 tridentata. w.	(br.) three-toothed.	—	1820.	—	—	
4 angustifolia. Jac.	(br.) narrow-leaved.	Bahama Isl.	—	—	S. ḥ.	Jacq. ic. 3. t. 636.
5 tenuis. w.	(br.) slender-leaved.	—	1823.	—	S. ḥ.	
6 media. w.	(br.) intermediate.	W. Indies.	—	—	S. ḥ.	
7 débilis. w.	(br.) weak.	—	1777.	—	S. ḥ.	Bot. mag. t. 1838.
8 integrifolia. w.	(br.) dwarf.	—	1768.	—	S. ḥ.	Comm. hort. 1. t. 58.
9 pygmæa. B.M. (g. br.)	least.	—	—	5.	S. ȳ.	Bot. mag. t. 1851.
10 furfuracea. w.	(br.) chaffy.	—	1691.	7. 8.	S. ḥ.	— t. 1741.
11 spiralis. B.P.	(br.) spiral.	N. S. W.	1796.	—	G. ȳ.	Trew. ehret. t. 61.
12 longifolia. w.	(br.) long-leaved.	C. B. S.	1820.	6. 7.	G. ḥ.	Jacq. frag. 1. t. 29.
13 lanuginosa. w.	(br.) woolly.	—	—	—	G. ḥ.	— 1. t. 30. 31.
14 Cycadis. w.	(br.) Cycas-like.	—	1775.	5. 6.	G. ḥ.	Thunb. act. ups. 2. t. 6.
15 hórrida. w.	(br.) glaucous spiny.	—	1800.	6. 8.	G. ḥ.	Jacq. frag. t. 27. 28.
16 latifolia. L.C.	(br.) broad-leaved.	—	—	—	G. ḥ.	
17 prunifera. L.C.	(br.) Plum-like.	—	—	—	S. ḥ.	
18 pùmila. L.C.	(ye.) small.	—	1812.	—	S. ȳ.	Bot. mag. t. 2006.
19 repanda. L.C.	(br.) repand.	—	—	—	S. ḥ.	
CYCAS. W.	CYCAS. Diœcia Polyandria. W.	E. Indies.	1700.	5. 6.	S. ḥ.	Bot. mag. t. 2826. 2827.
1 circinalis. w.	(br.) broad-leaved.	—	—	—	—	
2 glauca. L.en.	(br.) glaucous-leaved.	—	1814.	—	S. ḥ.	Linn. trans. 6. t. 29.
3 revoluta. w.	(br.) narrow-leaved.	China.	1737.	7. 8.	S. ḥ.	
4 squamosa. L.C.	(br.) scaly.	—	—	—	S. ḥ.	

ORDO CXIII.

HYDROCHARIDEÆ. *Brown prodr.* 344.

VALLISNERIA. L.	VALLISNERIA. Diœcia Diandria. L.					
spiralis. L.	(wh.) spiral.	S. Europe.	1818.	G.w. ȳ.	Mich.n.g.12.t.10.f.1.2.
HYDROCHARIS. L.	FROG-BIT. Diœcia Enneandria. L.					
Mórsus-rànæ. L.	(wh.) common.	Britain.	—	6. 7.	H.w. ȳ.	Eng. bot. v. 12. t. 808.
DAMAS'ONIUM. W.	DAMAS'ONIUM. Hexandria Polygynia. W.					
1 indicum. w.	(wh.) Indian.	E. Indies.	1800.	7. 9.	S.w. ȳ.	Bot. mag. t. 1201.
2 ovalifolium. B.P.	(wh.) oval-leaved.	N. S. W.	1820.	—	G.w. ȳ.	
STRATIOTES. W.	WATER-SOLDIER. Diœcia Dodecandria. L.					
aloïdes. w.	(wh.) Aloe-like.	England.	—	6. 7.	H.w. ȳ.	Eng. bot. v. 6. t. 379.

ORDO CXCIV.

ALISMACEÆ. Kth. synops. 1. p. 261.

SAGITTARIA. W.	ARROW-HEAD.	Monœcia Polyandria. L.						
1 sagittifolia. w.	(wh.) common.	England.	6. 8.	H.w. ȝ.	Eng. bot. v. 2. t. 84.		
2 latifolia. Ph.	(wh.) broad-leaved.	N. America.	1816.	—	H.w. ȝ.			
β flôre pleno.	(wh.) double-flowered.	—	—	—	H.w. ȝ.			
3 sinensis. B.M.	(wh.) Chinese.	China.	1812.	9. 11.	G.w. ȝ.	Bot. mag. t. 1731.		
4 obtusifolia. s.s.	(wh.) blunt-leaved.	—	1804.	7. 8.	G.w. ȝ.	Rheed. mal. 11. t. 45.		
5 Doniana.	(wh.) Don's.	Nepaul.	1820.	—	G.w. ȝ.			
hastata. D.P. non Ph.								
6 hastata. Ph.	(wh.) hastate-leaved.	N. America.	1816.	—	H.w. ȝ.			
7 obtusa. Ph.	(wh.) obtuse-leaved.	—	1822.	6. 9.	H.w. ȝ.			
8 heterophylla. Ph.	(wh.) various-leaved.	—	—	6. 8.	H.w. ȝ.			
9 lancifolia. w.	(wh.) lance-leaved.	W. Indies.	1787.	6. 7.	G.w. ȝ.	Bot. mag. t. 1792.		
10 angustifolia. B.R.	(wh.) narrow-leaved.	—	1827.	4. 8.	S.w. ȝ.	Bot. reg. t. 1141.		
11 falcata. Ph.	(wh.) falcate-leaved.	Carolina.	1811.	6. 7.	H.w. ȝ.			
12 rígida. B.M.	(wh.) brittle-leaved.	N. America.	1806.	—	H.w. ȝ.	Bot. mag. t. 1632.		
13 graminea. Ph.	(wh.) grass-leaved.	Carolina.	1812.	7. 8.	H.w. ȝ.			
14 nàtans. M.	(wh.) floating.	—	—	—	H.w. ȝ.			

ACTINOCA'RPUS. B.P. ACTINOCA'RPUS. Hexandria Hex-Octagynia.

1 minor. B.P.	(wh.) small.	N. S. W.	5. 8.	G.w. ȝ.			
2 Damasonium.	(wh.) common.	England.	6. 8.	H.w. ȝ.	Eng. bot. t. 1615.		

Alisma Damasonium, E.B. *Damasonium stellatum*, P.S.

ALISMA. B.P. WATER-PLANTAIN. Hexandria Polygynia. L.

1 Plantago. w.	(wh.) greater.	Britain.	6. 7.	H.w. ȝ.	Eng. bot. t. 837.		
2 lanceolata. With.	(wh.) spear-leaved.	—	—	—	H.w. ȝ.			
3 trivialis. Ph.	(wh.) blunt-leaved.	N. America.	1816.	—	H.w. ȝ.			
4 parnassifolia. w.	(wh.) Parnassus-leav'd.	Italy.	1824.	—	H.w. ȝ.			
5 nàtans. E.B.	(wh.) floating.	Wales.	7. 8.	H.w. ȝ.	Eng. bot. t. 775.		
6 ranunculoïdes. E.B. (w.) lesser.		Britain.	8.	H.w. ȝ.	—	t. 326.	
7 rédens. E.F.	(wh.) creeping.	Wales.	9. 10.	H.w. ȝ.	Cavan. ic. 1. t. 55.		

ORDO CXCV.

BUTOMEÆ. Kth. synops. 1. p. 260.

B'UTOMUS. L.	FLOWERING RUSH.	Enneandria Hexagynia. L.						
1 umbellatus. L.	(ro.) umbelled.	Britain.	6. 7.	H.w. ȝ.	Eng. bot. v. 10. t. 651.		
2 latifolius. D.P.	(wh.) broad-leaved.	Nepaul.	1823.	—	H.w. ȝ.			
LIMNOCH'ARIS. K.S.	LIMNOCH'ARIS.	Polyandria Polygynia. B.M.						
Plumieri. k.s.	(ye.) Plumier's.	S. America.	1822.	6. 10.	S. ♂.	Bot. mag. t. 2525.		
emarginata. H. et B. pl. æq. 1. t. 34.	<i>Alisma flava</i> . L.							

ORDO CXCVI.

JUNCAGINEÆ. Kth. synops. 1. p. 258.

SCHEUCHZ'ERIA. L.	SCHEUCHZ'ERIA.	Hexandria Trigynia. L.						
palustris. E.B. (br.gr.)	marsh.	England.	5. 6.	H.w. ȝ.	Eng. bot. v. 26. t. 1801.		
TRIGLO'CHIN. L.	ARROW-GRASS.	Hexandria Trigynia. L.						
1 elatum. N.	(gr.) long-spiked.	N. America.	1820.	5. 8.	H.w. ȝ.			
2 maritimum. E.B.	(gr.) sea-side.	Britain.	—	H.w. ȝ.	Eng. bot. v. 4. t. 255.		
3 palustre. E.B.	(gr.) marsh.	—	—	7. 8.	H.w. ȝ.	—	v. 6. t. 366.	
4 decipiens. B.P.	(gr.) New Holland.	N. S. W.	1820.	—	G. ȝ.			
5 Barrellieri. s.s.	(gr.) Barrelier's.	Italy.	1825.	—	H. ȝ.			
6 bulbosum. B.M. (pu.ye.)	bulbous-rooted.	C. B. S.	1806.	10.	G. ȝ.	Bot. mag. t. 1445.		

ORDO CXCVII.

FLUVIALES. *Kth. synops. 1. p. 131.*

POTAMOG'ETON. L.	POND-WEED.	Tetrandria Tetragynia. L.						
1 nātans. E.B.	(br.) broad-leaved.	Britain.	7. 8. H.w. 4.	Eng. bot. v. 26. t. 1822.			
2 heterophyllum. E.B.	(br.) various-leaved.	———	7. 9. H.w. 4.	———	v. 18. t. 1285.		
3 péroliātum. E.B.	(br.) perfoliate.	———	7. 8. H.w. 4.	———	v. 3. t. 168.		
4 dénsum. E.B.	(br.) close-leaved.	———	5. 7. H.w. 4.	———	v. 6. t. 397.		
5 flūtans. E.B.	(br.) long-leaved.	———	7. 9. H.w. 4.	———	v. 18. t. 1286.		
6 lūcens. E.B.	(br.) shining.	———	6. 8. H.w. 4.	———	v. 6. t. 376.		
7 lanceolātum. E.B.	(br.) spear-leaved.	England.	7. 8. H.w. 4.	———	v. 28. t. 1985.		
8 crispum. E.B.	(br.) curled.	Britain.	6. 7. H.w. 4.	———	v. 15. t. 1012.		
9 compréssum. E.B.	(br.) flat-stalked.	———	———	———	v. 6. t. 418.		
10 cuspidātum. E.F.	(br.) sharp-pointed.	———	7. H.w. 4.	Loes. pruss. t. 66.			
11 grāmīnum. E.B.	Grass-leaved.	———	7. 8. H.w. 4.	Eng. bot. v. 32. t. 2253.			
12 pusíllum. E.B.	(br.) small.	———	———	H.w. 4.	———	v. 3. t. 215.	
13 pectinātum. E.B.	(br.) Fennel-leaved.	———	6. 7. H.w. 4.	———	v. 5. t. 323.		
RU'PPIA. L.	RU'PPIA.	Tetrandria Tetragynia. L.						
marítima. E.B.	(br.) sea.	Britain.	8. 9. H.w. 4.	Eng. bot. v. 2. t. 136.			
APONOG'ETON. L.	APONOG'ETON.	Hex-Dodecandra Tri-Tetragynia.						
1 monostachyon. w.(wh.)	simple-spiked.	E. Indies.	1803.	8. 10. S.w. 4.	Andr. reposit. t. 406.			
2 distächyon. w.	(wh.) broad-leaved.	C. B. S.	1788.	5. 7. G.w.b.	Bot. mag. t. 1293.			
3 angustifòlium. w.(wh.)	narrow-leaved.	———	———	4. 9. G.w.b.	———	t. 1268.		
N'AJAS. L.	N'AJAS.	Monœcia Tetrandria. L.						
monospermá. w. (gr.)	single-seeded.	Europe.	1816.	7. 9. H.w. ⊖.	Mich.N.gen.11. t.8. f.2.			
ZANNICHELLIA. L.	WATER-WEED.	Monœcia Monandria. L.						
palústris. E.B.	(gr.) horned.	Britain.	7. 8. H.w. ⊖.	Eng. bot. v. 26. t. 1844.			
ZOSTERA. W.	SEA MATWEED.	Monandria Monogynia. L.						
marina. w.	(gr.) common.	Britain.	8. 9. H.w. 4.	Eng. bot. v. 7. t. 467.			
LEMNA. L.	DUCK-WEED.	Monœcia Diandria.						
1 trisulca. E.B.	(wh.) Ivy-leaved.	Britain.	5. 6. H.w. ⊖.	Eng. bot. v. 13. t. 926.			
2 minor. E.B.	(wh.) lesser.	———	6. 7. H.w. ⊖.	———	v. 16. t. 1095.		
3 gibba. E.B.	(wh.) thick-leaved.	———	———	H.w. ⊖.	———	v. 18. t. 1233.	
4 polyrhiza. E.B.	(wh.) greater.	———	7. 8. H.w. ⊖.	———	v. 35. t. 2458.		

ORDO CXCVIII.

PANDANEÆ. *Brown prodr. 340.*

PAND'ANUS. L.	SCREW-PINE.	Dioecia Monandria. L.						
1 l'avis. Lour.	(wh.) entire-leaved.	China.	1823.	S. h.			
2 odoratissimus. w.	(wh.) green-spined.	———	1771.	S. h.	Roxb. corom.1. t. 94--6.		
3 útilis. W.en.	(wh.) red-spined.	———	S. h.	Jacq. fragm. t. 14. f. 2.		
4 hūmilis. w.	(wh.) dwarf.	Mauritius.	S. h.			
5 latifòlius. L.C.	(wh.) broad-leaved.	E. Indies.	1820.	S. h.			
6 f'ostidus. H.B.	(wh.) strong scented.	———	1822.	S. h.			
7 marginatus. H.B.	(wh.) red-margined.	Mauritius.	1823.	S. h.			
8 furcatus. H.B.	(wh.) forked.	E. Indies.	———	S. h.	Rheed. mal. 2. t. 8.		
9 pedunculatus. B.P.	(wh.) peduncled.	N. Holland.	1825.	S. h.			
10 spiralis. B.P.	(wh.) spiral.	———	1805.	S. h.			
11 candelabrum. P.B.	(wh.) chandelier.	Guinea.	1822.	S. h.			
12 sessilis. Boj.	(wh.) sessile-fruited.	Pembu.	1825.	S. h.	Beauv. fl. owar. 1. t. 21.		
13 elegans. s.s.	(wh.) elegant.	Isle France.	1826.	S. h.			
14 muriatus. s.s.	(wh.) oblong-leaved.	Madagascar.	———	S. h.			
15 edulis. s.s.	(wh.) eatable.	———	1825.	S. h.			
16 amarylliñolius. H.B.	(wh.) Amaryllis-leav'd.	E. Indies.	1820.	S. h.			
17 inermis. H.B.	(wh.) spineless.	———	———	S. h.			
18 álbis. L.C.	(wh.) white-spined.	———	1823.	S. h.			
19 reflexus. L.C.	(wh.) reflexed-leaved.	———	1820.	S. h.			
20 longifolius. L.C.	(wh.) long-leaved.	———	1823.	S. h.			
21 turbinatus. L.C.	(wh.) top-shaped.	———	———	S. h.			
PHYTE'LEPHAS. R.P. PHYTE'LEPHAS.		Dioecia Polyandria. P.S.						
macrocárpa. R.P.	(wh.) large-fruited.	Peru.	1822.	S. h.			

ORDO CXCIX.

TYPHINÆ. *Kth. synops.* 1. p. 132.

TYPHA. L.	CAT'S-TAIL.	Monœcia Triandria. L.
1 latifolia. E.B.	(br.) great.	Britain. 7. H.w. ȝ. Eng. bot. v. 21. t. 1455.
2média. DC.	(br.) dwarf.	England. — H.w. ȝ. — v. 21. t. 1457.
minor. E.B.		
3 angustifolia. E.B.	(br.) lesser.	Britain. — H.w. ȝ. Eng. bot. v. 21. t. 1456.
4 minima. W.	(br.) least.	Europe. 1819. — H.w. ȝ.
5 dænatica. E.	(br.) Egyptian.	Egypt. 1825. — H.w. ȝ.
SPARG'ANIUM. L.	BUR-REED.	Monœcia Triandria. L.
1 ramosum. E.B.	(wh.) branched.	Britain. 7. 8. H.w. ȝ. Eng. bot. v. 11. t. 744.
2 simplex. E.B.	(wh.) unbranched.	— — H.w. ȝ. — v. 11. t. 745.
3 americanum. N.	(wh.) American.	N. America. 1823. — H.w. ȝ.
4 natans. E.B.	(wh.) floating.	England. 7. H.w. ȝ. Eng. bot. v. 4. t. 273.

ORDO CC.

AROIDEÆ. *Juss. gen.* 23.

Sect. I. ORONTIACEÆ. Brown prodr. 1. p. 337.

P'OTHOS. L.	P'OTHOS.	Tetrandria Monogynia. L.
1 scándens. R.S.	(gr.) climbing.	E. Indies. 1818. 4. 6. S. ȝ. Rheed. mal. 7. t. 40.
2 officinalis. F.I.	(st.) officinal.	— 1820. 4. 7. S. ȝ. —
3 Peépla. F.I.	(st.) Peeplee.	— — — S. ȝ. —
4 gigántea. F.I.	(pu.ye.) gigantic.	— 1824. S. ȝ. —
5 pertusa. F.I.	(gr.) perforated.	— — — S. ȝ. — Rheed. mal. 12. t. 20. 21.
6 lanceolata. R.S.	(vi.) lance-leaved.	Barbadoes. 1790. 4. 7. S. ȝ. Plum. amer. 47. t. 62.
7 acaulis. R.S.	(gr.) stemless.	W. Indies. — — — S. ȝ. Lodd. bot. cab. t. 483.
8 gráclilis. R.S.	(gr.) slender.	Trinidad. 1826. — — — S. ȝ. Rudg. pl. gui. 1. t. 32.
9 violacea. R.S. (g.wh.)	blue-fruited.	Jamaica. 1793. 4. 6. S. ȝ. Hook. exot. flor. t. 55.
10 cannæfolia. R.S. (wh.)	sweet-scented.	W. Indies. 1785. 4. 5. S. ȝ. Bot. mag. t. 603.
11 crassinérvia. R.S. (gr.)	thick-nerved.	S. America. 1796. 4. 6. S. ȝ. Jacq. ic. 3. t. 609.
12 cordáta. R.S.	(gr.) heart-leaved.	— 1770. 4. S. ȝ. Plum. ic. 26. t. 38.
13 sagittáta. B.M.	(g.wh.) arrow-leaved.	W. Indies. 1800. 8. S. ȝ. Bot. mag. t. 1584.
14 macrophylla. R.S. (gr.)	large-leaved.	— 1794. 5. 6. S. ȝ. Jacq. ic. 3. t. 610.
15 obtusifolia. R.S. (g.br.)	blunt-leaved.	Barbadoes. 1790. — — — S. ȝ.
16 coriácea. H.E.F. (g.br.)	leathery.	Brazil. 1824. 6. S. ȝ. Hook. ex. flor. t. 210.
17 Harrisii. H.E.F. (g.br.)	Harris's.	— — — S. ȝ. — — — t. 211.
18 caudáta. F.I.	(gr.) tailed.	E. Indies. 1824. S. ȝ. —
19 pinnátâ. W.	(gr.) wing-leaved.	— 1820. S. ȝ. — Rumph. am. 5. t. 183. f. 2.
20 décursiva. F.I.	(gr.) pinnatifid-leaved.	— 1824. S. ȝ. —
21 pinnatífida. F.I.	(gr.) pinnatifid.	— 1825. S. ȝ. —
22 heterophylla. F.I. (pu.)	various-leaved.	— 1822. S. ȝ. —
23 Lásia. F.I.	(re.) prickly.	— 1819. S. ȝ. —
24 digitáta. R.S.	(pu.) fingered.	Caracas. 1823. S. ȝ. Jacq. ic. 3. t. 611.
25 palmátâ. R.S. (g.pu.)	palmated.	S. America. 1803. 6. 7. S. ȝ. Plum. amer. t. 64. 65.
26 pentaphylla. R.S. (bl.)	five-leaved.	Cayenne. — 10. 11. S. ȝ. Bot. mag. t. 1375.
27 rubrinérvia. L.en. (pu.)	red-nerved.	S. America. 1820. S. ȝ. Bot. mag. t. 2953.
28 microphylla. B.M. (br.)	small-leaved.	Brazil. — 8. 10. S. ȝ. Bot. mag. t. 2953.
SYMPLOCA'RPU'S. N.	SCUNKWEED.	Tetrandria Monogynia. N.
fœtidus. N.	(d.pu.) fetid.	N. America. 1735. 3. 5. H. ȝ. Swt. br. fl. gar. t. 57.
Pòthos fœtida. B.M.	836. Dracónium fœtidum. w.	
DRACO'NTIUM. W.	DRAGON.	Heptandria Monogynia. L.
1 polyphyllum. w. (d.pu.)	purple-stalked.	India. 1759. 3. 6. S. ȝ. Bot. reg. t. 700.
2 spinosum. w. (d.pu.)	prickly.	Ceylon. — 4. 5. S. ȝ.
GYMNOSTACHYS. B.P.	GYMNOST'ACHYS.	Tetrandria Monogynia.
ancéps. B.P.	(gr.) flat-stemmed.	N. S. W. 1820. 6. 8. G. ȝ.
HOUTTUYNIA. J.	HOUTTU'YNIA.	Polyandria Polygynia. P.S.
cordáta. P.S.	(wh.) heart-leaved.	China. 1820. 6. 8. G. ȝ. Bot. mag. t. 2731.

CARLUDO'VICA. R.P.	CARLUDO'VICA.	Moneocia Polyandria. P.S.					
1 palmata. r.s.	(br.) palmate.	S. America.	1818.	S. $\ddot{\eta}$.		
<i>Sálmia palmata.</i> R.S.	<i>Ludòvia palmata.</i> P.S.						
2 latifolia. R.P.	(br.) broad-leaved.	Peru.	1819.	4. 6.	S. $\ddot{\eta}$.	Bot. mag. t. 2950. et 2951.	
3 angustifolia. R.P.	(br.) narrow-leaved.	S. $\ddot{\eta}$.		
4 palmæfolia. R.S. (br.)	Palm-leaved.	St. Domingo.	1826.	S. $\ddot{\eta}$.	Plum. ic. t. 39.	
<i>Sálmia palmæfolia.</i> w.							
5 jamaicensis. L.C. (br.)	Jamaica.	Jamaica.	1823.	S. $\ddot{\eta}$.		
CYCLAN'THUS. S.S.	CYCLAN'THUS.	Moneocia Monadelphia.					
1 Pluméri. s.s.	(gr.) Plumér's.	W. Indies.	1820.	S. $\ddot{\eta}$.		
2 bipartitus. s.s.	(gr.) cleft-leaved.	Trinidad.	S. $\ddot{\eta}$.		
A'CORUS. L.	SWEET-FLAG.	Hexandria Monogynia. L.					
1 Cálamus. s.s.	(gr.) common.	Britain.	6. 7.	H.w. $\ddot{\eta}$.	Eng. bot. v. 5. t. 356.	
2 terréstris. s.s.	(gr.) Chinese.	China.	1822.	—	H.w. $\ddot{\eta}$.		
3 gramineus. s.s.	(gr.) grass-leaved.	1786.	2.	H.w. $\ddot{\eta}$.	Smith spic. 15. t. 17.	
ORON'TIUM. L.	ORON'TIUM.	Hexandria Monogynia. L.					
aquaticum. s.s.	(st.) aquatic.	N. America.	1775.	6.	H.w. $\ddot{\eta}$.	Hook. ex. flor. t. 19.	
RO'HDEA. S.S.	RO'HDEA.	Hexandria Monogynia. S.S.					
japónica. s.s.	(ye.) Japan.	Japan.	1783.	1. 4.	H. $\ddot{\eta}$.	Bot. mag. t. 898.	
<i>Orón'tium japónicum.</i> B.M.							
TUPI'STRA. B.M.	TUPI'STRA.	Hexandria Monogynia. S.S.					
1 squálida. B.M.	(br.) brown-flowered.	E. Indies.	1810.	4. 6.	S. $\ddot{\eta}$.	Bot. reg. t. 704.	
2 nútans. B.R.	(br.) nodding-flowered.	1818.	11. 12.	S. $\ddot{\eta}$.	— t. 1223.	
ASPIDI'STRA. B.R.	ASPIDI'STRA.	Octandria Monogynia. S.S.					
lúrida. B.R.	(br. pu.) brown-flowered.	China.	1820.	2. 4.	S. $\ddot{\eta}$.	Bot. reg. t. 628.	
<i>punctata.</i> B.R. 977.							

Sect. II. AROIDEÆ VERÆ. Brown prodr. 335.

CA'LLA. K.S.	CA'LLA.	Heptandria Monogynia. L.					
1 palústris. w.	(wh.) marsh.	N. Europe.	1768.	7. 8.	H.w. $\ddot{\eta}$.	Bot. mag. t. 1831.	
2 pertusa. k.s.	(st.) perforated.	W. Indies.	1752.	4. 6.	S. $\ddot{\eta}$.	Jacq. schœn. 2. t. 184. 5.	
<i>Dracónium pertusum.</i> Jacq.							
3 aromática. B.M. (wh.) aromatic.	E. Indies.	1817.	—	S. $\ddot{\eta}$.	Bot. mag. t. 2279.		
4 occulta. B.C. (wh.) hidden-spiked.	China.	—	5. 7.	G. $\ddot{\eta}$.	Lodd. bot. cab. t. 12.		
RICHA'RDIÁ. Kth.	RICHA'RDIÁ.	Heptandria Monogynia.					
ethiopica. Kth. (wh.) Ethiopian.	C. B. S.	1731.	1. 5.	G. $\ddot{\eta}$.	Bot. mag. t. 832.		
<i>Cália ethiopica.</i> B.M.							
'ARUM. W.	'ARUM.	Moneocia Polyandria. L.					
1 crinitum. w. (d.pu.) hairy-sheathed.	Minorca.	1777.	3. 4.	F. $\ddot{\eta}$.	Bot. reg. t. 831.		
2 Dracúnculus. w. (pu.) common Dragon.	S. Europe.	1548.	6. 7.	H. $\ddot{\eta}$.	Moris. s. 13. t. 5. f. 46.		
3 Dracónium. w. (gr.) green Dragon.	N. America.	1759.	6.	H. $\ddot{\eta}$.	Bot. reg. t. 668.		
4 bulbiferum. B.M. (st. br.) bulb-bearing.	E. Indies.	1813.	3. 6.	S. $\ddot{\eta}$.	Bot. mag. t. 2508.		
5 venosum. w. (ye. pu.) purple-flowered.	1774.	3.	S. $\ddot{\eta}$.	Bot. reg. t. 1017.		
6 campanulatum. H.B. (pu.) bell-flowered.	E. Indies.	1816.	4. 6.	S. $\ddot{\eta}$.	Lodd. bot. cab. t. 820.		
7 triphyllum. w. (std.) three-leaved.	N. America.	1664.	5. 6.	H. $\ddot{\eta}$.	Bot. mag. t. 950.		
β Zebrinum. B.M. (std.) striped.	—	—	—	H. $\ddot{\eta}$.	Pl. sel. hort. ber. t. 8.		
8 pedánum. W. en. (bl. ye.) pedate-leaved.	S. America.	1815.	—	S. $\ddot{\eta}$.			
9 polyphyllum. L.en. (pu.) many-leaved.	1818.	—	S. $\ddot{\eta}$.			
10 ríngens. w. (pu.) gaping.	Japan.	1822.	—	G. $\ddot{\eta}$.			
11 atrorübens. w. (d.re.) purple-stalked.	N. America.	1758.	6. 7.	H. $\ddot{\eta}$.	Pluk. alm. t. 77. f. 5.		
12 ternátum. w. (pu.)	Japan.	1774.	5. 7.	F. $\ddot{\eta}$.	Rumph. amb. 5. t. 109.		
13 Colocásia. w. (br.) Egyptian.	Levant.	1551.	6. 8.	G. $\ddot{\eta}$.	Bot. reg. t. 450.		
14 orixéns. B.P. (re.) Orixian.	E. Indies.	1802.	8. 10.	S. $\ddot{\eta}$.	Herm. parad. t. 73.		
15 macrorhizon. w. (wh.) long-rooted.	—	1803.	—	S. $\ddot{\eta}$.	Rheed. mal. 11. t. 20.		
16 píctum. w. (d.pu.) white-veined.	S. Europe.	1820.	5. 6.	H. $\ddot{\eta}$.	Bot. mag. t. 339.		
17 divaricátum. w. (gr.) divaricated.	E. Indies.	1759.	6. 7.	S. $\ddot{\eta}$.	— t. 2324.		
18 trilobátum. w. (pu.) three-lobed.	Ceylon.	1714.	5. 6.	S. $\ddot{\eta}$.	Eng. bot. v. 19. t. 1298.		
β auriculátum. B.M. (pu.) eared.	—	—	—	S. $\ddot{\eta}$.			
19 maculátum. w. (gr.) common.	Britain.	—	5. 7.	H. $\ddot{\eta}$.	Bot. mag. t. 2432.		
20 orientále. M.B.	(br.) oriental.	Caucasus.	1818.	—	H. $\ddot{\eta}$.	Lodd. bot. cab. t. 281.	
21 itálicum. w.	(gr.) Italian.	Italy.	1683.	5. 6.	H. $\ddot{\eta}$.	Rheed. mal. 11. t. 17.	
22 viviparum. L.C.	(gr.) viviparous.	—	1817.	—	S. $\ddot{\eta}$.	Bocc. mus. 2. t. 50.	
23 minútum. w.	(re.) small.	E. Indies.	1812.	5. 7.	S. $\ddot{\eta}$.	Jacq. schœn. 2. t. 192.	
24 proboscídeum. w. (w.pu.) proboscis.	Apennines.	1825.	—	H. $\ddot{\eta}$.	Bot. reg. t. 512.		
25 Arisárum. w.	(gr.) Friar's Cowl.	S. Europe.	1596.	4. 6.	F. $\ddot{\eta}$.		
26 tenuifolium. w. (d.pu.) grass-leaved.	—	1570.	—	H. $\ddot{\eta}$.			

spirale. w. (pu.) spiral.	E. Indies.	1816.	4. 6. S. 4.	Bot. mag. t. 2220.
28 gramineum. s.s. (pu.) grass-leaved.	S. Europe.	1824.	— H. 4.	Lodd. bot. cab. t. 396.
29 flagelliforme. b.c. (pu.) whip-cord.	E. Indies.	—	— S. 4.	Jacq. amer. t. 152.
30 sagittifolium. L.en. arrow-leaved.	W. Indies.	1793.	5. 6. S. h.	Plum. ic. 26. t. 37.
31 hederaceum. w. (pu.) Ivy-leaved.	—	1816.	— S. h.	
32 obtusilobum. L.en. (pu.) blunt-lobed.	China.	—	— S. h.	
33 indicum. Lour. (pu.) Indian.	—	—	— S. h.	
34 ramosum. L.en. (pu.) branching.	—	—	— S. h.	
35 lingulatum. w. (ye.) tongue-leaved.	W. Indies.	1793.	— S. h.	
36 integrifolium. L.en. (gr.) entire-leaved.	—	1820.	— S. h.	
CAL'ADIUM. V.	CAL'ADIUM.	Monoezia Polyandria.	W.	
1 helleborifolium. w. (wh.) Hellebore-leaved.	Caracas.	1796.	6. 7. S. 4.	Jacq. ic. 3. t. 613.
2 pinnatifidum. w. (pu.) pinnatifid.	—	1817.	— S. 4.	Jacq. schœn. 2. t. 187.
3 bicolor. w. (wh.) two-coloured.	Brazil.	1773.	— S. 4.	Bot. mag. t. 2543.
4 nymphaeifolium. w. (sn.) Water-lily-leaved.	E. Indies.	1800.	— S. 4.	Rheed. mal. 11. t. 22.
5 esculentum. w. (g.pu.) esculent.	America.	1739.	6. 8. S. 4.	Sloan. jam. 1. t. 106. f. 1.
6 odoratum. B.R. (gr.) odorous.	E. Indies.	1818.	5. 7. S. 4.	Bot. reg. t. 641.
7 viviparum. H.P. (wh.) viviparous.	—	1816.	— S. 4.	Lodd. bot. cab. t. 281.
8 sagittifolium. w. (wh.) arrow-leaved.	W. Indies.	1710.	— S. 4.	Jacq. vind. 2. t. 157.
9 virginicum. H.E.F. (st.) Virginian.	N. America.	1759.	6. 7. H. 4.	Hook. ex. flor. t. 182.
'Arum virginicum. w.				
10 seguinum. w. (gr.) Dumb Cane.	America.	—	5. S. h.	Hook. ex. flor. t. 1.
11 maculatum. B.C. (gr.) spotted.	Trinidad.	1820.	4. 12. S. h.	Lodd. bot. cab. t. 608.
seguinum β maculatum.	Bot. mag. t. 2606.			
12 scandens. s.s. (wh.) climbing.	Guinea.	1822.	— S. h.	Beauv. fl. owar. t. 3.
13 eucullatum. p.s. (gr.) hooded.	China.	1816.	— S. h.	
14 xanthorrhizum. w. (st.) yellow-rooted.	—	—	4. 8. S. h.	Jacq. schœn. 2. t. 188.
15 grandifolium. w. (wh.) large-leaved.	Caracas.	1803.	3. 7. S. h.	Bot. mag. t. 2643.
16 arborescens. w. (wh.) tree.	W. Indies.	1759.	6. 7. S. h.	Plum. amer. 44. t. 60.
17 lacерum. w. (gr.) torn-leaved.	Caracas.	1822.	— S. h.	
18 tripartitum. w. (wh.) ternate-leaved.	—	1816.	— S. h.	Jacq. schœn. 2. t. 190.
19 auritum. (wh.) ear-leaved.	S. America.	1739.	6. 8. S. h.	— 2. t. 191.
20 pedatum. H.E.F. (ye.) pedate-leaved.	Brazil.	1824.	— S. h.	Hook. ex. flor. t. 206.
21 aculeatum. s.s. (wh.) prickly.	Surinam.	1822.	— S. h.	
AMBROS'INIA. W.	AMBROS'INIA.	Monoezia Monandria.	W.	
1 Bassii. w. (gr.pu.) flat oval-leaved.	S. Europe.	1823.	6. 8. H. 4.	Lam. ill. t. 737.
2 ciliaris. s.s. (gr.pu.) fringed-spathed.	E. Indies.	1824.	— S. 4.	

Sect. III. TACCACEÆ.

TA'CCA. W.	TA'CCA.	Hexandria Monogynia. L.		
1 pinnatifida. w. (pu.) Salep.	E. Indies.	1793.	6. 8. S. 4.	Lodd. bot. cab. t. 692.
2 áspera. H.B. (pu.) rough-stemmed.	—	1820.	— S. 4.	
3 lævis. H.B. (pu.) smooth.	—	—	— S. 4.	
4 integrifolia. B.M. (pu.) entire-leaved.	—	1810.	5. 7. S. 4.	Bot. mag. t. 1488.

ORDO CCI.

PIPERACEÆ. Kth. synops. 1. p. 103.

SAUR'URUS. W.	LIZARD'S-TAIL.	Heptandria Tetragynia. L.		
1 cérmuns. w. (wh.) drooping.	Virginia.	1759.	9. H.w. 4.	Lam. ill. t. 1.
2 lúcidus. s.s. (wh.) shining.	N. America.	1791.	8. 10. H.w. 4.	Jacq. ecl. t. 18.
3 chinénsis. Hort. (wh.) Chinese.	China.	1819.	— H.w. 4.	
PIPER. R.P.	PEPPER.	Diandria Trigynia. L.		
1 coriáceum. r.s. (gr.) coriaceous.	E. Indies.	1815.	7. 8. S. h.	Lodd. bot. cab. t. 128.
2 nitidum. r.s. (wh.) glossy.	Jamaica.	1793.	5. 6. S. h.	
3 peltatum. r.s. (gr.) peltated.	W. Indies.	1748.	— S. 4.	Plum. amer. t. 74.
4 umbellatum. r.s. (gr.) umbellated.	—	—	5. 7. S. h.	Flor. peruv. 1. t. 59. f. 1.
5 macrophyllum. r.s. (gr.) broad-leaved.	—	1800.	— S. h.	Sloan. hist. 1. t. 88. f. 1.
6 geniculatum. r.s. (gr.) jointed.	—	1826.	— S. h.	
7 aduncum. r.s. (gr.) hooked.	Jamaica.	1748.	5. S. h.	Jacq. ic. 2. t. 210.

PIPERACEÆ.

8 hirsutum. R.S. (br.) hairy-leaved.	Jamaica.	1793.	7.	S. h.
9 hispidum. R.S. (gr.) hispid.	S. America.	—	—	S. h.
10 tuberculatum. R.S. (gr.) tuberculate.	—	1816.	—	S. h.
11 ovatum. R.S. (gr.) oval-leaved.	Trinidad.	1824.	—	S. h.
12 Amalago. R.S. (gr.) rough-leaved.	Jamaica.	1759.	—	S. h.
13 Bétle. R.S. (gr.) Bétle.	E. Indies.	1804.	—	S. h.
14 nigrum. R.S. (gr.) black.	—	1790.	—	S. h.
15 triocicum. F.I. (gr.) glaucous-leaved.	—	1818.	—	S. h.
16 reticulatum. R.S. (gr.) netted.	W. Indies.	1748.	8.	S. h.
17 decumanum. R.S. (gr.) great.	Carthagena.	1768.	—	S. h.
18 Siriboa. R.S. (gr.) Siriboa.	E. Indies.	—	—	S. h.
19 longum. R.S. (gr.) long.	—	1788.	—	S. h.
20 glabrum. H.S.S. (gr.) smooth.	Campeachy.	1768.	—	S. h.
21 laurifolium. H.S.S. (gr.) Laurel-leaved.	Jamaica.	—	—	S. h.
22 racemosum. H.S.S. (gr.) racemed.	Campeachy.	—	—	S. h.
<i>Staphyle.</i> R.S.				
23 acutifolium. R.S. (gr.) acute-leaved.	Peru.	1823.	—	S. h.
24 columbinum. L.en. (gr.) spotted-stalked.	Brazil.	1820.	—	S. h.
25 plantagineum. R.S. (gr.) Plantain-leaved.	W. Indies.	—	—	S. h.
<i>mèdium.</i> Jacq.				
26 marginatum. L.en. (gr.) purple-edged.	S. America.	1811.	—	S. h.
27 unguiculatum. R.S. (gr.) claw-pointed.	Peru.	1822.	—	S. h.
<i>glaucescens.</i> Jacq. ecl. 1. t. 76.				
28 humile. Mill. (gr.) humble.	Jamaica.	1768.	—	S. h.
29 tomentosum. H.S.S. (gr.) oval-leaved downy.	VeraCruz.	—	—	S. h.
30 sidaefolium. L.en. (gr.) Sida-leaved.	S. America.	1821.	5. 7.	S. h.
<i>PEPEROMIA.</i> R.P. <i>PEPEROMIA.</i>				
1 brachiphylla. (gr.) short-leaved.	Diandria Trigynia.	—	—	S. h.
<i>Piper brachiphyllum.</i> R.S.				
2 amplexicaulis. R.S. (gr.) stem-clasping.	W. Indies.	1793.	6. 9.	S. ¼.
3 magnoliaefolia. R.S. (gr.) Magnolia-leaved.	—	—	1. 3.	S. ¼.
4 obtusifolia. R.S. (gr.) blunt-leaved.	S. America.	1739.	4. 7.	S. ¼.
5 clusiæfolia. B.M. (gr.) Clusia-leaved.	—	1739.	4. 7.	S. h.
6 cuneifolia. R.S. (gr.) wedge-leaved.	Caracas.	1809.	—	S. ¼.
7 alata. R.P. (gr.) winged.	S. America.	1812.	—	S. h.
8 acuminata. R.S. (gr.) acuminate.	W. Indies.	—	4. 9.	S. ¼.
9 maculosa. R.S. (gr.) spot-stalked.	St. Domingo.	1790.	9.	S. h.
10 pellucida. K.S. (gr.) pellucid.	S. America.	1748.	4. 9.	S. ⊙.
11 concinna. K.R. (gr.) small heart-leaved.	Jamaica.	1821.	—	S. h.
12 rotundifolia. K.S. (gr.) round-leaved.	S. America.	1823.	—	S. ¼.
13 hispidula. R.S. (gr.) annual hispid.	Jamaica.	1818.	8. 10.	S. ⊙.
14 nummularifolia. K.S. (gr.) money-wort-leaved.	—	—	4. 9.	S. ¼.
15 pubescens. R.S. (gr.) pubescent.	—	—	—	S. ¼.
16 subrotunda. H.S.S. (gr.) small Clusia-leav'd.	—	1811.	—	S. ¼.
17 repens. K.S. (gr.) creeping.	S. America.	1823.	—	S. ¼.
18 serpens. R.S. (gr.) trailing.	Jamaica.	—	—	S. ¼.
19 incana. H.E.F. (gr.) great downy.	Brazil.	1814.	2. 6.	S. h.
20 amplexifolia. L.en. (gr.) clasping-leaved.	Jamaica.	1822.	—	S. h.
21 talinifolia. K.S. (gr.) Talinum-leaved.	S. America.	—	4. 8.	S. h.
22 pulchella. R.S. (gr.) small-leaved.	Jamaica.	1778.	7. 10.	S. ¼.
23 pereskiaefolia. K.S. (gr.) Pereskia-leaved.	S. America.	1820.	5. 6.	S. ¼.
24 blanda. K.S. (gr.) villous.	Caracas.	1802.	5. 11.	S. ¼.
25 verticillata. R.S. (gr.) whorl-leaved.	Jamaica.	1816.	6. 10.	S. ⊙.
26 polystachia. H.E.F. (gr.) many-spiked.	W. Indies.	1775.	6. 7.	S. ¼.
27 quadrifolia. H.E.F. (gr.) four-leaved.	Santa Cruz.	1818.	5. 10.	S. ¼.
28 Haworthii. (gr.) Haworth's.	India.	1809.	9. 10.	S. ¼.
<i>Piper pubescens.</i> H.S. non Vahl.				
29 inaequifolia. R.P. (gr.) unequal-leaved.	Peru.	1800.	8. 10.	S. ¼.
30 stellata. R.S. (gr.) starry.	Jamaica.	1802.	5. 7.	S. ¼.
31 rubricaulis. L.en. (gr.) red-stalked.	W. Indies.	1815.	—	S. ¼.
32 rubella. H.E.F. (gr.) red-leaved.	Jamaica.	1820.	—	S. ¼.
33 trifolia. R.S. (gr.) three-leaved.	S. America.	1802.	6. 8.	S. ¼.
34 distachya. R.S. (gr.) two-rowed.	W. Indies.	1793.	—	S. ¼.
35 discolor. R.S. (gr.) two-coloured.	Jamaica.	1821.	7. 8.	S. h.
36 capensis. R.S. (gr.) Cape.	C. B. S.	1816.	—	G. h.
37 tricarinata. H.S.S. (gr.) triple-keeled.	—	1817.	—	S. h.

Jacq. ic. 2. t. 211.

Sloan. hist. 1. t. 87. f. 1.
Rheed. mal. 7. t. 15.
— 7. t. 12.

Plum. amer. t. 75.

Jacq. ic. 2. t. 215.
Rumph. am. 5. t. 117. f. 2.
Rheed. mal. 7. t. 14.

Flor. per. 1. t. 64. f.a.

Jacq. ic. 1. t. 8.

Jacq. ic. 2. t. 215.

Flor. per. 1. t. 57. f. b.

Lk. et Ot. ic. sel. 1. t. 6.

Jacq. ic. 2. t. 213.

Trew ehret. 54. t. 96.

Bot. mag. t. 2943.

Jacq. ic. 2. t. 214.

Flor. per. 1. t. 48. f. b.

Bot. mag. t. 1882.

Hook. ex. flor. t. 92.

Plum. amer. t. 72.

Plum. amer. t. 69.

Swartz icon. t. 4.

Hook. ex. flor. t. 66.

H. et B. n. gen. 1. t. 8.

Lodd. bot. cab. t. 574.

Hook. ex. flor. t. 67.

— t. 21.

Hook. ex. flor. t. 23.

— t. 22.

Flor. peruv. 1. t. 46. f. a.

Jacq. vind. 2. t. 217.

Nees. hort. ber. t. 8.

Hook. ex. flor. t. 58.

Plum. amer. t. 68.

— t. 67.

Lodd. bot. cab. t. 610.

ORDO CCII.

ORCHIDÆ. Juss. gen. 64.

O'RCHIS. B.P.

O'RCHIS. B.P.	O'RCHIS.	Gynandria Monandria. H.K.
1 Mòrio. w.	(pu.) meadow.	Britain. 4. 6. H.b. ¼. Eng. bot. v. 29. t. 2059.
2 longicórnus. w.	(pu.) flat-spurred.	Barbary. 1815. — H.b. ¼. Swt. br. fl. gar. t. 249.
3 pállens. w.	(st.) pale.	Switzerland. 1825. 5. 6. H.b. ¼. Jacq. aust. t. 45.
4 máscula. w.	(pu.) early purple.	Britain. 4. 5. H.b. ¼. Eng. bot. v. 9. t. 631.
5 palústris. w.	(li.) marsh.	Europe. 1824. — H.b. ¼. Jacq. ic. t. 181.
6 pauciflòra. T.N.	(ye.) few-flowered.	Italy. 1825. 6. 7. H.b. ¼. Ten. fl. nap. t. 88.
7 Nicodémi. T.N.	(pu.) Nicodemus's.	— — — — —
8 Cyrilli. T.N.	(ye.) Cyrillo's.	Naples. — — — — —
9 parviflòra. T.N.	(pu.) small-flowered.	Italy. — — — — —
10 ustuláta. w.	(pu.) dwarf.	England. 5. 6. H.b. ¼. Eng. bot. v. 1. t. 18.
11 militaris. w.	(pu.) military.	Switzerland. 1825. 4. 6. H.b. ¼. Swt. br. fl. gar. t. 163.
12 Rivinì. B.F.G.	(li.pu.) Rivinus's.	— — — — —
13 Smithii. B.F.G.	(ro.pu.) Smith's.	England. 5. 6. H.b. ¼. — — — — — t. 162.
	militaris. E.B. non w.	— — — — —
14 fúscia. w.	(br.re.) brown.	— — — — —
15 Simia. P.S.	(pa.bb.) Ape.	— — — — —
16 undulatífolia. B.B.	(li.) wave-leaved.	Italy. 1818. — H.b. ¼. Bot. reg. t. 375.
17 tephrosánthos. w.	(pa.) ash-coloured.	England. — H.b. ¼. Vaill. bot. t. 31.f.25-26.
18 acumináta. w. (li.wh.)	pointed-flower'd.	S. Europe. 1815. 4. 6. H.b. ¼. Bot. mag. t. 1932.
19 variegátâ. w.	(li.wh.) variegated.	Switzerland. 1825. — H.b. ¼. Hall. helv. t. 30.
20 globosa. w.	(pu.) round-spiked.	— 1792. 6. 7. H.b. ¼. Jacq. aust. 3. t. 265.
21 coriophóra. w.	(pu.) strong-scented.	S. Europe. 1825. 5. 6. H.b. ¼. Swt. br. fl. gar. t. 219.
22 hircina. w.	(gr.pu.) Lizard.	England. 6. 7. H.b. ¼. Eng. bot. v. 1. t. 34.
	Satyr'ium hircinum. E.B.	— — — — —
23 Robertiana. P.S.	(pu.) long-bracted.	S. Europe. 1818. 5. 6. H.b. ¼. Ten. fl. nap. t. 91.
24 longibracteáta. s.s.(pu.)	long-bracted.	— — — — —
25 quadripunctáta. T.N.	(pu.) four-spotted.	Italy. 1828. — H.b. ¼. Bot. reg. t. 357.
26 saccáta. T.N.	(pu.) saccate.	Sicily. — — — — —
27 papilionácea. w.	(pu.) papilionaceous.	S. Europe. 1788. 6. 7. H.b. ¼. Bot. mag. t. 1155.
28 pseudosambucina. T.N.	(pu.) Elder-smelling.	Italy. 1828. — H.b. ¼. Ten. flor. nap. t. 86.
	Ivy-leaved. — — — — —	— H.b. ¼. — — — — — t. 86. f. b.
29 sulphurea. B.M.	(st.) pale-yellow.	S. Europe. 1823. 5. 6. H.b. ¼. Bot. mag. t. 2569.
30 sambucina. w.	(ye.) Elder-scented.	Europe. 1825. 4. 5. H.b. ¼. Swt. br. fl. gar. t. 299.
31 Schleichéri. B.F.G.	(re.) Schleicher's.	Switzerland. — — — — —
	sambucina β rúbra. Sch. non Jacq.	— — — — — t. 199.
32 latifolia. w.	(pu.) broad-leaved.	Britain. 5. 6. H.b. ¼. Eng. bot. v. 33. t. 2308.
33 maculáta. w.	(li.pu.) spotted-palmette.	— — — — —
34 spectabilis. w. (wh.pu.)	showy American.	N. America. 1801. — H.b. ¼. Swt. br. fl. gar. t. 65.

ANACAMPTIS. R.O. ANACAMPTIS.

pyramidalis. r.o. (cr.) pyramidal. Britain. 6. 7. H.b. ¼. Eng. bot. v. 2. t. 110.

O'rchis pyramidalis. E.B.

GYMNADENIA. H.K. GYMNADENIA.

Gynandria Monandria. H.K.

1 conóspaea. H.K. (ro.) fragrant. Britain. 6. 7. H.b. ¼. Eng. bot. v. 1. t. 10.

O'rchis conóspaea. E.B.

2 odoratissima. R.O. (li.) sweet-scented. Switzerland. 1824. — H.b. ¼. Jacq. aust. 3. t. 264.

3 viridis. R.O. (gr.) Frog. Britain. — H.b. ¼. Eng. bot. v. 2. t. 94.

Satyr'ium viride. E.B. O'rchis viridis. H.K.

4 álbida. R.O. (wh.) small white. — — — — — H.b. ¼. Eng. bot. v. 8. t. 505.

A'CERAS. H.K. A'CERAS. Gynandria Monandria. H.K.

anthropophóra. H.K. (gr.) green-man. England. 6. H.b. ¼. Swt. br. fl. gar. t. 168.

HERMI'NIUM. H.K. HERMI'NIUM. Gynandria Monandria. H.K.

Monórchis. H.K. (st.) musk. England. 6. 7. H.b. ¼. Eng. bot. v. 1. t. 71.

Ophrys Monórchis. E.B.

CHAMO'RCHIS. R.O. CHAMO'RCHIS. Gynandria Monandria.

alpina. R.O. (wh.) Alpine. Switzerland. 1825. 6. 7. H.b. ¼. Jacq. vind. 1. t. 9.

Chamærépes alpina. s.s. Ophrys alpina. Jacq.

NIGRITELLA. R.O. NIGRITELLA. Gynandria Monandria.

angustifolia. R.O. (d.pu.) dark-flowered. N. Europe. 1759. 6. 7. H.b. ¼. Flor. dan. t. 998.

Satyr'ium nigrum. L. O'rchis nigra. Sw. Habenaria nigra. H.K.

PLATANTH'ERA. R.O. PLATANTH'ERA. Gynandria Monandria.

bifolia. R.O. (wh.) Butterfly. Britain. 5. 6. H.b. ¼. Eng. bot. v. 1. t. 22.

O'rchis bifolia. E.B. Habenaria bifolia. H.K.

ORCHIDÆ.

DISPERIS. W.	DISPERIS.	Gynandria Monandria. H.K.		
1 capensis. w. (cr.)	Cape.	C. B. S.	1816.	5. 6. F.b. 4.
2 cucullata. w. (li.)	cucullate.	—	1822.	6. 7. F.b. 4.
3 secunda. w. (st.)	side-flowering.	—	1797.	— F.b. 4.
CORYCIUM. W.	CORYCIUM.	Gynandria Monandria.		
1 orobanchoides. w. (ye.)	Orobanche-like.	C. B. S.	1825.	6. 8. F.b. 4.
2 crispum. w. (ye.)	curled-leaved.	—	—	5. 6. F.b. 4.
GLO'SSULA. B.R.	GLO'SSULA.	Gynandria Monandria.		Buxb. cent. 3. t. 11.
tentaculata. B.R. (gr.)	horned.	China.	1824.	5. 6. F.b. 4.
GOODYERA. H.K.	GOODY'ERA.	Gynandria Monandria. H.K.		Bot. reg. t. 862.
1 repens. H.K. (wh.)	creeping.	Scotland.	7. 8. H. 4.
<i>Satyr'ium repens.</i> E.B. <i>Neottia repens.</i> w.				Eng. bot. v. 5. t. 289.
2 tessellata. B.C. (wh.)	tessellated.	N. America.	1823.	6. 7. H. 4.
<i>Neottia repens.</i> Ph. <i>Goodyera pubescens.</i> β minor. B.M. 2540.				Lodd. bot. cab. t. 952.
3 pubescens. H.K. (wh.)	downy.	N. America.	1802.	— H. 4.
4 procera. H.E.F. (wh.)	Nepaul.	Nepaul.	1821.	7. 12. G. 4.
<i>Neottia procera.</i> B.R. 639.				Hook. ex. flor. t. 39.
5 discolor. B.R. (wh.)	red-leaved.	S. America.	1815.	10. 12. S. 4.
PELE'XIA. B.R.	PELE'XIA.	Gynandria Monandria. B.R.		Bot. reg. t. 271.
spiranthoides. B.R. (gr.)	decurrent-lipped.	W. Indies.	1823.	3. S. 4.
<i>Neottia adnata.</i> Sw.				Bot. reg. t. 985.
STENORHY'NCHUS. R.O. STENORHY'NCHUS.		Gynandria Monandria.		
1 speciosus. R.O. (sc.)	showy.	W. Indies.	1790.	3. 6. S. 4.
<i>Neottia speciosa.</i> Hook. ex. flor. t. 39.				Bot. mag. t. 1374.
2 orchioides. R.O. (pu.)	frosted-flowered.	Jamaica.	1806.	— S. 4.
3 plantagineus. H. (pu.)	Plantain-leaved.	W. Indies.	1825.	— S. 4.
<i>Neottia plantaginea.</i> Hooker.				Hook. ex. flor. t. 226.
4 aphylla. (br.li.)	leafless.	Trinidad.	—	— S. 4.
<i>Neottia aphylla.</i> B.M.				Bot. mag. t. 2797.
NEOT'TIA. H.K.	NEOT'TIA.	Gynandria Monandria. H.K.		
1 macrantha. (wh.)	great-flowered.	W. Indies.	1827.	3. 6. S. 4.
<i>Neottia grandiflora.</i> B.M. t. 2956. non t. 2730.				Bot. mag. t. 2956.
2 grandiflora. B.M. (gr.)	large-flowered.	Brazil.	1825.	2. 5. S. 4.
<i>Spiranthes grandiflora.</i> Bot. reg. t. 1043.				Bot. mag. t. 2730.
3 bicolor. B.R. (wh.)	two-coloured.	W. Indies.	1821.	7. 8. S. 4.
4 elata. w. (wh.)	tall.	—	1790.	4. 6. S. 4.
5 picta. H.K. (wh.)	painted-leaved.	Trinidad.	1805.	2. 4. S. 4.
6 cernua. H.K. (wh.)	nodding-flowered.	N. America.	1796.	8. 10. H. 4.
7 tortilis. w. (wh.)	twisting-spiked.	W. Indies.	1822.	6. 7. S. 4.
8 aestivalis. M. (wh.)	summer.	N. America.	—	— H. 4.
<i>tortilis.</i> Ph. non Sw.				
9 spiralis. H.K. (wh.)	Ladies-traces.	Britain.	8. 9. H.b. 4.
<i>Ophrys spiralis.</i> E.B.				Eng. bot. v. 8. t. 541.
10 pudica. (bh.)	modest.	China.	1819.	9. 12. H.b. 4.
<i>australis.</i> β . B.R. t. 602. <i>Spiranthes pudica.</i> Lindl. col.				Lindl. coll. t. 30.
11 australis. B.P. (wh.)	southern.	N. S. W.	1826.	— F.b. 4.
CALOCH'ILUS. B.P.	CALOCH'ILUS.	Gynandria Monandria. B.P.		
1 campéstris. B.P. (wh.)	field.	N. S. W.	1826.	4. 6. F.b. 4.
2 paludosus. B.P. (wh.)	marsh.	—	—	— F.b. 4.
PONTH'EVA. H.K.	PONTH'EVA.	Gynandria Monandria. H.K.		
1 glandulosa. H.K. (gr.)	glandular.	W. Indies.	1800.	1. 3. S. 4.
2 petiolata. B.R. (br.)	stalked.	—	1823.	7. 8. S. 4.
PRASOPHYLLUM. B.P. PRASOPHYLLUM.		Gynandria Monandria. B.P.		
1 elatum. B.P. tall.	N. S. W.	1826. F.b. 4.
2 flavum. B.P. (ye.)	yellow.	—	—	4. 6. F.b. 4.
3 striatum. B.P. (std.)	streaked.	—	— F.b. 4.
4 fuscum. B.P. (br.)	brown.	—	—	5. 6. F.b. 4.
5 pátens. B.P. spreading.	—	— F.b. 4.
6 rufum. B.P. (br.)	rufous.	—	—	5. 6. F.b. 4.
7 fimbriatum. B.P. fringed.	—	1826. F.b. 4.
CRYPTO'STYLIS. B.P. CRYPTO'STYLIS.		Gynandria Monandria. B.P.		
1 erecta. B.P. (br.)	upright.	N. S. W.	1826.	4. 6. F.b. 4.
2 longifolia. B.P. (br.)	long-leaved.	—	1822.	5. 6. F.b. 4.
<i>Malaxis subulata.</i> Lab.				Lab. n. holl. 2. t. 212.
ORTHO'CERAS. B.P.	ORTHO'CERAS.	Gynandria Monandria. B.P.		
strictum. B.P. upright.	N. S. W.	1826. F.b. 4.
DI'URUS. B.P.	DI'URUS.	Gynandria Monandria. B.P.		
1 maculata. B.P. (ye.sp.)	spotted.	N. S. W.	1823.	3. 5. F.b. 4.
2 aurea. B.P. (go.)	golden-flowered.	—	1810.	4. 5. F.b. 4.
				Sm. exot. bot. 1. t. 30.
				— 1. t. 9.

ORCHIDÆ.

3 pedunculata. B.P. (<i>ye.</i>) peduncled.	N. S. W.	1822.	2. 4. F.b. 4.	
4 sulphurea. B.P. (<i>st.</i>) sulphur-coloured.	_____	1826.	— F.b. 4.	
5 elongata. B.P. (<i>pu.</i>) elongated.	_____	—	— F.b. 4.	
THELYMITRA. B.P. THELYMITRA. Gynandria Monandria. B.P.				
1 ixoides. B.P. (<i>bl.sp.</i>) spotted blue.	N. S. W.	1810.	4. 6. F.b. 4.	
2 media. B.P. (<i>bl.</i>) middle.	_____	1820.	— F.b. 4.	
3 pauciflora. B.P. (<i>bl.</i>) few-flowered.	_____	1826.	— F.b. 4.	
4 angustifolia. B.P. (<i>bl.</i>) narrow-leaved.	_____	—	5. 6. F.b. 4.	
5 carneae. B.P. (<i>fl.</i>) flesh-coloured.	_____	1823.	— F.b. 4.	
6 venosa. B.P. (<i>wh.</i>) veined.	_____	1826.	3. 5. F.b. 4.	
LISTERA. H.K. LISTERA. Gynandria Monandria. H.K.				
1 ovata. H.K. (<i>gr.</i>) Tway-blade. Britain.	5. 6. H. 4.		Eng. bot. v. 22. t. 1548.
'Ophrys ovata. E.B. Epipactis ovata. w.				
2 cordata. H.K. (<i>br.gr.</i>) heart-leaved.	Britain.	6. 7. H. 4.	Eng. bot. v. 5. t. 358.
3 convallarioides. N. (<i>br.gr.</i>) small green.	N. America.	1826.	5. 6. H. 4.	Bart. fl. amer. t. 39. f. 1.
4 Nidus avis. (<i>br.</i>) Bird's-nest.	Britain.	5. H. 4.	Eng. bot. v. 1. t. 48.
'Ophrys Nidus avis. E.B. Epipactis Nidus avis E.F. Neottia Nidus avis. R.O.				
EPIPACTIS. H.K. EPIPACTIS. Gynandria Monandria. H.K.				
1 latifolia. w. (<i>pu.</i>) broad-leaved.	Britain.	7. 8. H. 4.	Eng. bot. v. 4. t. 269.
Serapias latifolia. E.B.				
2 palustris. w. (<i>pu.wh.</i>) marsh.	_____	H. 4.	Eng. bot. v. 4. t. 270.
3 pallens. w. (<i>wh.</i>) white.	_____	6. H. 4.	— v. 4. t. 271.
Serapias grandiflora. E.B.				
4 ensifolia. w. (<i>wh.</i>) narrow-leaved.	_____	H. 4.	Eng. bot. v. 7. t. 494.
5 rubra. w. (<i>pu.wh.</i>) purple.	_____	6. 7. H. 4.	— v. 7. t. 437.
ARETHUSA. H.K. ARETHUSA. Gynandria Monandria. H.K.				
1 bulbosa. H.K. (<i>pu.</i>) bulbous.	N. America.	5. 6. H.b. 4.	Bot. mag. t. 2204.
2? plicata. A.R. (<i>pu.</i>) plaited.	E. Indies.	1806.	6. 8. S.b. 4.	Andr. deposit. t. 321.
POGONIA. H.K. POGONIA. Gynandria Monandria. H.K.				
1 divaricata. H.K. (<i>pi.</i>) Lily-leaved.	N. America.	1787.	6. 7. H. 4.	Catesb. car. 1. t. 58.
2 ophioglossoides. B.R. (<i>pi.</i>) Adder's-tongue.	_____	1816.	— H. 4.	Bot. reg. t. 148.
3 pendula. B.R. (<i>pi.</i>) drooping.	_____	1824.	— H. 4.	— t. 906.
Arethusa pendula. w. parviflora. m. Triphora pendula. n.				
MICROTIS. B.P. MICROTIS. Gynandria Monandria. B.P.				
parviflora. B.P. (<i>gr.</i>) small-flowered.	N. S. W.	1826.	4. 6. F.b. 4.	
ACIANTHUS. B.P. ACIANTHUS. Gynandria Monandria. B.P.				
1 forniciatus. B.P. (<i>br.</i>) enclosed.	N. S. W.	1822.	4. 6. F.b. 4.	
2 exsertus. B.P. (<i>br.</i>) exserted.	_____	—	F.b. 4.	
3 caudatus. B.P. (<i>br.</i>) long-awned.	_____	1826.	— F.b. 4.	
CYRTOSTYLIS. B.P. CYRTOSTYLIS. Gynandria Monandria. B.P.				
reniformis. B.P. (<i>br.</i>) kidney-leaved.	N. S. W.	1824.	4. 5. F. 4.	
CHILOGLOTTIS. B.P. CHILOGLOTTIS. Gynandria Monandria. B.P.				
diphylla. B.P. (<i>br.</i>) two-leaved.	N. S. W.	1826.	4. 6. F.b. 4.	
PRESCOTIA. H.E.F. PRESCOTIA. Gynandria Monandria.				
plantaginifolia. H.E.F. (gr.) Plantain-lily. Rio Janeiro.	1822.	9. 4. S. 4.		
LI PARIS. Rich. LI PARIS. Gynandria Monandria. H.K.				Hook. ex. flor. t. 115.
1 foliosa. B.R. (<i>y.gr.</i>) leafy.	Mauritius.	1823.	9. S. 4.	
2 bituberculata. B.R. (<i>gr.br.</i>) two-warted.	Nepaul.	1822.	2. 3. S. 4.	Bot. reg. t. 882.
3 Loeselii. B.R. (<i>ye.</i>) Loesel's.	Britain.	7. H.b. 4.	Hook. ex. flor. t. 116.
Malaxis Loeselii. E.B.				Eng. bot. v. 1. t. 47.
4 longifolia. B.A. (<i>gr.</i>) long-leaved.	N. America.	1827.	— H.b. 4.	Bart. flor. amer. t. 75.
5 Correaea. N. (<i>ye.gr.</i>) two-leaved.	_____	1823.	5. 6. H.b. 4.	
6 liliifolia. B.R. (<i>pa.bl.</i>) Lily-leaved.	_____	1758.	6. 7. H.b. 4.	Bot. mag. t. 2004.
Malaxis liliiflora. B.M. Ophrys liliiflora. A.R. 65.				
7 reflexa. B.R. (<i>gr.</i>) reflexed.	N. Holland.	1820.	— G.b. 4.	Bot. reg. t. 1175.
8 elata. B.R. (<i>br.pu.</i>) tall.	Brazil.	—	S. 4.	
ERIOCHILUS. B.P. ERIOCHILUS. Gynandria Monandria. B.P.				
autumnalis. B.P. (<i>bh.</i>) autumnal.	N. S. W.	1823.	9. 10. G.b. 4.	Lab.n.holl. 2. t. 211. f. 2.
Epipactis cucullata. Lab.				
CALADENIA. B.P. CALADENIA. Gynandria Monandria. H.K.				
1 alba. B.P. (<i>wh.</i>) white.	N. S. W.	1810.	7. 8. G.b. 4.	
2 carneae. B.P. (<i>fl.</i>) flesh-coloured.	_____	1826.	6. 7. G.b. 4.	
3 coerulea. B.P. (<i>bl.</i>) blue-flowered.	_____	—	G.b. 4.	
4 alata. B.P. (<i>li.</i>) winged.	_____	1823.	6. 8. G.b. 4.	
5 testacea. B.P. fringe-flowered.	_____	1826. G.b. 4.	
LYPERANTHUS. B.P. LYPERANTHUS. Gynandria Monandria. B.P.				
1 suaveolens. B.P. (<i>d.br.</i>) sweet-scented.	N. S. W.	1822.	5. 7. F.b. 4.	
2 elliptica. B.P. (<i>d.br.</i>) elliptic-leaved.	_____	1826.	— F.b. 4.	
3 nigricans. B.P. (<i>bk.</i>) heart-leaved.	_____	—	F.b. 4.	

GLOSS'ODIA. B.P.	GLOSS'ODIA.	Gynandria Monandria. B.P.	
1 major. B.P.	(bl.) larger.	N. S. W. 1810. 6. 8. F.b. 4.	
2 minor. B.P.	(bl.) lesser.	1826. — F.b. 4.	
PTERO'STYLIS. B.P.	PTERO'STYLIS.	Gynandria Monandria. B.P.	
1 concinna. B.P.	(st.) neat.	N. S. W. 1826. 4. 6. F.b. 4.	
2 ophioglòssa. B.P.	(st.) adder's-tongue.	— 1822. 4. 6. F.b. 4.	
3 cùrta. B.P.	(st.) short-lipped.	— 1826. — F.b. 4.	
4 acuminata. B.P.	(st.) taper-pointed.	— 1823. 5. 7. F.b. 4.	
5 cucullata. B.P.	(st.) hooded.	V. Diem. Isl. 1823. 5. 7. F.b. 4.	
6 nútans. B.P.	(st.) nodding.	N. S. W. 1826. — F.b. 4.	
7 obtusa. B.P.	(st.) blunt-lipped.	— 1810. 7. 8. F.b. 4.	
8 reflexa. B.P.	(st.) reflexed.	— 1826. 4. 6. F.b. 4.	
9 grandiflora. B.P.	(st.) large-flowered.	— — — F.b. 4.	
10 longifolia. B.P.	(st.) long-leaved.	— 1823. 6. 8. F.b. 4.	
11 gibbosa. B.P.	(st.) gibbous.	— 1826. 5. 6. F.b. 4.	
CORYSA'NTHES. B.P.	CORYSA'NTHES.	Gynandria Monandria. B.P.	
1 fimbriata. B.P.	(br.) fringed.	N. S. W. 1822. 5. 7. F.b. 4.	
2 unguiculata. B.P.	(br.) pendulous-flowered.	— — — F.b. 4.	
3 bicalcarata. B.P.	(br.) two-spurred.	— — — F.b. 4. Sal. par. lond. t. 83.?	
<i>Corybas aconitiflorus.</i> P.L.?			
CALEYA. H.K.	CALEYA.	Gynandria Monandria. H.K.	
1 major. H.K.	(br.gr.) smooth-lipped.	N. S. W. 1810. 6. 7. F.b. 4.	
2 minor.	(br.gr.) tubercle-lipped.	— 1822. — F.b. 4.	
<i>Caleana minor.</i> B.P.			
CALOP'OGON. H.K.	CALOP'OGON.	Gynandria Monandria. H.K.	
<i>pusillus.</i> H.K.	tuberous-rooted.	N. America. 1771. 7. 8. H.b. 4. Swt. br. fl. gar. t. 115.	
<i>Limodorum tuberosum.</i> B.M. 116.	<i>Cymbidium pulchellum.</i> w.		
BL'ETIA. H.K.	BL'ETIA.	Gynandria Monandria. H.K.	
1 <i>Tankervilliae.</i> H.K. (wh. pu.) <i>LyTankerville's.</i> China. 1778. 3. 4. S. 4.		Bot. mag. t. 1924.	
<i>Limodorum Tankervilliae.</i> w.			
2 pallida. B.C.	(li.) pale-flowered.	China. 1820. 4. 8. S. 4.	
3 florida. H.K.	(pu.) purple.	W. Indies. 1786. 7. 8. S. 4.	
<i>Limodorum purpureum.</i> R.L.			
4 guineensis. G.D.	(pu.) Guinea.	Guinea. 1822. — S. 4.	
5 verecunda. H.K.	(pu.) tall.	W. Indies. 1733. 1. 5. S. 4.	
<i>Limodorum altum.</i> B.M. <i>Cymbidium verecundum.</i> w.		Bot. mag. t. 930.	
6 hyacinthina. H.K.	(pu.) hyacinthine.	China. 1802. 3. 9. G. 4.	
<i>Cymbidium hyacinthinum.</i> B.M.		Bot. mag. t. 1492.	
7 capitata. H.K.	(pu.) capitate.	W. Indies. 1795. 5. 7. S. 4.	
8 Woodfordii. B.M. (ye.or.) <i>Woodford's.</i>	Trinidad. 1820. 4. 8. S. 4.	Bot. mag. t. 2719.	
GEOD'ORUM. H.K.	GEOD'ORUM.	Gynandria Monandria. H.K.	
1 purpureum. H.K.	(pu.) purple.	E. Indies. 1800. 6. 8. S. 4.	
<i>Limodorum nútans.</i> R.C.		Roxb. cor. 1. t. 40.	
2 citrinum. H.K.	(st.) Lemon-coloured.	E. Indies. — 10. 12. S. 4.	
3 dilatatum. H.K.	(pi.) shovel-lipped.	— 5. 8. S. 4.	Andr. deposit. t. 626. Bot. reg. t. 675.
<i>Limodorum recurvum.</i> R.C. 1. t. 39.			
CALY'PSO. H.K.	CALY'PSO.	Gynandria Monandria. H.K.	
<i>americana.</i> H.K.	(pu.) American.	N. America. 1805. 5. 6. H.b. 4.	
<i>borealis.</i> H.E.F. t. 12.		Sal. par. lond. t. 89.	
MALA'XIS. Rich.	MALA'XIS.	Gynandria Monandria. H.K.	
<i>paludosa.</i> E.B.	(ye.) marsh.	England. — 7. H.b. 4.	
MICRO'STYLIS. N.	MICRO'STYLIS.	Gynandria Monandria. N.	
<i>ophioglossoides.</i> N.(gr.) Adder's-tongue.	N. America.	1820. 4. 5. H.b. 4.	
<i>β mexicana.</i> B.R. (gr.) Mexican.	Mexico.	1828. — H. 4.	
CORALLORRHIZA. H.K.	CORALLORRHIZA.	Gynandria Monandria. H.K.	
1 innata. H.K.	(st.) spurless.	Scotland. — 6. 7. H. 4.	
2 multiflora. H.E.F.	(st.) many-flowered.	N. America. 1824. 5. 6. H. 4.	
3 odontorrhiza. N. (w.pu.) tooth-rooted.		— 6. 7. H. 4.	
ST'ELIS. H.K.	ST'ELIS.	Gynandria Monandria. H.K.	
1 ophioglossoides. H.K. (st.) Adder's-tongue.	W. Indies.	1791. 5. 6. S. 4.	
2 micrantha. H.K.	(st.) small-flowered.	Jamaica. 1805. 11. 12. S. 4.	
RODRIGUEZIA. R.P.	RODRIGUEZIA.	Gynandria Monandria. B.R.	
<i>secunda.</i> K.S.	(re.) side-flowering.	S. America. 1819. 6. 12. S. 4.	
GOMEZA. B.M.	GOMEZA.	Gynandria Monandria. B.M.	
<i>recurva.</i> B.M.	(ye.) recurved.	Brazil. 1814. 5. 6. S. 4.	
NOTYLIA. B.R.	NOTYLIA.	Gynandria Monandria.	
<i>punctata.</i> B.R.	(wh.) dotted.	Trinidad. 1823. 7. 8. S. 4.	
<i>Pleurothallis punctata.</i> B.R. <i>Gomeza tenuiflora.</i> Lod. bot. cab. t. 806.		Bot. reg. t. 759.	

CIRRH'EA. B.R.	CIRR'EA. Gynandria Monandria.			Lodd. bot. cab. t. 936.
dépendens. B.R. (<i>ye.gr.</i>) pendent.	China. 1824. 6. 8. S. 4.			
<i>Cymbidium</i> <i>dépendens</i> . B.C.				
CYMB'DIUM. H.K.	CYM'R'DIUM. Gynandria Monandria. H.K.			
1 aloifolium. H.K. (<i>d.pu.</i>) Aloe-leaved.	China. 1789. 5. 6. S. 4.			
2 sinéuse. H.K. (<i>d.pu.</i>) Chinese.	—— 1793. 9. 10. S. 4.			Bot. mag. t. 387.
3 lancifolium. H.E.F. (<i>y.re.</i>) lance-leaved.	E. Indies. 1823. 5. 8. S. 4.			—— t. 888.
4 ensifolium. H.K. (<i>st.br.</i>) sword-leaved.	China. 1780. 6. 10. S. 4.			Hook. ex. flor. t. 51.
5 xiphifolium. B.R. (<i>gr.</i>) Xiphium-leaved	—— 1814. 5. 8. S. 4.			Bot. mag. t. 1751.
6 iridoïdes. D.P. (<i>wh.</i>) Iris-leaved.	Nepaul. 1828. — S. 4.			Bot. reg. t. 529.
7 suáve. B.P. (<i>gr.br.</i>) sweet-scented.	N. S. W. 1826. — G. 4.			
8 tripterum. w. (<i>wh.</i>) triangular-fruited.	Jamaica. 1790. 6. 7. S. 4.			Sm. ic. pict. t. 14.
ANISOP'E'TALUM. H.E.F.	ANISOP'E'TALUM. Gynandria Monandria.			
Careyànum. H.E.F. (<i>br.pu.</i>) Carey's.	Nepaul. 1823. 8. 10. G. 4.			
BRA'SSIA. H.K.	BRA'SSIA. Gynandria Monandria. H.K.			
1 maculata. H.K. (<i>w.st.pu.</i>) spotted.	Jamaica. 1806. 6. 7. S. 4.			
2 caudata. B.R. (<i>g.y.re.</i>) long-tailed.	W. Indies. 1823. — S. 4.			Bot. mag. t. 1691.
ANGUL'O'A. K.S.	ANGUL'O'A. Gynandria Monandria.			Bot. reg. t. 832.
grandiflora. K.S. (<i>ye.pu.</i>) large-flowered.	S. America. 1823. 6. 8. S. 4.			H. et B. pl. æq. 1. t. 27.
CATAS'ETUM. K.S.	CATAS'ETUM. Gynandria Monandria. B.M.			
1 tridentatum. H.E.F. (<i>ye.</i>) three-toothed.	Trinidad. 1823. 8. 12. S. 4.			Bot. mag. t. 2559.
2 Claveríngi. B.R. (<i>y.pu.</i>) Clavering's.	Brazil. — — S. 4.			Bot. reg. t. 840.
3 floribundum. H.E.F. (<i>y.pu.</i>) many-flowered.	—— — — S. 4.			Hook. ex. flor. t. 151.
4 semiapertum. H.E.F. (<i>gr.</i>) greenish-flowered.	—— 1825. 5. S. 4.			—— t. 213.
5 Hookeri. L.col. (<i>y.pu.</i>) Hooker's.	—— 1816. — — S. 4.			Lindl. coll. t. 40.
6 cristatum. B.R. (<i>g.wh.</i>) crested.	—— 1823. — — S. 4.			Bot. reg. t. 966.
CYRTOP'ODIUM. B.R.	CYRTOP'ODIUM. Gynandria Monandria. H.K.			
1 Andersónii. H.K. (<i>ye.</i>) Anderson's.	W. Indies. 1804. 4. 8. S. 4.			Bot. mag. t. 1800.
2 glutinosum. Mey. (<i>ye.</i>) gummy.	S. America. 1825. S. 4.			Bot. mag. t. 1814.
3 Woodfórdii. B.M. (<i>pu.</i>) Woodford's.	—— 1814. 10. S. 4.			Bot. reg. t. 573.
LISSOCH'ILUS. B.R.	LISSOCH'ILUS. Gynandria Monandria. B.R.			
speciosus. B.R. (<i>ye.</i>) showy.	C. B. S. 1818. 5. 6. S. 4.			Jacq. amer. t. 141.
ONCI'DIUM. H.K.	ONCI'DIUM. Gynandria Monandria. H.K.			Bot. mag. t. 777.
1 altissimum. w. (<i>ye.</i>) sharp-petaled.	W. Indies. 1793. 8. 9. S. 4.			
2 carthagénense. w. (<i>g.br.</i>) spread-eagle.	—— 1791. 5. 6. S. 4.			Bot. reg. t. 727.
	<i>Epidendrum undulatum</i> . B.M.			Sloan. hist. t. 148. f. 2.
3 lúridum. B.R. (<i>g.br.</i>) lurid.	Trinidad. 1818. 2. 6. S. 4.			Bot. reg. t. 1050.
4 variegátum. w. (<i>wh.pu.</i>) variegated.	Jamaica. 1825. 7. 8. S. 4.			—— t. 1007.
5 divaricatum. B.R. (<i>br.y.re.</i>) spreading.	Brazil. 1826. 9. 11. S. 4.			H. et B. n. gen. 1. t. 80.
6 pùbes. B.R. (<i>br.re.</i>) pubescent.	—— 1824. 3. 6. S. 4.			Bot. reg. t. 910.
7 ornithorhynchum. K.S. (<i>ye.</i>) long-beaked.	Mexico. 1826. 8. 10. S. 4.			Bot. mag. t. 2203.
8 Papilio. B.R. (<i>y.or pu.</i>) Butterfly.	Trinidad. 1823. 4. 6. S. 4.			Lindl. coll. t. 27.
9 flexuósom. B.M. (<i>ye.</i>) Zigzag.	Brazil. 1818. 6. 7. S. 4.			Bot. reg. t. 920.
10 barbátum. L.col. (<i>ye.</i>) bearded.	S. America. — 4. 5. S. 4.			Bot. mag. t. 1491.
11 pùmulum. B.R. (<i>ye.</i>) dwarf.	Brazil. 1824. 6. 7. S. 4.			—— t. 2773.
12 bifolium. H.K. (<i>ye.</i>) two-leaved.	S. America. 1811. 7. 8. S. 4.			Jacq. amer. t. 142.
13 pulchellum. B.M. (<i>bh.or.</i>) elegant.	Demerara. 1825. 6. 7. S. 4.			—— t. 131. f. 2.
14 triquétrum. H.K. (<i>ye.</i>) triangular-leaved.	Jamaica. 1793. 7. 8. S. 4.			
15 tetrapétalum. w. (<i>pu.ye.</i>) four-petaled.	—— 1824. — — S. 4.			
16 Cebolléta. w. (<i>br.</i>) awl-leaved.	Carthagena. — — S. 4.			
MACRAD'ENIA. B.R.	MACRAD'ENIA. Gynandria Monandria. B.R.			
lutescens. B.R. (<i>ye.</i>) yellowish.	Trinidad. 1821. 12. 2. S. 4.			Bot. reg. t. 612.
CŒLO'GYNE. B.R.	CŒLO'GYNE. Gynandria Monandria.			Bot. reg. t. 868.
1 fimbriata. B.R. (<i>st.</i>) fringed.	China. 1824. 9. S. 4.			
2 nítida. L.O.	(<i>ye.</i>) shining-leaved. E. Indies. 1822. S. 4.			Bot. reg. t. 989.
3 punctulata. L.O. (<i>ye.</i>) dot-flowered.	—— — — S. 4.			Lindl. coll. t. 2.
MEGACL'IUM. B.R.	MEGACL'IUM. Gynandria Monandria. B.R.			
falcatum. B.R. (<i>ye.re.</i>) falcate.	Sierra Leone. 1824. 6. 9. S. 4.			
TRIZ'EUXIS. L.col.	TRIZ'EUXIS. Gynandria Monandria.			
falcata. L.col. (<i>gr.</i>) falcate.	Trinidad. 1820. 2. 6. S. 4.			
PLEUROTHA'LLIS. H.K.	PLEUROTHA'LLIS. Gynandria Monandria. H.K.			
1 ruscifolia. H.K. (<i>st.</i>) Ruscus-leaved.	W. Indies. 1791. 5. 6. S. 4.			
2 racemiflora. H.E.F. (<i>st.</i>) racemed.	—— 1822. 7. 8. S. 4.			Hook. ex. flor. t. 197.
3 prolifera. B.R. (<i>ro.</i>) proliferous.	Brazil. 1825. 1. 12. S. 4.			—— t. 123.
4 foliosa. B.M. (<i>st.ye.</i>) leafy.	—— 2. 6. S. 4.			Bot. reg. t. 1298.
ORNITHOCE'PHALUS. BIRD'S-HEAD.	Gynandria Monandria.			Bot. mag. t. 2746.
gladiatus. H.E.F. (<i>gr.</i>) sword-leaved.	Trinidad. 1822. 5. 8. S. 4.			Hook. ex. flor. t. 127.
OCTOM'ERIA. H.K.	OCTOM'ERIA. Gynandria Monandria. H.K.			
1 graminifolia. H.K. (<i>st.</i>) Grass-leaved.	W. Indies. 1793. 4. 7. S. 4.			Bot. mag. t. 2764.

2 serratifolia, B.M. (<i>st. br.</i>) saw-leaved.	Brazil.	1826.	10. 12. S. 4.	Bot. mag. t. 2823.
3 spicata, D.P. (<i>wh.</i>) spiked.	Nepaul.	1823.	5. 8. S. 4.	
TRIBRA'CHIA. B.R. TRIBRA'CHIA.	Gynandria Monandria.			
péndula, B.R. (<i>gr.</i>) pendulous-spiked.	Sierra Leon.	1822.	6. 11. S. 4.	Bot. reg. t. 963.
SOPHRON'ITIS. B.R. SOPHRON'ITIS.	Gynandria Monandria.			
cérnua, B.R. (<i>ro.</i>) cernuous.	Brazil.	1826.	11. 12. S. 4.	Bot. reg. t. 1129.
GONG'ORA. R.P.	GONG'ORA.	Gynandria Monandria.		
1 atropurpurea, H.E.F. (<i>d. pu.</i>) dark-flowered.	Trinidad.	1824.	6. 7. S. 4.	Hook. ex. flor. t. 178.
2 speciosa, B.M. (<i>ye. or.</i>) beautiful.	Brazil.	1826.	4. 6. S. 4.	Bot. mag. t. 2755.
3 viridipurpurea, B.M. (<i>g. pu.</i>) green & purple.	—	1827.	6. 8. S. 4.	— t. 2978.
HETEROTA'XIS. B.R. HETEROTA'XIS.	Gynandria Monandria.			
crassifolia, B.R. (<i>ye.</i>) thick-leaved.	Jamaica.	1823.	6. 8. S. 4.	Bot. reg. t. 1028.
DIP'ODIUM. B.P.	DIP'ODIUM.	Gynandria Monandria. B.P.		
punctatum, B.P. (<i>pu.</i>) spotted.	N.S.W.	1822.	6. 8. G. 4.	Sm. exot. bot. 1. t. 12.
<i>Dendrobium punctatum.</i> Sm.				
CAMARI'DIUM. B.R. CAMARI'DIUM.	Gynandria Monandria. B.R.			
ochroleucum, B.R. (<i>st.</i>) pale-flowered.	Trinidad.	1823.	7. 10. S. 4.	Bot. reg. t. 844.
<i>Dendrobium? album.</i> H.E.F. t. 142.				
'ERIA. B.R.	'ERIA.	Gynandria Monandria. B.R.		
1stellata, B.R. (<i>st.</i>) starry.	Nepaul.	1818.	5. 6. S. 4.	Bot. reg. t. 904.
2 pubescens, B.R. (<i>ye.</i>) pubescent.	E. Indies.	1820.	3. 4. S. 4.	Hook. ex. flor. t. 124.
<i>Dendrobium pubescens.</i> H.E.F.				
3 rösea, B.R. (<i>bh.</i>) rose-coloured.	China.	1824.	10. S. 4.	Bot. reg. t. 978.
XYL'OBIVUM. B.R.	XYL'OBIVUM.	Gynandria Monandria.		
1 squáleum, B.R. (<i>ye. br.</i>) squálid.	Brazil.	1822.	7. 8. S. 4.	Bot. reg. t. 732.
<i>Dendrobium squáleum.</i> B.R. <i>Maxillaria squáleum.</i> Bot. mag. t. 2955.				
2 racemosum, (<i>ye. or.</i>) racemed.	Brazil.	1825.	6. 8. S. 4.	Bot. mag. t. 2789.
<i>Maxillaria racemosa.</i> B.M.				
MAXILL'ARIA. R.P. MAXILL'ARIA.	Gynandria Monandria. B.R.			
1 pallidiflora, B.M. (<i>st.</i>) pale-flowered.	St. Vincent's.	1826.	8. 11. S. 4.	Bot. mag. t. 2806.
2 ciliata, B.R. (<i>gr.</i>) fringe-lipped.	S. America.	1827.	5. 8. S. 4.	Bot. reg. t. 1206.
3 Barringtoniæ, B.R. (<i>gr.</i>) Barrington's.	W. Indies.	1790.	6. 8. S. 4.	Sm. ic. pict. t. 25.
<i>Dendrobium Barringtoniæ.</i> Hook. ex. flor. t. 119.				
4 Harrisoniæ, B.R. (<i>st. pu.</i>) Mrs. Harrison's.	Brazil.	1820.	4. 8. S. 4.	Bot. mag. t. 2927.
5 Parkéi, B.M. (<i>wh. pu.</i>) Parker's.	W. Indies.	1826.	— S. 4.	— t. 2729.
6 párvula, H.E.F. (<i>st. pu.</i>) small.	Brazil.	1824.	3. 5. S. 4.	Hook. ex. flor. t. 217.
7 aromática, H.E.F. (<i>ye.</i>) aromatic.	—	1825.	5. S. 4.	— t. 219.
SARCOCH'ILUS. B.P. SARCOCH'ILUS.	Gynandria Monandria. B.P.			
falcatus, B.P. (<i>pu.</i>) falcate.	N.S.W.	1821.	5. 8. G. 4.	
DENDROBIVUM. H.K. DENDROBIVUM.	Gynandria Monandria. H.K.			
1 speciosum, B.P. (<i>bh. re.</i>) showy.	N.S.W.	1801.	6. 8. G. 4.	Sm. exot. bot. 1. t. 10.
2 æmulum, B.P. (<i>wh.</i>) rival.	—	1822.	4. 2. G. 4.	Bot. mag. t. 2906.
3 rigidum, B.P. rigid.	N. Holland.	1824. S. 4.	
4 lingüiforme, B.P. (<i>w. st.</i>) tongue-leaved.	N. S. W.	1810.	4. 8. G. 4.	Sm. exot. bot. 1. t. 11.
5 teretifolium, B.P. (<i>pu.</i>) filiform-leaved.	—	1823.	— G. 4.	
6 longicórnus, B.R. (<i>bh.</i>) long-spurred.	E. Indies.	1828.	4. 6. S. ḥ.	Bot. reg. t. 1315.
7 secundum, B.R. (<i>ro.</i>) secund.	Sumatra.	—	— S. ḥ.	— t. 1291.
8 cucullatum, B.R. (<i>bh.</i>) cucullate.	E. Indies.	1815.	3. 5. S. ḥ.	Bot. reg. t. 548.
9 Pierárdi, H.E.F. (<i>bh.</i>) Pierard's.	—	—	12. 6. S. ḥ.	Hook. ex. flor. t. 9.
10 Calceolária, H.E.F. (<i>re. y.</i>) hollow-lipped.	—	1820.	— S. ḥ.	— t. 184.
11 fimbriatum, H.E.F. (<i>ye.</i>) fringed.	—	1818.	4. 6. S. ḥ.	— t. 71.
12 chrysanthum, B.R. (<i>ye.</i>) yellow-flower'd.	Nepaul.	1828.	2. 4. S. ḥ.	Bot. reg. t. 1299.
13 crumenatum, W. (<i>pu.</i>) pouched.	Java.	1822.	4. 6. S. ḥ.	Rumph.amb.6.t.47.f.2.
14 moniliforme, W. (<i>pu.</i>) glassy.	China.	1822.	6. 12. S. ḥ.	Bot. reg. t. 1314.
15 ancéps, B.R. (<i>st.</i>) two-edged.	E. Indies.	1826.	2. 9. S. ḥ.	— t. 1239.
ORNITHI'DIUM. H.K. ORNITHI'DIUM.	Gynandria Monandria. H.K.			
1 coccineum, H.K. (<i>sc.</i>) scarlet-flower'd.	W. Indies.	1790.	1. 12. S. 4.	Hook. ex. flor. t. 38.
<i>Cymbidium coccineum.</i> Bot. mag. t. 1437.				
2 reflexum, Gr. (<i>sc.</i>) reflexed.	W. Indies.	1825.	9. 12. S. 4.	
ISOCH'ILUS. H.K. ISOCH'ILUS.	Gynandria Monandria. H.K.			
1 linearis, H.K. (<i>re.</i>) linear-leaved.	W. Indies.	1791.	5. 7. S. 4.	Bot. reg. t. 745.
<i>Epidendrum lineare.</i> Jacq. amer. t. 131. f. 1. <i>Cymbidium lineare.</i> w.				
2 gramineoides, H.E.F. (<i>st.</i>) grass-like.	W. Indies.	1823.	— S. 4.	Hook. ex. flor. t. 196.
3 prolifér, H.K. (<i>wh.</i>) proliferous.	—	1793.	6. 11. S. 4.	Bot. reg. t. 825.
PHOLID'OTA. B.R. PHOLID'OTA.	Gynandria Monandria.			
imbricata, B.R. (<i>br. wh.</i>) imbricated.	Nepaul.	1824.	2. 8. S. 4.	Bot. reg. t. 1213.
POLYST'ACHIA. H.E.F. POLYST'ACHIA.	Gynandria Monandria.			
1 luteola, H.E.F. (<i>y. gr.</i>) smooth.	W. Indies.	1818.	7. 8. S. 4.	Hook. ex. flor. t. 103.
<i>Dendrobium polystachyum.</i> w.				

2 pubérula. B.R. (<i>y.gr.</i>) downy.	Sierra Leone. 1821.	8.	S. 2.	Bot. reg. t. 851.
BRASSAVOLA. H.K. BRASSAVOLA.	Gynandria Monandria. H.K.			Bot. mag. t. 543.
1 cucullata. H.K. (<i>wh.</i>) single-flowered. W. Indies.	1793.	6. 7.	S. 2.	—— t. 2878.
2 tuberculata. B.M. (<i>wh.st.</i>) tuberculated. Brazil.	1826.	6. 9.	S. 2.	
3 nodosa. Ly. (<i>wh.</i>) many-flowered. W. Indies.	1824.	10.	S. 2.	
BROUGHTONIA. H.K. BROUGHTONIA.	Gynandria Monandria. H.K.			Lodd. bot. cab. t. 793.
1 sanguinea. H.K. (<i>cr.</i>) blood-coloured. Jamaica.	1793.	6. 7.	S. 2.	
2 nítida. W.H. (<i>re.</i>) glossy.	E. Indies. 1824.	—	S. 2.	
ZYGOPE'TALON. B.M. ZYGOPE'TALON.	Gynandria Monandria.			Bot. mag. t. 2748.
1 Mackaii. B.M. (<i>gr.bl.</i>) Mackay's.	Brazil. 1825.	2. 5.	S. 2.	—— t. 2819.
2 rostratum. B.M. (<i>wh.pu.</i>) beaked.	Demarara.	—	9. 11.	S. 2.
CA'TTLEYA. L.col. CA'TTLEYA.	Gynandria Monandria. H.E.F.			Lindl. coll. t. 33.
1 labiata. L.col. (<i>vi.</i>) great-lipped.	S. America. 1818.	4. 10.	S. 2.	Hook. ex. flor. t. 186.
2 Loddigèsiæ. H.E.F. (<i>vi.</i>) Loddiges'.	1810.	—	S. 2.	
<i>Epidendrum violaceum.</i> Lodd. bot. cab. t. 337.				
3 Forbesii. B.R. (<i>ye.</i>) yellow.	Brazil. 1823.	9.	S. 2.	Bot. reg. t. 953.
4 crispa. B.R. (<i>wh.pu.</i>) curled-lipped.	Trinidad. 1820.	—	S. 2.	—— t. 1172.
5 intermédia. B.M. (<i>ro.</i>) intermediate.	Brazil. 1824.	3. 5.	S. 2.	Bot. mag. t. 2851.
CERATOCH'ILUS. B.C. CERATOCH'ILUS.	Gynandria Monandria.			Lodd. bot. cab. t. 1414.
1 grandiflorus. B.C. (<i>wh.</i>) large-flowered.	Trinidad. 1827.	3. 8.	S. 2.	Bot. mag. t. 2948-2949.
2 insignis. (<i>st.pu.</i>) splendid.	S. America. 1818.	9. 4.	S. 2.	
<i>Stanhopea insignis.</i> B.M.				
EPIDE'NDRUM. H.K. EPIDE'NDRUM.	Gynandria Monandria. H.K.			Bot. mag. t. 572.
1 cochleatum. w. (<i>d.pu.</i>) shell-flowered.	W. Indies. 1786.	1. 12.	S. 2.	—— t. 1669.
2 frágans. w. (<i>st.</i>) sweet-scented.	—	1778.	10.	Jacq. amer. t. 137.
3 secundum. w. (<i>re.</i>) pale-flowered.	—	1793.	6. 7.	Hook. ex. flor. t. 112.
4 polybulbon. w. (<i>wh.</i>) many-bulbed.	—	1822.	—	Lodd. bot. cab. t. 1084.
5 verrucosum. w. (<i>st.</i>) warted.	—	1824.	—	Hook. ex. flor. t. 50.
6 nútans. w. (<i>gr.</i>) nodding.	—	1794.	—	Bot. reg. t. 80.
7 umbellatum. w. (<i>st.</i>) umbelled.	Jamaica. 1793.	—	S. 2.	Lodd. bot. cab. t. 846.
8 diffusum. w. (<i>gr.</i>) diffuse.	W. Indies. 1818.	4. 5.	S. 2.	—— t. 887.
9 ancéps. B.C. (<i>gr.</i>) two-edged.	—	1820.	8. 10.	S. 2.
10 nocturnum. w. (<i>gr.</i>) night-smelling.	—	—	4. 6.	S. 2.
11 pallidiflorum. B.M. (<i>st.</i>) pale-flowered.	—	1828.	5. 6.	S. 2.
12 fuscatum. w. (<i>br.</i>) brown.	—	1790.	6. 7.	S. 2.
13 elongatum. w. (<i>re.</i>) long-stalked.	Caracas.	1798.	5. 8.	S. 2.
14 ellípticum. H.E.F. (<i>re.</i>) elliptic-leaved.	Brazil.	1824.	3. 8.	S. 2.
15 conópseum. H.K. (<i>st.</i>) Florida.	Florida.	1775.	8.	S. 2.
16 monophyllo. H.E.F. (<i>gr.</i>) one-leaved.	W. Indies.	1822.	7.	S. 2.
17 ciliare. B.R. (<i>wh.</i>) fringed.	—	1790.	6. 10.	S. 2.
18 cuspidatum. B.R. (<i>wh.</i>) pointed.	—	1808.	—	S. 2.
ENCYCLIA. B.M. ENCYCLIA.	Gynandria Monandria.			Bot. mag. t. 2980.
viridiflora. B.M. (<i>gr.</i>) green-flowered.	Brazil.	1826.	2. 6.	Bot. reg. t. 67.
FERNAND'ESIA. R.P. FERNAND'ESIA.	Gynandria Monandria.			Bot. mag. t. 611.
élegans. B.C. (<i>st.pu.</i>) elegant.	Trinidad.	1822.	4. 10.	S. 2.
<i>Lockhartia elegans.</i> Bot. mag. t. 2715.				Hook. ex. flor. t. 207.
CRYPTARRH'ENA. B.R. CRYPTARRH'ENA.	Gynandria Monandria. B.R.			Bot. mag. t. 109.
lunata. B.R. (<i>ye.</i>) crescent-lipped.	W. Indies.	1815.	5. 8.	Bot. reg. t. 784.
EUL'OPHIA. B.R. EUL'OPHIA.	Gynandria Monandria. B.R.			—— t. 783.
1 virens. B.R. (<i>gr.</i>) green.	E. Indies. 1823.	6. 8.	S. 2.	Bot. mag. t. 2831.
<i>Limodorum virens.</i> R.C.				
2 guineensis. B.R. (<i>pi.</i>) Guinea.	Sierra Leone. 1822.	11. 2.	S. 2.	Lodd. bot. cab. t. 1212.
3 grátilis. B.R. (<i>gr.</i>) slender	—	—	6. 9.	Bot. reg. t. 153.
4 streptopétala. B.R. (<i>ye.</i>) twisted-petaled.	C. B. S.	—	12. 6.	Roxb. corom. t. 1. 38.
5 ensata. B.R. (<i>st.</i>) sword-leaved.	—	1826.	7. 8.	Bot. reg. t. 686.
6 tristis. B.R. (<i>da.</i>) dark-flowered.	—	1825.	5. 6.	—— t. 742.
7 longicórnis. B.R. (<i>ye.</i>) long-spurred.	—	—	6. 8.	—— t. 1002.
8 barbata. B.R. (<i>ye.</i>) bearded.	—	1822.	—	—— t. 1147.
CALANTHE. B.R. CALANTHE.	Gynandria Monandria. B.R.			
1 veratrifolia. B.R. (<i>wh.</i>) Veratrum-leav'd.	E. Indies. 1819.	1. 9.	S. 2.	Bot. reg. t. 720.
2 sylvétris. Ly. (<i>wh.</i>) loose-spiked.	Madagascar. 1823.	6. 7.	S. 2.	
ANGRÆCUM. L.col. ANGRÆCUM.	Gynandria Monandria. B.R.			Bot. reg. t. 618.
1 maculatum. B.R. (<i>fl.</i>) spotted.	S. America. 1819.	10. 12.	S. 2.	—— t. 283.
2 falcatum. L.col. (<i>wh.</i>) falcate.	China. 1815.	—	S. 2.	
<i>Limodorum falcatum.</i> B.R.				
3 lúridum. Ly. (<i>br.</i>) lurid.	Sierra Leone. 1822.	7.	S. 2.	
IONOPSIS. K.S. IONOPSIS.	Gynandria Monandria.			
utricularioïdes. L.col. (<i>w.</i>) pale-flowered.	Trinidad. 1822.	6. 9.	S. 2.	Hook. ex. flor. t. 113.
<i>Iántha pálida.</i> H.E.F.				

ORDO CCIII.

SCITAMINEÆ. *Brown prodr.* 1. p. 305

- | | | | | | | |
|--|--|-------|--------------|---------------------------|-----------------------|--|
| GLO'BBA. L. | GLO'BBA. Monandria Monogynia. L. | | | | | |
| 1 marantina. w. | (ye.) marantine. E. Indies. | 1800. | 7. 8. S. 24. | | Exot. bot. 2. t. 103. | |
| 2 bulbifera. F.I. | (ye.) bulb-bearing. ————— | 1818. | — S. 24. | | | |
| 3 orixensis. F.I. | (or.) Orixian. ————— | 1819. | — S. 24. | | | |
| 4 versicolor. Ex.B. (y.e.) various-colour'd. | ————— | 1820. | — S. 24. | Sm. exot. bot. 2. t. 117. | | |
| <i>Hura. F.I.</i> | | | | | | |
| 5 péndula. F.I. | (ye.) pendulous-flower'd. ————— | 1822. | 6. 8. S. 24. | | | |
| 6 Careyana. F.I. | (ye.) Carey's. ————— | 1821. | — S. 24. | Rosc. scit. t. 110. | | |
| 7 sessiliflora. B.M. | (ye.) sessile-flower'd. ————— | 1807. | — S. 24. | Bot. mag. t. 1428. | | |
| MANTI'SIA. B.M. | MANTI'SIA. Monandria Monogynia. B.M. | | | | | |
| 1 saltatòria. B.M. (bl.ye.) Opera-girl's. | E. Indies. | 1808. | 6. S. 24. | Bot. mag. t. 1320. | | |
| <i>Glòbba subulata. F.I. purpurea. A.R.</i> | 615. <i>saltatòria. R. scit. t. 111.</i> | | | | | |
| 2 spathulata.R.S. (bl.ye.) spathulate. | E. Indies. | 1822. | 6. 8. S. 24. | | | |
| <i>Glòbba spathulata.F.I.</i> | | | | | | |
| CURC'UMA. W. | TURMERIC. Monandria Monogynia. L. | | | | | |
| 1 aromática. P.L.(ye.pi.) aromatic. | E. Indies. | 1797. | 4. 8. S. 24. | Sal. par. lond. t. 96. | | |
| <i>Zerumbet. F.I. Rheed. mal. v. 11. t. 7.</i> | | | | | | |

2 Zedoària. B.M. (<i>ye.pi.</i>) silky-leaved.	E. Indies.	1797.	4. 8.	S. 2.	Bot. mag. t. 1546.
3 zanthorrhiza. F.I. (<i>re.cr.</i>) yellow-rooted.	Amboyna.	1819.	4. 5.	S. 2.	Rosc. scitam. t. 108.
4 latifòlia. R.Sc. (<i>ye.pi.</i>) broad-leaved.	E. Indies.	—	—	S. 2.	— t. 104.
5 elàta. F.I. (<i>ye.pu.</i>) tall.	—	—	—	S. 2.	
6 cæsia. F.I. (<i>ye.re.</i>) clouded-leaved.	—	—	5. 6.	S. 2.	Rosc. scitam. t. 106.
7 æruginosa. F.I. (<i>ye.re.</i>) verdigrease.	—	1807.	4. 8.	S. 2.	— t. 105.
8 ferruginea. F.I. (<i>ye.re.</i>) rusty.	—	1819.	—	S. 2.	— t. 107.
9 rubescens. F.I. (<i>re.ye.</i>) red.	—	—	4. 6.	S. 2.	Rosc. scitam. t. 102.
10 comosa. F.I. (<i>ye.pi.</i>) close-flowered.	—	—	—	S. 2.	Bot. reg. t. 886.
11 leucorrhiza. F.I. (<i>yo.ro.</i>) white-rooted.	—	—	—	S. 2.	Rosc. scitam. t. 99.
12 angustifòlia. F.I. (<i>y.pu.</i>) narrow-leaved.	—	1822.	—	S. 2.	Roxb. corom. 2. t. 151.
13 lónga. F.I. (<i>st.bh.</i>) long-rooted	—	1759.	6. 8.	S. 2.	
14 Amáda. F.I. (<i>st.li.</i>) Mango-ginger.	—	1819.	—	S. 2.	Rosc. scitam. t. 100.
15 viridiflòra. F.I. (<i>gr.st.</i>) green-flowered.	—	1820.	—	S. 2.	
16 montána. F.I. (<i>ye.ro.</i>) mountain.	—	1823.	4. 6.	S. 2.	Rosc. scitam. t. 101.
17 reclinata. F.I. (<i>pi.</i>) reclining.	—	1824.	—	S. 2.	
18 petiolata. F.I. (<i>ye.pu.</i>) petiolate.	—	1822.	—	S. 2.	Bot. mag. t. 850.
19 rubricaulis. L.T. (<i>re.ye.</i>) red-stalked.	—	1818.	—	S. 2.	— t. 920.
20 amarissima. R.Sc. (<i>re.ye.</i>) bitter.	—	1822.	—	S. 2.	
KÆMPF'ERIA. L. KÆMPF'ERIA.	Monandria Monogynia. L.				
1 Galánga. w. (<i>pu.wh.</i>) officinal.	E. Indies.	1724.	6. 9.	S. 2.	
2 rotunda. w. (<i>pu.bh.</i>) round-rooted.	—	1768.	7. 8.	S. 2.	Bot. reg. t. 1212.
3 latifòlia. s.s. (<i>wh.</i>) broad-leaved.	—	1803.	4. 7.	S. 2.	Rosc. scitam. t. 94.
4 Roscoëana. B.R. (<i>wh.</i>) Roscoe's.	Ava.	1828.	—	S. 2.	Bot. reg. t. 173.
5 angustifòlia. F.I. (<i>pu.w.</i>) narrow-leaved.	E. Indies.	1797.	—	S. 2.	Rosc. scitam. t. 95.
6 pandurata. F.I. (<i>ro.</i>) fiddle-lipped.	—	1812.	—	S. 2.	— t. 93.
7 ovalifòlia. F.I. (<i>pu.wh.</i>) oval-leaved.	—	1819.	—	S. 2.	Rosc. scitam. t. 64.
8 marginata. r.s. (<i>pu.</i>) red-margined.	—	1820.	—	S. 2.	
ROSC'OEÀ. L.T. ROSC'OEÀ. Monandria Monogynia.	Monandria Monogynia. L.				
1 purpurea. L.T. (<i>pu.</i>) purple-flower'd. Nepaul.	—	1819.	7. 8.	S. 2.	
2 grácilis. L.T. (<i>pu.</i>) slender.	—	1820.	—	S. 2.	Rosc. scitam. t. 83.
3 elàtior. L.T. (<i>pu.</i>) tall.	—	—	—	S. 2.	Sm. exot. bot. 2. t. 112.
4 spicata. L.T. (<i>pu.</i>) spiked.	—	1821.	—	S. 2.	Bot. mag. t. 1426.
5 capitata. L.T. (<i>pu.</i>) headed.	—	1819.	—	S. 2.	
ZI'NGIBER. L.T. GINGER. Monandria Monogynia. H.K.	Monandria Monogynia. H.K.				
1 officinale. F.I. (<i>pu.st.</i>) narrow-leaved.	E. Indies.	1605.	6. 8.	S. 2.	
2 Zerúmbet. F.I. (<i>st.</i>) broad-leaved.	—	1690.	9. 11.	S. 2.	Roxb. corom. 2. t. 126.
3 Cassumunar. B.M. (<i>wh.</i>) downy-leaved.	—	1796.	8.	S. 2.	
<i>purpureum</i> . L.T. <i>Amomum Cliffordia</i> . And. repos. t. 555.					
4 róseum. F.I. (<i>ro.st.</i>) rose-coloured.	E. Indies.	1822.	6. 8.	S. 2.	Rosc. scitam. t. 88.
5 ligulatum. F.I. (<i>pi.ye.</i>) ligulate.	—	—	—	S. 2.	— t. 89.
6 rübēns. F.I. (<i>re.</i>) red-flowered.	—	1820.	5. 11.	S. 2.	Kæmpf. ic. Banks. t. 1.
7 squarrosum. F.I. (<i>wh.</i>) squarrose.	—	—	—	S. 2.	Rosc. scitam. t. 86.
8 panduratum. F.I. (<i>st.re.</i>) fiddle-leaved.	—	1819.	5. 8.	S. 2.	— t. 90.
9 Miòga. L.T. (<i>pi.</i>) Japanese.	Japan.	1796.	—	S. 2.	— t. 91.
10 chrysanthum. R.Sc. (<i>ye.</i>) yellow-flower'd. Nepaul.	—	1821.	—	S. 2.	
11 capitatum. F.I. (<i>ye.</i>) headed.	E. Indies.	1820.	6. 9.	S. 2.	
12 elàtum. F.I. (<i>ye.</i>) tall.	—	—	—	S. 2.	
AM'OMUM. F.I. AM'OMUM. Monandria Monogynia. L.	Monandria Monogynia. L.				
1 Afzélii. L.T. (<i>pi.</i>) sweet-scented. Sierra Leone.	1795.	5. 6.	S. 2.		Ann. bot. 1. t. 13.
2 grandiflorum. R.S. (<i>pi.w.</i>) large-flower'd.	—	—	6. 7.	S. 2.	Sm. exot. bot. t. 3.
3 sylvestre. R.S. (<i>ye.</i>) wood.	Jamaica.	1804.	5. 7.	S. 2.	Sloan. hist. 1.t.105.f.2.
4 Cardamomum. F.I. (<i>ye.</i>) bastard Cardamom. E. Indies.	1820.	4. 6.	S. 2.		Rumph. amb. 5.t.65.f.1.
5 angustifolium. F.I. (<i>re.y.</i>) narrow-leaved. Madagascar.	—	—	—	S. 2.	Sonnerat. it. 2. f. 137.
6 aculeatum. F.I. (<i>or.cr.</i>) bristly-capsuled. E. Indies.	1823.	6. 8.	S. 2.		Bot. reg. t. 929.
7 máximum. F.I. (<i>re.ye.</i>) great.	—	—	—	S. 2.	
8 dealbatum. F.I. (<i>wh.</i>) insipid-seeded.	—	1819.	—	S. 2.	
9 subulatum. F.I. (<i>ye.</i>) subulate-bracted.	—	—	5. 7.	S. 2.	
10 aromaticum. F.I. (<i>st.</i>) aromatic.	—	1823.	—	S. 2.	
11 sericeum. F.I. (<i>wh.pi.</i>) silky-leaved.	—	1819.	6. 8.	S. 2.	Blackw. t. 385.
12 grána-paradisi. w. (<i>re.</i>) Grains of Paradise. Madagascar.	—	2. 4.	S. 2.		
CO'STUS. F.I. CO'STUS. Monandria Monogynia. L.	Monandria Monogynia. L.				
1 speciosus. F.I. (<i>wh.</i>) large-flowered.	E. Indies.	1794.	8. 9.	S. 2.	Jacq. ic. 1. t. 1.
2 nepalensis. R.Sc. (<i>wh.</i>) Nepaul.	Nepaul.	1818.	—	S. 2.	Rosc. scitam. t. 80.
<i>speciosus</i> β. Bot. reg. 665.					
3 cylindricus. R.S. (<i>wh.re.</i>) cylindrical-spiked. Brazil.	1822.	5. 9.	S. 2.		Rosc. scitam. t. 78.
<i>Pisonis</i> . Bot. reg. t. 899.					
4 áfer. B.R. (<i>wh.</i>) African.	Sierra Leone.	—	—	S. 2.	Bot. reg. t. 683.
5 discolor. R.Sc. (<i>wh.</i>) two-coloured.	Maranham.	1823.	—	S. 2.	Rosc. scitam. t. 81.

6	maculatus. R.Sc. (<i>wh.</i>) spotted-stalked.	Sierra Leone. 1822.	5.	9.	S. 2.	Rosc. scitam. t. 82.	
7	lanatus. L.C. (<i>wh.</i>) woolly.	—	—	S. 2.		
8	spicatus. L.T. (<i>ye.</i>) spiky.	S. America. 1793.	6.	—	S. 2.	Rosc. scitam. t. 77.	
9	comosus. L.T. (<i>re.ye.</i>) tufted. 1815.	—	—	S. 2.	Jacq. ic. 3. t. 202.	
	Alpinia comosa. Jacq.						
10	spiralis. L.T. (<i>re.</i>) spiral.	5.	7.	S. 2.	Jacq. schœn. 1. t. 1.	
11	villossima. R.S. (<i>ye.</i>) villous.	W. Indies. 1823.	—	—	S. 2.	Jacq. fragm. t. 80.	
12	arabicus. L.T. (<i>wh.</i>) Arabian.	Arabia. 1752.	8.	—	S. 2.		
HELL'ENIA. B.P.	HELL'ENIA. Monandria Monogynia. W.						
	cœrulea. B.P. (<i>bh.</i>) blue-berried.	N. Holland. 1820.	5.	6.	G. 2.	Rosc. scitam. t. 76.	
ALPI'NIA. L.T.	ALPI'NIA. Monandria Monogynia. L.						
1	Allughas. F.I. (<i>wh.re.</i>) glossy-leaved.	E. Indies. 1796.	2.	3.	S. 2.	Rosc. scitam. t. 67.	
	Hellénia Allughas. Andr. deposit. t. 501.						
2	racemosa. L.T. (<i>wh.</i>) clustered.	W. Indies. 1752.	7.	9.	S. 2.	Plum. ic. 11. t. 20.	
3	Galânga. F.I. (<i>g.wh.</i>) Galanga-major.	E. Indies. 1818.	6.	10.	S. 2.	Rumph. amb. 5. t. 63.	
4	Roxburghii. (<i>w.cr.</i>) Roxburgh's. 1820.	—	—	S. 2.		
	bracteata. F.I. non L.T.						
5	malaccensis. F.I. (<i>w.y.cr.</i>) woolly-leaved. 1799.	4.	6.	S. 2.	Bot. reg. t. 328.	
6	nûtans. F.I. (<i>re.ye.</i>) nodding-flower'd. 1792.	—	—	S. 2.	Bot. mag. t. 1903.	
	Renealmia nûtans. A.R. 360.						
7	cérnua. B.M. (<i>y.wh.pu.</i>) drooping. 1790.	—	—	S. 2.	Bot. mag. t. 1900.	
8	auriculata. R.Sc. (<i>ye.re.</i>) eared. 1815.	—	—	S. 2.	Rosc. scitam. t. 74.	
9	diffissa. L.T. (<i>ye.</i>) cloven.	China. —	4.	6.	S. 2.	— t. 71.	
10	mütica. F.I. (<i>ye.re.</i>) spurless.	E. Indies. 1811.	4.	9.	S. 2.	— t. 69.	
11	calcarata. F.I. (<i>re.or.</i>) upright spiked.	China. 1804.	9.	—	S. 2.	Andr. deposit. t. 421.	
12	spicata. F.I. (<i>wh.pu.</i>) spiky.	E. Indies. 1823.	6.	9.	S. 2.		
13	striata. L.en. (<i>std.</i>) striated. 1818.	—	—	S. 2.		
14	bracteata. L.T. (<i>ye.re.</i>) bracted.	China. 1823.	5.	8.	S. 2.	Rosc. scitam. t. 70.	
	Roscoëna. R.S.						
15	magnifica. R.Sc. (<i>sc.</i>) magnificent.	Madagascar. 1826.	—	S. 2.	Rosc. scitam. t. 75.	
RENE'ALMIA. R.Sc.	RENE'ALMIA. Monandria Monogynia.						
1	occidentalis. (<i>pa.</i>) occidental.	Jamaica. 1793.	5.	7.	S. 2.	Gært. fruct. 1. t. 12.	
	Alpinia occidentalis. L.T.						
2	exaltata. R.Sc. (<i>re.</i>) tubular-flowered.	Demarara. 1820.	7.	8.	S. 2.	Rosc. scitam. t. 65.	
	Alpinia tubulata. Bot. reg. t. 777.						
3	punicea. (<i>cr.</i>) bright red.	E. Indies. —	—	—	S. 2.		
	Alpinia punicea. F.I.						
4	Cardamomum medium. (<i>re.</i>) 9-winged.	—	—	S. 2.		
5	linguiformis. F.I. (<i>re.</i>) tongue-lipped. 1825.	—	—	S. 2.		
ELETT'ARIA. L.T.	CARDAMON. Monandria Monogynia.						
	Cardamomum. L.T. (<i>w.pu.</i>) true.	E. Indies. 1815.	—	—	S. 2.	Linn. trans. 10. t. 4. 5.	
	Alpinia Cardamomum. F.I. Amomum Cardamomum. L.						
HEDY'CHIUM. F.I.	GARLAND-FLOWER. Monandria Monogynia. L.						
1	angustifolium. R.Sc. (<i>sc.</i>) narrow-leaved.	E. Indies. 1815.	5.	12.	S. 2.	Rosc. scitam. t. 60.	
2	aurantiacum. R.Sc. (<i>or.</i>) orange-coloured.	—	—	S. 2.	— t. 61.	
	angustifolium. B.R. 157. non Roxb.						
3	coccineum. R.Sc. (<i>sc.</i>) scarlet.	Nepaul.	1818.	5.	12.	S. 2.	Rosc. scitam. t. 58.
	angustifolium. B.M. non Roxb.						
4	longifolium. R.Sc. (<i>sc.</i>) long-leaved.	E. Indies. 1819.	6.	11.	S. 2.	Rosc. scitam. t. 59.	
5	câneum. R.Sc. (<i>re.</i>) flesh-coloured.	—	—	S. 2.	Bot. mag. t. 2637.	
6	flavum. R.Sc. (<i>ye.</i>) yellow. 1818.	—	—	S. 2.	Rosc. scitam. t. 49.	
7	flavescens. R.Sc. (<i>st.</i>) yellowish. 1821.	—	—	S. 2.	— t. 50.	
8	thyrsiforme. s.s. (<i>wh.</i>) thyrs-flowered. 1819.	7.	11.	S. 2.	— t. 56.	
	heteromallum. Bot. reg. t. 767.						
9	villousum. F.I. (<i>st.</i>) villous. 1815.	6.	11.	S. 2.	Rosc. scitam. t. 54.	
10	glaucum. R.Sc. (<i>wh.</i>) glaucous. 1820.	—	—	S. 2.	— t. 53.	
11	ellipticum. R.Sc. (<i>wh.</i>) elliptic-leaved.	—	5.	8.	— t. 55.	
12	acuminatum. R.Sc. (<i>w.or.</i>) taper-pointed. 1822.	—	—	S. 2.	Bot. mag. t. 2969.	
13	elatum. B.R. (<i>bh.</i>) tall.	Nepaul. 1818.	7.	12.	S. 2.	Bot. reg. t. 526.	
14	speciosum. F.I. (<i>ye.</i>) beautiful.	E. Indies. 1823.	8.	10.	S. 2.	Rosc. scitam. t. 52.	
15	máximum. R.Sc. (<i>wh.</i>) large-flowered. 1819.	5.	9.	S. 2.	Bot. mag. t. 708.	
16	coronarium. F.I. (<i>wh.</i>) sweet-scented. 1793.	7.	9.	S. 2.	Bot. reg. t. 771.	
17	Gardnerianum. R.Sc. (<i>ye.</i>) Gardner's.	Nepaul. 1818.	—	—	S. 2.	Bot. mag. t. 2300.	
18	spicatum. B.M. (<i>bh.</i>) spiky. 1820.	8.	11.	S. 2.		
19	gracile. F.I. (<i>wh.</i>) slender.	E. Indies. 1821.	—	—	S. 2.		

ORDO CCIV.

CANNEÆ. Brown prodr. 1. p. 307.

PHRY'NIUM. R.Sc.	PHRY'NIUM.	Monandria	Monogynia.	L.		
1 dichótomum. F.I. (wh.) forked.	E. Indies.	1807.	6. 9.	S. 2.	Rumph.	amb. 4. t. 7.
2 virgátum. F.I. (wh.) slender-stem'd.	1818.	—	S. 2.			
3 spicátum. F.I. (wh.) spiked.	—	—	S. 2.			
4 imbricátum. F.I. (l.) imbricated.	1824.	5. 8.	S. 2.			
5 grandiflórum. R.Sc. (st.) great-flowered.	Brazil.	1819.	6. 8.	S. 2.	Rosc.	scitam. t. 33.
6 parviflórum. F.I. (ye.) small-flowered.	E. Indies.	—	5. 9.	S. 2.	—	t. 34.
7 capitátum. F.I. (ro.) capitate.	1820.	—	S. 2.	Rheed.	mal. 11. t. 34.	
8 violáceum. R.Sc. (vi.) violet-coloured.	Brazil.	1823.	6. 8.	S. 2.	Rosc.	scitam. t. 37.
<i>Calathéa violácea.</i> Bot. reg. t. 961.						
9 comósium. R.Sc. (ye.) tufted.	Trinidad.	1812.	—	S. 2.	Rosc.	scitam. t. 35.
10 zebrinum. R.Sc. (pu.) Zebra plant.	Brazil.	1815.	4. 7.	S. 2.	—	t. 36.
<i>Maránta zebrina.</i> Bot. reg. t. 385.	<i>Calathéa zebrina.</i> B.R.					
11 flavescens.	(ye.) yellowish.	Brazil.	1823.	6. 8.	S. 2.	Bot. reg. t. 932.
<i>Calathéa flavescens.</i> B.R.						
12 Alloýa. R.Sc. (wh.) variegated.	Guiana.	1824.	—	S. 2.	Rosc.	scitam. t. 38.
13 Myrósmá. R.Sc. (wh.) Canna-like.	Surinam.	1825.	—	S. 2.	—	t. 39.
14 cylindricum. R.Sc. (st.) cylindrical.	Brazil.	—	2. 3.	S. 2.	—	t. 40.
15 setósum. R.Sc. bristly.	—	1826.	—	S. 2.	—	t. 41.
<i>Maránta secúnda.</i> Gr.						
16 Parkéria. R.Sc. (ye.) Parker's.	Grenada.	—	—	S. 2.	Rosc.	scitam. t. 42.
17 Casúpo. R.Sc. (ye.) two-coloured.	Trinidad.	1823.	6. 8.	S. 2.	—	t. 43.
<i>Calathéa discolor.</i> s. <i>Maránta Casúpo.</i> Jacq. frag. t. 63. f. 4.						
18 ellípticum. R.Sc. (st.) elliptic-leaved.	Guiana.	1823.	—	S. 2.	Rosc.	scitam. t. 44.
19 Rossii. L.C. (ye.) Ross's.	Brazil.	1825.	4. 7.	S. 2.	—	
20 villósum. L.C. (ye.) villous.	—	—	—	S. 2.	Jacq. ic. 2. t. 20.	
21 líticum.	(wh.) yellow-bracted.	Caracas.	1809.	6. 7.	S. 2.	
<i>Maránta lítica.</i> Jacq. <i>Jacquinii.</i> R.S.						
22 obliquum. L.C. (ye.) oblique-leaved.	E. Indies.	1824.	5. 9.	S. 2.	Bot. reg. t. 1020.	
23 angustifolium. L.C. (ye.) narrow-leaved.	—	—	—	S. 2.	—	t. 1210.
24 littorálē. Led. (ye.) sea-side.	Brazil.	1827.	—	S. 2.		
25 longibracteátum. (ti.) long-bracted.	—	1824.	6. 7.	S. 2.		
26 grandifolium. (ye.) large-leaved.	Brazil.	1825.	—	S. 2.		
27 maciléntum. B.R. (wh.) meagre.	—	—	—	S. 2.		
TH'ALIA. W.	TH'ALIA.	Monandria	Monogynia.	W.		
1 dealbáta. L.T. (da.pu.) mealy.	Carolina.	1791.	7. 9.	H. w. 2.	Bot. mag. t. 1690.	
2 geniculáta. R.Sc. (bl.) jointed.	Demerara.	1824.	6. 8.	S. 2.	Rosc. scitam. t. 45.	
CANNA. W.	INDIAN SHOT.	Monandria	Monogynia.	L.		
1 indica. R.Sc. (sc.) common.	W. Indies.	1570.	1. 12.	S. 2.	Rosc. scitam. t. 1.	
2 coccinea. L.T. (sc.) scarlet.	—	1731.	—	S. 2.	—	t. 11.
3 lútea. L.T. (ye.) yellow.	S. America.	1629.	—	S. 2.	—	t. 18.
4 orientális. R.Sc. (sc.) oriental.	E. Indies.	1570.	—	S. 2.	—	t. 12.
β flava.	(ye.) yellow Indian.	—	—	S. 2.	—	t. 13.
γ maculata.	(sc.ye.) spotted-flowered.	—	—	S. 2.	Bot. mag. t. 2085.	
5 pátens. B.R. (sc.ye.) spreading.	China.	1771.	7. 10.	H. 2.	Bot. reg. t. 576.	
6 speciosa. B.M. (sc.ye.) showy.	Nepaul.	1820.	—	H. 2.	Bot. mag. t. 2317.	
7 gigantea. B.R. (or.) gigantic.	S. America.	1788.	1. 12.	S. 2.	Bot. reg. t. 206.	
8 latifolia. R.Sc. (sc.) broad-leaved.	—	—	—	S. 2.	Rosc. scitam. t. 4.	
9 édulis. B.R. (sc.) eatable.	Peru.	1820.	6. 10.	S. 2.	Bot. reg. t. 775.	
10 discolor. B.R. (sc.) crimson-leaved.	Trinidad.	1827.	8. 12.	S. 2.	—	t. 1231.
11 Lambérti. B.R. (cr.) Mr. Lambert's.	—	1818.	5. 8.	S. 2.	—	t. 470.
12 exélsa. B.C. (sc.) tall.	Brazil.	1820.	1. 12.	S. 2.	Lodd. bot. cab. t. 743.	
13 limbáta. R.Sc. (sc.ye.) bordered.	—	1818.	—	S. 2..	Rosc. scitam. t. 9.	
α uro-vittata.	Lodd. bot. cab. t. 449.					
14 pállida. R.Sc. (st.) pale.	W. Indies.	1823.	—	S. 2.	Rosc. scitam. t. 19.	
<i>Lagunén sis.</i> Bot. reg. t. 1311.						
β punctata.	(ye.) spotted-flowered.	—	—	S. 2.	Rosc. scitam. t. 20.	
15 aurantiaca. R.Sc. (or.) orange-coloured.	Brazil.	—	—	S. 2.	—	t. 21.
16 compácta. R.Sc. (cr.) compact.	S. America.	—	—	S. 2.	—	t. 22.
17 pedunculáta. R.Sc. (ye.) stalked.	Brazil.	1820.	—	S. 2.	—	t. 8.
18 denudáta. R.Sc. (sc.ye.) naked-stalked.	—	—	—	S. 2.	—	t. 23.
19 sylvestris. R.Sc. (cr.) wood.	S. America.	1813.	—	S. 2.	—	t. 10.
20 lanuginosa. R.Sc. (sc.) woolly-stem'd.	Maranham.	1823.	—	S. 2.	—	t. 16.

- | | | | | | | | | | | |
|-----------------------|-------------------|-------------------|------------------|-------------------|--------------|-------|-----|--------|----------------------|------------------------|
| 21 | cárnea. | R.Sc. | (fl.) | flesh-coloured. | Brazil. | 1822. | 1. | 12. | S. 24. | Rosc. scitam. t. 15. |
| 22 | lanceoláta. | L.C. | (sc.) | spear-leaved. | _____ | 1820. | — | — | S. 24. | |
| 23 | culénta. | L.C. | (sc.) | esculent. | S. America. | — | — | — | S. 24. | |
| 24 | polymórfha. | L.C. | (sc.) | variable. | _____ | 1824. | — | — | S. 24. | |
| | iridífora. | B.M. | (cr.) | nodding-flower'd. | Peru. | 1816. | — | — | S. 24. | Bot. reg. 609. |
| 26 | fláccida. | R.Sc. | (ye.) | flaccid. | S. Carolina. | 1788. | — | — | S. 24. | Rosc. scitam. t. 6. |
| 27 | glaúca. | L.T. | (ye.) | glaucous. | S. America. | 1732. | — | — | S. 24. | Sm. exot. bot. t. 102. |
| | β rufa. | B.M. | (ye.) | brown. | _____ | | — | — | S. 24. | Bot. mag. t. 2302. |
| MARA'NTA. R.S. | | ARROW-ROOT | | Monandria | Monogynia. | L. | | | | |
| 1 | arundinácea. | R.S. (wh.) | Indian. | S. America. | 1732. | 7. | 8. | S. 24. | Bot. mag. t. 2307. | |
| 2 | índica. | R.S. | (wh.) | smooth-leaved. | India. | | — | — | S. 24. | Sloan. jam. t. 149. |
| 3 | divaricáta. | R.Sc. (wh.) | spreading. | Brazil. | 1825. | — | — | S. 24. | Rosc. scitam. t. 27. | |
| | β purpurea. | R.Sc. (pu.) | purple-flowered. | _____ | — | — | — | S. 24. | — t. 28. | |
| 4 | Tónchat. | R.S. | (pa.bl.) | ovate. | E. Indies. | 1819. | — | — | S. 24. | Rumph. amb. 4. t. 7. |
| 5 | angustifólia. | B.M. (li.) | narrow-leaved. | W. Indies. | 1820. | 7. | 8. | S. 24. | Bot. mag. t. 2398. | |
| 6 | cuspidáta. | R.Sc. (ye.) | cuspidate. | Sierra Leone. | 1822. | — | — | S. 24. | Rosc. scitam. t. 31. | |
| 7 | bicolor. | B.R. | (wh.) | two-coloured. | S. America. | 1823. | 1. | 12. | S. 24. | Bot. reg. t. 786. |
| 8 | gíbba. | R.S. | (wh.) | gibbous. | _____ | — | — | S. 24. | Rosc. scitam. t. 29. | |
| 9 | obliqua. | R.S. | (re.) | oblique. | Guiana. | 1803. | 10. | 1. | S. 24. | Rudg. gui. t. 11. |
| 10 | sylvática. | L.T. | (wh.) | wood. | W. Indies. | 1799. | 8. | 9. | S. 24. | |

ORDO CCV.

MUSACEÆ. Juss. gen. 61.

M'USA. L.	PLANTAIN-TREE.	Polygamia Monocelia. L.			
1 <i>radisiaca</i> . w. (<i>br.cr.</i>)	common.	India.	1690.	10. 12.	S. $\frac{1}{2}$.
2 <i>sapiéntum</i> . w. (<i>br.cr.</i>)	Banana-tree.	W. Indies.	1729.	10. 3.	S. $\frac{1}{2}$.
3 <i>maculáta</i> . w. (<i>br.pi.</i>)	spotted-fruited.	Mauritius.	1818.	—	S. $\frac{1}{2}$.
4 <i>rosacea</i> . w. (<i>ro.</i>)	rose-coloured.	—	1805.	3. 6.	S. $\frac{1}{2}$.
5 <i>coccínea</i> . w. (<i>sc.</i>)	scarlet-flower'd.	China.	1792.	12. 3.	S. $\frac{1}{2}$.
6 <i>ornáta</i> . F.I. (<i>li.or.</i>)	handsome.	E. Indies.	1823.	—	S. $\frac{1}{2}$.
7 <i>superba</i> . F.I. (<i>br.pa.</i>)	superb.	—	—	—	S. $\frac{1}{2}$.
8 <i>gláuca</i> . F.I. (<i>ye.</i>)	glaucous.	—	1824.	—	S. $\frac{1}{2}$.
URANIA. W.	URANIA.	Hexandria Monogynia. W.			
speciösa. w. (<i>wh.</i>)	Plantain-leaved.	Madagascar.	1813.	—	S. $\frac{1}{2}$.
<i>Ravenalia madagascariensis</i> . Jacq.				Jacq. schœn. 1.	t. 93.
STRELITZIA. H.K.	STRELITZIA.	Pentandria Monogynia. H.K.			
1 <i>augústa</i> . H.K. (<i>wh.</i>)	Plantain-like.	C. B. S.	1791.	2. 5.	S. $\frac{1}{2}$.
2 <i>reginæ</i> . H.K. (<i>ye.bl.</i>)	Canna-leaved.	—	1773.	4. 5.	I. $\frac{1}{2}$.
3 <i>ovata</i> . H.K. (<i>ye.bl.</i>)	oval-leaved.	—	1777.	2. 4.	I. $\frac{1}{2}$.
4 <i>húmilib.</i> L.en. (<i>ye.bl.</i>)	dwarf.	—	—	—	I. $\frac{1}{2}$.
5 <i>farinosa</i> . H.K. (<i>ye.bl.</i>)	mealy-stalked.	—	1795.	—	I. $\frac{1}{2}$.
6 <i>angustifólia</i> . H.K. (<i>ye.bl.</i>)	narrow-leaved.	—	1778.	5. 6.	I. $\frac{1}{2}$.
7 <i>parvifólia</i> . H.K. (<i>ye.bl.</i>)	small-leaved.	—	1796.	5. 7.	I. $\frac{1}{2}$.
<i>ß jáncea</i> . B.R. (<i>ye.bl.</i>)	rush-like.	—	—	—	I. $\frac{1}{2}$.
HELIC'ONIA. R.S.	HELIC'ONIA.	Pentandria Monogynia. L.			
1 <i>Bihai</i> . w. (<i>re.sn.</i>)	Plantain-leaved.	W. Indies.	1786.	4. 5.	S. $\frac{1}{2}$.
2 <i>humilis</i> , r.s. (<i>sc.</i>)	dwarf.	Caracas.	1798.	5. 6.	S. $\frac{1}{2}$.
3 <i>psittacórum</i> . r.s. (<i>ye.re.</i>)	Parrot-beaked.	W. Indies.	1797.	8. 9.	S. $\frac{1}{2}$.
4 <i>Swartziana</i> , r.s. (<i>or.</i>)	Swartz's.	Jamaica.	—	—	S. $\frac{1}{2}$.
<i>psittacórum</i> β . B.M.					
5 <i>dealbata</i> . L.C.	(<i>ye.</i>) mealy.	S. America.	1823.	—	S. $\frac{1}{2}$.
6 <i>hirsuta</i> . R.S.	(<i>ye.</i>) hairy-bracted.	—	—	—	S. $\frac{1}{2}$.
7 <i>buccinata</i> . F.I.	(<i>st.</i>) Indian.	E. Indies.	1820.	6. 8.	S. $\frac{1}{2}$.
				Rumph.amb. 5.	t. 62. f. 2.
				Trew. ehret. 3.	t. 18-20.
				—————	4. t. 21-23.
				Bot. reg. t. 706.	
				Andr. deposit. t. 47.	

ORDO CCVI.

IRIDEÆ. *Brown prodr.* 302

TIRIS. B.M.	FLOWER-DE-LUCE.	Triandria Monogynia. L.
1 susiana. R.S. (<i>br.bl.</i>)	Chalcedonian. Levant.	1596. 3. 4. H. 2.
2 livida. Trat. (<i>pu.bl.</i>)	livid.	H. 2.

IRIDEÆ.

3 florentina. r.s. (<i>wh.</i>) Florentine.	S. Europe.	1596.	5. 6. H. 2L.	Bot. mag. t. 671.
4 germánica. r.s. (<i>bl.</i>) German.	Germany.	1573.	— H. 2L.	— t. 670.
5 subbiflòra. B.M. (<i>bl.</i>) double-bearing.	Portugal.	1596.	4. 5. H. 2L.	— t. 1130.
6 páldia. r.s. (<i>pa.bl.</i>) pale Turkey.	Turkey.	—	5. 6. H. 2L.	— t. 685.
<i>nepalén sis.</i> Bot. reg. 818. non Don. prodr.				
7 longiflòra. r.s. (<i>pa.bl.</i>) long-flowered.	1824.	— H. 2L.	Swt. br. fl. gar. s.2. t. 36.
8 flavescens. r.s. (<i>st.</i>) pale yellow.	Switzerland.	1825.	— H. 2L.	Bot. mag. t. 2861.
9 sórdida. W.en. (<i>pa.ye.</i>) dull-flowered.	1818.	— H. 2L.	
<i>lutéscens.</i> Bot. mag. t. 2861. <i>nec aliorum.</i>				
10 sambucina.r.s. (<i>pu.wh.</i>) Elder-scented.	S. Europe.	1658.	6. H. 2L.	Bot. mag. t. 187.
11 squádens. r.s. (<i>pu.ye.</i>) brown-flowered.	1768.	— H. 2L.	— t. 787.
12 lúrida. r.s. (<i>da.pu.</i>) dark brown.	1758.	4. 5. H. 2L.	— t. 669. 986.
13 variegátæ. r.s. (<i>pu.ye.</i>) variegated.	Hungary.	1597.	5. 6. H. 2L.	— t. 16.
β <i>limbáta.</i> r.s. (<i>pu.ye.</i>) yellow-limbed.	N. America.	— H. 2L.	
14 neglécta. r.s. (<i>bl.wh.</i>) Horneman's.	1820.	6. H. 2L.	Bot. mag. t. 243.
15 viréscens. R.L. (<i>st.</i>) greenish-flowered.	1824.	5. 6. H. 2L.	Redout. lil. 5. t. 295.
16 Swértii. r.s. (<i>bl.wh.</i>) reflex-flowered.	1748.	— H. 2L.	— 6. t. 360.
17 plicátæ. r.s. (<i>bl.</i>) plaited-leaved.	Hungary.	1823.	— H. 2L.	Bot. mag. t. 870.
18 am'ena. R.L. (<i>pu.wh.</i>) pretty.	—	— H. 2L.	Redout. lil. 8. t. 336.
19 biflòra. r.s. (<i>ri.pu.</i>) two-flowered.	S. Europe.	1596.	4. 9. H. 2L.	Besl. h. eyst. 8. f. 1.
20 furcátæ. r.s. (<i>bl.</i>) forked.	Tauria.	1822.	3. 4. H. 2L.	Bot. reg. t. 801.
21 crístata. r.s. (<i>bl.ye.</i>) crested.	N. America.	1756.	5. H. 2L.	Bot. mag. t. 412.
22 fimbriáta. v. (<i>bl.</i>) fringed.	China.	1792.	5. 6. F. 2L.	Vent. cels. t. 9.
<i>chinénsis.</i> Bot. mag. t. 373.				
23 dichótoma.r.s. (<i>vi.pu.</i>) forked.	Dauria.	1784.	6. 8. H. 2L.	Swt. br. fl. gar. t. 96.
24 nudicáulis. r.s. (<i>pu.wh.</i>) naked-stalked.	1748.	5. 6. H. 2L.	
<i>aphy'lla. w.</i>				
25 arenária. r.s. (<i>ye.</i>) sand.	Hungary.	1802.	— H. 2L.	Bot. reg. t. 549.
26 lutéscens. r.s. (<i>ye.</i>) dwarf yellow.	Germany.	1748.	4. 5. H. 2L.	Redout. lil. 5. t. 263?
<i>pùmila.</i> var. <i>lutea.</i> Bot. mag. t. 1209.				
27 flavíssima. r.s. (<i>ye.</i>) bright yellow.	Siberia.	1814.	5. 6. H. 2L.	Jacq. ic. 3. t. 220.
28 pùmila. r.s. (<i>bl.</i>) dwarf.	Austria.	1596.	4. 5. H. 2L.	Bot. mag. t. 9.
29 violacea. (<i>vi.</i>) violet-coloured.	S. Europe.	— H. 2L.	— t. 1261.
<i>pùmila.</i> var. <i>violacea.</i> B.M.				
30 hungárica. r.s. (<i>bl.</i>) Hungarian.	Hungary.	1815.	5. H. 2L.	Swt. br. fl. gar. t. 74.
31 ibérica. r.s. (<i>vi.</i>) Iberian.	Iberia.	1821.	5. 6. H. 2L.	
32 Pseud-A'corus. r.s. (<i>ye.</i>) yellow-water.	Britain.	6. H. 2L.	Eng. bot. v. 9. t. 578.
β <i>pálidido-fláva.</i> B.M. (<i>ye.</i>) pale-yellow.	N. America.	1812.	— H. 2L.	Bot. mag. t. 2239.
33 Monniérii. r.s. (<i>ye.</i>) Le Monnier's.	Levant.	1820.	6. 7. H. 2L.	Redout. lil. 4. t. 236.
34 taúrica. B.C. (<i>ye.</i>) Taurian.	Tauria.	1826.	5. 6. H. 2L.	Lodd. bot. cab. t. 1506.
35 fœtidissima.r.s. (<i>br.bl.</i>) Gladwyn.	Britain.	6. H. 2L.	Eng. bot. v. 9. t. 596.
β <i>variegátæ.</i> (<i>br.bl.</i>) striped-leaved.	— H. 2L.	
36 versicolor. r.s. (<i>pu.ye.</i>) various-coloured.	N. America.	1732.	5. 6. H. 2L.	Bot. mag. t. 21.
37 fúlva. B.M. (<i>co.</i>) copper-coloured.	1812.	6. 7. H. 2L.	— t. 1496.
<i>cúprea.</i> Ph. published 1814. <i>fúlva.</i> B.M. published 1812.				
38 tridentatæ. Ph. (<i>bl.</i>) three-toothed.	N. America.	1824.	7. H. 2L.	Swt. br. fl. gar. t. 274.
<i>tripétala.</i> Bot. mag. t. 2886. scarcely the same species.				
39 virginica. r.s. (<i>bl.pu.</i>) Virginian.	Virginia.	1758.	6. 7. H. 2L.	Bot. mag. t. 703.
40 grácilis. Bolt. (<i>pu.ye.</i>) Bolton's.	N. America.	1812.	— H. 2L.	
<i>Boltoniána.</i> r.s.				
41 spuriæ. B.M. (<i>bl.</i>) spurious.	Siberia.	1759.	7. H. 2L.	Bot. mag. t. 58.
42 nótha. M.B. (<i>bl.</i>) hybrid.	1780.	— H. 2L.	Bieb.cent.pl.ros.2.t.77.
<i>halóphila.</i> Bot. mag. t. 875.				
43 ochroléuca. L. (<i>st.</i>) sulphur-coloured.	Levant.	1757.	— H. 2L.	Bot. mag. t. 61.
44 halóphila. Pal. (<i>st.</i>) long-leaved.	Siberia.	1780.	7. 9. H. 2L.	— t. 1131.
45 stenógyна. R.L. (<i>st.</i>) cream-coloured.	Caucasus.	1804.	7. H. 2L.	— t. 1515.
46 desertórum. Guld. (<i>bl.st.</i>) sweet-scented.	Russia.	1811.	— H. 2L.	— t. 1514.
<i>spatulátæ.</i> W.en.				
47 heterophylæ. s.s. (<i>st.</i>) various-leaved.	—	— H. 2L.	
48 lævigátæ. F. smooth.	Dahuria.	1828.	— H. 2L.	Swt. br. fl. gar. t. 68.
49 vérna. M. (<i>bl.</i>) vernal.	N. America.	1748.	4. 5. H. 2L.	Pall. it. 3. t. B. f. 1.
50 ventricosa.r.s. (<i>pa.bl.</i>) bellied.	Dahuria.	1800.	6. H. 2L.	Bot. mag. t. 50.
51 sibirica. r.s. (<i>bl.st.</i>) flacid-leaved.	Siberia.	1596.	5. 6. H. 2L.	Murr. c. goff. 7. t. 4.
52 flexuosa. r.s. (<i>wh.</i>) flexuose.	Germany.	— H. 2L.	
<i>sibirica. β alba.</i> B.M. t. 1163. Redoute <i>liliac.</i> t. 438. <i>acuta</i> β <i>alba.</i> W.enum.				
53 acuta. W.en. (<i>bl.</i>) straight-leaved.	Siberia.	— H. 2L.	
54 triflòra. W.en. (<i>bl.</i>) three-flowered.	Italy.	1816.	— H. 2L.	
55 haematóphylæ. F. (<i>bl.</i>) red-leaved.	Siberia.	1790.	4. 5. H. 2L.	Swt. br. fl. gar. t. 118.
<i>sanguínea.</i> r.s. <i>sibirica</i> γ <i>sanguínea.</i> Bot. mag. t. 1604.				

IRIDEÆ.

lacústris. n. (<i>pa.bl.</i>) marsh.	N. America.	1825.	5. 6. H. ȝ.	
5a prismática. R.S. (<i>bl.</i>) New Jersey.	—	1812.	— H. ȝ.	Bot. mag. t. 1504.
60 gramínea. R.S. (<i>bl.pu.</i>) grass-leaved.	Austria.	1597.	6. H. ȝ.	— t. 681.
61 húmilib. M.B. (<i>bl.</i>) pigmy.	Siberia.	1804.	5. 6. H. ȝ.	— t. 1123.
62 ruthénica. R.S. (<i>bl.</i>) Russian.	Russia.	1810.	4. 5. H. ȝ.	— t. 1393.
63 biglumis. R.S. (<i>bh.wh.</i>) two-glumed.	Siberia.	1829.	— H. ȝ.	Pall. it. 3. app. t. 6. f. 1.
64 tenuifolia. w. (<i>pa.bl.</i>) slender-leaved.	Dauria.	1796.	5. H. ȝ.	— 3. t. C. f. 2.
5 nepalensis. D.P. (<i>bl.</i>) Nepal.	Nepaul.	1824.	6. 7. H. ȝ.	Swt. br. fl. gar. s.2.t.11.
Pallásii. B.M. (<i>bl.</i>) Pallas's.	Siberia.	1820.	— H. ȝ.	
β chinensis. B.M. (<i>bl.</i>) Chinese.	China.	—	— H. ȝ.	Bot. mag. t. 2331.
67 longispátha. B.M. (<i>bl.</i>) long-spathed.	Russia.	1822.	— H. ȝ.	— t. 2528.
68 brachycúspis. B.M. (<i>bl.</i>) short-pointed.	Siberia.	1819.	6. H. ȝ.	— t. 2326.
69 celestina. N. (<i>bl.</i>) sky-blue.	N. America.	1824.	— F. ȝ.	
70 ténax. B.R. (<i>pu.</i>) tough-leaved.	California.	1827.	4. 5. H. ȝ.	Bot. reg. t. 1218.
71 tuberosa. R.S. (<i>vi.</i>) snake's-head.	Levant.	1597.	3. 4. H. ȝ.	Bot. mag. t. 531.
72 reticulata. R.S. (<i>bl.</i>) netted-rooted.	Iberia.	1821.	4. 5. H.b. ȝ.	
73 Xiphium. R.S. (<i>va.</i>) small bulbous.	Spain.	1596.	6. H.b. ȝ.	Bot. mag. t. 686.
74 lusitánica. R.S. (<i>ye.</i>) Portuguese.	Portugal.	—	4. 5. H.b. ȝ.	— t. 679.
75 Xiphioides. R.S. (<i>va.</i>) great bulbous.	Spain.	1571.	6. H.b. ȝ.	— t. 687.
76 caucásica. R.S. (<i>st.</i>) Caucasian.	Caucasus.	1821.	2. 3. H.b. ȝ.	Swt. br. fl. gar. t. 255.
77 alata. Lam. (<i>bl.</i>) winged.	S. Europe.	1801. H.b. ȝ.	Desf. atl. 1. t. 6?
scorpioides. Redout. liliac. t. 211.				
78 pérsica. R.S. (<i>bl.ye.</i>) Persian.	Persia.	1629.	3. H.b. ȝ.	Bot. mag. t. 1.
DI'ETES. Sal.	Di'ETES.	Triandria Monogynia.		
1 iridoídes. (<i>wh.bl.</i>) Iris-flowered.	C. B. S.	1758.	4. 8. F. ȝ.	Bot. mag. t. 693.
<i>Mor'ea iridoídes.</i> B.M. <i>Iris moraeóides.</i> Ker.				
2 catenulata. (<i>wh.bl.</i>) chain-dotted.	Madagascar.	1826.	— F. ȝ.	Bot. reg. t. 1074.
<i>Mor'ea catenulata.</i> B.R.				
MOR'EA. B.M.	MOR'EA.	Triandria Monogynia. L.		
1 Sisyrichtium. B.M. (<i>bl.</i>) Spanish nut.	S. Europe.	1597.	5. 6. F.b.	Bot. mag. t. 1407.
2 Tenoreána. B.F.G. (<i>bl.</i>) Tenore's.	Naples.	1824.	6. 7. F.b.	Swt. br. fl. gar. t. 110.
<i>Iris fugax.</i> Ten. flor. nap. t. 4. nec aliorum.				
3 minuta. B.R. (<i>ye.</i>) small.	C. B. S.	1825.	— F.b.	
4 papilionácea. B.R. (<i>ye.</i>) Butterfly.	—	1795.	5. 6. F.b.	Bot. mag. t. 750.
5 ciliata. B.R. (<i>ye.</i>) fringed-leaved.	—	1787.	4. 6. F.b.	— t. 1061.
6 barbigerá. Sal. (<i>ro.</i>) bearded.	—	—	4. 5. F.b.	— t. 1012.
7 críspa. B.R. (<i>std.</i>) short-spathed.	—	1803.	— F.b.	— t. 1284.
8 tristis. B.R. (<i>bl.ye.</i>) dull-coloured.	—	1768.	— F.b.	— t. 577.
9 tricolor. A.R. (<i>br.re.ye.</i>) three-coloured.	—	1809.	— F.b.	Andr. deposit. t. 83.
10 lúrida. B.R. (<i>d.pu.</i>) Mr. Griffin's.	—	1817.	6. 7. F.b.	Bot. reg. t. 312.
11 ramosa. B.R. (<i>ye.</i>) branching.	—	1789.	5. 6. F.b.	Bot. mag. t. 771.
12 plumaria. B.R. (<i>bl.</i>) feathered.	—	1825.	— F.b.	
13 viscária. B.R. (<i>br.</i>) bird-limed.	—	1800.	6. F.b.	Bot. mag. t. 587.
14 bituminosa. B.R. (<i>ye.</i>) clammy.	—	1787.	4. 5. F.b.	— t. 1045.
15 polystächya. B.R. (<i>bl.</i>) many-spiked.	—	1825.	6. 7. F.b.	
16 longiflora. B.R. (<i>ye.</i>) long-flowered.	—	1801.	6. 7. F.b.	Bot. mag. t. 712.
17 setacea. B.R. (<i>tl.</i>) bristle-leaved.	—	1825.	— F.b.	Thunb. Ir. t. 1. f. 1.
18 edulis. B.M. (<i>li.bl.</i>) long-leaved.	—	1792.	5. 6. F.b.	Bot. mag. t. 613.
19 odora. P.L. (<i>wh.</i>) sweet-scented.	—	—	— F.b.	Sal. par. lond. t. 10.
20 longifolia. A.R. (<i>st.</i>) pale-yellow.	—	1808.	— F.b.	Andr. deposit. t. 45.
<i>edulis y lutescens.</i> Bot. mag. t. 1238.				
21 angusta. B.R. (<i>st.</i>) rolled-leaved.	—	1790.	— F.b.	Bot. mag. t. 1276.
MA'RICA. H.K.	MA'RICA.	Triandria Monogynia. H.K.		
1 cérula. B.C. (<i>bl.</i>) blue.	Brazil.	1810.	4. 10. S. ȝ.	Bot. reg. t. 713.
2 Sabini. H.T. (<i>bl.</i>) Sabine's.	St. Thomas.	1822.	9. 11. S. ȝ.	Hort. trans. v. 6. pl. 1.
3 Northiana. H.K. (<i>wh.br.</i>) broad-stemmed.	Brazil.	1789.	4. 8. S. ȝ.	Bot. mag. t. 654.
<i>Mor'ea Northiana.</i> Andr. rep. t. 255.				
4 paludosa. B.M. (<i>wh.</i>) marsh.	Guiana.	1792.	7. 8. S. ȝ.	Bot. mag. t. 646.
5 semiaperta. B.C. (<i>ye.</i>) half-opened.	Brazil.	1820.	5. 7. S. ȝ.	Lodd. bot. cab. t. 685.
6 húmilib. B.C. (<i>wh.ye.</i>) dwarf.	—	1823.	— S. ȝ.	— t. 1081.
7 longifolia. L.O. (<i>st.</i>) long-leaved.	Brazil.	1830.	6. 10. S. ȝ.	Lk. et Ot. abb. t. 58.
8 martinicensis. H.K. (<i>ye.</i>) Martinico.	W. Indies.	1782.	5. 7. S. ȝ.	Bot. mag. t. 416.
<i>Iris martinicensis.</i> B.M. <i>Cipura martinicensis.</i> k.s. <i>Trimèzia lúrida.</i> Sal.				
HERBÉRTIA. B.F.G. HERBÉRTIA.	Monadelphia Triandria.			
1 pulchella. B.F.G. (<i>bl.</i>) plaited-leaved.	Maldonado.	1827.	6. 8. F.b.	Swt. br. fl. gar. t. 222.
2 pusilla. (<i>st.</i>) small.	Brazil.	1830.	— F.b.	Lk. et Ot. abb. t. 59.
<i>Ferraria pusilla.</i> Lk. et Otto.				
CYPELLA. B.M.	CYPELLA.	Monadelphia Triandria. B.M.		
Herbética. B.M. (<i>or.</i>) Mr. Herbert's.	Buenos Ayres.	1823.	6. 9. F.b.	Swt. br. fl. gar. s.2.t.33.
Tigrídia Herbética. B.M. t. 2599.	<i>Mor'ea</i> Herbética. E.R. t. 949.			

VIEUSSE'UXIA. DC. VIEUSSE'UXIA. Triandria-Monadelphia Monogynia.

1 tripetaloides. DC. (bh.) three petal-like.	C. B. S.	1802.	4. 5. F.b.	Bot. mag. t. 702.
<i>Moræa tripétala</i> . B.M. <i>Iris tripétala</i> . w.				
2 pavónia. DC. (or.gr.) Peacock.		1790.	5. 6. F.b.	Bot. mag. t. 1247.
<i>Moræa pavónia</i> . B.M. <i>Iris pavónia</i> . w.				
3 villósa. s.s. (pu.bl.) villous-leaved.	C. B. S.	1789.	4. 5. F.b.	Bot. mag. t. 571.
4 tricúspis. s.s. (pu.) trident-petaled.		1776.	6. F.b.	_____ t. 696.
5 glaúcopis. DC. (wh.bl.) white-flowered.			5. 7. F.b.	Redout. lil. 1. t. 42.
<i>Iris pavónia</i> . Bot. mag. t. 168. nec aliorum.				

6 Bellendéni. (ye.) Mr. Ker's. 1803. 6. 7. F.b. Bot. mag. t. 772.

Moræa tricúspis γ λεῖα. B.M. certainly a very distinct species.

7 ténnis. R.S. (or.) slender.	C. B. S.	1807.	5. 6. F.b.	Bot. mag. t. 1047.
8 spirális. R.S. (ye.) spiral-flowered.		1825.	— F.b.	De la Roch. dis. t. 5.
9 unguiculáris. R.S. (bh.) long-clawed.		1802.	— F.b.	Bot. mag. t. 593.

HOMERIA. V. HOMERIA. Monadelphia Triandria.

1 spicáta. (ye.or.pu.) spiked.	C. B. S.	1785.	5. 6. F.b.	Bot. mag. t. 1283.
<i>Moræa spicáta</i> . B.M.				

2 collina. (pi.) hill. 1768. — F.b. Bot. mag. t. 1033.

Moræa collina. B.M.

3 ochroleúca. Sal. (ye.) straw-coloured. — — — F.b. Bot. mag. t. 1103.

Moræa collina. γ. B.M. 1103.

4 lineáta. B.F.G. (pi.) lined-leaved. — — — F.b. Swt. br. fl. gar. t. 178.

5 porrifólia. B.F.G. (sc.) leek-leaved. — — — F.b. ic.

6 élégans. (or.gr.) elegant. — — — F.b. Jacq. schœn. I. t. 2.

7 aurantiaca. R.S. (or.) flaccid-petaled. — — — F.b. Bot. mag. t. 1612.

Moræa collina. (a) *miniata minor*. B.M. 1612.

8 miniáta. B.F.G. (pi.) spot-flowered. — — — F.b. Swt. br. fl. gar. t. 152.

HEXAGLOTTIS. Vt. HEXAGLOTTIS. Monadelphia Triandria.

1 flexuosa. (ye.) flexuose.	C. B. S.	1803.	5. 6. F.b.	Bot. mag. t. 695.
<i>longifolia</i> . Vent. <i>Moræa flexuosa</i> . B.M.				

2 virgáta. (ye.) slender. — — — F.b. Jacq. icon. 2. t. 228.

PARDA'NTHUS. S.S. PARDA'NTHUS. Triandria Monogynia. S.S.

1 chinénsis. s.s. (or.) Chinese.	China.	1759.	6. 7. H. 4.	Bot. mag. t. 171.
<i>Ixia chinénsis</i> . B.M. <i>Belamcanda chinénsis</i> . DC.				

2 nepalén sis. (or.) Nepal. Nepal. 1823. — H. 4. Bot. reg. t. 229.

BOBA'R'TIA. L. BOBA'R'TIA. Triandria Monogynia. L.

1 gladiáta. K.I. (ye.) hairy-stalked.	C. B. S.	1815.	6. 8. G. 4.	Bot. reg. t. 229.
<i>Márica gladiáta</i> . B.R. <i>Moræa gladiáta</i> . w.				

2 spathacea. L. (ye.) sheathed. — — — F.b. Thunb. dis. p. 9. t. 1.

SISYRINCHIUM. L. SISYRINCHIUM. Monadelphia Triandria.

1 latifólium. H.K. (wh.) broad-leaved.	W. Indies.	1737.	6. 8. S. 4.	Bot. mag. t. 655.
<i>Márica plicáta</i> . B.M. <i>Moræa plicáta</i> . w.				

2 bermudianum. w. (bl.) Iris-leaved. Bermudas. 1732. 5. 7. G. 4. Bot. mag. t. 94.

iridióides. B.M.

3 Nuttálli. (bl.) Nuttall's. N. America. 1823. — H. 4. Bot. mag. t. 2786.

bermudianum. N. nec aliorum.

4 chilénsis. B.M. (bl.) blue Chilian. Chile. 1826. 6. 7. F. 4. Bot. reg. t. 1067.

5 graminifólium. B.R. (ye.) grass-leaved. — — — F. 4. Bot. mag. t. 464.

6 flexuósium. B.R. (ye.) flexuose. — — — F. 4. Wild. h. ber. t. 91.

7 ancéps. w. (bl.) narrow-leaved. N. America. 1693. 6. 7. H. 4. — t. 92.

8 mucronátum. M. (bl.) pointed-flowered. — — — H. 4. Bot. mag. t. 2965.

9 convolutum. w. (ye.) convolute. S. America. 1816. 5. 7. F. 4. — t. 2116.

10 tenuifólium. w. (ye.) slender-leaved. Mexico. — — — F. 4. Bot. reg. t. 646.

11 lúteum. L.en. (ye.) yellow-flowered. — — — H. 4. Bot. reg. t. 1283.

12 califórnicum. H.K. (ye.) Californian. California. 1796. 5. 2. F. 4. Sm. ic. piet. t. 9.

Márica califórnicâ. B.M.

13 pedunculátum. B.M. (ye.) long-peduncled. Chile. 1827. 6. 8. F. 4. Bot. reg. t. 2965.

14 micránthum. w. (st.) small-flowered. Peru. 1822. 5. 9. F. 4. — t. 2116.

15 iridiólium. K.S. (st.vi.) Iris-leaved. S. America. — — — F. 4. Bot. reg. t. 646.

laxum. Bot. mag. t. 2312. *Márica iridiólia* B.R.

16 odoratissímum. B.R. (std.) sweet-scented. S. America. 1828. 6. 10. F. 4. Bot. reg. t. 646.

17 striatum. H.K. (ye.std.) streaked-flowered. Mexico. 1788. — — — F. 4. Sm. ic. piet. t. 9.

Márica striata. B.M.

18 palmífolium. w. (wh.) Palm-leaved. Brazil. 1799. 1. 5. S. 4. Bot. reg. t. 2965.

ORTHOSA'NTHUS. MORNING-FLOWER. Monadelphia Triandria.

multiflorus. s.f.a. (bl.) many-flowered. N. Holland. 1825. 5. 8. G. 4. Swt. flor. aust. t. 11.

LIBERTIA. S.S. LIBERTIA. Triandria-Monadelphia Monogynia.

1 paniculáta. s.s. (wh.) paniced. N. S. W. 1823. 5. 8. F. 4. Swt. br. fl. gar. t. 64.

2 grandiflora. (wh.) large-flowered. N. Zealand. 1822. 4. 9. H. 4. Renéalmia grandiflora. S.F.A. the old genus Renéalmia is restored by Roscoe.

3 pulchella. s.s. (wh.) neat. N. S. W. 1823. 4. 9. F. 4. Renéalmia pulchella. S.F.A.

PATERS'ONIA. B.P.	PATERS'ONIA.	Monadelphia Triandria. H.K.	
1 sericea. B.P.	(pu.) silky.	N. S. W.	1803. 5. 7. F. 4.
2 lanata. B.P.	(pu.) woolly-spathed.	N. Holland.	1825. 5. 8. F. 4.
3 longifolia. B.P.	(bl.) long-leaved.	N. S. W.	1822. 5. 7. F. 4.
4 media. B.P.	(bl.) middle.	—	1824. — F. 4.
5 glabrata. B.P.	(bl.) smooth.	—	1814. — F. 4.
6 longiscapa. S.F.A.	(bl.) long-stalked.	N. Holland.	1825. 5. 8. F. 4.
7 glauca. B.P.	(bl.) glaucous.	N. S. W.	1826. — F. 4.
WITS'ENIA. H.K.	WITS'ENIA.	Triandria Monogynia.	
1 maurea. H.K.	(or.) downy-flowered.	C. B. S.	1790. 11. 1. G. ½.
2 corymbosa. H.K.	(bl.) corymbose.	—	1803. 4. 9. G. ½.
3 ramosa. s.s.	(ye.bl.) branching.	—	1809. — G. ½.
4 partita. K.I.	(bl.) parted-flowered.	—	1822. — G. ½.
ARISTEA. Ker.	ARISTEA.	Triandria Monogynia. L.	
1 cyanea. H.K.	(bl.) woolly-headed.	C. B. S.	1759. 4. 6. G. ¼.
2 pusilla. B.M.	(bl.) flat-stemmed.	—	1806. 6. 7. G. ¼.
3 spiralis. H.K.	(wh.bl.) spiral-flowered.	—	1795. 4. 5. G. ¼.
4 capitata. H.K.	(bl.) tallest.	—	1790. 7. 8. G. ¼.
major. A.R.	160. <i>Mor'ea caerulea.</i> Th. mor. t. 2. f. 2.		
melaleuca. H.K.	(bl.bl.) three-coloured.	C. B. S.	1786. 5. 6. G. ¼.
FERR'ARIA. Ker.	FERR'ARIA.	Monadelphia Triandria. H.K.	
1 angustifolia.	(st.) narrow-leaved.	C. B. S.	1825. 4. 6. F.b.
2 antherosa. B.M.	(gr.) green variegated.	—	1800. 3. 7. F.b.
Ferrariola H.K.	viridiiflora. A.R.	t. 285.	
3 uncinata. B.F.G.	(br.bl.) hook-leaved.	—	1825. — F.b.
4 undulata. w.	(br.wh.) curled.	—	1755. — F.b.
5 atrata. L.C.	(da.) dark-flowered.	—	1825. — F.b.
6 divaricata. B.F.G.	(da.) spreading-anther'd.	—	— F.b.
7 obtusifolia. B.F.G.	(pu.br.) blunt-leaved.	—	— F.b.
TIGRIDIA. J.	TIGER-FLOWER.	Monadelphia Triandria. H.K.	
1 payonia. H.K.	(sc.) red-flowered.	Mexico.	1796. 5. 9. H.b.
Ferraria pavonia. A.R.	178. <i>Tigridia.</i> B.M. 532.		
2 conchiflora. B.F.G.	(ye.) yellow-flowered.	Mexico.	1824. — H.b.
GALAXIA. Ker.	GALAXIA.	Monadelphia Triandria. H.K.	
1 ovata. w.	(ye.) oval-leaved.	C. B. S.	1799. 5. 9. F.b.
2 grandiflora. A.R.	(ye.) large-flowered.	—	— F.b.
3 mucronularis. Sal.	(ro.) mucronated.	—	— F.b.
4 versicolor. Sal.	(ro.ye.) various-coloured.	—	— F.b.
5 graminea. w.	(ye.) narrow-leaved.	—	1795. 7. 8. F.b.
PEYRO'USIA.	PEYRO'USIA.	Triandria Monogynia. H.K.	
1 corymbosa. B.M.	(bl.) level-topped.	C. B. S.	1791. 5. 6. F.b.
2 falcatia. B.M.	(bl.) sickle-leaved.	—	1825. — F.b.
3 fasciculata. B.M.	(wh.) fascicled.	—	— F.b.
Galaxia plicata. Jacq.	<i>Ixia heterophylla.</i> Vahl.	Oviæda fasciculata. s.s.	
4 fissifolia. B.M.	(ro.) leafy-spiked.	C. B. S.	1809. 8. 9. F.b.
5 bracteata. B.M.	(wh.) bracteate.	—	1825. — F.b.
6 aculeata. B.F.G.	(wh.) prickly-stemmed.	—	5. 7. F.b.
7 anceps. B.F.G.	(bl.) flat-stemmed.	—	— F.b.
8 silenoides. B.M.	(pu.) Silene-like.	—	1822. — F.b.
9 Fabrichii. B.M.	(wh.) slender-branched.	—	1825. — F.b.
ANOMATH'ECA. Ker.	ANOMATH'ECA.	Triandria Monogynia. H.K.	
junccea. H.K.	(ro.) cut-leaved.	C. B. S.	1791. 4. 5. F.b.
Lapeyrouisia junccea. B.M.	<i>Gladiolus polystachyus.</i> A.R. 66.		
BABIANA. Ker.	BABIANA.	Triandria Monogynia. S.S.	
1 nana. s.s.	(bl.re.) dwarf.	C. B. S.	1807. 4. 6. F.b.
Gladiolus nanus. A.R.			Andr. deposit. t. 137.
2 rübro-cyanea. B.M.	(bl.re.) red and blue.	—	1794. — F.b.
3 villosa. B.M.	(cr.) dark red.	—	1778. 8. F.b.
<i>Ixia punicea.</i> Jacq. ic. 2. t. 287.			
4 obtusifolia. B.M.	(bl.) blunt-leaved.	—	1825. 5. 6. F.b.
<i>Ixia villosa.</i> Jacq. non H.K.			Jacq. ic. 2. t. 284.
5 purpurea. B.M.	(d.pu.) dark purple.	—	1806. — F.b.
6 stricta. B.M.	(bl.wh.) upright.	—	1757. — F.b.
7 angustifolia.	(bl.) narrow-leaved.	—	— F.b.
stricta (a) B.M.	<i>Ixia villosa</i> var. Jacq. frag. t. 14. f. 3.		
8 tenuiflora. B.F.G.	(bl.) slender-flowered.	C. B. S.	1825. — F.b.
9 sulphurea. B.M.	(st.) pale-flowered.	—	1803. — F.b.
10 mucronata. B.M.	(pu.ye.) mucronate.	—	1825. — F.b.
11 plicata. B.M.	(pa.bl.) sweet-scented.	—	1774. — F.b.
12 disticha. B.M.	(pa.bl.) two-ranked.	—	6. 7. F.b.
Gladiolus plicatus. Jacq. ic. t. 237. nec a/iorum.			Swt. br. fl. gar. ic.
			Bot. mag. t. 1053.
			Jacq. ic. 2. t. 253.
			Bot. mag. t. 576.
			— t. 626.

13	pállida. w.h.	(pa.) pale-flowered.	Hybrid.	6. 7.	F.b.	
14	ochroleúca.	(st.) cream-colour'd.	C. B. S.	1825.	5. 7.	F.b.	
15	spathácea. B.M. (bl.wh.)	stiff-leaved.	—	1801.	—	F.b.	Bot. mag. t. 638.
16	sambucina. B.M. (da.bl.)	Elder-scented.	—	1799.	4. 5.	F.b.	— t. 1019.
17	tubátæ. (wh.re.)	long-flowered.	—	1774.	—	F.b.	— t. 680.
	Gladíolus tubátus. Jacq. ic. 2. t. 264.						
18	tubiflóra. B.M. (wh.re.)	long-tubed.	—	—	—	F.b.	Bot. mag. t. 847.
19	ríngens. B.M. (pu.)	gaping-flowered.	—	1752.	5. 6.	F.b.	Lodd. bot. cab. t. 1006.
20	Thunbérghii. B.M. (pu.)	many-spiked.	—	1774.	—	F.b.	
	<i>Anthol'ya placida. w.</i>						
ANTHOL'YZA. Ker.	ANTHOL'YZA.	Triandria Monogynia. L.					
1	æthiòpica. B.M. (sc.or.)	flag-leaved.	C. B. S.	1759.	1. 4.	F.b.	Bot. mag. t. 561.
2	præálta. DC.	(sc.or.) tall.	—	—	—	F.b.	Redout. lit. 7. t. 387.
	æthiòpica. var. β. B.M.	1172. vittígera. Sal.					
3	montána. B.C.	(pu.) mountain.	—	1824.	5. 7.	F.b.	Lodd. bot. cab. t. 1022.
4	lúcidor. B.M.	(pu.) shining.	—	1825.	—	F.b.	
ANISA'NTHUS. B.F.G.	ANISA'NTHUS.	Triandria Monogynia.					
1	spléndens. B.F.G. (sc.)	splendid.	C. B. S.	1825.	4. 6.	F.b.	Swt. br. fl. gar. ic. ined.
2	Cunónia. B.F.G. (sc.)	scarlet.	—	1756.	5. 6.	F.b.	Bot. mag. t. 343.
	<i>Anthol'ya Cundónia. B.M. Gladíolus Cundónia. H.K.</i>						
PETA'MENES. Sal.	PETA'MENES.	Triandria Monogynia.					
	quadranguláris. (ye.sc.)	four-sided.	C. B. S.	1790.	4. 5.	F.b.	Bot. mag. t. 567.
	Gladíolus quadranguláris. B.M. abbreviatis.	Andr. repos. t. 166.					
WATS'ONIA. Ker.	WATS'ONIA.	Triandria Monogynia. H.K.					
1	aletróides. B.M. (ro.)	Aletris-flower'd.	C. B. S.	1774.	5. 7.	F.b.	Bot. mag. t. 441.
	β variegáta. B.M. (ro.)	variegated.	—	—	—	F.b.	— t. 533.
2	angústa. K.I.	(sc.) narrow-flowered.	—	1825.	—	F.b.	Jacq. ic. 2. t. 231.
3	húmisis. B.M.	(re.) lake-coloured.	—	1754.	—	F.b.	Bot. mag. t. 631.
4	Meriana. B.M.	(ro.) red-flowered.	—	1750.	—	F.b.	— t. 418.
5	iridifólia. Jac.	(fl.) Iris-leaved.	—	1795.	—	F.b.	Jacq. ic. 2. t. 234.
6	fúlgida. Sal.	(sc.) scarlet.	—	—	—	F.b.	Bot. mag. t. 600.
	<i>Anthol'ya fúlgens. A.R. 192.</i>						
7	róseo-álba. B.M. (w.bh.)	two-coloured.	—	7. 8.	F.b.	Bot. mag. t. 537.
	β variegáta. (wh.ro.)	variegated-flower'd.	—	—	F.b.	— t. 1193.
8	brevifólia. B.M.	(sc.) short-leaved.	—	1794.	5. 7.	F.b.	— t. 601.
9	rósea. B.M.	(ro.) pyramidal.	—	1803.	7. 8.	F.b.	— t. 1072.
10	strictiflóra. B.M.	(ro.) straight-flower'd.	—	1810.	6. 7.	F.b.	— t. 1406.
11	marginata. B.M.	(ro.) broad-leaved.	—	1774.	—	F.b.	— t. 608.
	β mìnór. B.M.	(ro.) shining-leaved.	—	1812.	—	F.b.	— t. 1530.
12	rùbens. K.I.	(re.) red-spathed.	—	1825.	—	F.b.	
13	punctátæ. H.K.	(bl.) dotted-flowered.	—	1800.	4. 5.	F.b.	Andr. repos. t. 177.
	<i>Ixia punctátæ. A.R.</i>						
14	plantagínea. B.M. (bl.)	Fox-tail.	—	1774.	6. 7.	F.b.	Bot. mag. t. 553.
15	spicátæ. B.M.	(bh.) hollow-leaved.	—	1791.	5.	F.b.	— t. 523.
	<i>Ixia fistulosa. B.M.</i>						
GLADÍOLUS. Ker.	CORN-FLAG.	Triandria Monogynia. L.					
1	viperátus. B.M. (gr.pu.)	perfumed.	C. B. S.	1787.	4. 5.	F.b.	Swt. br. fl. gar. t. 156.
2	alatus. B.M.	(sc.ye.) wing-flowered.	—	1795.	5. 6.	F.b.	— t. 187.
3	algoénsis. (co.pu.y.)	Algoa bay.	—	1824.	7.	F.b.	Bot. mag. t. 2608.
	alátus β algoénsis. B.M.						
4	namaquénsis. B.M. (sc.ye.)	thick-leaved.	—	1800.	5. 6.	F.b.	Bot. mag. t. 592.
5	permeabilis. R.S. (re.y.)	tall.	—	1825.	—	F.b.	De la Roch. dis. t. 2.
6	Watsónius. B.M. (sc.)	Watson's.	—	1791.	3. 5.	F.b.	Bot. mag. t. 450.
	β variegáta. (sc.ye.)	variegated-flower'd.	—	1801.	—	F.b.	— t. 569.
7	brevifólius. H.K.	(fl.) short-leaved.	—	1802.	—	F.b.	— t. 727.
8	hirútus. H.K.	(ro.) rose-coloured.	—	1795.	4. 6.	F.b.	— t. 574.
	róseus. Andr. reposit. t. 11.						
9	merianellus. K.I. (pu.ye.)	slender-flower'd.	—	—	—	F.b.	
10	púniceus. Lam.	(pu.) dark purple.	—	—	—	F.b.	Breyne cent. 24.t.12.f.1.
	villoso. K.I. Lamárekii. R.S.						
11	aphyllus. K.I.	(fl.) leafless.	—	—	—	F.b.	Bot. mag. t. 992.
12	versicolor. B.M. (re.ye.)	various-colour'd.	—	1794.	5. 6.	F.b.	Andr. reposit. t. 19.
13	lævis. K.I.	(br.re.st.) two-nerved.	—	1806.	—	F.b.	Bot. mag. t. 1042. ε.
	versicolor ε. binervis. B.M.						
14	Breyniánus. K.I. (br.y.)	Breynius's.	—	—	—	F.b.	Breyne afr. 2. t. 7. f. 1.
15	suavéolens. K.I. (br.y.)	sweet-scented.	—	1799.	—	F.b.	Bot. mag. t. 556.
16	elongátus. K.I. (bl.ye.)	long-leaved.	—	—	—	F.b.	Jacq. ic. 2. t. 244.
17	recurvus. B.M. (pu.ye.)	Violet-scented.	—	1758.	4. 5.	F.b.	Bot. mag. t. 578.
18	inflátus. K.I. (pa.pu.)	inflated.	—	1825.	—	F.b.	
19	grácilis. B.M. (pa.bl.)	slender.	—	1800.	—	F.b.	Bot. mag. t. 562.

20 débilis. B.M. (<i>wh.pu.</i>) weak.	C. B. S.	1822.	4. 5.	F.b.	Bot. mag. t. 2585.
21 trichonemifolius. B.M. (<i>ye.</i>) slender-leav'd.	1810.	5. 6.	F.b.	_____ t. 1483.	
22 cóncolor. P.L. (<i>st.</i>) pale-yellow.	1790.	—	F.b.	Sal. par. lond. t. 8.	
23 trítis. B.M. (<i>st.br.</i>) square-leaved.	1745.	—	F.b.	Bot. mag. t. 272.	
24 hyalinus. K.I. (<i>wh.st.pu.</i>) transparent.	1825.	—	F.b.	Jacq. ic. 2. t. 242.	
25 tenellus. K.I. (<i>st.br.</i>) feeble.	—	—	F.b.	_____ t. 248.	
26 Colvillii. B.F.G. (<i>re.ye.</i>) Colvill's.	Hybrid.	1823.	—	F.b.	Swt. br. fl. gar. t. 155.
27 cárneus. B.M. (<i>pa.pu.</i>) flesh-coloured.	C. B. S.	1796.	—	F.b.	Bot. mag. t. 591.
28 cuspidátus. B.M. (<i>w.pu.</i>) sharp-pointed.	—	1795.	4. 5.	F.b.	_____ t. 582.
29 angústus. B.M. (<i>st.re.</i>) narrow-leaved. <i>hastátus</i> . Thunb. non B.M.	—	1757.	—	F.b.	_____ t. 602.
30 vomérulus. K.I. (<i>bh.vi.</i>) spade.	—	1816.	—	F.b.	Bot. mag. t. 1564.
31 involútus. K.I. (<i>ro.pu.</i>) involute.	—	1757.	5. 6.	F.b.	Mill. ic. t. 286. f. 1.
32 èdulis. B.R. (<i>wh.pu.ye.</i>) eatable-rooted.	—	—	—	F.b.	Bot. reg. t. 169.
33 undulátus. B.M. (<i>st.pu.</i>) waved-flowered.	—	1760.	4. 5.	F.b.	Bot. mag. t. 647.
34 fasciátus. R.S. (<i>pi.pu.</i>) fasciate. <i>undulátus</i> β. Bot. mag.	—	—	—	F.b.	_____ t. 538.
35 blándus. B.M. (<i>bh.ro.</i>) fairest.	—	1774.	6.	F.b.	Bot. mag. t. 625.
36 álbidus. w. (<i>wh.</i>) white-flowered.	—	—	—	F.b.	_____ t. 648.
37 campanulátus. A.R. (<i>pu.</i>) bell-flowered.	—	1794.	5. 6.	F.b.	Andr. deposit. t. 188.
38 excélsus. K.I. (<i>pi.</i>) painted-flower'd. <i>blándus</i> δ. <i>excélsus</i> . B.M.	—	—	—	F.b.	Bot. mag. t. 1665.
39 trimaculátus. K.I. (<i>bh.pu.</i>) three-spotted.	—	—	—	F.b.	Lam. ill. 1. t. 32. f. 3.
40 flexuósus. B.M. (<i>bh.</i>) flexuous.	—	1825.	—	F.b.	Thunb. diss. t. 1. f. 1.
41 floribúnus. B.M. (<i>pi.w.</i>) large-flowered.	—	1788.	5. 7.	F.b.	Bot. mag. t. 610.
42 Millérii. B.M. (<i>wh.pu.</i>) Miller's.	—	1751.	4. 5.	F.b.	_____ t. 632.
43 cardinalis. B.M. (<i>sc.</i>) superb.	—	1789.	7. 8.	F.b.	_____ t. 135.
44 bockveldiánsis. W.H. (<i>co.</i>) Bockveld.	—	1826.	—	F.b.	
45 spoforthiánsis. W.H. (<i>va.</i>) Blando-cardinalis. Hybrid.	—	—	—	F.b.	
46 mitchamiánsis. W.H. (<i>va.</i>) Tristi-hirsutus.	—	—	—	F.b.	
47 rígidus. W.H. (<i>va.</i>) Tristi-blandus.	—	—	—	F.b.	
48 propínquus. W.H. (<i>var.</i>) Floribundo-blandus.	—	—	—	F.b.	
49 frágrans. W.H. (<i>va.</i>) Recurvo-tristis.	—	—	—	F.b.	
50 Haylockiánsis. W.H. (<i>va.</i>) Versicolori-blandus.	—	—	—	F.b.	
51 Herbertiánsis. (<i>va.</i>) Tristi-Spoforthianus.	—	—	—	F.b.	
52 odorátus. W.H. (<i>va.</i>) Hirsuto-Spoforthianus.	—	—	—	F.b.	
53 delicátus. W.H. (<i>va.</i>) Recurvo-blandus.	—	—	—	F.b.	
54 byzantínus. B.M. (<i>pu.</i>) Turkish. Turkey.	1629.	6. 7.	H.b.	Bot. mag. t. 874.	
55 communis. B.M. (<i>re.</i>) common. S. Europe.	1596.	—	H.b.	_____ t. 86.	
β cárneus. B.M. (<i>bh.</i>) pale-flowered.	—	—	—	H.b.	_____ t. 1575.
SPHÆROSPORA. SPHÆROSPORA. Triandria Monogynia.					
1 imbricata. (<i>re.wh.</i>) European. S. Europe.	1596.	6. 7.	H.b.	Bot. mag. t. 719.	
<i>Gladiolus ségetum</i> . B.M. <i>Gladiolus imbricatus</i> . K.I.					
2 triphylla. (<i>re.</i>) three-leaved. Greece.	1825.	—	H.b.	Flor. græc. 1. t. 38.	
SYNNOTIA. B.F.G. SYNNOTIA. Triandria Monogynia.					
1 variegáta. B.F.G. (<i>bl.ye.</i>) variegated-flower'd. C. B. S.	1825.	4. 6.	F.b.	Swt. br. fl. gar. t. 150.	
2 bicolor. B.F.G. (<i>ye.bl.</i>) two-coloured.	1786.	3. 4.	F.b.	Bot. mag. t. 548.	
<i>Ixia bicolor</i> . B.M. Sparáxis bicolor. Ker. <i>Gladiolus bicolor</i> . w.					
3 galeáta. B.F.G. (<i>vi.ye.</i>) helmet-flower'd.	1825.	—	F.b.	Jacq. ic. 2. t. 258.	
<i>Gladiolus galeatus</i> . Jacq.					
STREPTANTH'ERA. B.F.G. STREPTANTH'ERA. Triandria Monogynia.					
1 élégans. B.F.G. (<i>w.v.e.y.</i>) elegant.	C. B. S.	1825.	6. 7.	F.b.	Swt. br. fl. gar. t. 209.
2 cùprea. (<i>co.v.e.</i>) copper-coloured.	—	—	—	F.b.	
SPARÁXIS. Ker. SPARÁXIS. Triandria Monogynia. H.K.					
1 tricolor. H.K. (<i>cr.v.e.y.</i>) three-coloured. C. B. S.	1789.	5.	F.b.	Bot. mag. t. 381.	
2 versicolor. (<i>cr.bk.y.</i>) various-coloured.	1811.	—	F.b.	Swt. br. fl. gar. t. 160.	
3 Griffini. (<i>pu.v.e.y.</i>) Mr. Griffin's.	—	—	—	F.b.	Bot. mag. t. 1482. f. inf.
<i>tricolor</i> γ <i>violáceo-purpurea</i> . B.M.					
4 blánda. (<i>bh.</i>) blush-flowered.	—	—	—	F.b.	Bot. mag. t. 1482. f. med.
<i>tricolor</i> . δ. <i>subróseó-albida</i> . B.M.					
5 grandiflóra. H.K. (<i>pu.</i>) large-flowered.	1758.	4. 5.	F.b.	Bot. mag. t. 541.	
<i>a purpurea</i> . (<i>pu.</i>) velvet-flowered.	—	—	—	F.b.	Andr. deposit. t. 87.
β striáta. (<i>pu.st.</i>) streaked-flower'd.	—	—	—	F.b.	Bot. mag. t. 779.
6 Liliágo. (<i>wh.bh.</i>) Lily-flowered.	—	—	—	F.b.	Redout. lil. 2. t. 109.
<i>Ixia Liliágo</i> . R.L. Sparáxis grandiflóra γ <i>Liliágo</i> .	Bot. reg. t. 258.				
7 bulbifera. H.K. (<i>ye.</i>) bulb-bearing.	C. B. S.	1758.	5. 6.	F.b.	Bot. mag. t. 545.
8 frágrans. B.M. (<i>ye.</i>) fragrant.	—	1825.	—	F.b.	Jacq. ic. 2. t. 274.
9 anemonáflora. B.M. (<i>st.</i>) Anemone-flower'd.	—	—	—	F.b.	_____ 2. t. 273.
<i>Ixia anemonáflora</i> . Jacq.					

TRITONIA. Ker.	TRITONIA. Triandria Monogynia. H.K.		
1 críspa. h.k. (<i>ro.wh.</i>) curled-leaved.	C. B. S.	1787.	4. 5. F.b.
2 anígozanthiflóra. (<i>gr.</i>) Anigozanthus-fld.	_____	1825.	6. 8. F.b.
3 víridis. B.M.	(<i>gr.</i>) green-flowered.	1788.	7. F.b.
4 rósea. h.k. (<i>wh.ro.</i>) rosy.	_____	1793.	6. 7. F.b.
5 capénsis. B.M. (<i>st.pu.</i>) Cape.	_____	1811.	8. 10. F.b.
6 pálida. B.M. (<i>st.</i>) pale-flowered.	_____	1806.	— F.b.
7 longiflóra. B.M. (<i>st.re.</i>) long-flowered.	_____	1774.	6. 8. F.b.
<i>Hexaglottis longiflóra.</i> V. <i>Gladiolus longiflorus.</i> Jacq. <i>Ixia longiflóra.</i> B.M.	_____	—	
8 tenuiflóra. R.S. (<i>st.</i>) slender-flower'd.	C. B. S.	1811.	4. 6. F.b.
9 cónicolor. (<i>st.</i>) self-coloured.	_____	—	— F.b.
10 Rocheána. (<i>st.ro.</i>) bending-flowered.	_____	—	8. F.b.
11 pectinátata. B.M. (<i>bh.</i>) pectinated.	_____	1825.	5. 6. F.b.
12 striáta. B.M. (<i>st.vi.</i>) streaked.	_____	—	— F.b.
13 lineáta. B.M. (<i>st.br.</i>) pencilled.	_____	1774.	— F.b.
14 securígera. B.M. (<i>co.ye.</i>) copper-coloured.	_____	—	5. F.b.
15 fláva. B.M. (<i>ye.</i>) yellow.	_____	1780.	4. 6. F.b.
16 refrácta. B.R. (<i>ye.</i>) reflexed.	_____	1815.	5. 6. F.b.
17 squálida. B.M. (<i>pi.bh.</i>) sweet-scented.	_____	1774.	— F.b.
18 fenestráta. B.M. (<i>sn.</i>) open-flowered.	_____	1801.	— F.b.
19 purpúrea. K.I. (<i>pu.</i>) purple.	_____	1825.	— F.b.
20 crocátata. B.M. (<i>sn.</i>) Crocus-flowered.	_____	1758.	— F.b.
<i>Ixia crocátata.</i> B.M.	_____	—	
21 deóstata. B.M. (<i>sn.v.e.</i>) spotted-flower'd.	_____	1774.	— F.b.
22 miniáta. B.M. (<i>ye.</i>) late-flowered.	_____	1795.	8. F.b.
23 xanthospila. s.s. (<i>wh.</i>) yellow-spotted.	_____	1825.	5. 6. F.b.
MORPHÍXIA. K.I. MORPHÍXIA. Triandria Monogynia.			
1 lineáris. K.I. (<i>bh.</i>) slender.	C. B. S.	1796.	4. 5. F.b.
<i>Ixia lineáris.</i> R.S. <i>capillaris a gracíllima.</i> B.M. <i>Hyális grácilis.</i> Salisb.	_____	—	
2 capilláris. K.I. (<i>bh.</i>) wire-stemmed.	C. B. S.	1774.	— F.b.
<i>Hyális latifolia.</i> Salisb. <i>Ixia lácea.</i> Jacq. ic. 2. t. 281. <i>capillaris β strícta.</i>	_____	—	
3 aúlica. K.I. (<i>ro.</i>) rose-coloured.	C. B. S.	1774.	— F.b.
<i>Ixia aúlica.</i> H.K. <i>capillaris γ aúlica.</i> B.M. <i>Hyális aúlica.</i> Salisb.	_____	—	
4 incarnátata. K.I. (<i>fl.</i>) flesh-coloured.	C. B. S.	1774.	— F.b.
IXIA. Ker. IXIA. Triandria Monogynia. L.			
1 fucáta. B.M. (<i>bh.</i>) painted.	C. B. S.	6. 7. F.b.
2 aristáta. B.M. (<i>pi.</i>) salver-flowered.	_____	1800.	— F.b.
3 flexuósa. B.M. (<i>va.</i>) flexose.	_____	1757.	4. 5. F.b.
4 hybrida. B.M. (<i>wh.ro.</i>) spurious.	_____	—	— F.b.
5 leucántha. P.S. (<i>wh.</i>) white-flowered.	_____	1779.	5. 8. F.b.
6 pátens. B.M. (<i>cr.</i>) spreading-flower'd.	_____	—	4. 6. F.b.
7 crateróides. B.M. (<i>cr.</i>) crimson.	_____	1778.	— F.b.
8 monadélpha. B.M. (<i>bl.gr.</i>) monadelphous.	_____	1792.	— F.b.
9 curáta. A.R. (<i>or.</i>) shortened.	_____	—	— F.b.
monadélpha β. Bot. mag. t. 1378.	_____	—	
10 columelláris. B.M. (<i>pu.b.vé.</i>) variegated.	_____	1790.	6. 8. F.b.
11 cónica. B.M. (<i>or.vé.</i>) orange-color'd.	_____	1757.	4. 5. F.b.
12 fúscó-citrina. R.S. (<i>ye.br.</i>) yellow and brown.	_____	—	— F.b.
13 dúbia. R.S. (<i>ye.pu.</i>) doubtful.	_____	— F.b.
14 viridiflóra. R.S. (<i>gr.pu.</i>) green-flowered.	_____	1780.	5. 6. F.b.
15 maculáta. W. (<i>va.</i>) spot-flowered.	_____	—	— F.b.
16 ováta. A.R. (<i>wh.pu.</i>) ovate-spiked.	_____	—	— F.b.
17 capitáta. A.R. (<i>wh.bk.</i>) headed.	_____	—	— F.b.
18 ochroleúca. (<i>st.br.</i>) cream-coloured.	_____	1809.	— F.b.
19 erécta. H.K. (<i>wh.</i>) upright.	_____	1757.	— F.b.
α albiflóra. (<i>wh.</i>) white-flowered.	_____	—	— F.b.
β incarnátata. (<i>fl.</i>) flesh-coloured.	_____	—	— F.b.
γ lútea. (<i>ye.</i>) yellow-flowered.	_____	—	— F.b.
20 odoráta. (<i>ye.</i>) sweet-scented.	_____	—	— F.b.
21 retúsa. H.K. (<i>ro.</i>) retuse.	_____	1793.	2. 4. F.b.
<i>polystachia.</i> Bot. mag. t. 629.	_____	—	
22 scíllaris. B.M. (<i>ro.</i>) Squill-flowered.	_____	1787.	— F.b.
23 críspia. B.M. (<i>ro.</i>) curled-leaved.	_____	—	4. 5. F.b.
MELASPH'ÆRULA. Ker. MELASPH'ÆRULA. Triandria Monogynia. H.K.			
1 iridiflóra. B.F.G. (<i>wh.br.</i>) Iris-leaved.	C. B. S.	1825.	2. 4. F.b.
2 intermédia. (<i>pu.st.</i>) intermediate.	_____	1787.	4. 8. F.b.
gramínea. B.M. exct synonym.	_____	—	
3 gramínea. (<i>wh.br.</i>) grass-leaved.	_____	1825.	2. 5. F.b.
<i>Gladiolus gramineus.</i> Jacq. <i>Diásia graminifolia.</i> Red. lit. t. 163.	_____	—	
4 parviflóra. L.C. (<i>wh.</i>) small-flowered.	C. B. S.	1825.	— F.b.
			Lodd. bot. cab. t. 1444.

HESPERA'NTHA. Ker. **EVENING-FLOWER.** *Triandria Monogynia.* H.K.

1	cinnamômea. H.K. (w. br.)	curled-leaved.	C. B. S.	1787.	4. 5. F.b.	Bot. mag. t. 1054.
2	falcata. H.K. (wh. br.)	sickle-leaved.	—	—	— F.b.	— t. 566.
3	graminifolia. (wh. br.)	Grass-leaved.	—	1808.	8. 9. F.b.	— t. 1254.
	<i>pilosa</i> β <i>foliis nudis, floribus minoribus.</i> B.M.					
4	pilosâ. H.K. (wh. br.)	hairy.	C. B. S.	1811.	4. 5. F.b.	Bot. mag. t. 1475.
5	angûsta. B.M. (wh.)	narrow-leaved.	—	1825.	— F.b.	Jacq. ic. 2. t. 279.
	<i>Ixia angûsta.</i> Jacq.					
6	radiata. H.K. (wh. br.)	nodding-flowered.	—	1794.	4. 6. F.b.	Bot. mag. t. 573.
	<i>Ixia radiata.</i> Lam.					
GEISSORH'IZA.	Ker. TILE-ROOT. <i>Triandria Monogynia.</i> H.K.					
1	Rochæana. (bl. pu.)	plaid.	C. B. S.	1790.	5. F.b.	Bot. mag. t. 598.
2	monantha. (bl.)	one-flowered.	—	—	— F.b.	Houtt. syst. II. t. 78. f. 1.
3	setacea. B.M. (st. re.)	bristle-leaved.	—	1809.	6. 7. F.b.	Bot. mag. t. 1255.
4	obtusata. H.K. (st.)	pale-yellow.	—	1801.	5. F.b.	— t. 672.
5	vaginata. B.F.G. (ye.)	sheathed.	—	1825.	6. 9. F.b.	Swt. br. fl. gar. t. 138.
6	secunda. B.M. (bl.)	one-ranked.	—	1795.	5. 6. F.b.	Jacq. ic. 2. t. 277.
	<i>a cærulae.</i> (bl.) blue-flowered.		—	—	— F.b.	Bot. mag. t. 597.
7	sublutea. B.M. (st.)	yellowish.	—	1825.	— F.b.	— t. 1105.
	<i>Ixia sublutea.</i> Lam.					
8	humilis. K.I. (st. bh.)	dwarf.	—	1822.	— F.b.	
9	erecta. W.H. (st. bl.)	erect.	—	1824.	— F.b.	
10	imbricata. B.M. (st. re.)	imbricated.	—	—	— F.b.	
11	hirta. B.M. (pa. bl.)	hairy.	—	—	— F.b.	
12	ciliaris. Sal. (st.)	fringed.	—	— F.b.	
	<i>infléxa. K.I. Ixia infléxa. De la Roche.</i>					
13	excisa. B.M. (wh. pi.)	dwarf-cut-leaved.	—	1789.	4. 5. F.b.	Bot. mag. t. 584.
	<i>Ixia excisa.</i> B.M.					
SPATALA'NTHUS. B.F.G. RIBBON-FLOWER. <i>Monadelphia Triandria.</i>						
speciosus. B.F.G. (sc. y. ve.)	beautiful.	C. B. S.	1825.	8. 10. F.b.	Swt. br. fl. gar. t. 300.	
TRICHON'EMA. Ker. TRICHON'EMA. <i>Triandria Monogynia.</i> H.K.						
1	Bulbocodium. H.K. (pu.)	channel-leaved.	S. Europe.	1739.	3. 4. H.b.	Bot. mag. t. 265.
2	purpurascens. (pu.)	purple.	Naples.	1825.	— H.b.	Ten. fl. nap. v. I. t. 2.
	<i>Ixia purpurascens.</i> T.S.					
3	Colúmnæ. T.P. (st. li.)	Columna's.	—	—	— H.b.	Redout. lil. t. 88. f. A.
4	celestinum. (bl.)	blue.	N. America.	1820.	— H.b.	Bartr. it. t. 3.
	<i>Ixia celestina.</i> Ph.					
5	ròseum. B.M. (ro.)	rose-coloured.	C. B. S.	1808.	6. 7. F.b.	Bot. mag. t. 1225.
6	speciosum. B.M. (ro.)	crimson.	—	—	3. 4. F.b.	— t. 1476.
7	pùdicum. B.M. (ro.)	blush.	—	—	6. 8. F.b.	— t. 1244.
8	cruciatum. s.s. (ro.)	square-leaved.	—	1758.	5. 6. F.b.	Jacq. ic. 2. t. 290.
9	longifolium. Sal. (ro.)	small-flowered.	—	—	— F.b.	Bot. mag. t. 575.
	<i>cruciatum.</i> B.M. non Jacq.					
10	filifolium. K.I. (ye.)	slender-leaved.	—	1812.	— F.b.	Redout. lil. t. 251. f. 2.
11	recurvifolium. K.I. (st.)	recurve-flowered.	—	—	— F.b.	— t. 251. f. 1.
12	tortuosum. K.I. (ye.)	twisted-leaved.	—	1822.	— F.b.	
13	caulescens. B.M. (ye.)	caulescent.	—	1810.	6. 7. F.b.	Bot. mag. t. 1392.
14	chloroleucum. B.M. (st.)	milk-coloured.	—	1825.	5. 7. F.b.	Jacq. ic. 2. t. 270.
CR'OCUS. L. CR'OCUS. <i>Triandria Monogynia.</i> L.						
1	vérnus. E.B. (va.)	spring.	England.	2. 4. H.b.	Eng. bot. v. 5. t. 344.
	<i>β plámbeus.</i> H.T. (bl.)	lead-coloured.	—	—	— H.b.	Hort. t. v. 7. t. 11. 12. f. 10.
	<i>γ álbis mágior.</i> H.T. (wh.)	large-white.	—	—	— H.b.	— v. 7. t. 11. 12. f. 11.
	<i>δ leucorhy'nchus.</i> H.T. (w.)	white.	—	—	— H.b.	— v. 7. t. 11. 12. f. 12.
	<i>ε inflátus.</i> H.T. (bl.)	bellied.	—	—	— H.b.	— v. 7. t. 11. 12. f. 13.
	<i>ζ fucátus.</i> H.T. (li. pu.)	painted.	—	—	— H.b.	— v. 7. t. 11. 12. f. 14.
	<i>η pictus.</i> H.T. (li. pu.)	pencilled.	—	—	— H.b.	— v. 7. t. 11. 12. f. 15.
	<i>θ Andersoni.</i> H.T. (wh.)	Anderson's.	—	—	— H.b.	— v. 7. t. 11. 12. f. 16.
	<i>ι Sabini.</i> H.T. (bl.)	Sabine's.	—	—	— H.b.	— v. 7. t. 11. 12. f. 17.
	<i>κ Gloriánuſ.</i> H.T. (wh. bl.)	Queen of Portugal's.	—	—	— H.b.	— v. 7. t. 11. 12. f. 18.
	<i>λ pulchéllus.</i> H.T. (li. bl.)	pretty.	—	—	— H.b.	— v. 7. t. 11. 12. f. 19.
	<i>μ neapolítanus.</i> R.S. (std.)	Neapolitan.	Naples.	—	— H.b.	Bot. mag. t. 860.
	<i>ν obovátus.</i> R.S. (wh.)	obovate-flowered.	S. Europe.	—	— H.b.	— t. 2240.
2	striatus. L.en. (wh. vi.)	striated.	—	—	— H.b.	
3	versicolor. H.K. (wh. bl.)	party-coloured.	—	1629.	— H.b.	Bot. mag. t. 1110.
	<i>α purpùreus.</i> H.T. (pu.)	purple-flowered.	—	—	— H.b.	Hort. tr. v. 7. t. 11. 12. f. 6.
	<i>β plumósus.</i> H.T. (pu. w.)	feathered.	—	—	— H.b.	— v. 7. t. 11. 12. f. 7.
	<i>γ élégans.</i> H.T. (pu. wh.)	elegant.	—	—	— H.b.	— v. 7. t. 11. 12. f. 8.
	<i>δ urbánus.</i> H.T. (wh. li.)	homebred.	—	—	— H.b.	— v. 7. t. 11. 12. f. 9.

4	biflorus. H.K. (<i>wh.pu.</i>) Scotch.	Crimea.	1629.	2. 3. H.b.	Bot. mag. t. 845.
β	Parkinsoni. H.T. (<i>wh.bl.</i>) Parkinson's.	Italy.	1824.	— H.b.	Hort. tr. v. 7. t. 11. f. 4.
5	pusillus. R.S. (<i>wh.pu.</i>) small.	Corsica.	1828.	— H.b.	Swt. br. fl. gar. t. 106.
6	míminus. R.L. (<i>pu.wh.</i>) small.	Britain.	—	— H.b.	Redout. lili. t. 81.
7	præcox. E.B. (<i>pu.wh.</i>) early.	—	—	— H.b.	Eng. bot. t. 2645.
8	argenteus. H.T. (<i>pu.wh.</i>) silvery.	—	—	— H.b.	Hort. tr.v.7.t.11.12.f.8.
9	albiflorus. R.S. (<i>wh.</i>) white-flowered. Hungary.	—	—	2. 4. H.b.	Hop. et Hor. a. Meer. c. ic.
10	variegatus. R.S. (<i>ye.pu.</i>) variegated.	Levant.	1828.	— H.b.	M. B. cent. pl. rar. t. 1.
11	reticulatus. M.B. (<i>ye.vi.</i>) netted-bulbed.	Caucasus.	1825.	— H.b.	Bot. mag. t. 652.
12	susianus. H.K. (<i>ye.pu.</i>) Cloth of Gold.	Turkey.	1605.	— H.b.	Hort. trans. I. p.136. ic.
13	stellaris. H.T. (<i>ye.pu.</i>) starry cloth of gold. S.Europe.	1629.	—	— H.b.	Bot. mag. t. 1384.
14	sulphureus. B.M. (<i>st.</i>) pale yellow.	—	—	— H.b.	Hort. t. v.7. t.11.12. f.3.
15	lacteus. H.T. (<i>st.vi.</i>) milk-coloured.	—	—	— H.b.	Bot. mag. t. 2655.
β	penicillatus. H.T. (<i>st.vi.</i>) pencilled.	—	—	— H.b.	Eng. bot. t. 2646.
16	lagenæflorus. P.L. (<i>st.</i>) Gourd-flowered.	Greece.	—	— H.b.	Bot. mag. t. 45.
17	aureus. F.G. (<i>go.</i>) golden-flowered.	—	—	— H.b.	Bot. mag. t. 1111.
18	luteus. R.S. (<i>ye.</i>) common yellow. Levant.	1620.	—	— H.b.	Eng. bot. v. 5. t. 343.
	vénus. B.M. <i>neclitorum.</i>	—	—	— H.b.	Bot. mag. t. 1267.
19	mæsiacus. B.M. (<i>ye.</i>) great yellow.	Greece.	1629.	—	Bot. mag. t. 491.
20	sativus. R.S. (<i>vi.</i>) saffron.	England.	—	9. 10. H.b.	Eng. bot. v. 7. t. 491.
21	serotinus. R.S. (<i>vi.</i>) late-flowered.	S. Europe.	1629.	9. 11. H.b.	
22	Pallasi. M.B. (<i>vi.</i>) autumnal.	Tauria.	—	— H.b.	
	<i>autumnalis.</i> R.S.	—	—	— H.b.	
23	nudiflorus. E.B. (<i>vi.</i>) naked-flowered. England.	—	—	10. 11. H.b.	

ORDO CCVII.

HÆMODORACEÆ. Brown prodr. 299.

WACHENDO'RFIA. L. WACHENDO'RFIA. Triandria Monogynia. L.					
1	thyrsiflora. R.S. (<i>ye.</i>) tall-flowered.	C. B. S.	1759.	5. 6. G. 4.	Bot. mag. t. 1060.
2	paniculata. R.S. (<i>ye.</i>) paniced.	—	1700.	2. 5. F.b.	— t. 616.
3	Herberti. (<i>ye.</i>) Herbert's.	—	1823.	— F.b.	— t. 2610.
	<i>paniculata.</i> β. H. in B.M.	—	—	—	
4	hirsuta. R.S. (<i>ye.</i>) hairy.	—	1687.	4. 6. F.b.	Bot. mag. t. 614.
5	brevifolia. B.M. (<i>ye.</i>) short-leaved.	—	1795.	3. 4. F.b.	— t. 1166.
6	Breyniana. R.S. (<i>pu.ye.</i>) Breynius's.	—	1825.	— F.b.	Breyn. cent. f. 37.
7	graminea. s.s. (<i>ye.</i>) Grass-leaved.	—	1810.	6. G. 4.	
8	tenella. R.S. (<i>pu.</i>) slender-leaved.	—	1816.	— G. 4.	
XIPHI'DIUM. J.	XIPHI'DIUM. Triandria Monogynia. W.				
1	álbidum. s.s. (<i>wh.</i>) white.	W. Indies.	1787.	6. 8. S. 4.	Sw. fl. ind. oc. 1. t. 2.
	<i>floribundum.</i> Swartz.				
2	cœruleum. s.s. (<i>bl.</i>) blue.	—	1793.	— S. 4.	Aub. gui. 1. t. 11.
HÆMOD'ORUM. B.P. HÆMOD'ORUM. Triandria Monogynia. R.S.					
1	planifolium. B.P. (<i>st.</i>) flat-leaved.	N. S. W.	1810.	7. 11. F. 4.	Bot. mag. t. 1610.
2	teretifolium. B.P. (<i>st.</i>) round-leaved.	—	1822.	— F. 4.	
DIL'ATRIS. Ker.	DIL'ATRIS. Triandria Monogynia. R.S.				
1	corymbosa. w. (<i>pu.</i>) broad-petaled.	C. B. S.	1790.	5. 6. F. 4.	Sm. exot. bot. 1. t. 16.
	<i>umbellata.</i> L.				
2	viscosa. w. (<i>pu.</i>) clammy.	—	1795.	— F. 4.	
3	paniculata. w. (<i>pu.ye.</i>) paniced.	—	1825.	— F. 4.	
GYROTH'ECA. H.T. GYROTH'ECA. Triandria Monogynia.					
	tinctoria. (<i>re.</i>) red-rooted.	N. America.	1812.	6. 7. H. 4.	Mich. amer. 1. t. 4.
	<i>Lachnáthes tinctoria.</i> Ell. <i>Dilatris tinctoria.</i> Ph. <i>Heritiæ Gmelini.</i> M.				
LOPH'IOLA. B.M. LOPH'IOLA. Hexandria Monogynia. B.M.					
	aurea. B.M. (<i>ye.</i>) golden-flowered.	N. America.	1811.	5. 7. H. 4.	Bot. mag. t. 1596.
	<i>Conostylis americana.</i> Ph.				
LAN'ARIA. W.	LAN'ARIA. Hexandria Monogynia. W.				
	plumosa. w. (<i>pu.</i>) woolly.	C. B. S.	1787.	6. 7. G. 4.	
CONO'STYLIS. B.P. CONO'STYLIS. Hexandria Monogynia. S.S.					
1	aculeata. B.P. (<i>st.</i>) prickly-leaved.	N. Holland.	1824.	9. 2. F. 4.	Bot. mag. t. 2989.
2	serrulata. B.P. (<i>st.</i>) serrulate.	—	—	— F. 4.	
ANIGOZA'NTHOS. B.P. ANIGOZA'NTHOS. Hexandria Monogynia.					
1	flavida. B.P. (<i>br.gr.</i>) russet-green.	N. Holland.	1803.	5. 9. F. 4.	Bot. mag. t. 1151.
2	rufa. B.P. (<i>br.gr.</i>) rufous.	—	1824.	— F. 4.	Labill. voy. 1. t. 22.

ORDO CCVIII.

HYPOXIDEÆ. Brown.

HYPÖ'XIS. L.	HYPÖ'XIS.	Hexandria Monogynia. L.							
1 júncea. w.	(ye.) rushy.	Carolina.	1787.	6. 7.	H.b.				
2 hygrométrica. B.P.	(ye.) hairy-leaved.	N. Holland.	1820.	—	F.b.				
3 álba. w.	(wh.) white.	C. B. S.	1806.	—	F.b.				
4 lineáris. A.R.	(ye.) linear-leaved.	—	1792.	4. 5.	F.b.				
5 stellata. w.	(ye.) yellow starry.	—	1752.	4. 6.	F.b.				
6 elegans. P.S.	(wh.) white starry.	—	—	—	F.b.				
<i>stellata</i> β. B.M. <i>certainly a distinct species.</i>		—	—	—	—				
7 serrata. w.	(ye.) saw-leaved.	—	1788.	6. 7.	F.b.				
8 ovata. w.	(ye.) smooth-leaved.	—	1806.	2. 5.	F.b.				
9 veratrifolia. w.	(ye.) plaited-leaved.	—	1788.	6. 7.	F. 4.				
<i>plicata</i> . Jacq.		—	—	—	—				
10 carolinensis. M.	(ye.) Carolina.	Carolina.	1822.	—	F.b.				
11 decumbens. w.	(ye.) decumbent.	Jamaica.	1755.	6. 9.	S. 4.				
12 stellipilis. B.R.	(ye.) starry-haired.	C. B. S.	1821.	6. 12.	F. 4.				
13 obliqua. w.	(ye.) oblique-leaved.	—	1795.	6. 7.	F. 4.				
14 pratensis. B.P.	(ye.) field.	N. S. W.	1822.	—	F.b.				
15 erecta. s.s.	(ye.) upright.	N. America.	1752.	—	H.b.				
16 gracilis. Ot.	(ye.) slender.	S. America.	1828.	—	G.b.				
17 Sellou. Ot.	(ye.) Sello's.	Buenos Ayres.	1827.	—	F.b.				
18 sobolifera. w.	(ye.) creeping.	C. B. S.	1774.	6. 9.	F. 4.				
19 villosa. w.	(ye.) villous.	—	—	—	F. 4.				
20 obtusa. B.R.	(ye.) obtuse.	—	1816.	—	F. 4.				
21 aquatica. w.	(ye.) water.	—	—	1787.	6. 7.	F. 4.			
CURCULIGO. B.P.		CURCULIGO.	Hexandria Monogynia. S.S.						
1 plicata. B.R.	(ye.) plaited.	C. B. S.	1788.	6. 7.	F.b.				
<i>β glabra</i> .		(ye.) smooth-leaved.	—	1816.	—	F.b.			
2 orchioïdes. s.s.	(ye.) narrow-leaved.	E. Indies.	1800.	—	S. 4.				
3 brevifolia. s.s.	(ye.) short-leaved.	—	1804.	5. 7.	S. 4.				
4 latifolia. s.s.	(ye.) broad-leaved.	—	—	5. 8.	S. 4.				
5 sumatrana. B.C.	(ye.) Sumatra.	Sumatra.	1818.	—	S. 4.				
6 recurvata. s.s.	(ye.) recurved.	Bengal.	1805.	—	S. 4.				
BARBAC'ENIA. Vand.		BARBAC'ENIA.	Hexandria Monogynia.						
purpurea. B.M.		(pu.) purple-flower'd. Brazil.	1825.	8. 9.	S. ½.				
VELL'OSIA. Vand.		VELL'OSIA.	Polyandria Monogynia.						
squamata. P.B.	(bl.) scaly.	Brazil.	1829.	S. ½.				

ORDO CCIX.

AMARYLLIDEÆ. Brown prodr. p. 296.

STERNB'RGIA. W.K. STERNBERGIA. Hexandria Monogynia.

1 colchiciflora. B.R.	(ye.) Colchicum-flower'd. Hungary.	1816.	9. 10.	H.b.	W. et K. hung. 2. t. 157.
2 Clusiâna. B.R.	(ye.) Clusius's.	Constantinople...	—	H.b.	Clus. hist. 1. t. 163.
3 lutea. B.R.	(ye.) common yellow. S. Europe.	1596.	—	H.b.	Bot. mag. t. 290.
<i>Amaryllis lutea</i> . B.M.		—	—	—	
<i>β angustifolia</i> . (ye.) narrow-leaved.		—	—	—	H.b.
ZEPHYRA'NTHES. H.A. ZEPHYRANTHES.		Hexandria Monogynia.			
1 chloroleuca. H.A.	(st.) one-leaved.	5. 8. G.b.	Ker's rev. pl. 8. f. 1.
2 Atamásco. H.A.	(wh.) Atamasco Lily. N. America.	1629.	5. 6.	H.b.	Bot. mag. t. 239.
<i>Amayllis Atamasco</i> . B.M.		—	—	—	
3 tubispâtha. H.A.	tube-sheathed. S. America.	5. 7.	S.b.	Bot. mag. t. 1586.
<i>Amayllis tubispâtha</i> . B.M.		—	—	—	
4 cándida. B.M.	(wh.) white-flowered. Peru.	1822.	7. 10.	H.b.	Bot. mag. t. 2607.
<i>Amayllis cándida</i> . Bot. reg. t. 724.		—	—	—	
5 rôsea. B.R.	(ro.) rose-coloured. Havannah.	—	9. 10.	G.b.	Bot. reg. t. 821.

AMARYLLIDÆ.

6	striata. B.M. (bh.wh.) striped.	Mexico.	1824.	6. 8.	H.b.	Bot. mag. t. 2593.
7	verecunda. B.M. (bh.) modest.	—	—	—	H.b.	t. 2583.
8	carinata. B.M. (ro.) keeled-leaved.	—	—	—	H.b.	Swt. br. fl. gar. s. 2. t. 902.
9	grandiflora. B.R. (ro.) large-flowered.	—	1825.	—	H.b.	Bot. reg.excl.fol.t.902.
HABRA'NTHUS. B.M. HABRA'NTHUS.		Hexandria Monogynia.				
1	gracilifolius. B.M. (ro.) slender-leaved.	S. America.	1823.	9. 11.	G.b.	Bot. mag. t. 2464.
2	angustus. B.M. (pu.) narrow.	BuenosAyres.	1825.	—	G.b.	— t. 2639.
3	versicolor. B.M. (bh.) changeable.	S. America.	1823.	1. 4.	G.b.	— t. 2485.
4	lorifolius. B.M. (ro.) lorate-leaved.	—	1824.	G.b.	Bot. mag. t. 2597.
5	bifidus. B.M. (ro.) two-cleft.	BuenosAyres.	1825.	6. 9.	G.b.	Swt. br. fl. gar. s. 2.t.14.
6	spathaceus. B.M. (pu.) broad-spathed.	—	—	G.b.	s. 2.
7	robusta. W.H. (li.) robust.	—	1827.	7. 9.	F.b.	Bot. mag. t. 1125.
8	Andersoni. W.H. (ye.) Anderson's.	S. America.	1829.	5. 6.	F.b.	Bot. reg. t. 1148.
α aurea.	(ye.) golden.	—	—	—	F.b.	
β cuprea.	(co.) copper-coloured.	—	—	—	F.b.	
9	advena. W.H. (std.) streaked.	—	1807.	—	F.b.	
Amaryllis advena.	Bot. reg. t. 849.					
10	intermedia. W.H. (re.) intermediate.	Brazil.	1827.	1. 3.	S.b.	Bot. mag. t. 783.
	Amaryllis intermedia. B.R.					
BELLADO'NNA.	BELLADO'NNA LILY.	Hexandria Monogynia.				
1	purpurascens. (pu.) light purple.	C. B. S.	1712.	7. 9.	H.b.	Bot. reg. t. 714.
	Amaryllis Belladonna. L.					Ker's rev. t. 8. f. 2.
2	pallida. (pa.li.) pale-flowered.	—	—	F.b.	Bot. mag. t. 1450.
3	pudica. (ro.) modest.	—	1795.	5. 7.	F.b.	Bot. mag. t. 657.
	Amaryllis pudica. K.R.					Bot. reg. t. 352.
4	blanda. (bh.) charming.	—	1754.	—	F.b.	
AMARYLLIS. L.	AMARYLLIS.	Hexandria Monogynia. L.				
1	reticulata. w.(pu.wh.) netted-veined.	Brazil.	1777.	4. 5.	S.b.	
2	striatifolia. (pu.wh.) striped-leaved.	—	1815.	—	S.b.	
3	Sweertia. W.H. (va.) Striatifolia-Johnsoni. Colvill's.	1821.	1. 12.	S.b.		
	a brevifolia.c.c.(pu.wh.) short-leaved.	Colvill's.	1820.	—	S.b.	
	β supérbia.c.c.(cr.wh.) superb-flowered.	—	—	—	S.b.	
	γ obscura. c.c.(re.wh.) clouded.	—	—	—	S.b.	
	δ recurvata.c.c.(re.bh.) recurved-flowered.	—	—	—	S.b.	
	ε Wellsiiana.c.c.(re.wh.) Mr. Wells's.	—	—	—	S.b.	
	ζ Vallèti. (pu.wh.) Vallèti's.	—	—	—	S.b.	
	η rugosa. c.c.(pu.wh.) rugged-flowered.	—	—	—	S.b.	
	θ versicolor. (re.wh.) various-coloured.	—	—	—	S.b.	
	ι amena. (re.wh.) delightful.	—	—	—	S.b.	
	κ affinis. (re.wh.) likened.	—	—	—	S.b.	
	λ undulatflora. (pu.wh.) wave-flowered.	—	—	—	S.b.	
	μ palliiflora. (li.wh.) pale-flowered.	—	—	—	S.b.	
	ν discolor. (re.wh.) two-coloured.	—	—	—	S.b.	
	ξ expansa. (sc.wh.) expanded.	—	—	—	S.b.	
	ο variegata.c.c.(re.wh.) variegated-flowered.	—	—	—	S.b.	
	π nótha. (pu.wh.) bastard.	—	—	—	S.b.	
	ρ lineata. (re.wh.) line-petaled.	—	—	—	S.b.	
	σ modesta. (wh.re.) modest.	—	—	—	S.b.	
	τ patentissima. (sc.) spreading-flower'd.	—	—	—	S.b.	
	ν salmonea. (re.) salmon-coloured.	—	—	—	S.b.	
	φ bistrata. (cr.da.) two-lined.	—	—	—	S.b.	
	ψ ligulæflora. (re.wh.) strap-flowered.	—	—	—	S.b.	
	χ Annesleyana. (re.wh.) Lord Mountnorris's.	—	—	—	S.b.	
4	Colvillii. (va.) Reticulato-Johnsoni.	—	—	—	S.b.	Miss Chapman's ic.ined.
	a atrorubens.(da.pu.w.) dark-red.	—	—	—	S.b.	
	β purpurascens.(pu.re.) purplish.	—	—	—	S.b.	
	γ costata. (re.wh.) ribbed-leaved.	—	—	—	S.b.	
	δ dioica. (re.pu.) barren-anthered.	—	—	—	S.b.	
	ε cemula. (re.wh.) red and white.	—	—	—	S.b.	
	ζ nervifolia. (pu.re.) strong-nerved.	—	—	—	S.b.	
	η phœnicea. (da.pu.) dark purple red.	—	—	—	S.b.	
	θ stenantha.(pa.pu.w.) narrow-petaled.	—	—	—	S.b.	
	ι flaccida. (re.pu.w.) flaccid-leaved.	—	—	—	S.b.	
	κ bracteata. (re.pu.) large-bracted.	—	—	—	S.b.	
	λ párula. (pu.wh.) small-flowered.	—	—	—	S.b.	
5	præclara. W.H. (va.) Johnsoni-striatifolia. Colvill's.	—	—	—	S.b.	
	α pátens. (pu.wh.) spreading-flowered.	—	—	—	S.b.	
	β Colvilliæna.(re.p.w.) Colvill's.	—	—	—	S.b.	
	γ decora. (re.pu.) neat.	—	—	—	S.b.	

6	Johnsöni.	(sc. wh.) Reginæ-vittata. Johnson.	1810.	1. 12.	G.b.	Lodd. bot. cab. t. 159.
	spectabilis. B.C.	brasiliensis. Redout. liliac. c. ic.				
a	Scaramouch. W.H.	(sc. wh.) Mr. Herbert's. Mitcham.	1811.		G.b.	
β	Mater. W.H.	(sc. wh.) dame.			G.b.	
γ	Sorélia. W.H.	(sc. wh.) sister.			G.b.	
δ	quárta. W.H.	(sc. wh.) fourth.			G.b.	
ε	péndula. W.H.	(sc. wh.) pendulous.			G.b.	
ζ	Ponceaunéta. W.H.	(sc. wh.) related.			G.b.	
η	bárbe blanche. W.H.	(sc. wh.) white-bearded. Highclere.			G.b.	
7	reginæ. w.	(sc.) Mexican Lily. S. America.	1725.	5. 7.	S.b.	Bot. mag. t. 453.
8	formosa. w.h.	(va.) Striatifolio-reginæ. Colvill's.	1820.		S.b.	
a	grandiflora. (pu. wh.)	large-flowered.			S.b.	
β	pallescens. (pa. re. w.)	pale red and white.			S.b.	
γ	rubicunda. (ve. re.)	velvety red.			S.b.	
δ	coccinea. (sc.)	dark-veined scarlet.			S.b.	
ε	ringens. (re. wh.)	gaping-flowered.			S.b.	
ζ	carnescens. (fl. wh.)	flesh-coloured.			S.b.	
η	diluta. (pa. re. wh.)	stained-flowered.			S.b.	
θ	imbüta. (re.)	stained red.			S.b.	
ι	compacta. (pa. re.)	compact.			S.b.	
9	gloriosa.	(va.) Reticulato-reginæ.			S.b.	
a	sulcata. (re. wh.)	channelled-leaved.			S.b.	
β	rubescens. (re. wh.)	pale red.			S.b.	
γ	rðseo-ábla. (ro. wh.)	rosy and white.			S.b.	
δ	accèdens. (ve. re.)	velvety.			S.b.	
10	consanguínea. (sc. wh.)	Valleti-reginæ.	1824.		S.b.	
β	sericea. (pu.)	silky-flowered.			S.b.	
11	intermédia.	(re. wh.) Reginæ-equestris.	1820.		S.b.	Bot. mag. t. 2315.
12	spathacea. B.M.	(sc.) large-spathed. Hybrid.			S.b.	t. 129.
13	vittata. w.	(sc. wh.) striped. S. America.	1769.	4. 5.	S.b.	Lindl. coll. bot. ic.
β	latifolia. L.col.	(sc. wh.) broad-leaved.	1816.		G.b.	Coll. hort. rip. ic.
14	lineata. Colla.	(sc. wh.) lined-flowered.	1820.		S.b.	
	vittata γ. Harrisónæ.	Bot. reg. t. 988.				
15	magnifica.	(sc. wh.) Striatifolio-vittata. Colvill's.	1823.		S.b.	
16	púlchra. C.C.	(sc. wh.) Reticulato-vittata.			S.b.	
17	rütilla. K.R.	(sc.) fiery. Brazil.	1815.	1. 12.	S.b.	Bot. reg. t. 23.
18	spléndens. H.A.	(sc. wh.) Rutilo-equestri-vittata. Spofforth.	1819.		F.b.	Herb. append. p.52. ic.
a	Mars. W.H.	(sc. wh.) Mars.	1822.		F.b.	
β	Vesta. W.H.	(sc. wh.) Vesta.			F.b.	
γ	Bellona. W.H.	(sc. wh.) Bellona.			F.b.	
δ	álta. w.h.	(sc. wh.) tallest.			F.b.	
ε	ignescens.	(sc.) flame-coloured. Colvill's.			F.b.	
19	sanguínea. W.H.	(cr. wh.) Rutilo-vittata. Highclere.	1819.		G.b.	
20	micans. w.h.	(re. co.) Rutilo-croatica.	1820.		S.b.	
21	angústa.	(sc. pu.) Johnsoni-rutila. Colvill's.	1822.		S.b.	
22	rígida.	(re. co. g.) Crocato-rutila.			S.b.	
23	incl'ya. w.h.	(sc. wh.) Fulgido-vittata. Highclere.	1819.		S.b.	
24	Carnarvoni. W.H.	(sc. w.) Vittato-reginæ.			G.b.	
25	árdens. W.H.	(sc. co.) Crocato-reginæ. Spofforth.	1820.		S.b.	
a	salmònea. W.H.	(pa. re.) salmon-coloured.			S.b.	
β	charbon brulant.	(pa. r.) shining-flowered.			S.b.	
26	fúlgida. K.R.	(sc.) fulgid. Brazil.	1810.		S.b.	Bot. reg. t. 226.
27	subbarbáta.	(or. sc.) slightly bearded.	1823.		S.b.	Bot. mag. t. 2475.
28	flórida. w.h.	(sc. wh.) Fulgido-rutila. Spofforth.	1820.		S.b.	
29	május. w.h.	(sc. co.) Acuminato-rutila.			S.b.	
β	fratérna.	(sc.) brother. Colvill's.	1821.		S.b.	
30	venosa. w.h.	(sc. wh.) Vittato-Johnsoni. Highclere.			G.b.	
31	invérsa. w.h.	(sc. wh.) Johnsoni-vittata. Liverpool.	1812.		G.b.	
β	Sáturm. w.h.	(sc. wh.) Saturn.			G.b.	
γ	Pluto. w.h. (sc. wh.)	Pluto. Spofforth.	1814.		G.b.	
δ	Andromache. w.h. (sc.)	Andromache.			G.b.	
32	sidérea. w.h.	(sc.) Sanguineo-subbarbata. Spofforth.	1820.		S.b.	
33	stellata. w.h.	(sc.) Subbarbato-sanguinea.			S.b.	
34	renovata. w.h.	(sc. wh.) Splendente-fulgida.			S.b.	
35	expánса. w.h.	(sc. wh.) Equestri-florida.			S.b.	
36	Beatum. w.h.	(sc. co.) Johnsoni-acuminata. Wentworth.			S.b.	
β	Beatrice. w.h. (sc. co.)	Beatrice.			S.b.	
γ	tricolor.	(re. pu. gr.) three-coloured. Colvill's.	1821.		S.b.	
δ	macrántha.	(re. co.) large-flowered.			S.b.	

AMARYLLIDÆ.

			1821.	1.	12.	S.b.
ϵ <i>crispata</i> .	(re.co.) <i>curled-edged</i> .	Colvill's.	—	—	—	S.b.
ζ <i>tricuspis</i> .	(sc.or.) <i>three-pointed</i> .	—	—	—	—	S.b.
36 <i>lugubris</i> . w.h. (sc.or.)	<i>Styloso-Johnsoni</i> . Spofforth.	1814.	—	—	—	S.b.
α <i>sombrella</i> . w.h. (co.)	<i>sombre</i> .	—	—	—	—	S.b.
β <i>griselina</i> . w.h. (pa.)	<i>greyish</i> .	—	—	—	—	S.b.
γ <i>porporina</i> .w.h. (br.)	<i>brownish</i> .	—	—	—	—	S.b.
37 <i>tristis</i> . w.h. (sc.co.)	<i>Johnsoni-stylosa</i> .	—	—	—	—	S.b.
38 <i>Brookesii</i> .w.h.(sc.co.)	<i>Crocato-Johnsoni</i> . Brookes's.	1820.	—	—	—	S.b.
β <i>Creusa</i> . w.h. (sc.co.)	<i>Creusa</i> .	—	—	—	—	S.b.
39 <i>aurantiaca</i> .w.h. (or.)	<i>Reginæ-acuminata</i> .Spofforth.	—	—	—	—	S.b.
α <i>Camillus</i> .w.h.(or.sc.)	<i>Camillus</i> .	—	—	—	—	S.b.
β <i>Gustavus</i> .w.h.(or.sc.)	<i>Gustavus</i> .	—	—	—	—	S.b.
γ <i>amabilis</i> .	(sc.or.) <i>brilliant</i> .	Colvill's.	1821.	—	—	S.b.
δ <i>fulta</i> .	(co.) <i>coppery</i> .	—	—	—	—	S.b.
40 <i>platypétala</i> .	(sc.or.) <i>Acuminato-reginæ</i> .	—	—	—	—	S.b.
41 <i>Höddii</i> .	(sc.wh.) <i>Reginæ-fulgida</i> .	—	1820.	—	—	S.b.
42 <i>intermixta</i> .c.c.(re.wh.)	<i>Equestri-reginæ</i> .	—	—	—	—	S.b.
43 <i>Cooperi</i> . w.h. (co.)	<i>Crocato-acuminata</i> . Wentworth.	—	—	—	—	S.b.
β <i>transpârens</i> . (co.gr.)	<i>transparent</i> .	Colvill's.	1821.	—	—	S.b.
44 <i>élégans</i> . w.h. (sc.wh.)	<i>Formoso-vittata</i> . Highclere.	—	—	—	—	G.b.
45 <i>Burghcleriana</i> .w.h. (sc.wh.)	<i>Splendente-vittata</i> . Burghclere.	—	—	—	—	G.b.
46 <i>ancéps</i> . w.h. (sc.wh.)	<i>Florida-Johnsoni</i> . Spofforth.	—	—	—	—	G.b.
47 <i>concinna</i> .w.h. (sc.wh.)	<i>Formoso-Johnsoni</i> .	—	—	—	—	G.b.
α <i>Roundhead</i> . w.h. (sc.wh.)	<i>Roundhead</i> .	—	—	—	—	G.b.
β <i>Peter</i> . w.h.(sc.wh.)	<i>Peter</i> .	—	—	—	—	G.b.
γ <i>John</i> . w.h. (sc.wh.)	<i>John</i> .	—	—	—	—	G.b.
δ <i>Jeffrey</i> .w.h.(sc.wh.)	<i>Jeffrey</i> .	—	—	—	—	G.b.
48 <i>incónstans</i> .	(sc.or.) <i>Acuminato-Johnsoni</i> .Colvill's.	—	—	—	—	G.b.
α <i>venosa</i> .	(re.co.) <i>purple-veined</i> .	—	—	—	—	G.b.
β <i>lateritia</i> .	(re.co.) <i>brick-coloured</i> .	—	—	—	—	G.b.
γ <i>glaucescens</i> .	(re.co.) <i>glaucescents</i> .	—	—	—	—	G.b.
δ <i>cóncolor</i> .	(co.) <i>self-coloured</i> .	—	—	—	—	G.b.
ϵ <i>vâria</i> .	(re.co.) <i>variable-coloured</i> .	—	—	—	—	G.b.
ζ <i>crispiflora</i> .	(sc.) <i>scarlet curl-flowered</i> .	—	—	—	—	G.b.
η <i>pustulosa</i> .	(sc.co.) <i>pustulose-flowered</i> .	—	—	—	—	G.b.
θ <i>breviflora</i> .	(re.co.) <i>short-tubed</i> .	—	—	—	—	G.b.
ι <i>fulvo-cincta</i> .	(re.co.) <i>brown-edged</i> .	—	—	—	—	G.b.
κ <i>dilécta</i> .	(sc.) <i>bright-scarlet</i> .	—	—	—	—	G.b.
λ <i>tortuliflora</i> .	(sc.or.) <i>twisted-flowered</i> .	—	—	—	—	G.b.
μ <i>veniflora</i> .	(ye.re.) <i>veined-flowered</i> .	—	—	—	—	G.b.
ν <i>lineolata</i> .	(re.) <i>dark-lined</i> .	—	—	—	—	G.b.
ξ <i>basiflora</i> .	(re.pu.) <i>purple-bottomed</i> .	—	—	—	—	G.b.
\circ <i>imperialis</i> .	(sc.) <i>imperial</i> .	—	—	—	—	G.b.
π <i>punctulosa</i> .	(pa.sc.) <i>purple spotted</i> .	—	—	—	—	G.b.
ρ <i>unipárvula</i> .	(re.pu.) <i>lower small petaled</i> .	—	—	—	—	G.b.
σ <i>purpureo-rubens</i> .	(pu.re.) <i>purplish red</i> .	—	—	—	—	G.b.
τ <i>flexuosa</i> .	(pu.re.) <i>flexuose-petaled</i> .	—	—	—	—	G.b.
ν <i>deliciata</i> .	(sc.pu.) <i>purple-striped scarlet</i> .	—	—	—	—	G.b.
49 <i>crocata</i> . K.R.	(sn.) <i>saffron-coloured</i> . Brazil.	1810.	4.	6.	S.b.	Bot. reg. t. 38.
50 <i>stylosa</i> .	(co.re.) <i>long-styleled</i> . Maranham.	1821.	2.	6.	S.b.	Bot. mag. t. 2278.
	<i>maranensis</i> . B.R. 719. <i>Hippeastrum stylosum</i> . B.M.					
51 <i>acuminata</i> . B.R. (co.) <i>acute-flowered</i> . Maranham.	—	—	4.	8.	S.b.	Bot. reg. t. 534.
	<i>Hippeastrum pulverulentum</i> . Bot. mag. t. 227.					
52 <i>spuria</i> .	(sc.co.) <i>Acuminato-reginæ</i> . Colvill's.	1822.	—	—	S.b.	
α <i>sordida</i> .	(co.) <i>dull-copper</i> .	—	—	—	S.b.	
β <i>carinata</i> .	(re.co.) <i>keeled-flowered</i> .	—	—	—	S.b.	
53 <i>striatiflora</i> .	(sc.wh.) <i>Acuminato-Sweetii</i> . β .	1823.	—	—	S.b.	
β <i>consobrina</i> .	(pi.pu.) <i>cousin</i> .	—	—	—	S.b.	
γ <i>diversicolor</i> .	(re.w.pu.) <i>different-coloured</i> .	—	—	—	S.b.	
54 <i>nobile</i> . w.h. (st.pu.)	<i>Acuminato-striatifolia</i> .Spofforth.	1820.	—	—	S.b.	
55 <i>venusta</i> .w.h.(sc.pu.w.)	<i>Fulgido-striatifolia</i> .	—	—	—	S.b.	
56 <i>dilécta</i> . w.h. (sc.pu.)	<i>Equestri-striatifolia</i> .Colvill's.	—	—	S.b.		
57 <i>Go'weni</i> . w.h.(pu.co.)	<i>Striatifolio-acuminata</i> ,Highclere.	—	—	S.b.		
α <i>Cavalier</i> .w.h.(pu.co.)	<i>Cavalier</i> .	—	—	S.b.		
β <i>Biondelta</i> .w.h.(pu.c.)	<i>Biondelta</i> .	—	—	S.b.		
γ <i>Agatha</i> .w.h.(pu.co.)	<i>Agatha</i> .	—	—	S.b.		
58 <i>Harrisoni</i> .w.h.(sc.pu.)	<i>Striatifolio-stylosa</i> . Harrison's.	1824.	—	—	S.b.	
59 <i>triplex</i> .w.h.(sc.pu.w.)	<i>Goweni-vittatum</i> . Highclere.	1820.	—	—	S.b.	

60	puschéríma. w.h. (p.co.)	Sweetii-pulverulenta. Spofforth.	1824.	4.	8.	S.b.		
α	Vezzosa. (pu.re.co.)	Vezzosa.	—	—	—	S.b.		
β	Gratian. (pu.re.co.)	Gratian.	—	—	—	S.b.		
γ	Amorosetta. (pu.re.co.)	Amorosetta.	—	—	—	S.b.		
δ	Salamander. (pu.re.co.)	changing-coloured.	—	—	—	S.b.		
ϵ	Calisto. (pu.re.co.)	Calisto.	—	—	—	S.b.		
61	bélla. w.h. (pu.re.w.)	Sweetii-Carnarvoni.	—	—	—	S.b.		
62	discolor. w.h. (sc.br.)	Johnsoni-tristis.	—	—	—	S.b.		
63	variata. w.h. (sc.pu.)	Formoso-inversum.	—	—	—	S.b.		
64	quintuplex. w.h. (pu.re.)	Harrisoni-tristis.	—	1826.	—	S.b.		
65	exímia. w.h. (va.)	Goweni-Highcleriána. Highclere.	—	—	—	S.b.		
66	equéstris. K.R.	(sc.) Barbadoes Lily. W. Indies.	1710.	7.	10.	S.b.	Bot. mag. t. 305.	
β	májor. B.R.	(sc.) larger.	—	—	—	S.b.	Bot. reg. t. 234.	
γ	plena.	(sc.) double-flowered.	—	1809.	—	S.b.		
67	miniáta. R.P.	(se.) vermillion.	S. America.	—	—	S.b.	Lindl. coll. bot. t. 11.	
68	solandriflóra. L.col.(st.)	Solandra-flowered.	—	1820.	4.	6.	S.b.	
α	chloroleúca.	(st.) dull-coloured.	—	—	—	S.b.		
β	rúbro-striáta.	(st.) red-striped.	—	—	—	S.b.	Bot. mag. t. 2573.	
γ	purpurásca. (st.pu.)	purple-tubed.	—	—	—	S.b.		
δ	striáta.	(pu.st.) purple-striped.	—	—	—	S.b.	Bot. reg. t. 876.	
69	Haylöckii. w.h. (st.pu.)	Solandrifloro-acuminata.	—	—	—	S.b.		
70	Herbérti.	(va.) Solandrifloro-stylosa. Spofforth.	—	—	—	S.b.		
α	lóngia.	(va.) long-flowered.	—	—	—	S.b.		
β	carnélla.	(fl.) flesh-coloured.	—	—	—	S.b.		
γ	magnífica.	(va.) magnificent.	—	—	—	S.b.		
71	Highcleriána. w.h. (va.)	Styloso-Solandriflora. Highclere.	—	—	—	S.b.		
β	picta.	(va.) painted lady. Spofforth.	—	—	—	S.b.		
γ	Dómínie.	(va.) Dominie.	—	—	—	S.b.		
72	calyptráta. K.R.	(gr.) green-flowered. Brazil.	1816.	5.	8.	S.b.	Bot. reg. t. 164.	
73	aúlica. K.R.	(sc.gr.) crowned.	—	—	1.	6.	S.b.	
74	psittácina. K.R.	(sc.gr.) parrot-like.	—	—	1.	12.	S.b.	
75	Griffini. w.h. (sc.gr.)	Psittacino-Johnsoni. Griffin's.	1820.	—	—	S.b.		
α	Napoleóna. w.h. (sc.gr.)	Napoleon's.	—	—	—	S.b.		
β	Victor. w.h. (sc.gr.)	Victor.	—	—	—	S.b.		
γ	Alompra. w.h. (sc.gr.)	Alompra.	—	—	—	S.b.		
δ	retinérvia.	(sc.gr.) netted-nerved.	Colvill's.	1822.	—	S.b.		
ϵ	recurvíflora.	(sc.gr.) recurve-pointed.	—	—	—	S.b.		
ζ	campanulátia. (re.gr.)	bell-flowered.	—	—	—	S.b.		
η	quadrilineáta. (sc.pu.g.)	four-lined.	—	—	—	S.b.		
θ	albocíntia. (wh.sc.)	white-edged.	—	—	—	S.b.		
ι	quadrícolor. (sc.pu.g.w.)	four-coloured.	—	—	—	S.b.		
76	affíne.	(sc.gr.) Psittacino-reginæ.	—	—	—	S.b.		
α	canaliculáta.	(sc.gr.) channel-leaved.	—	—	—	S.b.		
β	membranácea.	(re.pu.g.) membranous-edged.	—	—	—	S.b.		
γ	purpurostriáta.	(re.pu.) purple-striped.	—	—	—	S.b.		
δ	latifida.	(re.pu.g.) broad-segmented.	—	—	—	S.b.		
77	tortuosa.	(sc.gr.) Psittacino-equestris.	—	—	—	S.b.		
78	multistriáta.	(sc.wh.) Psittacino-Valleti.	—	—	—	S.b.	Miss Chapman ic.	
β	retiflora.	(sc.w.g.) netted-flowered.	—	—	—	S.b.		
79	pretiosum. w.h. (sc.gr.)	Psittacino-Griffini. Highclere.	1824.	—	—	S.b.		
80	sínistrum. w.h. (sc.w.g.)	Griffini-vittatum.	—	—	—	S.b.		
SPREIK'ELIA. H.A.	JACOBÉA LILY.	Hexandria Monogynia.						
formosissima. H.A. (cr.)	crimson.	S. America.	1658.	5.	6.	F.b.	Bot. mag. t. 47.	
<i>Amaryllis formosissima.</i> B.M.								
LYCORIS. H.A.	LYCORIS.	Hexandria Monogynia.						
1	aúrea. H.A.	(ye.) golden.	China.	1777.	8.	12.	G.b.	Bot. reg. t. 611.
	<i>Amaryllis aúrea.</i> B.R.							
2	radiáta. H.A.	(pi.) Snowdrop-leaved.	—	1758.	9.	10.	G.b.	Bot. reg. t. 596.
	<i>Amaryllis radiáta.</i> B.R.							
NER'INE. H.A.	NER'INE.	Hexandria Monogynia.						
1	curvifólia. H.A.	(sc.) curve-leaved.	C. B. S.	1794.	7.	9.	F.b.	Bot. mag. t. 725.
	<i>Amaryllis curvifolia.</i> B.M.	<i>Fothergilla.</i> Andr. deposit. t. 163.						
2	corúsca. H.A.	(sc.) brilliant.	C. B. S.	1809.	—	—	F.b.	Bot. mag. t. 1089.
3	rósea. B.M.	(ro.) rose-coloured.	—	—	—	—	F.b.	— t. 2124.
4	venústa. H.A.	(ro.) poppy-coloured.	—	1806.	6.	8.	F.b.	— t. 1090.
5	sarniénsis. H.A.	(re.) Guernsey Lily.	—	1659.	9.	10.	F.b.	— t. 294.
6	pulchéllea. H.A.	(re.) pretty.	—	1820.	7.	9.	F.b.	— t. 2407.
7	flexuósä. H.A.	(pa.re.) zigzag.	—	1795.	9.	10.	F.b.	Bot. reg. t. 172.
8	húmiliis. H.A.	(re.) small.	—	—	7.	10.	F.b.	Bot. mag. t. 726.

AMARYLLIDEÆ.

9 undulâta. H.A. (<i>pa.re.</i>) waved-flower'd. C. B. S.	1767.	7. 10. F.b.	Bot. mag. t. 369.
10 versicolor. H.A. (<i>li.</i>) changeable-col'd. Hybrid.	1815.	— F.b.	Herb. append. ic.
IXIOL'RION. H.A. IXIOL'RION. Hexandria Monogynia.			
1 montânum. r.s. (<i>bl.</i>) mountain. Persia.	1829. H.b.	Redout. liliac. t. 241.
2 tatâricum. r.s. (<i>bl.</i>) Tartarian.	Tartary.	1822. 4. 6. H.b.	Pallit. 3.ap. 85. t. D.f. 1.
<i>Amaryllis tatârica. M.B.</i>			
STRUM'ARIA. Ker. STRUM'ARIA. Hexandria Monogynia. L.			
1 truncâta. w. (<i>wh.bh.</i>) truncated. C. B. S.	1795.	4. 5. F.b.	Jacq. ic. 2. t. 357.
2 rubellâ. w. (<i>li.</i>) pale red. —	—	5. 6. F.b.	— 2. t. 358.
3 angustifolia. w. (<i>wh.li.</i>) narrow-leaved. —	—	4. 5. F.b.	— 2. t. 359.
4 linguefolia. w. (<i>wh.</i>) tongue-leaved. —	1812.	— F.b.	— 2. t. 356.
5 filifolia. B.R. (<i>wh.</i>) fine-leaved. —	1774.	10. 12. F.b.	Bot. reg. t. 440.
6 spirâlis. B.M. (<i>ro.wh.</i>) spiral-stalked. —	—	4. 8. F.b.	Bot. mag. t. 1883.
7 crîspa. B.M. (<i>ro.li.</i>) curled-flowered. —	1790.	— F.b.	— t. 1363.
<i>Amaryllis crîspa. Jacq. schoen. t. 72.</i>			
8 undulâta. w. (<i>wh.ro.</i>) waved. —	1825.	— F.b.	Jacq. ic. t. 360.
9 stellâris. B.M. (<i>pi.</i>) starry. —	1794.	10. 11. F.b.	Jacq. schoen. 1. t. 71.
10 gemmâta. B.M. (<i>wh.</i>) gynandrous. —	1810.	8. 10. F.b.	Bot. mag. t. 1620.
IMHO'FIA. H.A. IMHO'FIA. Hexandria Monogynia.			
marginâta. H.A. (<i>pu.</i>) red margined. C. B. S.	1795.	9. 10. F.b.	Jacq. schoen. 1. t. 65.
<i>Brunsvigia marginâta. H.K. Amaryllis marginâta. Jacq.</i>			
BUPH'ONE. H.A. BUPH'ONE. Hexandria Monogynia.			
1 disticha. H.A. (<i>pu.</i>) distichous. C. B. S.	1823.	9. 10. F.b.	Patters. iter. t. 1.
2 toxicâria. H.A. (<i>li.</i>) Poison-bulb. —	1774.	— F.b.	Bot. reg. t. 567.
β obtusifolia. H.A. (<i>li.</i>) wave-leaved. —	1816.	— F.b.	
3 ciliâris. H.A. (<i>pa.pu.</i>) fringe-leaved. —	1752.	6. 8. F.b.	Bot. reg. t. 1153.
<i>Brunsvigia ciliâris. B.R. Amaryllis ciliâris. L. Hæmânthus ciliâris. w.</i>			
AMMO'CHARIS. H.A. AMMO'CHARIS. Hexandria Monogynia.			
1 corântica. H.A. (<i>pu.</i>) Mr. Burchell's. C. B. S.	1816.	7. 8. F.b.	Bot. reg. t. 139.
<i>Amaryllis corântica. B.R.</i>			
β pallida. B.R. (<i>li.</i>) pale-flowered. —	—	— F.b.	Bot. reg. t. 1219.
2 falcâta. H.A. (<i>pu.</i>) sickle-leaved. —	1774.	5. 6. F.b.	Bot. mag. t. 1443.
<i>Brunsvigia falcâta. B.M. Amaryllis falcâta. H.K. Crinum falcatum. Jacq.</i>			Jacq. vind. 3. t. 60.
BRUNSV'I'GIA. Ker. BRUNSV'I'GIA. Hexandria Monogynia. H.K.			
1 laticôma. (<i>pa.pu.</i>) broad-headed. C. B. S.	1818.	3. 6. F.b.	Bot. reg. t. 497.
<i>lucida. H.A. Amaryllis laticôma. B.R.</i>			
2 minor. B.R. (<i>ro.</i>) lesser. —	1820.	— F.b.	Bot. reg. t. 954.
3 multiflora. B.M. (<i>sc.gr.</i>) many-flowered. —	1752.	6. 8. F.b.	Bot. mag. t. 1619.
<i>Amaryllis orientâlis. w.</i>			
β rubricaulis. H.A. (<i>cr.</i>) red-stemmed.	1812.	— F.b.	Jacq. schoen. 1. t. 39.
4 Josephinæ. B.R. (<i>sc.</i>) Josephine's.	1814.	— F.b.	Redout. lit. t. 370-372.
β minor. B.R. (<i>ro.</i>) smaller.	—	— F.b.	Bot. reg. t. 192, 193.
γ striâta. B.M. (<i>pu.std.</i>) striped-flowered.	1823.	— F.b.	Bot. mag. t. 2578.
5 striâta. H.K. (<i>pu.</i>) striated.	1795.	9. 10. F.b.	Jacq. schoen. 1. t. 70.
6 rádula. H.K. (<i>pu.</i>) rasp-leaved.	1790.	4. 8. F.b.	— t. 68.
PHYCE'LLA. B.R. PHYCE'LLA. Hexandria Monogynia.			
1 ignea. B.R. (<i>sc.ye.</i>) fiery. Chili.	1824.	8. 10. G.b.	Bot. reg. t. 809.
<i>beta glauca. (sc.ye.) glaucous-leaved.</i>	—	— G.b.	Bot. mag. t. 2687.
2 cyrtanthoïdes. B.R. (<i>sc.ye.</i>) Cyrtanthus-like.	1821.	6. 8. G.b.	— t. 2399.
<i>Amaryllis cyrtanthoïdes. B.M.</i>			
GRIFFI'NIA. B.R. GRIFFI'NIA. Hexandria Monogynia. B.R.			
1 hyacinthina. B.R. (<i>bl.</i>) largest.	Brazil.	1815. 6. 9. S.b.	Bot. reg. t. 163.
<i>Amaryllis hyacinthina. B.R.</i>			
2 intermédia. B.R. (<i>bl.</i>) intermediate.	—	1825. 5. 7. S.b.	Bot. reg. t. 990.
3 parviflora. B.R. (<i>bl.</i>) small flowered.	—	1820. — S.b.	— t. 511.
HÆMA'NTHUS. W. BLOOD-FLOWER. Hexandria Monogynia. L.			
1 coccineus. s.s. (<i>sc.</i>) salmon-colour'd. C. B. S.	1629.	8. 10. F.b.	Bot. mag. t. 1075.
2 coarctatus. s.s. (<i>sc.</i>) compressed.	—	1795. 2. 3. F.b.	Bot. reg. t. 181.
3 tigrinus. s.s. (<i>sc.</i>) Tiger-spotted.	—	1790. 12. 2. F.b.	Bot. mag. t. 1705.
4 rotundifolius. s.s. (<i>sc.</i>) round-leaved.	—	— 6. 10. F.b.	— t. 1618.
5 crassipes. s.s. (<i>sc.</i>) thick-leaved.	—	1816. — F.b.	
6 hyalocârpus. s.s. (<i>re.</i>) clear-berried.	—	— F.b.	Jacq. schoen. 4. t. 409.
7 quadriâlvis. s.s. (<i>re.</i>) four-valved.	—	1790. 9. 10. G.b.	Bot. mag. t. 1523.
8 humilis. s.s. (<i>re.</i>) small.	—	1825. — F.b.	
9 Pumilio. s.s. (<i>re.</i>) dwarf.	—	1789. 8. 9. F.b.	Jacq. schoen. 1. t. 61.
10 pubescens. s.s. (<i>wh.</i>) villous-leaved.	—	1774. — F.b.	Bot. reg. t. 382.
11 albiflôs. w. (<i>wh.</i>) smooth-broad-lid.	—	1791. 4. 8. F.b.	— t. 984.
12 cárneus. s.s. (<i>ro.</i>) flesh-coloured.	—	1819. 6. 8. F.b.	— t. 509.

13 lanceæfoliūs. s.s. (re.) spear-leaved.	C. B. S.	1794.	9. 10. F.b.	Jacq. schœn. 1. t. 60.
14 carinātūs. P.S. (re.) keel-leaved.	—	1759.	8. 9. F.b.	
15 amaryllōfides. s.s. (re.) Amaryllis-like.	—	1825.	— F.b.	
16 moschātūs. s.s. (re.) musk-scented.	—	1816.	— F.b.	
17 sanguíneus. s.s. (sc.) crimson.	—	1820.	6. 8. F.b.	
18 punīceus. s.s. (sc.) waved-leaved.	—	1722.	5. 9. G.b.	Bot. mag. t. 1315.
19 multiflōrus. s.s. (sc.) many-flowered.	Sierra Leon.	1783.	— S.b.	— t. 1995 & 961.
CRINUM. W.H. CRINUM. Hexandria Monogynia.				
1 pedunculātūm. B.P. (wh.) New Holland. N. S. W.	—	1790.	4. 8. G.b.	
2 exaltātūm. H.A. (wh.) high.	E. Indies.	1820.	— S.b.	Bot. reg. t. 52.
3 canaliculātūm. K.R. (wh.) channelled-leaved.	—	1810.	— S.b.	
4 brachyāndrum. K.R. (wh.) short-anthered. N. Holland.	1819.	—	G.b.	
5 brevīlimbūm. H.A. (wh.) short-limbed.	Pacific Isl.	1820.	6. 8. S.b.	
6 prōcerum. H.A. (bh.) tall.	Rangoon.	1822.	— S.b.	Bot. mag. t. 2684.
7 sīnicum. H.A. (wh.) Chinese.	China.	1819.	— S.b.	
8 anōmalum. H.A. (wh.) plaited-leaved.	—	1822.	— S.b.	
9 asiāticum. K.R. (wh.) Poison bulb.	—	1732.	— S.b.	Bot. mag. t. 1073.
10 plicātūm. B.M. (wh.) plicate-leaved.	—	1823.	4. 6. S.b.	— t. 2908.
11 declinātūm. H.A. (wh.) declined.	E. Indies.	1819.	4. 8. S.b.	— t. 2231.
12 bracteātūm. w. (wh.) short-leaved.	Mauritius.	1810.	6. 8. S.b.	Bot. reg. t. 179.
brevifolium. H.B. published 1814. bracteātūm. w. 1799.	Jacq. schœn. 4. t. 495 in 1804.			
β angustifolium. H.A. (wh.) narrow-leaved.	—	—	S.b.	
13 erubēscens. K.R. (re.wh.) blush-coloured.	S. America.	1789.	— S.b.	Bot. mag. t. 1232.
α rubrīlībum. (re.) red-limbed.	—	—	S.b.	
β major. H.A. (re.wh.) larger.	—	—	S.b.	
γ minor. H.A. (re.wh.) lesser.	—	—	S.b.	
δ corant'ynum. W.H. (bh.) Corantyne.	Berbice.	1820.	— S.b.	
ε aquāticum. W.H. (wh.) water.	Brazil.	1823.	— S.b.	
ζ glābrum. ábum. W.H. (wh.) shining-leaved.	Maranham.	1820.	— S.b.	
η glābrum rùbrum. W.H. (re.wh.) smooth red.	—	1823.	— S.b.	
θ undulātūm. W.H. (wh.) wave-leaved.	Brazil.	—	S.b.	Hook. ex. flor. t. 200.
Crinum undulātūm. Hooker.				
14 cruentūm. B.R. (re.) red-flowered.	S. America.	1810.	— S.b.	Bot. reg. t. 171.
15 sumatrānum. K.R. (wh.) Sumatra.	Sumatra.	—	S.b.	— t. 1049.
16 macrocarpōn. H.A. (wh.) large-fruited.	Pegu.	1820.	— S.b.	
17 élégans. H.A. (wh.) elegant.	—	—	9. 11. S.b.	Bot. mag. t. 2592.
18 venūstūm. H.A. pretty.	E. Indies.	—	... S.b.	
19 canalifolium. H.A. (wh.) channelled.	—	—	6. 9. S.b.	
β major. W.H. (wh.) larger.	—	1825.	— S.b.	
20 longifolium. K.R. (wh.) long-leaved.	—	1810.	— S.b.	
21 lorifolium. K.R. (wh.) lorate-leaved.	—	1819.	— S.b.	
22 americānum. W. (wh.) American.	S. America.	1752.	7. 9. S.b.	
23 Commelinī. K.R. (wh.) Commelin's.	—	1798.	6. 9. S.b.	Bot. mag. t. 1034.
24 strīctum. B.M. (wh.) upright.	S. America?	1820.	8. 10. S.b.	Jacq. schœn. 2. t. 202.
25 defīxum. K.R. (wh.) marsh.	E. Indies.	1810.	4. 8. S.b.	Bot. mag. t. 2635.
26 ensifolium. K.R. (bh.) sword-leaved.	—	1820.	— S.b.	— t. 2208.
27 am'ōnum. K.R. (wh.) delightful.	—	1810.	— S.b.	— t. 2301.
28 multiflōrum. Df. (wh.) many-flowered.	—	1822.	— S.b.	
29 verecündum. H.A. (bh.) blush.	E. Indies.	1820.	6. 9. S.b.	
30 hūmīle. B.M. (wh.) humble.	—	1816.	4. 8. S.b.	Bot. mag. t. 2636.
31 confertūm. B.M. (wh.) crowded-flowered.	N. Holland.	1822.	6. 7. G.b.	— t. 2522.
32 angustifolium. B.P. (wh.) rough-edged.	—	1823.	— G.b.	
33 arenārium. B.M. (wh.) sand.	—	1818.	5. 8. G.b.	Bot. mag. t. 2355.
β blandum. B.M. (bh.) blush-flowered.	—	1821.	— G.b.	— t. 2531.
34 purpurascens. W.H. (pu.) purplish.	Fernando Po.	1826.	— S.b.	
35 erythrophyllum. W.H. (pu.) red-leaved.	Rangoon.	1824.	— S.b.	
36 amábile. K.R. (pu.wh.) beautiful.	Sumatra.	1810.	1. 12. S.b.	Bot. mag. t. 1605.
37 augūstūm. K.R. (pu.wh.) august.	Mauritius.	1819.	4. 8. S.b.	— t. 2397.
38 mauritiānum. B.C. (wh.) Mauritius.	—	1816.	6. 9. S.b.	Lodd. bot. cab. t. 650.
39 submersum. B.M. (pu.wh.) lake.	Brazil.	1823.	8. 10. S.b.	Bot. mag. t. 2463.
40 scábrum. B.M. (cr.wh.) rough-leaved.	S. America.	1810.	1. 12. S.b.	— t. 2180.
41 zeylānicum. H.A. (cr.wh.) Ceylon.	Ceylon.	1771.	7. 8. S.b.	— t. 1171.
Amaryllis zeylānicā. B.M.				
42 latifolium. H.A. (bh.) broad-leaved.	E. Indies.	1806.	7. 9. S.b.	Bot. reg. t. 1297.
43 longistylum. W.H. (bh.) long-styled.	—	1819.	6. 10. S.b.	
44 insigne. W.H. (cr.wh.) noble.	—	—	4. 8. S.b.	Bot. reg. t. 579.
45 Careyānum. B.M. (bh.) Carey's.	Mauritius.	1820.	1. 12. S.b.	Bot. mag. t. 2166.
46 speciosum. B.M. (w.ro.) specios.	E. Indies.	1819.	6. 9. S.b.	— t. 2217.
47 moluccānum. B.M. (w.bh.) Molucca.	Moluccas.	—	— S.b.	— t. 2292.

48 giganteum. A.R. (<i>wh.</i>) gigantic. <i>petiolatum.</i> H.A. <i>Amaryllis gigantea.</i> Bot. mag. t. 92.	SierraLeone. 1792.	7. 10.	S.b.	Andr. rep. t. 169.
49 Broussoneti. H.A. (<i>wh.pu.</i>) Broussonet's.	—	—	—	S.b.
50 yuccæides. H.A. (<i>re.wh.</i>) Yucca-flowered.	1740.	6. 10.	S.b.	Bot. mag. t. 2121.
51 distichous. H.A. (<i>re.wh.</i>) distichous. <i>Amaryllis ornata a.</i> B.M.	—	—	—	Bot. mag. t. 1253.
52 Lâncei. W.H. (<i>re.wh.</i>) Lance's.	Surinam. 1825.	—	S.b.	Bot. mag. t. 2133.
53 flaccidum. B.M. (<i>wh.</i>) flaccid.	N. S. W. 1820.	7. 9.	G.b.	Bot. reg. t. 426.
<i>Amaryllis australis.</i> Bot. reg. t. 426.	—	—	—	
54 Förbesi. W.H. (<i>re.wh.</i>) Forbes's.	Delagoa Bay. 1824?	—	S.b.	Bot. reg. t. 303.
55 longiflorum. H.A. (<i>re.wh.</i>) long-flowered. W. Indies.	1816.	5. 8.	S.b.	Bot. reg. t. 661.
<i>Amaryllis longiflora</i> β <i>longiflora.</i> B.R.	—	—	—	
56 capense. H.A. (<i>wh.</i>) common Cape.	C. B. S. 1752.	7. 11.	H.b.	Bot. mag. t. 623.
<i>Amaryllis longiflora.</i> B.M.	—	—	—	
β minor. W.H. (<i>re.wh.</i>) lesser.	Highclere. 1819.	—	G.b.	Bot. reg. t. 2688.
57 riparium. H.A. (<i>cr.wh.</i>) bank.	—	1816.	6. 9.	Bot. reg. t. 2352.
58 revolutum. H.A. (<i>wh.re.</i>) revolute. <i>Amaryllis revoluta.</i> A.—B.R.	—	1774.	8. 10.	Hort. trans. ic.
59 crassifolium. H.A. (<i>wh.ro.</i>) thick-leaved. <i>Amaryllis revoluta.</i> B.—B.R.	—	—	—	
60 algoneum. W.H. (<i>re.wh.</i>) Algoa-bay.	C. B. S. 1826.	—	G.b.	
61 caffrum. W.H. (<i>re.wh.</i>) Caffre.	—	1825.	—	
62 aquaticum. H.A. (<i>ro.bh.</i>) water.	—	1816.	5. 8.	G.b.
63 Goweniænum. W.H. (<i>bh.</i>) zeylanico-capense. Highclere. 1813.	—	—	6. 9.	G.b.
64 Herberthianum. (<i>re.wh.</i>) Scabro-capense. Spofforth.	1818.	—	—	G.b.
<i>Spofforthianum.</i> Herbert MSS.	—	—	—	
65 fertile. W.H. (<i>wh.</i>) Canaliculato-capense.	—	—	—	G.b.
β <i>grandiflorum.</i> W.H. (<i>wh.</i>) larger-flowered.	—	—	—	G.b.
66 ingens. W.H. (<i>wh.</i>) Pedunculato-capense. Highclere.	—	—	—	G.b.
67 erectius. W.H. (<i>bh.</i>) Corantyno-capense. Spofforth.	—	—	—	G.b.
68 Highclerianum. W.H. (<i>wh.</i>) Asiatico-capense. Highclere	—	—	—	G.b.
69 elongatum. W.H. (<i>wh.</i>) Brevisolio-capense. Spofforth.	1819.	—	—	G.b.
70 modestum. W.H. (<i>bh.</i>) Specioso-capense.	—	—	—	G.b.
71 pudicum. W.H. (<i>bh.</i>) Careyano-capense.	—	—	—	G.b.
72 cùpimum. W.H. (<i>re.wh.</i>) Cruento-capense.	—	—	—	G.b.
73 tortuosum. W.H. (<i>wh.</i>) Defixo-capense.	—	—	—	G.b.
74 Félix. W.H. (<i>bh.</i>) Latifolio-capense.	—	1828.	—	G.b.
75 micropermum. W.H. (<i>bh.</i>) Revoluto-capense.	—	—	—	G.b.
76 ègrègium. W.H. (<i>re.wh.</i>) Pedunculato-riparium.	—	1826.	—	G.b.
77 præstans. W.H. (<i>re.wh.</i>) Canaliculato-riparium.	—	—	—	G.b.
78 vittatum. W.H. (<i>re.wh.</i>) Scabro-exaltatum.	—	1818.	—	S.b.
79 Digweedi. W.H. (<i>re.wh.</i>) Scabro-americanum. Highclere.	1826.	—	—	S.b.
80 Coopéri. W.H. (<i>re.wh.</i>) Specioso-longifolium. Wentworth	—	—	—	S.b.
81 amatum. W.H. (<i>re.wh.</i>) Zeylanico-pedunculatum. Highclere.	—	—	—	S.b.
82 Baconi. W.H. (<i>re.wh.</i>) Zeylanico-erubescens. Mr. Bacon's.	—	—	—	S.b.
83 candidum. W.H. (<i>wh.</i>) Americano-bracteatum. Spofforth.	—	—	—	S.b.
84 rubricaulé. W.H. (<i>bh.</i>) Erubescens-bracteatum.	—	—	—	S.b.
85 pùlchrum. W.H. (<i>re.wh.</i>) Scabro-erubescens. Highclere.	—	—	—	S.b.
86 bulbulosum. W.H. (<i>bh.</i>) Capensi-erubescens.	—	—	—	G.b.
87 duplex. W.H. (<i>bh.</i>) Americano-erubescens. Spofforth.	—	—	—	S.b.
β minor W.H. (<i>bh.</i>) lesser.	Highclere.	—	—	S.b.
88 flaccidulum. W.H. (<i>bh.</i>) Specioso-defixum. Spofforth.	—	—	—	S.b.
89 promissum. W.H. (<i>re.wh.</i>) Forbesi-Careyanum.	—	—	—	S.b.
90 mirabile. W.H. (<i>wh.</i>) Flaccido-canaliculatum.	—	—	—	S.b.
91 ùnicum. W.H. (<i>wh.</i>) Flaccido-bracteatum.	—	—	—	S.b.
92 triplex. W.H. (<i>wh.</i>) Careyano-fertile.	—	—	—	S.b.
93 divérsum. W.H. (<i>wh.</i>) Careyano-tortuosum.	—	—	—	S.b.
94 quâdruplex. W.H. (<i>wh.</i>) Candido-fertile.	—	—	—	S.b.
VALLOTA. H.A. VALL'OTA. Hexandria Monogynia.	—	—	—	
purpurea. H.A. (<i>re.</i>) purple-stalked. C. B. S. 1774.	5. 8.	F.b.	Jacq. schœn. 1. t. 62.	
<i>Amaryllis purpurea.</i> w. <i>elata.</i> Jacq.	—	—	—	
α major. (<i>re.</i>) larger.	—	—	—	Bot. mag. t. 1430.
β minor. (<i>re.</i>) lesser.	—	—	—	Bot. reg. t. 552.
GASTRON'EMA. H.A. GASTRON'EMA. Hexandria Monogynia.	—	—	—	Bot. mag. t. 2291.
clavatum. H.A. (<i>re.wh.</i>) clavate.	C. B. S. 1816.	5. 8.	F.b.	
<i>Cyrtanthus uniflorus.</i> Bot. reg. t. 168. <i>Amaryllis Pumilio.</i> H.K.	—	—	—	
CYRTANTHUS. W. CYRTA'NTHUS. Hexandria Monogynia. W.	—	—	—	
1 obliquus. w. (<i>or.gr.</i>) oblique-leaved. C. B. S. 1774.	5. 8.	G.b.	Bot. mag. t. 1133.	
2 striatus. B.M. (<i>ye.sc.</i>) striped.	1823.	—	G.b.	— t. 2534.

3 vittatus. s.s.	(std.) ribbon.	C. B. S.	1823.	5. 8. G.b.	
4 spiralis. B.R.	(sc.) spiral-leaved.	1790.	—	G.b.	Bot. reg. t. 167.
5 collinus. B.R.	(sc.) hill.	1816.	—	G.b.	— t. 162.
6 pallidus. B.M.	(fl.) pale-flowered.	1822.	—	G.b.	Bot. mag. t. 2471.
7 ventricosus. w.	(sc.) bellied.	1770.	—	G.b.	Jacq. schoen. I. t. 76.
8 angustifolius. w.	(sc.) narrow-leaved.	1774.	5. 6. G.b.		Bot. mag. t. 271.
9 odorus. B.R.	(sc.) sweet-scented.	1818.	5. 7. G.b.		Bot. reg. t. 503.
COBU'RGHIA. B.F.G.	Hexandria Monogynia.				
incarnata. B.F.G.	(sc.) splendid.	Quito.	1820.	8. 9. F.b.	Swt. br. fl. gar. s.2. t.17.
STENOME'SSON. H.A.	STENOME'SSON.	Hexandria Monogynia.			
1 flavum. B.M.	(ye.) slender-toothed.	Peru.	1823.	5. 11. G.b.	Bot. mag. t. 2641.
Chrysiphala flava. B.R.	778.				
2 curvidentatum. B.M.	(ye.) curve-toothed.	1824.	—	G.b.	Bot. mag. t. 2640.
3 pauciflorum.	(ye.) few-flowered.	1822.	—	G.b.	Hook. ex. flor. t. 132.
Chrysiphala pauciflora. H.E.F.					
CHLIDA'NTHUS. H.A.	CHLIDA'NTHUS.	Hexandria Monogynia.			
frâgrans. H.A.	(ye.) yellow fragrant.	S. America.	1821.	6. 8. G.b.	Bot. reg. t. 640.
EUCROSIA. B.R.	EUCROSIA.	Hexandria Monogynia.			
bico. B.R.	(or.re) two-coloured.	Cape Horn.	1816.	4. 5. S.b.	Bot. reg. t. 207.
CALOS'EMMA. B.P.	CALOSTE'MMA.	Hexandria Monogynia.			
1 luteum. B.R.	(ye.) yellow.	N. Holland.	1819.	8. 10. F.b.	Bot. reg. t. 421.
2 purpureum. B.P.	(pu.) purple.	—	—	—	— t. 422.
3 album. B.P.	(wh.) white-flowered.	—	1824.	—	G.b.
E'URYCLES. Sal.	E'URYCLES.	Hexandria Monogynia.			
1 nuda.	(wh.) naked stamen'd.	E. Indies.	1822.	5. 7. S.b.	
Crinum nervosum. H.B.					
2 alata.	(wh.) wing-stamen'd.	N. Holland.	1821.	—	S.b.
Pancratium australasicum. B.R.					Bot. reg. t. 715.
3 coronata.	(wh.) crowned.	E. Indies.	1759.	—	S.b.
Pancratium amboinense. B.M.	Prôphys amboinensis. H.A.				Bot. mag. t. 1419.
Three very distinct species, all flowered at Mr. Colvill's.					
HYMENOCALLIS. H.A.	HYMENOCALLIS.	Hexandria Monogynia.			
1 speciosa. K.R.	(wh.) large-flowered.	W. Indies.	1759.	5. 9. S.b.	Bot. mag. t. 1453.
Pancratium speciosum. B.M.					
2 ovata. K.R.	(wh.) broad-leaved.	—	—	—	S.b.
3 amânea. K.R.	(wh.) handsome.	Guiana.	1790.	—	S.b.
4 frâgrans. L.T.	(wh.) fragrant.	W. Indies.	—	—	S.b.
5 guianensis. B.R.	(wh.) Guiana.	Guiana.	1818.	—	S.b.
6 caribaea. K.R.	(wh.) Caribbean.	W. Indies.	1730.	—	S.b.
Pancratium caribaeum. B.M.					
7 declinata. B.C.	(wh.) declining.	Brazil.	—	—	S.b.
8 undulata. K.R.	(wh.) wave-leaved.	S. America.	—	—	S.b.
9 expansa. B.M.	(wh.) expanded.	W. Indies.	1818.	—	S.b.
10 pedalis. B.C.	(wh.) foot-flowered.	Brazil.	1815.	—	S.b.
11 pátens. K.R.	(wh.) spreading.	W. Indies.	—	—	S.b.
12 littoralis. K.R.	(wh.) sea side.	S. America.	1758.	—	S.b.
13 Dryândri. K.R.	(wh.) Dryander's.	—	—	—	S.b.
14 disticha. H.A.	(wh.) two-ranked.	—	—	—	S.b.
15 angusta. K.R.	(wh.) narrow-leaved.	—	—	—	S.b.
16 tenuiflora. H.A.	(wh.) slender-flowered.	—	—	—	S.b.
17 crassifolia. H.A.	(wh.) thick-leaved.	S. America.	—	—	S.b.
18 acutifolia.	(wh.) acute-leaved.	Mexico.	1824.	—	G.b.
Pancratium mexicanum. B.R. nec aliorum.	Hymenocallis litoralis. ð acutifolia. B.M. 2621.				Bot. reg. t. 940.
19 rotata. H.A.	(wh.) large-crowned.	Carolina.	1803.	7. 9. G.b.	Bot. mag. t. 827.
20 mexicana. K.R.	(wh.) Mexican.	Mexico.	1732.	8. G.b.	Ker's review. pl.3. f.2.
21 caroliniana. H.A.	(wh.) Carolina.	Carolina.	1759.	6. 7. G.b.	— pl. 3. f. 1.
22 Staplesi. W.H.	(wh.) Mr. Staples's.	Mexico.	1826.	—	G.b.
23 ovalifolium. W.H.	(wh.) oval-leaved.	S. America.	1820.	—	S.b.
PANCRATIUM. H.A.	PANCRATIUM.	Hexandria Monogynia.			
1 maritimum. K.R.	sea Daffodil.	S. Europe.	1597.	5. 7. G.b.	Bot. reg. t. 161.
carolinianum. B.R. 927.	nec aliorum.				
2 canariense. B.R.	Canary.	Canaries.	1815.	6. 7. G.b.	Bot. reg. t. 174.
3 illyricum. K.R.	Ilyrian.	S. Europe.	1615.	5. 6. G.b.	Bot. mag. t. 718.
4 verecundum. K.R.	Narcissus-leav'd.	E. Indies.	1776.	6. 8. G.b.	Bot. reg. t. 413.
5 zeylanicum. K.R.	one-flowered.	Ceylon.	1752.	—	S.b.
ISMENE. Sal.	ISMENE.	Hexandria Monogynia.			
1 nûtans. H.A.	(wh.) nodding.	Brazil.	1810.	6. 8. S.b.	Bot. mag. t. 1561.
Pancratium calathinum. B.M. non R.L.					
2 calathinum. H.A.	(wh.) cup-flowered.	—	—	6. 7. S.b.	Bot. reg. t. 215.

3	Amáncæs. H.A. (ye.) Narcissus-flower'd. Peru.	1804.	6. 7. S.b.	Bot. mag. t. 1224.
	<i>Pancratium Amáncæs.</i> B.M. <i>Narcissus Amáncæs.</i> R.P. 3. t. 283. f.a.			
NARCISSUS. L.	NARCISSUS. Hexandria Monogynia. L.			
'AJAX. H.S.S.				
1	obvallâris. H.S.S. (ye.) Sibthorp's. S. Europe.	3. 4. H.b.	Bot. mag. t. 1301. f.inf.	
2	bicolor. L. (wh.ye.) two-coloured. Spain. 1629. 4. 5. H.b.	——	t. 1187.	
	'Ajax lorifolius. H.S.S.			
3	tubæflorus. s.p. (wh.ye.) tube-flowered. S. Europe.	3. 5. H.b.		
	'Ajax bicolor. H.S.S.			
4	moschatus. B.M. (wh.st.) greater white. Spain. 1629. 4. H.b.	Bot. mag. t. 924.		
5	álbus. L.T. (wh.) white flowered. ——— 1759. 3. 4. H.b.	——	t. 1300.	
	moschatus. δ. B.M. 'Ajax álbus. H.S.S.			
6	cérnus. R.B. (wh.) drooping-flowered. 1828. ——— H.b.			
	β plenus. (wh.) double-flowered. ——— H.b.			
7	abscissus. H.S.S. (ye.) clipt trunk. ——— H.b.			
8	màjor. B.M. (ye.) large. Spain. 1629. ——— H.b.			
9	propinquus. s.p. (ye.) great jagged. S. Europe. ——— H.b.			
	màjor β. B.M. 'Ajax propinquus. H.S.S.			
10	Telamónius. H.S.S. (ye.) Telamonian. ——— ——— H.b.			
	β plenus. H.S.S. (ye.) double Daffodil. ——— ——— H.b.			
	γ grandiplenus. H.S.S. (ye.) great double. ——— ——— H.b.			
11	nobilis. H.S.S. (ye.) noble. ——— ——— H.b.			
12	spurius. H.S.S. (ye.) spurious. England. ——— H.b.			
13	serratus. H.S.S. (st.ye.) serrate. ——— ——— H.b.			
	β suavis. H.S.S. (st.ye.) sweet-scented. ——— ——— H.b.			
14	Pseudo-Narcissus. (st.ye.) Daffodil. ——— 3. 4. H.b.			
	β scoticus. H.S.S. (st.ye.) Scotch Daffodil. Scotland. ——— H.b.			
	γ plenus. H.S.S. (ye.) double French. ——— H.b.			
	δ plenissimus. H.S.S. (ye.) rose double. ——— H.b.			
15	lobulâris. H.P. (ye.) lobe-crowned. England. ——— H.b.			
16	pùmilus. s.p. (ye.) small wedge-fld. S. Europe. 3. H.b.			
	'Ajax cuneiflorus. H.S.S.			
17	mìnòr. B.M. (ye.) small. Spain. 1629. 3. 5. H.b.			
	α mìnimus. H.S.S. (ye.) smallest. ——— ——— H.b.			
	β mìnòr. H.S.S. (ye.) small. ——— ——— H.b.			
	γ mèdius. H.S.S. (ye.) middle-sized. ——— ——— H.b.			
	δ conspicuus. H.S.S. (ye.) conspicuous. ——— ——— H.b.			
	ε angustus. H.S.S. (ye.) narrow-leaved. ——— ——— H.b.			
	DIOM'EDES. H.S.S.			
18	Macléaii. B.M. (st.ye.) Mac Leay's. Levant. 4. 5. H.b.			
	Diomèdes màjor. H.P.			
19	Sabini. B.R. (st.ye.) Sabine's. ——— ——— H.b.			
	Diomèdes mìnòr. H.P.			
	CORBUL'ARIA. H.S.S.			
20	tenuifolius. L.T. (ye.) slender-leaved. Biscay. 1760. 3. 4. H.b.			
21	lobulâtus. H.S.S. (ye.) lobed. ——— ——— H.b.			
22	obèsus. s.p. (ye.) inflated. Portugal. 1796. ——— H.b.			
	β minor. H.S.S. (ye.) smaller. ——— ——— H.b.			
23	álbicans. H.S.S. (wh.) Trompet Marin. 1629. ——— H.b.			
24	Bulbocodium. w. (ye.) hoop petticoat. S. Europe. ——— ——— H.b.			
	QUELTIA. H.S.S.			
25	aurântius. H.S.S. (ye.) orange rimmed. ——— ——— H.b.			
	β lûteo-aurântius. (wh.ye.) Butter and eggs. ——— ——— H.b.			
26	incomparâbilis. B.M. (st.ye.) peerless. Portugal. ——— 4. 5. H.b.			
27	sulphureus. (st.) self coloured peerless. 1629. ——— H.b.			
	Quél'tia còncolor. H.S.S.			
	β plenus. (st.) double orange crown. ——— ——— H.b.			
28	semipartitus. H.S.S. (st.ye.) cloven-cupped. 1818. 3. 4. H.b.			
29	Quél'tia. (wh.st.) peerless white. 4. H.b.			
	Quél'tia álba. H.S.S.			
	β níveo-aurântius. (wh.st.) Orange Phænix. ——— ——— H.b.			
30	montanus. B.R. (wh.) mountain. Pyrenees. 1620. ——— H.b.			
31	galanthifolius. H.S. (wh.) Snowdrop-leav'd. 1720. 5. H.b.			
32	cápax. H.S.S. (wh.) capacious. Levant. ——— H.b.			
	calathinus. R.L. Quél'tia cápax. H.S.S.			
33	Redoutei. (wh.) long-tubed. ——— ——— H.b.			
	calathinus. R.L. non L. nec aliorum.			

SCHISANTHES. H.S.S.

34	orientalis. L. (st.ye.) oriental.	Levant.	1778.	4. 5. H.b.	Bot. mag. t. 948.
	incomparabilis. β. B.M. <i>Schisánthes orientális</i> . H.S.S.				
35	ochroleucus. H.R. (st.) drooping.	Portugal.	1620.	4. H.b.	Bot. mag. t. 48.
	triándrus. B.M. non L. <i>Ganymèdes céruus</i> . H.S.S. <i>ochroleucus</i> . H.R.				
36	triándrus. L. (wh.) white drooping. S. Europe.	1629.	— H.b.	Park. parad. t. 93. f. 2.	
	<i>Ganymèdes álbus</i> . H.R.				
37	nútans. H.S.S. (ye.) yellow cupped.	1789.	3. 5. H.b.	Bot. mag. t. 945.	
	trílobus. B.M. <i>Ganymèdes nútans</i> . H.S.S.				
38	cóncolor. H.S.S. (st.) self-coloured.	1626.	4. H.b.		
39	striátulus. H.S.S. (ye.st.) faint-striped.	...	— H.b.		
40	pulchellus. H.S.S. (ye.wh.) white-cupped.	1620.	— H.b.		
	<i>PHYLOGYNE</i> . H.S.S.				
41	rugulósus. H.S.S. (ye.) wrinkle-cupped.	...	— H.b.		
42	odòrus. L. (ye.) starry-flowered. Corsica.	1720.	3. H.b.	Bot. mag. t. 934.	
	<i>calathinus</i> . α. B.M. <i>Phylógyne odòra</i> . H.S.S.				
43	interjéctus. H.S.S. (ye.) great curled cup. S. Europe.	...	4. H.b.		
44	trílobus. L. (ye.) three-lobed. Spain.	1629.	3. H.b.	Bot. mag. t. 78.	
	<i>odòrus</i> . B.M. <i>Phylógyne trilòba</i> , H.S.S. <i>calathina</i> . Sal.				
45	heminalis. H.S.S. (ye.) lesser curled cup.	...	4. H.b.	Park. parad. t. 93. f. 4.	
46	párvulus. (ye.) small.	1629.	— H.b.		
	<i>Phylógyne minor</i> . H.S.S.				
β reginæ.	(ye.) Queen Ann's Jonquil.	—	— H.b.		
	<i>HERMIONE</i> . H.S.S.				
47	Jonquilla. B.M. (ye.) Jonquil.	Spain.	1596.	4. 5. H.b.	Bot. mag. t. 15.
β plénus.	(ye.) double Jonquil.	—	— H.b.		
48	símilis. H.S.S. (ye.) lesser Jonquil.	S. Europe.	1626.	— H.b.	
49	bifrons. B.M. (ye.) double-faced.	1807.	4. H.b.	Bot. mag. t. 1186.	
50	compréssus. L.T. (ye.) Jasmine Jonquil.	1790.	3. 4. H.b.		
51	primulinus. H.S. (ye.) Cowslip-cupped.	—	— H.b.	Bot. mag. t. 1299.	
	<i>bifrons</i> . β. B.M. <i>Hermione primulina</i> . H.S.S.				
52	calathinus. L. (ye.) many-flowered. Levant.	—	— H.b.	Rudb. elys. 2. p. 60.f.5.	
53	tereticáulis. L.T. (st.) cream-coloured.	—	4. H.b.	Bot. mag. t. 1298?	
54	citrinus. H.S.S. (wh.ye.) Citroniere.	—	— H.b.	t. 946.	
	<i>orientális</i> . γ. B.M.				
55	floribundus. H.S.S. (wh.ye.) grand monarque.	—	— H.b.		
56	Trewiánus. B.M. (wh.ye.) Bazelman major.	—	— H.b.	Bot. mag. t. 940.	
	<i>Hermione grandiflora</i> . H.S.S.				
57	crenulatus. H.S. (wh.ye.) Bazelman minor.	—	— H.b.	Park. parad. t. 81. f. 5.	
58	Tazetta. L. (wh.ye.) French Daffodil.	1626.	— H.b.	— t. 81. f. 3.	
59	dúbius. P.S. (wh.) doubtful.	S. France.	—	Redout. lil. t. 429.	
60	pátulus. DC. (wh.ye.) spreading.	Isl. Hyères.	—		
61	fstulósus. H.S. (wh.ye.) pipe-stemmed.	Levant.	—		
62	cerinus. H.S. (wh.ye.) waxen cupped.	—	5. H.b.		
63	Lùna. H.S.S. (wh.) broad white.	—	4. F.b.		
64	papyràtius. B.M. (wh.) paper white.	Italy.	—	2. 3. F.b.	Bot. mag. t. 947.
	<i>Hermione papyràtia</i> . H.S.S.				
65	únicolor. T.N. (wh.) one-coloured.	Naples.	—	— H.b.	Ten. fl. nap. 1. t. 26.
66	pr'æcox. T.N. (st.ye.) early-flowered.	—	— H.b.	— 1. t. 27.	
67	ítalicus. B.M. (st.ye.) Italian.	Italy.	—	— F.b.	Bot. mag. t. 1188.
	<i>Hermione itálica</i> . H.S.S.				
a semiplénus.	(wh.) semidouble.	—	3. F.b.		
β plénus.	(wh.) double Roman.	—	— F.b.		
68	gracilis. B.R. (pa.ye.) slender.	—	4. H.b.	Bot. reg. t. 816.	
69	teniòr. B.M. (pa.ye.) slender 2-flowered.	Italy.	1626.	— H.b.	Bot. mag. t. 379.
70	latifolius. H.S.S. (ye.) broad-l'd. orange.	Levant.	—	4. 5. H.b.	
71	multiflorus. H.S.S. (ye.) common yellow.	—	— H.b.	Bot. mag. t. 925.	
	<i>Tazetta</i> . B.M. non L.				
72	intermèdius. R.L. (pa.ye.) intermediate.	Pyrenees.	—	— H.b.	Redout. lil. t. 426.
	<i>orientális</i> . δ. Bot. mag. t. 1026. <i>Hermione multiflora</i> . β àurea. H.S.S.?				
	<i>NARCISSUS</i> . H.S.S.				
73	poéticus. L. (wh.) Poet's.	Greece.	1620.	4. F.b.	Park. parad. t. 75. f. 3.
74	angustifolius. B.M. (wh.) narrow-leaved.	Switzerland.	1626.	— H.b.	Bot. mag. t. 193.
75	majális. B.M. (wh.) May-flowering.	England?	—	5. H.b.	Eng. bot. v. 4. t. 275.
	<i>poéticus</i> . E.B. non L. <i>patelláris</i> . s.p.				
β exsérthus. H.S.S. (wh.) long-stamened.	—	—	— H.b.		
γ plénus. H.S.S. (wh.) double-flowered.	—	—	— H.b.	Park. parad. t. 75. f. 1.	
76	biflorus. B.M. (wh.) two-flowered.	—	4. 5. H.b.	Eng. bot. v. 4. t. 276.	

77 níveus. W.en. (<i>wh.</i>) snowy white.	S. Europe.	4. 5. H.b.	
<i>CHLORA'STER.</i> H.P.				
78 viridiflórus. B.M. (<i>gr.</i>) green-flowered.	Barbary.	1629.	9. 10. F.b.	Bot. mag. t. 1687.
<i>Chloráster fissus.</i> H.P.				
79 intéger. H.P. (<i>gr.</i>) entire-cupped.	—	—	F.b.	Park. parad. t. 93. f. 6.
80? obsolétus. H.S.S. (<i>w.ye.</i>) leafy autumnal.	Spain.	—	—	— 90.t.89.f.4.
81? serotinus. w. (<i>wh.</i>) late-flowered.	Barbary.	—	—	Clus. hist. t. 252.
<i>GALA'NTHUS.</i> L. SNOWDROP.	Hexandria Monogynia. L.			
1 nivalis. w. (<i>wh.</i>) common.	Britain.	1. 3. H.b.	Eng. bot. v. 1. t. 19.
2 plicátus. B.R. (<i>wh.</i>) plaited.	S. Europe.	1818.	2. 4. H.b.	Bot. reg. t. 545.
<i>LEUCOJUM.</i> L. SNOWFLAKE.	Hexandria Monogynia. L.			
1 vérnum. w. (<i>wh.</i>) spring.	Germany.	1596.	2. 4. H.b.	Bot. mag. t. 46.
2 carpáthicum. (<i>wh.</i>) Carpathian.	Carpathian mt. 1816.	—	H.b.	— t. 1993.
3 aestivum. w. (<i>wh.</i>) summer.	England.	4. 5. H.b.	— t. 1210.
4 pulchéllum. P.L. (<i>wh.</i>) neat.	—	—	H.b.	Sal. par. lond. t. 74.
<i>'ACIS.</i> P.L. 'Acis.	Hexandria Monogynia.			
1 rósea. S.F.G. (<i>ro.</i>) rose-coloured.	S. Europe.	1826.	8. 9. H.b.	Swt. br. fl. gar. t. 297.
2 grandiflóra. S.F.G. (<i>bh.</i>) large-flowered.	—	1828.	—	Redout. liliac. t. 217.
3 trichophy'lla. S.F.G. (<i>wh.</i>) many-flowered.	Barbary.	1812.	1. 2. F.b.	Bot. reg. t. 544.
<i>Leucójum trichophyl'lum.</i> B.R.				
4 autumnális. P.L. (<i>bh.</i>) autumnal.	Portugal.	1629.	9.	Bot. mag. t. 960.
<i>Leucójum autumnálle.</i> B.M.				
<i>GETHY'LLIS.</i> L. GETHY'LLIS.	Hexandria Monogynia. L.			
1 spirális. w. (<i>wh.</i>) spiral-leaved.	C. B. S.	1780.	6. 7. G.b.	Bot. mag. t. 1088.
2 ciliáris. w. (<i>wh.</i>) fringed.	—	1788.	—	Jacq. schœn. t. 79.
3 villósa. w. (<i>wh.</i>) hairy.	—	1787.	—	
4 lanceoláta. w. (<i>wh.</i>) spear-leaved.	—	1790.	—	
<i>ALSTROEM'ERIA.</i> W. ALSTROEM'ERIA.	Hexandria Monogynia. L.			
1 Pelegrina. w. (<i>re.bh.</i>) spotted flower'd. Peru.	—	1753.	6. 9. H. 4.	Bot. mag. t. 139.
β álbida. (<i>wh.gr.</i>) white flowered.	—	1820.	—	Swt. br. fl. gar. t. 267.
2 Símsii. s.s. (<i>sc.ye.</i>) red-flowered.	Chili.	1822.	—	Hook. ex. flor. t. 181.
<i>pulchella.</i> B.M. 2353. Hook. ex. flor. t. 64.				
3 Hookéri. (<i>ro.</i>) rose-coloured.	Chili.	1822.	—	
rósea. H.E.F. non R.P.			H. 4.	
4 Flós Martini. B.R. (<i>w.p.y.</i>) St. Martin's.	—	—	—	Bot. reg. t. 731.
púlchra. Bot. mag. t. 2421. tricolor.	Hook. ex. flor. t. 65.			Lodd. bot. cab. t. 1497.
5 bicolor. B.C. (<i>ye.wh.</i>) two-coloured.	Chili.	1826.	—	Swt. br. fl. gar. s.2. t. 15.
6 psittácina. s.s. (<i>sc.gr.</i>) Parrot-like.	Mexico.	1829.	7. 9. H. 4.	
7 acutifólia. Ot. (<i>or.</i>) sharp-leaved.	S. America.	—	9. 10. F. 4.	
8 pállida. Ot. (<i>pa.</i>) pale-flowered.	Chili.	—	—	
9 Curtisiána. R.S. (<i>wh.sc.</i>) striped-flowered.	Brazil.	1776.	2. 4. S. 4.	Bot. mag. t. 125.
<i>Lígtu.</i> B.M. nec aliorum.				
10 ovata. s.s. (<i>sc.</i>) oval-leaved.	Peru.	1824. H. 4. —	Cavan. ic. 1. t. 76.
11 Salsilla. w. (<i>pu.gr.</i>) climbing.	—	1806.	6. 7. H. 4. —	Fenill. per. 2.p. 713. t. 6.
12 édulis. Tus. (<i>ro.gr.</i>) eatable-rooted.	W. Indies.	1801.	6. 9. S. 4. —	Andr. deposit. t. 649.
<i>Salsilla.</i> Bot. mag. t. 1613. non L.				
13 hirtella. k.s. (<i>pi.gr.</i>) hairy.	Mexico.	1824. H. 4. —	Swt. br. fl. gar. t. 228.
<i>CL'IVIA.</i> B.R. CL'IVIA.	Hexandria Monogynia.			
nóbilis. B.R. (<i>sc.gr.</i>) scarlet.	C. B. S.	1820.	7. 10. G. 4.	Bot. reg. t. 1182.
<i>Imatophy'llum Aitoni.</i> Bot. mag. t. 2856.				
<i>DORYA'NTHES.</i> B.P. DORYA'NTHES.	Hexandria Monogynia. S.S.			
excélsa. B.P. (<i>sc.</i>) gigantic.	N. S. W.	1800.	7. 10. G. 4.	Bot. mag. t. 1685.

ORDO CCX.

HEMEROCALLIDÆ. Brown prodr. 1. p. 295.

HEMEROCA'LLIS. S.S. DAY LILY. Hexandria Monogynia. L.

1 graminea. B.M. (<i>ye.</i>) narrow-leaved.	Siberia.	1759.	6. 7. H. 4.	Bot. mag. t. 873.
2 fláva. B.M. (<i>ye.</i>) yellow.	—	1596.	—	— t. 19.
3 speciosa. W.H. (<i>co.</i>) specious.	Jamaica.	1816.	—	
4 disticha. B.F.G. (<i>co.</i>) fan-like.	China.	1798.	5. 7. H. 4.	Swt. br. fl. gar. t. 28.
5 fúlva. B.M. (<i>co.</i>) copper-coloured.	Levant.	1596.	6. 8. H. 4.	Bot. mag. t. 64.

HEMEROCALLIDEÆ.

517

FUNKIA. S.S.	FUNKIA. Hexandria Monogynia. S.S.				
1 álba. <i>subcordata</i> . s.s.	(wh.) white-flowered. Japan.	1790.	8. 9. H. 4.	Bot. mag. t. 1433.	
2 cœrulea. <i>ovata</i> . s.s.	(wh.) blue-flowered. <i>Hemerocallis cœrulea</i> . B.M.	—	6. 8. H. 4.	Bot. mag. t. 894.	
AGAPANTHUS. W.	AFRICAN LILY. Hexandria Monogynia. L.				
1 umbellatus. w.	(bl.) large-flowered. C. B. S.	1692.	1. 12. G. 4.	Bot. mag. t. 500.	
β variegatus.	(bl.) striped-leaved.	—	— G. 4.		
2 præcox. W.en.	(bl.) early-flowering.	—	1. 4. G. 4.	Lodd. bot. cab. t. 42.	
3 minor. B.C.	(bl.) small.	—	— G. 4.		
POLIANTHES. L.	POLIANTHES. Hexandria Monogynia. L.				
1 gracilis. L.en.	(wh.) slender. Brazil.	1822.	8. 10. S. 4.	Bot. reg. t. 63.	
2 tuberosa. w.	(wh.) common.	E. Indies.	1629.	— S. 4.	
β flôræ pleno.	(wh.) double-flowered.	—	— S. 4.		
BLANDFORDIA. B.P.	BLANDFORDIA. Hexandria Monogynia. S.S.				
1 nobilis. B.P.	(sc.ye.) noble. N. S. W.	1803.	7. 8. G. 4.	Bot. reg. t. 286.	
2 grandiflora. B.P.	(sc.ye.) large-flowered.	—	1824.	— G. 4.	
3 punicea.	(cr.) large-leaved. N. Holland.	1812.	— G. 4.	Lab. nov. hol. 1. t. 111.	
<i>Alétris punicea</i> . Labillardiere.					
TRITOMA. B.M.	TRITOMA. Hexandria Monogynia. S.S.				
1 Uvária. H.K.	(or.) great. C. B. S.	1707.	8. 9. H. 4.	Bot. mag. t. 758.	
2 Burchellii. W.H.	(or.) Mr. Burchell's.	—	1816.	— H. 4.	
3média. H.K.	(sc.or.) lesser.	—	1789.	12. 6. H. 4.	
4 pumila. H.K.	(sc.or.) least.	—	1774.	9. 11. F. 4.	
VELTHEIMIA. B.M.	VELTHEIMIA. Hexandria Monogynia. W.				
1 viridifolia. w. (sc.or.) green-leaved.	C. B. S.	1768.	11. 4. G.b.	Bot. mag. t. 501.	
<i>Alétris capensis</i> . B.M.					
2 intermédia.	(ro.ye.) subglauous.	—	1819.	— G.b.	
3 glauca. w.	(ro.ye.) glaucous.	—	1781.	1. 4. G.b.	
ALETRIS. W.	COLIC-ROOT. Hexandria Monogynia. L.				
1 farinosa. w.	(wh.) common. N. America.	1768.	6. H. 4.	Bot. mag. t. 1418.	
2 aurea. Ph.	(wh.) golden-tipped.	—	1811.	7. 8. H. 4.	Willd. hort. ber. t. 8.
SANSEVITERA. W.	SANSEVITERA. Hexandria Monogynia. W.				
1 glauca. H.S.	(wh.) glaucous.	Sierra Leon.	6. 10. S. 4.	
2 polyphylla. H.S.	(wh.) upright-glaucous.	—	—	S. 4.	
3 guineensis. H.S.	(wh.) Guinea.	Guinea.	1690.	6. 11. S. 4.	
4 lætevirens. H.S.	(wh.) light green.	—	—	S. 4.	
5 longiflora. B.M.	(wh.) long-flowered.	—	—	7. 8. S. 4.	
6 spicata. H.S.	(wh.) spiked.	—	—	S. 4.	
7 fulvocincta. H.S.s.	(wh.) brown-edged.	Brazil.	1818.	6. 9. S. 4.	
8 zeylanica. H.S.	(wh.) Ceylon.	Ceylon.	1731.	6. 11. S. 4.	
9 lanuginosa. w.	(wh.) woolly.	E. Indies.	—	S. 4.	
10 ensifolia. H.S.	(wh.) sword-leaved.	—	—	S. 4.	
11 grandicuspis. H.S.s.	(wh.) large-pointed.	—	—	S. 4.	
12 pumila. H.S.	(wh.) dwarf.	C. B. S.	1796.	— S. 4.	
13 stenophylla. L.en.	(wh.) slender-leaved.	—	1818.	6. 8. S. 4.	
14 cárnea. A.R.	(ro.) flesh-coloured.	China.	1792.	4. 10. H. 4.	
TULBAGIA. W.	TULBAGIA. Hexandria Monogynia. L.			Andr. deposit. t. 361.	
1 alliacea. w.	(gr.) Narcissus-leaved. C. B. S.	1774.	5. 7. G. 4.	Bot. mag. t. 806.	
2 affinis. L.en.	(gr.) linear-leaved.	—	1820.	— G. 4.	
3 cepacea. w.	(gr.) Onion-scented.	—	1795.	4. G. 4.	
BRODIAEA. L.T.	BRODIAEA. Triandria Monogynia. L.T.				
1 grandiflora. L.T.	(bl.) large-flowered. Georgia.	1806.	6. H.b.	Bot. reg. t. 1183.	
<i>Hoekera coronaria</i> . Sal. par. lond. t. 98.					
2 congesta. L.T.	(bl.) close-headed.	—	—	5. H.b.	
PHYLLOMA. B.M.	PHYLLOMA. Hexandria Monogynia.				
1 aloiflorum. B.M.	(ye.) Aloe-like.	Bourbon.	1766.	4. 6. D.S. h.	
<i>Lomatophyllum borbonicum</i> . w.	<i>Draconia marginata</i> . H.K.			Bot. mag. t. 1585.	
2 macrum. L.en.	(ye.) lean.	Mauritius.	1817.	6. 7. D.S. h.	
3 rufocinctum.	(ye.) rosy-edged.	E. Indies.	1818.	— D.S. h.	
<i>Aloe rufocincta</i> . H.S.s.					
PACHIDENDRON. H.R.	PACHIDENDRON. Hexandria Monogynia. H.R.				
1 africanum. H.R.	(ye.) African.	C. B. S.	1731.	7. D.G. h.	
<i>a latum</i> . H.R.	(ye.) broad-leaved.	—	—	D.G. h.	
β angustum. H.R.	(ye.) narrow-leaved.	—	—	D.G. h.	
<i>Aloe pseudo-africana</i> . L.en.	<i>Aloe africana</i> . β. angustior. B.M.			Bot. mag. t. 2517.	
2 principis. H.R.	(ye.) revolving red-spined.	—	—	D.G. h.	
3 angustifolium. H.R.	(ye.) narrow-leaved.	—	1812.	— D.G. h.	
4 ferox. H.R.	(st.re.) great hedgehog.	—	1759.	4. 5. D.G. h.	
<i>Aloe ferox</i> . B.M.		2 L 3		Bot. mag. t. 1975.	

5	pseudo-ferox. H.R. (<i>st.</i>)	false hedgehog.	C. B. S.	1759.	4.	D.G. $\frac{1}{2}$.	Comm. praejud. t. 20.
6	supral'æve. H.R. (<i>ye.</i>)	upright hedgehog.	_____	1731.	D.G. $\frac{1}{2}$.	
A'LOE. H.R.	ALOE.	Hexandria Monogynia. L.					
1	dichótoma. H.S.	smooth-stemmed.	C. B. S.	1780.	D.G. $\frac{1}{2}$.	
2	purpurásens. H.S. (<i>re.gr.</i>)	purple.	_____	1789.	7.	D.G. $\frac{1}{2}$.	Bot. mag. t. 1474.
3	soccotrína. H.S. (<i>sc.g.</i>)	soccotrine.	_____	1731.	4.	D.G. $\frac{1}{2}$.	_____ t. 472.
4	spicáta. H.R. (<i>wh.gr.</i>)	white spotted.	_____	1795.	D.G. $\frac{1}{2}$.	
5	plúridens. H.P.	many-toothed.	_____	1822.	D.G. $\frac{1}{2}$.	
6	arborésrens. H.S. (<i>sc.</i>)	tree.	_____	1731.	11.	D.G. $\frac{1}{2}$.	Bot. mag. t. 1306.
7	gráclis. H.P.	slender-stalked.	_____	1822.	D.G. $\frac{1}{2}$.	
8	flavispina. H.S. (<i>sc.</i>)	yellow-spined.	_____	1790.	8.	D.G. $\frac{1}{2}$.	
9	mitræfórmis. H.S. (<i>sc.</i>)	mitre.	_____	1732.	D.G. $\frac{1}{2}$.	Bot. mag. t. 1270.
10	nóbilis. H.S. (<i>sc.</i>)	great mitre.	_____	1800.	7.	D.G. $\frac{1}{2}$.	
11	Commelini. H.R. (<i>sc.</i>)	Commeline's.	_____	1813.	D.G. $\frac{1}{2}$.	
	β mitræfórmis. H.R.	yellowish-toothed.	_____	D.G. $\frac{1}{2}$.	
12	dístans. H.S. (<i>sc.gr.</i>)	small mitre.	_____	1732.	8.	D.G. $\frac{1}{2}$.	Bot. mag. 1362.
	γ depréssâ. H.R. (<i>sc.gr.</i>)	depressed.	_____	D.G. $\frac{1}{2}$.	
13	xanthacántha. H.S.s. (<i>sc.</i>)	yellow mitre.	_____	1817.	D.G. $\frac{1}{2}$.	
14	vulgâris. DC. (<i>ye.or.</i>)	yellow-flowered. W. Indies.	barbadénsis. H.S.	1596.	8.	D.S. $\frac{1}{2}$.	DC. plant. gras. t. 27.
15	chinénsis. H.S.s. (<i>ye.</i>)	Chinese.	China.	1817.	D.G. $\frac{1}{2}$.	
16	cæsia. H.S.s. (<i>sc.</i>)	cæsian tree.	C. B. S.	1818.	D.G. $\frac{1}{2}$.	
	β elátior. H.S.s. (<i>sc.</i>)	tall.	_____	D.G. $\frac{1}{2}$.	
17	frutéscens. H.S.s. (<i>sc.</i>)	lesser tree.	_____	D.G. $\frac{1}{2}$.	
18	albispina. H.S. (<i>sc.</i>)	white-spined.	_____	1796.	D.G. $\frac{1}{2}$.	
19	glauca. H.S. (<i>sc.gr.</i>)	glaucous red-spined.	_____	1731.	1.	D.G. $\frac{1}{2}$.	
	α mähr. H.S. (<i>sc.gr.</i>)	larger.	_____	D.G. $\frac{1}{2}$.	
	β rhodacántha. B.M. (<i>sc.gr.</i>)	lesser red-spined.	_____	D.G. $\frac{1}{2}$.	Bot. mag. t. 1278.
	γ spinösior. H.R. (<i>sc.gr.</i>)	long-spined.	_____	1820.	D.G. $\frac{1}{2}$.	
20	lineata. H.S. (<i>sc.gr.</i>)	striped red-spined.	_____	1789.	D.G. $\frac{1}{2}$.	
	α vîridis. H.R. (<i>sc.gr.</i>)	greenest-leaved.	_____	D.G. $\frac{1}{2}$.	
	β glauçescens. H.R. (<i>sc.</i>)	glaucous.	_____	D.G. $\frac{1}{2}$.	
21	dorsális. R.S. (<i>ye.</i>)	spiny-keeled.	_____	1826.	D.G. $\frac{1}{2}$.	
22	depréssâ. H.S. (<i>sc.ye.</i>)	depressed.	_____	1731.	8.	D.G. $\frac{1}{2}$.	Bot. mag. t. 1332.
23	brevifòlia. H.S. (<i>sc.gr.</i>)	short-leaved.	_____	6.	D.G. $\frac{1}{2}$.	Bot. reg. t. 996.
24	prolífera. H.R. (<i>or.y.</i>)	proliferous.	_____	1818.	D.G. $\frac{1}{2}$.	
25	sérra. DC. (<i>re.gr.</i>)	saw-leaved.	_____	D.G. $\frac{1}{2}$.	Jacq. eclog. t. 61.
26	striátâ. H.S. (<i>re.gr.</i>)	streaked.	paniculâta. Jacquin.	1795.	D.G. $\frac{1}{2}$.	Jacq. fragm. t. 68.
27	striátula. H.P. (<i>sc.</i>)	stripe-sheathed.	_____	1820.	D.G. $\frac{1}{2}$.	
28	ciliâris. H.P. (<i>sc.gr.</i>)	fringe-leaved.	_____	D.G. $\frac{1}{2}$.	
29	tenuíor. H.P. (<i>ye.</i>)	green-sheathed.	_____	1819.	D.G. $\frac{1}{2}$.	
30	albocínta. H.S.s. (<i>sc.gr.</i>)	pearl edged.	_____	1812.	D.G. $\frac{1}{2}$.	
31	serrulâta. H.S. (<i>pi.</i>)	serrulated.	_____	1789.	7.	D.G. $\frac{1}{2}$.	Bot. mag. t. 1415.
32	obscûra. H.S. (<i>sc.or.</i>)	great soap.	_____	1727.	6.	D.G. $\frac{1}{2}$.	— t. 1323.
	α picta. a. B.M.	—	_____	1819.	D.G. $\frac{1}{2}$.	
	β glauçior. H.R. (<i>sc.or.</i>)	glaucous-leaved.	_____	D.G. $\frac{1}{2}$.	
	γ magnidens. H.R. (<i>sc.or.</i>)	large-toothed.	_____	D.G. $\frac{1}{2}$.	
33	pallésrens. H.R. (<i>pi.gr.</i>)	pale serrulate.	_____	1820.	D.G. $\frac{1}{2}$.	
34	latifòlia. H.S. (<i>sc.or.</i>)	broad-leaved.	_____	1795.	7.	D.G. $\frac{1}{2}$.	Bot. mag. t. 1346.
35	grandidentâta. R.S. (<i>sc.gr.</i>)	great-toothed.	_____	D.G. $\frac{1}{2}$.	
36	saponâria. H.S. (<i>sc.or.</i>)	common soap.	_____	1727.	D.G. $\frac{1}{2}$.	Bot. mag. t. 1460.
	β luteo-striâta. H.R. (<i>sc.</i>)	yellow-striped.	_____	1821.	D.G. $\frac{1}{2}$.	
37	microcántha. H.S.s. (<i>re.</i>)	small-spined.	_____	1819.	D.G. $\frac{1}{2}$.	Bot. mag. t. 2272.
38	vîrens. H.S. (<i>sc.</i>)	apple green.	_____	1790.	8.	D.G. $\frac{1}{2}$.	— t. 1355.
39	suberécta. H.S. (<i>sc.ye.</i>)	lesser hedgehog.	_____	1789.	3.	D.G. $\frac{1}{2}$.	
	β semiguttâta. H.R. (<i>sc.gr.</i>)	half-warted.	_____	1819.	D.G. $\frac{1}{2}$.	
40	acuminâta. H.S. (<i>sc.ye.</i>)	middle hedgehog.	hùmilis. B.M. non H.S.	1795.	D.G. $\frac{1}{2}$.	Bot. mag. t. 757.
41	echinâta. H.R. (<i>sc.gr.</i>)	great tubercled.	_____	1821.	3.	D.G. $\frac{1}{2}$.	
42	tuberculâta. H.S. (<i>sc.gr.</i>)	tuberced.	_____	1796.	D.G. $\frac{1}{2}$.	
43	aristata. H.P. (<i>sc.gr.</i>)	bearded many-leaved.	_____	1819.	D.G. $\frac{1}{2}$.	
44	subtuberculâta. H.P. (<i>se.</i>)	rough & smooth.	_____	D.G. $\frac{1}{2}$.	
45	hùmilis. H.S. (<i>sc.gr.</i>)	dwarf hedgehog.	_____	1731.	D.G. $\frac{1}{2}$.	
46	incúrvâ. H.S. (<i>sc.gr.</i>)	incurved hedgehog.	hùmilis β incúrvâ. B.M.	1796.	D.G. $\frac{1}{2}$.	DC. plant. gras. t. 39.
47	tenuifòlia. H.R. (<i>sc.gr.</i>)	thin-leaved.	_____	1821.	D.G. $\frac{1}{2}$.	
48	variegâta. H.S. (<i>sc.gr.</i>)	partridge-breast.	_____	1720.	3.	D.G. $\frac{1}{2}$.	Bot. mag. t. 513.

RHIPIDODE'NDRON. W.	RHIPIDODE'NDRON.	Hexandria Monogynia.	
1 plicátile. H.R. (<i>re.gr.</i>) fan.	Africa.	1723.	6. 7. D.G. ½.
<i>dístichum</i> . w. <i>A'lœ plicátilis</i> . B.M.		1820.	— D.G. ½.
β major. H.R. (<i>re.</i>) larger fan.	C. B. S.	1780.	— D.G. ½.
2 dichotomum. w. (<i>re.gr.</i>) dichotomous.	TONGUE-ALOE.		Patt. journ. t. 3. 4. 5.
GASTERIA. H.S.	Hexandria Monogynia. H.S.		
1 oblíqua. H.S. (<i>ro.w.g.</i>) broad-marbled.	C. B. S.	1759.	6. 8. D.G. ½.
<i>maculata</i> . H.P. <i>A'lœ língua</i> , a. B.M.		1759.	Bot. mag. t. 979.
2 púchra. H.S. (<i>ro.gr.</i>) narrow-marbled.		1759.	Bot. mag. t. 765.
<i>A'lœ maculata</i> . B.M.			
3 pícta. H.P. (<i>ro.gr.</i>) painted.		1824.	— D.G. ¼.
<i>A'lœ Bowieana</i> . R.S.			
4 formosa. H.P. (<i>sc.gr.</i>) beautiful.		1826.	— D.G. ¼.
5 fasciata. H.P. (<i>ro.gr.</i>) marbled.		1820.	— D.G. ¼.
β láixa. H.P. (<i>ro.gr.</i>) loose-leaved.		—	— D.G. ¼.
6 retáta. H.P. (<i>ro.gr.</i>) chequered.		1826.	— D.G. ¼.
<i>A'lœ dict'yodes</i> . R.S.			
7 nigricans. H.S. (<i>re.gr.</i>) dark-leaved.		1790.	6. 7. D.G. ¼.
8 marmorata. H.P. (<i>re.g.</i>) marble-leaved.		—	— D.G. ¼.
9 crassiòlia. H.P. (<i>re.gr.</i>) thick-leaved.		1820.	— D.G. ¼.
10 brevífolia. H.S. (<i>re.gr.</i>) short-leaved.		1809.	7. 8. D.G. ¼.
α lätewrens. H.P. (<i>re.gr.</i>) pale-green.		—	— D.G. ¼.
β perviriðis. H.P. (<i>re.gr.</i>) darker-green.		—	— D.G. ¼.
11 obtusifolia. H.R. (<i>re.g.</i>) blunt-leaved.		—	— D.G. ¼.
12 móllis. H.R. (<i>re.gr.</i>) soft muddy-leaved.		1819.	— D.G. ¼.
13 subnígricans. H.P. (<i>re.g.</i>) dark-leaved.		1820.	— D.G. ¼.
β glábrior. H.P. (<i>re.gr.</i>) smoother.		—	— D.G. ¼.
14 dísticha. H.P. (<i>re.gr.</i>) distichous.		1819.	— D.G. ¼.
α minor. H.P. (<i>re.gr.</i>) lesser.		—	— D.G. ¼.
β major. H.P. (<i>re.gr.</i>) larger.		—	— D.G. ¼.
15 conspurcata. H.R. (<i>re.g.</i>) dirtied.		1796.	3. 11. D.G. ¼.
16 angulata. H.P. (<i>re.gr.</i>) angular.		—	— D.G. ¼.
17 sulcata. H.P. (<i>re.w.g.</i>) furrowed.		1791.	— D.G. ¼.
18 excavata. H.P. (<i>re.gr.</i>) excavated.		1824.	— D.G. ¼.
19 angustifolia. H.S. (<i>re.g.</i>) narrow.		1731.	— D.G. ¼.
20 l'avis. H.P. (<i>re.gr.</i>) smooth.		1820.	— D.G. ¼.
21 subverrucosa. H.P. (<i>re.g.</i>) large-warted.		—	— D.G. ¼.
α grandipunctata. H.R. large-dotted.		—	— D.G. ¼.
β parvipunctata. H.R. small-dotted.		—	— D.G. ¼.
22 verrucosa. H.S. (<i>re.g.</i>) pearl.		1731.	3. 11. D.G. ¼.
β latifolia. H.R. (<i>re.g.</i>) broad-leaved.		1820.	— D.G. ¼.
23 intermédia. H.S. (<i>re.g.</i>) intermediate.		1790.	— D.G. ¼.
β asperrima. H.R. (<i>re.g.</i>) rough-leaved.		1820.	— D.G. ¼.
γ l'ævior. H.P. (<i>re.gr.</i>) smoother.		—	— D.G. ¼.
δ lóngior. H.P. (<i>re.gr.</i>) longer-leaved.		—	— D.G. ¼.
24 répens. H.R. (<i>re.gr.</i>) creeping-rooted.		—	— D.G. ¼.
25 párvula. H.P. (<i>re.gr.</i>) smallest.		1824.	— D.G. ¼.
26 décipiens. H.P. (<i>re.gr.</i>) deceiving.		1823.	7. 10. D.G. ¼.
27 carimata. H.S. (<i>sc.gr.</i>) rough-keeled.		1731.	6. 7. D.G. ¼.
28 strigata. H.P. (<i>re.gr.</i>) rough strigate.		1824.	7. 10. D.G. ¼.
29 lètipunctata. H.P. (<i>re.gr.</i>) pale-spotted.		1821.	— D.G. ¼.
30 subcarinata. H.S.s. (<i>re.</i>) slightly-keeled.		1820.	6. 7. D.G. ¼.
β striata. H.R. (<i>re.gr.</i>) striated.		—	— D.G. ¼.
31 undata. H.P. (<i>re.gr.</i>) waved.		1825.	— D.G. ¼.
32 glábra. H.S. (<i>re.gr.</i>) smooth-keeled.		1796.	6. 7. D.G. ¼.
β minor. H.R. (<i>re.gr.</i>) smaller.		1820.	— D.G. ¼.
33 nítida. B.M. (<i>sc.gr.</i>) shining.		—	— D.G. ¼.
α parvipunctata. H.R. small dotted.		—	— D.G. ¼.
β grandipunctata. H.R. large-dotted.		—	— D.G. ¼.
34 trigona. H.R. (<i>li.gr.</i>) trigonal.		—	— D.G. ¼.
35 obtusa. H.P. (<i>re.gr.</i>) smooth obtuse.		—	— D.G. ¼.
36 acinacifolia. H.R. (<i>ro.g.</i>) long sword-leaved.		1818.	— D.G. ¼.
37 nitens. H.P. (<i>re.gr.</i>) sparse-spotted.		1819.	— D.G. ¼.
38 venusta. H.P. (<i>re.gr.</i>) charming.		1822.	— D.G. ¼.
39 pluripunctata. H.P. (<i>re.gr.</i>) many-spotted.		—	— D.G. ¼.
40 ensifolia. H.P. (<i>re.gr.</i>) long sword-leaved.		1820.	— D.G. ¼.
41 cándicans. H.R. (<i>re.</i>) rough marbled white.		1818.	7. 8. D.G. ¼.
42 línta. H.P. (<i>re.</i>) smeared.		1822.	— D.G. ¼.
43 bicolor. H.P. (<i>re.</i>) half-marbled.		—	— D.G. ¼.

HEMEROCALLIDÆ.

BOWTEA. H.P.	Bow'EA. Hexandria Monogynia.
1 africana. H.P. (<i>ye.</i>) African.	C. B. S. 1820. 4. 10. D.G. ¼.
2 myriacántha. H.P. (<i>ro.g.</i>) many-spined.	— D.G. ¼.
HAWO'RTHIA. H.S.S. Hawo'RTHIA.	Hexandria Monogynia. H.S.
1 viscósa. H.S. (<i>wh.re.</i>) clammy.	C. B. S. 1732. 5. 7. D.G. ¼. Bot. mag. t. 814.
<i>A'loe viscósa.</i> B.M.	
α mājor. (<i>wh.re.</i>) largest.	— D.G. ¼.
β mīnor. (<i>wh.re.</i>) lesser.	— D.G. ¼.
γ parvifòlia. (<i>wh.re.</i>) smaller-leaved.	— D.G. ¼.
2 áspéra. H.S. (<i>wh.re.</i>) rough-triangular.	1795. — D.G. ¼.
3 indurátâ. H.R. (<i>wh.re.</i>) hard branched.	1819. — D.G. ¼.
4 subtortuosa. R.S. (<i>wh.re.</i>) twisted triangular.	1818. — D.G. ¼.
5 torquâta. H.P. (<i>wh.re.</i>) long-twisted.	1824. — D.G. ¼.
6 concínna. H.S.S. (<i>wh.re.</i>) neat triangular.	— D.G. ¼.
7 cordifolia. H.S.S. (<i>wh.re.</i>) heart-leaved.	— D.G. ¼.
8 asperiúscula. H.S.S. (<i>wh.re.</i>) roughish.	1818. — D.G. ¼.
9 tortuosa. H.S. (<i>wh.re.</i>) twisted.	1794. 5. 9. D.G. ¼. Bot. mag. t. 1337.
<i>A'loe rígida.</i> B.M. non DC.	
β curta. H.R. (<i>wh.re.</i>) short twisted.	1817. — D.G. ¼.
γ tortella. H.R. (<i>wh.re.</i>) lesser twisted.	— D.G. ¼.
10 expânsa. H.S. (<i>wh.re.</i>) spreading-leaved.	1795. 7. 10. D.G. ¼.
11 rígida. H.R. (<i>wh.gr.</i>) great rigid.	1824. — D.G. ¼. DC. plant. gras. t. 62.
12 hy'bida. H.R. (<i>wh.re.</i>) bastard pearl.	1821. 5. 9. D.G. ¼.
13 scâbra. H.R. (<i>wh.re.</i>) rugged.	1818. — D.G. ¼.
14 sórdida. H.R. (<i>wh.re.</i>) sordid.	— D.G. ¼.
15 recârva. H.R. (<i>wh.re.</i>) recurved.	1795. 8. D.G. ¼. Bot. mag. t. 1353.
16 nígricans. H.P. (<i>wh.re.</i>) granulated black.	1820. 5. 9. D.G. ¼.
17 párvâ. H.P. (<i>wh.re.</i>) dwarf.	— D.G. ¼.
18 tessellâta. H.P. (<i>wh.re.</i>) dark chequered.	1822. — D.G. ¼.
19 aspérulea. H.P. (<i>wh.re.</i>) pale rough.	— D.G. ¼.
20 multifâria. H.P. (<i>wh.re.</i>) many-leaved.	— D.G. ¼.
21 retûsa. H.S. (<i>wh.gr.</i>) smooth cushion.	1720. — D.G. ¼. Bot. mag. t. 455.
<i>A'loe retûsa.</i> B.M.	
22 mütica. H.R. (<i>wh.gr.</i>) blunt cushion.	1819. — D.G. ¼.
23 mirâbilis. H.S.S. (<i>wh.ro.</i>) rough cushion.	1795. — D.G. ¼. Bot. mag. t. 1354.
24 túrgida. H.S.S. (<i>wh.ro.</i>) turgid cushion.	1818. — D.G. ¼.
25 l'âte-virens. H.S.S. (<i>wh.re.</i>) light green.	— D.G. ¼.
26 papillôsa. H.S.S. (<i>wh.re.</i>) papillose.	— D.G. ½.
β semipapillôsa. H.R. partly warted. — D.G. ½.
27 erécta. H.R. (<i>wh.re.</i>) upright pearl.	1818. — D.G. ¼. DC. plant. gras. t. 57.
<i>margaritifera.</i> DC.	
28 fasciata. H.S.S. (<i>wh.re.</i>) barred pearl.	— D.G. ¼.
29 subulâta. R.S. (<i>wh.re.</i>) awl-pointed.	— D.G. ¼.
30 claripérla. H.P. (<i>wh.re.</i>) bright-pearled.	1824. — D.G. ¼.
31 granâta. H.S.S. (<i>wh.re.</i>) small pearl.	1732. — D.G. ¼. Dill. elth. t. 16. f. 18.
<i>A'loe margaritifera. γ. mímina.</i> Dill.	
32 brêvis. H.S.S. (<i>wh.ro.</i>) short-leaved.	1810. — D.G. ¼. Bot. mag. t. 1360.
<i>A'loe margaritifera. γ. mímina.</i> B.M.	
33 minor. H.S. (<i>pu.gr.</i>) lesser pearl. 5. 8. D.G. ¼. Bot. mag. t. 815.
34 margaritifera. H.S.S. (<i>wh.</i>) pearl.	1725. — D.G. ¼. Bradl. succ. 3. t. 21.
35 semiglabrâta. H.S.S. (<i>w.gr.</i>) half-smoothed.	1810. — D.G. ¼.
36 rádula. H.S. (<i>ro.wh.</i>) raspy pearl.	1805. 6. 9. D.G. ¼. Jacq. schoen. t. 35.
37 attenuâta. H.S. (<i>wh.re.</i>) chalky pearl.	1790. 5. 8. D.G. ¼. Bot. mag. t. 1345.
<i>A'loe rádula.</i> B.M. non Jacq.	
38 Reinwârdti. H.R. (<i>wh.re.</i>) chequered-pearl.	1821. — D.G. ¼.
39 semimargaritifera. (<i>wh.</i>) semi-pearly.	1801. — D.G. ¼.
α máxima. H.S.S. (<i>wh.re.</i>) largest.	— D.G. ¼. Comm. hort. 2. t. 10.
β mājor. H.S.S. (<i>wh.re.</i>) large.	— D.G. ¼.
γ mīnor. H.S.S. (<i>wh.re.</i>) smaller.	1817. — D.G. ¼.
δ multiperlata. (<i>wh.re.</i>) many pearly.	1818. — D.G. ¼.
40 coartâta. H.P. (<i>wh.re.</i>) upright dull-spotted.	1822. — D.G. ¼.
41 álbicans. H.S. (<i>wh.re.</i>) white-edged.	1795. 9. 10. D.G. ¼. Bot. mag. t. 1452.
42 l'avis. H.R. (<i>wh.re.</i>) narrow-bordered.	1820. — D.G. ¼. Comm. hort. 2. t. 7.
43 ramifera. H.R. (<i>wh.re.</i>) branching white.	— D.G. ¼.
44 virésens. H.R. (<i>wh.re.</i>) bordered green.	— D.G. ¼.
β minor. H.R. (<i>wh.re.</i>) lesser.	— D.G. ¼.
45 arachnoïdes. H.S. (<i>lh.gr.</i>) cobweb.	1727. 8. 9. D.G. ¼. Bot. mag. t. 756.
46 translucens. H.S.S. (<i>bh.</i>) transparent.	1795. 5. 8. D.G. ¼. — t. 1417.
<i>A'loe arachnoïdes β translucens.</i> B.M.	

47	setatæ. H.S.S. (<i>wh.re.</i>) bristle-edged.	C. B. S.	1818.	5. 8.	D.G. ȝ.
	β nigricans. H.R. (<i>wh.re.</i>) blackish.	—	—	—	D.G. ȝ.
	γ media. H.R. (<i>wh.re.</i>) middle.	—	—	—	D.G. ȝ.
	δ major. H.R. (<i>wh.re.</i>) larger.	—	—	—	D.G. ȝ.
48	pallida. H.R. (<i>wh.re.</i>) pale green.	—	—	—	D.G. ȝ.
49	atrovirens. H.R. (<i>wh.bh.</i>) dark green. <i>A'loe atrovirens.</i> DC. arachnoïdes pumila. B.M.	—	1752.	5.	D.G. ȝ. Bot. mag. t. 1361.
50	chlorocántha. H.R. (<i>w.re.</i>) green-toothed.	—	1820.	5. 8.	D.G. ȝ.
51	angustifòlia. H.P. (<i>w.re.</i>) slender-leaved.	—	—	6. 7.	D.G. ȝ.
52	reticulatæ. H.S. (<i>wh.</i>) netted. <i>A'loe arachnoïdes</i> γ reticulata. B.M.	—	1794.	5. 8.	D.G. ȝ. Bot. mag. t. 1314.
53	cymbiformis. H.S. (<i>w.pu.</i>) boat-leaved. concava. H.R.	—	1795.	6. 8.	D.G. ȝ. Bot. mag. t. 802.
54	altilínea. H.P. (<i>wh.re.</i>) ridge-lined.	—	1821.	—	D.G. ȝ.
55	obtusa. H.P. (<i>wh.re.</i>) small blunt-leaved.	—	—	—	D.G. ȝ.
56	planifòlia. H.P. (<i>wh.re.</i>) flat oval-leaved.	—	—	—	D.G. ȝ.
57	mucronatæ. H.R. (<i>w.pu.</i>) membrane-pointed. C. B. S.	1818.	—	—	D.G. ȝ.
58	límvida. H.R. (<i>w.pu.</i>) limpid.	—	—	—	D.G. ȝ.
59	aristatæ. H.R. (<i>w.pu.</i>) pin-pointed.	—	—	—	D.G. ȝ.
60	cuspidatæ. H.R. (<i>w.pu.</i>) sharp-pointed.	—	—	—	D.G. ȝ.
61	denticulatæ. H.R. (<i>w.re.</i>) pale pin-pointed.	—	—	—	D.G. ȝ.
A'PICRA. H.S.S. A'PICRA. Hexandria Monogynia. H.S.S.					
1	spiralis. H.S.S. (<i>wh.g.</i>) great spiral.	C. B. S.	1790.	8. 9.	D.G. ḥ.
2	spirella. H.S.S. (<i>wh.g.</i>) small spiral.	—	1808.	—	D.G. ḥ.
3	bullulata. H.S.S. (<i>st.gr.</i>) blistered.	—	1818.	—	D.G. ḥ. Jacq. frag. t. 109.
4	pentagona. H.S.S. (<i>wh.</i>) five sided.	—	1731.	6. 8.	D.G. ḥ. Bot. mag. t. 1338.
5	quinquangularis. R.S. (<i>w.st.</i>) five-angled.	—	—	D.G. ḥ.
6	rigida. w. (<i>wh.re.</i>) gunpowdered. pseudo-rigida. H.S.S.	—	1818.	—	D.G. ḥ. Jacq. frag. t. 108.
7	aspera. H.S.S. (<i>wh.st.</i>) rough triangled.	—	—	—	D.G. ḥ.
	α minor. H.S.S. (<i>wh.st.</i>) lesser.	—	—	—	D.G. ḥ.
	β major. H.S.S. (<i>wh.st.</i>) larger.	—	1795.	—	D.G. ḥ.
8	bicarinata. H.S.S. (<i>w.st.</i>) double-keeled.	—	1819.	—	D.G. ḥ.
9	nigra. H.P. (<i>wh.</i>) rough black.	—	1820.	5. 8.	D.G. ḥ.
10	imbricata. H.R. (<i>st.</i>) rough-flowered. <i>A'loe spiralis.</i> B.M. non H.S.	—	1731.	6. 7.	D.G. ḥ. Bot. mag. t. 1455.
11	foliolosa. H.P. (<i>wh.</i>) many small-leaved.	—	1795.	6. 8.	D.S. ḥ. Bot. mag. t. 1352.

ORDO CCXI.

DIOSCOREÆ. Brown prodr. 1. p. 294.

RAJ'ANIA. W.	RAJ'ANIA. Dioccia Hexandria. L.				
1	hastata. w. (<i>gr.</i>) halberd-leaved.	W. Indies.	1822.	7. 8.	S. ȝ. —. Plum. amer. 84. t. 98.
2	cordata. w. (<i>gr.</i>) heart-leaved.	—	1786.	—	S. ȝ. —. Plum. ic. 155. f. 1.
3	quinquefolia. w. (<i>gr.</i>) five-leaved.	—	1820.	—	S. ȝ. —. — 155. f. 2.
DIOSC'OREA. W.	YAM. Dioccia Hexandria. L.				
1	pentaphylla. w. (<i>gr.</i>) five-leaved.	E. Indies.	1768.	S. ȝ. —. Rheed. mal. 7. t. 35.
2	triphylla. w. (<i>gr.</i>) three-leaved.	—	1818.	S. ȝ. —. Jacq. ic. t. 627.
3	quinqueloba. w. (<i>gr.</i>) five-lobed.	Japan.	1822.	7. 9.	G. ȝ. —. Kämpf. ic. t. 15.
4	brasiliensis. w. (<i>gr.</i>) Brazilian.	Brazil.	1818.	—	S. ȝ. —.
5	heterophylla. H.B. (<i>gr.</i>) various-leaved.	E. Indies.	—	S. ȝ. —.
6	aculeata. w. (<i>gr.</i>) prickly-stemmed.	—	1803.	7. 9.	S. ȝ. —. Rheed. mal. 7. t. 37.
7	rubella. H.B. (<i>re.g.</i>) reddish.	—	—	—	S. ȝ. —.
8	purpurea. H.B. (<i>pu.g.</i>) purple.	—	1816.	—	S. ȝ. —.
9	angustifolia. s.s. (<i>gr.</i>) narrow-leaved.	Peru.	1821.	—	S. ȝ. —.
10	nummularia. w. (<i>gr.</i>) Money-wort.	E. Indies.	1820.	S. ȝ. —. Rumph. amb. 5. t. 162.
11	alata. w. (<i>gr.</i>) wing-stalked.	—	1739.	S. ȝ. —. Rheed. mal. 7. t. 38.
12	bulbifera. w. (<i>gr.</i>) bulb-bearing.	—	1692.	7. 9.	S. ȝ. —. Sal. par. lond. t. 17.
13	crispata. H.B. (<i>gr.</i>) curl-stalked.	—	1818.	—	S. ȝ. —.
14	altissima. s.s. (<i>gr.</i>) tallest.	Martinica.	1821.	—	S. ȝ. —. Plum. ic. 117. f. 2.
15	coriacea. s.s. (<i>gr.</i>) leathery.	S. America.	1818.	—	S. ȝ. —.
16	anguina. H.B. (<i>gr.</i>) snake.	E. Indies.	1803.	—	S. ȝ. —.
17	pulchella. H.B. (<i>gr.</i>) pretty.	—	1819.	S. ȝ. —.

DIOSCOREÆ.

18 atropurpurea. H.B. (<i>pu.g.</i>) dark purple.	C. B. S.	1820.	S. $\frac{1}{4}$.—.
19 globosa. H.B. (<i>gr.</i>) globular.	_____	1818.	S. $\frac{1}{4}$.—.
20 fasciculata. H.B. (<i>gr.</i>) bundled.	_____	1817.	S. $\frac{1}{4}$.—.
21 glabra. H.B. (<i>gr.</i>) smooth.	_____	1803.	7. 9.	S. $\frac{1}{4}$.—.
22 sativa. w. (<i>gr.</i>) common.	W. Indies.	1733.	—	S. $\frac{1}{4}$.—. Rheed. mal. 8. t. 51.
23 piperifolia. w. (<i>gr.</i>) Pepper-leaved.	S. America.	1817.	S. $\frac{1}{4}$.—. Plum. ic. 117. f. 1.
24 nepalensis. (<i>gr.</i>) Nepaul.	Nepaul.	1816.	7. 8.	H. $\frac{1}{4}$.—.
25 cinnamomifolia. B.M. (<i>g.</i>) Cinnamon-leav'd. Brazil.	1826.	11. 12.	S. $\frac{1}{4}$.—.	Bot. mag. t. 2825.
26 quaternata. Ph. (<i>re.g.</i>) four-leaved.	N. America.	1811.	7.	H. $\frac{1}{4}$.—.
27 villosa. Ph. (<i>re.g.</i>) pubescent.	_____	1752.	8.	H. $\frac{1}{4}$.—. Jacq. ic. 3. t. 626.
28 oppositifolia. L. (<i>gr.</i>) opposite-leav'd.	E. Indies.	1803.	S. $\frac{1}{4}$.—. Petiv. gaz. t. 31. f. 6.
TESTUDIN'ARIA. B.T. TESTUDIN'ARIA.	Dicecia Hexandria.			
1 Elephantiæ. B.R. (<i>gr.</i>) Elephant's-foot. C. B. S.	1774.	7. 8.	G. $\frac{1}{4}$.—.	Bot. reg. t. 92.
Tamus Elephantiæ. Bot. mag. t. 1347.				
2 montana. B.T. (<i>gr.</i>) glaucous-leaved.	_____	1816.	—	G. $\frac{1}{4}$.—.

ORDO CCXII.

TAMEÆ. Brown prodr. 294 in obs.

TAMUS. L.

1 communis. w.

BLACK BRYONY. Dicecia Hexandria. L.

(gr.) common. England. 5. 8. H. $\frac{1}{4}$.—. Eng. bot. t. 91.

2 crética. w.

(gr.) Cretan.

Candia.

1739.

7. 8. H. $\frac{1}{4}$.—.

ORDO CCXIII.

SMILACEÆ. Brown prodr. 292.

SMILAX. L.

SMILAX. Dicecia Hexandria. L.

1 áspéra. s.s. (<i>w.gr.</i>) rough bindweed.	S. Europe.	1648.	9.	H. $\frac{1}{2}$.—. Schk. handb. 3. t. 328.
2 auriculata. (<i>w.gr.</i>) ear-leaved.	_____	—	—	H. $\frac{1}{2}$.—. Pluk. alm. t. 110. f. 3.
2 nigra. s.s. (<i>w.gr.</i>) black.	_____	1817.	—	H. $\frac{1}{2}$.—.
3 catalonica. s.s. (<i>w.gr.</i>) Catalonian.	_____	—	—	H. $\frac{1}{2}$.—.
4 mauritanica. s.s. (<i>w.gr.</i>) red-berried.	_____	1820.	8. 10.	F. $\frac{1}{2}$.—.
5 excelsa. s.s. (<i>w.gr.</i>) tall.	Syria.	1739.	8. 9.	F. $\frac{1}{2}$.—. Buxb. cent. 1. t. 32.
6 horrida. s.s. (<i>w.gr.</i>) straight-spined.	N. America.	1820.	—	H. $\frac{1}{2}$.—.
7 zeylanica. s.s. (<i>w.gr.</i>) Ceylon.	E. Indies.	1778.	—	S. $\frac{1}{2}$.—. Rumph. amb. 5. t. 161.
8 quadrangularis. s.s. (<i>w.g.</i>) square-stalked.	N. America.	1812.	6. 8.	H. $\frac{1}{2}$.—. Wats. dend. brit. t. 109.
9 longifolia. s.s. (<i>w.gr.</i>) long-leaved.	Cayenne.	1823.	—	S. $\frac{1}{2}$.—.
10 Watsoni. (<i>w.gr.</i>) Watson's.	N. America.	1811.	7. 8.	H. $\frac{1}{2}$.—. Wats. dend. brit. t. 110.
longifolia. W.D. non Rich.				
11 Sarsaparilla. s.s. (<i>w.gr.</i>) medicinal.	_____	1664.	—	H. $\frac{1}{2}$.—. Wats. dend. brit. t. 111.
12 brasiliensis. s.s. (<i>w.gr.</i>) Brazilian.	Brazil.	1820.	—	S. $\frac{1}{2}$.—.
13 hastata. s.s. (<i>w.gr.</i>) halberd-leaved.	Carolina.	1822.	—	F. $\frac{1}{2}$.—. Pluk. alm. t. 111. f. 3.
14 China. s.s. (<i>w.gr.</i>) Chinese.	China.	1759.	G. $\frac{1}{2}$.—. Kämpf. amoen. t. 782.
15 acuminata. s.s. (<i>w.gr.</i>) sharp-pointed.	W. Indies.	1822.	6. 7.	S. $\frac{1}{2}$.—. Plum. ic. 83.
16 glauca. B.M. (<i>wh.</i>) glaucous-leaved.	N. America.	1811.	5. 7.	H. $\frac{1}{2}$.—. Bot. mag. t. 1846.
17 australis. B.P. (<i>w.gr.</i>) oblong-leaved.	N. S. W.	1815.	—	G. $\frac{1}{2}$.—.
18 rotundifolia. s.s. (<i>w.g.</i>) round-leaved.	N. America.	1760.	7. 8.	H. $\frac{1}{2}$.—.
19 laurifolia. s.s. (<i>w.gr.</i>) Laurel-leaved.	_____	1739.	—	H. $\frac{1}{2}$.—. Catesb. car. 1. t. 15.
20 tamnoïdes. s.s. (<i>w.gr.</i>) Tamus-leaved.	_____	—	6. 7.	H. $\frac{1}{2}$.—. — 1. t. 52.
21 caduca. s.s. (<i>w.gr.</i>) deciduous.	_____	1759.	—	H. $\frac{1}{2}$.—.
22 havanensis. s.s. (<i>w.gr.</i>) Havannah.	S. America.	1823.	S. $\frac{1}{2}$.—. Jacq. amer. t. 179. f. 102.
23 ovalifolia. D.P. (<i>w.gr.</i>) oval-leaved.	Nepaul.	1822.	H. $\frac{1}{2}$.—.
24 Bona-nox. s.s. (<i>w.gr.</i>) fringed.	N. America.	1739.	6. 7.	H. $\frac{1}{2}$.—. Pluk. alm. t. 111. f. 1.
25 maculata. D.P. (<i>w.gr.</i>) spotted.	Nepaul.	1819.	H. $\frac{1}{2}$.—.
26 prolifera. H.B. (<i>w.gr.</i>) proliferous.	E. Indies.	—	S. $\frac{1}{2}$.—.
27 glabra. H.B. (<i>w.gr.</i>) smooth.	_____	1820.	S. $\frac{1}{2}$.—.
28 Alpinia. s.s. (<i>w.gr.</i>) Grecian.	Greece.	—	6. 8.	F. $\frac{1}{2}$.—.
29 latifolia. B.P. (<i>w.gr.</i>) broad-leaved.	N. Holland.	1791.	F. $\frac{1}{2}$.—.
30 lanceolata. Ph. (<i>w.gr.</i>) spear-leaved.	N. America.	1785.	5. 6.	H. $\frac{1}{2}$.—. Catesb. car. 2. t. 84.

31	herbacea. s.s. (gr.)	herbaceous.	N. America.	1669.	7.	H. 4. —.	Bot. mag. t. 1920.
32	peduncularis. s.s. (gr.)	long-peduncled.	—	1812.	5. 7.	H. 4. —.	
33	globifer. s.s. (w.gr.)	Surinam.	Surinam.	1819.	S. h. —.	
34	macrophylla. s.s. (w.gr.)	large-leaved.	W. Indies.	1822.	S. h. —.	
35	Pseûdo-chîna. s.s. (w.gr.)	Bastard China.	S. America.	1739.	5. 6.	S. h. —.	Sloan. jam. 1. t. 143. f. 1.
36	glyciphylla. B.P. (w.gr.)	Botany-bay Tea.	N. S. W.	1815.	—	S. h. —.	
37	canariensis. s.s. (w.gr.)	Canary.	Canaries.	1816.	—	F. h. —.	
38	pùbera. s.s. (w.gr.)	downy.	N. America.	1806.	—	H. h. —.	
39	rùbens. w.d. (pi.)	pink-flowered.	—	1811.	7. 8.	H. h. —.	Wats. dend. brit. t. 108.
RIPOG'ONUM. B.P.	RIPOG'ONUM.	Hexandria Monogynia.					
álfifm. B.P.	(wh.) white-flowered.	N. S. W.	1820.	4. 6.	G. h. —.		
RU'SCUS. L.	BUTCHER'S-BROOM.	Dicæcia Monadelphia. L.					
1	aculeatus. w. (w.gr.)	prickly.	England.	12. 6.	H. h.	Eng. bot. v. 8. t. 560.
2	laxus. L.T.	(w.gr.) loose.	Europe.	1. 6.	H. h.	
3	Hypophyllum. w. (g.y.)	thick-leaved.	Italy.	1640.	5. 6.	H. h.	Bot. mag. t. 2049.
	β latifolius. w. (g.y.)	broad-leaved.	S. Europe.	—	—	H. h.	Dill. elt. t. 251. f. 303.
4	Hypoglossum. w. (g.y.)	double-leaved.	Italy.	1596.	4. 5.	H. h.	Schk. handb. 3. t. 340.
5	andrógynum. w. (g.y.)	climbing.	Canaries.	1713.	—	G. h. —.	Bot. mag. t. 1898.
6	reticulatus. w. (g.y.)	netted-leaved.	C. B. S.	1816.	—	G. h. —.	
7	volutilis. w. (g.y.)	many-nerved.	—	—	—	G. h. —.	
8	racemosa. w. (g.wh.)	Alexandrian Laurel.	Portugal.	1713.	6.	H. h.	Wats. dend. brit. t. 145.
POLYGONATUM. D.	SOLOMON'S-SEAL.	Hexandria Monogynia.					
1	verticillatum. R.L. (w.g.)	whorl-leaved.	Scotland.	6. 7.	H. 4.	Redout. liliac. t. 244.
	Convallaria verticillata.	Eng. bot. v. 2. t. 128.					
2	bracteatum. Th. (wh.g.)	bracted.	Switzerland.	1827.	—	H. 4.	
3	leptophyllum. D.P.	slender-leaved.	Nepaul.	1816.	6. 7.	H. 4.	
4	oppositifolium. D.P.	opposite-leaved.	—	—	—	H. 4.	Lodd. bot. cab. t. 640.
	Convallaria oppositifolia. B.C.						
5	vulgare. R.L. (w.gr.)	common.	England.	5. 6.	H. 4.	Redout. liliac. t. 258.
	Convallaria Polygomatum.	Eng. bot. v. 4. t. 280.					
6	pubescens. Ph. (w.g.)	pubescent.	N. America.	1812.	—	H. 4.	Willd. hort. ber. t. 45.
7	canaliculatum. Ph. (w.g.)	channel-leaved.	—	—	6.	H. 4.	
8	angustifolium. Ph. (st.g.)	narrow-leaved.	—	1824.	5. 6.	H. 4.	
9	multiflorum. R.L. (w.g.)	many-flowered.	Britain.	—	H. 4.	Eng. bot. v. 4. t. 279.
10	latifolium. R.L. (st.)	broad-leaved.	Europe.	1802.	—	H. 4.	Redout. liliac. t. 243.
11	macrophyllum. (w.g.)	large-leaved.	N. America.	1823.	—	H. 4.	
	latifolium. Ph. nec aliorum.						
12	polyanthemum. M.B. (w.)	many-flowered.	Caucasus.	—	—	H. 4.	
SMILACINA. Df.	SMILACINA.	Hexandria Monogynia. Ph.					
1	racemosa. Df. (wh.)	cluster-flowered.	N. America.	1640.	5. 6.	H. 4.	Redout. liliac. t. 230.
	Convallaria racemosa.	Bot. mag. t. 899.					
2	stellata. Df. (wh.)	star-flowered.	—	1633.	—	H. 4.	Bot. mag. t. 185.
3	trifolia. Df. (wh.)	three-leaved.	—	1812.	6. 7.	H. 4.	Gmel. sib. 1. t. 6.
4	canadensis. Ph. (wh.)	Canadian.	—	—	—	H. 4.	Redout. lil. t. 216. f. 1.
5	bifolia. Df. (wh.)	least.	N. Europe.	1596.	5. 6.	H. 4.	Bot. mag. t. 510.
	Convallaria bifolia. B.M.	Majanthemum bifolium.	Redout. liliac. t. 216. f. 2.				
6	borealis. B.M. (g.y.)	oval-leaved.	N. America.	1778.	5. 6.	H. 4.	Bot. mag. t. 1403.
7	umbellata. Df. (wh.)	umbel-flowered.	—	—	—	H. 4.	— t. 1155.
OPHIOPOGON. B.M.	SNAKE'S-BEARD.	Hexandria Monogynia. B.M.					
1	japonicus. B.M. (wh.)	Japan.	Japan.	1784.	6.	F. 4.	Bot. mag. t. 1063.
2	spicatus. B.R. (bl.)	spiked.	China.	1821.	8. 10.	F. h.	Bot. reg. t. 593.
3	intermedius. D.P. (wh.)	intermediate.	Nepaul.	1824.	—	F. 4.	
CONVALLARIA. D.	LILY OF THE VALLEY.	Hexandria Monogynia. L.					
majális. E.B.	(wh.) common.	Britain.	5. 6.	H. 4.	Eng. bot. v. 15. t. 1035.	
a ábla.	(wh.) white-flowered.	—	—	—	H. 4.		
β ròsea.	(ro.) rose-coloured.	—	—	—	H. 4.		
γ plèna.	(wh.) double-flowered.	—	—	—	H. 4.		
STREPTOPUS. M.	STREPTOPUS.	Hexandria Monogynia. Ph.					
1	amplexifolius. R.L. (g.y.)	heart-leaved.	Europe.	1752.	5.	H. 4.	W. et K. hung. t. 167.
2	distortus. M. (g.y.)	distorted.	N. America.	—	—	H. 4.	
3	ròseus. M. (ro.)	rose-coloured.	—	1806.	6. 7.	H. 4.	Bot. mag. t. 1489.
	Uvularia ròsea. B.M.						
4	lanuginosus. M. (g.y.)	woolly.	—	1812.	—	H. 4.	Bot. mag. t. 1490.
5	simplex. D.P. (br.gr.)	simple-stalked.	Nepaul.	1822.	—	H. 4.	
MED'EOLA. S.S.	INDIAN CUCUMBER.	Hexandria Trigynia. L.					
virginica. s.s. (gr.)	Virginian.	Virginia.	1759.	6.	H. 4.	Bot. mag. t. 1316.	
GYROMÍA. N.	GYROMÍA. N.						
TRI'LLIUM. L.	TRI'LLIUM.	Hexandria Trigynia. L.					
1	séssile. s.s. (da.pu.)	sessile-flowered.	N. America.	1759.	4. 5.	H. 4.	Bot. mag. t. 40.

2	petiolatum. s.s. (<i>d.pu.</i>)	Plantain-leaved. N.America.	1811.	4. 5. H. 4.	
3	erythrocárpum. M. (<i>w.re.</i>)	painted-flowered. <i>pictum. Ph.</i>	—	5. 6. H. 4.	Swt. br. fl. gar. t. 212.
4	undulatum. s.s. (<i>wh.</i>)	wave-flowered.	—	—	H. 4.
5	pusílum. s.s. (<i>pu.</i>)	dwarf. Carolina.	1812.	—	F. 4.
6	ovátum. s.s. (<i>pa.pu.</i>)	purple-flowered. N.America.	—	—	F. 4.
7	cérnum. s.s. (<i>wh.</i>)	drooping-flowered.	1758.	4. 5. H. 4.	Bot. mag. t. 954.
8	eréctum. w. (<i>d.pu.</i>)	stinking.	1759.	—	H. 4.
9	péndulum. s.s. (<i>wh.</i>)	white-flowered. <i>erethocárpum. B.M. non M.</i>	1805.	—	H. 4.
10	obovátum. s.s. (<i>d.ro.</i>)	obovate-petaled.	1824.	—	H. 4.
11	grandiflórum. s.s. (<i>wh.</i>)	large-flowered. <i>erythrocárpum. B.M. non M.</i>	1799.	4. 6. H. 4.	Sal. par. lond. t. 1.
12	stylósum. N. (<i>ro.</i>)	long-styled. Carolina.	1823.	—	F. 4.
P'ARIS. L.	HERB-PARIS.	Octandria Tetragynia. L.			
1	quadrifólia. E.B. (<i>gr.</i>)	True-love-or-1-berry. England.	5. 6. H. 4.	Eng. bot. v. 1. t. 7.
2	verticillata. M.B. (<i>gr.</i>)	whorl-leaved. Siberia.	1825.	—	H. 4.
3	polyphylla. D.P. (<i>gr.</i>)	many-leaved. Nepaul.	1816.	—	H. 4.

ORDO CCXIV.

ASPHODELEÆ. Brown prodr. 1. p. 274.

ASPHO'DELUS. L. ASPHODEL. Hexandria Monogynia. L.

1	clavátus. H.B. (<i>wh.</i>)	club-seeded.	E. Indies.	1808.	7. 10. S. ⊖.
2	fistulósus. s.s. (<i>wh.</i>)	Onion-leaved.	S. Europe.	1596.	6. 9. H. 4.
3	intermédius. L.en. (<i>wh.</i>)	intermediate.	Canaries.	1824.	—
4	álbus. s.s. (<i>wh.</i>)	upright.	S. Europe.	5. 7. H. 4.
5	æstivus. s.s. (<i>wh.</i>)	summer.	Portugal.	1819.	—
6	asiáticus. R.S. (<i>wh.</i>)	Asiatic.	Levant.	H. 4.
7	ramósus. s.s. (<i>wh.</i>)	branched.	S. Europe.	1551.	—
8	prolífér. M.B. (<i>wh.</i>)	proliferous.	Armenia.	1824.	6. 10. H. ⊖.
9	tenuior. s.s. (<i>ye.gr.</i>)	slender-leaved.	Caucasus.	1823.	7.
10	créticus. s.s. (<i>ye.gr.</i>)	Cretan.	Candia.	1821.	6. 7. H. 4.
11	libúrnicus. R.S. (<i>ye.gr.</i>)	oriental.	Levant.	—	—
12	tauricus. s.s. (<i>ye.gr.</i>)	Taurian.	Tauria.	1812.	5. 7. H. 4.
13	capillaris. R.S. (<i>ye.gr.</i>)	awl-leaved.	—	—	H. 4.
14	lúteus. s.s. (<i>ye.</i>)	common yellow. Sicily.	—	1596.	5. 6. H. 4.

EREM'URUS. S.S. EREM'URUS. Hexandria Monogynia. S.S.

spectabilis. s.s. (<i>ye.</i>)	channel-leaved. Caucasus.	1800.	5. 6. H. 4.
Asphodelus altàicus. w.			

CZA'CKIA. S.S. ST. BRUNO'S LILY. Hexandria Monogynia. S.S.

Liliástrum. s.s. (<i>wh.</i>)	Savoy Spiderwort. Switzerland.	1629.	5. 6. H. 4.
Anthéricum Liliástrum. B.M. Hemerocális Liliástrum. W.en.			

PHALA'NGIUM. J. PHALA'NGIUM. Hexandria Monogynia.

1	Liliágo. B.M. (<i>wh.</i>)	grass-leaved.	S. Europe.	1596.	5. 6. H. 4.
	α major. B.M. (<i>wh.</i>)	larger.	—	—	H. 4.
	β mìnór. B.M. (<i>wh.</i>)	lesser	—	—	H. 4.
2	ramosum. B.M. (<i>wh.</i>)	branching.	—	1570.	H. 4.

Anthéricum ramosum. w.

3	græcum. Lam. (<i>wh.</i>)	Grecian.	Greece.	1828.	—
4	nepalense. B.R. (<i>wh.</i>)	Nepaul.	Nepaul.	1822.	6. 10. F. 4.
5	glauicum. P.S. (<i>wh.</i>)	glaucous.	S. America.	1816.	—
6	pomeridiánum. (<i>wh.</i>)	afternoon-flower'd.	—	1818.	G. 4.

Anthéricum pomeridiánum. B.R. Scilla pomeridiána. DC.

ANTHE'RICUM. ANTHE'RICUM. Hexandria Monogynia. L.

1	hirsútum. R.S. (<i>wh.</i>)	hairy-leaved.	—	1822.	6. 10. G. 4.
2	blepharóphoron. R.S. (<i>w.pu.</i>)	fringed.	—	—	G. 4.
3	bipedunculatum. (<i>wh.</i>)	two-flowered.	—	1825.	4. 5. G. 4.
4	triflórum. w. (<i>wh.</i>)	three-flowered.	—	1782.	8. 10. G. 4.
5	undulátum. w. (<i>wh.</i>)	wave-leaved.	—	1825.	4. 6. G. 4.
6	pilosum. w. (<i>wh.</i>)	pilose.	—	—	G. 4.
7	longifolium. w. (<i>wh.</i>)	long-leaved.	—	—	G. 4.
8	graminifolium. w. (<i>wh.</i>)	grass-leaved.	—	1794.	6. 7. G. 4.
9	vespertinum. w. (<i>w.re.</i>)	afternoon-flower'd.	—	1803.	5. 9. G. 4.

Bot. mag. t. 954.
t. 470.
W. hort. ber. 1. t. 35.

Eng. bot. v. 1. t. 7.

Bot. mag. t. 984.

Blackw. t. 238.

Bot. mag. t. 799.

Hoff.com.m.l.p.t.43.

Bot. mag. t. 2626.

Lodd. bot. cab. t. 915.

Jacq. f. eclog. t. 77.

Redout. liliac. t. 470.

— t. 380.

Bot. mag. t. 773.

Swt. br. fl. gar. t. 188.

Bot. mag. t. 318.

Redout. lil. t. 269.

Bot. mag. t. 1635.

— t. 914.

— t. 1055.

Flor. græc. t. 336.

Bot. reg. t. 998.

Bot. reg. t. 564.

Jacq. ic. 2. t. 410.

Jacq. ic. 2. t. 416.

— t. 413.

— t. 411.

Bot. mag. t. 1040.

10	Jacquinianum. R.S.(w.) <i>Jacquin's.</i>	C. B. S.	1795.	5. 9.	G. ¼.	Jacq. ic. 2. t. 412.
11	divaricatum. R.S. (<i>wh.</i>) <i>divaricate.</i>	_____	_____	_____	G. ¼.	Jacq. schœn. 4. t. 414.
12	revolutum. w. (<i>wh.</i>) <i>curled-flowered.</i>	_____	1731.	9. 12.	G. ¼.	Bot. mag. t. 1044.
13	floribundum. w. (<i>st.</i>) <i>thick-spiked.</i>	_____	1774.	3. 4.	G. ¼.	
14	squamēum. w. (<i>wh.</i>) <i>scaly.</i>	_____	1820.	_____	G. ¼.	
15	filiiforme. w. (<i>wh.gr.</i>) <i>thread-leaved.</i>	_____	1774.	4.	G. ¼.	
16	elongatum. R.S. (<i>wh.g.</i>) <i>long-racemed.</i>	_____	1822.	_____	G. ¼.	
17	flexifolium. w. (<i>wh.</i>) <i>flexuous-leaved.</i>	_____	1795.	5. 6.	G. ¼.	
BULBINE. L. <i>BULB'INE.</i> Hexandria Monogynia. L.						
1	hispida. L. (<i>bh.gr.</i>) <i>hairy-leaved.</i>	C. B. S.	1774.	5. 6.	D.G. ¼.	Jacq. ic. 2. t. 409.
2	canaliculata. s.s. (<i>wh.</i>) <i>channel-leaved.</i>	_____	_____	4. 5.	D.G. ¼.	Bot. mag. t. 1124.
3	semibarbata. s.s. (<i>ye.</i>) <i>half bearded.</i>	N. S. W.	1820.	4. 8.	D.G. ¼.	Lodd. bot. cab. t. 320.
<i>Anthéricum semibarbatum. B.P.</i>						
4	australis. s.s. (<i>ye.</i>) <i>bulbous.</i>	_____	1819.	_____	D.G.b.	
<i>Anthéricum bulbosum. B.P.</i>						
5	ánnua. s.s. (<i>ye.</i>) <i>annual.</i>	C. B. S.	1731.	5. 6.	H. ⊙.	Bot. mag. t. 1451.
<i>Anthéricum ánnuum. B.M.</i>						
6	incurva. R.S. (<i>ye.</i>) <i>incurved-leaved.</i>	_____	1822.	_____	D.G. ¼.	
7	longiscapa. s.s. (<i>ye.</i>) <i>glaucous-leaved.</i>	_____	1759.	4. 8.	D.G.b.	Bot. mag. t. 1339.
8	favosa. R.S. (<i>ye.</i>) <i>conical-rooted.</i>	_____	1824.	_____	D.G. ¼.	
9	asphodeloides. s.s. (<i>ye.</i>) <i>upright-leaved.</i>	_____	1759.	6. 8.	D.G. ½.	Jacq. vind. t. 181.
10	rostrata. s.s. (<i>ye.</i>) <i>glaucous shrubby.</i>	_____	1812.	3. 8.	D.G. ½.	Jacq. ic. 2. t. 403.
11	frutescens. w. (<i>ye.</i>) <i>shrubby.</i>	_____	1702.	_____	D.G. ½.	Bot. mag. t. 816.
12	pugioniformis. s.s. (<i>ye.</i>) <i>dagger-leaved.</i>	_____	1793.	4. 6.	D.G.b.	t. 1454.
13	bisulcata. H.P. (<i>ye.</i>) <i>two-channelled.</i>	_____	1825.	_____	D.G.b.	
14	triquetra. s.s. (<i>wh.</i>) <i>triquetrous.</i>	_____	_____	_____	D.G. ¼.	
15	scabra. s.s. (<i>wh.</i>) <i>rough.</i>	_____	_____	_____	D.G. ¼.	
16	dubia. R.S. (<i>wh.</i>) <i>doubtful.</i>	_____	1822.	_____	D.G. ¼.	
17	ciliata. s.s. (<i>wh.re.</i>) <i>fringed.</i>	_____	_____	_____	D.G. ¼.	
18	cauda-felis. R.S. (<i>wh.</i>) <i>Cats-tail.</i>	_____	1825.	_____	D.G. ¼.	
19	planifolia. R.S. (<i>ro.w.</i>) <i>two-coloured.</i>	S. Europe.	1824.	_____	H. ¼.	Desf. atl. 1. t. 90.
<i>Phalángium bicolor. Redout. liliac. t. 215.</i>						
20	mesembryanthoides. H.P. (<i>ye.</i>) <i>Figwort-like.</i>	C. B. S.	1824.	_____	D.G. ¼.	
21	crispa. R.S. (<i>ye.</i>) <i>curled-leaved.</i>	_____	1825.	_____	D.G. ¼.	
22	præmorsa. s.s. (<i>ye.</i>) <i>bitten-rooted.</i>	_____	1816.	_____	D.G.b.	Jacq. ic. 2. t. 406.
23	graminea. H.R. (<i>ye.</i>) <i>Grass-leaved.</i>	_____	1819.	6. 8.	D.G.b.	
24	alooides. R.S. (<i>ye.</i>) <i>Aloe-leaved.</i>	_____	1732.	4. 8.	D.G.b.	Bot. mag. t. 1317.
<i>Anthéricum aloides. B.M.</i>						
25	nutans. s.s. (<i>st.gr.</i>) <i>nodding.</i>	_____	1812.	_____	D.G. ¼.	Jacq. ic. 2. t. 407.
26	latifolia. s.s. (<i>ye.gr.</i>) <i>broad-leaved.</i>	_____	_____	_____	D.G. ¼.	2. t. 408.
ARTHROP'ODIUM. B.P. <i>ARTHROP'ODIUM.</i> Hexandria Monogynia. S.S.						
1	cirrhatum. B.R. (<i>wh.</i>) <i>large-flowered.</i>	N. Zealand.	1819.	5. 8.	F. ¼.	Bot. reg. t. 709.
2	paniculatum. B.P. (<i>w.pu.</i>) <i>panicked.</i>	N. S. W.	1800.	6. 9.	F. ¼.	Bot. mag. t. 1421.
<i>minus.</i> Bot. reg. 866. non B.P. <i>Anthéricum milleflorum.</i> Redout. liliac. t. 58.						
3	minus. B.P. (<i>wh.pu.</i>) <i>simple-racemed.</i>	_____	1819.	_____	F. ¼.	
4	pendulum. s.s. (<i>vi.</i>) <i>pendulous-flower'd.</i> N. Holland.	1817.	_____	_____	F. ¼.	
<i>Anthéricum pendulum. W.en.</i>						
5	strictum. B.P. (<i>wh.pu.</i>) <i>upright.</i>	V. Diem. Isl.	1826.	_____	G. ¼.	
6	fimbriatum. B.P. (<i>wh.</i>) <i>fringe-flowered.</i>	N. S. W.	1825.	_____	F. ¼.	
STYPA'NDRA. B.P. <i>STYPA'NDRA.</i> Hexandria Monogynia.						
1	glauca. B.P. (<i>bl.</i>) <i>glaucous.</i>	N. S. W.	1819.	4. 6.	F. ¼.	
2	cæspitosa. B.P. (<i>bl.</i>) <i>turfy.</i>	_____	1824.	_____	F. ¼.	
3	umbellata. B.P. (<i>bl.</i>) <i>umbel-flowered.</i>	_____	1820.	_____	F. ¼.	
CÆSIA. B.P. <i>CÆSIA.</i> Hexandria Monogynia. S.S.						
<i>vittata.</i> B.P. (<i>wh.</i>) <i>nodding-flowered.</i> N. S. W.						
TRICO'RYNE. B.P.	TRICO'RYNE. Hexandria Monogynia. S.S.		1816.	7. 8.	F. ¼.	
1	simplex. B.P. (<i>ye.</i>) <i>simple-stalked.</i>	N. S. W.	1819.	7. 9.	H. ⊙.	
2	elatior. B.P. (<i>ye.</i>) <i>tall.</i>	_____	1825.	_____	G. ¼.	
CHLOROPHYTUM. B.P. <i>CHLOROPHYTUM.</i> Hexandria Monogynia. S.S.						
1	inornatum. B.M. (<i>gr.</i>) <i>green-flowered.</i>	Sierra Leone.	6. 8.	S. ¼.	Bot. mag. t. 1071.
2	orchidatum. B.R. (<i>wh.</i>) <i>Orchis-like.</i>	_____	1822.	1. 12.	S. ¼.	Bot. reg. t. 813.
3	elatum. B.P. (<i>wh.</i>) <i>tall.</i>	C. B. S.	1751.	8. 9.	G. ¼.	Redout. lil. t. 191.
<i>Anthéricum elatum. h.k. Phalángium elatum. R.L.</i>						
PUSCHK'INIA. M.B. <i>PUSCHK'INIA.</i> Hexandria Monogynia. M.B.						
<i>scilloides.</i> M.B. (<i>w.bl.</i>) <i>Squill-like.</i> Iberia.						
<i>Adámsia scilloides.</i> Lindl. coll. bot. t. 24.						
ALBUCA. Ker.	ALBUCA. Hexandria Monogynia. L.					
1	altissima. w. (<i>wh.gr.</i>) <i>tall.</i>	C. B. S.	1780.	4. 5.	F.b.	Jacq. ic. 1. t. 63.
2	cornuta. R.L. (<i>wh.gr.</i>) <i>three-horned.</i>	_____	1812.	_____	F.b.	Redout. liliac. t. 70.
3	major. w. (<i>ye.gr.wh.</i>) <i>great.</i>	_____	1759.	_____	F.b.	Bot. mag. t. 804.

4 mīnor. w. (<i>ye.gr.</i>) small.	C. B. S.	1768.	5. 6. F.b.	Bot. mag. t. 720.
5 flāccida. w. (<i>ye.gr.</i>) flaccid.	1791.	— F.b.	Jacq. ic. 2. t. 442.	
6 viridiflōra. w. (<i>ye.gr.</i>) green-flowered.	1794.	6. 7. F.b.	Bot. mag. t. 1656.	
7 coarctāta. w. (<i>ye.</i>) channel-leaved.	1774.	5. 6. F.b.	Andr. reposit. t. 450.	
8 fastigiāta. w. (<i>wh.</i>) level-topped.	—	— F.b.	Jacq. ic. 2. t. 442.	
9 caudāta. w. (<i>wh.gr.</i>) upright-flowered.	1791.	5. 7. F.b.	Bot. mag. t. 1481.	
10 setōsa. w. (<i>gr.ye.</i>) bristly-rooted.	1795.	— F.b.	Jacq. ic. 2. t. 441.	
11 aurēa. w. (<i>ye.gr.</i>) golden-flowered.	1825.	— F.b.	Redout. liliac. t. 195.	
12 abyssinica. w. (<i>w.y.g.</i>) African.	—	— F.b.	Jacq. schœn. 1. t. 81.	
13 frāgrans. w. (<i>ye.gr.</i>) sweet-scented.	1791.	6. 7. F.b.	Jacq. ic. 2. t. 445.	
14 viscōsa. w. (<i>w.y.g.</i>) clammy-leaved.	1779.	5. 6. F.b.	— 2. t. 439.	
15 spirālis. w. (<i>st.gr.</i>) spiral-leaved.	1795.	6. F.b.	Bot. mag. t. 1329.	
16 vittāta. B.M. (<i>ye.gr.</i>) ribbon.	1802.	7. 8. F.b.	Bot. mag. t. 1046.	
17 parviflōra. R.S. (<i>ye.gr.</i>) small-flowered.	1812.	— F.b.		
18 physodes. B.M. (<i>w.pu.</i>) dingy-flowered.	1804.	6. 7. F.b.		
19 nematōdes. (<i>w.pu.</i>) slender-leaved.	1812.	— F.b.		
<i>Anthéricum nematōdes.</i> R.S.				
20 exuvīāta. B.M. (<i>w.pu.</i>) Adder skin.	1795.	5. 7. F.b.	Bot. mag. t. 871.	
<i>Anthéricum exuvīatum.</i> Jacq. ic. 2. t. 415.				
21 anthericōides. (<i>ye.gr.</i>) Anthericum-like.	1788.	8. F.b.		
<i>Anthéricum abuceoides.</i> w.				
22 filiflōra. Ker. (<i>wh.gr.</i>) filiform-leaved.	1825.	7. 8. F.b.	Jacq. ic. 2. t. 414.	
23 fūgax. B.R. (<i>wh.pu.</i>) fugacious.	1795.	4. 6. F.b.	Bot. reg. t. 311.	
<i>Anthéricum frāgrans.</i> Jacq. schœn. 1. t. 86.				
ORNITHO'GALUM. L. STAR OF BETHLEHEM.	Hexandria Monogynia. L.			
1? uniflōrum. w. (<i>ye.</i>) one-flowered.	Siberia.	1781.	5. 6. H.b.	Nov.act.pet.18.t.6.f.3.
2 ixioïdes. H.K. (<i>wh.</i>) Ixia-like.	California.	1796.	— H.b.	Swt. br. fl. gar. s.2.t.42.
3 montānum. T.N. (<i>w.g.</i>) mountain.	Naples.	1825.	— H.b.	Eng. bot. v. 2. t. 130.
4 collinum. R.S. (<i>wh.gr.</i>) hill.	Sicily.	1829.	— H.b.	Swt. br. fl. gar. s.2.t.58.
5 umbellatūm. E.B. (<i>w.g.</i>) common.	England.	4. 6. H.b.	Ten. fl. nap. v. 1. t. 34.
6 refractūm. W.en. (<i>w.g.</i>) reflexed.	S. Europe.	1825.	— H.b.	Swt. br. fl. gar. s.2.
7 tenuifolium. R.S. (<i>w.g.</i>) slender-leaved.	Sicily.	1830.	— H.b.	Jacq. ic. 2. t. 426.
8 excāpum. T.N. (<i>w.g.</i>) short-stalked.	Naples.	1823.	— H.b.	— 2. t. 428.
9 fimbriātum. M.B. (<i>w.g.</i>) fringed.	Tauria.	1820.	2. 3. H.b.	Bot. reg. t. 235.
10 comōsum. w. (<i>wh.gr.</i>) short-spiked.	Austria.	1596.	6. 8. H.b.	Jacq. coll. 2. t. 18. f. 13.
11 cōnicum. w. (<i>wh.</i>) conical.	C. B. S.	1818.	— F.b.	
12 nīveum. w. (<i>wh.</i>) snowy.	—	1774.	— F.b.	
13 ovātum. w. (<i>wh.gr.</i>) oval-leaved.	—	1816.	4. 8. F.b.	
14 notātum. R.S. (<i>ye.br.</i>) brown-marked.	—	1825.	6. 8. F.b.	
<i>maculātum.</i> Jacq. non Thunb.				
15 ciliātum. w. (<i>wh.</i>) ciliated.	—	1818.	4. 6. F.b.	
16 crenulātum. w. (<i>st.</i>) crenulate.	—	1825.	6. 8. F.b.	
17 maculātum. R.S. (<i>sn.br.</i>) spotted-flowered.	—	—	F.b.	Jacq. ic. 2. t. 433.
18 secūndum. R.S. (<i>st.br.</i>) one-ranked.	—	1816.	— F.b.	
19 graminifolium. R.S. (<i>w.g.</i>) grass-leaved.	—	—	F.b.	Jacq. ic. 2. t. 427.
20 ténellum. w. (<i>st.gr.</i>) slender.	—	1816.	5. 8. F.b.	
21 pilōsum. w. (<i>wh.</i>) hairy-leaved.	—	—	F.b.	
22 revolutūm. w. (<i>wh.gr.</i>) revolute-flowered.	—	1795.	4. 8. F.b.	Bot. reg. t. 315.
23 corymbosum. B.R. (<i>wh.</i>) corymbose.	Peru.	1822.	— H.b.	— 906.
24 Rudolphi. R.S. (<i>wh.</i>) Rudolph's.	C. B. S.	1812.	— F.b.	Jacq. ecl. t. 20.
<i>tenuifolium.</i> Redout. liliac. t. 312. non Guss.				
25 prasiūm. B.R. (<i>gr.</i>) green-flowered.	—	1816.	6. 7. F.b.	Bot. reg. t. 158.
26 vīrens. B.R. (<i>gr.</i>) greenish.	—	1823.	— F.b.	— 814.
27 suavēolens. w. (<i>gr.ye.</i>) fragrant.	—	1816.	— F.b.	Jacq. ic. 2. t. 431.
28 barbātum. w. (<i>ye.gr.</i>) bearded.	—	1795.	5. 7. F.b.	Jacq. schœn. 1. t. 91.
29 polyphyllum. w. (<i>g.y.</i>) many-leaved.	—	1825.	— F.b.	Jacq. ic. 2. t. 430.
30 rupēstre. w. (<i>wh.gr.</i>) rock.	—	1795.	5. 8. F.b.	Bot. mag. t. 792.
31 juncifolium. w. (<i>w.pu.</i>) Rush-leaved.	—	1794.	7. 8. F.b.	Jacq. ic. 2. t. 429.
32 fuscātum. w. (<i>br.gr.</i>) brown-flowered.	—	1825.	4. 6. F.b.	Andr. reposit. t. 260.
33 odorātum. w. (<i>gr.ye.</i>) sweet-scented.	—	1795.	5. 6. F.b.	Jacq. ic. 2. t. 425.
34 elātum. A.R. (<i>wh.</i>) Egyptian.	Egypt.	1804.	6. 7. H.b.	Bot. mag. t. 876.
35 pyramidāle. w. (<i>wh.</i>) pyramidal.	Spain.	1752.	— H.b.	Jacq. vind. 1. t. 29.
36 latifolium. w. (<i>wh.</i>) broad-leaved.	Egypt.	1629.	— H.b.	Bot. mag. t. 805.
37 longibracteātum. w. (<i>w.</i>) long-bracted.	C. B. S.	1812.	7. 12. F.b.	Jacq. schœn. 1. t. 888.
38 caudātum. w. (<i>wh.gr.</i>) long-spiked.	—	1774.	2. 8. F.b.	Redout. liliac. t. 417.
39 scilloides. w. (<i>wh.gr.</i>) Squill-like.	—	1795.	6. 7. F.b.	Bot. mag. t. 2510.
40 trig'num. R.L. (<i>wh.gr.</i>) three styled.	1825.	— F.b.	Eng. bot. v. 7. t. 499.
41 narbonése. w. (<i>w.g.</i>) Narbonne.	S. Europe.	1810.	7. 8. H.b.	
42 pyrenāicum. w. (<i>st.</i>) spiked.	England.	6. 7. H.b.	

43	sulfureum. R.S.	(st.) straw-coloured. Hungary.	1818.	6. 10. H.b.	Bot. mag. t. 2623.
	<i>Anthéricum sulphureum.</i> Walds. et Kit. pl. hung. 1.	t. 95. <i>Phalángium sulfureum.</i> Poiret.			Renealm. spec. t. 90.
44	stachyoïdes. w.	(w.g.) close-spiked. S. Europe.	1771.	4. 7. H.b.	Bot. mag. t. 1134.
45	láctea. w.	(st.gr.) milk white. C. B. S.	1796.	6. 7. F.b.	Eng. bot. v. 28. t. 1997.
46	nútans. w.	(wh.gr.) nodding. Britain.	4. 5. H.b.	Jacq. ic. 2. t. 435.
47	coarctatum. w.	(wh.) close-flowered. C. B. S.	1804.	6. 7. F.b.	Jacq. ic. 2. t. 190.
48	aureum. w.	(go.) golden. ———	1790.	— F.b.	Jacq. ic. 2. t. 436.
49	flavissimum. Jacq. (ye.) great yellow. ———	—	—	— F.b.	— t. 438.
50	miniátum. Jacq. (ye.) stained. ———	—	—	— F.b.	Bot. mag. t. 1164.
51	thyroïdes. w. (wh.br.) thyrs-flowered. ———	1757.	—	— F.b.	Bot. reg. t. 316.
	α álbum. (wh.br.) white-flowered. ———	—	—	— F.b.	— t. 305.
52	β flavescens. (st.br.) pale-yellow. ———	—	—	— F.b.	
53	bicolor. R.S.	(wh.br.) two-coloured. ———	1816.	— F.b.	
54	tuberòsum. R.S.	(ye.) large-rooted. ———	1787.	— F.b.	Bot. mag. t. 728.
55	arábicum. w.	(wh.) great-flowered. Egypt.	1629.	3. 4. F.b.	
	brachystachys. f. (wh.) short-spiked. Dahuria.	1821.	—	— H.b.	
56	Bérgii. R.S.	(wh.gr.) Bergius's. ———	1816.	— F.b.	
57	marítimum. Lam. (wh.) officinal Squill. S. Europe.	1629.	4. 5. F.b.	Bot. mag. t. 918.	
	<i>Squilla.</i> B.M. <i>Scilla marítima.</i> w.				
58	concínum. S.P.	(wh.) neat. Portugal.	1797.	— H.b.	Bot. mag. t. 953.
59	unifòlium. B.M.	(wh.) one-leaved. Gibraltar.	1805.	6. 7. F.b.	— t. 935.
G'AGEA. S.A.B.	G'AGEA.	Hexandria Monogynia.			
1	fascicularis. s.a.b.	(ye.) bunch-rooted. Britain.	3. 4. H.b.	Bot. mag. t. 1200.
	lutea. B.M. 1200.	<i>Ornithógalum lúteum.</i> Eng. bot. v. 1. t. 21. non L.			
2	glaúca. B.F.G.	(ye.) glaucous-leaved. Switzerland.	1825.	—	Swt. br. fl. gar. t. 177.
3	bracteoláris. s.a.b. (ye.) sheathing-bracted.	—	— H.b.	— t. 158.
	<i>Ornithógalum lúteum.</i> L. non E.B. v. s. in herb. linn.				
4	mínima. R.S.	(ye.) corn-field. Europe.	1825.	— H.b.	Swt. br. fl. gar. s.2.t.29.
	stelláris. s.a.b.	<i>Ornithógalum arvénse.</i> s.s. <i>Sternbergii.</i> Hop. v. s. in herb. linn.			
5	spathacea. R.S.	(ye.) sheathed. Germany.	1759.	5. H.b.	Flor. græc. t. 331.
6	reticulata. R.S.	(ye.) netted-sheathed. Tauria.	1829.	— H.b.	Pall. it. 3. ap. t. D. f. 2.
7	pusilla. R.S.	(ye.) small. Bohemia.	1823.	— H.b.	Reichenb. ic. t. 117.
8	Clusiána. R.S.	(ye.) Clusius's.	S. Europe.	
9	podólica. R.S.	(ye.) Polish.	Poland.	1827.	— H.b.
10	erubescens. R.S.	(y.bh.) blush.	—	— H.b.	
11	Liotárdi. R.S.	(ye.) fistulous-leaved. S. Europe.	1825.	— H.b.	Sternb. denks. 2. t. 3.
	<i>Ornithógalum fistulòsum.</i> Redout. liliac. t. 221.				
12	pygmæa. R.S.	(ye.) pigmy.	Spain.	1826.	— H.b.
13	arvénse. R.S.	(ye.) villous-stalked. Europe.	— H.b.	Flor. græc. t. 332.
	<i>Ornithógalum villòsum.</i> M.B.				
14	bohémicum. R.S. (ye.g.)	Bohemian.	Bohemia.	1828.	— H.b.
	saxatilis. R.S.	(gr.ye.) rock.	Germany.	—	Sturm. flor. t. 23.
16	bulbifera. R.S.	(ye.g.) bulbiferous.	Tauria.	1829.	— H.b.
BARNA'R'DIA. B.R.	BARNA'R'DIA.	Hexandria Monogynia.			
1	scilloides. B.R.	(ro.) Squill-like.	China.	1824.	9. 10. H.b.
2	japónica. R.S.	(pu.) Japan.	Japan.	1821.	6. 8. H.b.
	<i>Onithógalum japonicum.</i> w.				
LLO'YDIA. Sal.	LLO'YDIA.	Hexandria Monogynia.			
1	serotina.	(wh.std.) mountain.	Wales.	6. 7. H.b.
	<i>Anthéricum serotinum.</i> E.B. <i>Phalángium serotinum.</i> Redout. liliac. t. 270.				Eng. bot. v. 12. t. 793.
2	striata.	(wh.std.) stripe-flowered.	Siberia.	1789.	5. 7. H.b.
	<i>Ornithógalum striatum.</i> s.s.				<i>Gagea serotina.</i> Ker.
SCI'LLA. W.	SCI'LLA.	Hexandria Monogynia. L.			
1	non scripta. L.en. (bl.)	Harebells.	Britain.	3. 6. H.b.
	<i>nútans.</i> E.B. <i>Hyacinthus non scriptus.</i> L.				Eng. bot. v. 6. t. 377.
2	cérnua. R.S.	(ro.) flesh-coloured.	Spain.	— H.b.
3	pátula. R.S.	(bl.) spreading.	S. Europe.	1633.	4. 6. H.b.
	<i>campanulata.</i> B.M. t. 1102. non t. 128.				Redout. liliac. t. 225.
4	campanulata. w.	(bl.) bell-flowered.	Spain.	—	5. 6. H.b.
5	corymbosa. s.s.	(ro.) Cape.	C. B. S.	1793.	8. 12. F.b.
	<i>Massónia corymbosa.</i> B.M. <i>Hyacinthus corymbosus.</i> L.				Bot. mag. t. 128.
6	brevifolia. R.S.	(wh.) short-leaved.	—	1822.	1. 3. F.b.
7	brachyphylla. R.S. (bh.)	blush.	—	1811.	— F.b.
	<i>Scilla brevifolia.</i> B.M. non Thunb.				Bot. mag. t. 1468.
8	sibirica. H.K.	(bl.) Siberian.	Siberia.	1796.	2. 3. H.b.
9	am'œnula. R.S.	(bl.) pretty.	Russia.	1822.	3. 4. H.b.
10	am'œna. R.S.	(bl.) nodding.	Levant.	1596.	— H.b.
11	pùmila. R.S.	(pa.bl.) small.	Portugal.	1819.	4. 5. H.b.
	<i>monophyllos.</i> Link ex R.S.				Andr. rep. t. 365.

12 bifòlia. w.	(va.) two-leaved.	England.	2. 4.	H.b.	Eng. bot. v. 1. t. 24.
α cærulea.	(bl.) blue-flowered.	—	—	H.b.	
β álba.	(wh.) white-flowered.	—	—	H.b.	
13 cárnea.	(ro.) rosy.	—	—	—	H.b.	
14 præcox. w.	(bl.) early-flowering.	1790.	—	5. 7.	H.b.	
15 Lílio-hyacínthus.w.	(bl.) Lily-rooted.	S. Europe.	1597.	—	H.b.	
16 vérra. w.	(bl.) vernal.	Britain.	4. 5.	H.b.	Redout. lil. t. 205.
17 umbellàta. s.s.	(bl.) umbel-flowered.	Pyrenees.	1817.	—	H.b.	Eng. bot. v. 1. t. 23.
18 itálica. s.s.	(bl.) Italian.	Italy.	1605.	4. 7.	H.b.	Redout. liliac. t. 166.
19 autumnális. w.	(pu.) autumnal.	England.	8.	H.b.	Bot. mag. t. 663.
20 intermèdia. R.S.	(bl.) intermediate.	Sicily.	1829.	—	H.b.	Eng. bot. v. 2. t. 78.
21 obtusifòlia. R.S. (vi.bl.)	blunt-leaved.	Barbary.	1828.	—	H.b.	Redout liliac. 4. t. 190.
22 undulàta. R.S.	(li.) wave-leaved.	S. Europe.	—	—	H.b.	Desf. atl. 1. t. 47.
23 lingulàta. R.S.	(bl.) tongue-leaved.	Barbary.	—	—	H.b.	Redout liliac. t. 321.
24 vincentína. R.S.	(bl.) spreading-flowered.	Portugal.	1827.	—	H.b.	
25 peruviana. w.	(va.) corymbose.	Spain.	1607.	5.	H.b.	Bot. mag. t. 749.
α cærulea.	(bl.) blue-flowered.	—	—	—	H.b.	— t. 749.
β álba.	(wh.) white-flowered.	—	—	—	H.b.	
26 Cupaniána. R.S.	(bl.) Cupani's.	Sicily.	1826.	—	H.b.	Cup.panph.sic.5.t.20.1.
27 lusitánica. B.M. (pa.bl.)	Portugal.	Portugal.	1777.	—	H.b.	Bot. mag. t. 1999.
28 hyacinthóides. w. (bl.)	Hyacinth-like.	Madeira.	1585.	8.	H.b.	— t. 1140.
29 esculénta. B.M. (pa.bl.)	Quamash.	N. America.	1811.	5. 7.	H.b.	— t. 1574.
<i>Phalángium Quamash. Ph.</i>						
HYACINTHUS. S.S.						
<i>Hyacinth. Hexandria Monogynia. L.</i>						
1 orientális. w.	(bl.) common.	Levant.	1596.	3. 5.	H.b.	Bot. reg. t. 995.
α cæruleus.	(bl.) blue-flowered.	—	—	—	H.b.	
β lýricus.	(bl.) Lyra grandis.	—	—	H.b.	
γ rüber.	(re.) red-flowered.	—	1596.	—	H.b.	
δ flávus.	(ye.) yellow-flowered.	—	—	—	H.b.	
ε álbus.	(wh.) white-flowered.	—	—	—	H.b.	
ζ semiplénus.	(va.) semidouble.	—	—	—	H.b.	
η plénus.	(va.) double-flowered.	—	—	—	H.b.	
θ mítplex.	(va.) full-flowered.	—	—	—	H.b.	
ι atrocæruleus.	(bl.) Pourpre fonce.	—	—	—	H.b.	
κ sceptrifòrmis. (pa.bl.)	Porcelaine sceptre.	—	—	—	H.b.	
λ sanguinoléntus. (re.)	Waterloo.	—	—	—	H.b.	
μ ornatus.	(pi.) L'honneur d'Amsterdam.	—	—	—	H.b.	
ν ranunculifòrus. (ye.)	yellow Ophir.	—	—	—	H.b.	
2 amethystinus. w. (bl.)	Amethyst-coloured.	S. Europe.	1759.	4. 5.	H.b.	
UROPETALON. B.R. UROPE'TALON.						
<i>Hexandria Monogynia. B.R.</i>						
1 viride. B.R.	(gr.) green-flowered.	C. B. S.	1774.	8.	F.b.	
<i>Hyacinthus víridis. L. Zucágna víridis. Th. Lachenalia víridis. w.</i>						
2 glaucum. B.R. (gr.br.)	glaucous-leaved.	C. B. S.	1816.	7. 8.	F.b.	
3 críspum. B.R. (br.gr.)	curled-leaved.	—	—	—	F.b.	
4 longifòlium. B.R. (br.gr.)	long-leaved.	—	1822.	—	F.b.	
5 hyacinthóides. R.S. (re.)	Hyacinth-like.	—	—	—	F.b.	
6 serotinum. B.R. (bl.)	late-flowering.	Spain.	1629.	6. 8.	F.b.	
7 fúlvum. H.S.L. (re.bl.)	tile-red.	Mogador.	1808.	6. 9.	F.b.	
<i>Scilla serotina. β. B.M. Hyacinthus fúlvus. Cav.</i>						
BELLEV'ALIA. Lap. BELLEV'ALIA.						
<i>Hexandria Monogynia.</i>						
româna.	(wh.) Roman.	Italy.	1596.	5.	H.b.	Bot. mag. t. 939.
<i>operculata. Lap. Scilla româna. B.M. Hyacinthus românum. L.</i>						
MUS'CARI. S.S.						
<i>GRAPE HYACINTH. Hexandria Monogynia. S.S.</i>						
1 moschátum. s.s.	(br.) musk.	Levant.	1596.	4. 5.	H.b.	
2 ambrosiacum. R.L.	(wh.) strong-scented.	—	—	H.b.	Bot. mag. t. 734.
3 macrocarpum. B.F.G.	(ye.) large-capsuled.	Turkey.	1812.	—	H.b.	Redout. liliac. t. 132.
<i>moschátum β. fárum. Bot. mag. t. 1565.</i>						
4 comósium. s.s. (br.pu.)	purple.	S. Europe.	1596.	—	H.b.	
β monstròsum.	(pu.) feathered.	—	—	—	H.b.	
5 ciliátum. s.s.	(w.pu.) ciliated.	Tauria.	1818.	—	H.b.	
6 pállens. s.s.	(pa.bl.) pale-flowered.	—	—	—	H.b.	
7 glaucum. B.R.	(pa.) glaucous-leaved.	Persia.	1825.	—	H.b.	
8 racemósium. s.s.	(bl.) starch.	Britain.	—	H.b.	
9 commutátum. R.S. (d.bl.)	confused.	Sicily.	—	H.b.	
10 parvifòrum. R.S. (bl.)	small-flowered.	—	1827.	—	H.b.	
11 botryóides. s.s.	(va.) bunch-flower'd.	Italy.	1596.	—	H.b.	
α azùreum. B.F.G.	(bl.) bright-blue.	—	—	—	H.b.	
β pálidum. B.F.G.	(pa.) pale blue.	—	—	—	H.b.	
γ álbum. B.F.G.	(wh.) white-flowered.	—	—	—	H.b.	

LACHEN'ALIA. W.	LACHEN'ALIA.	Hexandria Monogynia. W.	C. B. S.	1774.	3. 4.	F.b.	
1 pénula. w. (<i>y.re.g.</i>) pendulous.	<i>pendulosa.</i>					F.b.	Jacq. ic. 2. t. 400.
<i>a viridifolia.</i> (<i>y.re.g.</i>) green-leaved.						F.b.	Bot. mag. t. 590.
β maculata. (<i>y.re.g.</i>) spotted-leaved.						F.b.	
2 quadrícolor. B.M. (<i>va.</i>) four-coloured.				1780.	3. 5.	F.b.	Jacq. ic. 2. t. 388.
<i>a viridifolia.</i> (<i>y.sc.g.</i>) green-leaved.						F.b.	Bot. mag. t. 588.
β colorata. (<i>y.sc.g.</i>) coloured-leaved.						F.b.	—— t. 1097.
3 lutéola. Jacq. (<i>ye.</i>) yellow-flowered.				1774.	4. 5.	F.b.	Jacq. ic. 2. t. 395.
<i>a viridifolia.</i> (<i>ye.</i>) green-leaved.						F.b.	Bot. mag. t. 1020.
β maculata. (<i>ye.</i>) spotted-leaved.						F.b.	—— t. 1704.
4 tricolor. B.M. (<i>y.re.g.</i>) three-coloured.						F.b.	—— t. 82.
5 rúbida. w. (<i>re.</i>) dotted-flowered.				1803.	9. 10.	F.b.	—— t. 993.
6 punctata. s.s. (<i>w.re.</i>) spotted-flowered.				1825.	6. 8.	F.b.	Jacq. ic. 2. t. 397.
7 flava. A.R. (<i>ye.</i>) bright-yellow.				1805.	5. 6.	F.b.	Andr. deposit. t. 77.
8 isopétala. w. (<i>pa.br.</i>) equal-flowered.				1804.		F.b.	Jacq. ic. 2. t. 401.
9 hírtia. R.S. (<i>w.bl.</i>) hairy-leaved.				1825.		F.b.	
10 sessiliflora. A.R. (<i>bh.</i>) sessile-flowered.				1804.		F.b.	Andr. rep. t. 460.
11 unifòlia. w. (<i>bl.re.</i>) one-leaved.				1795.	3. 4.	F.b.	Bot. mag. t. 766.
12 rósea. A.R. (<i>ro.</i>) rose-coloured.				1800.	4. 5.	F.b.	Andr. rep. t. 296.
13 pusilla. w. (<i>wh.</i>) dwarf.				1825.		F.b.	Jacq. ic. 2. t. 385.
14 refléxia. R.S. (<i>ye.</i>) reflex-leaved.						F.b.	
15 bifòlia. B.M. (<i>ro.wh.</i>) cow-leaved.				1813.	3. 4.	F.b.	Bot. mag. t. 1611.
16 contáminata. w. (<i>w.br.</i>) contaminated.				1774.	2. 4.	F.b.	—— t. 1401.
17 purpúrea. w. (<i>pu.w.</i>) purple.				1819.	4. 5.	F.b.	Jacq. ic. 2. t. 393.
18 violacea. w. (<i>g.vi.</i>) violet.				1795.	3. 4.	F.b.	—— t. 394.
19 lúcida. B.M. (<i>w.br.</i>) glossy-leaved.				1798.	3. 5.	F.b.	Bot. mag. t. 1372.
20 racemosa. B.M. (<i>wh.</i>) racemed.				1811.	5.	F.b.	—— t. 1517.
21 frágans. w. (<i>wh.</i>) sweet-scented.				1798.	3. 5.	F.b.	Jacq. schœn. t. 82.
22 unicolor. B.M. (<i>li.</i>) self-coloured.				1806.	5. 6.	F.b.	Bot. mag. t. 1373.
23 anguinea. S.F.G. (<i>w.g.</i>) snake.				1825.	4. 5.	F.b.	Swt. br. fl. gar. t. 179.
24 purpureo-cerulea. (<i>bl.</i>) purple-blue.				1798.		F.b.	Bot. mag. t. 745.
25 pustulata. w. (<i>w.st.</i>) blistered.				1790.	1. 4.	F.b.	—— t. 817.
26 liliiflora. w. (<i>wh.</i>) Lily-flowered.				1820.	4. 6.	F.b.	Jacq. ic. 2. t. 387.
27 orthopetalala. R.S. (<i>w.gr.</i>) wedge-flowered.				1825.		F.b.	—— 2. t. 383.
28 pátila. w. (<i>wh.ro.</i>) spreading-flower'd.				1795.	4. 5.	F.b.	—— 2. t. 384.
29 nervosa. B.M. (<i>br.pu.</i>) nerved-leaved.				1810.	6.	F.b.	Bot. mag. t. 1497.
30 angustifolia. w. (<i>wh.</i>) narrow-leaved.				1793.	4. 5.	F.b.	—— t. 735.
31 hyacinthoides. w. (<i>wh.</i>) Hyacinth-flower'd.				1812.	3. 4.	F.b.	Jacq. ic. 2. t. 382.
32 pálida. w. (<i>pa.bl.</i>) pale-flowered.				1782.		F.b.	Bot. reg. t. 314.
33 mediána. R.S. (<i>st.gr.</i>) intermediate.						F.b.	Jacq. ic. t. 392.
34 mutabilis. S.F.G. (<i>ye.</i>) changeable.				1825.	2. 5.	F.b.	Swt. br. fl. gar. ser. 2.
35 orchioides. w. (<i>ye.</i>) Orchis-like.				1752.	2. 4.	F.b.	Bot. mag. t. 1269.
36 glauca. w. (<i>wh.bl.</i>) sea-green.				1795.	3. 4.	F.b.	Jacq. ic. 2. t. 391.
DRI'MIA. S.S.	DRI'MIA.	Hexandria Monogynia. S.S.	C. B. S.	1774.	8. 10.	F.b.	
1 lanceolata. R.S. (<i>y.gr.</i>) spear-leaved.							Andr. deposit. t. 299.
<i>Lachenalia reflexa.</i> Andr. rep. t. 299.							
2 lanceæfòlia. B.M. (<i>g.pu.</i>) lance-leaved.				1800.	9. 10.	F.b.	Bot. mag. t. 643.
3 longipedunculata. (<i>g.pu.</i>) long-stalked.						F.b.	Schr. pl. rar. mon. t. 100.
4 acuminata. B.C. (<i>g.br.</i>) taper-leaved.						F.b.	Lodd. bot. cab. t. 1041.
5 Gawléri. R.S. (<i>gr.pu.</i>) revolute-flowered.				1774.		F.b.	Bot. mag. t. 1380.
<i>Hyacinthus revolutus.</i> H.K.							
6 ovalifolia. R.S. (<i>gr.pu.</i>) oval-leaved.				1812.		F.b.	Lodd. bot. cab. t. 278.
7 undulata. s.s. (<i>g.pu.</i>) wave-leaved.				1825.	8. 10.	F.b.	Jacq. ic. 2. t. 376.
8 purpurascens. s.s. (<i>pu.</i>) purplish.						F.b.	
9 ciliáris. s.s. (<i>g.pu.</i>) ciliated.					F.b.	Bot. mag. t. 1444.
10 pusilla. s.s. (<i>gr.</i>) dwarf.				1793.	5. 6.	F.b.	Jacq. ic. 2. t. 374.
11 media. s.s. (<i>w.pu.</i>) semicylindrical-l'd.				1818.	7. 8.	F.b.	—— 2. t. 375.
12 elata. s.s. (<i>g.w.</i>) tall.				1799.	10. 11.	F.b.	Bot. mag. t. 822.
13 altissima. s.s. (<i>wh.</i>) tallest.				1791.	8. 9.	F.b.	—— t. 1074.
ERIOSPERMUM. W.	ERIOSPERMUM.	Hexandria Monogynia. W.	C. B. S.	1795.	6. 8.	F.b.	
1 parvifolium. s.s. (<i>wh.</i>) small-leaved.				1800.		F.b.	Jacq. ic. 2. t. 422.
2 latifolium. s.s. (<i>wh.</i>) broad-leaved.						F.b.	—— 2. t. 420.
3 Bellendèni. (<i>wh.</i>) Mr. Ker's.						F.b.	Bot. mag. t. 1382.
<i>latifolium a.</i> B.M. non Jacq.							
4 lanceæfòlium. s.s. (<i>st.</i>) lance-leaved.				1795.		F.b.	Jacq. ic. 2. t. 421.
5 pubescens. s.s. (<i>wh.</i>) pubescent.				1820.		F.b.	Bot. reg. t. 578.
6 lanuginosum. s.s. (<i>wh.</i>) woolly-leaved.						F.b.	Jacq. schœn. 3. t. 265.
7 folioliferum. B.M. (<i>w.g.</i>) leaflet-bearing.				1806.		F.b.	Andr. deposit. t. 521.
8 paradoxicum. B.M. (<i>w.g.</i>) paradoxical.				1820.		F.b.	Jacq. col. sup. 81. t. 1.

MASS'ONIA. S.S.	MASS'ONIA. Hexandria Monogynia, W.						
1 grandiflora. B.R. (<i>wh.</i>) honey-bearing.	C. B. S.	1825.	1. 4.	F.b.			Bot. reg. t. 958.
2 cándida. B.T. (<i>wh.</i>) white-flowered.		1816.	2. 5.	F.b.			— t. 694.
longifolia β. cándida. B.R.							
3 longifolia. s.s. (<i>wh.</i>) long-leaved.							F.b.
4 latifolia. s.s. (<i>g.pu.</i>) broad-leaved.		1775.	3. 4.	F.b.			Bot. mag. t. 848.
5 cordata. s.s. (<i>wh.</i>) heart-leaved.		1820.					F.b.
6 muricata. s.s. (<i>wh.</i>) prickly-leaved.		1790.	4. 5.	F.b.			Bot. mag. t. 559.
7 scabra. H.K. (<i>wh.</i>) shagreen-leaved.			1. 4.	F.b.			Andr. deposit. t. 220.
pustulata. Bot. mag. t. 642.							
8 echinata. s.s. (<i>wh.</i>) rough-leaved.			5.	F.b.			
9 pauciflora. s.s. (<i>wh.</i>) few-flowered.							F.b.
10 angustifolia. s.s. (<i>wh.</i>) narrow-leaved.		1775.	3. 4.	F.b.			Bot. mag. t. 736.
11 undulata. s.s. (<i>wh.</i>) wave-leaved.		1791.	4.	F.b.			
12 ensifolia. s.s. (<i>pi.</i>) trumpet-flowered.		1790.	9. 2.	F.b.			Bot. mag. t. 554.
violacea. A.R. 46. Agapanthus ensifolius. w. Hyacinthus bifolius. Cav.							
EUCOMIS. S.S.	EUCOMIS. Hexandria Monogynia. W.						
1 nana. s.s. (<i>gr.</i>) dwarf.	C. B. S.	1774.	5.	F.b.			Bot. mag. t. 1495.
2 bifolia. s.s. (<i>gr.st.</i>) two-leaved.		1792.	4. 5.	F.b.			— t. 840.
3 purpureocaulis. s.s. (<i>gr.</i>) purple-stalked.		1794.	3. 4.	F.b.			Andr. deposit. t. 369.
4 régia. s.s. (<i>g.wh.</i>) tongue-leaved.		1702.	3. 5.	F.b.			Dill. elth. t. 92. f. 109.
5 undulata. s.s. (<i>g.wh.</i>) wave-leaved.		1760.					Bot. mag. t. 1083.
6 punctata. s.s. (<i>g.wh.</i>) spotted.		1783.	7.	F.b.			— t. 913.
7 striata. H.K. (<i>g.wh.</i>) streaked.		1790.	6. 12.	F.b.			— t. 1539.
A'LLIUM. L.	GARLIC. Hexandria Monogynia. L.						
1 sativum. G.D. (<i>re.</i>) common.	Sicily.	1548.	6. 9.	H.b.			Lobel. ic. t. 158.
2 Ophioscôrodon. G.D. (<i>br.</i>) Rocambole.	Crete.	1596.	8.	H.b.			Clus. his. p. 191. t. 210.
3 contravérsum. G.D. (<i>pu.</i>) simple-rooted.	Egypt.	1823.		H.b.			Cord. hist. p. 144. f. 1.
Ophioscôrodon. L.en.							
4 arenarium. G.D. (<i>re.</i>) sand.	Britain.	6. 8.	H.b.			Eng. bot. v. 19. t. 1358.
5 Scorodopràsum. G.D. (<i>re.</i>) wild Rocambole.	Denmark.	1596.	7.	H.b.			Flor. dan. t. 219. et 1455.
6 littoreum. G.D. (<i>re.</i>) Italian sand.	Italy.	1824.	8.	H.b.			
monspessulànum. W.							
7 vineale. G.D. (<i>ro.</i>) Crow.	Britain.	6. 8.	H.b.			Eng. bot. v. 28. t. 1974.
8 compáctum. G.D. (<i>ro.</i>) two-headed crow.	France.	1819.		H.b.			Moris. sect. 4. t. 15. f. 3.
9 Púrshii. G.D. (<i>da.pu.</i>) Pursh's crow.	N. America.	1824.		H.b.			
vineale. Ph. nec aliorum.							
10 proliferum. C.D. (<i>wh.</i>) tree Onion.	Egypt.	7.	H.b.			Bot. mag. t. 1469.
Cépa β bulbifera. B.M.							
11 Pórrum. G.D. (<i>gr.pu.</i>) Leek.	Switzerland.	1562.	5. 7.	H.b.			Plenck. ic. t. 421.
12 Ampelopràsum. G.D. (<i>pa.</i>) blue Leek.	England.	7. 8.	H.b.			Eng. bot. v. 24. t. 1657.
13 rotundum. G.D. (<i>pu.</i>) round-headed.	S. Europe.	1819.	6. 7.	H.b.			Clus. hist. p. 195. t. 220.
14 Hallérrii. G.D. (<i>pa.pu.</i>) Haller's.		1818.		H.b.			Flor. græc. t. 312.
15 rubicundum. G.D. (<i>re.</i>) marbled leek.	C. B. S.	7. 8.	F.b.			Bot. mag. t. 1560.
ampelopràsum β. B.M.							
16 Waldsteini. G.D. (<i>d.pu.</i>) Waldstein's.	Smyrna.	1824.	6. 7.	H.b.			W. et K. hung. t. 82.
17 exsertum. G.D. (<i>ro.</i>) long-stamen'd.	Russia.	1819.	7. 8.	H.b.			
18 erectum. G.D. (<i>wh.</i>) upright leek.	C. B. S.	1824.		F.b.			Moris. sect. 4. t. 15. f. 4.
19 acutifòrum. G.D. (<i>re.</i>) sharp-flowered.	France.	—		H.b.			
20 Synnotii. G.D. (<i>li.</i>) Synnot's.	C. B. S.	1825.	6. 7.	F.b.			
21 verrucosum. G.D. (<i>li.</i>) warted-flower'd.				F.b.			
22 multiflorum. G.D. (<i>vi.</i>) many-flowered.	Barbary.	1826.	7. 8.	F.b.			
23 descendens. G.D. (<i>d.pu.</i>) early flowering.	Switzerland.	1766.	6.	H.b.			Flor. græc. t. 316.
24 sphærocéphalon. G.D. (<i>pu.</i>) late-flowered.	Germany.	1759.	7.	H.b.			Bot. mag. t. 1764.
25 pruinatum. G.D. (<i>li.</i>) frosted.	Portugal.	1819.	6. 8.	H.b.			
26 carneum. G.D. (<i>li.</i>) blush-coloured.	France.	1816.	6. 7.	H.b.			Moris. sect. 4. t. 14. f. 4.
27 ascalonicum. G.D. (<i>li.</i>) Shallot.	Palestine.	1548.		H.b.			
β chinense. G.D. (<i>li.</i>) Chinese Shallot.	China.		H.b.			
28 fissile. (<i>gr.</i>) Scallion.	Asia minor.		H.b.			Matt. opus. p. 420. ic.
Cépa fissilis. Matt.							
29 Cépa. G.D. (<i>wh.gr.</i>) Onion.	6.	H. ♂. b.			Flor. græc. t. 326.
β aggregatum. G.D. (<i>w.g.</i>) Potatoe Onion.			H.b.			
30 Schœnopràsum. G.D. (<i>pu.</i>) Chives.	Britain.	5. 6.	H.b.			Eng. bot. v. 34. t. 2411.
31 foliòsum. G.D. (<i>pu.</i>) leafy Chives.	France.	1820.		H.b.			Redout. lit. 4. t. 214.
32 sibiricum. G.D. (<i>li.</i>) Siberian Chives.	Siberia.	1778.	6. 8.	H.b.			Bot. mag. t. 1141.
33 fistulòsum. G.D. (<i>gr.</i>) Welsh Onion.	—	1629.	6. 7.	H. ♂. b.			— t. 1230.
β alticum. Pallas. (<i>gr.</i>) Altaiic.	Altay.	1823.		H.b.			Pall. it. 2. n. 108. t. R.
34 cepæfórmæ. G.D. (<i>w.g.</i>) Onion-like.	1824.		H.b.			Bot. mag. t. 1408.
35 obliquum. G.D. (<i>st.</i>) oblique-leaved.	Siberia.	1759.		H.b.			

36 ramòsum. G.D.	(st.) branchingChives. Siberia.	1824.	6.	H.b.	
37 angùstum. G.D.	(pu.) narrow-leaved.	—	—	H.b.	
38 Pallasiò. G.D.	(pu.) Pallas's Chives. Russia.	—	H.b.	Murr.c.got.6.1775.t.3.
39 pusilum. G.D.	(li.) small.	Italy.	1824.	6. 7.	Redout. liliac. 2. t. 118.
40 rubéllum. G.D.	(ro.) reddish.	Siberia.	1825.	7. 8.	H.b.
41 reticulatum. G.D.(wh.)	netted.	N. America.	1811.	—	H.b.
42 oxypétalum. G.D.(bh.)	acute-petal'd.	Switzerland.	1822.	7.	H.b.
43 oléraceum. G.D.(st.pu.)	purple-striped.	England.	—	Eng. bot. v. 7. t. 488.
44 carinátum. G.D.(st.br.)	keeled.	—	5. 6.	— v. 24. t. 1658.
45 f'cétidum. w.	(da.pu.) twisted.	Siberia.	1823.	—	H.b.
46 violáceum. G.D.	(vi.) violet-flowered.	Hungary.	1819.	7.	H.b.
fléxum. w.k.					W. et K. hung.3. t.278.
47 áasperum. G.D.	(vi.) rough-leaved.	S. Europe.	1820.	—	H.b.
carinátum. R.L. nec aliorum.					Red. lil. 7. t. 368.
48 longispáthum. G.D.(g.pu.)	long-spathed.	—	1779.	6. 7.	H.b.
pállens. Bot. mag. t. 1420.					Red. lil. 6. t. 316.
49 paniculátum. G.D.(bh.)	panicked.	—	1780.	—	H.b.
50 pállens. G.D. (pa.wh.)	pale-flowered.	—	1779.	—	H.b.
51 tenuiflórum. G.D. (ro.)	slender-flowered.	Italy.	1824.	—	H.b.
52 pulchéllum. G.D. (vi.)	beautiful.	Russia.	1819.	8.	H.b.
paniculátum. R.L. colorátum. s.s.					Redout. lil. 5. t. 252.
53 flávum. G.D.	(ye.) yellow-leafy.	S. Europe.	1759.	6. 7.	H.b.
54 caucásicum. G.D. (ro.)	Caucasian.	Caucasus.	1810.	—	H.b.
55 rupéstre. G.D.	(ro.) small Caucasian.	—	1824.	—	H.b.
56 globosum. G.D.	(li.) globular-headed.	Siberia.	1820.	7.	H.b.
57 montánum. G.D. (ro.)	mountain.	Greece.	—	—	6. 8. H.b.
58 parviflórum. G.D.(br.)	small-flowered.	S. Europe.	1781.	6. 7.	H.b.
59 moschátum. G.D.(w.pu.)	small-leaved.	Hungary.	1805.	7. 8.	H.b.
setáceum. W.K.					W. et K. hung. 1. t. 68.
60 Stelleriánum. G.D. (g.w.)	Steller's.	Siberia.	1829.	—	H.b.
61 saxátile. G.D.	(wh.) rock.	Caucasus.	1825.	6. 7.	H.b.
62 stríctum. G.D.	(ro.) upright.	Siberia.	1820.	7.	H.b.
63 lineáre. G.D.	(re.) linear-leaved.	—	1819.	7. 8.	H.b.
64 nútans. G.D.	(fl.) great nodding.	—	1785.	—	H.b.
65 senescens. G.D.	(li.) Narcissus-leaved.	—	1596.	7.	H.b.
66 Andersoni. G.D.	(li.) Andersson's.	—	1819.	—	H.b.
67 spuriúm. G.D.	(li.) spurious.	—	—	—	H.b.
68 gláucum. G.D. (pu.)	glaucous.	—	1596.	6. 7.	H.b.
senescens. B.M. non L. baicalense. W.en.					Bot. mag. t. 1150.
69 angulósium. G.D. (pu.)	angular-stalked.	—	1739.	—	H.b.
70 acutángulum. G.D. (pu.)	acute-angled.	—	1816.	—	H.b.
71 danubiále. G.D. (pu.)	short-leaved.	—	—	—	H.b.
72 láxum. G.D. (pu.)	loose umbelled.	Russia.	1824.	—	H.b.
73 serotinum. G.D. (pu.)	late-flowering.	Switzerland.	—	8.	H.b.
74 álbidum. G.D. (wh.)	whitish-flower'd.	Caucasus.	1823.	6. 8.	H.b.
75 flavéscentis. G.D. (st.)	half naked.	Siberia.	1819.	8.	H.b.
denudátum. R.L.					Redout. lil. 6. t. 357.
76 rubens. G.D.	(pu.) red-rooted.	S. Europe.	1805.	5.	H.b.
bisúlcum. Redout. liliac. t. 286.					Bot. mag. t. 1318.
77 prostrátum. G.D. (li.)	small trailing.	Siberia.	6.	H.b.
78 congéstum. G.D. (pu.)	crowded-flower'd.	—	—	H.b.
79 pedemontánum. G.D.(ro.)	Piedmont.	Piedmont.	1820.	7. 8.	H.b.
80 narcissiflórum. G.D.(ro.)	large-flowered.	France.	1818.	—	H.b.
81 suavéolens. G.D. (w.pu.)	sweet-scented.	—	1801.	6. 8.	H.b.
82 ochroleúcum. G.D.(st.)	cream-colour'd.	Hungary.	1816.	7.	H.b.
83 cérnuum. G.D. (bh.)	drooping.	N. America.	1806.	6. 7.	H.b.
84 stellátum. G.D.	(li.) starry-flowered.	—	1811.	—	H.b.
85 caroliniánum. G.D.(li.)	Carolina.	—	1819.	—	H.b.
86 paradóxum. G.D. (wh.)	paradoxical.	Siberia.	1823.	4.	H.b.
87 scorzonérifolium. G.D. (ye.)	Scorzonera-leaved.	S. Europe.	—	6. 7.	H.b.
88 canadénse. G.D.	(ro.) Canadian.	N. America.	1739.	—	H.b.
89 mutábilis. G.D.	(bh.) changeable.	—	7.	H.b.
90 mágicum. G.D.	(st.) broad-leaved.	Spain.	1596.	6. 7.	H.b.
91 ambiguum. G.D.	(ro.) ambiguous.	Levant.	—	5. 6.	H.b.
incarnátum. s.s. róseum β.	Bot. mag. t. 978.	—			Flor. græc. t. 327.
92 am'œnum. G.D.	(ro.) fine red-flowered.	Italy.	1820.	—	H.b.
cárneum. T.N.					Ten. fl. nap. t. 28.
93 róseum. G.D.	(ro.) rose-coloured.	Spain.	1752.	6.	H.b.
illyricum. G.D.	(pi.) Illyrian.	Austria.	6. 7.	H.b.

95 subhirsutum. G.D. (wh.) hairy. β obtusum. (wh.) blunt-flowered.	S. Europe.	1596.	6.	H.b.	Cyr. pl. r. nap. 2. t. 3. Bot. mag. t. 774.
96 Clusiánum. G.D. (wh.) Clusius's.	—	1803.	6. 8.	H.b.	Clus. hist. 1. p. 192.
97 Chamae-Môly. G.D. (w.) dwarf-Moly.	—	1774.	2. 4.	F.b.	Bot. mag. t. 1203.
98 majâle. G.D. (wh.) large Moly.	Italy.	1825.	6.	F.b.	Ten. fl. nap. 1. t. 28.
99 brachystêmon. R.L. (w.) short-stamened.	S. France.	1826.	—	H.b.	Redout. liliac. 7. t. 374.
100 longifólium. G.D. (pu.) long-leaved.	Mexico.	—	8. 10.	F.b.	Bot. reg. t. 1034.
101 Cowâni. B.R. (wh.) Cowan's.	Chili.	1824.	5.	F.b.	— t. 758.
102 neapolitânum. G.D. (w.) Neapolitan. álbium. Bivon.	Italy.	—	4. 5.	H.b.	Swt. br. fl. gar. t. 201.
103 triquétrum. G.D. (wh.) triquetrous.	S. Europe.	1759.	5. 6.	H.b.	Bot. mag. t. 869.
104 médium. G.D. (wh.) intermediate.	Hungary.	1824.	5.	H.b.	Ten. fl. nap. 1. t. 31.
105 pendulînum. G.D. (w.) pendulous.	Italian.	—	4. 5.	H.b.	Bot. mag. t. 1148.
106 nigrum. G.D. (wh.) black. mágicum. B.M. β rubescens. (bh.) pale red-flowered.	S. Europe.	1596.	6.	H.b.	Flor. græc. t. 323. W. et K. hung. 1. t. 17.
107 atropurpureum. G.D. (d.pu.) dark purple.	Hungary.	1825.	—	H.b.	Bot. mag. t. 499.
108 Môly. G.D. (ye.) large yellow.	S. Europe.	1604.	—	H.b.	Eng. bot. v. 2. t. 122.
109 tricoccum. G.D. (wh.) three-seeded.	N. America.	1770.	7.	H.b.	Redout. liliac. 8. t. 482.
110 ursinum. G.D. (wh.) Ramson.	Britain.	—	4. 5.	H.b.	Bot. mag. t. 1222.
111 sulcátum. G.D. (wh.) furrowed.	—	1823.	—	H.b.	Redout. lili. 2. t. 98.
112 Victoriaâlis. G.D. (st.) long-rooted.	Austria.	1739.	5.	H.b.	Bot. mag. t. 1142.
113 odòrum. G.D. (wh. pu.) sweet-flowered. tatáricum. Bot. mag.	Siberia.	1787.	6. 7.	H.b.	t. 1129.
114 inodòrum. B.M. (bh.) fragrant. frâgrans. Vent. cels. t. 26. grâcile. w.	Carolina.	1776.	3. 4.	H.b.	Bot. mag. t. 1524&1035.
115 striatum. G.D. (bh.w.) striated. Ornithógalum gramineum. B.M. 2419.	America.	1800.	5. 6.	H.b.	Pall. it. 2. ap. t. 2.
116 cásipum. G.D. (bh.) Caspian. Amaryllis cásipca. w. Crinum cásipcum. Pal.	Astracan.	1820.	4. 5.	H.b.	
117 Akâka. F. Persian.	Persia.	1829.	—	H.b.	
SOWERB'ÆA. B.P. SOWERB'ÆA. júncea. B.P. (ro.) Rush-leaved.	Hexandria Monogynia. N.S. W.	1792.	5. 7.	G. 4.	Bot. mag. t. 1104.
LAXMA'NNIA. B.P. grâciliis. B.P. (wh.) slender.	LAXMA'NNIA. Hexandria Monogynia. N.S. W.	1822.	5. 7.	G. 4.	
APHYLLA'NTHES. L. LILY PINK. monspeliensis. L. (bh.) Rush-like.	Hexandria Monogynia. France.	1791.	6. 7.	H. 4.	Bot. mag. t. 1132.
THYSAN'OTUS. B.P. THYSAN'OTUS. (bh.) tuberous-rooted. N.S. W.	Hexandria Monogynia. S.S.	1822.	7. 9.	G. 4.	
1 tuberòsus. B.P. (bh.) tall.	—	—	—	G. 4.	
2 elâtior. B.P. (bh.) equal-anthered.	V. Diem. Isl.	—	—	G. 4.	
3 isantherus. B.P. (bh.) rush-like.	N. S. W.	—	—	G. 4.	
4 júnceus. B.P. (bh.) Chlamyspôrum junçifolium.	Par. lond. t. 103.	—	—	G. 4.	
CYANE'LLA. L. CYANE'LLA. (bh.) 1-flowered white. C. B. S.	Hexandria Monogynia. L.	1825.	4. 6.	F.b.	
1 álba. s.s. (wh.) yellow-flowered.	—	1768.	—	F.b.	
2 lútea. s.s. (ye.) Orchidiformis. s.s. (bh.) Orchis-like.	—	1788.	5. 12.	F.b.	
3 lineâta. B.T. (std.) lined.	—	1825.	4. 8.	F.b.	
4 odoratissima. B.R.	—	1816.	—	F.b.	
5 capénsis. s.s. (pu.) purple-flowered.	—	1768.	—	F.b.	
CONANTH'ERA. R.P. CONANTH'ERA. bifolia. R.P. (bh.) two-leaved.	Hexandria Monogynia. P.S. Chile.	1823.	3. 5.	F.b.	
CUMMI'NGIA. D.D. CUMMI'NGIA. campanulâta. D.D. (bh.) Hyacinth-flowered. Chile.	Hexandria Monogynia.	1823.	4. 6.	F.b.	
Conanthèra campanulâta. Hook. ex. flor. t. 214. bifolia. B.M. 2496. nec aliorum.	—	—	—	—	
LEUCOCO'RYNE. B.R. LEUCOCO'RYNE.	Hexandria Monogynia.	—	—	—	
1 odorâta. B.R. (wh.) sweet-scented.	Chile.	1826.	8. 9.	F. 4.	
2 ixioïdes. B.R. (bh.) Ixia-like.	—	1821.	7. 9.	F.b.	
Brod'æa ixioïdes. B.M.	—	—	—	—	
TRITEL'EIA. B.R. TRITEL'EIA.	Hexandria Monogynia.	—	—	—	
grandiflora. B.R. (bh.) large-flowered.	N. America.	1827.	7. 8.	H. 4.	
HESPEROSCO'R'DUM. B.R. MISSOURI HYACINTH.	Hexandria Monogynia.	—	—	—	
hyacíathinum. B.R. (bh.) largest-flowered.	N. America.	1827.	7.	H.b.	
Brod'æa grandiflora. Pursh non Smith.	—	—	—	—	
DIANE'LLA. B.P. DIANE'LLA.	Hexandria Monogynia. W.	—	—	—	
1 ensifolia. B.M. (wh.) whitish-flowered.	E. Indies.	1731.	8.	S. 4.	
Dracæna ensifolia. H.K.	—	—	—	—	
2 nemorösa. Lam. (bh.) blue Indian.	—	—	6. 8.	S. 4.	
3 corùlea. B.P. (bh.) rough-edged.	N. S. W.	1783.	5. 8.	F. 4.	Rumph. amb. 5. t. 73.
					Bot. mag. t. 505.

4 longifolia. B.P.	(bl.) long-leaved.	N. S. W.	1820.	5. 8. F. ȝ.	Bot. reg. t. 734.
5 lœvis. B.P.	(bl.) smooth-leaved.	_____	1823.	— F. ȝ.	
6 revoluta. B.P.	(bl.) revolute-margined.	_____	1822.	— F. ȝ.	
7 divisoria. B.P.	(bl.) divaricata.	N. Holland.	1805.	7. 8. G. ȝ.	
8 strumosa. B.R.	(bl.) strumous.	_____	1822.	5. 8. G. ȝ.	Bot. reg. t. 751.
LUZUR'AGA. B.P.	Luzur'aga.	Hexandria Monogynia. S.S.			
1 cymosa. B.P.	(pu.) cymose.	N. S. W.	1820.	4. 6. G. ȝ.	
2 montana. B.P.	(pu.) mountain.	_____	1823.	— G. ȝ.	
EUSTR'EPhUS. B.P.	Eustr'Ephus.	Hexandria Monogynia. S.S.			
1 latifolius. B.P.	(pu.) broadest-leaved.	N. S. W.	1800.	6. 9. G. ȝ. —	Bot. mag. t. 1245.
2 angustifolius. B.P.	(pu.) narrow-leaved.	N. Holland.	1816.	— G. ȝ. —	
MYRSIPHYLLUM. S.S. MYRSIPHYLLUM.	Hexandria Trigynia. S.S.				
1 asparagoïdes. s.s. (wh.)	broadest-leaved.	C. B. S.	1702.	10. 3. G. ȝ. —	Redout. liliac. t. 442.
Médeola asparagoïdes. L.					
2 angustifolium. s.s. (bh.)	narrow-leaved.	_____	1752.	12. 3. G. ȝ. —	Redout. liliac. t. 393.
HERR'ERIA. R.P.	Herr'eria.	Hexandria Monogynia.			
1stellata. R.P.	(ye.) prickly-stalked.	Chile.	1825.	6. 9. G. ȝ. —	Flor. peruv. 3. t. 303. f.a.
2 parviflora. B.R.	(gr.) small-flowered.	Brazil.	_____	S. ȝ. —	Bot. reg. t. 1042.
ASPA'RAGUS. L.	Aspa'ragus.	Hexandria Monogynia. L.			
1 amarus. s.s.	(ye.) bitter.	S. France.	1820.	6. 8. H. ȝ.	Redout. liliac. t. 446.
2 officinalis. s.s.	(gr.) common.	England.	— H. ȝ.	Eng. bot. v. 5. t. 339.
3 tenuifolius. DC.	(ye.) slender-leaved.	Hungary.	— H. ȝ.	Redout. liliac. t. 434.
sylvaticus. W. et K. hung. 3. t. 201.					
4 declinatus. s.s.	(st.) long-leaved.	C. B. S.	1759.	— F. ȝ.	
5 decumbens. s.s. (wh.)	decumbent.	_____	1792.	5. F. ȝ.	Jacq. schoen. 1. t. 97.
6 flexuosus. s.s. (wh.)	flexuose.	_____	7. 8. F. ȝ. —	
7 davuricus. s.s.	(st.) Daurian.	Dauria.	1819.	6. 8. H. ȝ.	
8 caspius. R.S.	(gr. wh.) Caspian.	Caucasus.	1821.	— H. ȝ.	
9 maritimus. s.s. (gr. wh.)	sea-side.	Siberia.	1820.	— H. ȝ.	
10 tricarinatus. DC.	(g.y.) three-keeled.	_____	— G. ȝ. —	Redout. liliac. t. 451.
11 verticillatus. s.s. (g.y.)	whorl-leaved.	Caucasus.	1752.	— H. ȝ. —	Buxb. cent. 5. ap. t. 37.
12 umbellatus. R.S.	(g.w.) umbel-flowered.	Canaries.	1816.	— G. ȝ.	
13 grandiflorus. R.S.	(wh.) large-flowered.	Teneriffe.	1828.	— G. ȝ. —	
14 longifolius. F.	(gr. ye.) long-leaved.	Russia.	1827.	— H. ȝ.	
15 acutifolius. s.s. (ye.)	needle-leaved.	Spain.	1640.	— H. ȝ.	Flor. græc. t. 337.
16 racemosus. s.s. (wh.)	branching.	E. Indies.	1808. S. ȝ.	
17 sarmentosus. s.s. (wh.)	linear-leaved.	Ceylon.	1710.	8. S. ȝ.	Redout. liliac. t. 460.
18 falcatus. s.s.	(wh.) sickle-leaved.	E. Indies.	1792. S. ȝ.	Burm. zeyl. t. 13 f. 2.
19 retrofractus. s.s. (wh.)	Larch-leaved.	Africa.	1759.	8. 9. G. ȝ.	Pluk. alm. t. 375. f. 3.
20 capensis. s.s. (wh. re.)	Cape.	C. B. S.	1691.	4. 5. G. ȝ.	Jacq. schoen. 3. t. 266.
21 aethiopicus. w.	(wh.) angular-stalked.	_____	1816. G. ȝ.	
22 scandens. w.	(wh. vi.) climbing.	_____	1795.	5. 6. G. ȝ. —	
23 asiaticus. s.s.	(wh.) Asiatic.	Asia.	1759.	— G. ȝ.	Pluk. alm. t. 14. f. 4.
24 subulatus. s.s.	(wh.) awl-leaved.	C. B. S.	1811.	— G. ȝ.	
25 Nivenianus. R.S. (w. pu.)	Niven's.	_____	—	— G. ȝ.	
26 lanceus. R.S.	(wh.) spear-leaved.	_____	—	— G. ȝ. —	
27 laricinus. R.S.	(wh.) Larch-leaved.	_____	1816.	— G. ȝ.	
28 tetragonus. R.S.	(wh.) square-leaved.	_____	1822.	— G. ȝ.	
29 albus. s.s.	(wh.) white.	Spain.	1540.	5. 7. F. ȝ.	Moris. s. I. t. 1. f. 3.
30 aphyllus. s.s.	(ye.) prickly.	S. Europe.	1640.	— F. ȝ.	Flor. græc. t. 338.
31 horridus. s.s.	(ye.) long-spined.	_____	1819.	— F. ȝ.	t. 239.
32 Broussoneti. L.en.	(st.) Broussonet's.	Canaries.	1815.	— F. ȝ.	Jacq. eclog. 2. t. 40.
CHARLWO'ODIA. S.F.A.	CHARLWO'ODIA.	Hexandria Monogynia.			
1 congesta. S.F.A.	(pu.) crowded-panicked.	N. Holland.	1821.	5. 7. S. ȝ.	Swt. flor. aust. t. 18.
2 stricta. S.F.A.	(bl.) loose-panicked.	_____	—	— S. ȝ.	Bot. reg. t. 956.
Dracaena stricta.	Bot. mag. t. 2575.				
DRACÆNA. W.	DRAGON-TREE.	Hexandria Monogynia. L.			
1 Draco. s.s.	(wh.) common.	E. Indies.	1640. S. ȝ.	Blackw. t. 358.
2 ensifolia. H.S.	(wh.) sword-leaved.	1800. S. ȝ.	
3? indivisa. s.s.	(wh.) undivided.	N. Zealand.	1819.	6. 8. G. ȝ.	
4? australis. s.s.	(wh.) Forster's.	_____	—	— G. ȝ.	Bot. mag. t. 2835
5 marginata. s.s.	(wh.) red-margined.	Madagascar.	1816. S. ȝ.	
6 ellata. W. enum.	—				
6 arborea. s.s.	(wh.) tree.	Africa.	1810.	3. 5. S. ȝ.	
7 frâgrans. s.s.	(wh.) fragrant.	_____	1768.	2. 5. S. ȝ.	Bot. mag. t. 1081.
8 férrea. L.	(wh.) dark-purple.	China.	1771.	3. 4. S. ȝ.	t. 2053.
9 terminata. w.	(wh.) red-leaved.	India.	1812.	4. 8. S. ȝ.	Lodd. bot. cab. t. 1224.
10 umbraculifera. s.s. (wh.) umbrella.	—	Mauritius.	1788.	4. 6. S. ȝ.	t. 289.
11 cérua. s.s.	(st. li.) drooping.	—	—	— S. ȝ.	Jacq. schoen. 1. t. 96.

ASPHODELEÆ.

12 mauritiâna. s.s. (<i>wh.</i>) Mauritius.	Mauritius.	1816. S. $\frac{1}{2}$.	
13 undulâta. s.s. (<i>wh.</i>) wave-leaved.	C. B. S.	— G. $\frac{1}{2}$.	
14 ovâta. B.M. (<i>li.</i>) oval-leaved.	Sierra Leon.	1790.	8. 9. S. $\frac{1}{2}$.	Bot. mag. t. 1180.
15 nodôsa. L.C. (<i>wh.</i>) knotted.	—	1822. S. $\frac{1}{2}$.	
16 reflexâ. s.s. (<i>li.</i>) reflex-leaved.	Madagascar.	1816.	5. 8. S. $\frac{1}{2}$.	Redout. liliac. t. 92.
17 surculôsa. B.R. (<i>st.</i>) trailing.	Sierra Leone.	1822.	— S. $\frac{1}{2}$.	Bot. reg. t. 1169.
CORDYL'INE. B.P. CORDYL'INE.	Hexandria Monogynia.			
1 hemichry'sa. P.S. (<i>wh.</i>) golden-leaved.	Bourbon.	1823. S. $\frac{1}{2}$.	
2 parviflôra. s.s. (<i>wh.</i>) small-flowered.	Mexico.	1828. S. $\frac{1}{2}$.	H. et B.n. gen. 3. t. 674.
PHO'RMIUM. W. FLAX-LILY.	Hexandria Monogynia. W.			
ténax. w. (<i>ye.</i>) Iris-leaved.	N. Zealand.	1788.	8. F. $\frac{1}{2}$.	Redout.liliac.t.448-449.
XANTHORRHŒA.B.P. GUM-TREE.	Hexandria Monogynia. S.S.			
1 arbòrea. B.P. (<i>wh.</i>) tree.	N. S. W.	1828. G. $\frac{1}{2}$.	
2 australis. B.P. (<i>wh.</i>) southern.	V. Diem. Isl.	1820. F. $\frac{1}{2}$.	
3 hâstile. B.P. (<i>wh.</i>) yellow gum.	N. S. W.	1803.	4. 5. F. $\frac{1}{2}$.	
4 mèdia. B.P. (<i>wh.</i>) intermediate.	—	1819.	— F. $\frac{1}{2}$.	
5 mînor. B.P. (<i>wh.br.</i>) small.	—	1804.	4. 6. F. $\frac{1}{2}$.	
6 bracteâta. B.P. (<i>wh.</i>) long-bracted.	—	1810.	4. 5. F. $\frac{1}{2}$.	
ROXBUR'GHIA. W. ROXBUR'GHIA.	Octandria Monogynia.			
1 gloriôsides. r.c.(<i>st.pu.</i>)cordate-leaved.	E. Indies.	1803.	4. 6. S. $\frac{1}{2}$. Roxb. corom. 1. t. 32.	
2 viridiflôra. Ex.B.(<i>st.cr.</i>)elliptic-leaved.	—	—	— S. $\frac{1}{2}$. Sm. exot. bot. 1. t. 57.	
gloriôsa. Bot. mag. t. 1500.				

ORDO CXXV.

BROMELIACEÆ. Juss. gen. 49.

AG'AVE. L.	AG'AVE. Hexandria Monogynia. L.			
1 americâna. H.S. (<i>gr.</i>) large American. S.America.	1640.	8. 10. D.G. $\frac{1}{2}$.	Andr. deposit. t. 438.	
2 Milléri. H.S. (<i>gr.</i>) Miller's.	—	1768.	— D.G. $\frac{1}{2}$.	
3 flaccida. H.S. (<i>gr.</i>) flaccid.	—	1790.	— D.S. $\frac{1}{2}$.	
4 lûrâta. B.M. (<i>gr.</i>) Vera Cruz.	Vera Cruz.	1731.	6. 7. D.S. $\frac{1}{2}$.	Bot. mag. t. 152.
5 brachystächys. s. (<i>gr.</i>) short-spiked. <i>spicâta. R.L. non Cavan.</i>	Mexico.	1826. D.G. $\frac{1}{2}$.	Redout. liliac. t. 485.
6 spicâta. s. (<i>gr.ye.</i>) spiked.	Havannah.	1825. D.S. $\frac{1}{2}$.	Cav.h.reg.mad.ic.ined.
7 angustifolia. H.S. (<i>gr.</i>) narrow-leaved.	St. Helena.	1770. D.S. $\frac{1}{2}$.	
8 yuccafolia. H.S.s. (<i>y.g.</i>) Yucca-leaved.	S. America.	1817. D.S. $\frac{1}{2}$.	Redout.liliac.t.328,329.
9 mexicâna. H.S.s. (<i>gr.</i>) Mexican.	Mexico.	1818. D.G. $\frac{1}{2}$.	
10 Karâtto. H.S. (<i>gr.</i>) Karatto.	S. America.	1768. D.S. $\frac{1}{2}$.	
11 vivipara. H.S. (<i>gr.ye.</i>) viviparous.	—	1731.	8. 10. D.S. $\frac{1}{2}$.	Comm. præl. t. 15.
12 polyacânta. H.R. (<i>g.y.</i>) black-spined.	—	1800. D.S. $\frac{1}{2}$.	
13 virginica. H.S. (<i>gr.ye.</i>) Virginian.	N. America.	1765.	9. F. $\frac{1}{2}$.	Bot. mag. t. 1157.
FURCR'EA. V.	FURCR'EA. Hexandria Monogynia. H.S.			
1 gigântea. (<i>gr.wh.</i>) gigantic. <i>Agâve fa'cida. w.</i>	S. America.	1690.	9. 1. D.S. $\frac{1}{2}$.	DC. plant. gras. t. 126.
2 tuberosa. H.K. (<i>gr.wh.</i>) tuberous.	—	1739.	8. 9. D.S. $\frac{1}{2}$.	Comm. hort. 2. f. 19.
3 cubénsis. H.S. (<i>wh.gr.</i>) Cuba.	—	— D.S. $\frac{1}{2}$.	Jacq.am.pic.t.260.f.25.
4 rígida. H.S. (<i>gr.ye.</i>) Vera Cruz.	—	1768. D.S. $\frac{1}{2}$.	Spin. cat. 1812. c. ic.
5 australis. H.S. entire-leaved.	N. Holland.	1811. D.S. $\frac{1}{2}$.	
6 madagascariénis. H. Madagascar.	Madagascar.	1818. D.S. $\frac{1}{2}$.	
7 cantâla. H.S.s. Canton.	China.	— D.G. $\frac{1}{2}$.	
LITT'EA. H.S.S.	LITT'EA. Hexandria Monogynia. H.S.S.			
geminiiflôra. H.S.S. (<i>gr.</i>) twin-flowered.	S. America.	1810. S. $\frac{1}{2}$.	Bot. reg. t. 1145.
Agâve geminiiflôra. Ker. <i>Buonapártea júncea.</i> H.S. nec aliorum.				
ANANA'SSA. B.H.	PINE-APPLE. Hexandria Monogynia. L.			
1 sativa. B.R. (<i>bl.</i>) common.	S. America.	1690.	1. 12. S. $\frac{1}{2}$.	Bot. mag. t. 1554.
2 semiserrâta. s.s. (<i>bl.</i>) semiserrate.	—	— S. $\frac{1}{2}$.	
3 lúcida. W.en.	—	— S. $\frac{1}{2}$.	
4 débilis. B.R. (<i>bl.sc.</i>) red-bracted.	—	1826.	— S. $\frac{1}{2}$.	
5 bracteâta. B.R. (<i>bl.sc.</i>) red-bracted.	Jamaica.	1785.	9. 10. S. $\frac{1}{2}$.	Bot. reg. t. 1081.
BILLBE'RGLIA. Th.	BILLBE'RGLIA. Hexandria Monogynia.			
1 zebrina. B.M. (<i>ye.</i>) white-barred.	Brazil.	1825.	6. 7. S. $\frac{1}{2}$.	Bot. mag. t. 2686.
2 fasciâta. B.R. (<i>bl.</i>) banded.	—	—	8. 10. S. $\frac{1}{2}$.	Bot. reg. t. 1130.

3 iridifolia. B.R. (<i>ye.bl.</i>) drooping.	Brazil.	1826.	3. 5. S. 4.	Bot. reg. t. 1068.
4 amena. B.R. (<i>ye.bl.</i>) pale-flowered.	S. America.	1810.	3. 11. S. 4.	Lodd. bot. cab. t. 76.
<i>Bromelia pallida</i> . Bot. reg. t. 344.				
5 pyramidalis. B.R. (<i>cr.</i>) pyramidal.	Brazil.	2. 4. S. 4.	Bot. reg. t. 203.
<i>Bromelia nudicaulis</i> . B.R. <i>pyramidalis</i> . Bot. mag. t. 1732.				
β bicolor. B.R. (<i>cr.bl.</i>) two-coloured.	S. America.	— S. 4.	Bot. reg. t. 1181.
6 nudicaulis. B.R. (<i>st.</i>) naked-stalked.	Trinidad.	1822.	4. 6. S. 4.	Hook. ex. flor. t. 143.
7 cruentata. B.M. (<i>bl.</i>) blood-stained.	Brazil.	1824.	— S. 4.	Bot. mag. t. 2892.
BROM'ELIA. B.R. BROM'ELIA. Hexandria Monogynia.				
1 melanantha. B.R. (<i>bk.</i>) black-flowered.	Trinidad.	1823.	5. 7. S. 4.	Bot. reg. t. 766.
2 fastuosa. L.col. (<i>pu.</i>) noble.	S. America.	1815.	8. 10. S. ½.	Lindl. col. bot. t. 1.
3 lingulata. s.s. (<i>ye.</i>) tongue-leaved.	—	1759.	5. 6. S. ½.	Plum. ic. t. 64. f. 1.
4 Karatas. s.s. (<i>ro.</i>) upright-leaved.	W. Indies.	1739.	— S. ½.	Jacq. vind. t. 1. t. 31. 32.
5 humiliis. s.s. (<i>bl.</i>) dwarf.	—	1789.	3. S. 4.	Jacq. ic. t. 60.
6 Pinguin. s.s. (<i>re.</i>) broad-leaved.	—	1699.	3. 4. S. ½.	Jacq. am. ed. pict. t. 91.
7 chrysanthra. s.s. (<i>ye.</i>) yellow-flowered.	S. America.	1820.	— S. ½.	Jacq. scheen. t. 1. t. 55.
8 exudans. B.C. (<i>ye.</i>) clammy.	—	1816.	— S. 4.	Lodd. bot. cab. t. 801.
9 paniculigera. s.s. (<i>sc.</i>) paniced.	W. Indies.	1822.	— S. 4.	
10 sylvestris. s.s. (<i>cr.</i>) wild.	S. America.	1820.	7. 8. S. 4.	Bot. mag. t. 2392.
11 Acanga. s.s. (<i>ye.</i>) recurved.	—	1824.	— S. 4.	Pis. bras. t. 91.
12 sessiliflora. L.C. sessile-flowered.	—	—	— S. 4.	
GUZMA'NNIA. R.P. GUZMA'NNIA. Hexandria Monogynia. S.S.				
tricolor. R.P. (<i>g.st.ro.</i>) three-coloured.	S. America.	1818.	5. 6. S. 4.	Lind. coll. bot. t. 8.
PITCA'RNIA. S.S. PITCA'RNIA. Hexandria Monogynia. L.				
1 staminea. s.s. (<i>sc.</i>) long-stamened.	S. America.	1823.	1. 3. S. 4.	Bot. mag. t. 2411.
2 integrifolia. s.s. (<i>sc.</i>) entire-leaved.	W. Indies.	1800.	8. S. 4.	— t. 1462.
3 latifolia. s.s. (<i>sc.</i>) broad-leaved.	—	1785.	8. 9. S. 4.	— t. 856.
4 bracteata. R.L. (<i>sc.</i>) large bracted-red.	—	1799.	4. 5. S. 4.	— t. 2813.
5 sulphurea. A.R. (<i>st.</i>) yellow-flowered.	—	1797.	6. 8. S. 4.	Andr. deposit. t. 249.
bracteata β sulphurea. Bot. mag. t. 1416.				
6 suavolens. B.R. (<i>st.</i>) sweet-scented.	Brazil.	1826.	6. 7. S. 4.	Bot. reg. t. 1069.
7 albiflos. B.M. (<i>wh.</i>) white-flowered.	—	1824.	9. 12. S. 4.	Bot. mag. t. 2642.
8 bromeliacea. s.s. (<i>sc.</i>) Pine-apple-leaved.	Jamaica.	1781.	6. 8. S. 4.	— t. 824.
9 angustifolia. s.s. (<i>sc.</i>) narrow-leaved.	Santa Cruz.	1777.	12. 1. S. 4.	— t. 1547.
10 furfuracea. B.M. (<i>ro.</i>) mealy.	S. America.	1818.	7. 9. S. 4.	— t. 2657.
11 flammæa. B.R. (<i>sc.</i>) flame-coloured.	Brazil.	1823.	11. 12. S. 4.	— t. 1092.
12 media. L.C. (<i>sc.</i>) intermediate.	S. America.	—	7. 9. S. 4.	
13 adiflora. L.C. (<i>sc.</i>) Iris-flowered.	—	—	— S. 4.	
14 humiliis. L.C. (<i>sc.</i>) dwarf.	—	—	— S. 4.	
POURR'ETIA. R.P. POURR'ETIA. Hexandria Monogynia. S.S.				
1 lanuginosa. R.P. (<i>gr.</i>) woolly.	S. America.	1826.	4. 8. S. 4.	Flor. peruv. 3. t. 256.
2 pyramidata. K.S. (<i>st.</i>) pyramidal.	—	1822.	4. 6. S. 4.	— 3. t. 257.
3 coarctata. R.P. (<i>ye.</i>) crowded-spiked.	Chili.	—	— S. 4.	Feuil. per. 3. t. 39.
TILLA'NDSIA. L. TILLA'NDSIA. Hexandria Monogynia. L.				
1 usneoides. s.s. (<i>ye.gr.</i>) moss-like.	W. Indies.	1823.	4. 8. S. 4.	Sloan. hist. t. 122. f. 2-3.
2 Bartramii. s.s. Bartram's.	Carolina.	1825.	— S. 4.	
3 recurvata. s.s. (<i>pu.</i>) recurve-leaved.	Jamaica.	1793.	7. S. 4.	Sloan. jam. 1. t. 121. f. 1.
4 xiphioides. B.R. (<i>wh.</i>) Xiphium-like.	BuenosAyres.	1815.	— S. 4.	Bot. reg. t. 105.
5 serrata. s.s. (<i>ye.</i>) saw-leaved.	Jamaica.	1793.	6. 7. S. 4.	Plum. ic. 63. t. 75. f. 1.
6 lingulata. s.s. (<i>ye.</i>) tongue-leaved.	—	1776.	— S. 4.	Jacq. amer. t. 62.
7 acaulis. B.R. (<i>wh.</i>) stemless.	Brazil.	1826.	7. 9. S. 4.	Bot. reg. t. 1157.
8 psittacina. B.M. (<i>or.y.</i>) gaudy-flowered.	—	1827.	4. 8. S. 4.	Bot. mag. t. 2841.
9 setacea. s.s. (<i>st.</i>) bristle-leaved.	Jamaica.	1822.	— S. 4.	
10 stricta. s.s. (<i>bl.</i>) stiff-leaved.	Brazil.	1810.	11. S. 4.	Bot. mag. t. 1529.
11 bulbosa. H.E.F. (<i>bl.</i>) bulbous.	Trinidad.	1823.	8. 9. S. 4.	Hook. ex. flor. t. 173.
12 aloifolia. H.E.F. (<i>st.</i>) Aloe-leaved.	—	1824.	11. 12. S. 4.	— t. 205.
13 pulchra. H.E.F. (<i>wh.</i>) pretty.	—	1822.	10. 11. S. 4.	— t. 154.
14 ánceps. B.C. (<i>bl.</i>) two-edged.	W. Indies.	1819.	7. S. 4.	Lodd. bot. cab. t. 771.
15 utriculata. s.s. (<i>st.</i>) bottle.	S. America.	1793.	7. 8. S. 4.	
16 tenuifolia. s.s. (<i>sc.</i>) slender-leaved.	W. Indies.	1822.	— S. 4.	Plum. ic. 232. f. 2.
17 acutifolia. L.en. (<i>st.</i>) acute-leaved.	Brazil.	1815.	— S. 4.	Bot. reg. t. 749.
flexuosa β pallida. B.R.				
18 flexuosa. s.s. (<i>sc.</i>) flexuose.	W. Indies.	1790.	— S. 4.	Jacq. amer. t. 63.
19 nūtans. s.s. (<i>bl.</i>) nodding-spiked.	—	1793.	— S. 4.	
20 fasciculata. s.s. (<i>st.</i>) bundled.	—	1822.	— S. 4.	
21 niftida. H.E.F. (<i>wh.</i>) shining-leaved.	Jamaica.	1823.	11. 12. S. 4.	Hook. ex. flor. t. 218.
22 polystachya. s.s. (<i>st.</i>) many-spiked.	W. Indies.	1820.	7. 9. S. 4.	
23 angustifolia. s.s. (<i>st.</i>) narrow-leaved.	—	1822.	— S. 4.	
24 paniculata. s.s. (<i>st.</i>) paniced.	—	1817.	— S. 4.	Plum. ic. t. 237.

25 coarctata. G.C. crowded.	W. Indies. 1822.	S. 4.
26 compræssa. G.C. compressed.	— — —	S. 4.
27 græcilis. G.C. slender.	— — —	S. 4.
28 ramosa. G.C. branching.	— — —	S. 4.
29 rígida. G.C. rigid.	— — —	S. 4.

BUONAPARTEA. R.P. BUONAPARTEA. *Hexandria Monogynia*.júnea. R.P. (*bl.*) rush-leaved. Peru. 1800. S. 4. Flor. peruv. 3. t. 242.

ORDO CCXVI.

TULIPACEÆ. *Kth. synops. 1. p. 292.*

YU'CCCA. L.	ADAM'S-NEEDLE. Hexandria Monogynia. L.		
1 conspœcia. H.S.S. (<i>wh.</i>) broad rough-edged.	1810. 8. 10. F. h.		
2 aloifolia. H.S.S. (<i>wh.</i>) Aloe-leaved.	1696. — G. h.	Bot. mag. t. 1700.	
3 serrulata. H.S.S. (<i>wh.</i>) narrow saw-leaved.	1808. F. h.		
4 Dracónis. H.S.S. (<i>wh.</i>) drooping-leaved.	1732. 8. 9. F. h.	Dill. elt. t. 324. f. 417.	
5 crenulata. H.S.S. (<i>wh.</i>) rough-edged oblique.	1818. F. h.		
6 arcuata. H.S.S. (<i>wh.</i>) arcuated.	1817. F. h.		
7 tenuifolia. H.S.S. (<i>wh.</i>) slender-leaved.	— F. h.		
8 concava. H.S.S. (<i>wh.</i>) hollow-thready.	1816. 7. 8. H. 4.		
9 filamentosa. H.S.S. (<i>wh.</i>) common-thready.	1675. 7. 10. H. 4.		
10 flaccida. H.S.S. (<i>wh.</i>) flacid-thready.	1816. — H. 4.		
11 puberula. H.P. (<i>wh.</i>) pubescent-stemmed.	1822. — H. 4.	Swt. br. fl. gar. t. 251.	
12 glaucescens. H.S.S. (<i>wh.</i>) glaucescent.	1815. 7. 8. H. 4.	— t. 53.	
13 rectírva. P.L. (<i>wh.</i>) recurved-leaved.	1795. 8. 9. H. h.	Sal. par. lond. t. 31.	
14 glauca. B.M. (<i>st.</i>) glaucous-leaved.	1812. — H. 4.	Bot. mag. t. 2662.	
15 strícta. B.M. (<i>gr. pu.</i>) straight-leaved.	1811. — H. h.	— t. 2222.	
16 angustifolia. H.S.S. (<i>gr.</i>) narrow-leaved.	— — H. h.	— t. 2236.	
17 supérbia. H.S.S. (<i>wh.</i>) superb. 8. 10. G. h.	Andr. deposit. t. 473.	
gloriæsa. A.R. nec aliorum.			
18 acuminata. S.F.G. (<i>br. w.</i>) tapering-flowered. 7. 9. H. h.	Swt. br. fl. gar. t. 195.	
19 gloriæsa. H.S.S. (<i>wh.</i>) handsome.	1596. 7. 8. H. h.	Bot. mag. t. 1260.	
20 rufocincta. H.S.S. (<i>wh.</i>) rufous-edged.	1816. — H. 4.		
21 obliqua. H.S.S. (<i>wh.</i>) oblique-leaved.	1808. — H. 4.		
β mähr. H.S.S. (<i>wh.</i>) larger.	— — H. 4.		
22 spinosa. K.S. (<i>pu. or.</i>) spiny-leaved.	1829. G. h.		
TULIPA. L.	TULIP. Hexandria Monogynia. L.		
1 altaica. R.S. (<i>ye.</i>) Altaic.	Altay mount. 1822. 4. 5. H.b.		
2 saxatilis. R.S. (<i>ye.</i>) rocky.	Crete. 1827. — H.b.	Reich. ic. rar. t. 396.	
3 pâtens. R.S. (<i>wh. g. y.</i>) spreading-flowered.	Siberia. 1829. — H.b.		
4 stellata. B.M. (<i>bh. wh.</i>) starry.	E. Indies. 1827. 3. 4. G.b.	Bot. mag. t. 2762.	
5 montana. B.R. (<i>cr.</i>) mountain.	Persia. 1826. — H.b.	Bot. reg. t. 1106.	
6 biflora. s.s. (<i>wh. vi. ye.</i>) two-flowered.	Russia. 1806. 4. H.b.	— t. 535.	
7 tricolor. F. three-coloured.	— 1823. H.b.		
8 Celsiana. s.s. (<i>ye. re.</i>) Cels's.	Levant. 6. 7. H.b.	Bot. mag. t. 717.	
Breyniæna. B.M. nec aliorum.			
9 Biebersteiniana. R.S. (<i>y. pu.</i>) Bieberstein's.	Siberia. 1820. — H.b.		
rêpens. F. ?			
10 sylvestris. s.s. (<i>ye.</i>) wild.	England. 4. 5. H.b.	Eng. bot. v. 1. t. 63.	
11 türctica. Roth. (<i>ye.</i>) Florentine.	S. Europe. — H.b.	Swt. br. fl. gar. t. 186.	
12 Clusiæna. R.S. (<i>sc. wh.</i>) Clusius's.	Sicily. 1636. 5. 6. H.b.	Bot. mag. t. 1390.	
13 maculata. R.S. (<i>ye. gr.</i>) spotted-flowered.	Spain. — H.b.		
14média. R.S. (<i>sc. wh.</i>) intermediate. 1828. — H.b.		
15 cornuta. R.S. (<i>va.</i>) horn-petaled.	Levant. 1816. 5. H.b.		
16 undulata. (<i>sc. ve.</i>) wave-leaved.	Whitley's. 1820. 5. 6. H.b.	Bot. reg. t. 127.	
17 óculus solis. s.s. (<i>sc. ve.</i>) Sun's-eye.	S. Europe. 1816. — H.b.	Redout. liliac. t. 219.	
18 præcox. s.s. (<i>sc. gr.</i>) early-flowering.	Naples. 1825. 3. 4. H.b.	Swt. br. fl. gar. t. 157.	
19 pérsica.	(<i>sc. ve.</i>) Persian.	1826. — H.b.	Bot. reg. t. 1143.
20 suavolens. s.s. (<i>sc. ye.</i>) Van-Thol.	S. Europe. 1603. — H.b.	Bot. mag. t. 839.	
21 pubescens. s.s. (<i>va.</i>) wool. woolly-stalked. — H.b.	Swt. br. fl. gar. t. 78.	
22 Gesneriana. s.s. (<i>va.</i>) common.	Levant. 1577. 4. 5. H.b.	Bot. reg. t. 380.	
α coccinea.	(<i>sc. ve.</i>) scarlet-flowered.	Constantinople. 1816. — H.b.	Bot. mag. t. 1135.
β versicolor.	(<i>va.</i>) various-coloured.	Levant. — — H.b.	
γ simploniensis. (<i>st. pu.</i>) Semplon. — H.b.	Swt. flor. guid. t. 53.	

δ <i>Alexandrīna</i> . (st. pu. ve.) <i>Alexandrina</i>	H.b.	Swt. flor. gnid. t. 138.
ε <i>Burnardīana</i> . (st. pu. ve.) <i>Burnard's agitator</i>	—	H.b.	— t. 132.
ζ <i>Collingwōodii</i> . (y. pu. ve.) <i>Lord Collingwood</i>	—	H.b.	— t. 85.
η <i>Leopoldī</i> . (ye. ve.) <i>Prince Leopold</i>	—	H.b.	— t. 128.
θ <i>Geōrgii</i> . (ye. ve.) <i>George the Fourth</i>	—	H.b.	— t. 89.
ι <i>platyántha</i> . (ye. ve.) <i>Platonia</i>	—	H.b.	— t. 77.
κ <i>imperiális</i> . (ye. ve.) <i>Emperor of Austria</i>	—	H.b.	— t. 103.
λ <i>Miguelī</i> . (ye. ve.) <i>Don Miguel</i>	—	H.b.	— t. 19.
μ <i>Polyphēmi</i> . (ye. ve.) <i>Lawrence's Polyphemus</i>	—	H.b.	— t. 7.
ν <i>sanguífera</i> . (ye. cr. ve.) <i>Bataille d'Eyleau</i>	—	H.b.	— t. 4.
ξ <i>Duckettiana</i> . (y. sc. ve.) <i>Sir George Duckett</i>	—	H.b.	— t. 66.
ο <i>Canningiāna</i> . (y. sc. ve.) <i>Strong's Canning</i>	—	H.b.	— t. 135.
π <i>règia</i> . (ye. sc. ve.) <i>Strong's King</i>	—	H.b.	— t. 42.
ρ <i>Titiána</i> . (ye. sc. ve.) <i>Titian</i>	—	H.b.	— t. 109.
σ <i>Jūno</i> . (wh. sc. pu.) <i>Rose Juno</i>	—	H.b.	— t. 56.
τ <i>Daphneánā</i> . (wh. cr.) <i>Rose Daphne</i>	—	H.b.	— t. 83.
υ <i>retusa</i> . (wh. cr.) <i>Walworth</i>	—	H.b.	— t. 26.
φ <i>fulmínea</i> . (wh. sc.) <i>Bartlett's thunderbolt</i>	—	H.b.	— t. 95.
χ <i>Hilliána</i> . (wh. sc.) <i>Lord Hill</i>	—	H.b.	— t. 82.
ψ <i>fulgens</i> . (w. ro.) <i>Rose brillante</i>	—	H.b.	— t. 5.
ω <i>Creweæ</i> . (wh. sc.) <i>Sherwood's Lady Crewe</i>	—	H.b.	— t. 153.
aa <i>Victoriána</i> . (w. ro. cr.) <i>Princess Victoria</i>	—	H.b.	— t. 38.
ββ <i>Stróngii</i> . (w. cr. ve.) <i>Strong's high Admiral</i>	—	H.b.	— t. 11.
γγ <i>puélla</i> . (wh. ro.) <i>Pucelle d'Orleans</i>	—	H.b.	— t. 50.
օօ <i>principissæ</i> . (w. ro. cr.) <i>Strong's Duchess of Kent</i>	—	H.b.	— t. 75.
εε <i>púrpureo-ròsea</i> . (w. ro. pu.) <i>Rose camusa de craix</i>	—	H.b.	— t. 116.
ζζ <i>iridiflóra</i> . (wh. pu.) <i>Strong's rainbow</i>	—	H.b.	— t. 124.
ηη <i>Hollandiána</i> . (w. pu. ve.) <i>Lord Holland</i>	—	H.b.	— t. 106.
օօ <i>Ludoviána</i> . (wh. ve.) <i>Louis XVI</i>	—	H.b.	— t. 150.
ιι <i>lancastriensis</i> . (w. pu. ve.) <i>Lancashire hero</i>	—	H.b.	— t. 119.
κκ <i>marginátā</i> . (wh. ve.) <i>Davey's Trafalgar</i>	—	H.b.	— t. 145.
λλ <i>Penelopéana</i> . (w. ve.) <i>Penelope</i>	—	H.b.	— t. 57.
μμ <i>Jenkinsi</i> . (w. ve.) <i>Goldham's Earl of Liverpool</i>	—	H.b.	— t. 16.
νν <i>Fredericiána</i> . (w. ve.) <i>Willmer's Duke of York</i>	—	H.b.	— t. 121.
ξξ <i>Buonaparteána</i> . (w. ve.) <i>Buonaparte</i>	—	H.b.	— t. 72.
օօ <i>augústa</i> . (wh. ve.) <i>Grande Monarque</i>	—	H.b.	— t. 92.
ππ <i>Lampsoniána</i> . (w. ve.) <i>Lampson</i>	—	H.b.	— t. 100.
ρρ <i>Aglaíana</i> . (wh. pu.) <i>Aglaia</i>	—	H.b.	— t. 87.
σσ <i>Alexándri</i> . (wh. pu.) <i>Violet Alexander</i>	—	H.b.	— t. 30.
ττ <i>virgínea</i> . (wh. ve.) <i>Lady of the Lake</i>	—	H.b.	— t. 62.
vv <i>Winefrediána</i> . (w. v.) <i>Winefred</i>	—	H.b.	— t. 141.
φφ <i>Daveyána</i> . (wh. ve.) <i>Daveyana</i>	—	H.b.	— t. 35.
χχ <i>Maria</i> . (wh. ve. pu.) <i>Goldham's Maria</i>	—	H.b.	— t. 48.
ψψ <i>majéstā</i> . (wh. ve.) <i>Majestueuse</i>	—	H.b.	— t. 39.
ωω <i>álbida</i> . (wh. ve.) <i>Gloria alborum</i>	—	H.b.	— t. 74.
aaa <i>Charlotteána</i> . (w. pu.) <i>Princess Charlotte's cenotaph</i>	—	H.b.	— t. 23.
βββ <i>pléna</i> . (va.) <i>double-flowered</i> . Levant. 1577. 4. 5. H.b. —	—	—	—	—
γγγ <i>laciniátā</i> . (ye. re. pu.) <i>Parrot</i> . —	—	—	H.b.	—

FRITILLARIA. L. FRITILLARY. Hexandria Monogynia, L.

1 <i>præcox</i> . (wh.) <i>early white</i> . Europe.	3. 4.	H.b.	
Meleagris β præcox. P.S.			
2 Meleagris. E.B. (pu. wh.) <i>chequered</i> . Britain.	3. 5.	H.b.	Eng. bot. v. 9. t. 622.
3 meleagroïdes. R.S. (d. vi.) <i>Altaic</i> . Altay. 1827.	—	H.b.	
4 tenella. M.B. (br. pu.) <i>slender</i> . Caucasus. 1824.	—	H.b.	Bot. mag. t. 1216.
racemosa. a minor. B.M. <i>orientalis</i> . Adam.			
5 nervosa. s.s. (d. pu.) <i>nerved-leaved</i> . —	4. 5.	H.b.	Bot. mag. t. 853.
latifolia. β minor. B.M.			
6 lutea. M.B. (ye. pu.) <i>large yellow</i> . —	1812.	—	Bot. mag. t. 1538.
7 latifolia. w. (pu. br.) <i>broad-leaved</i> . —	1604.	—	— t. 1207.
8 ruthénica. s.s. (pu.) <i>Russian</i> . —	1820.	—	H.b.
9 messanénisia. s.s. (pu. ye.) <i>Italian</i> . Italy. 1825.	—	H.b.	
10 lusitanica. s.s. (vi. ye.) <i>Portugal</i> . Portugal. —	—	—	H.b.
11 nigra. B.M. (d. pu. ye.) <i>dark-flowered</i> . Pyrenees. 1596. 5. H.b.			Bot. mag. t. 664.
pyrenæica. B.M. nec ariorum. aquitánica. Clusius.			
12 racemosa. R.S. (br. pu.) <i>cluster-flowered</i> . Russia. 1605. 5. 6. H.b.			Bot. mag. t. 952.
13 pyrenæica. L. (pu.) <i>Pyrenean</i> . S. Europe.	4. 5.	H.b.	Flor. græc. t. 328.
14 tulipifolia. M.B. (pu. vi.) <i>Tulip-leaved</i> . Caucasus. 1816.	—	H.b.	
15 obliqua. B.M. (pu.) <i>oblique-leaved</i> . —	4.	H.b.	Bot. mag. t. 857.
16 pérsica. B.M. (pu.) <i>Persian</i> . Persia. 1596. 4. 5. H.b.			— t. 1537.
β minor. B.M. (pu.) <i>lesser</i> . —	—	H.b.	— t. 962.

17	verticillatæ. F.	(pu.) acute-leaved.	Siberia.	1820.	4. 5. H.b.
18	scándens. F.	(st.) blunt-leaved.	1827.	—	H.b.
19	leucántha. F.	(wh.) white-flowered.	Altay.	—	H.b.
20	Imperialis. w.	(y.re.) Crown Imperial.	Persia.	1596.	3. 4. H.b.
	α rúbra.	(re.) red-flowered.	—	—	H.b.
	β flava.	(ye.) yellow-flowered.	—	—	H.b.
	CYCLOBO'THRA. S.F.G.	CYCLOBO'THRA.	Hexandria Monogynia.		
1	barbáta. s.F.G.	(ye.) bearded.	Mexico.	1827.	7. 8. H.b.
	Fritillaria barbata. K.S.				
2	purpurea. S.F.G.	(pu.y.) purple-flowered.	—	—	H.b.
	CALOCHO'R'TUS. Ph.	CALOCHO'R'TUS.	Hexandria Monogynia.		
1	macrocápus. B.R.	(pu.) long-fruited.	Columbia.	1826.	7. 8. H.b.
2	nítidus. D.	(pu.ye.) glossy.	—	—	H.b.
	AMBL'RION. Raf.	AMBL'RION.	Hexandria Monogynia.		
1	pùdicum.	(ye.) dwarf.	N. America.	1825. H.b.
	Lilium pùdicum. Ph.				
2	lanceolátum.	(pu.) spear-leaved.	—	1823. H.b.
	Fritillaria lanceolata. Ph.	Lilium affine. R.S.			
3	camtschaticénsē. (pu.)	Camtschatka.	Camtschatka.	1759.	5. H.b.
	Lilium camtschaticénsē. L.T.				
4	álbum.	(wh.) white-flowered.	N. America.	1825.	4. H.b.
	Fritillaria alba. N.				
	LI'LUM. L.	LILY.	Hexandria Monogynia. L.		
1	cárdium. w.	(wh.) common white.	Levant.	1596.	6. 7. H.b.
	β striatum.	(wh.re.) striped-flowered.	—	—	H.b.
	γ spicátum.	(wh.) monstrous-flowered.	—	—	H.b.
	δ variegátum.	(wh.) striped-leaved.	—	—	H.b.
2	peregrinum. R.S.	(wh.) drooping-flowered.	—	—	H.b.
3	lanceolínum. R.S.	(wh.) spear-leaved.	Japan.	1824. F.b.
4	longiflórum. R.S.	(wh.) long-flowered.	—	1819.	7. 9. F.b.
	β suavéolens. R.S.	(wh.) one-flowered.	—	—	F.b.
5	japonicum. R.S.	(wh.) Japan.	—	1804.	— F.b.
6	bullíferum. R.S.	(or.) bulb-bearing.	Italy.	1596.	— H.b.
	β umbellátum. B.M.	(or.) umbel-flowering.	—	—	H.b.
7	cróceum. R.S.	(sn.) saffron-color'd.	S. Europe.	—	— H.b.
8	latifólium. L.en.	(or.) broad-leaved.	—	—	H.b.
9	pubéscens. R.S.	(or.) pubescent.	—	—	H.b.
10	spectabíle. F.	(or.) showy Siberian.	Siberia.	1754.	— H.b.
	pensylvánicum. B.M.	t. 872. daúricum. B.M.	1210. in note.		
11	monadélp'hum. B.M.	(y.) monadelphous.	Caucasus.	1800.	6. 7. H.b.
12	cóncolor. s.s.	(sc.or.) self-coloured.	—	1806.	7. 8. F.b.
13	pùmilum. s.s.	(sc.) dwarf.	Dahuria.	1816.	6. 8. H.b.
	linífolium. Horn.	tenuífolium. R.S.			
14	andinum. N.	(sc.) umbel-flowered.	Andes.	1811.	7. 8. H.b.
	umbellátum. Ph.	philadélp'hicum β andinum. B.R.			
15	philadélp'hicum. W.	(or.) Philadelphian.	N. America.	1757.	— H.b.
16	Catesb'æi. w.	(sc.y.) Catesby's.	Carolina.	1787.	7. 9. H.b.
17	carolinianum. M.	(ye.) Carolina.	—	1820.	— H.b.
18	canadénse. B.M.	(ye.) Canadian.	N. America.	1629.	7. 8. H.b.
	β penduliflórum. R.L.	red-flowered.	—	—	H.b.
	canadénse β . rúbra.	Bot. mag. t. 858.			
19	supérbum. w.	(ye.) superb.	—	1727.	6. 8. H.b.
20	Mártagon. w.	(va.) Turk's-cap.	Germany.	1596.	6. 7. H.b.
	α glábrum.	(pu.) smooth-stalked.	—	—	H.b.
	β pubéscens.	(pu.) hairy-stalked.	—	—	H.b.
	γ pállidum. H.	(li.) pale-flowered.	—	—	H.b.
	δ élatum. H.	(pu.) tall.	—	—	H.b.
	ϵ dorsipúnctum. H.	(pu.) spotted backed.	—	—	H.b.
	ζ purpureum. H.	(pu.) purple.	—	—	H.b.
	η perpurpureum. H.	(pu.) dark purple.	—	—	H.b.
	θ ocelláre. H.	(pu.) eyed.	—	—	H.b.
	ι petioláre. H.	(pu.) petioled.	—	—	H.b.
	κ purpùreo-plénüm. H.	(pu.) double purple.	—	—	H.b.
	λ álbum. H.	(wh.) white-flowered.	—	—	H.b.
	μ álbo-plénüm.	(wh.) double white.	—	—	H.b.
21	tigrinum. B.M.	(or.bh.) tiger-spotted.	China.	1804.	7. 9. H.b.
22	pyrenáicum. s.s.	(ye.) Pyrenean.	Pyrenees.	1596.	6. 7. H.b.
23	chaledónicum. w.	(sc.) scarlet Martagon.	Levant.	—	— H.b.
24	pompónium. w.	(sc.) scarlet pomponie.	Siberia.	1629.	5. 6. H.b.

Bot. mag. t. 194.
— t. 1215.

Swt. br. fl. gar. t. 273.

Swt. br. fl. gar. ser. 2.t. 20.

Bot. reg. t. 1152.

Pursh. fl. amer. t. 8.

Bot. mag. t. 278.

Hayn. getr. dars. 8.t. 27.

Bot. reg. t. 560.

Bot. mag. t. 1591.

— t. 36.

— t. 1018.

Redout. liliac. t. 210.

Swt. br. fl. gar. t. 75.

Bot. mag. t. 1405.

— t. 1165.

Bot. reg. t. 132.

Bot. reg. t. 504.

Bot. mag. t. 519.

— t. 259.

Bot. reg. t. 580.

Bot. mag. t. 800.

Redout. lit. t. 105.

Bot. mag. t. 936.

Redout. lit. t. 146.

Bot. mag. t. 1634.

— t. 893.

Bot. mag. t. 1237.

Redout. liliac. t. 145.

Bot. mag. t. 30.

— t. 971.

25 pulchéllum. f.	pretty.	Dahuria.	1829.	H.b.
GLORIOSA. L.	GLORIOSA. Hexandria Monogynia. L.				
1 viréscens. B.M. (gr.)	greenish-flowered.	Mozambique.	1822.	7. 8. S. 4. —.	Bot. mag. t. 2539.
2 supérba. L.	(or.) superb.	E. Indies.	1690.	— S. 4. —.	Bot. reg. t. 77.
3 simplex. L.	(bl.) simple-leaved.	Senegal.	1756.	— S. 4.	
	<i>Methonica gloriösa</i> . Sal.				
ERYTHR'ONIUM. L.	Dog's TOOTH VIOLET.	Hexandria Monogynia. L.			
1 Déns canis. w. (pu.) common.	Europe.	1596.	3. 4. H.b.	Redout. liliac. t. 194.	
2 americanum. B.M. (ye.) yellow-flower'd.	N. America.	1665.	4. 5. H.b.	Bot. mag. t. 1113.	<i>lanceolatum</i> . Ph.
3 álibidum. n. (wh.) white American.	—	1825.	—	H.b.	

ORDO CCXVII.

MELANTHACEÆ. *Brown prod. 272.*

BULBO'CODIUM. W.	BULBO'CODIUM.	Hexandria Monogynia. L.				
1 vérrum. s.s. (ii.)	spring-flowering.	Spain.	1629.	2. 3. H.b.	Bot. mag. t. 153.	
2 versicolor. s.s. (li.ye.)	various-colour'd.	Russia.	1820.	9. 10. H.b.	Bot. reg. t. 571.	
	<i>Cóchicum versicolor</i> . B.R.					
CO'LCHICUM. S.S.	MEADOW SAFFRON.	Hexandria Trigynia. L.				
1 byzantinum. s.s. (li.)	broad-leaved.	Levant.	1629.	9. 10. H.b.	Bot. mag. t. 1122.	
2 autumnale. E.B.	(li.) common.	Britain.	—	— H.b.	Eng. bot. v. 2. t. 133.	
β álbum.	(wh.) white flowered.	—	—	— H.b.		
γ multipléx.	(li.) double-flowered.	—	—	— H.b.		
3 crocifòrum. B.M.	(li.) Crocus-flowered.	—	—	— H.b.	Bot. mag. t. 2673.	
4 variegátum. w. (std.)	chequer-flowered.	Greece.	1629.	8. 10. H.b.	— t. 1028.	
5 arenárium. s.s. (ii.)	sand.	Hungary.	1816.	9. 10. H.b.	W. et K. hung. 2. t. 179.	
6 umbrósium. f.	(li.) wood.	Siberia.	—	— H.b.	Bot. reg. t. 541.	
7 montánum. s.s. (li.)	mountain.	S. Europe.	1818.	— H.b.		
UVUL'ARIA. S.S.	UVUL'ARIA.	Hexandria Monogynia. L.				
1 perfoliáta. s.s. (gr.ye.)	perfoliate.	N. America.	1710.	5. 6. H. 4.	Sm. exot. bot. I. t. 49.	
2 fláva. s.s. (ye.)	yellow-flowered.	—	—	— H. 4.	— 1. t. 50.	
3 lanceoláta. H.K.	(ye.) spear-leaved.	—	1710.	7. 8. H. 4.	Cornut. canad. t. 41.	
4 grandiflóra. H.K.	(ye.) large yellow.	—	1802.	5. 6. H. 4.	Sm. exot. bot. I. t. 51.	
5 sessilifólia. s.s. (st.)	sessile-leaved.	—	1790.	6. H. 4.	— 1. t. 52.	
6 pubérula. s.s. (ye.)	pubescent.	—	1818.	— H. 4.	Swt.br.fl.gar.s.2.t.21.	
DISP'ORUM. D.P.	Disp'ORUM.	Hexandria Monogynia.				
1 parvifòrum. D.P.	(ye.) small-flowered.	Nepaul.	1823.	6. 8. F. 4.		
2 Pitsútum. D.P.	(ye.) umbel-flowered.	—	1826.	— F. 4.		
3 fúlvum. Sal. (br.pu.)	brown-flowered.	China.	1801.	6. 11. F. 4.	Bot. mag. t. 916.	
	<i>Uvularia chinensis</i> . B.M.					
SCHELHAMM'ERA. B.P.	SCHELHAMM'ERA.	Hexandria Monogynia. S.S.				
1 unduláta. B.P.	(ro.) wave-leaved.	N. S. W.	1825.	6. 9. F. 4.	Bot. mag. t. 2712.	
2 multiflóra. B.P.	(ro.) many-flowered.	—	1826.	— F. 4.		
BURCHA'R'DIA. B.P.	BURCHA'R'DIA.	Hexandria Monogynia. S.S.				
	umbelláta. B.P.	(wh.) umbel-flowered.	N. S. W.	1825.	6. 8. F. 4.	
ANGUILL'ARIA. B.P.	ANGUILL'ARIA.	Hexandria Trigynia.				
1 dioica. B.P.	(br.) dioecious.	N. S. W.	1826.	— F. 4.		
2 biglandulósa. B.P.	(br.) two-glandled.	—	1822.	6. 8. F. 4.		
3 indica. B.P.	(d.pu.) Indian.	E. Indies.	—	— F. 4.		
	<i>Melánthium indicum</i> . L.					
ORNITHOGLO'SSUM. S.S.	BIRD'S-TONGUE.	Hexandria Trigynia. S.S.				
1 undulátum. B.F.G.(g.pu.)	wave-leaved.	C. B. S.	1825.	8. 10. F.b.	Swt. br. fl. gar. t. 131.	
2 víride. s.s. (gr.br.)	green-flowered.	—	1788.	10. 11. F.b.	Sal. par. lond. t. 54.	
	<i>Melánthium víride</i> . Bot. mag. t. 994.					
ANDROCY'MBIUM. W.	ANDROCY'MBIUM.	Hexandria Trigynia. S.S.				
1 melanthioídes. s.s. (gr.)	narrow-leaved.	C. B. S.	1825.	8. 10. F.b.		
2 euromoídes. s.s. (gr.)	Tulip-leaved.	—	1794.	3. 5. F.b.	Swt. br. fl. gar. t. 165.	
	<i>Melánthium euromoídes</i> . B.M. 641.					
3 volutáre. B.T.	(gr.) rolled-leaved.	—	1816.	— F.b.		
MELA'NTHIUM. W.	MELA'NTHIUM.	Hexandria Trigynia. L.				
1 júnceum. w. (li.)	Rush-leaved.	C. B. S.	1788.	6. 11. F.b.	Bot. mag. t. 558.	
2 secundum. w. (wh.)	side-flowering.	—	1812.	— F.b.	Lam. ill. f. 269. f. 2.	

3 capénsē. w. (wh.)	spotted-flowered.	C. B. S.	1768.	5. 6. F.b.	
4 ciliatum. w. (wh.)	fringed.		1825.	— F.b.	
5 gramineum. s.s. (wh.)	grassy.	Barbary.	1823.	— F.b.	
KO'LBEA. R.S.	Ko'LBEA.	Hexandria Trigynia.			Cavan. icon. t. 587, f. 1.
Breyniâna. R.S. (ye.)	Breynius's.	C. B. S.	1787.	6. 7. F.b.	Bot. mag. t. 767.
Melânthium uniflorum. w.					
WURMB'EA. W.	WURMB'EA.	Hexandria Trigynia. W.			
1 campanulata. w. (wh.)	bell-flowered.	C. B. S.	1788.	5. 6. F.b.	Bot. mag. t. 1291.
Melânthium monopétalum. B.M.					
2 purpurea. H.K. (pu.)	purple.		—	— F.b.	Bot. mag. t. 694.
Melânthium spicatum. B.M.	Wurmbèa capensis.	Andr. deposit.	t. 221.		
3 pumila. w. (wh.)	dwarf.	C. B. S.	1800.	5. 6. F.b.	
4 longiflora. w. (wh.)	long-flowered.		—	— F.b.	
TOFIELDIA. L.T.	TOFIELDIA.	Hexandria Trigynia. S.S.			
1 palustris. L.T. (st.)	Scotch.	Britain.	7. 8. H. 4.	Eng. bot. v. 8. t. 536.
2 racemosa. Hop. (st.)	racemed.	Germany.	1827.	— H. 4.	Linn. trans. v. 12. t. 8. f. 2.
3 alpina. L.T. (st.)	Alpine.	Europe.	7. 8. H. 4.	Redout. lil. t. 256.
palustris. DC.					
4 pubens. L.T. (st.)	downy.	N. America.	1790.	4. 5. H. 4.	
5 glutinosa. L.T. (ye.)	glutinous.		1827.	— H. 4.	
XEROPHYLLUM. M.	XEROPHYLLUM.	Hexandria Trigynia. S.S.			
1 gramineum. s.s. (wh.)	grass-leaved.	N. America.	1812.	5. 6. H. 4.	
Heloniâs graminea. B.M.					
2 tenuax. s.s. (wh.)	tough-leaved.		1811.	— H. 4.	Pursh. amer. I. t. 9.
3 asphodeloides. s.s. (wh.)	Asphodel-like.		1765.	— H. 4.	Bot. mag. t. 748.
Heloniâs asphodeloides. B.M.					
HELONIAS. W.	HELONIAS.	Hexandria Trigynia. L.			
1 bullata. w. (pu.)	broad-leaved.	N. America.	1758.	4. 5. H. 4.	Bot. mag. t. 747.
latifolia. M.					
2 erythrosperma. M. (wh.)	channel-leaved.		1770.	6. H. 4.	Bot. mag. t. 803.
l'æta. B.M.	Melânthium l'ætum. w.				
3 angustifolia. M. (wh.)	narrow-leaved.		1812.	— H. 4.	Bot. mag. t. 1540.
ZIGAD'ENUS. M.	ZIGAD'ENUS.	Hexandria Monogynia. S.S.			
1 glaberrimus. w. (wh.)	smooth-leaved.	N. America.	1811.	5. 6. H. 4.	Mich. amer. I. t. 22.
Heloniâs glaberrima. Bot. mag. t. 1680.					
2 bracteatus. (wh.)	bracted.		—	— H. 4.	Bot. mag. t. 1703.
Heloniâs bracteata. B.M.					
3 élégans. Ph. (wh.)	elegant.		1830.	6. 7. H. 4.	
NOL'INA. M.	NOL'INA.	Hexandria Trigynia. S.S.			
georgiana. M. (wh.)	Georgian.	Georgia.	1812.	7. 8. H. 4.	Pluk. mant. t. 342. f. 1.
LEIMA'NTHIUM. W.	LEIMA'NTHIUM.	Polygamia Monococcia.			
virginicum. w. (wh.)	Virginian.	N. America.	1768.	6. 7. H. 4.	Bot. mag. t. 985.
Heloniâs virginica. B.M.	Melânthium virginicum. Ph.				
2 monoicum. (wh.)	monoecious.		1817.	— H. 4.	
Melânthium monoicum. Ph.					
3 hy'bridum. (wh.)	hybrid.		1822.	— H. 4.	
Melânthium hy'bridum. Ph. racemosum. M.					
PELIOSA'NTHES. A.R. PELIOSA'NTHES.		Hexandria Monogynia.			
1 humiliis. A.R. (gr.)	small.	E. Indies.	1809.	5. 6. S. 4.	Bot. mag. t. 1532.
2 Tête. A.R. (gr.)	large.		1807.	4. S. 4.	t. 1302.
CYM'ATION. S.S.	CYM'ATION.	Hexandria Trigynia. R.S.			
laevigatum. s.s. (bl.)	smooth.	C. B. S.	1824.	4. 5. G.b.	
Lichensteinia laevigatu. w.					
VER'ATRUM. L.	WHITE HELLEBORE.	Polygamia Monococcia. L.			
1 nigrum. w. (da.pu.)	dark-flowered.	Siberia.	1596.	6. 7. H. 4.	Bot. mag. t. 963.
2 album. w. (wh.)	white-flowered.	S. Europe.	1548.	6. 8. H. 4.	Flor. dan. t. 1120.
3 Lobeliânum. W.en.(wh.)	Lobel's.		1818.	— H. 4.	
4 víride. w. (gr.)	green-flowered.	N. America.	1742.	7. 8. H. 4.	Bot. mag. t. 1096.
5 parviflorum. M. (gr.)	small-flowered.		1809.	— H. 4.	
6 angustifolium. Ph. (gr.)	narrow-leaved.		1823.	— H. 4.	
CHAMÆLI'RUM. W.	CHAMÆLI'RUM.	Diœcia Hexandria.			
carolinianum. w. (st.)	spiked-flowered.	N. America.	1759.	7. 8. H. 4.	Bot. mag. t. 1062.
Heloniâs lutea. B.M.	dioica. Ph. Ophyostachys virginica.	Redout. liliac. t. 464.			

ORDO CCXVIII.

PONTEDEREÆ. *Kth. synops.* 1. p. 273.**PONTED'ERIA.** L. PONTED'ERIA. Hexandria Monogynia. L.

- 1 cordàta. s.s. (*bl.*) heart-leaved. N. America. 1759. 6. 8. H.w. 4. Bot. mag. t. 1156.
 2 azùrea. s.s. (*pu.*) large-flowered. Jamaica. 1822. — S.w. 4. — t. 2932.
 3 lanceolàta. s.s. (*bl.*) lanceolate. Carolina. 8. 9. G.w. 4. Lodd. bot. cab. t. 613.
 4 angustifòlia. s.s. (*bl.*) narrow-leaved. N. America. 1806. 6. 8. H.w. 4.
 5 crassipè. s.s. * (*bl.*) thick-footstalked. Brazil. 1822. — S.w. 4. Mart. n. pl. bras. t. 4.
 6 dilatata. s.s. (*bl.*) spreading. E. Indies. 1806. 5. S.w. 4. Andr. deposit. t. 490.

HETERANTH'ERA. R.S. HETERANTH'ERA. Triandria Monogynia. W.

- 1 reniformis. R.S. (*wh.*) kidney-leaved. S. America. 1812. 6. 7. G.w. 4. Flor. per. 1. t. 71. f. a.
 2 acùta. w. (*wh.*) acute-leaved. N. America. — — H.w. 4.
 3 limosa. R.S. (*bl.*) oval-leaved. W. Indies. 1822. — S.w. 4. Mich. amer. 1. t. 5. f. 1.

LEPTA'NTHUS. M. LEPTA'NTHUS. Triandria Monogynia. Ph.

- gramineus. M. (*ye.*) grass-like. N. America. 1823. 7. 8. H.w. 4. Hook. ex. flor. t. 94.
Heteranthera graminea. Ph. Schölleria graminifolia. w.

ORDO CCXIX.

COMMELINEÆ. *Brown prodr.* 1. p. 268.**DICHORISA'NDRA.** S.S. DICHORISA'NDRA. Hexandria Monogynia. S.S.

- 1 thyrsiflòra. s.s. (*bl.*) bunch-flowered. Brazil. 1820. 8. 10. S. 4. Bot. reg. t. 682.
 2 grátilis. s.s. (*bl.*) slender. — — S. 4.
 3 oxypétala. B.M. (*pu.*) sharp-flowered. — — 1825. 6. 9. S. 4. Bot. mag. t. 2721.
 4 pubérula. s.s. (*bl.*) pubescent. — — S. 4.

CAMP'ELIA. K.S. CAMP'ELIA. Hexandria Monogynia. S.S.

- Zanònìa. k.s. (*wh.*) Gentian-leaved. S. America. 1759. 7. 12. S. 4. Redout. lil. 4. t. 192.
Commelinæ Zanònìa. R.L. *Tradescantia Zanònìa.* w. *Zanònìa bibracteata.* Cram.

TRADESCA'NTIA. L. SPIDERWORT. Hexandria Monogynia. L.

- 1 fuscata. s.s. (*bl.*) rusty. S. America. 1820. 8. 12. S. 4. Bot. reg. t. 482.
 2 malabárica. s.s. (*pu.*) Grass-leaved. E. Indies. 1776. 7. 8. S. 4. Rheed. mal. 9. t. 63.
 3 tuberòsa. s.s. (*bl.*) tuberous-rooted. — — 1817. — S. 4.
 4 speciòsa. s.s. (*pu.*) handsome. Mexico. 1825. — S. 4.
 5 discolor. s.s. (*wh.*) purple-leaved. S. America. 1783. 4. 9. S. 4. Bot. mag. t. 1192.
 6 cordifòlia. s.s. (*pu.*) heart-leaved. Jamaica. 1822. — S. 4.
 7 erécta. s.s. (*bl.*) upright. Mexico. 1794. 7. 8. H. 4. Bot. mag. t. 1340.
 8 undáta. s.s. (*bl.*) wave-leaved. S. America. 1817. 8. 10. H. 4.
 9 pulchella. s.s. (*pu.*) pretty. Mexico. 1825. — G. 4.
 10 crassifòlia. s.s. (*pu.*) thick-leaved. — — 1796. 7. 10. H. 4. Bot. mag. t. 1598.
 11 subáspera. s.s. (*bl.*) Lyon's. N. America. 1812. 5. 10. H. 4. — t. 1597.
 12 virginica. s.s. (*bl.*) Virginian. — — 1629. 5. 8. H. 4. Park. parad. t. 151. f. 4.
 α cærulea. (*bl.*) blue-flowered. — — — H. 4. Bot. mag. t. 105.
 β rúbra. (*re.*) red-flowered. — — — H. 4.
 γ álbida. (*wh.*) whitish-flowered. — — — H. 4.
 δ nivea. (*wh.*) snowy white. — — 1824. — H. 4.
 ε pilosa. B.R. (*bl.*) hairy. — — 1827. — H. 4. Bot. reg. t. 1055.
 13 congesta. H.E. (*bl.*) close-flowered. — — 1825. — H. 4.
 14 rosea. s.s. (*ro.*) rose-coloured. Carolina. 1802. 5. 10. H. 4. Swt. br. fl. gar. t. 181.
 15 multiflòra. s.s. (*wh.*) many-flowered. Jamaica. 1824. — S. 4.
 16 crássula. B.M. (*wh.*) white-flowered. Brazil. 1826. 12. 1. S. 4. Bot. mag. t. 2935.
 17 procumbens. s.s. (*wh.*) procumbent. Caracas. 1823. — S. 4. Jacq. ic. 2. t. 355.
 multiflòra. Jacq. non w.
 18 parviflòra. s.s. (*bl.*) small-flowered. Peru. 1822. — H. 4. Flor. per. 3. t. 272. f. 2.
 19 latifòlia. s.s. (*vi.*) broad-leaved. Lima. 1820. 9. 11. H. 4. — 3. t. 272.
 20 divaricàta. s.s. (*bl.*) spreading. Trinidad. — — 5. 8. S. 4. Aub. gui. 1. t. 12.
 21 geniculàta. s.s. (*wh.*) knotted. W. Indies. 1783. 7. 8. S. 4. Jacq. amer. t. 64.
 22 paniculàta. s.s. (*bl.*) panicle. E. Indies. 1824. — S. 4.

CYAN'OTIS. D.P.	CYAN'OTIS.	Hexandria Monogynia.
1 axillâris. D.P.	(bl.) axil-flowered.	E. Indies. 1822. 8. 10. S. ⊙.
2 cristâta. D.P.	(bl.) crest-bunched.	Ceylon. 1770. 7. 9. S. ⊙.
Tradescântia cristâta. B.M.		
3 barbâta. D.P.	(bl.) bearded.	Nepaul. 1824. — H. ⊙.
CARTON'EMA. B.P.	CARTON'EMA.	Hexandria Monogynia.
spicatum. B.P.	(ye.) spiked.	New Holland. 1822. 7. 8. S. ¼.
CALLI'SIA. L.	CALLI'SIA.	Triandria Monogynia. L.
rêpens. L.	(gr.) creeping.	W. Indies. 1776. 6. 7. S. ¼.
ANEIL'EMA. B.P.	ANEIL'EMA.	Triandria Monogynia.
1 biflôra. B.P.	(bl.) two-flowered.	N. S. W. 1820. 7. 8. F. ¼.
2 acuminâta. B.P.	(bl.) sharp-pointed.	—— 1822. — F. ¼.
3 longifòlia. H.E.F.	(bl.) long-leaved.	Zanzibar. 1823. 8. 9. S. ¼.
4 sínica. B.R.	(bl.) Chinese.	China. 1820. 5. 6. G. ¼.
5 nudiflôra. B.P. (bl. pu.)	naked-flowered.	E. Indies. — 8. 10. S. ⊙.
6 spirâta. B.P.	(bl.) spear-leaved.	—— 1783. 7. 8. S. ⊙.
Commelina spirâta. L.		
7 affinis. B.P.	(bl.) rough-angled.	N. S. W. 1825. — F. ¼.
COMMEL'INA. B.P.	COMMEL'INA.	Triandria Monogynia. L.
1 angustifòlia. R.S.	(bl.) narrow-leaved.	Carolina. 1812. 7. 8. F. ¼.
2 truncâta. R.S.	(pa.bl.) truncate-bracted.	—— 1822. 7. 9. H. ¼.
3 cyánea. B.P.	(bl.) bright blue.	N. S. W. 1820. — F. ¼.
4 dianthifòlia. DC.	(bl.) Pink-leaved.	—— 1816. — S. ¼.
5 longicaùlis. R.S.	(bl.) long-stalked.	Caracas. 1806. 8. S. ¼.
6 hirtella. R.S.	(bl.) hairy.	N. America. 1820. 7. 8. H. ¼.
7 virgíñica. R.S.	(bl.) Virginian.	—— 1779. — H. ¼.
8 africana. R.S.	(ye.) yellow-flowered.	C. B. S. 1759. 5. 10. G. ¼.
9 fasciculâta. R.S.	(bl.) woolly tubered.	Lima. 1817. — H. ¼.
10 communis. R.S.	(bl.) common.	America. 1732. 6. 7. H. ⊙.
11 polygamâ. R.S.	(bl.) polygamous.	Japan. — — H. ¼.
12 caroliniana. R.S.	(bl.) Carolina.	Carolina. — — H. ¼.
13 dùbia. R.S.	(bl.) dubious.	—— 1819. 7. 9. H. ¼.
14 tuberôsa. R.S.	(bl.) tuberous-rooted.	Mexico. 1732. — H. ¼.
15 cœlestis. R.S.	(bl.) sky-blue.	—— 1813. 7. 10. H. ¼.
16 pálida. R.S.	(bl.) pale-flowered.	Mexico. 1822. — H. ¼.
rûbens. Redoute liliac.	7. t. 367.	
17 turbinâta. R.S.	(bl.) top-shaped.	Santa Cruz. 1824. — S. ¼.
18 obliqua. R.S.	(bl.) oblique-leaved.	S. America. 1827. — S. ¼.
19 erecta. R.S.	(bl.) upright.	Virginia. 1732. 8. 9. H. ¼.
20 mollis. R.S.	(bl.) soft.	Caracas. 1804. — H. ¼.
21 bengalensis. R.S.	(bl.) Bengal.	Bengal. 1794. 6. S. ¼.
22 parviflora. L.en.	(bl.) small-flowered.	—— 1823. — S. ¼.
23 deficiens. B.M.	(bl.) deficient.	Brazil. 1825. 1. 12. S. ¼.

Rheed. mal. 10. t. 13.
Bot. mag. f. 1435.

Jacq. amer. t. 11.

Hook. ex. flor. t. 204.
Bot. reg. t. 659.
Rheed. mal. 9. t. 63.

Redout. lil. 7. t. 390.
Jacq. ic. 2. t. 294.

Pluk. alm. t. 174. f. 4.
Bot. mag. t. 1431.
Flor. per. 1. t. 72. f. b.
Redout. lil. 4. t. 206.
Kæmpf. am. p. 888. ic.

Redout. lil. 6. t. 359.
Andr. reposit. t. 399.
Swt. br. fl. gar. t. 3.
Willd. hort. ber. 2.t.87,

Dill. elt. 94. t. 77. f. 88.
Jacq. ic. 2. t. 293.
Burm. ind. t. 7. f. 3.

Bot. mag. t. 2644.

ORDO CCXX.

PALMÆ. Brown prodr. p. 266.

CHAMÆD'OREA. W. CHAMÆD'OREA. Diœcia Hexandria. W.

1 grâcilis. W.	(st.) slender.	Caracas. 1803. S. ½.
2 frâgrans. M.P.	(st.) fragrant.	Peru. 1823.

Jacq. scheen. 2. t. 247-8.
Martius Palm. ic.

S'ABAL. P.S. S'ABAL. Hexandria Trigynia. P.S.

1 Adansoni. B.M.	(st.) dwarf.	Florida. 1810. 6. 8. S. ¼.
Córypha minor.	Jacq. vind. 3. t. 8. <i>Chamærops acâulis</i> . M.	
2 Blackburniana. L.C.	(st.) Blackburn's.	S. America. 1824. S. ¼.
3 graminifolia. L.C.	(st.) grass-leaved.	—— 1822. S. ¼.
4 umbraculifera. L.C.	(st.) umbrella.	—— — — S. ¼.

Bot. mag. t. 1434.

RHA'PIS. W. RHA'PIS. Polygamia Monœcia. W.

1 flabelliformis. W.	(st.) creeping-rooted.	China. 1774. 8. S. ¼.
2 arundinacea. W.	(st.) simple-leaved.	Carolina. 1765. 9. G. ½.

Bot. mag. t. 1371.

CHAMÆROPS. W. CHAMÆROPS. Polygamia Dicecia. W.

1 excelsa. L.C.	(st.) tall.	Nepaul. 1822. S. ½.
2 grâcilis. L.C.	(st.) slender.	S. America. — — S. ½.
3 guianensis. L.C.	(st.) Guiana.	Guiana. 1824. S. ½.
4 humiliis. S.S.	(st.) dwarf fan palm.	S. Europe. 1731. 2. 3. G. ½.

Andr. reposit. t. 599.

5 serrulata. s.s.	(st.) saw-leaved.	N. America.	1809.	G. h.
6 Hy'strix. s.s.	(st.) Porcupine.	Georgia.	1801.	G. h.
7 Palmetto. s.s.	(st.) smooth-stem'd.	Carolina.	1809.	G. h.
LIVIST'ONA. B.P.	LIVIST'ONA.	Hexandria Monogynia.	S.S.		
1 inermis. B.P.	(st.) spineless.	N. Holland.	1823.	S. h.
2 humilis. B.P.	(st.) dwarf spiny.	—	—	S. h.
LODOI'CEA. S.S.	DOUBLE COCOA-NUT.	Dicecia Monadelphia.			
Sechellârum. B.M.(st.)	Seychelle's Island.	Seychelle's Isl.	1827.	S. h.
LAT'ANIA. W.	LAT'ANIA.	Dicecia Monadelphia. W.			
1 rufra. w.	(st.) red.	Mauritius.	1788.	S. h.
2 glaucophylla. L.C.	(st.) glaucous-leaved.	—	1821.	S. h.
3 borbonica. w.	(st.) Bourbon.	Bourbon.	—	S. h.
chinensis. Jacq.	Livistonæ species? B.P.				Jacq. frag. 1. t. 11.
CO'RYPHA. L.	FAN PALM.	Hexandria Monogynia.	W.		
1 australis. B.P.	(st.) New Holland.	N. S. W.	1822.	S. h.
2 elata. H.B.	(st.) tall.	E. Indies.	1820.	S. h.
3 glaucescens. L.C.	(st.) glaucescent.	—	—	S. h.
4 umbraculifera. w.	(st.) great.	—	1742.	S. h.
5 Után. s.s.	(st.) Molucca.	—	1822.	S. h.
TAL'IERA. S.S.	TAL'IERA.	Hexandria Monogynia.	S.S.		
bengalensis. s.s.	(st.) Bengal.	E. Indies.	1822.	S. h.
Corypha Taliera. H.B.					
PH'ENIX. W.	DATE PALM.	Dicecia Hexandria.	W.		
1 dactylifera. w.	(st.) common.	Levant.	1597.	G. h.
2 sylvestris. H.B.	(st.) wood.	E. Indies.	1820.	S. h.
3 paludosa. H.B.	(st.) marsh.	—	—	S. h.
4 reclinata. w.	(st.) reclining.	C. B. S.	1792.	G. h.
5 farinifera. w.	(st.) small.	E. Indies.	1800.	S. h.
6 acaulis. s.s.	(st.) stalkless.	—	1820.	S. 4.
7 leonensis. L.C.	(st.) African.	Sierra Leon.	—	S. h.
8 pygmæa. L.C.	(st.) pigmy.	—	—	S. 4.
S'AGÜS. W.	SAGO PALM.	Monoclea Hexandria.	W.		
1 Rúmpii. w.	(st.) Rumphius's.	E. Indies.	1800.	S. h.
2 vinifera. P.S.	(st.) prickly-leaved.	Africa.	1820.	S. h.
3 Rúffia. w.	(st.) Madagascar.	Madagascar.	—	S. h.
4 pedunculata. Poir.	(st.) peduncled.	—	—	S. h.
HYP'H'ENE. G.	HYP'H'ENE.	Dicecia Hexandria.			
coriacea. G.	(st.) leathery.	Egypt.	1822.	S. h.
Cucifera thebäica. Del.					
AR'ECA. W.	CAEPRAGE-TREE.	Monoclea Monadelphia.	W.		
1 Cátechu. w.	(st.) medicinal.	E. Indies.	1690.	S. h.
2 humilis. w.	(st.) dwarf.	—	1814.	S. h.
3 rufra. w.	(st.) red.	Isl. France.	1823.	S. h.
4 crinita. w.	(st.) hairy.	—	1819.	S. h.
5 oleracea. w.	(st.) esculent.	W. Indies.	1656.	S. h.
6 triandra. H.B.	(st.) three-anthered.	E. Indies.	1819.	S. h.
7 lutescens. w.	(st.) yellowish.	Mauritius.	—	S. h.
8 exilis. L.C.	(st.) slender.	—	1822.	S. h.
9 Mánicot. L.C.	(st.) Manicot.	—	—	S. h.
10 montaña. L.C.	(st.) mountain.	—	1819.	S. h.
ENOCA'RPUS. M.P.	ENOCA'RPUS.	Monoclea Hexandria.			
Batáua. M.P.	(st.) American.	S. America.	1822.	S. h.
EUTE'RPE. G.	EUTE'RPE.	Polygamia Monoclea.			
1 globosa. G.	(st.) globular.	E. Indies.	1819.	S. h.
Arèca spicata. Lam.					
2 pisifera. G.	(st.) Pea-bearing.	—	—	S. h.
WALLI'CHIA. S.S.	WALLI'CHIA.	Polygamia Monoclea.			
caryotoïdes. s.s.	(st.) Caryota-like.	E. Indies.	1818.	S. h.
EL'AIS. W.	OILY PALM.	Dicecia Hexandria.	W.		
1 guineensis. w.	(st.) Guinea.	Guinea.	1730.	S. h.
2 melanococca. G.	(st.) black-fruited.	S. America.	1820.	S. h.
3 pernambucana. L.C.	(st.) Pernambuca.	—	—	S. h.
SYA'GRUS. M.P.	SYA'GRUS.	Polygamia Monoclea.			
cocoïdes. M.P.	(st.) Cocoa-like.	Brazil.	1823.	S. h.
EL'ATE. W.	EL'ATE.	Monoclea Hexandria.			
sylvestris. w.	(st.) prickly-leaved.	E. Indies.	1763.	4. 6.	S. h.
ACROC'OMIA. M.P.	ACROC'OMIA.	Monoclea Hexandria.			
1 sclerocarpa. M.P.	(st.) dry-fruiting.	Brazil.	1822.	S. h.
2 aculeata. L.C.	(st.) prickly.	W. Indies.	1796.	S. h.
Côcos aculeata. w.					

3 fusiformis.	(st.) great Macaw-tree.	W. Indies.	1731. S. h.
Cocos fusiformis. w.				
4 mīnor. L.C.	(st.) lesser.	S. America.	1823. S. h.
COCOS. W.	COCOA-NUT-TREE.	Monecia Hexandria. W.		
1 nucifera. w.	(st.) common.	E. Indies.	1690. S. h.
2 flexuosa. M.P.	(st.) flexuous.	Brazil.	1823. S. h.
3 plumosa. L.C.	(st.) feathered.	W. Indies.	— S. h.
MAXIMIL'ANA. M.P.	MAXIMILI'ANA.	Polygamia Monocia.		
rēgia. M.P.	(st.) membranous-fl'd.	Brazil.	1822. S. h.
THR'INAX. W.	THR'INAX.	Hexandria Monogynia. W.		
parviflora. w.	(st.) small.	Jamaica.	1778. S. h.
ASTROC'ARYUM. M.P.	ASTROC'ARYUM.	Monocia Hexandria.		
1 acaule. M.P.	(st.) stalkless.	Brazil.	1823. S. 4.
2 campéstre. M.P.	(st.) field.	—	1824. S. h.
3 aculeatum. s.s.	(st.) prickly.	Guiana.	— S. h.
4 vulgare. M.P.	(st.) common.	Brazil.	1823. S. h.
5 Murumūru. M.P.	(st.) tall.	—	— S. h.
GEON'OMA. W.	GEON'OMA.	Dicecia Hexandria.		
1 simplicifrons. w.	(st.) simple-fronded.	S. America.	1823. S. 4.
2 macrostachys. M.P.	(st.) large-spiked.	—	— S. 4.
3 acaulis. M.P.	(st.) stalkless.	Brazil.	1822. S. 4.
4 pinnatifrons. w.	(st.) pinnated.	S. America.	— S. 4.
5 Schottiana. L.C.	(st.) Schott's.	Brazil.	— S. 4.
MAURITIA. L.	MAURITIA.	Dicecia Hexandria. W.		
1 flexuosa. w.	(st.) flexuous.	W. Indies.	1819. S. h.
2 armata. M.P.	(st.) spiny.	Brazil.	1822. S. h.
3 vinifera. M.P.	(st.) straight-stem'd.	—	— S. h.
LICUALA. W.	LICUALA.	Hexandria Monogynia. W.		
1 spinosa. w.	(st.) spiny.	E. Indies.	1802. S. h.
2 peltata. H.E.	(st.) peltate.	—	1819. S. h.
CA'LAMUS. W.	CA'LAMUS.	Hexandria Monogynia. S.S.		
1 vērus. s.s.	(st.) true.	E. Indies.	1820. S. h.
2 niger. s.s.	(st.) black.	—	— S. h.
3 rūdētum. s.s.	(st.) common.	China.	1812. S. h.
4 Zalācca. s.s.	(st.) Java.	E. Indies.	— S. h.
5 álbus. P.S.	(st.) common.	—	— S. h.
6 Drāco. s.s.	(st.) Dragon.	—	1819. S. h.
SEAFO'RTHIA. B.P.	SEAFO'RTHIA.	Polygamia Monocia.		
élégans. B.P.	(st.) elegant.	N. Holland.	1822. S. h.
CARY'OTA. W.	CARY'OTA.	Monocia Polyandria. W.		
1 ürens. w.	(st.) torn-leaved.	E. Indies.	1788. S. h.
2 hórida. w.	(st.) horrid.	S. America.	1820. S. h.
DIPLOTH'EMIUM. M.P.	DIPLOTH'EMIUM.	Polygamia Monocia.		
1 marítimum. M.P.	(st.) sea-side.	Brazil.	1823. S. 4.
2 campéstre. M.P.	(st.) field.	—	— S. 4.
ATT'ALEA. K.S.	ATT'ALEA.	Monocia Polyandria.		
1 hūmilis. M.P.	(st.) dwarf.	Brazil.	1823. S. h.
2 cómpta. M.P.	(st.) decked.	—	— S. h.
3 funifera. M.P.	(st.) rope-bearing.	—	— S. h.
4 excésa. M.P.	(st.) lofty.	—	— S. h.
5 speciosa. M.P.	(st.) handsome.	—	1824. S. h.
6 Róssii. L.C.	(st.) Ross's.	—	— S. h.
7 spectabilis. L.C.	(st.) showy.	—	— S. h.
GOM'UTUS. S.S.	GOM'UTUS.	Monocia Polyandria.		
sacharifer. s.s.	(st.) sugar-bearing.	E. Indies.	1820. S. h.
MANICARIA. G.	MANIC'ARIA.	Polygamia Monocia.		
saccifera. g.	(st.) bag-like.	E. Indies.	1822. S. h.
BORA'SSUS. W.	BORA'SSUS.	Dicecia Hexandria. W.		
flabellifōrmis. w.	(st.) fan-leaved.	E. Indies.	1771. S. h.
N'IPA. W.	N'IPA.	Monocia Monadelphia. W.		
frūticans. w.	(st.) shrubby.	E. Indies.	1822. S. h.
DESMO'NCUS. M.P.	DESMO'NCUS.	Monocia Monadelphia.		
1 polyacánthus. M.P.	(st.) many-spined.	Brazil.	1822. S. h.
2 orthacánthus. M.P.	(st.) long-spined.	—	— S. h.
3 americanus. L.C.	(st.) American.	S. America.	1823. S. h.
4 dūbius. L.C.	(st.) doubtful.	—	— S. h.
BA'CTRIS. W.	BA'CTRIS.	Monocia Hexandria. W.		
1 pectinata. M.P.	(st.) pectinated.	Brazil.	1825. S. h.
2 cuspidata. M.P.	(st.) sharp-pointed.	—	— S. h.

Roxb. corom. 1. t. 73.
Mart. palm. ic.

Mart. palm. ic.

Mart. palm. ic.
_____ ic.

Rumph. amb. 1. t. 9.

Rumph. amb. 5. t. 54.
_____ 5. t. 52.

Gært. fr. 2. t. 139. f. 2.
Rumph. amb. 5. f. 52.
_____ 5.t.58.f.1

Rheed. mal. 1. t. 11.

Mart. palm. ic.
_____ ic.

Mart. palm. ic.
_____ ic.

Mart. palm. ic.
_____ * ic.

Mart. palm. ic.
_____ 2. c.

Gært. sem. 2. t. 176.

Roxb. corom. 1. t. 71.72.

Rumph. amb. 1. t. 16.

Mart. palm. ic.
_____ ic.

Mart. palm. ic.

3 minor. w.	(st.) lesser.	S. America.	1691.	S. h.	Jacq. amer. t. 171. f. 1.
4 macracantha. M.P.	(st.) large-spined.	Brazil.	1822.	S. h.	Mart. palm. ic.
5 mäjor. w.	(st.) greater.	Carthagena.	1800.	S. h.	Jacq. amer. t. 171. f. 2.
6 caryotafolia. M.P.	(st.) Caryota-leaved.	Brazil.	1825.	S. h.	Mart. palm. ic.

ORDO CCXXI.

JUNCEÆ. Brown prodr. 1. p. 257.

KI'NGIA. L.T.	GRASS-TREE.	Hexandria Monogynia.				
australis. L.T.	(st.) southern.	N. Holland.	1830.	G. h.	
XER'OTES. B.P.	XER'OTES.	Dicecia Hexandria.				
1 flexifolia. B.P.	(st.) bending-leav'd.	N. S. W.	1819.	6. 8.	F. 4.	Thunb. diss. drac. 17. f. 2.
2 mucronata. B.P.	(st.) mucronated.	—	1823.	—	F. 4.	
3 filiformis. B.P.	(st.) filiform.	—	—	—	F. 4.	Th. dis. drac. n. 1. ic.
<i>Drac'ana filiformis.</i> Th.						
4 denticulata. B.P.	(st.) toothed.	—	1825.	—	F. 4.	
5 laxa. B.P.	(st.) loose-flowered.	—	1819.	—	F. 4.	
6 rigida. B.P.	(st.) stiff-leaved.	N. Holland.	1796.	6. 7.	F. 4.	Lodd. bot. cab. t. 798.
<i>Lomandra rigida.</i> Lab. n. hol. 1. t. 120.						
7 montana. B.P.	(st.) mountain.	N. S. W.	1824.	—	F. 4.	
8 longifolia. B.P.	(st.) long-leaved.	—	1796.	—	F. 4.	Lab. nov. hol. 1. t. 19.
<i>Lomandra longifolia.</i> Lab.						
9 Hy'strix. B.P.	(st.) Porcupine.	—	1823.	—	F. 4.	
10 'æmula. B.P.	(st.) rival.	—	—	—	F. 4.	
11 echinata. M.C.	(st.) bristly.	N. Holland.	1824.	—	F. 4.	
PLE'EA. M.	PLE'EA.	Enneandria Trigynia. Ph.				
tenuifolia. M.	(br.) slender-leaved.	Carolina.	1812.	5. 8.	F. 4.	Bot. mag. t. 1956.
NARTH'ECIUM. E.B. NARTH'ECIUM.		Hexandria Monogynia.				
1 ossifragum. E.B. (ye.) Lancashire Asphodel.	Britain.	—	7. 8.	H. 4.		Eng. bot. v. 8. t. 535.
2 americanum. B.M. (ye.) American.	N. America.	1811.	—	H. 4.		Bot. mag. t. 1505.
LUZ'ULA. DC.	WOODRUSH.	Hexandria Monogynia. S.S.				
1 pilosa. R.S.	(br.) spring.	Britain.	—	3. 5.	H. 4.	Eng. bot. v. 11. t. 730.
2 flavescens. s.s.	(st.) yellowish.	S. Europe.	1817.	—	H. 4.	Host. gram. 3. t. 94.
3 Forstéri. DC.	(br.) Forster's.	England.	—	5. 6.	H. 4.	Eng. bot. v. 18. t. 1293.
4 glabrata. s.s.	(br.) smooth.	Switzerland.	1825.	—	H. 4.	Desv. j. bot. 1. t. 5. f. 3.
5 albida. DC.	(wh.wh.) white.	—	1815.	—	H. 4.	Leers. herb. t. 13. f. 6.
6 lutea. DC.	(st.) yellow.	—	—	—	H. 4.	
7 spadicea. DC. (da.br.) light red.	Europe.	1816.	—	H. 4.	Vill. delph. 2. t. 6.	
8 parviflora. s.	(br.) small-flowered.	Switzerland.	1824.	—	H. 4.	Rost. junc. p. 26. t. 1. f. 1.
9 nivea. DC.	(wh.) snowy.	—	1770.	—	H. 4.	Scheuch. gram. t. 7. f. 7.
10 sylvatica. s. (br.wh.) wood.	Britain.	—	5.	H. 4.	Eng. bot. v. 11. t. 737.	
<i>Juncus sylvaticus.</i> E.B.						
11 pallescens. s.	(br.) palish.	N. Europe.	1828.	—	H. 4.	Linn. fl. lap. t. 10. f. 2.
12 hyperborea. s.	(br.) northern.	—	—	—	H. 4.	
13 arcuata. s. (wh.) fringe-bracted.	Scotland.	—	7.	H. 4.	Wahl. lapp. t. 4.	
14 spicata. DC.	(br.) spiked.	—	—	—	H. 4.	Eng. bot. v. 17. t. 1176.
15 congesta. s.s.	(br.) close-headed.	Britain.	—	6.	H. 4.	Purt. suppl. t. 9.
<i>Juncus liniger.</i> Purt.						
16 sudética. DC.	(br.) Swedish.	Sweden.	1824.	—	H. 4.	Host. gram. 4. t. 99.
17 pediformis. DC.	(br.) foot-shaped.	S. Europe.	1822.	5. 6.	H. 4.	
18 campéstris. DC.	(br.) field.	Britain.	—	4. 5.	H. 4.	Eng. bot. v. 10. t. 672.
19 multiflora. s.s.	(br.) many-flowered.	Europe.	1822.	5. 6.	H. 4.	
JU'NCUS. DC.	RUSH.	Hexandria Monogynia. L.				
1 multicapitatus. s. (br.) many-headed.	Scotland.	—	6. 7.	H. 4.		
<i>polycéphalus.</i> E.F. non Mich.						
2 fusco-áter. s.	(br.) dark brown.	Europe.	1825.	—	H. 4.	Eng. bot. v. 30. t. 2144.
3 obtusiflorus. L.T. (br.) whorled.	Britain.	—	7. 8.	H. 4.	—	v. 30. t. 2143.
4 lampocarpus. L.T. (br.) shining-fruited.	—	—	—	H. 4.	—	v. 4. t. 238.
5 acutiflorus. L.T. (br.) sharp-flowered.	—	—	6. 7.	H. 4.	—	
<i>articulatus.</i> E.B.						
6 Holoschænus. B.P. (br.) stripe-seeded.	N. S. W.	1824.	—	F. 4.		
7 subverticillatus. L.T. (br.) whorled.	Britain.	—	7. 8.	H. 4.	Flor. dan. t. 817.	
8 tenuis. s.	(gr.pu.) slender.	N. America.	1826.	—	H. 4.	
9 castaneus. E.B.	(br.) black-spiked.	Scotland.	—	7.	H. 4.	Eng. bot. v. 13. t. 900.

10 <i>Jacquini. s.</i> (br. pu.) <i>Jacquin's.</i>	S. Europe.	1824.	7.	H. 4.	Jacq. fl. aust. t. 221.
<i>biglumis.</i> Jacq. vind. t. 4. f. 2. nec aliorum.	Scotland.	8.	H. 4.	Eng. bot. v. 13. t. 898.
11 <i>biglumis.</i> E.B. (br.) two-flowered.	Britain.	7.	H. 4.	_____ v. 13. t. 899.
12 <i>triglumis.</i> E.B. (br.) three-flowered.	—	5. 7.	H. 4.	Weig. obs. t. 2. f. 5.
13 <i>capitatus.</i> s.s. (br.g.) headed.	—	7.	H. 4.	Flor. dan. t. 1099.
14 <i>supinus.</i> L.T. (br.) dwarf.	—	7.	H. 4.	Eng. bot. v. 12. t. 801.
15 <i>uliginosus.</i> E.B. (br.) little bulbous.	—	7. 8.	H. 4.	_____ v. 21. t. 1482.
16 <i>trifolius.</i> E.B. (br.) three-leaved.	Scotland.	—	H. 4.	_____ v. 31. t. 2174.
17 <i>Gesneri.</i> E.F. (g.br.) <i>Gesner's.</i>	—	7.	H. 4.	
<i>gracilis.</i> E.B. non B.P.					
18 <i>bufonius.</i> E.B. (pa.) toad.	Britain.	6. 8.	H. 4.	Eng. bot. v. 12. t. 802.
19 <i>cœnosus.</i> L.T. (br.) mud.	—	7. 8.	H. 4.	Flor. dan. t. 431.
20 <i>compræssus.</i> L.T. (g.br.) round-fruited.	—	—	H. 4.	Eng. bot. v. 13. t. 934.
<i>bulbosus.</i> E.B.					
21 <i>squarrosum.</i> E.B. (br.) Goose-corn.	—	6. 7.	H. 4.	Eng. bot. v. 13. t. 933.
22 <i>Tenagēja.</i> s.s. (g.br.) annual.	Europe.	1816.	—	H. 4.	Flor. dan. t. 1160.
23 <i>aristulatus.</i> M. (br.) awned.	N. America.	1820.	—	H. 4.	
24 <i>plebeius.</i> B.P. (pa.pu.) linear-leaved.	N. S. W.	1816.	—	F. 4.	
25 <i>planifolius.</i> B.P. (br.pu.) flat-leaved.	—	1822.	7. 8.	F. 4.	
26 <i>pauciflorus.</i> B.P. (pa.) few-flowered.	—	—	—	F. 4.	
27 <i>vaginatus.</i> B.P. (st.) sheathed.	—	1823.	—	F. 4.	
28 <i>polycéphalus.</i> M. (gr.) paniced.	N. America.	1828.	—	H. 4.	
29 <i>macrostèmon.</i> s. (gr.) large-stamened.	—	1826.	—	H. 4.	
30 <i>filiformis.</i> E.B. (gr.) least.	Britain.	8.	H. 4.	Eng. bot. v. 17. t. 1175.
31 <i>pallidus.</i> B.P. (pa.) pale-flowered.	N. Holland.	1829.	—	F. 4.	Flor. dan. t. 1095.
32 <i>árticus.</i> w. (da.br.) arctic.	Scotland.	7. 8.	H. 4.	
33 <i>bálticus.</i> s. (br.) Baltic.	Baltic.	1821.	—	H. 4.	
34 <i>effusus.</i> E.B. (gr.) soft.	Britain.	5. 8.	H. 4.	Eng. bot. v. 12. t. 836.
35 <i>conglomeratus.</i> E.B. (br.) common.	—	6. 7.	H. 4.	_____ v. 12. t. 835.
36 <i>glaucus.</i> E.B. (br.) glaucous.	—	7. 8.	H. 4.	_____ v. 10. t. 665.
37 <i>marítimus.</i> E.B. (br.) sea-side.	—	8.	H. 4.	_____ v. 24. t. 1725.
38 <i>acutus.</i> E.B. (g.br.) great sharp sea.	—	7. 8.	H. 4.	_____ v. 23. t. 1614.
39 <i>globulosus.</i> E. (br.) globular.	Egypt.	1829.	—	H. 4.	
? FLAGELL'ARIA. L. FLAGELL'ARIA.	Hexandria Trigynia. L.				
índica. L. (st.) Indian.	India.	1782.	6. 7.	S. h.	Redout. ill. t. 257.
? PHIL'YDRUM. B.P. PHIL'YDRUM.	Monandria Monogynia. B.M.				
lanuginosum. B.P. (ye.) woolly.	N. S. W.	1801.	6. 7.	S. w. ♂.	Bot. mag. t. 783.
? BURMA'NNIA. L. BURMA'NNIA.	Triandria Monogynia.				
1 distächya. B.P. (bl.) two-spiked.	N. S. W.	1822.	6. 8.	G.w. 4.	Burm. zeyl. t. 20. f. 1.
2 biflora. L. (bl.) two-flowered.	Virginia.	9.	G.w. 4.	
? GILLIESIA. B.R. GILLIESIA.	Hexandria Monogynia.				
gramínea. B.R. (st.) Grass-like.	S. America.	1824.	5. 7.	G. 4.	Bot. reg. t. 992.

ORDO CCXXII.

RESTIACEÆ. Brown prodr. p. 243.

X'YRIS. B.P.	X'YRIS. Triandria Monogynia. L.				
1 l'avis. B.P. (ye.) smooth.	N. S. W.	1819.	6. 8.	G. 4.	
2 bracteata. B.P. (ye.) bracteate.	—	1825.	—	G. 4.	
3 júnccea. B.P. (ye.) rushy.	—	1822.	—	G. 4.	
4 gracilis. B.P. (ye.) slender.	V. Diem. Isl.	1821.	—	G. 4.	
5 operculata. B.P. (ye.) rush-leaved.	N. S. W.	1804.	6. 7.	G. 4.	
6 Pürshii. (ye.) Pursh's. <i>índica.</i> Ph. non Vahl.	N. America.	1825.	—	F. 4.	
7 índica. w. (ye.) Indian.	E. Indies.	1822.	5. 7.	S. 4.	Rheed. mal. 9. t. 71.
8 ancéps. R.S. (ye.) flat-edged.	Malabar.	1824.	—	S. 4.	
9 americana. R.S. (bl.) blue-flowered.	Trinidad.	1825.	—	S. 4.	Aubl. gnian. 40. t. 15.
10 caroliniana. R.S. (ye.) Carolina.	N. America.	1812.	—	G. 4.	
11 capén sis. R.S. (ye.) Cape.	C. B. S.	1822.	—	G. 4.	
12 brevifolia. M. (ye.) short-leaved.	Carolina.	1812.	6. 8.	F. 4.	
ERIOCA'ULON. L. PIPEWORT.	Triandria Trigynia. L.				
1 septangulare. E.B. (wh.) jointed.	Scotland.	9.	H.w. 4.	Eng. bot. v. 11. t. 733.
2 australe. B.P. (wh.) New Holland.	N. S. W.	1822.	8. 10.	G.w. 4.	

DEVÁ'UXIA. B.P.	DEVA'UXIA.	Monandria Monogynia.				
1 Patersóni. B.P. (gr.)	Paterson's.	N. S. W.	1825.	6. 7. F. 4.	Linn. trans. v. 10. t. 12. f. 2	
Centrolépis 'æmula.	L.T.					
2 Billardieri. B.P. (gr.)	Billardiere's.			F. 4.	Linn. trans. v. 10. t. 12. f. 1	
Centrolépis cuspidigera.	L.T. fasciculáris.	Lab. nov. holl. 1. p. 7. t. 1.				
HYPOL'ENA. B.P.	HYPOL'ENA.	Dicecia Triandria.				
1 fastigiata. B.P. (br.) crowded.		N. S. W.	1821.	6. 8. F. 4.		
2 exsúca. B.P. (br.) round-stalked.				F. 4.		
LEPTOCA'RPU. B.P.	LEPTOCA'RPU.	Dicecia Triandria.				
1 ténax. B.P. (br.) tough.		N. S. W.	1824.	6. 8. F. 4.	Lab. nov. holl. 2. t. 229.	
Schændum ténax fémina.	Lab.					
2 imbricátus. s.s. (br.) imbricated.	C. B. S.			G. 4.		
EL'EGIA. W.	EL'EGIA.	Dicecia Triandria. W.				
1 júncea. w. (br.) Rush-like.	C. B. S.	1789.	7. 8. F. 4.	Rottb. gram. 10. t. 3. f. 2.		
2 racemosa. w. (br.) racemed.		1804.	5. 6. F. 4.	Lam. ill. t. 884. f. 4.		
LEPYR'ODIA. B.P.	LEPYR'ODIA.	Dicecia Triandria. W.				
gracilis. B.P. (br.) slender.	N. S. W.	1823.	5. 6. F. 4.			
WILLDEN'OVIA. S.S.	WILLDEN'OVIA.	Dicecia Triandria. W.				
1 striata. s.s. (br.) striated.	C. B. S.	1816.	6. 8. F. 4.	Th.act.hol. 1790. t. 2. f. 1.		
2 tères. s.s. (br.) round-stalked.		1790.	F. 4.	1790. t. 2. f. 2.		
THAMNOCHO'R'TUS. B.P.	THAMNOCHO'R'TUS.	Dicecia Triandria.				
1 dichótomas. B.P. (br.) forked.	C. B. S.	1816.	6. 8. F. 4.	Rottb. gram. t. 2. f. 1.		
2 scarioös. s.s. (br.) scariose.		1822.	G. 4.	Berg. cap. t. 5. f. 8.		
RE'STIO. B.P.	ROPE-GRASS.	Dicecia Triandria. L.				
1 australis. B.P. (br.) southern.	V. Diem. Isl.	1812.	4. 6. F. 4.			
2 gracilis. B.P. (br.) slender.	N. S. W.	1823.	6. 8. F. 4.			
3 complanatus. B.P. (br.) flat-stemmed.			F. 4.			
4 tectorum. s.s. (br.) thatch.	C. B. S.	1793.	5. 6. F. 4.	Rottb. gram. 10. t. 3. f. 2.		
5 vaginatus. s.s. (br.) sheathed.		1816.	F. 4.			
6 fastigiatus. B.P. (br.) crowded.	N. S. W.	1824.	F. 4.			
7 virgatus. s.s. (br.) twiggy.	C. B. S.	1816.	F. 4.	Rottb. gram. t. 1. f. 2.		
8 paniculatus. s.s. (br.) paniced.			F. 4.	—	t. 2. f. 3.	
9 lateriflòrus. B.P. (br.) side-flowering.	N. S. W.	1819.	F. 4.	Labil. nov. hol. 2. t. 228.		
Caloròphus elongata.	Lab.					
10 tetraphyllus. B.P. (br.) four-leaved.		1825.	F. 4.	Labil. nov. hol. 2. t. 226.		

ORDO CCXXIII.

CYPERACEÆ. Brown prodr. 1. p. 212.

CAREX. L.	SEDGE.	Monœcia Triandria L.					
1 dioica. w.	(br.) dioecious.	Britain.	5. 6. H. 4.	Eng. bot. v. 8. t. 543.		
2 Davalliana. w.	(br.) Davall's.		H. 4.	—	v. 30. t. 2123.	
3 pauciflòra. E.B.	(st.) few-flowered.		6.	H. 4.	—	v. 29. t. 2041.
4 pyrenáica. s.s.	(br.) Pyrenean.	Pyrenees.	1816.	—	H. 4.	Schk. caric. 5.t.D.f.15.	
5 pulicaris. E.B.	(br.) Flea.	Britain.	6. 7. H. 4.	Eng. bot. v. 15. t. 1051.		
6 Fraseriana. H.K.	(wh.) Fraser's.	N. America.	1809.	4. 6. H. 4.	Bot. mag. t. 1391.		
7 cyperoides. s.s.	(br.) Bohemian.	Bohemia.	1801.	6. 7. H. 4.	Schk. caric. t. A. f. 5.		
8 f'œtida. s.s.	(br.) stinking.	Switzerland.	1791.	7. 8. H. 4.	—	t. Hh. f. 96.	
9 incúrva. E.B.	(br.) curved.	Scotland.	H. 4.	Eng. bot. v. 13. t. 927.		
10 stenophylla. s.s.	(br.) slender-leaved.	Europe.	1824.	—	H. 4.	Schk. caric. t. G. f. 32.	
11 chordorrhiza. s.s.	(br.) creeping-rooted.	Germany.	—	H. 4.	—	t. G. f. 31.	
12 baldensis. s.s.	(br.) three-style.	Mount Baldo.	1829.	5. 7. H. 4.	—	t. Y. f. 81.	
13stellulata. E.B.	(br.) little prickly.	Britain.	5. 6. H. 4.	Eng. bot. v. 12. t. 806.		
14 rosea. s.s.	(ro.) Rose.	N. America.	1812.	—	H. 4.	Schk. caric. Zzz. f. 179.	
15 retroflexa. s.s.	(br.) retroflexed.	Pensylvania.	1826.	—	H. 4.	—	t. Kkk. f. 140.
16 triceps. s.s.	(br.) three-headed.	N. America.	1825.	—	H. 4.	—	t. Zzz. f. 180.
17 f'œnea. s.s.	(br.) short-spiked.		1819.	—	H. 4.		
18 muricata. E.B.	(br.) greater prickly.	Britain.	—	H. 4.	Eng. bot. v. 16. t. 1097.	
19 brizoïdes. s.s.	(br.) Briza-like.	Germany.	1815.	5. 7. H. 4.	Host. gram. 36. t. 47.		
20 argyroglochin. s.s.	(wh.) silver-spiked.		1823.	—	H. 4.	Sch. caric. t. Xxx. f. 174.	
21 straminea. s.s.	(st.) slender-stalked.	N. America.	1803.	6. 7. H. 4.	—	t. Hhf. f. 132.	
22 stipata. s.s.	(br.) sharp-scaled.		1820.	—	H. 4.	—	
23 Muhlenbergii. s.s. (gr.)	Muhlenberg's.		1824.	—	H. 4.	t. Yyy. f. 178.	

CYPERACEÆ.

24 Schrèberi. s.s.	(br.) Schreber's.	Germany.	1800.	6. 7. H. 4.	Host. gram. I. t. 46.
25 ovalis. E.B.	(gr., br.) oval-spiked.	Britain.	— H. 4.	Eng. bot. v. 5. t. 306.
26 cûta. E.B.	(wh.) white.	6. H. 4.	— v. 20. t. 1386.
27 divisa. E.B.	(bk.) bracteated.	5. 7. H. 4.	— v. 16. t. 1096.
28 sparganioides. s.s.(br.)	Sparagnum-like.	N. America.	1822.	— H. 4.	Schk. caric. t. Lll. f. 142.
29 arenaria. E.B.	(br.) sand.	Britain.	6. 7. H. 4.	Eng. bot. v. 13. t. 928.
30 intermédia. E.B.	(br.) soft brown.	5. 7. H. 4.	— v. 29. t. 2042.
31 schœnoïdes. w.	(br.) Scheenus-like.	Austria.	1819.	— H. 4.	Host. gram. I. t. 45.
32 lagopodioides. w.	(br.) Hare's-foot.	N. America.	1805.	6. 7. H. 4.	Schk. cari. t. Yyy. f. 177.
33 elongata. E.B. (br., gr.)	elongated.	England.	5. 6. H. 4.	Eng. bot. v. 27. t. 1920.
34 remota. E.B.	(wh.) remote.	Britain.	— H. 4.	— v. 12. t. 832.
35 axillaris. E.B.	(br.) axillary.	England.	— H. 4.	— v. 14. t. 993.
36 divulsa. E.B.	(wh.) gray.	Britain.	5. H. 4.	— v. 9. t. 629.
37 nemorosa. s.s.	(gr.) shady wood.	Germany.	5. 7. H. 4.	Eng. bot. v. 5. t. 307.
38 vulpina. E.B.	(gr.) great spiky.	Britain.	5. 8. H. 4.	
39 appressa. B.P.	(gr.) close-spiked.	N. S. W.	1802.	— F. 4.	
40 inversa. B.P.	(gr.) inverted.	1820.	— F. 4.	
41 scoparia. w.	(gr.) broom.	N. America.	1812.	6. 7. H. 4.	Schk. cari. t. Xxx. f. 175.
42 multiflora. w.	(gr.) many-flowered.	1819.	— H. 4.	— t. Lll. f. 144.
43 paradoxia. w.	(gr.) paradoxical.	Germany.	—	— H. 4.	— t. E. f. 21.
44 teretiuscula. E.B. (br.)	lesser-panicked.	Britain.	5. 6. H. 4.	Eng. bot. v. 15. t. 1065.
45 paniculata. E.B.	(br.) great paniced.	England.	6. 7. H. 4.	— v. 15. t. 1064.
46 tenella. E.F.	(br.) slender.	Scotland.	6. H. 4.	Schk. caric. t. P. f. 104.
47 norvægica. w.	(gr.) Norway.	Norway.	1823.	— H. 4.	— t. S. f. 66.
48 atrata. E.B.	(bk.) black.	Britain.	6. 7. H. 4.	Eng. bot. v. 29. t. 2044.
49 bicolor. w.	(br., ye.) two-coloured.	Switzerland.	1823.	— H. 4.	Sch. car. t. Aaaa. f. 181.
50 pedunculata. w.	(gr.) peduncled.	N. America.	—	5. 7. H. 4.	— t. Ggg. f. 131.
51 longifolia. B.P.	(gr.) long-leaved.	N. S. W.	—	— F. 4.	
52 thuringiaca. w.	(gr.) Thuringian.	Hungary.	—	— H. 4.	
53 Buxbaumii. w.	(br.) Buxbaum's.	S. Europe.	1819.	5. 6. H. 4.	Schk. cari. t. Ppp. f. 155.
54 glareosa. w.	(br.) narrow-leaved.	Norway.	—	— H. 4.	— t. X. f. 76.
55 clandestina. E.B. (wh.)	dwarf silvery.	Britain.	5. H. 4.	— t. Aaa. f. 97.
56 digitata. E.B.	(pu.) fingered.	England.	5. 6. H. 4.	Eng. bot. v. 30. t. 2124.
57 plantaginea. w.	(br.) broad-leaved.	N. America.	1805.	— H. 4.	— v. 9. t. 615.
58 pilulifera. E.B.	(br.) round-headed.	Britain.	4. 6. H. 4.	Schk. caric. t. U. f. 70.
59 montana. s.s.	(br.) mountain.	Europe.	1819.	5. 6. H. 4.	Eng. bot. v. 13. t. 885.
<i>collina. w.</i>					
60 ciliata. w.	(br.) fringed.	Germany.	—	— H. 4.	Schk. caric. t. I. f. 42.
61 præcox. E.B.	(st.) vernal.	Britain.	4. H. 4.	Eng. bot. v. 16. t. 1099.
62 tomentosa. E.B.	(st.) downy-fruited.	England.	6. H. 4.	— v. 29. t. 2046.
63 extensa. E.B.	(gr.) long-bracted.	Britain.	— H. 4.	— v. 12. t. 833.
64 flava. E.B.	(ye.) yellow.	5. 6. H. 4.	— v. 18. t. 1294.
65 Odéri. E.B.	(ye.) Eder's.	England.	6. 7. H. 4.	— v. 25. t. 1773.
66 alba. w.	(wh.) bristle-leaved.	Germany.	1819.	— H. 4.	Host. gram. I. t. 59.
67 Mielichóferi. E.B.	(gr.) loose-spiked.	Scotland.	—	7. 8. H. 4.	Eng. bot. v. 32. t. 2293.
68 ferruginea. w.	(br.) ferruginous.	Europe.	1820.	6. 8. H. 4.	Schk. caric. t. M. f. 48.
69 saxatilis. w.	(st.) rock.	N. Europe.	—	— H. 4.	— t. I. f. 40.
70 rigidia. E.B.	(bk.) rigid.	Britain.	6. 7. H. 4.	Eng. bot. v. 29. t. 2047.
71 pùlla. E.B.	(bk., br.) russet.	Scotland.	7. H. 4.	— v. 29. t. 2045.
72 cæspitosa. E.B.	(gr.) tufted bog.	Britain.	5. 6. H. 4.	— v. 21. t. 1507.
73 stricta. E.B.	(br.) straight-leaved.	4. 5. H. 4.	— v. 13. t. 914.
74 alpétris. w.	(br.) Alpine.	Europe.	1804.	5. 6. H. 4.	Schk. caric. t. G. f. 35.
75 nitida. w.	(br.) glossy.	Austria.	1805.	— H. 4.	Host. gram. I. t. 71.
76 umbrösa. w.	(gr.) shade.	1818.	6. 7. H. 4.	— t. 69.
77 granularis. w.	(gr.) grain-seeded.	N. America.	1807.	— H. 4.	Sch. caric. t. Vvv. f. 169.
78 anceps. w.	(gr.) two-edged.	1805.	7. 8. H. 4.	— t. Fff. f. 128.
79 demissa. s.s.	(br.) low.	Norway.	1825.	— H. 4.	
80 binervis. E.B.	(bk.) green-ribbed.	Britain.	6. H. 4.	Eng. bot. v. 18. t. 1235.
81 distans. E.B.	(br.) loose.	— H. 4.	— v. 18. t. 1234.
82 fulva. E.B.	(ye., br.) tawny.	6. 7. H. 4.	— v. 18. t. 1295.
83 loliae. w.	(br.) Ray-grass-like.	Sweden.	1810.	— H. 4.	Schk. car. t. Pp. f. 104.
84 lucorum. W.en.	(gr.) grove.	N. America.	1825.	— H. 4.	
85 Micheliæ. w.	(br.) Micheli's.	Europe.	1818.	— H. 4.	Host. gram. I. t. 72.
86 pendula. E.B.	(gr.) great pendulous.	Britain.	5. 6. H. 4.	Eng. bot. v. 33. t. 2315.
87 strigosa. E.B.	(gr.) loose pendulous.	— H. 4.	— v. 14. t. 994.
88 depauperata. E.B. (gr.)	starved-wood.	England.	6. 7. H. 4.	— v. 16. t. 1098.
89 pilosa. s.s.	(gr.) hairy.	Germany.	1819.	— H. 4.	Schk. caric. t. M. f. 49.
90 panicea. E.B.	(bk.) Pink-leaved.	Britain.	5. 7. H. 4.	Eng. bot. v. 21. t. 1505.
91 oligocarpa. w.	(gr.) few-seeded.	N. America.	1825.	— H. 4.	Schk. caric. t. Vvv. f. 178.

92 rostrata. w.	(gr.) beaked.	N. America.	1818.	5. 7. H. 4.	Schk. caric. t. Hhh. f. 134
93 conglobata. w.	(gr.) globular-fruited.	Hungary.	—	— H. 4.	
94 sylvatica. E.B.	(gr.) wood.	Britain.	5. 6. H. 4.	Eng. bot. v. 14. t. 995.
<i>Driméja. w.</i>					
95 Pseudocypérus. w.	(gr.) Bastard Cypress.	—	6. 7. H. 4.	Eng. bot. v. 4. t. 242.
96 júncea. s.s.	(gr.) rush-like.	N. America.	1823.	— H. 4.	
97 pallescens. E.B.	(g.ye.) pale.	Britain.	4. 6. H. 4.	Eng. bot. v. 31. t. 2185.
98 limosa. E.B.	(ye.ye.) green and gold.	—	6. H. 4.	— v. 29. t. 2043.
99 rariflora. E.B.	(bk.) loose-flowered.	Scotland.	— H. 4.	— v. 35. t. 2516.
100 capillaris. E.B.	(br.) capillary.	Britain.	7. 8. H. 4.	— v. 29. t. 2069.
101 lavigata. E.B.	(gr.) smooth-stalked.	—	5. 6. H. 4.	— v. 20. t. 1387.
102 ustulata. E.B.	(bk.) scorched alpine.	Scotland.	6. 7. H. 4.	— v. 34. t. 2404.
103 flexuosa. w.	(br.) bending.	N. America.	1807.	— H. 4.	Schk. car. t. Ddd. f. 124.
104 recúrva. E.B.	(bk.br.) glaucous heath.	England.	5. 6. H. 4.	Eng. bot. v. 21. t. 1506.
β Micheliana. E.B.	(bk.) smooth-fruited.	—	— H. 4.	— v. 32. t. 2236.
105 nútans. w.	(br.) nodding.	Austria.	1815.	6. 7. H. 4.	Host. gram. I. t. 83.
106 acuminata. w.	(br.) taper-pointed.	Istria.	1819.	— H. 4.	— 1. t. 97.
107 hírta. E.B.	(br.) hairy.	Britain.	5. 6. H. 4.	Eng. bot. v. 10. t. 685.
108 filiformis. E.B.	(bk.) slender-leaved.	—	6. 7. H. 4.	— v. 13. t. 904.
109 acuta. E.B.	(bk.) slender-spiked.	—	5. 6. H.w. 4.	— v. 9. t. 580.
110 paludosa. E.B.	(bk.) lesser common.	—	— H.w. 4.	— v. 12. t. 807.
111 riparia. E.B.	(bk.) great common.	—	4. 6. H.w. 4.	— v. 9. t. 579.
112 vesicaria. E.B.	(bk.) short-spiked.	—	5. 6. H. 4.	— v. 11. t. 779.
113 ampullacea. E.B.	(st.) slender-beaked.	—	— H. 4.	— v. 11. t. 780.
114 bullata. w.	(st.) blistered.	N. America.	1811.	6. H. 4.	Schk. caric. Uuu. f. 166.
115 hordeiformis. w.	(gr.) Barley-like.	France.	1805.	6. 7. H. 4.	
116 crinita. w.	(gr.) haired.	N. America.	1807.	— H. 4.	Schk. caric. t. Eee. f. 125.
117 aquatilis. w.	(gr.) water.	Lapland.	1819.	— H.w. 4.	
118 secalina. w.	(br.) rye-like.	Europe.	1820.	— H. 4.	Schk. caric. t. Ddd. f. 121
119 salina. w.	(br.) salt marsh.	Norway.	6. H. 4.	
120 eriophora. F.	(wh.) wool-bearing.	Dahuria.	1828.	— H. 4.	
121 pentastachys. F.	(br.) five-spiked.	Kamtschatka.	1821.	— H. 4.	
122 córsica. Ot.	(br.) Corsican.	Corsica.	1825.	— H. 4.	
KOBR'ESIA. W.	KOBR'ESIA.	Monœcia Triandria. W.			
caricina. w.	(gr.) Carex-like.	Britain.	8. H. 4.	Eng. bot. v. 20. t. 1410.
<i>Schænus monoicus. E.B.</i>	<i>Carex hybrida. s.s.</i>				
SCL'ERIA. W.	SCL'ERIA.	Monœcia Triandria. W.			
1 Flagellum. s.s.	(gr.) climbing.	W. Indies.	1822. S. 4.	Gært. sem. I. p. 13. t. 2.
2 latifolia. s.s.	(gr.) broad-leaved.	—	1824. S. 4.	
3 verticillata. s.s.	(gr.) whorled.	N. America.	1825.	6. 7. H. 4.	
4 hirtella. s.s.	(gr.) hairy.	—	1822.	— H. 4.	
5 auricoma. Ot.	(ye.) golden-locks.	—	1826.	7. 9. H. 4.	
6 ciliata. w.	(gr.) fringed.	—	1823.	— H. 4.	
SCH'ENUS. B.P.	Bog-RUSH.	Triandria Monogynia. L.			
1 nigricans. w.	(bk.) black-headed.	Britain.	7. H.w. 4.	Eng. bot. v. 16. t. 1121.
2 pusillus. R.S.	(br.) small.	W. Indies.	1824.	7. 8. S. 4.	Swtz. fl. ind. oc. ic. t. 6.
3 mucronatus. R.S. (br. pu.) clustered.		S. Europe.	1781.	4. 5. H.w. 4.	Flor. græc. I. t. 43.
4 imberbis. B.P.	(bk.) beardless.	N. S. W.	1818.	— F. 4.	
5 melanostachys. B.R. (br.) black-spiked.		—	1822.	6. 8. F. 4.	
CHÆTO'SPORA. B.P.	CHÆTO'SPORA.	Triandria Monogynia. S.S.			
1 ferruginea. s.s.	(br.) brown.	Switzerland.	1818.	7. 8. H.w. 4.	Host. gram. 4. t. 71.
<i>Schænus ferrugineus. R.S.</i>					
2 turbinata. B.P.	(br.) turbinate.	N. S. W.	1822.	— F. 4.	
3 paludosa. B.P.	(br.) marsh.	—	1824.	— F.w. 4.	
LEPIDOSPE'RMA. B.P.	LEPIDOSPE'RMA.	Triandria Monogynia. S.S.			
1 gladiata. B.P.	(br.) flat-stemmed.	N. Holland.	1819.	7. 8. F.w. 4.	Lab. nov. hol. I. t. 12.
2 concava. B.P.	(br.) concave.	—	1824.	— F. 4.	
3 exaltata. B.P.	(gr.) tall.	—	1816.	— H. 4.	
DUL'CHIUM. P.S.	DUL'CHIUM.	Triandria Monogynia. S.S.			
1 spathaceum. P.S.	(br.) sheathed.	N. America.	1818.	7. 8. H.w. 4.	Pluk. alm. t. 301. f. 1.
2 canadense. Ph.	(gr.) Canadian.	Canada.	1826.	— H. 4.	
RHYNCHO'SPORA. R.S.	RHYNCHO'SPORA.	Triandria Monogynia. S.S.			
1 aurea. R.S.	(ye.) golden.	W. Indies.	1825.	7. 8. S. 4.	Rottb. gram. t. 21. f. 1.
2 laxa. R.S.	(br.) loose.	N. America.	1827.	— H.w. 4.	
3 álba. R.S.	(wh.) white-headed.	Britain.	8. H.w. 4.	Eng. bot. v. 14. t. 985.
<i>Schænus álbis. E.B.</i>					
4 fusca. R.S.	(br.) brown-headed.	—	— H.w. 4.	Eng. bot. v. 22. t. 1575.
5 distans. Ph.	(br.) filiform-leaved.	N. America.	1820.	7. 8. H.w. 4.	
6 comata. R.S.	(br.) leafy.	Brazil.	—	— S.w. 4.	

CYPERACEÆ.

CL'ADIUM. B.P.	CL'ADIUM.	Diandria Monogynia. S.S.					
1 Mariscus. B.P. (br.) prickly. germánicum. R.S. Schænus Mariscus. E.B.	England.	7. 8. H.w. 24.	Eng. bot. v. 14. t. 950.			
2 occidentale. R.S. (br.) West Indian.	W. Indies.	1824.	—	S.w. 24.			
3 glomeratum. B.P. (br.) close-headed.	N. S. W.	1819.	—	F.w. 24.			
4 juncinum. B.P. (br.) rushy.	—	1823.	—	F.w. 24.			
5 schoenoides. B.P. (br.) Bogrush-like. Schænus acutus. Lab.	—	—	—	F.w. 24.	Lab. nov. hol. 1. t. 18.		
FIGMARI'STYLIS. B.P. FIMBRI'STYLIS.	Triandria Monogynia. R.S.						
1 ánnua. R.S. (br.) annual.	S. Europe.	1818.	7. 9. H.w. 24.	Host. gram. 3. t. 63.			
2 puberula. R.S. (br.) pubescent.	N. America.	1820.	7. 8. H. 24.				
3 dichotoma. R.S. (br.) forked.	E. Indies.	1817.	7. 9. S.w. 24.	Rottb. gram. t. 13. f. 1.			
4 autumnalis. R.S. (br.) autumnal.	N. America.	1822.	7. 9. H. 24.				
5 ferruginea. R.S. (br.) rusty.	Jamaica.	1825.	—	S.w. 24.	Sloan. hist. 1. t. 77. f. 1.		
ABILDG'AARDIA. B.P. ABILDG'AARDIA.	Triandria Monogynia. R.S.						
monostachya. B.P. (wh.) one-spiked.	W. Indies.	1819.	6. 8. S. 24.	Sloan. jam. 1. t. 79. f. 2.			
ELEO'CHARIS. B.P. ELEO'CHARIS.	Triandria Monogynia. R.S.						
1 acuta. B.P. (br.) acute-scaled.	N. S. W.	1819.	7. 8. F.w. 24.				
2 geniculata. R.S. (br.) jointed.	America.	—	—	F.w. 24.	Sloan. hist. 1. t. 83. f. 3.		
3 plantaginea. R.S. (br.) Plantain-like.	E. Indies.	1826.	7. 9. S.w. 24.	Rottb. gram. t. 15. f. 2.			
4 palustris. R.S. (br.) marsh.	Britain.	7. H.w. 24.	Eng. bot. v. 2. t. 131.			
Scirpus palustris. E.B.							
5 ovata. R.S. (br.) ovate.	Germany.	1818.	7. 8. H.w. 24.	Host. gram. 3. t. 56.			
6 multicaulis. E.F. (br. pu.) many-stalked. Scirpus multicaulis. E.B.	Britain.	7. H.w. 24.	Eng. bot. v. 17. t. 1187.			
7 acicularis. R.S. (br.) upright needle.	—	—	—	H.w. 24.	Eng. bot. v. 11. t. 749.		
8 obtusa. R.S. (br.) obtuse-scaled.	N. America.	1818.	7. 8. H.w. 24.				
9 glaucescens. R.S. (br.) glaucous.	—	1819.	—	H.w. 24.			
10 tenuis. R.S. (br.) slender.	—	—	—	H.w. 24.			
11 capitata. R.S. (br.) headed.	America.	1824.	—	H.w. 24.	Rottb. gram. t. 15. f. 3.		
12 intermedia. R.S. (br.) intermediate.	Pennsylvania.	1826.	—	H.w. 24.			
ISOL'EPIS. B.P.	Isol'epis.	Triandria Monogynia. R.S.					
1 fluitans. R.S. (br.) floating.	Britain.	7. 8. H.w. 24.	Eng. bot. v. 3. t. 216.			
Scirpus fluitans. E.B.							
2 gracilis. (br.) knotted.	N. Holland.	1820.	—	G.w. 24.			
nodosa. B.P. Scirpus gracilis. Linn. trans. 10. t. 15. f. 2.							
3 setacea. R.S. (br. gr.) bristle-like.	Britain.	—	H.w. 24.	Eng. bot. v. 24. t. 1693.		
4 Vahlii. R.S. (br. gr.) Vahl's.	S. America.	1828.	—	S.w. 24.			
5 Micheliæna. R.S. (st.) Micheli's.	S. Europe.	—	—	H.w. 24.	Host. gram. 3. t. 69.		
6 Holoschenus. R.S. (br.) round-headed.	Britain.	—	H.w. 24.	Eng. bot. v. 23. t. 1612.		
β australis. R.S. (br.) southern.	S. Europe.	—	H.w. 24.	Pluk. phyt. t. 40. f. 5.		
7 romana. w. (br.) Roman.	England.	—	H.w. 24.	Jacq. aust. 5. t. 448.		
8 complanata. R.S. (br.) flat-stemmed.	E. Indies.	1820.	—	S.w. 24.			
9 angularis. R.S. (br.) angular-stemmed.	—	1827.	—	S.w. 24.			
10 curvifolia. R.S. (br.) curve-leaved.	—	—	—	S.w. 24.			
11 pubigera. Schr. (br.) pubescent.	—	—	—	S.w. 24.			
12 Ehrenbergii. Ot. (br.) Ehrenberg's.	Egypt.	1825.	—	H.w. 24.			
SCI'RPU'S. B.P.	CLUB-RUSH.	Triandria Monogynia. L.					
1 cæspitosus. E.B. (br.) scaly-stalked.	Britain.	7. H.w. 24.	Eng. bot. v. 15. t. 1029.			
2 campéstris. R.S. (pu. st.) field.	S. Europe.	1824.	7. 9. H.w. 24.	Hayn. term. bott. t. 25. f. 4.			
3 pauciflorus. E.B. (br.) chocolate-headed.	—	8. H.w. 24.	Eng. bot. v. 16. t. 1122.			
Bæothryon. R.S.							
4 pungens. R.S. (br.) pungent.	Europe.	8. 9. H.w. 24.				
5 mucronatus. R.S. (gr. br.) mucronated.	Britain.	—	H.w. 24.	Scheuch. agr. t. 9. f. 14.		
6 Luzulæ. R.S. (br.) clustered.	E. Indies.	1776.	7. 9. S.w. 24.				
7 lacustris. E.B. (br.) tall.	Britain.	7. H.w. 24.	Eng. bot. v. 10. t. 666.			
8 glaucus. E.B. (br.) glaucous.	England.	7. 8. H.w. 24.	—	v. 33. t. 2321.		
9 carinatus. s.s. (br.) compressed.	Britain.	7. H.w. 24.	—	v. 11. t. 791.		
Scirpus compressus. E.B.							
10 rufus. s.s. (br.) rusty.	—	6. 7. H.w. 24.	Eng. bot. v. 15. t. 1010.			
Scirpus rufus. E.B.							
11 triquetus. E.B. (re. br.) triangular.	England.	8. H.w. 24.	Eng. bot. v. 24. t. 1694.			
12 carinatus. E.B. (br.) blunt-edged.	—	7. 8. H.w. 24.	—	v. 28. t. 1983.		
13 maritimus. E.B. (br.) salt marsh.	Britain.	7. 9. H.w. 24.	—	v. 8. t. 542.		
14 sylvaticus. E.B. (br. gr.) wood.	—	—	—	H.w. 24.	v. 13. t. 919.		
15 radicans. R.S. (br. gr.) rooting.	Germany.	—	H.w. 24.	Schk. in Ust. an. 4. t. 1.		
16 atrovirens. R.S. (da. g.) dark green.	N. America.	1818.	—	H.w. 24.			
17 quinquangularis. R.S. (br.) five-angled.	E. Indies.	1823.	—	S.w. 24.			
18 thyrsiflorus. F. (br.) thyrsse-flowered.	Siberia.	1821.	7. 9. H.w. 24.				

C'YPERUS. L.	C'YPERUS.	Triandria Monogynia. L.
1 Aitòni. R.S. (br.) Aiton's.	S. America.	C. B. S.
2 articulátus. R.S. (st.) jointed.	Jamaica.	1820. ——
3 viscósum. R.S. (gr.br.) clammy.	C. B. S.	1781. 5. 8.
4 tenellus. R.S. (gr.br.) slender.	Levant.	1819. 7. 9.
5 mucronátus. R.S. (wh.) pointed.	Hungary.	1781. 7. 8.
6 pannónicus. R.S. (g.w.) dwarf.	Guinea.	1822. 7. 9.
7 margarítaceus. R.S. (wh.) pearl.	E. Indies.	1816. ——
8 monocéphalus. F.I. (da.br.) one-headed.	1827.	— S.w. 4.
9 pygmæus. R.S. (gr.) dwarf.	Jamaica.	1825. ——
10 filiformis. R.S. (br.) filiform.	—	— S.w. 4.
11 tenuis. R.S. (gr.) slender.	—	— S.w. 4.
12 polystachys. R.S. (br.) many-spiked.	E. Indies.	1819. ——
13 bráneus. R.S. (br.) brown-spiked.	W. Indies.	1826. ——
14 brizæus. R.S. (br.) Briza-like.	—	1824. 7. 8.
15 diándrus. R.S. (br.) two-stamened.	N. America.	1827. ——
16 Purshii. R.S. (br.wh.) Pursh's.	—	1816. ——
17 Háspan. R.S. (br.) digitate-spiked.	Trinidad.	1824. ——
18 alternifòlius. R.S. (br.) alternate-leaved.	Madagascar.	1781. ——
19 élégans. R.S. (re.) elegant.	Jamaica.	1801. 5. 9.
20 compréssus. R.S. (pu.st.) compressed.	S. America.	1819. ——
21 vegetus. R.S. (br.) smooth.	America.	1790. ——
compréssus. Jacq. non w.	—	G.w. 4.
22 formosus. R.S. (ye.) handsome.	N. America.	1827. 7. 9.
23 Luzulæ. R.S. (gr.br.) compact.	W. Indies. — S.w. 4.
polycéphalus. Lam. non L.en.	—	Rottb. gram. t. 13. f. 2.
24 confertus. R.S. (ye.) crowded-spiked.	W. Indies.	1825. 7. 9.
25 squarrosus. R.S. (br.) globe-headed.	E. Indies.	1828. ——
26 paniculatus. R.S. (br.) paniced.	W. Indies.	1804. 5. 7.
27 flavescens. R.S. (br.ye.) yellowish.	Germany.	1776. 6. 9.
28 fuscus. R.S. (br.) brown.	Fulham.	—
29 conglomeratus. R.S. (br.) many-flowered.	Java.	1820. 7. 9.
30 erubescens. L.en. (bh.) blush.	—	— S.w. 4.
31 tuberosus. R.S. (g.pu.) tuberous-rooted.	N. America.	1827. ——
32 rotundus. R.S. (g.pu.) round-rooted.	Africa.	1828. ——
33 gláber. R.S. (br.) smooth.	S. Europe.	1820. ——
34 australis. R.S. (br.) southern.	—	— H. 4.
35 Tenoreánus. R.S. (ye.) Tenore's.	Naples.	1825. ——
36 punctatus. R.S. (br.gr.) spotted-seeded.	E. Indies.	1820. ——
37 Ornithòpus. R.S. (st.pu.) bird's-foot.	St. Domingo.	1824. ——
38 esculentus. R.S. (st.) Rush-nut.	S. Europe.	1597. 7.
39 Hy'dra. M. (gr.wh.) American.	N. America.	1812. H. 4.
40 Pangórei. R.S. (br.) Ceylon.	Ceylon.	1823. 7. 9.
41 bádius. R.S. (br.) saw-leaved.	Algiers.	1800. F.w. 4.
42 pátilus. R.S. (br.st.) spreading.	Hungary.	1818. ——
43 lóngus. E.B. (br.gr.) sweet.	England.	—
44 pròcerus. R.S. (br.wh.) tall rough.	E. Indies.	1826. 7. 9.
45 tenuiflorus. R.S. (br.) slender-flower'd.	S. Europe.	1818. 7. 8.
46 Mónti. R.S. (pu.gr.) spreading-spiked.	—	— H.w. 4.
47 Paramáttá. R.S. (br.pu.) tuberous-rooted.	—	— H.w. 4.
48 spéctabilis. Schreb. (pu.) showy.	—	1827. —— H.w. 4.
49 diándrus. R.S. (st.br.) diandrous.	N. America.	1828. —— H.w. 4.
50 planifòlius. R.S. (br.) flat-leaved.	W. Indies.	1824. —— S.w. 4.
51 Linkiànus. R.S. (br.gr.) Link's.	S. Europe.	1819. 6. 8.
polycéphalus. L. enum. non Lam.	—	H.w. 4.
52 strigósum. R.S. (br.st.) bristle-spiked.	W. Indies.	1786. 7. 9.
53 Michauxiànus. R.S. (g.w.) Michaux's.	N. America.	1812. —— H.w. 4.
54 flavicòmum. R.S. (ye.) yellow-spiked.	—	— H.w. 4.
55 Iria. R.S. (st.gr.) tall.	E. Indies.	1802. 7. — S.w. 4.
56 dístans. R.S. (br.st.) distant.	W. Indies.	1820. 7. 8. S.w. 4.
57 tegétum. R.S. (br.) creeping-rooted.	E. Indies.	1822. —— S.w. 4.
58 giganteus. R.S. (st.) gigantic.	Trinidad.	1824. —— S.w. 4.
59 fastigiatus. R.S. (ye.) lofty.	E. Indies.	1800. 5. 8. S.w. 4.
60 alopecuroides. R.S. (g.b.) fox-tailed.	C. B. S.	1804. —— G.w. 4.
61 cruentus. R.S. (pu.) purple-spotted.	Arabia.	1820. —— S.w. 4.
62 glomeratus. R.S. (br.) round-headed.	Italy.	1804. —— H. 4.
63 ligulatus. R.S. (br.gr.) ligulate.	C. B. S.	1820. —— G.w. 4.
64 diffòrmis. R.S. (br.) deformed.	Levant.	1819. 6. 8. S.w. 4.
65 protractus. L.en. (br.) protracted.	Egypt.	1820. —— H.w. 4.

66 dives. s.s. (ye.) large-bracted.	Egypt. 1820.	7.	H.w.○.	Delil. aëg. 149. t. 4. f. 3.
67 chlorostachys. Stev. (gr.) green-spiked.	Siberia. 1823.	—	H.w.‡.	
68 multíplex. Lk. (br.) many-spiked. 1829.	—	H.w.‡.	
69 scáber. Lag. (br.) rough.	Spain. 1823.	—	H.w.○.	
70 xanthócoma. Ot. (ye.) yellow-haired.	Sicily. 1829.	—	H.w.‡.	
PAPYRUS. S.S. PAPER-RUSH.	Triandria Monogynia. S.S.			
1 antiquòrum. s.s. (br.y.) common.	Egypt. 1802.	7. 9.	S.w.‡.	Michel. gen. pl. t. 19.
Cyperus Pap'rus. R.S.				
2 laxiflòrus. s.s. (br.) loose-flowered.	Madagascar. 1822.	—	S.w.‡.	
3 odorátus. s.s. (st.) fragrant.	W. Indies. 1819.	—	S.w.‡.	Rudg. guian. p. 17. t. 20.
MARI'SCUS. R.S. MARI'SCUS.	Triandria Monogynia. R.S.			
1 capilláris. R.S. (st.br.) capillary.	Trinidad. 1824.	7. 10.	S.w.‡.	
2 aggregátus. R.S. (g.br.) clustered. 1820.	7. 9.	S.w.‡.	
3 l'avis. B.P. (br.g.) smooth.	N. S. W. 1819.	6. 8.	G.w.‡.	
4 ovuláris. R.S. (br.g.) oval-spiked.	Carolina. 1818.	—	F.w.‡.	Pluk. alm. t. 91. f. 4.
5 retrofráctus. R.S. (st.pu.) retrofracted.	N. America. 1827.	—	H.w.‡.	— t. 415. f. 4.
6 umbellátus. R.S. (br.g.) umbelled.	E. Indies. 1789.	—	S.w.‡.	Rottb. gram. t. 4. f. 2.
7 elátus. R.S. (br.g.) tall. 1805.	—	S.w.‡.	Jacq. ic. 2. t. 300.
8 confléxus. L.en. (br.g.) contracted.	Brazil. 1819.	7. 8.	S.w.‡.	
KYLL'INGA. W. KYLL'INGA.	Triandria Monogynia. L.			
1 monocéphala. R.S. (wh.) one-headed.	India. 1793.	6. 7.	S.w.‡.	Rottb. gram. t. 4. f. 4.
2 crucifòrmis. R.S. (wh.) cross-formed.	E. Indies. 1825.	—	S.w.‡.	Swartz. hort. gott. ic.
3 intermédia. B.P. (wh.) intermediate.	N. S. W. 1822.	—	G.w.‡.	
4 triceps. R.S. (wh.pu.) three-headed.	E. Indies. 1776.	9. 11.	S.w.‡.	Rottb. gram. t. 4. f. 6.
5 uncináta. L.en. (gr.) hook-seeded.	Brazil. 1820.	7. 8.	S.w.‡.	
6 odoráta. R.S. (gr.) fragrant.	S. America. 1822.	—	S.w.‡.	
7 polycéphala. L.en. (wh.) many-headed.	Brazil. 1820.	—	S.w.‡.	
FUIR'ENA. B.P. FUIR'ENA.	Triandria Monogynia. L.			
1 umbellátta. R.S. (br.) umbelled.	W. Indies. 1822.	6. 8.	S.w.‡.	Rottb. gram. t. 19. f. 3.
2 glomeráta. R.S. (pu.g.) crowded.	E. Indies. 1827.	—	S.w.○.	— t. 17. f. 1.
ELYNA. R.S. EL'YNA.	Triandria Monogynia. R.S.			
spicáta. R.S. (br.) simple-spiked.	Europe. 1819.	6. 8.	H.w.‡.	Vill. delph. 2. t. 6.
Cárex Bellárdi. Host. gram. 4. t. 77.				
TRICHO'PHORUM. P.S. TRICO'PHORUM.	Triandria Monogynia. P.S.			
1 cyperinum. P.S. (br.st.) cyperine.	N. America. 1802.	5. 9.	H.w.‡.	Pluk. mant. t. 419. f. 3.
2 alpinum. P.S. (br.st.) Alpine.	Scotland.	7.	H.w.‡.	Eng. bot. v. 5. t. 311.
Eriophorum alpinum. E.B.				
ERIO'PHORUM. L. COTTON-GRASS.	Triandria Monogynia. L.			
1 vaginátum. E.B. (br.) Hare's-tail.	Britain.	3. 4.	H.w.‡.	Eng. bot. v. 13. t. 873.
2 capitátum. E.B. (br.) round-headed.	Scotland.	8. 9.	H.w.‡.	— v. 34. t. 2387.
3 grálice. E.B. (bk.g.) slender.	7. 8.	H.w.‡.	— v. 34. t. 2402.
4 angustifólium. (pu.g.) narrow-leaved.	Britain.	4. 5.	H.w.‡.	— v. 8. t. 564.
5 virginicum. R.S. (pu.g.) Virginian.	N. America. 1802.	5. 8.	H.w.‡.	Pluk. alm. t. 299. f. 4.
6 polystachion. E.B. (pu.g.) broad-leaved.	Britain.	4.	H.w.‡.	Eng. bot. v. 8. t. 563.
7 pubescens. E.F. (bk.) pubescent.	4. 5.	H.w.‡.	

ORDO CCXXIV.

GRAMINEÆ. Juss. gen. p. 28.

REMIR'EA. R.S.	REMIR'EA.	Triandria Monogynia.		
marítima. R.S.	sea-side.	America. 1822.	7. 8.	G.w.‡.
NA'R'DUS. R.S.	MAT-GRASS.	Triandria Monogynia. L.		Aubl. guian. 1. t. 16.
strícta. E.B.	upright.	Britain.	6. 7.	H.w.‡.
ORYZO'PSIS. M.	ORYZO'PSIS.	Triandria Monogynia. R.S.		Eng. bot. v. 5. t. 290.
asperifòlia. M.	rough-leaved.	N. America. 1822.	7. 8.	H.w.‡.
LYG'EUM. R.S.	LYG'EUM.	Triandria Monogynia. L.		Mich. amer. 1. t. 9.
Spártum. R.S.	rush-leaved.	Spain. 1776.	5. 6.	H.w.‡.
CORNU'COPIÆ. R.S.	HORN OF PLENTY.	Triandria Monogynia. L.		Schreb. gram. t. 52.53.
cucullátum. R.S.	hooded.	Levant. 1788.	8.	H.○.
CE'NCHRUS. R.S.	CE'NCHRUS.	Triandria Monogynia. L.		Flor. græc. 1. t. 51.
1 tribulóides. R.S.	spinous.	N. America. 1818.	5. 8.	H.○.
2 echinátus. R.S.	rough-spiked.	W. Indies. 1691.	8. 12.	S.♂.
3 australis. B.P.	New Holland.	N. S. W. 1822.	6. 8.	H.○.
				Sloan. hist. 1. t. 65. f. 1.
				Cavan. ic. 5. p. 39. t. 462.

PENNIS'ETUM. R.S.	PENNIS'ETUM.	Triandria Monogynia. R.S.				
1 setōsum. R.S.	setose.	W. Indies.	1804.	7. 8. H.○.		
2 polystāchym. R.S.	many-spiked.	E. Indies.	—	— H.○.	Rumph. amb. 6.t.7. f.2.	
3 cenchroïdes. R.S.	ciliated.	C. B. S.	1777.	7. 8. H.○.	Giseck. ic. 1. t. 23.	
<i>Cénchrus ciliáris. L.</i>						
4 orientálē. R.S.	oriental.	Levant.	1819.	— F. 2.		
5 violáceum. R.S.	violet-spiked.	Senegal.	1829.	— S.○.		
6 compréssum. B.P.	flat-stemmed.	N. S. W.	1820.	— H. 2.		
7 holcoïdes. R.S.	Holcus-like.	E. Indies.	1823.	— S.○.		
<i>Pánicum holcoïdes. F.I.</i>						
8 barbátum. R.S.	bearded.	—	—	— S.○.		
9 dichótomum. R.S.	forked.	Egypt.	1824.	— H. 2.	Delil. ægyp. t. 8. f. 3.	
10 nepalénse. s.s.	Nepaul.	Nepaul.	1822.	— H.○.		
11 ásperum. R.S.	rough-leaved.	C. B. S.	1823.	— G. 2.		
PENICILL'ARIA. R.S.	PENICILL'ARIA.	Polygamia Monœcia. W.				
1 spicáta. R.S.	spiked.	E. Indies.	1592.	6. 7. H.○.	Pluk. alm. t. 32. f. 4.	
2 alopecuroïdes.	Fox-tail.	Jamaica.	1739.	7. 10. H.○.	— t. 92. f. 5.	
<i>ciliáta. W.en. cylindrica. R.S.</i>	<i>Pánicum alopecuroïdes. L.</i>	<i>Alopecurus indicus. L.</i>				
SET'ARIA. R.S.	SET'ARIA.	Triandria Digynia. R.S.				
1 sericea. R.S.	silky.	W. Indies.	1780.	5. 9. H.○.		
2 verticilláta. R.S.	rough.	England.	—	7. 8. H.○.	Eng. bot. v. 13. t. 874.	
3 viridis. R.S.	green.	—	—	— H.○.	— v. 13. t. 875.	
<i>Pánicum viride. E.B.</i>	<i>Pennisétum viride. B.P.</i>					
4 pùmila. R.S.	dwarf.	—	1819.	— H.○.		
5 macrochæta. R.S.	long-spiked.	—	—	— H.○.		
6 purpuráscens. R.S.	purplish.	S. America.	1822.	— H.○.		
7 intermédia. R.S.	intermediate.	E. Indies.	1819.	8. 10. H.○.		
8 Weinmánni. R.S.	Weinman's.	Teneriffe.	1816.	— H.○.		
9 glaúca. R.S.	glaucous.	S. Europe.	1771.	7. 8. H.○.	Host. gram. 2. t. 16.	
10 helvòla. R.S.	pale red.	E. Indies.	1822.	8. 10. H.○.		
11 corrugáta. R.S.	rugged-flowered.	Carolina.	1828.	— H.○.		
12 erubésens. R.S.	blush.	S. America.	1822.	— H.○.		
13 geniculáta. R.S.	jointed.	Antilles.	1805.	7. 8. H.○.		
14 marítima. R.S.	sea-side.	—	1818.	— H.○.		
15 germánica. R.S.	German.	Germany.	1548.	— H.○.	Host. gram. 2. t. 15.	
16 itálica. R.S.	Italian.	Italy.	1616.	— H.○.	Rumph. amb. 3.t.76.f.2.	
<i>Pánicum itálicum.</i>	Host. gram. 4. t. 14.	<i>Pennisétum itálicum. B.P.</i>				
17 compósita. R.S.	compound.	S. America.	1823.	— H.○.		
18 cylindrica. R.S.	cylindrical.	—	—	— H.○.		
19 setósa. R.S.	bristly.	W. Indies.	1822.	— H.○.		
20 polystachia. R.S.	many-spiked.	Brazil.	1824.	— S. 2.		
21 tenacíssima. R.S.	tough.	—	1819.	— H.○.		
22 Pennisétum. R.S.	Indian.	E. Indies.	1820.	6. 8. S.○.		
23 scándens. R.S.	climbing.	—	—	— H.○.		
24 caudáta. R.S.	tailed.	Trinidad.	1824.	— H.○.		
25 muricáta. R.S.	sea-side.	N. America.	1819.	— H.○.		
26 aurícoma. Lk.	golden-headed.	—	1826.	— H.○.		
ANTHOXA'NTHUM. R.S.	SPRING-GRASS.	Diandria Digynia. L.				
1 odorárum. E.B.	sweet-scented.	Britain.	—	5. H. 2.	Eng. bot. v. 9. t. 647.	
β láxum.	loose-spiked.	—	—	— H. 2.	Loes. ic. 22.	
γ pubéscens.	downy-spiked.	—	—	— H. 2.		
δ ramðsum.	branching-spiked.	—	—	— H. 2.		
2 amárum. R.S.	bitter.	S. Europe.	1810.	7. H. 2.	Trin. ic. gram. t. 14.	
3 grácile. R.S.	slender.	Italy.	1825.	7. 8. H.○.	— t. 13.	
4 ovátum. R.S.	oval-spiked.	Spain.	1824.	— H.○.		
SPART'INA. R.S.	SPART'INA.	Triandria Monogynia. R.S.				
1 cynosuroïdes. R.S.	Dog's-tail.	N. America.	1781.	8. 9. H. 2.	Linn. fil. fasc. 1. t. 9.	
<i>Dáctylis cynosuroïdes. L.</i>	<i>Trachynòtia cynosuroïdes. M.</i>	<i>Limnetis cynosuroïdes. P.S.</i>				
2 stricta. R.S.	upright.	Britain.	—	8. H. 2.	Eng. bot. v. 6. t. 380.	
<i>Dáctylis stricta. E.B.</i>	<i>Limnetis pungens. P.S.</i>					
3 glàbra. R.S.	smooth.	N. America.	1827.	— H. 2.		
4 alterniflòra. R.S.	alternate-flower'd.	S. Europe.	—	— H. 2.		
5 polystachia. R.S.	many-spiked.	N. America.	1781.	8. 9. H. 2.		
6 pùmila. R.S.	dwarf.	—	1820.	— H. 2.		
7 pectináta.	pectinated.	—	1827.	— H. 2.		
8 júncea. R.S.	rushy.	—	1781.	7. 8. H. 2.		
9 pátens. R.S.	spreading.	—	—	— H. 2.		
10 geniculáta. R.S.	jointed.	Java.	1822.	— S. 2.	Burm. ind. t. 12. f. 3.	
11 arundinácea. R.S.	Reed-like.	Tristan d'Acunha.	1820.	— G. 2.		

ZO'YSIA. R.S.	ZO'YSIA.	Triandria Digynia. W.			
púngens. R.S.	pungent.	E. Indies.	1819.	6. 8. S. 4.
LE'ERSIA. B.P.	LE'ERSIA.	Triandria Digynia. W.			
1 oryzoides. s.s.	Rice-like.	S. Europe.	1793.	7. 8. H.w. 4.
2 virginica. s.s.	Virginian.	N. America.	—	— H.w. 4.
3 austrális. B.P.	southern.	N. S. W.	1819.	— H.w. 4.
4 lenticularis. s.s.	round-glumed.	N. America.	1825.	— H.w. 4.
5 brasiliensis. R.S.	Brazilian.	Brazil.	1826.	— S.w. 4.
PER'OTIS. R.S.	PER'OTIS.	Triandria Digynia. W.			
latifolia. R.S.	broad-leaved.	E. Indies.	1777.	8. 9. S. 4.
ALOPECURUS. R.S.	FOXTAIL-GRASS.	Triandria Digynia. L.			Rheed. mal. 12. t. 62.
1 agréstis. E.B.	slender.	Britain.	7. 8. H. 4.
2 utriculatus. R.S.	bladdered.	Italy.	1777.	— H. 4.
3 echinatus. R.S.	rough.	C. B. S.	1822.	— G. 4.
4 pratensis. E.B.	meadow.	Britain.	5. H. 4.
5 nigricans. R.S.	blackish.	Europe.	1815.	6. 7. H. 4.
Tauntoniensis. Sinc. gr. wob. p. 232.					
6 trivialis. s.o.	long-awned.	Bohemia.	1827.	— H. 4.
7 scaber. s.o.	scabrous.	—	—	— H. 4.
8 Giesekeanus. R.S. ovatus. F.D. non Forst.	Gieseke's.	N. Europe.	1828.	— H. 4.	Flor. dan. t. 1565.
9 geniculatus. E.B.	floating.	Britain.	5. 8. H.w. 4.
10 fulvus. E.B.	orange-spiked.	England.	6. H.w. 4.
11 alpinus. E.B.	Alpine.	Scotland.	5. 6. H. 4.
12 ramosus. R.S.	branching.	N. America.	1823.	— H. 4.
13 arundinaceus. R.S.	Reed-like.	—	1819.	— H. 4.
14 macrostachyos. s.s.	large-spiked.	Barbary.	1824.	6. 8. H. 4.
15 bulbosus. E.B.	bulbous.	England.	7. H. 4.
16 colobochinoës. R.S.	Siberian.	Siberia.	1826.	6. 7. H. 4.
17 Gerardi. Trin.	Gerard's.	S. Europe.	1820.	6. 8. H. 4.
Phleum Gerardi. R.S.					Jacq. ic. t. 301.
PHLEUM. R.S.	CAT'S-TAIL-GRASS.	Triandria Digynia. L.			
1 pratense. R.S.	common.	Britain.	7. H. 4.
a majus. S.G.W.	greater.	—	—	— H. 4.
β minus. S.G.W.	lesser.	—	—	7. 8. H. 4.
2 nodosum. w.	knotted.	—	—	7. 9. H. 4.
3 Bertoloniï. R.S.	Bertoloni's.	France.	1818.	— H. 4.
4 alpinum. R.S.	alpine.	Scotland.	7. H. 4.
5 commutatum. R.S.	confused.	Piedmont.	1819.	— H. 4.
6 felinum. R.S.	Grecian.	Greece.	1822.	— H. 4.
7 echinatum. R.S.	hedgehog.	Dalmatia.	—	8. 9. H. 4.
ACHNODO'NTON. R.S.	ACHNODO'NTON.	Triandria Digynia. R.S.			Host. gram. 3. t. 11.
1 Bellárdi. R.S.	bulbous.	Spain.	1798.	6. 7. H. 4.
Phálaris Bellárdii. w. bulbosa. L.					
2 tenué. R.S.	slender.	S. Europe.	1826.	— H. 4.
3 arenaria. L.en.	sand.	Britain.	6. 7. H. 4.
Phálaris arenaria. E.B.	Phléum arenarium. F.D. t. 915.	Chilochlòa arenaria. R.S.			Host. gram. 4. t. 222.
CHILOCHL'OA. R.S.	CHILOCHL'OA.	Triandria Digynia. R.S.			
1 Boehméri. R.S.	Phalaris-like.	England.	7. 9. H. 4.
Phálaris phleoides. E.B.	Phléum phalaroides. Kœl.	Baehméri. w.			Eng. bot. v. 7. t. 459.
2 cuspidata. R.S.	sharp-pointed.	Tyrol.	1824.	7. 8. H. 4.
3 Micheli. L.en.	Micheli's.	Scotland.	6. 7. H. 4.
Phléum Micheli. E.B.					Eng. bot. v. 82. t. 2265.
4 áspéra. R.S.	rough.	England.	7. H. 4.
Phléum paniculatum. E.B.	Phléum áspерум. Jacq. ic. t. 14.	Phálaris áspéra. w.			Eng. bot. v. 15. t. 1077.
5 ánnua. R.S.	annual.	Caucasus.	1825.	7. 8. H. 4.
SCHMI'DTIA. S.S.	SCHMI'DTIA.	Diandria Digynia. S.S.			
súttilis. s.s.	close umbellled.	Bohemia.	1824.	6. 8. H. 4.
Coleánthus súttilis. R.S.					Tratt. fl. aus. t. 451.
POLYP'OGON. R.S.	POLYP'OGON.	Triandria Digynia. W.			
1 monspeliensis. R.S.	chaffy.	Britain.	7. 8. H. 4.
Agrostis panicea. E.B.					Eng. bot. v. 24. t. 1704.
2 littoralis. E.F.	sea-side.	—	—	Eng. bot. v. 18. t. 1251.
3 Lagáscae. R.S.	Lagasca's.	Spain.	1824.	— H. 4.
4 marítimus. R.S.	shore.	S. Europe.	1822.	— H. 4.
5 glomeratus. R.S.	clustered.	N. America.	—	7. 8. H. 4.
6 tatáricus. F.	Tartarian.	Tartary.	1825.	— H. 4.
CER'ESIA. P.S.	CER'ESIA.	Triandria Digynia. P.S.			
élégans. P.S.	elegant.	Peru.	1818.	8. 9. H. 4.
Páspalum membranàceum. s.s.					Jacq. frag. t. 86.

PA'SPALUM. W.	PA'SPALUM.	Triandria Digynia. L.		
1 disséctum. s.s.	dissected.	S. America. 1822.	8. 10. H. ⊖.	
2 scrobiculatum.s.s.	dimpled.	E. Indies. 1778.	7. 9. S. Ȑ.	Hout. n. his. t. 89. f. 3.
3 stoloniferum. s.s.	purple.	Peru. 1794.	— G. Ȑ.	Jacq. ic. 2. t. 302.
4 paniculatum. s.s.	panicked.	W. Indies. 1782.	— S. ⊖.	Sloan. hist. 1. t. 72. f. 2.
5 virgatum. s.s.	twiggy.	— 1822.	— S. Ȑ.	— 1. t. 69. f. 2.
6 l'æve. M.	smooth.	N. America. 1818.	— H. Ȑ.	Lecomt. pasp. t. 91.
7 mucronatum.R.S.	mucronate.	— 1825.	— H. ⊖.	Lecomt. pas. t. 91.
8 Lecomteanum. R.S.	Lecomte's.	— 1827.	— H. Ȑ.	
9 latifolium. R.S.	broad-leaved.	— —	— H. Ȑ.	
10 platense. R.S.	Monte Video.	Monte Video. 1824.	— H. Ȑ.	
11 difforme. R.S.	deformed.	N. America. 1818.	7. 9. H. Ȑ.	Lecomt. pas.t.91.p.284.
12 pusillum. R.S.	small.	St. Thomas. 1822.	— S. Ȑ.	
13 conjugatum. s.s.	conjugate.	W. Indies. —	— S. Ȑ.	Gært.fruct.2. t. 80. f.4.
14 notatum. s.s.	two-spiked.	— 1776.	7. S. ȝ.	Swartz obs. t. 2. f. 1.
15 vaginatum. R.S.	sheathed.	— 1824.	— S. Ȑ.	
16 platycaulis. R.S.	flat-stalked.	— —	— S. Ȑ.	
17 molle. s.s.	soft.	St. Thomas. 1822.	— S. Ȑ.	
18 Kora. R.S.	smooth-flowered.	E. Indies. 1818.	7. 8. S. Ȑ.	
19 longiflorum. R.S.	long-flowered.	— 1824.	— S. Ȑ.	
20 setaceum. M.	bristly.	N. America. —	— H. Ȑ.	
21 pilosum. R.S.	hairy.	S. America. 1826.	— S. Ȑ.	
22 incertum. R.S.	uncertain.	— 1812.	— H. Ȑ.	
23 dubium. s.s.	doubtful.	— 1818.	— H. Ȑ.	
24 glabrum. s.s.	smooth.	W. Indies. 1823.	— S. Ȑ.	
25 tenellum. w.	elegant.	— 1817.	— S. Ȑ.	
elegans. s.s.				
26 laxum. R.S.	loose.	S. America. 1819.	— S. Ȑ.	
27 inaequale. L.en.	unequal-valved.	Manilla. 1822.	— S. Ȑ.	
AMPHIP'OGON. B.P. AMPHIP'OGON.		Triandria Digynia. S.S.		
strictus. B.P.	upright.	N. S. W. 1823.	5. 6. G. Ȑ.	
GASTRI'DIUM. R.S.	GASTRI'DIUM.	Triandria Digynia. R.S.		
1 australis. R.S.	yellow.	England. —	8. H. Ȑ.	Eng. bot. v. 16. t. 1107.
<i>Milium lendigerum. E.B.</i>				
2 muticum. R.S.	beardless.	Sicily. 1819.	7. 8. H. ⊖.	
CHÆT'URUS. R.S.	CHÆT'URUS.	Triandria Digynia. R.S.		
fasciculatus. R.S.	fascicled.	S. Europe. 1815.	7. 9. H. ⊖.	
MILIUM. R.S.	MILLET-GRASS.	Triandria Digynia. L.		
1 nigricans. R.S.	Guinea maize.	Peru. 1824.	6. 8. H. ⊖.	
2 effusum. E.B.	common.	Britain. —	6. 7. H. Ȑ.	Eng. bot. v. 16. t. 1106.
3 vernale. R.S.	vernal.	Caucasus. 1819.	4. 5. H. ⊖.	
4 scabrum. R.S.	rough.	S. Europe. 1826.	— H. ⊖.	
5 confertum. R.S.	close-flowered.	Italy. —	— H. Ȑ.	
6 amphicarpon. R.S.	earth-fruited.	N. America. 1827.	— H. Ȑ.	
7 gallécium. R.S.	capillary.	Gallicia. 1819.	6. 8. H. Ȑ.	Pursh. am. sept. 2. t. 2.
8 velutinum. R.S.	downy.	Mexico. 1823.	— H. ⊖.	
9 filiforme. Lag.	slender.	New Spain. 1824.	— H. Ȑ.	
10 microstérnum. R.S.	small-seeded.	— 1822.	— H. ⊖.	
11 phleoides.	Phleum-like.	E. Indies. 1820.	— S. Ȑ.	
PIPTATH'ERUM. R.S. PIPTATH'ERUM.		Triandria Digynia. R.S.		
1 paradóxum. R.S.	black-seeded.	France. 1771.	6. 7. H. ⊖.	Host. gram. 3. t. 23.
<i>Agróstis paradóxa. L.</i> <i>Milium paradóxum. w.</i> <i>Uráchna viréscens. Trin.</i>				
2 cœrulescens. R.S.	blueish.	Barbary. 1819.	— H. Ȑ.	Desf. atl. 1. t. 12.
3 multiflorum. R.S.	many-flowered.	S. Europe. 1778.	— H. Ȑ.	Host. gram. 3. t. 45.
<i>Milium arundinaceum. F.G. t. 66.</i> <i>Agróstis miliacea. L.</i>				
4 punctatum. R.S.	dotted.	W. Indies. 1824.	— S. Ȑ.	
CALAMAGRO'STIS. S.S. CALAMAGRO'STIS.		Triandria Digynia. S.S.		
1 epigéjos. s.s.	wood.	Britain. —	7. H. Ȑ.	Eng. bot. v. 6. t. 403.
2 intermédia. R.S.	intermediate.	Carlsruhe. 1827.	— H. Ȑ.	
3 littorea. s.s.	sea-side.	Switzerland. 1824.	7. 8. H. Ȑ.	Schrad.fl. ger.1.t.4.f.2.
4 láxa. Host.	loose.	— —	— H. Ȑ.	Host. gram. 4. t. 23.
5 glauca. R.S.	glaucous.	N. America. —	— H. Ȑ.	
6 Halleriána. s.s.	Haller's.	Switzerland. 1819.	— H. Ȑ.	Schrad.fl. ger.1.t.4.f.3.
<i>Arundo pseudophragmites. R.S.</i>				
7 stricta. s.s.	upright.	Scotland. —	7. 8. H. Ȑ.	Eng. bot. v. 30. t. 2160.
8 varia. Host.	various.	S. Europe. 1813.	— H. Ȑ.	Host. gram. 4. t. 47.
9 acutiflora. DC.	sharp-flowered.	Germany. 1819.	— H. Ȑ.	
10 sylvatica. DC.	wood.	Europe. 1813.	— H. Ȑ.	Host. gram. 4. t. 49.
<i>arundinacea. Roth.</i> <i>Agróstis arundinacea. L.</i> <i>pyramidalis. Host.</i>				

GRAMINEÆ.

11 purpurascens. Br.	purpleish.	N. America.	1827.	7. 8. H. 4.
12 Schwäbii. s.s.	Saxon.	Saxony.	1828.	— H. 4.
13 Langsdórfii. s.s.	Langsdorff's.	1823.	— H. 4.
14 confinis. s.s.	hairy-flowered.	N. America.	1820.	— H. 4.
15 montana. Host.	mountain.	Europe.	—	— H. 4.
16 Höstii. R.S.	Host's.	—	—	— H. 4.
<i>sylvatica</i> . Host. nec <i>alitorum</i> .					Host. gram. 4. t. 46.
17 canadensis. N.	Canadian.	N. America.	1826.	— H. 4.
18 lanceolata. s.s.	common.	England.	7. H. 4.
<i>Arundo Calamagróstis</i> . E.B.	<i>Agróstis</i>	<i>lanceolata</i> . R.S.			Eng. bot. v. 30. t. 2159.
19 speciosa. Host.	beautiful.	Germany.	1813.	7. 8. H. 4.
<i>Agróstis Calamagróstis</i> . R.S.					Host. gram. 4. t. 45.
STIPA. R.S.		FEATHER-GRASS.	Triandria Digynia. L.		
1 pennata. E.B.	common.	Britain.	7. 8. H. 4.
2 tortilis. R.S.	twisted-awned.	S. Europe.	1823.	— H. ⊙.
3 júnea. R.S.	rushy.	France.	1772.	— H. 4.
4 gigantea. R.S.	gigantic.	Spain.	1823.	— H. 4.
5 Lagáscæ. R.S.	Lagasca's.	—	1822.	— H. 4.
6 capillata.	capillary.	S. Europe.	—	— H. 4.
7 tenacissima. R.S.	tough.	—	1822.	— H. 4.
8 Aristella. R.S.	awned.	Montpelier.	1806.	— H. 4.
9 Redóskii. L.A.	Redowski's.	Altay.	1828.	— H. 4.
10 mollis. B.P.	soft.	N. S. W.	1824.	— F. 4.
11 conferta. R.S.	close paniced.	—	1819.	— H. 4.
12 sibirica. R.S.	Siberian.	Siberia.	1777.	— H. 4.
AGRA'ULUS. P.B.	AGRA'ULUS.	Triandria Digynia. R.S.			
1 canina. P.B.	brown.	Britain.	7. H. 4.
<i>Agróstis canina</i> . E.B.	<i>Trichódium caninum</i> . R.S.				Eng. bot. v. 26. t. 1856.
2 alpina. P.B.	Alpine.	S. Europe.	1818.	7. 8. H. 4.
3 rupéstris. R.S.	rock.	Pyrenees.	—	— H. 4.
4 neglecta. R.S.	neglected.	Hungary.	1822.	— H. 4.
5 flavescens.	yellowish.	S. Europe.	—	— H. 4.
6 setacea.	bristly.	Britain.	—	— H. 4.
<i>Agróstis setacea</i> . E.B.	<i>Trichódium setáceum</i> . R.S.				
7 rúbra. R.S.	red.	Lapland.	1820.	— H. 4.
8 hybrida. R.S.	hybrid.	Switzerland.	—	— H. 4.
9 diffusa. L.en.	spreading.	—	1823.	— H. 4.
TRICH'ODIUM. M.		WINTER-GREEN-GRASS.	Triandria Digynia.		
1 decumbens. R.S.	decumbent.	N. America.	1786.	6. 7. H. 4.
2 laxiflorum. R.S.	loose-flowered.	—	1816.	— H. 3.
3 elegans. R.S.	elegant.	France.	—	7. 9. H. ⊙.
AGRO'STIS. R.S.		BENT-GRASS.	Triandria Digynia. R.S.		
1 stolonifera. R.S.	creeping.	Britain.	7. 8. H. 4.
α latifolia. S.G.W.	Fiorin grass.	—	—	—	— H. 4.
β angustifolia. S.G.W.	narrow-leaved.	—	—	—	— H. 4.
γ aristata. S.G.W.	awned.	—	—	—	— H. 4.
δ nemoralis. S.G.W.	wood.	—	—	—	— H. 4.
ϵ palustris. S.G.W.	marsh.	—	—	—	— H. 4.
2 dulcis. Df.	sweet.	S. France.	1824.	— H. 4.
3 stolonizans. R.S.	stoloniferous.	Poland.	1827.	— H. 4.
4 diffusa. R.S.	spreading.	Europe.	1816.	— H. 4.
5 neglecta. R.S.	neglected.	Hungary.	—	— H. 4.
6 capillaris. R.S.	capillary.	S. Europe.	—	— H. ⊙.
7 tenella. R.S.	slender.	Switzerland.	1824.	— H. 4.
8 álba. R.S.	white rooted.	Britain.	7. H. 4.
β purpurascens.	purplish.	—	—	— H. 4.
γ vivipara.	viviparous.	—	—	— H. 4.
9 párvula. R.S.	small.	Hungary.	1826.	— H. 4.
10 tenuifolia. R.S.	slender-leaved.	Caucasus.	1819.	7. 8. H. 4.
11 pátula. R.S.	knotted spreading.	Switzerland.	1824.	— H. 4.
12 valentina. R.S.	Valentia.	Spain.	—	— H. 4.
13 dóbria. R.S.	doubtful.	Germany.	1818.	— H. 4.
14 gigantea. R.S.	gigantic.	Caucasus.	—	— H. 4.
15 frondosa. R.S.	frondose.	Naples.	1824.	— H. 4.
16 vulgáris. R.S.	fine.	Britain.	—	— H. 4.
17 ambigua. R.S.	ambiguous.	Germany.	1820.	— H. 4.
18 Júressi. R.S.	close paniced.	—	—	— H. 4.
19 virgínica. R.S.	Virginian.	N. America.	—	— H. 4.
20 brasiliensis. R.S.	Brazilian.	Brazil.	1828.	— S. 4.

Eng. bot. v. 19. t. 1356.
Desf. atl. 1. t. 31. f. 1.
Flor græc. 1. t. 85.Desf. atl. 1. t. 28.
Host. gram. 3. t. 5.
Desf. atl. 1. t. 30.
Host. gram. 4. t. 34.
Ledeb. ic. t. 98.

Ledeb. ic. t. 99.

Eng. bot. v. 26. t. 1856.
Host. gram. 3. t. 49.Host. gram. 4. t. 52.
Eng. bot. v. 17. t. 1188.Fraser mon. c. ic.
Mich. amer. 1. t. 8.
Lois journ. bot. 2.t.8.f.1.Eng. bot. v. 22. t. 1532.
Sinc. gr. wob. p. 225. ic.
— p. 346. ic.
— p. 345. ic.

Host. gram. 4. t. 55.

Sm. ic. ined. 3. t. 54.
Schad. fl. ger. 1. t. 5. f. 1.
Sinc. gr. wob. p. 342. ic.
Eng. bot. v. 17. t. 1189.
Leers fl. herb. t. 4. f. 3.

Trin. icon. gr. t. 35.

Leers fl. herb. t. 4. f. 4.

Eng. bot. v. 24. t. 1671.
Høst. gram. t. 56.
Trin. ic. gram. t. 29.
Lab. nov. holl. I. t. 23.

21 marítima. R.S. sea-side.	S. Europe.	1825.	7. 8. H. 4.	
22 Forstéri. R.S. Forster's.	N. Zealand.	1822.	— H. 4.	
<i>Avéna filiformis.</i> Forst.					
23 Billardieri. B.P. Labillardiere's.	N. Holland.	1806.	— H. 4.	Lab. nov. holl. 1. t. 31.
24 plebèia. B.P. plebeian.	N. S. W.	1822.	— H. 4.	
25 æmula. B.P. flat-leaved.	—	1806.	— H. 4.	
26 retrofrácta. R.S. broad-leaved.	—	—	4. 5. F. 4.	
27 hirsùta. R.S. hairy.	Teneriffe.	1810.	— F. 4.	
28 Spica véniti. E.B. silky.	England.	6. 7. H. ⊙.	Eng. bot. v. 14. t. 951.
29 pállida. DC. pale.	France.	1818.	— H. ⊙.	
30 interrúpta. R.S. interrupted.	S. Europe.	—	— H. ⊙.	Host. gram. 3. t. 48.
31 purpùrea. R.S. purple.	—	1820.	— H. ⊙.	
32 pauciflóra. R.S. few-flowered.	Switzerland.	1824.	6. 8. H. 4.	Schrad.fl. ger. 1. t. 3. f. 2.
33 décumbens. Host. decumbent.	Hungary.	—	— H. 4.	Host. gram. 4. t. 54.
SPOROBOLUS. B.P.	SPOROBOLUS.	Triandria	Digynia.		
1 diándrus. R.S. diandrous.	E. Indies.	1820.	6. 8. S. ⊙.	
2 pulchélus. R.S. pretty.	N. Holland.	1828.	— F. 4.	
3 indicus. B.P. Indian.	India.	1773.	8. 10. H. ⊙.	Slean. hist. 1. t. 73. f. I.
4 purpurascens. R.S. purplish.	Jamaica.	1820.	— S. ⊙.	
5 tenacíssimus. L.en. tough.	W. Indies.	1801.	— S. 4.	Jacq. ic. 1. t. 16.
6 elongátus. B.P. elongated.	N. S. W.	1822.	— H. ⊙.	
7 micránthus. small-flowered.	Monte Video.	1825.	— H. ⊙.	
COLPODIUM. Trin.	COLPODIUM.	Triandria	Digynia.		
1 Stevénii. Trin. various-coloured.	Caucasus.	1823.	7. 8. H. 4.	
<i>Agróstis versicolor.</i> Stev.					
2 latifólium. Br. broad-leaved.	Melville Isl.	1821.	— H. ⊙.	
TRICHOCHL'OA. Tr.	TRICHOCHL'OA.	Triandria	Digynia.		
1 tenuiflóra. slender-flowered.	N. America.	1820.	7. 8. H. 4.	Willd. hort. ber. t. 12.
<i>longiseta.</i> Tr.	<i>Agróstis tenuiflóra.</i> w.	<i>Cinna tenuiflóra.</i> L.en.			
2 mexicana. Tr. Mexican.	America.	1780.	6. 9. H. 4.	
3 foliòsa. Tr. leafy.	N. America.	1819.	7. 8. H. 4.	
<i>Agróstis filiformis.</i> W.en.	<i>Cinna filiformis.</i> L.en.				
4 sobolífera. Tr. soboliferous.	N. America.	1819.	7. 8. H. 4.	
5 microspérma. DC. small-seeded.	Mexico.	1818.	8. 10. H. ⊙.	
6 capillaris. DC. capillary.	Carolina.	—	— H. 4.	
7 expánsta. DC. expanded.	—	1822.	— H. 4.	P.B.agros. nov. t. 8. f. 2.
<i>purpurea.</i> P.B.					
8 polypògon. DC. many-bearded.	—	1812.	— H. 4.	P.B.agros. nov. t. 8. f. 3.
CINNA. L.	CINNA.	Monandria	Digynia.		
1 arundinácea. L. Reed-like.	Canada.	1789.	7. H. 4.	Schreb. gram. t. 49.
2 crinita. Tr. rough-keeled.	N. Zealand.	1822.	— F. 4.	Lab. nov. hol. 2. t. 263.
3 monándra. R.S. monandrous.	N. America.	1826.	— H. ⊙.	
KNAPPIA. E.B.	KNAPPIA.	Triandria	Digynia.		
agrostidea. E.B. small.	Wales.	3. 4. H. ⊙.	Eng. bot. v. 16. t. 1127.
<i>Agróstis minima.</i> w.	<i>Chamagróstis minima.</i> R.S.	<i>Sturmia verna.</i> P.S.			
COLOBA'CHNE. R.S.	COLOBA'CHNE.	Triandria	Digynia.	R.S.	
vaginàta. R.S. sheathed.	Levant.	1818.	4. H. 4.	Flor. græc. t. 64.
<i>Alopecúrus angustifolius.</i> F.G.	<i>Polypogon vaginatus.</i> w.				
HELEOCHL'OA. Host.	HELEOCHL'OA.	Triandria	Monogynia.		
1 alopecuroídes. R.S. Foxtail-like.	Germany.	1816.	7. 8. H. ⊙.	Host gram. 1. t. 29.
2 phalaroídes. P.B. Phalaris-like.	Caucasus.	1827.	— H. ⊙.	
3 schoenoídes. R.S. Schœnus-like.	S. Europe.	1783.	— H. ⊙.	Jacq. ic. 1. t. 15.
CRYPSIS. Host.	CRYPSIS.	Triandria	Monogynia.		
aculeata. R.S. prickly.	S. Europe.	1783.	— H. ⊙.	Host. gram. 1. t. 31.
LAPPAGO. W.	LAPPAGO.	Triandria	Digynia.	W.	
1 racemòsa. w. branching.	S. Europe.	1771.	— H. ⊙.	Flor. græc. t. 101.
<i>Trágus racemòsus.</i> R.S.	<i>Cénchrus racemòsus.</i> L.				
2 aliena. s.s. Brazilian.	Brazil.	1826.	6. 8. S. ⊙.	
3 biflóra. F.I. two-flowered.	E. Indies.	1824.	— S. 4.	
MUHLENBERGIA. W.	MUHLENBERGIA.	Triandria	Digynia.	W.	
diffusa. R.S. spreading.	N. America.	1816.	5. 6. H. 4.	Schreb. gram. t. 51.
CHÈTARIA. R.S.	CHÈTARIA.	Triandria	Digynia.	R.S.	
1 adscensionis. R.S. West Indian.	W. Indies.	1822.	7. 8. S. 4.	Sloan. hist. 1. t. 2. f. 56.
<i>Arístida adscensionis.</i> w.					
2 Hystrix. R.S. Porcupine.	E. Indies.	—	... S. 4.	Pluk. alm. t. 191. f. 5.
3 oligántha. R.S. long-glumed.	N. America.	1826.	— H.w. 4.	
4 pállens. R.S. pale.	Chile.	1825.	— H. 4.	
5 divaricata. R.S. spreading.	S. America.	1818.	7. 8. G. 4.	Jacq. eclog. gram. t. 6.

GRAMINEÆ.

6 gigantea. R.S.	gigantic.	Teneriffe. 1824.	7. 8. F. ¼.	
7 cœrulescens. R.S.	blueish.	Canaries. 1818.	— H. ⊙.	Desf. atl. t. 21. f. 2.
CURTOPOGON. R.S. CURTOPOGON.	Triandria Dignya.	dichotomus. R.S. forked.	N. America. 1826.	6. 8. H. ♂.
<i>Aristida dichotoma</i> . Ph.				Spr. act. pet. 2. t. 6.
ARISTIDA. R.S. ARISTIDA.	Triandria Dignya. L.			
1 obtusa. R.S.	bluntawned.	Egypt. 1828.	6. 8. H. ⊙.	Delil. ægyp. t. 13. f. 2.
2 vagans. R.S.	spreading-paniced.	N. S.W. 1820.	7. 8. F. ¼.	Cavan. ic. 5. t. 471.
HORDEUM. R.S. BARLEY.	Triandria Dignya. L.			
1 vulgare. R.S.	spring.	Levant.	7. H. ⊙.	Host. gram. 3. t. 34.
β giganteum.	giant.	—	— H. ⊙.	Viborg cereal. t. 1.
2 cœlestis. R.S.	naked-seeded.	Levant.	— H. ⊙.	Viborg cereal. t. 2.
3 nigrum. R.S.	black-seeded.	—	— H. ♂.	Flor. græc. I. t. 98.
4 hexastichon. R.S.	winter.	—	— H. ⊙.	
5 bulbosum. R.S.	bulbosum.	S. Europe. 1770.	— H. ¼.	
6 jubatum. R.S.	long-bearded.	N. America. 1782.	7. 8. H. ♂.	
7 Sieberianum. R.S.	Sieber's.	Egypt. 1828.	— H. ⊙.	
8 distichum. R.S.	common.	Tartary.	7. H. ⊙.	Viborg cereal. t. 3.
β imberbe. R.S.	beardless.	—	— H. ⊙.	
9 nudum. R.S.	naked seeded.	—	— H. ⊙.	
10 Zeocriton. R.S.	Battledore.	—	— H. ⊙.	Arduin. sag. t. II. f. 4.
11 murinum. R.S.	wall.	Britain.	4. 8. H. ⊙.	Viborg cereal. t. 4.
12 pratense. R.S.	meadow.	—	— H. ¼.	Eng. bot. v. 28. t. 1971.
13 maritimum. E.B.	sea-side.	—	— H. ♂.	— v. 6. t. 409.
14 nepalense.	Nepaul.	Nepaul. 1825.	7. 8. H. ⊙.	— v. 17. t. 1205.
15 Hystrix. R.S.	hedgehog.	Spain. 1817.	— H. ⊙.	
16 bifarium. R.S.	two-ranked.	—	— H. ⊙.	
ÆGOP'OGON. R.S. ÆGOP'OGON.	Triandria Dignya. R.S.			
1 pusillus. R.S.	small.	S. America. 1822.	7. 8. H. ⊙.	Cav. h. r. mad. 1. t. 5. f. 2.
2 trisetus. R.S.	three awned.	—	— H. ⊙.	— 1. t. 5. f. 3.
MICROCHLOA. B.P. MICROCHLOA.	Triandria Dignya. R.S.			
setacea. B.P.	bristly.	Tropics. 1806.	7. H. ⊙.	Roxb. corom. 2. t. 132.
MELINIS. S.S. MELINIS.	Triandria Dignya. S.S.			Nees Hor. ber. 54. t. 7.
minutiflora. s.s.	glutinous.	Brazil. 1822.	6. 7. S. ¼.	
<i>Tristegis glutinosa</i> . Nees. <i>Suárdia picta</i> . Schrank h. mon. 11. t. 58.				
DIGITARIA. R.S. DIGITARIA.	Triandria Dignya. R.S.			Eng. bot. v. 12. t. 849.
1 sanguinalis. R.S.	slender-spiked.	Britain.	8. H. ⊙.	
<i>Panicum sanguinale</i> . E.B.				Jacq. obs. 3. t. 30.
2 bicornis. R.S.	two-horned.	E. Indies. 1825.	8. 9. H. ⊙.	
3 affinis. R.S.	related.	S. America. 1820.	7. S. H. ⊙.	
4 ægyptiaca. R.S.	Egyptian.	Egypt. 1794.	— H. ⊙.	
5 filiformis. s.s.	filiform.	N. America. 1781.	— H. ⊙.	
6 Roxburghii. s.s.	Roxburgh's.	E. Indies. 1823.	— H. ⊙.	
<i>Panicum filiforme</i> . F.I.				
7 villosa. s.s.	villous.	N. America. 1781.	7. 9. H. ⊙.	
8 debilis. s.s.	weak.	Barbary. 1824.	— H. ⊙.	
9 glabra. R.S.	smooth.	S. Europe. 1817.	— H. ⊙.	Schrad. germ. 1. t. 3. f. 6.
<i>Syntherisma glabrum</i> . Schrad.				
10 paspalodes. R.S.	trailing.	N. America. —	— H. ⊙.	Host. gram. 4. t. 15.
11 ciliaris. R.S.	fringed.	S. Europe. 1804.	— H. ⊙.	
12 barbata. R.S.	bearded.	E. Indies. 1824.	— H. ⊙.	
13 biformis. R.S.	two-formed.	Canaries. 1820.	— H. ⊙.	
14 marginata. R.S.	margined.	Brazil. 1825.	— H. ⊙.	
15 linearis. R.S.	linear-spiked.	India. 1804.	— H. ⊙.	
16 inaequalis. s.s.	hairy-sheathed.	Manila. 1824.	7. 8. S. ⊙.	Burm. ind. 25. t. 10. f. 3.
17 serotina. M.	late-flowered.	N. America. —	— H. ⊙.	
<i>Paspalum serotinum</i> . R.S.				
18 Andersoni. Ot.	Anderson's.	MonteVideo. 1825.	6. 8. F. ¼.	
19 eriogona. Schr.	woolly-angled.	S. America. —	— S. ¼.	
20 mollissima. Schr.	soft.	—	— S. ¼.	
21 piligera. Lk.	hairy.	MonteVideo. —	— H. ⊙.	
22 violascens. Lk.	violet.	—	— H. ⊙.	
CYNODON. R.S. CYNODON.	Triandria Dignya. R.S.			
1 Dactylon. R.S.	creeping.	England.	7. 8. H. ¼.	
β indicus.	Doob-grass.	India. 1820.	— S. ¼.	Eng. bot. v. 12. t. 850.
2 linearis. R.S.	linear-leaved.	E. Indies. 1796.	— S. ¼.	Sinc. gr. wob. p.290. ic.
3 stellatus. R.S.	starry.	St. Helena. 1823.	— S. ¼.	
4 virgatus. R.S.	twiggy.	E. Indies. 1826.	— S. ¼.	

5 præcox. R.S. early-flowered.	N. America.	1823.	6. 7. H. ¼.	
<i>Digitaria sanguinalis</i> . M.					
BRACHYEL'YTRUM. R.S.	BRACHYEL'YTRUM.	Triandria Digynia.			
aristatum. R.S. awned.	N. America.	1827.	6. 7. H. ¼.	
<i>Muhlenbergia erecta</i> . Ph.	<i>Dilep'yrum aristosum</i> . M.				Schreb. gram. t. 50.
PHALARIS. R.S.	CANARY-GRASS.	Triandria Digynia. R.S.			
1 canariensis. R.S. common.	Britain.	6. 8. H. ⊙.	Eng. bot. v. 19. t. 1310.
2 microstachya. DC. small-spiked.	N. America.	1818.	— H. ⊙.	Host. gram. 2. t. 38.
3 aquatica. R.S. water.	Egypt.	1778.	6. 7. H. ⊙.	Barrel. ic. 700. f. 2.
4 nitida. R.S. glossy.	Sicily.	1824.	— H. ⊙.	
5 commutata. R.S. changed.	Italy.	1823.	— H. ⊙.	
6 capensis. R.S. Cape.	C. B. S.	1804.	— G. ¼.	
7 bulbosa. R.S. bulb-bearing.	Spain.	1824.	— H. ¼.	
8 variegata. R.S. variegated.	Sicily.	1825.	— H. ¼.	
9 caerulescens. R.S. blueish.	S. Europe.	1824.	— H. ⊙.	Buxb. cent. 4. t. 53.
10 nodosa. R.S. knotted.	—	1825.	— H. ¼.	Flor. græc. t. 56.
11 paradoxa. R.S. paradoxical.	—	—	— H. ⊙.	— 1. t. 58.
12 appendiculata. R.S. appended.	Egypt.	1823.	— H. ⊙.	
DIGRA'PHIS. Tr.	DIGRA'PHIS.	Triandria Digynia.			
1 arundinacea. Tr. Reed-like.	Britain.	7. H.w. ¼.	Eng. bot. v. 6. t. 402.
<i>Arundo colorata</i> . E.B.	<i>Phalaris arundinacea</i> . L.	<i>Baldingeria</i> . Fl. wet.			
β variegata. striped-leaved.	—	—	— H. ¼.	
2 americana. s.s. American.	N. America.	— H. ¼.	
LAG'URUS. R.S.	HARE'S-TAIL-GRASS.	Triandria Digynia. L.			
ovatus. E.B. oval-spiked.	Guernsey.	6. H. ⊙.	Eng. bot. v. 19. t. 1334.
ECHINOP'OGON. R.S.	ECHINOP'OGON.	Triandria Digynia.			
ovatus. R.S. ovate panicle.	N. Holland.	1820.	7. 8. H. ¼.	Lab. nov. holl. 1. t. 21.
<i>Agrrostis ovata</i> . Lab.					
HIEROCHL'OA. P.B.	HIEROCHL'OA.	Triandria Digynia. R.S.			
1 borealis. R.S. northern.	Scotland.	5. 6. H. ¼.	Sinc. gr. wob. p. 167. ic.
<i>Holcus odoratus</i> . F.D.	t. 963. <i>repens</i> .	Host. gram. 3. t. 3.			
2 australis. R.S. southern.	S. Europe.	1777.	6. 7. H. ¼.	Host. gram. 1. t. 4.
3 frægrans. R.S. fragrant.	N. America.	—	— H. ¼.	
4 alpina. R.S. Alpine.	Melville Isl.	1827.	— H. ¼.	Flor. dan. t. 1508.
5 pauciflora. R.S. few-flowered.	—	—	— H. ¼.	
CATABR'OSA. R.S.	CATABR'OSA.	Triandria Digynia. R.S.			
1 aquatica. R.S. water.	Britain.	5. 6. H.w. ¼.	Eng. bot. v. 22. t. 1557.
<i>Aira aquatica</i> . E.B.					
2 viridula. R.S. close-panicle.	—	1718.	— H.w. ¼.	
3 humilius. Tr. dwarf.	Caucasus.	1824.	6. 7. H. ¼.	
AI'RA. R.S.	HAIR-GRASS.	Triandria Digynia. L.			
1 flexuosa. E.B. waved.	Britain.	7. 8. H. ¼.	Eng. bot. v. 22. t. 1519.
2 uliginosa. R.S. marsh.	Germany.	1825.	— H. ¼.	Reich. ic. r. C. 2. t. 150.
3 versicolor. R.S. various-color'd.	S. Europe.	1823.	— H. ¼.	
4 caryophyllea. E.B. silver.	Britain.	—	— H. ⊙.	Eng. bot. v. 12. t. 812.
5 elegans. R.S. elegant.	Switzerland.	1824.	— H. ⊙.	Host. gram. 4. t. 35.
6 pulchella. R.S. pretty.	Spain.	1820.	— H. ⊙.	
7 pallens. R.S. pale.	N. America.	—	— H. ¼.	
8 arundinacea. R.S. Reed-like.	Cumana.	1823.	— S. ¼.	
9 semineutra. s.s. doubtful.	Hungary.	1819.	6. 7. H. ¼.	
DESCHA'MPSIA. R.S.	DESCHA'MPSIA.	Triandria Digynia. R.S.			
1 cæspitosa. R.S. turfy.	Britain.	8. H. ¼.	Eng. bot. v. 21. t. 1453.
<i>Aira cespitosa</i> . E.B.					
2 Biebersteiniæna. R.S.	Bieberstein's.	Caucasus.	1823.	6. 8. H. ¼.	
3 refracta. R.S. reflexed.	Spain.	1822.	— H. ¼.	
4 alpina. R.S. Alpine.	Scotland.	5. 6. H. ¼.	Eng. bot. v. 30. t. 2102.
<i>Aira larvigata</i> . E.B.	<i>alpina</i> . L.	<i>Avena alpina</i> . Tr.			
5 brevifolia. R.S. short-leaved.	Melville Isl.	1827.	— H. ¼.	
6 böttnerica. Tr. bristle-leaved.	Bottnia.	1824.	— H. ¼.	
<i>Aira böttnerica</i> . R.S.	<i>montana</i> . R.S.				
SCHI'MMUS. R.S.	SCHI'MMUS.	Triandria Digynia. R.S.			
1 Gouani. Tr. Gouan's.	Montpelier.	1820.	6. 7. H. ¼.	
<i>Aira media</i> . Gouan.	<i>Deschampsia media</i> . et <i>discalcolor</i> . R.S.				
2 marginatus. R.S. margined.	Spain.	1781.	— H. ⊙.	Cavan. ic. 1. t. 44. f. 2.
<i>Festuca calycina</i> . w.	<i>Kæleria calycina</i> . DC.				
3 fasciculatus. P.B. fascicled.	St. Domingo.	1827.	— S. ⊙.	
<i>Digitaria horizontalis</i> . w.					
DUPO'NTIA. R.S.	DUPO'NTIA.	Triandria Digynia.			
Fischéri. R.S. Fischer's.	Melville Isl.	1827.	6. 8. H. ¼.	

GRAMINEÆ.

CORYNE'PHORUS. R.S.	CORYNE'PHORUS.	Triandria Digynia. R.S.				
1 canescens. R.S.	canescent.	England.	7. 8. H. 2.			Eng. bot. v. 17. t. 1190.
<i>Aira canescens.</i> E.B.						
2 articulatus. R.S.	jointed.	S. Europe. 1824.	— H. 2.			Desf. atl. 1. t. 13.
PERIBA'LLIA. Tr.	PERIBA'LLIA.	Triandria Digynia. R.S.				
hispanica. Tr.	Spanish.	Spain. 1820.	6. 7. H. 2.			Cavan. ic. 1. t. 44. f. 1.
<i>Aira involucrata.</i> Cav. <i>Airopsis involucrata.</i> R.S.						
AIRO'PSIS. R.S.	AIRO'PSIS.	Triandria Digynia. R.S.				
1 globosa. R.S.	globe-flowered.	S. Europe. 1823.	6. 8. H. 2.			Cavan. ic. 3. t. 274. f. 1.
2 obtusata. R.S.	blunt-flowered.	N. America. —	6. 7. H. 2.			
3 brevifolia. R.S.	short-leaved.	Missouri. 1818.	— H. 2.			
4 agrostidea. R.S.	Agrostis-like.	S. Europe. 1823.	— H. 2.			DC. ic. rar. t. 1.
<i>Poa agrostidea.</i> DC. <i>Aira agrostidea.</i> Lois.						
MELICA. R.S.	MELIC-GRASS.	Triandria Digynia. L.				
1 ciliata. R.S.	ciliated.	Europe. 1771.	7. H. 2.			Flor. græc. 1. t. 70.
2 nūtans. R.S.	mountain.	Britain.	6. 7. H. 2.			Eng. bot. v. 15. t. 1059.
3 uniflora. R.S.	wood.	—	5. 6. H. 2.			— v. 15. t. 1058.
4 minuta. R.S.	small.	S. Europe. 1825.	— H. 2.			Cavan. ic. 2. t. 175. f. 2.
5 Bauhinia. R.S.	Italian.	Italy. 1806.	6. 7. H. 2.			Host. gram. 4. t. 23.
6 pyramidalis. R.S.	pyramidal.	Barbary. 1804.	— H. 2.			Barrel. ic. t. 96. f. 1.
7 speciosa. R.S.	handsome.	N. America. 1812.	— H. 2.			Schreb. gram. t. 54.
<i>glabra.</i> Ph. <i>rarijöra.</i> Schreb.						
8 altissima. R.S.	tallest.	Siberia. 1770.	7. 8. H. 2.			Host. gram. 2. t. 9.
9 diffusa. R.S.	spreading.	N. America. 1824.	— H. 2.			
10 pálida. R.S.	pale.	Quito. 1825.	— S. 2.			
11 exasperata. R.S.	exasperate.	—	— H. 2.			
12 digitata. R.S.	fingered.	E. Indies. 1824.	— S. 2.			
13 refracta. R.S.	reflexed.	—	— S. 2.			
14 latifolia. R.S.	broad-leaved.	—	— S. 2.			
MOLINIA. P.B.	MOLINIA.	Triandria Digynia. R.S.				
1 cœrulea. P.B.	purple.	Britain.	8. H. 2.			Eng. bot. v. 11. t. 750.
<i>varia.</i> s.s. <i>Melica cœrulea.</i> E.B. <i>Enodium cœruleum.</i> R.S.						
2 sylvatica. R.S.	wood.	Europe. 1824.	— H. 2.			
DIN'EB'A. R.S.	DIN'EB'A.	Triandria Digynia. R.S.				
1 americanæ. R.S.	American.	S. America. 1822.	6. 8. S. 2.			Jacq. fragm. t. 121. f. 1.
2 arábica. R.S.	Arabian.	Levant. 1816.	8. 9. H. 2.			
3 divaricata. R.S.	spreading.	Egypt. 1823.	— H. 2.			
WANGENHEIMIA. Tr.	WANGENHEIMIA.	Triandria Digynia. R.S.				
Líma. Tr.	imbriated.	Spain. 1776.	7. 8. H. 2.			Cavan. ic. 1. t. 91.
<i>Cynosurus Líma.</i> Cav. <i>Dinèba Líma.</i> R.S.						
SA'CCHARUM. R.S.	SUGAR-CANE.	Triandria Digynia. L.				
1 officinárum. R.S.	common.	India. 1597.	... S. 2.			Rumph. amb. 5. t. 74. f. 1.
2 sinense. F.I.	Chinese.	China. 1822.	... S. 2.			Roxb. corom. 3. t. 232.
3 prōcerum. F.I.	lofty.	E. Indies.	... S. 2.			
4 violaceum. R.S.	violet-coloured.	W. Indies. 1816.	... S. 2.			Sloan. hist. 1. t. 70. f. 1.
5 caudátum. R.S.	tailed.	—	... S. 2.			Rheed. mal. 12. t. 46.
6 contráctum. R.S.	contracted.	S. America. 1823.	... S. 2.			
7 exaltátum. F.I.	tall.	E. Indies. 1826.	... S. h.			
8 polystachyum. R.S.	many-spiked.	W. Indies.	... S. h.			
9 Mánja. F.I.	slender.	E. Indies. 1805.	... S. 2.			
IMPER'ATA. R.S.	IMPER'ATA.	Triandria Digynia. R.S.				
1 arundinacea. R.S.	Reed-like.	N. Holland. 1820.	7. 8. F. 2.			Cyril. ic. fasc. 2. t. 11.
2 Thunbergii. R.S.	Thunberg's.	E. Indies. 1822.	... S. h.			Rheed. mal. 12. t. 46.
3 Koenigi. R.S.	Koenig's.	W. Indies. 1818.	... S. h.			
4 spontanea. R.S.	spontaneous.	E. Indies.	... S. h.			
TRICHOLÆNA. R.S.	TRICHOLÆNA.	Triandria Digynia. R.S.				
micrantha. R.S.	small-flowered.	S. Europe. 1816.	7. 8. H. 2.			Jacq. ecl. gram. t. 34.
<i>Saccharum Teneriffæ.</i> Jacq.						
ERIA'NTHUS. R.S.	ERIA'NTHUS.	Triandria Digynia. R.S.				
1 saccharoides. R.S.	Sugar-cane-like.	N. America. 1816.	7. 8. H. 2.			
<i>Saccharum giganteum.</i> Ph.						
2 brevihárbita. R.S.	short-bearded.	—	1822.	— H. 2.		
3 strictus. R.S.	upright.	—	1828.	— H. 2.		
RIPIDIUM. Tr.	RIPIDIUM.	Polygamia Monœcia. Tr.				
1 Ravennæ. Tr.	rough.	S. Europe. 1816.	7. 8. H. 2.			Host. gram. 3. t. 1.
<i>Andropogon Ravennæ.</i> Host. <i>Saccharum Ravenna.</i> w.						
2 strictum. Tr.	upright.	Hungary. 1802.	— H. 2.			Host. gram. 2. t. 2.
<i>Saccharum adpresso.</i> Kit. <i>Andropogon strictus.</i> Host.						

ANDROP'OGON. R.S. ANDROP'OGON.	Polygamia Monœcia. L.			
1 alopecuroides. R.S.	Foxtail. America. 1818.	G. 4.	Sloan. hist. 1. t. 14.	
2 miliförmis. R.S.	Millet-like. E. Indies. —	S. 4.		
<i>miliaceus</i> . F.I. non Forsk.				
3 argenteus. DC.	silvery. Mexico. 1822.	G. 4.	DC. hort. m. ined. t. 68.	
4 saccharoides. R.S.	Sugar-cane-like. Jamaica. 1818. —	S. 4.		
5 laguroides. R.S.	hares-tail-like. New Spain. 1820.	S. 4.	DC. h. mons. ined. t. 69.	
6 hirtus. R.S.	hairy. S. Europe. 1802. 7. 8. S. 4.		Host. gram. 4. t. 1.	
7 angustifolius. R.S.	narrow-leaved. —	1818. —	Jacq. aust. 4. t. 384.	
8 Ischænum. R.S.	woolly. —	1768. —	Gerard. gall. p. 107. t. 4.	
9 serratus. R.S.	saw-leaved. Japan. 1816.	G. 4.		
<i>laxus</i> . w.				
10 comosus. R.S.	tufted. Egypt. 1824.	H. 4.		
11 annulatus. R.S.	annular. —	—	H. 4.	Forsk. egyp. t. 7.
12 furcatus. R.S.	forked. N. America. 1818. 7. 8. H. 4.			
CHRYSOP'OGON. Tr. CHRYSOP'OGON.	Polygamia Monœcia.			
1 Gryllus. Tr.	purple-flowered. S. Europe. 1791. 6. 7. H. 4.		Flor. græc. 1. t. 67.	
<i>Andropogon Gryllus</i> . F.G. <i>Holcus Gryllus</i> . B.P.				
2 aciculatus. Tr.	sharp-pointed. E. Indies. 1825. 6. 8. S. 4.		Rheed. mal. 12. t. 43.	
LEPEOCE'RCIS. Tr. LEPEOCE'RCIS.	Polygamia Monœcia.			
serrata. Tr.	serrated. E. Indies. 1812. 6. 8. S. 4.			
<i>Andropogon serratum</i> . w.				
POLLINIA. R.S.	POLLINIA. Polygamia Monœcia.			
1 distachya. R.S.	two-spiked. S. Europe. 1805. 7. 8. H. 4.		Flor. græc. 1. t. 69.	
2 striata. R.S.	nerve-glumed. E. Indies. 1793. 8. S. 4.			
3 undata. R.S.	waved. Mauritius. 1823. 7. 9. S. 4.		Jacq. ic. 3. t. 361.	
CYMBOP'OGON. R.S. CYMBOP'OGON.	Polygamia Monœcia.			
1 elegans. R.S.	elegant. E. Indies. 1816. 6. 8. S. 4.			
2 prostratus.	prostrate. —	—	S. 4.	
<i>glandulosus</i> . R.S. <i>Anthistiria prostrata</i> . F.I. <i>Andropogon prostratus</i> . L.				
3 Schœnanthus. R.S.	Lemon-grass. E. Indies. 1786.	S. 4.	Rumph. amb. 5. t. 72. f. 2.	
<i>Andropogon Schœnanthus</i> . w.				
4 arundinaceus. R.S.	Reed-like. E. Indies. 1820. 6. 8. S. 4.			
HETEROP'OGON. R.S. HETEROP'OGON.	Polygamia Monœcia.			
1 Allionii. R.S.	Allioni's. S. Europe. 1816. 7. 8. H. 4.		Allion. fl. ped. t. 91. f. 4.	
2 contortus. R.S.	twisted. E. Indies. 1779. 7. 9. S. 4.		Schk. hand. t. 342. f. a.	
<i>Andropogon contortus</i> . w.				
APL'UDA. R.S.	APL'UDA. Polygamia Monœcia. L.			
1 aristata. R.S.	awned. E. Indies. 1820.	S. 4.	Schreb. gram. 2. t. 42.	
2 villosa. R.S.	villous. —	—	S. 4.	
ANATHERUM. R.S.	ANATH'ERUM. Polygamia Monœcia.			
1 muricatum. R.S.	muricate. E. Indies. 1820.	S. 4.		
<i>Andropogon muricatus</i> . Retz. <i>squarrösus</i> . w.				
2 virginicum. R.S.	Virginian. N. America. 1819. 7. 8. H. 4.		Sloan. jam. 1. t. 68. f. 2?	
3 bicorne. R.S.	two-horned. W. Indies. 1825. —	S. 4.	Sloan. hist. 1. t. 15.	
4 caudatum. R.S.	tailed. —	1826. —	S. 4.	
5 müticum. R.S.	smooth-spiked. C. B. S. 1794. 7. 9. G. 4.			
CALAM'INA. R.S.	CALAM'INA. Polygamia Monœcia.			
1 gigantea. R.S.	gigantic. Isl. Luzon. 1822. 7. 8. G. 4.		Cavan. ic. 5. t. 458.	
2 mütica. R.S.	awnless. E. Indies. 1816. —	S. 4.		
<i>Apluda mütica</i> . L.				
CYMBA'CHNE. R.S.	CYMBA'CHNE. Polygamia Monœcia.			
ciliata. R.S.	fringed. E. Indies. 1824. 7. 8. S. 4.			
ANTHISTIRIA. R.S.	ANTHISTIRIA. Polygamia Monœcia. L.			
1 ciliata. R.S.	fringed. Jamaica. 1821. 8. 10. S. ⊖.		Cavan. ic. 5. t. 459.	
2 barbata. R.S.	bearded. Japan. 1816. —	G. ⊖.	Desf. j. ph. 40. p. 294. t. 2.	
3 polystachya. R.S.	many-spiked. E. Indies. 1824. —	S. ⊖.		
4 heterocloia. R.S.	jointed. —	—	S. ⊖.	
5 australis. B.P.	New Holland. N. Holland. 1818. —	F. 4.		
SO'RGHUM. R.S.	SO'RGHUM. Polygamia Monœcia.			
1 vulgare. R.S.	Indian Millet. India. 1596. 7. H. ⊖.		Host. gram. 4. t. 2.	
2 nigrum. R.S.	black-glumed.	—	Ard. sag. pad. 1. t. 5. f. 1.	
3 bicolor. R.S.	two-coloured. Persia. 1731. —	H. ⊖.	Mieg. ac. helv. 8. t. 4. f. 4.	
4 cernuum. R.S.	nodding.	—	Host. gram. 4. t. 3.	
5 rübens. R.S.	red. Africa. 1817. —	H. ⊖.		
6 nervosum. R.S.	nerved-leaved. China. 1827. —	H. ⊖.		
7 saccharatum. R.S.	yellow-seeded. India. 1759. 7. 8. S. ♂.		Host. gram. 4. t. 4.	
8 caffrorum. R.S.	Cape. C. B. S. 1816. —	H. ⊖.	Ard. sag. pad. 1. t. 1. f. 1.	
<i>Arduini</i> . Jacq. ecl. gram. t. 18.				

9 avenáceum. R.S.	Oat-like.	C. B. S. 1816.	7. 9. H. ⊖.	
10 halepénse. R.S.	panicled.	Syria. 1691.	7. 8. H. ⊖.	Flor. græc. 1. t. 68.
11 elongátum. R.S.	long-panicled.	N. Holland. 1823.	— F. ⊖.	
HO'LCUS. R.S.	SOFT-GRASS.	Polygamia Monœcia. L.		
1 mollis. R.S.	creeping.	Britain.	7. 8. H. ⊖.	
2 lanátus. R.S.	meadow.	—	6. 7. H. ⊖.	Eng. bot. v. 17. t. 1170.
ARRHENATH'ERUM. R.S. ARRHENATH'ERUM.		Polygamia Monœcia.		— v. 17. t. 1169.
1 avenáceum. R.S.	Oat-like.	Britain.	6. 7. H. ⊖.	Eng. bot. v. 12. t. 813.
Hólcus avenáceus. E.B. Avéna elatiōr.	F.D. t. 165.			
β mūticum.	awnless.	Scotland.	— H. ⊖.	
2 bulbōsum. Presl.	bulbous.	Germany.	— H. ⊖.	Host. gram. 4. t. 30.
Hólcus bulbōsum. Host. Avéna bulbōsa. w.				
ATHEROP'OGON. R.S. ATEROP'OGON.	Polygamia Monœcia.			
apludoídes. R.S.	Apluda-like.	N. America. 1808.	6. 7. H. ⊖.	Jacq. ecl. gram. t. 7.
Chlōris cartipendula. M.				
CHL'ORIS. R.S.	CHL'ORIS.	Polygamia Monœcia. L.		
1 foliōsa. R.S.	leafy.	St. Thomas. 1822.	7. 9. S. ⊖.	
2 ciliāta. R.S.	ciliated.	Jamaica. 1779.	— S. ⊖.	
3 radiāta. R.S.	many spiked.	W. Indies. 1739.	8. 9. S. ⊖.	Hayn. term. bot. t. 40. f. 2
4 pálida. R.S.	pale-spiked.	S. France. 1816.	— H. ⊖.	
5 barbāta. R.S.	bearded.	E. Indies. 1777.	6. 7. S. ⊖.	Jacq. ecl. gram. 1. t. 8.
6 Roxburghiāna. R.S.	Roxburgh's.	—	1824.	
7 inflāta. R.S.	inflated.	California. —	7. 8. S. ⊖.	
8 fasciculāta. R.S.	crowded.	Brazil. —	— H. ⊖.	
9 carib'ea. R.S.	Caribbean.	W. Indies. 1827.	— S. ⊖.	
10 polydáctyla. R.S.	many-spiked.	—	1818.	
11 elegans. R.S.	elegant.	Mexico. 1825.	— H. ⊖.	Jacq. ecl. gram. 1. t. 9.
12 comprēss'a. R.S.	compressed.	S. France. 1820.	6. 8. H. ⊖.	H. et B. n. gen. l. t. 49.
13 megastachya. R.S.	long-spiked.	N. Holland. 1825.	— F. ⊖.	
14 ventricōsa. B.P.	swelled.	N. S. W. 1818.	— G. ⊖.	
15 truncāta. B.P.	truncated.	—	1822.	
16 pubescens. R.S.	pubescent.	Peru. 1824.	— H. ⊖.	
17 filifōrmis. R.S.	filiform.	—	1825.	
18 distichophylla. R.S.	fan-leaved.	Chile. 1822.	— H. ⊖.	
19 retūsa. R.S.	retuse-glumed.	—	—	
20 dolichostachya. R.S.	long-spiked.	Philippine Isl. 1824.	7. 8. H. ⊖.	
21 crinita. R.S.	crowded-spiked.	—	—	
22 rufescens. R.S.	rufescent.	—	—	
23 tenella. R.S.	slender.	E. Indies. 1825.	— S. ⊖.	
24 grácilis. s.s.	slender.	S. America. 1824.	— S. ⊖.	
25 Durandiana. R.S.	Durand's.	—	—	
EUST'ACHYS. R.S.	EUST'ACHYS.	Polygamia Monœcia.		
1 petr'ea. R.S.	flat-stalked.	Jamaica. 1779.	7. 8. S. ⊖.	Jacq. ecl. gram. 1. t. 11.
Chlōris petr'ea. w.				
2 submútica. R.S.	Mexican.	Mexico. 1827.	— H. ⊖.	H. et B. pl. am. 1. t. 59.
CHRYSURUS. R.S.	CHRYSURUS.	Polygamia Monœcia.		
1 aureus. s.s.	golden-spiked.	Levant. 1770.	7. H. ⊖.	Host. gram. 3. t. 4.
cynosurōides. R.S. Cynosurus aūreus. F.		G. I. t. 79. Lamárekia aūrea. DC.		
2 echinātus. R.S.	rough.	England.	8. H. ⊖.	Eng. bot. v. 19. t. 1333.
Cynosurus echinātus. E.B.				
3 effusus. R.S.	spreading.	S. Europe. 1824.	7. 8. H. ⊖.	Desf. atl. 1. t. 17.
4 elegans. R.S.	elegant.	—	—	
ISCH'ÆMUM. S.S.	ISCH'ÆMUM.	Polygamia Monœcia. L.		
1 mūticum. R.S.	awnless.	E. Indies. 1824.	6. 8. S. ⊖.	Rheed. mal. 12. t. 49.
2 conjugátum. R.S.	yoked.	—	1826.	
3 austrāle. B.P.	southern.	N. S. W. 1822.	7. 8. F. ⊖.	
4 rugosum. s.s.	rough.	E. Indies. 1791.	— S. ⊖.	Sal. stirp. rar. 1. t. 1.
MEO'SCHIUM. R.S.	MEO'SCHIUM.	Polygamia Monœcia.		
1 aristātum. R.S.	awned.	E. Indies. 1808.	6. 7. S. ⊖.	
Isch'ænum aristatum. s.s.				
2 barbātum. R.S.	bearded.	Java. —	— S. ⊖.	Houtt. syst. 12. t. 93. f. 4.
ROTTB'OELLIA. R.S. ROTTBO'ELLI.	Triandria Digynia. L.			
1 exaltāta. R.S.	tall.	E. Indies. 1806.	— S. ♂.	
2 perforāta. R.S.	perforated.	—	1822.	Roxb. corom. t. 182.
3 glábra. R.S.	smooth.	—	—	
HEMA'RTHRIA. B.P.	HEMA'RTHRIA.	Triandria Digynia.		
comprēss'a. B.P.	flat-spiked.	N. S. W. 1822.	6. 7. G. ⊖.	
LODICUL'ARIA. P.B.	LODICUL'ARIA.	Triandria Digynia.		
fasciculāta. P.B.	fascicled.	S. Europe. 1826.	6. 7. H. ⊖.	P. Beauv. agros. t. 11. f. 6.
Rottbøellia fasciculāta. Desf. atl. 1. t. 36.				

STENOTA'PHRUM. Tr.	STENOTA'PHRUM.	Polygamia Monœcia.				
1 glâbrum. Tr.	smooth.	Carolina.	1826.	6. 7. H. 4.		
2 dimidiatum.	creeping-rooted.	W. Indies.	1822.	— S. 4.	Schrank.	h.monac. t. 98.
<i>americanum.</i> Schk.						
3 complanatum. Sck.	flat-spiked.	E. Indies.	1824.	— S. 4.	Burm. ind. t. 8. f. 5.	
LEPTURUS. B.P.	HARD-GRASS.	Triandria Digynia.				
1 incurvatus. Tr.	sea-side.	Britain.	7. H. ⊙.	Eng. Bot. v. 11. t. 760.	
<i>Rottboellia incurvata.</i> E.B.	<i>Ophiurus incurvatus.</i> R.S.					
2 filiformis. Tr.	thready.	S. Europe.	1800.	— H. 4.	Barrel. ic. t. 6.	
3 cylindricus. Tr.	cylindrical.	Portugal.	1806.	— H. ⊙.	Giorn. pis. 4. f. 4-8.	
<i>Rottboellia cylindrica.</i> w. <i>subulata.</i> DC.	<i>Monéma subulata.</i> R.S.					
PHOL'URUS. Tr.	PHOL'URUS.	Triandria Digynia.				
pannonicus. Tr.	Hungarian.	Hungary.	1804.	— H. ⊙.	Host. gram. 1. t. 24.	
MONE'RMA. R.S.	MONE'RMA.	Monandria Digynia.				
monandra. R.S.	monandrous.	Spain.	1804.	— H. ⊙.	Cavan. ic. t. 39. f. 1.	
<i>Rottboellia monandra.</i> Cav.	<i>Nárdus aristata.</i> L.	<i>Psilurus nardoídes.</i> Tr.				
MANIS'URIS. R.S.	MANIS'URIS.	Polygamia Monœcia. W.				
granularis. R.S.	round grained.	E. Indies.	1784.	6. 7. S. ⊙.	Roxb. corom. 2. t. 118.	
ANTHE'PHORA. R.S.	ANTHE'PHORA.	Triandria Monogynia.				
1 elegans. R.S.	elegant.	Jamaica.	1776.	8. 9. S. ⊙.	Schreb. gram. t. 44.	
<i>Tripsacum hermaphroditum.</i> L.	<i>Cénchrus levigatus.</i> s.s.					
2 villosa. R.S.	villous.	W. Indies.	1824.	— S. ⊙.		
TRI'PSACUM. S.S.	TRI'PSACUM.	Monœcia Triandria. L.				
1 dactyloides. s.s.	rough-seeded.	Virginia.	1640.	8. H. 4.	Lam. ill. t. 750.	
2 monostachyon. s.s.	single-spiked.	N. America.	1815.	— H. 4.	Willd. hort. ber. t. 1.	
ECHINOCHLO'A. R.S.	ECHINOCHLO'A.	Triandria Digynia. R.S.				
1 stagnina. R.S.	pond.	E. Indies.	1802.	7. 8. H.w. ⊙.		
2 commutata. R.S.	Hungarian.	Hungary.	—	— H.w. ⊙.	Host. gram. 3. t. 51.	
<i>Pánicum stagninum.</i> Host.						
3 intermédia. R.S.	intermediate.	—	1816.	— H. ⊙.		
4 erythrosperma. R.S.	red-seeded.	—	1825.	— H. ⊙.		
5 hispida. R.S.	hispid.	Egypt.	1829.	— H. ⊙.		
6 Crus-Córví. R.S.	Crow's-foot.	E. Indies.	1781.	— H. ⊙.		
7 Crus-Gálí. R.S.	Cock's-foot.	England.	— H. ⊙.	Eng. bot. v. 13. t. 876.	
<i>Pánicum Crus-Gálí.</i> E.B.						
8 echinata. R.S.	bristly.	N. America.	1825.	— H. ⊙.	Jacq. ecl. gram. t. 20.	
9 setígera. R.S.	setigerous.	E. Indies.	1820.	— S. ⊙.		
ORTHOP'OGON. B.P.	ORTHOP'OGON.	Triandria Digynia. S.S.				
1 hirtellus. s.s.	hairy.	W. Indies.	1795.	6. 7. S. ♂.		
2 setárius. s.s.	bristly.	—	1823.	— S. ⊙.		
3 undulatifolius. s.s.	wave-leaved.	S. Europe.	1795.	6. 7. H. ⊙.	Host. gram. 3. t. 52.	
<i>Pánicum hirtellum.</i> Host. non L.						
4 Burmanni. R.S.	Burman's.	E. Indies.	1819.	— S. ⊙.	Burm. ind. t. 12. f. 1.	
5 africanus.	African.	Africa.	1822.	7. 9. S. ⊙.	Pal. fl. d'ow. 2. t. 67. f. 1.	
<i>Oplismenus africanus.</i> R.S.						
6 compósitus. s.s.	compound spiked.	E. Indies.	1819.	— S. ⊙.	Linn.pfl.sys.12.t.96.f.1.	
7 fláccidus. B.P.	flaccid.	N. S. W.	1824.	— H. ⊙.		
8 imbecillus. B.P.	feeble.	—	—	— H. ⊙.		
PA'NICUM. R.S.	PANIC-GRASS.	Triandria Digynia. R.S.				
1 colónum. R.S.	purple.	E. Indies.	1699.	7. 8. H. ⊙.	Ehret pict. t. 3. f. 3.	
2 fluitans. R.S.	floating.	—	1825.	— S.w. ⊙.		
3 brizoídes. R.S.	Briza-like.	—	1801.	6. 7. S. ♂.	Jacq. gram. fasc. t. 2.	
4 flavidum. R.S.	yellowish.	—	1823.	— S. ♂.		
5 mólle. R.S.	soft.	W. Indies.	1824.	— H. ⊙.		
6 Michauxianum. R.S.	Michaux's.	N. America.	—	— H. ⊙.		
<i>Pánicum mólle.</i> Mich. non Swartz.						
7 fasciculatum. R.S.	fascicled.	Jamaica.	1801.	— H. ⊙.		
8 pláticum. R.S.	plaited.	E. Indies.	1824.	6. 8. S. 4.		
9 palmifolium. R.S.	Palm-leaved.	—	1804.	— S. 4.	Jacq. ecl. gram. t. 1.	
10 neuródies. R.S.	nerved-leaved.	Nepaul.	1823.	— G. 4.		
<i>nervosum.</i> F.I. non Lam.						
11 costatum. F.I.	ribbed.	Mauritius.	1805.	— S. 4.		
12 carthaginése. R.S.	Carthagena.	S. America.	1818.	— S. 4.		
13 conglomerátum. R.S.	conglomerate.	E. Indies.	—	— S. 4.		
14 índicum. F.I.	Indian.	—	—	— S. 4.		
15 hispídulum. R.S.	hispid.	—	1804.	7. 8. H. ⊙.		
16 frumentáceum. F.I.	cultivated.	—	1810.	— H. ⊙.		
17 dichótومum. R.S.	forked.	N. America.	1817.	— H. ⊙.	Sp.m.act.pet.1810.t.4.	
18 coloratum. R.S.	coloured.	Egypt.	1771.	7. 9. H. ⊙.	Jacq. ic. 1. t. 58 ?	

GRAMINEÆ.

19 répens. R.S.	slender.	S. Europe. 1777.	7. 9. H.○.	Flor. græc. 1. t. 61.
20 proliferum. R.S.	proliferous.	N. America. 1816.	— H.○.	
21 diffórmē. R.S.	deformed.	E. Indies. 1825.	— S. 4.	
22 numidiānum. R.S.	Numidian.	Barbary. 1816.	— H.○.	Desf. atl. 1. t. 11.
23 miliáceum. R.S.	Millet.	E. Indies. 1596.	— H.○.	Host. gram. 2. t. 20.
24 ténue. R.S.	slender.	—	— S. 2.	
25 tenuissimum. R.S.	very slender.	Brazil. 1827.	— S. 2.	
26 attenuátum. R.S.	attenuated.	—	— H.○.	
27 gongylôdes. R.S.	branching-flowered.	—	— H. 4.	
28 miliáre. R.S.	Millet-like.	E. Indies. 1824.	— H.○.	
29 muricatum. R.S.	prickly.	—	— H.○.	
30 capillâre. R.S.	hair-pancled.	America. 1758.	6. 8. H.○.	Host. gram. 4. t. 16.
β minor. R.S.	smaller.	—	— H.○.	
31 tenellum. R.S.	slender-branch'd.	Sierra Leon. 1820.	— S.○.	
32 grossârium. R.S.	cultivated.	W. Indies. 1810.	— H.○.	
33 máximum. R.S.	Guinea-grass.	Tropics. 1822.	— S. 4.	Jacq. ic. 1. t. 13.
34 apérrium. R.S.	rough.	S. America. 1824.	7. 9. H.○.	
35 fuscum. R.S.	brown.	Jamaica. 1822.	— S. 2.	
36 latifolium. R.S.	broad-leaved.	N. America. 1765.	8. 9. H. 4.	Moris. his. 3. t. 5. f. 4.
37 macrocarpon. R.S.	large-fruited.	—	— H. 4.	
38 microcarpon. R.S.	small-fruited.	—	— H. 4.	
39 diffusum. R.S.	diffuse.	W. Indies. 1823.	— S.○.	
40 oryzoides. R.S.	Rice-like.	—	— S.○.	
41 clandestinum. R.S.	hidden-flowered.	N. America. 1802.	7. H. 4.	Spr. m. act. pet. 2. t. 5.
42 virgâtum. R.S.	twiggy.	—	8. 9. H. 4.	
43 ánceps. R.S.	flat-edged.	—	7. 8. H. 4.	
44 uliginosum. R.S.	marsh.	E. Indies. 1819.	— S.w. 4.	
45 nitidum. R.S.	glossy.	N. America. 1811.	— H.○.	
46 arundinâceum. R.S.	Reed-like.	Jamaica. 1819.	— S. 4.	
47 tripheron. R.S.	brown-seeded.	E. Indies. 1820.	6. 8. S.○.	
<i>tenellum. F.I. nec aliorum. Roxburghii. s.s.</i>				
48 brevitolium. R.S.	short-leaved.	—	7. 8. S. 4.	Pluk. alm. t. 189. f. 4.
49 divaricatum. R.S.	divaricate.	Jamaica. —	— S. 4.	Jacq. schœn. 1. t. 25.
50 interrúptum. R.S.	interrupted.	E. Indies. 1820.	— S. 4.	
51 arboréscens. R.S.	arborescent.	—	3. 4. S. h.	
52 pâtens. R.S.	spreading.	—	7. 8. S. 4.	Burm. ind. t. 10. f. 3.
53 altissimum. R.S.	lofty.	—	— H.○.	
54 eriogonum. R.S.	woolly-angled.	N. Holland. 1824.	— G. 4.	
55 marginatum. B.P.	margin-leaved.	N. S. W. 1822.	— F. 4.	
56 bicolor. B.P.	two-coloured.	—	— F. 4.	
ISA'CHNE. B.P.	Isa'chne. Triandria Dignynia. R.S.	—	6. 8. H.w. 4.	
australis. B.P.	New Holland. N. S. W. 1820.	—		
MONA'CHNE. R.S.	MONA'CHNE. Triandria Dignynia. R.S.	—		
unilaterális. R.S.	one-sided.	Jamaica. 1819.	6. 8. S. 4.	Sloan. hist. 1. t. 14. f. 2.
<i>Milium villosum. w.</i>				
AXO'NOPUS. R.S.	AXO'NOPUS. Triandria Dignynia. R.S.	—		
1 digitátus. R.S.	fingered.	Jamaica. 1818.	6. 8. H.○.	
2 cimicinus. R.S.	spotted.	E. Indies. 1778.	7. 9. H.○.	
<i>Pánicum cimicinum. w. Milium cimicinum. L.</i>				
BECKMA'NNIA. R.S.	BECKMA'NNIA. Triandria Dignynia. R.S.	—		
erucæfórmis. R.S.	linear-spiked.	Europe. 1773.	7. H.○.	Host. gram. 3. t. 6.
SEC'ALE. R.S.	RYE. Triandria Dignynia. R.S.	—		
1 cereale. R.S.	common.	Tauria. —	6. 7. H.○.	
α hybrénum. R.S.	winter.	—	— H. 4.	Blackw. t. 424.
β vérrnum. R.S.	spring.	—	— H.○.	Host. gram. 2. t. 48.
γ compôsítum. R.S.	compound-spiked.	—	— H.○.	
2 frágile. R.S.	brittle.	—	— H.○.	
3 villosum. F.G.	villus.	S. Europe. 1790.	— H.○.	Flor. græc. t. 97.
<i>Triticum villosum. R.S.</i>				
4 sylvéstre. Host.	wood.	Hungary. 1816.	— H. 4.	Host. gram. 4. t. 2.
5 créticum. w.	Cretan.	Candia. —	— H. 4.	
TETRAP'OGON. S.S.	TETRAP'OGON. Polygamia Monococcia.	—		Desf. atl. 2. t. 255.
villosus. s.s.	villus.	Barbary. 1818.	7. 8. H.○.	
TRITICUM. R.S.	WHEAT. Triandria Dignynia. L.	—		
1 æstivum. R.S.	summer.	Baschiros. —	6. 7. H.○.	Host. gram. 3. t. 26.
α ambigens. LG. β Bâhi. LG. γ Báncos. LG. δ Câlot. LG. ε cándidum. LG. ζ compáctum. LG.				
η Lamárckii. LG. θ Rôyo. LG. ι truncátulum. LG. κ variegátum. LG. λ confértum. LG.				
μ hùmíle. LG. ν médium. LG. ξ mágus. LG. \o Pástor. LG. π versicolor. LG.				

2	hybérnum. R.S.	Lammas.	6. 7. H. ♂.	Host. gram. 3. t. 26. f. 8
	<i>a candidissimum.</i> LG.	β compáctum. LG.	γ Morisóni. LG.	δ Pôlo. LG.	ϵ Tessieri. LG.	
3	sibíricum. R.S.	Siberian.	Siberia.	— H. ♂.	
4	velutínūm. R.S.	velvet-spiked.	— H. ♂.	Rod. Schub.cer.n.4.f.4.
5	compósitum. R.S.	Egyptian.	Egypt. 1799.	— H. ♂.	Host. gram. 3. t. 27.
	<i>a aureum.</i> LG.	β compositum. LG.	γ edéntulum. LG.	δ prismáticum. LG.	ϵ rúbrum. LG.	
6	Linnaeánūm. R.S.	Linnean.	Egypt.	6. 7. H. ♂.	
	<i>a Caballero.</i> LG.	β dùbium. LG.	γ Lajusticia. LG.	δ depilatum. LG.	ϵ píceo-cæruleum. LG.	
7	túrgidum. R.S.	turgid.	6. 7. H. ♂.	Host. gram. 3. t. 28.
	<i>a Collado.</i> LG.	β compósitum. LG.	γ montanum. LG.	δ Tournefórtii. LG.	ϵ Caüms. LG.	
	ζ Lapiedra. LG.					
8	fastuósūm. R.S.	thick-grained.	6. 7. H. ♂.	
	<i>a albellum.</i> LG.	β A'sso. LG.	γ Cannabáte. LG.	δ Cástro. LG.	ϵ Echeándia. LG.	
	η Alvarez. LG.	θ Lobellia. LG.				
9	Gærtneriánum. R.S.	Gærtner's.	H. ♂.	
	<i>a Bernal.</i> LG.	β cávum. LG.	γ ciliósum. LG.	δ Grælls. LG.	ϵ Oliet. LG.	
	η speñoides. LG.	θ truncatūm. LG.	ι Vasallo. LG.	κ yépes. LG.	λ Halleri. LG.	
	ν Valcárcel. LG.	ξ Xixóna. LG.	\circ Cleménte. LG.	π Colóna. LG.	μ Párck. LG.	
	τ Moráles. LG.	υ Plákenet. LG.	ϕ Sammártin. LG.	χ Theophrásti. LG.	σ Gleichen. LG.	
10	platystachyūm. R.S.	flat-spiked.	6. 7. H. ♂.	
11	cochlearé. R.S.	spoon-like.	— H. ♂.	
	<i>a aristátum.</i> LG.	β mucronatum. LG.				
12	Cevállos. R.S.	Trigo Moro.	— H. ♂.	
13	dúrum. R.S.	Barbary.	— H. ♂.	
14	villósūm. Host.	villous.	— H. ♂.	Host. gram. 4. t. 6.
15	sículum. R.S.	Sicilian.	— H. ♂.	
16	hordeifórme. R.S.	Barley form.	— H. ♂.	Host. gram. 4. t. 5.
17	compáctum. R.S.	compact.	— H. ♂.	— 4. t. 7.
18	polónicūm. R.S.	Polish.	1692.	— H. ♂.	— 3. t. 31.
19	monocóccon. R.S.	one-grained.	1648.	— H. ♂.	— 3. t. 32.
	<i>a glábrum.</i> LG.	β pubescens. LG.				
20	dicóccum. R.S.	two-grained.	— H. ♂.	Rod. Schub.cer.n.15.f.2
	<i>a álbum.</i>	white-spiked.	— H. ♂.	
	β rúfum.	brown-spiked.	— H. ♂.	
21	atrátum. R.S.	dark-spiked.	— H. ♂.	Host. gram. 4. t. 8.
22	tricóccum. R.S.	three-grained.	— H. ♂.	Rod. Schub.cer.n.17.f.3
23	Cienfuégos. R.S.	tooth-valved.	— H. ♂.	
24	Bauhini. R.S.	truncate-valved.	— H. ♂.	
25	Spélta. R.S.	Spelt.	— H. ♂.	Host. gram. 3. t. 29.
	Zéa. Host.					
	<i>a Astúrum.</i> LG.	β rufescens. LG.	γ submúticum. LG.	δ álbum. LG.	ϵ cyáneum. LG.	
26	venulósūm. R.S.	veined.	Egypt.	6. 7. H. ♂.	
27	ichyostachyūm. R.S.	slender-spiked.	— H. ♂.	
	AGRÓPYRÚM. R.S.	COUCH-GRASS.	Triandria Dignynia. R.S.			
1	júnceum. R.S.	rushy.	England.	6. 7. H. Ȑ.	Eng. bot. v. 12. t. 814.
	<i>Triticum júnceum.</i> E.B.					
2	acútum. R.S.	acute-glumed.	France. 1816.	— H. Ȑ.	
3	littorále. Host.	sea-side.	Mediterranean. —	— H. Ȑ.	Host. gram. 4. t. 9.
4	rígidum. R.S.	rigid.	S. Europe. 1805.	— H. Ȑ.	
5	intermèdium. Host.	intermediate.	Germany.	—	— H. Ȑ.	Host. gram. 2. t. 22.
6	glaúcum. R.S.	glaucous.	Switzerland. 1824.	— H. Ȑ.	— 4. t. 10?
7	rupéstre. R.S.	rock.	Siberia. 1822.	— H. Ȑ.	
8	obtusiflórum. R.S.	blunt-flowered.	— H. Ȑ.	
9	púngens. R.S.	pungent.	S. Europe. 1824.	— H. Ȑ.	
10	densiflórum. R.S.	close-flowered.	Siberia.	— H. Ȑ.	
11	gigánteum. R.S.	gigantic.	S. Europe. 1805.	— H. Ȑ.	Host. gram. 2. t. 23.
	<i>Triticum elongátum.</i> Host.					
12	répens. R.S.	common.	Britain.	7. 8. H. Ȑ.	Eng. bot. v. 13. t. 909.
	<i>a arvénse.</i> R.S.	field.	— H. Ȑ.	Trattin. tab. t. 566.
	β subulátum. R.S.	subulate-glumed.	— H. Ȑ.	Schreb. gram. t. 26. f. 2.
	γ dumetórum. R.S.	many-flowered.	— H. Ȑ.	— II.IV.f.3.
	δ Vaillantiánum. R.S.	Vaillant's.	— H. Ȑ.	— f. 6.
	ϵ Leersiánūm. R.S.	Leers's.	— H. Ȑ.	Leers fl. herb. t. 12. f. 4.
	ζ capilláre. R.S.	hair-awned.	— H. Ȑ.	
13	sibíricum. R.S.	Siberian.	Siberia. 1825.	— H. Ȑ.	
14	dasýanthum. R.S.	thick-flowered.	Russia.	— H. Ȑ.	
15	caninum. R.S.	bearded.	Britain.	— H. Ȑ.	Eng. bot. v. 20. t. 1372
16	dístichum. R.S.	two-ranked.	C. B. S. 1816.	— G. Ȑ.	
17	prostrátum. R.S.	prostrate.	Siberia. 1780.	6. H. ♂.	Jacq. vind. 3. t. 44.

18 orientâle. R.S. hairy-spiked. <i>Secale orientale</i> . w.	Levant.	1807.	6. 7. H. ♂.	Will.n.act.ber.2.t.4.f.3.
19 pátulum. Tr. spreading. <i>Triticum squarrösüm</i> . R.S. <i>Buonapártis</i> . Spr.	Egypt.	1800.	— H. ♂.	
20 imbricatum. R.S. imbricated.	Caucasus.	1824.	6. 8. H. ♀.	Gmel. sib. 1. n. 51. t. 23.
21 pectinifórmē. R.S. comb-like.	Tauria.	—	— H. ♀.	Buxb. cent. 1. t. 50. f. 3.
22 cristátum. R.S. crested.	—	—	— H. ♀.	Eng. bot. v. 32. t. 2267.
23 desertórum. R.S. desert.	Altay.	1822.	— H. ♀.	Ledeb. ic. alt. t. 246.
24 angustifólium. R.S. narrow-leaved.	Siberia.	—	— H. ♀.	
25 muricátum. R.S. bristly.	—	1825.	— H. ♀.	
26 pectinátum. R.S. pectinated.	V. Diem. Isl.	1822.	— H. ♀.	
27 variegátum. R.S. variegated.	—	1818.	— H. ♀.	
L'OLIUM. R.S.	DARNEL. <i>Triandria Digynia</i> . L.			
1 peréne. R.S. Rye-grass.	Britain.	5. 6. H. ♀.	Eng. bot. v. 5. t. 315.
α vulgare. R.S. common.	—	— H. ♀.	Host. gram. 1. t. 25.
β tenué. R.S. slender.	—	— H. ♀.	Sinc. gra.wob. p.211. ic.
γ compósitum. R.S. compound.	—	— H. ♀.	Scheuch. agr. 1. f. 7. D.
δ ramösüm. R.S. branching.	—	— H. ♀.	Leers gram. t. 12. f. 1.
ε Russelliánum. H.G.W. <i>Russell's</i> .	—	— H. ♀.	Sinc. gra.wob. p. 216. ic.
ζ Whitwoorthii. H.G.W. <i>Whitworth's</i> .	—	— H. ♀.	
η Stickneyanum. H.G.W. <i>Stickney's</i> .	—	— H. ♀.	
ϑ paniculátum. H.G.W. <i>panicked</i> .	—	— H. ♀.	
ι monströsüm. H.G.W. <i>monstrous-flowered</i> .	—	— H. ♀.	
κ viviparum. R.S. viviparous.	—	— H. ♀.	
λ multiflòrum. R.S. many-flowered.	—	—	— H. ♀.	
μ hùmile. R.S. dwarf.	—	— H. ♀.	
2 multiflòrum. R.S. many-flowered.	Europe.	1816.	— H. ♂.	Vaill. par. t. 17. f. 3.
3 arvénse. R.S. beardless.	England.	7. H. ♂.	Eng. bot. v. 16. t. 1125.
4 canadénse. Panz. Canadian.	Canada.	1822.	— H. ♂.	
5 complanátum. L.en. rigid.	Switzerland.	1825.	— H. ♂.	
<i>rígidum</i> . R.S.				Eng. bot. v. 16. t. 1124.
6 temuléntum. R.S. bearded.	Britain.	7. 8. H. ♂.	
7 speciosum. R.S. pretty.	Tauria.	1824.	— H. ♂.	
Æ'GILOPS. R.S.	HARD-GRASS.			
1 caudáta. R.S. tailed.	Polygamia Monœcia. L.			
2 cylindrica. R.S. cylindrical.	Candia.	1820.	7. 8. H. ♂.	Host. gram. 2. t. 7.
3 ováta. R.S. oval-spiked.	Hungary.	1805.	6. 7. H. ♂.	Flor. græc. 1. t. 93.
4 triariastàta. R.S. three-awned.	S. Europe.	1683.	— H. ♂.	Host. gram. 2. t. 6.
<i>triuncialis</i> . Host. non w.	—	1739.	7. 8. H. ♂.	Schreb. gram. t.10. f. 1.
5 triuncialis. R.S. long-spiked.	—	—	— H. ♂.	Schreb.gram.2.t.27.f.2.
6 comòsa. R.S. tufted.	Greece.	1828.	— H. ♂.	
7 squarrösa. w. rough-spiked.	Levant.	1794.	6. 7. H. ♀.	
<i>Triticum squarrösüm</i> . R.S.				
8 Hy'strix. N. hedgehog.	Missouri.	1819.	7. 8. H. ♀.	
E'LYMUS. R.S.	LYME-GRASS.			
1 arenárius. E.B. sand.	Triandria Digynia. L.			
2 geniculátus. E.B. pendulous.	Britain.	4. 6. H. ♀.	Eng. bot. v. 24. t. 1672.
3 Delileánus. R.S. Delile's.	England.	7. H. ♀.	— v.23. t. 1586.
4 sabulösus. R.S. glaucous.	Egypt.	1827.	— H. ♂.	Delil. egypt. t. 13. f. 1.
5 giganteus. R.S. gigantic.	Siberia.	1806.	6. 7. H. ♀.	
6 mexicanus. R.S. Mexican.	N. America.	1790.	7. 8. H. ♀.	Jacq. ecl. gr. t. 19.
7 sibiricus. R.S. Siberian.	Mexico.	1823.	6. 8. H. ♀.	Schreb.gram.1.t.21.f.1.
8 móllis. R.S. soft.	Siberia.	1758.	6. 7. H. ♀.	
9 racemosus. R.S. racemed.	Kamtschatka.	1829.	— H. ♀.	
10 téner. R.S. tender.	—	1816.	— H. ♀.	
11 philadélpicus. R.S. Philadelphian.	Siberia.	1801.	— H. ♀.	
12 canadénsis. R.S. Canadian.	N. America.	1790.	7. 8. H. ♀.	Moris. s. 8. t. 2. f. 10.
13 glaucifòlius. R.S. glaucous.	—	1699.	— H. ♀.	
14 villösus. R.S. villous.	—	1816.	— H. ♀.	
15 virginicus. R.S. Virginian.	—	1802.	6. 7. H. ♀.	
16 striátus. R.S. striated.	—	1781.	— H. ♀.	
β minor. R.S. lesser.	—	1790.	— H. ♀.	
17 europæus. R.S. European.	England.	—	— H. ♀.	Eng. bot. v. 19. t. 1317.
18 crinitus. R.S. long-awned.	Smyrna.	1806.	7. 8. H. ♂.	Host. gram. 1. t. 27.
19 intermèdus. R.S. intermediate.	Iberia.	1819.	7. 9. H. ♂.	Buxb. cent. 1. t. 52.
20 altáicus. F. Altaic.	Altay.	1829.	— H. ♀.	
21 pauciflorus. R.S. few-flowered.	—	1824.	— H. ♂.	
22 Cáput Medúsa. R.S. Portuguese.	Portugal.	1784.	— H. ♂.	Schreb.gram.2.t.24.f.2.
23 júnceus. R.S. rushy.	Siberia.	1823.	7. 8. H. ♀.	

24 hordeiformis. R.S. Barley-like.	1816.	7. 8. H.○.	
25 Hy'strix. R.S. rough. Levant. 1770.	—	H.○.	Jacq. ic. 2. t. 305.
Asprilla Hy'strix. W.en.			
26 pseudo-hy'strix. R.S. rough American. N. America. 1818.	—	H.○.	
ENNEAP'OGON. R.S. ENNEAP'OGON. Triandria Digynia. R.S.			
1 nigricans. R.S. blackish. N. S. W. 1822.	7. 8. F. 2.		
2 pallidus. R.S. pale. ——— 1824.	—	F. 2.	
Pappóphorum pallidum. B.P.			
3 phleoides. R.S. Phleum-like. S. America. 1822.	—	F. 2.	
PAPPO'PHORUM. B.P. PAPPO'PHORUM. Triandria Digynia. R.S.			
1 alopecuroideum. R.S. Foxtail-like. S. America. 1825.	6. 8. S. 2.		
2 laguroides. R.S. Harestail-like. W. Indies. 1826.	—	S. 2.	
ECHIN'ARIA. R.S. ECHIN'ARIA. Triandria Digynia. R.S.			
capitata. R.S. headed. S. Europe. 1771.	5. 8. H.○.		
Cénchrus capitatus. F.G. Séssleria echinata. Host. gram. 3. t. 8.			
SESL'ERIA. R.S. SESL'ERIA. Triandria Digynia. R.S.			
1 elongata. R.S. long-spiked. Austria. 1805.	6. 7. H. 2.		
2 cylindrica. R.S. cylindrical. S. Europe. 1818.	—	H. 2.	
3 tenuifolia. R.S. slender-leaved. ———	4. 6. H. 2.		
4 cœrulea. R.S. blue. Britain.	4. 5. H. 2.		
5 albicans. R.S. whitish. Carpathian Mt. 1826.	—	H. 2.	
6 nítida. R.S. glossy. Italy. 1825.	5. 6. H. 2.		
7 tenella. R.S. weak. Switzerland. 1810.	—	H. 2.	
8 sphærocéphala. R.S. round-headed. ———	—	H. 2.	
9 disticha. R.S. distichous. ——— 1824.	—	H. 2.	
Poa disticha. Host.			
TRI'ODIA. B.P. TRI'ODIA. Triandria Digynia. R.S.			
decumbens. R.S. decumbent. Britain.	7. 8. H. 2.		
Poa decumbens. E.B. Festuca decumbens. w. Danthonia decumbens. DC.			
REDTOP-GRASS. Triandria Digynia. R.S.			
TR'IDENS. R.S. quinquéfida. R.S. five-cleft. N. America. 1816.	7. 8. H. 2.		
Triodìa cupræxa. Jacq. Windsöria poeñfornis. n. Poa quinquéfida. Ph.			
DANTH'ONIA. DC. DANTH'ONIA. Triandria Digynia. R.S.			
1 longifolia. B.P. long-leaved. N. S. W. 1823.	7. 8. F. 2.		
2 semiannularis. B.P. semiannual. N. Holland. 1822.	—	F. 2.	
Arundo semiannularis. Lab.			
3 pilosa. B.P. hairy. N. S. W. 1823.	—	F. 2.	
4 provincialis. R.S. Provence. S. Europe. 1818.	6. 7. H. 2.		
5 spicata. R.S. spiked. N. America. 1823.	—	H. 2.	
6 strigosa. R.S. bristle-pointed. Britain.	—	H. 2.	
Avena strigosa. E.B.			
7 macrantha. R.S. large-flowered. C. B. S. 1816.	—	H.○.	
8 curvifolia. R.S. curve-leaved. ———	—	G. 2.	
9 calycina. R.S. glossy-calyxed. ——— 1822.	—	G. 2.	
KÖEL'ERIA. R.S. KÖEL'ERIA. Triandria Digynia. R.S.			
1 cristata. R.S. crested. Europe.	6. 8. H. 2.		
β glabra. R.S. smooth. Britain.	7. 8. H. 2.		
Aira cristata. E.B. Poa cristata. w.			
2 grandiflora. R.S. great-flowered. S. Europe. 1824.	6. 8. H. 2.		
3 glauca. DC. glaucous. Germany. 1816.	—	H. 2.	
4 pensylvanica. DC. Pensylvanian. N. America. 1818.	—	H. 2.	
5 nítida. R.S. glossy. ——— 1828.	—	H. 2.	
6 macrantha. R.S. large-flowered. Russia. 1824.	—	H. 2.	
7 albescens. R.S. whitish. S. France. 1829.	—	H. 2.	
8 valesiaca. R.S. Swiss. Switzerland. 1824.	—	H. 2.	
9 setacea. R.S. bristly. S. Europe. 1825.	—	H. 2.	
10 tuberosa. P.S. tuberous. ——— 1802.	7. 8. H. 2.		
11 hirsuta. R.S. hairy. Switzerland. 1824.	—	H. 2.	
12 villosa. R.S. villous. S. Europe. 1800.	6. 7. H. 2.		
Aira pubescens. w. Phalaris pubescens. Lam.			
13 phleoides. R.S. cat's-tail. Portugal. 1802.	7. 8. H.○.		
14 hispida. R.S. hispid. S. Europe. 1819.	6. 7. H.○.		
15 brachystachya. R.S. short-spiked. ———	—	H.○.	
16 parviflora. L.en. small-flowered. S. Europe. ———	—	H.○.	
Avena parviflora. Desf. Trisetum parviflorum. R.S.			
17 egyptiaca. Lk. Egyptian. Egypt. 1826.	—	H.○.	
18 laxa. Th. loose-flowered. ——— 1829.	—	H.○.	
AVENA. R.S. OAT-GRASS. Triandria Digynia. L.			
1 præcox. R.S. early-flowering. Britain.	5. 6. H.○.		
Aira præcox. E.B.			
	2 O 4		
		Eng. bot. v. 18. t. 1296.	

GRAMINEÆ.

2 brevis. R.S. short.	Germany.	1804.	5. 6. H.○.	Host. gram. 3. t. 42.
3 álba. R.S. white.	France.	6. 7. H.○.	Host. gram. 2. t. 59.
4 sativa. R.S. common.	S. America.	— H.○.	
<i>α melanospérma.</i>	<i>black Oat.</i>	—	— H.○.	
<i>β leucospermum.</i>	<i>white Oat.</i>	—	— H.○.	
5 orientális. R.S. Tartarian.	Levant.	1798.	— H.○.	Host. gram. 3. t. 44.
6 nuda. R.S. naked.	— H.○.	— 3. t. 43.
7 fátua. R.S. wild.	Britain.	8. H.○.	Eng. bot. v. 31. t. 2221.
8 hirsúta. R.S. hairy.	S. Europe.	—	6. 8. H.○.	
9 stérilis. R.S. bearded.	—	1640.	7. 8. H.○.	Jacq. ic. 1. t. 23.
10 sempervirens. R.S. evergreen.	Pyrenees.	1816.	6. 7. H.‡.	Host. gram. 3. t. 41.
11 fállax. R.S. fallacious.	S. Europe.	1824.	— H.‡.	
12 pállens. R.S. pale.	Portugal.	—	— H.‡.	
13 praténensis. R.S. narrow-leaved.	Britain.	— H.‡.	Eng. bot. v. 17. t. 1204.
14 cantábrica. R.S. evergreen.	—	1825.	— H.‡.	
15 bromoídes. R.S. Bromus-like.	S. Europe.	—	— H.‡.	
16 planicúlmis. R.S. flat-stalked,	Hungary.	1822.	— H.‡.	Schrad.fl. ger. 1.t.6.f.2.
<i>latifolia.</i> Host. gram. 4. t. 32.					
17 versicolor. R.S. various-colored.	Europe.	1816.	6. 8. H.‡.	Host. gram. 2. t. 52.
GAUD'INIA. R.S. GAUD'INIA.	Triandria Digynia. R.S.			
frágilis. R.S. brittle.	S. Europe.	1770.	7. 8. H.○.	Host. gram. 2. t. 54.
<i>Avéna frágilis.</i> Host.					
TRIS'ETUM. R.S.	TRISE'TUM.	Triandria Digynia. R.S.			
1 tenuis. R.S. slender.	Europe.	1804.	7. 8. H.○.	Host. gram. 2. t. 55.
<i>striatum.</i> P.S. <i>Avéna tenuis.</i> Host.					
2 pensylvanicum. R.S. Pennsylvanian.	N. America.	1785.	6. 7. H.○.	
3 micránthum. R.S. small-flowered.	S. Europe.	1823.	— H.○.	
4 negléctum. R.S. neglected.	—	—	— H.○.	Savi fl. pis. 1. t. 1. f. 4.
5 alopecúrus. R.S. Foxtail.	—	1816.	— H.○.	W.berl.mag.2.t.8. f.3.
6 Loeflingiánum. R.S. Loefling's.	S. Europe.	1770.	— H.○.	
7 condensátum. R.S. close-flowered.	Sicily.	1824.	— H.○.	
8 pilósüm. R.S. hairy.	Iberia.	—	— H.○.	
9 pubéscens. R.S. pubescent.	Britain.	—	7. 8. H.‡.	Eng. bot. v. 23. t. 1640.
<i>Avéna pubéscens.</i> E.B.					
10 rígidum. R.S. rigid.	Caucasus.	1825.	— H.‡.	
11 alpinum. R.S. Alpine.	Scotland.	—	7. H.‡.	Eng. bot. v. 30. t. 2141.
<i>Avéna planicúlmis.</i> E.B. non Schrad.					
12 carpáthicum. R.S. Carpathian.	Carp. mount.	1818.	7. 8. H.‡.	Host. gram. 4. t. 31.
13 flavéscens. R.S. golden.	Britain.	—	6. 7. H.‡.	Eng. bot. v. 14. t. 952.
<i>pratense.</i> P.S. <i>Avéna flavéscens.</i> E.B.					
14 alpéstre. R.S. mountain.	S. Europe.	1816.	— H.‡.	Host. gram. 3. t. 39.
15 brevifólium. R.S. short-leaved.	—	—	— H.‡.	— 3. t. 40.
16 argéntium. R.S. silver.	Switzerland.	1791.	— H.‡.	— 2. t. 53.
17 distichophyllum. fan-leaved.	—	1796.	— H.‡.	Vill. delph. 2. t. 4. f. 4.
18 airoides. R.S. Aira-like.	—	1800.	— H.‡.	Flor. dan. t. 228.
<i>Avéna airoides.</i> DC. <i>Aira subspicata.</i> W.					
19 glabrátum. R.S. smooth.	Portugal.	1818.	7. 8. H.○.	
CENTOTH'ECA. R.S.	CENTOTH'ECA.	Triandria Digynia. R.S.			
lappácea. R.S. Bur.	India.	1773.	7. S.○.	Pal.deBeau.ag.t.14.f.7.
<i>Cénchrus lappácea.</i> L.					
CABR'ERA. LG.	CABR'ERA.	Triandria Digynia.			
chrysoblépharis. LG.	golden-spiked.	S. America.	1822.	6. 7. S.‡.	
P'OA. R.S.	MEADOW-GRASS.	Triandria Digynia. L.			
1 praténensis. R.S. smooth-stalked.	Britain.	—	5. 6. H.‡.	Eng. bot. v. 15. t. 1073.
2 angustifólia. DC. narrow-leaved.	—	—	5. H.‡.	Sinc.gra.wob.p.184.ic.
3 humilis. R.S. short-blueish.	—	—	5. 6. H.‡.	Eng. bot. v. 14. t. 1004.
<i>subcarúlea.</i> E.B.					
4 víridis. R.S. green.	N. America.	1812.	— H.‡.	
5 hy'brida. R.S. hybrid.	Switzerland.	1824.	— H.‡.	
6 Torreyána. R.S. Torrey's.	N. America.	1825.	— H.‡.	Eng. bot. v. 15. t. 1072.
7 trivialis. E.B. common.	Britain.	—	6. 8. H.‡.	Eng. bot. v. 16. t. 1141.
8 Gamelinii. R.S. Gamelin's.	Germany.	1825.	— H.‡.	
9 ánnua. E.B. annual.	—	—	1. 12. H.○.	
10 cespítosa. R.S. tufted.	N. Zealand.	1822.	6. 8. H.‡.	Eng. bot. v. 15. t. 1071.
11 grácilis. R.S. slender.	Siberia.	1818.	— H.‡.	Host. gram. 3. t. 15.
12 bulbosa. R.S. bulbous.	England.	—	5. 6. H.‡.	
13 láxá. R.S. loose.	Scotland.	—	7. H.‡.	
<i>flexuosa.</i> Eng. bot. v. 16. t. 1123.					
14 distichophylla. R.S. fan-leaved.	Switzerland.	1829.	— H.‡.	Host. gram. 4. t. 26.

GRAMINEÆ.

4 rígida. L.en. hard.	England.	6. 7. H.○.	Eng. bot. v. 20. t. 1371.
<i>Pòa rígida.</i> E.B. <i>Glycèria rígida.</i> E.F. <i>Megastachya rígida.</i> R.S.	Barbary. 1816.	— H.○.	Desf. atl. 1. t. 22.
5 articulàta. L.en. jointed.	Europe.	— H.○.	Cyrill. neap. fasc. 2. t. 2.
6 marítima. sea-side.			
<i>dichotoma.</i> L.en. <i>Triticum marítimum.</i> L. <i>Festúca marítima.</i> DC.			
HYDROCHLOA. Har. HYDROCHLOA. Triandria Digynia.			
1 distans. Har. distant.	Britain.	7. H.w. 4.	Eng. bot. v. 14. t. 986.
<i>Pòa distans.</i> E.B. <i>Glycèria distans.</i> E.F.			
2 aquática. Har. water.	—	— H.w. 4.	Eng. bot. v. 19. t. 1315.
<i>Pòa aquática.</i> E.B. <i>Glycèria aquática.</i> E.F.			
3 arundinácea. Reed-like.	Caucasus. 1825.	— H.w. 4.	
<i>Pòa arundinacea.</i> M.B.			
4 marítima. Har. sea-side.	Britain.	— H.w. 4.	Eng. bot. v. 16. t. 1140.
<i>Pòa marítima.</i> E.B. <i>Glycèria marítima.</i> E.F.			
GLYCERIA. B.P. SWEET-GRASS. Triandria Digynia. R.S.			
flútans. B.P. floating.	Britain.	5. 8. H. 4.	Eng. bot. v. 22. t. 1520.
<i>Pòa flútans.</i> E.B. <i>Festúca flútans.</i> w.			
ERAGROSTIS. R.S. ERAGROSTIS. Triandria Digynia. R.S.			
1 poæoídes. R.S. Poa-like.	Italy. 1699.	7. H.○.	Flor. græc. t. 73.
<i>Pòa Eragróstis.</i> F.G.			
2 tephrosánthos. R.S. Martinica.	Martinica. 1824.	7. 9. S.○.	
3 pilosa. R.S. hairy.	Italy. 1804.	7. 8. H.○.	Host. gram. 2. t. 68.
4 verticillata. R.S. whorled.	S. Europe. 1816.	— H.○.	Cavan. ic. 1. t. 93.
5 punctata. R.S. spotted.	E. Indies. —	— S.○.	
6 ægyptiaca. R.S. Egyptian.	Egypt. 1812.	— H.○.	Descr. de l'Egypt. t. 10.
7 purpurascens. R.S. purplish.	—	— H.○.	
8 despiciens. R.S. doubtful.	—	— H.○.	
9 tenella. R.S. slender.	E. Indies. 1781.	— S.○.	Rheed. mal. 12. t. 41.
10 cynosuroídes. R.S. Dogstail.	Egypt. 1824.	— H.○.	Descr. de l'Egypt. t. 10.
<i>Pòa cynosuroídes.</i> w.			
11 capillaris. L.en. hair-paniced.	N. America. 1781.	10. 11. H.○.	Moris. s. 3. t. 6. f. 33.
<i>Pòa capillaris.</i> R.S.			
12 mexicana. L.en. Mexican.	Mexico. 1816.	9. 10. H.○.	
13 abyssinica. L.en. Abyssinian.	Abyssinia. 1824.	— H.○.	Jacq. icon. 1. t. 17.
14 plumosa. R.S. feathered.	E. Indies. 1826.	— S.○.	Rheed. mal. 12. t. 41.
MEGASTACHYA. R.S. MEGASTACHYA. Triandria Digynia. R.S.			
1 Eragrostis. R.S. Love-grass.	S. Europe. 1776.	7. 8. H.○.	Schreb. gram. t. 38.
<i>Pòa megastachya.</i> p.s. <i>Briza Eragrostis.</i> w. <i>Eragrostis major.</i> Host. gram. 4. t. 24.			
2 nigricans. R.S. blackish.	S. America. 1823.	— S. 4.	
3 amabilis. R.S. purple.	E. Indies. 1802.	— S.○.	L. pfl. syst. 12. t. 91. f. 2.
<i>Pòa amabilis.</i> L.			
4 spectabilis. R.S. showy.	N. America. 1816.	— H.○.	
5 pulchella. R.S. pretty.	Tauria. 1825.	— H.○.	
6 elongata. R.S. long-paniced.	E. Indies. 1812.	— S. 4.	Jacq. ecl. gr. t. 3.
7 ciliaria. R.S. ciliated.	Jamaica. 1776.	— S.○.	Jacq. ic. 2. t. 304.
8 rupestris. R.S. rock.	E. Indies. 1823.	— S.○.	
9 Boryana. R.S. Bory's.	Mauritius. 1822.	— S.○.	
10 riparia. R.S. crowded.	E. Indies. 1824.	— S.○.	
11 mucronata. R.S. mucronated.	Africa. 1822.	7. 9. S. 4.	Beauv. fl. ow. 1. t. 4.
12 polymorpha. R.S. variable.	Tropics. 1826.	— S. 4.	
DACTYLIS. R.S. COCK'S-FOOT-GRASS. Triandria Digynia. L.			
1 glomerata. R.S. rough.	Britain.	6. 7. H. 4.	Eng. bot. v. 5. t. 335.
2 glaucescens. R.S. glaucous.	Europe. 1816.	— H. 4.	
3 hispánica. R.S. Spanish.	Spain. 1814.	— H. 4.	
4 glauca. R.S. glaucous.	—	— H. 4.	
5 capitata. R.S. headed.	Sardinia. 1824.	— H. 4.	
6 altaica. R.S. Altaic.	Altay. 1826.	— H. 4.	
7 marítima. R.S. sea-side.	Caucasus. 1823.	— H. 4.	
8 littoralis. R.S. shore.	S. Europe. 1816.	— H. 4.	
9 repens. R.S. creeping.	—	— H. 4.	Lam. ill. t. 45. f. 5.
10 adscendens. R.S. ascending.	C. B. S. 1823.	— G. 4.	Desf. atl. 1. t. 15.
11 lagopodioídes. R.S. harefoot.	E. Indies. 1826.	— S. 4.	Burm. ind. t. 12. f. 2.
12 brevifolia. R.S. short-leaved.	—	— S. 4.	
13 cynosuroídes. R.S. Dog's-tailed.	—	— S. 4.	
14 aristata. E. awned.	Egypt. 1828.	— H.○.	
BRIZA. R.S. QUAKING-GRASS. Triandria Digynia. R.S.			
1 minor. R.S. small.	England.	7. H. 4.	Eng. bot. v. 19. t. 1316.
2 vires. R.S. green.	S. Europe. 1800.	7. 8. H.○.	Hayn. term. bot. t. 25. f. 6
3 media. E.B. common.	Britain.	5. 6. H. 4.	Eng. bot. v. 5. t. 340.

4 Clùssi. R.S. Clusius's.	S. Europe.	1823.	5. 6. H. ¼.	Fouc. j. bot. 3. t. 24. f. 2.
5 geniculàta. R.S. jointed.	C. B. S.	1816.	7. 8. H. ¼.	Flor. græc. 1. t. 75.
6 elàtior. R.S. tall.	Greece.	1788.	— H. ¼.	— 1. t. 76.
7 màxima. R.S. great-spiked.	S. Europe.	1633.	6. 7. H. ¼.	Scheuch. agr. 4. t. 4. f. 10.
8 hùmilius. R.S. dwarf.	Tauria.	1818.	— H. ¼.	
9 rùbra. R.S. red.	S. Europe.	—	— H. ¼.	
CYNOS'URUS. R.S.	DOGSTAIL-GRASS. Triandria Digynia. L.	Britain.	8. H. ¼.	Eng. bot. v. 5. t. 316.
cristatus. E.B. crested.				
UNIOLA. R.S.	SEASIDE OAT. Triandria Digynia. L.	N. America.	1793.	6. 7. H. ¼.	Catesb. car. 1. t. 32.
1 paniculàta. R.S. paniced.	—	1809.	— H. ¼.	
2 latifòlia. R.S. broad-leaved.	—	1790.	7. H. ¼.	
3 spicata. R.S. spiked.	—	1789.	6. 7. H. ¼.	
4 distichophylla. R.S. two-ranked.	—			
Festuca distichophylla. Ph.					
CERATOCHL'OA. R.S.	CERATOCHL'OA. Triandria Digynia. R.S.				
unioloides. R.S. Uniola-like.	N. America.	1796.	6. 7. H. ¼.	Willd. h. ber. t. 3.
Festuca unioloides. w.					
GRAPHEPHORUM. R.S.	GRAPHEPHORUM. Triandria Digynia.				
melicoides. R.S. Melic-like.	N. America.	1824.	6. 8. H. ¼.	
A'ira melicoides. M.					
LEPTOCHL'OA. R.S.	LEPTOCHL'OA. Triandria Digynia. R.S.				
1 cynosuroides. R.S. filiform.	E. Indies.	1818.	8. 9. H. ¼.	
2 pròcera. R.S. tall.	Brazil.	1823.	— S. ¼.	Jacq. ecl. gram. t. 4.
3 filiformis. R.S. slender.	S. America.	1818.	— H. ¼.	Sloan. jam. 1. t. 70. f. 2.
4 virgata. R.S. fine-spiked.	—	1727.	7. 9. H. ¼.	
5 tenerrima. R.S. slenderest.	China.	1820.	— H. ¼.	
6 domingensis. L.en. close-spiked.	W. Indies.	—	6. 8. S. ¼.	
ELEUS'INE. R.S.	ELEUS'INE. Triandria Digynia. R.S.				
1 coracàna. R.S. thick-spiked.	E. Indies.	1714.	7. 9. H. ¼.	Schreb. gram. 2. t. 35.
2 strícta. R.S. upright.	—	1822.	— H. ¼.	Rheed. mal. 12. t. 69.
3 índica. R.S. Indian.	India.	1714.	— H. ¼.	
4 rígida. R.S. rigid.	Monte Video.	1826.	— H. ¼.	
5 verticillata. R.S. whorled.	E. Indies.	1823.	— H. ¼.	
6 calycina. R.S. calycine.	—	—	— H. ¼.	
7 cereális. E. Rye-like.	Egypt.	1827.	— H. ¼.	
8 ovális. E. oval-spiked.	—	—	— H. ¼.	
9 tristächya. E. three-spiked.	—	—	— H. ¼.	
DACTYLOCTENIUM. DACTYLOCTENIUM. Triandria Digynia. R.S.					
ægyptiacum. R.S. Egyptian.	Egypt.	1770.	7. 9. H. ¼.	Moris. s. 8. t. 3. f. 7.
Cynosurus ægyptius. L.	Eleusine ægyptiaca. P.S.				
RHABDOCHL'OA. R.S.	RHABDOCHL'OA. Triandria Digynia. R.S.				
1 cruciàta. R.S. cross-spiked.	W. Indies.	1818.	7. 9. H. ¼.	Sloan. his. 1. t. 69. f. 1.
Chlòris cruciàta. w.	Agróstis cruciàta. L.				
2 mucronata. R.S. mucronate.	N. America.	1820.	— H. ¼.	Lam. ill. t. 48. f. 2.
3 virgata. R.S. slender.	W. Indies.	—	— H. ¼.	Jacq. icon. 1. t. 22.
4 domingensis. R.S. West Indian.	—	1820.	— S. ¼.	Jacq. ecl. gr. t. 12.
CHOND'R'OSIUM. K.S.	CHOND'R'OSIUM. Triandria Digynia. R.S.				H. et B. n. gen. 1. t. 57.
1 procumbens. Desv. procumbent.	Philippines.	1816.	7. 9. H. ¼.	
Antinochlòa procumbens. R.S.	Atheropògon procumbens. Jacq.				
2 ténue. K.S. slender.	Mexico.	1823.	— H. ¼.	
3 prostratum. prostrate.	—	—	— H. ¼.	
Antinochlòa prostrata. R.S.	Bouteloua prostrata. Lag.				
4 hirsutum. hairy.	—	—	— H. ¼.	
CORYCA'R'PUS. Zea.	CORYCA'R'PUS. Diandria Digynia. S.S.				
arundinaceus. s.s. Reed-like.	N. America.	1810.	6. 7. H. ¼.	Mich. amer. 1. t. 10.
Festuca diándra. M.	Diarrhèa americana. R.S.				
DIPLO'CHNE. R.S.	DIPLO'CHNE. Triandria Digynia. R.S.				
fasciculàris. R.S. fascicled.	N. America.	1818.	6. 8. H. ¼.	
Festuca polystachya. M.					
SCHEDON'ORUS. R.S.	SCHEDON'ORUS. Triandria Digynia. R.S.				
1 pratensis. R.S. meadow.	Britain.	6. 7. H. ¼.	Eng. bot. v. 23. t. 1592.
Festuca pratensis. E.B.					
2 phœnicoides. R.S. creeping-rooted.	Switzerland.	1824.	— H. ¼.	Eng. bot. v. 23. t. 1593.
3 elàtior. R.S. tall.	Britain.	— H. ¼.	
Festuca elàtior. E.B. et Sinc. gr. wob. p. 361. ic.					
4 cerulescens. R.S. blueish.	Barbary.	1818.	6. 7. H. ¼.	Host. gram. 3. t. 20.
5 spadiceus. R.S. Gerard's.	Italy.	1775.	4. 6. H. ¼.	— 1. t. 9.
6 inérmis. R.S. naked-spiked.	Hungary.	1816.	6. 7. H. ¼.	Eng. bot. v. 14. t. 1005.
7 calamarius. R.S. Reed-like.	Scotland.	7. 8. H. ¼.	

GRAMINEÆ.

8 decíduus.	deciduous.	England.	6. 7. H. 4.	Eng. bot. v. 32. t. 2266.
<i>Festúca decídua.</i> E.B.				
9 exaltátus. R.S.	high upright.	Sicily. 1827.	— H. 4.	
10 sylváticus. R.S.	wood.	Germany. 1804.	7. 8. H. 4.	Host. gram. 2. t. 78.
11 serótinus. R.S.	late-flowering.	S. Europe. 1818.	7. 9. H. 4.	— 2. t. 92.
12 loliáceus. R.S.	spiked.	England.	6. 7. H. 4.	Eng. bot. v. 26. t. 1821.
13 Scheuchzéri. R.S.	Scheuchzer's.	S. Europe. 1822.	— H. 4.	
14 violáceus. R.S.	violet coloured.	Europe.	— H. 4.	Host. gram. 4. t. 61.
15 nútans. R.S.	nodding.	Hungary. 1823.	— H. 4.	
16 nigréscens. R.S.	blackish.	Switzerland.	— H. 4.	Host. gram. 2. t. 81.
17 poëfórmis. R.S.	Poa-like.	— 1818.	— H. 4.	— 2. t. 91.
18 pùmilus. R.S.	dwarf.	S. Europe.	— H. 4.	
19 nítidus. R.S.	glossy.	Hungary. 1827.	— H. 4.	
20 tenélus. R.S.	slender.	N. America. 1804.	7. 8. H. ⊙.	
<i>Festúca tenélla.</i> Ph. <i>Brachypodium festucoides.</i> Len.				
FEST'UCA. R.S.	FESCUE-GRASS.	Triandria Digynia. L.		
1 cynosuroídes. R.S.	dogs-tail.	Barbary. 1825.	6. 8. H. ⊙.	Desf. atl. 1. t. 21.
2 Fénas. R.S.	tough.	Spain. 1822.	— H. 4.	
<i>Pòtēnax.</i> L.en.				
3 ovina. R.S.	Sheep's.	Britain.	6. H. 4.	Eng. bot. v. 9. t. 585.
4 brachyphyllea. R.S.	short-leaved.	Melville Isl. 1827.	— H. 4.	
<i>brevifolia.</i> Br. non Muhlenberg.				
5 tenuifólia. L.en.	slender-leaved.	Britain.	— H. 4.	
6 cæsia. E.B.	gray.	England.	6. 7. H. 4.	Eng. bot. v. 27. t. 1917.
7 vivipara. E.B.	viviparous.	Britain.	7. H. 4.	— v. 19. t. 1355.
8 glauca. R.S.	glaucous.	S. Europe.	6. 7. H. 4.	
9 pálens. R.S.	pale.	Germany. 1816.	— H. 4.	Host. gram. 2. t. 88.
10 strícta. Host.	upright.	—	— H. 4.	— 2. t. 86.
11 hirsuta. Host.	hairy.	—	— H. 4.	— 2. t. 85.
12 Hallér. R.S.	Haller's.	Europe. 1820.	— H. 4.	
13 cùrvula. R.S.	curved.	Switzerland.	— H. 4.	
14 alpina. R.S.	Alpine.	—	— H. 4.	
15 amethy'stina. R.S.	blue.	S. Europe. 1804.	— H. 4.	
16 picta. R.S.	painted.	Hungary. 1826.	— H. 4.	
17 váría. R.S.	variable.	S. Europe. 1823.	— H. 4.	Host. gram. 2. t. 90.
18 vallésiaca.	Vallesian.	Vallesia.	— H. 4.	
19 pannónica. R.S.	Hungarian.	Hungary.	— H. 4.	Host. gram. 4. t. 62.
20 xanthina. R.S.	yellow.	Switzerland. 1818.	— H. 4.	— 3. t. 19.
<i>flavescens.</i> Host. non Bellardi.				
21 flavescens. DC.	yellowish.	—	— H. 4.	
22 duriúscula. E.B.	hard.	Britain.	— H. 4.	Eng. bot. v. 7. t. 470.
23 dumetórum. S.G.W.	pubescent.	—	— H. 4.	
24 glábra. S.G.W.	smooth.	—	6. H. 4.	Sinc. gr. wob. p. 180. ic.
25 hordeiformis. S.G.W.	Barley-like.	—	— H. 4.	— p. 159. ic.
26 cámbrica. S.G.W.	Welsh.	Wales.	— H. 4.	— p. 157. ic.
27 heterophylla. R.S.	various-leaved.	S. Europe. 1812.	6. 7. H. 4.	Host. gram. 3. t. 18.
28 rùbra. E.B.	red creeping.	Britain.	7. H. 4.	Eng. bot. v. 29. t. 2056.
29 vagináta. R.S.	sheathed.	Hungary. 1804.	6. 7. H. 4.	
30 Kitabeliána. R.S.	Kitaibel's.	— 1818.	— H. 4.	
<i>pubescens.</i> L.en. non Zea.				
31 láxa. R.S.	loose.	Switzerland. 1824.	— H. 4.	Host. gram. 2. t. 80.
32 decolorans. R.S.	discoloured.	—	— H. 4.	
<i>láxa.</i> Gaud. nec aliorum.				
33 nútans. R.S.	nodding.	N. America. 1805.	— H. 4.	
34 rùbens. R.S.	red.	S. Europe. 1776.	— H. ⊙.	Flor. græc. t. 83.
<i>Bròmus rùbens.</i> F.G.				
35 scábra. R.S.	rough.	C. B. S. 1826.	— G. 4.	
36 bracteàta. DC.	leafy-bracted.	—	— H. 4.	
37 l'ævis. f.	smooth.	Russia. 1806.	7. H. 4.	
38 grandiflora. Ph.	large-flowered.	N. America. 1812.	6. 7. H. 4.	
39 aspérrima. Ot.	roughest.	—	1825.	
40 rupéstris. Ot.	rock.	—	— H. 4.	
MYGALURUS. L.en. MOUSTAIL-GRASS.	Triandria Digynia. L.en.			
1 bromoídes. L.en.	barren.	Britain.	6. H. ⊙.	Eng. bot. v. 20. t. 1411.
2 caudátus. L.en.	wall.	England.	— H. ⊙.	— v. 20. t. 1412.
<i>Festúca Myurus.</i> E.B.				
3 alopecuroídes. L.en.	Fox-tail.	Barbary. 1799.	6. 8. H. ⊙.	Desf. atl. 1. t. 25.
4 ciliata. Lk.	fringed.	Portugal. 1827.	— H. ⊙.	Host. gram. 4. t. 65.
5 uniglumis. L.en.	single-husked.	England.	— H. ⊙.	Eng. bot. v. 20. t. 1430.

6 deliciatulus, L.en.	delicate.	Spain.	1817.	7. 8. H.○.	
7 geniculatus, L.en.	jointed.	S. Europe.	1793.	6. 7. H.○.	
<i>Bròmus geniculatus</i> , R.S. <i>Festuca stipoides</i> , R.S.					
BRACHYP'ODIUM , R.S. BRACHYP'ODIUM , Triandria Digynia, R.S.					
1 pinnatum, R.S.	spiked heath.	Britain.	6. 8. H.○.	
<i>Bròmus pinnatus</i> , E.B. <i>Triticum pinnatum</i> , DC. <i>Festuca pinnata</i> , E.F.					Eng. bot. v. 11. t. 730.
2 rupéstre, R.S.	rock.	S. Europe.	1816.	— H.○.	Host. gram. 4. t. 17.
3 strigosum, R.S.	strigose.	Tauria.	1824.	— H.○.	Host. gram. 4. t. 18.
4 cæpitosum, R.S.	tufted.	S. Europe.	—	— H.○.	Flor. græc. t. 84.
5 ramosum, R.S.	branched.	—	1818.	— H.○.	
6 phenicoïdes, R.S.	pointed-leaved.	S. France.	1824.	— H.○.	
7 obtusifolium, R.S.	blunt-leaved.	Spain.	1818.	— H.○.	
8 Tenoriánum, R.S.	Tenore's.	Naples.	1824.	— H.○.	
9 Barrelieri, R.S.	Barrelier's.	—	1827.	— H.○.	
10 longifolium, R.S.	long-leaved.	Barbary.	1818.	— H.○.	
11 sylvaticum, R.S.	wood.	Britain.	— H.○.	Eng. bot. v. 11. t. 729.
<i>Bròmus sylvaticus</i> , E.B. <i>Triticum sylvaticum</i> , DC. <i>Festuca sylvatica</i> , E.F.					
12 distachyum, R.S.	two-spiked.	S. Europe.	1772.	— H.○.	Host. gram. 1. t. 20.
13 megastachyum, R.S.	long-spiked.	—	1826.	— H.○.	
14 macrostachyum, R.S.	large-spiked.	—	—	— H.○.	
15 ásperum, R.S.	stiff-leaved.	Spain.	1818.	— H.○.	Jacq. icon. 2. t. 303.
16 unioloides, L.en.	Uniola-like.	Sicily.	1758.	7. 8. H.○.	
17 brevisétum, R.S.	short-awned.	—	1825.	— H.○.	Vivian. fr. fl. it. t. 26. f. 1.
18 frágile, R.S.	brittle.	Europe.	1824.	— H.○.	— t. 23. f. 2.
19 Halléri, R.S.	Haller's.	S. Europe.	1818.	— H.○.	
20 hispánicum, Viv.	slender-stem'd.	—	—	— H.○.	
<i>tenuículum</i> , R.S.					
21 tenuifórum, R.S.	slender-flowered.	—	1823.	— H.○.	Host. gram. 2. t. 26.
22 Nárdus, L.en.	slender.	—	—	— H.○.	
<i>tenellum</i> , R.S. <i>Triticum Nárdus</i> , DC.					
23 loliaceum, R.S.	spiked sea.	Britain.	6. 7. H.○.	Eng. bot. v. 4. t. 221.
<i>Triticum loliaceum</i> , E.B.					
24 Pòa, R.S.	meadow-grass.	France.	1824.	— H.○.	Pluk. phyt. t. 32. f. 7.
25 biunciale, R.S.	small.	S. Europe.	1818.	6. 8. H.○.	Viv. fl. it. fr. t. 24.
26 unilateralè, R.S.	one-sided.	—	1824.	— H.○.	Bocc. mus. t. 57. f. 2.
27 mexicànum, Bes.	Mexican.	Mexico.	1816.	7. 9. H.○.	
<i>Festuca mexicana</i> , R.S.					
TRICH'ÆTA , R.S. TRICH'ÆTA , Triandria Digynia, R.S.					
ovata, R.S.	oval-spiked.	Spain.	1824.	6. 8. H.○.	Cavan. ic. 6. t. 591. f. 2.
<i>Bròmus ovatus</i> , Cav. <i>Trisetum ovatum</i> , P.S.					
BR'OMUS , R.S. BROME-GRASS , Triandria Digynia, R.S.					
1 secalinus, R.S.	smooth Rye.	England.	6. 8. H.○.	Eng. bot. v. 17. t. 1171.
2 elongátus, R.S.	elongated.	Switzerland.	1823.	— H.○.	Host. gram. 1. t. 11.
3 commutátus, R.S.	downy Rye.	Britain.	— H.○.	Eng. bot. v. 27. t. 1884.
4 velutinus, R.S.	hairy-headed.	—	— H.○.	
5 Triniánus, R.S.	Trinius's.	Gilan.	1826.	— H.○.	Trin. m.ac.pet. 1813. t. 9.
6 mollis, R.S.	soft.	Britain.	5. 7. H.○.	Eng. bot. v. 15. t. 1078.
7 confertus, R.S.	close-spiked.	Iberia.	1825.	— H.○.	Jacq. ecl. gr. t. 14.
8 racemosus, R.S.	smooth.	England.	6. 8. H.○.	Eng. bot. v. 15. t. 1079.
9 erectus, E.B.	upright.	—	— H.○.	— v. 7. t. 471.
10 Biebersteinii, R.S.	Bieberstein's.	Caucasus.	1825.	— H.○.	
11 australis, B.P.	southern.	N. S. W.	1823.	— H.○.	
12 lanceolátus, R.S.	spear-leaved.	Canaries.	1798.	— H.○.	Buxb. cent. ap. t. 19.
13 lanuginosus, R.S.	woolly.	S. Europe.	1824.	— H.○.	
14 variegátus, R.S.	variegated.	Caucasus.	—	— H.○.	Eng. bot. v. 27. t. 1885.
15 squarrosus, R.S.	corn.	England.	— H.○.	
16 vestitus, R.S.	clothed.	C. B. S.	1816.	— G.○.	Jacq. ecl. gr. t. 15.
17 wolgensis, R.S.	Wolga.	Caucasus.	—	— H.○.	
18 laxus, R.S.	loose.	—	—	— H.○.	Desf. atl. 1. t. 25.
19 alopecuroïdes, R.S.	Fox-tail.	S. Europe.	1799.	— H.○.	
<i>contortus</i> , Desf.					
20 pùrgans, R.S.	purging.	Canada.	1793.	— H.○.	
21 pubescens, R.S.	pubescent.	N. America.	1816.	— H.○.	Eng. bot. v. 17. t. 1172.
22 ásper, E.B.	hairy wood.	England.	— H.○.	
23 hirsutissimus, Cyr.	very hairy.	S. Europe.	1822.	— H.○.	
24 pendulinus, R.S.	pendulous.	New Spain.	1823.	— H.○.	
25 glaucus, R.S.	glaucous.	Pyrenees.	—	— H.○.	Lap. fl. pyren. t. 53. bis.
26 altissimus, R.S.	tallest.	N. America.	1812.	— H.○.	

27	giganteus. R.S.	gigantic.	Britain.	7. 8. H. 4.	Eng. bot. v. 26. t. 1820.
28	triflorus.	three-flowered.	—	—	H. 4.	v. 27. t. 1918.
	<i>Festuca triflora</i> . E.B.						
29	montanus. R.S.	mountain.	Switzerland.	1827.	H. ⓠ.	
30	longiflorus. R.S.	long-flowered.	—	1818.	H. 4.	
31	stenophylloides. R.S.	slender-leaved.	—	1824.	H. 4.	
32	angustifolius. M.B.	narrow-leaved.	Caucasus.	—	H. 4.	
33	ciliatus. R.S.	ciliated.	N. America.	1802.	6. 8. H. 4.	Eng. bot. v. 28. t. 1984.
34	arvensis. R.S.	field.	Britain.	—	7. H. 4.	— v. 15. t. 1030.
35	sterilis. R.S.	barren.	—	—	6. 7. H. ⓠ.	Flor. græc. t. 82.
36	tectorum. R.S.	nodding.	Europe.	1776.	6. 8. H. ♂.	Eng. bot. v. 14. t. 1006.
37	matritensis. R.S.	wall.	Britain.	—	6. H. ⓠ.	
	<i>diandrus</i> . E.B.						
38	ligusticus. R.S.	contracted.	Italy.	1824.	6. 8. H. ⓠ.	Barrel. ic. 76. n. 2.
39	rigidus. R.S.	rigid.	S. France.	1822.	H. ⓠ.	Host. gram. 1. t. 18.
40	pilosus. R.S.	hairy wall.	S. Europe.	—	H. ⓠ.	
41	jubatus. R.S.	loose paniced.	Italy.	1824.	H. ⓠ.	
42	máximus. R.S.	great.	Morocco.	1804.	H. ⓠ.	Desf. atl. 1. t. 26.
43	scoparius. R.S.	broom.	S. Europe.	1824.	H. ⓠ.	Cavan. ic. 1. t. 45. f. 2.
44	humilis. R.S.	low.	Spain.	—	H. ⓠ.	— ic. 6. t. 589. f. 2.
45	scaberrimus. R.S.	roughest.	Italy.	—	H. ⓠ.	
46	coarctatus. H.H.	close-flowered.	—	1826.	H. 4.	
47	parviflorus. D.F.	small-flowered.	—	1824.	H. 4.	
48	tortilis. Presl.	twisted.	Sicily.	1825.	H. 4.	
49	caucasicus. F.	Caucasean.	Caucasus.	—	H. 4.	
ROSTR'ARIA. Tr.	ROSTR'ARIA.	Triandria Digynia.					
pubescens. Tr.	pubescens.	—	1824.	6. 8. H. ⓠ.			
<i>Bròmus dactyloides</i> . Roth.	<i>Dáctylis pungens</i> . Horn.						
PHRAGMITES. Tr.	REED.	Triandria Digynia. R.S.					
communis. Tr.	common.	Britain.	...	7. 8. H.w. 4.	Eng. bot. v. 6. t. 401.		
Árundo phragmites. E.B.							
ARUNDO. S.S.	REED-GRASS.	Triandria Digynia. S.S.					
1 Dómax. s.s. manured.	S. Europe.	1648.	7. 8. H. 4.			
β versicolor. striped.	—	—	H. 4.	Host. gram. 4. t. 38.		
2 ténax. s.s. tough.	—	1823.	H. 4.	Moris. s. 8. t. 8. f. 9.		
3 festucacea. W.en.	Fescue-like.	Germany.	—	H.w. 4.	Cyril. neap. f. 2. t. 12.		
PSAMMA. R.S.	SEA MATWEED.	Triandria Trigynia. R.S.					
arenaria. R.S. common.	Britain.	—	6. 7. H. 4.	Eng. bot. v. 8. t. 520.		
Árundo arenaria. E.B.							
ARUNDIN'ARIA. R.S.	CANE-BRAKE.	Triandria Monogynia. S.S.					
1 glaucescens. R.S. glaucous.	E. Indies.	1818. S. h.			
<i>Ludolfia glaucescens</i> . w.							
OL'YRA. W.	OL'YRA.	Monococcia Triandria. L.					
paniculata. w. broad-leaved.	W. Indies.	1783.	7. S. 4.			
Z'EA. W.	INDIAN CORN.	Monococcia Triandria. L.					
1 Máys. w. common.	America.	1562.	6. 7. H. ⓠ.	Sloan. jam. 1. t. 64. f. 2.		
2 Curáguia. s.s. saw-leaved.	Chile.	1820.	H. ⓠ.	Lam. ill. t. 749.		
COTIX. W.	JOB'S-TEARS.	Monococcia Triandria. L.					
1 Láchryma. s.s. common.	E. Indies.	1596.	6. 7. S. 4.			
2 agrestis. s.s. round-fruited.	—	1812.	S. 4.	Bot. mag. t. 2479.		
3 exaltata. s.s. tall.	China.	1816.	S. 4.	Rumph. amb. 6. t. 9. f. I.		
4 arundinacea. s.s. reed-like.	Mexico.	1824.	S. 4.	Jacq. ic. gram.		
5 K'oenigii. s.s. Koenig's.	E. Indies.	1818.	S. 4.			
6 gigantea. H.B. gigantic.	Bengal.	1820.	H. 4.			
BAMB'USA. W.	BAMBOO-CANE.	Hexandria Monogynia. W.					
1 arundinacea. w. common.	India.	1730. S. h.	Roxb. corom. 1. t. 79.		
2 verticillata. w. whorl-flowered.	—	1802. S. h.	— 1. t. 80.		
MELOCA'NNA. Tr.	MELOCA'NNA.	Hexandria Monogynia. S.S.					
bambusoïdes. Tr. berry bearing.	E. Indies.	1818. S. h.			
<i>Bambusa baccifera</i> . Roxb.							
PH'ARUS. W.	PH'ARUS.	Monococcia Hexandria. W.					
1 latifolius. w. broad-leaved.	Jamaica.	1793.	7. 8. S. 4.	Brown. jam. t. 38. f. 3.		
2 angustifolius. D.L. narrow-leaved.	Trinidad.	1821.	S. 4.			
EHRHA'RTA. W.	EHRHA'RTA.	Hexandria Monogynia. S.S.					
1 panicea. w.	Panic-grass.	C. B. S.	1790.	5. 7. G. 4.	Sm. ined. 1. t. 9.		
2 melicoides. s.s.	Melica-like.	—	1816.	G. 4.			
MICROL'ÆNA. B.P.	MICROL'ÆNA.	Tetrandria Digynia. B.P.					
stipoïdes. B.P. Stipa-like.	N. S. W.	1822.	7. 8. F. 4.	Lab. n. hol. 1. t. 118.		
<i>Ehrharta stipoïdes</i> . Lab.							

POTAMOPHILA. B.P.	POTAMOPHILA.	Polygamia Monœcia. B.P.
parviflora. B.P. small-flowered.	N. S. W. 1820. 6. 7. F.w. 4.
ORYZA. W.	RICE.	Hexandria Digynia. L.
sativa. w. common.	Ethiopia. 1596. 7. S.w. ⊙.
ZIZANIA. W.	CANADA RICE.	Monœcia Hexandria. W.
1 aquatica. H.K. water.	N. America. 1790. 7. 9. H.w. 4. Linn. trans. 7. t. 13.
2 miliacea. M. Millet-like.	Carolina. 1816. — H.w. 4.
3 fluítans. M. floating.	N. America. 1824. 7. 8. H.w. 4.
PARIGANA. W.	PARIGANA.	Monœcia Polyandria. W.
campéstris. w.	Cayenne.	1803. S. I. Aubl. gui. 2. t. 337.

** SUBCLASSIS II. CRYPTOGAMEÆ. DC.

ORDO CCXXV.

CHARACEÆ. Kth. synops. v. 1. p. 102.

CH'ARA. L.	CH'ARA.	Monœcia Monandria. W.				
1 vulgáris. E.B. common.	Britain.	7. 8.	H.w. ⊙.	Eng. bot. v. 5. t. 336.	
2 hispida. E.B. prickly.	—	6. 9.	H.w. ⊙.	—	v. 7. t. 463.
NITELLA. Ag. NITELLA.	Monœcia Monandria.				
1 tomentosa. w. tomentose.	6. 7.	H.w. ⊙.	Moris. s. 15. t. 4. f. 9.	
2 fléxilis. Ag. smooth.	Britain.	7. 8.	H.w. ⊙.	—	
3 opáca. Ag. opaque.	—	—	H.w. ⊙.	Eng. bot. v. 15. t. 1070.	
Chàra fléxilis. E.B.						
4 translúcens. Ag. great transparent.	—	—	H.w. ⊙.	Eng. bot. v. 26. t. 1855.	
5 nidifica. Ag. proliferous.	—	—	H.w. ⊙.	—	v. 24. t. 1703.
6 grácilis. Ag. slender.	—	—	H.w. ⊙.	—	v. 30. t. 2140.

ORDO CCXXVI.

EQUISETACEÆ. DC. fl. fr. ed. 3. v. 2. p. 28.

EQUIS'ETUM. W.	HORSE-TAIL.	Cryptogamia Gonopterides. W.				
1 arvénse. w. (br.ye.) corn.	Britain.	3. 4. H. 4.	Eng. bot. v. 29. t. 2020.			
2 fluviátile. w. (br.ye.) great water.	—	4. 5. H. 4.	—	v. 29. t. 2022.		
3 umbrosum. W.en. (br.ye.) shade.	Germany. 1818.	— H. 4.				
4 sylváticum. w. (br.ye.) wood.	Britain.	— H. 4.	Eng. bot. v. 27. t. 1874.			
5 limosum. w. (br.ye.) smooth naked.	—	6. 7. H. 4.	—	v. 13. t. 929.		
6 palustre. w. (br.ye.) marsh.	—	— H. 4.	—	v. 29. t. 2021.		
7 praténse. w. (br.ye.) meadow.	Germany. 1820.	— H. 4.				
8 scirpóides. w. (br.ye.) American.	N. America. 1812.	— H. 4.				
9 hyémale. w. (br.ye.) Shave grass.	Britain.	7. 8. H. 4.	Eng. bot. v. 13. t. 915.			
10 variegátum. w. (br.ye.) variegated.	Scotland.	6. 7. H. 4.	—	v. 28. t. 1987.		
11 filiforme. (br.ye.) thready.	N. America. 1800.	7. 8. H. 4.				

ORDO CCXXVII.

MARSILEACEÆ. Brown prodr. p. 166.

PILUL'ARIA. W.	PILLWORT.	Cryptogamia Hydropterides. W.				
globulifera. w. (br.)	Pepper-grass.	Britain.	6. 9. H.w. 4.	Eng. bot. v. 8. t. 521.		
ISO'ETES. W.	QUILLWORT.	Cryptogamia Hydropterides. W.				
lacústris. w. marsh.	Britain.	5. 10. H.w. 4.	Eng. bot. v. 16. t. 1084.		

MARSILEACEÆ.

SALVÍNIA. W. nātans. w.	SALVÍNIA. Cryptogamia Hydropterides. W. floating. Europe. 1818. 5. 8. H.w. 4. Schkuhr filic. t. 173.
MARSI'LEA. W. quadrifolia. w.	MARSI'LEA. Cryptogamia Hydropterides. W. four-leaved. S. Europe. 1820. 6. 8. H.w. 4. J.mem.ac.sci.par.1740.

ORDO CCXXVIII.

LYCOPODIACEÆ. *Kth. synops. 1. p. 95.*

LYCOP'ODIUM. W. CLUB-MOSS.	Cryptogamia Stachyopterides. W.
1 caroliniánum. w. (br.) Carolina.	Carolina. 1812. 7. 8. H. 4.
2 clavátum. w. (br.) common.	Britain. — H. 4.
3 complanátum. w. (br.) ArborVitæ-leav'd.N.America.	1770. — H. 4.
4 alpinum. w. (br.) Savin-leaved.	Britain. 8. H. 4.
5 dendróideum. w. (br.) fan.	N. America. 1770. 7. H. 4.
6 annotinum. w. (br.) interrupted.	Britain. 6. 8. H. 4.
7 inundátum. w. (br.) marsh. — H.w. 4.
8 alopecuroídes. w. (br.) Fox-tail.	N. America. 1816. — H. 4.
9 Selaginoídes. w. (br.) prickly.	Britain. 8. H. 4.
10 ornithopodioídes. w. (br.) Bird's-claw. 1812. 6. 8. H. 4.
11 helvéticum. w. (br.) Swiss.	Switzerland. 1779. — H. 4.
12 rupéstre. w. (br.) rock.	N. America. 1818. — H. 4.
13 denticulátum. w. (br.) toothed.	Switzerland. 1779. 7. H. 4.
14 depréssum. w. (br.) depressed.	C. B. S. 1816. 7. 8. G. 4.
15 Selágo. w. (br.) Fir.	Britain. 6. 8. H. 4.
16 apódum. w. (br.) distichous.	N. America. 1815. 7. H. 4.
17 lucidulum. w. (br.) glossy. 1822. — H. 4.
18 dénum. B.P. (br.) dense.	N. S. W. — 7. 8. G. 4.
PSIL'OTUM. B.P. PSIL'OTUM. Cryptogamia Stachyopterides. W.	
triquétrum. B.P. (ye.) triangular.	N. S. W. 1793. 7. 8. G. 4.
<i>Lycopodium nūdum.</i> L. <i>Bernhárdia dichótoma.</i> w.	

Dill. musc. t. 62. f. 5.
Eng. bot. v. 4. t. 224.
Flor. dan. t. 78.
Eng. bot. v. 4. t. 234.
Hook. ex. flor. t. 7.
Eng. bot. v. 24. t. 1727.
—— v. 4. t. 239.
Schkuhr filic. t. 160.
Eng. bot. v. 16. t. 1148.
Dill. musc. t. 66. f. 1.B.
—— t. 64. f. 2.
—— t. 63. f. 11.
—— t. 66. f. 1. A.
Eng. bot. v. 4. t. 233.
Dill. musc. t. 64. f. 3.
Schkuhr filic. t. 159.
Lab.nov.hol.2.t.251.f.1
Schkuhr filic. t. 165. b.

ORDO CCXXIX.

FILICES. *Juss. gen. 14.*SUBORDO I. *OPHIOGLOSSÆ.* Brown prodr. p. 163.

OPHIOGLO'SSUM. W. ADDER'S-TONGUE.	Cryptogamia Stachyopterides. W.
1 vulgátum. E.B. (br.) common.	Britain. 5. 6. H. 4.
2 bulbósum. w. (br.) bulbous-rooted.	N. America. — H. 4.
3 lusítanicum. w. (br.) Portugal.	Portugal. — H. 4.
4 gramíneum. w. (br.) grass-like.	N. Holland. 1822. — G. 4.
5 costátum. B.P. (br.) ribbed.	N. S. W. — — G. 4.
6 reticulátum. w. (br.) netted.	W. Indies. 1793. — S. 4.
7 petiolátum. w. (br.) petiolate. 1822. 1. 3. S. 4.
BOTRY'CHIUM. W. MOONWORT.	Cryptogamia Stachyopterides. W.
1 Lunària. w. (br.ye.) common.	Britain. 5. 6. H. 4.
<i>Osmunda lunaria.</i> E.B.	
2 fumarioídes.w. (br.ye.) Fumitory-leaved.	Carolina. 1806. 7. 8. H. 4.
3 oblíquum. w. (br.ye.) oblique.	N. America. 1821. 6. 7. H. 4.
4 disséctum. w. (br.ye.) cut-leaved. 1806. 7. H. 4.
5 virginicum. w. (br.ye.) Virginian. 1790. 8. H. 4.
6 austrále. B.P. (br.ye.) southern.	N. S. W. 1823. 7. 8. H. 4.
7 gráciele. Ph. (br.ye.) slender.	N. America. 1824. — H. 4.

Schkuhr fil. t. 157.
Pluk. mant. t. 427. f. 7.
Schkuhr fil. t. 158.
—— t. 156.

SUBORDO II. OSMUNDACEÆ. Brown prodr. p. 161.

LYG'ODIUM. B.P.	SNAKE'S-TONGUE.	Cryptogamia Schismatopterides. W.
1 polymóphum. k.s. (br.y.) variable.	S. America.	1816. 5. 9. S. 4. — Cav. ic. 6. t. 595. f. 1.
<i>Ugēna polymóphha.</i> Cav. <i>Hydroglóssum hirsutum.</i> w.		
2 scándens. Sw. (br.y.) climbing.	E. Indies.	1793. 5. 9. S. 4. — Lodd. bot. cab. t. 742.
3 microphy'lum. B.P. (br.y.) small-leaved.	N. Holland.	1818. — S. 4. — Cavan. icon. 6.t.595.f.2.
4 venástum. Sw. (br.y.) elegant.	S. America.	1812. — S. 4. —
5 circinátum. Sw. (br.y.) rounded.	W. Indies.	1818. — S. 4. —
6 palmátum. Sw. (br.y.) palmate.	N. America.	1823. — S. 4. — Schkuhr filic. t. 140.
<i>Hydroglóssum palmátum.</i> w.		

ANE'IMIA. W.	ANE'IMIA.	Cryptogamia Schismatopterides. W.
1 fraxinifòlia. Rad. (br.y.) Ash-leaved.	Brazil.	1829. 9. 12. S. 4. —
2 hírta. w. (br.y.) hairy-leaved.	W. Indies.	1824. — S. 4. —
3 collinà. Rad. (br.y.) hill.	Brazil.	1829. — S. 4. —
4 hirsuta. w. (br.y.) hairy.	W. Indies.	1794. S. 4. —
5 laciñiata. (br.y.) jagged.	—	— S. 4. —
6 adiantifòlia. w. (br.y.) Maiden-hair-leav'd.	—	1793. 8. 9. S. 4. —
7 lanceolata. w. (br.y.) spear-leaved.	—	1820. — S. 4. —

OSMU'NDA. W.	OSMU'NDA.	Cryptogamia Schismatopterides. W.
1 cinnamómea. w. (br.) woolly.	N. America.	1772. 6. H. 4. —
2 regális. w. (br.y.) flowering fern.	Britain. 7. 8. H. 4. —
3 Claytoniàna. w. (br.y.) Clayton's.	N. America.	1772. 8. H. 4. —
4 interrupta. w. (br.y.) interrupted.	— 6. 7. H. 4. —
5 spectabilis. w. (br.y.) showy.	—	1811. 7. H. 4. —
6 humilis. w. (br.y.) dwarf.	—	1823. 6. 8. H. 4. —

TODEA. W.	TODEA.	Cryptogamia Schismatopterides. W.
africana. w. (br.y.) African.	C. B. S.	1805. 5. 8. G. 4. —

SCHIZ'ÆA. W.	SCHIZ'ÆA.	Cryptogamia Schismatopterides. W.
1 penicillata. w. (br.y.) pencilled.	S. America.	1816. 6. 7. S. 4. —
2 pusilla. Ph. (br.y.) small.	N. America.	1825. — H. 4. —
3 rupéstris. B.P. (br.y.) rock.	N. S. W.	1822. — G. 4. —
4 bifida. B.P. (br.y.) bifid.	—	— G. 4. —

SUBORDO III. GLEICHENEÆ. Brown prodr. p. 160.

GLEICH'ENIA. B.P.	GLEICH'ENIA.	Cryptogamia Poropterides. W.
1 pubéscens. k.s. (br.) pubescent.	Vera Cruz.	1822. 6. 8. S. 4. —
<i>Merténsia pubéscens.</i> w.		
2 Spelúnca. B.P. (br.) various-leaved.	N. S. W.	— — G. 4. —
3 Hermánni. B.P. (br.) Hermann's.	India.	1829. S. 4. —
4 microphylla. B.P. (br.) small-leaved.	N. S. W.	1825. 6. 8. G. 4. —
5 flabellata. B.P. (br.) fan-leaved.	—	1823. — G. 4. —

MARATTIA. W.	MARATTIA.	Cryptogamia Poropterides. W.
alata. w. (br.y.) winged.	Jamaica.	1793. 5. 8. S. 4. — Sm. ic. ined. t. 46.

SUBORDO IV. PARKERIACEÆ. Hook et Grev. ic. fil. 97.

PARK'ERIA. H.F.	PARK'ERIA.	Cryptogamia Poropterides.
1 pteridoïdes. H.F. (br.) Brake-like.	W. Indies.	1822. 5. 10. S. 4. — H. et G. ic. fil. t. 97.
2 Lockhárti. H.F. (br.) Lockhart's.	Trinidad.	1824. — S. 4. —
CERATO'PTERIS. S.S.	CERATO'PTERIS.	Cryptogamia Poropterides.
thalictróides. s.s. (br.) Thalictrum-like.	E. Indies.	1818. 6. 10. S. 4. — Pluk. alm. t. 215. f. 3.
<i>Ellobocárpus oleráceus.</i> Kaulfuss.		

SUBORDO V. POLYPODIACEÆ. Brown prodr. p. 145.

ACRO'STICHUM. W.	ACRO'STICHUM.	Cryptogamia Filices. L.
1 viscosum. H.F. (br.y.) viscous.	W. Indies.	1826. 6. 10. S. 4. — H. et G. ic. fil. t. 61.
2 simplex. w. (br.y.) simple.	Jamaica.	1793. 5. 7. S. 4. — Lodd. bot. cab. t. 709.
3 apódum. H.F. (br.y.) stemless.	W. Indies.	1824. — S. 4. — H. et G. ic. fil. t. 99.

FILICES.

4 crinítum. w. (<i>br.ye.</i>) hairy.	W. Indies. 1793.	5. 7. S. 4.	H. et G. ic. fil. t. 1.
5 alcicórne. B.P. (<i>br.</i>) Elk's-horn.	N. S. W. 1808.	8. 10. G. 4.	Bot. reg. t. 262-3.
6 Stemària. Beau. (<i>br.</i>) Stag's-horn.	Guinea. 1822.	5. 8. S. 4.	Beau. fl. d'ow. v.1. pl. 2.
7 appendiculátum. H. (<i>br.</i>) appendaged.	W. Indies. 1824.	6. 8. S. 4.	Hook. ex. flor. t. 108.
8 àureum. w. (<i>br.ye.</i>) golden.	America. 1820.	7. 8. G. 4.	Schkuhr fil. t. 1.
NOTHOLÆNA. B.P. NOTHOLÆNA.	Cryptogamia Filices.		
1 Marántæ. B.P. (<i>br.ye.</i>) Swiss.	Switzerland. 1824.	6. 8. H. 4.	Spreng. crypt. t.2. f.18.
<i>Acrostichum Marántæ. w.</i>			
2 distans. B.P. (<i>br.ye.</i>) distant.	N. S. W. 1823.	— G. 4.	
3 lanuginòsa. s.s. (<i>br.ye.</i>) woolly.	Madeira. 1778.	8. 9. G. 4.	Desf. atl. 2. t. 256.
<i>Acrostichum velleum. H.K.</i>			
HEMIONITIS. W. HEMIONITIS.	Cryptogamia Filices. W.		
palmata. w. (<i>br.ye.</i>) palmated.	W. Indies. 1793.	6. 8. S. 4.	Hook. ex. flor. t. 33.
GYMNOGRA'MMA. GYMNOGRA'MMA.	Cryptogamia Filices.		
1 pedata. s.s. (<i>br.ye.</i>) pedate.	New Spain. 1822.	6. 9. S. 4.	Swtz. syn. fil. t. 1. f. 3.
2 rufa. s.s. (<i>br.ye.</i>) rusty-haired.	Jamaica. 1793.	6. 8. S. 4.	Schkuhr fil. t. 17. 21.
<i>Hemionitis rufa. w.</i>			
3 tartárea. s.s. (<i>br.ye.</i>) whitened.	S. America. 1817.	— S. 4.	Willd. hort. ber.1. t.40.
<i>Acrostichum tartáreum. Swtz. Hemonitis dealbata. w.</i>			
4 peruviana. s.s. (<i>br.ye.</i>) Peruvian.	Peru. 1822.	6. 8. G. 4.	Plum. fil. t. 144.
5 trifoliáta. s.s. (<i>br.ye.</i>) three-leaved.	Jamaica. 1810.	— S. 4.	Willd. hort. ber.1. t. 41.
6 calomeláno. s.s. (<i>br.ye.</i>) mealy.	W. Indies. 1790.	7. 8. S. 4.	
7 chrysophylla. s.s. (<i>br.ye.</i>) golden-leaved.	— 1824.	— S. 4.	
8 sulphurea. s.s. (<i>br.ye.</i>) sulphur-colored.	Jamaica. 1808.	— S. 4.	
9 leptophylla. s.s. (<i>br.ye.</i>) slender-leaved.	S. Europe. 1824.	5. 8. H. 4.	
10 chærophylla. s.s. (<i>br.ye.</i>) Chervil-leaved.	Brazil. 1825.	6. 8. S. 4.	
MENISCUM. W. MENISCUM.	Cryptogamia Filices. W.		
1 triphyllum. s.s. (<i>br.ye.</i>) three-leaved.	E. Indies. 1820.	6. 8. S. 4.	H. et G. ic. fil. t. 120.
2 sorbitolum. s.s. (<i>br.ye.</i>) Service-leaved.	S. America. 1825.	— S. 4.	Plum. fil. t. 110.
3 reticulatum. w. (<i>br.ye.</i>) netted.	Martinico. 1793.	6. 8. S. 4.	Hook. ex. flor. t. 78.
XIPHO'PTERIS. K.F. SWORD FERN.	Cryptogamia Filices.		
1 serrulata. K.F. (<i>br.ye.</i>) saw-leaved.	W. Indies. 1823.	6. 8. S. 4.	
<i>Grammitis serrulata. w.</i>			
2 myosuroides. s.s. (<i>br.ye.</i>) Mousetail-like.	— 1824.	— S. 4.	
GRAMMITIS. Sw. GRAMMITIS.	Cryptogamia Filices. W.		
1 australis. B.P. (<i>br.ye.</i>) southern.	N. S. W. 1822.	6. 8. G. 4.	H. et G. ic. fil. t. 43.
2 linearis. s.s. (<i>br.ye.</i>) linear-leaved.	Jamaica. 1823.	— S. 4.	Schkuhr crypt. t. 7.
3 lanceolata. s.s. (<i>br.ye.</i>) spear-leaved.	Mauritius. 1824.	— S. 4.	H. et G. ic. fil. t. 62.
4 elongata. s.s. (<i>br.ye.</i>) long-leaved.	W. Indies. —	— S. 4.	Lodd. bot. cab. t. 15.
5 furcata. H.F. (<i>br.ye.</i>) forked-leaved.	Trinidad. 1825.	— S. 4.	
CETERACH. W. CETERACH.	Cryptogamia Filices.		
officinárum. w. (<i>br.</i>) common.	Britain.	5. 10. H. 4.	
<i>Scopolendrium Ceterach. Eng. bot. v. 18. t. 1244.</i>			
POLYBOTRYA. S.S. POLYBOTRYA.	Cryptogamia Filices.		
1 vivipara. H.E.F. (<i>br.ye.</i>) viviparous.	W. Indies. 1823.	5. 8. S. 4.	Hook. ex. flor. t. 107.
2 cervina. s.s. (<i>br.ye.</i>) hart's-tongue.	— 1824.	— S. 4.	H. et G. ic. fil. t. 81.
NIPHOBOLUS. K.F. NIPHOBOLUS.	Cryptogamia Filices.		
1 pertusus. s.s. (<i>br.ye.</i>) bored.	China. 1817.	7. 10. G. 4.	Hook. ex. flor. t. 162.
<i>Polypodium pertusum. Hooker.</i>			
2 rupéstris. s.s. (<i>br.ye.</i>) rock.	N. S. W. 1824.	— G. 4.	H. et G. ic. fil. t. 93.
3 conflúentes. s.s. (<i>br.ye.</i>) confluent.	— 1822.	— G. 4.	Thunb. jap. t. 38.
4 Língua. s.s. (<i>br.ye.</i>) tongue-like.	Japan. 1817.	— G. 4.	Plum. fil. 105. t. 121.
POLYPODIUM. W. POLYPODY.	Cryptogamia Filices. L.		
1 sérpens. w. (<i>ye.</i>) trailing.	W. Indies. 1822.	3. 8. S. 4.	Plum. fil. t. 137.
2 angustifolium. w. (<i>ye.</i>) narrow-leaved.	S. America. —	— S. 4.	Plum. fil. t. 118.
3 lanceolatum. w. (<i>ye.</i>) spear-leaved.	— 1824.	— S. 4.	Schkuhr fil. t. 8. c. p.
4 attenuatum. B.P. (<i>ye.</i>) attenuated.	N. S. W. 1823.	— G. 4.	Plum. fil. 114. t. 130.
5 piloselloides. w. (<i>ye.</i>) Mouse-ear.	W. Indies. 1793.	— S. 4.	H. et G. ic. fil. t. 125.
6 vaccinifolium. s.s. (<i>ye.</i>) Whortle-leaved.	Brazil. 1826.	— S. 4.	Plum. fil. 107. t. 127.
7 teniòsum. s.s. (<i>ye.</i>) jointed.	S. America. 1815.	— S. 4.	— 115. t. 131.
8 lycopodioides. w. (<i>ye.</i>) Club-moss.	W. Indies. 1793.	— S. 4.	Thunb. ic. jap. t. 18.
9 phyllitidis. w. (<i>ye.</i>) Hart's-tongue.	— 1824.	— S. 4.	Lab. nov. hol. 2. t. 240.
10 iroïdes. s.s. (<i>ye.</i>) Iris-leaved.	E. Indies. 1824.	— S. 4.	Plum. fil. 59. t. 76.
11 crassifolium. w. (<i>ye.</i>) thick-leaved.	W. Indies. 1816.	— S. 4.	
12 repens. w. (<i>ye.</i>) creeping.	S. America. —	— S. 4.	
13 lineare. s.s. (<i>ye.</i>) linear-leaved.	Japan. 1822.	— G. 4.	
14 Billardieri. B.P. (<i>ye.</i>) La Billardiere's.	V. Diem. Isl. 1823.	— G. 4.	
15 àureum. w. (<i>ye.</i>) golden.	W. Indies. 1742.	3. 5. S. 4.	
16 phymatodes. w. (<i>ye.</i>) warted.	— 1816.	— S. 4.	

17 quercifolium. s.s. (<i>ye.</i>)	Oak-leaved.	E. Indies.	1824.	3. 5. S. 24.	Rumph. amb. 6. t. 36.
18 curvatum. s.s. (<i>ye.</i>)	curve-leaved.	Jamaica.	1822.	— S. 24.	Eng. bot. v. 16. t. 1149.
19 vulgare. E.B.	(<i>ye.</i>) common.	Britain.	5. 10. H. 24.	Bolt. fil. t. 2. f. 5. a.
β cāmbricum.	(<i>ye.</i>) Welsh.	Wales.	— H. 24.	Plum. fil. t. 77.
20 virginianum. w.	(<i>ye.</i>) Virginian.	N. America.	7. H. 24.	H. et G. ic. fil. t. 56.
21 Scoulerii. H.F.	(<i>ye.</i>) Scouler's.	—	1827.	— H. 24.	Lodd. bot. cab. t. 748.
22 pectinatum. w.	(<i>ye.</i>) comb-leaved.	W. Indies.	1793.	6. 9. S. 24.	H. et G. ic. fil. t. 42.
23 Plumula. w.	(<i>ye.</i>) feather.	S. America.	1822.	5. 8. S. 24.	Plum. fil. 66. t. 85.
24 scolopendrioïdes. H.F.	(<i>ye.</i>) hart's-tongue-l.	Jamaica.	1824.	— S. 24.	— 85. t. 102. f.A.
25 tenuifolium. w.	(<i>ye.</i>) slender-leaved.	S. America.	1822.	— S. 24.	Schkuhr fil. t. 11. b.
26 asplenifolium. w.	(<i>ye.</i>) Spleenwort-leav'd.	Martinico.	1790.	— S. 24.	— p. 11. t. 10.
27 incanum. w.	(<i>ye.</i>) hoary.	America.	1811.	7. H. 24.	
28 trichomanoides. w.	(<i>ye.</i>) Trichomanes-like.	W. Indies.	1822.	5. 8. S. 24.	
29 decumanum. s.s. (<i>ye.</i>)	robust.	Brazil.	1818.	— S. 24.	
30 juglandifolium. w.	(<i>ye.</i>) Walnut-leaved.	S. America.	1822.	— S. 24.	
31 proliferum. s.s. (<i>ye.</i>)	proliferous.	Brazil.	1829.	9. 12. S. 24.	
32 crenatum. w.	(<i>ye.</i>) crenated.	S. America.	1824.	— S. 24.	
33 fraxinifolium. w.	(<i>ye.</i>) Ash-leaved.	Caracas.	1822.	5. 8. S. 24.	
34 tenellum. B.P.	(<i>ye.</i>) slender.	N. S. W.	—	5. 7. S. 24.	
35 tetragonum. s.s. (<i>ye.</i>)	four-sided.	W. Indies.	1826.	6. 9. S. 24.	Eng. bot. v. 31. t. 2224.
36 Phegopteris. E.B.	(<i>ye.</i>) sun-fern.	Britain.	6. 7. H. 24.	Pluk. alm. t. 284. f. 2.
37 hexagonopterum. w.	(<i>ye.</i>) triangular.	N. America.	1811.	— H. 24.	
38 pruinatum. w.	(<i>ye.</i>) white-leaved.	Jamaica.	1793.	5. 8. S. 24.	Sloan. jam. 1. t. 57. f. 3.
39 effusum. w.	(<i>ye.</i>) spreading.	—	1769.	11. S. 24.	Eng. bot. v. 9. t. 616.
40 Dryopteris. w.	(<i>ye.</i>) tender-branched.	Britain.	6. 9. H. 24.	
41 connectile. w.	(<i>ye.</i>) connected.	N. America.	1812.	7. H. 24.	Eng. bot. v. 22. t. 1525.
42 calcareum. w.	(<i>ye.</i>) rigid-branched.	Britain.	— H. 24.	

WO'ODSIA. L.T.

WO'ODSIA. Cryptogamia Filices. L.

1 pubescens. s.s. (<i>br.ye.</i>)	pubescent.	Brazil.	1826.	1. 12. S. 24.	
2 glabélla. s.s. (<i>br.ye.</i>)	smooth.	N. America.	1827.	6. 9. H. 24.	Eng. bot. v. 29. t. 2023.
3 hyperborea. L.T. (<i>br.ye.</i>)	hairy Alpine.	Scotland.	7. 9. H. 24.	Schkuhr fil. t. 19.
4 ilvensis. L.T. (<i>br.ye.</i>)	rock.	N. America.	1812.	6. 7. H. 24.	
5 vestita. Spr. (<i>br.ye.</i>)	clothed.	—	1816.	— H. 24.	H. et G. ic. fil. t. 68.
6 Perriniâna. H.F. (<i>br.ye.</i>)	Perrin's.	—	1825.	8. 11. G. 24.	

Alsophila Perriniâna. s.s.

PLEOPELTIS. K.S. PLEOPELTIS.

Cryptogamia Filices.

1 ensifolia. H.E.F. (<i>br.y.</i>)	sword-leaved.	C. B. S.	1823.	5. 6. S. 24.	Hook. ex. flor. t. 62.
2 latifolia. L.C. (<i>br.ye.</i>)	broad-leaved.	—	—	— S. 24.	
3 nuda. D.P. (<i>br.ye.</i>)	naked.	Nepaul.	—	— S. 24.	Hook. ex. flor. t. 63.
4 angusta. s.s. (<i>br.ye.</i>)	narrow-leaved.	Mexico.	1826.	— S. 24.	
ASPIDIUM. B.P. SHIELD-FERN.	Cryptogamia Filices. W.				
1 Hookeri. (<i>br.ye.</i>)	Hooker's.	W. Indies.	1812.	6. 9. S. 24.	H. et G. ic. fil. t. 96.
prolifica. H.F. non B.P.					

2 rhizophyl'lum. H.F. (<i>br.y.</i>)	rooting-leaved.	Jamaica.	1816.	— S. 24.	H. et G. ic. fil. t. 59.
3 trifoliatum. w. (<i>br.ye.</i>)	three-leaved.	W. Indies.	1769.	4. 8. S. 24.	Jacq. ic. 3. t. 638.
4 cicutarium. w. (<i>br.ye.</i>)	Cicuta-leaved.	N. America.	1823.	— H. 24.	Pluk. alm. t. 289. f. 4.
5 pátent. w. (<i>br.ye.</i>)	pubescent.	W. Indies.	1784.	7. 9. S. 24.	Schkuhr fil. t. 334.
6 dentatum. w. (<i>br.ye.</i>)	toothed.	Scotland.	7. H. 24.	Eng. bot. v. 23. t. 1588.
Cyáthea dentata. E.B.					
7 Halleri. w. (<i>br.ye.</i>)	Haller's.	Switzerland.	1824.	4. 8. H. 24.	
8 bulbiferum. w. (<i>br.ye.</i>)	bulbiferous.	N. America.	1638.	7. 8. H. 24.	
9 rhæticum. w. (<i>ye.</i>)	stone.	Switzerland.	— H. 24.	
10 frágile. w. (<i>ye.</i>)	brittle.	Britain.	6. 8. H. 24.	Eng. bot. v. 23. t. 1587.
11 régium. w. (<i>ye.</i>)	laciniated.	—	6. H. 24.	v. 3. t. 163.
Cyáthea incisa. E.B.					
12 æmulum. w. (<i>ye.</i>)	dwarf.	Madeira.	1779. F. 24.	
13 alpinum. w. (<i>ye.</i>)	Alpine.	S. Europe.	1825.	6. 8. H. 24.	Jacq. ic. 3. t. 642.
14 montanum. w. (<i>ye.</i>)	mountain.	—	—	— H. 24.	
15 irriguum. E.B.	brook.	Britain.	6. H. 24.	Eng. bot. v. 31. t. 2199.
16 atomarium. w. (<i>ye.</i>)	small.	N. America.	1822.	6. 8. H. 24.	

NEPHR'ODIUM. B.P. NEPHR'ODIUM.	Cryptogamia Filices.				
1 Lonchitidis. H.S.L. (<i>ye.</i>)	rough Alpine.	Britain.	5. 8. H. 24.	Eng. bot. v. 12. t. 797.
Aspidium Lonchitidis. E.B.					
2 auriculatum. H.S.L. (<i>ye.</i>)	eared.	E. Indies.	1793.	7. S. 24.	
3 acrostichoïdes. M. (<i>ye.</i>)	Acrostichum-like.	N. America.	1824.	5. 8. H. 24.	Schkuhr fil. t. 30.
4 mucronatum. s.s. (<i>ye.</i>)	mucronate.	Jamaica.	1826.	— S. 24.	
5 exaltatum. B.P. (<i>ye.</i>)	lofty.	—	1793.	5. 8. S. 24.	Schkuhr fil. t. 32. b.
6 unatum. B.P. (<i>ye.</i>)	smooth united.	E. Indies.	—	— S. 24.	Burm. zeyl. t. 44. f. 1.
7 propinquum. B.P. (<i>ye.</i>)	pubescent.	—	—	— S. 24.	

FILICES.

8 mólle. B.P.	(ye.) soft.	N. S. W.	1822.	5. 8. G. 4.	Jacq. ic. t. 640.
9 sesquipedále. s.s.	(ye.) black-spotted.	W. Indies.	1824.	— S. 4.	
10 hirsútulum. s.s.	(ye.) hairy.	E. Indies.	1823.	— S. 4.	
11 èdule. D.P.	(ye.) eatable-rooted.	Nepaul.	1826.	— G. 4.	
12 semicordátum. s.s.	(ye.) half cordate.	W. Indies.	1822.	— S. 4.	
13 noveboracéns. H.S.L.	(ye.) river-side.	N. America.	1812.	7. H. 4.	Schkuhr fil. t. 46.
14 pectinátum.	(ye.) comb-like.	1823.	5. 8. S. 4.	— t. 296.
<i>Aspidium trapezoides</i> . Schk.					
15 Oreópteris. H.S.L.	(ye.) Heath.	Britain.	7. H. 4.	Eng. bot. v. 15. t. 1019.
16 Thelypteris. H.S.L.	(ye.) Lady fern.	7. 8. H. 4.	— v. 15. t. 1018.
17 sérra. s.s.	(ye.) saw-like.	Jamaica.	1819.	— S. 4.	
18 parasiticum. w.	(ye.) parasitical.	E. Indies.	1824.	— S. 4.	
19 Goldiánium. H.F.	(ye.) Goldie's.	N. America.	1822.	— H. 4.	H. et G. ic. fil. t. 102.
20 cristátum. H.S.L.	(ye.) crested heath.	Britain.	6. 7. H. 4.	Eng. bot. v. 30. t. 2125.
21 aculeátum. H.S.L.	(ye.) common prickly.	6. 8. H. 4.	— v. 22. t. 1562.
22 pubéscens. H.F.	(ye.) pubescent.	Jamaica.	1824.	— S. 4.	H. et G. ic. fil. t. 162.
23 marginál. H.S.L.	(ye.) marginal-flower'd.	N. America.	1772.	6. 9. H. 4.	Schkuhr fil. t. 45. b.
24 Félix-más. H.S.L.	(ye.) Male fern.	Britain.	6. 8. H. 4.	Eng. bot. v. 21. t. 1458.
β Smithii.	(ye.) Smith's.	— H. 4.	— v. 28. t. 1949.
<i>Aspidium cristátum</i> . E.B. 1949. non 2. 25.					
γ eròsum. Ph.	(ye.) jagged.	N. America.	1812.	— H. 4.	Schkuhr fil. t. 45.
25 lancastriénse. s.s.	(ye.) Lancaster.	1825.	— H. 4.	
26 frágans. H.F.	(ye.) fragrant.	Europe.	— H. 4.	H. et G. ic. fil. t. 70.
27 platyphyllum. s.s.	(ye.) flat-leaved.	S. America.	1826.	— S. 4.	
28 lobátum. H.S.L.	(ye.) close-leaved.	England.	6. 9. H. 4.	Eng. bot. v. 22. t. 1563.
29 intermédium. w.	(ye.) intermediate.	N. America.	1816.	— H. 4.	
30 spinulósium. H.S.L.	(ye.) crested prickly.	Britain.	6. 8. H. 4.	Eng. bot. v. 21. t. 1460.
31 dilatátum. H.S.L.	(ye.) great crested.	— H. 4.	— v. 21. t. 1461.
32 obtúsium. w.	(ye.) blunt leaved.	N. America.	1827.	— H. 4.	Schkuhr fil. t. 21.
33 rigidum. w.	(ye.) rigid.	Germany.	1816.	— H. 4.	
34 elongátum. H.S.L.	(ye.) cut-leaved.	Madeira.	1799.	7. 8. F. 4.	
35 villósium. H.S.L.	(ye.) villous.	W. Indies.	1793.	7. S. 4.	
36 decompósito. B.P.	(ye.) decomound.	N. S. W.	1820.	6. 8. F. 4.	Schkuhr fil. t. 46. b.
37? Báromez. Baromez.	Tartary.	1824. S. 4.	
ALLANT'ODIA. B.P.	ALLANT'ODIA.	Cryptogamia	Filices.		
1 umbrósia. B.P.	(ye.) Madeira.	Madeira.	1779.	6. 9. F. 4.	Schkuhr fil. t. 61.
2 axillaris. s.s.	(ye.) slender.	— F. 4.	
<i>Polypodium axillare</i> . H.K. <i>Aspidium axillare</i> . Sw.					
3 australis. B.P.	(ye.) southern.	V. Diem. Isl.	1820.	— F. 4.	
4 ténera. B.P.	(ye.) membranaceous.	N. S. W.	—	— F. 4.	
ASPL'ENIUM. B.P.	SPLENWORT.	Cryptogamia	Filices.	L.	
1 acútum. s.s.	(br.) acute.	Canaries.	1818.	4. 7. F. 4.	
2 fissum. s.s.	(br.) cleft-leaved.	Hungary.	1825.	— H. 4.	
3 fontánum. B.P.	(st.) smooth rock.	England.	6. 8. H. 4.	Eng. bot. v. 29. t. 2024.
<i>Aspidium fontánum</i> . E.B.					
4 Félix-fémina. B.P.	(st.) female fern.	Britain.	6. 9. H. 4.	Eng. bot. v. 21. t. 1459.
5 Michaùxi. s.s.	(st.) Michaux's.	N. America.	1812.	— H. 4.	
6 Athyrium. s.s.	(st.) American.	1820.	— H. 4.	Schkuhr filic. t. 78.
<i>Nephrodium asplenoides</i> . M.					
7 Hallèri. s.s.	(st.) Haller's.	Switzerland.	1825.	— H. 4.	
8 Adiantum-nigrum. w.	(br.) black.	Britain.	4. 10. H. 4.	Eng. bot. v. 28. t. 1950.
9 lanceolátum. E.B.	(br.) lanceolate.	England.	6. 10. H. 4.	— v. 4. t. 240.
10 rádicans. s.s.	(br.) rooting-branched.	W. Indies.	1822.	— S. 4.	
11 bissectum. s.s.	(br.) bifid.	Jamaica.	1821.	— S. 4.	
12 montánum. w.	(br.) mountain.	N. America.	1812.	7. H. 4.	
13 frágans. w.	(br.) fragrant.	Jamaica.	1793.	— S. 4.	
14 Ruta murária. w.	Wall-rue.	Britain.	6. 10. H. 4.	Eng. bot. v. 3. t. 150.
15 cicutárium. H.S.	(br.) Cicuta-like.	S. America.	1822.	— S. 4.	Plum. fil. t. 48. a.
16 formosum. w.	(br.) handsome.	Santa Cruz.	—	— S. 4.	
17 prémorsum. w.	(br.) snip-leaved.	Jamaica.	1793.	— S. 4.	
18 striátum. w.	(br.) striated.	W. Indies.	—	6. 8. S. 4.	
19 thelypteróides. w.	Ladyfern-like.	N. America.	1823.	7. H. 4.	
20 sulcátum. s.s.	(br.) furrowed.	W. Indies.	1827.	— S. 4.	Schkuhr fil. t. 76. b.
21 pùmulum. s.s.	(br.) dwarf.	1823.	— S. 4.	Plum. fil. t. 46.
22 canariénse. s.s.	(br.) Canary Island.	Canaries.	1824.	— G. 4.	— t. 66. A.
23 rhizóphorum. w.	(br.) root-bearing.	Jamaica.	1793.	6. 8. S. 4.	Sloan. j. 1. t. 29. 30. f. 1.
24 víride w.	(br.) green.	Britain.	6. 9. H. 4.	Eng. bot. v. 32. t. 2257.
25 melanocáulon. w.	(br.) black-stalked.	N. America.	1812.	7. H. 4.	
26 Trichomanes. w.	(br.) Maiden hair.	Britain.	5. 10. H. 4.	Eng. bot. v. 8. t. 576.

27	germánicum. w.	alternate-leaved. Scotland.	6.	10.	H. 4.	Eng. bot. v. 32. t. 2258.
	<i>alternifolium</i> . E.B. <i>Bréynii</i> . Sw.						
28	ébèneum. w.	(br.) ivory striped.	N. America.	1779.	9.	H. 4.	Lodd. bot. cab. t. 5.
	<i>trichomanoides</i> . M.						
29	bipartitum. s.s.	(br.) two-parted.	W. Indies.	1822.	—	S. 4.	
30	dimidiatum. s.s.	(br.) narrow-leaved.	—	1827.	—	S. 4.	
31	aúritum. s.s.	(br.) eared.	—	1829.	9.	12.	S. 4.
32	zamiaefolium. s.s.	(br.) Zamia-leaved.	S. America.	1818.	6.	10.	S. 4.
33	falcatum. B.P.	(br.) sickle-leaved.	N. S. W.	1825.	6.	7.	F. 4.
34	monánthemum. w.	(br.) one-flowered.	C. B. S.	1790.	7.	G. 4.	
35	reséctum. s.s.	(br.) cut-leaved.	Mauritius.	1820.	—	S. 4.	
36	marinum. w.	(br.) sea-side.	Britain.	6.	10.	H. 4.
37	dentatum. H.F.	(br.) tooth-leaved.	Jamaica.	1824.	—	S. 4.	
38	flabellifolium. w.	(br.) fan-leaved.	N. S. W.	1825.	6.	7.	G. 4.
39	déntex. w.	(br.) toothed.	C. B. S.	1790.	6.	10.	G. 4.
40	diffórmē. B.P.	(br.) deformed.	N. S. W.	1822.	6.	7.	F. 4.
41	Shephérdii. s.s.	(br.) Shepherd's.	Jamaica.	1824.	6.	9.	S. 4.
42	angustifolium. s.s.	(br.) narrow-leaved.	N. America.	1812.	—	H. 4.	
43	septentrionale. w.	(br.) forked.	Britain.	6.	10.	H. 4.
44	palmatum. s.s.	(br.) palmated.	S. Europe.	1816.	7.	10.	F. 4.
45	attenuatum. B.P.	(br.) slender.	N. S. W.	1825.	6.	7.	F. 4.
46	rhizophyl'num. w.	(br.) rooting-leaved.	N. America.	1880.	—	H. 4.	
47	serratum. w.	(br.) saw-leaved.	W. Indies.	1793.	—	S. 4.	
48	Nidus. B.P.	(br.) nest.	N. Holland.	1822.	—	F. 4.	
CÆNO'PTERIS. S.S. CÆNO'PTERIS. Cryptogamia Filices.							
1	Odontites. s.s.	(br.) long-leaved.	N. Holland.	1822.	6.	10.	G. 4.
	<i>Asplenium Odontites</i> . B.P. <i>Dárea Odontites</i> . w.						Thunb.n.ac.pet.t.E.f.1.
2	appendiculata. s.s.	(br.) loose.	V. Diem. Isl.	—	—	G. 4.	
3	rhizophylla. s.s.	(br.) rooting-leaved.	W. Indies.	1827.	—	S. 4.	
SCOLOPE'NDRIUM. W. HART'S-TONGUE. Cryptogamia Filices. W.							
1	officinárum. w.	(br.ye.) common.	Britain.	7.	8.	H. 4.
	β críspum.	(br.ye.) curled-leaved.	—	—	—	H. 4.	
	γ undulatum.	(br.ye.) waved-leaved.	—	—	—	H. 4.	
	δ multífidum.	(br.ye.) clustered.	—	—	—	H. 4.	
	ε ramosum.	(br.ye.) branching.	—	—	—	H. 4.	
2	Hemionítis. s.s.	(br.) Mule's fern.	Canaries.	1779.	6.	7.	G. 4.
	<i>Asplenium Hemionitis</i> . L.						Schkuhr crypt. t. 66.
DO'ODIA. B.P. Do'ODIA. Cryptogamia Filices.							
1	áspera. B.P.	(br.ye.) rough-stalked.	N. S. W.	1808.	3.	9.	G. 4.
2	caudáta. B.P.	(br.ye.) tailed.	—	1822.	—	G. 4.	
WOODWA'RDIЯ. W. Woodwa'rdia. Cryptogamia Filices. W.							
1	onocleoides. w.	(br.ye.) Onoclea-like.	N. America.	1812.	8.	10.	H. 4.
2	virginica. w.	(br.ye.) Virginian.	—	1774.	8.	9.	H. 4.
3	rádicans. w.	(br.ye.) rooting-leaved.	Madeira.	1779.	9.	10.	G. 4.
BLE'CHNUM. W. BLE'CHNUM. Cryptogamia Filices. W.							
1	brasiliense. s.s.	(br.ye.) Brazilian.	Brazil.	1828.	9.	11.	S. 4.
2	cartilagineum. B.P.	(br.) cartilaginous.	N. S. W.	1822.	6.	8.	F. 4.
3	denticulatum. s.s.	(br.ye.) toothed.	Teneriffe.	1826.	6.	9.	F. 4.
4	lævigatum. B.P.	(br.) smooth.	N. S. W.	1820.	3.	9.	G. 4.
5	angustifolium. s.s.	(br.ye.) narrow-leaved.	Trinidad.	1824.	—	S. 4.	
6	longifolium. s.s.	(br.) long-leaved.	—	1826.	—	S. 4.	
7	glandulosum. s.s.	(br.) glandular.	Brazil.	1823.	—	S. 4.	
8	occidentale. w.	(br.) American.	S. America.	1777.	—	S. 4.	
9	australe. w.	(br.) Cape.	C. B. S.	1691.	—	G. 4.	
10	polypodioides. s.s.	(br.) Polypody-leav'd.	Brazil.	1829.	8.	12.	S. 4.
11	Lancéola. s.s.	(br.) spear-leaved.	—	—	—	S. 4.	
LOM'ARIA. W. LOM'ARIA. Cryptogamia Filices. W.							
1	nuda. w.	(br.) naked.	V. Diem. Isl.	1822.	4.	8.	F. 4.
	<i>Onoclea nuda</i> . Lab. <i>Stegania nuda</i> . B.P.						Lab. n. hol. 2. t. 246.
2	Spicant. s.s.	(br.) northern.	Britain.	7.	9.	H. 4.
	<i>Osmunda Spicant</i> . L. <i>Bléchnum boreale</i> . E.B.						Eng. bot. v. 17. t. 1159.
3	onocleoides. s.s.	(br.) Onoclea-like.	Jamaica.	1824.	—	S. 4.	
4	sorbifolia. s.s.	(br.) Service-leaved.	W. Indies.	1793.	—	S. 4.	
	<i>Acróstichum sorbifolium</i> . L.						Plum. fil. t. 117. dextr.
5	longifolia. s.s.	(br.) long-leaved.	—	1810.	6.	9.	S. 4.
6	procera. s.s.	(br.) lofty.	N. Holland.	1822.	4.	8.	F. 4.
	<i>Bléchnum procerum</i> . Lab. <i>Stegania procera</i> . B.P.						Lab. n. hol. 2. t. 247.
ONOCL'EА. W. ONOCL'EА. Cryptogamia Filices. L.							
1	sensibilis. w.	(br.) sensitive fern.	Virginia.	1799.	8.	H. 4.	Schkuhr fil. t. 102.
2	obtusilobata. Ph.	(br.) blunt-lobed.	N. America.	1812.	7.	H. 4.	— t. 103.

FILICES.

- | | | |
|--|----------------------------|----------------------------|
| STRUTHIO'PTERIS. W. | STRUTHIO'PTERIS. | Cryptogamia Filices. |
| 1 germanica. s.s. (br.) German. | Europe. | 1760. 7. 8. H. 4. |
| Osmunda Struthiopteris. L. | Onoclea Struthiopteris. w. | |
| 2 pensylvánica. w. (br.) Pensylvanian. | N. America. | 1812. — H. 4. |
| Onoclea nodulosa. Schkuhr. | | Schkuhr fil. t. 104. |
| VITTARIA. W. | VITTARIA. | Cryptogamia Filices. W. |
| lineata. w. (br.) linear-leaved. | America. | 1793. S. 4. |
| ANTR'OPHYUM. S.S. ANTR'OPHYUM. | Cryptogamia Filices. | |
| lanceolatum. s.s. (br.) spear-leaved. | W. Indies. | 1793. 6. 10. S. 4. |
| Hemionitis lanceolata. L. | | Schkuhr fil. t. 6. |
| DIPL'AZIUM. W. | DIPL'AZIUM. | Cryptogamia Filices. W. |
| 1 plantagineum. s.s. (br.) Plantain-leav'd. | W. Indies. | 1822. S. 4. |
| 2 grandifolium. w. (br.) large-leaved. | Jamaica. | 1793. 8. 10. S. 4. |
| 3 juglandifolium. s.s. (br.) Walnut-leaved. | — | 1825. S. 4. |
| 4 seraporiense. s.s. (br.) East Indian. | E. Indies. | 1827. S. 4. |
| 5 auriculatum. s.s. (br.) eared. | S. America. | 1820. 8. 10. S. b. |
| 6 barbadense. w. (br.) Barbadoes. | W. Indies. | 1822. S. 4. |
| 7 malabáricum. s.s. (br.) Malabar. | E. Indies. | 1818. 6. 8. S. b. |
| Asplenium ambiguum. Swartz. | | |
| 8 arboréscens. s.s. (br.) arborescent. | Mauritius. | 1824. S. b. |
| 9 esculentum. w. (br.) eatable. | Ceylon. | 1822. S. b. |
| CRYPTOGRA'MMA. Br. CRYPTOGRA'MMA. | Cryptogamia Filices. | |
| 1 acrostichoïdes. Br. (st.) Acrostichum-like. N. America. | 1827. | 7. 9. H. 4. |
| Allosorus acrostichoïdes. s.s. | | H. et G. ic. fil. t. 29. |
| 2 crispá. Br. (st.) curled. | Britain. | — H. 4. |
| Allosorus crispus. s.s. Pteris crispia. E.B. Osmunda crispia. L. | | Eng. bot. v. 17. t. 1160. |
| PTERIS. W. | BRAKE. | Cryptogamia Filices. L. |
| 1 pedata. w. (st.) pedate. | W. Indies. | 1816. 5. 8. S. 4. |
| 2 palmata. w. (br. st.) palmated. | — | 1823. 6. 7. S. 4. |
| 3 argentea. w. (st. br.) silvery. | Siberia. | — H. 4. |
| 4 longifolia. w. (st. br.) long-leaved. | W. Indies. | 1770. 7. 9. S. 4. |
| 5 grandifolia. w. (st. br.) large-leaved. | — | 1793. S. 4. |
| 6 serrulata. w. (st. br.) various-leaved. | India. | 1770. 8. 9. S. 4. |
| 7 denticulata. s.s. (st. br.) toothed. | Brazil. | 1824. 5. 9. S. 4. |
| 8 crética. w. (st. br.) Cretan. | Candia. | 1818. 5. 7. G. 4. |
| 9 atropurpurea. w. (br.) dark purple. | N. America. | 1770. 8. 9. H. 4. |
| 10 nemoralis. W. en. (st. br.) wood. | Africa. | 1823. 5. 8. S. 4. |
| 11 biaurita. s.s. (st. br.) two-eared. | W. Indies. | 1824. — S. 4. |
| 12 Plumérii. s.s. (st. br.) Plumier's. | S. America. | 1818. — S. 4. |
| 13 arguta. w. (st. br.) sharp-notched. | Madeira. | 1778. 8. 9. F. 4. |
| 14 umbrósa. B.P. (st. br.) shade. | N. S. W. | 1823. 5. 8. F. 4. |
| 15 cordata. w. (st. br.) heart-leaved. | Mexico. | 1825. — G. 4. |
| 16 sagittata. w. (st. br.) arrow-leaved. | S. America. | 1820. — S. 4. |
| 17 vespertilioñis. B.P. (st.) Bat-winged. | N. S. W. | 1823. — F. 4. |
| 18 trémula. B.P. (st. br.) trembling. | — | 1822. — F. 4. |
| 19 esculenta. B.P. (st. br.) esculent. | — | 1815. — F. 4. |
| 20 gracilis. w. (st. br.) slender. | Canada. | 1823. 7. H. 4. |
| 21 hastata. w. (st. br.) hastate. | W. Indies. | — 5. 8. S. 4. |
| 22 aculeata. w. (st. br.) prickly-stem'd. | — | 1793. S. b. |
| 23 falcatá. B.P. (st. br.) falcate. | N. S. W. | 1820. 5. 7. F. 4. |
| 24 caudata. w. (st. br.) American. | N. America. | 1777. 9. 12. H. 4. |
| 25 aquilina. E.B. (st. br.) common. | Britain. | — 7. 8. H. 4. |
| 26 lanuginosa. w. (st. br.) woolly. | Bourbon. | 1818. — S. 4. |
| 27 intramarginalis. K.F. (br.) Mexican. | Mexico. | 1828. 7. 12. S. 4. |
| TÆN'ITIS. W. | TÆN'ITIS. | Cryptogamia Filices. W. |
| 1 lanceolata. s.s. (br.) spear-leaved. | W. Indies. | 1818. 6. 9. S. 4. |
| 2 angustifolia. s.s. (br.) narrow-leaved. | — | 1823. — S. 4. |
| 3 graminifolia. H.E.F. (br.) grass-leaved. | Trinidad. | 1820. 6. 8. S. 4. |
| 4 furcata. s.s. (br.) forked. | — | 1824. — S. 4. |
| 5 chinensis. s.s. (br.) Chinese. | China. | 1828. — G. 4. |
| ADIA'NTUM. W. | MAIDENHAIR. | Cryptogamia Filices. L. |
| 1 reniforme. w. (br.) kidney-leaved. | Madeira. | 1699. 6. 9. G. 4. |
| 2 obliquum. s.s. (br.) oblique-leaved. | W. Indies. | 1826. 4. 8. S. 4. |
| 3 macrophyllum. w. (br.) large-leaved. | Jamaica. | 1793. — S. 4. |
| 4 lunulatum. s.s. (br.) lunulate. | E. Indies. | 1826. — S. 4. |
| 5 deltoidium. s.s. (br.) deltoid-leaved. | Jamaica. | 1824. — S. 4. |
| 6 denticulatum. s.s. (br.) toothed-leaved. | W. Indies. | 1825. — S. 4. |
| 7 serrulatum. w. (br.) saw-leaved. | Jamaica. | 1823. 6. 8. S. 4. |
| 8 radiatum. w. (br.) radiated. | W. Indies. | 1776. 4. 8. S. 4. |
| 9 pedatum. w. (br.) Canadian. | N. America. | 1640. 8. 9. H. 4. |
| | | Sloan. j. 17. t. 36. f. 2. |
| | | Plum. fil. t. 100. |
| | | Schkuhr fil. t. 115. |

0 hispidulum. B.P. (br.) bristly.	N. Holland.	1822.	5. 8. G. 4.	
1 striatum. s.s. (br.) streaked.	W. Indies.	—	— S. 4.	
2 villosum. w. (br.) hairy-stalked.	Jamaica.	1775.	6. 9. S. 4.	Schkuhr fil. t. 120.
3 pulverulentum.w.(br.) dusty.	W. Indies.	1793.	— S. 4.	— t. 119.
4 triangularis. m.s.s. (br.) triangular.	Trinidad.	1824.	— S. 4.	
1 Capillus-vis. w.(br.) true.	Britain.	5. 9. H. 4.	Eng. bot. v. 22. t. 1564.
1 trapeziformis. (br.) rhomb-leaved.	—	1793.	— S. 4.	Schkuhr fil. t. 122.
1 pentadactylis. (br.) five-fingered.	Brazil.	1828.	— S. 4.	H. et G. ic. fil. t. 98.
1 cuneata. (br.) wedge-leaved.	—	1824.	— S. 4.	— ic. fil. t. 30.
1 as. " (br.) likened.	N. S. W.	1823.	5. 8. G. 4.	Lab.n.hol.2.t.248.f.2.
trigona.		1822.	— G. 4.	
2 formosa. (br.) handsome.	Jamaica.	1793.	7. S. 4.	Pluk. alm. t. 254. f. 1.
2 tenerum. (br.) tender.	CHEILA'N S. W. CHEILA'NTHES.	Cryptogamia Filices. W.		
ferruginea. s.s. (br.) ferruginous.	Peru.	1816.	5. 8. S. 4.	
gracilis. (br.) slender.	Canada.	1827.	— H. 4.	
pteroidea. s.s. (br.) Pteris-like.	C. B. S.	1775.	7. 9. G. 4.	Hout. n. hist. t. 96. f. 3.
odora. s.s. (br.) sweet-scented.	S. Europe.	1824.	— F. 4.	
suavolens. s.s. (br.) scented.	Madeira.	1778.	— G. 4.	
fragrans. s.s. (br.) fragrant.	E. Indies.	—	— S. 4.	Sw. syn. fil. t. 3. f. 6.
micropilla. s.s. (br.) small-leaved.	W. Indies.	1822.	5. 8. S. 4.	
vestita. s.s. (br.) clothed.	N. America.	1812.	7. H. 4.	Schkuhr fil. t. 124.
hirta. s.s. (br.) hairy.	C. B. S.	1816.	5. 8. G. 4.	
lendigera. (br.) chaffy.	S. America.	1822.	— S. 4.	
tenuifolia. B. (br.) slender-leaved.	N. S. W.	1824.	— G. 4.	
spectabilis. s.s. (br.) showy.	Brazil.	1829.	9. 12. S. 4.	
repens. s.s. (br.) creeping.	W. Indies.	1824.	6. 10. S. 4.	
LONCH'ITI LONCH'ITIS.	Cryptogamia Filices. W.			
1 pedata. (br.) pedate.	Jamaica.	1793.	6. 8. S. 4.	Brown. jam. 89. t. 1.
Pteris p. (br.)	W. Indies.	—	— S. 4.	Schkuhr fil. t. 86.
hirsuta. s.s. (br.)	LINDS'EAE. Cryptogamia Filices. W.			
1 reniformis. (br.) leaved.	Trinidad.	1826.	5. 7. S. 4.	Linn. trans. v.3. t.7.f.1.
2 linearis. (br.) leaved.	N. S. W.	1820.	4. 6. G. 4.	Sw. fil. 118. t. 3. f. 3.
3 falcatula. L.T. (br.) leaved.	—	1824.	5. 8. S. 4.	Linn. trans. v. 3.t.7.f.1.
4 trapeziformis. L.T. (br.) leaved.	W. Indies.	1825.	— S. 4.	— v. 3. t. 9.
5 microphylla. B.P. (br.) leaved.	N. S. W.	1820.	4. 6. G. 4.	
DAVAL'IA. W. LIA. Cryptogamia Filices. W.				
1 pyxidata. B.P. (ye.) s. (br.) N. S. W.	1808.	4. 9. G. 2.		
2 canariensis. w. (ye.) (br.) Hare-s-toot fern. Canaries.	1699.	— G. 4.	Jacq. ic. 1. t. 200.	
3 gibberosa. s.s. (br.ye.) gibbous-flower'd. N. Zealand.	1825.	— G. 4.	Swt. flor. aust. t. 31.	
4 fumarioidea. s.s. (br.ye.) Fumitory-like.	W. Indies.	1828.	6. 10. S. 4.	
5 flaccida. B.P. (br.ye.) flaccid.	N. Holland.	1826.	— G. 4.	
BALA'NTIUM. K.F. BALA'NTIUM. Cryptogamia Filices.				
Culcita. K.S. (br.ye.) shining-leaved.	Madeira.	1779.	9. 10. F. 4.	
Dicks'onia Culcita. w.				
DICKS'ONIA. W. DICKS'ONIA. Cryptogamia Filices. W.				
1 pilosiuscula. w. (br.ye.) hairy.	N. America.	1811.	7. 9. H. 4.	Schkuhr fil. t. 131.
2 arborescens. w. (br.ye.) tree.	S. Helena.	1786.	6. 12. S. 2.	Jacq. ic. t. 201.
3 dissicata. w. (br.ye.) cut-leaved.	Jamaica.	1793.	9. 10. S. 4.	
4 adiantoides. s.s. (br.st.) Adiantum-like.	W. Indies.	1828.	8. 12. S. 4.	
CYATH'EAE. B.P. CYATH'EAE. Cryptogamia Filices. W.				
1 arborea. w. (br.ye.) tree.	W. Indies.	1793.	— S. 4.	Plum. fil. 1. t. 1. 2.
2 excelsa. s.s. (br.ye.) tall.	Mauritius.	1825.	— S. 2.	
ALSO'PHILA. B.P. ALSO'PHILA. Cryptogamia Filices.				
1 australis. B.P. (br.st.) southern.	N. S. W.	1820.	— G. 2.	
2 aspera. s.s. (br.ye.) rough.	W. Indies.	1824.	— S. 2.	
HEMIT'ELIA. B.P. HEMIT'ELIA. Cryptogamia Filices.				
multiflora. s.s. (br.ye.) many-flowered.	Jamaica.	1824.	— S. 2.	
TRICHO'MANES. W. TRICHO'MANES. Cryptogamia Filices. W.				
1 membranaceum. w. (br.) membranaceous.	W. Indies.	1820.	5. 8. S. 4.	Hook. ex. flor. t. 76.
2 floribundum. s.s. (br.ye.) many-flowered.	—	1825.	— S. 4.	H. et G. ic. fil. t. 9.
3 brevisetum. H.K. (br.ye.) short-styled.	Britain.	5. 6. H. 4.	Eng. bot. v. 20. t. 1417
Hymenophyl'lum alatum. E.B.				
4 alatum. s.s. (br.ye.) wing-leaved.	W. Indies.	1824.	— S. 4.	H. et G. ic. fil. t. 11.
5 crispum. s.s. (br.ye.) curled.	—	1828.	— S. 4.	— ic. fil. t. 12.
HYMENOPHYLLUM. FILMY-LEAF. Cryptogamia Filices. W.				
1 polyanthos. s.s. (br.ye.) many-flowered.	W. Indies.	1824.	6. 9. S. 4.	H. et G. ic. fil. t. 128.
2 tunbridgeense. w. (br.) Tunbridge.	Britain.	5. 6. H. 4.	Eng. bot. v. 3. t. 162.
3 hirsutum. s.s. (br.ye.) hairy.	Trinidad.	1823.	5. 9. S. 4.	H. et G. ic. fil. t. 84.

ADDENDA et CORRIGENDA.

RANUNCULACEÆ. p. 1.

CLE'MATIS. p. 1.	VIRGIN'S BOWER.
grandiflora. p. 1. (gr.) large-flowered.	Sierra Leone. 1822.
<i>chlorantha.</i> B. Reg.	
43 viornoïdes. Jac. (li.) Viorna-like.	N. America. 1826.
ANEM'ONÆ. p. 3.	ANEMONE.
26 Richardsoni. H.A. (ye.) Richardson's.	N. America. 1827.
27 deltoidea. H.A. (wh.) deltoid.	Columbia. —
RANU'NCULUS. p. 4.	CROWFOOT.
98 rhomboïdeus. H.A. (ye.) rhomboid-leav'd.	N. America. 1825.
99 glaberrimus. H.A. (ye.) smoothest.	— 1827.
100 ovális. H.A. (ye.) oval-leaved.	— —
101 brevicáulis. H.A. (ye.) short-stalked.	— —
102 cardiophy'llus. H.A. (ye.) heart-leaved.	— —
103 pygmaeus. DC. (ye.) pygmy.	— —
104 Eschscholtzii. H.A. (ye.) Eschscholtz's.	— —
105 pedatifidus. DC. (ye.) pedatifid.	— —
106 recurvátus. DC. (ye.) recurved.	— —
107 Schlechtendàlli. H.A. (ye.) Schlechtendahl's. <i>fasciculàris.</i> Schlec. non Muhl.	— —
CA'LTHA. p. 7.	MARSH MARYGOLD.
9 biflora. H.A. (wh.) two-flowered.	N. America. 1827.
10 leptosépala. H.A. (wh.) slender-sepaled.	— —
CO'PTIS. p. 7.	CO'PTIS.
2 asplenifolia. H.A. (wh.) Fern-leaved.	N. America. 1827.
ACON'ITUM. p. 9.	MONKSHOOD.
129 Nuttallii. (pa.) Nuttall's.	N. America. 1828.
<i>pállidum.</i> N. non R.A.	
130 glábrum. DC. (pa.) smooth. 1829.
	— — H. 2.
	5. 6. H. 2.
	6. 9. H. 2.
	4. 7. H. 2.
	— — H. 2.
	Edin. phi. jou. 6.t. 11.f.l.
	Hook. fl. b. am. t. 5. A.
	— — — t. 6. B.
	— — — t. 7. A.
	Bot. mag. t. 2999.
	Wahl. lapp. t. 8. f. 1.
	Schlec. an. s. 2. t. 2.
	Hook. fl. b. am. t. 10.
	6. 7. H. 2.
	Hook. fl. b. am. t. 11.
	7. 8. H. 2.

DILLENIACEÆ. p. 12.

TRACHYTE'LLA. DC.	TRACHYTE'LLA.	Polyandria Monogynia.
Actæa. DC.	(wh.) spear-leaved.	China. 1826. 6. 8. H. 2.

MAGNOLIACEÆ. p. 13.

TALA'UMA. DC.	TALA'UMA.	Polyandria Polygynia.
Candóllii. Bl. (st.) De Candolle's.	Java.	1827. 4. 6. S. 2.

Magnolia odoratissima. Hort.

BERBERIDEÆ. p. 16.

BERBERIS. p. 16.	BARBERRY.
19 caroliniana. N. (ye.) Carolina.	Carolina. 1828. 5. 7. H. 2.
20 crataégina. DC. (ye.) Hawthorn-like.	Levant. 1829. H. 2.
21 glumacea. DC. (ye.) glumaceous.	N. America. 1827. H. 2.
A'CHLYS. DC.	A'CHLYS.
triphyllea. DC. (wh.) three-leaved.	N. America. 1827. 4. 6. H. 2.
EPIM'EDIUM. p. 16.	BARRENWORT.
2 hexándrum. H.A. (li.) hexandrous.	N. America. 1827. 4. 5. H. 2.
	Hook. fl. b. am. t. 12.
	Hook. fl. b. am. t. 13.

NYMPHÆACEÆ. p. 17.

NYMPHÆA. p. 17.	WATER-LILY.					
odoràta. p. 18.	sweet-scented.	N. America.				
β <i>máxima</i> . n. (wh.) largest.	—	1828.	7. 8.	H.w. $\frac{1}{4}$.		
20 sanguínea. n. (re.) bloody.	—	—	—	H.w. $\frac{1}{4}$.		
N'UPHAR. p. 18.	N'UPHAR.					
advéna. p. 18.	striped-flowered.	N. America.				
β <i>lutea</i> . n. (ye.) yellow-flowered.	—	1828.	—	H.w. $\frac{1}{4}$.		

SARRACENIÆ. p. 18.

SARRACENIA. p. 18.	SIDE-SADDLE-FLOWER.					
1 minor. n.	(pu.) small.	Carolina.	1829.	6. 7.	H. $\frac{1}{4}$.	Swt. br. fl. gar. s. 2.

PAPAVERACEÆ. p. 18.

ARGEM'ONE. p. 19.	ARGEM'ONE.					
5 intermédia.	(wh.) intermediate.	Mexico.	1828.	7. 10.	H. $\frac{1}{4}$.	
SANGUIN'ARIA. p. 19.	PUCCOON.					
2 grandiflora. Ros. (wh.) large-flowered.	N. America.	1812.	3. 5.	H. $\frac{1}{4}$.	Rosc. flor. ill. sea. t. 8.	
RCÈM'ERIA. p. 19.	RCÈM'ERIA.					
vermiculata. Lehm. (re.) red-flowered.	Persia.	1829.	6. 9.	H. $\frac{1}{2}$.		
GLA'UCIUM. p. 19.	HORN POPPY.					
5 rùbrum. DC.	(re.) red-flowered.	Levant.	—	—	H. $\frac{1}{2}$.	Flor. græc. t. 488.
6 pérsicum. DC.	(re.ve.) red and velvet.	Persia.	—	—	H. \odot .	

CRUCIFERÆ. p. 21.

NASTURTIUM. p. 22.	NASTURTIUM.					
1 nátans. DC.	(st.) floating.	N. America.	1827.	5. 9.	H.w. $\frac{1}{4}$.	Deles. ic. sel. 2. t. 15.
BRA'YA. p. 22.	BRA'YA.					
3 pilosa. H.A.	(pu.) hairy.	N. America.	—	4. 6.	H. $\frac{1}{4}$.	Hook.fl.b.am.t.17.f.A.
ARABIS. p. 22.	WALL-CRESS.					
47 retrofracta. Gr. (bh.) bending.	N. America.	—	4. 7.	H. $\frac{1}{4}$.		
Turritis retrofracta. H.A.						
48 pátila. W.K.	(pu.) spreading.	Hungary.	—	—	H. $\frac{1}{4}$.	W. et K. hung. 1. t. 59.
49 lævigata. DC.	(wh.) smooth.	N. America.	—	—	H. $\frac{1}{4}$.	
MACROP'ODIUM. p. 2.	MACROP'ODIUM.					
2 laciniatum. H.A. (wh.) cut-leaved.	N. America.	—	6. 9.	H. \odot .	Hook.bot.mis.p.III.ic	
PA'RRYA. p. 24.	PA'RRYA.					
3 macrocarpa. Br. (pu.) large-podded.	N. America.	—	5. 6.	H. $\frac{1}{4}$.	Hook. fl. b. am. t. 15.	
DR'ABA. p. 25.	WHILLOW-GRASS.					
35 crassifolia. Gr. (ye.) thick-leaved.	N. America.	—	4. 5.	H. $\frac{1}{4}$.		
ERY'SIMUM. p. 30.	ERY'SIMUM.					
crepidifolium. s.s. (ye.) Crepis-leaved.	N. America.	1829.	5. 6.	H. $\frac{1}{2}$.		
ÆTHION'Ema. p. 32.	ÆTHION'Ema.					
5 membranaceum. DC. (li.) membranous-podded.	Persia.	—	6. 8.	H. $\frac{1}{4}$.	Swt. br. fl. gar. s. 2. t. 69.	
SIN'APIS. p. 34.	MUSTARD.					
30 amplexicaulis. DC. (ye.) stem-clasping.	Algiers.	—	6. 9.	H. \odot .	Desf. atl. 2. t. 153.	

CAPPARIDEÆ. p. 37.

CLE'OME. p. 37.	CLE'OME.					
20 speciosissima. B.R. (pu.) pretty.	Mexico.	1827.	8. 11.	H. \odot .	Bot. reg. t. 1312.	
CA'PPARIS. p. 38.	CAPER-TREE.					
acutifolia. (wh.) sharp-pointed.	S. America.	—	6. 8.	S. h.	Bot. reg. t. 1320.	
acuminata. Bot. reg. nec aliorum.						

VIOLARIEÆ. p. 43.

VIOILA. p. 43.	VIOLET.					
89 pedatifida. G.D. (bl.) pedatifid.	N. America. 1827.	5. 7. H. ¼.				
90 attenuata. B.F.G. (wh.) attenuated.	—	1812.	4. 8. H. ¼.			Swt. br. fl. gar. s. 2.
	<i>lanceolata</i> . Pursh et Nuttall <i>non</i> Lin.					
91 præmorsa. B.R. (ye.) bitten-rooted.	—	1827.	4. 6. H. ¼.			Bot. reg. t. 1254.
HYMENANTH'ERA. DC. HYMENANTH'ERA.	Monadelphia Pentandria.					
dentata. DC. (ye.) toothed.	N. S. W. 1822.	3. 5. G. $\frac{1}{2}$.				

POLYGALÆ. p. 46.

POLY'GALA. p. 46.	MILKWORT.					
42 rubella. Ph. (ro.) pale red.	N. America. 1828.	6. 7. H. ¼.				

FRANKENIACEÆ. p. 48.

FRANK'ENIA. p. 48.	SEA HEATH.					
9 corymbosa. DC. (ro.) corymb-flowered.	Barbary. 1829.	6. 8. F. $\frac{1}{2}$.			Desf. atl. 1. t. 93.	

CARYOPHYLLEÆ. p. 48.

DIA'NTHUS. p. 49.	PINK.					
97 ciliatus. Lehm. (pi.) fringed.	1829.	6. 8. H. ¼.			
98 sicus. s.s. (pi.) Sicilian.	Sicily.	—	— H. ¼.			
AREN'ARIA. p. 56.	SANDWORT.					
68 glomerata. DC. (wh.) close-headed.	Tauria.	1829.	7. 9. H. \odot .			
SPERG'ULA. p. 55.	SPURREY.					
9 pilifera. DC. (wh.) piliferous.	Corsica.	1829.	6. 8. H. ¼.			

LINEÆ. p. 58.

LINUM. p. 58.	FLAX.					
41 altaicum. F. (bl.) Altay.	Altay.	1829.	6. 8. H. ¼.			

MALVACEÆ.

MA'LVA. p. 59.	MALLOW.					
72 Munroana. B.R. (sc.) Munro's.	N. America. 1827.	5. 10. H. ¼.				Bot. reg. t. 1306.
HIBISCUS. p. 61.	HIBISCUS.					
86 Telfairiae. (va.) Mrs. Telfair's.	Hybrid.	1825.	1. 12. S. $\frac{1}{2}$.			
α roseus.	(ro.) rose-coloured.	—	—	S. $\frac{1}{2}$.		
β lilacinus.	(li.) pink.	—	—	S. $\frac{1}{2}$.		Bot. mag. t. 2891.
γ carneus.	(fl.) flesh-coloured.	—	—	S. $\frac{1}{2}$.		
δ fulvescens.	(bf.) buff-coloured.	—	—	S. $\frac{1}{2}$.		
ϵ ochroleucus.	(st.) straw-coloured.	—	—	S. $\frac{1}{2}$.		
S'IDA. p. 63.	S'IDA.					
59 repanda. DC. (st.) repanded.	1829.	6. 8. S. $\frac{1}{2}$.			

BUTTNERIACEÆ. p. 66.

RE'EVESIA. B.R.	RE'EVESIA.	Polyadelphia Polyandria.				
thyrsidea. B.R. (wh.) thyrs-flowered.	China.	1818.	1. 7. G. $\frac{1}{2}$.			Bot. reg. t. 1236.
STER'CULIA. p. 66.	STER'CULIA.					
16 lanceolata. B.R. (br.) spear-leaved.	—	1822.	5. 6. G. $\frac{1}{2}$.			Bot. reg. t. 1256.
DOMB'EYA. p. 69.	DOMB'EYA.					
4 angulata. DC. (ro.) angle-leaved.	Mauritius.	1826.	4. 8. S. $\frac{1}{2}$.			Bot. mag. t. 2905.

HYPERICINEÆ. p. 76.

HYPERICUM. p. 76. ST. JOHN'S-WORT.
 77 Geblèri. Led. (ye.) Gebler's. Altay. 1829. 6. 8. H. b.

GUTTIFERÆ. p. 78.

GARCI'NIA. p. 78. GARCINIA.
 5 purpùrea. h.b. (pu.) purple. E. Indies. 1828. S. h.

GERANIACEÆ. p. 88.

PHYMATA'NTHUS. p. 89. WART-FLOWER.

6 intertinctus. s.g. (re.v.) stained.	Hybrid.	1827.	2. 10.	G. h.	Swt. ger. ser. 2. t. 54.
PELARG'ONIUM. p. 92. STORK'S-BILL.					
504 nodosum. s.g. (cr.v.) knotted-stalked.	Hybrid.	1824.	4. 11.	G. h.	Swt. ger. ser. 2. t. 68.
505 bipinnatifidum. s.g. (pu.) bipinnatifid.	—	1827.	—	G. h.	— ser. 2. t. 62.
506 Darnleyanum. s.g. (cr.) Earl of Darnley's.	—	1826.	—	G. h.	— ser. 2. t. 63.
507 pullaceum. s.g. (sc.v.) dark brown.	—	1827.	5. 10.	G. h.	— ser. 2. t. 76.
508 contiguum. s.g. (sc.) contiguous.	—	1828.	—	G. h.	— ser. 2. t. 73.
509 iconicum. s.g. (sc.bk.) figured.	—	—	—	G. h.	— ser. 2. t. 88.
510 Bluntianum. s.g. (sc.ve.) Miss Blunt's.	—	—	—	G. h.	— ser. 2. t. 79.
511 eriophorón. s.g. (sc.ve.) wool-bearing.	—	—	—	G. h.	— ser. 2. t. 90.
512 obtusidentatum. s.g. (sc.) blunt-toothed.	—	—	—	G. h.	— ser. 2. t. 92.
513 melancholicum. s.g. (sc.ve.) dark-edged.	—	1827.	—	G. h.	— ser. 2. t. 53.
514 Spéculum. s.g. (re.ve.) looking-glass.	—	—	—	G. h.	— ser. 2. t. 52.
515 anacamptón. s.g. (re.ve.) recurved-calyx'd.	—	—	—	G. h.	— ser. 2. t. 64.
516 Peytòniæ. s.g. (re.ve.) Lady Peyton's.	—	—	—	G. h.	— ser. 2. t. 46.
517 ursinum. s.g. (re.) shaggy-calyxed.	—	1828.	—	G. h.	— ser. 2. t. 94.
518 suffusum. s.g. (pi.re.) suffused.	—	1827.	—	G. h.	— ser. 2. t. 47.
519 laxulum. s.g. (ro.ve.) loose-umbelled.	—	1828.	—	G. h.	— ser. 2. t. 75.
520 conchylállum. s.g. (vi.ve.) violet-purple.	—	—	—	G. h.	— ser. 2. t. 95.
521 porphy'reon. s.g. (pu.) bright-purple.	—	—	—	G. h.	— ser. 2. t. 89.
522 mollifolium. s.g. (pi.ve.) soft-leaved.	—	1827.	—	G. h.	— ser. 2. t. 77.
523 Colleyanum. s.g. (pu.) Colley's.	—	1828.	—	G. h.	— ser. 2. t. 83.
524 Littleanum. s.g. (pu.li.) Little's.	—	1827.	—	G. h.	— ser. 2. t. 72.
525 dædáleum. s.g. (pi.re.) variable-coloured.	—	—	—	G. h.	— ser. 2. t. 100.
526 Veitchianum. s.g. (pu.) Veitch's.	—	—	—	G. h.	— ser. 2. t. 81.
527 implicátum. s.g. (pu.) implicated.	—	—	—	G. h.	— ser. 2. t. 86.
528 insculptum. s.g. (li.ve.) engraved-petal'd.	—	1828.	—	G. h.	— ser. 2. t. 65.
529 lanósum. s.g. (li.ve.) wool-bearing.	—	—	—	G. h.	— ser. 2. t. 84.
530 flabellifolium. s.g. (std.) fan-leaved.	—	—	—	G. h.	— ser. 2. t. 48.
531 cordifórmē. s.g. (bh.pu.) heart-shaped.	—	1827.	—	G. h.	— ser. 2. t. 49.
532 præclarum. s.g. (wh.pu.) clear-coloured.	—	—	—	G. h.	— ser. 2. t. 67.
533 clathrátum. s.g. (pi.ve.) burred-petaled.	—	—	—	G. h.	— ser. 2. t. 85.
534 instrátum. s.g. (wh.ve.) spreading-spotted.	—	1828.	—	G. h.	— ser. 2. t. 78.
535 Yeatmaniánum. s.g. (pu.ve.) Miss Yeatman's.	—	1827.	—	G. h.	— ser. 2. t. 59.
536 mirábole. s.g. (li.ve.) admirable.	—	—	—	G. h.	— ser. 2. t. 57.
537 látum. s.g. (pi.pu.) genteel.	—	—	—	G. h.	— ser. 2. t. 91.
538 commíxtum. s.g. (re.ve.) mingled.	—	—	—	G. h.	— ser. 2. t. 71.
539 compáctum. s.g. (bh.re.) compact.	—	1828.	—	G. h.	set. 2. t. 70.
540 exquisítum. s.g. (w.pu.) dainty-flowered.	—	1824.	—	G. h.	ser. 2. t. 55.
541 politum. s.g. (bh.re.) polished.	—	1827.	—	G. h.	ser. 2. t. 87.
542 Atkinsianum. s.g. (bh.pu.) Mr. Atkins's.	—	1828.	—	G. h.	ser. 2. t. 99.
543 atrovírens. s.g. (bh.pu.) dark green-leav'd.	—	1827.	—	G. h.	— ser. 2. t. 74.
544 adventitium. s.g. (pu.ve.) adventitious.	—	1826.	—	G. h.	— ser. 2. t. 80.
545 perámplum. s.g. (li.) very large-flower'd.	—	—	—	G. h.	— ser. 2. t. 61.
546 glabréscens. s.g. (w.pu.) smoothish-leav'd.	—	1824.	—	G. h.	— ser. 2. t. 45.
547 Hillianum. s.g. (w.pu.) Hill's.	—	1828.	—	G. h.	— ser. 2. t. 51.
548 pallidulum. s.g. (pi.) pale pink.	—	—	—	G. h.	— ser. 2. t. 97.
549 urbánum. s.g. (pi.) home bred.	—	—	—	G. h.	— ser. 2. t. 50.
550 dissimile. s.g. (re.l.ve.) dissimilar.	—	—	—	G. h.	— ser. 2. t. 60.
551 fastuosum. s.g. (re.ve.) fastuous.	—	—	—	G. h.	— ser. 2. t. 93.
552 succuléntum. s.g. (sc.ve.) succulent-leav'd.	—	1827.	—	G. h.	— ser. 2. t. 69.
553 Drákeæ. s.g. (cr.ve.) Mrs. Drake's.	—	1828.	—	G. h.	— ser. 2. t. 96.

ADDENDA et CORRIGENDA.

554 Annesleyànum. s.g. (re.ve.) Miss Annesley's. Hybrid.	1828.	5. 10.	G. h.	Swt. ger. ser. 2. t. 56.
555 Gloriànum. s.g. (cr.ve.) Queen of Portugal's.	—	—	G. h.	— ser. 2. t. 82.
556 Kenrickæ. s.g. (re.ve.) Mrs. Kenrick's.	—	—	G. h.	— ser. 2. t. 58.
557 nùtans. s.g. (sc.ve.) nodding-flowered.	1827.	—	G. h.	— ser. 2. t. 66.
558 Sweetiànum. Sm. (cr.ve.) Sweet's.	1829.	—	G. h.	—
559 staphysagroides. s.g. (pu.) Stavesacre-l'd. C. B. S.	1827.	—	G. h.	Swt. ger. ser. 2. t. 98.

BALSAMINEÆ. p. 101.

IMP'ATIENS. p. 102. TOUCH-ME-NOT.				
pálida. p. 102. pale-flowered.	N. America.	1812.	6. 10.	H. ⊖.
<i>Nòli-tángere.</i> E.B. nec aliorum.				Eng. bot. v. 14. t. 937.
7 boreális. (ye.) northern.	Denmark.	—	H. ⊖.
<i>Nòli-tángere.</i> F.D. non L.				Flor. dan. t. 582.
Nòli-tángere. L. (y.re.) English.	England.	—	H. ⊖.

OXALIDEÆ. p. 102.

O'XALIS. p. 102. WOOD-SORREL.				
113 tortuosa. B.R. (ye.) twisted.	Chile.	1825.	6. 8.	G. h.
114 floribunda. Leh. (ro.) many-flowered.	Brazil.	1829.	4. 9.	F. ȳ.
úrbica. Hil. (li.) Martius's.		1828.	5. 9.	F. ȳ.

Martiàna. Zuc. p. 102. ex S.S. Lodd. bot. cab. ic.

ZYGOPHYLLEÆ. p. 104.

LA'RREA. p. 104. LA'RREA.				
2 divaricata. DC. (ye.) spreading-lobed.	BuenosAyres.	1829.	6. 9.	G. h.

RUTACEÆ. p. 105.

APLOPHYLLUM. p. 105. APLOPHYLLUM.				
patavina, tuberculata, villosa, linifolia et dahurica, lege patavinum, tuberculatum, villosum, linifolium, et dahuricum.				
9 Buxbaumii. J.R. (ye.) Buxbaum's.	Levant.	1828.	6. 8.	H. ȳ.
<i>Ruta Buxbaumii.</i> DC.				Buxb. cent. 2. t. 28. f.2.

DICTAMNUS. p. 105. FRAXINELLA.

3 angustifolius. s.f.g. (pu.) narrow-leaved.	Siberia.	1829.	5. 7.	H. ȳ.
				Swt. br. fl. gar. ser. 2.

CELASTRINEÆ. p. 109.

EUONYMUS. p. 109. SPINDLE-TREE.				
16 nànus. DC. (st.) dwarf.	Caucasus.	1829.	5. 7.	H. h.
17 gròssus. F.I. (wh.) large.	Nepaul.	1828.	6. 8.	F. h.

RHAMNEÆ. p. 112.

TRAV'OA. H.M.	TRAV'OA.	Pentandria Monogynia.		
quinquenervia. H.M. (w.) five-nerved.	Chile.	1827.	5. 7.	F. h.

HOMALINEÆ. p. 116.

BLACKWE'LLIA. p. 11. BLACKWE'LLIA.				
7 padifolia. B.R. (wh.) Padus-leaved.	China.	1826.	8. 9.	G. h.

Bot. reg. t. 1308.

CASSUVIEÆ. p. 117.

MELANORRH'ŒA. W. MELANORRH'ŒA.	Polygamia Dioecia.			
usitata. w.i. (re.) useful.	E. Indies.	1828.	S. h.
HOLIGA'RNA. DC.	HOLIGA'RNA.	Polygamia Monœcia.		Wall. ind. pl. t. 11. 12.
longifolia. DC. (wh.) long-leaved.	E. Indies.	1828.	Roxb. corom. 3. t. 282.

BURSERACEÆ. p. 120.

IC'ICA. DC.	IC'ICA.	Octo-Decandria Monogynia.				
1 enneándria. DC. (<i>wh.</i>) nine-stamened.	Guiana.	1825.	S. h.	Aubl. guian. 1. t. 134.	
2 decándra. DC. (<i>wh.</i>) ten-stamened.	—	—	S. h.	Aubl. guian. 1. t. 132.	
3 altíssima. DC. (<i>wh.</i>) tallest.	—	1822.	S. h.		

LEGUMINOSÆ. p. 122.

Subordo I. PAPILIONACEÆ.

S'OPHORA. p. 122.	S'OPHORA.					
12 velutina. B.R. (<i>pu.</i>) velvety.	China.	1826.	6. 7. G. h.	Bot. reg. t. 1185.		
BAPTI'SIA. p. 123.	BAPTI'SIA.					
10 minor. Lehm. (<i>bl.</i>) lesser.	N. America.	6. 7. H. Ʌ.			
ANTHYLLIS. p. 132.	KIDNEY-VETCH.					
22 polycéphala. DC. (<i>ye.</i>) many-headed.	Barbary.	1829.	6. 7. H. Ʌ.	Desf. atl. 2. t. 195.		
HOSA'CKIA. B.R.	HOSA'CKIA.	Diadelphia Decandria.				
1 bicolor. B.R. (<i>y.w.</i>) two-coloured.	N. America.	1823.	7. 9. H. Ʌ.	Bot. reg. t. 1257.		
<i>Lòtus pinnatùs.</i> Bot. mag. t. 2913.						
2 Purshiana. B.R. (<i>ye.</i>) Pursh's.	—	1827.	— H. Ʌ.			
3 decumbens. B.R. (<i>ye.</i>) decumbent.	—	—	— H. Ʌ.			
4 parviflòra. B.R. (<i>ye.</i>) small-flowered.	—	—	— H. Ʌ.			
TEPHROSIA. p. 141.	TEPHROSIA.					
38 chinensis. H.C. (<i>li.</i>) Chinese.	China.	1822.	5. 8. G. h.			
ASTRA'GALUS. p. 145.	MILLE VETCH.					
120 succulentus. B.R. (<i>pu.</i>) succulent.	N. America.	1827.	6. 7. H. Ʌ.	Bot. reg. t. 1324.		
KENN'EDYA. p. 156.	KENN'EDYA.					
monophylla. p. 157.	simple-leaved.	N. S. W.	1790.	3. 8. G. h. — Vent. malm. t. 106.		
β longeracemosa. (<i>pi.</i>) long-racemed.	—	1828.	— G. h. — Bot. reg. t. 1336.			
LUP'INUS. p. 159.	LUPINE.					
34 pulchellus. (<i>bl.</i>) handsome.	Mexico.	—	6. 10. F. h.	Swt. br. fl. gar. s. 2. t. 67.		

Subordo III. MIMOSEÆ.

AC'ACIA. p. 164.	AC'ACIA.					
209 gravéolens. L.C. (<i>ye.</i>) strong-scented.	N. Holland.	1825.	3. 5. G. h.	Lodd. bot. cab. t. 1460.		
uncinata. p. 165. (<i>st.</i>) hooked.	N. Holland.	1823.	8. 1. G. h.	Bot. reg. t. 1332.		

210 álbida. B.R. (<i>ye.</i>) white spiny.	Peru.	1824.	4. 6. G. h.	— t. 1317.		
--	-------	-------	-------------	------------	--	--

Subordo IV. CÆSALPINEÆ.

CASTANOCA'Rpus.	MORETON BAY CHESTNUT.	Decandria Monogynia.				
australis.	(<i>wh.</i>) southern.	N. Holland.	1829. G. h.		
CA'SSIA. p. 169.	CA'SSIA.					

116 brevifolia. DC. (<i>ye.</i>) short-leaved.	Madagascar.	1824.	6. 7. S. h.			
--	-------------	-------	-------------	--	--	--

ROSACEÆ. p. 178.

R'UBUS. p. 187.	BRAMBLE AND RASPBERRY.					
85 flávus. D.P. (<i>wh.</i>) yellow-fruited.	Nepaul.	1827. F. h.			
86 setosus. DC. (<i>wh.</i>) bristly-stalked.	N. America.	—	5. 7. H. h.			
87 nutkánus. DC. (<i>wh.</i>) Nootka sound.	—	—	— H. h.			
DR'YAS. p. 190.	DR'YAS.					
3 Drummondii. B.M. (<i>ye.</i>) Drummond's.	—	—	4. 8. H. Ʌ.	Bot. mag. t. 2972.		
POTENTILLA. p. 190.	CINQUEFOIL.					
nívea. p. 190. (<i>ye.</i>) snowy-leaved.	N. Europe.	1816.	6. 8. H. Ʌ.			
β macrophylla. B.M. (<i>ye.</i>) large-leaved.	N. America.	1827.	— H. Ʌ.	Bot. mag. t. 2982.		
101 grácilis. B.M. (<i>ye.</i>) slender.	—	—	— H. Ʌ.	— t. 2984.		
102 Hopwoodiana. s.F.G. (<i>ro.s.</i>) Hopwood's. Recto-formosa.	1829.	6. 9. H. Ʌ.		Swt. br. fl. gar. s. 2. t. 61.		

COMBRETACEÆ. p. 195.

POIVREA. p. 196. Por'VREA.

6 Afzéllii. D.D. (sc.) Afzelius's. Sierra Leone. 1824. 7. 10. S. h. Bot. mag. t. 2944.

Combrétum grandiflorum. B.M. not G.D. in Linn. trans., which is a true *Combrétum*, with 8 stamens; we have adopted M. Decandolle's division of the genus, as Mr. Don informed us, he found the same distinctions as given by Decandolle, and was inclined to divide them himself, but afterwards arranged them in distinct sections, by which they might be readily divided, one genus having irregular convolute cotyledons, and the other straight plaited ones; this circumstance he mentioned to us before his paper was published, and before M. Decandolle's volume had arrived.

VOCHYSIEÆ. p. 197.

QUALEA. M.P.

QUALEA. Monandria Monogynia.

violacea. M.P. (vi.) violet-coloured. Brazil. 1824. S. h. Mart. pl. bras. 1. t. 81.

ERISMA. DC.

ERISMA. Monandria Monogynia.

floribunda. DC. (bl.) many-flowered. Guiana. 1825. S. h. Tratt. tab. t. 105.

ONAGRARIEÆ. p. 197.

FU'CHSIA. p. 197.

FU'CHSIA.

11 thymifolia. DC. (re.) Thyme-leaved. Mexico. 1829. 4. 10. F. h. Swt. br. fl. gar. s. 2. t. 25.

MELASTOMACEÆ. p. 203.

PLER'OMA. p. 204.

PLER'OMA.

3? villosum. DC. (pu.) villous. S. America. 1820. 5. 19. S. h. Bot. mag. t. 2630.
Melastoma villosa. B.M.

PASSIFLOREÆ. p. 216.

PASSIFLORA. p. 216. PASSION-FLOWER.

65 Cavanillesii. DC. (co.) Cavanille's. W. Indies. 1822. 7. 10. S. h. Cavan. dis. 10. t. 273.

66 Andersoni. DC. (st.) Andersson's. St. Vincent. 1823. — S. h.

67 Raddiana. DC. (st. pu.) Raddi's. Brazil. 1829. — S. h.

68 cephalaima. DC. crowded-leaved. 1826. S. h. Bory. an. gen. 2. t. 22. f. 2.

69 obscura. H.C. (gr. st.) obscure. S. America. 1823. 2. 10. S. h.

CACTEÆ. p. 235.

MAMMILLARIA. p. 235. MAMMILLARIA.

23 fulvispina. H.P. (re.) brown-spined. Brazil. 1822. 8. 10. D.S. h.

24 pulchra. H.B.R. (pu.) pretty. Mexico. 1828. 6. 10. D.S. h. Bot. reg. t. 1329.

25 depræssa. DC. (wh. re.) depressed. S. America. 1800. 6. 8. D.S. h. DC. diss. t. 2. f. 2.
discolor. DC. dis. non Haw. *Spinii*. Colla hort. rip. ie.

discolor. H.S. (wh. re.) two-coloured. S. America. — — D.S. h.

26 stellaris. H.S. (ro.) starry. — 1816. 6. 9. D.S. h. Lodd. bot. cab. t. 79.
non pusilla. DC. ex. H.P.

ECHINOCA'CTUS. p. 235. ECHINOCA'CTUS.

22 cornigerus. DC. (wh.) horn-spined. Mexico. 1830. 6. 9. D.S. h. DC. diss. cact. t. 7.

23 nobilis. H.P. (wh.) noble. — 1796. — D.S. h.

24 Hy'strix. H.P. (wh.) Porcupine. S. America. 1818. — D.S. h.

CEREUS. p. 236. CEREUS.

74 magnus. H.P. (wh.) great 12-angled. St. Domingo. 1830. 6. 10. D.S. h.

75 ferrox. H.P. (wh.) fierce upright. Brazil. 1828. — D.S. h.

76 Aethiops. H.P. (wh.) black-spined. — — D.S. h.

77 setiger. H.P. small quadrangular. 1826. D.S. h.

78 undatus. H.P. triangular China. 1824. D.S. h.

79 crispatus. H.P. curly-edged. Brazil. 1826. D.S. h.

80 Colvillii. (ro.) Colvill's. Grandiflora-speciosissimus. — D.S. h.

OPU'NTIA. p. 237.	INDIAN FIG.					
38 longispina. H.P. (ye.) long-spined.	Brazil.	1826.	6. 10.	D.S. $\frac{1}{2}$.		
39 glomerata. H.P. (ye.) long flat-spined.	—	—	—	D.S. $\frac{1}{2}$.		
40 rosea. DC. (ro.) rose-coloured.	Mexico.	1830.	D.S. $\frac{1}{2}$.	DC. diss. cact. t. 15.	
41 imbricata. DC. (ro.) imbricated.	S. America.	1820.	D.S. $\frac{1}{2}$.		
<i>Cereus imbricatus</i> . p. 237.						
42 cylindrica. DC. (ro.) cylindrical.	Peru.	1779.	D.S. $\frac{1}{2}$.		
<i>Cereus cylindricus</i> . p. 237.						
RHIPS'ALIS. p. 238.	RHIPS'ALIS.					
8 cereúscula. H.P.	Cereus-like.	Brazil.	1828.	D.S. $\frac{1}{2}$.	

GROSSULARIEÆ. p. 238.

R'IBES. p. 238.	CURRENT AND GOOSEBERRY.					
45 punctatum. DC. (st.) spotted.	Chile.	1825.	3. 5.	H. $\frac{1}{2}$.	Bot. reg. t. 1278.	

ESCALLONEÆ. p. 239.

ESCALL'ONIA. p. 239.	ESCALL'ONIA.					
5 bifida. Ot. (pi.) bifid.	Mexico.	1827.	F. $\frac{1}{2}$.		
6 viscósa. Ot. (wh.) viscous.	Mendoza.	1829.	G. $\frac{1}{2}$.		

SAXIFRAGEÆ. p. 239.

SAXI'FRAGA. p. 239.	SAXIFRAGE.					
102 spicata. L.T. (sp.) spiked.	N. America.	1827.	5. 6.	H. $\frac{1}{2}$.		
103 arguta. L.T. (sp.) deep-notched.	—	—	—	H. $\frac{1}{2}$.		
104 nudicaulis. L.T. (wh.) naked-stalked.	—	—	—	H. $\frac{1}{2}$.		
105 ferruginea. Gr. (wh.) ferruginous.	—	—	—	H. $\frac{1}{2}$.		
106 pyrolifolia. L.T. (wh.) Pyrola-leaved.	—	—	—	H. $\frac{1}{2}$.		
107 tricuspidata. L.T. (ye.) three-pointed.	Greenland.	1827.	—	H. $\frac{1}{2}$.		
HEUCH'ERA. p. 242.	HEUCH'ERA.					
1 micrantha. B.R. (gr.) small-flowered.	Columbia.	—	6. 7.	H. $\frac{1}{2}$.	Bot. reg. t. 1302.	
2 Richardsoni. D. (gr.) Richardson's.	N. America.	—	—	H. $\frac{1}{2}$.		
TIARE'LLA. p. 242.	TIARE'LLA.					
5 colorans. Gr. (wh.) coloured.	N. America.	—	4. 6.	H. $\frac{1}{2}$.		
MITE'LLA. p. 242.	MITE'LLA.					
5 trifida. Gr. (wh.) trifid.	N. America.	—	5. 7.	H. $\frac{1}{2}$.		
DRUMMO'NDIA. DC. DRUMMO'NDIA.	Pentandria Digynia.					
<i>Mitella pentandra</i> . Bot. mag. t. 2933.						

UMBELLIFERÆ. p. 243.

ERY'NGIUM. p. 244.	ERYNGO.	Pentandria Digynia.				
36 Andersoni. LG. (bl.) Anderson's.	8. 9.	H. $\frac{1}{2}$.		
DISCOPLE'URA. p. 245.	DISCOPLE'URA.					
2 capillacea. DC. (wh.) capillary.	N. America.	1824.	7. 9.	H. \odot .		
<i>A'mmi capillaceum</i> . M.						
FER'ULA. p. 250.	GIANT-FENNEL.					
21 linearis. Cerv. (ye.) linear-leaved.	Mexico.	1828.	8. 10.	F. $\frac{1}{2}$.		
ANTHRISCUS. p. 254.	ROUGH CHERVILL.					
9 torquata. DC. (wh.) twisted.	France.	1726.	6. 7.	H. δ .	Jacq. aust. 1. t. 63.	
<i>Chærophyllum bulbosum</i> . Jacq.						
ARPI'TIUM. Neck.	ARPI'TIUM.	Pentandria Digynia.				
simplic. (wh.) simple-stalked.	Switzerland.	1824.	6. 8.	H. $\frac{1}{2}$.	Jacq. misc. 2. t. 2.	
<i>Laserpitium simplex</i> . L. Gaya simplex. Gaud. fl. helv. 2. t. 6. non <i>Gaya</i> . Kth.						

ARALIACEÆ. p. 256.

ADA'MIA. W.I.	ADA'MIA.	Didynamia Angiospermia.				
cyanea. w.i.	(bl.) blue-flowered.	E. Indies.	1829.	S. $\frac{1}{2}$.	Wall. ind. t. 26.

CAPRIFOLIACEÆ. p. 257.

SYMPH'ORIA. p. 258. ST. PETER'S-WORT.
3 montâna. k.s. (li.) mountain. Mexico. 1829. 7. 10. H. $\frac{1}{2}$. H. et B. nov. gen. 3.t. 296

LORANTHEÆ. p. 259.

A'UCUBA. W. A'UCUBA. Diœcia Tetrandria.
japónica. (pu.) blotch-leaved. Japan. 1783. 5. 7. H. $\frac{1}{2}$. Bot. mag. t. 1197.
LORA'NTHUS. p. 259. LORA'NTHUS.
2 terréstris.terrestrial. N. Holland. 1830. G. $\frac{1}{2}$.

RUBIACEÆ. p. 260.

ASPER'ULA. p. 266. WOODRUFF.
25 nítida. F.G. (br.) glossy. Greece. 1829. 7. 9. H. $\frac{1}{2}$. Flor. græc. t. 124.

DIPSACEÆ. p. 269.

ASTEROC'EPHALUS. p. 270. STARHEAD.
paucisétus. Jac. (st.) few-bristled. S. Europe. 1827. 7. 9. H. $\frac{1}{2}$.

COMPOSITÆ. p. 271.

CR'EPIS. p. 273. CR'EPIS.
33 macrorhiza. B.M. (ye.) large-rooted. Madeira. 1829. 7. 9. H. $\frac{1}{2}$. Bot. mag. t. 2988.
CHUQUIR'AGA. p. 280. CHUQUIR'AGA.
2 spinosa. L.T. (ye.) spiny. Peru. 1825. 4. 5. F. $\frac{1}{2}$.
Bacazia spinosa. p. 286.
JU'NGIA. L. JU'NGIA. LABIATIFLORÆ. Syngenesia Æqualis.
paniculata. D.D. (ye.) paniced. Chile. 1825. 7. 9. H. $\frac{1}{2}$. DC. an. mus. 19. t. 7.
Dumerilia paniculata. p. 260.
CENTA'UREA. p. 285. CENTAURY.
141 argùta. s.s. (st.) sharp notched. Canaries. 1829. — F. $\frac{1}{2}$.
142 decípiens. Thu. (pu.) deceiving. France. — — H. $\frac{1}{2}$.
VERN'ONIA. p. 289. VERN'ONIA.
14 tomentosa. Ót. (pu.) tomentose. Brazil. 1827. 9. 11. S. $\frac{1}{2}$.
DELI'SLIA. S.S. DELI'SLIA. Syngenesia Necessaria.
Bertèrii. s.s. (ye.) Bertero's. Isl. Magdalén. 1829. 6. 8. G. \odot . Mart. dec. 47. f. 1.
Milleria biflora. p. 313.
CALE'NDULA. p. 313. MARYGOLD.
22 tomentosa. s.s. (ye.) tomentose. C. B. S. 1828. — H. \odot .
CLADA'NTHUS. p. 319. CLADA'NTHUS.
2 canescens. s.F.G. (ye.) canescent. Canaries. 1829. 3. 8. F. $\frac{1}{2}$. Swt. br. fl. gar. ser. 2.

LOBELIACEÆ. p. 322.

LOB'ELIA. p. 322. LOB'ELIA.
84 purpùrea. B.R. (pu.) purple. Chile. 1825. 8. 9. G. $\frac{1}{2}$. Bot. reg. t. 1325.

CAMPANULACEÆ. p. 325.

LIGHTFO'OTIA. p. 325. LIGHTFO'OTIA.
Loddigesii. A.DC. (pa.) Loddiges'. C. B. S. 1824. 6. 8. G. $\frac{1}{2}$. Lodd. bot. cab. t. 1038.
tenella. B.C.
PLATYC'ODON. A.DC. PLATYC'ODON. Pentandria Monogynia.
grandiflorum. DC. (bl.) great-flowered. Siberia. 1782. 6. 8. H. $\frac{1}{2}$. Bot. mag. t. 252.
Campánula grandiflora. B.M. *Wahlenbergia grandiflora*. p. 325.

WAHLENBERGIA. A.DC. WAHLENBERGIA.

3 Kitaibèlii. A.DC. (vi.) Kitaibel's.	Hungary.	1823.	6. 8. H. ȝ.	W. et K. hung. 2. t. 154.
Campánula graminifòlia. p. 327.	W. et K. hung. non Flor. græc.			
4 tenuifòlia. A.DC. (vi.) slender-leaved.	Hungary.	1820.	5. 9. H. ȝ.	W. et K. hung. 2. t. 155.
5 capénisia. A.DC. (vi.) Cape.	C. B. S.	1803.	7. 10. H. ⊙.	Bot. mag. t. 782.
Campánula capénisia. B.M. Roëlla decurrentis. Andr. rep. t. 238.				
6 cérrua. A.DC. (wh.) nodding-flowered.	C. B. S.	1804.	7. 9. G. ♂.	
7 lineàris. A.DC. (wh.) linear-leaved.	—	1822.	7. 10. H. ⊙.	
8 procumbens. A.DC. (bl.) procumbent.	—	1824.	— H. ⊙.	Alph. DC. camp. t. 15.
9 hederacea. A.DC. (pa. bl.) Ivy-leaved.	England.	5. 6. H. ȝ.	Eng. bot. v. 2. t. 73.
Campánula hederacea. p. 328. Aikinia hederacea. Salisb.				
10 arvática. (bl.) Spanish.	Spain.	1825.	5. 7. H. ȝ.	Pluk. phyt. t. 23. f. 1.
Campánula arvática. LG.				
11 gráclis. A.DC. (bl.) slender.	N. S. W.	1794.	4. 8. G. ♂.	Bot. mag. t. 691.
Campánula gráclis. B.M.				
12 strícta. (bl.) upright.	—	—	— G. ♂.	Sm. exot. bot. t. 45.
Campánula strícta. Exot. bot. non Linn. erécta. p. 326.				
13 capillaris. B.C. (bl.) capillary.	N. Holland.	1824.	— G. ♂.	Lodd. bot. cab. t. 1406.
14 littoralis. Lab.	(bl.) sea side.	V. Diem. Isl.	1820.	Lab. nov. holl. 1. t. 70.
15 quadrifida. A.DC. (bl.) four-cleft.	N. S. W.	1822.	— G. ♂.	
16 saxícola. A.DC. (bl.) small rock.	V. Diem. Isl.	1825.	5. 10. G. ♂.	
17 Siebérii. A.DC. (bl.) Sieber's.	N. S. W.	1826.	— G. ♂.	
18 undulata. A.DC. (bl.) waved.	C. B. S.	1822.	5. 7. G. ȝ.	
Campánula undulata. p. 326.				
19 hispídula. A.DC. (bl. wh.) hispid-calyxed.	—	1816.	6. 9. H. ⊙.	Comm. hort. 2. t. 37.
Campánula hispídula. p. 326.				
20 capillacea. A.DC. (wh.) capillary.	C. B. S.	1822.	5. 7. H. ȝ.	Alph. DC. camp. t. 17.
lobeloides. A.DC. (bh.) Lobelia-like.	Canaries.	1777.	7. 8. H. ⊙.	
péndula. p. 323.				

SYMPHIA'NDRA. A.DC. SYMPHIA'NDRA. Monadelphia Pentandria.

péndula. A.DC. (st.) pendulous.	Caucasus.	1824.	6. 8. H. ȝ.
Campánula péndula. p. 327. n. 102.			

SPECUL'ARIA. A.DC. VENUS-LOOKING-GLASS. Pentandria Monogynia.

1 Spéculum. A.DC. (bl.) common.	S. Europe.	1596.	5. 8. H. ⊙.	Bot. mag. t. 102.
Campánula Spéculum. B.M. Primatocárpus Spéculum. L'H. Lagôusia arvénensis.				Dur.
β álba. (wh.) white-flowered.	—	—	H. ⊙.	
2 hirsuta. A.DC. (bl.) hairy.	Italy.	1822.	5. 8. H. ⊙.	Ten. fl. nap. t. 19.
3 hy'brid'a. A.DC. (bl.) corn.	England.	— H. ⊙.	Eng. bot. v. 6. t. 375.
Campánula hy'brid'a. E.B. Primatocárpus hy'bridus. L'H.				
4 falcata. A.DC. (bl.) falcate.	Italy.	1822.	— H. ⊙.	Ten. fl. nap. t. 20.
5 pentagónia. A.DC. (bl.) large-flowered.	Turkey.	1686.	— H. ⊙.	Bot. reg. t. 56.
Campánula pentagónia. B.R.				
6 perfoliata. A.DC. (bl.) perfoliate.	N. America.	1680.	— H. ⊙.	Moris. s. 5. t. 2. f. 23.
MU'SCHIA. A.DC. MU'schia. Pentandria Monogynia.				
aùrea. A.DC. (ye.) golden-flowered.	Madeira.	1777.	7. 9. G. ȝ.	Bot. reg. t. 57.
Campánula aùrea. p. 326. n. 38.				
α latifòlia. A.DC. (ye.) broad-leaved.	—	—	G. ȝ.	Duham. ed. nov. t. 41.
β angustifòlia. A.DC. (ye.) narrow-leaved.	—	—	G. ȝ.	Jacq. schœn. 4. t. 472.
PETROMAR'ULA. A.DC. PETROMAR'ULA. Pentandria Monogynia.				
pinnata. A.DC. (bl.) winged-leaved.	Candia.	1640.	6. 8. F. ȝ.	Flor. græc. t. 220.
Phyteuma pinnatum. p. 328.				
JAS'IONE. p. 828. JAS'IONE. Monadelphia Pentandria.				
4 hùmilis. A.DC. (bl.) dwarf.	S. France.	1824.	6. 7. H. ȝ.	

ERICEÆ. p. 330.

ANDRO'MEDA. p. 331. ANDRO'MEDA.

23 tetragóna. s.s. (bh.) four-sided.	N. America.	1827. H. ȝ.	Pall. ross. 2. t. 73. f. 4.
--------------------------------------	-------------	-------	------------	-----------------------------

EBENACEÆ. p. 349.

DIO'SPYROS. p. 349. DATE PLUM.

19 Mabòla. B.R. (st.) Mabola-tree.	Philippines.	1822.	4. 5. S. ȝ.	Bot. reg. t. 1139.
------------------------------------	--------------	-------	-------------	--------------------

APOCINEÆ. p. 353.

PACHYP'ODIUM. B.R. **PACHYP'ODIUM.** Pentandria Monogynia.
succuléntum. (*wh.re.*) succulent. C. B. S. 1820. 4. 6. D.G. $\frac{1}{2}$. Bot. reg. t. 1321.
tuberòsum. B.R. *Echites succulenta.* Jacq. fragm. t. 117. see p. 355.

ASCLEPIADEÆ. p. 356.

ASCLEPIAS. p. 360. **SWALLOW-WORT.**
33 Greeniàna. N. Green's. N. America. 1828. 6. 8. H. $\frac{1}{2}$.

POLEMONIACEÆ. p. 367.

POLEM'ONIUM. p. 367. GREEK VALERIAN.
9 pulcherrimum. B.M. (*bl.*) bright blue. N. America. 1827. 7. 8. H. $\frac{1}{2}$. Bot. mag. t. 2979.
10 moschàtum. Gr. (*bl.*) musk-scented. — — — H. $\frac{1}{2}$.
GI'LIA. p. 368. **Gr'lia.**
3 pungens. B.M. (*bl.*) pungent-leaved. N. America. 1827. 6. 8. H. \odot . Bot. mag. t. 2977.
BONPLA'NDIA. Cav. **BONPLA'NDIA.** Pentandria Monogynia.
geminiflòra. Cav. (*bl.*) twin-flowered. New Spain. 1813. 5. 12. S. \odot . Bot. reg. t. 92.
Caldàsia heterophylla. w. non *Caldàsia*. DC.

BORAGINEÆ. p. 374.

ONO'SMA. p. 375.	ONO'SMA.					
16 Gmelini. L.A.	(<i>st.</i>) Gmelin's.	Altay.	1829.	4. 8. H. $\frac{1}{2}$.	Ledeb. ic. alt. t. 280.	
17 polyphyllum. L.A. (<i>st.</i>)	crowded-leav'd.	Tauria.	—	7. 8. H. $\frac{1}{2}$.	— t. 24.	
18 rigidum. L.A. (<i>pa.ye.</i>)	rigid.	—	1826.	— H. $\frac{1}{2}$.	— t. 238.	

HYDROPHYLLEÆ. p. 380.

E'UTOCA. p. 380. **E'UTOCA.**
3 sericea. B.M. (*bl.*) silky. N. America. 1827. 5. 7. H. $\frac{1}{2}$. Bot. mag. t. 3003.
To Franklinii, p. 380, add Bot. mag. t. 2985.

SOLANEÆ. p. 380.

CE'LSIA. p. 380.	CE'LSIA.					
9 heterophylla. P.S. (<i>ye.</i>) various-leaved.	1829.	7. 9. F. δ .			
SALPIGLÓ'SSIS. p. 383.	SALPIGLO'SSIS.					
5 intermédia.s. F.G. (<i>br.ye.</i>) intermediate.	Chile.	1829.	6. 9. F. $\frac{1}{2}$.	Swt. br. fl. gar. s. 2.		

SCROPHULARINÆ. p. 389.

SCROPHUL'ARIA. p. 391. FIGWORT.
51 lanàta. F. (*br.*) woolly. Siberia. 1828. 7. 11. H. $\frac{1}{2}$.
MI'MULUS. p. 394. MONKEY-FLOWER.
9 propinquus. B.R. (*ye.*) related. N. America. 1827. 4. 10. H. $\frac{1}{2}$. Bot. reg. t. 1330.
CALCEO'LARIA. p. 396. SLIPPERWORT.
15 Herbertiàna. B.R. (*ye.*) Mr. Herbert's. Chile. 1828. 5. 10. G. $\frac{1}{2}$. Bot. reg. t. 1313.

LABIATÆ. p. 399.

ST'ACHYS. p. 406.	HEDGE-NETTLE.					
49 sericea.	(<i>li.</i>) silky.	Nepaul.	1830.	7. 9. H. $\frac{1}{2}$.		
O'CYMMUM. p. 413.	BASIL.					
21 montanum. B.M. (<i>wh.</i>) mountain.	W. Indies.	1825.	5. 8. S. \odot .	Bot. mag. t. 2996.		
HY'PTIS. p. 414.	Hy'PTIS.					
14 madagascarién sis. (<i>bl.</i>) Madagascar.	Madagascar.	1828.	6. 8. S. δ .			

VERBENACEÆ. p. 415.

LANT'ANA. p. 417. LANT'ANA.
25 mollis. Gr. (ro.) soft. Mexico. 1826. 4. 8. G. h.

ACANTHACEÆ. p. 419.

RUE'LLIA. p. 420. RUE'LLIA.
28 picta. B.R. (bl.) painted. St. Domingo. 1826. 4. 8. S. h. Lodd. bot. cab. t. 1448.
JUSTI'CIA. p. 421. JUSTI'CIA.
60 guttata. B.R. (ye.sp.) spotted. E. Indies. 1828. 4. 8. S. ¼. Bot. reg. t. 1334.

PRIMULACEÆ. p. 423.

DODEC'ATHEON. p. 423. AMERICAN COWSLIP.
1 Mèadia. p. 423. Mead's. Virginia. 1744. 4. 6. H. ¼. Bot. mag. t. 12.
 α lilacinum. (li.) common lilac. — — — — — H. ¼. Trew Ehret. t. 12.
 β albiflorum. (wh.) white-flowered. 1824. — — — — — H. ¼. Lodd. bot. cab. t. 1489.
 γ elegans. S.F.G. (ro.) rose-coloured. 1827. 4. 7. H. ¼. Swt. br. fl. gar. s.2.t.60.
 δ giganteum. (li.) gigantic. — — — — — H. ¼.
2 integrifolium. (vi.) entire-leaved. N. America. 1829. — — — — — H. ¼. Pluk. alm. t. 79. f. 6.
ANDROS'ACE. p. 425. ANDROS'ACE.
17 carinata. Tor. (st.) keeled. N. America. 1827. 4. 7. H. ¼.
18 linearis. Gr. (bh.) linear-leaved. — — — — — H. ¼.

BEGONIACEÆ. p. 437.

BEG'ONIA. p. 437. BEG'ONIA.
40 longipes. B.M. (wh.) long-stalked. Mexico. 1828. 3. 8. S. h. Bot. mag. t. 3001.

POLYGONEÆ. p. 438.

POLYG'ONUM. p. 438. PERSICARIA.
61 injucundum. B.R. (st.) unattractive. 1825. 5. 6. G. h. Bot. reg. t. 1250.
RHE'UM. p. 440. RHUBARB.
15 leucorizum. F. (st.) white-rooted. Siberia. 1827. 5. 6. H. ¼.

AROIDEÆ. p. 479.

P'OTHOS. p. 479. P'OTHOS.
29 longifolia. R.S. (br.) long-leaved. Brazil. 1828. 9. 10. S. ¼.
30 reflexa. Ot. (br.) reflexed. — — — — — S. ¼.
31 reptans. Ot. (br.) creeping. — — — — — S. h.
32 rubescens. Ot. (br.) reddish. — — — — — S. ¼.

ORCHIDEÆ. p. 483.

APLE'CTRUM. N. APLE'CTRUM. Gynandria Monandria.
hyèmale. n. (br.) winter. N. America. 1827. H. ¼.
RODRIGU'EZIA. p. 487. RODRIGU'EZIA.
2 planifolia. H.C. (re.) flat-leaved. S. America. 1826. 6. 12. S. ¼.

CANNEÆ. p. 494.

PHRY'NIUM. p. 494. PHRY'NIUM.
28 coloratum. B.M. (or.) orange-coloured. Brazil. 1828. 4. 5. S. ¼. Bot. mag. t. 3010.
2 Q 2

MUSACEÆ. p. 495.

M'USA. p. 495. chinénsis.	PLANTAIN-TREE. (sc.) dwarf Chinese. China.	1829. S. 4.
------------------------------	---	-------	------------

IRIDEÆ. p. 495.

TRICHON'EMA. p. 503.	TRICHON'EMA.			
ramiflòra.	(pu.) branching-flower'd. Naples.	1830.	5. 6. H.b.	
<i>Ixia ramiflòra</i> , T.N.				
CR'OCUS. p. 503.	CR'OCUS.			
24 Thomàsii. T.N.	(bl.) Thomas's.	Naples.	1830. H.b.
25 odòrus. T.N. sweet-scented.	—	— H.b.
26 suavèolens. T.N. fragrant.	—	— H.b.
27 Imperàti. s.s.	(vi.) Imperati's.	—	—	2. 4. H.b.

Ten. mem. croc. t. 3.

AMARYLLIDEÆ. p. 505.

ZEPHYRA'NTHES. p. 505.	ZEPHYRA'NTHES.			
10 mesochlòa. w.h. (bh.) green-centred.	S. America.	1828.	6. 9. F.b.	
BRUNSVI'GIA. p. 510.	BRUNSWICK-LILY.			
7 grandiflòra. B.R. (pi.) large-flowered.	C. B. S.	1827.	8. 9. F.b.	
PHYCE'LLA. p. 510.	PHYCE'LLA.			
3 Herbertiàna. B.R. (re.y.) Mr. Herbert's. Chile.		1825.	5. 7. F.b.	
4 corúsca. Gr. (sc.or.) glossy.	—	—	—	F.b.
NARCI'SSUS. p. 514.	NARCI'SSUS.			
82 neglèctus. Ten. (wh.y.) neglected.	Naples.	1830.	4. 5. H.b.	

Bot. reg. t. 1335.

Bot. reg. t. 1341.

ASPHODELEÆ. p. 524.

SCI'LLA. p. 527.	SQUILL.			
30 odoràta. r.s.	(bl.) sweet-scented.	Portugal.	1818.	5. 6. H.b.
31 pratén sis. r.s.	(bl.) meadow.	Hungary.	1827.	— H.b.
32 ròsea, Leh.	(ro.) rose-coloured.	Numidia.	—	— H.b.
33 mauritànica. r.s. (bl.) Mauritanian.	Africa.	1819.	—	H.b.
DRI'MIA. p. 513.	DRI'MIA.			
14 villòsa, B.R. (g.st.) villous.	C. B. S.	1826.	5. 6. F.b.	
A'LLIU.M. p. 530.	GARLICK.			
118 euósmon. Ot. (br.) good-scented.	S. America.	1829.	5. 6. F.b.	
119 glandulòsum. Ot. (br.) glandular.	—	—	—	F.b.
120 nudicaulé. Leh. naked-stalked.	Altay.	—	H.b.
121 guttátum. G.D. (w.) spotted.	Odessa.	1819.	7. 8. H.b.	
122 Fischeri.	Fischer's.	Siberia.	1829. H.b.
123 acútum. s.s.	(li.) acute.	1819.	7. 8. H.b.
124 intermédia. G.D. (w.) intermediate.	S. Europe.	1827.	—	H.b.
125 fúscum. W.K.	(br.) brown-flowered. Hungary.	1820.	—	H.b.
126 juncifòlium. G.D. (w.) Rush-leaved.	Chile.	1826.	—	H.b.
127 striatéllum. G.D. (st.) striped.	—	1823.	—	H.b.
128 cinéreum. Leh. (st.) grey.	Siberia.	1829.	—	H.b.
MI'LLA. C.I.	MI'LLA.	Hexandria Monogynia.		
biflòra. c.i.	(wh.) two-flowered.	Mexico.	1826.	5. 6. F. 4.

W. et K. hung. t. 189.

Bot. reg. t. 1346.

W. et K. hung. t. 241.

Bot. mag. t. 2419.

Cavan. icon. 2. t. 196.

BROMELIACEÆ. p. 534.

POURR'ETIA. p. 535.	POURR'ETIA.			
4 cérùlea, Miers. (bl.) blue.	Chile.	1827.	4. 8. S. 4.	
5 rubricàulis. Mier. (bl.re.) red-stemmed.	—	—	—	S. 4.
6 magnispátha. Col. (g.w.) large-spathed.	S. America.	1829.	—	S. 4.
CARAGU'ATA. Plum.	CARAGU'ATA.	Hexandria Monogynia.		
lingulàta. Ly.	(ye.) tongue-leaved.	Jamaica.	1776.	6. 7. S. 4.
Tillàndsia lingulàta. p. 535.				

Coll. h. rip. t. 19.

Jacq. amer. t. 62.

BUONAPA'RTEA. p. 536. **BUONAPA'RTEA.**
2 grátilis. slender. Mexico. 1828. G. 4.

TULIPACEÆ. p. 536.

YU'CCA.	p. 536.	ADAM'S-NEEDLE.					
23 grátilis. Lk.	(wh.) slender.	Mexico.	1829.	G. 4.		
ERYTHR'ONIUM.	p. 539.	DOG'S-TOOTH VIOLET.					
Déns cànis. L.	(pu.) common.	Europe.	1596.	3. 4.	H.b.	Swt. br. fl. gar. s. 2.	
Déns cànis.	Redoute liliac. t. 193, non Bot. mag.						
4 bifida. s.F.G.	(ro.) cleft-stigma'd.	Switzerland.	—	H.b.	Swt. br. fl. gar. s. 2.	
β albiflora.	(wh.) white-flowered.	—	—	H.b.	—	s. 2.

MELANTHACEÆ. p. 539.

ZIGAD'ENUS. p. 540. **ZIGAD'ENUS.** Hexandria Trigynia.
3 élégans. Ph. (wh.) elegant. N. America. 1828. 5. 6. H. 4.

COMMELINEÆ. p. 541.

TRADESCA'NTIA.	p. 541.	SPIDERWORT.					
23 crásipes.	(bl.) thick-stalked.	Mexico.	1827.	5. 10.	H. 4.		
24 diurética. M.P.	(bl.) diuretic.	Brazil.	1825.	6. 8.	S. 4.		
COMMELINA.	p. 542.	COMMELINA.					
24 cucullátâ. H.C.	(bl.) hooded.	Brazil.	1825.	6. 9.	G. ⊙.		
25 caripénsis. H.S.	(bl.) rough-leaved.	Trinidad.	1826.	6. 8.	S. 4.		

RESTIACEÆ. p. 546.

ERIOCA'ULON. p. 546. PIPEWORT.
3 decangulaire. Ph. (u.h.) ten-angled. N. America. 1826. 8. 10. H.w. 4. Lodd. bot. cab. t. 1310.
4 fasciculátum. R.S. (wh.) fascicled. Guiana. 1825. — S.w. 4. Lam. enc. 3. t. 50. f. 3.

CYPERACEÆ. p. 547.

CA'REX.	p. 547.	SEDGE.					
123 rupéstris. s.s.	(br.) rock.	Switzerland.	1819.	6. 7.	H. 4.	All. ped. t. 92. f. 1.	
124 curvula. s.s.	(br.) curved.	S. Europe.	1824.	—	H. 4.	Schk. caric. t. D. f. 17.	
125 lobâta. w.	(br.) lobed.	Switzerland.	1819.	5. 7.	H. 4.	— t. D. f. 18.	
126 répens. s.s.	(br.) creeping.	—	1823.	6. 7.	H. 4.	— t. I. f. 11.	
127 leporina. w.	(br.) hare.	—	1819.	—	H. 4.	— t. Fff. f. 1. 29.	
128 lóngipes. D.P.	(br.) long-stalked.	Nepaul.	1823.	—	H. 4.		
129 microstâchya. s.s. (br.)	small-spiked.	N. Europe.	1822.	5. 6.	H. 4.	Schk. caric. t. 6. f. 11.	
130 mucronâta. s.s. (br.)	bristle-pointed.	S. Europe.	1819.	6. 7.	H. 4.	— t. K. f. 44.	
131 phaeostâchys. E.F. (br.)	brown-spiked.	Scotland.	6.	H. 4.		
132 stictocárpa. E.F. (br.)	dotted-fruited.	—	6. 7.	H. 4.		
133 angustifólia. E.F. (br.)	narrow-leaved.	—	—	H. 4.		
134 podocárpa. s.s. (br.)	foot-podded.	N. America.	1827.	—	H. 4.		
135 brachystâchys. s.s. (br.)	short-spiked.	S. Europe.	1819.	—	H. 4.	Schk. car. t. P. f. 58.	
SCH'ENUS.	p. 549.	BOG-RUSH.					
6 ferrugineus. R.S.	(br.) rusty.	Europe.	1829.	7. 8.	H.w. 4.	Host. gram. 4. t. 71.	
DICHROM'ENA.	R.S.	DICHROM'ENA.	Triandria Monogynia.				
leucocéphala. R.S. (wh.)	white-spiked.	W. Indies.	1822.	9. 12.	S.w. 4.		
Sch'enus stellátus. w.							

FIMBRI'STYLIS.

6 diphylla. R.S. (br.) two-leaved. E. Indies. 1822. 6. 8. S.w. 4.

ABILDGAA'RDIÀ.

R.S. ABILDGAA'RDIÀ.

2 tristâchya. R.S. (wh.) three-spiked. E. Indies. 1824. 6. 8. S. 4.

ISO'LEPIS.

p. 550. ISO'LEPIS.

13 supina. R.S. (br.) trailing. Europe. 1826. 7. 8. H.w. 4.

Host. gram. 3. t. 64.

2 Q 3

ADDENDA et CORRIGENDA.

SCI'RPU'S. p. 550.	CLUB-RUSH.							
19 elongátus. D.P. (br.) elongated.	Nepaul.	1825.	7. 9.	H.w. 4.				
20 articulátus. R.S. (br.) jointed.	E. Indies.	1826.	8. 10.	S.w. 4.	Rheed. mal. 12. t. 71.			
CYPERUS. p. 551.	CYPERUS.							
71 nilóticus. R.S. (br.) Nile.	Egypt.	1810.	8. 10.	H. 4.				
72 compáctus. R.S. (br. pu.) compact.	China.	1819.	7. 8.	H. 4.				
73 kyllingæoídes. R.S. (br.) Kyllinga-like.	N. America.	1828.	—	H. 4.				
74 dùbius. w. (br.) doubtful.	E. Indies.	1802.	7. 9.	S. 4.				
75 leucocéphalus. R.S. (wh.) white-headed.	—	1819.	—	S. 4.				
76 trisíucus. D.P. (br.) three-furrowed.	Nepaul.	1830.	—	H. 4.				
77 carícinus. D.P. (br.) Carex-like.	—	1826.	—	H. 4.				
78 rígidus. R.S. (re.wh.) rigid.	Madagascar.	—	—	S. 4.				
HYP'OE LYTRUM. R. HYP'OE LYTRUM.	Triandria Monogynia.							
argénteum. R.S. (wh. pu.) silvery.	E. Indies.	1824.	7. 8.	S. 4.				
MARI'SCUS. p. 552.	MARI'SCUS.							
9 paníceus. R.S. (br. st.) Panic-like.	E. Indies.	1820.	6. 7.	S.w. 4.	Rottb. gram. t. 4. f. 1.			
KYLL'INGA. p. 552.	KYLL'INGA.							
8 brevifólia. R.S. (gr.) short-leaved.	E. Indies.	1817.	5. 6.	S.w. 4.	Rottb. gram. t. 4. f. 3.			

FILICES. p. 576.

LYG'ODIUM. p. 577.	SNAKE'S-TONGUE.							
7 volúbile. s.s. (br.) twining.	W. Indies.	1819.	5. 9.	S. 4. —.	Sloan. jam. 1. t. 46. f. 1.			
8 hastátum. w. (br.) halberd-leaved.	Maranham.	1820.	—	S. 4. —.				
ANEIMIA. p. 577.	ANEIMIA.							
8 Phyllítidis. s.s. (br.) flat-leaved.	Trinidad.	1824.	5. 8.	S. 4. —.	Plum. fil. t. 156.			
SCHIZ'EA. p. 577.	SCHIZ'EA.							
5 élégans. s.s. (br.) elegant.	Trinidad.	1825.	5. 8.	S. 4.	Vahl. symb. 2. t. 50.			
GLEICH'ENIA. p. 577.	GLEICH'ENIA.							
6 pectinátata. s.s. (br.) pectinated.	S. America.	1824.	6. 8.	S. 4.				
DAN'EA. S.S.	DAN'EA.	Cryptogamia Poropterides.						
aláta. s.s. (br.) winged.	W. Indies.	1823.	6. 8.	S. 4.	Plum. fil. t. 109.			
ACRO'STICHUM. p. 577.	ACRO'STICHUM.							
9 longifólium. s.s. (br.) long-leaved.	W. Indies.	1827.	5. 10.	S. 4.				
10 glandulósom. H.F. (br.) glandular.	Jamaica.	1825.	—	S. 4.				
11 fimbriátum. s.s. (br.) fringed.	S. America.	1824.	—	S. 4.				
NOTHOL'ÆNA. p. 578.	NOTHOL'ÆNA.							
4 piloselloídes. s.s. (br. ye.) simple-leaved.	E. Indies.	1822.	6. 8.	S. 4.	Houtt. pflanz. t. 96. f. 2.			
MENI'SCIUM. p. 578.	MENI'SCIUM.							
4 prolíferum. s.s. (br.) prolíférous.	E. Indies.	1820.	5. 6.	S. 4.				
NIPHO'BOLUS. p. 578.	NIPHO'BOLUS.							
5 adnáscentis. s.s. (br.) woolly-leaved.	E. Indies.	1824.	4. 8.	S. 4.				
ASPL'ENIUM. p. 580.	SPLEENWORT.							
49 viviparum. H.K. (br.) viviparous.	Mauritius.	1820.	6. 9.	S. 4.				
50 bulbíferum. s.s. (br.) bulb-bearing.	N. Zealand.	—	—	G. 4.	Schk. crypt. t. 79.			
51 diffórmē. B.P.	(br.) two-formed.	N. Holland.	1823.	—	G. 4.			
52 Petrárchæ. nc.	(br.) Petrarch's.	France.	1819.	—	H. 4.			
53 obtusátum. s.s. (br.) obtuse.	N. Zealand.	1824.	—	F. 4.	H. et G. ic. fil. t. 152.			
54 cultrifólium. s.s. (br.) keen-leaved.	W. Indies.	1820.	—	S. 4.	Lab. nov. hol. 2. t. 242.			
55 brasiliénse. s.s. (br.) Brazilian.	Brazil.	1822.	—	S. 4.	Plum. fil. t. 3. f. 1.			
DOO'DIA. p. 581.	Doo'DIA.							
3 médiá. B.P.	(br.) middle-sized.	N. Holland.	1823.	3. 9.	G. 4.			
BLE'CHNUM. p. 581.	BLE'CHNUM.							
12 pectinátum. s.s. (br.) pectinate.	S. America.	1827.	5. 8.	S. 4.				
13 serrulátum. s.s. (br.) saw-leaved.	N. America.	1818.	—	H. 4.				
14 striátum. s.s. (br.) striated.	N. Holland.	1824.	—	F. 4.				
LOM'ARIA. p. 581.	LOM'ARIA.							
7 falcatá. s.s. (br.) falcate.	—	1823.	5. 8.	F. 4.	Lab. nov. hol. 2. t. 248.			
DIPLAZIUM. p. 582.	DIPL'AZIUM.							
10 castaneafólium. s.s. (br.) Chestnut-leaved.	Guinea.	1824.	8.	S. 4.				
PTERIS. p. 582.	BRAKE.							
28 sagittafólia. s.s. (br.) arrow-leaved.	Brazil.	1825.	6. 8.	S. 4.				
29 rotundifólia. s.s. (br.) round-leaved.	N. Zealand.	1824.	7. 8.	F. 4.				
30 élégans. s.s. (br.) elegant.	E. Indies.	—	8. 10.	S. 4.				
31 discolor. s.s. (br.) discoloured.	Brazil.	1825.	—	S. 4.				
32 heterophyl'a. s.s. (br.) various-leaved.	W. Indies.	—	—	S. 4.				
33 leptophyl'a. s.s. (br.) slender-leaved.	Brazil.	—	—	S. 4.				
34 chinénsis. L.C. (br.) Chinese.	China.	—	—	G. 4.	Plum. fil. t. 37.			

35 crenulata. L.C. (br.) crenulated.	1827.	7. 8. S. 4.	
36 Cervantèsii. H.L. (br.) Cervantes'. Mexico. 1824.	—	G. 4.	
ADIA'NTUM. p. 582. MAIDENHAIR.			
22 varium. s.s. (br.) variable. S. America. 1820.	7. 9.	S. 4.	
23 ternatum. s.s. (br.) ternate. ———	—	S. 4.	
24 patens. w. (br.) spreading. Brazil. 1824.	8. 10.	S. 4.	
25 rhomboideum. s.s. (br.) rhomboid. S. America. 1820.	6. 9.	S. 4.	
CHEILA'NTHES. p. 583. CHEILA'NTHES.			
14 caudata. B.P. (br.) tailed. N. Holland. 1824.	6. 8.	F. 4.	
LINDSÆA. p. 583. LINDSÆA.			
6 media. B.P. (br.) intermediate. N. Holland. 1823.	4. 6.	F. 4.	
DAVA'LLIA. p. 583. DAVA'LLIA.			
6 elegans. s.s. (br.) elegant. N. Holland. 1824.	4. 9.	G. 4.	Schk. crypt. t. 127.
7 dubia. B.P. (br.) doubtful. ——— 1826.	—	G. 4.	
8 concavatensis. L.C. (br.) Concavado. Brazil. 1823.	—	S. 4.	
CIB'OTIUM. S.S. CIB'OTIUM. Cryptogamia Filices.			
Billardieri. s.s. (br.) Labillardiere's. N. Holland. 1824.	G. h.	

VIOLARIEÆ. p. 43.

ANCHIE'TIA. Hil.	ANCHIE'TIA.	Pentandria Monogynia.	
pyrifolia. Hil. (wh.) Pear-leaved. Brazil.	1822.	S. h. — Mart. bras. t. 16.
Noisettia pyrifolia. Martius. see our p. 43.			

TILIACEÆ. p. 70

LUH'EA. W.	LUH'EA.	Polyadelphia Polyandria.	
paniculata. M.P. (ro.) paniced.	Brazil.	1828. S. h. Mart. bras. t. 62.

RUBIACEÆ. p. 260.

LIPO'STOMA. D.D.	LIPO'STOMA.	Tetrandria Monogynia.	
campanuliflora. D.D. (bl.) bell-flowered.	Brazil.	1825.	2. 10. S. 4. Bot. mag. t. 2840.
Hedyotis campanuliflora. p. 263. Eginetia capitata. Gr.			

PROTEACEÆ. p. 443.

SI'MSIA. B.P.	SI'MSIA.	Tetrandria Monogynia.	
anethifolia. B.P. (ye.) Dill-leaved.	N. Holland.	1825. G. h.
ANAD'ENIA. B.P.	ANAD'ENIA.	Tetrandria Monogynia.	
pulchella. B.P. (ro.) pretty.	N. Holland.	1824. G. h.
GUEVINA. Mol.	GUEVINA.	Tetrandria Monogynia.	
Avellana. L.T. (st.) variable-leaved. Chile.	1826. G. h.	Feuil. per. 3. t. 33.
Quádria heterophylla. Flor. per. 1. t. 99. f. b.			

NATURAL ORDER NOT DETERMINED.

BRE'XIA. B.R.	BRE'XIA.	Pentandria Monogynia.	
1 spinosa. B.R. (wh.) spiny-leaved.	Madagascar.	1815.	7. 8. S. h. Bot. reg. t. 872.
2 madagascariensis. (wh.) Madagascar.	—	—	S. h. — t. 730.
Venâna madagascariensis. Lamarck.			
3 chrysophylla. (wh.) golden-leaved.	—	1820. S. h.
LAUROPHY'LLUS. Th. LAUROPHY'LLUS.	Polygamia Dioccia.		
capensis. w. (gr.) Cape.	C. B. S.	1801. G. h.
PORANTH'ERA. L.T. PORANTH'ERA.	Pentandria Trigynia.		
ericifolia. L.T. (st.) Heath-leaved.	N. S. W.	1825.	5. 8. G. h. Linn. trans. 10.t. 22.f. 2

- PHLO'X.** p. 368. **LYCHNIDEA.** N. America. 1827. 5. 7. H. 4. Bot. reg. t. 1351.
36 speciosa. B.R. (ro.) showy.
- STERC'ULIA.** **STERC'ULIA.** Tragacantha. B.R. (st.) Tragacanth-tree. Sierra Leone. 1822. 5. 6. S. h. Bot. reg. t. 1353.
- SCI'LLA.** p. 527. **SQUILL.** 34 plumbbea. B.R. (bl.) lead-coloured. C. B. S.? 1812. 4. 6. F.b. Bot. reg. t. 1355.
- LOB'ELIA.** p. 322. **LOB'ELIA.** 85 Krausii. B.M. (sc.) Krauss's. St. Domingo. 1828. 1. 3. S. 4. Bot. mag. t. 3012.
- VANGUERIA.** p. 260. **VANGU'ERIA.** 3 velutina. B.M. (gr.) velvety. Madagascar. 1828. 5. 6. S. h. Bot. mag. t. 3014.
- BRACHYSTE'LMA.** p. 359. **BRACHYSTE'LMA.** 2 spathulatum. B.R. (br.) spatulate-leaved. C. B. S. 1824. 8. 10. D.G. 4. Bot. reg. t. 1113.
3 crispum. B.M. (br.y.) wave-leaved. — 1829. — D.G. 4. Bot. mag. t. 3016.
- CEROP'EGIA.** p. 359. **CEROP'EGIA.** 9 elegans. B.M. (pu.) elegant. E. Indies. 1828. 8. 10. S. 7. — Bot. mag. t. 3015.
- ENCY'CLIA.** p. 490. **ENCY'CLIA.** 2 patens. B.M. (br.) spreading-flowered. Brazil. 1829. 4. 8. S. 4. Bot. mag. t. 3013.
- Omitted at page 378.
- COLD'ENIA.** L. **COLD'ENIA.** Tetrandria Tetragynia. procumbens. L. (wh.) procumbent. E. Indies. 1699. 7. 8. S. ⊙. Lamarck. illus. t. 89.
- Page 75.
- G'ELA.** Lour. **G'ELA.** Octandria Monogynia lanceolata. B.C. (st.) spear-leaved. China. 1820. 4. 8. G. h. Lodd. bot. cab. t. 938.
Ximènia? lanceolata. DC. *Sèlas lanceolata*. s.s.
- Page 364.
- OPHIORRH'IZA.** L. **SNAKE-ROOT.** Mungos. L. (wh.) petiolate. E. Indies. 1820. 5. 8. S. 4.

INDEX GENERUM.

	Page.		Page.		Page.		Page.
<i>'Abies</i>	475	<i>Adámsia</i>	525	<i>Agrimónia</i>	192	<i>Alternanthera</i>	432
<i>ABIE' TINÆ</i>	474	<i>Adansónia</i>	65	<i>Agropyrum</i>	565	<i>Alternanthera</i>	432
<i>Abildgárdia</i>	550, 597	<i>Adélia</i>	459	<i>Agrostémma</i>	54	<i>Alth'æa</i>	60
<i>Abròma</i>	67	<i>Adélia</i>	460	<i>Agrostis</i>	556	<i>Alysicárpus</i>	152
<i>Abrònìa</i>	430	<i>Adenándra</i>	105	<i>Agrostis</i>	554, 555, 556,	<i>ALYSS'INEÆ</i>	24
<i>A'brus</i>	156	<i>Adenándra</i>	106	<i>Agrostis</i>	557, 559, 571	<i>Aly'ssum</i>	25
<i>Abùta</i>	16	<i>Adenanthèra</i>	164	<i>Agynéja</i>	458	<i>Aly'ssum</i>	24, 25, 31
<i>Abùtilon</i>	64	<i>Adenánthos</i>	446	<i>Ailánthus</i>	121	<i>Aly'xia</i>	353
<i>Acácia</i>	164, 589	<i>Adenárium</i>	57	<i>Aira</i>	559	<i>AMARANTHACEÆ</i>	
<i>Acácia</i>	162, 163	<i>Adenocárpus</i>	131	<i>Aira</i>	559, 560, 567,	<i>Amaránthus</i>	430
<i>Ac'æna</i>	192	<i>Adenóphora</i>	325	<i>Airópsis</i>	568,	<i>Amaránthus</i>	430
<i>Acálypha</i>	456	<i>Adenóstyles</i>	302	<i>Aitonia</i>	569, 571	<i>AMARYLLI'DEÆ</i>	
<i>ACANTHACEÆ</i>	419	<i>Adenotricha</i>	304	<i>Airópsis</i>	560	<i>Amaryllis</i>	505
<i>Acánthus</i>	419	<i>Adésmia</i>	149	<i>Ajuga</i>	560	<i>Amaryllis</i>	506
<i>Acárnà</i>	285	<i>Adiántum</i>	582, 599	<i>Ajuga</i>	560	<i>Amaryllis</i>	506
<i>'Acer</i>	82	<i>Adina</i>	263	<i>Ajuga</i>	560	<i>Amárobius</i>	417
<i>A'cer</i>	82	<i>Adlúmia</i>	20	<i>Ajuga</i>	560	<i>Amblírion</i>	538
<i>A'ceras</i>	483	<i>Adónis</i>	4	<i>Ajuga</i>	560	<i>Ambrosía</i>	481
<i>Acérates</i>	361	<i>Adóxa</i>	242	<i>Ajuga</i>	560	<i>Amelánchier</i>	177
<i>Acerátium</i>	71	<i>Adyséton</i>	25	<i>Ajuga</i>	560	<i>Amélus</i>	298
<i>ACERI'NEÆ</i>	82	<i>E'giceras</i>	348	<i>Ajuga</i>	560	<i>Amélus</i>	298
<i>Achania</i>	61	<i>E'gilops</i>	566	<i>ALANG'IEÆ</i>	206	<i>Amerimnum</i>	162
<i>Achillæa</i>	319	<i>E'ginétia</i>	599	<i>Alángium</i>	206	<i>Amerimnum</i>	162
<i>Achillæa</i>	316	<i>E'giphila</i>	416	<i>Albúca</i>	525	<i>Amethy'stea</i>	403
<i>Achimenes</i>	390	<i>E'gle</i>	75	<i>Alchímilla</i>	192	<i>Ammánbia</i>	201
<i>A'chlys</i>	584	<i>E'gopódium</i>	245	<i>Alcina</i>	313	<i>A'mmi</i>	245, 246, 248, 591
<i>Achnodónton</i>	554	<i>E'gopógon</i>	558	<i>Alde'a</i>	380	<i>AMM'INEÆ</i>	244
<i>A'chras</i>	349	<i>E'ollánthus</i>	414	<i>Aldrovánda</i>	46	<i>Anmóbius</i>	291
<i>Achyrránthes</i>	432	<i>E'ránthes</i>	491	<i>Alétris</i>	517	<i>Anmócharis</i>	510
<i>Achyrránthes</i>	431, 432	<i>Aèrides</i>	491	<i>Alétris</i>	517	<i>Ammyrsine</i>	344
	433	<i>E'rua</i>	432	<i>Aleurites</i>	458	<i>Amómum</i>	492
<i>Achyronìa</i>	127	<i>E'schynómene</i>	149	<i>Aleurites</i>	458	<i>Amómum</i>	492, 493
<i>Achyrophorus</i>	276	<i>E'schynómene</i>	143, 148,	<i>ALH'AGEÆ</i>	15	<i>Amòdora</i>	86
<i>Aciánthus</i>	486			<i>Allamánda</i>	354	<i>Amórpha</i>	142
<i>Acicárpha</i>	271	<i>E'sculus</i>	83	<i>Allantòdia</i>	580	<i>AMPELI'DEÆ</i>	86
<i>Acidóton</i>	456	<i>E'sculus</i>	83	<i>Alliária</i>	30	<i>Ampelópsis</i>	87
<i>Acilépis</i>	284	<i>E'theiléma</i>	420	<i>Alliònia</i>	430	<i>Amphérèphis</i>	288
<i>Aciótis</i>	204	<i>E'thionémâ</i>	32, 585	<i>Alliònia</i>	429	<i>Amphicárpa</i>	156
<i>'Acis</i>	516	<i>E'thónia</i>	275	<i>A'llium</i>	530, 596	<i>Amphilòphium</i>	366
<i>Acisanthèra</i>	202	<i>E'thusa</i>	247	<i>Allosòrus</i>	582	<i>Amphipógon</i>	555
<i>Acmadènia</i>	106	<i>E'thusa</i>	245	<i>A'línus</i>	468	<i>A'mphodus</i>	157
<i>Acmélla</i>	307	<i>Afzélia</i>	171	<i>A'loe</i>	518	<i>Amsónia</i>	355
<i>Acména</i>	212	<i>Afzélia</i>	86, 396	<i>A'loe</i>	518	<i>AMYDAL'INEÆ</i>	173
<i>Acnida</i>	436	<i>Agapánthus</i>	517	<i>Aloy'sia</i>	418	<i>Amy'dalus</i>	173
<i>Aconitum</i>	9, 584	<i>Agapánthus</i>	530	<i>Alpínia</i>	493	<i>AMYRI'DEÆ</i>	119
<i>A'corus</i>	480	<i>Agastachys</i>	446	<i>Alpínia</i>	493	<i>Am'yris</i>	119
<i>Acrocéphalus</i>	414	<i>Agasy'lis</i>	252	<i>Alpínia</i>	493	<i>Am'yris</i>	119
<i>Acrocomia</i>	543	<i>Agath'æa</i>	305	<i>Alpínia</i>	493	<i>Anábasis</i>	434
<i>Acroglöchin</i>	434	<i>A'gathis</i>	474	<i>Alsíne</i>	55, 56	<i>Anacámpseros</i>	219
<i>Acróstichum</i>	577, 598	<i>Agathélpis</i>	415	<i>Alpecúrus</i>	554	<i>Anacámpseros</i>	225
<i>Acróstichum</i>	578, 581	<i>Agathósma</i>	106	<i>Alpecúrus</i>	553, 557	<i>Anacámpsis</i>	483
<i>Acrótriche</i>	346	<i>Agathýrsus</i>	277	<i>Also'deia</i>	45	<i>Anacárdium</i>	117
<i>Act'æa</i>	11	<i>Agàti</i>	143	<i>ALSO'DINEÆ</i>	45	<i>Anacyclus</i>	318
<i>Act'æa</i>	11	<i>Agave</i>	534	<i>Alsóphila</i>	583	<i>Anadenia</i>	599
<i>Actinélla</i>	322	<i>Agave</i>	534	<i>Alsóphila</i>	579	<i>Anagállis</i>	425
<i>Actinocárpus</i>	477	<i>Agenòra</i>	276	<i>Alstónia</i>	355	<i>Anag yris</i>	122
<i>Actinóchloa</i>	571	<i>Agerátum</i>	301	<i>Alstreméria</i>	516	<i>Ananássia</i>	534
<i>Actinómeris</i>	309	<i>Agerátum</i>	301				
<i>Actinophyllum</i>	256	<i>Agláia</i>	75				
<i>Actinòtus</i>	243	<i>Agónis</i>	209				
<i>Ac'ynos</i>	409	<i>Agraulus</i>	556				
<i>Adámia</i>	591	<i>Agrimónia</i>	192				

INDEX GENERUM.

Page.	Page.	Page.	Page.	Page.	
Anántherix	361	Anthéricum	524	Arduína	353
Anarrhinum	390	Anthéricum	524, 525,	Areca	543
Anastática	28	Anthistíria	526	Areça	543
Anastática	28	Anthistíria	561	Aremónia	192
ANASTATI'CEÆ	28	Anthistíria	561	Arenária	56, 856
Anathérum	561	Anthocercis	380	Arenária	56, 57
Anchiétia	599	A'nthodon	79	Arethúsa	486
ANCHON'IEÆ	32	Antholòma	79	Aretíta	425
Anchùsa	377	Anthol'ya	500	Argània	349
Anchùsa	376, 377, 378	Anthol'ya	500	Argemône	19
Ancistrocárpus	433	Anthospermum	265	Argemône	19
Andersónia	345	Anthoxântum	553	Argyreia	373
Andira	168	Anthriscus	254, 591	Argyroch'æta	314
Andráchne	459	Anthriscus	254	Aristea	499
Andreóskia	29	Anthyl'lis	132, 589	Aristida	558
Androcýmbium	539	Anthyl'lis	128, 147, 152	Aristida	557, 558
Andromáchia	298	Antidésma	460	Aristolochia	451
Andrómeda	331, 593	ANTIDE'SMEÆ	460	Aristotélia	117
Andrómeda	331, 332	Antiphyl'la	241	Armeniaca	174
Andropôgon	561	Antirrhinum	390	Armèria	426, 427
Andropôgon	560, 561	Antirrhinum	390	Armoràcia	26
Androsâce	425	Antróphyum	582	A'rñica	299
Androsâce	425	Atótus	124	A'rñica	299
Andros'ænum	76	Apalatða	172	Arnopôgon	279
Andr'yala	274	Apárgia	274	Arnóseris	275
Andr'yala	274	Apárgia	275	AROI'DEÆ	475
Aneiléma	542	Apéiba	71	Arònìa	177
Aneíma	577, 598	Apñanes	192	Arpitum	591
Anemône	3	Apñanochilus	400	Arracácia	255
Anemône	3	Apñelândra	420	Arrhenatherum	562
Anesorhiza	247	Apñeléxis	293	Artabôtrys	15
Anéthum	251	Apñyllanthes	532	Artèdia	253
Anéthum	247	A'picra	521	Artemisia	316
Angélica	249	'Arios	157	Arthropodium	525
Angélica	248, 249, 251	'Arios	157	Arthrostémma	204
ANG'ELICEÆ	249	'Arium	245	Arthrostémma	203
Angelònia	394	Apłectrum	595	Artocárpus	462
Angiánthus	288	Aplophy'lum	105, 588	Arum	480
Angóphora	209	Apłuda	561	Arum	481
Angr'æcum	490	Apłuda	561	Arundinària	574
Anguillària	539	APOCI'NEÆ	353	Arundo	574
Angulòa	488	Apócynum	356	Arundo	555, 556, 559,
Angúria	216	Apócynum	361	567, 569, 574	
Anigozánthus	504	Aponogéton	478	Arungàna	76
Anisântus	500	AQUIFOL'ACEÆ	110	ASA'RINEÆ	451
Anisochilus	414	Aquilària	117	A'sarum	451
Anisòdus	383	AQUILARI'NEÆ	117	Ascaricida	289
Anísomeles	403	Aquilegia	8	ASCLEPI'ADEÆ	355
Anísomeles	407	A'rabis	22, 485	Asclépias	360, 394
Anisopétalum	488	ARAL'ACEÆ	21	Asclépias	360, 361
Anìsum	246, 247	A'rachis	167	A'scyrum	77
Anneslèa	17	A'rachis	149	Asimina	14
Anneslèa	163	Aràlia	256	Aspálathus	129
Annòna	14, 15	Aràlia	256, 257	Aspáragus	533
Anòda	63	ARAL'ACEÆ	256	Asperugo	377
Anòda	63	Araucària	474	Asperugo	377
Anodóntea	25	A'rbutus	331	Aspérula	266, 592
Anomathèca	499	A'rbutus	330	Aspérula	264
Anóna	14	Archangélica	249	ASPHOD'ELEÆ	524
ANON'ACEÆ	14	Archemôra	251	Asphodelus	524
Anópterus	239	A'rectium	280	Asphodelus	524
Anredêra	434	A'rectium	284	Aspidistra	480
Antennària	291	Arctopus	255	Aspídium	579
ANTHEMI'DEÆ	315	Arctostáphylos	330	Aspídium	579, 580
A'ñthemis	318	Arctothéca	312	Asplénium	580, 598
A'ñthemis	317, 318, 319,	Arctòtis	312	Asplénium	581, 582
.	321	Arctòtis	312, 313	Aspéllea	567
Anthéphora	563	Ardisia	348	Assónia	69

	Page.		Page.		Page.		Page.	
Balbisia	308	Beslèria	329	Botry'ceras	447	Buchnèra	394	
Ballòta	405	Bèta	436	Botry'chium	576	Buchnèra	394	
<i>Ballota</i>	405, 414	Betònica	406	<i>Bouteloua</i>	571	Buchólia	432	
Balsamína	101	BET'ULINEÆ	467	Bouvárdia	263	Bucìda	195	
BALSAMI'NEÆ	101	Bétula	467	Bowïëa	520	Búddlea	389	
Balsamíta	315	Beurrèria	379	Bowlésia	243	Buffónia	54	
Ba'samodéndron	120	Bídens	311	Brabéjum	446	Bulbine	525	
Baltimòra	314	<i>Bidens</i>	306	Brachyach'yris	298	Bulbocòdium	539	
Bambùsa	574	Bifòris	256	Brachyel'ytrum	559	Bulbocòdium	539	
<i>Bambusa</i>	574	Bigelòvia	265	Brachyl'æna	289	Bulliàrda	222	
Banistèria	81	Bignònia	365	Brachypòdium	573	Bumálida	109	
BANISTERI'EÆ	81	<i>Bignonia</i>	353, 365, 366	<i>Brachypodium</i>	572	Bumélia	348	
Banksia	447	BIGNONIACEÆ		Brachy'iris	298	Bunchòsia	81	
Bàphìa	162	Billardièra	48	Brachyséma	123	BUN'I ADEÆ	36	
Baptisia	123, 589	Billbèrgia	534	Brachystéma	359, 600	Bùniás	36	
<i>Baptisia</i>	122	<i>Billottia</i>	210	Bradleja	458	<i>Bünias</i>	28, 35	
Barbacènia	505	Biophytum	102	Brandésia	433	Bùniu'm	246	
Barbarèa	22	Biscutélla	27	Brasènia	17	<i>Bunium</i>	245, 246, 247,	
Barkhaùsia	278	<i>Biscutella</i>	28	Brassavòla	490	249		
Barlierà	420	Biserrula	148	Brásia	488	Buonapártea	536, 597	
Barnárdia	527	Bivon'æa	32	Brássica	33	<i>Buonapártea</i>	534	
Baròsma	105	Bíxa	40	<i>Brassica</i>	31, 34, 35	Búphone	510	
Barringtònia	213	BIXI'NEÆ	40	BRASSI'CEÆ	33	Buphtál'mum	309	
<i>Barringtònia</i>	213	Blackwéllia	116, 588	Bráya	22, 485	<i>Buphtál'mum</i>	309	
BARRINGTONI'EÆ	213	Bl'æría	332	Bréxia	599	Bupleúrum	246	
Bartholina	484	Bl'æría	332	Bridèlia	459	<i>Bupleúrum</i>	247, 251	
Bartònia	218	Bléchnum	581	Brignòlia	247	Burchárdia	539	
Bártzia	395	Bléchnum	581	Briza	570	Burchellia	262	
<i>Bártzia</i>	395	Bléchnum	420	<i>Briza</i>	570	Burglária	111	
Baryósma	168	Blépharis	419	Brodí'æa	517	Burmánnia	546	
Basélla	443	Blephília	404	<i>Brodíæa</i>	532	Bursària	48	
<i>Basélla</i>	434	Blètia	487	Bromgiártia	168	Burséra	120	
Bássia	349	Blighia	84	<i>Bromelia</i>	535	BURSERACEÆ	120	
Bastárdia	64	Blitum	436	<i>Bromélia</i>	535	Burtònia	124	
Bátschia	374	Blumenbächia	218	BROMELIACEÆ		Butea	161	
Bauèra	243	Bobártia	497			BUT'OMEÆ	477	
Bauhínia	172	Boccónia	19	Brómus	573	Butòmüs	477	
Beaufòrtia	207	<i>Boccónia</i>	19	Brósimum	462	BUUTNERIACEÆ		
Beaumóntia	356	Böebéra	305	Bross'æa	332			
Beckmánnia	564	Böehmèria	462	Brotéra	288	BUTTN'ERIEÆ	66	
Béscia	345	Bæóbótrys	347	<i>Bròtera</i>	69, 287, 414	Búxus	456	
Befària	345	Boerhaàvia	430	Broughtònia	490	<i>Búxus</i>	472	
Begònìa	437, 595	Boerháavia	430	Broussonètia	462	B'yblis	46	
BEGONTACEÆ	437	Boldòa	430	<i>Broussonètia</i>	122, 462	Byrsónima	80	
Behrínia	274	Boltònia	298	Browállia	394	Bystropògon	401	
<i>Bejària</i>	345	BOMB'ACEÆ	65	Brównia	168	Cabómba	17	
Bélis	475	Bómbar	66	Brownlowia	71	Cabréra	568	
Belladónnna	506	Bómbar	66	Brúcea	121	Cacàlia	302	
Bellevalìa	528	Bomplàndia	594	<i>Brúcea</i>	121	<i>Cacàlia</i>	302, 303	
Bellidiástrum	299	Bontìa	419	Brugmánsia	383	Cächrys	255	
<i>Bellínia</i>	383	BORAGI'NEÆ	375	Brunellìa	121	<i>Cächrys</i>	244, 249, 252,	
Béllis	299	Boràgo	378	Brunfelsia	389	255		
Bélliùm	299	Boràgo	378	Brúnia	115	CA'CTEÆ	235	
Benincàsà	214	Boràssus	544	<i>Brúnia</i>	115	Cáctus	235, 236, 237, 238	
Bérardia	116	Borbònìa	126	BRUNTACEÆ	115	Cacoùcia	196	
Bérardia	284	<i>Borbònìa</i>	126, 127	Brunníchia	438	Càdia	169	
BERBERI'DEÆ	16	Borkhousènia	389	Brunsvigia	510, 596	Cælestina	301	
Bérberis	16, 584	Borònia	107	<i>Brunsvigia</i>	510	Cænopteris	581	
Berchémia	112	Bòrya	460	B'r'ya	162	Cæsalpínia	168	
Berkheyà	312	Bòscia	38	Bryonia	214	<i>Cæsalpínia</i>	168, 169	
Bergènia	289	Bòsea	436	<i>Bryònia</i>	214	CEASALPI'NEÆ	167	
Bergèra	75	Bossi'æa	126	Bryophyllum	224	C'asia	525	
Bérgia	55	Boswéllia	120	Búbon	251	Cæsulia	288	
Beringèria	405			Búbon	247, 248, 250	Cajànus	159	
Bernhärdia	576			Bubròma	67	Cakile	28	
Bérrya	71			Búcco	106	<i>Cakile</i>	36	
Berteròa	24			Buchanània	117	CAKILI'NEÆ	28	
Berthollètia	213							
Berzèlia	115							

INDEX GENERUM.

Page.	Page.	Page.	Page.	Page.	
Caladènia	486	CAMELLI'EAE ..	73	Carrichtèra	35
Calàdium	481	Camerària	355	Cárthamus	284
Calamagròstis ..	555	Camerària	354, 355	Cárthamus	284
Calamína	561	Campánula	324	Cartonèma	542
Calamíntha	409	Campánula 324, 328, 593		Càrum	245
Calámpelis	366	CAMPANULACEAE ..		Càrum	248
Cálamus	544 324, 592		Caruncularia	358
Calandrínia	219	Campèlia	541	Cárya	465
Calánthe	490	Camphorósma	437	Caryòcar	83
Calàthea	494	Campomanésia	210	CARYOPHYLLEAE ..	
Calceolària	396, 494	CAMPYLOSPERMÆ 48	
Caldàsia	369, 594 253		Caryophyllus	212
Càlea	307	Campy'lia	90	Caryòta	544
Càlea	289, 306, 307	Canarina	324	Caseària	116
Caleácte	307	Canàrium	120	Cássia	169, 589
Caléndula	313, 592	Canavàlia	159	CASSI'EAE ..	168
Calepina	35	Candóllea	13	Cassine	110
Caleàna	487	Canellà	79	Cassinia	291
Caleÿa	487	Cànnà	494	Cassinia	288
Càlla	480	Cànnabis	463	CASSUVITÆ ..	117
Càlla	480	CA'NNEAE ..	494	Cassuvium	117
Calicárpa	416	Cànthium	261	Cass'ytha	441
Calícoma	243	Càntua	368	Castàlia	18
Calligonus	440	Caperònìa	456	Catànæa	467
Calliòpèa	275	Capnophyllum	251	Castanocàrpus ..	589
Calliòpsis	311	CAPPARI'DEAE ..	37	Castillèja	395
Calisàce	251	CAPPAR'EAE ..	38	Casuarina	472
Callisia	542	Càpparis	38, 585	Catabòrsa	559
Callistàchys	123	Càpparis	38, 39	Catálpa	365
Callistèma	297	Caprària	389	Catanànche	280
Callistèmon	209	Caprària	389	Catàppa	195
Callitrichè	201	CAPRIFOLITACEAE ..		Catasétum	488
CALLITRICH'I'NEAE ..	201	Caprifòlium	258	Catesb'aea	262
Càllitris	473	Capsélla	27	Cathartocàrpus ..	169
Càlluna	332	Càpsicum	384	Cattlèya	490
Calochilus	485	Caragàna	144	Catùrus	462
Calochòrtus	538	Caraguàta	596	CAUCALI'NEAE ..	253
Calodèndron	105	Caràlia	197	Caúcalis	253
Calóphaca	144	Carallùma	359	Caúcalis	253, 254
CALOPHY'LLEAE ..	78	Caràpa	86	Caulophyllum	16
Calophy'llum ..	78	Caràpa	264	Ceanòthus	113
Calopògon	487	Carapichea	264	Cecropia	464
Calòrophus	547	Cardámíne	23	Cedrela	86
Calostèmma	513	Cardámíne	23, 24	CEDR'EAE ..	86
Calothàmnus	207	Cardiospérnum ..	83	Cèdrus	475
Calòtis	294	Cardòpatum	287	CELASTR'INEAE ..	109
Calòtropis	361	Carduncéllus	284	Celàstrus	110
Cátha	7, 584	Càrduu	281	Celàstrus	110
CALYCA'NTHEAE ..		Càrduu	282, 283	Celàstrus	110
..... 194		CARDU'ACEAE ..	280	Celàstrus	110
Calycànthus	194	Càrex	547, 597	Celòsia	431
Calycànthus	195	Càrex	552	Celòsia	431, 433
CALYCEREA ..	271	Carèya	213	Célisia	380, 594
CALYCIFL'ORAE ..	109	Cargillia	349	Célisia	394
Calycòtome	131	Càrica	216	Céltis	464
Calymènia	429	Caríssa	353	Céltis	464
Caly'psò	487	Caríssa	353	Cénchrus	552
Calyprànthes	211	Carlina	284	Cénchrus 553, 557, 563,	
Calyprànthes	211, 212	Carlina	281 567, 568	
Caly'ptrion	43	Carlowízia	284	Cènia	318
Calystègia	370	Carludòvica	480	Centolòphium	248
Calythrix	206	Carmich'ælia	139	Centaureà	285, 592
Camaridiúm	489	Carolínia	66	Centaureà 281, 285, 287	
Cambessèdia	117	Carpèsium	290	Centothèca	568
Camelinà	31	Càrpinus	467	Centrànthus	268
CAMELLI'NEAE ..	31	Carpodòntus	256	Centrocàrpha	310
Caméllia	73	Carpodòntus	77	Centrolèpis	547
Caméllia	73	Carpogògon	159	Centrospérnum ..	321

Centúnculus	426	Cephaëlis	264	CEPHALANTHEAE ..	263
Cephalòphora	315	Cephalòtus	192	Cephalanthus	263
Cephalòtus	192	Ceranthèra	45	Cephalària	270
Ceranthèra	45	Cerastium	57	Cephalòphora	315
Cerastium	57	Cerastium	54, 55	Cerastium	54, 55
Cerastium	54, 55	Cérasus	174	Cérasus	174
Cératòla	460	Ceratiola	460	Cératòpterus	577
Ceratiola	460	Ceratocàrpus	436	Ceratosànthes	215
Ceratocàrpus	436	Ceratocéphalus	4	Ceratospérnum	437
Ceratocéphalus	4	Ceratochilus	490	Cérbera	354
Ceratochilus	490	Ceratòchloa	571	Cérbera	354
Ceratòchloa	571	Ceratònia	169	CERCODIANAE ..	200
Ceratònia	169	Ceratopétalum	243	CÉRCODIANAE ..	200
Ceratopétalum	243	CERATOPHYLLEAE ..		CÉRATOPHYLLEAE ..	201
CERATOPHYLLEAE 201		Ceratophyllum	201
Ceratophyllum	201	Ceratópterus	577	Ceratophyllum	201
Ceratópterus	577	Ceratosànthes	215	Ceratospérnum	437
Ceratosànthes	215	Ceratospérnum	437	Cérès	554
Ceratospérnum	437	Cérès	554	Cérès	554
Cérès	554	Cérès	590	Cérès	590
Cérès	590	Cérès	591	Cérès	591
Cérès	591	Cérinthe	375	Ceropègia	359, 600
Cérinthe	375	Ceropègia	359, 600	Cestrinus	284
Cercocàrpus	193	Cestrinus	284	Céstrum	388
Cercocàrpus	193	Cestrinus	284	Céterach	578
Cercòdia	201	Cestrinus	284	Chærophyllum	254
CERCODIANAE ..	200	Cestrinus	284	Chærophyllum	254, 255
CERCODIANAE ..	200	Cestrinus	284 591	
CERCODIANAE ..	200	Cestrinus	284	Chætanàthera	280
Chætanàthera	280	Cestrinus	284	Chætarìa	557
Chætarìa	557	Cestrinus	284	Chætocàlyx	141
Chætarìa	557	Cestrinus	284	Chætogàstra	204
Chætocàlyx	141	Cestrinus	284	Chætony'chia	220
Chætogàstra	204	Cestrinus	284	Chætospora	549
Chætony'chia	220	Cestrinus	284	Chætòrus	555
Chætospora	549	Cestrinus	284	Chaillettia	117
Chætòrus	555	Cestrinus	284	CHAILLETTIA	117
Chaillettia	117	Cestrinus	284	Chaiturus	407
CHAILLETTIA	117	Cestrinus	284	Chamædòrea	542
Chaiturus	407	Cestrinus	284	CHAMÆLAUCIEAE ..	542
Chaiturus	407	Cestrinus	284	Chamælaùcium	207
Chamædòrea	542	Cestrinus	284	Chamælaùcium	207
CHAMÆLAUCIEAE ..	542	Cestrinus	284	Chamælírium	540
Chamædòrea	542	Cestrinus	284	Chamænérion	197
Chamælaùcium	207	Cestrinus	284	Chamærèpes	483
Chamælaùcium	207	Cestrinus	284	Chamærops	542
Chamælírium	540	Cestrinus	284	Chamærops	542
Chamænérion	197	Cestrinus	284	Chamærops	542
Chamærèpes	483	Cestrinus	284	Chamagòròtis	557
Chamærops	542	Cestrinus	284	Chamagòròtis	557
Chamærops	542	Cestrinus	284	Chamissòa	432
Chamagòròtis	557	Cestrinus	284	Chamórchis	483
Chamagòròtis	557	Cestrinus	284	Chaptàlia	280
Chamissòa	432	Cestrinus	284	Chaptàlia	280
Chamórchis	483	Cestrinus	284	Chàra	575
Chaptàlia	280	Cestrinus	284	Chàra	575

Page.	Page.	Page.	Page.
<i>Chàra</i> 575	<i>Cibòtium</i> 599	<i>COBÆ'ACEÆ</i> 367	<i>CòniuM</i> 247, 251, 255
<i>CHAR'ACEÆ</i> 575	<i>Cicèa</i> 456	<i>Cobúrgia</i> 513	<i>CONNAR'ACEÆ</i>
<i>Chardinia</i> 294	<i>Cicéndia</i> 364	<i>Coccocy'pselum</i> 262 121
<i>Chariloódia</i> 533	<i>Cicer</i> 153	<i>Coccoey'pselum</i> 262	<i>Connàrus</i> 121
<i>Chasmónia</i> 405	<i>Cicer</i> 154	<i>Coccóloba</i> 438	<i>Conocárpus</i> 196
<i>Cheilánthes</i> .. 583, 599	<i>CICHOR'ACEÆ</i> 271	<i>Cócculus</i> 15	<i>Conocéphalus</i> 462
<i>Cheiránthus</i> 21	<i>Cichórium</i> 279	<i>Cócculus</i> 15	<i>Conospérnum</i> 446
<i>Cheiránthus</i> 22, 29, 30	<i>Cicónium</i> 100	<i>Cochleária</i> 26	<i>Conostégia</i> 205
..... 31, 32, 37	<i>Cicúta</i> 244	<i>Cochleária</i> 26, 31	<i>Conóstylis</i> 504
<i>Cheirínia</i> .. 29, 30, 31	<i>Ciliária</i> 240	<i>Cochlospérmum</i> 73	<i>Conringia</i> 31
<i>Cheirostémón</i> 66	<i>Cimicifuga</i> 11	<i>Cócos</i> 544	<i>Convallária</i> 523
<i>Chelidónium</i> 19	<i>CINCH'ONEÆ</i> 262	<i>Cócos</i> 543, 544	<i>Convallária</i> 523
<i>Chelidónium</i> 19	<i>Cinchóna</i> .. 260, 262	<i>Codárium</i> 172	<i>CONVOLVUL'A-</i>
<i>Chelône</i> 393	<i>Cineraria</i> 304	<i>Còdon</i> 383	CEÆ 369
<i>Chenôlea</i> 434	<i>Cineraria</i> .. 304, 305	<i>Codonóphora</i> 329	<i>Convólvulus</i> 369
<i>CHENOP'ODEÆ</i> 433	<i>Cína</i> 557	<i>Cœlogyne</i> 488	<i>Convólculus</i> 370, 371,
<i>Chenopódium</i> .. 434	<i>Cinnamòdum</i> 441	<i>CÉLOSPÉ'RMEÆ</i> 256 372, 373
<i>Chenopódium</i> 431, 434	<i>Circæa</i> 200	<i>Coffea</i> 264	<i>Con'zya</i> 289
<i>Cherleria</i> 57	<i>CIRCÆ'EÆ</i> 200	<i>Coffea</i> 264	<i>Con'zya</i> 289, 294
<i>Chilóchloa</i> 554	<i>Cirrh'aea</i> 488	<i>COFFE'ACEÆ</i> 263	<i>Coókia</i> 75
<i>Chilóchloa</i> 554	<i>Cirsium</i> 282	<i>Còix</i> 574	<i>Copaféra</i> 171
<i>Chiloglottis</i> 486	<i>Cissámpelos</i> 15	<i>Colbértia</i> 13	<i>Cóptis</i> 7, 584
<i>Chilópsis</i> 365	<i>Cissámpelos</i> 16	<i>Cólichicum</i> 539	<i>Corallorrhiza</i> 487
<i>Chimáphila</i> 341	<i>Cissus</i> 86	<i>Cólichicum</i> 539	<i>Corbulària</i> 514
<i>Chimonánthus</i> .. 195	<i>Císsus</i> 87	<i>Coldènia</i> 600	<i>Córchorus</i> 70
<i>Chiococca</i> 264	<i>CISTI'NEÆ</i> 40	<i>Coleánthus</i> 554	<i>Córchorus</i> 194
<i>Chionánthus</i> 351	<i>Cistus</i> 40	<i>Colebroókia</i> 400	<i>Córdia</i> 379
<i>Chiranthodéndron</i> 66	<i>Cistus</i> 41, 42	<i>Coleonémä</i> 106	<i>Córdia</i> 380
<i>Chirónia</i> 363	<i>Citharéxylum</i> .. 417	<i>Coleus</i> 413	<i>Cordylíne</i> 534
<i>Chlamy'sporum</i> .. 532	<i>Citrus</i> 75	<i>Coll'aea</i> 140	<i>Cordylocárpus</i> .. 36
<i>CHLEN'ACEÆ</i> 72	<i>Cladánthus</i> .. 319, 592	<i>Collétia</i> 113	<i>Corema</i> 460
<i>Chlidánthus</i> 513	<i>Cládium</i> 550	<i>Collétia</i> 113	<i>Coreópsis</i> 311
<i>Chloánthes</i> 417	<i>Cladóstachys</i> 91	<i>Collínisia</i> 390	<i>Coreópsis</i> 307, 309, 311
<i>Chlòra</i> 363	<i>Clárkia</i> 199	<i>Collinsónia</i> 412	<i>CORIA'NDREÆ</i> .. 256
..... 259	<i>Clausénia</i> 75	<i>Collòmia</i> 368	<i>Coriándrum</i> 256
<i>CHLORA'NTHEÆ</i>	<i>Clavija</i> 353	<i>Colobáchné</i> 557	<i>Coriándrum</i> 256
<i>Chloránthus</i> 259	<i>Claytonia</i> 220	<i>Cologânia</i> 140	<i>CORIARI'EÆ</i> .. 109
<i>Chloráster</i> 516	<i>CLEMATI'DEÆ</i> .. 1	<i>Colophónia</i> 120	<i>Coriària</i> 109
<i>Chlòris</i> 562	<i>Clématis</i> 1, 584	<i>Colopídium</i> 557	<i>Còrion</i> 256
<i>Chlòris</i> 562, 571	<i>Cléome</i> 37, 585	<i>Colubrìna</i> 113	<i>Còris</i> 425
<i>Chloróphytum</i> 525	<i>Cléome</i> .. 31, 37, 38	<i>Columélia</i> 308	<i>Corispérnum</i> 436
<i>Chloroxýlon</i> 86	<i>CLE'OMEÆ</i> 37	<i>Columnéa</i> 390	<i>Cornucòpiæ</i> 552
<i>Chomélia</i> 264	<i>Cleoñia</i> 410	<i>Cùlùria</i> 189	<i>Córnus</i> 257
<i>Chondrilla</i> 277	<i>Clerodéndrum</i> .. 415	<i>Colùtea</i> 144	<i>Cornùtia</i> 417
<i>Chondròsea</i> 240	<i>Cléthora</i> 332	<i>Colùtea</i> 144	<i>Cornùtia</i> 417
<i>Chondròsum</i> 571	<i>Cleyéra</i> 72	<i>Comándra</i> 450	<i>COROLLIF'OREÆ</i>
<i>Chorísmo</i> 274	<i>Clidémia</i> 205	<i>Comarópsis</i> 189 347
<i>Chorispora</i> 28	<i>Clifortia</i> 193	<i>Cómárum</i> 192	<i>Coronilla</i> 148
<i>Chorizéma</i> 123	<i>Clifontia</i> 332	<i>COMBRET'ACEÆ</i>	<i>Coronilla</i> 143
<i>Chorozéma</i> 123	<i>Clintonia</i> 409 195	<i>CORONIL'LEÆ</i> .. 148
<i>Chrístia</i> 149	<i>Clinopódium</i> 324	<i>Combrétum</i> 196	<i>Corónopus</i> 31
<i>Chrysanthéllum</i> .. 308	<i>Clintonia</i> 324	<i>Combrétum</i> 196, 590	<i>Corréa</i> 107
<i>Chrysánthemum</i> 320	<i>Clítòria</i> 140	<i>Comespérma</i> 47	<i>Corrigiola</i> 220
<i>Chrysánthemum</i> 321,	<i>Clítòria</i> 140	<i>Commeliná</i> .. 542, 597	<i>Cortùsa</i> 423
..... 322	<i>CLIT'ORIÆ</i> 138	<i>Commeliná</i> .. 541, 542	<i>Córybas</i> 487
<i>Chrysiphala</i> 513	<i>Clívia</i> 516	<i>COMMEL'INEÆ</i> .. 541	<i>Corycéarpus</i> 571
<i>CHRYSOBAL'A-</i>	<i>Clúsia</i> 78	<i>Commersónia</i> 67	<i>Cory'ciúm</i> 485
<i>NEÆ</i> 173	<i>CLUSI'EÆ</i> 78	<i>Commersónia</i> 67	<i>Corydális</i> 20
<i>Chrysobálanus</i> .. 173	<i>Cluy'tia</i> 459	<i>Comocládia</i> 119	<i>Córylus</i> 467
<i>Chrysócoma</i> 298	<i>Clypéola</i> 25	<i>ComPO'SITÆ</i> .. 271	<i>Corynélia</i> 144
<i>Chrysócoma</i> .. 291, 298	<i>Cnemidóstachys</i> .. 460	<i>Comptónia</i> 472	<i>Coryn'éphorus</i> 560
<i>Chrysophyllum</i> .. 349	<i>Cneòrum</i> 121	<i>Conanthera</i> 532	<i>Córypha</i> 543
<i>Chrysopogón</i> 561	<i>Cnéstis</i> 121	<i>Cónchium</i> 447	<i>Córypha</i> 542, 543
<i>Chrysóps</i> 291	<i>Cnicus</i> .. 282, 283, 287	<i>Condália</i> 112	<i>Corysánthes</i> 487
<i>Chrysópsis</i> 295	<i>Cnídium</i> 248	<i>Condylocárpus</i> 252	<i>Coscínum</i> 15
<i>ChrysospléniuM</i> .. 242	<i>Cnídium</i> .. 248, 249	<i>CONIFERÆ</i> .. 473	<i>Cósmos</i> 310
<i>Chrysúrus</i> 562	<i>Cnidósculus</i> 457	<i>Conioselinum</i> 249	<i>Cósmea</i> 311
<i>Chuquirága</i> .. 280, 592	<i>Cob'aea</i> 367	<i>Cònium</i> 255	<i>Cosmélia</i> 345

	Page.		Page.		Page.		Page.	
Cossignia	84	CUCURBITACEÆ	213	Cyphia	324	Deschampsia	559	
Cóstus	492	Cullumia	312	Cypripedium	491	Desmánthus	163	
Cotoneáster	176	CUMI'NEÆ	252	Cyrilla	332	Desmochæta	432	
Cótula	318	Cumínium	252	Cyrilla	390	Desmochæta	432	
Cótula	318	Cummíngia	352	Cyrtánthus	512	Desmódiuim	149	
Cotylédon	224	Cùnila	404	Cyrtánthus	512	Desmódiuim	149	
Cotylédon	225	Cùnila	404	Cyrtopodium	488	Desmóncus	544	
Coulteria	168	Cunninghamia	475	Cyrtostylis	486	DET'ARIEÆ	173	
Coumaroúna	168	Cunónia	243	Cysticápnos	20	Detarium	173	
Courouúpita	213	CUNONIACEÆ	242	CYTINÆ	452	Devaúxia	547	
Courséria	143	Cupánia	84	Cytisus	130	Devérra	247	
Coutaria	262	Cùpheia	202	Cytisus	123, 126, 131,	Diálium	172	
Coutouúbea	364	CUPRE'SSINEÆ	473 143, 144, 159	Dianélia	532		
Crámbe	35	Cupréssus	474	Czákia	524	Diánthus	49, 586	
Craniolària	367	Cupréssus	474	Dacry'dium	473	Diapénsia	342	
Craniótome	407	CUPULIFERÆ	465	Dactylanthes	453	Diapénsia	342	
Crássula	222	Curatélla	13	Dáctylis	570	Diarrhëna	571	
Crássula 222, 223, 224,	226	Curcúlico	505	Dáctylis	553, 574	Diászia	394	
CRASSULACEÆ	222	Curtísia	491	Dactylocténium	571	Diásia	502	
Cratægus	175, 176, 177	Curtógyne	223	Dáhlia	310	Diastella	444	
Cratæva	38	Curtopogon	558	D'aemia	361	Dicéra	72	
Creodus	259	Cuscuta	373	Dáis	449	Dicérlma	151	
Crepidària	452	CUSP'ARIEÆ	108	Dalbergia	161	Dichilus	128	
Crépis	273, 592	Cussónia	256	Dalbergia	142	Dichóndra	373	
Crèpis	273, 274, 275,	Cyámus	17	DALBERGL'IEÆ	161	Dichorisándra	126	
..... 276, 278, 279		Cyanélia	532	Dàlea	141	Dichromèna	597	
Crescéntia	389	Cyanotis	542	Dàlea	141	Dicksónia	583	
Crescéntia	389	Cyànum	286	Dalechámpia	456	Dicksónia	583	
Créssa	373	Cyat'hèa	583	Dalibárda	189	Diclíptera	422	
Crinum	511	Cyat'hèa	579	Dalibárda	189	Diclytra	20	
Crinum	513, 532	Cyathôdes	346	Dalrymplea	109	DICOTYLED'ONÆ		
Cristária	63	CYC'ADEÆ	476	Dalrymplea	109	NEÆ	1	
Críthmum	249	Cycas	476	Damasónium	476	Dictamnus	105, 588	
Críthmum	251	Cyclamen	423	Damasónium	477	Didélta	312	
Critónia	299	Cyclánthus	400	Dámmara	474	Didiscus	243	
Cròcus	503, 596	Cyclobóthra	538	Dampière	325	DIDYMOCA'RPE		
Crossândra	420	Cyclòpia	123	Danæa	598	367		
Crotalària	127	Cydónia	178	Danthònia	567	Didymocárpus	367	
Crotalària 123, 126, 128		Cylipògon	141	Danthònia	567	Diervílla	258	
Crótón	457	Cylísta	160	Daphné	448	Diètes	497	
Cròton	456, 457, 458	Cymátion	549	Daréa	581	Digitális	392	
Crówea	107	Cymbáchna	561	Darlingtònia	163	Digitaria	558	
Crozophora	457	Cymbária	390	Darwinia	207	Digitaria	559	
Crucianélia	265	Cymbidium	488	Dasytémon	222	Digráphis	559	
CRUCIFERÆ	21	Cymbidium 487, 488, 489		Datisca	39	Didálris	504	
Crùdya	172	Cymbopogon	561	DATISCEÆ	39	Dilatris	504	
Crupina	285	Cymbùrus	418	Datûra	383	Dilepyrum	559	
Cry'psis	557	Cymínosma	107	Datûra	383	Dillénia	18	
Cryptándra	114	Cymópterus	252	Datûra	383	Dillénia	18	
Cryptarrhèna	490	Cynánchezum	361	Daubentònia	144	DILLENIACEÆ	12	
Cryptocárta	441	Cynánchezum 360, 361,	362	DAUCI'NEÆ	253	Dillwy'nia	124	
CRYPTOGAMEÆ	575 361	362	Davallia	583, 599	Dillwy'nia	124	
Cryptográmma	582	Cynoglossum	378	Davílesia	125	Dimácría	91	
Cryptólepis	355	Cynoglossum	378	Davílesia	124	Dimocárpus	84	
Cryptospérmum	268	Cynométra	171	Décodon	202	Dimórpha	171	
Cryptostégia	356	Cynosùrus	571	Decumária	206	Dinéba	560	
Cryptostémma	312	Cynosùrus	560, 562, 569,	Deeringia	432	Dinéba	560	
Cryptóstylis	485	Cynosùrus	571	Delima	13	Diinetus	373	
Cryptotænia	246	Cynosùrus	561	DELIMACEÆ	12	Dióclea	159	
Cucífera	543	Cynosùrus	561	Delíslia	592	Diòdia	265	
Cucíbalus	51	Cynosùrus	571	Delphínum	8	Diomèdea	309	
Cuculábalus	51, 53	Cypérus	551, 598	Dendrobium	489	Diomèdes	514	
Cuculária	197	Cypérus	552	Dendrobium	489	Dionæa	46	
Cúcumis	214			Dentária	24	Dioscòrea	521	
Cúcumis	214			Dentella	263	DIOSCOR'EÆ		
Cúcumis	215			Dermásea	240	Diósma	106	
Cúcúrbita	215			Derris	161	Diósma 105, 106, 116		
				Deschampsia	559	DIO'SMEÆ	105	

INDEX GENERUM.

607

Page.	Page.	Page.	Page.
Diosp'yros .. 349, 593	Dryándra .. 448	Embóthrium .. 447	Eriosynáphe .. 250
Diòtis .. 437	Dr'yas .. 190, 589	Embóthrium .. 447	Erísmá .. 590
Diphylélia .. 16	Dr'yas .. 189	Embryópteris .. 350	Eríthalis .. 260
Dipláchna .. 571	Drymária .. 55	Emericá .. 356	Ernódea .. 264
Diplázium .. 582, 598	Dr'ypis .. 54	'Emex .. 440	Erodén'drum .. 443, 444
DIPLECO'L OBEÆ .. 36	Duchésnia .. 189	Emiléa .. 302	Eròdium .. 89
Diplochítia .. 205	Dulichium .. 549	EMPE'TREÆ .. 460	Eróphila .. 26
Diplócoma .. 299	Dumásia .. 140	Empétrum .. 460	Erpétion .. 45
Diplólepis .. 361	Dumerília .. 280	Empétrum .. 460	Erúca .. 35
Diplopáppus .. 294	Dumerília .. 592	Empleúrum .. 107	Erucária .. 36
Diplophyllum .. 399	Dupón'tia .. 559	Enarthrocárpus .. 36	ERUCARIEÆ .. 36
Diplostégium .. 204	Duránta .. 417	Encélia .. 308	Ervília .. 154
Diplostéphium .. 295	Duroía .. 261	Enchy'læna .. 434	Er'vum .. 154
Diplotáxis .. 34	Duvállia .. 356	Ency'clia .. 490	Ery'ngium .. 244, 591
Diplothémium .. 544	Duvaú'a .. 119	ENDO'GENÆ .. 476	Ery'simum .. 30, 585
Dipódium .. 489	Dyssòdia .. 305	Enkiánthus .. 331	Ery'simum 22, 29, 30, 31
DIPS'ACEÆ .. 269	EBEN'ACEÆ .. 349	Enneapògon .. 567	Erythr'aea .. 363
Dípsacus .. 269	E'benus .. 152	Enodiúm .. 560	Erythrina .. 160
Díptera .. 240	Ecastaphyllum .. 162	Enslènia .. 360	Erythrìna .. 161
Dípterix .. 168	Eccremocárpus .. 66	Entáda .. 162	Erythrol'æna .. 283
Dirca .. 448	Eccremocárpus .. 66	Enteléa .. 70	Erythrónium .. 539, 597
Dísa .. 484	Echevêria .. 225	EPACRI'DEÆ .. 345	Erythrophlémum .. 163
Disándra .. 394	Echinácea .. 310	E'pacris .. 45	ERYTHROXYLEÆ .. 80
Dischídia .. 360	Echinária .. 567	E'pacris .. 345, 346	Erythróxylum .. 80
Dischísma .. 414	Echinocáctus .. 235, 590	Epáltes .. 315	Erythróxylon .. 80
Discopleúra .. 245, 591	Echinóchloa .. 563	Ephédra .. 474	ESCALL'ONEÆ .. 239
Disénuma .. 217	Echinóphora .. 255	Ephiélis .. 84	Escallónia .. 239, 591
Disophylla .. 400	Echinopógon .. 559	Epidéndrum .. 490	Eschscholtzia .. 19
Dispérès .. 485	Echinops .. 287	Epidéndrum .. 488, 489,	E'sula .. 454, 455
Dissórum .. 539	ECHINOPS'I'DEÆ .. 287 490	Ethùlia .. 315
Dissoléna .. 354	Echinospér'mum .. 377	Epig'aea .. 342	Eucaly'ptus .. 208
Diùrus .. 485	Echites .. 355	Epilobium .. 198	Euchétes .. 106
Dodártia .. 390	Echites .. 356	Epilobium .. 197, 198	Euchilus .. 124
Dodecáttheon .. 423, 595	Echium .. 375	Epimédium .. 16, 584	Euchròma .. 395
Dodon'aea .. 85	Eclípta .. 308	Epipáctis .. 486	Eucléa .. 457
DODON'A'ACEÆ .. 85	Edwárd'sia .. 122	Epipítis .. 486	EUCLÍ'DEÆ .. 28
Dólichos .. 158	Ehrétia .. 379	Epiphy'llum .. 237	Euclídiúm .. 28
Dólichos .. 158, 159	Ehrhárta .. 574	EQUISETACEÆ .. 575	Eucomis .. 530
Doliocárpus .. 12	Ehrhárta .. 574	Equisétum .. 575	Eucròsia .. 513
Dombéya .. 69, 586	Ekebér'gia .. 86	Eragróstis .. 570	Eudésmia .. 208
Dombéya .. 367, 475	ELÆA'GNEÆ .. 451	Eragróstis .. 570	Eugénia .. 212
DOMBEY'ACEÆ .. 69	Eláeagnus .. 451	Eránthēmum .. 422	Eugénia .. 211, 212, 213
Dóndia .. 244	ELÆOCAR'PÆ .. 71	Eránthemum .. 415, 420	EUHEDYS'AREÆ .. 148
Dónia .. 298, 299	Elaeocárpus .. 71	Eranthus .. 7	EULEPTOSPE'RMEÆ .. 208
Doódia .. 531, 598	Elaeodéndron .. 110	Eremùrus .. 524	Eulóphia .. 490
Doódia .. 149	Elaeodéndron .. 111, 349	Eria .. 489	EUONY'MEÆ .. 109
Dorónicum .. 299	Eláis .. 543	Erica .. 332	Eupat'OREÆ .. 299
Dorónicum .. 298, 299	Eláphrium .. 120	Erica .. 332, 341	Eupatórium .. 300
Dorstémia .. 464	Elátè .. 543	ERICEÆ .. 330	Eupatòrium .. 291, 301
Doryánthes .. 516	Elatérium .. 215	Erigénia .. 243	Euphórbia .. 453
Dory'cnium .. 137	Elátine .. 55	Erigeron .. 294	Euphórbia .. 452
Drába .. 25, 585	Elégia .. 547	Erigeron .. 290	EUPHORBIACEÆ .. 452
Drába .. 25, 26	Eleiotis .. 152	Erinus .. 394	Euphòria .. 84
Drac'æna .. 533	Eleócharis .. 550	Eriobót'ra .. 176	Euphrásia .. 394
Drac'æna 517, 532, 533	Elephántopus .. 288	Eriocália .. 243	Eury'a .. 72
Dracocéphalum .. 408	Elettària .. 492	Eriocaúlon .. 546, 597	Eury'ale .. 17
Dracocéphalum .. 406	Eleusine .. 571	Eriocéphalus .. 317	Eury'cycles .. 513
Dracóntium .. 479	Eleusine .. 571	Eriochilus .. 486	Eury'spermum .. 443
Dracóntium .. 479, 480	Elich'r ysum .. 292, 293	Eriócoma .. 314	Eury'coma .. 121
Dracophyl'um .. 345	Ellisia .. 380	Eriodéndron .. 66	Eustachys .. 562
Drepánia .. 275	Ellobocárpus .. 577	Eriogonum .. 440	Eustégia .. 361
Drepanocárpus .. 161	Elödea .. 76	Eriol'æna .. 69	Eustoma .. 364
Drepanophyl'um .. 245	Elschóltzia .. 400	Erióphorum .. 552	Euphòria .. 84
Drummón'dia .. 591	Elymus .. 566	Erióphorum .. 552	Euphrásia .. 394
Drímia .. 529	El'yna .. 552	Eriophyl'um .. 322	Eury'a .. 72
Drímys .. 13	Elytrária .. 420	Eriospermum .. 529	Eury'ale .. 17
Dróséra .. 45	Embélia .. 348	Eriostém'on .. 107	Eury'cycles .. 513
DROSER'ACEÆ .. 45	E'mblica .. 459	Eriostém'on .. 407	Eury'spermum .. 443

INDEX GENERUM.

	Page.		Page.		Page.		Page.	
Eustrephus	533	Fragária	189	Gaúra	198	Glossòdia	497	
Eutáxia	124	Fragária	192	Gay'a	64	Glòssula	485	
Eutérpe	543	Franciscea	389	Gay'a	591	Glottidium	143	
Eúthales	324	Frankènia	48, 586	Gazània	312	Gloxínia	329	
Euthàmia	298	FRANKENTACÆ	48	Geissomèria	420	Glycéria	570	
Eútoca	380, 594	Fransèria	315	Geissorhiza	503	Glycèria	569, 570	
Euxènia	315	Frasèra	363	Gèla	600	Glycine	140	
Euzòmum	35	Fráxinus	350	Gelònium	457	Glycine	127, 141, 149,	
'Evax	294	Fráxinus	351	Gelsémium	353 156, 157, 168		
Evòdia	107	Frezièra	72	Genetyl'lis	207	Glycosmis	75	
Evòlvulus	373	FRÉZI'EAE	72	Geniòsporum	414	Glycrrhiza	141	
E'xacum	364	Frièsia	72	Genipa	261	Gmelina	417	
E'xacum	364	Fritillària	537	Genista	129	Gnaphàlium	291	
Excàcaria	459	Fritillària	538	Genista	131	Gnaphàlium	291, 292,	
Exocàrpus	450	Fúchzia	197, 590	GENI'STEAE	125 293, 308		
EXOG'ENAE	1	FUCHSI'EAE	197	Gentiàna	362	Gnètum	464	
Exostémma	262	Fuirèna	552	GENTIANEAE	362	Gnídia	449	
Fàba	153	Fumària	20	Geodòrum	487	Godóya	78	
Fabràcia	210	Fumària	20	GEOFFREAE	167	Goldbächia	32	
Fabràcia	209	FUMARIAE	20	Geoffroya	168	Gomèza	487	
Fagàra	107, 120	Fúnkia	517	Geónoma	544	Gómphia	109	
Fagelia	157	furcr'ea	534	Geófila	264	Gomphocárpus	361	
Fagonia	104	Fusànum	450	Georgina	310	Gomphocárpus	361	
Fagr'aea	353	Gärtnera	81	GERANTACEAE	88	Gompholòbiu	124	
Fagus	467	Gàgea	527	GERANI'EAE	88	Gompholòbiu	123	
Falcària	245	Gàgea	527	Geraniùm	88	Gomphréna	432	
Fálkia	373	Gagñeòbina	163	Geraniùm	88, 90, 91,	Gomphréna	432	
Farsètia	24	Gaillardia	310 92, 93, 94, 97, 98, 99, 92, 93, 94, 97, 98, 99,	Gomutòs	544	
Farsètia	24	GALACINEAE	345	Galactìes	100, 101	Gongòra	489	
Fèdia	268	Galáctia	140	Galactites	285	Goniocárpus	200	
Fèdia	269	Galactites	285	Galactodéndron	462	Gonolòbus	360	
Ferdinàndia	318	Galanthus	516	Galanthus	516	Gonostémou	358	
Fernandésia	490	Galarh'œus	455	Gálox	345	Gonostémou	357	
Fernèlia	262	Gáloxia	499	Galáxia	499	Goodènia	324	
Ferònia	75	Galáxia	499	GESNER'EAE	329	Goodènia	324, 325	
Ferrària	499	Gálxia	499	Gethy'lis	516	GOODEN'OVIAE	324	
Ferrària	497, 499	G'ALEAE	265	Getònia	196	Goòdia	126	
Ferreòla	349	Galedùpa	161	Géum	190	Goòdia	126	
Férula	250, 591	Galegà	141	Géum	189, 190	Goodyèra	485	
Férula	249, 250, 253	Galèga	142, 144, 156	Ghinia	418	Gordònia	73	
Festúca	572	Galenia	437	Gilia	368, 594	GORDONI'EAE	73	
Festúca	559, 567, 570,	Galeobdolon	405	Gilia	368	Gortèria	312	
..... 571, 572, 573, 574	Galeópsis	405	Gilia	368	Gossypium	63		
Feuillæa	215	Galinsògea	308	Gillénia	194	Gouània	114	
Fibigia	24	Galinsògea	308	Gilliësia	546	Gr'æmia	315	
Ficària	7	Galinòsògea	308	Ginginsia	220	GRAMINAE	552	
FICO'I'DEAE	227	Galipèa	108	Ginkgo	473	Grammáthes	223	
Ficus	460	Gàlium	266	Ginòria	202	Grammitis	578	
Fièldia	366	Gàlium	266	Gisèkia	433	Grammitis	578	
Filàgo	294	Galphímia	81	Githàgo	54	GRAN'ATEAE	195	
FILICES	576	Garcinia	78, 587	Gladiolus	500	Grángea	318	
Fimbristylis	550, 597	GARCINIÆ	78	Gladiolus	499, 500, 501,	Grangèria	173	
Fischèria	358	Gardènia	261 502	Graphéphorum	571		
Flacoúrtia	40	Gardènia	260, 261	Glaúcium	19, 585	Gratiola	396	
FLACOURTANEAE	40	GARDENIÆ	260	Glaúx	428	Gratiola	396	
Flagellària	546	Gardòquia	409	Glechòma	408	Grenvillea	96	
Flavèria	313	Garidélla	8	Gleditschia	168	Grevillea	446	
Flemíngia	152	Garùga	120	Gleichènia	577, 598	Grèwia	70	
Flindérsia	86	Gastèria	519	GLEICHEN'EAE	577	Grèwia	206	
Florestina	306	Gastònia	256	Glínus	234	Grias	213	
FLUVIALES	478	Gastrídium	555	Glòbba	491	GRI'ELEAE	89	
Fœniculum	247	Gastrocárpha	280	Glòbba	491	Gríelum	89	
Fœniculum	245	Gastrolòbium	124	Globularia	426	Griffinia	510	
Fontanèsia	351	Gastronéma	512	Gaudichaudia	81	Grindèlia	298	
Forskòhlea	463	Gaudichaudia	81	GLOBUL'ARIE	426	Grislea	202	
Fothergilla	473	Gaulthèria	331 426	Gronòvia	216		
Fothergilla	205	Gaulthèria	330	Gloriòsa	539	GROSSULARIEAE	238	

INDEX GENERUM.

609

Page.	Page.	Page.	
Guaícum	104	<i>Hebenstreitia</i>	414, 415
Guàrea	86	<i>Hedeòma</i>	404
Guattèria	15	<i>Hédera</i>	256
Guazúma	67	<i>Hédéra</i>	87
Guipínia	27	<i>Hedy'chium</i>	493
Guettárda	260	HEDYOTÍDEÆ	262
GUETTA'RDEÆ	260	<i>Hedyotis</i>	262
Guevina	599	<i>Hedyótis</i>	599
Guilandina	168	<i>Hedy'pnio</i> s	275
Guilandina	168	<i>Hedy'pniois</i>	275
Gundélia	288	HEDYSAREÆ	148
Gunnèra	12	<i>Hedy'sarum</i>	151
Gustàvia	213	<i>Hedy'sarum</i>	127, 140, 149, 150, 151, 152, 153
GUUTTIFERÆ	78	<i>Heimia</i>	202
Guzmánnia	533	<i>Helénium</i>	307
Gymnædia	483	<i>Heleóchloa</i>	557
Gymnémâma	360	<i>Heliánthemum</i>	41
Gymnocladus	168	HELIA'NTHEÆ	306
Gymnogrâmma	578	<i>Heliánthus</i>	309
Gymnolómâa	309	<i>Heliánthus</i>	312
Gymnóstachys	479	<i>Helichrysum</i>	292
Gymnóstylis	317	<i>Helichrysum</i>	294
Gynandrópsis	37	<i>Helicónia</i>	495
Gypsóphila	48	<i>Helicóteres</i>	65
Gypsóphila	51	<i>Heliocárpus</i>	70
Gyrocarpus	442	<i>Heliófila</i>	36
Gyromâia	523	HELIOPHILÆ	36
Gyrotémon	71	<i>Heliotrópium</i>	374, 379
Habenària	484	<i>Helleborinæ</i>	485
Habenària	483	HELLÉBOREÆ	7
Hablitzia	434	<i>Hellénia</i>	493
Habrántus	506	<i>Hellénia</i>	493
Hacquetia	244	<i>Heliotrópium</i>	374, 379
Hæmadictyon	356	<i>Helleborinæ</i>	485
Hæmânthus	510	HELIOPHYLLÆ	7
Hæmatóxylon	169	<i>Hippocratea</i>	79
HÆMODORACEÆ	504	HIPPOCRATE'A-	
Hæmodôrum	504	CEÆ	79
Hægea	221	<i>Hippocrépis</i>	148
Hækeâ	446	<i>Hippomâne</i>	458
Halésia	347	<i>Hippomarâthrum</i>	255
Halimodéndron	144	<i>Hippóphæ</i>	451
Hallèria	389	<i>Hippophæ</i>	451
Hália	127	HIPPURÍDEÆ	201
Halochâris	282	<i>Hippuris</i>	201
Halocuémum	437	<i>Híptage</i>	81
HALOR'AGEÆ	200	HIP'TAGEÆ	81
Halorágis	200	<i>Hippocrépis</i>	148
<i>Halorágis</i>	201	<i>Hir'æa</i>	81
HAMAMELIDÆ	473	<i>Hírculus</i>	240
Hamaméllis	473	<i>Hirtélla</i>	173
Hamelia	473	<i>Hòarea</i>	90
Hamelia	260	<i>Hoffmanséggia</i>	169
HAMEL'ACEÆ	260	<i>Hóitzia</i>	368
Hamiltónia	451	<i>Hólcus</i>	562
Hamiltónia	263	<i>Hólcus</i>	559, 561, 562
Hapalostéphium	273	<i>Holigárna</i>	588
Hardwicâlia	169	<i>Holmskioldia</i>	417
Haróngâ	76	<i>Holsteum</i>	55
Harpalyce	274	<i>Holsteum</i>	55
Harrâchia	420	<i>Homálium</i>	116
Harrisònia	361	<i>Homéría</i>	498
Hartògia	111	<i>Hoódia</i>	359
Hasselquâstia	252	<i>Hookèra</i>	517
Hastingia	417	<i>Hòpea</i>	347
Havètia	78	<i>Hórdeum</i>	558
Hawórthia	520	<i>Horkèlia</i>	189
Hebenstreitia	414	<i>Hormínum</i>	409
		<i>Hornemânnia</i>	393, 394
		<i>Hosáckia</i>	589
		<i>Hósta</i>	417
		<i>Hottònia</i>	425
		Houstònìa	361
		<i>Houttuynia</i>	479
		<i>Hòvea</i>	125
		<i>Hovènia</i>	113
		<i>Hóya</i>	359
		<i>Hudsònia</i>	43
		<i>Huérnia</i>	359
		<i>Hugònâia</i>	72
		<i>Hùmea</i>	291
		<i>Hùmulus</i>	463
		<i>Hunnemânnia</i>	19
		<i>Hùra</i>	458
		<i>Hutchinsia</i>	27
		<i>Hutchinsia</i>	27
		<i>Hyacinthus</i>	528
		<i>Hyacinthus</i>	527, 528, 529, 530
		<i>Hyâlis</i>	502
		<i>Hyænânche</i>	457
		<i>Hydrângea</i>	242
		<i>Hydrâstis</i>	4
		HYDROCHARI-	
		DEÆ	476
		HYDROCARYE	200
		<i>Hydrocharis</i>	476
		<i>Hydrochloa</i>	570
		<i>Hydrocotyle</i>	243
		<i>Hydrocotylineæ</i>	243
		<i>Hydroglòssum</i>	577
		<i>Hydrolea</i>	369
		HYDROLEACEÆ	369
		HYDROPELTÍDEÆ	17
		<i>Hydropeltis</i>	17
		HYDROPHYLLÆ	380
		<i>Hydrophyllum</i>	380
		<i>Hydrophyllum</i>	380
		<i>Hygróphila</i>	420
		<i>Hymenæa</i>	172
		<i>Hymenanâtha</i>	586
		<i>Hymenocâllis</i>	513
		<i>Hymenocârpus</i>	132
		<i>Hymenodictyon</i>	262
		<i>Hymenogyne</i>	234
		<i>Hymenopâppus</i>	306
		<i>Hymenophyllum</i>	583
		<i>Hymenophyllum</i>	583
		<i>Hyoscyamus</i>	382
		<i>Hyoscyamus</i>	382
		<i>Hyoséris</i>	276
		<i>Hyoséris</i>	275, 276
		<i>Hypécoum</i>	19
		<i>Hyperanthâra</i>	168
		HYPERICI'NEÆ	76
		<i>Hypéricum</i>	76
		<i>Hiph'æne</i>	543
		<i>Hypoëalyptus</i>	128
		HYPOCHÆRÍDEÆ	276
		<i>Hypoëstes</i>	422
		<i>Hypol'æna</i>	547
		<i>Hypopithys</i>	342
		HYPOXI'DEÆ	505

INDEX GENERUM.

Page.	Page.	Page.	Page.
Hy'ptis 414, 594	Jacarânda 366	Kundmânnia 247	Lawsônia 202
Hyssopus 402	Jacksônia 124	Kûlnia 299	Laxmânnia 532
<i>Hyssopus</i> 402	Jacquínia 348	Kuhnistèria 141	Lebekâria 129
Iántha 490	JACOBÆ 301	Ky'dia 69	Lebékchia 139
Ibérîs 27	Jambolífera 107	Kyllínga 552, 598	Lebretònia 61
<i>Ibérîs</i> 27	Jambôsa 212	LABÍTATÆ 399	Lecanocárpus 431
<i>Ibbetsônia</i> 123	Janiphæ 458	LABIATIFLÓRÆ 280	Lechêa 43
Ichnoçárpus .. 356	Jasione 328, 593	Lâblab 159	Lechenaúltia 324
I'cica 589	JASMI'NEÆ 352	Lachenàlia 529	LECYTHÍDEÆ 213
Ilex 111	Jasminum 352	Lachn'æa 449	Lécythis 213
<i>Ilex</i> 110, 111	Jasònia 294	Lachnâthes 504	Lécythis 213
ILLECEBREÆ .. 220	Jâtrophæ 457	Lactùca 277	Ledeburia 246
Illécerebrum 220	Jâtrophæ 457, 458	Lactùca 277	Lèdum 344
ILLécerebrum 220, 221, 431,	Jeffersonia 17	LACT'UCEÆ 277	Lèdum 344
..... 432	Jenkinsônia 100	L'ætia 40	Leeâ 87
Illicium 13	Johannia 280	Lagáscea 288	LEE'ACEÆ 87
Imatophyl'lum .. 516	Jôhnia 80	Lagónychia 288	Leérzia 554
Imbricâria 349	Joliffia 215	Lagenària 213	LEGUMIN'OSÆ 122
Imhöfia 510	Jonèsia 169	Lagerstr'œmia 202	Leimânthium 540
Impâtiens ... 102, 588	Jonidium 45	LAGERSTR'ŒMIEÆ 202	Leiophy'llum 344
Impâtiens 101	Jonidium 43, 45	Lagéttâ 448	Lémâna 478
Imperâta 560	Jossínia 211	Lagocœcia 255	Léns 154
Imperatòria 251	JUGLA'NDEÆ 465	Lagony'chium 164	LENTIBUL'ARIEÆ 422
Imperatòria 249, 250,	Júglans 465	Lagoséris 273	Leonôtis 404
..... 251	Júglans 465	Lagoséris 278	Leóntice 16
Indigófera 139	JUNCAGI'NEÆ 477	Lagunculària 196	Leóntodon 274
In'ga 163	JU'NCEÆ 545	Lagunéa 65	Leóntodon 275
<i>In'ga</i> 163	Júncus 545	Lagun'æa 62	Leontopódium 291
Inocárpus 348	Júncus 545	Lagûrus 559	Leonùrus 405
Inùla 290	Júngia 592	Lahâya 221	Leonùrus 407
Inùla 291	Juniperus 474	Lamárckia 110, 562	Lepechímia 409
Ionópsis 490	Jussi'æa 199	Lambértia 447	Lepeocérctis 561
Ipom'œa 370	JUSSI'ÆÆ 199	Lâmium 406	Lepidagáthis 420
Ipom'œa 373	Justícia 421, 595	Lâmium 406	LEPIDI'NEÆ 31
Ipomópsis 368	Justícia 420, 422	Lanària 504	Lepidium 31
Ipomópsis 368	Kämpferia 492	Lancisia 318	Lépidium 27
Iresine 433	Kalanchöe 224	Lantâna 417, 595	Lapeyroúsia 499
Iresine 433	Kâlmia 342	Lapeyroúsia 499	LAPL'ACEÆ 73
IRI'DEÆ 495	Kalosântes 224	Lappa 281	Leptândra 396
'Iris 495	Kaulfüssia 305	Lappâgo 557	Leptândus 541
'Iris 497, 498	Kennedyâ 156, 589	Lapsâna 274	Leptâsæo 240
Isâchne 564	Kernéra 26	Lapsâna 275, 276	Leptocarp'æa 22
Isânthus 399	Kérria 194	Larbréa 55	Leptocárpus 547
ISATI'DEÆ 32	Kiggelâria 40	Lârix 475	Leptocaúlis 245
Isâtis 32	KIGGEL'ARIEÆ 40	Lârrea 104, 588	Leptochloa 571
Isch'ænum 562	Kingia 545	Laserpitium 252	Leptomèria 450
Isch'ænum 562	Kirganèlia 459	Laserpitium 249, 252,	LEPTOSPE'RMEÆ 207
Iséertia 260	Kitaibélia 60 355, 591	Leptospérnum 210
Ismène 513	Kleinöhövia 67	Lasiândra 204	Leptospérnum 209
Isnârdâ 200	Kleïnia 301	Lasiospérnum 279	Leptostélma 299
Isocárpha 307	Knâppia 557	Lasiospérnum 308	Leptùrus 563
Isochilus 489	Knaútia 271	Lasiopétalum 67	Lepyròdia 547
Isoëtes 575	Knaútia 271	Lasiopétalum 67	Léria 280
Isôlepis 550, 597	Knowltonia 4	Lasiopétalum 67	Lespedèza 152
Isopétalum 101	Knóxia 265	Latânia 543	Lespedèza 149
Isopogón 442	Kobrësia 549	Lathr'æa 422	Lessértia 144
Isop'yrum 7	Kochia 434	Lâthyrus 155	Lestibudèisia 431
Isótoma 324	Kochia 434, 435	Lâthyrus 143, 154	Lett'sòmia 72
I'tea 345	Kœlèria 567	Lâugeria 260	Lett'sòmia 373
I'tea 48, 332	Kœlèria 559	LAURI'NÆ 441	Leucadéntron 442
'Iva 314	Kœlpinia 275	Laurophyl'lus 599	Leucadéntron 444
I'xia 502	Kœlreutèria 85	Laúrus 141	Leucas 404
I'xia 498, 499, 500,	Kœnigia 440	Laúrus 141	Leucéria 280
..... 501, 502	Kólbea 540	Lavândula 408	Leucocôryne 532
Ixiolirion 510	Kòniga 25	Lavândula 414	Leucôjum 516
Ixòdia 291	Krigia 276	Lavatèra 60	Leucopodón 346
Ixòra 263	Krigia 276	Lavènia 308	Leucoscéptrum 402
Ixòra 263	Krubera 252	LAVOIS'E'REÆ 203	Leucospérnum 444

Page.	Page.	Page.	Page.
Leucostemma	293	Loddigèisia	128
Leüzea	285	Lodiculària	562
Leüzea	282	Lodoicea	543
Levisticum	249	Lœflingia	222
Lewisia	227	Logània	364
Leysèra	301	Loiseleuria	344
Leysèra 281, 307, 308		Lòlium	566
Liatris	289	Lomàndra	545
Libanòtis	248	Lomària	581, 598
Libértia	498	Lomàtia	447
Lichtensteinia	247	Lomatophy'lum'	517
Lichtensteinia	540	Lonchitis	583
Licuàla	544	Lonchocárpus	143
Lidbékcia	318	Longchámpisia	308
Lidbékcia	318	Lonicera	258
Lightfoótiá	325, 592	Lonicera	258, 259
Ligularia	240	Lopèzia	200
Ligústicum	248	Lophánthus	402
Legústicum 245, 248,		Lophiola	504
.... 249, 250, 255		Lophíra	259
Ligústrum	352	LOPHIREÆ	259
Lílium	538	Lophoscíadium	253
Lílium	538	Lophospérnum	390
Límeum	220	Lopímia	63
Limnétis	553	LORA'NTHEÆ	259
Limnócharis	477	Loránthus	259, 592
Limodórum 487, 490		Loreya	205
Limònìa	75	L'OTEAE	125
Limònìa	75	Lòtus	137
Memosélla	394	Lòtus 132, 137, 138,	
Linària	390 139	
Lincònìa	116	Loúrea	149
Lindérnia	394	Loureira	460
Lindérnia	394	Lówea	178
Linds'æa 583, 599		Lubinìa	425
L'INEÆ	58	Lucùlia	260
Linn'æa	259	Lucúma	349
Linocièra	351	Lùdia	40
Linum	58, 586	Lùdia	40
Línnum	58	Ludólfia	574
Lipària	127	Ludòvia	480
Lipària	127	Ludwigia	200
Lipàris	486	Ludwigia	200
Lipóstoma	599	Lúffa	214
Líppia	419	Luhéa	599
Líppia	418	Lunnitzéra	413, 414
Liquidámbar	472	Lunària	24
Liriódendron	14	Lunària	24
Lisiánthus	364	Lupináster	137
Lisiánthus	364	Lupinus	159, 589
Lissánthe	346	Lùzula	545
Lissochilus	488	Luzuriàga	532
Listéra	486	Ly'chnis	54
Líta	363	Ly'chnis	52, 53, 54
Lithospérnum	374	Lycium	389
Litt'æa	534	Lycopérsicum	385
Litorélla	429	LYCOPODIACEÆ	
Livistòna	543 576	
Llóydia	527	Lycopodium	576
Loàsa	218	Lycopodium	576
Loàsa	218	Lycópsis	377
LO'ASEÆ	218	Lycópsis	376, 377
Lobàdium	119	Lycópus	399
Lobària	240	Lycòris	509
Lobèlia	322, 592	Lygèum	552
Lobèlia	324, 325	Lygodésmia	277
LOBELIACEÆ 322		Lygòdium	577, 598
Lockhártia	490	Lyonia	331
		Lyónsia	356
		Lyperánthus	486
		Lysimàchia	425
		Lysimàchia	425
		Lysinéma	345
		LYTHRARIÆ	201
		L'ythrum	202
		L'ythrum	202
		Màba	349
		Macbridea	409
		Macleáya	19
		Maclùra	462
		Macradènia	488
		Macránthus	156
		Macrocnémum	262
		Macrocnémum	261
		Macrolòbium	172
		Macropódium	23, 585
		Macróstilos	106
		Macrotòrops	122
		Macrotòrys	11
		Màdia	306
		M'æsa	347
		Magnòlia	14
		Magnòlia	484
		MAGNOLIACEÆ	13
		Magydaris	255
		Mahérnia	68
		Mahónia	16
		Mahùrea	78
		Maièta	205
		Mairànìa	330
		Majánthemum	523
		Malabàila	251
		Malachodédon	73
		Malachra	60
		Maláxis	487
		Maláxis	485, 486
		Malcòmia	28
		Malloccòca	70
		Malochìchia	159
		Málope	58
		Malpighia	80
		Malpighia	80, 81
		MALPIGHIACEÆ	
		Málva	59, 586
		MALVACEÆ	58
		Malvavíscus	61
		Mamméa	78
		Mammillaria	235, 590
		Mandràgora	383
		Manéttia	263
		Mangifera	117
		Mangifera	117
		Mangílìa	348
		Manicària	544
		Manisùris	563
		Mantisía	491
		Manùlea	394
		Máppa	458
		Maránta	495
		Maránta	494
		Marátia	577
		Marcgràvia	79
		MARCGRAVIA-CEÆ	79
		Mariálva	78
		Márica	497
		Márica	498
		Márla	78
		Mariscus	552, 598
		Márlea	213
		Marrùbium	408
		Marrùbium	405
		Marsdènia	360
		Marshallia	306
		Marsilea	576
		MARSILEACEÆ	576
		Marty'nia	367
		Massónia	530
		Massònìa	527
		Matàxa	308
		Mátayba	84
		Mathiola	21
		Matricària	319
		Matiòla	260
		Máttia	378
		Mauràndia	390
		Maurìa	119
		Maurítia	544
		Maxillària	489
		Maxillària	489
		Maximiliàna	544
		Máytenu	110
		Mazus	393
		Meconópsis	19
		Medèola	523
		Medicàgo	132
		Medusea	453
		Megacarp'a	28
		Megaclíniu	488
		Mégaseu	239
		Megastàchy	570
		Melaleùca	207
		Melaleùca	210
		MELALEUCEÆ	207
		Melampyrum	395
		Melampòdium	313
		Melananthèra	306
		Melanorrh'ea	588
		Melanoselinu	253
		Melanosticta	169
		MELANTHACEÆ	
	 539	
		Melánthium	539
		Melánthium	539, 540
		Melasph'ærula	502
		Melástoma	204
		Melástoma 203, 204, 205,	206
		MELASTOMACEÆ	
	 203	
		Melhània	69
		Mélia	85
		MELIACEÆ	85
		Meliáñthus	104
		Mélica	560
		Mélica	560
		Mélichrus	346
		Melicòcca	84
		Melicope	107
		Melicytus	40
		Melilotus	134

INDEX GENERUM.

Page.	Page.	Page.	Page.
<i>Melildtus</i> 134	<i>Mílum</i> 555	<i>Morónbæa</i> 79	<i>Narthècium</i> 545
<i>Melinis</i> 558	<i>Mílum</i> 555, 564	<i>Morphíxia</i> 502	<i>Nastúrtium</i> 21, 585
<i>Mélissa</i> 409	<i>Mílla</i> 596	<i>Mòrus</i> 462	<i>Naucléa</i> 263
<i>Mélissa</i> 409	<i>Millèria</i> 313	<i>Mòrus</i> 462	<i>Nauenbúrgia</i> 288
<i>Melittis</i> 409	<i>Millingtònìa</i> 366	<i>Moschósma</i> 413	<i>Nébélia</i> 116
<i>Melocáctus</i> 235	<i>Mimètes</i> 444	<i>Mougeóttia</i> 68	<i>Negúnido</i> 82
<i>Melocánna</i> 574	<i>Mimòsa</i> 163	<i>Mouríria</i> 195	<i>Néja</i> 299
<i>Melochia</i> 67	<i>Mimòsa</i> 163, 164, 165,	<i>Mozinna</i> 460	<i>Nélitrís</i> 210
<i>Melodinus</i> 356	166, 167	<i>Mucùna</i> 159	<i>Nelsónia</i> 422
<i>Melóthria</i> 215	<i>MIMÓSEÆ</i> 163	<i>Muhlenbérghia</i> 557	<i>Nelúmbium</i> 17
<i>MEMECY'LEE</i> 195	<i>Mímulus</i> 394, 594	<i>Muhlenbérghia</i> 559	<i>NELUMB'ONÆ</i> 17
<i>Memécylon</i> 195	<i>Mimúsops</i> 349	<i>MUL'INEÆ</i> 243	<i>Nectoúxia</i> 383
<i>Meniducus</i> 25	<i>Minuártia</i> 221	<i>Mulléra</i> 142	<i>Néctris</i> 17
<i>Meniscíum</i> 578, 598	<i>MINUA'RTIEÆ</i> 221	<i>Mündia</i> 47	<i>Negrétia</i> 159
<i>MENISPERM'A-CÆ</i> 15	<i>Mirabilis</i> 429	<i>Muntingia</i> 71	<i>Nemésia</i> 390
<i>Menispérnum</i> 15	<i>Mirbélia</i> 125	<i>Muráltia</i> 47	<i>Nemopánthes</i> 111
<i>Menispérnum</i> 15	<i>Miscopetalum</i> 240	<i>Murráya</i> 75	<i>Nemophila</i> 380
<i>Méntha</i> 399	<i>Mitchélla</i> 263	<i>Murucúnia</i> 218	<i>Neóttia</i> 485
<i>Méntha</i> 400	<i>Mítella</i> 242	<i>Musa</i> 495, 596	<i>Neóttia</i> 485, 486
<i>MENTHO'I'DEÆ</i> 399	<i>Mítella</i> 242, 591	<i>MUS'ACEÆ</i> 495	<i>Nepénthes</i> 452
<i>Mentzélia</i> 218	<i>Mitrasácmé</i> 364	<i>Muscária</i> 241	<i>Nepéta</i> 407
<i>Menyánthes</i> 364	<i>Mniárum</i> 221	<i>Muscári</i> 528	<i>Nepéta</i> 402, 409
<i>Menyánthes</i> 364	<i>Modécca</i> 218	<i>Mussæ'nda</i> 260, 261, 262	<i>NEP'E'TEÆ</i> 404
<i>Menziésia</i> 341	<i>Mehríngia</i> 55	<i>Müsschia</i> 593	<i>Nephélium</i> 84
<i>Meóschium</i> 562	<i>Moé'nchia</i> 55	<i>Mutisia</i> 280	<i>Nephródium</i> 579
<i>Mercuriális</i> 452	<i>Mogiphánes</i> 432	<i>M'yagrum</i> 32	<i>Nephrodíum</i> 580
<i>Meriania</i> 203	<i>Moldenháweria</i> 169	<i>M'yagrum</i> 31, 36	<i>Nerine</i> 509
<i>Meriania</i> 204	<i>Molin'ea</i> 84	<i>Magalurus</i> 572	<i>Nérium</i> 354
<i>Merténsia</i> 464	<i>Molínia</i> 560	<i>Mygíndia</i> 111	<i>Nérium</i> 354, 355
<i>Merténsia</i> 577	<i>Mólilia</i> 221	<i>Mylocárium</i> 332	<i>Nes'ea</i> 202
<i>Mesembryánthemum</i> 227	<i>Mollúgo</i> 55	<i>MYOPO'RINÆ</i> 419	<i>Nes'ea</i> 202
<i>Mesembryánthemum</i> 234	<i>Molopospérnum</i> 255	<i>Myóporum</i> 419	<i>Néslia</i> 31
<i>MESEMBRYANTH'E-MÆ</i> 227	<i>Moltkia</i> 375	<i>Myóseris</i> 278	<i>Nestléra</i> 308
<i>Méspilus</i> 175	<i>Moluccélla</i> 405	<i>Myosotis</i> 377	<i>Neurocárпum</i> 140
<i>Méspilus</i> 176, 177	<i>Moluccélla</i> 405	<i>Myosótis</i> 377	<i>Neurol'ena</i> 307
<i>Messerschmídia</i> 378	<i>Momórdica</i> 215	<i>Myosúrus</i> 4	<i>Neurolòma</i> 24
<i>Messerschmídia</i> 378	<i>Momórdica</i> 214	<i>My'rcea</i> 210	<i>Nicándra</i> 383
<i>Metalásia</i> 292	<i>Monáchna</i> 564	<i>Myrica</i> 472	<i>Nicolsónia</i> 149
<i>Metastélmá</i> 362	<i>Monánthes</i> 226	<i>Myrica</i> 472	<i>Nicotiana</i> 382
<i>Methónica</i> 539	<i>Monárdia</i> 403	<i>Myricária</i> 203	<i>Niebühría</i> 39
<i>Metrosidéros</i> 209	<i>Monárda</i> 404	<i>MYR'ICEÆ</i> 472	<i>Nigélia</i> 8
<i>Metrosidéros</i> 207, 209	<i>Monárda</i> 404	<i>Myriophyl'lum</i> 201	<i>Nigrítella</i> 483
<i>Méum</i> 249	<i>MONA'RDEÆ</i> 403	<i>Myrística</i> 442	<i>Nintóða</i> 258
<i>Méum</i> 245, 247	<i>Monérma</i> 563	<i>MYRISTI'CEÆ</i> 442	<i>Nipa</i> 544
<i>Meyéra</i> 308	<i>Monérma</i> 563	<i>MYR'SINE</i> 348	<i>Niphobolus</i> 578
<i>Michaúxia</i> 325	<i>D'ONEÆ</i> 476	<i>Myrsine</i> 348	<i>Nissolia</i> 142
<i>Michélia</i> 13	<i>Monodóra</i> 14	<i>MYRSI'NEÆ</i> 347	<i>Nítélla</i> 575
<i>Micónia</i> 205	<i>Monópsis</i> 324	<i>Myrsiphyl'lum</i> 533	<i>Nitraria</i> 235
<i>Micóniæ</i> 205	<i>Monótopca</i> 346	<i>MYRTACEÆ</i> 206	<i>Nivènia</i> 445
<i>Micranthë</i> 78	<i>Monótropa</i> 342	<i>MYRT'EÆ</i> 210	<i>Nócca</i> 288
<i>Micrántes</i> 240	<i>MONOTRÒPEÆ</i> 342	<i>Myrtus</i> 210	<i>Noisetta</i> 43
<i>Microcèle</i> 364	<i>Monsónia</i> 88	<i>Myrtus</i> 210	<i>Nolàna</i> 383
<i>Micróchloa</i> 558	<i>Monsónia</i> 88	<i>Myrtus</i> 211, 212	<i>Nolina</i> 540
<i>Micròcos</i> 70	<i>Montia</i> 220	<i>Nagéia</i> 472	<i>Nonatélia</i> 260
<i>Micròdon</i> 415	<i>Montinia</i> 197	<i>Nájas</i> 472	<i>Nònea</i> 276
<i>Microl'æna</i> 574	<i>MONTÍ'NIEÆ</i> 197	<i>Náma</i> 369	<i>Norántea</i> 79
<i>Microlícia</i> 203	<i>Mor'æa</i> 497	<i>Nandina</i> 16	<i>NORA'NTÆ</i> 79
<i>Microlòma</i> 362	<i>Mor'æa</i> 497	<i>Nap'æa</i> 64	<i>Notel'æa</i> 351
<i>Micropus</i> 294	<i>Mor'æa</i> 497, 498	<i>Naravélia</i> 2	<i>Nothol'æna</i> 578, 598
<i>Microstachys</i> 460	<i>Moricándia</i> 34	<i>Narcíssus</i> 514, 596	<i>Notóceras</i> 22
<i>Micróstilis</i> 487	<i>Morina</i> 269	<i>Narcíssus</i> 515	<i>Nóthria</i> 48
<i>Microtéa</i> 433	<i>Morinda</i> 263	<i>Nárdus</i> 552	<i>NOTORH'IZÆ</i> 28
<i>Micròtis</i> 486	<i>Morínga</i> 168	<i>Nárdus</i> 563	<i>Noty'lia</i> 487
<i>Mikánia</i> 300	<i>Morisónia</i> 39		<i>Núphar</i> 18, 585

INDEX GENERUM.

613

Page.	Page.	Page.	Page.
Nyctérium .. 387	Opothèca .. 432	Oxystélma .. 361	Patrísia .. 40
Nymph'æa .. 17, 585	Opópanax .. 249	Oxy'tropis .. 145	PATRÍSIEÆ .. 40
Nymph'æa .. 18	Oporínia .. 275	Oxy'tropis .. 146	Paullinia .. 84
NYMPHÆACEÆ .. 17	Opuñtia .. 237, 591	Ozothámnus .. 291	Paullinia .. 84
Ny'ssa .. 451	OPUNTIACEÆ .. 235	Pachidéndron .. 517	PAULL'NIEÆ .. 83
Obésia .. 358	O'rbea .. 357	Pachíra .. 66	Pavéttä .. 263
O'chna .. 108	ORCHIDEÆ .. 483	Pachypódium .. 594	Pávia .. 83
O'chna .. 109	O'rchis .. 483	Pachyrhizus .. 159	Pavónia .. 61
OCHNACEÆ .. 108	O'rchis .. 483, 484	Pachysándra .. 456	Pectinária .. 357
Ochròma .. 66	Oreoselinum .. 250	Pachysándra .. 456	Pécis .. 306
Ochròsia .. 354	Oríganum .. 402	Pæderia .. 265	Pécis .. 308
O'chrus .. 155	Orláya .. 253	Pæderota .. 396	PEDALI'NEÆ .. 367
Ochthódium .. 28	Ormocárpus .. 149	Pælimbía .. 250	Pedálium .. 367
Ocôtea .. 441	Ormòsia .. 122	Pæónia .. 12	Pediculàris .. 395
Otoméria .. 488	Ornithidium .. 489	PÆONIACEÆ .. 11	Pedilánthus .. 452
O'cymum .. 413, 594	Ornithocéphalus .. 488	Palafóxiá .. 301	Péganum .. 105
O'cymum .. 413, 414	Ornithogálum .. 526	Palávia .. 63	Péganum .. 105
OXYMO'DEÆ .. 413	Ornithogálum 527, 532	Paliúrus .. 112	Pékea .. 213
Odontites .. 247	Ornithoglóssum .. 539	Pallasia .. 308	PELARGONI'EÆ .. 89
Edèra .. 288	Orníthopus .. 148	PA'LME .. 542	Pelargónium .. 92, 587
Edmánnia .. 126	Orníthopus .. 148	Pánax .. 256	Pelargónium 89, 90, 91,
Enánthe .. 247	Ornítrophe .. 84	Pancrátium .. 513 92, 100, 101
Enánthe .. 247, 251	O'rnuś .. 351	Pancrátium .. 513	Peléxia .. 485
Enocárpus .. 543	Orobánchez .. 422	PAND'ANEÆ .. 478	Peliosánthes .. 540
Enoplía .. 112	OROBANCH'EÆ .. 422	Pandárus .. 478	Peltária .. 25
Enothèra .. 198	O'robus .. 156	Pánicum .. 563	Pen'ea .. 346
Œnothèra .. 200	O'robus 143, 155, 156	Pánicum 553, 558, 563,	PEN'E'ACEÆ .. 346
OLACI'NEÆ .. 74	ORONTIACEÆ .. 479 564	Penicillària .. 553
'Olax .. 74	Oróntiūm .. 480	Papáver .. 18	Pennisétum .. 553
Oldenlândia .. 262	Oróntiūm .. 480	PAPAVER'ACEÆ .. 18	Pennisétum .. 553
'Olea .. 351	Oróstachys .. 225 18	Pentadésma .. 78
'Olea .. 110, 351	Ortégia .. 221	PAPILION'ACEÆ .. 122	Pentápetes .. 69
OLEINÆ .. 350	Orthoceras .. 485	Pappophórum .. 567	Pentaphyl'on .. 137
Olínia .. 115	Orthopógon .. 563	Pappophórum .. 567	Pentáptera .. 196
Olísbea .. 197	Orthósanthus .. 498	Pap'yruś .. 552	Pentatáxis .. 292
Olivèria .. 255	Orthosíphon .. 413	Pardánthus .. 498	PENTH'OREÆ .. 227
Ol'yla .. 574	ORTHOSEPR'MEÆ .. 243	Pariàna .. 575	Penthórum .. 227
Omalánthus .. 456	Orvála .. 406	Parietária .. 463	Pentstémon .. 383
Omphálea .. 458	O'r'ya .. 575	Parinárium .. 173	Péntzia .. 316
Omphalòbium .. 121	Oryzópsis .. 552	Pàris .. 524	Peperòmia .. 482
Omphalòdes .. 378	Osbéckia .. 204	Parivòa .. 171	Péplis .. 201
ONAGR'ARIEÆ .. 197	OSBECKIA .. 204	Parkeria .. 577	Peraltea .. 168
Oncídium .. 488	Osmítæ .. 311	PARKERI'ACEÆ .. 577	Perdicíum .. 280
Onobròma .. 284	Osmorhíza .. 255	Párkia .. 163	Perdicíum .. 280
Onobròma .. 284	Osmünd'a .. 577	Parkinsónia .. 169	Peréksia .. 238
Onobr'ychis .. 151	Osmunda 577, 581, 582	Parnássia .. 46	Pergulària .. 359
Onocleà .. 581	Osmund'ACEÆ .. 577	Paróchetus .. 159	Pergulària .. 356
Onocleà .. 581, 582	Oss'æa .. 205	Paronychia .. 220	Peribália .. 560
Onónis .. 131	Osteospérnum .. 314	PARÓNYCH'EÆ .. 220	Perilla .. 400
Onopórdum .. 283	Ostéricum .. 249	PAROPS'I'EÆ .. 216	Periplòca .. 356
Onopórdum .. 284	O'strya .. 467	Párrya .. 24, 485	Periplòca .. 356
Onoséris .. 278	OSYRI'DEÆ .. 450	Parsónsia .. 356	Peripterá .. 63
Onósma .. 375, 594	Os'yris .. 450	Parthénium .. 314	Peritoma .. 37
Onosmòdium .. 375	Otánthus .. 315	Pascália .. 308	Perlèbia .. 255
Operculària .. 268	Othónna .. 305	Páspalum .. 555	Perojða .. 346
Operculària .. 268	Otídia .. 90	Páspalum .. 554, 558	Perótis .. 554
OPERCULARI'EÆ .. 268	Otóptera .. 140	Passerína .. 449	Pérsæa .. 441
OPIOGLO'SSEÆ .. 576	Oútea .. 172	Passerína .. 449	Persoónia .. 446
Ophioglóssum .. 576	Oviédia .. 499	Passerína .. 449	Persoónia .. 306
Ophioglóssum .. 576	OXALI'DEÆ .. 102	Passiflóra .. 216, 590	Petáclite .. 292
Ophiopóngon .. 523	O'xalis .. 102, 588	Passiflóra .. 217, 218	Petalòdma .. 195
Ophiorhiza .. 600	Oxyánthus .. 261	PASSIFL'OREÆ .. 216	Petalostémum .. 141
Ophiostachys .. 540	Oxybáphus .. 429	Pastinàca .. 251	Petámenes .. 500
Ophióxylon .. 354	Oxycóceus .. 330	Pastinàca .. 249, 251	Petivéria .. 433
Ophiurus .. 563	Oxylóbium .. 123	Patagónium .. 149	Petr'æa .. 417
Ophrys .. 484	Oxypétalum .. 361	Patagónula .. 380	Petròbium .. 306
Ophrys .. 483, 485	Oxy'ria .. 439	Patersónia .. 499	Petrocális .. 25
Opisménus .. 563	Oxy'spora .. 203	Patrímia .. 268	

INDEX GENERUM.

Page.	Page.	Page.	Page.	Page.	Page.
Petromarula .. 593	PHYTOLACEÆ 433	Podalyria .. 123	Portuláca .. 219		
Pétróphila .. 443	Piaránthus .. 259	Podalyria .. 123	Portuláca .. 219		
Petroselínium .. 245	Picòtia .. 378	Podánthes .. 358	Portulacária .. 220		
Pettùnia .. 383	Picrámnia .. 119	Podánthus .. 315	PORTULACEÆ 219		
Peùce .. 475	Picridium .. 279	Podocárpus .. 473	Posoquèria .. 261		
PEUVED'ANEÆ .. 249	Picris .. 276	Podólepis .. 299	Póssira .. 162		
Peucédanum .. 250	Pictétia .. 148	Podolòbium .. 123	Potamogéton .. 478		
Peucédanum 249, 252	Pilea .. 463	PODOPHY'LLEÆ	Potamóphila .. 575		
Peyroúisia .. 499	Pileánthus .. 207		Potentília .. 190, 589		
Pfáffia .. 433	Pilulària .. 575	Podophyllum .. 17	POTENTÍLLÆ .. 187		
Phàca .. 145	Pimeléa .. 450	Podophyllum .. 17	Potérium .. 193		
Phàca 143, 144, 145, 147	Pimpinélla .. 246	Podópterus .. 440	Pòthos .. 479, 595		
Phacélia .. 380	Pimpinélla .. 245	Podospermum .. 279	Pòthos .. 479		
Phalángium .. 524	Pincknýea .. 262	Podostigma .. 361	Poupártia .. 120		
Phalángium 525, 527,	Pinguícula .. 422	Pogónia .. 486	Pourrétia .. 535, 596		
..... 528	Pinus .. 475	Pogónia .. 486	Pozóa .. 243		
Phálaris .. 559	Pinus .. 474, 475	Pogostémón .. 400	PRASI'EÆ .. 413		
Phálaris 554, 559, 567	Piper .. 481	Poinciana .. 169	Prásium .. 413		
PHANEROG'ANEÆ 476	Piper .. 482	Poirétia .. 149	Prasophyllum .. 485		
Pharnâceum .. 55	PIPERACEÆ 481	Poit'æa .. 143	Prémna .. 417		
Pharnâceum .. 220	Piptánthus .. 122	Póivrea .. 196, 590	Prenánthes .. 274		
Phàrus .. 574	Piptátherum .. 555	Polanísia .. 38	Prenánthes .. 274, 277		
PHASEOLEÆ .. 156	Piquéría .. 301	POLEMONTACEÆ	Prescótia .. 486		
Phaséolus .. 157	Piriquéta .. 218	Polemónium .. 367, 594	Prestónia .. 355		
Phaylópsis .. 420	Piscídia .. 143	Polianthes .. 517	Priestleya .. 127		
Phébalium .. 107	Pisónia .. 430	Pollíchia .. 221	Primula .. 423		
Phellândrium 247, 249	Pistácia .. 119	POLLICHI'EÆ .. 221	Primula .. 425		
PHILADELPHEÆ	Pistorinía .. 225	Pollínia .. 561	PRIMULACEÆ .. 423		
..... 206	Pisum .. 154	Polybótria .. 578	Prinos .. 111		
Philadélfus .. 206	Pisum .. 155	Polycarp'a .. 221	Prismatocárpus .. 328		
Philly'rea .. 351, 352	Pitcaírnia .. 535	POLYCARP'EÆ .. 221	Priva .. 418		
Philothèca .. 107	Pittósporum .. 48	Polycárpon .. 221	Próckia .. 40		
Philóxerous .. 432	Pittósporum .. 48	Polycénia .. 414	Pròcris .. 462		
Philóxerous .. 432	Plagiánthus .. 65	Polycnémum .. 437	Proiphys .. 513		
Philydrum .. 546	Plagiòlbium .. 126	Poly'gala .. 46, 586	Prosperinàca .. 201		
Phlém .. 554	Planéra .. 464	Poly'gala .. 47	Prósopis .. 164		
Phlém .. 554	Plantagin'EÆ 428	POLYG'ALEÆ .. 46	Prostánthera .. 414		
Phlòmis .. 405	Plantàgo .. 428	Polygonátum .. 523	Prôtea .. 443		
Phlòmis .. 404, 405	Platanthèra .. 483	POLYGO'NEÆ .. 438	Prôtea 442, 443, 444, 445		
Phlóx .. 368, 599	PLAT'ANEÆ .. 472	Poly'gonum .. 438, 595	PROTEACEÆ .. 442		
Ph'œnix .. 543	Platylòbium .. 126	Poly'gonum .. 434, 439	Prunélla .. 410		
Phœnócoma .. 293	Platylòbium .. 126	Poly'mnia .. 313	Prúnus .. 174		
Pholidòta .. 489	Platypétalum .. 24	POLYPODIA'CEÆ	Prúnus .. 173, 174		
Pholiùrus .. 563	Platy'pteris .. 306	Poly'podium .. 577	Psámma .. 574		
Phórmium .. 534	Platy'stlys .. 155	Poly'podium .. 578, 580	Pseudosophòra .. 122		
Photinia .. 176	Plectránthus .. 413	Poly'pôgon .. 554	Psìàdia .. 299		
Phragmitès .. 574	Plectránthus .. 413	Poly'spora .. 72	Psidium .. 210		
Phry'yma .. 413	Plectronia .. 110	Polystachya .. 489	Psilòtum .. 576		
Phry'nium .. 494, 595	Pleópeltis .. 545	POM'ACEÆ .. 175	Psilurus .. 563		
Phycélla .. 510, 596	Pleròma .. 204, 590	Pomadérris .. 113	Psophocárpus .. 159		
Phyl'lica .. 114	Pleurándra .. 13	Pomària .. 169	Psorálea .. 138		
Phyl'lica .. 114, 115	Pleuróthallis .. 488	Pombália .. 45	Psorálea .. 141		
Phyllánthus .. 458	Pleurothállis .. 487	Poncelétia .. 345	Psychíne .. 35		
Phyllánthus .. 459	Plómaca .. 265	Póngamia .. 161	PSYCHÍNEÆ .. 35		
Phyll'lis .. 265	Plucoméria .. 355	PONTED'EREÆ .. 541	Psychòtria .. 264		
Phylloclàdus .. 473	Poà .. 568	Pontedèria .. 541	Psychòtria .. 260, 264		
Phyllòdium .. 151	Pleurospérum .. 255	Ponthièva .. 485	Ptèlea .. 121		
Phyllolòbium .. 159	Pleurothállis .. 488	Portenschläglia .. 110	PTELE'ACEÆ .. 121		
Phyllòma .. 517	Pleurothállis .. 487	Poràna .. 373	Ptélidium .. 110		
Phyllogýne .. 515	Ploméria .. 265	Porána .. 373	Ptéris .. 582, 598		
Phymatánthus 89, 587	Plukénétia .. 456	Populus .. 471	Ptéris .. 582, 583		
Physàlis .. 383	PLUMBAGI'NEÆ	Poràna .. 373	Pterocárpus .. 161		
Physospérum .. 255	Plumbàgo .. 427	Poranthèra .. 599	Pterocárpus .. 161, 162		
Physostégia .. 406	Plumèria .. 355	Porlièra .. 104	Pterocéphalus .. 271		
Phytéléphas .. 478	Pòa .. 567, 569, 570	Portenschláglia .. 110	Pteroneúrum .. 24		
Phyteúma .. 328	Pocóckia .. 134	Portlândia .. 262	Pterònìa .. 285		
Phyteúma .. 593		Portlândia .. 262	Pterospérum .. 69		
Phytolacca .. 433					

INDEX GENERUM.

615

Page.	Page.	Page.	Page.
Pterostylis 487	Retanilla 113	Róthia 274	SANTALACEÆ 450
Pterygodium .. 484	Rétzia 369	Róthia 156	Sántalum 450
Ptilostéphium .. 308	Rhabdóchloa 571	Rottboéllia 562	Santolina 315
Ptychotís 245	Rhagadiolus 275	Rottboéllia 562, 563	Santolina 315
Puerária 140	Rhagadiolus 275	Rottlèra 457	Sanvitália 308
Pulicária 291	Rhágodia 434	Róurea 121	SAPINDACEÆ 83
Pulmonária 375	RHA'MNEÆ .. 112	Roxbúrgchia 534	Sápindus 84
Pulmonária 374	Rhámnuus 112	Royéna 350	Sápium 458
Pulsatilla 3	Rhámnuus 112, 113, 464	Rúbia 266	Sápium 458
Pulten'æa 124	Rhápis 542	RUBIACEÆ .. 260	Saponária 51
Pulten'æa 123	Rhapóntica 287	Rúbus 187, 589	Saponária 51
Púnica 195	Rhéum 440	Rúbus 189	SAP'OTEÆ .. 348
Pupátiu 432	Rhéxia 203	Rudbékia 310	Sáraca 169
Purgosea 223	Rhéxia .. 203, 204	Rudbékia .. 307, 310	Sáracha 383
Púrshia 193	RHEXI'EÆ .. 203	Rudólphia 161	Sarcánthus 491
Puschkínia 525	Rhinánthus 395	Ruellia 420, 595	Sarcocápnos 20
Pycnánthemum .. 401	Rhipidodéndron .. 519	Ruellia 420	Sarcocaudon 88
Pycnostächys .. 414	RHIPSAL'DEÆ .. 238	Ruizia 69	Sarcocéphalus 263
Pyréthrum 321	Rhípsalis 238, 591	Ruizia 84	Sarcochilus 489
Pyrola 341	RHIZOB'OLEÆ .. 83	Rulíngia 67	Sarcocécca 456
Pyrulária 451	Rhizóphora 197	Rulíngia 219	Sarcólobus 360
Pyrus 177	RHIZOPH'OREÆ	Rúmex 439	Sarcophylum 129
Pyrus.. 176, 177, 178	Rhodiola 197	Rúmex 439, 440	Sarcostémma 361
Pyxidanthéra 342	Rhododéndron .. 342	Rúmia 244	Sarothra 77
Quádria 599	Rhodòra 343	Rúmia 251	Sarracénia .. 18, 585
Quálea 590	RHODOR'ACEÆ .. 342	Rúppia 478	SARRACENTEÆ 18
Quássia 108	Rhopália 447	RúsCUS 523	SATUREÍ'NÆ .. 401
Quélta 514	Rhús 117	Russelia 390	Satureja 401
Quérkus 465	Rhýchosia 157	Rúta 105	Satyrium 484
Querària 221	Rhynchóspora .. 549	Rúta 105	Satyrium .. 483, 485
Querària 220	Ribes 238, 591	Rútta 105	Sauraúja 72
QUER'ACEÆ .. 221	Richárdia 480	Rútta 105	SAURAÚ'JEÆ .. 72
Quisqualis 196	Richardsónia .. 265	Rútscheria 79	Saurúrus 481
Quivisia 85	Rícinos 458	Ryánea 40	Saussurea 281
Radiola 58	Ricótia 24	Sábá .. 542	Sauvagésia 45
Ráfnia 126	Riedléia 68	Sabbátia 363	Saxífraga .. 239, 591
Ráfnia 123, 126	Rindéra 578	Sabicea 260	SAXIFR'AGEÆ .. 239
Rajânia 521	Ripídium 560	Sáccharum 560	Scabiôsa 269
Ramónda 382	Ripgónum 523	Sáccharum 560	Scabiôsa 269, 270, 271
Rândia 260	Ritchièa 48	Sagerétia 112	Sca'volá 324
RANUNCUL'ACEÆ .. 1	Rittéra 162	SALIC'ARIEÆ .. 201	Scália 299
Ranúnculus 4	Rívina 433	SALI'CEÆ .. 468	SCANDIC'INEÆ .. 254
Ranúnculus .. 7	Robértia 276	Salicórnia 436	Scándix 254
RAPH'ANEÆ .. 35	Robertsónia .. 239, 240	Salisbúria 473	Schæfféria 115
Raphánus 36	Robínia 143	Sálix 468	Schedonorus 571
Raphiolépis .. 176	Robínia 141, 142, 143,	Sáimea 307	Schelhamméra .. 539
Rapístrum 36 144, 168	Sálmia 480	Scheuchzéria .. 477
Rapístrum .. 31	Róchea 223	Salomónia 47	Schinus 119
Raspália 115	Rodigia 276	Salpiánthus 430	Schinus 119
Rauwólphia 353	Rodigia 276	Salpíglóssis .. 383, 594	Schiveréckia .. 25
Rauwólphia 353	Rodrigüezia .. 487	Salsóla 434	Schiz'æa .. 577, 598
Ravenália 495	Roëlla 328	Salsóla 434, 435	Schizándra 16
Reaumúria 235	Roëlla 327	Sálvia 410	SCHIZA'NDREÆ .. 16
REAU'MURIA .. 235	Rœmèria 19, 585	Sálvinia 576	SCHIZÁNTHUS 396
Redoutéa 63	Rœmèria 571	Sámara 348	SCHIZOPE'TALEÆ .. 37
Reevésia 586	Roepéria 104	Sambucus 257	Schizopétalum .. 37
Relhânia 307	Róhdea 480	Samélus 426	Schkuhria 306
Remírea 552	Rolándra 287	Sam'yda 116	Schmáltzia 119
Renanthéra .. 491	Rondeletia 262	SAMY'DEÆ .. 116	Schmidélia 84
Reneálmia 493	Rósa 178	Sandóricum 85	Schmidítia 554
Reneálmia .. 493, 498	ROS'ACEÆ .. 178	Sanguinária .. 19, 585	Schœnòdum 547
Requiénia 132	Roscóea 492	Sanguisórba 193	Sch'enus .. 549, 597
Reséda 39	Roséa 433	SANGUISO'RBEÆ .. 192	Sch'enus .. 549, 550
RESED'ACEÆ .. 39	Rosmarínus 404	Sanícula 244	Schœpfia 347
RESTI'ACEÆ .. 546	Rostrária 574	SANIC'ULEÆ .. 243	Schóllerá .. 330, 541
Réstio 547		Sanseviéra 517	Schotia 172
			Schoús'bæa .. 196

INDEX GENERUM.

Page.	Page.	Page.	Page.	Page.	
Schránka	163	Seriola	276	SMILA'CEÆ	522
Schrebera	111	Seriola	276	Smilacina	523
Schubertia	474	Seriphium	288	Smilax	522
Schúltzia	245	Serissa	264	Smíthia	149
Schwénckia	396	Serpícula	200	SMY'RNEÆ	255
Scilla	527, 596, 599	Serrátula	281	Smyrnium	255
Scilla	527, 528	Serrátula 281, 282, 283		Smyrnium	245, 251
Sciadophyllum	256	Serrùria	445	Sobolewskia	32
Scírus	550, 598	Sersalísia	349	Sója	158
Scírus	550	SESAMEÆ	367	Solándra	388
SCITAMINEÆ	491	Sésamum	367	SOLANEÆ	380
SCLERA'NTHEÆ	221	Sesbània	143	Solánum	385
Scleránthus	221	Sesbània	143	Solánum	385, 387
Sclerà	549	Séseli	248	Soldanélla	423
Sclerocárpus	312	Séseli	245, 248	Soldevilla	276
Scleróchloa	569	SESELI'NEÆ	247	Sólea	45
Sclerosciádium	247	Seslèria	567	Solidàgo	297
Sclerothámnus	124	Seslèria	567	Solidàgo	298, 307
Scolopéndrium	581	Sesúvium	234	Soliva	317
Scolopéndrium	578	Setària	553	Sónchus	278
Scólymus	280	Séthia	80	Sónchus	277, 279
Scopària	389	Seymèria	396	Sonneràtia	210
Scopòlia	382	Seymóuria	92	Sophòra	122, 589
Scopòlia	121	Shephérdia	451	Sophòra	122, 123
Scorpiurus	148	Sherárdia	266	SOPH'OREÆ	122
Scorzonèra	278	Sherárdia	267	Sophronítis	489
Scorzonèra	279	Sibbaldia	192	Sórbus	177
SCORZON'EREÆ	278	Sibthórpia	394	Sorghum	561
Scóttia	126	Sícos	215	Sorocéphalus	445
Scrophulària 391, 594		Sida	63, 586	Soulángia	114
SCROPHULA'RÍ-		Sida	63, 64, 65	Southwéllia	66
NÆ	389	Sideránthus	298	Sowerb'æa	532
Scutellària	410	Sideritis	407	Spanalzánia	192
Scútia	113	Siderodéndron	264	Spanánthe	243
Scyphánthus	218	Sideróxylon	349	Sparáxis	501
Scytália	84	Sideróxylon 115, 348, 349		Sparáxis	501
Seafórtia	544	Siegesbékia	308	Sparagñium	479
Seb'æa	364	Sievérsia	190	Sparaganóphorus	315
Secale	564	Sievérsia	189	Sparrmánnia	70
Secale	566	Silàus	249	Spartina	553
Secamòn	356	Silène	51	Spartiánthus	129
Sèchium	215	SIL'ENEÆ	48	Spártium	129
Securidàca	47	Siler	252	Spártium 129, 130, 131	
Securígera	148	Siler	252	Spatialánthus	503
Securinèga	460	SILER'INEÆ	252	Spatálla	445
Sédum	225	Sílphiwm	314	Spathélia	121
Sèdum	225	Sílybum	282	Spathódea	366
SELAGI'NEÆ	414	Simàba	108	Spatulária	240
Selàgo	415	Simarùba	108	Speculária	593
Selàgo	415	SIMAR'UBEÆ	108	Spennéra	203
Selinum	249	Símsia	599	Spergùla	55, 586
Selinum 248, 249, 250,		Símsia	307, 311	Spermacòce	265
..... 251		Sínàpis	34, 585	Spermacòce	265
Sellòa	301	Sínàpis	35	SPERMACOCEÆ	265
Semicárpus	117	Singàna	79	Spermadíctyon	263
Semperfívum	226	Símíngia	329	Spermáxyrum	75
Semperfívum	226	Siphonánthus	414	Sphácele	406
Senàcia	48	Siphònìa	458	Sphæránthus	288
Senàcia	110	Síson	245	Sphæròlóbium	124
Senebiéra	31	Síson 245, 246, 247, 252		Sphæroph'ysa	144
Senecíllis	304	SISY'MBREÆ	28	Sphæròspora	501
Senècio	302	Sisy'mbrium	29	Spallerocárpus	255
Senècio	305	Sisy'mbrium 22, 29, 35		Sphenógyne	312
Séptas	222	Sisyrínchium	498	Sphenótoma	345
Seràpias	484	Sium	246	Spielmánnia	418
Seràpias	486	Sium 245, 246, 247, 251		Spiégelia	364
Seriàna	83	Sloànea	71	Spiłánthes	306
Seríngia	67	Smeathmánnia	216	Spiłánthes	306, 307
				Spinàcia	436
				Spir'æa	194
				Spir'æa	194
				SPIR'EACEÆ	193
				Spirálepis	292
				Spiranthera	108
				Spiranthes	485
				SPIRO'L'OBÆ	36
				SPONDIACEÆ	120
				Spóndias	120
				Sporòbolus	557
				Spreikélia	509
				Sprengélia	345
				Staàvia	115
				Stächys	406, 594
				Stächys	406, 407
				Stachytárphe	418
				STACKHOUSEÆ	
				Stackhóusia	452
				Stæhelina	285
				Stæhelina	281
				Stanhópea	490
				Stanlèya	31
				Stapélia	357
				Stapélia 356, 357, 358,	
					359
				Staphyléa	109
				STAPHYLEACEÆ	109
				Stárkea	298
				Státice	426
				Stauracánthus	129
				Stegànìa	581
				Stélis	487
				Stellària	55
				Stellària	55
				Stellària	449
				Stemòdia	389
				Stenanthèra	346
				Stenocárpus	447
				Stenochilus	419
				Stenomésson	513
				Stenorhy'nchus	485
				Stenospérmum	209
				Stenotáphrum	563
				Stephània	39
				Stercùlia	66, 599
				STERCUL'EÆ	66
				Stereóxylon	239
				Sterigma	32
				Steriphoma	39
				Sternbékia	79
				Sternbérghia	505
				Stèvia	22
				Stèvia	300
				Stèvia	301, 306
				Stilàgo	460
				Stillíngia	458
				Stípa	556
				Stizolòbium	159
				Stob'æa	284
				St'œbe	288
				Stokesia	285
				Stratiòtes	476
				Stravàdiu	213
				Strélitzia	495
				Streptanthèra	501
				Stréptium	418

Page.	Page.	Page.	Page.
Streptocárpus 397	Támarix 203	Thelymitra 486	Tradescántia 541, 542
Stréptopus 523	Támarix 203	Thenárdia 355	Tragacántha 147
Stróbos 475	TAMEÆ 522	Theobróma 67	Trágia 460
Sstrophanthus 354	Tamónia 418	Theophrásta 353	Trágium 246
Strumária 510	Támus 522	Thermópsis 123	Tragopogón 279
Struthiolá 449	Tanacétum 316	Thésium 450	Tragopýrum 439
Struthiópteris 582	Tanacétum 316	Thèsium 450	Trágus 557
Strychnos 353	Tanacétum 389	Thespésia 63	Trápa 200
Stuártia 73	TARAX'ACEÆ 274	Thláspi 27	Tratteníckia 306
Stuártia 73	Tarchonánthus 315	Thláspi 27, 31, 32	Travóa 588
Stúrmia 557	Tasmánnia 13	THLÁSPI'DEÆ 27	Trérsia 453
Stylándra 361	Tauschériá 32	Thomásia 67	TREMA'NDREÆ 47
STYLI'DEÆ 324	Taxánthema 427	Thrínax 544	Treviránia 390
Stylidium 324	Taxódium 474	Thrúnica 275	Tréwia 557
Stylosánthes 149	TAXI'NEÆ 473	Thryális 81	Tríánthema 219
Stypándra 525	Táxus 473	Thúja 474	Tribráchia 489
Styphélia 346	Táxus 473	Thúja 473	Tribulus 104
Styphélia 346	Tecòma 366	Thunbérzia 419	Trichæta 573
STYPH'ELLEÆ 346	Tectóna 418	Thýmra 409	Trichéra 269
STYRACI'NEÆ 347	Teédia 389	Thýmra 409	Trichília 85
St'yrax 347	Teesdália 27	THYMEL'ÆÆ 448	TRICHI'LIEÆ 85
Suaèda 435	TELEPHI'EÆ 220	Th'ymus 401	Trichocéphalus 114
Suárdia 558	Téléphiúm 220	Th'ymus 399, 409	Trichóchloa 557
Subulária 37	Telfária 215	Thysanótus 532	Trichodésma 378
SUBUL'ARIEÆ 37	Tellíma 242	Thyselinum 250, 251	Trichódium 556
Succísa 269	Telopéa 447	Tiarélla 242	Trichódium 556
Succívia 35	Templetónia 126	Tiarélla 242	Tricholæna 560
Sutherlándia 144	Tenória 247, 251	Tiarídium 374	Trichómunes 583
Swainsóna 144	Tephrosía 141, 589	Tibouchina 204	Trichonémia 503, 596
Swártzia 162	Tephrosía 156	Ticoréa 108	Trichóphorum 552
SWARTZI'EÆ 162	Tephrothámnus 126	Tigárea 193	Trichophyllum 322
Sweetia 156	Terámnus 156	Tigrídia 499	Trichorhiza 491
Swértia 363	TEREBINTH'ACEÆ 247	Tigrídia 497	Trichosánthes 215
Swieténia 86	Terminália 195	Tilia 71	Trichostéma 403
Swieténia 86	TERMIN'ALIEÆ 195	Tiliacóra 15	Trichostéma 399
S'yagrus 543	Ternstremia 72	Till'æa 222	Tricoryne 525
Symphíandra 593	TERNSTRÖEMIA'-CEE 72	Tillándsia 535	Tridactylites 241
Symphónia 79	Terpnánthus 108	Tithónia 312	Tridax 308
SYMPH'ONIEÆ 79	Testudinária 522	Tithymalus 453, 454	Tridens 567
Symphóría 258, 592	Tetrácerá 12	Tittmánnia 116	Tridéntea 358
Sy'mphytum 375	Tetradíum 121	Tittmánnia 393, 394	Trientális 425
Sympíèza 332	Tetragónia 234	Tocoça 205	TRIFOLI'EÆ 132
Symplocárpus 479	Tetragónia 201	Toddália 121	Trifolium 135
SYMPLOCI'NEÆ 347	Tetragonolóbus 138	Tódea 577	Trifolium 134, 137
Sy'mplocos 347	Tetragonothéca 307	Tofieldia 540	Triglóchin 477
Synándra 404	Tetramérium 264	Tólpis 275	Trigonélla 134
Syncárpha 281	Tetranémá 149	Tonsélla 79	Trigónia 80
Synedrélla 308	Tetranémá 441	Tontélea 79	Trigónis 84
Synnótia 501	Tetranthéra 441	TORDYLI'NEÆ 252	Trífilum 523
Syntherísmo 558	Tetrapógon 564	Tordylíum 252	Trinia 245
Sy'tenia 30	Tetrápteris 81	Tordylíum 251, 252	Triodiá 567
Syringa 352	Tetrahéca 47	Torénia 393	Triodiá 567
Szy'gium 211	Tetrazy'gia 205	Torilis 254	Triópteris 81
Tabernæmontána 354	Tefúcrum 402	Torilis 253	Trióstéum 259
Tácca 481	Thalámia 473	Tormentilla 190	Triphásia 75
TACC'ACEÆ 481	THALAMIFL'ORÆ 1	Törtula 418	Triphora 486
Tachigália 169	Thàlia 494	Tournefórtia 378	Tripiñnária 389
Tacsónia 218	Thalíctrum 2	Tournefórtia 374, 379	Triplaris 440
Tænitís 582	Thamnéa 116	Tourrézia 367	Tripsacum 563
Tagétes 305	Thamnochórtus 547	Toucharòa 172	Tripsacum 563
Taliéra 543	Thápsia 252	Trachélium 328	Triptilión 280
Talaúma 584	Thápsia 248	Trachymène 243	Trisetum 568
Talínum 219	THAPS'I'EÆ 252	Trachymène 243	Trisétum 567, 573
Tamaríndus 169	Thásپium 248	Trachynòtia 553	Tristánia 207
TAMARISCI'NEÆ 203	Thásپium 245	Trachyspérnum 245	Tristegis 558
	Théa 73	Trachytélla 584	Triteleia 532
	Thely'gonum 463	Tradescántia 541, 597	

INDEX GENERUM.

Page.	Page.	Page.	Page.
Triticum 564	VACCINIÆ 329	Viola 43, 45	Wulfenia 396
Triticum 564, 565, 566, 573	Vaccinium 329	VIOLARIEÆ	Wurmbea 540
Tritoma 517	Vaccinium 330	Virgilia 122	W'ylia 254
Tritonia 502	Valantia 266	Virgilia 122	Xanthium 314
Triumfetta 70	Valantia 266, 267	Viscaria 53	Xanthochymus 78
Trixago 395	Valeriana 268	Viscum 259	Xanthorrhæa 534
Trixis 280	VALERIANÆ 268	Vismia 76	Xanthosia 243
Trixis 280	Valerianella 269	Vismaga 245	Xenopöma 419
Trizeixis 488	Vallaris 356	Visnea 350	Xeranthemum 294
Trochiscanthes .. 248	Vallésia 353	Vitex 416	Xeranthemum 292, 293, 294
Trochocarpa .. 346	Vallisneria 476	Vitex 417	Xerophyllum 540
Tróllius 7	Vallotta 512	Vitis 87	Xeròtes 545
Tromótriche 358	Vanda 491	Vitis 87	Ximenësia 307
TROPÆOLEÆ 101	Vanda 491	Vittaria 582	Ximènia 75
Tropæolum .. 101	Vandellia 394	Voandzeia 168	Ximènia 104
Trophis 450	Vangueria 260, 600	Vochysia 197	Xiphidium 504
Tróximon 276	Vanilla 491	VOCHYSIEÆ	Xiphópteris 578
Tróximon 276	Varronia 379, 380	Vohiria 363	Xylòbium 489
Tulbagia 517	Vascoa 126	Volkamèria 416	Xylomèlum 447
Tùlipa 536	Vauánthes 223	Volkamèria 415, 416	Xylophylla 459
TULIPACEÆ 536	Velèzia 54	Vouápá 172	Xylopìa 15
Tupistra 480	Vélia 35	Wachendorfia 504	Xylosteum 258
Turgènia 254	Vélia 35	Wahlenbergia 325, 593	Xyris 546
Turnèra 218	VE'LLEÆ 35	Waldsteinia 189	Xysmalòbium 361
TURNERIACEÆ 218	Velleia 324	Waldsteinia 189	Yucca 536, 597
Turpínia .. 109	Velleia 324	Walkéra 109	Zacintha 276
Turpínia .. 109, 149	Vellòsia 505	Wallchia 543	Zaluzània 318
Turræa 85	Veltheimia 517	WALLICH'IEÆ	Zàmia 476
Turritis 22	Venâna 599	Wallerthia 249	Zannichéllia 478
Turritis 22, 23, 34, 585	Ventenatia 324, 346	Wallerthia 417	Zandòria 540
Tussilago 302	Ventilago 114	Walthèria 68	Zapània 418
Tussilago 280	Verâtrum 540	Wangenheimia 560	Zanthonhùza 12
Tylóphora 589	Verbascum 380	Watsonia 500	Zanthoxylum 107
Typha 479	Verbascum 382	Webera 261	Zèa 574
TYPHINEÆ 479	Verbena 418	Webera 261	Zephyránthes 505
Ugèna 577	Verbena 418	Wedèlia 309	Zexmènia 309
'Ulex 129	VERBENACEÆ 415	Wedèlia 313	Zièrea 107
Ulmus 464	Verbesina 308	Weinmannia 242	Zietènia 407
Ulmus 464	Verbesina 308	Wendländia 15	Zigadénus 540
UMBELLIFERÆ	Vernonia 289, 592	Westónia 156	Zilla 35
..... 243	Vernonia 289	Westríngea 402	ZILL'EÆ 35
Umbilicus 225	VERNONIACEÆ 288	Whitleya 383	Zíngiber 492
Uncària 263	Verónica 396	Wibeliu 274	Zinnia 307
Uniola 571	Verónica 396, 399	Wiborgia 308	Zizània 575
Unònà 15	Verticórdia 207	Wigándia 369	Zizia 245
Urâchne 555	Vesicaria 24	Wildenòvia 547	Ziziphora 404
Urânia 495	Véstia 383	Willemètia 113	Zizyphus 112
Urâria 149	Vibòrgia 128	Willughbèia 356	Zizyphus 112, 464
Uraspérnum .. 255	Viburnum 257	Windsòria 567	Zœgea 285
Urêna 60	Vicia 153	Wintèra 13	Zórnia 149
Uropétalon 528	Vicia .. 143, 154, 156	Winterâna 13	Zórnia 150
Urtica 462	Vici'EÆ 153	WINTEREÆ	Zosímia 252
Urtica 462	Vieusseùxia 498	Wistèria 157	Zostèra 478
URTI'CEÆ 460	Vigna 159	Witheringia 388	Zóysia 554
Urvil'a 83	Viguéra 310	Witsénia 499	Zuccagnia 528
Utriculària 423	Villârsia 364	Wittlesbáckia 73	Zuringèra 108
Uvària 15	Vilmorínia 140	Woodfórdia 202	Zygotépàlum 496
Uvària 15	Viminària 124	Woódsia 579	ZYGOPHYLLEÆ
Uvularia 539	Vînca 354	Woodwárdia 581 104
Uvularia 523, 539	VINI'FERÆ	Wórmia 13	Zygophyllum 104
Vaccària 51	Viola 43, 586	Wrightia 354	Zygophyllum 104

ENGLISH INDEX.

Page.	Page.	Page.	Page.
Abele-tree 471	Balsam-herb 421	Birch-tree 467	Burnet 193
Achania 61	Balsam of Capevi 171	Bird Cherry 175	Burnet Saxifrage 246
Adam's needle 536	Balsam of Peru 122	Bird's-eye Primrose 424	Bur Parsley 253
Adder's-tongue 576	Balsam of Tolu 122	Bird's-foot 148	Bur-reed 479
African Almond 446	Balsam-seed 122	Bird's-foot Trefoil 134,	Butcher's Broom 523
African fleabane 315	Balsam-tree 78, 220 137	Butter and eggs 514
African Lily 517	Balsam-wood 122	Bird's-head 488	Butter-bur 302
African Marygold 305	Bamboo-cane 574	Bird's-nest 486	Butterfly-plant 483, 488
Agrimony 192	Bane-berries 11	Bird Pepper 384	Butter-nut 83
Akee-tree 84	Banana tree 495	Bird's-tongue 304, 539	Butter-tree 78, 349
Alaternus 112	Barbadoes Gooseberry	Birthwort 451	Butterwort 422
Alder 468	Barberry 16, 584	Bishop's-weed 245	Button-tree 196
Alexander's 255	Barley 558	Bitter-sweet 385	Button-weed 265
Alkekengi 384	Barren-wort 16, 584	Bitter Vetch 156	Button-wood 263
Alligator Pear 441	Base Rocket 39	Bitter-wood 15	Cabbage 33
All-seed 58, 221, 435	Basil 413	Blackberry 188	Cabbage-tree 543
Allspice 212	Bastard Balm 409	Black Bryony 522	Cajuputi-tree 207
Allspice-tree 194	Bastard Cabbage-tree	Black Knapweed 285	Calabash-tree 389
Allum-root 242 168	Black Parsley 253	Calaba-tree 78
Almond 173	Bastard Cardamom 492	Black Saltwort 428	Calamint 409
Almond Peach 173	Bastard Cedar 67, 86	Bladder Ketmia 62	Calathian Violet 362
Aloe-wood 117	Bastard Cress 27	Bladder-nut 109	Calf's-snout 390
Althea-frutex 61	Bastard Cypress 549	Bladder Senna 144	Caltrops 104
Amaranth 430	Bastard Flower-fence	Bleaberry 329	Cammock 132
Amber-tree 265 164	Blessed Thistle 287	Campion 51
Amboyna pitchtree 474	Bastard Hare's-ear 265	Bloodwort 19	Camphire-tree 441
American birds-eye 424	Bastard Indigo 142	Blue-bottle 286	Canada Rice 575
American Cowslip 423	Bastard Mangeneel 355	Bog-rush 549	Canary Tea 63
Anchovy Pear 213	Bastard Orpine 459	Bonacie-bark 449	Candleberry Myrtle 472
Andaman red wood 161	Bastard Pimpernell 426	Borage 378	Candy-tuft 27
Angelica 249	Bastard Quince 177	Borecole 33	Cane-brake 574
Angelica-tree 256	Bastard Saffron 284	Box-thorn 388	Canterbury bells 327
Anise 246	Bastard Vervain 418	Box-tree 456	Cape Aster 305
Aniseed-tree 13	Bastard Vetch 145	Brake 582	Cape Holly 110
Anota 40	Bawd-money 249	Bramble 178	Cape Jasmine 261
Ape Orchis 483	Bay-tree 441	Brasiletto 168	Cape Phillyrea 110
Apple 177	Bead-tree 85	Bread-fruit 462	Caper-bush 38, 585
Apple-berry 48	Bean 153	Bread-nut 462	Caper Spunge 454
Apricot 174	Bean-caper 104	Bread-root 138	Carambola-tree 102
Arbor-vita 474	Bean-trefoil 122	Broccoli 33	Caraway 245
Archangel 406	Bear-berry 330	Brome-grass 573	Cardamon 493
Arracacha 255	Bearbind 370	Brook-lime 398	Cardinal-flower 323
Arrow-grass 477	Bear's-breech 419	Brookweed 426	Cardoon 284
Arrow-head 477	Bear's-ear Sanicle 423	Broom 131	Carline thistle 284
Arrow-root 495	Bedstraw 266	Broom-rape 422	Carnation 50
Artichoke 284	Bengal Quince 75	Brown Knapweed 286	Carob-tree 169
Asarabacca 451	Beetle 467	Brussels sprouts 33	Carrión-flower 357
Ash-tree 350	Bee Ophrys 484	Bryony 214	Carrot 253
Asparagus 533	Beet 436	Buck-bean 364	Cascarilla 457
Aspen 472	Belladonna Lily 506	Buckler Mustard 27	Cashew-nut 117
Asphodel 524	Bell-flower 325	Buck's-eye tree 83	Cassava 458
Atamasco Lily 505	Bengal Quince 75	Buckthorn 112	Cassia 441
Auricula 425	Benjamin-tree 441	Buck-wheat 438	Cassioberry-bush 258
Avens 190	Bent-grass 556	Buck-wheat tree 332	Castor-oil nut 458
Awl-wort 37	Berry-bearing Alder	Buffalo-clover 136	Catalpa-tree 365
Azarole 176 113	Bugle 403	Catchfly 51
Bahama red-wood 113	Betle 482	Bugloss 377	Cat-chop 228
Balm 409	Betony 406	Bugwort 11	Catmint 407
Balm of Gilead 409	Bilimbi-tree 102	Bullace-tree 174	Cat's-ear 276
Balsam 101	Billberry 329	Bullet-grape 87	Cat's-tail 479
Balsam Apple 215	Bindweed 369	Burdock 280	Cat's-tail grass 554

ENGLISH INDEX.

Page.	Page.	Page.	Page.
Cat Thyme 403	Cornelian Cherry 257	Dragon's-head 408	Fox-chop 227
Cauliflower 33	Corn Cockle 54	Dragon-tree 533	Foxglove 392
Cedar 474, 475	Corn-flag 500	Dropwort 194	Fox-Grape 87
Cedar of Goa 474	Cornish Moneywort 394	Drumstick-tree 169	Foxtail-grass 554
Cedar of Lebanon 475	Costmary 315	Duck-weed 478	Fraxinella 105, 588
Celandine 19	Cotton-grass 552	Duck's-foot 17	French Honeysuckle 156
Celery 245	Cotton-Rose 294	Dumb Cane 481	French Marygold 305
Centaury 285, 592	Cotton-Thistle 283	Dwarf fan palm 542	French Sorrel 440
Chamomile 318	Cotton-tree 63	Dyer's green-weed 130	Friar's-cowl 480
Charlock 34	Couch-grass 565	Dyer's-weed 39	Fringe-tree 351
Chaste-tree 416	Cowbane 244	Earth-nut 167, 246	Fritillary 537
Chaw-stick 114	Cow-berry 330	Earth Pea 155	Frog-bit 476
Cheese-rennet 267	Cow-grass 135	Egg-plant 387	Frog Orchis 483
Cherimolli 14	Cowitch 159	Egyptian-thorn 166	Fumitory 20
Cherry-tree 174	Cow Parsnip 251	Elastic gum tree 458	Furze 129
Chervil 254	Cowrie Pine 471	Elder 257	Fustick-wood 463
Chestnut 467	Cowslip 423	Elecampane 290	Gambier-tree 263
Chestnut Oak 466	Cow-tree 462	Elephant Apple 75	Garden Balsam 421
Chicasaw 174	Cow-wheat 395	Elephant-foot 288	Garland flower 493
Chickling Vetch 155	Crab 177	Elephant's-foot 522	Garlic 530
Chick Pea 153	Cranberry 330	Elm-tree 464	Garlic-Pear 38
Chickweed 55	Crane's-bill 88	Enchanter's Nightshade 200	Genip-tree 261
Chile Pine 474	Creeping Cereus 236	Endive 280	Gentian 362
China Aster 297	Cress Rocket 35	English Mercury 434	Germanander 402
Chinese varnish 118	Crosswort 265	Ermine-chop 228	German madwort 377
Chives 530	Crown Imperial 538	Eryngo 244, 591	Giant Fennel 250, 591
Chocho 215	Crow-foot 4, 584	Evening-flower 503	Ginger 492
Chocolate-nut 67	Crow-garlic 530	Evening Primrose 198	Gingerbread Plum 173
Christmas Rose 7	Crow's-foot-grass 563	Evergreen Thorn 175	Ginseng 256
Christ's thorn 112	Cuckold-tree 166	Everlasting 291	Gladwyn 496
Cicely 254	Cuckoo-flower 23	Everlasting Pea 155	Glasswort 436
Cinnamon 441	Cucumber 214	Eye-bright 394	Glastonbury-thorn 176
Cinquefoil 190, 589	Cumin 252	Fan Palm 543	Globe Amaranth 432
Citron 75	Cup-flower 218	Feather-grass 556	Globe-flower 7
Clary 412	Currant 238, 591	Felwort 363	Globe Thistle 287
Clearing-nut 353	Custard Apple 14	Female Fern 580	Goat's-beard 194, 279
Cleavers 267	Cypress 474	Fennel 247	Goat's Rue 141
Cloth of Gold 504	Daffodil 514	Fenugreek 134	Goat's Thorn 147
Clove 49	Dahlia 310	Fescue-grass 572	Gold of pleasure 31
Clover 136	Dahoon Holly 111	Feverfew 321	Golden-pert 396
Clove-tree 212	Daisy 299	Feverwort 259	Golden Poley 403
Clown's all heal 407	Dammar Pine 474	Fiddle-wood 417	Golden-rod 297
Club-moss 576	Dandelion 274	Fig Marygold 227	Golden-rod tree 436
Club-rush 550	Darnel 567	Fig-tree 460	Golden Saxifrage 242
Cochineal-fig 237	Date Palm 543	Figwort 391	Golden Thistle 280
Cockscomb 431	Date Plum 349	Filbert 467	Goldy-locks 298
Cocksfoot-grass 570	Day Lily 516	Filmy-leaf 583	Gooseberry 238
Cockshead 152	Deadly Carrot 215	Fiorin-grass 556	Goosefoot 434
Cockspur-thorn 175	Deadly Nightshade 383	Fir-tree 475	Gourd 213
Cocoa-nut-tree 544	Dead Nettle 405	Fishbone Thistle 283	Gout-weed 235
Cocoa-Plum 173	Deodar 475	Fish-poison 142	Grains of Paradise 492
Codlins and cream 198	Deptford Pink 49	Flit Pea 126	Grape 87
Coffee-tree 264	Devil's-bit 269	Flax 58, 586	Grape Hyacinth 528
Cogwood-tree 441	Dewberry 188	Flax Lily 534	Grass-tree 545
Coleseed 33	Dill 251	Feebleane 289	Great Burnet 193
Colewort 35	Dittany of Crete 402	Fleawort 291	Greater Knapweed 286
Colic-root 517	Dock 439	Flixweed 29	Greek Valerian 367
Colpoon-tree 110	Dodder 373	Flower de luce 495	Green Dragon 480
Coltsfoot 302	Dog-chop 227	Flower-fence 169	Green-man 483
Columbine 8	Dog's-bane 356	Flowering Ash 351	Green Quinoa 435
Comfrey 375	Dog's Cabbage 463	Flowering Fern 577	Gromwell 374
Common Cress 31	Dog's-tail grass 553	Flowering Raspberry 188	Ground Ivy 408
Common Dragon 480	Dog's-tooth Violet 539,	Flowering Rush 477	Ground Pine 403
Common Orpine 225 597	Fly Honeysuckle 258	Groundsel 302
Conrajerva-root 464	Dogwood 257	Fly Ophrys 484	Groundsel-tree 289
Coral-tree 160	Doob-grass 558	Fool's Parsley 247	Guaco 300
Coriander 256	Double Cocoa-nut 543		Guava 210
Cork-tree 466	Dragon 479		Guilder Rose 257

Page.	Page.	Page.	Page.
Guernsey Lily 509	Indian-corn 574	Lettuce 277	Meadow Rue 2
Gum Arabic 166	Indian Cress 101	Licca-tree 108	Meadow Saffron 539
Gum Cistus 41	Indian Cucumber 523	Lignum vitæ 104	Meadow Saxifrage 248
Gum Succory 277	Indian Fig 237, 591	Lilac 352	Meadow-sweet 194
Gum-tree 534	Indian Lotus 17	Lily 538	Medick 132
Hair-grass 559	Indian Madder 262	Lily of the valley 523	Medlar 175
Hairy green-weed 130	Indian Millet 561	Lily Pink 532	Medusa's-head 453
Hard-grass 563, 567	Indian Shot 494	Lily-thorn 262	Melic-grass 560
Hare's-ear 246	Indigo 139	Lime 75	Melilot 134
Hare's-foot Fern 583	Ipecacuanha 454	Lime-tree 71	Melon 214
Hare's-foot Trefoil 135	Irish Heath 341	Lion's-ear 404	Melon Thistle 235
Hare's-tail Grass 559	Iron-tree 264	Lion's-foot 274, 291	Mercury 452
Hart's-tongue 581	Iron-wood 209, 349	Lion's-leaf 16	Mexican Lily 507
Hartwort 252	Ironwort 408	Liquorice 141	Mezereum 448
Hassagay-tree 111	Ivy 256	Lizard Orchis 483	Michaelmas daisy 296
Hatchet Vetch 148	Jaca-tree 464	Lizard's-tail 481	Mignonette 39
Hawkweed 271	Jack in a box 442	Lobel's Catchfly 53	Milfoil 319
Hawthorn 176	Jacobea Lily 509	Locust-tree 143, 172	Milk Parsley 249
Heart's-ease 45	Jalap 371	Logwood 169	Milk Thistle 282
Heart-seed 83	Jamaica Birch 120	London-pride 240	Milk Vetch 145, 589
Heath 332	Jamaica Dogwood 143	London Rocket 29	Milkwort 46, 586
Hedgehog 134	Jamaica Ebony 162	Looking-glass plant 66	Milk-wood 462
Hedge Hyssop 396	Jambolan-tree 211	Loose-strife 425	Mint 399
Hedge Mustard 29	Japan Allspice 195	Loquat 176	Misseltoe 259
Hedge Nettle 406	Jasmine 352	Lord Anson's Pea 155	Mocker-nut 465
Hellebore 7	Jerusalem Artichoke 309	Lousewort 395	Moldavian balm 409
Hemlock 255	Jerusalem Sage 405	Lovage 249	Molucca balm 405
Hemp 463	Job's-tears 574	Love Apple 385	Moly 530
Hemp Agrimony 300	Jonquill 515	Love-grass 570	Monkey-flower 394
Hen and chicken 299	Judas-tree 172	Love lies bleeding 431	Monk's Basil 413
Henbane 382	Juniper 474	Lucerne 133	Monkshood 9, 584
Henbit 406	Jupiter's-beard 132	Lungwort 375	Moon-seed 15
Herb Paris 524	Kanguru Vine 86	Lupine 159, 589	Moonwort 576
Herb Robert 89	Kentucky Coffee-tree 368	Lychnidea 368	Moreton bay Chestnut 589
Heron's-bill 89	Kidney-bean 157	Lyme-grass 566	Morning-flower 498
Hickory-nut 465	Kidney Vetch 132, 589	Madder 266	Moschatel 242
Hog Plum 120	Knawel 221	Madwort 25	Motherwort 405
Hogweed 430	Knotted Marjoram 402	Maidenhair 582	Mountain Ash 177
Holly 111	Knot-grass 220, 438	Maidenhair-tree 473	Mountain Rose 168, 203
Hollyhock 60	Laburnum 130	Maiden Plum 119	Mountail Sorrel 439
Honesty 24	Lace-bark 448	Mahogany-tree 86	Mouse-chop 228
Honey-berry 84	Lady Fern 580	Malabar nightshade 433	Mouse-ear 271
Honey-flower 104	Ladies-mantle 192	Malabar-nut 420	Mouse-ear Chickweed 57
Honeysuckle 258	Ladies-slipper 491	Malay Apple 212	Mouse-tail 4
Honeywort 375	Ladies-smock 23	Male Fern 580	Mouse-tail grass 572
Hooded Milfoil 423	Ladies-traces 485	Mallow 59, 586	Moving-plant 150
Hop 463	Lamb's-chop 228	Mamme-tree 78	Moxa 317
Hop Hornbeam 467	Lamb's Lettuce 269	Manchineel-tree 458	Mudwort 394
Horehound 408	Lamp-wick 405	Mandrake 383	Mugwort 317
Hornbeam 467	Lancashire Asphodel 545	Mango-ginger 492	Mulberry 462
Horn Gourd 215	Lance-wood 15	Mango-tree 117	Mule Fern 581
Horn Poppy 19, 585	Larch 475	Mangosteen 78	Mullein 380
Hornwort 201	Larkspur 8	Manna Ash 351	Musk Melon 215
Horse Chestnut 83	Laserwort 252	Maple 82	Musk Okro 62
Horse Radish 26	Laurel 175	Mare's-tail 201	Musk Thistle 281
Horse-radish tree 168	Laurestine 258	Marjoram 402	Mustard 34, 585
Horse-shoe Vetch 148	Lavender 408	Marrowfat Pea 154	Myrobalan 174
Horse-thistle 282	Lavender-cotton 315	Marsh Mallow 60	Myrrh 255
Hottentot Cherry 110	Lendwort 427	Marsh Marygold 584	Myrtle 211
Hottentot Fig 230	Leather-wood 448	Marsh Pennywort 243	Mysore-thorn 169
Hound's-tongue 378	Lee Chee 84	Marsh Thistle 282	Naseberry-tree 349
Houseleek 226	Leek 530	Martagon 538	Navelwort 224
Humble plant 162	Lemon 75	Marygold 313, 592	Navew 33
Hyacinth 528	Lemon-grass 561	Marvel of Peru 429	Nectarine 173
Hyæna-poison 457	Lentil 154	Masterwort 250, 251	Nettle 462
Hyssop 402	Leopard's-bane 299	Mastick-tree 119	Nettle-tree 464
Ice-plant 232		May Apple 17	
Indian buckwheat 438		Meadow-grass 568	

ENGLISH INDEX.

Page.		Page.		Page.		Page.		Page.	
New Jersey Tea	113	Pepper	481	Redtop-grass	567	Screw-tree	65		
New Zealand cork-tree	70	Peppermint	399	Red-water tree	163	Scrubby Oak	259		
New Zealand Spruce	473	Peppermint-tree	208	Reed	574	Scurvy-grass	26		
New Zealand Tea	210	Pepper Vine	87	Reed-grass	574	Sea-buckthorn	451		
Nicker-tree	198	Perriwinkle	354	Rest-harrow	131	Sea Chickweed	57		
Nightshade	385	Petty whin	130	Rhubarb	440	Sea Daffodil	513		
Nipplewort	274	Pheasant's eye	4	Ribbon-flower	503	Sea-egg	134		
Norfolk Island Pine	475	Phillyrea	351, 352	Rib-grass	428	Sea Heath	48, 586		
Nutmeg	442	Physic-nut	457	Rice	575	Sea Holly	244		
Nut-tree	467	Picotee	49, 50	Rocambole	530	Sea Kale	35		
Oak	465	Pigeon Pea	159	Rocket	29	Sea Lavender	427		
Oat-grass	567	Pigeon Plum	173	Rock-Rose	40	Sea Matweed	478, 574		
Oats	568	Pig-nut	465	Rope-grass	547	Sea Parsnp	255		
Ogeechee-Lime	451	Pilewort	7	Rose	178	Sea Rocket	28		
Oil-nut	451	Pillwort	575	Rose Acacia	143	Seaside Balsam	457		
Oil-seed	308	Pimpnell	425	Rose Campion	54	Seaside grape	438		
Oily-grain	367	Pine Apple	534	Rosemary	404	Seaside Laurel	458		
Oily-Palm	543	Pine-tree	475	Rose of Jericho	28	Seaside Oat	571		
Old man's beard	279	Pink	49, 50, 586	Rose-root	225	Sedge	547		
Oleander	354	Pin-pillow	237	Rough Chervil	254, 591	Self-heal	410		
Oleaster	451	Pilewort	546	Rough Plum	173	Sengreen	25		
Olibanum-tree	120	Wishamia	355	Rue	105	Senna	170		
Olive	351	Pistachia-tree	118	Rupture-wort	220	Sensitive Fern	581		
Olive-bark tree	195	Pitcher-leaf	192	Rush	545	Sensitive-plant	162		
Olive-wood	110	Pitcher-plant	452	Rush-nut	551	Septfoil	190		
Omime-root	414	Plane-tree	472	Rye	564	Serpent-wood	353		
Onion	513	Plantain	428	Rye-grass	566	Service-tree	177		
Opera girls	491	Plantain-tree	495	Sacred Bean	17	Shaddock	75		
Orache	435	Plank-plant	126	Sage	410	Shallot	530		
Orange-Gourd	216	Plowman's spikenard	289	Sago-Palm	543	Sheep's-beard	279		
Orange-tree	75	Plum-tree	174	Saintfoin	151	Sheep's Scabious	328		
Orpine	220	Poet's Cassia	450	St. Andrew's cross	77	Sheep's Sorrel	440		
Osage Apple	462	Poison bulb	510, 511	St. Barnaby's thistle	287	Shepherd's club	380		
Oswego Tea	404	Poison-nut	353, 354	St. Bruno's Lily	524	Shepherd's Cress	27		
Otaheite Apple	120	Poison Oak	118	St. John's bread	169	Shepherd's purse	27		
Otaheite Chestnut	348	Poley	403	St. John's wort	76, 587	Shield Fern	579		
Otaheite Myrtle	460	Polyody	578	St. Martin's flower	516	Shore-weed	429		
Ox-eye	309	Pomegranate	195	St. Peter's-wort	258, 592	Shrubby-horsetail	472		
Ox-eye Daisy	320	Pond-weed	478	Salep	481	Shrubby Trefoil	121		
Ox-lip	424	Poplar	478	Salt-tree	144	Shumee-tree	166		
Ox-tongue	276	Poppy	18	Saltwort	434	Siberian bird's-eye	424		
Paeony	12	Portugal Laurel	175	Samphire	249	Siberian Crab	177		
Palma Christi	458	Potatoe	385	Sandal-wood	450	Side-saddle flower	18, 585		
Palms	542, 545	Primrose	423	Sandbox-tree	457	Silk cotton tree	66		
Palo de vacca	462	Prince's feather	431	Sand Myrtle	344	Silver Fir	475		
Panic-grass	563	Privet	352	Sandwort	56, 586	Silver-tree	442		
Papaw-tree	216	Puccoon	98, 585	Sanicle	244	Single-seeded Cucum- ber	215		
Paper Mulberry	462	Pumpkin	215	Sapota	349	Skirret	246		
Paper Rush	552	Purslane	219	Sappan fruit	163	Skull-cap	410		
Paraguay Tea	111	Purslane-tree	220	Sarsaparilla	522	Slipper Spurge	452		
Parsley	245	Pyracantha	175	Satin-wood	86	Slipperwort	396		
Parsley-piert	192	Quaking-grass	570	Savanna-flower	355	Sloe-tree	174		
Parsnip	251	Quamash	528	Savin	474	Small Burdock	314		
Pasque-flower	3	Quillwort	575	Savory	401	Snail-flower	157		
Passion-flower	216, 590	Quince	178	Savoy	33	Snake Cucumber	214		
Patience Dock	439	Radish	36	Savoy Spiderwort	524	Snake Gourd	215		
Pea	154	Ragged Robin	54	Sawfly Ophrys	484	Snakeroot	11, 452		
Peach	173	Ragwort	303, 305	Sawwort	281	Snake's-beard	523		
Pear	177	Rambootan	84	Saxifrage	239, 591	Snake's-tongue	577		
Pearlwort	54	Ramoone-tree	450	Scabious	269	Snake's-weed	438		
Pecan-nut	465	Rampion	326	Scallion	530	Snakewood	464		
Pellitory	463	Ranson	532	Scammony	369	Snakewort	300		
Pellitory of Spain	318	Rape	34	Scarlet-runner	157	Snap-dragon	390		
Penny-cress	27	Raspberry	188	Scorpion-grass	377	Sneezewort	319		
Pennyroyal	400	Red Cedar	474	Scorpion Senna	148	Snowball-tree	257		
Pennyroyal-tree	401	Red Quinoa	435	Scotch bird's-eye	424				
Pennywort	225	Red Sandal-wood	161	Scotch Laburnum	130				
				Screw-Pine	478				

ENGLISH INDEX.

623

Page.	Page.	Page.
Snowberry 258	Stork's-bill 92, 587	Tree Mallow 60
Snowberry-bush 264	Stramonium 383	Tree Onion 530
Snowdrop 516	Strapwort 220	Trefoil 135
Snowdrop-tree 347	Strawberry 89	Tricolor 430
Snowflake 516	Strawberry-blite 436	Trompet Marin 514
Soap-berry 84	Strawberry-tree 331	True Chervil 254
Soapwort 51	Succory 279	Trumpet-flower 365
Soft-grass 562	Sugar-cane 560	Trumpet Honeysuckle 259
Soldier-wood 163	Sulphurwort 250	Tuberose 517
Solomon's seal 523	Sumach 117	Tulip 536
Sorrel 440	Summer Cypress 434	Tulip-tree 14
Sorrel-tree 331	Sundew 45	Tupelo 451
Sour-gourd 65	Sunflower 309	Turk's-cap 235
Soursop 14	Sun-Rose 41	Turmeric 491
Southernwood 316	Supple Jack 84	Turnep 33
South-sea Myrtle 210	Swallow-wort 360	Turnep Cabbage 33
South-sea Tea 111	Sweet Bay 441	Turnsole 374
Sow-thistle 278	Sweet Briar 179	Turpentine-tree 119
Soy 158	Sweet-flag 480	Twayblade 486
Spanish Broom 129	Sweet Gale 472	Umbrella-wort 429
Spanish Cress 31	Sweet-grass 570	Valerian 268
Spanish Elm 379	Sweet-gum tree 472	Velvet-leaf 15
Spanish-nut 497	Sweet Maudlin 319	Velvet Tamarind 172
Sparrow-wort 449	Sweet Pea 155	Venus Flytrap 46
Spear-mint 399	Sweet Potatoe 371	Venus looking-glass 593
Speedwell 396	Sweet Scabious 270	Venus Navelwort 378
Spider Ophrys 484	Sweet-sop 14	Venus's comb 254
Spiderwort 541	Sweet Sultan 285	Vervain 418
Spignel 248	Sweet William 49	Vetch 153
Spinage 436	Sword Fern 578	Vine 87
Spindle-tree 109, 588	Syrian Rue 105	Violet 43, 586
Spleenwort 580	Syringa 206	Viper's Bugloss 375
Sponge-tree 166	Tacamahac 472	Viper's-grass 278
Spread-eagle 488	Tallow-tree 458	Virginian creeper 87
Spring-grass 553	Tamarind 169	Virginian Hemp 436
Spruce Fir 475	Tamarisk 203	Virginian Silk 356
Spurge 453	Tangier Pea 155	Virgin's Bower 1
Spurge Laurel 449	Tansy 316	Wall-cress 22, 585
Spurless Violet 45	Tare 154	Wallflower 21
Spurrey 55, 586	Tarragon 317	Wall Rocket 34
Squash 215	Teak-wood 418	Wall Rue 580
Squill 527	Teasel 269	Walnut-tree 465
Squirting Cucumber 215	Tea-tree 73	Wamppee-tree 75
Staff tree 110	Thistle 281	Waratah 447
Star Apple 345	Thorn Apple 383	Wart-flower 89, 587
Starch Hyacinth 528	Thrift 426	Wartwort 455
Starfish 357	Throatwort 328	Water-caltrops 200
Star-flower 117	Tick-seed 436	Water Chickweed 220
Starhead 270, 592	Tiger-chop 228	Water Cress 22
Star of Bethlehem 526	Tiger-flower 499	Water Dropwort 247
Star of the earth 429	Tile-root 503	Water Hemlock 247
Starry Scabious 285	Toad-flax 390	Water Horehound 398
Star-thistle 287	Tobacco 382	Water-leaf 380
Starwort 295	Tonquin Bean 168	Water Lily 17, 585
Sterile Strawberry 192	Toothpick 245	Water Melon 214
Stichwort 55	Toothwort 24, 422	Water Milfoil 201
Stinking Horehound 405	Touch me not 102, 588	Water Parsnep 246
Stock 21	Tower Mustard 22	Water Pepper 439
Stonecrop 225	Traveller's joy 1	Water Plantain 477
Storax 347	Treacle Mustard 25	Water Purslane 201
	Tree Celadine 19	Zebra-plant 494

N.B. Mr. SWEET having now finished several of his Botanical Works, and many of his Friends having expressed a wish, that he would Cultivate some handsome, rare, and choice Plants for Sale, finding them at present so difficult to attain, has some intention of commencing in that way; and will be obliged to any of his Friends for their Orders, which will be punctually Executed, and the Plants sent with their proper Names.

WORKS

ON

BOTANY, AGRICULTURE, & RURAL ECONOMY,

Now Publishing, or in the Press.

SUPERIOR BOTANICAL WORK.

In Monthly Numbers, with Eight Coloured Plates, Royal 8vo. price 4s.

EDWARD'S BOTANICAL REGISTER; or, Ornamental Flower Garden and Shrubbery. No. VIII. Third Volume of the New Series, for October, 1830. Continued by JOHN LINDLEY, Esq. F.R.S., Professor of Botany in the University of London, &c. &c.

Each Number contains Eight finely coloured Portraits, (to equal Drawings,) of the handsomest Plants and Shrubs grown in the Gardens and Public and Private Establishments of this Country, accompanied by their History, Mode of Treatment in Cultivation, Propagation, &c.

The thirteen Volumes completed, may be had either together or separately, price 2*l.* 9*s.* each, or by Numbers, 4*s.* each.

"What we consider the most valuable feature of this Work, and what distinguishes it peculiarly in its class, is the judicious selection of its subjects, and the constant introduction of greenhouse and hardy flowers and shrubs, in preference to those which are to be preserved only in the hot-house.—*Literary Gazette.*

"Mr. Douglas has sent home, to the Horticultural Society, more new and beautiful hardy herbaceous plants, from North America, than were ever before introduced by an individual from any country. These are beautifully figured in the Botanical Register, and are the more valuable as being fit for the open garden in every part of the Island.—*Loudon's Magazine of Botany, &c.*

"The Botanical Register, from containing most or all of the new plants introduced by the Horticultural Society, from the great care with which its plates are executed, and the judicious remarks on culture and general habit by Mr. Lindley, is, in consequence, the superior publication."—*Idem.*

"The Botanical Register has now, therefore, our unqualified approbation, and we shall lose no opportunity of saying so."—*Gardener's Magazine, No. XXVI. for June, 1830, page 279.*

BY THE FIRST AUTHORITY.

*In Royal 8vo. with Plates coloured to equal Drawings, (to be continued Monthly,) price 5*s.* No. XXXVI, for October 1, 1830.*

THE POMOLOGICAL MAGAZINE, or Figures and Descriptions of the most important Varieties of Fruit cultivated in Great Britain for the Dessert; with any Information resulting from successful Practice calculated to improve their Cultivation, pointing out the best or most proper Aspects, Situations, &c. Conducted by two Gentlemen intimately connected with the Horticultural Society of London.

"Handsomely, we may say splendidly got up, with forty-eight accurately drawn and finely coloured Plates. We cannot too highly recommend this work, as valuable to consult in the choice of fruits, and beautiful to look upon."—*Literary Gazette, Dec. 13, 1828.*

Valuable Works now Publishing by J. RIDGWAY.

" This is one of the most pleasing publications that we have met with for some time past, and is well worthy the attention of those whose delight is in the cultivation of Fruit. The various specimens of the Fruits given are coloured, and exceedingly well executed." — *Morning Herald*, Feb. 5, 1829.

Volumes I. and II., handsomely bound in Cloth, Price 3*l.* 3*s.* each.

*New Edition, Royal 8vo. Nineteen Plates, 2*s.**

ROSARUM MONOGRAPHIA; or, a Botanical History of Roses. To which is added an APPENDIX, for the Use of Cultivators, in which the most remarkable Garden Varieties are systematically arranged: with Nineteen Plates; Eighteen beautifully Coloured. By JOHN LINDLEY, Esq. F.R.S. L.S. and G.S., Professor of Botany at the University of London.

A TREATISE ON BULBOUS ROOTS; containing a Botanical Arrangement and Description of the Plants heretofore included under the Genera AMARYLLIS, CYRTANTHUS, CRINUM, and PANCRATIUM; with general Observations and Directions for their Cultivation. By the Hon. and Rev. WILLIAM HERBERT. In royal 8vo. with coloured Plates. Price 5*s.*

SWEET'S GERANIACEÆ; or beautifully coloured Figures of the finest Geraniums; with their History, and plain Directions for their Cultivation, Propagation, &c. 5 vols. royal 8vo. price 3*l.* 16*s.* each, or by Numbers, 3*s.* each.

THE FLORIST'S GUIDE; publishing in Monthly Numbers, at 3*s.* Each Number contains Four correctly and handsomely coloured Figures of the Flowers most esteemed by Florists, such as Tulips, Ranunculus', Auriculas, Hyacinths, Carnations, Pinks, Roses, &c. with the most approved methods of cultivation, directions for raising new varieties from seeds, &c. No. 40 was published on the 1st of October, 1830.

This Work is to be completed in Two Volumes, when it will comprise a Treatise for the Cultivation and Management of every Florist Flower.

THE FLORA AUSTRALASICA. By ROBERT SWEET, F.L.S. Published in One handsome royal octavo Volume, containing fifty-six elegantly coloured Figures, of the most beautiful and curious Plants, natives of New Holland, and the South Sea Islands, (generally called Botany Bay Plants;) they are well adapted for a Greenhouse or Conservatory, and many will survive the Winters in the open ground, with a very little protection. Price £2. 2*s.* 6*d.* in Fourteen Numbers, Index, &c.; or Single Numbers, at 3*s.* each.

SWEET's CISTINEÆ: An Account of the Family of CISTUS, or Rock-Rose, illustrated by handsomely coloured Figures, with the best Method of Cultivation and Propagation. The greater part of the species succeed well in the open borders, or in rock-work, and produce their showy flowers nearly all the Summer; the others may be protected through the Winter in a garden frame, or in the Greenhouse. Completed in one handsome Volume, royal octavo, price £4. 8*s.* bound in cloth, or, if preferred, the Work may be had in Numbers, 3*s.* each.

Valuable Works now Publishing by J. RIDGWAY.

Sweet's Catalogue of Plants, Second Edition, with the Colour of the Flowers, &c. Price One Guinea.

SWEET'S HORTUS BRITANNICUS; or, a Catalogue of all the Plants grown in the Gardens or found wild in Great Britain, arranged according to the Natural Orders to which they belong, with the addition of the Linnean Class and Order to each Genus; their Systematic and English Names, when first cultivated, the Colour of the Flowers, Accentuations, whether Trees, Shrubs, Evergreens, Climbers, Perennials, Bulbs, Water-plants, Biennials, or Annuals, references to Figures, and numerous Synonyms, with many other improvements. The whole brought down to the present time, and contains many hundreds of Plants not yet enumerated in any Catalogue of this country.

* * * The rapid Sale of the last Edition proves the preference of the Arrangement in Natural Orders to that of the Linnean; and, in the present, the Author has spared neither labour nor expense to render it worthy the pre-eminence with which that was honoured.

In Monthly Royal 8vo. Numbers, price 3s.

The BRITISH FLOWER GARDEN and Ornamental Shrubbery.
By ROBERT SWEET, F.L.S. No. XVII. for October 1. New Series.

Each Number contains four full and highly coloured Figures, with Dissections of the more essential parts, of the most beautiful, rare, curious, or interesting Herbaceous Plants or Flowering Shrubs, that are cultivated in the open Borders, Shrubberies, &c. in the Gardens of this Country; with a full Account of the best and most successful Mode of Management in Culture and Propagation, the Soils and Situations most suitable for them. Twenty-five Numbers will form each Volume, and three Volumes will complete the Series. The First Series, in Three Volumes, may be had, price 3*l.* 16*s.* each.

"The British Flower Garden ought to be in the hands of all Cultivators of hardy, flowering Plants."—*Loudon's Gardener's Mag.*

In the Press, Fifth Edition, nearly ready.

SWEET'S HOTHOUSE and GREENHOUSE MANUAL;
or, BOTANICAL CULTIVATOR. This Volume contains a full account of the best method of Cultivation and Propagation of all Plants grown in the Hothouses, Greenhouses, Frames, or in the open ground of the Gardens, or Plantations in this Country; with a full account of the Management of Bulbs, and the Parasitical and other Orchideous Plants. Price 12. boards.

"Indeed, what Mr. Sweet has said on the culture of Bulbs and Epiphates in the last edition of his Botanical Cultivator, may be considered as the *ultimatum* on this subject for the British Gardener."—*Loudon's Gardener's Magazine.*

IRIDEARUM GENERA, cum Ordinis Charactere Naturali, Specierum Enumeratione, Synonymisque. Auctore JOHN BELDEN KER, Esq. price 6*s.* 6*d.*

THE GARDENER'S REMEMBRANCER, & APIARIAN'S MONTHLY CALENDAR; The Gardener's Vegetable Seed Calendar, and Fruit Calendar; beautifully printed in Typolithography, and ornamented, on imperial drawing paper. It shews, at one view, every thing necessary to be done in Rural Affairs, to ensure a pro-

per succession of Crops, &c. with proper directions for the Management and Propagation of Bees, &c. &c. Price 5s. 6d.

"The work we have at present in hand contains, on one sheet, every direction necessary to the practical gardener, and will be found invaluable to lady florists." *Athenaeum.*

CUSTOMS AND HUSBANDRY OF COUNTIES.

In Two Volumes, with Plates, price 15s. each.

THE PRESENT STATE OF THE TENANCY OF LAND IN GREAT BRITAIN, showing the principal Customs and Practices between Incoming and Outgoing Tenants, and the most usual method under which Land is now held in the several Counties; with Notices of the Husbandry and Implements in use; from an actual Survey in 1827, 1828, and 1829. By LEWIS KENNEDY, and T. B. GRAINGER.

** The Second Volume comprises, the **HIGHLAND AND GRAZING DISTRICTS**. Also, Facts relative to the present state of the Woolgrowers, the general Mode of Management of Sheep Farms, with a Brief History of Sheep in Great Britain. And may be had separately.

"Interesting, not only to the Land-owner and Farmer, but, where such matters are under legal investigation, by suggesting important questions, it is calculated greatly to promote an expeditious and equitable decision on the subjects under consideration."—*Times.*

SHEEP HUSBANDRY.

EARL STANHOPE on **SHEEP HUSBANDRY**; being a Letter to the Owners and Occupiers of Sheep Farms; with the Evidence taken before a Select Committee of the House of Lords, appointed to take into consideration the State of the British Wool Trade; classed under different Heads. Price 4s.

PRACTICAL OBSERVATIONS on the National Advantages of the Arable System of **SHEEP HUSBANDRY**, and on the Improvement of **BRITISH FINE WOOL**, without deteriorating the Value of the Carcass. By JOSHUA KIRBY TRIMMER. Price 3s.

OBSERVATIONS on the Disease which has lately been so destructive to Sheep, called Bane, or Coath, particularising the Causes, and minutely describing the Modes of effecting its Cure; and pointing out those means which ought to be adopted to prevent its recurrence. By JOHN DAVEY. Read at the Annual Meeting of the Bath and West of England Society, and ordered to be printed. 2s. 6d.

DAIRY HUSBANDRY.

THE COMPLETE HISTORY of **DAIRY HUSBANDRY**; and particularly with reference to the System pursued for many years at the celebrated Establishment at Willow Bank, near Glasgow; including Improved Modes for Ventilating Stables, Reclaiming Waste Lands, Management of Hedge Row Fences, and other Information highly important to the Farmer and Rural Economist. By WILLIAM HARLEY. One Volume, 8vo. Copper-plates. 21s. bound in Cloth.

Valuable Works now Publishing by J. RIDGWAY.

Second Edition, Price 3s. 6d.

IMPROVED SHORT HORNS, and their Pretensions stated ; being an Account of this celebrated Breed of Cattle, derived from Authentic Sources. To which is added, An Enquiry as to their Value for general purposes, placed in competition with the improved Herefords. By the Rev. HENRY BERRY. Dedicated to Sir CHARLES MORGAN, Bart. M. P.

MANGEL WURTZEL, *Price 6d.*

REPORT of the COMMITTEE of the DONCASTER AGRICULTURAL ASSOCIATION, on the Advantages of **MANGEL WURTZEL**, as a Fallow Crop; founded on Returns received in answer to Queries issued by the Committee.

BONE MANURE.

A REPORT of the COMMITTEE of the DONCASTER AGRICULTURAL ASSOCIATION on the Advantages of Bones as a Manure, founded on Returns received in answer to certain Queries issued by the Committee.

Returns were received from forty-nine practical Farmers of every variety of Soil in the County.

" This Report is so important, is so ably drawn up, and embodies such a compendium of practical information, that we feel it our duty to the Farming Interest of Britain to recommend its perusal."—*Farmers' Journal.*

IMPROVEMENT OF MEADOW AND PASTURE LANDS.

HORTUS GRAMINEUS WOBURNENSIS; or, an Account of the Results of various Experiments on the Produce and Fattening Properties of different Grasses, and other Plants, used as the Food of the more valuable domestic Animals; carried on for a number of years at Woburn Abbey, by command of the Duke of Bedford, and, being now matured, are presented for the benefit of the agricultural community. Illustrated with numerous Specimens of the Plants and Seeds, in Sixty Plates, with Practical Observations on their natural habits, the Soils best suited to their Growth, with Instructions for the Preservation of their Seed; pointing out the kinds most profitable for Permanent Pasture, Irrigated Meadows, Dry or Upland Pasture, and the Alternate Husbandry. Accompanied with the discriminating Character of the Species and Varieties. By G. SINCLAIR, F.L.S. and F.H.S., Gardener to his Grace the Duke of Bedford, &c. &c. Third Edition. Price £1. 10s. plain, and coloured £2. 2s. under the Author's immediate inspection.

" The Hortus Gramineus Woburnensis does honour to our country." ***—*Monthly Critical Gazette.*

Second Edition, in 8vo. with Engravings, 9s. boards,

THE BRITISH FARMER; to which is added, **THE PLOUGH-MAN'S GUIDE**. By JOHN FINLAYSON, Esq.

" One of the most valuable Tracts on Agriculture that has appeared for a number of years.—*Loudon's Gardeners' Magazine.*

Valuable Works now Publishing by J. RIDGWAY.

AN ESSAY ON THE WEEDS OF AGRICULTURE; with Practical Remarks on their Destruction. By the late B. HOLDITCH, Esq. and G. SINCLAIR, F.L.S. F.H.S. Author of *Hortus Gramineus Woburnensis*. Second edition. 3s. 6d.

"If we were at liberty to follow our inclinations, we should treat this pamphlet, not according to its bulk, but its importance, and follow Mr. Holditch, and his editor, (for it is only the 1st chap. that is strictly, in its present form at least, attributable to the former,) through their respective chapters on "Weeds which infest Samples of Corn; Fallow Weeds; Weeds which are principally objectionable, as they encumber the Soil, or whose Roots are annual, and whose Seeds pass the corn-sieve; Weeds that never rise in the crop, nor come into the sickle; Pasture Weeds, &c. &c." * * * The passages we have marked in our progress, as worthy of quotation, might form a valuable little Manual for the Practical Farmer."—*Monthly Magazine*.

A SYSTEM FOR MANAGING HEAVY AND WET LANDS WITHOUT SUMMER FALLOWS; under which a considerable Farm in Hertfordshire is kept perfectly clean, and made productive. By THOMAS GREG, Esq. Third Edition, with an Appendix, pointing out how the infant Turnip may be protected from Insects. To which are added, Extracts from some Letters written by the Earl of Thanet, proving the infallibility of the Plan; as presented to the Board of Agriculture, and now published at their request. Price 5s. with Plates.

ELLIOT'S FARMER'S ACCOUNT BOOK for 1831; being the result of many years extensive practice. By Double Entry in one Book, the arrangements are clear and concise; printed and ruled ready for filling up, under the heads of Day Labour, Task Labour, Bills, Debts owing and owed, Live Stock, Green Crops, Corn, Hay, and Straw, Rent, Taxes, Cash, &c. Illustrated by a Michaelmas Valuation, and a Weekly Account, filled up for examples, &c. Price 1l. 1s. for one year, and 10s. 6d. for a half year.

"No Farmer desirous of knowing the exact state of his expences, ought to be without this Work."

PONTEY'S PROFITABLE PLANTER; a Practical Treatise on Planting Forest Trees in every description of Soil and Situation. 1 vol. 8vo. plates, 10s. 6d. boards. Fourth edition.

"Pontey, one of the most extensive and successful Planters now living, and also a Landscape Gardener of no small distinction." — *Vide Sir Henry Steuart, Bart. on Planting*, page 62.

PONTEY'S FOREST PRUNER, or Timber Owner's Assistant. 1 vol. 8vo. plates, 12s. Fourth edition.

PONTEY'S RURAL IMPROVER; a Practical Treatise on the Management of such Rural Scenes and Objects as are necessary to promote the Comfort, Convenience, and Embellishment of the Residences of the higher Ranks of Society. With plates, 1 vol. 4to. 2l. 2s.

* * * The above three Works are by W. PONTEY, Planter and Forest Pruner to the Duke of Bedford.

THE FRUIT GROWER'S INSTRUCTOR; or a Practical Treatise on Fruit Trees, from the Nursery to Maturity; to which is added an effectual Remedy for, and Prevention from the American Blight, which causes the Canker in Apple Trees. By G. BLISS, Nurseryman. 6s.

" This is a Practical Treatise on the Culture and Preservation of Fruit Trees in general, and more particularly the Apple Tree, by one of the most extensive growers in the kingdom. The best modes of Grafting, Pruning, and Budding, are laid down with the utmost clearness. A very ample Catalogue of our Fruit Trees, and a description of the appearance, flavour, and qualities of their production, is also given." * * * * " But the most valuable part of this volume is that which relates to the Canker, and the remedy which Mr. Bliss has discovered for this destructive blight." * * * — *Monthly Critical Gazette.*

THE BRITISH FARMER'S (QUARTERLY) MAGAZINE, No. XVI. published on August 1, 1830, contains—

Branch I. Original Communications: Portrait, Short-horned Bull.—On the Improvement of Waste Land, by Mr. Aiton, (concluded.)—On the Advantages of Shallow Sowing, (concluded.)—Remarks on Ditto.—Mr. Gray's Letter to the late Lord Castlereagh, on the Currency, (concluded.)—Remarks on Mr. Taylor's Letter on the Wheat Fly, by Delta.—Mr. Gray's further Observations on "Observer."—Mr. Berry, on the Improved Breeds of Cattle, in Reply to Mr. Hayward.—On the Anglesea Breed of Cattle, by Mr. Berry.—On the Choice of a Farm.—Mr. Gray's Address to Electors of Great Britain and Ireland.—On Tithes.

Review of Agricultural Publications: Billington's Facts and Observations on Forest Pruning.—Colonization.—Irish Cottagers, by Mr. Doyle.—Webster's Dictionary.—The British Co-operator.—Oracle of Health.

Branch II. Agricultural Intelligence: Miscellaneous Extracts and Observations.—Agricultural Quarterly Reports: For England.—For Scotland.—Corn Trade,—Smithfield, Mrak Lane.—Wool Trade.—Provision Trade.—Hop Trade.—State of Ireland.—Hamburg Prices of Grain.—Principal Fairs.

Branch III. Sporting Intelligence.

No. XVII. will be published on November 1st.

The Three Volumes of this Work, (comprising Nos. I. to XIII.) illustrated by Eleven finely engraved Portraits of celebrated Cattle, Ground Plan and Sections of a New Barn and Offices, on improved and tried principles; may be had, price 2l. 18s., neatly half bound, in calf; or in Numbers, at 4s. each.

A PRACTICAL TREATISE on BREEDING, REARING, and FATTENING DOMESTIC POULTRY, &c. After Plans pursued with advantage and profit in France. Second edition. 5s.

A SERIES of FACTS, HINTS, OBSERVATIONS, and EXPERIMENTS, on the different Modes of raising YOUNG PLANTATIONS of OAK, " For future Navies," from the Acorn, Seedling, and larger Plants; shewing the Difficulties and Objections that have occurred in the Practical Part; with Remarks upon the Fencing, Draining, Pruning, and Training Young Trees; and Experimental Remarks upon Fruit Trees. By WILLIAM BILLINGTON, C.H.S. Superintendent of His Majesty's Land in the Forest of Dean, in the County of Durham, &c. &c. 12s.

SAMUEL CHILD'S, *EVERY MAN HIS OWN BREWER.*

A PRACTICAL TREATISE, explaining the Art and Mystery of Brewing PORTER, ALE, TWOPENNY, and TABLE BEER; intended to reduce the Expenses of Families. By SAMUEL CHILD, Common Brewer. A new Edition, 1s.

* * Upwards of *Eleven Thousand* of this little Tract have been sold.

1832

Valuable Works now Publishing by J. RIDGWAY.

THE ENGLISH RACE HORSE.

A TREATISE ON THE CARE, TREATMENT, AND TRAINING, OF THE ENGLISH RACE HORSE; with important Details applicable to bettering the Condition of Horses in general. By R. DARVILL, V. S. to the Seventh Hussars. Illustrated by Plates, 1 vol. 8vo. 1*l.* 1*s.*

"Never before was such a book written in any language, so replete with those minute but indispensable particulars of practice, and by a writer who has personally performed his part throughout the whole of the practice. This is the true book of reference for every stud and training groom, and every jockey.—*Vide Lawrence on the Horse*, p. 297; also the *Sporting Magazine and British Farmer's Magazine*.

THE VETERINARY TABLET; or a Concise View of all the Diseases of the Horse; with their Cause, Symptoms, and most approved and effectual Modes of Cure. By a Veterinary Surgeon. 1*s. 6d.*

THE YOUNG HORSEWOMAN'S COMPENDIUM OF THE MODERN ART OF RIDING. In progressive Lessons; designed to give a secure and graceful Seat on Horseback; at the same time so effectually to form the Hand, that, in a short time, perfect command of the Horse may be obtained. By EDWARD STANLEY, Royal Artillery, and late of Vernon's Establishment, near Hyde Park; with illustrative Plates, 10*s.* boards.

"But we have said enough of this Manual, and have only to add, that it is a very sensible and judicious Publication."—*Literary Gazette*.

Preparing for Publication, by Subscription, and to be ready early in the Spring of 1831, dedicated, by express permission, to the Most Noble the Marquess of Bute,

OBSERVATIONS on the present Defective State of ENGLISH TIMBER; the Causes which retard its Growth, together with the means which ought to be immediately adopted to prevent the further premature decay of this valuable description of property, of which its owners in general seem not aware. By P. LAUDER, of Cardiff, Member of the Bristol, Glamorgan, and Monmouth Horticultural Societies, &c. &c.

The Work will be in octavo, with numerous illustrative Plates; the price One Guinea, to be paid on delivery of the Work.

Nearly Two Hundred Names have already been received; and amongst them some of the principal Members of the Peerage and Baronetage. A List of Subscribers will accompany the Work.

Also, to be ready early in the Spring of 1831,

The FOSSIL FLORA of GREAT BRITAIN; or, Figures and Descriptions of the Vegetable Remains found in a Fossil state in this Country. By JOHN LINDLEY, Esq. F.R.S. L.S. & G.S.; and WILLIAM HUTTON, Esq., F.G.S., &c. To appear in Quarterly Parts. Royal 8vo. Price 6*s.* Each Part will contain Ten Copper-plates, and about forty pages of Letter-press.

As a correct list of Subscribers' names is purposed being printed in the Work, Noblemen, Gentlemen, and Ladies, are requested to forward them, written as they would wish them to appear, addressed to the Publishers, Messrs. RIDGWAY, Piccadilly.













